ANCHORED FOAMS AND ANNULAR HOMOLOGY

ROSTISLAV AKHMECHET AND MIKHAIL KHOVANOV

ABSTRACT. We describe equivariant SL(2) and SL(3) homology for links in the solid torus via foam evaluation. The solid torus is replaced by 3-space with a distinguished line in it. Generators of state spaces for annular webs are represented by foams with boundary that may intersect the distinguished line; intersection points, called anchor points, contribute additional terms, reminiscent of square roots of the Hessian, to the foam evaluation. Both oriented and unoriented SL(3) foams are treated in the paper.

Contents

| 1. Introduction | 1 |
|--|----|
| 2. $SL(2)$ anchored homology | 3 |
| 2.1. Anchored surfaces and their evaluations | 3 |
| 2.2. State spaces | 10 |
| 2.3. Annular link homology | 15 |
| 3. Unoriented $SL(3)$ anchored homology | 19 |
| 3.1. Unoriented anchored $SL(3)$ foams and their evaluations | 20 |
| 3.2. Skein relations | 25 |
| 3.3. State spaces | 27 |
| 3.4. Remark on Lee's theory | 31 |
| 3.5. Unlabeled anchor points and bigon decomposition | 33 |
| 4. Oriented $SL(3)$ anchored homology | 37 |
| 4.1. Oriented $SL(3)$ foams and their evaluations | 38 |
| 4.2. Oriented anchored $SL(3)$ foams and their evaluations | 43 |
| 4.3. Skein relations | 47 |
| 4.4. State spaces | 50 |
| 4.5. Annular $SL(3)$ -link homology | 54 |
| References | 55 |

1. INTRODUCTION

Asaeda-Przytycki-Sikora [APS04] homology of links in the solid torus has led to a number of interesting developments [Rob13b, GN14, BG15, GLW18, GLW17, BPW19, Akh20] and extensions of their work to SL(N) and GL(N) link homology in the solid torus [QR18, QW21, QRS18].

GL(N) and SL(N) link homology theories are closely related to foam evaluation. This connection was made the most transparent by the work of Robert and Wagner [RW20], who wrote down a combinatorial formula for GL(N) closed foam evaluation that allows to

Date: May 3, 2021.

build GL(N) link homology from the ground up, bypassing categorical approaches to the latter. A variation of their formula was used to evaluate unoriented SL(3) foams [KR21], giving a combinatorial approach to some of the structures discovered by Kronheimer and Mrowka [KM19b].

In this paper we extend foam evaluation framework to build equivariant SL(2) and SL(3) state spaces for annular webs and, consequently, equivariant SL(2) and SL(3) homology for links in the solid torus. Our construction complements earlier work [QR18, QW21] on the subject. The same approach allows to define state spaces for unoriented SL(3) annular webs, extending the construction in [KR21].

In the APS (Asaeda-Przytycki-Sikora) annular homology and its equivariant and SL(N) generalizations, one first defines state spaces for annular SL(2) and SL(N) webs, where annular SL(2) webs are just collections of embedded circles in an annulus.

Our idea is to think of an open solid torus as the complement to a line L in \mathbb{R}^3 , chosen for convenience to be the z-axis. An annular SL(N) web Γ is then placed into the xy-plane with (0,0) removed. To define its state space $\langle \Gamma \rangle$, we consider SL(N) foams F in the half-space \mathbb{R}^3_- bounded by the xy-plane such that Γ is the boundary of F. These foams may intersect the z-axis, and we refer to the intersection points as anchor points and to such foams as anchored foams. Anchor points additionally carry a label from 1 to N, and we modify foam evaluation by adding a new type of factors associated to anchor points.

In this paper we treat N = 2 and N = 3 cases, with modified evaluations given by formulas (2) and (88), respectively, also see (39) for the unoriented SL(3) anchored foam evaluation.

Anchored foam evaluation take values in the ring of polynomials rather than the ring of symmetric polynomials. One starts with an admissible coloring c of facets of a foam F, as usual. An anchor point labeled i lying on a facet of color j contributes $\delta_{i,j}\sqrt{\pm f'(x_i)}$ to the evaluation $\langle F, c \rangle$, where, in the SL(3) case as an example, $f(x) = (x - x_1)(x - x_2)(x - x_3)$ is the polynomial of degree three with roots x_1, x_2, x_3 . The full evaluation $\langle F \rangle$ is given by summing over $\langle F, c \rangle$ for all admissible colorings c. We check integrality property of these evaluations, with $\langle F \rangle$ a polynomial in x_1, x_2, x_3 , in the SL(3) case.

Given evaluations of anchored closed foams, one can form state spaces for annular webs. We show that this modified evaluation, with anchor points contributing $\delta_{i,j}\sqrt{\pm f'(x_i)}$, perfectly matches the structure of state spaces of annular homology, in SL(2) and SL(3) cases. The construction also allows us to define unoriented SL(3) homology for annular trivalent graphs, extending [KR21] to the annular framework.

With state spaces at hand, it is straightforward to define annular SL(2) and SL(3) link homology, by analogy with [Kho00, APS04, BN05, Akh20] in the SL(2) setting, with [KR21] in the unoriented SL(3) setting, and with [Kho04, MV07, RW20] in the oriented SL(3) setting. State spaces and link homology carry additional gradings coming from intersection points of foams with the z-axis. We show that the result matches equivariant SL(2) homology [Akh20] of the first author. A simple modification of the construction (truncating the ground ring by sending x_i 's to 0 upon evaluation) gives a foam approach to the original APS homology. We expect that the non-equivariant variant of our SL(3) construction recovers N = 3 case of the homology in [QR18]. It seems that the equivariant annular SL(3) homology, as described in the present paper, is new.

Section 2 describes SL(2) homology via anchored foams. The evaluation is defined in Section 2.1, which also contains the skein relations for anchored SL(2) foams. The state spaces are studied in Section 2.2. The state space of n circles in the annulus is a free module of rank 2^n over the ground ring R_{α} of polynomials in two variables, see Theorem 2.11. The numbers of contractible and essential circles control the bigraded rank. This section also discusses categories of *anchored* and *annular* cobordisms. Annular cobordisms between annular SL(2) webs are disjoint from the z-axis, while anchored cobordism may intersect it.

Theorem 2.19 identifies the annular cobordism functor with that constructed in [Akh20]. Consequently, equivariant annular SL(2) link homology [Akh20] can be rederived via anchored foams. To obtain the original APS homology, one can use anchored foam evaluation, combined with the homomorphism $R_{\alpha} \longrightarrow \mathbb{Z}$ taking α_1, α_2 to 0 to get state spaces and cobordism maps in the APS theory.

Section 3 constructs the state spaces for the annular unoriented SL(3) foam theory, extending the construction of [KR21]. We start with the evaluation (Section 3.1), followed by skein relations on annular foams (Section 3.2) and properties of state spaces (Section 3.3). Section 3.4 describes similarities between anchor points contributions and Lee's theory, given by inverting the discriminant in the ground ring. Similar to the planar case [KR21], we don't know a way to describe the state space of an annular web when regions of valency at most four, allowing an inductive simplification, are absent.

In Section 4 we describe annular equivariant SL(3) link homology, based on anchored (annular) oriented SL(3) foams. This homology extends Mackaay-Vaz [MV07] equivariant SL(3) homology of links in \mathbb{R}^3 , also see [Kho04, MN08, Cla09, Rob13a] for the non-equivariant homology in \mathbb{R}^3 . We start with a review of oriented SL(3) foams in Section 4.1 and then follow a similar route to that of the earlier sections.

We expect that our construction admits a generalization to SL(N) homology for all N via an extension of the Robert-Wagner formula [RW20] to the anchored case.

Acknowledgments: M.K. was partially supported by NSF grant DMS-1807425 while working on the paper. R.A. was supported by the Jefferson Scholars Foundation. R.A. would like to thank his advisor Slava Krushkal for encouraging him to pursue this project.

2. SL(2) Anchored homology

2.1. Anchored surfaces and their evaluations. Consider the integral polynomial ring $R_{\alpha} = \mathbb{Z}[\alpha_1, \alpha_2]$ in two variables α_1, α_2 . Define a grading on R_{α} by setting

(1)
$$\deg(\alpha_1) = \deg(\alpha_2) = 2.$$

Denote by τ the nontrivial involution of $\{1, 2\}$. It is given by $\tau(i) = 3 - i$, for $i \in \{1, 2\}$. Also denote by τ the induced involution of R_{α} which permutes α_1, α_2 , so that $\tau(\alpha_i) = \alpha_{\tau(i)} = \alpha_{3-i}$. Let R be the τ -invariant subring of R_{α} , which consists of symmetric polynomials in α_1, α_2 . The subring R is itself a polynomial ring, $R = \mathbb{Z}[E_1, E_2]$, where E_1, E_2 are elementary symmetric polynomials in α_1, α_2 ,

$$E_1 = \alpha_1 + \alpha_2, \quad E_2 = \alpha_1 \alpha_2.$$

Degrees of E_1 and E_2 are 2 and 4, respectively.

Let $L \subset \mathbb{R}^3$ denote the z-axis, $L = (0,0) \times \mathbb{R}$. Let $S \subset \mathbb{R}^3$ be a closed, smoothly embedded surface which intersects L transversely. The surface S may be decorated by dots, disjoint from L, that can otherwise float freely on components of S. The intersection points $S \cap L$ are called *anchor points*. Fix a labeling ℓ , which is a map from the set of anchor points to $\{1, 2\}$,

$$\ell \colon S \cap L \to \{1, 2\}.$$

Order the anchor points by $1, \ldots, 2k$, read from bottom to top, so that the labeling ℓ consists of a choice $\ell(j) \in \{1, 2\}$ for each $1 \leq j \leq 2k$. We will define an evaluation

$$\langle S \rangle \in R_{\alpha}$$

for S with the fixed labeling ℓ , which is omitted from the notation.

Let $\operatorname{Comp}(S)$ denote the set of connected components of S. A coloring of S is a function $c: \operatorname{Comp}(S) \to \{1, 2\}$, and we denote by $\operatorname{adm}(S)$ the set of colorings of S. Surface S has $2^{|\operatorname{Comp}(S)|}$ colorings. Fix a coloring c. For i = 1, 2, let $d_i(c)$ denote the number of dots on components colored i. Let S_2 denote the union of the 2-colored components. For $1 \leq j \leq 2k$, let c(j) denote the color of the j-th anchor point, induced by c, which may in general be different from the fixed label $\ell(j)$. Define

(2)
$$\langle S, c \rangle = (-1)^{\chi(S_2)/2} \frac{\alpha_1^{d_1(c)} \alpha_2^{d_2(c)} \left(\prod_{j=1}^{2k} (\alpha_{c(j)} - \alpha_{\ell(j)})\right)^{1/2}}{(\alpha_1 - \alpha_2)^{\chi(S)/2}}$$

Note that $\chi(S_2)$ is even since S_2 is a closed surface in \mathbb{R}^3 . Let us explain the square root in the above equation.

Each component S' of S intersects L at an even number of points p_1, \ldots, p_{2m} , which can be ordered as encountered along L, from bottom to top. Suppose S' is colored by c(S') = j, and moreover S' contains an anchor point labeled j. Then the product $\prod_{j=1}^{2m} (\alpha_{c(j)} - \alpha_{\ell(j)}) = 0$, since it contains a term $\alpha_j - \alpha_j = 0$, and the entire evaluation $\langle S, c \rangle = 0$. Thus, the evaluation (2) is only nonzero when the anchor points on a component S' colored j are all labeled by the complementary color $\tau(j)$. In this case, each component contributes an even number of factors of either $\alpha_1 - \alpha_2$ or $\alpha_2 - \alpha_1$ to the product $\prod_{j=1}^{2m} (\alpha_{c(j)} - \alpha_{\ell(j)})$, and we define the square root to be $(\alpha_1 - \alpha_2)^m$ or $(\alpha_2 - \alpha_1)^m$, respectively. If S' has no anchor points, this term is 1 and can be removed from the product.

Note that the evaluation is the product of evaluations of individual components,

(3)
$$\langle S, c \rangle = \prod_{S' \in \text{Comp}(S)} \langle S', c(S') \rangle.$$

Thus, if S' is colored 1 by c' = c(S'), has 2k anchor points all labeled 2 and carries d dots, then

(4)
$$\langle S', c' \rangle = \alpha_1^d (\alpha_1 - \alpha_2)^{k - \chi(S')/2}$$

If S' is colored 2 by c' = c(S'), has 2k anchor points all labeled 1 and carries d dots, then

(5)
$$\langle S', c' \rangle = (-1)^{\chi(S')/2+k} \alpha_2^d (\alpha_1 - \alpha_2)^{k-\chi(S')/2} = \alpha_2^d (\alpha_2 - \alpha_1)^{k-\chi(S')/2}$$

Otherwise, if one of the anchor points has the same label as the color of S', the evaluation $\langle S', c' \rangle = 0$ and $\langle S, c \rangle = 0$.

Define the evaluation of S by

(6)
$$\langle S \rangle = \sum_{c} \langle S, c \rangle$$
,

where the sum is over all colorings of S. Note that if $S \cap L = \emptyset$, then $\langle S \rangle$ agrees with the evaluation in [RW20, KR20b]. Also note that $\langle S \rangle = 0$ if a component of S has two anchor points with different labels 1, 2.

We have

(7)
$$\langle S \rangle = \prod_{S' \in \operatorname{Comp} S} \langle S' \rangle,$$

that is, evaluation of S is the product of evaluations over connected components of S.

We can rewrite $\langle S \rangle$ as follows. First, suppose S is connected, carrying d dots, with $2k \ge 0$ anchor points. For i = 1, 2, let c_i denote the coloring of S by i. Define

(8)
$$\langle S, c_1 \rangle = \frac{\alpha_1^d ((\alpha_1 - \alpha_{\ell(1)}) \cdots (\alpha_1 - \alpha_{\ell(2k)}))^{1/2}}{(\alpha_1 - \alpha_2)^{\chi(S)/2}},$$

(9)
$$\langle S, c_2 \rangle = (-1)^{\chi(S)/2} \frac{\alpha_2^d ((\alpha_2 - \alpha_{\ell(1)}) \cdots (\alpha_2 - \alpha_{\ell(2k)}))^{1/2}}{(\alpha_1 - \alpha_2)^{\chi(S)/2}}$$

Again, square roots in the above equations are taken in the natural way. If S has oppositely labeled anchor points then both (8) and (9) are zero. If all anchor points are labeled 1, then (8) is zero, whereas (9) is equal to

$$\langle S, c_2 \rangle = (-1)^{\chi(S)/2} \frac{\alpha_2^d (\alpha_2 - \alpha_1)^k}{(\alpha_1 - \alpha_2)^{\chi(S)/2}}$$

On the other hand, if all anchor points are labeled by 2 then (9) is zero and (8) equals

$$\frac{\alpha_1^d (\alpha_1 - \alpha_2)^k}{(\alpha_1 - \alpha_2)^{\chi(S)/2}}.$$

Then for connected S with anchor points, we have

$$\langle S \rangle = \langle S, c_1 \rangle + \langle S, c_2 \rangle,$$

where at most one of the summands $\langle S, c_i \rangle$ is nonzero.

Clearly the evaluation is multiplicative under disjoint union. That is, if $S = S_1 \sqcup \cdots \sqcup S_n$, then

$$\langle S \rangle = \langle S_1 \rangle \cdots \langle S_n \rangle.$$

Remark 2.1. Unlike closed foam evaluations appearing elsewhere [RW20, KR21, KK20, KR20b, RW16], our evaluation does not in general produce a symmetric function. The following examples illustrate this.

Example 2.2. Let S be a sphere intersecting L in two points with labels i and j and carrying d dots. If $i \neq j$, then each coloring c yields $\langle S, c \rangle = 0$. If both anchor points are labeled 1, then only coloring S by 2 contributes to the sum, and we have

$$\langle S \rangle = \langle S, c_2 \rangle = -\frac{\alpha_2^d(\alpha_2 - \alpha_1)}{\alpha_1 - \alpha_2} = \alpha_2^d.$$

On the other hand, if both anchor points are labeled 2, then

$$\langle S \rangle = \langle S, c_1 \rangle = \alpha_1^d.$$

This is summarized pictorially in (10). Both signs are positive since $k + \chi(\mathbb{S}^2)/2 = 1 + 1 = 2$ is even.



Note that these evaluations are not symmetric in α_1, α_2 .

Example 2.3. More generally, let S be a genus g surface with d dots and 2k anchor points. If k = 0 (that is, if S is disjoint from L) then the evaluation is

$$\langle S \rangle = \frac{\alpha_1^d + (-1)^{g-1} \alpha_2^d}{(\alpha_1 - \alpha_2)^{1-g}}.$$

On the other hand, if k > 0, then

(11)
$$\langle S \rangle = \begin{cases} \alpha_2^d (\alpha_2 - \alpha_1)^{k+g-1} & \text{if } \ell(1) = \dots = \ell(2k) = 1, \\ \alpha_1^d (\alpha_1 - \alpha_2)^{k+g-1} & \text{if } \ell(1) = \dots = \ell(2k) = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 2.4. For any anchored surface $S \subset \mathbb{R}^3$ with d dots and 2k anchor points, its evaluation $\langle S \rangle$ is a homogeneous polynomial in α_1 and α_2 of degree $-\chi(S) + 2d + 2k$.

Proof. If S does not intersect L, then this follows from Example 2.3. Suppose that S intersects L. It suffices to verify the statement for connected surfaces. If S intersects L, then the statement follows from (11), since k > 0.

We recall the following notation from [KR20b]. For i = 1, 2, we allow surfaces to carry decorations (i) consisting of i inscribed in a small circle. They must be disjoint from L and are allowed to float freely along the connected component on which they appear. We call these *shifted* dots. Diagrammatically, a shifted dot (i) is the difference between a dot and α_i ,

(12)
$$(i) = -\alpha_i$$

Lemma 2.5. Let S be an anchored foam and let $S \cup (i)$ denote the anchored foam obtained by placing a shifted dot (i) on some component S' of S. Then

$$\left\langle S \cup (i) \right\rangle = \begin{cases} 0 & \text{if } S' \text{ has an anchor point labeled } \tau(i) \\ (-1)^i (\alpha_1 - \alpha_2) \left\langle S \right\rangle & \text{if all anchor points on } S' \text{ are labeled } i. \end{cases}$$

Proof. This is clear from the definitions.

Lemma 2.5 is summarized diagrammatically in the relations (13).



Alternatively, the skein relations (13) may written compactly as in (14).

(14)
$$\bullet *^{i} = \tau(\alpha_{i}) *^{i}$$

Lemma 2.6. The local relations (15), (16), and (17) hold.

(15)
$$\boxed{\bullet \bullet} = E_1 \boxed{\bullet} - E_2 \boxed{}$$

(16)
$$= \underbrace{\bullet}_{F_{1}} + \underbrace{\bullet}_{F_{2}} - E_{1}$$

Proof. The relation (15) is straightforward. Let us now verify equation (16), which is proved in the same way as for non-anchored foams, see [KR20b, Lemma 3.5]. Let S denote the surface on the left, and let F denote the surface obtained by surgering S as shown on the right. Denote by F^t (resp. F^b) the surface obtained from F by placing an additional dot on the top (resp. bottom) depicted disk. Note that anchor points, as well as their labels, are



FIGURE 1. Local models for colorings of F. Shaded indicates color 1 and solid white indicates color 2.

the same for F^t, F^b , and F. Colorings of F, F^t , and F^b are in a canonical bijection. There are four local models for a coloring of F, illustrated in Figure 1.

Let c be a coloring of F of the type shown in Figure 1c, with the corresponding coloring of F^t and F^b still denoted by c. We have

$$\langle F^t, c \rangle = \alpha_1 \langle F, c \rangle \qquad \langle F^b, c \rangle = \alpha_2 \langle F, c \rangle,$$

hence $\langle F^t, c \rangle + \langle F^b, c \rangle - E_1 \langle F, c \rangle = 0$. A similar calculation holds for a coloring c of Figure 1d type.

There is a natural bijection between colorings of S and colorings of F of Figures 1a and 1b types. Let c be a coloring of F of Figure 1a type, and continue to denote by c the corresponding coloring of S. Then

$$\chi(F) = \chi(S) + 2 \qquad \qquad \chi(F_2(c)) = \chi(S_2(c)),$$

$$\langle F^t, c \rangle = \alpha_1 \langle F, c \rangle \qquad \qquad \langle F^b, c \rangle = \alpha_1 \langle F, c \rangle,$$

so we have

$$\langle F^t, c \rangle + \langle F^b, c \rangle - E_1 \langle F, c \rangle = (\alpha_1 - \alpha_2) \langle F, c \rangle = \langle S, c \rangle.$$

Finally, if c is a coloring of F of the Figure 1b type, then

$$\chi(F) = \chi(S) + 2 \qquad \langle F^t, c \rangle = \alpha_2 \langle F, c \rangle,$$

$$\chi(F_2(c)) = \chi(S_2(c)) + 2 \qquad \langle F^b, c \rangle = \alpha_2 \langle F, c \rangle,$$

which yields

$$\langle F^t, c \rangle + \langle F^b, c \rangle - E_1 \langle F, c \rangle = (\alpha_2 - \alpha_1) \langle F, c \rangle = -(\alpha_2 - \alpha_1) \frac{\langle S, c \rangle}{\alpha_1 - \alpha_2} = \langle S, c \rangle$$

We now address equation (17), where anchor points are present. Let S denote the surface on the left-hand side of the equality. Let F^1 , F^2 denote the two anchored foams obtained by surgery on S in which the new anchor points are both labeled 1 or 2, respectively, so that (17) reads $\langle S \rangle = \langle F^1 \rangle + \langle F^2 \rangle$. For each i = 1, 2 there are four local models for a coloring of F^i , shown in Figure 2. Colorings c in Figure 2c and Figure 2d evaluate to zero for both i = 1, 2,

$$\left\langle F^{1},c\right\rangle =\left\langle F^{2},c\right\rangle =0$$

and they don't correspond to any colorings of S. There is a natural bijection between colorings of S and colorings of F^i of the types in Figures 2a and 2b.

Let c be a coloring of S in which the depicted region of S in (17) is colored 1, with the corresponding colorings of F^1 and F^2 still denoted c. We have immediately that $\langle F^1, c \rangle = 0$.



FIGURE 2. Local models for colorings of F^i . Shaded indicates color 1 and solid white indicates color 2.

On the other hand,

$$\chi(F^2) = \chi(S) + 2, \qquad \chi(F_2^2(c)) = \chi(S_2(c)),$$

and F^2 has two additional anchor points compared to S, both labeled 2 and their regions colored 1. Therefore

$$\langle F^1, c \rangle + \langle F^2, c \rangle = \langle F^2, c \rangle = (\alpha_1 - \alpha_2) \frac{\langle S, c \rangle}{\alpha_1 - \alpha_2} = \langle S, c \rangle.$$

Now let c be a coloring of S in which the depicted region of (17) is colored 2, and continue to denote by c the corresponding colorings of F^1 and F^2 . Then $\langle F^2, c \rangle = 0$. Since

$$\chi(F^1) = \chi(S) + 2, \qquad \chi(F_2^1(c)) = \chi(S_2(c)) + 2,$$

and F^1 contains two more anchor points labeled 1 and colored 2 than S does, we obtain

$$\langle F^1, c \rangle + \langle F^2, c \rangle = \langle F^1, c \rangle = -(\alpha_2 - \alpha_1) \frac{\langle S, c \rangle}{\alpha_1 - \alpha_2} = \langle S, c \rangle.$$

Relation $\langle S \rangle = \langle F^1 \rangle + \langle F^2 \rangle$ in Figure (17) follows.

Equation (16) can also be written using shifted dots,



Corollary 2.7. The local relation (19) holds.



Proof. This can be seen by applying the neck-cutting relation (16) near the depicted contractible circle and evaluating the resulting anchored sphere according to the formula (10).

| | label 1 | label 2 |
|--------|---------|---------|
| i odd | 1 | -1 |
| i even | -1 | 1 |

FIGURE 3. The contribution of the *i*-th anchor point on S to adeg(S).

2.2. State spaces. Following [BHMV95, KR20b], we can apply the universal construction to the evaluation described above. Let $\mathcal{P} = \mathbb{R}^2 \setminus (0,0)$ denote the punctured plane. Given a collection C of disjoint simple closed curves in \mathcal{P} , let Fr(C) denote the free R_{α} -module with a basis consisting of properly embedded compact surfaces $S \subset \mathbb{R}^2 \times (-\infty, 0]$ with $\partial S = C$ and which are transverse to the ray $L_- := (0,0) \times (-\infty, 0]$. The intersection $S \cap L_-$ is a 0-submanifold of L_- and consists of finitely many points. Moreover, each such surface Smust carry a labeling, a map

$$\ell = \ell_S : S \cap L_- \longrightarrow \{1, 2\}$$

from the set of its intersection points with the ray L_{-} (its anchor points) to $\{1, 2\}$. For a basis element $S \in Fr(C)$, let $\overline{S} \subset \mathbb{R}^2 \times [0, \infty)$ denote its reflection through the plane \mathbb{R}^2 . Labels of anchor points do not change upon reflection. For two basis elements $S, S' \in Fr(C)$ denote by $\overline{SS'}$ the closed anchored surface obtained by gluing \overline{S} to S' along their common boundary C.

Define a bilinear form

(20)
$$(-,-)\colon \operatorname{Fr}(C) \times \operatorname{Fr}(C) \to R_{\alpha}$$

by setting $(S, S') = \langle \overline{S}S' \rangle$. A direct computation shows that the form is symmetric, since for a closed surface T the evaluation satisfies $\langle \overline{T} \rangle = \langle T \rangle$.

Define the state space of C, denoted $\langle C \rangle$, to be the quotient of Fr(C) by the kernel

$$\{x \in \operatorname{Fr}(C) \mid (x, y) = 0 \text{ for all } y \in \operatorname{Fr}(C)\}\$$

of this bilinear form. For a basis element $S \in Fr(C)$, we will write [S] to denote its equivalence class in $\langle C \rangle$.

Equip the ground ring R_{α} with a bigrading by placing α_1, α_2 in bidegree (2,0). We extend this bigrading (qdeg, adeg) to Fr(C) as follows. For a basis element $S \in Fr(C)$ with d dots and m anchor points, set the quantum grading $qdeg(S) \in \mathbb{Z}$ to be

(21)
$$q \deg(S) = -\chi(S) + 2d + m.$$

Note that if S is a closed surface, then $\langle S \rangle \in R_{\alpha}$ is a homogeneous polynomial of degree qdeg(S), following the degree convention (1).

Next, let $\ell(1), \ldots, \ell(m)$ denote the labels of the anchor points of S, ordered from bottom to top, and define the annular grading $\operatorname{adeg}(S) \in \mathbb{Z}$ by setting

(22)
$$\operatorname{adeg}(S) = \sum_{i=1}^{m} (-1)^{i+\ell(i)}$$

In other words, if the *i*-th anchor point p_i is labeled 1, then it contributes 1 to adeg if *i* is odd and -1 if *i* is even. Likewise, if p_i has label 2 then it contributes -1 if *i* is odd and 1 if *i* is even, see also Figure 3. Multiplication by α_1, α_2 increases (qdeg, adeg)-bidegree by (2,0).



FIGURE 4. The (qdeg, adeg) bidegrees of some anchored surfaces whose boundary consists of two non-contractible circles.

Example 2.8. Let C consist of two non-contractible circles. The bidegree (qdeg, adeg) of the four anchored surfaces in Fr(C) whose underlying surface consists of two disks each intersecting L_{-} once are recorded in Figure 4.

Lemma 2.9. Let S be an anchored surface. Then $\langle S \rangle = 0$ or $\operatorname{adeg}(S) = 0$.

Proof. If some component of S has anchor points with different labels then $\langle S \rangle = 0$. Assume that all anchor points on any component of S are labeled identically. We also assume that S intersects L, otherwise $\operatorname{adeg}(S) = 0$ is immediate. As usual, order the anchor points p_1, \ldots, p_m from bottom to top.

Take a generic half-plane P in \mathbb{R}^3 containing the anchor line L, so that $P \cap S$ consists of finitely many arcs (with boundary on L) and circles (disjoint from L). For any arc a in $P \cap S$ with boundary $\partial a = \{p_i, p_j\}$, necessarily i and j have opposite parities, and moreover $\ell(p_i) = \ell(p_j)$ by assumption. Therefore the total contribution of the anchor points p_i and p_j to $\operatorname{adeg}(S)$ is zero. Summing over all arcs in $P \cap S$ yields the statement of the lemma.

The subspace ker((,)) \subset Fr(C) respects this bigrading on Fr(C). Consequently, the bigrading descends to the state space $\langle C \rangle$.

Note that the relations (16) and (17) are bi-homogeneous. Let $S \in Fr(C)$ be a basis element of the form $S = S_1 \sqcup S_2$ where $S_1, S_2 \in Fr(C)$ are anchored surfaces with S_2 closed. Then in $\langle C \rangle$ we have

$$[S] = \langle S_2 \rangle [S_1], \quad \langle S_2 \rangle \in R_{\alpha}.$$

Moreover, the relation (23) is bi-homogeneous. That it is homogeneous with respect to qdeg follows from the fact that $\langle S_2 \rangle \in R_{\alpha}$ is a polynomial of degree $qdeg(S_2)$. Lemma 2.9 ensures that $adeg(S_2) = adeg(\langle S_2 \rangle) = 0$, so $adeg(S) = adeg(S_1)$.

Given a bigraded module $M = \bigoplus_{(i,j) \in \mathbb{Z}^2} M_{i,j}$ over a commutative domain such that each $M_{i,j}$ has finite rank, define its graded rank to be

$$\operatorname{grank}(M) = \sum_{i,j} \operatorname{rank}(M_{i,j}) q^i a^j.$$

Lemma 2.10. Let $C \subset \mathcal{P}$ be a single circle. Then the state space $\langle C \rangle$ is a free R_{α} -module of rank 2. Moreover, we have

$$\operatorname{grank}(\langle C \rangle) = \begin{cases} q + q^{-1} & \text{if } C \text{ is contractible} \\ a + a^{-1} & \text{if } C \text{ is non-contractible.} \end{cases}$$



FIGURE 5. Basis elements for the state space of a single circle C. The first two surfaces form a basis if C is contractible, and the last two form a basis if C is non-contractible.

Proof. We consider two cases. If C is contractible, then by applying the neck-cutting relation (16) near C and evaluating closed anchored surfaces as in equation (23), we see that $\langle C \rangle$ is spanned by the two elements S and S_{\bullet} shown in Figures 5a and 5b. Bidegrees of S and S_{\bullet} are (-1,0) and (1,0), respectively. Computing the matrix of the bilinear form (20) for these elements yields

$$\begin{pmatrix} \overline{S}S & \overline{S}S_{\bullet} \\ \overline{S}_{\bullet}S & \overline{S}_{\bullet}S_{\bullet} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & E_1 \end{pmatrix},$$

which is invertible, thus S, S_{\bullet} constitute a basis for $\langle C \rangle$.

Now suppose C is non-contractible. Applying the neck-cutting relation (17) near C and evaluating closed anchored surfaces shows that the two elements S_1, S_2 depicted in Figures 5c and 5d span $\langle C \rangle$. Bidegrees of S_1 and S_2 are (0, 1) and (0, -1), respectively. The matrix of the bilinear form is

$$\begin{pmatrix} \overline{S_1}S_1 & \overline{S_1}S_2\\ \overline{S_2}S_1 & \overline{S_2}S_2 \end{pmatrix} = \begin{pmatrix} 1 & 0\\ 0 & 1 \end{pmatrix},$$

hence S_1, S_2 are linearly independent and constitute a basis of $\langle C \rangle$.

Theorem 2.11. Let $C \subset \mathcal{P}$ consist of n contractible circles and m non-contractible circles. Then the state space $\langle C \rangle$ is a free R_{α} -module of rank 2^{n+m} . Moreover, we have

grank
$$(\langle C \rangle) = (q + q^{-1})^n (a + a^{-1})^m$$
.

Proof. Consider a 2^{n+m} element set B(C) of basis vectors of Fr(C) consisting of surfaces S satisfying

- Each component of S is a disk.
- Each disk in S with contractible boundary is disjoint from L_{-} and carries either zero or one dot.
- Each disk in S with non-contractible boundary intersects L_{-} exactly once, and its intersection point may be labeled by either 1 or 2.

That B(C) spans $\langle S \rangle$ follows from applying the two neck-cutting relations (16) and (17) near the circles in C and evaluating closed anchored surfaces. Linear independence of B(C) and the statement regarding graded rank follow from the computations in Lemma 2.10. \Box

Elements of the basis B(C) constructed above are standard generators. For such a $\Sigma \in B(C)$ with d dots and anchor points labeled ℓ_1, \ldots, ℓ_m , we have

(24)
$$\operatorname{qdeg}(\Sigma) = -n + 2d, \quad \operatorname{adeg}(\Sigma) = \sum_{i=1}^{m} (-1)^{i+\ell(i)}.$$

Let $C_0, C_1 \subset \mathcal{P}$ be two collections of disjoint circles in the punctured plane. An *anchored* cobordism from C_0 to C_1 is a smoothly and properly embedded compact surface $S \subset \mathbb{R}^2 \times [0, 1]$ with boundary $\partial S = C_0 \sqcup C_1$, such that $C_i \subset \mathbb{R}^2 \times \{i\}$, i = 0, 1. Moreover, S is required to intersect the arc $L_{[0,1]} := (0,0) \times [0,1]$ transversely and come equipped with a labeling of these intersection points (called *anchor points*), which is a map

$$\ell = \ell_S \colon S \cap L_{[0,1]} \to \{1,2\}$$

from the set of its anchor points to $\{1, 2\}$. Anchored cobordisms are allowed to carry dots which can float on components but cannot jump to a different component.

We compose anchored cobordisms in the usual manner. For anchored cobordisms $S_1: C_0 \rightarrow C_1, S_2: C_1 \rightarrow C_2$, let $S_2S_1: C_0 \rightarrow C_2$ denote the anchored cobordism obtained by gluing along the common boundary C_1 and re-scaling. Labels of anchor points of S_2S_1 are inherited from labels of S_1 and S_2 .

As above, if an anchored cobordism S from C_0 to C_1 has m anchor points and carries d dots, define

$$qdeg(S) = -\chi(S) + 2d + m.$$

Let $\ell(1), \ldots, \ell(m)$ denote the labels of anchor points of S, ordered from bottom to top, and let n be the number of non-contractible circles in C_0 . Set

$$\operatorname{adeg}(S) = (-1)^n \sum_{i=1}^m (-1)^{i+\ell(i)}.$$

Remark 2.12. If $C_0 = \emptyset$, then S is a basis element of $Fr(C_1)$, and moreover the two degrees qdeg(S), adeg(S) defined above for anchored cobordisms agree with the definitions in (21) and (22) for elements of $Fr(C_1)$.

An anchored cobordism S from C_0 to C_1 induces an R_{α} -linear map

$$S: \operatorname{Fr}(C_0) \to \operatorname{Fr}(C_1)$$

defined on the basis by gluing along the common boundary C_0 . The definition of state spaces via universal construction immediately implies that we have an induced map

$$(25) \qquad \langle S \rangle : \langle C_0 \rangle \to \langle C_1 \rangle .$$

Lemma 2.13. Let $S_1: C_0 \to C_1$ and $S_2: C_1 \to C_2$ be anchored cobordisms. Then

$$\operatorname{qdeg}(S_2S_1) = \operatorname{qdeg}(S_2) + \operatorname{qdeg}(S_1), \quad \operatorname{adeg}(S_2S_1) = \operatorname{adeg}(S_2) + \operatorname{adeg}(S_1).$$

In particular, $\langle S_1 \rangle : \langle C_0 \rangle \to \langle C_1 \rangle$ is a map of bidegree $(qdeg(S_1), adeg(S_1))$.

Proof. The first equality involving qdeg is straightforward. Let n and m denote the number of non-contractible circles in C_0 and C_1 respectively, and let k denote the number of anchor points of S_1 . We have

$$\operatorname{adeg}(S_2S_1) = \operatorname{adeg}(S_1) + (-1)^{n+m+k} \operatorname{adeg}(S_2).$$

Note n + m + k is even, since it is equal to the number of anchor points of the closed surface obtained by gluing disks to all boundary circles of S_1 .

The final statement concerning the bidegree of $\langle S_1 \rangle$ follows from interpreting generators of $\langle C_0 \rangle$ as anchored cobordisms $\emptyset \to C_0$, as in Remark 2.12.



FIGURE 6. Elementary saddles involving non-contractible circles.

Definition 2.1. An annular cobordism is an anchored cobordism $S \subset \mathbb{R}^2 \times [0, 1]$ which is disjoint from the arc $L_{[0,1]} = (0,0) \times [0,1]$. An elementary annular cobordism is one with a single non-degenerate critical point with respect to the height function $\mathbb{R}^2 \times [0,1] \to [0,1]$.

Elementary annular cobordisms consist of a union of a product cobordism with a cup, cap, or saddle. Every annular cobordism may be obtained by composing finitely many elementary ones. Cup and cap annular cobordisms always have contractible boundary. There are four types of elementary annular saddles involving at least one non-contractible circle, illustrated in Figure 6. In the next four examples we write down the maps assigned to these four cobordisms in the standard bases of state spaces, as defined in the proof of Theorem 2.11. We also use the notation of shifted dots from (12).

Example 2.14. Figure 6a map. The calculation for this map follows at once from the skein relation (14).



Example 2.15. Figure 6b map. This calculation follows easily from the skein relation (19).



Example 2.16. Figure 6c map. A convenient way to perform this calculation is to use neckcutting with shifted dots (18) near the contractible circle and then simplify using the relations (13).

$$(28) \qquad \overbrace{1}^{\times} \longrightarrow \overbrace{1}^{\times} 1 \qquad \overbrace{1}^{1} \longrightarrow \overbrace{2}^{\times} \longrightarrow \overbrace{2}^{\times} 2 \qquad \textcircled{2}$$

Example 2.17. Figure 6d map. The neck-cutting relation (17) is helpful here. For the dotted cup we also use (14) to simplify further.



Recall the involution τ of R_{α} that transposes α_1, α_2 and extend it to an antilinear involution, also denoted τ , of the free R_{α} -module Fr(C) as follows. Involution τ on Fr(C) sends a surface S to the same surface with the labeling ℓ of anchor points reversed and acts on linear combinations by

$$\tau\left(\sum_{i}\lambda_{i}S_{i}\right) = \sum_{i}\tau(\lambda_{i})\tau(S_{i}).$$

For a closed surface S we have, by direct computation, $\langle \tau(S) \rangle = \tau(\langle S \rangle)$, showing compatibility of the two involutions. If S, in addition, carries shifted dots, involution τ reverses their labels, so that $\tau(\underline{0}) = \underline{0}$ and $\tau(\underline{0}) = \underline{0}$. Involution τ descends to an involution, also denoted τ , on $\langle C \rangle$. Annular degree is negated under τ , $\operatorname{adeg}(\tau(S)) = -\operatorname{adeg}(S)$, for an anchored cobordism S.

2.3. Annular link homology. Let ACob denote the category whose objects consist of collections of finitely many disjoint simple closed curves in the punctured plane \mathcal{P} . A morphism from C_0 to C_1 in ACob is an anchored cobordism from C_0 to C_1 , up to ambient isotopy fixing the boundary point-wise and mapping $L_{[0,1]}$ to itself. Let ACob' denote the subcategory of ACob with the same objects as ACob but whose morphisms are isotopy classes of annular cobordisms, disjoint from the anchor line L. The composition of annular cobordisms is again annular.

Let R_{α} -ggmod denote the category of bigraded R_{α} -modules and homogeneous maps (of any bidegree) between them. We have a functor

$$\langle - \rangle$$
: ACob $\rightarrow R_{\alpha}$ - ggmod

which sends a collection of circles $C \subset \mathcal{P}$ to the state space $\langle C \rangle$ and sends an anchored cobordism S from C_0 to C_1 to the map $\langle S \rangle : \langle C_0 \rangle \to \langle C_1 \rangle$ as in (25). By Lemma 2.13, $\langle S \rangle$ is

a map of bidegree (qdeg(S), adeg(S)). We can restrict to the category of annular cobordisms to get a functor

$$\langle - \rangle' : \operatorname{ACob}' \to R_{\alpha} - \operatorname{ggmod},$$

which assigns to an annular cobordism S a map $\langle S \rangle' = \langle S \rangle$ of bidegree (qdeg(S),0). The restriction $\langle - \rangle'$ does not change the state space assigned to a collection of circles $C \subset \mathcal{P}$.

On the other hand, a functor

$$\mathcal{G}_{\alpha}$$
: ACob' $\rightarrow R_{\alpha}$ - ggmod

was introduced in [Akh20]. We briefly recall \mathcal{G}_{α} below.

Consider the algebra

$$A_{\alpha} = R_{\alpha}[X]/((X - \alpha_1)(X - \alpha_2)).$$

It is a free R_{α} -module with basis $\{1, X\}$. The trace $\epsilon_{\alpha} \colon A_{\alpha} \to R_{\alpha}$ given by $1 \mapsto 0, X \mapsto 1$ makes A_{α} into a Frobenius algebra, which defines a (1 + 1)-dimensional TQFT, a functor \mathcal{F}_{α} from the category of dotted cobordisms to the category of R_{α} -modules. A dot on a cobordism is interpreted as multiplication by $X \in A_{\alpha}$. Define a grading on A_{α} by setting

(30)
$$qdeg(1) = -1, \quad qdeg(X) = 1.$$

With this grading, a cobordism S with d dots is assigned by \mathcal{F}_{α} a map of degree $-\chi(S) + 2d$. Alternatively, the TQFT \mathcal{F}_{α} is the result of applying the universal construction to the closed surface evaluation (6) when restricted to surfaces disjoint from L and collections of contractible circles in \mathcal{P} . See [KR20b] for further details about the Frobenius pair (R_{α}, A_{α}) .

Let $C \subset \mathcal{P}$ be a collection of n contractible and m non-contractible circles. Define the bigraded R_{α} -module $\mathcal{G}_{\alpha}(C)$ as follows. As an R_{α} -module, we set $\mathcal{G}_{\alpha}(C) = \mathcal{F}_{\alpha}(C) = A_{\alpha}^{\otimes (n+m)}$. Define the *annular grading*, denoted adeg, on $\mathcal{F}_{\alpha}(C)$ as follows.

Every tensor factor A_{α} corresponding to a contractible circle is concentrated in annular degree zero. Order the non-contractible circles in C from outermost (furthest from the puncture) to innermost. Introduce the notation

(31)
$$v_0 = 1, \quad v_1 = X - \alpha_1, \\ v'_0 = 1, \quad v'_1 = X - \alpha_2.$$

Both $\{v_0, v_1\} = \{1, X - \alpha_1\}$ and $\{v'_0, v'_1\} = \{1, X - \alpha_2\}$ constitute an R_{α} -basis for A_{α} . Set

(32)
$$\operatorname{adeg}(v_0) = \operatorname{adeg}(v'_0) = -1, \quad \operatorname{adeg}(v_1) = \operatorname{adeg}(v'_1) = 1.$$

The annular grading on non-contractible circle is defined by assigning the homogeneous basis $\{v_0, v_1\}$ or $\{v'_0, v'_1\}$ to the corresponding tensor factor of A_{α} in an alternating manner with respect to nesting in \mathcal{P} , with the convention that the outermost circle is assigned $\{v_0, v_1\}$.

It is convenient to distinguish between the modules assigned to different types of circles in \mathcal{P} . Let V_{α} and V'_{α} denote the R_{α} -modules A_{α} with bases $\{v_0, v_1\}$ and $\{v'_0, v'_1\}$, respectively. The notation A_{α} will be reserved for the module assigned to a contractible circle, with basis $\{1, X\}$.

The R_{α} -module $\mathcal{G}_{\alpha}(C)$ also carries a quantum grading qdeg inherited from (30). Define a modified quantum grading qdeg' on $\mathcal{G}_{\alpha}(C)$ by

$$qdeg' = qdeg - adeg.$$

We will consider $\mathcal{G}_{\alpha}(C)$ as a bigraded R_{α} -module with bigrading (qdeg', adeg). Bidegrees are recorded in Figure 7.

| | 1 | X | v_0 | v_1 | v_0' | v_1' |
|-------|----|---|-------|-------|--------|--------|
| qdeg' | -1 | 1 | 0 | 0 | 0 | 0 |
| adeg | 0 | 0 | -1 | 1 | -1 | 1 |

FIGURE 7. The (qdeg', adeg)-bidegrees of relevant elements, where $\{1, X\}$ is a basis for a contractible circle and $\{v_0, v_1\}, \{v'_0, v'_1\}$ are bases for non-contractible circles.

Remark 2.18. The modified quantum grading qdeg' appears elsewhere in the literature and is more natural in the context of annular link homology. In [GLW18] this grading was denoted j'. Similarly, the annular link homology defined in [BPW19] carries the modified quantum grading.

We now define \mathcal{G}_{α} on annular cobordisms. For an annular cobordism $S \subset \mathbb{R}^2 \times [0, 1]$, if the boundary of S is contractible in \mathcal{P} then $\mathcal{G}_{\alpha}(S) = \mathcal{F}_{\alpha}(S)$, where \mathcal{F}_{α} is the TQFT corresponding to the Frobenius algebra A_{α} as above. Formulas for the maps assigned by \mathcal{G}_{α} to the four elementary cobordisms in Figure 6 are recorded below. If other essential circles are present, then due to parity the formulas may be slightly different from those below. To obtain the full set of formulas, one interchanges $v_0 \leftrightarrow v'_0$, $v_1 \leftrightarrow v'_1$, and $\alpha_1 \leftrightarrow \alpha_2$.

$$V_{\alpha} \otimes A_{\alpha} \xrightarrow{(A)} V_{\alpha} \qquad \qquad V_{\alpha} \otimes V'_{\alpha} \xrightarrow{(B)} A_{\alpha} \\ v_{0} \otimes 1 \mapsto v_{0} \qquad \qquad v_{0} \otimes v'_{0} \mapsto 0 \\ (34) \qquad v_{1} \otimes 1 \mapsto v_{1} \qquad (35) \qquad v_{1} \otimes v'_{0} \mapsto X - \alpha_{1} \\ v_{0} \otimes X \mapsto \alpha_{1}v_{0} \qquad \qquad v_{0} \otimes v'_{1} \mapsto X - \alpha_{2} \\ v_{1} \otimes X \mapsto \alpha_{2}v_{1} \qquad \qquad v_{1} \otimes v'_{1} \mapsto 0 \end{cases}$$

 $\begin{array}{cccc} (36) & V_{\alpha} \stackrel{(C)}{\longrightarrow} V_{\alpha} \otimes A_{\alpha} & & A_{\alpha} \stackrel{(D)}{\longrightarrow} V_{\alpha} \otimes V_{\alpha}' \\ v_{0} \mapsto v_{0} \otimes (X - \alpha_{2}) & & (37) & 1 \mapsto v_{0} \otimes v_{1}' + v_{1} \otimes v_{0}' \\ v_{1} \mapsto v_{1} \otimes (X - \alpha_{1}) & & X \mapsto \alpha_{1} v_{0} \otimes v_{1}' + \alpha_{2} v_{1} \otimes v_{0}' \end{array}$

Theorem 2.19. The functors $\langle - \rangle'$: ACob' $\rightarrow R_{\alpha}$ -ggmod and \mathcal{G}_{α} : ACob' $\rightarrow R_{\alpha}$ -ggmod are naturally isomorphic via bidegree-preserving maps.

Proof. Let $C \subset \mathcal{P}$ be a collection of circles. We will define an R_{α} -linear, bidegree preserving isomorphism $\Phi_C \colon \langle C \rangle \to \mathcal{G}_{\alpha}(C)$ and show that it is natural with respect to annular cobordisms.

Let *n* and *m* denote the number of contractible and non-contractible circles in *C*, respectively. Fix an ordering $1, \ldots, n$ on the contractible circles in *C*. The R_{α} -module $\mathcal{G}_{\alpha}(C)$ is free with basis given by elements of the form

$$y_1 \otimes \cdots \otimes y_n \otimes z_1 \otimes \cdots \otimes z_m,$$

where each y_i is in $\{1, X\}$, specifying a basis element of the *i*-th contractible circle, and each z_j is in either $\{v_0, v_1\}$ or $\{v'_0, v'_1\}$, depending on nesting, specifying basis elements of the non-contractible circles. The ordering of factors $z_1 \otimes \cdots \otimes z_m$ corresponding to noncontractible circles is from outermost to innermost as usual, so that the first factor z_1 labels the outermost non-contractible circle.



FIGURE 8. An example of the isomorphism Φ_C when C consists of one contractible circle and two non-contractible circles. Basis elements Σ of $\langle C \rangle$ are drawn with the corresponding basis element $\Phi_C(\Sigma) \in \mathcal{G}_{\alpha}(C)$ written underneath.

We now define the isomorphism $\Phi_C: \langle C \rangle \to \mathcal{G}_{\alpha}(C)$. Recall the standard basis B = B(C) for $\langle C \rangle$ defined in the proof of Theorem 2.11. For $\Sigma \in B$ with anchor points labeled ℓ_1, \ldots, ℓ_m , read from bottom to top, set

$$\Phi_C(\Sigma) = y_1 \otimes \cdots \otimes y_n \otimes z_1 \otimes \cdots \otimes z_m,$$

where $y_i = 1$ if the corresponding cup in Σ is undotted and $y_i = X$ if the corresponding cup in Σ is dotted. The generators z_i of non-contractible circles are determined using the rule

$$z_j = \begin{cases} v_1 & \text{if } j \text{ is odd and } \ell_j = 1\\ v_0 & \text{if } j \text{ is odd and } \ell_j = 2\\ v'_0 & \text{if } j \text{ is even and } \ell_j = 1\\ v'_1 & \text{if } j \text{ is even and } \ell_j = 2 \end{cases}$$

See Figure 8 for an example of the assignment Φ_C when n = 1, m = 2. By comparing the bidegree formula (24) for Σ with the bidegree of $\Phi_C(\Sigma)$ (see Figure 7), we see that Φ_C is a bidegree-preserving isomorphism. Recall that we use the modified quantum grading (33) for $\mathcal{G}_{\alpha}(C)$.

Now let $S: C_1 \to C_2$ be an annular cobordism. To complete the proof, we check that the square

$$\begin{array}{ccc} \langle C_1 \rangle & \stackrel{\Phi_{C_1}}{\longrightarrow} & \mathcal{G}_{\alpha}(C_1) \\ \langle S \rangle & & & \downarrow \mathcal{G}_{\alpha}(S) \\ \langle C_2 \rangle & \stackrel{\Phi_{C_2}}{\longrightarrow} & \mathcal{G}_{\alpha}(C_2) \end{array}$$

commutes. If all the boundary circles of S are contractible, then commutativity of the square is straightforward. Otherwise, if S has at least one non-contractible boundary circle, it suffices to consider the case where S is one of the elementary annular cobordisms depicted



(A) Seam points where three facets meet. (B) A seam vertex where six facets meet.

FIGURE 9. Local model of a pre-foam near singular points. The singular graph s(F) is drawn bold.

in Figure 6. Formulas for these maps were recorded in Examples 2.14 - 2.17. Comparing with the formulas (34) - (37) completes the proof.

Let $\mathbb{A} := S^1 \times [0, 1]$ denote the annulus. For an oriented link $L \subset \mathbb{A} \times [0, 1]$ in the thickened annulus, a generic projection of L onto $\mathbb{A} \times \{0\}$ yields a link diagram D in the interior of \mathbb{A} . Identifying the interior of \mathbb{A} with the punctured plane \mathcal{P} , we may form the cube of resolutions of D in the usual way, with all smoothings drawn in \mathcal{P} . Diagrams representing isotopic annular links are related by Reidemeister moves away from the puncture. By standard arguments [Kho00, BN05], applying the functor $\langle -\rangle'$: ACob' $\to R_{\alpha}$ – ggmod to the cube of resolutions yields a bigraded chain complex $\langle D \rangle$ whose chain homotopy type is an invariant of L. Theorem 2.19 implies that the resulting annular homology is isomorphic to that of [Akh20].

3. Unoriented SL(3) anchored homology

We recall definitions and notations from [KR21], including that of (unoriented) SL(3) foams and refer the reader to [KR21, Section 2.1] for more details.

Definition 3.1. A (closed) SL(3) pre-foam is a compact 2-dimensional CW complex equipped with a PL-structure such that each point has an open neighborhood that is either an open disk, the product of a tripod and an open interval (Figure 9a), or the cone over the 1-skeleton of a tetrahedron (Figure 9b). Points of the first type are called *regular*, those of the second are called *seam points*, and those of the third are called *seam vertices*. A (closed) SL(3)foam is a closed SL(3) pre-foam together with a PL embedding into \mathbb{R}^3 .

We will simply write *pre-foam* and *foam* in place of *closed* SL(3) (*pre-)foam*. For a prefoam F, denote by v(F) the set of seam vertices and by s(F) the set of seam points and seam vertices. The subspace s(F) is a 4-valent graph which may contain closed loops. Connected components of $s(F) \setminus v(F)$ are called *seams*.

The subspace $F \setminus s(F)$ is a (not necessarily compact) surface, and a connected component of $F \setminus s(F)$ will be called a *facet* of F. The (finite) set of facets of F is denoted f(F). Facets of pre-foams may be decorated by a finite number of dots, which are allowed to float freely on their facets but may not cross seams or enter seam vertices.

A coloring of a pre-foam F is a map

$$c\colon f(F)\to\{1,2,3\}.$$



FIGURE 10. The local model for a pre-admissible coloring near a seam point.

That is, a coloring assigns 1, 2 or 3 to each facet of F. A coloring is called *pre-admissible* if the three facets meeting at each seam of F have distinct colors, see Figure 10. For a pre-admissible coloring c and $1 \leq i, j \leq 3, i \neq j$, let $F_{ij}(c)$ denote the union of facets colored i or j. The pre-admissibility condition guarantees that each $F_{ij}(c)$ is a closed surface (see [KR21, Proposition 2.2]).

A coloring c is called *admissible* if each $F_{ij}(c)$ is orientable. For a foam F (that is, a prefoam embedded in \mathbb{R}^3), every pre-admissible coloring is admissible, since $F_{ij}(c)$ is a closed surface in \mathbb{R}^3 .

3.1. Unoriented anchored SL(3) foams and their evaluations. Fix a field k of characteristic 2. In this section the following commutative rings will be used.

- $R'_x = \Bbbk[x_1, x_2, x_3]$ is the ring of polynomials in three variables.
- $R_x = \mathbb{k}[E_1, E_2, E_3]$ the subring of R'_x that consists of symmetric polynomials in x_1, x_2, x_3 , with generators E_i being elementary symmetric polynomials:

$$E_1 = x_1 + x_2 + x_3,$$

$$E_2 = x_1 x_2 + x_1 x_3 + x_2 x_3,$$

$$E_3 = x_1 x_2 x_3.$$

- $R''_x = R'_x[(x_1+x_2)^{-1}, (x_2+x_3)^{-1}, (x_1+x_3)^{-1}]$ is a localization of R'_x given by inverting $x_i + x_j$, for $1 \le i < j \le 3$.
- $\widetilde{R}'_x = \Bbbk[\sqrt{x_1}, \sqrt{x_2}, \sqrt{x_3}]$ is the extension of R'_x obtained by introducing square roots of x_1, x_2, x_3 .
- $\widetilde{R}''_x = \mathbb{k}[\sqrt{x_1}, \sqrt{x_2}, \sqrt{x_3}, (x_1 + x_2)^{-1}, (x_2 + x_3)^{-1}, (x_1 + x_3)^{-1}]$ is a localization of \widetilde{R}'_x given by inverting $x_i + x_j$, for $1 \le i < j \le 3$.

All five of these rings are graded by setting $deg(x_i) = 2$, i = 1, 2, 3. Inclusions of the above rings are summarized in the following diagram.

$$(38) \qquad \qquad \widetilde{R}'_x \subset \widetilde{R}''_x \\ \cup \qquad \cup \\ R_x \subset R'_x \subset R''_x \end{cases}$$

We follow the notation established in [KR21] for these rings with the additional subscript x to distinguish from the notation in Section 2.

Definition 3.2. An anchored SL(3) foam F is an SL(3) foam $F' \subset \mathbb{R}^3$ that may intersect the line L at finitely many points away from the singular graph s(F') of F'. Thus each intersection point belongs to some facet f of F', and intersection of facets with L are required to be transverse. Denote by $p(F) = F \cap L$ the set of intersection points (anchor points) of F. Intersection points carry labels in $\{1, 2, 3\}$; that is, F comes equipped with a fixed map

$$\ell \colon p(F) \to \{1, 2, 3\}.$$

It is convenient to order anchor points p_1, \ldots, p_m from bottom to top, with labels $\ell_i = \ell(p_i)$, $i = 1, \ldots, m$.

We now refine the notion of admissible coloring of a foam to that of admissible coloring of an anchored foam F. Consider an anchored foam F with the underlying foam F'. A coloring $c \in \operatorname{adm}(F')$ induces a coloring of anchor points in F', by assigning to each point the color of its facet. We say that c is admissible if that's exactly the labeling of anchor points of F, that is, $\ell(p) = c(f)$ for each anchor point p in a facet f, and then set $c(p) = \ell(p)$.

In this way, the set of admissible colorings of F' is in a bijection with the set of admissible colorings of anchored foams F that become F' upon forgetting the labeling of anchor points:

$$\operatorname{adm}(F') \cong \prod_{F} \operatorname{adm}(F)$$

Various constructions with SL(3) foams in [KR21] extend directly to anchored foams. In particular, bicolored surfaces $F_{ij}(c)$ are well-defined, associated to an admissible coloring c. We will also call an admissible coloring simply a coloring. We will use i, j, k to denote the three elements of $\{1, 2, 3\}$, not necessarily in that order.

We refine [KR21, Definition 2.9] for anchored foams.

Definition 3.3. Let F be an anchored foam, $c \in \operatorname{adm}(F)$ be an admissible coloring, and Σ a connected component of $F_{ij}(c)$ which is disjoint from L. Define a coloring c' of F which swaps the colors i and j on facets of Σ , and leaves all other facets colored according to c. We say that c and c' are related by an ij-Kempe move along Σ . Note that since Σ has no anchor points, c' is still an admissible coloring of F.

Kempe moves can be done on components Σ of $F_{ij}(c)$ that intersect L as well, but the resulting anchored foam F_0 is different from F due to carrying different labels on anchor points on Σ .

For $k \in \{1, 2, 3\}$, denote by k', k'' its two complementary elements, so that $\{k, k', k''\} = \{1, 2, 3\}$. Let F be an anchored foam with labeling ℓ . Let $c \in \operatorname{adm}(F)$ be an admissible coloring. For an anchor point $p \in p(F)$ lying on a facet $f \in f(F)$, we set $c(p) = c(f) = \ell(p)$; that is, c(p) is the color of the facet, according to c, on which p lies, which equals $\ell(p)$ since c is admissible. For $1 \leq i \leq 3$, let $d_i(c)$ denote the number of dots on facets colored i. For $1 \leq i \neq j \leq 3$, let $F_{ij}(c)$ be the union of facets of F colored i or j. The space $F_{ij}(c)$ is a closed surface in \mathbb{R}^3 and hence has even Euler characteristic. Set

(39)
$$\langle F, c \rangle = \frac{P(F, c)}{Q(F, c)}$$

where

(40)
$$P(F,c) = \prod_{i=1}^{3} x_i^{d_i(c)} \cdot \left(\prod_{p \in p(F)} (x_{c(p)} + x_{\ell(p)'}) (x_{c(p)} + x_{\ell(p)''}) \right)^{1/2}$$

(41)
$$Q(F,c) = \prod_{1 \le i < j \le 3} (x_i + x_j)^{\chi(F_{ij}(c))/2}$$

The product of the two terms under the square root, for a given anchor point p, is equal to

$$(x_1 + x_2)(x_1 + x_3)$$
 if $c(p) = 1$,
 $(x_2 + x_1)(x_2 + x_3)$ if $c(p) = 2$,
 $(x_3 + x_1)(x_3 + x_2)$ if $c(p) = 3$.

Remark 3.1. This product is the inverse of the square decoration \Box in [KR21, Section 4.1]. The square decoration was used to study a separable version of the unoriented SL(3) theory, with the discriminant $\mathcal{D} = (x_1 + x_2)(x_1 + x_3)(x_2 + x_3)$ inverted, which is a version of the Lee theory. Here, we use the defect line L rather than freely floating square dots in [KR21, Section 4.1] in the opposite way, to add factors to the evaluation rather than divide by terms in the discriminant.

Remark 3.2. If c is an admissible coloring of the underlying foam F' of F but not of the anchored foam F, then the evaluation (39) is still defined and equal to zero:

(42)
$$\langle F, c \rangle = 0, \quad c \in \operatorname{adm} F' \setminus \operatorname{adm} F.$$

This holds since, for some $p \in p(F)$, its color c(p) differs from its label $\ell(p)$, so that $x_{c(p)} + x_{c(p)} = 0$ appears under the square root in (40) and P(F, c) = 0. Thus,

$$(x_{c(p)} + x_{\ell(p)'})(x_{c(p)} + x_{\ell(p)''}) = \begin{cases} (x_{\ell(p)} + x_{\ell(p)'})(x_{\ell(p)} + x_{\ell(p)''}) & \text{if } c(p) = \ell(p) \\ 0 & \text{otherwise.} \end{cases}$$

Define the *evaluation of* F to be

(43)
$$\langle F \rangle = \sum_{c \in \operatorname{adm}(F)} \langle F, c \rangle$$

Alternatively, we can sum over the larger set of $c \in \operatorname{adm}(F')$, due to (42).

Let us explain the square root in equation (40). The equality $\sqrt{x+y} = \sqrt{x} + \sqrt{y}$ holds in a commutative ring of characteristic 2, so $\langle F, c \rangle$ is in the ring \widetilde{R}''_x , see (38). We will show in Proposition 3.8 that, in fact, no square roots appear, so that $\langle F, c \rangle \in R''_x$. Likewise, in Proposition 3.9 we show that $\langle F \rangle \in R'_x$.

The evaluation (43) is multiplicative with respect to disjoint union and does not depend on a particular embedding of F into $M = (\mathbb{R}^3, L)$ as long as anchor points on F and their labels are specified.

If an anchored foam F is a disjoint union of anchored foams $F_1 \sqcup \cdots \sqcup F_k$, then

$$\langle F \rangle = \langle F_1 \rangle \cdots \langle F_k \rangle$$

If F is disjoint from L, then $\langle F \rangle$ is equal to the evaluation in [KR21, Section 2.3].

Example 3.3. Let F be a 2-sphere \mathbb{S}^2 with two anchor points and d dots. Its evaluation is zero unless both points have the same label $i \in \{1, 2, 3\}$, in which case there is only admissible coloring c which colors F by i. Let $j, k \in \{1, 2, 3\}$ denote the complementary elements to i. The surfaces $F_{ij}(c), F_{ik}(c)$ are 2-spheres, while $F_{jk}(c) = \emptyset$. Then the evaluation is

$$\langle F \rangle = \frac{x_i^d \left((x_i + x_j)^2 (x_i + x_k)^2 \right)^{1/2}}{(x_i + x_j)(x_i + x_k)} = x_i^d.$$

Example 3.4. More generally, let F be a genus g surface carrying d dots and 2n > 0 anchor points. It evaluates to zero unless all points are labeled by the same $i \in \{1, 2, 3\}$. In this case, letting $j, k \in \{1, 2, 3\}$ be the complementary elements to i, the evaluation is

$$\langle F \rangle = \frac{x_i^d \left((x_i + x_j)(x_i + x_k) \right)^n}{\left((x_i + x_j)(x_i + x_k) \right)^{1-g}} = x_i^d \left((x_i + x_j)(x_i + x_k) \right)^{n+g-1}$$

Example 3.5. Consider the theta foam F whose facets each intersect L once, with anchor points labeled $i, j, k \in \{1, 2, 3\}$ and facets carrying d_1, d_2 , and d_3 dots, as shown in (44).

In an admissible coloring of the underlying foam, the three facets must have distinct colors, so $\langle F \rangle = 0$ if i, j, k are not distinct. If i, j, k are distinct, then there is one admissible coloring c which colors the top, middle, and bottom facets, respectively, by i, j, and k. The surfaces $F_{ij}(c), F_{ik}(c), F_{jk}(c)$ are 2-spheres, and the evaluation is

$$\langle F \rangle = x_i^{d_1} x_j^{d_2} x_k^{d_3}.$$

Remark 3.6. Note that the evaluation of an anchored foam is in general not a symmetric function in x_1, x_2, x_3 , whereas in [KR21] the evaluation is always an element of R_x .

Let us call a sequence $\ell \in \{1, 2, 3\}^m$ pre-admissible if the following holds. Let u_1, u_2, u_3 be three non-zero elements of the abelian group $\mathbb{Z}/2 \times \mathbb{Z}/2$. Sequence ℓ is pre-admissible iff

(45)
$$\sum_{i=1}^{m} u_{\ell(i)} = 0 \in \mathbb{Z}/2 \times \mathbb{Z}/2$$

Proposition 3.7. If an anchored foam F has an admissible coloring, the sequence ℓ of its anchor points is pre-admissible.

Proof. Consider a generic intersection of F with a half-plane in \mathbb{R}^3 bounding L. This intersection is a trivalent graph Γ in the half-plane. Coloring c of F induces a coloring c' of edges of Γ such that around each trivalent vertex of Γ the colors of the three edges are distinct (Tait coloring). On the boundary points (one-valent vertices) of Γ the coloring is given by labeling ℓ . The sum on the left hand side of (45) is zero since it can alternatively be written as the sum of triples of vectors $u_1 + u_2 + u_3 = 0$ over all trivalent vertices of Γ . Each inner edge of Γ , that bounds two trivalent vertices, contributes $u_i + u_i = 0$ to the sum, where i is the color of the edge. An edge with one or both endpoints on the boundary contributes the sum of u_i 's, over its boundary points.

For an anchored foam F and $1 \le i \le 3$, let an(i) denote the number of anchor points of F with label i (the dependence on F is omitted).

Proposition 3.8. For an anchored foam F and an admissible coloring c, we have $\langle F, c \rangle \in R''_x$.

Proof. Recall the rings R''_x and \tilde{R}''_x defined in (38). It's clear that $\langle F, c \rangle$ belongs to the larger ring \tilde{R}''_x .

The expression in (39) under the square root is equal to

$$(x_1 + x_2)^{\operatorname{an}(1) + \operatorname{an}(2)} (x_2 + x_3)^{\operatorname{an}(2) + \operatorname{an}(3)} (x_1 + x_3)^{\operatorname{an}(1) + \operatorname{an}(3)}.$$

For $1 \leq i < j \leq 3$, the integer $\operatorname{an}(i) + \operatorname{an}(j)$ is even since it is equal to the number of intersection points of the closed surface $F_{ij}(c)$ with L, see also Proposition 3.7. Consequently, taking the square root produces integral exponent of $x_i + x_j$, implying that $\langle F, c \rangle$ is in R''_x . \Box

Using the above notation, the square root term in (40) is equal to

(46)
$$\widetilde{Q}(F,c) := \prod_{1 \le i < j \le 3} (x_i + x_j)^{(\operatorname{an}(i) + \operatorname{an}(j))/2},$$

so that formula (39) can be rewritten as

(47)
$$\langle F, c \rangle = \prod_{i=1}^{3} x_i^{d_i(c)} \prod_{1 \le i < j \le 3} (x_i + x_j)^{(\operatorname{an}(i) + \operatorname{an}(j) - \chi(F_{ij}(c)))/2}.$$

Proposition 3.9. For an anchored foam F, we have $\langle F \rangle \in R'_x = \mathbb{k}[x_1, x_2, x_3]$.

Proof. Proof of Theorem 2.17 in [KR21] extends with minor changes to this case. Note that the evaluation is no longer a symmetric function. We must show that positive powers of $x_i + x_j$, $1 \le i < j \le 3$, do not appear in the denominator of $\langle F \rangle$. Let us specialize to i = 1, j = 2. Denominators $x_1 + x_2$ in the evaluations $\langle F, c \rangle$ may appear only from the components of $F_{12}(c)$ that are two-spheres. If a 2-sphere does not intersect L, the proof in [KR21] works in this case as well. Suppose a 2-sphere component Σ of $F_{12}(c)$ intersects Lin an(1) points colored 1 and an(2) points colored 2 (necessarily in the corresponding facets of F carrying those colors under c). These points contribute

$$(x_1 + x_2)^{\operatorname{an}(1) + \operatorname{an}(2)} (x_1 + x_3)^{\operatorname{an}(1)} (x_2 + x_3)^{\operatorname{an}(2)}$$

to the expression under the square root, and $an(1) + an(2) \ge 2$, allowing to cancel the denominator term $x_1 + x_2$ that Σ contributes. Summing over all admissible colorings and otherwise following the arguments in [KR21, Theorem 2.17] implies the result.

Remark 3.10. Contributions of anchor points to the evaluation $\langle F, c \rangle$ can be interpreted as follows. Consider polynomial $f(x) = (x - x_1)(x - x_2)(x - x_3) \in R'_x[x]$. Then

$$f'(x) = (x - x_2)(x - x_3) + (x - x_1)(x - x_3) + (x - x_1)(x - x_2)$$

and

$$f'(x_1) = (x_1 - x_2)(x_1 - x_3),$$

$$f'(x_2) = (x_2 - x_1)(x_2 - x_3),$$

$$f'(x_3) = (x_3 - x_1)(x_3 - x_2).$$

Contribution of an anchor point p with a label $i = \ell(p)$ to the evaluations $\langle F, c \rangle$ and $\langle F \rangle$ is then $\sqrt{f'(x_i)}$, the square root of the derivative of f at the root x_i of the polynomial

f. In characteristic two signs do not matter, but this observation hints how to extend the evaluation to characteristic 0.

Since the labels i_1, \ldots, i_m of anchor points are fixed in a given F, these marked points contribute the same term,

$$\sqrt{f'|_{L\cap F}} := \left(\prod_{r=1}^m f'(x_{i_r})\right)^{1/2}$$

and we have

(48)
$$\langle F, c \rangle = \sqrt{f'|_{L \cap F}} \cdot \langle F', c \rangle, \quad \langle F \rangle = \sqrt{f'|_{L \cap F}} \cdot \langle F' \rangle,$$

where F' is the foam F viewed as a regular foam with anchored points and their labels ignored. When coloring c of F is not compatible with labels of anchor points, though, we should define $\sqrt{f'|_{L\cap F}} = 0$ to match the formula $\langle F, c \rangle = 0$.

Also notice that, switching to characteristic 0 and from the matrix factorization viewpoint [KR08], f(x) = w'(x) is the derivative of the potential $w(x) = x^4/4 - E_1 x^3/3 + E_2 x^2/2 - E_3 x$, so that the contributions of anchor points are given by square roots of the second derivative $\sqrt{w''(x_i)}$ at critical points of w, analogous to the square root of the Hessian factor that appears, for example, in the steepest descent method formulas.

3.2. Skein relations. In this subsection we record several local relations satisfied by the evaluation of anchored SL(3) foams. We start with the following proposition concerning the relations in [KR21, Section 2.5], which should be understood as occurring away from the anchor line L.

Proposition 3.11. The twelve local relations in Propositions 2.22 – 2.33 from [KR21] hold.

Proof. The arguments in [KR21] apply without modification.

We will use shifted dots in this section, as in (12). For $1 \le i \le 3$, we allow anchored foams to carry decorations of the form $(i) = \bullet + x_i$ on a facet. They are required to be disjoint from L, float freely on their facets, but cannot move past seams or seam vertices.

(49)
$$(i) = + x_i$$

For an anchored foam F carrying (i) on some facet $f \in f(F)$, any coloring $c \in \operatorname{adm}(F)$ which colors f by i evaluates to zero, $\langle F, c \rangle = 0$. An anchor point labeled i has the same effect as placing $\sqrt{(j)(k)} = \sqrt{(i')(i'')}$ on the facet on which it lies (recall our conventions that $\{1, 2, 3\} = \{i, j, k\} = \{i, i', i''\}$). See also equation (53) and the discussion in Section 3.4.

We also have relations involving the anchor line.

Lemma 3.12. The local relations (50), (51), (52), (53), and (54) hold.



In the last two equations, $\{i, j, k\} = \{1, 2, 3\}.$

Proof. Let us verify equation (50); the other three relations are easier to check and the proof is left to the reader. Denote by F the anchored foam on the left-hand side, and by G^1, G^2, G^3 the three foams on the right-hand side, with the superscript corresponding to the labels of the depicted anchor points. For $1 \le i \le 3$, let $\operatorname{adm}_i(F)$ be the set of admissible colorings of F in which the depicted tube is colored by i. Admissible colorings of G^i must color the two disks by i, so there is a natural bijection $\operatorname{adm}_i(F) \cong \operatorname{adm}(G^i)$.

i

k

For $c \in \operatorname{adm}_i(F)$, let $c' \in \operatorname{adm}(G^i)$ denote the corresponding coloring. We will show that

$$\langle F, c \rangle = \left\langle G^i, c' \right\rangle,$$

which completes the proof.

The anchored foam G^i carries two more anchor points, both labeled *i*, than *F* does, while the dot placement for G^i and *F* is the same, so

$$P(G, c') = (x_i + x_j)(x_i + x_k)P(F, c),$$

where $\{i, j, k\} = \{1, 2, 3\}$. On the other hand,

$$\chi(G_{ij}^{i}(c')) = \chi(F_{ij}(c)) + 2, \quad \chi(G_{ik}^{i}(c')) = \chi(F_{ik}(c)) + 2, \quad \chi(G_{jk}^{i}(c')) = \chi(F_{jk}(c)),$$

which yields

$$Q(G,c') = (x_i + x_j)(x_i + x_k)Q(F,c)$$

Thus $\langle F, c \rangle = \langle Q, c' \rangle$ as desired. Summing over all admissible colorings of F we get

$$\langle F \rangle = \langle G^1 \rangle + \langle G^2 \rangle + \langle G^3 \rangle,$$

completing the proof.

3.3. State spaces. We generalize the notion of webs and cobordisms between them from [KR21, Section 3.1] in the presence of the anchor line L.

Definition 3.4. A web is a trivalent graph Γ which is PL-embedded into the punctured plane $\mathcal{P} = \mathbb{R}^2 \setminus \{(0,0)\}$. We allow webs to have closed loops with no vertices. A anchored foam with boundary V is the intersection of a closed anchored foam $F \subset \mathbb{R}^3$ with a thickened plane $\mathbb{R}^2 \times [0,1]$ such that $F \cap (\mathcal{P} \times \{i\})$, i = 0,1 is a web (in particular, F is disjoint from the two points (0,0,0) and (0,0,1)), and dots of F are disjoint from $\mathbb{R}^2 \times \{i\}$ and from L. Foams with boundary are considered equivalent if there is an orientation-preserving homeomorphism of $\mathbb{R}^2 \times [0,1]$ taking one to the other which fixes the boundary of $\mathbb{R}^2 \times [0,1]$ pointwise and maps the line segment $L_{[0,1]} := \{(0,0)\} \times [0,1]$ to itself.

For a foam with boundary V, let

$$p(V) = V \cap L_{[0,1]}$$

denote its intersection points with the anchor line, called *anchor points*. Each anchor point is required to carry a label in $\{1, 2, 3\}$.

We view V as a cobordism from the web $\partial_0 V := V \cap (\mathbb{R}^2 \times \{0\})$ to the web $\partial_1 V := V \cap (\mathbb{R}^2 \times \{1\})$. A closed foam is then a cobordism from the empty web to itself. We will often refer to foams with boundary simply as *foams* when the meaning is clear from context. Composition WV of foams V, W with $\partial_1 V = \partial_0 W$ is defined in the natural way. We obtain a category AFoam of webs and anchored foams.

The category AFoam has a contravariant involution ω which is the identity on webs and which sends a foam to its reflection about $\mathbb{R}^2 \times \{1/2\}$, preserving the labels of anchor points. As for closed foams, denote by s(V) and v(V) the singular graph and singular vertices, respectively, of a foam with boundary V. Define the *degree* of V to be

(55)
$$\deg(V) = 2(|d(V)| + |p(V)| - \chi(V)) - \chi(s(V)),$$

where d(V) is the set of dots on V.

The definition of admissible colorings extend naturally to anchored foams with boundary. An admissible coloring induces a Tait coloring on the boundary webs. If a foam with boundary V has an admissible coloring c, then by [KR21, Remark 2.8],

(56)
$$\deg(V) = 2|d(V)| + 2|p(V)| - \left(\chi(V_{12}(c)) + \chi(V_{13}(c)) + \chi(V_{23}(c))\right).$$

It follows that for a closed foam F, its evaluation $\langle F \rangle \in R'_x$ is a homogeneous polynomial of degree deg(F).

Lemma 3.13. For composable foams V and W, we have

 $\deg(WV) = \deg(W) + \deg(V).$



FIGURE 11. Direct sum decompositions from [KR21, Section 3.3], where the depicted regions do not contain the puncture.

Proof. This follows from [KR21, Proposition 3.1] and |p(WV)| = |p(W)| + |p(V)|.

We now define state spaces for webs via universal construction and the evaluation formula (43). For a web Γ , let

 $Fr(\Gamma)$

denote the free R'_x -module generated by all anchored foams V from the empty web to Γ . Define a bilinear form

$$(-,-)\colon\operatorname{Fr}(\Gamma)\times\operatorname{Fr}(\Gamma)\to R'_{\mathfrak{s}}$$

by $(V, W) = \langle \omega(V)W \rangle$. This bilinear form is symmetric since $\langle F \rangle = \langle \omega(F) \rangle$ for any closed anchored foam F. Define the state space $\langle \Gamma \rangle := \operatorname{Fr}(\Gamma)/\ker((-, -))$ to be the quotient of $\operatorname{Fr}(\Gamma)$ by the kernel

$$\ker((-,-)) = \{ x \in \operatorname{Fr}(\Gamma) \mid (x,y) = 0 \text{ for all } y \in \operatorname{Fr}(\Gamma) \}$$

of the bilinear form. Note that (-, -) is degree-preserving, so its kernel and the state space $\langle \Gamma \rangle$ are graded R'_x -modules.

An anchored foam $V \colon \Gamma_0 \to \Gamma_1$ naturally induces a map

$$\langle V \rangle : \langle \Gamma_0 \rangle \to \langle \Gamma_1 \rangle$$

of degree deg(V), defined by sending the equivalence class of a basis element $U \in \operatorname{Fr}(\Gamma_0)$ to the class of the composition VU. This is functorial with respect to composition of anchored foams, $\langle WV \rangle = \langle W \rangle \langle V \rangle$ for composable anchored foams with boundary V and W.

Remark 3.14. For a web Γ and basis elements $V_1, V_2 \in Fr(\Gamma)$, an admissible coloring of the closed foam $\omega(V_2)V_1$ induces a Tait coloring of Γ . Thus $\langle \Gamma \rangle = 0$ if Γ has no Tait colorings; see also [KR21, Proposition 3.16].

Proposition 3.15. The local¹ isomorphisms in Propositions 3.12 - 3.15 from [KR21], also shown in Figure 11, hold.

Proof. Proposition 3.11 guarantees that the explicit isomorphisms defined in [KR21] hold in the anchored setting as well. \Box

¹Here local means that the webs involved in the isomorphisms are identical outside of a disk which is disjoint from the puncture, and in this disk they are related as in the figures accompanying the statements of the propositions.

Proposition 3.16. Let $\Gamma \subset \mathcal{P}$ be a web with a non-contractible circle C which bounds a disk in $\mathbb{R}^2 \setminus \Gamma$, and let $\Gamma' = \Gamma \setminus C$ be the web obtained by removing C. Then there is an isomorphism

$$\langle \Gamma \rangle \cong \langle \Gamma' \rangle \oplus \langle \Gamma' \rangle \oplus \langle \Gamma' \rangle$$

given by the maps shown in (57).



Proof. This follows from Example 3.3 and the relation (50). Note that there are no grading shifts in the three copies of $\langle \Gamma' \rangle$.

It is an interesting and nontrivial problem to identify the state spaces $\langle \Gamma \rangle$. In the construction in [KR21] without the anchor line, state spaces can be simplified using the relations in [KR21, Section 3.3], see Figure 11. In particular, bipartite webs always contain a contractible circle, bigon, or square, so the state space in the bipartite case is a free module of graded rank equal to the Kuperberg bracket [Kup96], normalized as in [Kho04]; see also [KR21, Proposition 3.17, Proposition 4.15]. The simplest web which cannot be simplified using the relations in Figure 11 and for which the state space is unknown is the dodecahedral graph, as explored in [Boo19, KR20a], and, on the gauge theory side, in [KM16, KM19b, KM19a].

One may also ask to identify state spaces in the presence of the anchor line and the modified evaluation considered in this paper. Propositions 3.15 and 3.16 give some ways to simplify state spaces. In general, we are not able to decompose the bigon, square, and triangle regions in Figure 11 if they contain the puncture. An extended evaluation, obtained by introducing additional types of intersection points of L with a foam, is discussed in Section 3.5. The following lemma addresses reducibility of smallest webs.

Lemma 3.17. Let $\Gamma \subset \mathbb{R}^2$ be a connected, planar, trivalent graph with no edges connecting a vertex to itself².

- (1) If Γ is bipartite, then Γ has at least two bounded faces with at most four edges each.
- (2) If at most one of the bounded faces of Γ has fewer than five edges, then Γ has at least eight vertices.

²A graph with such an edge has trivial state space, see Remark 3.14.

Proof. Let v, e, f denote the number of vertices, edges, and faces (including the unbounded face) of Γ , respectively. Label the faces $1, \ldots, f$, and for $1 \leq i \leq f$, let r_i denote the number of edges comprising the boundary of the *i*-th face. We have

(58)
$$\sum_{i=1}^{f} r_i = 2e = 3v,$$

where the second equality holds since Γ is trivalent.

We first prove statement (1). Since Γ is bipartite, each r_i is even. Suppose for the sake of contradiction that at most one bounded face of Γ has four or fewer edges. Then equation (58) implies

$$\sum_{i=1}^{f} r_i > 6(f-2),$$

so 12 > 6f - 3v. On the other hand, an Euler characteristic computation gives

$$12 = 6(f - e + v) = 6f - 3v,$$

which is a contradiction.

Let us now address statement (2). From equation (58) we obtain

$$3v \ge 5(f-2) + 4 = 5f - 6$$

since, by assumption, there are (f-2) faces with at least 5 edges each, and the remaining two faces each have at least two edges. This together with an Euler characteristic computation gives $f \ge 6$, and it follows that $v \ge 8$.

Corollary 3.18. Let $\Gamma \subset \mathcal{P}$ be a bipartite web. Then $\langle \Gamma \rangle$ is a free R'_x -module of rank equal to the number of Tait colorings of Γ .

Proof. By statement (1) of Lemma 3.17, any such web has either an innermost non-contractible circle or a region, not containing the puncture, which either bounds a closed loop, or is a bigon or square face. Thus state space can be reduced using Propositions 3.15 and 3.16, and since the resulting web remains bipartite we can continue the procedure. \Box

It is natural to ask what is the simplest web for which the state space cannot be reduced using Propositions 3.15 and 3.16. By Statement (2) of Lemma 3.17, such a web has at least eight vertices. The web shown in Figure 12 has precisely eight vertices and cannot be simplified using our local relations. We have not identified the state space of this web, but it can be approached via the 4-periodic (and, in general, non-exact) complex described in [KR21, Section 4.3]. It can be applied along any of the four edges of Figure 12 web near either the marked or the infinite point. One of the other three webs in the complex contains a loop and has trivial homology, but additional computations are needed to identify the state space due to non-exactness of the complex.

An annular graph $\Gamma \subset \mathcal{P}$ is called *reducible* if its state space can be reduced to a sum of those for the empty annular graph by recursively applying relations (A)-(D) in Figure 11 and relation in Proposition 3.16. It may make sense to also allow reductions to annular graphs without Tait colorings (including graphs with loops), since such graphs have trivial state spaces.

A reducible annular graph allows an identification of its state space with a suitable free graded R_x -module by recursively applying the above state sum decompositions. As a special



FIGURE 12. The simplest non-reducible web in the punctured plane.

case, we have the following decomposition formula for collections of simple closed curves in an annulus.

Proposition 3.19. Let $\Gamma \subset \mathcal{P}$ consist of *n* contractible circles and *m* non-contractible circles. Then the state space $\langle \Gamma \rangle$ is a free R'_x -module of graded rank $3^m(q^2 + 1 + q^{-2})^n$.

In particular, for a reducible Γ , the graded rank of the free R'_x -module $\langle \Gamma \rangle$ can be computed recursively.

Anchored foams and state spaces carry an additional $(\mathbb{Z}/2 \times \mathbb{Z}/2)$ -grading as follows. Recall that u_1, u_2, u_3 denote the nonzero elements of $\mathbb{Z}/2 \times \mathbb{Z}/2$. For a foam V with (possibly empty) boundary, define

$$\operatorname{adeg}(V) = \sum_{p \in p(V)} u_{\ell(p)}.$$

We call adeg the *annular degree*. Clearly adeg is additive under disjoint union and composition.

The annular degree extends to a $(\mathbb{Z}/2 \times \mathbb{Z}/2)$ -grading on $Fr(\Gamma)$, for a web $\Gamma \subset \mathcal{P}$, by setting the ground ring R'_x to be concentrated in annular degree zero. Proposition 3.7 implies that $\langle F \rangle = 0$ or $\operatorname{adeg}(\langle F \rangle) = 0$ for any closed foam F. It follows that (-, -) preserves annular degree, so adeg descends to a $(\mathbb{Z}/2 \times \mathbb{Z}/2)$ -grading on the state space $\langle \Gamma \rangle$. The annular grading is the unoriented version of the grading on state spaces of annular oriented webs by the integral weight lattice of \mathfrak{sl}_3 , see Section 4.4, even though the action of the latter is lacking on the equivariant annular state spaces.

In [KR21, Section 4] the authors consider localization of the unoriented SL(3) theory given by inverting the discriminant $\mathcal{D} = (x_1 + x_2)(x_1 + x_3)(x_2 + x_3)$. This localization results in a significant simplification of the theory, making it separable, so to speak. In particular, a suitable 4-term sequence of web state spaces in [KR21, Section 4.3] is exact.

This localization easily extends to the annular case. The corresponding 4-term sequence is exact in the annular case as well. The ground ring for that theory is $R'_{\mathcal{D}} := \Bbbk[x_1, x_2, x_3, \mathcal{D}^{-1}]$, with \Bbbk a characteristic two field. The analogue of [KR21, Proposition 4.13] holds: the localized state space of an annular web Γ is a projective $R'_{\mathcal{D}}$ -module of rank equal to the number of Tait colorings of Γ . The latter is the number of edge colorings of Γ into three colors so that at each vertex the colors are distinct. Proof of this result in [KR21] easily adapts to the annular case, with the modification that the region around the marked point can be inductively simplified, if necessary, by reducing to the other three terms in the exact sequence, until it has a single edge (a loop around the marked point).

3.4. Remark on Lee's theory. Recall the function

(59)
$$f(x) = (x+x_1)(x+x_2)(x+x_3) = x^3 + E_1 x^2 + E_2 x + E_3$$

(in characteristic 2 signs do not matter) with coefficients in the ring R_x and roots in $R'_x \supset R_x$. One can form the quotient ring $A := R'_x[x]/(f(x))$, naturally isomorphic to the homology of a contractible circle in our theory. Let

(60)
$$\mathcal{D} = (x_1 + x_2)(x_1 + x_3)(x_2 + x_3) = E_1 E_2 + E_3$$

be the discriminant. Consider the localization

(61)
$$R'_{\mathcal{D}} := R'_x[\mathcal{D}^{-1}], \quad A_{\mathcal{D}} := R'_{\mathcal{D}} \otimes_{R'_x} A$$

Introduce idempotents $e_1, e_2, e_3 \in A_D$:

(62)
$$e_i := \frac{(x+x_j)(x+x_k)}{(x_i+x_j)(x_i+x_k)}, \quad \{i,j,k\} = \{1,2,3\}.$$

We have

(63)
$$1 = e_1 + e_2 + e_3, \quad e_i e_j = \delta_{i,j} e_i.$$

These idempotents decompose ring $A_{\mathcal{D}}$ into the direct product

(64)
$$A_{\mathcal{D}} \cong R'_{\mathcal{D}} e_1 \times R'_{\mathcal{D}} e_2 \times R'_{\mathcal{D}} e_3 \cong R'_{\mathcal{D}} \times R'_{\mathcal{D}} \times R'_{\mathcal{D}}$$

An idempotent e_i can be visualized as floating on a facet of a foam F, in the localized theory. These idempotents allow to decompose an evaluation of a foam F with n facets into 3^n terms by summing over all ways to place each of these three idempotents onto facets of F. Each term is straightforward to compute and equals zero unless the idempotents define a Tait coloring (an admissible coloring) of F.

Idempotent e_i bears a close relation to an anchor point labeled *i*. The anchor point *p* on a facet *f* contributes the term $\sqrt{f'(x_{c(f)})} = \sqrt{(x_{c(f)} + x_j)(x_{c(f)} + x_k)}$ to the evaluation $\langle F, c \rangle$. The square of this term is either 0 (if $i \neq c(f)$) or the denominator of e_i , if i = c(f), for any coloring *c* of *F*.

Comparing e_i and an anchor point p labeled i, when coloring c associates color $c(f) \neq i$ to the facet f carrying e_i or p, both evaluations are zero. When c(f) = i, the idempotented dot e_i contributes 1 to the evaluation, while the anchor point contributes $\sqrt{f'(x_i)}$. Denominator of e_i is $f'(x_i)$.

One can try to unify e_i and anchor points p by considering anchor lines and circles L in \mathbb{R}^3 possibly intersecting a foam F. Intersection points (anchor points) carry labels $i \in \{1, 2, 3\}$ and a circle anchor points labeled i is the idempotent e_i . Then a "small" circle intersecting a facet f at two points, both labeled i, can also be converted into e_i . Notice that once e_i are allowed, integrality is lost and an evaluation of such a foam may contain denominators which are products of $x_i + x_j$.

For a different generalization, instead of a single line $L \subset \mathbb{R}^3$ consider a 1-manifold Lproperty embedded in \mathbb{R}^3 , say a finite union of lines and circles, possibly knotted. All anchor points (intersection points with L) on a foam F carry labels, with the usual contribution to the evaluation, as in formula (40). Integrality Theorem 4.12 still holds for such generalized evaluation. In particular, given k points on a plane, one can define various state spaces for webs Γ embedded in the plane and disjoint from these marked points. Also note that for $k \geq 2$ punctures, bipartite graphs are in general not reducible, which makes it harder to understand corresponding state spaces in the oriented SL(3) case.



(A) A Type 1 anchor point marked \circ and carry- (B) A Type 2 anchor point marked * with label ing no label. $i \in \{1, 2, 3\}.$

FIGURE 13. Type 1 and Type 2 anchor points.

Remark 3.20. A handle next to but disjoint from an anchor line can be written as a sum of three lower genus terms intersecting the line, see equation (52), which follows from the formula

$$m \circ \Delta(1) = (x_1 + x_2)(x_1 + x_3) + (x_1 + x_2)(x_2 + x_3) + (x_1 + x_3)(x_2 + x_3) = f'(x_1) + f'(x_2) + f'(x_3)$$

3.5. Unlabeled anchor points and bigon decomposition. Direct sum decompositions for webs Γ containing a bigon, triangle, or square face which do not contain the puncture are given in Proposition 3.15. On the other hand, Proposition 3.16 describes how to simplify a web containing an innermost non-contractible circle. In order to have direct sum decompositions for more general regions containing the puncture we introduce additional types of intersections of the anchor line L with a foam and modify the evaluation $\langle - \rangle$.

In addition to anchor points, which carry labels in $\{1, 2, 3\}$ as in Definition 3.2, we allow finitely many transverse intersections of L with a foam F away from the singular graph s(F), and we do not require labels. We will call the usual (labeled) anchor points Type 2, and the new (unlabeled) anchor points Type 1. In the figures, we denote Type 2 anchor points by an asterisk * as usual, along with a label in $\{1, 2, 3\}$, and Type 1 anchor points will be indicated by a small unshaded circle \circ . Figure 13 illustrates the convention. Let $p_1(F)$ and $p_2(F)$ denote the set of Type 1 and Type 2 anchor points, respectively (using the notation in Section 3.1, $p(F) = p_2(F)$). The definition of admissible coloring remains the same.

We modify the evaluation in the presence of Type 1 points as follows. Let $c \in \operatorname{adm}(F)$. For $p \in p_1(F)$ lying on some facet $f \in f(F)$, let c(p) := c(f) denote the coloring of the facet on which p lies. Also recall that for $i \in \{1, 2, 3\}$, we write i', i'' and j, k to denote the two complementary elements, so that $\{1, 2, 3\} = \{i, j, k\} = \{i, i', i''\}$.

Define

(65)
$$\widetilde{Q}_{\circ}(F,c) = \prod_{p \in p_1(F)} \sqrt{x_{c(p)'} + x_{c(p)''}},$$

(66)
$$P_{\circ}(F,c) = P(F,c) \cdot \widetilde{Q}_{\circ}(F,c),$$

(67)
$$\langle F, c \rangle_{\circ} = \frac{P_{\circ}(F, c)}{Q(F, c)},$$

(68)
$$\langle F \rangle_{\circ} = \sum_{c \in \operatorname{adm}(F)} \langle F, c \rangle_{\circ}.$$

where P(F,c) and Q(F,c) are as defined in (40) and (41). In other words, a Type 1 point p on an *i*-colored facet contributes a factor of $\sqrt{x_j + x_k}$ to the evaluation $\langle F, c \rangle_{\circ}$.

Remark 3.21. Type 1 intersection points are related to the triangle decoration from [KR21, Section 4.1]. Precisely, the contribution of a Type 1 point p to the square root in (65) equals the inverse of placing a triangle decoration on the facet where p lies. See relation (71), as well as Remark 3.1 for a related discussion.

Note that Type 1 intersection point contributes half the degree of a Type 2 point to the degree of the evaluation and, thus, to the degree of a cobordism represented by a foam with boundary.

Example 3.22. Consider a 2-sphere F carrying d dots and intersecting L in two Type 1 anchor points, as shown in (69).



(69)

(70)

For $1 \leq i \leq 3$, let $c_i \in \operatorname{adm}(F)$ color F by i. Then

$$\langle F, c_i \rangle_{\circ} = \frac{x_i^d (x_j + x_k)}{(x_i + x_j)(x_i + x_k)} , \langle F \rangle_{\circ} = \langle F, c_1 \rangle_{\circ} + \langle F, c_2 \rangle_{\circ} + \langle F, c_3 \rangle_{\circ} = \frac{x_1^d (x_2 + x_3)^2 + x_2^d (x_1 + x_3)^2 + x_3^d (x_1 + x_2)^2}{(x_1 + x_2)(x_1 + x_3)(x_2 + x_3)} = \frac{x_1^d (x_2^2 + x_3^2) + x_2^d (x_1^2 + x_3^2) + x_3^d (x_1^2 + x_2^2)}{(x_1 + x_2)(x_1 + x_3)(x_2 + x_3)}.$$

Thus $\langle F \rangle_{\circ} = 0$ if d = 0, 2, and $\langle F \rangle_{\circ} = 1$ if d = 1. For $d \geq 3$, the last expression above equals the ratio of the antisymmetrizer with exponent (d, 2, 0) and antisymmetrizer with exponent (2, 1, 0) (up to adding signs, which does not matter in characteristic 2). Thus $\langle F \rangle_{\circ}$ equals the Schur function $s_{\lambda}(x_1, x_2, x_3)$ for the partition $\lambda = (d - 2, 1, 0)$ when $d \geq 3$.

Example 3.23. Consider a 2-sphere F carrying d dots and intersecting L in one Type 1 anchor point and one Type 2 anchor point, as shown in (70).



Then F has one admissible coloring, and

$$\langle F \rangle_{\circ} = \frac{x_i^d \sqrt{(x_i + x_j)(x_i + x_k)(x_j + x_k)}}{(x_i + x_j)(x_i + x_k)} = \frac{x_i^d \sqrt{x_j + x_k}}{\sqrt{(x_i + x_j)(x_i + x_k)}}$$

From Example 3.23 we see that the evaluation $\langle F \rangle_{\circ}$ in general has denominators and square roots, so we can only conclude that

$$\langle F \rangle_{\circ} \in \widetilde{R}_{\circ} := \mathbb{k}[x_1, x_2, x_3, (x_1 + x_2)^{-1/2}, (x_2 + x_3)^{-1/2}, (x_1 + x_3)^{-1/2}]$$

Note that \widetilde{R}_{\circ} is a subring of \widetilde{R}''_x , see Section 3.1 and diagram (38).

We use \widetilde{R}_{\circ} as the ground ring of the theory. Evaluations of closed anchored foams F with two types of anchor points belong to this ring. We define the state space $\langle \Gamma \rangle_{\circ}$ of a trivalent graph $\Gamma \subset \mathcal{P}$ using this evaluation and following the general recipe of Section 3.3. The state space is a graded \widetilde{R}_{\circ} -module, but, due to the presence of invertible elements $(x_i + x_j)^{1/2}$ of degree 1, grading carries little information, and for many purposes one can downsize and consider the degree zero part $\langle \Gamma \rangle_{\circ}^{0}$ of the state space, which is a module over the degree 0 subring $\widetilde{R}_{\circ}^{0}$ of \widetilde{R}_{\circ} .

This theory is functorial and foams with top and bottom boundary and anchor points of those two different types induce maps between the corresponding state spaces. Various direct sum decompositions that hold for the unoriented SL(3) theory $\langle -\rangle$ hold for this theory as well.

We also have local relations involving Type 1 intersection points.

Lemma 3.24. The local relations³ (71), (72), (73), and (74) hold for the theory $\langle - \rangle_{\circ}$.



³To clarify relation (72): the first term on the right-hand side of the equality has a Type 1 anchor point on each of two front-facing half-bubbles, while the second term has a Type 1 anchor point on each of the two back-facing half-bubbles.



Proof. Relation (71) is straightforward and left to the reader. Let us verify relation (72). Denote by F the foam on the left-hand side of the equality, and denote by F^1 and F^2 the two foams on the right-hand side. There is a natural identification $\operatorname{adm}(F^1) = \operatorname{adm}(F^2)$.

Let $c \in \operatorname{adm}(F^1)$ be a coloring in which the front two half-bubble facets are differently colored, say the top front half-bubble is colored j, the bottom front half-bubble is colored k, and the remaining "big" facet is colored i. Continue to denote by $c \in \operatorname{adm}(F^2)$ the corresponding coloring of F^2 . The top Type 1 intersection point of F^1 contributes $\sqrt{x_i + x_j}$, while the to $\langle F^1, c \rangle$ and the bottom Type 1 intersection point of F^1 contributes $\sqrt{x_i + x_j}$, while the contributions of these points to $\langle F^2, c \rangle$ are reversed. Thus in characteristic two we have

$$\langle F^1, c \rangle + \langle F^2, c \rangle = 0.$$

Next, the admissible colorings of F are in natural bijection with the admissible colorings of F^1 (and of F^2) in which the front half-bubbles of F^1 are colored the same. Let $c \in \operatorname{adm}(F)$, and let $c' \in \operatorname{adm}(F^1) \cong \operatorname{adm}(F^2)$ denote the corresponding colorings. Suppose that c' colors the front half-bubbles of F^1 by j, the "big" facet by i, and the back half-bubbles by k. Then

$$\left\langle F^{1},c'\right\rangle = \frac{x_{i}+x_{k}}{x_{j}+x_{k}}\left\langle F,c
ight
angle \quad ext{and} \quad \left\langle F^{2},c'
ight
angle = \frac{x_{i}+x_{j}}{x_{j}+x_{k}}\left\langle F,c
ight
angle,$$

from which we obtain

$$\langle F, c \rangle = \langle F^1, c' \rangle + \langle F^2, c' \rangle,$$

which completes the proof of relation (72).

We now address the relation (73). Let G denote the foam on the left-hand side of the equation, and let G' denote the foam on the right-hand side. Let $c \in \operatorname{adm}(G)$, and assume c colors the "big" facet of G by i, the front bubble by j, and the back bubble by k. Let $c' \in \operatorname{adm}(G)$ denote the coloring which is identical to c except the front and back bubbles are colored by k and j, respectively. Let $c' \in \operatorname{adm}(G')$ denote the coloring of G' in which the depicted facet is colored i, and the remaining facets are colored according to c (equivalently, c'). We claim that

$$\langle G, c \rangle + \langle G, c' \rangle = \langle G', c'' \rangle,$$



FIGURE 14. Isomorphisms which simplify a bigon region containing the puncture, for the theory $\langle - \rangle_{o}$. In the top map, the top foam has a Type 1 point on the front half-bubble, and the bottom foam has a Type 1 point on the back half-bubble. In the bottom map, the first foam has a Type 1 point on the front half-bubble, and the second foam has a Type 1 point on the back half-bubble.

which completes the proof. To verify the above equality, observe that

$$\langle G, c \rangle = \frac{x_i + x_k}{x_j + x_k} \langle G', c'' \rangle$$
 and $\langle G, c' \rangle = \frac{x_i + x_j}{x_j + x_k} \langle G', c'' \rangle$

The proof of relation (74) is similar and left to the reader.

The previous lemma allows us to simplify the state space $\langle \Gamma \rangle_{\circ}$ assigned to a web $\Gamma \subset \mathcal{P}$ with a bigon region containing the puncture.

Proposition 3.25. The two maps shown in Figure 14 are mutually inverse isomorphisms between state spaces of graphs in the theory $\langle - \rangle_{\circ}$.

Proof. This follows from the relations in Lemma 3.24.

4. Oriented SL(3) anchored homology

In this section we recall oriented SL(3) foams, which were introduced in [Kho04] in the context of sl(3) link homology. An equivariant analogue was defined in [MV07], see also [MN08, Cla09, MPT14, Mac09, Rob13a] for various aspects of SL(3) foams and link homology. In Section 4.1 we define an evaluation of oriented SL(3) foams via colorings in the style of Robert-Wagner [RW20] and show in Theorem 4.22 that our evaluation agrees with that

of [MV07]. In Section 4.2 we deform the evaluation in the presence of the anchor line L. In Theorem 4.12 we show that our evaluation is always a polynomial.

To avoid introducing new notation, in this section we will reuse the notation for various rings from Section 3.

- $R'_x = \mathbb{Z}[x_1, x_2, x_3]$ is the ring of polynomials in three variables.
- $R_x = \mathbb{Z}[E_1, E_2, E_3]$ the subring of R'_x that consists of symmetric polynomials in x_1, x_2, x_3 , with generators E_i being elementary symmetric polynomials:

$$E_1 = x_1 + x_2 + x_3,$$

$$E_2 = x_1 x_2 + x_1 x_3 + x_2 x_3,$$

$$E_3 = x_1 x_2 x_3.$$

- $R''_x = R'_x[(x_1 x_2)^{-1}, (x_2 x_3)^{-1}, (x_1 x_3)^{-1}]$ is a localization of R'_x given by inverting $x_i - x_j$, for $1 \le i < j \le 3$.
- R̃'_x = R'_x[√x₁ x₃, √x₂ x₃, √x₁ x₃] is the extension of R'_x obtained by introducing square roots of √x_i x_j, for 1 ≤ i < j ≤ 3.
 R̃''_x = R̃'_x[(x₁ x₂)⁻¹, (x₂ x₃)⁻¹, (x₁ x₃)⁻¹] is a suitable localization of the ring R̃'_x.

All five of these rings are graded by setting $\deg(x_1) = \deg(x_2) = \deg(x_3) = 2$. Inclusions of the above rings are summarized in the following diagram.

4.1. Oriented SL(3) foams and their evaluations. We begin by recalling the definition of oriented SL(3) foams from [Kho04, Section 3.2].

Definition 4.1. A (closed) oriented SL(3) pre-foam F consists of the following data.

- An orientable surface F' with connected components F_1, \ldots, F_k and a partition of the boundary components of F' into triples. The underlying CW structure of F is obtained by identifying the three circles in each triple. The image of the three circles in each triple becomes a single circle in F, called a *singular circle*. The image of the surfaces F_i are called *facets*. Three facets meet at each singular circle.
- For each singular circle Z, we fix a cyclic ordering of the three facets meeting at Z. There are two possible choices of cyclic ordering for each Z.
- Each facet may carry some number of dots, which are allowed to float freely along the facet but cannot cross singular circles.

A oriented SL(3) foam is a pre-foam as above equipped with an embedding into \mathbb{R}^3 , along with an orientation on each facet such that any two of the three facets meeting at each singular circle are incompatibly oriented, as shown in Figure 15a. Each singular circle Zacquires an induced orientation, see Figure 15b. This induced orientation on Z specifies a cyclic ordering of the three facets meeting at Z by following the left-hand rule, Figure 15c, and we require this to match the cyclic ordering specified by the pre-foam F.

Note that unlike unoriented foams considered in Section 3, the oriented SL(3) pre-foams in the present section do not contain singular vertices. When there is no risk of confusion



(A) Orientations of three facets (B) The induced orientation of a (C) The induced cyclic ordering. meeting at a singular circle.

FIGURE 15

between the foams introduced in the Definition 4.1 and those of Section 3, in this section we will simply write (pre-)foam rather than oriented SL(3) (pre-)foam.

For a pre-foam F, let $\Theta(F)$ denote the set of its singular circles and $\theta(F) = |\Theta(F)|$ the number of singular circles. Each $Z \in \Theta(F)$ has a neighborhood homeomorphic to the product of a circle \mathbb{S}^1 and a tripod. Let f(F) denote the set of facets of F. We use the definitions of pre-admissible and admissible colorings of pre-foams and foams from Section 3 in the present situation. For a pre-foam F, $\operatorname{adm}(F)$ denotes the set of admissible colorings of F. Note that if F is a foam, every pre-admissible coloring is also admissible.

Fix a pre-foam F and an admissible coloring $c \in \operatorname{adm}(F)$. For $1 \leq i \neq j \leq 3$, bicolored surfaces $F_{ij}(c)$ consist of all facets colored i or j; each $F_{ij}(c)$ is a closed, orientable surface. For $1 \leq i \leq 3$, let $F_i(c)$ be the surface consisting of all facets of F which are colored i by c; the surface $F_i(c)$ is orientable and has $\theta(F)$ boundary components. Denote by $\overline{F}_i(c)$ the closed surface obtained by gluing disks along boundary components of $F_i(c)$. We have

(76)
$$\chi(\overline{F}_i(c)) = \chi(F_i(c)) + \theta(F), \ 1 \le i \le 3$$
$$\chi(F_{ij}(c)) = \chi(F_i(c)) + \chi(F_j(c)), \ 1 \le i < j \le 3.$$

The three facets meeting at each singular circle are colored by i, j, k, where as before we use i, j, k to denote the three elements of $\{1, 2, 3\}$. We now define quantities $\theta^{\pm}(c), \theta_{ij}^{\pm}(c)$ associated with the set of singular circles $\Theta(F)$ and the admissible coloring c.

Definition 4.2. Let F be a pre-foam with admissible coloring c, and let $1 \le i < j \le 3$. A singular circle $Z \in \Theta(F)$ is *positive* with respect to (i, j) if the cyclic ordering of the colors of the three facets meeting at Z is $(i \ k \ j)$. If F is a foam, then an equivalent formulation is as follows: when looking along the orientation of Z with the facet colored k drawn below, the *i*-colored facet is to the left of the *j*-colored facet. Otherwise, we say Z is negative with respect to (i, j). See Figure 16a for a pictorial definition. Let $\theta_{ij}^+(c)$ (resp. $\theta_{ij}^-(c)$) denote the number of positive (resp. negative) circles with respect to (i, j). We have

$$\theta_{ij}^+(F,c) + \theta_{ij}^-(F,c) = \theta(F).$$

We say that a singular circle Z is *positive* with respect to c if the colors of the three facets meeting at Z are (1 2 3) in the cyclic ordering, and otherwise Z is negative, see Figures 16b and 16c. Let $\theta^+(F,c)$ (resp. $\theta^-(F,c)$) denote the number of positive (resp. negative) circles in F with respect to c. We have

(77)
$$\theta^+(F,c) + \theta^-(F,c) = \theta(F).$$

We will often omit F from the notation and simply write θ , $\theta_{ij}^{\pm}(c)$, and $\theta^{\pm}(c)$.



(A) A positive (i, j)-circle, where (B) A positive singular circle. (C) A negative singular circle. i < j.

FIGURE 16

We now define the evaluations $\langle F, c \rangle$ and $\langle F \rangle$. For a pre-foam $F, c \in \text{adm}(F)$, and $1 \leq i \leq 3$, let $d_i(c)$ denote the number of dots on facets colored *i*. Define

(78)
$$P(F,c) = \prod_{i=1}^{3} x_i^{d_i(c)}$$

(79)
$$Q(F,c) = \prod_{1 \le i < j \le 3} (x_i - x_j)^{\chi(F_{ij}(c))/2}$$

(80)
$$s(F,c) = \sum_{i=1}^{3} i\chi(\overline{F}_i(c))/2 + \sum_{1 \le i < j \le 3} \theta_{ij}^+(c)$$

Set

(81)
$$\langle F, c \rangle = (-1)^{s(F,c)} \frac{P(F,c)}{Q(F,c)},$$

(82)
$$\langle F \rangle = \sum_{c \in \operatorname{adm}(F)} \langle F, c \rangle.$$

A priori, the evaluations $\langle F, c \rangle$ and $\langle F \rangle$ lie in the ring R''_x (see diagram (75)).

In what follows, we use the symbol \equiv to mean equality modulo 2. Note that

(83)
$$\sum_{i=1}^{3} i\chi(\overline{F}_i(c))/2 \equiv \frac{\chi(\overline{F}_1(c)) + \chi(\overline{F}_3(c))}{2},$$

since $\chi(\overline{F}_2(c))$ is even. Moreover, from (76) we obtain

(84)
$$\sum_{i=1}^{3} i\chi(\overline{F}_{i}(c))/2 \equiv \theta + \sum_{i=1}^{3} i\chi(F_{i}(c))/2$$

Lemma 4.1. For a pre-foam F and $c \in adm(F)$, we have

$$\sum_{1 \le i < j \le 3} \theta_{ij}^+(c) \equiv \theta^+(c)$$

It follows that

(85)
$$s(F,c) \equiv \sum_{i=1}^{3} i\chi(F_i(c))/2 + \theta^-(c).$$

Proof. Let $Z \in \Theta(F)$. Observe that if Z is positive with respect to c, then it contributes only to $\theta_{13}^+(c)$. Likewise, if Z is negative then it contributes to both $\theta_{12}^+(c)$ and $\theta_{23}^+(c)$ but not to $\theta_{13}^+(c)$, which verifies the first equality. The second equality follows from equations (84) and (77).

Example 4.2. Let F be a 2-sphere \mathbb{S}^2 with d dots. For $1 \leq i \leq 3$, let $c_i \in \operatorname{adm}(F)$ color F by i. We have

$$\begin{split} \langle F \rangle &= \langle F, c_1 \rangle + \langle F, c_2 \rangle + \langle F, c_3 \rangle \\ &= -\frac{x_1^d}{(x_1 - x_2)(x_1 - x_3)} + \frac{x_2^d}{(x_1 - x_2)(x_2 - x_3)} - \frac{x_3^d}{(x_1 - x_3)(x_2 - x_3)} \\ &= \frac{-x_1^d(x_2 - x_3) + x_2^d(x_1 - x_3) - x_3^d(x_1 - x_2)}{(x_1 - x_2)(x_2 - x_3)(x_1 - x_3)} \\ &= -s_{(d-2,0,0)}(x_1, x_2, x_3) = -h_{d-2}(x_1, x_2, x_3) = -\sum_{i+j+k=d-2} x_1^i x_2^j x_3^k, \end{split}$$

where $s_{(d-2,0,0)}(x_1, x_2, x_3)$ is the Schur function of the partition (d-2, 0, 0), and $h_{d-2}(x_1, x_2, x_3)$ is the complete symmetric function of degree d-2. In particular $\langle F \rangle = 0$ if d = 0 or d = 1, and $\langle F \rangle = -1$ if d = 2.

Example 4.3. Let F be the theta foam shown in (86).



Given any $c \in \operatorname{adm}(F)$, each capped-off surface $\overline{F}_i(c)$ and each bicolored surface $F_{ij}(c)$ is a 2-sphere. In particular,

$$s(F,c) \equiv \theta^+(c)$$

For $\sigma \in S_3$, let $c(\sigma) \in \operatorname{adm}(F)$ denote the coloring which colors the top facet by $\sigma(1)$, the middle facet by $\sigma(2)$, and the bottom facet by $\sigma(3)$. We have

$$\langle F \rangle = \sum_{\sigma \in S_3} \langle F, c(\sigma) \rangle = \frac{\sum_{\sigma \in S_3} (-1)^{\theta^+(c(\sigma))} x_{\sigma(1)}^{d_1} x_{\sigma(2)}^{d_2} x_{\sigma(3)}^{d_3}}{(x_1 - x_2)(x_1 - x_3)(x_2 - x_3)},$$

and moreover

$$\theta^+(c(\sigma)) \equiv |\sigma|,$$

where $|\sigma|$ is the length of σ .

Therefore if $d_1 \ge d_2 \ge d_3$, we have

$$\langle F \rangle = s_{(d_1 - 2, d_2 - 1, d_3)}(x_1, x_2, x_3),$$

the Schur function with partition $(d_1 - 2, d_2 - 1, d_3)$. In particular, $\langle F \rangle = 0$ if d_1, d_2, d_3 are not distinct. If d_1, d_2, d_3 are distinct and $d_1 + d_2 + d_3 \leq 3$, then up to cyclic permutation

there are two choices, for which the evaluation is recorded in (87).



The symmetric group S_3 naturally acts on $\operatorname{adm}(F)$ and on the five rings in the diagram (75). The following lemma is analogous to [RW20, Lemma 2.16].

Lemma 4.4. Let F be a pre-foam, $c \in \operatorname{adm}(F)$, and $\sigma \in S_3$. Then

 $\sigma(\langle F, c \rangle) = \langle F, \sigma(c) \rangle \,.$

Proof. We may assume that σ is a transposition $(i \ i + 1)$ for i = 1, 2. We have

$$\sigma(P(F,c)) = P(F,\sigma(c)), \qquad \sigma(Q(F,c)) = (-1)^{\chi(F_{i(i+1)}(c))/2}Q(F,\sigma(c)).$$

Let $k \in \{1, 2, 3\} \setminus \{i, i + 1\}$. Note that a singular circle Z is positive with respect to c if and only if Z is negative with respect to $\sigma(c)$, so

$$\theta^+(c) + \theta^+(\sigma(c)) = \theta = \theta^-(c) + \theta^-(\sigma(c)).$$

Moreover, we have

$$F_i(c) = F_{i+1}(\sigma(c)), \ F_{i+1}(c) = F_i(\sigma(c)), \ F_k(c) = F_k(\sigma(c)).$$

Therefore

$$\begin{split} s(F,c) - s(F,\sigma(c)) &= \frac{\chi(F_{i+1}(c)) - \chi(F_i(c))}{2} + \theta^-(c) - \theta^-(\sigma(c)) \\ &\equiv \frac{\chi(F_{i+1}(c)) - \chi(F_i(c))}{2} + \theta \\ &\equiv \frac{\chi(F_{i+1}(c)) + \chi(F_i(c))}{2} \\ &\equiv \frac{\chi(F_{i(i+1)}(c))}{2}, \end{split}$$

which completes the proof.

Corollary 4.5. The evaluation $\langle F \rangle$ is a symmetric rational function.

Later we will prove that $\langle F \rangle$ is in fact a polynomial, see Corollary 4.13.

Lemma 4.6. Let $i \in \{1,2\}$, let F be a pre-foam, and let $c \in \operatorname{adm}(F)$ be an admissible coloring. Suppose $c' \in \operatorname{adm}(F)$ is obtained from c by an (1,2)-Kempe move along a surface $\Sigma \subset F_{12}(c)$. Then

$$s(F,c) \equiv s(F,c') + \frac{\chi(\Sigma)}{2}$$

Proof. Note that this is analogous to [RW20, Lemma 2.19]. Letting $\theta(\Sigma)$ denote the number of seam circles on Σ , we have

$$\theta^{-}(c) + \theta^{-}(c') \equiv \theta(\Sigma) \equiv \chi(F_1(c) \cap \Sigma).$$

Note also that

$$\chi(F_1(c)) - \chi(F_1(c')) = \chi(F_1(c) \cap \Sigma) - \chi(F_2(c) \cap \Sigma), \chi(F_2(c)) - \chi(F_2(c')) = \chi(F_2(c) \cap \Sigma) - \chi(F_1(c) \cap \Sigma).$$

We compute:

$$s(F,c) - s(F,c') \equiv \frac{\chi(F_1(c)) - \chi(F_1(c'))}{2} + \frac{2(\chi(F_2(c)) - \chi(F_2(c')))}{2} + \theta(\Sigma)$$
$$\equiv \frac{\chi(F_2(c) \cap \Sigma) - \chi(F_1(c) \cap \Sigma)}{2} + \chi(F_1(c) \cap \Sigma)$$
$$\equiv \frac{\chi(\Sigma)}{2}.$$

4.2. Oriented anchored SL(3) foams and their evaluations. In this section we introduce (oriented) anchored SL(3) foams and their evaluations.

Definition 4.3. An oriented anchored SL(3) foam F is an oriented foam $F' \subset \mathbb{R}^3$ that may intersect the anchor line L at finitely many points away from the singular circles of F', so that each intersection point belongs to some facet of F', and moreover these intersections are required to be transverse. Denote by $p(F) = F' \cap L$ the set of intersection points (anchor points) of F. The anchor points carry labels in $\{1, 2, 3\}$; that is, F comes equipped with a fixed map

$$\ell \colon p(F) \to \{1, 2, 3\}.$$

Fix an anchored foam F and an admissible coloring c of the underlying foam F'. Each anchor point $p \in p(F)$ lying on a facet f inherits a color c(p) := c(f). As in Section 3, we say that c is an admissible coloring of the anchored foam F if for each $p \in p(F)$, the color of pequals the label of p, that is, $c(p) = \ell(p)$. Denote by adm(F) the set of admissible colorings of F.

For $i \in \{1, 2, 3\}$, let i', i'' denote the complementary elements, so that $\{i, i', i''\} = \{1, 2, 3\}$. Define the evaluations

(88)
$$\langle F, c \rangle = (-1)^{s(F,c)} \frac{P(F,c)}{Q(F,c)} \left(\prod_{p \in p(F)} (-1)^{c(p)-1} (x_{c(p)} - x_{\ell(p)'}) (x_{c(p)} - x_{\ell(p)''}) \right)^{1/2}$$

(89)
$$\langle F \rangle = \sum_{c \in \operatorname{adm}(F)} \langle F, c \rangle ,$$

where P(F,c), Q(F,c) and s(F,c) are as defined in (78), (79), and (80), respectively.

Let us explain the square root in equation (88). We have $c(p) = \ell(p)$ for every anchor point $p \in p(F)$. If p is labeled i, then it contributes

$$(-1)^{i-1}(x_i - x_j)(x_i - x_k)$$

to the product under the square root. More concretely, the product of the two terms under the square root, for a fixed anchor point p, is equal to

$$(x_1 - x_2)(x_1 - x_3)$$
 if $c(p) = 1$,
 $(x_1 - x_2)(x_2 - x_3)$ if $c(p) = 2$,
 $(x_1 - x_3)(x_2 - x_3)$ if $c(p) = 3$.

Let an(i) be the number of anchor points p with c(p) = i. Then for $1 \le i < j \le 3$ the sum an(i) + an(j) is even, which follows from Proposition 3.7.

We define the square root as the product

(90)
$$\widetilde{Q}(F,c) := \prod_{1 \le i < j \le 3} (x_i - x_j)^{(\mathrm{an}(i) + \mathrm{an}(j))/2}$$

and rewrite formula (88) as

(91)
$$\langle F, c \rangle := (-1)^{s(F,c)} \frac{P(F,c)Q(F,c)}{Q(F,c)}$$
$$= (-1)^{s(F,c)} P(F,c) \prod_{1 \le i < j \le 3} (x_i - x_j)^{(\operatorname{an}(i) + \operatorname{an}(j) - \chi(F_{ij}(c)))/2}.$$

Note that $\widetilde{Q}(F,c)$ depends only on the labels of anchor points and not on the coloring c, as long as c respects labels of anchor points (otherwise, the evaluation is 0). Consequently, it can also be denoted $\widetilde{Q}(F)$. Alternatively, it may be useful to allow more general colorings c, with $\widetilde{Q}(F,c) = 0$ for c not compatible with the labels of anchor points.

Recall diagram (75) and the surrounding discussion for notations of various rings. The above formula implies the following proposition.

Proposition 4.7. The evaluation $\langle F, c \rangle$ is an element of R''_x .

Remark 4.8. As discussed in Remark 3.2, if c is an admissible coloring of the underlying foam F' but not of the anchored foam F, then the evaluation (88) is still well-defined and equal to zero. Even if we don't restrict the notion of admissible colorings of an anchored foam to those which color anchor points according to their labels, additional terms in the evaluation will each be 0, not contributing anything.

Example 4.9. Let F be a 2-sphere \mathbb{S}^2 carrying d dots and intersecting L twice. Then $\langle F \rangle = 0$ unless both anchor points are labeled by $i \in \{1, 2, 3\}$. In this case, there is one admissible coloring c which colors F by i. We see that $s(F, c) \equiv i$, and the evaluation is

$$\langle F \rangle = (-1)^i x_i^d.$$

Example 4.10. Consider the theta foam F whose facets each intersect L exactly once, shown in (92). There is one admissible coloring c, and we have

$$\langle F \rangle = \langle F, c \rangle = \begin{cases} x_i^{d_1} x_j^{d_2} x_k^{d_3} & \text{if } (i, j, k) = (1, 3, 2) \text{ or a cyclic permutation,} \\ -x_i^{d_1} x_j^{d_2} x_k^{d_3} & \text{if } (i, j, k) = (1, 2, 3) \text{ or a cyclic permutation.} \end{cases}$$



The symmetric group S_3 acts on all five of the rings in diagram (75). Recall also that S_3 acts on the set of admissible colorings of an un-anchored foam (i.e., those considered in Section 4.1). However, for an anchored foam $F, c \in \text{adm}(F)$, and $\sigma \in S_3$, the coloring $\sigma(c)$ is in general not admissible for F.

Consider instead the anchored foam $\sigma(F)$ defined as follows. The underlying foam of $\sigma(F)$ agrees with the underlying foam of F. If anchor points of F are labeled by $\ell: p(F) \rightarrow \{1,2,3\}$, then the anchor points of $\sigma(F)$ are labeled by $\sigma(l): p \mapsto \sigma(\ell(p))$. Note that σ provides a bijection $\operatorname{adm}(F) \cong \operatorname{adm}(\sigma(F))$ via $c \mapsto \sigma(c)$. The following lemma says that the evaluations $\langle F \rangle$ and $\langle \sigma(F) \rangle$ differ by a sign, and moreover the sign depends only on σ and on labels of anchor points of F.

Lemma 4.11. For an anchored foam $F, c \in adm(F)$, and $\sigma \in S_3$, we have

$$\sigma\left(\langle F, c \rangle\right) = (-1)^{\varepsilon(F,\sigma)} \left\langle \sigma(F), \sigma(c) \right\rangle,$$

where

(92)

(93)
$$\varepsilon(F,\sigma) = \sum_{\substack{1 \le i < j \le 3, \\ \sigma(i) > \sigma(j)}} \frac{\operatorname{an}(i) + \operatorname{an}(j)}{2}.$$

It follows that

$$\sigma\left(\langle F\rangle\right) = (-1)^{\varepsilon(F,\sigma)}\left\langle\sigma(F)\right\rangle.$$

Proof. By Lemma 4.4, we have

$$\sigma\left((-1)^{s(F,c)}\frac{P(F,c)}{Q(F,c)}\right) = (-1)^{s(\sigma(F),\sigma(c))}\frac{P(\sigma(F),\sigma(c))}{Q(\sigma(F),\sigma(c))}$$

It is clear that

$$\sigma(\widetilde{Q}(F)) = (-1)^{\varepsilon(F,\sigma)} \widetilde{Q}(\sigma(F)),$$

and the first equality follows. For the second equality, we have

$$\sigma\left(\langle F \rangle\right) = \sum_{c \in \operatorname{adm}(F)} \sigma\left(\langle F, c \rangle\right)$$
$$= (-1)^{\varepsilon(F,\sigma)} \sum_{c \in \operatorname{adm}(F)} \langle \sigma(F), \sigma(c) \rangle$$
$$= (-1)^{\varepsilon(F,\sigma)} \langle \sigma(F) \rangle.$$

For $1 \leq i \neq j \leq 3$, consider the ring

$$R_{ij}'' := R_x'[(x_i - x_k)^{-1}, (x_j - x_k)^{-1}].$$

Each R''_{ij} is a subring of R''_x . A permutation $\sigma \in S_3$ sends R''_{ij} isomorphically onto $R''_{\sigma(i)\sigma(j)}$. We are now ready for the main result of this section.

Theorem 4.12. The evaluation $\langle F \rangle$ of an anchored foam is an element of R'_x , the polynomial ring in variables x_1, x_2, x_3 .

Proof. The proof is similar to that of [KR21, Theorem 2.17] and [RW20, Proposition 2.18]. By Lemma 4.11, it suffices to show that $\langle F \rangle \in R''_{12}$ for any anchored foam F. This is because we may take a permutation $\sigma \in S_3$ sending 1 to i and 2 to j, and consider the anchored foam $\sigma^{-1}(F)$. Then $\langle \sigma^{-1}(F) \rangle \in R''_{12}$ implies that

$$\pm \langle F \rangle = \pm \left\langle \sigma(\sigma^{-1}(F)) \right\rangle = \pm \sigma \left(\left\langle \sigma^{-1}(F) \right\rangle \right) \in R_{ij}'',$$

where the first equality comes from Lemma 4.11. It follows that

$$\langle F \rangle \in R_{12}'' \cap R_{23}'' \cap R_{13}'' = R_x'$$

Let us show that $\langle F \rangle \in R_{12}''$. Partition $\operatorname{adm}(F)$ into equivalence classes as follows. For $c \in \operatorname{adm}(F)$, the class C_c containing c consists of colorings obtained from c by performing a sequence of (1, 2) Kempe moves along surfaces in $F_{12}(c)$ which are disjoint from L. If $F_{12}(c)$ has n connected components, $k \geq 0$ of which are disjoint from L, then C_c consists of 2^k elements. We will show that

$$\sum_{c' \in C_c} \langle F, c' \rangle \in R_{12}'',$$

which will conclude the proof.

Write $\Sigma := F_{12}(c)$ as a disjoint union

$$\Sigma = \Sigma' \cup \Sigma_1 \cup \cdots \cup \Sigma_k,$$

where each Σ_a , $a = 1, \ldots, k$ is connected and disjoint from L, and where each component of Σ' intersects L. For i = 1, 2 and $a = 1, \ldots, k$, let $t_i(a)$ denote the number of dots on *i*-colored facets (according to c) of Σ_a , and let t_3 denote the number of dots on 3-colored facets (according to c) of F. We claim that

$$(94) \quad \sum_{c'\in C_c} \langle F, c' \rangle = \frac{x_3^{t_3} \cdot \prod_{a=1}^k \left(x_1^{t_1(a)} x_2^{t_2(a)} + (-1)^{\chi(\Sigma_a)/2} x_2^{t_1(a)} x_1^{t_2(a)} \left(\frac{(x_1 - x_3)^{\ell_{\Sigma_a}(c)/2}}{(x_2 - x_3)} \right) \right) \cdot \widetilde{Q}(F)}{(x_1 - x_2)^{\chi(\Sigma)/2} (x_1 - x_3)^{\chi(F_{13}(c))/2} (x_2 - x_3)^{\chi(F_{23}(c))/2}}$$

where

• $\ell_{\Sigma_a}(c) \in 2\mathbb{Z}$ is an even integer such that

 $\chi(F_{13}(c')) = \chi(F_{13}(c)) - \ell_{\Sigma_a}(c) \text{ and } \chi(F_{23}(c')) = \chi(F_{23}(c)) + \ell_{\Sigma_a}(c)$

for the coloring $c' \in C_c$ which is obtained from c by a (1, 2) Kempe move along Σ_a . See [KR21, Lemma 2.12 (3)] for details regarding this integer.

• $\tilde{Q}(F)$ is the contribution from the anchor points of F, equation (90).

To verify the claimed equality, expand the product to obtain 2^k terms, each of which corresponds to one of the 2^k colorings in C_c . That the sign is correct follows from Lemma 4.6.

Finally, we argue that $(x_1 - x_2)^{\chi(\Sigma)/2}$ divides the numerator of (94). Positive contributions to $\chi(\Sigma)$ come from 2-sphere components of Σ . Each Σ_a which is a 2-sphere contributes

one to the exponent $\chi(\Sigma)/2$. On the other hand, the corresponding factor in the product in the numerator of (94) is divisible by $x_1 - x_2$. The remaining positive contributions to $\chi(\Sigma)/2$ come from 2-sphere components of Σ' . Such a component Σ_0 contains at least two anchor points, each labeled 1 or 2, so the contribution from Σ_0 can be cancelled with terms in $\widetilde{Q}(F)$.

Corollary 4.13. If F is a pre-foam or a foam which is disjoint from L, then $\langle F \rangle \in R_x$, the ring of symmetric polynomials in x_1, x_2, x_3 .

Proof. This follows from Lemma 4.11 and Theorem 4.12.

4.3. Skein relations. In this section we record several local relations involving oriented anchored SL(3) foams.

Lemma 4.14. The local relations (95), (96), (97), and (98) hold for anchored foams. Seam lines are drawn in bold in relation (98) to clarify the picture.

Proof. Proofs of these four relations are similar to Propositions 2.33, 2.22, 2.23, and 2.24 in [KR21], respectively, with the caveat that we must keep track of the sign (80). Moreover, S_3 symmetry is used in [KR21] to simplify the calculations. Anchor points and their labels are the same for the foams depicted in each of these four relations, so Lemma 4.11 implies that we may use S_3 symmetry in a similar manner.

Y, *i*

We verify relation (96) and leave the remaining three relations to the reader. Let F denote the foam appearing on the left-hand side of the equality. The six foams on the right-hand

side are identical except for placement of dots. We denote them by G^1, \ldots, G^6 , so that the relation reads

$$\langle F \rangle = -\left(\left\langle G^1 \right\rangle + \left\langle G^2 \right\rangle + \left\langle G^3 \right\rangle \right) + E_1\left(\left\langle G^4 \right\rangle + \left\langle G^5 \right\rangle \right) - E_2\left\langle G^6 \right\rangle.$$

Admissible colorings of G^1, \ldots, G^6 are in canonical bijection. For $c \in \operatorname{adm}(G^1)$, let

$$\langle G, c \rangle := -\left(\left\langle G^1, c \right\rangle + \left\langle G^2, c \right\rangle + \left\langle G^3, c \right\rangle \right) + E_1\left(\left\langle G^4, c \right\rangle + \left\langle G^5, c \right\rangle \right) - E_2\left\langle G^6, c \right\rangle.$$

There are two types of colorings of G^1 : those which color the two depicted disks the same, and those which color them differently. Those of the first type are in canonical bijection with colorings of F.

Suppose $c \in \operatorname{adm}(G^1)$ colors both disks the same color, say i, and denote by $c \in \operatorname{adm}(G^2) \cong \cdots \cong \operatorname{adm}(G^6)$ and $c' \in \operatorname{adm}(F)$ the corresponding colorings. We will show that $\langle F, c' \rangle = \langle G, c \rangle$. We may assume i = 1. Then

$$\langle G^1, c \rangle = \langle G^2, c \rangle = \langle G^3, c \rangle = x_1^2 \langle G^6, c \rangle, \quad \langle G^4, c \rangle = \langle G^5, c \rangle = x_1 \langle G^6, c \rangle,$$

which yields

$$\langle G, c \rangle = -3x_1^2 \langle G^6, c \rangle + 2E_1 x_1 \langle G^6, c \rangle - E_2 \langle G^6, c \rangle = -(x_1 - x_2)(x_1 - x_3) \langle G^6, c \rangle.$$

To compare this with $\langle F, c' \rangle$, observe that

$$\chi(F_1(c')) + 2 = \chi(G_1^6(c)), \quad \chi(F_2(c')) = \chi(G_2^6(c)), \quad \chi(F_3(c')) = \chi(G_3^6(c)),$$

which implies $s(F, c') \equiv s(G, c) + 1$. Moreover, we have

$$\chi(F_{12}(c')) + 2 = \chi(G_{12}^6(c)), \quad \chi(F_{13}(c')) + 2 = \chi(G_{13}^6(c)), \quad \chi(F_{23}(c')) = \chi(G_{23}^6(c)).$$

Therefore

$$\langle G^6, c \rangle = -\frac{\langle F, c' \rangle}{(x_1 - x_2)(x_1 - x_3)}$$

which verifies $\langle F, c' \rangle = \langle G, c \rangle$.

To complete the proof, suppose that c colors the top depicted disk by i and the bottom disk by j, with $i \neq j$. We have

$$\begin{split} \left\langle G^{1},c\right\rangle &=x_{i}^{2}\left\langle G^{6},c\right\rangle ,\ \left\langle G^{2},c\right\rangle &=x_{i}x_{j}\left\langle G^{6},c\right\rangle ,\ \left\langle G^{3},c\right\rangle &=x_{j}^{2}\left\langle G^{3},c\right\rangle ,\\ \left\langle G^{4},c\right\rangle &=x_{i}\left\langle G^{6},c\right\rangle ,\ \left\langle G^{5},c\right\rangle &=x_{j}\left\langle G^{6},c\right\rangle . \end{split}$$

Therefore $\langle G, c \rangle = 0$, which concludes the proof.

Lemma 4.15. Let F be an anchored foam. Denote by $F_{n,m}$ the anchored foam obtained from F by adding a bubble (disjoint from L) to some facet in F, with the two new facets carrying n and m dots respectively, such that the facet with n dots directly precedes the facet with m dots in the cyclic ordering. Let F_n denote the foam obtained from F by adding n dots to the same facet. This is shown in (99). Then



Remark 4.16. The relations in Lemmas 4.14 and 4.15 also hold for pre-foams.

Similar to the SL(2) and unoriented SL(3) setting, for oriented SL(3) foams we allow shifted dots $(i) = \bullet - x_i \ (1 \le i \le 3)$ on a facet.

$$(i)$$
 = $-x_i$

They must be disjoint from L and are allowed to float freely on their facets but cannot cross seam lines.

Lemma 4.17. The local relations shown in (100), (101), and (102) hold.



In the last equation we assume j < k.

Proof. We verify equation (100) and leave the remaining relations to the reader. The argument is similar to that of relation (50) in Lemma 3.12, so we will be brief. Let F denote the foam on the left-hand side, and let G^1, G^2, G^3 denote the three foams on the right-hand side, with superscript corresponding to labels of the anchor points. For $1 \le i \le 3$, let $\operatorname{adm}_i(F)$ consist of all admissible colorings of F which color the depicted tube by i. There is a natural bijection $\operatorname{adm}_i(F) \cong \operatorname{adm}(G^i)$.



FIGURE 17. The orientations at each trivalent vertex of an oriented SL(3) web must be either all outgoing or all incoming.

Given $c \in \operatorname{adm}_i(F)$, let $c' \in \operatorname{adm}(G^i)$ denote the corresponding coloring. Arguing as in the proof of Lemma 3.12, we obtain

$$\langle F, c \rangle = \pm \langle G^i, c' \rangle.$$

It remains to show that the above sign is equal to $(-1)^i$. We have

$$\chi(F_j(c)) = \chi(G_j^i(c')), \ \chi(F_k(c)) = \chi(G_k^j(c')), \ \chi(F_i(c)) = \chi(G_i^i(c')) - 2, \ \theta^{\pm}(F,c) = \theta^{\pm}(G^i,c'),$$

so $s(F,c) \equiv s(G^i,c') + i$ as needed.

4.4. State spaces. In this section we define state spaces associated to oriented SL(3) webs. Much of this is analogous to notions in Section 3.3.

Definition 4.4. An oriented SL(3) web is a planar trivalent graph $\Gamma \subset \mathcal{P}$ in the punctured plane, which may have closed loops with no vertices. Moreover, edges and loops of Γ carry orientations such that each vertex is either a source or a sink, as shown in Figure 17. In this section we will simply write web rather than oriented SL(3) web.

The definition of an anchored foam with boundary in the oriented setting is analogous to that of Definition 3.4. The singular graph of a foam with boundary V is a union of finitely many arcs (with boundary in $\mathbb{R}^2 \times \{0, 1\}$) and circles (disjoint from $\mathbb{R}^2 \times \{0, 1\}$). Intersection points of V with $L_{[0,1]}$ (anchor points) must be disjoint from the singular graph and carry labels in $\{1, 2, 3\}$. Facets of V are required to carry orientations satisfying the convention in Figure 15a near singular points. As usual, we will use the left-hand rule to specify these orientations and cyclic orderings by orienting each singular circle and arc, as shown in Figures 15b and 15c.

As in Section 3.3, let $\partial_i V := V \cap (\mathbb{R}^2 \times \{0\})$ for i = 0, 1. The orientation of facets of V induces an orientation on $\partial_0 V$ and $\partial_1 V$ via the convention in Figure 18. We view V as a cobordism from the oriented web $\partial_0 V$ to the oriented web $\partial_1 V$. Composition WV of foams V, W with $\partial_1 V = \partial_0 W$ is defined in the natural way.

Denote by $p(V) = V \cap L_{[0,1]}$ the set of anchor points of V and by |d(V)| the number of dots. The degree of V is defined to be

(103)
$$\deg(V) = 2(|d(V)| + |p(V)| - \chi(V)) + \chi(\partial V).$$

Degree is clearly additive under composition and is compatible with the grading on R'_x , in the sense that if V is a closed foam, then $\deg(V) = \deg(\langle V \rangle)$.

As in Definition 2.1, by an *annular foam* we mean a foam (with boundary) which is disjoint from L. The composition of two annular foams is again annular.

There is an involution ω defined by reflecting a foam with boundary through $\mathbb{R}^2 \times \{1/2\}$. We have $\partial_1 V = \partial_0(\omega(V))$ and $\partial_0 V = \partial_1(\omega(V))$ for any foam with boundary V. Given a web



FIGURE 18. Our convention for the induced orientation on the webs $\partial_0 V$ (bottom) and $\partial_1 V$ (top).





FIGURE 19. Local relations for state spaces of oriented SL(3) webs, where the depicted regions do not contain the puncture.

 $\Gamma \subset \mathcal{P}$, let $Fr(\Gamma)$ denote the free R'_x -module generated by foams with boundary V from the empty web to Γ (that is, $\partial_0 V = \emptyset$, $\partial_1 V = \Gamma$). Define a bilinear form

(-,-): $\operatorname{Fr}(\Gamma) \times \operatorname{Fr}(\Gamma) \to R'_x$

by $(V, W) = \omega(V)W$. This bilinear form is symmetric since $\langle F \rangle = \langle \omega(F) \rangle$ for any closed foam F. The state space $\langle \Gamma \rangle$ is the quotient of $Fr(\Gamma)$ by the kernel

$$\ker((-,-)) = \{ x \in \operatorname{Fr}(\Gamma) \mid (x,y) = 0 \text{ for all } y \in \operatorname{Fr}(\Gamma) \}$$

of the bilinear form,

$$\langle \Gamma \rangle := \operatorname{Fr}(\Gamma) / \ker((-,-)).$$

The state space $\langle \Gamma \rangle$ inherits the grading from $Fr(\Gamma)$ since (-, -) is degree-preserving. A foam with boundary V from Γ_0 to Γ_1 naturally induces a map

$$\langle V \rangle : \langle \Gamma_0 \rangle \to \langle \Gamma_1 \rangle$$

of degree deg(V), defined by sending the equivalence class of a basis element $U \in \operatorname{Fr}(\Gamma_0)$ to the equivalence class of $VU \in \operatorname{Fr}(\Gamma_1)$. This assignment is functorial with respect to composition of foams, $\langle WV \rangle = \langle W \rangle \langle V \rangle$ for composable V, W.

Lemma 4.18. The three local isomorphisms shown in Figure 19 hold.

Proof. The arguments for relations (a), (b), and (c) of the figure are analogous to Propositions 7, 9, and 8, respectively, of [Kho04]. The relevant relations are Lemmas 4.14 and 4.15. \Box

Proposition 4.19. Let $\Gamma \subset \mathcal{P}$ be a web with a non-contractible circle C which bounds a disk in $\mathbb{R}^2 \setminus \Gamma$, and let $\Gamma' = \Gamma \setminus C$ be the web obtained by removing C. Then there is an isomorphism

$$\langle \Gamma \rangle \cong \langle \Gamma' \rangle \oplus \langle \Gamma' \rangle \oplus \langle \Gamma' \rangle$$

given by the maps shown in (104) (orientation of the circle is omitted).



Proof. This follows from Example 4.9 and the neck-cutting relation (100).

Theorem 4.20. For any web $\Gamma \subset \mathcal{P}$, the state space $\langle \Gamma \rangle$ is a free graded R'_x -module of rank equal to the number of Tait colorings of Γ . Moreover, if Γ is contractible, then the graded rank of $\langle \Gamma \rangle$ equals the Kuberberg polynomial [Kup96] of Γ , normalized as in [Kho04, Section 2].

Proof. Lemma 3.17 (1) guarantees that we can reduce $\langle \Gamma \rangle$ to a direct sum of empty webs by recursively applying the local isomorphisms in Lemma 4.18 and Proposition 4.19. It is then clear that the rank equals the number of Tait colorings.

If Γ is contractible then $\langle \Gamma \rangle$ can be simplified using only the isomorphisms in Lemma 4.18. Upon taking graded ranks, these isomorphisms recover the recursive relations for computing the Kuperberg polynomial.

Theorem 4.20 does not address the graded rank of state spaces of non-contractible webs. These may be computed recursively. As a special case, if Γ consists of *n* contractible and *m* non-contractible circles, then $\langle \Gamma \rangle$ is free of graded rank $3^m (q^2 + 1 + q^{-2})^n$.

Given a web $\Gamma \subset \mathcal{P}$, we can forget the puncture and the anchor line L and apply the universal construction to the evaluation (82). Precisely, let $\operatorname{Fr}(\Gamma)_{\text{forget}}$ denote the free R_x -module generated by all foams with boundary Γ (forgetting the anchor line). By Corollary 4.13, we can define the bilinear form (-, -): $\operatorname{Fr}(\Gamma)_{\text{forget}} \times \operatorname{Fr}(\Gamma)_{\text{forget}} \to R_x$ and the corresponding state space $\langle \Gamma \rangle_{\text{forget}}$ in the usual way. Thus we obtain state spaces for webs in \mathbb{R}^2 , functorial with respect to foams in $\mathbb{R}^2 \times [0, 1]$. These state spaces and maps induced by foams are graded via equation (103), where |p(V)| = 0.



FIGURE 20. The oriented intersection number between a facet and L.

Proposition 4.21. For a contractible web $\Gamma \subset \mathcal{P}$, there is a degree-preserving isomorphism

$$\langle \Gamma \rangle \cong \langle \Gamma \rangle_{\text{forget}} \,,$$

natural with respect to foams with contractible boundary and which are disjoint from L.

Proof. This follows from Theorem 4.20.

On the other hand, Mackaay-Vaz [MV07] define an evaluation $\langle - \rangle_{\rm MV}$ for oriented SL(3) pre-foams and use it to define an equivariant (also called *universal*) version of the sl(3) link homology introduced in [Kh004]. They work over the ground ring $\mathbb{Z}[a, b, c]$ and associate a state space $\langle \Gamma \rangle_{\rm MV}$ to each web $\Gamma \subset \mathbb{R}^2$ via the universal construction applied to their pre-foam evaluation $\langle - \rangle_{\rm MV}$. To compare with our situation, identify $\mathbb{Z}[a, b, c]$ with the ring $R_x = \mathbb{Z}[E_1, E_2, E_3]$ of symmetric functions in x_1, x_2, x_3 via a ring isomorphism φ defined by $\varphi(a) = E_1, \varphi(b) = -E_2, \varphi(c) = E_3$.

Theorem 4.22. For any closed pre-foam F, we have

 $\left\langle F\right\rangle =\varphi\left(\left\langle F\right\rangle _{\mathrm{MV}}\right).$

It follows that there are isomorphisms $\langle \Gamma \rangle_{\text{forget}} \cong \langle \Gamma \rangle_{\text{MV}} \otimes_{\mathbb{Z}[a,b,c]} R_x$ for any web $\Gamma \subset \mathbb{R}^2$, natural with respect to maps induced by foams with boundary.

Proof. The evaluation $\langle - \rangle_{\rm MV}$ is defined by applying the local relations (3D), (CN), (S), and (Θ) in [MV07, Section 2.1] to reduce any foam to an element of $\mathbb{Z}[a, b, c]$. Under the change of variables $a \mapsto E_1$, $b \mapsto -E_2$, $c \mapsto E_3$, these four relations hold for our evaluation $\langle - \rangle$ by relation (95), relation (96), Example 4.2, and Example 4.3. The statement follows.

As in the SL(2) and unoriented SL(3) setting considered earlier in the paper, we can define an additional grading on oriented SL(3) foams and state spaces. Define the abelian group

(105)
$$\Lambda = \mathbb{Z}w_1 \oplus \mathbb{Z}w_2 \oplus \mathbb{Z}w_3/(w_1 + w_2 + w_3),$$

on three generators and one relation. Λ is a free abelian group of rank two.

Orient the anchor line L from bottom to top. For an anchored foam V with boundary and $p \in p(V)$ an anchor point lying on some facet f, let $s(p) \in \{\pm 1\}$ denote the oriented intersection number between f and L (s(p) does not depend on the label of p), see Figure 20 for the convention. Define the *annular degree of* V to be

(106)
$$\operatorname{adeg}(V) = \sum_{p \in p(V)} s(p) w_{\ell(p)} \in \Lambda.$$

Proposition 4.23. If F is a closed anchored foam with an admissible coloring c, then adeg(F) = 0.

Proof. The proof is similar to that of Proposition 3.7. The intersection of F with a generic half-plane that bounds L is an oriented web Γ with boundary points on L. An admissible coloring c of F induces a Tait coloring of Γ . The boundary points (one-valent vertices) of Γ are colored according to their label. The sum in (106) may be rewritten as the sum of terms $\pm(w_1 + w_2 + w_3) = 0$ over all trivalent vertices of Γ , where the sign is +1 if all edges are incoming and -1 if all edges are outgoing. Each *i*-colored inner edge e of Γ bounds two trivalent vertices and contributes $\pm(w_i - w_i) = 0$ since e is oriented towards one of its boundary vertices and away from the other. The remaining edges, with one or both endpoints on L, contribute precisely $\operatorname{adeg}(F)$.

Let $\Gamma \subset \mathcal{P}$ be an (annular oriented) SL(3) web. An anchored foam $F \subset \mathbb{R}^3_-$ with $\partial F = \Gamma$ has a well-defined degree $\operatorname{adeg}(F) \in \Lambda$ via (106). Furthermore, we equip the coefficient ring R'_x with an Λ -grading, with all elements of degree 0. This makes free R'_x -module $\operatorname{Fr}(\Gamma)$ into an Λ -graded module, and Proposition 4.23 implies that the kernel of the bilinear form on $\operatorname{Fr}(\Gamma)$ is Λ -graded as well. Consequently, the grading descends to an Λ -grading on the state space $\langle \Gamma \rangle$. A foam V with boundary induces a map $\langle V \rangle : \langle -\partial_0 \Gamma \rangle \to \langle \partial_1 \Gamma \rangle$ which changes adeg by $\operatorname{adeg}(V)$. If V has no anchor points, it induces an annular degree 0 map between the state spaces of its boundaries. The state space of a contractible web is concentrated in annular degree zero.

A-grading on $\langle \Gamma \rangle$ is the analogue of grading on finite-dimensional SL(3) representations by the weight lattice. In fact, in the non-equivariant version of our construction, where all x_i 's are set to 0 upon closed foam evaluation (and state spaces are defined accordingly, over a ground field rather than the ring R'_x), the state space $\langle \Gamma \rangle$ is naturally an \mathfrak{sl}_3 -representation. We also refer the reader to Queffelec-Rose [QR18] for the construction of sutured annular \mathfrak{sl}_n -homology, with state spaces of annular webs carrying an \mathfrak{sl}_n -action. In the equivariant case, it is not clear how to define an \mathfrak{sl}_3 -action or what's the substitute for it.

Denote by AFoam_{or} the category whose objects consist of oriented SL(3) webs in \mathcal{P} and whose morphisms are R'_x -linear combinations of anchored cobordisms between webs. Morphism spaces in this category are triply graded via (qdeg, adeg). The state space construction assembles into a functor

$$\langle - \rangle$$
: AFoam_{or} $\rightarrow R'_x - g_3 \mod$

landing in the category of triply-graded R'_x -modules.

This functor respects the trigradings on the hom spaces in the two categories. Restricting to the subcategory of annular cobordisms and their linear combinations, the induced maps have annular degree 0.

4.5. Annular SL(3)-link homology. Let $L \subset \mathbb{A} \times [0, 1]$ be a link in the thickened annulus. Projecting onto $\mathbb{A} \times \{0\} = \mathbb{A}$ and identifying the interior of \mathbb{A} with the punctured plane \mathcal{P} , we obtain a link diagram $D \subset \mathcal{P}$. Following [Kho04, MV07], form the cube of resolutions of D, with each resolution web drawn in the punctured plane. Applying the functor $\langle - \rangle$: AFoam_{or} $\rightarrow R'_x - g_3$ mod yields a chain complex C(D) of $\mathbb{Z} \oplus \Lambda$ -graded R'_x -modules, with the \mathbb{Z} -grading given by deg, equation (103), and the Λ -grading given by adeg, equation (106). Degree shifts in the cube of resolutions are applied only to the \mathbb{Z} -degree deg. Diagrams in \mathcal{P} representing isotopic annular links are related by Reidemeister moves away from the puncture. Proofs of Reidemeister invariance in [MV07] are local, and all local relations (away from L) on foams in [MV07] also hold for our evaluation $\langle - \rangle$ by (96), Example 4.2, and Example 4.3. It follows that the chain homotopy class of C(D) is an invariant of the annular link L. We define equivariant annular SL(3) homology as cohomology groups H(C(D)).

Moreover, foams between webs appearing in the cube of resolutions are disjoint from L. Thus the differential preserves annular degree throughout the complex. Consequently, equivariant annular sl(3) link homology carries a homological grading as well as an internal $\mathbb{Z} \oplus \Lambda$ -grading (deg, adeg). Cohomology groups H(C(D)) are trigraded R'_x -modules.

References

- [Akh20] Rostislav Akhmechet. Equivariant annular Khovanov homology. 2020. Preprint: arXiv:2008.00577.
- [APS04] Marta M. Asaeda, Józef H. Przytycki, and Adam S. Sikora. Categorification of the Kauffman bracket skein module of *I*-bundles over surfaces. *Algebr. Geom. Topol.*, 4:1177–1210, 2004.
- [BG15] John A. Baldwin and J. Elisenda Grigsby. Categorified invariants and the braid group. Proc. Amer. Math. Soc., 143(7):2801–2814, 2015.
- [BHMV95] Christian Blanchet, Nathan Habegger, Gregor Masbaum, and Pierre Vogel. Topological quantum field theories derived from the Kauffman bracket. *Topology*, 34(4):883–927, 1995.
- [BN05] Dror Bar-Natan. Khovanov's homology for tangles and cobordisms. *Geom. Topol.*, 9:1443–1499, 2005.
- [Boo19] David Boozer. Computer bounds for Kronheimer-Mrowka foam evaluation. 2019. To appear in *Experimental Mathematics*. Preprint: arXiv:1908.07133.
- [BPW19] Anna Beliakova, Krzysztof K. Putyra, and Stephan M. Wehrli. Quantum link homology via trace functor I. Invent. Math., 215(2):383–492, 2019.
- [Cla09] David Clark. Functoriality for the \mathfrak{su}_3 Khovanov homology. Algebr. Geom. Topol., 9(2):625–690, 2009.
- [GLW17] J. Elisenda Grigsby, Anthony M. Licata, and Stephan M. Wehrli. Annular Khovanov-Lee homology, braids, and cobordisms. *Pure Appl. Math. Q.*, 13(3):389–436, 2017.
- [GLW18] J. Elisenda Grigsby, Anthony M. Licata, and Stephan M. Wehrli. Annular Khovanov homology and knotted Schur-Weyl representations. *Compos. Math.*, 154(3):459–502, 2018.
- [GN14] J. Elisenda Grigsby and Yi Ni. Sutured Khovanov homology distinguishes braids from other tangles. *Math. Res. Lett.*, 21(6):1263–1275, 2014.
- [Kho00] Mikhail Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000.
- [Kho04] Mikhail Khovanov. sl(3) link homology. Algebr. Geom. Topol., 4:1045–1081, 2004.
- [KK20] Mikhail Khovanov and Nitu Kitchloo. A deformation of Robert-Wagner foam evaluation and link homology. 2020. Preprint: https://arxiv.org/abs/2004.14197.
- [KM16] Peter B. Kronheimer and Tomasz S. Mrowka. Exact triangles for SO(3) instanton homology of webs. J. Topol., 9(3):774–796, 2016.
- [KM19a] Peter B. Kronheimer and Tomasz S. Mrowka. A deformation of instanton homology for webs. Geom. Topol., 23(3):1491–1547, 2019.
- [KM19b] Peter B. Kronheimer and Tomasz S. Mrowka. Tait colorings, and an instanton homology for webs and foams. J. Eur. Math. Soc. (JEMS), 21(1):55–119, 2019.
- [KR08] Mikhail Khovanov and Lev Rozansky. Matrix factorizations and link homology. *Fund. Math.*, 199(1):1–91, 2008.
- [KR20a] Mikhail Khovanov and Louis-Hadrien Robert. Conical sl(3) foams. 2020. Preprint: arXiv:2011.11077.
- [KR20b] Mikhail Khovanov and Louis-Hadrien Robert. Link homology and Frobenius extensions II. 2020. Preprint: arXiv:2005.08048.
- [KR21] Mikhail Khovanov and Louis-Hadrien Robert. Foam evaluation and Kronheimer-Mrowka theories. Adv. Math., 376:107433, 2021.
- [Kup96] Greg Kuperberg. Spiders for rank 2 Lie algebras. Comm. Math. Phys., 180(1):109–151, 1996.
- [Mac09] Marco Mackaay. sl(3)-foams and the Khovanov-Lauda categorification of quantum sl(k). 2009. Preprint: arXiv:0905.2059.

R. AKHMECHET AND M. KHOVANOV

- [MN08] Scott Morrison and Ari Nieh. On Khovanov's cobordism theory for \mathfrak{su}_3 knot homology. J. Knot Theory Ramifications, 17(9):1121–1173, 2008.
- [MPT14] Marco Mackaay, Weiwei Pan, and Daniel Tubbenhauer. The \mathfrak{sl}_3 -web algebra. Math. Z., 277(1-2):401-479, 2014.
- [MV07] Marco Mackaay and Pedro Vaz. The universal sl₃-link homology. *Algebr. Geom. Topol.*, 7:1135–1169, 2007.
- [QR18] Hoel Queffelec and David E. V. Rose. Sutured annular Khovanov-Rozansky homology. *Trans. Amer. Math. Soc.*, 370(2):1285–1319, 2018.
- [QRS18] Hoel Queffelec, David E. V. Rose, and Antonio Sartori. Annular Evaluation and Link Homology. 2018. Preprint: arXiv:1802.04131.
- [QW21] Hoel Queffelec and Paul Wedrich. Khovanov homology and categorification of skein modules. *Quantum Topol.*, 12(1):129–209, 2021.
- [Rob13a] Louis-Hadrien Robert. Sur l'homologie \$13 des enchevêtrement; algèbre de Khovanov-Kuperberg. PhD thesis, Université Paris 7 – Denis Diderot, July 2013.
- [Rob13b] Lawrence P. Roberts. On knot Floer homology in double branched covers. *Geom. Topol.*, 17(1):413–467, 2013.
- [RW16] David E. V. Rose and Paul Wedrich. Deformations of colored \mathfrak{sl}_N link homologies via foams. Geom. Topol., 20(6):3431–3517, 2016.
- [RW20] Louis-Hadrien Robert and Emmanuel Wagner. A closed formula for the evaluation of foams. *Quantum Topol.*, 11(3):411–487, 2020.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE VA 22904-4137 *Email address:* ra5aq@virginia.edu

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK NY 10027 *Email address:* khovanov@math.columbia.edu