

LOCAL COHOMOLOGY FOR GORENSTEIN HOMOLOGICALLY SMOOTH DG ALGEBRAS

X.-F. MAO AND H. WANG

ABSTRACT. In this paper, we introduce the theory of local cohomology and local duality to Noetherian connected cochain DG algebras. We show that the notion of local cohomology functor can be used to detect the Gorensteinness of a homologically smooth DG algebra. For any Gorenstein homologically smooth locally finite DG algebra \mathcal{A} , we define a group homomorphism $\text{Hdet} : \text{Aut}_{dg}(\mathcal{A}) \rightarrow k^\times$, called the homological determinant. As applications, we present a sufficient condition for the invariant DG subalgebra \mathcal{A}^G to be Gorenstein, where \mathcal{A} is a homologically smooth DG algebra such that $H(\mathcal{A})$ is a Noetherian AS-Gorenstein graded algebra and G is a finite subgroup of $\text{Aut}_{dg}(\mathcal{A})$. Especially, we can apply this result to DG down-up algebras and non-trivial DG free algebras generated in two degree-one elements.

INTRODUCTION

Local cohomology is an indispensable tool in almost all branches of analytic and algebraic geometry as well as in commutative and combinatorial algebra. Since the flourishing of non-commutative algebraic geometry in 1990s, people have done a lot to develop a theory of local cohomology in the context of non-commutative cases. In [Yek], Yekutieli introduced balanced dualizing complexes for non-commutative graded algebras by using local cohomology functors. After his work, Jørgensen [Jor1] generalized the theory of local cohomology and local duality theorem to non-commutative positively graded Noetherian algebras which are quotients of Artin-Schelter Gorenstein algebras. Later, Van den Bergh gave a more general local duality formula in [VDB]. In [JoZ], local cohomology functor was used to define homological determinants for AS-Gorenstein algebras. In the invariant theory of Hopf algebra action (and formerly group actions) on non-commutative graded algebras, the notion of homological determinant usually serves as an effective tool in precise analysis (cf. [CWZ, JoZ, JZ, KKZ1, LMZ1]).

In this paper, we are concerned with connected cochain DG k -algebras. Any connected graded algebra can be seen as a connected cochain DG algebra with zero differential. The motivation of this paper is to introduce an analogous local cohomology theory under the background of DG homological algebra. Let \mathcal{A} be a connected cochain DG algebra with maximal DG ideal $\mathfrak{m} = \mathcal{A}^{\geq 1}$. For any DG \mathcal{A} -module M , its subset

$$\Gamma_{\mathfrak{m}}(M) = \{m \in M \mid \mathcal{A}^{\geq n}m = 0, \text{ for some } n \geq 1\}$$

is a DG \mathcal{A} -submodule of M . We actually define a left exact covariant \mathcal{A} -linear functor

$$\Gamma_{\mathfrak{m}}(-) = \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, -)$$

2010 *Mathematics Subject Classification.* Primary 16E45, 16E65, 16W20, 16W50.

Key words and phrases. Gorenstein DG algebra, homologically smooth DG algebra, local cohomology, homological determinant.

from the category $\mathcal{C}(\mathcal{A})$ of left DG \mathcal{A} -modules to itself. It is natural to define its right derived functor $R\Gamma_{\mathfrak{m}}(-)$ as the local cohomology functor. We emphasize that our definitions of torsion DG modules and the local cohomology functor are quite different from the definitions in [Jor2]. Due to the differences between DG homological functors and hyper-homological functors in computation, it seems difficult to get a complete generalization of the original theorem of local duality as in [Jor1, Theorem 2.3]. For example, we don't know whether the torsion functor preserves the homotopically injective property of DG modules. However, under some mild hypothesis, we prove the following local duality theorem (see Theorem 3.1) in DG context.

Theorem A. Let M be a compact DG module over a locally finite connected cochain DG algebra \mathcal{A} . Then $R\Gamma_{\mathfrak{m}}(M)' \cong R\mathrm{Hom}_{\mathcal{A}}(M, R\Gamma_{\mathfrak{m}}(\mathcal{A})')$ in $\mathcal{D}(\mathcal{A}^p)$.

Note that we add compact condition on the DG module M , and we define the Matlis dual of a graded k -vector space V as

$$V' = \mathrm{Hom}_k(V, k) = \bigoplus_{i \in \mathbb{Z}} \mathrm{Hom}_k(V^i, k).$$

Using the duality formula of the theorem above, we give characterizations of Gorensteinness for Noetherian homologically smooth connected cochain DG algebras. This is the following theorem (see Theorem 3.6).

Theorem B. Let \mathcal{A} be a Noetherian homologically smooth connected cochain DG algebra. Then \mathcal{A} is Gorenstein if and only if $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \mathcal{A}' \otimes_{\mathcal{A}} L$ in $\mathcal{D}(\mathcal{A}^e)$, where L is an invertible DG bi-module over \mathcal{A} .

Applying the characterization of Gorensteinness for Noetherian homologically smooth connected cochain DG algebras in the theorem above, we give a proof of the following interesting theorem (see Theorem 3.9).

Theorem C. Assume that \mathcal{A} is a Noetherian homologically smooth and Gorenstein connected cochain DG algebra. Then for any $M \in \mathcal{D}^c(\mathcal{A})$, $R\Gamma_{\mathfrak{m}}(M)'$ is a compact DG \mathcal{A}^p -module. And there is a pair of contravariant equivalences of categories,

$$\mathcal{D}^c(\mathcal{A}) \begin{array}{c} \xrightarrow{R\Gamma_{\mathfrak{m}}(-)'} \\ \xleftarrow{R\Gamma_{\mathfrak{m} \circ p}(-)'} \end{array} \mathcal{D}^c(\mathcal{A}^p).$$

People have done a lot of work to extend the rich theory of commutative Gorenstein rings to DG algebras (cf. [FHT1, AF, DGI, FJ, FIJ, MW2]). In this paper, we study the Gorensteinness of invariant DG subalgebras of some nice Gorenstein DG algebras. In the literature, there have been many papers on the behavior of the Gorenstein property under the passage to the subring of invariants (cf. [Wat1, Wat2, Hin, JZ, KKZ1]). However, it is infeasible to mimic the proof there since the techniques and tools used in commutative and non-commutative graded setting are not available in DG context. For this, we need to develop some tools. Let \mathcal{A} be a Gorenstein, homologically smooth and locally finite connected cochain DG algebra. We define homological determinant for any DG algebra automorphism of \mathcal{A} in Definition 4.2. Suppose G is a finite subgroup of the automorphism group of a connected cochain DG algebra \mathcal{A} . The invariant DG subalgebra is defined to be

$$\mathcal{A}^G = \{x \in \mathcal{A} \mid g(x) = x, \text{ for all } g \in G\}.$$

Then $H(G) : \{H(g) \mid g \in G\}$ is a finite subgroup of $\mathrm{Aut}_{gr} H(\mathcal{A})$ and the graded fixed subalgebra $H(\mathcal{A})^{H(G)}$ of $H(\mathcal{A})$ is isomorphic to $H(\mathcal{A}^G)$ (see Proposition 4.4). Moreover, we prove the following theorem (see Theorem 4.5).

Theorem D. Let \mathcal{A} be a homologically smooth locally finite connected cochain DG algebra such that $H(\mathcal{A})$ is a Noetherian AS-Gorenstein graded algebra. If G is

a finite subgroup of $\text{Aut}_{dg}(\mathcal{A})$ such that the homological determinant of each $\sigma \in G$ is 1, then the fixed DG subalgebra \mathcal{A}^G is a Gorenstein DG algebra.

In addition, we find that the result in Theorem D can be successfully applied to DG down-up algebras and some non-trivial DG free algebras (see Corollary 4.6 and Corollary 4.7).

To close the introduction, let us run through some notation and conventions. Let W be a graded k -vector space. For any $j \in \mathbb{Z}$, the j -th suspension $\Sigma^j W$ of W is a graded vector space defined by $(\Sigma^j W)^i = W^{i+j}$. We assume that the reader is familiar with basics on DG algebras and DG modules. Most of the notations and conventions on them we take in this paper are same as in [MW1, MW2]. Let \mathcal{A} be a connected cochain DG algebra. We take the following notations:

- (1) $\mathcal{A}^\#$ is the underlying graded algebra of \mathcal{A} ;
- (2) $\text{Aut}_{dg}(\mathcal{A})$ is the group of all DG algebra automorphisms of \mathcal{A} ;
- (3) \mathcal{A}^p is the opposite DG algebra of \mathcal{A} ;
- (4) \mathcal{A}^e is the enveloping DG algebra of \mathcal{A} ;
- (5) $\mathfrak{m}_{\mathcal{A}}$ is the maximal DG ideal $\mathcal{A}^{>0}$ of \mathcal{A} ;
- (6) $H(\mathcal{A})$ is the cohomology graded algebra of \mathcal{A} ;
- (7) $\mathcal{C}(\mathcal{A})$ is the category of left DG modules over \mathcal{A} ;
- (8) $\mathcal{D}(\mathcal{A})$ is the derived category of $\mathcal{C}(\mathcal{A})$;
- (9) $\mathcal{D}(\mathcal{A})$ is the full triangulated subcategory of $\mathcal{D}(\mathcal{A})$ consisting of compact DG \mathcal{A} -modules.

Let z be a graded cocycle element in \mathcal{A} , we write $[z]$ for the cohomology class in $H(\mathcal{A})$ represented by z . For any $M, N \in \mathcal{C}(\mathcal{A})$ and $\sigma \in \text{Aut}_{dg}(\mathcal{A})$, We denote ${}^\sigma M$ by an object in $\mathcal{C}(\mathcal{A})$ such that ${}^\sigma M$ has the same underlying complex of k -vector spaces as M and its multiplication is defined by $a \cdot m = \sigma(a)m$ for any $a \in \mathcal{A}$ and $m \in M$. A k -linear chain map $f : M \rightarrow N$ is called a σ -linear chain map if it gives a morphism of DG \mathcal{A} -modules, $f : M \rightarrow {}^\sigma N$, i.e. if $f(am) = \sigma(a)f(m)$.

Finally, we recall some properties of DG algebras, which will be mentioned in this paper. Let \mathcal{A} be a connected cochain DG algebra.

- (1) If $\mathcal{A}^\#$ is a Noetherian graded algebra, then \mathcal{A} is called Noetherian;
- (2) If $\dim_k \mathcal{A}^i < \infty, \forall i \in \mathbb{Z}$, then \mathcal{A} is called locally finite;
- (3) If $\dim_k H(R\text{Hom}_{\mathcal{A}}(k, \mathcal{A})) = 1$, then \mathcal{A} is called left Gorenstein (cf. [FHT1]); Similarly, \mathcal{A} is called right Gorenstein, if $\dim_k H(R\text{Hom}_{\mathcal{A}^p}(k, \mathcal{A})) = 1$;
- (4) If ${}_{\mathcal{A}}k$, or equivalently ${}_{\mathcal{A}^e}\mathcal{A}$, admits a minimal semi-free resolution with a semi-basis concentrated in degree 0, then \mathcal{A} is called Koszul (cf. [HW]);
- (5) If ${}_{\mathcal{A}}k$, or equivalently the DG \mathcal{A}^e -module \mathcal{A} is compact, then we say that \mathcal{A} is a homologically smooth DG algebra (cf. [MW3, Corollary 2.7]).

Note that each Noetherian connected cochain DG algebra is locally finite. While the converse is generally not true. In general, a left Gorenstein connected cochain DG algebra may be not right Gorenstein. However, a homologically smooth DG algebra is left Gorenstein if and only if it is right Gorenstein (see [MW2, Remark 6.6]). Beside this, we have the following remark for a homologically smooth DG algebra (see [MW2]).

Remark 0.1. *Let \mathcal{A} be a homologically smooth connected cochain DG algebra and $M \in \mathcal{D}(\mathcal{A})$. Then*

- (1) $R\text{Hom}_{\mathcal{A}}(k, \mathcal{A}) \xrightarrow{\sim} \Sigma^i {}_{\mathcal{A}}k$ if and only if $R\text{Hom}_{\mathcal{A}^p}(k, \mathcal{A}) \xrightarrow{\sim} \Sigma^i {}_{\mathcal{A}}k$;
- (2) $M \in \mathcal{D}^c(\mathcal{A})$ if $\dim_k H(M) < \infty$;
- (3) $H(\mathcal{A})$ is not bounded above if \mathcal{A} is not quasi-isomorphic to k .

1. SOME BASICS ON DG MODULES AND DG ALGEBRAS

In this section, we give some lemmas on DG modules and DG algebras, which will be used in the later sections.

Lemma 1.1. *Let I be a semi-injective DG \mathcal{A}^e -module. Then I is semi-injective both as a DG \mathcal{A} -module and as a DG \mathcal{A}^{op} -module.*

Proof. Since $\text{Hom}_{\mathcal{A}}(-, I) \cong \text{Hom}_{\mathcal{A}}(-, \text{Hom}_{\mathcal{A}^e}(\mathcal{A}^e, I)) \cong \text{Hom}_{\mathcal{A}^e}(\mathcal{A} \otimes -, I)$, it is easy to check that I is a semi-injective DG \mathcal{A} -module. Similarly, we can prove that I is also a semi-injective DG \mathcal{A}^{op} -module. \square

Lemma 1.2. *Let \mathcal{A} and \mathcal{B} be two connected cochain DG algebras. Then for any DG \mathcal{A} -module P , DG \mathcal{B} -module Q and DG $\mathcal{A} \otimes \mathcal{B}$ -module N , the chain map*

$$\begin{aligned} \phi : \text{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, N) &\rightarrow \text{Hom}_{\mathcal{A}}(P, \text{Hom}_{\mathcal{B}}(Q, N)) \\ f &\mapsto \phi(f) : x \mapsto f_x : y \mapsto f(x \otimes y) \end{aligned}$$

is an isomorphism, whose inverse is

$$\begin{aligned} \psi : \text{Hom}_{\mathcal{A}}(P, \text{Hom}_{\mathcal{B}}(Q, N)) &\rightarrow \text{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, N) \\ g &\mapsto \psi(g) : x \otimes y \mapsto g(x)(y). \end{aligned}$$

Proof. For any $f \in \text{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, N)$ and $x \in P$, we should prove the following statements

- (1) $f_x \in \text{Hom}_{\mathcal{B}}(Q, N)$;
- (2) $\phi(f) \in \text{Hom}_{\mathcal{A}}(P, \text{Hom}_{\mathcal{B}}(Q, N))$;
- (3) ϕ is a chain map.

For any $a, b \in B$, $x \in P$ and $y \in Q$, we have $f_x \in \text{Hom}_{\mathcal{B}}(Q, N)$ since

$$\begin{aligned} f_x(by) &= f(x \otimes by) = f[(1 \otimes b)(x \otimes y)] \\ &= (1 \otimes b)f(x \otimes y) = (1 \otimes b)f_x(y) = bf_x(y). \end{aligned}$$

Similarly, $\phi(f)(ax) = a\phi(f)(x)$ since

$$\begin{aligned} \phi(f)(ax) &= f_{ax} : y \mapsto f(ax \otimes y) = (a \otimes 1)f(x \otimes y) \\ a\phi(f)(x) &= af_x : y \mapsto af(x \otimes y) = (a \otimes 1)f(x \otimes y). \end{aligned}$$

Furthermore,

$$\begin{aligned} [\partial_{\text{Hom}} \circ \phi(f)](x)(y) &= [\partial_{\text{Hom}_{\mathcal{B}}(Q, N)} \circ f_x - (-1)^{|f|}\phi(f)(\partial_P(x))](y) \\ &= \partial_N \circ f_x(y) - (-1)^{|f_x|}f_x[\partial_Q(y)] - (-1)^{|f|}f(\partial_P(x) \otimes y) \\ &= \partial_N(f(x \otimes y)) - (-1)^{|f|+|x|}f[x \otimes \partial_Q(y)] - (-1)^{|f|}f[\partial_P(x) \otimes y] \end{aligned}$$

and

$$\begin{aligned} [\phi \circ \partial_{\text{Hom}}(f)](x)(y) &= [\partial_{\text{Hom}}(f)]_x(y) = [\partial_N \circ f - (-1)^{|f|}f \circ \partial_{P \otimes Q}]_x(y) \\ &= \partial_N(f(x \otimes y)) - (-1)^{|f|}f[\partial_P(x) \otimes y + (-1)^{|x|}x \otimes \partial_Q(y)] \\ &= \partial_N(f(x \otimes y)) - (-1)^{|f|}f[\partial_P(x) \otimes y] - (-1)^{|f|+|x|}f[x \otimes \partial_Q(y)]. \end{aligned}$$

So ϕ is a chain map. Similarly, we can show that ψ is chain map. Furthermore,

$$\begin{aligned} \psi \circ \phi(f)(x \otimes y) &= \phi(f)(x)(y) = f_x(y) = f(x \otimes y), \\ \phi \circ \psi(g)(x \otimes y) &= \psi(g)_x(y) = \psi(g)(x \otimes y) = g(x)(y), \end{aligned}$$

for any $f \in \text{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, N)$ and $g \in \text{Hom}_{\mathcal{A}}(P, \text{Hom}_{\mathcal{B}}(Q, N))$, $x \in P$ and $y \in Q$. Thus $\phi \circ \psi = \text{id}$ and $\psi \circ \phi = \text{id}$. \square

Lemma 1.3. *Let \mathcal{A} and \mathcal{B} be two homologically smooth connected cochain DG algebras. Then*

- (1) $\mathcal{A} \otimes \mathcal{B}$ is a homologically smooth connected cochain DG algebra;
- (2) $\mathcal{A} \otimes \mathcal{B}$ is Gorenstein if both \mathcal{A} and \mathcal{B} are Gorenstein DG algebras.

Proof. (1) Let P and Q be the minimal semi-free resolutions of ${}_{\mathcal{A}}k$ and ${}_{\mathcal{B}}k$ respectively. Then $P \otimes Q$ is a minimal semi-free resolution of ${}_{\mathcal{A} \otimes \mathcal{B}}k$. By Lemma 1.2, we have

$$H(\mathrm{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, k)) \cong H(\mathrm{Hom}_{\mathcal{A}}(P, \mathrm{Hom}_{\mathcal{B}}(Q, k))).$$

Since ${}_{\mathcal{A}}k$ and ${}_{\mathcal{B}}k$ are compact by the assumption, both P and Q have a finite semi-basis. This implies that $\dim_k H(\mathrm{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, k)) < \infty$. Hence $\mathcal{A} \otimes \mathcal{B}$ is also a homologically smooth connected cochain DG algebra.

(2) By (1), $\mathcal{A} \otimes \mathcal{B}$ is homologically smooth. Let P and Q be the minimal semi-free resolutions of ${}_{\mathcal{A}}k$ and ${}_{\mathcal{B}}k$ respectively. Then $P \otimes Q$ is a minimal semi-free resolution of ${}_{\mathcal{A} \otimes \mathcal{B}}k$. We have

$$\begin{aligned} H(\mathrm{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, \mathcal{A} \otimes \mathcal{B})) &\stackrel{(a)}{\cong} H(\mathrm{Hom}_{\mathcal{A}}(P, \mathrm{Hom}_{\mathcal{B}}(Q, \mathcal{A} \otimes \mathcal{B}))) \\ &\stackrel{(b)}{\cong} H(\mathrm{Hom}_{\mathcal{A}}(P, \mathcal{A} \otimes \mathrm{Hom}_{\mathcal{B}}(Q, \mathcal{B}))) \\ &\stackrel{(c)}{\cong} H(\mathrm{Hom}_{\mathcal{A}}(P, \mathcal{A}) \otimes \mathrm{Hom}_{\mathcal{B}}(Q, \mathcal{B})) \\ &\stackrel{(d)}{\cong} H(\mathrm{Hom}_{\mathcal{A}}(P, \mathcal{A})) \otimes H(\mathrm{Hom}_{\mathcal{B}}(Q, \mathcal{B})), \end{aligned}$$

where (a) is by Lemma 1.2, (b) and (c) are obtained by the fact that both P and Q have a finite semi-basis, and (d) is by the well known Künneth formula. On the other hand, $\dim_k H(\mathrm{Hom}_{\mathcal{A}}(P, \mathcal{A})) = \dim_k H(\mathrm{Hom}_{\mathcal{B}}(Q, \mathcal{B})) = 1$ since \mathcal{A} and \mathcal{B} are Gorenstein. Thus $\dim_k H(\mathrm{Hom}_{\mathcal{A} \otimes \mathcal{B}}(P \otimes Q, \mathcal{A} \otimes \mathcal{B})) = 1$. This implies that $\mathcal{A} \otimes \mathcal{B}$ is also Gorenstein. \square

Definition 1.4. A DG \mathcal{A}^e -module L is called an invertible DG \mathcal{A} -bimodule if there is some DG \mathcal{A}^e -module L^\vee such that $L \otimes_{\mathcal{A}} L^\vee \cong L^\vee \otimes_{\mathcal{A}} L \cong \mathcal{A}$ over \mathcal{A}^e .

For any $\phi \in \mathrm{Aut}_{dg}(\mathcal{A})$ and $i \in \mathbb{Z}$, we define an invertible DG bimodule $\Sigma^i \mathcal{A}(\phi)$ as follows. As a left DG \mathcal{A} -module, $\Sigma^i \mathcal{A}(\phi)$ is the DG \mathcal{A} -module $\Sigma^i \mathcal{A}$ with a generator $e = 1$ in degree $-i$. The right multiplication is given by the rule $ea = (-1)^{i \cdot |a|} \phi(a)e$, $a \in \mathcal{A}$. One sees that $\Sigma^i \mathcal{A}(\phi) \otimes_{\mathcal{A}} \Sigma^u \mathcal{A}(\psi) \cong \Sigma^{i+u} \mathcal{A}(\phi \circ \psi)$, for any $\psi \in \mathrm{Aut}_{dg}(\mathcal{A})$.

Lemma 1.5. *Let $L \cong \Sigma^i \mathcal{A}(\phi)$ be an invertible DG bimodule over a connected cochain DG algebra \mathcal{A} , for some $i \in \mathbb{Z}$ and $\phi \in \mathrm{Aut}_{dg}(\mathcal{A})$. Then $\mathrm{Hom}_{\mathcal{A}}(L, L) \cong \mathcal{A}$ as a DG \mathcal{A}^e -module.*

Proof. We have the following isomorphisms of DG \mathcal{A}^e -modules:

$$\mathrm{Hom}_{\mathcal{A}}(L, L) \cong \mathrm{Hom}_{\mathcal{A}}(L, \mathcal{A}) \otimes_{\mathcal{A}} L \cong \mathcal{A}(\phi^{-1}) \otimes_{\mathcal{A}} \mathcal{A}(\phi) \cong \mathcal{A}.$$

\square

By [Yek, Proposition 1.10], any invertible graded bimodule over a connected graded algebra R can be denoted by $\Sigma^i R(\phi)$, for some $i \in \mathbb{Z}$ and $\phi \in \mathrm{Aut}_{gr}(R)$. We attempt to obtain a similar result in DG context. However, the following counter example indicates that not all invertible DG \mathcal{A} -bimodules have the form $\Sigma^i \mathcal{A}(\psi)$, $i \in \mathbb{Z}$ and $\psi \in \mathrm{Aut}_{dg}(\mathcal{A})$.

Example 1.6. Let $\mathcal{A} = k[x]/(x^2)$ with $\partial_{\mathcal{A}} = 0$ and $|x| = 1$. Assume that L and L^\vee are DG \mathcal{A} -bimodules such that

$$\begin{aligned} L^\# &= \mathcal{A}^\# e \text{ with } e \cdot \bar{x} = \bar{x}e, \partial_L(e) = \bar{x}e \\ L^{\vee\#} &= \mathcal{A}^\# f \text{ with } f \cdot \bar{x} = \bar{x}f, \partial_{L^\vee}(f) = -\bar{x}f. \end{aligned}$$

Then $\partial_\otimes(e \otimes f) = \bar{x}e \otimes f - e \otimes \bar{x}f = e \cdot \bar{x} \otimes f - e \otimes \bar{x}f = 0$ in $L \otimes_{\mathcal{A}} L^\vee$. One see that the morphisms

$$\begin{aligned} \theta : L \otimes_{\mathcal{A}} L^\vee &\rightarrow \mathcal{A} \\ e \otimes f &\mapsto 1_{\mathcal{A}} \end{aligned}$$

and

$$\begin{aligned} \omega : L^\vee \otimes_{\mathcal{A}} L &\rightarrow \mathcal{A} \\ f \otimes e &\mapsto 1_{\mathcal{A}} \end{aligned}$$

are both isomorphisms of DG \mathcal{A}^e -modules. Therefore, L and L^\vee are invertible DG \mathcal{A} -bimodules. However, L is not isomorphic to $\mathcal{A}(\phi)$ for any $\phi \in \text{Aut}_{dg}(\mathcal{A})$, since $\partial_L(e) = \bar{x}e \neq 0$.

Lemma 1.7. Let M a DG \mathcal{A} -module such that $\dim_k H(M) = 1$. Then $M \cong \Sigma^i_{\mathcal{A}} k$ in $\mathcal{D}(\mathcal{A})$, for some integers $i \in \mathbb{Z}$.

Proof. Since $\dim_k H(M) = 1$, we have $H(M) = \Sigma^i k$ for some integers $i \in \mathbb{Z}$. For each $j \in \mathbb{Z}$, there is a decomposition of k -vector spaces $M^j = B^j \oplus H^j \oplus C^j$ with $B^j = \text{Im } \partial^{j-1}$ and $\text{Ker } \partial^j = B^j \oplus H^j$. Define $\tau^{\geq i}(M)$ as the DG \mathcal{A} -submodule of M such that

$$[\tau^{\geq i}(M)]^j = \begin{cases} 0, & \text{if } j \leq i-1, \\ C^i \oplus H^i, & \text{if } j = i, \\ M^j, & \text{if } j > i. \end{cases}$$

The inclusion map $\iota : \tau^{\geq i}(M) \rightarrow M$ is a quasi-isomorphism of DG \mathcal{A} -modules. Let $o(\tau^{\geq i}(M))$ be the DG \mathcal{A} -submodule of $\tau^{\geq i}(M)$ such that

$$o(\tau^{\geq i}(M))^m = \begin{cases} C^i, & \text{if } m = i, \\ [\tau^{\geq i}(M)]^m & \text{if } m \neq i. \end{cases}$$

Obviously, $o(\tau^{\geq i}(M))$ is quasi-trivial. Hence the canonical morphism

$$\tau^{\geq i}(M) \rightarrow \tau^{\geq i}(M)/o(\tau^{\geq i}(M)) = \Sigma^i_{\mathcal{A}} k$$

is a quasi-isomorphism. Therefore, $M \cong \Sigma^i_{\mathcal{A}} k$ in $\mathcal{D}(\mathcal{A})$. \square

Lemma 1.8. Let M be a DG \mathcal{A}^e -module such that ${}_{\mathcal{A}}M \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ and $M_{\mathcal{A}} \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$. Then there is an invertible DG \mathcal{A} -bimodule $L = \Sigma^i \mathcal{A}(\psi)$ for some $\psi \in \text{Aut}_{dg}(\mathcal{A})$ such that $M = L \otimes_{\mathcal{A}} \mathcal{A}'$.

Proof. By the assumption, we have ${}_{\mathcal{A}}(\Sigma^i M') \cong {}_{\mathcal{A}}\mathcal{A}$ and $(\Sigma^i M')_{\mathcal{A}} \cong \mathcal{A}_{\mathcal{A}}$. Let e be the generator of ${}_{\mathcal{A}}(\Sigma^i M')$. Then for any $a \in \mathcal{A}$, there exists unique $l_a \in \mathcal{A}$ such that $e \cdot a = l_a e$. This induces a map $\phi : \mathcal{A} \rightarrow \mathcal{A}$ such that $\phi(a) = l_a$. It is straightforward to check that ϕ is an isomorphism of the graded algebra \mathcal{A} . Furthermore, ϕ is a

chain map since

$$\begin{aligned}
\partial_{\mathcal{A}}(\phi(a))e &= \partial_{\mathcal{A}}(l_a)e \\
&= \partial_{\Sigma^i M'}(l_a e) \\
&= \partial_{\Sigma^i M'}(e \cdot a) \\
&= e \cdot \partial_{\mathcal{A}}(a) \\
&= l_{\partial_{\mathcal{A}}(a)}e = \phi(\partial_{\mathcal{A}}(a))e,
\end{aligned}$$

for any $a \in \mathcal{A}$. So $\phi \in \text{Aut}_{dg} \mathcal{A}$ and $\Sigma^i M' \cong \mathcal{A}(\phi)$ as a DG \mathcal{A}^e -module. Therefore, the DG \mathcal{A} -bimodule

$$\begin{aligned}
M &\cong \Sigma^i(\mathcal{A}(\phi))' \cong \Sigma^i \text{Hom}_k(\mathcal{A} \otimes_{\mathcal{A}} \mathcal{A}(\phi), k) \\
&\cong \Sigma^i \text{Hom}_{\mathcal{A}}(\mathcal{A}(\phi), \text{Hom}_k(\mathcal{A}, k)) \\
&\cong \Sigma^i \text{Hom}_{\mathcal{A}}(\mathcal{A}(\phi), \mathcal{A} \otimes_{\mathcal{A}} \mathcal{A}') \\
&\cong \Sigma^i \text{Hom}_{\mathcal{A}}(\mathcal{A}(\phi), \mathcal{A}) \otimes_{\mathcal{A}} \mathcal{A}' \\
&\cong \Sigma^i \mathcal{A}(\phi^{-1}) \otimes_{\mathcal{A}} \mathcal{A}'.
\end{aligned}$$

Let $\psi = \phi^{-1}$ and $L = \Sigma^i \mathcal{A}(\psi)$. Then the DG \mathcal{A} -bimodule L is invertible and $M \cong L \otimes_{\mathcal{A}} \mathcal{A}'$. \square

2. LOCAL COHOMOLOGY FUNCTOR IN DG SETTING

Assume that M is a DG module over a connected cochain DG algebra \mathcal{A} . Its subset

$$\Gamma_{\mathfrak{m}}(M) = \{m \in M \mid \mathcal{A}^{\geq n} m = 0, \text{ for some } n \geq 1\}$$

is actually a DG \mathcal{A} -submodule. Indeed, for any $m \in \Gamma_{\mathfrak{m}}(M)$ and $a \in \mathcal{A}$, there exists an integer $n \geq 1$ such that $\mathcal{A}^{\geq n} m = 0$ and hence $\mathcal{A}^{\geq n}(am) \subseteq \mathcal{A}^{\geq n} m = 0$. Then $am \in \Gamma_{\mathfrak{m}}(M)$. Moreover, we have

$$0 = \partial_M(\mathcal{A}^{\geq n} m) = \partial_{\mathcal{A}}(\mathcal{A}^{\geq n})m + \mathcal{A}^{\geq n} \partial_M(m)$$

and $\partial_{\mathcal{A}}(\mathcal{A}^{\geq n})m \subseteq \mathcal{A}^{\geq n} m = 0$. So $\partial_M(m) \in \Gamma_{\mathfrak{m}}(M)$. We denote $\Gamma_{\mathfrak{m}}(M)$ by the torsion submodule of M . Let $f : M \rightarrow N$ be a morphism of DG \mathcal{A} -modules. Then $f(\Gamma_{\mathfrak{m}}(M)) \subseteq \Gamma_{\mathfrak{m}}(N)$ since f is \mathcal{A} -linear. Hence there exists a morphism of DG \mathcal{A} -modules $\Gamma_{\mathfrak{m}}(f) : \Gamma_{\mathfrak{m}} M \rightarrow \Gamma_{\mathfrak{m}} N$ which agrees with f on each element of $\Gamma_{\mathfrak{m}}(M)$. Furthermore, if $g : M \rightarrow N, h : N \rightarrow L$ are morphisms of DG \mathcal{A} -modules and $a \in \mathcal{A}$, then

$$\begin{cases} \Gamma_{\mathfrak{m}}(h \circ f) = \Gamma_{\mathfrak{m}}(h) \circ \Gamma_{\mathfrak{m}}(f), \\ \Gamma_{\mathfrak{m}}(f + g) = \Gamma_{\mathfrak{m}}(f) + \Gamma_{\mathfrak{m}}(g), \\ \Gamma_{\mathfrak{m}}(\text{id}_M) = \text{id}_{\Gamma_{\mathfrak{m}}(M)}. \end{cases}$$

Thus $\Gamma_{\mathfrak{m}}(-)$ is a covariant functor from $\mathcal{C}(\mathcal{A})$ to itself. Since the torsion functor

$$\Gamma_{\mathfrak{m}}(-) = \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, -),$$

it is left exact. By the definition of torsion functor above, it is clear that the functors $\text{Hom}_{\mathcal{A}}(M, -)$ and $\text{Hom}_{\mathcal{A}}(M, \Gamma_{\mathfrak{m}}(-))$ are naturally isomorphic, if M is a finite length DG \mathcal{A} -module. We write $R\Gamma_{\mathfrak{m}}(-)$ as the right derived functor of $\Gamma_{\mathfrak{m}}(-)$.

For any object M in $\mathcal{C}(\mathcal{A})$, its local cohomology $H_{\mathfrak{m}}(M)$ is defined to be the graded $H(\mathcal{A})$ -module $H(R\Gamma_{\mathfrak{m}}(M))$. Similarly, we have functors $\Gamma_{\mathfrak{m}^{op}}$ and $\Gamma_{\mathfrak{m}_{\mathcal{A}^e}}$ on $\mathcal{C}(\mathcal{A}^{op})$ and $\mathcal{C}(\mathcal{A}^e)$ respectively. We use the same notations for some variants of these functors. For example, $\Gamma_{\mathfrak{m}}$ defines a functor from $\mathcal{C}(\mathcal{A} \otimes \mathcal{B})$ to $\mathcal{C}(\mathcal{A} \otimes \mathcal{B})$, where \mathcal{B} is another connected cochain DG algebra.

Lemma 2.1. *Let \mathcal{A} be a homologically smooth and locally finite connected cochain DG algebra. Then $R\Gamma_{\mathfrak{m}}(-)$ commutes with direct limits.*

Proof. Since \mathcal{A} is homologically smooth, ${}_{\mathcal{A}}k$ is a compact DG \mathcal{A} -module. For any integer $n \geq 1$, $\mathcal{A}/\mathcal{A}^{\geq n}$ is finite dimensional. Hence $\mathcal{A}/\mathcal{A}^{\geq n}$ is a compact DG \mathcal{A} -module by Remark 0.1. Let F be a minimal semi-free resolution of $\mathcal{A}/\mathcal{A}^{\geq n}$. Then F admits a finite semi-basis. Hence $R\mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, -) = \mathrm{Hom}_{\mathcal{A}}(F, -)$ commutes with direct limits. This implies that $R\Gamma_{\mathfrak{m}}(-) = \varinjlim_{n \geq 1} R\mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, -)$ commutes with direct limits. \square

Lemma 2.2. *Assume that \mathcal{A} is a homologically smooth locally finite connected cochain DG algebra. If M is a direct limit of DG \mathcal{A} -modules with finite dimensional cohomology, then the canonical map*

$$\iota : R\Gamma_{\mathfrak{m}}(M) \rightarrow M$$

is a quasi-isomorphism.

Proof. By Lemma 2.1, it suffices to show that the canonical map $R\Gamma_{\mathfrak{m}}(M) \rightarrow M$ is a quasi-isomorphism, if M is a DG \mathcal{A} -module with finite dimensional cohomology. Since \mathcal{A} is homologically smooth, $k_{\mathcal{A}}$ is compact. By Remark 0.1, M' is a compact DG \mathcal{A}^p -module. There is a minimal semi-free resolution $\gamma : F_{M'} \rightarrow M'$ such that $F_{M'}$ has a finite semi-basis. This implies $\Gamma_{\mathfrak{m}}((F_{M'})') = (F_{M'})'$. Since $H(M)$ is finite dimensional, $\theta : M \rightarrow M'$ defined by $\theta(m)(f) = (-1)^{|f| \cdot |m|} f(m)$ is a quasi-isomorphism. So $\mathrm{Hom}_k(\gamma, k) \circ \theta : M \rightarrow (F_{M'})'$ is a semi-injective resolution of M . By the following commutative diagram

$$\begin{array}{ccc} H(R\Gamma_{\mathfrak{m}}(M)) & \xrightarrow{H(\iota)} & H(M) \\ \downarrow = & & \downarrow H(\mathrm{Hom}_k(\gamma, k) \circ \theta) \\ H(\Gamma_{\mathfrak{m}}((F_{M'})')) & \xrightarrow{=} & H((F_{M'})'), \end{array}$$

we get that $H(\iota)$ is an isomorphism. Hence ι is a quasi-isomorphism. \square

Proposition 2.3. *Assume that E is a semi-injective DG $\mathcal{A} \otimes \mathcal{B}$ -module. For any DG \mathcal{B} -module Q , $\mathrm{Hom}_{\mathcal{B}}(Q, E)$ is a semi-injective DG \mathcal{A} -module, and $\Gamma_{\mathfrak{m}_{\mathcal{B}}}(E)$ is a direct limit of semi-injective DG \mathcal{A} -modules.*

Proof. By Lemma 1.2, one sees that $\mathrm{Hom}_{\mathcal{A}}(-, \mathrm{Hom}_{\mathcal{B}}(Q, E)) = \mathrm{Hom}_{\mathcal{A} \otimes \mathcal{B}}(- \otimes Q, E)$. This implies that $\mathrm{Hom}_{\mathcal{B}}(Q, E)$ is a semi-injective DG \mathcal{A} -module. Hence $\Gamma_{\mathfrak{m}_{\mathcal{B}}}(E) = \varinjlim_{n \geq 1} \mathrm{Hom}_{\mathcal{B}}(\mathcal{B}/\mathcal{B}^{\geq n}, E)$ is a direct limit of semi-injective DG \mathcal{A} -modules. \square

3. LOCAL DUALITY THEOREM AND ITS APPLICATIONS

The notion of Gorenstein DG algebras was first introduced by Félix-Halperin-Thomas (cf. [FHT1]). In algebraic topology, Gorenstein DG algebras are algebraic model of Gorenstein spaces. In the last twenty years, Gorenstein DG algebras have been intensively studied in literature (cf. [FJ, FHT1, FIJ, Gam, MW2]). This section gives a characterization of Gorensteinness for homologically smooth connected cochain DG algebras by using local cohomology functor.

Theorem 3.1 (Local Duality Theorem). *Let M be an object in $\mathcal{D}^c(\mathcal{A})$. Then*

$$R\Gamma_{\mathfrak{m}}(M)' = R\mathrm{Hom}_{\mathcal{A}}(M, R\Gamma_{\mathfrak{m}}(\mathcal{A})')$$

in $\mathcal{D}(\mathcal{A}^p)$.

Proof. Since M is compact, it admits a minimal semi-free resolution F_M with a finite semi-basis. Let $\eta : \mathcal{A} \xrightarrow{\sim} I$ be a semi-injective resolution of the DG \mathcal{A}^e -module \mathcal{A} . Since

$$\mathrm{Hom}_{\mathcal{A}}(-, I) \cong \mathrm{Hom}_{\mathcal{A}}(-, \mathrm{Hom}_{\mathcal{A}^e}(\mathcal{A}^e, I)) \cong \mathrm{Hom}_{\mathcal{A}^e}(\mathcal{A} \otimes_k -, I),$$

$_{\mathcal{A}}I$ is semi-injective. This implies that $I \otimes_{\mathcal{A}} F_M$ is a semi-injective DG \mathcal{A} -module since F_M is a semi-free DG \mathcal{A} -module with a finite semi-basis. Since

$$R\mathrm{Hom}_{\mathcal{A}}(M, R\Gamma_{\mathfrak{m}}(\mathcal{A})') = \mathrm{Hom}_{\mathcal{A}}(F_M, \Gamma_{\mathfrak{m}}(I)') \cong (\Gamma_{\mathfrak{m}}(I) \otimes_{\mathcal{A}} F_M)'$$

and

$$R\Gamma_{\mathfrak{m}}(M)' = R\Gamma_{\mathfrak{m}}(F_M)' = (\Gamma_{\mathfrak{m}}(I \otimes_{\mathcal{A}} F_M))' \cong (\Gamma_{\mathfrak{m}}(I) \otimes_{\mathcal{A}} F_M)',$$

we have

$$R\Gamma_{\mathfrak{m}}(M)' = R\mathrm{Hom}_{\mathcal{A}}(M, R\Gamma_{\mathfrak{m}}(\mathcal{A})')$$

in $\mathcal{D}(\mathcal{A}^p)$ □

Remark 3.2. For any compact DG \mathcal{A}^p -module N , we can similarly prove that

$$R\Gamma_{\mathfrak{m}^p}(N)' = R\mathrm{Hom}_{\mathcal{A}^p}(N, R\Gamma_{\mathfrak{m}^p}(\mathcal{A})')$$

in $\mathcal{D}(\mathcal{A})$.

Lemma 3.3. Assume that \mathcal{A} is a left Gorenstein locally finite connected cochain DG algebra. Then $R\Gamma_{\mathfrak{m}}(\mathcal{A})$ is a direct limit of DG \mathcal{A}^e -modules with finite dimensional cohomology, and $H_{\mathfrak{m}}(\mathcal{A})'$ is bounded below.

Proof. Let I be a semi-injective resolution of \mathcal{A} as a DG \mathcal{A}^e -module. Since \mathcal{A} is left Gorenstein, we have

$$H(R\mathrm{Hom}_{\mathcal{A}}(k, \mathcal{A})) = H(\mathrm{Hom}_{\mathcal{A}}(k, I)) = \Sigma^j k$$

for some $j \in \mathbb{Z}$. For any $n \geq 1$, $H(\mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I))$ is finite dimensional since $\mathcal{A}/\mathcal{A}^{\geq n}$ is finite dimensional. Hence $\Gamma_{\mathfrak{m}}(I) = \varinjlim_{n \geq 1} \mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I)$ is a direct limit of DG \mathcal{A}^e -modules with finite dimensional cohomology.

Since $\mathcal{A}/\mathcal{A}^{\geq n}$ is concentrated in degrees from 0 to $n-1$ for each $n \geq 1$, $H(\mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I))$ is concentrated in degrees from $-j+1-n$ to $-j$. Hence $H(\Gamma_{\mathfrak{m}}(I))' \cong H(\varinjlim_{n \geq 1} \mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I))' \cong (\varinjlim_{n \geq 1} H(\mathrm{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I)))'$ is concentrated in degrees $\geq j$. □

Proposition 3.4. Let \mathcal{A} be a homologically smooth locally finite connected cochain DG algebra. Then \mathcal{A} is Gorenstein if and only if $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^i({}_{\mathcal{A}}\mathcal{A})'$ in $\mathcal{D}(\mathcal{A}^p)$.

Proof. By Lemma 2.2, we have $H_{\mathfrak{m}}(k) = k$. If \mathcal{A} is Gorenstein, then there exists $i \in \mathbb{Z}$ such that $R\mathrm{Hom}_{\mathcal{A}}(k, \mathcal{A}) \cong \Sigma^i k_{\mathcal{A}}$ in $\mathcal{D}(\mathcal{A}^p)$. By Theorem 3.1,

$$\begin{aligned} H_{\mathfrak{m}}(k)' &\cong H(R\mathrm{Hom}_{\mathcal{A}}(k, R\Gamma_{\mathfrak{m}}(\mathcal{A})')) \\ &\cong H(R\mathrm{Hom}_{\mathcal{A}}(k, \mathcal{A})^L \otimes_{\mathcal{A}} R\Gamma_{\mathfrak{m}}(\mathcal{A})') \\ &\cong H(\Sigma^i k^L \otimes_{\mathcal{A}} R\Gamma_{\mathfrak{m}}(\mathcal{A})'). \end{aligned}$$

Hence $H(\Sigma^i k^L \otimes_{\mathcal{A}} R\Gamma_{\mathfrak{m}}(\mathcal{A})') \cong k$. By Lemma 3.3 and [MW1, Proposition 2.4], there is a minimal semi-free resolutions $_{\mathcal{A}}F \xrightarrow{\sim} {}_{\mathcal{A}}(R\Gamma_{\mathfrak{m}}(\mathcal{A})')$. Since

$$H(\Sigma^i k^L \otimes_{\mathcal{A}} R\Gamma_{\mathfrak{m}}(\mathcal{A})') \cong \Sigma^i k \otimes_{\mathcal{A}} F = k,$$

we have $F = \Sigma^{-i} {}_{\mathcal{A}}\mathcal{A}$. This implies that $R\Gamma_{\mathfrak{m}}(\mathcal{A})_{\mathcal{A}} \cong \Sigma^i({}_{\mathcal{A}}\mathcal{A})'$ in $\mathcal{D}(\mathcal{A}^p)$.

Conversely, if $R\Gamma_{\mathfrak{m}}(\mathcal{A})_{\mathcal{A}} \cong \Sigma^i({}_{\mathcal{A}}\mathcal{A})'$ in $\mathcal{D}(\mathcal{A}^p)$, then by Theorem 3.1, we have

$$H(R\mathrm{Hom}_{\mathcal{A}}(k, \mathcal{A})) \cong H(R\mathrm{Hom}_{\mathcal{A}}(k, \Sigma^i R\Gamma_{\mathfrak{m}}(\mathcal{A})')) \cong \Sigma^i H_{\mathfrak{m}}(k) = \Sigma^i k.$$

Since \mathcal{A} is homologically smooth, \mathcal{A} is Gorenstein by Remark 0.1. □

Remark 3.5. For any homologically smooth locally finite connected cochain DG algebra \mathcal{A} , we can similarly prove that \mathcal{A} is Gorenstein if and only if $R\Gamma_{\mathfrak{m}^p}(\mathcal{A}) \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ in $\mathcal{D}(\mathcal{A})$, by Remark 3.2 and Remark 0.1.

Theorem 3.6. Let \mathcal{A} be a Noetherian, homologically smooth connected cochain DG algebra. Then \mathcal{A} is Gorenstein if and only if $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong L \otimes_{\mathcal{A}} \mathcal{A}'$ in $\mathcal{D}(\mathcal{A}^e)$ for some invertible DG \mathcal{A} -bimodule $L = \Sigma^i \mathcal{A}(\phi)$ with $\phi \in \text{Aut}_{dg}(\mathcal{A})$.

Proof. Assume that $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong L \otimes_{\mathcal{A}} \mathcal{A}'$ in $\mathcal{D}(\mathcal{A}^e)$, where $L = \Sigma^i \mathcal{A}(\phi)$ with $\phi \in \text{Aut}_{dg}(\mathcal{A})$. Then

$$\begin{aligned} R\text{Hom}_{\mathcal{A}}(k, R\Gamma_{\mathfrak{m}}(\mathcal{A})') &\cong R\text{Hom}_{\mathcal{A}}(k, \text{Hom}_k(L \otimes_{\mathcal{A}} \mathcal{A}', k)) \\ &\stackrel{(1)}{\cong} R\text{Hom}_{\mathcal{A}}(k, \text{Hom}_{\mathcal{A}^p}(L, \mathcal{A}')) \\ &\stackrel{(2)}{\cong} R\text{Hom}_{\mathcal{A}}(k, \text{Hom}_{\mathcal{A}^p}(L, \mathcal{A})) \\ &\stackrel{(3)}{\cong} R\text{Hom}_{\mathcal{A}}(k, \mathcal{A}) \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}^p}(L, \mathcal{A}), \end{aligned}$$

where (1), (2) and (3) are obtained by the fact that $k \in \mathcal{D}^c(\mathcal{A})$. On the other hand, we have $R\text{Hom}_{\mathcal{A}}(k, R\Gamma_{\mathfrak{m}}(\mathcal{A})') \cong R\Gamma_{\mathfrak{m}}(k)'$ by Theorem 3.1. By Lemma 2.2 and Lemma 1.7, $(R\Gamma_{\mathfrak{m}}(k)')_{\mathcal{A}} \cong k_{\mathcal{A}}$ in $\mathcal{D}(\mathcal{A}^p)$. Therefore,

$$R\text{Hom}_{\mathcal{A}}(k, \mathcal{A}) \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}^p}(L, \mathcal{A}) \cong k_{\mathcal{A}}$$

in $\mathcal{D}(\mathcal{A}^p)$. Since $L = \Sigma^i \mathcal{A}(\phi)$, we have $R\text{Hom}_{\mathcal{A}}(k, \mathcal{A}) \cong k_{\mathcal{A}} \otimes_{\mathcal{A}} \Sigma^i \mathcal{A}(\phi)$. Thus \mathcal{A} is Gorenstein.

Conversely, if \mathcal{A} is Gorenstein, we should prove that $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong L \otimes_{\mathcal{A}} \mathcal{A}'$ in $\mathcal{D}(\mathcal{A}^e)$ for some invertible DG \mathcal{A} -bimodule L . By Lemma 1.8, it suffices to prove that $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ in $\mathcal{D}(\mathcal{A}^p)$ and $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ in $\mathcal{D}(\mathcal{A})$.

By Proposition 3.4, $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ in $\mathcal{D}(\mathcal{A}^p)$, for some $i \in \mathbb{Z}$. Let $\mathcal{A} \xrightarrow{\sim} I$ be a semi-injective resolution of \mathcal{A} . As a graded $\mathcal{A}^{\#}$ -module, $\text{Hom}_{\mathcal{A}}(k, I)^{\#} \cong \bigoplus_{j \in J} \Sigma^{u_j}({}_{\mathcal{A}^{\#}}k)$. Since \mathcal{A} is Gorenstein, $\dim_k H(\text{Hom}_{\mathcal{A}}(k, I)) = 1$. This implies that there is a graded cocycle element $e \in \text{Hom}_{\mathcal{A}}(k, I)$ of degree i such that $\text{Hom}_{\mathcal{A}}(k, I) \cong {}_{\mathcal{A}}ke \oplus {}_{\mathcal{A}}X$ as a DG \mathcal{A} -module, where X is a quasi-trivial DG \mathcal{A} -module with $\mathfrak{m}X = 0$. Therefore, $(\mathcal{A}_{\mathcal{A}})^{\#'} \otimes ke \oplus ((\mathcal{A}_{\mathcal{A}})^{\#'} \otimes X^{\#})$ is an injective envelope of $\text{Hom}_{\mathcal{A}}(k, I)^{\#}$. Now let $E^{\#}$ be an injective $\mathcal{A}^{\#}$ -module such that $\text{Hom}_{\mathcal{A}}(k, I)^{\#}$ is an essential graded $\mathcal{A}^{\#}$ -submodule of $E^{\#}$. Then $E^{\#} \cong (\mathcal{A}_{\mathcal{A}})^{\#'} \otimes ke \oplus ((\mathcal{A}_{\mathcal{A}})^{\#'} \otimes X^{\#})$ and $E^{\#} \subseteq \Gamma_{\mathfrak{m}}(I)^{\#}$. The graded $\mathcal{A}^{\#}$ -module $I^{\#}$ can be decomposed as $I^{\#} \cong E^{\#} \oplus J^{\#}$, for some injective $\mathcal{A}^{\#}$ -module $J^{\#}$. Since $\text{Hom}_{\mathcal{A}}(k, I)^{\#} \cap J^{\#} = 0$, it follows that $\Gamma_{\mathfrak{m}^{\#}}(J^{\#}) = 0$. Hence $\Gamma_{\mathfrak{m}}(I)^{\#} = \Gamma_{\mathfrak{m}^{\#}}(I^{\#}) \subseteq E^{\#}$. Therefore,

$$\Gamma_{\mathfrak{m}}(I)^{\#} = \mathcal{A}^{\#'} \otimes ke \oplus (\mathcal{A}^{\#'} \otimes X^{\#}).$$

We have

$$\Gamma_{\mathfrak{m}}(I)^{\#'} \cong (ke)' \otimes \mathcal{A}_{\mathcal{A}^{\#}} \oplus ((\mathcal{A}_{\mathcal{A}})^{\#'} \otimes X^{\#})'.$$

Since

$$\Gamma_{\mathfrak{m}}(I) = \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I),$$

$\text{Hom}_{\mathcal{A}}(k, I)$ is a DG \mathcal{A} -submodule of $\Gamma_{\mathfrak{m}}(I)$. Hence $\partial_{\Gamma_{\mathfrak{m}}(I)}(e) = 0$. So $\partial_{\Gamma_{\mathfrak{m}}(I)'}|_{(ke)'} = 0$. For any $a \in \mathcal{A}$, we have $\partial_{\Gamma_{\mathfrak{m}}(I)'}(e' \otimes a) = (-1)^{|e'|} e' \otimes \partial_{\mathcal{A}}(a)$ since $\Gamma_{\mathfrak{m}}(I)'$ is a DG \mathcal{A}^p -module. This implies that $(ke)' \otimes \mathcal{A}_{\mathcal{A}}$ is a DG \mathcal{A}^p -submodule of $\Gamma_{\mathfrak{m}}(I)'$. Let

$$\iota : (ke)' \otimes \mathcal{A}_{\mathcal{A}} \rightarrow \Gamma_{\mathfrak{m}}(I)'$$

be the inclusion map. Since $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ in $\mathcal{D}(\mathcal{A}^p)$, $H_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^i H(\mathcal{A})'$ as a graded k -vector space. It is easy to check that $H(\iota)$ is an isomorphism as a

morphism of graded k -vector space. This implies that ι is a quasi-isomorphism as a morphism of DG \mathcal{A}^p -modules. Therefore, the composition of DG morphisms

$$\Gamma_{\mathfrak{m}}(I) \rightarrow \Gamma_{\mathfrak{m}}(I)'' \xrightarrow{\iota'} (\mathcal{A}_{\mathcal{A}})' \otimes (ke)''$$

is also a quasi-isomorphism. Clearly, the DG \mathcal{A} -module

$$(\mathcal{A}_{\mathcal{A}})' \otimes (ke)'' \cong (\mathcal{A}_{\mathcal{A}})' \otimes ke.$$

So $\Gamma_{\mathfrak{m}}(I) \simeq \Sigma^i(\mathcal{A}_{\mathcal{A}})'$ as a DG \mathcal{A} -module, for some $i \in \mathbb{Z}$. By Lemma 1.8, $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong L \otimes_{\mathcal{A}} \mathcal{A}'$ in $\mathcal{D}(\mathcal{A}^e)$, for some invertible DG \mathcal{A} -bimodule $L = \Sigma^i \mathcal{A}(\phi)$ with $\phi \in \text{Aut}_{dg}(\mathcal{A})$. \square

Remark 3.7. By considering $\dim_k H(k^L \otimes_{\mathcal{A}} k)$, one sees that \mathcal{A} is homologically smooth if and only if \mathcal{A}^p is homologically smooth. It follows that \mathcal{A}^{op} is also a homologically smooth and Gorenstein DG algebra when the conditions in Theorem 3.4 hold. By Theorem 3.4, we have $R\Gamma_{\mathfrak{m}^{op}}(\mathcal{A}^p) \cong Q \otimes_{\mathcal{A}^{op}} (\mathcal{A}^p)' \cong \mathcal{A}' \otimes_{\mathcal{A}} Q$ in $\mathcal{D}(\mathcal{A}^e)$, where $Q = \Sigma^i \mathcal{A}(\phi)$ is an invertible DG \mathcal{A} -bimodule with $\phi \in \text{Aut}_{dg}(\mathcal{A})$.

Proposition 3.8. Let \mathcal{A} be a Noetherian, homologically smooth and Gorenstein connected cochain DG algebra. Then

$$R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong R\Gamma_{\mathfrak{m}_{\mathcal{A}^e}}(\mathcal{A}) \cong R\Gamma_{\mathfrak{m}^{op}}(\mathcal{A}^p)$$

in $\mathcal{D}(\mathcal{A}^e)$.

Proof. Let $\mathcal{A} \xrightarrow{\sim} I$ be a semi-injective resolution of the DG \mathcal{A}^e -module \mathcal{A} . By Lemma 1.1, I is semi-injective both as a DG \mathcal{A} -module and as a DG \mathcal{A}^{op} -module. Since \mathcal{A} is a Noetherian homologically smooth Gorenstein connected cochain DG algebra, we have $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong L \otimes_{\mathcal{A}} \mathcal{A}'$ in $\mathcal{D}(\mathcal{A}^e)$ by Theorem 3.6, for some invertible DG \mathcal{A} -bimodule L . For any $n \geq 1$, let $F_{\mathcal{A}^{op}/(\mathcal{A}^p)^{\geq n}}$ be the minimal semi-free resolution of the DG right \mathcal{A} -module $\mathcal{A}^p/(\mathcal{A}^p)^{\geq n}$. Then

$$\begin{aligned} R\Gamma_{\mathfrak{m}_{\mathcal{A}^e}}(\mathcal{A}) &\cong \Gamma_{\mathfrak{m}_{\mathcal{A}^e}}(I) \\ &= \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^e}((\mathcal{A}^e)/(\mathcal{A}^e)^{\geq n}, I) \\ &= \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^e}((\mathcal{A}^e)/(\mathcal{A}^{\geq n} \otimes \mathcal{A}^p + \mathcal{A} \otimes (\mathcal{A}^p)^{\geq n}), I) \\ &= \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^e}((\mathcal{A}/\mathcal{A}^{\geq n}) \otimes (\mathcal{A}^p/(\mathcal{A}^p)^{\geq n}), I) \\ &= \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^{op}}(\mathcal{A}^p/(\mathcal{A}^p)^{\geq n}, \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq n}, I)) \\ &= \varinjlim_{n \geq 1} \varinjlim_{m \geq 1} \text{Hom}_{\mathcal{A}^{op}}(\mathcal{A}^p/(\mathcal{A}^p)^{\geq n}, \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq m}, I)) \\ &\cong \varinjlim_{n \geq 1} \varinjlim_{m \geq 1} \text{Hom}_{\mathcal{A}^{op}}(F_{\mathcal{A}^{op}/(\mathcal{A}^p)^{\geq n}}, \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq m}, I)) \\ &\cong \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^{op}}(F_{\mathcal{A}^{op}/(\mathcal{A}^p)^{\geq n}}, \varinjlim_{m \geq 1} \text{Hom}_{\mathcal{A}}(\mathcal{A}/\mathcal{A}^{\geq m}, I)) \\ &\cong \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^{op}}(F_{\mathcal{A}^{op}/(\mathcal{A}^p)^{\geq n}}, L \otimes_{\mathcal{A}} \mathcal{A}') \\ &\cong \varinjlim_{n \geq 1} \text{Hom}_{\mathcal{A}^{op}}(\mathcal{A}^p/(\mathcal{A}^p)^{\geq n}, L \otimes_{\mathcal{A}} \mathcal{A}') \\ &= L \otimes_{\mathcal{A}} \mathcal{A}' \cong R\Gamma_{\mathfrak{m}}(\mathcal{A}) \end{aligned}$$

in $\mathcal{D}(\mathcal{A}^e)$. Similarly, we can prove that $R\Gamma_{\mathfrak{m}^e}(\mathcal{A}) \cong R\Gamma_{\mathfrak{m}^{op}}(\mathcal{A}^p)$ in $\mathcal{D}(\mathcal{A}^e)$.

□

Theorem 3.9. *Let \mathcal{A} be a Noetherian, homologically smooth and Gorenstein connected cochain DG algebra. Then for any $M \in \mathcal{D}^c(\mathcal{A})$, $R\Gamma_{\mathfrak{m}}(M)'$ is a compact DG \mathcal{A}^p -module. And there is a quasi-inverse contravariant equivalence of categories,*

$$\mathcal{D}^c(\mathcal{A}) \begin{array}{c} \xrightarrow{R\Gamma_{\mathfrak{m}}(-)'} \\ \xleftarrow{R\Gamma_{\mathfrak{m}^{op}}(-)'} \end{array} \mathcal{D}^c(\mathcal{A}^p).$$

Proof. Let F_M be a minimal semi-free resolution of M . Since M is compact, F_M has a finite semi-basis. By Theorem 3.1 and Theorem 3.6, we have

$$\begin{aligned} R\Gamma_{\mathfrak{m}}(M)' &\cong R\mathrm{Hom}_{\mathcal{A}}(M, R\Gamma_{\mathfrak{m}}(\mathcal{A})') \\ &\cong \mathrm{Hom}_{\mathcal{A}}(F_M, (L \otimes_{\mathcal{A}} \mathcal{A}')') \\ &\cong \mathrm{Hom}_{\mathcal{A}}(F_M, \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}'')) \\ &\cong \mathrm{Hom}_{\mathcal{A}}(F_M, \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A})) \\ &\cong \mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}) \otimes_{\mathcal{A}} \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}), \end{aligned}$$

where L is an invertible DG bimodule over \mathcal{A} . By Lemma 1.4, the DG \mathcal{A}^e -module $L \cong \Sigma^i \mathcal{A}(\phi)$, for some $i \in \mathbb{Z}$ and $\phi \in \mathrm{Aut}_{dg} \mathcal{A}$. This implies that

$$\mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}) \otimes_{\mathcal{A}} \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}) \in \mathcal{D}^c(\mathcal{A}^p)$$

since F_M admits a finite semi-basis. So $R\Gamma_{\mathfrak{m}}(M)'$ is a compact DG \mathcal{A}^p -module. To prove the quasi-inverse contravariant equivalence of categories, it suffices to show that $R\Gamma_{\mathfrak{m}^{op}}(R\Gamma_{\mathfrak{m}}(M)')' \cong M$ in $\mathcal{D}(\mathcal{A})$. By Remark 3.2,

$$R\Gamma_{\mathfrak{m}^{op}}(R\Gamma_{\mathfrak{m}}(M)')' \cong R\mathrm{Hom}_{\mathcal{A}^p}(R\Gamma_{\mathfrak{m}}(M)', R\Gamma_{\mathfrak{m}^{op}}(\mathcal{A}^p)')$$

in $\mathcal{D}(\mathcal{A})$. By Remark 3.7, we have $R\Gamma_{\mathfrak{m}^{op}}(\mathcal{A}^p) \cong \mathcal{A}' \otimes_{\mathcal{A}} Q$ in $\mathcal{D}(\mathcal{A}^e)$, for some invertible DG \mathcal{A} -bimodule Q . Proposition 3.8 indicates that $\mathcal{A}' \otimes_{\mathcal{A}} Q \cong L \otimes_{\mathcal{A}} \mathcal{A}'$ in $\mathcal{D}(\mathcal{A}^e)$. Therefore,

$$\begin{aligned} R\Gamma_{\mathfrak{m}^{op}}(R\Gamma_{\mathfrak{m}}(M)')' &\cong \mathrm{Hom}_{\mathcal{A}^p}(\mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}) \otimes_{\mathcal{A}} \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}), (\mathcal{A}' \otimes_{\mathcal{A}} Q)') \\ &\cong \mathrm{Hom}_{\mathcal{A}^p}(\mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}) \otimes_{\mathcal{A}} \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}), (L \otimes_{\mathcal{A}} \mathcal{A}')') \\ &\cong \mathrm{Hom}_{\mathcal{A}^p}(\mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}) \otimes_{\mathcal{A}} \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}), \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}'')) \\ &\cong \mathrm{Hom}_{\mathcal{A}^p}(\mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}), \mathrm{Hom}_{\mathcal{A}^p}(\mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}), \mathrm{Hom}_{\mathcal{A}^p}(L, \mathcal{A}))) \\ &\cong \mathrm{Hom}_{\mathcal{A}^p}(\mathrm{Hom}_{\mathcal{A}}(F_M, \mathcal{A}), \mathcal{A}) \\ &\cong F_M. \end{aligned}$$

So $R\Gamma_{\mathfrak{m}^{op}}(R\Gamma_{\mathfrak{m}}(M)')' \cong M$ in $\mathcal{D}(\mathcal{A})$. □

4. HOMOLOGICAL DETERMINANT

In this section, we will introduce and study the notion of homological determinant for group actions on Gorenstein homologically smooth DG algebras. For these, we need the following lemma first.

Lemma 4.1. *Let \mathcal{A} be a Gorenstein homologically smooth locally finite connected cochain DG algebra. For any $\sigma \in \mathrm{Aut}_{dg}(\mathcal{A})$, consider the morphism of DG \mathcal{A}^p -modules*

$$\Sigma^{-i} \sigma^{-1} : \Sigma^{-i} \mathcal{A}_{\mathcal{A}} \rightarrow \Sigma^{-i} \mathcal{A}_{\mathcal{A}},$$

which is σ^{-1} -linear. The Matlis dual map $(\Sigma^{-i} \sigma^{-1})' : \Sigma^i {}_{\mathcal{A}} \mathcal{A}' \rightarrow \Sigma^i {}_{\mathcal{A}} \mathcal{A}'$ is σ -linear. Hence there exists a scalar $c \in k^{\times}$ such that the $H(\sigma)$ -linear map

$$H_{\mathfrak{m}}(\sigma) : H_{\mathfrak{m}}(\mathcal{A}) \rightarrow H_{\mathfrak{m}}(\mathcal{A})$$

is equal to

$$cH[(\Sigma^{-i}\sigma^{-1})'] : \Sigma^i_{H(\mathcal{A})}[H(\mathcal{A})]' \rightarrow \Sigma^i_{H(\mathcal{A})}[H(\mathcal{A})]'$$

Proof. By Theorem 3.6, $H_{\mathfrak{m}}(\mathcal{A})$ is equal to $\Sigma^i_{H(\mathcal{A})}[H(\mathcal{A})]'$, for some $i \in \mathbb{Z}$. Acting $H_{\mathfrak{m}}(-)$ on the \mathcal{A} -linear map $\sigma : \mathcal{A}\mathcal{A} \rightarrow {}^{\sigma}\mathcal{A}$, we get the $H(\mathcal{A})$ -linear map

$$H_{\mathfrak{m}}(\sigma) : H_{\mathfrak{m}}(\mathcal{A}) \rightarrow H_{\mathfrak{m}}({}^{\sigma}\mathcal{A}) = {}^{H(\sigma)}H_{\mathfrak{m}}(\mathcal{A}).$$

By Theorem 3.6, we may write this as

$$H_{\mathfrak{m}}(\sigma) : \Sigma^i_{H(\mathcal{A})}[H(\mathcal{A})]' \rightarrow \Sigma^{iH(\sigma)}_{H(\mathcal{A})}[H(\mathcal{A})]'$$

By Matlis duality, one sees that the only $H(\mathcal{A})$ -linear maps from $\Sigma^i_{H(\mathcal{A})}[H(\mathcal{A})]'$ to $\Sigma^{iH(\sigma)}_{H(\mathcal{A})}[H(\mathcal{A})]'$ are of the form $cH[(\Sigma^{-i}\sigma^{-1})']$, where $c \in k$.

Finally, we want to see $c \neq 0$. This is easy, since $H_{\mathfrak{m}}(\sigma)H_{\mathfrak{m}}(\sigma^{-1}) = \text{id}_{H_{\mathfrak{m}}(\mathcal{A})}$. \square

Lemma 4.1 enables us to give the following definition.

Definition 4.2. Assume that \mathcal{A} is a Gorenstein, homologically smooth, locally finite connected cochain DG algebra. For each $\sigma \in \text{Aut}_{dg}(\mathcal{A})$, we have $H_{\mathfrak{m}}(\sigma) = cH[(\Sigma^{-i}\sigma^{-1})']$ by Lemma 4.1, for some $c \in k^{\times}$ and $i \in \mathbb{Z}$. We write $\text{Hdet}_{\mathcal{A}}\sigma = c^{-1}$, and call it the homological determinant of σ . It is easy for one to see that $\text{Hdet}_{\mathcal{A}}$ defines a group homomorphism $\text{Aut}_{dg}(\mathcal{A}) \rightarrow k^{\times}$.

Proposition 4.3. Let \mathcal{A} be a homologically smooth locally finite connected cochain DG algebra such that $H(\mathcal{A})$ is a Noetherian AS-Gorenstein algebra. Then for any $\sigma \in \text{Aut}_{dg}\mathcal{A}$, we have $\text{Hdet}_{\mathcal{A}}\sigma = \text{hdet}_{H(\mathcal{A})}H(\sigma)$.

Proof. Since $H(\mathcal{A})$ is a Noetherian AS-Gorenstein algebra, we have

$$H_{\mathfrak{m}_{H(\mathcal{A})}}^i(H(\mathcal{A})) \cong \begin{cases} 0, & \text{for } i \neq d, \\ H(\mathcal{A})H(\mathcal{A})'(l), & \text{for } i = d, \end{cases}$$

by [JoZ, Lemma 2.1]. Here d and l are the injective dimension and the Gorenstein index of $H(\mathcal{A})$, respectively. So

$$\text{Ext}_{H(\mathcal{A})}^i(k, H(\mathcal{A})) \cong \begin{cases} 0, & \text{for } i \neq d, \\ k(l), & \text{for } i = d, \end{cases}$$

By [Gam, Proposition 1], \mathcal{A} is a Gorenstein DG algebra. By Theorem 3.6, $R\Gamma_{\mathfrak{m}}(\mathcal{A}) \cong \Sigma^j(\mathcal{A}\mathcal{A})'$ in $\mathcal{D}(\mathcal{A}^p)$ for some $j \in \mathbb{Z}$. Then by Theorem 3.1,

$$\begin{aligned} H(R\text{Hom}_{\mathcal{A}}(k, \mathcal{A})) &\cong H(R\text{Hom}_{\mathcal{A}}(k, \Sigma^j R\Gamma_{\mathfrak{m}}(\mathcal{A})')) \\ &\cong \Sigma^j H_{\mathfrak{m}}(k) = \Sigma^j k. \end{aligned}$$

By [KM, Theorem 4.7], there is a natural spectral sequence

$$E_2^{p,q} = \text{Ext}_{H(\mathcal{A})}^{p,q}(k, H(\mathcal{A})) \Rightarrow H^{p+q}(R\text{Hom}_{\mathcal{A}}(k, \mathcal{A})).$$

Hence $j = d - l$. Let $c = \text{Hdet}_{\mathcal{A}}\sigma$. Then

$$H_{\mathfrak{m}}(\sigma) = c^{-1}H[(\Sigma^{l-d}\sigma^{-1})'] : \Sigma^{d-l}_{H(\mathcal{A})}[H(\mathcal{A})]' \rightarrow \Sigma^{d-l}_{H(\mathcal{A})}[H(\mathcal{A})]'$$

by Definition 4.2. We have

$$(1) \quad H_{\mathfrak{m}}(\sigma) = c^{-1}H[(\Sigma^{l-d}\sigma^{-1})'] = c^{-1}\Sigma^{d-l}[H(\sigma^{-1})]' = c^{-1}\Sigma^{d-l}[(H(\sigma))^{-1}]'.$$

By [JoZ, 2.2 and 2.3], $H(\sigma) : H(\mathcal{A}) \rightarrow H(\mathcal{A})$ induces a $H(\sigma)$ -linear map

$$H_{\mathfrak{m}_{H(\mathcal{A})}}^d(H(\sigma)) : {}_{H(\mathcal{A})}H(\mathcal{A})'(l) \rightarrow {}_{H(\mathcal{A})}H(\mathcal{A})'(l).$$

By the definition of the homological determinant of $H(\sigma)$, we have

$$(2) \quad H_{\mathfrak{m}_{H(\mathcal{A})}}^d(H(\sigma)) = [\text{hdet}_{H(\mathcal{A})}H(\sigma)]^{-1}\{[(H(\sigma))^{-1}]'(l)\}.$$

By (1) and (2), we have $\text{Hdet}_{\mathcal{A}}\sigma = c = \text{hdet}_{H(\mathcal{A})}H(\sigma)$. \square

Proposition 4.4. *Let \mathcal{A} be a connected cochain DG algebra. Suppose that G is a finite subgroup of $\text{Aut}_{dg}\mathcal{A}$. Then*

- (1) $\mathcal{A}^G = \{a \in \mathcal{A} | g(a) = a, \forall g \in G\}$ is a connected cochain DG subalgebra of \mathcal{A} ;
- (2) $H(G) : \{H(g) | g \in G\}$ is a finite subgroup of $\text{Aut}_{gr}H(\mathcal{A})$;
- (3) The graded fixed subalgebra $H(\mathcal{A})^{H(G)}$ of $H(\mathcal{A})$ is isomorphic to $H(\mathcal{A}^G)$.

Proof. (1) It is straightford to show that \mathcal{A}^G is a subcomplex which is closed under the addition and multiplication of \mathcal{A} .

(2) For any $g_1, g_2 \in G$, we have $H(g_1)H(g_2) = H(g_1 \circ g_2) \in H(G)$. Hence $H(G)$ is a subgroup of $\text{Aut}_{gr}H(\mathcal{A})$.

(3) Let $\iota : \mathcal{A}^G \rightarrow \mathcal{A}$ be the inclusion morphism of DG algebras. It induces a morphism of graded algebras $H(\iota) : H(\mathcal{A}^G) \rightarrow H(\mathcal{A})$. For any $[z] \in \ker(H(\iota))$, we have $z \in \mathcal{A}^G$, $\partial_{\mathcal{A}^G}(z) = 0$ and $H(\iota)([z]) = [z] = 0$ in $H(\mathcal{A})$. There exists $a \in \mathcal{A}$ such that $\partial_{\mathcal{A}}(a) = z$. Let $x = \frac{1}{|G|-1} \sum_{g \in G} g(a)$. For any $\sigma \in G$, we have

$$\sigma(x) = \frac{1}{|G|-1} \sum_{g \in G} \sigma \circ g(a) = \frac{1}{|G|-1} \sum_{g' \in G} g'(a) = x$$

and

$$\begin{aligned} \partial_{\mathcal{A}^G}(x) &= \partial_{\mathcal{A}^G} \left[\frac{1}{|G|-1} \sum_{g \in G} g(a) \right] \\ &= \frac{1}{|G|-1} \sum_{g \in G} \partial_{\mathcal{A}} \circ g(a) \\ &= \frac{1}{|G|-1} \sum_{g \in G} g \circ \partial_{\mathcal{A}}(a) \\ &= \frac{1}{|G|-1} \sum_{g \in G} g(z) = \frac{1}{|G|-1} \sum_{g \in G} z = z. \end{aligned}$$

Then $[z] = 0$ in $H(\mathcal{A}^G)$ and $H(\iota)$ is a monomorphism. Hence $H(\mathcal{A}^G) \cong \text{im}(H(\iota))$. For any $[z] \in H(\mathcal{A}^G)$, we have $\partial_{\mathcal{A}^G}(z) = \partial_{\mathcal{A}}(z) = 0$ and $g(z) = z$, for any $g \in G$. Hence $H(g)([z]) = [g(z)] = [z]$, for any $g \in G$. This implies that $[z] \in H(\mathcal{A})^{H(G)}$. So $\text{im}(H(\iota)) \subseteq H(\mathcal{A})^{H(G)}$. Conversely, let $[b] \in H(\mathcal{A})^{H(G)}$. Then $[b] = H(g)([b]) = [g(b)]$ for any $g \in G$. So there exists $x_g \in \mathcal{A}$ such that $g(b) - b = \partial_{\mathcal{A}}(x_g)$. Set

$$y = \frac{1}{|G|-1} \sum_{g \in G} g(b).$$

For any $\sigma \in G$, we have

$$\sigma(y) = \sigma \left[\frac{1}{|G|-1} \sum_{g \in G} g(b) \right] = \frac{1}{|G|-1} \sum_{g \in G} \sigma \circ g(b) = \frac{1}{|G|-1} \sum_{g' \in G} g'(b) = y.$$

Furthermore,

$$\begin{aligned} \partial_{\mathcal{A}}(y) &= \partial_{\mathcal{A}} \left[\frac{1}{|G|-1} \sum_{g \in G} g(b) \right] \\ &= \frac{1}{|G|-1} \sum_{g \in G} \partial_{\mathcal{A}} \circ g(b) \\ &= \frac{1}{|G|-1} \sum_{g \in G} g(\partial_{\mathcal{A}}(b)) = 0 \end{aligned}$$

and

$$\begin{aligned}
\lceil \iota(y) \rceil &= \lceil y \rceil = \lceil \frac{1}{|G|^{-1}} \sum_{g \in G} g(b) \rceil \\
&= \lceil \frac{1}{|G|^{-1}} \sum_{g \in G} (z + \partial_{\mathcal{A}}(x_g)) \rceil \\
&= \lceil b + \partial_{\mathcal{A}}(\frac{1}{|G|^{-1}} \sum_{g \in G} x_g) \rceil = \lceil b \rceil.
\end{aligned}$$

Then $H(\mathcal{A})^{H(G)} \subseteq \text{im}(H(\iota))$ and hence $H(\mathcal{A})^{H(G)} = \text{im}(H(\iota)) \cong H(\mathcal{A}^G)$. \square

Theorem 4.5. *Let \mathcal{A} be a homologically smooth locally finite connected cochain DG algebra such that $H(\mathcal{A})$ is a Noetherian AS-Gorenstein graded algebra. If G is a finite subgroup of $\text{Aut}_{dg}(\mathcal{A})$ such that the homological determinant of each $\sigma \in G$ is 1, then the fixed DG subalgebra \mathcal{A}^G is a Gorenstein DG algebra.*

Proof. By Proposition 4.3, $\text{hdet}_{H(\mathcal{A})} H(\sigma) = \text{Hdet}_{\mathcal{A}}(\sigma) = 1$ for any $\sigma \in G$. Hence $H(\mathcal{A})^{H(G)}$ is AS-Gorenstein by [JoZ, Theorem 3.3]. Since $H(\mathcal{A}^G) \cong H(\mathcal{A})^{H(G)}$ by Proposition 4.4, we conclude that $H(\mathcal{A}^G)$ is AS-Gorenstein. This implies that \mathcal{A}^G is a Gorenstein DG algebra by [Gam, Proposition 1]. \square

DG down-up algebras are introduced and systematically studied in [MHLX]. Recall that a connected cochain DG algebra \mathcal{A} is called down-up if $\mathcal{A}^\#$ is a graded down-up algebra $A(\alpha, \beta)$, which is a graded associative algebra generated by degree-one elements x and y over k with the relations given by

$$x^2 y = \alpha x y x + \beta y x^2 \quad \text{and} \quad x y^2 = \alpha y x y + \beta y^2 x.$$

When $\beta \neq 0$, $A(\alpha, \beta)$ is Noetherian. By [MHLX, Theorem A], the differential of a Noetherian DG down-up algebra can be described as follows:

- (a) $\partial_{\mathcal{A}} = 0$ if $1 + \alpha - \beta \neq 0$ or $1 + \alpha - \beta = 0, \beta^3 \neq 1$;
- (b) $\partial_{\mathcal{A}}$ is defined by either

$$\begin{cases} \partial_{\mathcal{A}}(x) = c y^2 \\ \partial_{\mathcal{A}}(y) = d x^2 \\ c \cdot d = 0, \quad c, d \in k \end{cases} \quad \text{or} \quad \begin{cases} \partial_{\mathcal{A}}(x) = 2d x^2 + c x y + c y x - \frac{c^2}{d} y^2 \\ \partial_{\mathcal{A}}(y) = -\frac{d^2}{c} x^2 + d x y + d y x + 2c y^2 \\ c \in k^\times, d \in k^\times \end{cases}$$

if $1 + \alpha - \beta = 0, \beta^3 = 1$ and $\beta \neq 1$;

- (c) $\partial_{\mathcal{A}}$ is defined by $\begin{cases} \partial_{\mathcal{A}}(x) = c_1 x^2 + c_2(x y + y x) + c_3 y^2 \\ \partial_{\mathcal{A}}(y) = d_1 x^2 + d_2(x y + y x) + d_3 y^2 \\ c_i, d_i \in k, i = 1, 2, 3 \end{cases}$ if $\alpha = 0$ and $\beta = 1$.

Therefore, we can divide Noetherian DG down-up algebras with non-trivial differentials into the following three cases:

Case 1. $\mathcal{A}^\# = A(\xi - 1, \xi)$, Case 2. $\mathcal{A}^\# = A(\xi^2 - 1, \xi^2)$, Case 3. $\mathcal{A}^\# = A(0, 1)$,

where ξ is a fixed primitive cubic root of unity. A Noetherian DG down-up algebra of the third case is not isomorphic to any Noetherian DG down-up algebras of the other two cases since we have $A(0, 1) \not\cong A(\xi - 1, \xi)$ and $A(0, 1) \not\cong A(\xi^2 - 1, \xi^2)$ by [BR, Corollary 6.2]. By [MHLX, Theorem B], the set of non-trivial Noetherian DG down-up algebras belonging to Case 1 or Case 2 can be classified into 3 different isomorphism classes represented by

$$(1) \mathcal{A}_1 \text{ with } \mathcal{A}_1^\# = A(\xi - 1, \xi) \text{ and } \begin{cases} \partial_{\mathcal{A}_1}(x) = y^2 \\ \partial_{\mathcal{A}_1}(y) = 0; \end{cases}$$

- (2) \mathcal{A}_2 with $\mathcal{A}_2^\# = A(\xi - 1, \xi)$ and $\begin{cases} \partial_{\mathcal{A}_2}(x) = 0 \\ \partial_{\mathcal{A}_2}(y) = x^2; \end{cases}$
- (3) \mathcal{A}_3 with $\mathcal{A}_3^\# = A(\xi - 1, \xi)$ and $\begin{cases} \partial_{\mathcal{A}_2}(x) = 2x^2 + xy + yx - y^2 \\ \partial_{\mathcal{A}_2}(y) = -x^2 + xy + yx + 2y^2. \end{cases}$

By [MHLX, Proposition 5.5],

$$\begin{aligned} H(\mathcal{A}_1) &= \frac{k\langle [xy+yx], [y] \rangle}{\left(\begin{array}{c} \xi[y][xy+yx] - [xy+yx][y] \\ [y^2] \end{array} \right)}, \\ H(\mathcal{A}_2) &= \frac{k\langle [xy+yx], [x] \rangle}{\left(\begin{array}{c} [x][xy+yx] - \xi[xy+yx][x] \\ [x^2] \end{array} \right)}, \\ \text{and } H(\mathcal{A}_3) &= k[[xy+yx]^3] \end{aligned}$$

are all AS-Gorenstein. Furthermore, $\mathcal{A}_1, \mathcal{A}_2$ and \mathcal{A}_3 are Calabi-Yau DG algebras by [MHLX, Proposition 6.1, Corollary 6.2 and Proposition 6.4]. For Case 3, $H(\mathcal{A})$ is AS-Gorenstein by Proposition 5.7, and \mathcal{A} is Calabi-Yau by [MHLX, Theorem 6.11]. Therefore, any DG down-up DG algebra \mathcal{A} is homologically smooth and $H(\mathcal{A})$ is a Noetherian AS-Gorenstein graded algebra. Then we have the following corollary.

Corollary 4.6. *Let \mathcal{A} be a Noetherian DG down-up algebra. If G is a finite subgroup of $\text{Aut}_{dg}(\mathcal{A})$ such that the homological determinant of each $\sigma \in G$ is 1, then the fixed DG subalgebra \mathcal{A}^G is a Gorenstein DG algebra.*

In [MXYA], the notion of DG free algebra is introduced. Recall that a connected cochain DG algebra \mathcal{A} is called DG free if $\mathcal{A}^\#$ is a free algebra $k\langle x_1, x_2, \dots, x_n \rangle$ generated by degree 1 elements. By [MXYA, Theorem 2.4], $\partial_{\mathcal{A}}$ is in one to one correspondence with a set of crisscross ordered n -tuples of $n \times n$ matrices. More precisely, there exists a crisscross ordered n -tuple (M^1, M^2, \dots, M^n) of $n \times n$ matrices such that $\partial_{\mathcal{A}}$ is defined by

$$\partial_{\mathcal{A}}(x_i) = (x_1, x_2, \dots, x_n) M^i \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, \forall i \in \{1, 2, \dots, n\}.$$

Here, (M^1, M^2, \dots, M^n) with

$$M^i = (c_1^i, c_2^i, \dots, c_n^i) = \begin{pmatrix} r_1^i \\ r_2^i \\ \vdots \\ r_n^i \end{pmatrix}$$

is called crisscross if

$$\sum_{l=1}^n [c_j^l r_l^i - c_i^l r_j^l] = (0)_{n \times n}, \forall i, j \in \{1, 2, \dots, n\}.$$

When $n = 2$, we classify all isomorphism classes of DG free algebras. It is proved in [MXYA] that all non-trivial DG free algebras generated by 2 degree-one elements are Koszul Calabi-Yau DG algebras, which are obviously homologically smooth. Furthermore, their cohomology graded algebras were all Noetherian and AS-Gorenstein by [MXYA, Proposition 5.1]. Applying Theorem 4.5, we get the following corollary.

Corollary 4.7. *Let \mathcal{A} be a non-trivial DG free algebra with 2 degree-one generators. If G is a finite subgroup of $\text{Aut}_{dg}(\mathcal{A})$ such that the homological determinant of each $\sigma \in G$ is 1, then the fixed DG subalgebra \mathcal{A}^G is a Gorenstein DG algebra.*

ACKNOWLEDGMENTS

X.-F. Mao is supported by the National Natural Science Foundation of China (No. 11871326).

REFERENCES

- [AF] L. L. Avramov and H. V. Foxby, Locally gorenstein homomorphisms, *Amer. J. Math.* 114 (1992), 1007-1047.
- [AFH] L. L. Avramov, H. V. Foxby and S. Halperin, "Differential Graded Homological Algebra," in preparation.
- [Ap] D. Apassov, Dualizing modules over differential graded rings, pp. 41-47 in "Complexes and Differential Graded Modules," ph. d. thesis, Lund University, 1999.
- [BR] G. Benkart and T. Roby, Down-Up algebras, *J. Algebra* 209, (1998) 305-344. Addendum: *J. Algebra* 213 (1999), 378.
- [CWZ] K. Chan, C. Walton and J.J. Zhang, Hopf actions and Nakayama automorphisms, *J. Algebra* 409 (2014), 26-53.
- [DGI] W. Dwyer, J. P. C. Greenlees and S. Iyengar, "Duality in algebra and topology" *Adv. Math.* 200 (2006), 357-402.
- [FJ] A. J. Frankild and P. Jørgensen, Gorenstein Differential Graded Algebras, *Israel J. Math.* 135 (2003), 327-354.
- [FHT1] Y. Félix, S. Halperin, and J. C. Thomas, Gorenstein spaces, *Adv. Math.* 71 (1988), 92-112.
- [FHT2] Y. Félix, S. Halperin and J. C. Thomas, "Rational Homotopy Theory", *Grad. Texts in Math.* 205, Springer, Berlin, 2000.
- [FIJ] A. Frankild, S. Iyengar and P. Jørgensen, Dualizing differential graded modules and gorenstein differential graded algebras, *J. London Math. Soc.* (2) 68 (2003), 288-306.
- [Gam] H. Gammelin, Gorenstein space with nonzero evaluation map, *Trans. Amer. Math. Soc.* 351 (1999), 3433-3440.
- [HOZ] J.-W. He, F. Van Oystaeyen and Y. Zhang, Hopf algebra actions on differential graded algebras and applications, *Bulletin of Belgian Math. Soc.* 18 (2011), 99-111.
- [Hin] V. A. Hinic, On the Gorenstein property of the ring of invariants of a Gorenstein ring, *Izv. Akad. Nauk SSSR Ser. Mat.* Tom 40 (1976), 47-53.
- [HW] J.-W. He and Q.-S. Wu, Koszul differential graded algebras and BGG correspondence, preprint.
- [Iv] B. Iversen, Amplitude inequalities for complexes, *Ann. Sci. Ecole Norm. Sup.* 10 (1977), 547-558.
- [Ke] B. Keller, Deriving DG categories, *Ann. Sci. École Norm. Sup.* (4) 27 (1994), no. 1, 63-102.
- [KKZ1] E. Kirkman, J. Kuzmanovich and J.J. Zhang, Gorenstein subrings of invariants under Hopf algebra actions, *J. Algebra*, 322 (2009), 3640-3669.
- [KM] I. Kriz and J. P. May, "Operads, Algebras, Modules and Motives," *Astérisque* 233 (1995).
- [Jor1] P. Jørgensen, Local cohomology for non-commutative graded algebras, *Comm. in Algebra* 25 (1997), no. 2, 575-591.
- [Jor2] P. Jørgensen, Duality for cochain DG algebras. *Sci. China Math.*, 2013, 56: 79-89
- [JoZ] P. Jørgensen and J.J. Zhang, Gourmet's guide to Gorensteinness, *Adv. Math.*, 151(2000), 313-345.
- [JZ] N. Jing and J.J. Zhang, Gorensteinness of invariant subrings of quantum algebras, *J. Algebra*, 221 (1999), 669-691.
- [LMZ1] J.-F. Lv, X.-F. Mao and J.J. Zhang, Nakayama automorphism and applications, *Trans. Amer. Math. Soc.*, 369 (2017), 2425-2460.
- [LPWZ2] D.-M. Lu, J.H. Palmieri, Q.-S. Wu and J.J. Zhang, *Koszul equivalences in A_∞ -algebras*, *New York J. Math.* 14 (2008), 325-378.
- [Mon] S. Montgomery. Hopf algebras and their actions on rings, volume 82 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1993
- [MHLX] X.-F. Mao, J.-W. He, M. Liu and J.-F. Xie, Calabi-Yau properties of non-trivial Noetherian DG down-up algebras, *J. Algebra Appl.* 17(5) (2018):1850090-45.
- [MW1] X.-F. Mao and Q.-S. Wu, Homological invariants for connected DG algebra, *Comm. Algebra* 36 (2008), 3050-3072.
- [MW2] X.-F. Mao and Q.-S. Wu, Compact DG modules and Gorenstein DG algebras, *Sci. china Ser. A* 52 (2009), 648-676.
- [MW3] X.-F. Mao and Q.-S. Wu, Cone length for DG modules and global dimension of DG algebras, *Comm. Algebra*, 39 (2011), 1536-1562.

- [MXYA] X.-F. Mao, J.-F. Xie, Y.-N. Yang and Almire. Abia, Isomorphism problem and homological properties of DG free algebras, *Comm. Algebra*, 47 (10), 2019, 4031-4060
- [MYY] X.-F. Mao, Y.-N. Yang and C.-C. Ye, A sufficient condition for a connected DG algebra to be Calabi-Yau, *Comm. Algebra*, 47(8), 2019, 3280-3296.
- [MWYZ] X.-F. Mao, X.-T. Wang, Y.-N. Yang and M.-Y. Zhang, Calabi-Yau properties of 3-dimensional DG Sklyanin algebras, arxiv: math. RA/2009.03524V1
- [MWZ] X.-F. Mao, X. Wang and M.-Y. Zhang, DG Algebra structures on the quantum affine n -space $\mathcal{O}_{-1}(k^n)$, *J. Algebra*, 594, (2022), 389-482.
- [VDB] M. Van den Bergh, Existence Theorems for dualizing complexes over Non-commutative graded and filtered rings, *J. Algebra* 195 (1997), 662-679.
- [Wat1] K. Watanabe, Certain invariants subrings are Gorenstein, *Osaka J. Math.* 11 (1974) 1-8.
- [Wat2] K. Watanabe, Certain invariants subrings are Gorenstein II, *Osaka J. Math.* 11 (1974) 379-388.
- [Yek] A. Yekutieli, Dualizing complexes over noncommutative graded algebra, *J. Algebra* 153 (1992), 41-84.

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, CHINA
Email address: xuefengmao@shu.edu.cn

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, CHINA
Email address: happywang97@shu.edu.cn