DUAL CURVATURE MEASURES ON CONVEX FUNCTIONS AND ASSOCIATED MINKOWSKI PROBLEMS

NIUFA FANG AND JIAZU ZHOU*

ABSTRACT. In this paper, the q-th dual curvature measure is extended to convex functions and the associated Minkowski problem is posed. A special case includes the q-th dual curvature measure of convex bodies which defined by Huang, Lutwak, Yang and Zhang. Existence for the functional dual Minkowski problem is showed when $q \leq 0$ and the uniqueness part is obtained with some assumptions.

1. INTRODUCTION

The classical Brunn-Minkowski theory originated with the thesis of Brunn in 1887 and is in its essential parts with the creation of Minkowski around the turn of the century. Mixed volumes are important geometric concepts, which include quermassintegrals, surface area measures, curvature measures and mixed area measures. An interesting problem related to surface area measures is the classical Minkowski problem that is crucial important in the Brunn-Minkowski theory:

What are necessary and sufficient conditions for a finite Borel measure μ on the unite sphere S^{n-1} so that μ is the surface area measure of a convex body in \mathbb{R}^n ?

The Minkowski problem was solved in multiple ways by Minkowski, Aleksandrov, and Fenchel and Jessen [55, 62]. Landmark contributions to establishing regularity for the Minkowski problem are due to Lewy [40], Nirenberg [56], Pogorelov [57, 58], Cheng and Yau [12], Caffarelli [9], Guan and Ma [32] and others.

If the measure has a density f with respect to the Lebesgue measure of the unit sphere S^{n-1} , the Minkowski problem is equivalent to the study of solution to the following Monge-Ampère equation on the unit sphere

$$\det(h_{ij} + \delta_{ij}h) = f,$$

where h is the unknown function on S^{n-1} to be found, h_{ij} is the covariant derivative of h with respect to an orthonormal frame on S^{n-1} and δ_{ij} is the Kronecker delta.

The L_p Brunn-Minkowski theory, a generalization of the classical Brunn-Minkowski theory, attracted increasing interest in the recent years. Lutwak [48] introduced the

²⁰¹⁰ Mathematics Subject Classification. 52A40, 26B25, 26D10.

Key words and phrases. Convex functions; functional dual curvature measures; dual quermassintegrals, functional dual Minkowski problem.

The first author is supported in part by NSFC (No.12001291).

^{*}The corresponding author is supported in part by NSFC (No.12071318).

 L_p surface area measure, a Borel measure on convex body in \mathbb{R}^n with the origin in its interior. The L_p surface area measure is an important concept in the L_p Brunn-Minkowski theory, and it provides an integral representation for the L_p mixed volume. By the L_p cosine transform of the L_p surface area measure, Lutwak, Yang and Zhang [49] defined the L_p Petty projection body and established the well-known L_p Petty projection inequality. Finding the necessary and sufficient conditions for a given measure to guarantee that it is the L_p surface area measure is the existence problem called the L_p Minkowski problem posed in [48]. Solving the L_p Minkowski problem requires solving a degenerate and singular Monge-Ampère type equation on the unit sphere. The L_p Minkowski problem has been solved for $p \geq 1$ [13, 38, 48, 51], and for the critical case p < 1 [7, 63, 64, 72–75]. The solution of the L_p Minkowski problem and the L_p Petty projection inequality are the key tools used for establishing the L_p affine Sobolev inequality and analogues [14, 33, 34, 50, 68].

A theory analogous to the Brunn-Minkowski theory was introduced in [45]. It demonstrates a remarkable duality in convex geometry, and thus is called the *dual Brunn-Minkowski theory*. The important work of the duality between projection bodies and intersection bodies exhibited in [46]. The dual Brunn-Minkowski theory has attracted extensive attention since the intersection body helped achieving a major breakthrough in the solution of the celebrated Busemann-Petty problem. In the past decades, the dual Brunn-Minkowski theory has a rapid growth [28–30, 47, 67, 69, 77].

Very recently, Huang, Lutwak, Yang and Zhang [35] introduced *dual curvature measures*. This new family of measures miraculously connects the cone-volume measure and Aleksandrov's integral curvature. The associated Minkowski problems of dual curvature measures are called *dual Minkowski problems*. Since the groundbreaking work of Huang, Lutwak, Yang and Zhang, the intrinsic partial differential equations that arise within the dual Brunn-Minkowski theory which wait a full 40 years after the birth of the dual theory to emerge. Some significant breakthroughs of dual Minkowski problems have been achieved [6,8,11,31,36,37,41,52,65,66,70,71,76] but many cases remain open.

Motivated by works of Huang, Lutwak, Yang and Zhang [35], we will introduce the *functional dual curvature measures* on the setting of convex functions. The new functional dual curvature measures include dual curvature measures introduced in [35] as a special case. All works in this paper are characterized as the *functional dual Brunn-Minkowski theory*, a new branch in convex geometric analysis.

It is well known that the classical isoperimetric inequality is equivalent to the Sobolev inequality. This equivalence shows a close connection between geometry and analysis. The geometric inequalities and geometric problems of functions received considerable attention. Actually an analytic inequality contains more information than its corresponding geometric inequality. During the past decades, many analytic inequalities with geometric background were found [1–4,10,15,16,22–24,26,27,42,54, 59] basically from the viewpoint of integral and convex geometry.

The algebraic structure (i.e., "Minkowski sum" and "scalar multiplication") on the set of convex bodies is the cornerstone in the classical Brunn-Minkowski theory. By using the *infimal convolution* and *right scalar multiplication* of convex functions, Colesanti and Fragalà [17] introduced the "sum" and "scalar multiplication" of log-concave functions. Colesanti and Fragalà's work belongs to the *functional Brunn-Minkowski theory*. We recall that the infimal convolution and the right scalar multiplication. Let $\varphi, \psi : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ be convex functions. The *infimal convolution* of φ and ψ is defined by [60],

$$\varphi \Box \psi(x) = \inf_{y \in \mathbb{R}^n} \{ \varphi(x - y) + \psi(y) \} \quad \forall x \in \mathbb{R}^n,$$

and the right scalar multiplication, φt , is defined by,

$$(\varphi t)(x) = t\varphi\left(\frac{x}{t}\right), \text{ for } t > 0.$$

In order to study the functional dual Minkowski problems, the crucial step is to properly define the functional dual quermassintegrals. For a non-negative convex function $\varphi : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ and $q \in \mathbb{R}$, the (n-q)-th dual quermassintegral $\widetilde{W}_{n-q}(\varphi)$ is defined by

$$\widetilde{W}_{n-q}(\varphi) = \int_{\mathbb{R}^n} \varphi(x)^{-q} d\gamma_n(x), \qquad (1.1)$$

where $d\gamma_n(x) = (2\pi)^{-\frac{n}{2}} e^{-\frac{|x|^2}{2}} dx$ denotes the Gaussian probability measure. We will prove that the (n-q)-th dual quermassintegral, $\widetilde{W}_{n-q}(\varphi)$, of the convex function φ includes the (n-q)-th dual quermassintegral of a convex body.

For $q \in \mathbb{R} \setminus \{0\}$, the (n-q)-th dual mixed quermassintegral, $\widetilde{W}_{n-q}(\varphi, \psi)$, of non-negative convex functions φ and ψ is defined by

$$\widetilde{W}_{n-q}(\varphi,\psi) = \frac{1}{q} \lim_{t \to 0^+} \frac{\widetilde{W}_{n-q}(\varphi \Box(\psi t)) - \widetilde{W}_{n-q}(\varphi)}{t}.$$
(1.2)

The Minkowski type inequality for the functional dual mixed quermassintegrals can be stated as follows:

Minkowski type inequality. If $q \leq 0$ and φ, ψ are non-negative convex functions in \mathbb{R}^n , then

$$\widetilde{W}_{n+1-q}(\varphi,\psi) \ge \frac{\widetilde{W}_{n+1-q}(\psi)^{\frac{1}{1-q}} - \widetilde{W}_{n+1-q}(\varphi)^{\frac{1}{1-q}}}{\widetilde{W}_{n+1-q}(\varphi)^{\frac{q}{1-q}}} + \widetilde{W}_{n+1-q}(\varphi,\varphi),$$
(1.3)

with equality if and only if $\varphi(x) = \psi(x - x_0)$ for some $x_0 \in \mathbb{R}^n$ when q = 0, and $\varphi(x) = \psi(x)$ when q < 0.

It should be pointed out that $\widetilde{W}_{n+1-q}(\varphi, \varphi)$ does not agree with $\widetilde{W}_{n+1-q}(\varphi)$ but it satisfies

$$q\widetilde{W}_{n-q}(\varphi,\varphi) = (n-q)\widetilde{W}_{n-q}(\varphi) - \int_{\mathbb{R}^n} |x|^2 \varphi(x)^{-q} d\gamma_n(x)$$

for $q \in \mathbb{R} \setminus \{0\}$.

If $q \leq 0$ and φ, ψ satisfy some additional conditions (see details in Theorem 3.1), we prove that the functional dual mixed quermassintegral has the following integral representation:

$$\frac{d}{dt}\widetilde{W}_{n+1-q}(\varphi\Box(\psi t))\Big|_{t=0^+} = \int_{\mathbb{R}^n} \psi^*(x)d\widetilde{C}_q(\varphi, x),$$
(1.4)

where ψ^* is the *Legendre transform* of ψ (defined in Section 2) and the Borel measure $\widetilde{C}_q(\varphi, \cdot)$ on \mathbb{R}^n is defined by

$$\widetilde{C}_q(\varphi, \cdot) = (\nabla \varphi)_{\sharp}(\varphi^{-q} d\gamma_n).$$

When limited to a subclass of convex functions, $\tilde{C}_q(\varphi, \cdot)$ reduces to the *q*-th dual curvature measure of a convex body. The *q*-th dual curvature measure of the convex function φ is also called the *functional q-th dual curvature measure*. It leads naturally to the following Minkowski problem involving the functional *q*-th dual curvature measure.

The functional dual Minkowski problem. Let μ be a finite Borel measure on \mathbb{R}^n and $q \in \mathbb{R}$. Find the necessary and sufficient conditions on μ so that it is the q-th dual curvature measure $\tau \widetilde{C}_q(\varphi, \cdot)$ of a convex function φ in \mathbb{R}^n , where the constant $\tau \in \mathbb{R}$.

We will show that when the measure μ has a density function $g : \mathbb{R}^n \to \mathbb{R}$, the partial differential equation for the functional dual Minkowski problem is a Monge-Ampère type equation:

$$(2\pi)^{-\frac{n}{2}} f(\nabla f^*)^{-q} e^{-\frac{|\nabla f^*|^2}{2}} \det(\nabla^2 f^*) = g \quad \text{on} \quad \Omega,$$
(1.5)

where f is the unknown convex function in \mathbb{R}^n to be found, ∇f and $\nabla^2 f$ denote respectively the gradient vector and the Hessian matrix of f with respect to an orthonormal frame in \mathbb{R}^n , and f^* denotes the Legendre transform of f (see the detailed definition in Section 2), $\Omega = \{x \in \mathbb{R}^n : f(x) < +\infty\} \cap \{x \in \mathbb{R}^n : f^*(x) < +\infty\}$.

We solve the functional dual Minkowski problem when $q \leq 0$.

Existence of solution to the functional dual Minkowski problem. Let μ be a non-zero finite Borel measure with $\int_{\mathbb{R}^n} |x| d\mu(x) < \infty$ on \mathbb{R}^n and $q \leq 0$. If μ is not supported in a lower-dimensional subspace, then there exists a constant $\tau \in \mathbb{R}$ and a non-negative, convex function φ with finite $\int_{\mathbb{R}^n} \varphi(x)^{1-q} d\gamma_n(x)$ such that $\mu(\cdot) = \tau \widetilde{C}_q(\varphi, \cdot)$. Moreover, if φ is essentially-continuous then $\widetilde{C}_q(\varphi, \cdot)$ is not supported in a lower-dimensional subspace.

We remark that a convex function $\varphi : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ is *essentially-continuous* if φ is lower semi-continuous and if the set of points where φ is discontinuous has zero \mathcal{H}^{n-1} -measure, where \mathcal{H}^{n-1} is the (n-1)-dimensional Hasusdorff measure.

By the Minkowski type inequality (1.3), we establish the uniqueness of the solution to the functional dual Minkowski problem.

Uniqueness of the solution to the functional dual Minkowski problem. Let $q \leq 0$ and let φ_1, φ_2 be non-negative convex functions in \mathbb{R}^n with $\varphi_1(0) = \varphi_2(0) = 0$. If there exist $c_1, c_2 > 0$ such that $\varphi_1^* - c_2\varphi_2^*$ and $\varphi_2^* - c_1\varphi_1^*$ are convex, then $\widetilde{C}_{q-1}(\varphi_1, \cdot) = \widetilde{C}_{q-1}(\varphi_2, \cdot)$ implies that φ_1 and φ_2 agree up to a translation when q = 0, and $\varphi_1 = \varphi_2$ almost everywhere when q < 0.

In [35], authors proved that the 0-th dual curvature measure is the Aleksandrov's integral curvature, and the related Minkowski problem is called Aleksandrov problem. Therefore, the existential part can be viewed as the solution to the *functional Aleksandrov problem* when q = 0. Similarly, the functional *n*-th dual curvature measure, $\tilde{C}_n(\varphi, \cdot)$, is the functional cone-volume measure, and the corresponding Minkowski problem is the *functional log-Minkowski problem*. Unfortunately, the uniqueness part does not imply these two special cases. In this paper, we only consider the functional dual Minkowski problem for the case $q \leq 0$.

2. Preliminaries

2.1. The setting of convex functions.

A function $\varphi : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ is convex if for every $x, y \in \mathbb{R}^n$ and $\lambda \in [0, 1]$

$$\varphi((1-\lambda)x + \lambda y) \le (1-\lambda)\varphi(x) + \lambda\varphi(y).$$

Let

$$\operatorname{dom}(\varphi) = \{ x \in \mathbb{R}^n : \varphi(x) \in \mathbb{R} \}.$$

We say that φ is proper if dom $(\varphi) \neq \emptyset$. The Legendre transform of φ is the convex function defined by

$$\varphi^*(y) = \sup_{x \in \mathbb{R}^n} \{ \langle x, y \rangle - \varphi(x) \} \qquad \forall y \in \mathbb{R}^n.$$
(2.1)

Clearly, $\varphi(x) + \varphi^*(y) \ge \langle x, y \rangle$ for all $x, y \in \mathbb{R}^n$. There is an equality if and only if $\varphi(x) < +\infty$ and $y \in \partial \varphi(x)$, the subdifferential of φ at x define by

$$\partial \varphi(x) = \{ z \in \mathbb{R}^n : \varphi(y) \ge \varphi(x) + \langle z, y - x \rangle \text{ for all } y \in \mathbb{R}^n \}.$$
(2.2)

Hence,

$$\varphi^*(\nabla\varphi(x)) + \varphi(x) = \langle x, \nabla\varphi(x) \rangle.$$

The following elementary property was proved in [60],

$$\varphi^*(0) = -\inf\varphi. \tag{2.3}$$

For convenient, we set

$$\mathcal{L} = \left\{ \varphi : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\} \middle| \quad \varphi \text{ proper, convex, } \lim_{|x| \to +\infty} \varphi(x) = +\infty \right\},$$

and \mathcal{L}_o , all non-negative convex functions of the subset of \mathcal{L} .

A function $\varphi \in \mathcal{L}$ is *lower semi-continuous*, if the subset $\{x \in \mathbb{R}^n : \varphi(x) > t\}$ is an open set for any $t \in (-\infty, +\infty]$. The function φ^* is always convex and lower semi-continuous. If φ is a lower semi-continuous convex function, then φ^* is also a lower semi-continuous convex function, and $\varphi^{**} = \varphi$. It should be noted that for a general function φ one always has $\varphi^{**} \leq \varphi$.

For $\varphi, \psi \in \mathcal{L}$, the *infimal convolution* is defined by

$$\varphi \Box \psi(x) = \inf_{y \in \mathbb{R}^n} \{ \varphi(x - y) + \psi(y) \} \quad \forall x \in \mathbb{R}^n,$$
(2.4)

and the *right scalar multiplication* is defined by,

$$(\varphi t)(x) = t\varphi\left(\frac{x}{t}\right), \quad \text{for} \quad t > 0.$$
 (2.5)

A function $f : \mathbb{R}^n \to \mathbb{R}$ is *log-concave* if $f = e^{-\varphi}$ with $\varphi \in \mathcal{L}$. Then we have the following Prékopa-Leindler inequality [62]:

Let 0 < t < 1 and f, g, h be non-negative integrable functions in \mathbb{R}^n satisfying

$$h((1-t)x + ty) \ge f(x)^{1-t}g(y)^t,$$

for all $x, y \in \mathbb{R}^n$. Then

$$\int_{\mathbb{R}^n} h(x)dx \ge \left(\int_{\mathbb{R}^n} f(x)dx\right)^{1-t} \left(\int_{\mathbb{R}^n} g(x)dx\right)^t.$$
(2.6)

Equality holds in (2.6) if and only if f and g are log-concave functions, and $f(x) = g(x - x_0)$ for some $x_0 \in \mathbb{R}^n$.

The following properties will be useful late [17, 60].

Proposition 2.1. Let $\varphi, \psi \in \mathcal{L}$. Then

(1) $(\varphi \Box \psi)^* = \varphi^* + \psi^*;$ (2) $(\varphi t)^* = t\varphi^*, t > 0.$

Theorem A. [60, Theorem 10.9] Let C be a relatively open convex set, and let f_1, f_2, \dots , be a sequence of finite convex functions on C. Suppose that the real number sequence $f_1(x), f_2(x), \dots$, is bounded for each $x \in C$. It is then possible to select a subsequence of f_1, f_2, \dots , which converges uniformly on closed bounded subsets of C to some finite convex function f.

2.2. Dual curvature measures.

Let \mathbb{R}^n be equipped with the usual Euclidean norm $|\cdot|$. A subset K of \mathbb{R}^n is called a convex body if it is a compact convex set with non-empty interior. The set of all convex bodies that contain the origin in their interiors is denoted by \mathcal{K}_o^n . For $x, y \in \mathbb{R}^n$, let $\langle x, y \rangle$ denote the standard inner product of x and y. For a convex body K in \mathbb{R}^n , we write ∂K for the boundary of K.

For $K, L \in \mathcal{K}_o^n$ and real t, s > 0, the Minkowski combination, tK + sL, is defined by

$$tK + sL = \{tx + sy : x \in K, y \in L\}.$$

The support function $h_K(\cdot) = h(K, \cdot) : \mathbb{R}^n \to \mathbb{R}$ of a convex body K is defined by

$$h(K, x) = \max\{\langle x, y \rangle : y \in K\}, \ x \in \mathbb{R}^n.$$

The convex body K is uniquely determined by its support function $h(K, \cdot)$. It is obvious that for t > 0, the support function of the convex body $tK = \{tx : x \in K\}$ satisfies

$$h(tK, \cdot) = th(K, \cdot).$$

For $K \in \mathcal{K}_o^n$, its radial function $\rho(K, \cdot) : \mathbb{R}^n \setminus \{o\} \to [0, \infty)$ is defined by

$$o(K, x) = \max\{t \ge 0 : tx \in K\}.$$

The radial function is denoted also by $\rho_K(x)$.

For a convex body K, its gauge function $\|\cdot\|_K$ is defined by

$$||x||_{K} = \min\{t \ge 0 : x \in tK\}.$$
(2.7)

The *polar body* K° of K is defined by

$$K^{\circ} = \{ x \in \mathbb{R}^n : \langle x, y \rangle \le 1 \text{ for all } y \in K \}.$$

If $K \in \mathcal{K}_o^n$, then

$$\|x\|_{K} = h_{K^{\circ}}(x) = \frac{1}{\rho_{K}(x)}.$$
(2.8)

It is obvious that

$$||x||_{K} = 1 \quad \text{whenever} \quad x \in \partial K. \tag{2.9}$$

Let $K \in \mathcal{K}_o^n$ and $q \in \mathbb{R}$. The q-th dual curvature measure $\widetilde{C}_q(K, \cdot)$ is defined by (see, e.g., [35])

$$\widetilde{C}_q(K,\omega) = \frac{1}{n} \int_{\boldsymbol{\alpha}_K^*(\omega)} \rho_K^q(u) du, \qquad (2.10)$$

for each Borel set $\omega \subset S^{n-1}$. Here $\boldsymbol{\alpha}_{K}^{*}(\omega)$ denotes the set of directions $u \in S^{n-1}$ such that the boundary point $\rho_{K}(u)u$ belongs to $\nu_{K}^{-1}(\omega)$, where ν_{K} is the Gaussian map which defined on the boundary ∂K of K, and ν_{K}^{-1} is the inverse Gaussian map. Equivalently, for a bounded Borel function $g: S^{n-1} \to \mathbb{R}$,

$$\int_{S^{n-1}} g(v) d\widetilde{C}_q(K, v) = \frac{1}{n} \int_{S^{n-1}} g(\alpha_K(u)) \rho_K^q(u) du,$$
(2.11)

where $\alpha_K(u) = \nu_K(\rho_K(u)u)$. Moreover, if K is strictly convex, then

$$\int_{S^{n-1}} g(v) d\widetilde{C}_q(K, v) = \frac{1}{n} \int_{\partial K} g(\nu_K(x)) h_K(\nu_K(x)) |x|^{q-n} d\mathcal{H}^{n-1}(x), \qquad (2.12)$$

where \mathcal{H}^{n-1} is the (n-1)-dimensional Hausdorff measure. We note that the total measure of the q-th dual curvature measure is the (n-q)-th dual quermassintegral $\widetilde{W}_{n-q}(K)$, i.e.,

$$\widetilde{W}_{n-q}(K) = \frac{1}{n} \int_{S^{n-1}} \rho_K(u)^q du.$$
(2.13)

It was proved in [35] that the *n*-th dual curvature measure of a convex body is the cone volume measure of this convex body, while the zeroth dual curvature measure

of a convex body is Aleksandrov's integral curvature of the polar body of this body, i.e.,

$$\widetilde{C}_n(K,\cdot) = \frac{1}{n} h_K(\cdot) dS_K(\cdot), \qquad (2.14)$$

$$\widetilde{C}_0(K,\cdot) = \mathcal{H}^{n-1}(\boldsymbol{\alpha}_{K^{\circ}}(\cdot)).$$
(2.15)

The associated Minkowski problems of the cone volume measure and the Aleksandrov's integral curvature measure are called *log-Minkowski problem* (see, e.g., [7, 63, 64, 72]) and *Aleksandrov problem* (see, e.g., [62]), respectively.

3. Mixed dual quermassintegrals of convex functions

In this section, we will define the dual quermassintegrals for convex functions that include the dual quermassintegrals for convex bodies. Inspired by works of [35], we investigate the functional dual mixed quermassintegrals.

3.1. Integral formula of the functional dual mixed quermassintegrals.

Let $q \in \mathbb{R} \setminus \{0\}$, and $\varphi \in \mathcal{L}_o$. The (n-q)-th dual quermassintegral is defined as

$$\widetilde{W}_{n-q}(\varphi) = \int_{\mathbb{R}^n} \varphi(x)^{-q} d\gamma_n(x).$$
(3.1)

Let q < n and $K \in \mathcal{K}_o^n$. By the polar coordinates of x, (2.13) and (2.8), we have

$$\widetilde{W}_{n-q}(\|x\|_{K^{\circ}}) = \int_{\mathbb{R}^{n}} \|x\|_{K^{\circ}}^{-q} d\gamma_{n}(x)$$

$$= (2\pi)^{-\frac{n}{2}} \int_{0}^{+\infty} e^{-\frac{r^{2}}{2}} r^{n-q-1} dr \int_{S^{n-1}} \|u\|_{K^{\circ}}^{-q} du$$

$$= n(2\pi)^{-\frac{n}{2}} 2^{\frac{n-q}{2}-1} \Gamma\left(\frac{n-q}{2}\right) \widetilde{W}_{n-q}(K).$$

In this sense, our definition (3.1) extended dual quermassintegrals of convex bodies to the functional version.

For real $q \neq 0$, the normalized dual quermassintegral $\overline{W}_{n-q}(\varphi)$ is defined as

$$\overline{W}_{n-q}(\varphi) = \left(\int_{\mathbb{R}^n} \varphi(x)^{-q} d\gamma_n(x)\right)^{\frac{1}{q}},\tag{3.2}$$

and, for q = 0, by

$$\overline{W}_n(\varphi) = \exp\left(-\int_{\mathbb{R}^n} \log\varphi(x) d\gamma_n(x)\right).$$
(3.3)

The functional dual mixed quermassintegrals are also introduced.

Definition 3.1. Let $q \in \mathbb{R} \setminus \{0\}$, and let $\varphi, \psi \in \mathcal{L}_o$. The (n - q)-th dual mixed quermassintegral of φ and ψ is defined as

$$\widetilde{W}_{n-q}(\varphi,\psi) = \frac{1}{q} \lim_{t \to 0^+} \frac{\widetilde{W}_{n-q}(\varphi \Box(\psi t)) - \widetilde{W}_{n-q}(\varphi)}{t}$$

In particular, when $\varphi = \psi$, we have

Lemma 3.1. Let $q \in \mathbb{R} \setminus \{0\}$, and let $\varphi \in \mathcal{L}_o$. If $\widetilde{W}_{n-q}(\varphi)$ is finite, then $q\widetilde{W}_{n-q}(\varphi,\varphi) = (n-q)\widetilde{W}_{n-q}(\varphi) - \int_{\mathbb{D}^n} |x|^2 \varphi(x)^{-q} d\gamma_n(x).$

Proof. From the definitions of infimal convolution (2.4) and right scalar multiplication (2.5), we have

$$\begin{split} \varphi \Box[\varphi t](x) &= \inf_{y \in \mathbb{R}^n} \left\{ \varphi(x-y) + t\varphi\left(\frac{y}{t}\right) \right\} \\ &\leq (1+t)\varphi\left(\frac{x}{1+t}\right), \end{split}$$

and the convexity of φ deduces

$$\frac{1}{1+t}\varphi(x-y) + \frac{t}{1+t}\varphi\left(\frac{y}{t}\right) \ge \varphi\left(\frac{x}{1+t}\right),$$

for any $x, y \in \mathbb{R}^n$ and t > 0. This means that $\varphi \Box(\varphi t) = \varphi(1+t)$. A direct calculation yields

$$\begin{split} & \frac{\widetilde{W}_{n-q}(\varphi\Box(\varphi t)) - \widetilde{W}_{n-q}(\varphi)}{t} \\ &= \frac{(2\pi)^{-\frac{n}{2}}}{t} \left[\int_{\mathbb{R}^n} (1+t)^{-q} \varphi\left(\frac{x}{1+t}\right)^{-q} e^{-\frac{|x|^2}{2}} dx - \int_{\mathbb{R}^n} \varphi(x)^{-q} e^{-\frac{|x|^2}{2}} dx \right] \\ &= \frac{(2\pi)^{-\frac{n}{2}}}{t} \left[\int_{\mathbb{R}^n} (1+t)^{n-q} \varphi(x)^{-q} e^{-\frac{(1+t)^2|x|^2}{2}} dx - \int_{\mathbb{R}^n} \varphi(x)^{-q} e^{-\frac{|x|^2}{2}} dx \right] \\ &= (2\pi)^{-\frac{n}{2}} \left[\frac{(1+t)^{n-q} - 1}{t} \right] \int_{\mathbb{R}^n} \varphi(x)^{-q} e^{-\frac{(1+t)^2|x|^2}{2}} dx \\ &+ (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} \varphi(x)^{-q} e^{-\frac{|x|^2}{2}} \left[\frac{e^{-\frac{(t^2+2t)|x|^2}{2}} - 1}}{t} \right] dx. \end{split}$$

By the monotone convergence theorem, let $t \to 0^+$, we obtain

$$q\widetilde{W}_{n-q}(\varphi,\varphi) = (n-q)\widetilde{W}_{n-q}(\varphi) - \int_{\mathbb{R}^n} |x|^2 \varphi(x)^{-q} d\gamma_n(x).$$

Let

$$\mathcal{L}' = \left\{ \varphi \in \mathcal{L} \middle| \quad \varphi \in \mathcal{C}^2_+, \ \lim_{|x| \to +\infty} \frac{\varphi(x)}{|x|} = +\infty \right\}.$$

Here C^2_+ used for the set of functions whose Hessian matrix are positive definite at each point. Let \mathcal{L}'_o denote the set of the non-negative convex function $\varphi \in \mathcal{L}'$ and there exists constants a, b > 0 such that $\varphi(x) \leq b|x|^{1+a}$ when |x| is big enough for $x \in \mathbb{R}^n$.

In order to give the integral representation of the functional dual mixed quermassintegral we need the following three lemmas.

Lemma 3.2. [17, Lemma 2.5] Let $\varphi \in \mathcal{L}$. There exist constants $a, b \in \mathbb{R}$ with a > 0 such that

$$\varphi(x) > a|x| + b,$$

for $x \in \mathbb{R}^n$.

Lemma 3.3. [17, Lemma 4.10] Let $\varphi, \psi \in \mathcal{L}'$ and, for any t > 0, set $\varphi_t = \varphi \Box(\psi t)$. Assume that $\psi(0) = 0$, then

(1)
$$\varphi_1(x) \leq \varphi_t(x) \leq \varphi(x), \quad \forall x \in \mathbb{R}^n, \ \forall t \in [0, 1];$$

(2) $\forall x \in \{x \in \mathbb{R}^n : \varphi(x) < +\infty\}, \quad \lim_{t \to 0^+} \varphi_t(x) = \varphi(x);$
(3) $\forall E \subset \{x \in \mathbb{R}^n : \varphi(x) < +\infty\}, \quad \lim_{t \to 0^+} \nabla \varphi_t(x) = \nabla \varphi(x) \text{ uniformly on } E.$

Lemma 3.4. [17, Lemma 4.11] Let $\varphi, \psi \in \mathcal{L}'$ and, for any t > 0, set $\varphi_t = \varphi \Box(\psi t)$. Assume that $\psi(0) = 0$, then for $x \in int(\{x \in \mathbb{R}^n : \varphi(x) < +\infty\})$

$$\frac{d}{dt}\varphi_t(x) = -\psi^*(\nabla\varphi_t(x)).$$

The next lemma will be used later.

Lemma 3.5. Let q < 0 and $\varphi \in \mathcal{L}'_o$. If $0 < \int_{\mathbb{R}^n} \varphi^{2-q}(x) d\gamma_n(x) < +\infty$, then

$$\int_{\mathbb{R}^n} \varphi^*(\nabla \varphi(x)) \varphi^{-q}(x) d\gamma_n(x) < +\infty.$$

Proof. By a direct calculation, we have

$$\begin{split} &\int_{\mathbb{R}^n} \varphi^*(\nabla\varphi(x))\varphi^{-q}(x)d\gamma_n(x) \\ &= \int_{\mathbb{R}^n} (\langle x, \nabla\varphi(x) \rangle - \varphi(x))\varphi^{-q}(x)d\gamma_n(x) \\ &= \frac{1}{1-q} \int_{\mathbb{R}^n} \langle x, \nabla\varphi^{1-q}(x) \rangle d\gamma_n(x) - \int_{\mathbb{R}^n} \varphi^{1-q}(x)d\gamma_n(x) \\ &= \frac{(2\pi)^{-\frac{n}{2}}}{1-q} \int_{\mathbb{R}^n} \operatorname{div}(\varphi^{1-q}e^{-\frac{|x|^2}{2}}x)dx + \frac{1}{1-q} \int_{\mathbb{R}^n} |x|^2 \varphi(x)^{1-q} d\gamma_n(x) \\ &\quad -\frac{n+q-1}{1-q} \int_{\mathbb{R}^n} \varphi(x)^{1-q} d\gamma_n(x). \end{split}$$

Since there exist constants a > 0, b > 0 such that $\varphi(x) \leq b|x|^{1+a}$ when |x| is big enough for $x \in \mathbb{R}^n$, hence

$$\int_{\mathbb{R}^n} \operatorname{div}(\varphi^{1-q} e^{-\frac{|x|^2}{2}} x) dx$$

= $\lim_{r \to +\infty} \int_{rB} \operatorname{div}(\varphi^{1-q} e^{-\frac{|x|^2}{2}} x) dx$
= $\lim_{r \to +\infty} r e^{-\frac{r^2}{2}} \int_{\partial(rB)} \varphi^{1-q}(x) d\mathcal{H}^{n-1}(x)$
= 0.

The Hölder inequality yields

$$\int_{\mathbb{R}^n} |x|^2 \varphi(x)^{1-q} d\gamma_n(x) \le \left[\int_{\mathbb{R}^n} \varphi(x)^{2-q} d\gamma_n(x) \right]^{\frac{1-q}{2-q}} \left[\int_{\mathbb{R}^n} |x|^{2(2-q)} d\gamma_n(x) \right]^{\frac{1}{2-q}}.$$

The finiteness of $\int_{\mathbb{R}^n} \varphi^*(\nabla \varphi(x)) \varphi^{-q}(x) d\gamma_n(x)$ follows from the above facts, $\widetilde{W}_{n+1-q}(\varphi)$ is finite and $\int_{\mathbb{R}^n} |x|^{2(2-q)} d\gamma_n(x)$ is a constant depend on n and q.

A convex function φ is an *admissible perturbation* for the convex function ψ if there exists a constant c > 0 such that the function

$$\varphi^* - c\psi^* \tag{3.4}$$

is convex.

The following theorem provides an integral formula of the (n+1-q)-th functional dual mixed quermassintegral.

Theorem 3.1. Let $q \leq 0$ and $\varphi, \psi \in \mathcal{L}'_o$ and assume that φ is an admissible perturbation for ψ . If $\varphi(0) = \psi(0) = 0$ and $\widetilde{W}_{n+2-q}(\varphi)$ is finite, then $\widetilde{W}_{n+1-q}(\varphi, \psi) \in [0, +\infty]$ and

$$\widetilde{W}_{n+1-q}(\varphi,\psi) = \int_{\mathbb{R}^n} \psi^*(\nabla\varphi(y))\varphi(y)^{-q}d\gamma_n(y)$$

=:
$$\int_{\mathbb{R}^n} \psi^*(x)d\widetilde{C}_q(\varphi,x).$$
 (3.5)

Proof. The proof is divided as several steps.

Step 1: $\widetilde{W}_{n+1-q}(\varphi, \psi) \in [0, +\infty].$

Let φ, ψ be non-negative convex functions in \mathbb{R}^n and, for any t > 0, set $\varphi_t = \varphi \Box(\psi t)$. Since $\psi(0) = 0$, then Lemma 3.3 (1) yields that the function φ_t is pointwise decreasing with respect to t.

Because of Lemma 3.3 (1) and (2), so that for every $x \in \mathbb{R}^n$ there exists $\bar{\varphi}(x) = \lim_{t \to 0^+} \varphi_t(x)$ and it holds $\bar{\varphi}(x) \leq \varphi(x)$. By the monotone convergence theorem and $q \leq 0$, we have

$$\widetilde{W}_{n+1-q}(\bar{\varphi}) = \lim_{t \to 0^+} \widetilde{W}_{n+1-q}(\varphi_t) \le \widetilde{W}_{n+1-q}(\varphi).$$

Let us consider separately the two cases $\widetilde{W}_{n+1-q}(\bar{\varphi}) < \widetilde{W}_{n+1-q}(\varphi)$ and $\widetilde{W}_{n+1-q}(\bar{\varphi}) = \widetilde{W}_{n+1-q}(\varphi)$.

If $\widetilde{W}_{n+1-q}(\bar{\varphi}) < \widetilde{W}_{n+1-q}(\varphi)$, then

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q}(\varphi_t) - \widetilde{W}_{n+1-q}(\varphi)}{t} = -\infty.$$

If $\widetilde{W}_{n+1-q}(\bar{\varphi}) = \widetilde{W}_{n+1-q}(\varphi)$, we further distinguish the following two subcases:

$$\exists t_0 > 0 \text{ such that } \widetilde{W}_{n+1-q}(\varphi_{t_0}) = \widetilde{W}_{n+1-q}(\varphi),$$

or

$$\widetilde{W}_{n+1-q}(\varphi_t) < \widetilde{W}_{n+1-q}(\varphi) \text{ for all } t > 0.$$

Lemma 3.3 (1) declares that $\widetilde{W}_{n+1-q}(\varphi_t)$ is a monotone decreasing function of t, so that $\widetilde{W}_{n+1-q}(\varphi_t) = \widetilde{W}_{n+1-q}(\varphi)$ for every $t \in [0, t_0]$ in the former subcase. This means

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q}(\varphi_t) - \widetilde{W}_{n+1-q}(\varphi)}{t} = 0.$$

Now, we consider the latter subcase. Since $\widetilde{W}_{n+1-q}(\varphi_t)$ is a decreasing convex function of t when $q \leq 0$, hence

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q}(\varphi_t) - \widetilde{W}_{n+1-q}(\varphi)}{t} \in [-\infty, 0].$$
(3.6)

Combining the above facts and the definition of functional dual quermassintegrals, we conclude that $\widetilde{W}_{n+1-q}(\varphi, \psi) \in [0, +\infty]$.

Step 2: For t > 0, it holds

$$\widetilde{W}_{n+1-q}(\varphi_t) - \widetilde{W}_{n+1-q}(\varphi) = \int_0^t \Phi(s) ds, \qquad (3.7)$$

where

$$\Phi(s) = \int_{\mathbb{R}^n} \psi^*(x) d\widetilde{C}_q(\varphi_s, x).$$
(3.8)

By Lemma 3.4, it holds

$$\lim_{h \to 0} \frac{\varphi_{t+h}^{1-q}(x) - \varphi_t^{1-q}(x)}{h} = (q-1)\psi^*(\nabla\varphi_t(x))\varphi_t^{-q}(x),$$
(3.9)

for $x \in \mathbb{R}^n$. From the facts $q \leq 0$ and $\psi(0) = 0$, Lemma 3.3 (1) and (3), we infer that for every $s \in [0, 1]$, the non-negative function $\psi^*(\nabla \varphi_s(x))\varphi_s^{-q}(x)$ are bounded above

by some continuous function independent of s. Then, by the pointwise convergence in (3.9) and dominated convergence theorem, we have

$$\frac{1}{q-1}\lim_{h\to 0}\frac{\widetilde{W}_{n+1-q}(\varphi_{t+h})-\widetilde{W}_{n+1-q}(\varphi_t)}{h}$$
$$=\frac{1}{q-1}\lim_{h\to 0}\int_{\mathbb{R}^n}\frac{\varphi_{t+h}^{1-q}(x)-\varphi_t^{1-q}(x)}{h}d\gamma_n(x)$$
$$=\int_{\mathbb{R}^n}\psi^*(\nabla\varphi_t(x))\varphi_t^{-q}(x)d\gamma_n(x).$$

Moreover,

$$\widetilde{W}_{n+1-q}(\varphi_t) - \widetilde{W}_{n+1-q}(\varphi) = \int_0^t \int_{\mathbb{R}^n} \psi^*(x) d\widetilde{C}_q(\varphi_s, x) ds.$$

Step 3: The function Φ defined in (3.8) takes finite values at every s > 0.

Let s > 0. Since $\varphi(0) = 0$, hence $\varphi^* \ge 0$ and

$$s\Phi(s) \leq \int_{\mathbb{R}^n} (\varphi^*(x) + s\psi^*(x)) d\widetilde{C}_q(\varphi_s, x)$$
$$= \int_{\mathbb{R}^n} \varphi^*_s(x) d\widetilde{C}_q(\varphi_s, x).$$

Lemma 3.5 ensures $\Phi(s) < +\infty$ when s > 0.

Let now s = 0. Since ψ is an admissible perturbation for φ , by its definition (3.4) and Lemma 3.2, there exist constants a, c > 0 and $b \in \mathbb{R}$ such that

$$(\varphi^* - c\psi^*)(y) \ge a|y| + b,$$
 (3.10)

for $y \in \mathbb{R}^n$. Hence,

0

$$\Phi(0) = \int_{\mathbb{R}^n} \psi^*(x) d\widetilde{C}_q(\varphi, x)
\leq c^{-1} \int_{\mathbb{R}^n} \varphi^*(x) d\widetilde{C}_q(\varphi, x) - \frac{b}{c} \widetilde{W}_{n-q}(\varphi) - \frac{a}{c} \int_{\mathbb{R}^n} |\nabla \varphi(x)| \varphi^{-q}(x) d\gamma_n(x)
\leq c^{-1} \int_{\mathbb{R}^n} \varphi^*(x) d\widetilde{C}_q(\varphi, x) - \frac{b}{c} \widetilde{W}_{n-q}(\varphi).$$

The finiteness of $\Phi(0)$ follows from Lemma 3.5, and a > 0, c > 0.

Step 4: The function Φ defined in (3.8) is continuous at every s > 0, and it is continuous from the right at s = 0.

Let $y = \nabla \varphi_s(x)$. Then we have

$$\Phi(s) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} \psi^*(y) \varphi_s(\nabla \varphi_s^*(y)) e^{-\frac{|\nabla \varphi_s^*(y)|^2}{2}} \det(\nabla^2 \varphi_s^*(y)) dy.$$

By the convexity of $\phi = \varphi^* - c\psi^*$ for some c > 0, one gets

$$\nabla^2 \phi = \nabla^2 \varphi^* - c \nabla^2 \psi^*,$$

and it is a symmetric positive semi-definite matrix. This implies that

$$\nabla^2 \psi^* = c^{-1} \nabla^2 \varphi^* - c^{-1} \nabla^2 \phi.$$

According to the fact that

$$\det(A+B)^{1/n} \ge \det(A)^{1/n} + \det(B)^{1/n},$$

for symmetric positive semi-definite matrices A and B, we infer that

$$det(\nabla^{2}\varphi_{s}^{*}(y)) = det(\nabla^{2}\varphi^{*}(y) + s\nabla^{2}\psi^{*}(y)) = det((1 + c^{-1}s)\nabla^{2}\varphi^{*}(y) - c^{-1}s\nabla^{2}\phi) \leq (1 + c^{-1}s)^{n} det(\nabla^{2}\varphi^{*}(y)).$$
(3.11)

Obviously,

$$|\nabla\varphi_s^*(y)|^2 = \left|\nabla\varphi^*(y) + s\nabla\psi^*(y)\right|^2$$

is a continuous function with respect to s. Hence,

$$\lim_{s \to 0} |\nabla \varphi_s^*(y)|^2 = \lim_{s \to 0} |\nabla \varphi^*(y)|^2,$$

i.e., for any $0 < \varepsilon < 1$ there exists an $s_0 > 0$ small enough such that

$$-\varepsilon < |\nabla \varphi_s^*(y)|^2 - |\nabla \varphi^*(y)|^2 < \varepsilon$$
(3.12)

when $0 < s < s_0$. On the other hand, by the convexity of $\phi = \varphi^* - c\psi^*$ for some c > 0 and $\varphi(0) = \psi(0) = 0$, we have

$$\varphi_{s}(\nabla\varphi_{s}^{*}(y)) = \langle y, \nabla\varphi_{s}^{*}(y) \rangle - \varphi_{s}^{*}(y) \\
= (1 + \frac{s}{c}) \Big(\langle y, \nabla\varphi^{*}(y) \rangle - \varphi^{*}(y) \Big) - \frac{s}{c} \Big(\langle y, \nabla\phi(y) \rangle - \phi(y) \Big) \\
= (1 + \frac{s}{c}) \varphi(\nabla\varphi^{*}(y)) - \frac{s}{c} \phi^{*}(\nabla\phi(y)) \\
\leq (1 + \frac{1}{c}) \varphi(\nabla\varphi^{*}(y)).$$
(3.13)

From (3.11), (3.12), (3.13) and (3.10) for $s \in (0, \min\{s_0, 1\})$ we have

$$\begin{split} \psi^{*}(y)\varphi_{s}(\nabla\varphi_{s}^{*}(y))^{-q} \det(\nabla^{2}\varphi_{s}^{*}(y))e^{-\frac{|\nabla\varphi_{s}^{*}(y)|^{2}}{2}} \\ &\leq (1+c^{-1})^{n-q}e\psi^{*}(y)\varphi(\nabla\varphi^{*}(y))^{-q} \det(\nabla^{2}\varphi^{*}(y))e^{-\frac{|\nabla\varphi^{*}(y)|^{2}}{2}} \\ &\leq (1+c^{-1})^{n-q}c^{-1}e\psi^{*}(y)\varphi(\nabla\varphi^{*}(y))^{-q} \det(\nabla^{2}\varphi^{*}(y))e^{-\frac{|\nabla\varphi^{*}(y)|^{2}}{2}} \\ &\quad -\frac{b}{c}\varphi(\nabla\varphi^{*}(y))^{-q} \det(\nabla^{2}\varphi^{*}(y))e^{-\frac{|\nabla\varphi^{*}(y)|^{2}}{2}}. \end{split}$$

Let
$$h_1(y) = (2\pi)^{-\frac{n}{2}} (1+c^{-1})^n c^{-1} e \varphi^*(y) \varphi(\nabla \varphi^*(y))^{-q} \det(\nabla^2 \varphi^*(y)) e^{-\frac{|\nabla \varphi^*(y)|^2}{2}}$$
 and $h_2(y) = \max\{0, -\frac{b}{c}\}(2\pi)^{-\frac{n}{2}} \varphi(\nabla \varphi^*(y))^{-q} \det(\nabla^2 \varphi^*(y)) e^{-\frac{|\nabla \varphi^*(y)|^2}{2}}$. Then

$$\int_{\mathbb{R}^n} h_1(y) dy = (2\pi)^{-\frac{n}{2}} (1+c^{-1})^n c^{-1} e \int_{\mathbb{R}^n} \varphi^*(y) \varphi(\nabla \varphi^*(y))^{-q} \det(\nabla^2 \varphi^*(y)) e^{-\frac{|\nabla \varphi^*(y)|^2}{2}} dy$$

$$= (1+c^{-1})^n c^{-1} \int_{\mathbb{R}^n} \varphi^*(\nabla \varphi(x)) \varphi(x)^{-q} d\gamma_n(x),$$
and

and

$$\int_{\mathbb{R}^n} h_2(y) dy = \max\{0, -\frac{b}{c}\} \widetilde{W}_{n+1-q}(\varphi)$$

From Lemma 3.5, we obtain the integrability of the non-negative function $h_1 + h_2$ with respect to the Gaussian probability measure γ_n . Therefore, the desired claim follows from dominated convergence theorem.

Step 5: Equality (3.5) holds.

Combining with (3.7), the finiteness (which proved in Step 3) and continuity (which proved in Step 4) of $\Phi(s)$ for s > 0, we have

$$\Phi(s) = \frac{d}{dt} \widetilde{W}_{n+1-q}(\varphi_t) \Big|_{t=s}.$$
(3.14)

The continuity from the right of Φ at s = 0 implies

$$\lim_{s \to 0^+} \Phi(s) = \Phi(0) = \int_{\mathbb{R}^n} \psi^*(x) d\widetilde{C}_q(\varphi, x).$$
(3.15)

Then, equality (3.5) follows from (3.7), (3.14) and (3.15).

3.2. Minkowski version inequality.

There is a Minkowski inequality for the functional dual mixed quermassintegral.

Lemma 3.6. Let $\varphi, \psi \in \mathcal{L}$. For any $t \in (0,1)$ and $x, y \in \mathbb{R}^n$, then

$$[\varphi(1-t)]\Box[\psi t]((1-t)x+ty) \le (1-t)\varphi(x) + t\psi(y),$$
(3.16)

with equality if and only if $\varphi(x) = \psi(x - x_0)$ for some $x_0 \in \mathbb{R}^n$.

Proof. For any $x, y \in \mathbb{R}^n$ and $t \in (0, 1)$, we have

$$\begin{split} & [\varphi(1-t)]\Box[\psi t]((1-t)x+ty) \\ &= \inf_{z \in \mathbb{R}^n} \left\{ (1-t)\varphi\left(\frac{(1-t)x+ty-z}{1-t}\right) + t\psi\left(\frac{z}{t}\right) \right\} \\ & \leq (1-t)\varphi(x) + t\psi(y), \end{split}$$

where in the last inequality we have used z = ty. This is the desired inequality.

The inequality (3.16) is equivalent to

$$e^{-[\varphi(1-t)]\Box[\psi t]((1-t)x+ty)} \ge (e^{-\varphi(x)})^{1-t} (e^{-\psi(y)})^t.$$

Then, the Prékopa-Leindler inequality (2.6) deduces

$$\int_{\mathbb{R}^n} e^{-[\varphi(1-t)]\Box[\psi t](x)} dx \ge \left(\int_{\mathbb{R}^n} e^{-\varphi(x)} dx\right)^{1-t} \left(\int_{\mathbb{R}^n} e^{-\psi(y)} dx\right)^t.$$
(3.17)

If there is an equality in (3.16), then Hölder inequality yields

$$\int_{\mathbb{R}^n} e^{-[\varphi(1-t)]\Box[\psi t](x)} dx = \int_{\mathbb{R}^n} \left(e^{-\varphi(x)}\right)^{1-t} \left(e^{-\psi(x)}\right)^t dx.$$
$$\leq \left(\int_{\mathbb{R}^n} e^{-\varphi(x)} dx\right)^{1-t} \left(\int_{\mathbb{R}^n} e^{-\psi(y)} dx\right)^t.$$

This means that there is an equality in the Prékopa-Leindler inequality (3.17). The equality condition of Prékopa-Leindler inequality guarantees that convex functions φ and ψ agree up to a translation. Conversely, it is not hard to check that there is an equality in (3.16) if $\varphi(x) = \psi(x - x_0)$ for some $x_0 \in \mathbb{R}^n$.

We are now in the position to obtain the following Brunn-Minkowski inequality for the functional dual quermassintegrals.

Theorem 3.2. Let $\varphi, \psi \in \mathcal{L}_o$. If $\overline{W}_{n-q}(\varphi)$ and $\overline{W}_{n-q}(\psi)$ are finite and $q \leq -1$, for any $t \in (0, 1)$, then

$$\overline{W}_{n-q}([\varphi(1-t)]\Box[\psi t]) \le (1-t)\overline{W}_{n-q}(\varphi) + t\overline{W}_{n-q}(\psi)$$
(3.18)

with equality if and only if $\varphi(x) = \psi(x - x_0)$ with $x_0 \in \mathbb{R}^n$ when q = -1 and $\varphi = \psi$ when q < -1.

Proof. Since $q \leq -1$, (3.16) and Minkowski's inequality, we have

$$\overline{W}_{n-q}([\varphi(1-t)]\Box[\psi t]) \leq \left(\int_{\mathbb{R}^n} ((1-t)\varphi(x) + t\psi(y))^{-q} d\gamma_n(x)\right)^{\frac{1}{-q}} \leq (1-t)\overline{W}_{n-q}(\varphi) + t\overline{W}_{n-q}(\psi).$$

The equality conditions follow from the equality conditions of inequality (3.16) and Minkowski's inequality.

Similar to the proof of Theorem 3.2, we also obtain the case of q > -1 and $q \neq 0$. Corollary 3.1. Let $\varphi, \psi \in \mathcal{L}_o$. If $\overline{W}_{n-q}(\varphi)$ and $\overline{W}_{n-q}(\psi)$ are finite and q > -1 and $q \neq 0$, then

$$\overline{W}_{n-q}([\varphi(1-t)]\Box[\psi t]) \ge (1-t)\overline{W}_{n-q}(\varphi) + t\overline{W}_{n-q}(\psi)$$
with equality if and only if $\varphi = \psi$.
$$(3.19)$$

The following lemma is useful.

Lemma 3.7. Let $q \leq 0$ and $\varphi, \psi \in \mathcal{L}_o$. Then

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q}([\varphi(1-t)]\Box[\psi t]) - \widetilde{W}_{n+1-q}(\varphi)}{t} = (q-1)\widetilde{W}_{n+1-q}(\varphi,\psi) - (q-1)\widetilde{W}_{n+1-q}(\varphi,\varphi).$$

Proof. Let $a(t) = \frac{t}{1-t}$, for $t \in (0, 1)$. From the definition of infimal convolution (2.4), we have

$$\begin{split} \left[\varphi \Box [\psi a(t)]\right](1-t)(x) \\ &= (1-t)\left[\varphi \Box [\psi a(t)]\right]\left(\frac{x}{1-t}\right) \\ &= (1-t)\inf_{y\in\mathbb{R}^n}\left\{\varphi\left(\frac{x-(1-t)y}{1-t}\right) + \frac{t}{1-t}\psi\left(\frac{(1-t)y}{t}\right)\right\} \\ &= \inf_{y\in\mathbb{R}^n}\left\{(1-t)\varphi\left(\frac{x-y}{1-t}\right) + t\psi\left(\frac{y}{t}\right)\right\} \\ &= [\varphi(1-t)]\Box [\psi t](x), \end{split}$$

and

$$\frac{\widetilde{W}_{n+1-q}([\varphi(1-t)]\Box[\psi t]) - \widetilde{W}_{n+1-q}(\varphi)}{t} = \frac{\widetilde{W}_{n+1-q}([\varphi\Box[\psi a(t)]](1-t)) - \widetilde{W}_{n+1-q}(\varphi\Box[\psi a(t)))}{t} + \frac{\widetilde{W}_{n+1-q}(\varphi\Box[\psi a(t)) - \widetilde{W}_{n+1-q}(\varphi)}{t}.$$
(3.20)

We set $\varphi_{a(t)} = \varphi \Box [\psi a(t)]$, and

$$b_t(s) = \widetilde{W}_{n+1-q} \big(\varphi_{a(t)}(1-s) \big),$$

for $s \in (0, 1)$. By a direct calculation,

$$\frac{d}{ds}b_{t}(s) = (2\pi)^{-\frac{n}{2}}\frac{d}{ds}(1-s)^{1-q}\int_{\mathbb{R}^{n}}\left(\varphi_{a(t)}\left(\frac{x}{1-s}\right)\right)^{1-q}e^{-\frac{|x|^{2}}{2}}dx$$

$$= (2\pi)^{-\frac{n}{2}}\frac{d}{ds}(1-s)^{n+1-q}\int_{\mathbb{R}^{n}}\left(\varphi_{a(t)}\left(x\right)\right)^{1-q}e^{-\frac{(1-s)^{2}|x|^{2}}{2}}dx$$

$$= -(n+1-q)(2\pi)^{-\frac{n}{2}}(1-s)^{n-q}\int_{\mathbb{R}^{n}}\left(\varphi_{a(t)}\left(x\right)\right)^{1-q}e^{-\frac{(1-s)^{2}|x|^{2}}{2}}dx$$

$$+ (2\pi)^{-\frac{n}{2}}(1-s)^{n+1-q}\int_{\mathbb{R}^{n}}\left(\varphi_{a(t)}\left(x\right)\right)^{1-q}e^{-\frac{(1-s)^{2}|x|^{2}}{2}}(1-s)|x|^{2}dx.$$

Then for every fixed $t \in (0,1)$, by Lagrange theorem we have that there exists a $\bar{s} \in (0,1)$ such that

$$\frac{\widetilde{W}_{n+1-q}\left(\left[\varphi\Box[\psi a(t)]\right](1-t)\right) - \widetilde{W}_{n+1-q}(\varphi\Box[\psi a(t))]}{t} = b'_t(\bar{s}) \\
= -(n+1-q)(2\pi)^{-\frac{n}{2}}(1-\bar{s})^{n-q} \int_{\mathbb{R}^n} \left(\varphi_{a(t)}(x)\right)^{1-q} e^{-\frac{(1-\bar{s})^2|x|^2}{2}} dx \\
+ (2\pi)^{-\frac{n}{2}}(1-\bar{s})^{n+1-q} \int_{\mathbb{R}^n} \left(\varphi_{a(t)}(x)\right)^{1-q} e^{-\frac{(1-\bar{s})^2|x|^2}{2}}(1-\bar{s})|x|^2 dx$$

By Lemma 3.3, as $t \to 0^+$ functions $\varphi_{a(t)}$ converge decreasing to a convex function $\tilde{\varphi}$. Then, by monotone convergence theorem, $\bar{s} \to 0^+$ and $a(t) \to 0^+$ as $t \to 0^+$, we obtain

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q} \left(\left[\varphi \Box \left[\psi a(t) \right] \right] (1-t) \right) - \widetilde{W}_{n+1-q} \left(\varphi \Box \left[\psi a(t) \right] \right)}{t}$$
$$= -(n+1-q) \widetilde{W}_{n+1-q} \left(\widetilde{\varphi} \right) + \int_{\mathbb{R}^n} \widetilde{\varphi} \left(x \right)^{1-q} |x|^2 d\gamma_n(x).$$
(3.21)

Concerning the second addendum in (3.20), Definition 3.1 shows immediately that

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q}(\varphi \Box [\psi a(t)) - \widetilde{W}_{n+1-q}(\varphi)]}{t} = (q-1)\widetilde{W}_{n+1-q}(\varphi, \psi).$$
(3.22)

Recall that the functions $\varphi_{a(t)}$ converge decreasing to some convex positive function $\tilde{\varphi}$ as $t \to 0^+$. In order to obtain the desired equality, we may distinguish the two cases of $\widetilde{W}_{n+1-q}(\tilde{\varphi}) \leq \widetilde{W}_{n+1-q}(\varphi)$ and $\widetilde{W}_{n+1-q}(\tilde{\varphi}) = \widetilde{W}_{n+1-q}(\varphi)$. If $\widetilde{W}_{n+1-q}(\tilde{\varphi}) \leq \widetilde{W}_{n+1-q}(\varphi)$, the limit in (3.22) becomes $-\infty$, and the limit in (3.21) is finite, hence

$$\lim_{t \to 0^+} \frac{\widetilde{W}_{n+1-q}([\varphi(1-t)]\Box[\psi t]) - \widetilde{W}_{n+1-q}(\varphi)}{t} = -\infty$$

and the result of the lemma holds true. If $\widetilde{W}_{n+1-q}(\tilde{\varphi}) = \widetilde{W}_{n+1-q}(\varphi)$, then $\tilde{\varphi} = \varphi$ almost everywhere, so that the right hand side of (3.21) is $(1-q)\widetilde{W}_{n+1-q}(\varphi,\varphi)$. The lemma follows by summing up (3.21) and (3.22).

By Lemma 3.7, we obtain the following Minkowski inequality for the functional dual mixed quermassintegrals.

Theorem 3.3. Let $\varphi, \psi \in \mathcal{L}_o$. If $q \leq 0$, then

$$\widetilde{W}_{n+1-q}(\varphi,\psi) \ge \frac{\overline{W}_{n+1-q}(\psi) - \overline{W}_{n+1-q}(\varphi)}{\overline{W}_{n+1-q}(\varphi)^q} + \widetilde{W}_{n+1-q}(\varphi,\varphi),$$
(3.23)

with equality if and only if $\varphi(x) = \psi(x-x_0)$ with $x_0 \in \mathbb{R}^n$ when q = 0 and $\varphi(x) = \psi(x)$ when q < 0.

Proof. Let

$$\Upsilon(t) = \widetilde{W}_{n+1-q}([\varphi(1-t)]\Box[\psi t])^{\frac{1}{1-q}}.$$

From the Brunn-Minkowski style inequality (3.19), we noticed that $\Upsilon(t)$ is convex on [0, 1], hence

$$\Upsilon(t) \le \Upsilon(0) + t[\Upsilon(1) - \Upsilon(0)], \qquad (3.24)$$

for $t \in [0, 1]$. As a consequence, the derivative of the function Υ at t = 0 satisfies

$$\Upsilon'(0) \le \Upsilon(1) - \Upsilon(0). \tag{3.25}$$

Lemma 3.7 deduces that

$$\Upsilon'(0) = \widetilde{W}_{n+1-q}(\varphi)^{\frac{q}{1-q}} \left[\widetilde{W}_{n+1-q}(\varphi,\varphi) - \widetilde{W}_{n+1-q}(\varphi,\psi) \right].$$

Inequality (3.25) can be rewritten as

$$\widetilde{W}_{n+1-q}(\varphi)^{\frac{q}{1-q}} \left[\widetilde{W}_{n+1-q}(\varphi,\varphi) - \widetilde{W}_{n+1-q}(\varphi,\psi) \right] \\
\leq \widetilde{W}_{n+1-q}(\psi)^{\frac{1}{1-q}} - \widetilde{W}_{n+1-q}(\varphi)^{\frac{1}{1-q}}.$$
(3.26)

Finally, assume that $\varphi(x) = \psi(x - x_0)$ with $x_0 \in \mathbb{R}^n$ when q = 0 and $\varphi(x) = \psi(x)$ when q < 0, then (3.23) holds with equality sign. Conversely, assume that (3.23) holds with equality sign. By inspection of the above proof one sees immediately that also inequality (3.25), and hence inequality (3.24) and inequality (3.19), must hold with equality sign. This entails that (3.19) holds as an equality, and therefore, φ and ψ agree up to a translation when q = 0 and $\varphi(x) = \psi(x)$ when q < 0.

4. DUAL CURVATURE MEASURES FOR CONVEX FUNCTIONS

Theorem 3.1 provides a suitable way to define dual curvature measures for convex functions, and we will investigate the functional dual curvature measure in this section.

Definition 4.1. Let $q \in \mathbb{R}$ and $\varphi \in \mathcal{L}_o$. If $0 < \int_{\mathbb{R}^n} \varphi(x)^{-q} d\gamma_n(x) < +\infty$, then $\widetilde{C}_q(\varphi, \cdot)$, the q-th dual curvature measure of φ , is defined as

$$\int_{\mathbb{R}^n} g(z) d\widetilde{C}_q(\varphi, z) = \int_{\mathbb{R}^n} g(\nabla \varphi(x)) \varphi(x)^{-q} d\gamma_n(x), \qquad (4.1)$$

for every bounded continuous function $g: \mathbb{R}^n \to \mathbb{R}$. Here $d\gamma_n(x) = (2\pi)^{-\frac{n}{2}} e^{-\frac{|x|^2}{2}} dx$ is the Gaussian probability measure.

If φ is a \mathcal{C}^2 -smooth and strictly convex function, then the map

$$\nabla \varphi : \{ x \in \mathbb{R}^n : \varphi(x) < +\infty \} \to \{ x \in \mathbb{R}^n : \varphi^*(x) < +\infty \}$$

is smooth and bijective. By the formulas (see, e.g., [53]),

$$x = \nabla \varphi^*(\nabla \varphi(x))$$
 and $\nabla^2 \varphi^*(\nabla \varphi(x)) = (\nabla^2 \varphi(x))^{-1}$ (4.2)

for the points of $\{x \in \mathbb{R}^n : \varphi(x) < +\infty\}$ at which its Hessian $\nabla^2 \varphi$ exists and is invertible, we have

$$\begin{split} \int_{\Omega} g(x)d\widetilde{C}_{q}(\varphi,x) &= (2\pi)^{-\frac{n}{2}} \int_{\Omega} g(\nabla\varphi(x))\varphi(x)^{-q} e^{-\frac{|x|^{2}}{2}} dx \\ &= (2\pi)^{-\frac{n}{2}} \int_{\Omega} g(\nabla\varphi(x))\varphi(\nabla\varphi^{*}(\nabla\varphi(x)))^{-q} e^{-\frac{|\nabla\varphi^{*}(\nabla\varphi(x))|^{2}}{2}} dx \\ &= (2\pi)^{-\frac{n}{2}} \int_{\Omega} g(z)\varphi(\nabla\varphi^{*}(z))^{-q} e^{-\frac{|\nabla\varphi^{*}(z)|^{2}}{2}} \det(\nabla^{2}\varphi^{*}(z)) dz, \end{split}$$

where $\Omega = \{x \in \mathbb{R}^n : \varphi(x) < +\infty\} \cap \{x \in \mathbb{R}^n : \varphi^*(x) < +\infty\}$. Hence, if φ is a \mathcal{C}^2 -smooth and strictly convex function, then $\widetilde{C}_q(\varphi, \cdot)$ is absolutely continuous with respect to the *n*-dimensional Hausdorff measure and the Radon-Nikodym derivative is

$$\frac{d\hat{C}_q(\varphi, x)}{dx} = (2\pi)^{-\frac{n}{2}}\varphi(\nabla\varphi^*(x))^{-q}e^{-\frac{|\nabla\varphi^*(x)|^2}{2}}\det(\nabla^2\varphi^*(x)) \quad \text{on} \quad \Omega.$$

The following lemma shows that the Borel measure $\widetilde{C}_q(\varphi, \cdot)$ includes the Borel measure $\widetilde{C}_q(K, \cdot)$ of [35].

Lemma 4.1. Let $K \in \mathcal{K}_o^n$ be strictly convex and q < n. Then the Borel measure $\widetilde{C}_q(||x||_K, \cdot)$ includes the q-th dual curvature measure, $\widetilde{C}_q(K, \cdot)$, of K.

Proof. Let $K \in \mathcal{K}_o^n$ be strictly convex and $\varphi(x) = ||x||_K$. Let \overline{V}_K denote the normalized cone measure of K, which is given by

$$d\overline{V}_K(z) = \frac{\langle z, \nu_K(z) \rangle}{nV(K)} d\mathcal{H}^{n-1}(z) \quad \text{for} \quad z \in \partial K.$$

Here V(K) denotes the *n*-dimensional volume of K. For $x \in \mathbb{R}^n$, we write x = rz, with $z \in \partial K$, then $dx = nV(K)r^{n-1}drd\overline{V}_K(z)$. Since the map $x \mapsto \nabla ||x||_K$ is 0-homogeneous, hence

$$\begin{split} &\int_{\mathbb{R}^n} g(x) d\widetilde{C}_q(\varphi, x) \\ &= \int_{\mathbb{R}^n} g(\nabla \varphi(x)) \varphi(x)^{-q} d\gamma_n(x) \\ &= (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} g(\nabla \|x\|_K) \|x\|_K^{-q} e^{-\frac{|x|^2}{2}} dx \\ &= n(2\pi)^{-\frac{n}{2}} V(K) \int_0^\infty \int_{\partial K} g(\nabla \|z\|_K) r^{n-q-1} e^{-\frac{r^2|z|^2}{2}} d\overline{V}_K(z) dr \\ &= (2\pi)^{-\frac{n}{2}} 2^{\frac{n-q-2}{2}} nV(K) \int_0^\infty e^{-r} r^{\frac{n-q}{2}-1} dr \int_{\partial K} g(\nabla \|z\|_K) |z|^{q-n} d\overline{V}_K(z) \\ &= (2\pi)^{-\frac{n}{2}} 2^{\frac{n-q-2}{2}} \int_0^\infty e^{-r} r^{\frac{n-q}{2}-1} dr \int_{\partial K} g(\nabla \|z\|_K) |z|^{q-n} h_K(\nu_K(z)) d\mathcal{H}^{n-1}(z). \end{split}$$

Because of (see, e.g., [62]),

$$\nabla \|z\|_{K} = \frac{\nu_{K}(z)}{\|\nu_{K}(z)\|_{K^{\circ}}},$$

for $z \in \partial K$, then

$$\int_{\mathbb{R}^n} g(x) d\widetilde{C}_q(\varphi, x) = (2\pi)^{-\frac{n}{2}} 2^{\frac{n-q-2}{2}} \int_0^\infty e^{-r} r^{\frac{n-q}{2}-1} dr \int_{\partial K} g\left(\frac{\nu_K(z)}{\|\nu_K(z)\|_{K^\circ}}\right) |z|^{q-n} h_K(\nu_K(z)) d\mathcal{H}^{n-1}(z).$$

This means that the Borel measure $\widetilde{C}_q(||x||_K, \cdot)$ takes the following form: For any Borel subsets $I \subseteq [0, \infty)$ and $\Omega \subseteq \partial K$,

$$\widetilde{C}_q(\varphi, I \times \Omega) = \nu_1(I)\nu_2(\Omega).$$

The measure ν_1 is independent of the choice of K and it is a measure with density

$$(2\pi)^{-\frac{n}{2}}2^{\frac{n-q-2}{2}}e^{-r}r^{\frac{n-q}{2}-1}.$$

The measure ν_2 is a measure on ∂K which implies the *q*-th dual curvature measure, $\widetilde{C}_q(K, \cdot)$, of K. In fact, if $\mathcal{R}(x) = \frac{x}{|x|}$ for $x \in \mathbb{R}^n \setminus \{0\}$, then

$$\mathcal{R}_{*}(\nu_{2})$$

is referred to as the dual curvature measure, $\widetilde{C}_q(K, \cdot)$, of K, and $\mathcal{R}_*(\nu_2)$ is the pushforward of ν_2 via \mathcal{R} . This shows that the dual curvature measure of a convex body can be recovered from the functional dual curvature measure of a particular convex function.

The next lemma will be used later.

Lemma 4.2. [21, Lemma 3] Let $\varphi : \mathbb{R}^n \to [0, +\infty]$ be an essentially-continuous, convex function. Fix a vector $0 \neq \theta \in \mathbb{R}^n$ and let $H = \theta^{\perp} \subset \mathbb{R}^n$ be the hyperplane orthogonal to θ . Then, for \mathcal{H}^{n-1} -almost every $y \in H$, the function

$$\mathbb{R} \to \mathbb{R} \cup \{+\infty\}$$
$$t \mapsto \varphi(y + t\theta).$$

is continuous on \mathbb{R} , and locally-Lipschitz in the interior of the interval in which it is finite.

The support $\text{Supp}(\mu)$ of a measure μ in \mathbb{R}^n is a closed set that contains all points $x \in \mathbb{R}^n$ with $\mu(U) > 0$ for any open set U containing x.

Lemma 4.3. Let $\varphi : \mathbb{R}^n \to [0, +\infty]$ be an essentially-continuous, convex function and $q \leq 0$. If

$$0 < \int_{\mathbb{R}^n} \varphi^{1-q}(x) d\gamma_n(x) < +\infty \quad and \quad \varphi(0) = 0,$$

then the q-th dual curvature measure of φ is not supported in any hyperplane.

Proof. Assume that $\operatorname{Supp}(\widetilde{C}_q(\varphi, \cdot)) \subseteq \theta^{\perp}$ for a unit vector $\theta \in S^{n-1}$. Without loss of generality, assume $\theta = e_n$, where $e_n = (0, \dots, 0, 1)$. For $x \in \mathbb{R}^n$, we write x = (y, t) with $y \in \mathbb{R}^{n-1}$ and $t = x_n \in \mathbb{R}$. Suppose $\varphi(x) > 0$ for $x \in \mathbb{R}^n$, then

$$0 = \int_{\mathbb{R}^n} |z_n| d\widetilde{C}_q(\varphi, z)$$

=
$$\int_{\mathbb{R}^n} \left| \frac{\partial \varphi(x)}{\partial x_n} \right| \varphi(x)^{-q} d\gamma_n(x)$$

=
$$\frac{1}{1-q} \int_{\mathbb{R}^n} \left| \frac{\partial \varphi^{1-q}(x)}{\partial x_n} \right| d\gamma_n(x).$$

For almost every $(y, t) \in \mathbb{R}^n$,

$$\frac{\partial \varphi^{1-q}(y,t)}{\partial t} = 0$$

Since $q \leq 0$, the function φ^{1-q} is also convex. According to Lemma 4.2, for almost any $y \in \mathbb{R}^{n-1}$, the function $t \to \varphi^{1-q}(y,t)$ is continuous in \mathbb{R} and locally-Lipschitz in the interior of the interval $\{t : \varphi(y,t) < +\infty\}$. Therefore, for almost any $y \in \mathbb{R}^{n-1}$, the convex function $t \to \varphi^{1-q}(y,t)$ is constant in \mathbb{R} . The condition $\varphi(0) = 0$ implies that $\varphi \equiv 0$ in \mathbb{R} and so that the integral $\int \varphi^{1-q}(y,t) e^{-\frac{t^2}{2}} dt$ equals to 0. By Fubini theorem, the function φ^{1-q} can not have a finite, non-zero integral. This is a contradiction. \Box

From Jensen's inequality we know that

$$-q \mapsto \left(\int_{\mathbb{R}^n} \varphi(x)^{-q} d\gamma_n(x)\right)^{-\frac{1}{q}}$$

is strictly monotone increasing, unless φ is constant on \mathbb{R}^n . Hence, the condition in Lemma 4.3 includes the assumption in the Definition 4.1 when $q \leq 0$.

A function Z defined on a lattice $(\mathcal{C}, \lor, \land)$ and taking values in an abelian semigroup is called a *valuation* if

$$Z(f \lor g) + Z(f \land g) = Z(f) + Z(g), \tag{4.3}$$

for all $f, g \in C$. A function Z defined on a set $S \subset C$ is called a valuation if (4.3) holds whenever $f, g, f \lor g, f \land g \in S$.

For S the space of convex bodies, \mathcal{K}^n , in \mathbb{R}^n with \vee denoting union and \wedge intersection, the notion of valuation is classical. The classical valuation played a critical role in Dehn's solution of Hilbert's Third Problem and have been a central focus in convex geometric analysis. In the 1930s, Blaschke classified the real-valued valuations on convex bodies that are SL(n) invariant. Blaschke's work was greatly extended by Hadwiger in his famous classification of continuous, rigid motion invariant valuations and characterization of elementary mixed volumes.

In [35], authors proved that dual curvature measures are valuations, i.e.,

$$\widetilde{C}_q(K,\cdot) + \widetilde{C}_q(L,\cdot) = \widetilde{C}_q(K \cap L, \cdot) + \widetilde{C}_q(K \cup L, \cdot),$$

for each $K, L \in \mathcal{K}_o^n$ such that $K \cup L \in \mathcal{K}_o^n$.

Recently, valuations were defined on function spaces. For a space S of real-valued functions we denote by $f \vee g$ the pointwise maximum of f and g while $f \wedge g$ denotes their pointwise minimum. The characteristic functions χ_K, χ_L of convex bodies K and L satisfy

$$\chi_{K\cup L} = \max\{\chi_K, \chi_L\} \text{ and } \chi_{K\cap L} = \min\{\chi_K, \chi_L\},\$$

for all $K, L \in \mathcal{K}_o^n$ such that $K \cup L \in \mathcal{K}_o^n$. Valuations on function spaces can be seen as a generalization of valuations on \mathcal{K}_o^n . For the classification of valuations on function spaces one can refer to [18-20, 43, 44].

We are ready to prove that the q-th dual curvature measure $\widetilde{C}_q(\varphi, \cdot)$ is a valuation.

Lemma 4.4. Let $\varphi, \psi \in \mathcal{L}_o$, and $q \in \mathbb{R}$. If $\min\{\varphi, \psi\}$ is convex, then

$$\widetilde{C}_q(\varphi, \cdot) + \widetilde{C}_q(\psi, \cdot) = \widetilde{C}_q(\min\{\varphi, \psi\}, \cdot) + \widetilde{C}_q(\max\{\varphi, \psi\}, \cdot).$$
(4.4)

Proof. If $\min\{\varphi, \psi\}$ is convex, then for almost every $x \in \mathbb{R}^n$,

$$\nabla(\max\{\varphi,\psi\})(x) = \begin{cases} \nabla\varphi(x) & \text{when } \varphi(x) > \psi(x) \\ \nabla\psi(x) & \text{when } \varphi(x) < \psi(x) \\ \nabla\varphi(x) = \nabla\psi(x) & \text{when } \varphi(x) = \psi(x) \end{cases},$$

and

$$\nabla(\min\{\varphi,\psi\})(x) = \begin{cases} \nabla\varphi(x) & \text{when } \varphi(x) < \psi(x) \\ \nabla\psi(x) & \text{when } \varphi(x) > \psi(x) \\ \nabla\varphi(x) = \nabla\psi(x) & \text{when } \varphi(x) = \psi(x) \end{cases}.$$

Therefore,

$$\begin{split} &\int_{\mathbb{R}^n} g(y) d\widetilde{C}_q(\max\{\varphi,\psi\},y) + \int_{\mathbb{R}^n} g(y) d\widetilde{C}_q(\min\{\varphi,\psi\},y) \\ &= \int_{\mathbb{R}^n} g(\nabla(\max\{\varphi,\psi\})(y))(\max\{\varphi,\psi\})^{-q} d\gamma_n(y) \\ &\quad + \int_{\mathbb{R}^n} g(\nabla(\min\{\varphi,\psi\})(y))(\min\{\varphi,\psi\})^{-q} d\gamma_n(y) \\ &= \int_{\mathbb{R}^n \cap \{\varphi \geq \psi\}} g(\nabla(\max\{\varphi,\psi\})(y))(\max\{\varphi,\psi\})^{-q} d\gamma_n(y) \\ &\quad + \int_{\mathbb{R}^n \cap \{\varphi < \psi\}} g(\nabla(\max\{\varphi,\psi\})(y))(\max\{\varphi,\psi\})^{-q} d\gamma_n(y) \\ &\quad + \int_{\mathbb{R}^n \cap \{\varphi \geq \psi\}} g(\nabla(\min\{\varphi,\psi\})(y))(\min\{\varphi,\psi\})^{-q} d\gamma_n(y) \\ &\quad + \int_{\mathbb{R}^n \cap \{\varphi < \psi\}} g(\nabla(\min\{\varphi,\psi\})(y))(\min\{\varphi,\psi\})^{-q} d\gamma_n(y) \\ &\quad + \int_{\mathbb{R}^n \cap \{\varphi \geq \psi\}} g(\nabla\varphi(y))\varphi^{-q} d\gamma_n(y) + \int_{\mathbb{R}^n \cap \{\varphi < \psi\}} g(\nabla\psi)(y))\psi^{-q} d\gamma_n(y) \\ &\quad + \int_{\mathbb{R}^n \cap \{\varphi \geq \psi\}} g(\nabla\psi(y))\psi^{-q} d\gamma_n(y) + \int_{\mathbb{R}^n \cap \{\varphi < \psi\}} g(\nabla\varphi(y))\varphi^{-q} d\gamma_n(y) \\ &\quad = \int_{\mathbb{R}^n} g(y)d\widetilde{C}_q(\varphi,y) + \int_{\mathbb{R}^n} g(y)\psi^{-q} d\widetilde{C}_q(\psi,y), \end{split}$$

for all continuous integrable function g. Hence, we obtain the desired valuation property (4.4).

5. Dual Minkowski problems for convex functions

5.1. A minimization problem.

We write $L_1(\mu)$ to denotes the class of μ -integrable functions $f : \mathbb{R}^n \to [0, \infty]$, i.e.,

$$L_1(\mu) = \left\{ f : \int_{\mathbb{R}^n} f(x) d\mu(x) < +\infty \right\}.$$

Let μ be a finite Borel measure on \mathbb{R}^n and $q \in \mathbb{R}$. We set

$$\Psi_{\mu}(f) = \frac{1}{|\mu|} \int_{\mathbb{R}^n} f(x) d\mu(x) - e^{-\overline{W}_{n+1-q}(f^*)}, \qquad (5.1)$$

for $f \in L_1(\mu)$ with f(0) = 0. Here $|\mu|$ denotes the total measure of μ and $\overline{W}_{n+1-q}(f^*)$ is the normalized dual quermassintegral, namely,

$$\overline{W}_{n+1-q}(f^*) = \left[\int_{\mathbb{R}^n} (f^*(x))^{1-q} d\gamma_n(x) \right]^{\frac{1}{1-q}}.$$

We should point out that the restriction f(0) = 0 guarantees that $f^*(x) \ge 0$, since (2.3) and the fact $f^{**} \le f$.

We consider the minimization problem,

$$\inf_{f} \left\{ \Psi_{\mu}(f) : 0 < \overline{W}_{n+1-q}(f^{*}) < \infty, f \in L_{1}(\mu) \text{ and } f(0) = 0 \right\}.$$
(5.2)

If the convex function φ_0 solves the infimum in (5.2), then the pre-given measure μ is exactly the *q*-th dual curvature of φ_0^* . We need the following result which has been provided in [5] and a special case one can found in [17]. For a complete proof one can see in the recent paper [61].

Lemma 5.1. Let φ be a lower semi-continuous convex function in \mathbb{R}^n and let g be a bounded continuous function in \mathbb{R}^n . If $\varphi_t(x) = \varphi(x) + tg(x)$ for sufficiently small t (i.e., |t| < a for sufficiently small a) and $x \in \mathbb{R}^n$, then

$$\frac{d}{dt}\varphi_t^*(x)\Big|_{t=0} = -g(\nabla\varphi^*(x))$$

at any point $x \in \mathbb{R}^n$ in which φ^* is differentiable.

The next shows that if φ_0 solves the optimal problem (5.2) then the pre-given measure μ is the q-th dual curvature measure of φ_0^* .

Lemma 5.2. Let μ be a finite Borel measure on \mathbb{R}^n that is not supported in a lower dimensional subspace with $\int_{\mathbb{R}^n} |x| d\mu(x) < \infty$ and $q \leq 0$. If there exists a lower semi-continuous and convex μ -integrable function $\varphi_0 : \mathbb{R}^n \to \mathbb{R}$ such that

$$\Psi_{\mu}(\varphi_0) = \inf_{f} \left\{ \Psi_{\mu}(f) : f \in L_1(\mu), 0 < W_{n+1-q}(f^*) < \infty, f(0) = 0 \right\},\$$

then

$$\mu = \tau \widetilde{C}_q(\varphi_0^*, \cdot),$$

with $\tau = \frac{|\mu|}{\widetilde{W}_{n-q}(\varphi_0^*)}$.

Proof. We assume that there exists a φ_0 with $0 < \overline{W}_{n+1-q}(\varphi_0^*) < \infty$ such that it is a minimizer for $\Psi_{\mu}(f)$, i.e.,

$$\Psi_{\mu}(\varphi_0) = \inf_{f} \left\{ \Psi_{\mu}(f) : f \in L_1(\mu), 0 < \overline{W}_{n+1-q}(f^*) < \infty, f(0) = 0 \right\}.$$
(5.3)

For any μ -integrable continuous, bounded function $g : \mathbb{R}^n \to \mathbb{R}$ with g(0) = 0, we set $\varphi_t(x) = \varphi_0(x) + tg(x)$. Since g is bounded, if |g| < M then we can select a $t_0 > 0$ such that

$$\varphi_0 - t_0 M < \varphi_t < \varphi_0 + t_0 M, \tag{5.4}$$

and

$$\frac{1}{|\mu|} \int_{\mathbb{R}^n} \varphi_t(x) d\mu(x) \le \frac{1}{|\mu|} \int_{\mathbb{R}^n} \varphi_0(x) d\mu(x) + t_0 M,$$

for all $t \in [-t_0, t_0]$. This means that there exists $t_0 > 0$ such that φ_t is μ -integrable on \mathbb{R}^n and $\varphi_t(0) = 0$ for all $t \in [-t_0, t_0]$. Inequalities (5.4) imply that

$$\varphi_0^* - t_0 M < \varphi_t^* < \varphi_0^* + t_0 M, \tag{5.5}$$

for all $t \in [-t_0, t_0]$. Hence, the Minkowski inequality deduces

$$\overline{W}_{n+1-q}(\varphi_t^*) \le \overline{W}_{n+1-q}(\varphi_0^*) + t_0 M < +\infty.$$

Because of φ_0 is a minimizer of the optimal problem (5.3),

$$\Psi_{\mu}(\varphi_0) \leq \Psi_{\mu}(\varphi_t) \text{ for all } t \in [-t_0, t_0].$$

Hence,

$$\left. \frac{d}{dt} \Psi_{\mu}(\varphi_t) \right|_{t=0} = 0.$$

By Lemma 5.1 deduces

$$\frac{d}{dt}\varphi_t^*(x)\Big|_{t=0} = -g(\nabla\varphi_0^*(x))$$

at any point $x\in\mathbb{R}^n$ in which φ_0^* is differentiable. By the dominated convergence theorem, we have

$$\begin{aligned} \left. \frac{d}{dt} \Psi_{\mu}(\varphi_t) \right|_{t=0} &= \left. \frac{d}{dt} \left(\frac{1}{|\mu|} \int_{\mathbb{R}^n} \varphi_t(x) d\mu(x) \right) \right|_{t=0} - \frac{d}{dt} \left(e^{-\overline{W}_{n+1-q}(\varphi_t^*)} \right) \Big|_{t=0} \\ &= \left. \frac{1}{|\mu|} \int_{\mathbb{R}^n} g(x) d\mu(x) \right. \\ &\left. -\overline{W}_{n+1-q}(\varphi_0^*)^q e^{-\overline{W}_{n+1-q}(\varphi_0^*)} \int_{\mathbb{R}^n} g(\nabla \varphi_0^*(x)) (\varphi_0^*(x))^{-q} d\gamma_n(x) \right] \end{aligned}$$

for any μ -integrable continuous, bounded function $g: \mathbb{R}^n \to \mathbb{R}$. This means

$$\mu(\cdot) = |\mu| \overline{W}_{n+1-q}(\varphi_0^*)^q e^{-\overline{W}_{n+1-q}(\varphi_0^*)} \widetilde{C}_q(\varphi_0^*, \cdot).$$
(5.6)

By taking the integration from both sides of (5.6), one sees that

$$\frac{1}{\widetilde{W}_{n-q}(\varphi_0^*)} = \overline{W}_{n+1-q}(\varphi_0^*)^q e^{-\overline{W}_{n+1-q}(\varphi_0^*)}.$$

This finishes the proof.

Next, we prove that the infimum in (5.2) is attained. The following lemma can be found in [21].

Lemma 5.3. Let μ be a non-zero finite Borel measure on \mathbb{R}^n that is not supported in a lower-dimensional subspace, and let M be the interior of $\operatorname{conv}(\operatorname{Supp}(\mu))$. If $x_0 \in M$, then there exists $C_{\mu,x_0} > 0$ with the following property: For any non-negative, μ -integrable, convex function $\varphi : \mathbb{R}^n \to [0, \infty]$,

$$\varphi(x_0) \le C_{\mu,x_0} \int_{\mathbb{R}^n} \varphi d\mu$$

26

Proposition 5.1. Let μ be a finite Borel measure on \mathbb{R}^n with $\int_{\mathbb{R}^n} |x| d\mu(x) < \infty$ that is not supported in a lower dimensional subspace. If $q \leq 0$, then there exists a nonnegative convex, lower semi-continuous and μ -integrable function $\varphi : \mathbb{R}^n \to \mathbb{R}$ with $\varphi(0) = 0$ and $\overline{W}_{n+1-q}(\varphi^*)$ is finite such that

$$\Psi_{\mu}(\varphi) = \inf_{f} \left\{ \Psi_{\mu}(f) : 0 < \overline{W}_{n+1-q}(f^{*}) < \infty, f \in L_{1}(\mu), f(0) = 0 \right\}.$$
 (5.7)

Proof. Since $f^{**} \leq f$ and $f^{***} = f^*$ for any function f, it yields

$$\Psi_{\mu}(f^{**}) \leq \Psi_{\mu}(f).$$

We can just consider the infimum of Ψ_{μ} by restricting our attention to the non-negative, convex and lower semi-continuous functions.

Let $\tilde{f}(x) = |x|$ and set $c_{\mu} = \Psi_{\mu}(\tilde{f})$. Then c_{μ} is a finite real number. Let $\varphi_1, \varphi_2, \cdots$ be a minimizing sequence μ -integrable convex functions with $\varphi_i(0) = 0$ $(i = 1, 2, \cdots)$, i.e.,

$$\lim_{i \to \infty} \Psi_{\mu}(\varphi_i) = \inf_{f} \left\{ \Psi_{\mu}(f) : 0 < \overline{W}_{n+1-q}(f^*) < \infty, f(0) = 0 \right\} \le c_{\mu}.$$
 (5.8)

Furthermore, we may remove finitely many elements from the sequence $\{\varphi_i\}$ and assume that for all i,

$$\Psi_{\mu}(\varphi_i) \le c_{\mu}.\tag{5.9}$$

Since $\overline{W}_{n+1-q}(\varphi_i^*) > 0$ for all *i*, therefore

$$\frac{1}{|\mu|} \int_{\mathbb{R}^n} \varphi_i d\mu \le c_\mu + 1, \tag{5.10}$$

for all i. Let M be the interior of $conv(Supp(\mu))$. By Lemma 5.3 and (5.10), we have

$$\varphi_i(x_0) \le (c_\mu + 1)C_{\mu,x_0},$$
(5.11)

for all *i* and any $x_0 \in M$. Theorem A ensures that there exists a subsequence $\{\varphi_{i_j}\}_{j=1,2,\dots}$ that converges pointwise in M to a convex function $\varphi : M \to \mathbb{R}$. Additionally, φ is non-negative in M with $\varphi(0) = 0$.

We extend the definition of φ by setting $\varphi(x) = +\infty$ for $x \notin \overline{M}$. We still need to define φ for $x \in \partial M$. Set

$$\varphi(x) = \lim_{\lambda \to 1^{-}} \varphi(\lambda x), \quad x \in \partial M.$$
 (5.12)

Since the function $\lambda \to \varphi(\lambda x)$ is non-decreasing for $\lambda \in (0, 1)$, hence the limit in (5.12) always exists in $[0, +\infty]$. We need to show that φ is μ -integrable. For any $x \in \overline{M}$, the point $\lambda x \in M$ for $0 < \lambda < 1$, and by the pointwise convergence in M,

$$\varphi(\lambda x) = \lim_{j \to \infty} \varphi_{i_j}(\lambda x), \tag{5.13}$$

for any $x \in \overline{M}$. Since the measure μ does not support in a lower-dimensional subspace, so that $0 \in \text{Supp}(\mu) \subseteq \overline{M}$, then from (5.13), (5.10), for any $0 < \lambda < 1$, by the

convexity and $\varphi_i(0) = 0$,

$$\begin{aligned} \varphi_{i_j}(\lambda x) &\leq \lambda \varphi_{i_j}(x) + (1-\lambda)\varphi_{i_j}(0) \\ &\leq \varphi_{i_j}(x), \end{aligned}$$

by Fatou's Lemma yields

$$\int_{\mathbb{R}^{n}} \varphi(\lambda x) d\mu(x) \leq \liminf_{j \to \infty} \int_{\mathbb{R}^{n}} \varphi_{i_{j}}(\lambda x) d\mu(x) \\
\leq \liminf_{j \to \infty} \int_{\mathbb{R}^{n}} \varphi_{i_{j}}(x) d\mu(x).$$
(5.14)

Recall that we have $\varphi(\lambda x) \nearrow \varphi(x)$ as $\lambda \to 1^-$. From the monotone convergence theorem and (5.14),

$$\int_{\mathbb{R}^n} \varphi(x) d\mu(x) = \lim_{\lambda \to 1^-} \int_{\mathbb{R}^n} \varphi(\lambda x) d\mu(x)$$

$$\leq \liminf_{j \to \infty} \int_{\mathbb{R}^n} \varphi_{i_j}(x) d\mu(x)$$

$$< +\infty.$$

Hence, φ is μ -integrable.

By the continuity of φ in M, we can choose a dense sequence x_1, x_2, \cdots in M such that

$$\varphi^*(y) = \sup_{x \in \mathbb{R}^n} \{ \langle x, y \rangle - \varphi(x) \}$$

=
$$\sup_{x \in M} \{ \langle x, y \rangle - \varphi(x) \}$$

=
$$\sup_{i \ge 1} \{ \langle x_i, y \rangle - \varphi(x_i) \},$$

for $y \in \mathbb{R}^n$. For $j \geq 1$, we set $\tilde{\varphi}_j^*(y) = \max_{1 \leq i \leq j} \{ \langle x_i, y \rangle - \varphi(x_i) \}$. We notice that $\tilde{\varphi}_j^* \nearrow \varphi^*$, then the increment of function t^{1-q} when $q \leq 0$ and the monotone convergence theorem deduce

$$\overline{W}_{n+1-q}(\varphi^*) = \lim_{j \to \infty} \overline{W}_{n+1-q}(\tilde{\varphi}_j^*).$$

For any $\varepsilon > 0$, there exists j_0 such that

$$\left|\overline{W}_{n+1-q}(\varphi^*) - \overline{W}_{n+1-q}(\tilde{\varphi}_{j_0}^*)\right| \leq \frac{\varepsilon}{2}.$$

By the fact $\varphi_{i_j} \to \varphi$ pointwise on the set $\{x_1, \dots, x_{j_0}\}$, then for sufficiently large j, we have

$$|\varphi_{i_j}^*(x) - \tilde{\varphi}_{j_0}^*(x)| \le \frac{\varepsilon}{2},\tag{5.15}$$

for all $x \in \mathbb{R}^n$. Since q < 0, then Minkowski's inequality deduces

$$\overline{W}_{n+1-q}(\varphi^*) \leq \overline{W}_{n+1-q}(\tilde{\varphi}_{j_0}^*) + \frac{\varepsilon}{2} \\
\leq \liminf_{j \to \infty} \left(\int_{\mathbb{R}^n} (\varphi_{i_j}^*(x) + \frac{\varepsilon}{2})^{1-q} d\gamma_n(x) \right)^{\frac{1}{1-q}} + \frac{\varepsilon}{2} \\
\leq \liminf_{j \to \infty} \overline{W}_{n+1-q}(\varphi_{i_j}^*) + \varepsilon.$$

This means

$$\overline{W}_{n+1-q}(\varphi^*) \leq \liminf_{j \to \infty} \overline{W}_{n+1-q}(\varphi^*_{i_j}) < +\infty$$

and

$$-e^{-\overline{W}_{n+1-q}(\varphi^*)} \leq \liminf_{j \to \infty} -e^{-\overline{W}_{n+1-q}(\varphi^*_{i_j})}.$$
(5.16)

Together (5.16) with (5.14), we have

$$\Psi_{\mu}(\varphi) \le \liminf_{i \to \infty} \Psi_{\mu}(\varphi_i) \tag{5.17}$$

Therefore, there exists a non-negative, μ -integrable, convex function φ with $\varphi(0) = 0$ such that

$$\Psi_{\mu}(\varphi) = \inf_{f} \left\{ \Psi_{\mu}(f) : f \in L_{1}(\mu), f(0) = 0 \right\}.$$

This completes our proof.

We are now in the position to prove the existence of solution to the functional dual Minkowski problem.

Theorem 5.1. Let μ be a non-zero finite Borel measure on \mathbb{R}^n with $\int_{\mathbb{R}^n} |x| d\mu(x) < \infty$ which is not supported in a lower-dimensional subspace and $q \leq 0$. There exists a non-negative convex function φ with $\varphi(0) = 0$ and $\int_{\mathbb{R}^n} \varphi(x)^{1-q} d\gamma_n(x)$ is finite such that $\mu(\cdot) = \tau \widetilde{C}_q(\varphi, \cdot)$.

Proof. The proof follows from Lemma 5.2 and Proposition 5.1.

5.2. Uniqueness of the dual functional Minkowski problem.

In this subsection we will show that if the dual curvature measures of two convex functions are equal under some asymptions, then the convex functions are also equal almost everywhere in \mathbb{R}^n . The tool we will be used is the Minkowski inequality (3.23), similar to the uniqueness part of the L_p Minkowski problem for convex bodies.

We are ready to prove the uniqueness of the solution to the functional dual Minkowski problems.

Theorem 5.2. Let $q \leq 0$ and $\varphi_1, \varphi_2 \in \mathcal{L}'_o$ with $\varphi_1(0) = \varphi_2(0) = 0$. Assume that φ_1 is an admissible perturbation for φ_2 and φ_2 is also an admissible perturbation for φ_1 . If $\widetilde{C}_{q-1}(\varphi_1, \cdot) = \widetilde{C}_{q-1}(\varphi_2, \cdot)$, then $\varphi_1(x) = \varphi_2(x - x_0)$ for some $x_0 \in \mathbb{R}^n$ when q = 0 and $\varphi_1(x) = \varphi_2(x)$ when q < 0.

Proof. Firstly notice that the equality $\widetilde{C}_{q-1}(\varphi_1, \cdot) = \widetilde{C}_{q-1}(\varphi_2, \cdot)$ implies $\overline{W}_{n+1-q}(\varphi_1) = \overline{W}_{n+1-q}(\varphi_2)$. By the fact $\overline{W}_{n+1-q}(\varphi_1) = \overline{W}_{n+1-q}(\varphi_2)$ and (3.23), we have

$$W_{n+1-q}(\varphi_1,\varphi_2) \ge W_{n+1-q}(\varphi_1,\varphi_1), \tag{5.18}$$

and

$$\widetilde{W}_{n+1-q}(\varphi_2,\varphi_1) \ge \widetilde{W}_{n+1-q}(\varphi_2,\varphi_2).$$
(5.19)

On the other hand, since $\widetilde{C}_{q-1}(\varphi_1, \cdot) = \widetilde{C}_{q-1}(\varphi_2, \cdot)$, then Theorem 3.1 yields

$$\widetilde{W}_{n+1-q}(\varphi_1,\psi) = \widetilde{W}_{n+1-q}(\varphi_2,\psi),$$

for any convex function ψ with $\psi(0) = 0$. In particular, taking $\psi = \varphi_1$ or $\psi = \varphi_2$, one sees that

$$\widetilde{W}_{n+1-q}(\varphi_1,\varphi_1) = \widetilde{W}_{n+1-q}(\varphi_2,\varphi_1), \qquad (5.20)$$

and

$$\widetilde{W}_{n+1-q}(\varphi_2,\varphi_2) = \widetilde{W}_{n+1-q}(\varphi_1,\varphi_2).$$
(5.21)

Equalities (5.20) and (5.21) guarantee the inequalities (5.18) and (5.19) both hold with equality. Finally, the equality condition of (3.23) has make sure that $\varphi_1(x) = \varphi_2(x - x_0)$ for some $x_0 \in \mathbb{R}^n$ when q = 0 and $\varphi_1(x) = \varphi_2(x)$ when q < 0.

References

- D. Alonso-Gutiérrez, B. Merino, C. H. Jiménez and R. Villa, Rogers-Shephard inequality for log-concave functions, J. Funct. Anal., 271 (2016) 3269-3299.
- [2] D. Alonso-Gutiérrez, B. Merino, C. H. Jiménez and R. Villa, John's ellipsoid and the integral raito of a log-concave function, J. Geom. Anal., 28 (2018) 1182-1201.
- [3] S. Artstein-Avidan, B. Klartag and V. Milman, The Santaló point of a function, and a functional form of Santaló inequality, Mathematika, 51 (2004) 33-48.
- [4] S. Artstein-Avidan, B. Klartag, C. Schütt and E. Werner, Functional affine-isoperimetry and an inverse logarithmic Sobolev inequality, J. Funct. Anal., 262 (2012) 4181-4204.
- [5] R. J. Berman and B. Berndtsson, Real Monge-Ampére equations and Kähler-Ricci solitons on toric log Fano varieties, Ann. Fac. Sci. Toulouse Math., 22 (2013), no. 4, 649-711.
- [6] K. Böröczky, M. Henk and H. Pollehn, Subspace concentration of dual curvature measures of symmetric convex bodies, J. Differ. Geom., 109 (2018) 411-429.
- [7] K. Böröczky, E. Lutwak, D. Yang and G. Zhang, The logarithmic Minkowski problem, J. Amer. Math. Soc., 26 (2013) 831-852.
- [8] K. Böröczky, E. Lutwak, D. Yang, G. Zhang and Y. Zhao, The dual Minkowski problem for symmetric convex bodies, arXiv:1703.06259v1.
- [9] L. Caffarelli, Interior W^{2,p}-estimates for solutions of the Monge-Ampère equation, Ann. Math., 131 (1990) 135-150.
- [10] U. Caglar, M. Fradelizi, O. Guedon, J. Lehec, C. Schuett and E. Werner, Functional versions of L_p-affine surface area and entropy inequalities, Int. Math. Res. Not., 2016 (4) (2016) 1223-1250.
- [11] C. Chen, Y. Huang and Y. Zhao, Smooth solutions to the L_p dual Minkowski problem, Math. Ann., (2018). https://doi.org/10.1007/s00208-018-1727-3
- [12] S. Y. Cheng and S. T. Yau, On the regularity of the solution of the *n*-dimensional Minkowski problem, Comm. Pure Appl. Math., 29 (1976) 495-516.

- [13] K. Chou and X. Wang, The L_p -Minkowski problem and the Minkowski problem in centro-affine geometry, Adv. Math., 205 (2006) 33-83.
- [14] A. Cianchi, E. Lutwak, D. Yang and G. Zhang, Affine Moser-Trudinger and Morrey-Sobolev inequalities, Calc. Var. Partial Differ. Equ., 36 (2009) 419-436.
- [15] A. Colesanti, Functional inequalities related to the Rogers-Shephard inequality, Mathematika, 53 (2006) 81-101.
- [16] A. Colesanti, Brunn-Minkowski inequalities for variational functionals and related problems, Adv. Math., 194 (2005) 105-140.
- [17] A. Colesanti and I. Fragalà, The first variation of the total mass of log-concave functions and related inequalities, Adv. Math., 244 (2013) 708-749.
- [18] A. Colesanti, N. Lombardi and L. Parapatits, Translation invariant valuations on quasi-concave functions, Studia Math., 243 (2018), no.1, 79 - 99.
- [19] A. Colesanti, M. Ludwig and F. Mussnig, Valuations on convex functions, Int. Math. Res. Not., rnx189, https://doi.org/10.1093/imrn/rnx189.
- [20] A. Colesanti, M. Ludwig and F. Mussnig, Minkowski valuations on convex functions, Calc. Var. Partial Differ. Equ., 56 (2017) no. 6, Art. 162, 29pp.
- [21] D. Cordero-Erausquin and B. Klartag, Moment measures, J. Funct. Anal., 268 (2015) 3834-3866.
- [22] N. Fang, D. Xing and D. Ye, Geometry of log-concave functions: the L_p Asplund sum and the L_p Minkowski problem, arXiv:2006.16959v1.
- [23] N. Fang, W. Xu, J. Zhou and B. Zhu, The sharp convex mixed Lorentz-Sobolev inequality, Adv. Appl. Math., (accepted).
- [24] N. Fang and J. Zhou, LYZ ellipsoid and Petty projection body for log-concave functions, Adv. Math., 340 (2018) 914-959.
- [25] M. Fradelizi, Sections of convex bodies through their centroid, Arch. Math., 69 (1997), no. 6, 515-522.
- [26] M. Fradelizi and M. Meyer, Some functional forms of Blaschke-Santaló inequality, Math. Z., 256 (2007) 379-395.
- [27] M. Fradelizi and M. Meyer, Some functional inverse Santaló inequality, Adv. Math., 218 (2008) 1430-1452.
- [28] R. Gardner, On the Busemann-Petty problem concerning central sections of centrally symmetric convex bodies, Bull. Amer. Math. Soc., 30 (1994) 222-226.
- [29] R. Gardner, Intersection bodies and the Busemann-Petty problem, Trans. Amer. Math. Soc., 342 (1994) 435-445.
- [30] R. Gardner, A positive answer to the Busemann-Petty problem in three dimensions, Ann. Math., 140 (1994) 435-447.
- [31] R. Gardner, D. Hug, W. Weil, S. Xing and D. Ye, General volumes in the Orlicz-Brunn-Minkowski theory and a related Minkowski Problem I, Calc. Var. Partial Differ. Equ., (2019) 58: 12. https://doi.org/10.1007/s00526-018-1449-0.
- [32] P. Guan and X. Ma, The Christoffel-Minkowski problem I: convexity of solutions of a Hessian equation, Invent. Math., 151 (2003) 553-577.
- [33] C. Haberl and F. Schuster, Asymmetric affine L_p Sobolev inequalities, J. Funct. Anal., 257 (2009) 641-658.
- [34] C. Haberl, F. Schuster and J. Xiao, An asymmetric affine Pólya-Szegö principle, Math. Ann., 352 (3) (2012) 517-542.
- [35] Y. Huang, E. Lutwak, D. Yang and G. Zhang, Geometric measures in the dual Brunn-Minkowski theory and their Minkowski problems, Acta Math., 216 (2016) 325-388.
- [36] Y. Huang, E. Lutwak, D. Yang and G. Zhang, The L_p Aleksandrov problem for L_p integral curvature, J. Differ. Geom., 110 (2018) 1-29.
- [37] Y. Huang and Y. Zhao, On the L_p dual Minkowski problem, Adv. Math., 332 (2018) 57-84.

- [38] D. Hug, E. Lutwak, D. Yang and G. Zhang, On the L_p Minkowski problem for polytopes, Discrete Comput. Geom., 33 (2005) 699-715.
- [39] B. Klartag, Logarithmically-concave moment measures I, in Geometric aspects of functional analysis, volume 2116 of lecture Notes in Math., Springer, Berlin, (2014).
- [40] H. Lewy, On differential geometry in the large. I. Minkowski problem, Trans. Amer. Math. Soc., 43 (1938) 258-270.
- [41] Q. Li, W. Sheng and X. Wang, Flow by Gauss curvature to the Aleksandrov and dual Minkowski problems, J. Eur. Math. Soc., (accepted for publication).
- [42] Y. Lin, Affine Orlicz Pólya-Szegö principle for log-concave functions, J. Funct. Anal., 273 (2017) 3295-3326.
- [43] M. Ludwig, Valuations on function spaces, Adv. Geom., 11 (2011) 745-756.
- [44] M. Ludwig, Valuations on Sobolev spaces, Amer. J. Math., 134 (2012) 827-842.
- [45] E. Lutwak, Dual mixed volumes, Pacific J. Math., 58 (1975) 531-538.
- [46] E. Lutwak, Intersection bodies and dual mixed volumes, Adv. Math., 71 (1988) 232-261.
- [47] E. Lutwak, Centroid bodies and dual mixed volumes, Proc. London Math. Soc., 60 (1990) 365-391.
- [48] E. Lutwak, The Brunn-Minkowski-Firey theory I: Mixed volumes and the Minkowski problem, J. Differ. Geom., 38 (1993) 131-150.
- [49] E. Lutwak, D. Yang and G. Zhang, L_p affine isoperimetric inequalities, J. Differ. Geom., 56 (2000) 111-132.
- [50] E. Lutwak, D. Yang and G. Zhang, Sharp affine L_p Sobolev inequalities, J. Differ. Geom., 62 (2002) 17-38.
- [51] E. Lutwak, D. Yang and G. Zhang, On the L_p Minkowski problem, Trans. Amer. Math. Soc., 356 (2004) 4359-4370.
- [52] E. Lutwak, D. Yang and G. Zhang, L_p dual curvature measures, Adv. Math., 329 (2018) 85-132.
- [53] R. McCann, A convexity principle for interacting gases, Adv. Math., 128 (1997) 153-179.
- [54] V. Milman and L. Rotem, Mixed integral and related inequalities, J. Funct. Anal., 264 (2013) 570-604.
- [55] H. Minkowski, Allgemeine Lehrsätze über die convexen polyeder, Gött. Nachr., 1897 (1897) 198-219.
- [56] L. Nirenberg, The Weyl and Minkowski problems in differential geometry in the large, Comm. Pure Appl. Math., 6 (1953) 337-394.
- [57] A. Pogorelov, A regular solution of the n-dimensional Minkowski problem, Dokl. Akad. Nauk SSSR, 199 (1971), 785-788.
- [58] A. Pogorelov, The Minkowski multidimensional problem, Winston, Washington, 1978.
- [59] L. Rotem, Support functions and mean width for α -concave functions, Adv. Math., 243 (2013) 168-186.
- [60] T. Rockafellar, Convex Analysis, Princeton University Press (Princeton, 1970).
- [61] L. Rotem, Surface area measures of log-concave functions, arXiv: 2006.16933.
- [62] R. Schneider, Convex Bodies: The Brunn-Minkowski Theory, Encyclopedia of Mathematics and its Applications, 44, Cambridge University Press, Cambridge, (2014).
- [63] A. Stancu, The discrete planar L_0 -Minkowski problem, Adv. Math., 167 (2002) 160-174.
- [64] A. Stancu, On the number of solutions to the discrete two-dimesional L_0 -Minkowski problem, Adv. Math., 180 (2003) 290-323.
- [65] H. Wang, N. Fang and J. Zhou, Continuity of the solution to the dual Minkowski problem for negative indices, Proc. Amer. Math. Soc., 147 (2019) 1299-1312.
- [66] S. Xing and D. Ye, The general dual Orlicz-Minkowski problem, Indiana Univ. Math. J., (accepted for publication).
- [67] G. Zhang, Centered bodies and dual mixed volumes, Trans. Amer. Math. Soc., 345 (1994) 777-801.
- [68] G. Zhang, The affine Sobolev inequality, J. Differ. Geom., 53 (1999) 183-202.

- [69] G. Zhang, A positive solution to the Busemann-Prtty problem in \mathbb{R}^4 , Ann. Math., 149 (1999) 535-543.
- [70] Y. Zhao, The dual Minkowski problem for negative indices, Calc. Var. Partial Differ. Equ., 56 (2017) 1124-x.
- [71] Y. Zhao, Existence of solutions to the even dual Minkowski problem, J. Differ. Geom., 110 (2018) 543-572.
- [72] G. Zhu, The logarithmic Minkowski problem for polytopes, Adv. Math., 262 (2014) 909-931.
- [73] G. Zhu, The centro-affine Minkowski problem for polytopes, J. Differ. Geom., 101 (2015) 159-174.
- [74] G. Zhu, The L_p Minkowski problem for polytopes for 0 , J. Funct. Anal., 269 (2015) 1070-1094.
- [75] G. Zhu, L_p Minkowski problem for polytopes for p < 0, Indiana Univ. Math. J., 66 (2017), no. 4, 1333-1350.
- [76] B. Zhu, S. Xing and D. Ye, The dual Orlicz-Minkowski problem, J. Geom. Anal., 28 (2018) 3829-3855.
- [77] B. Zhu, J. Zhou and W. Xu, Dual Orlicz-Brunn-Minkowski theory, Adv. Math., 264 (2014) 700-725.

SCHOOL OF MATHEMATICS, HUNAN UNIVERSITY, CHANGSHA, HUNAN 300071, CHINA. *Email address:* fangniufa@hnu.edu.cn

School of Mathematics and Statistics, Southwest University, Chongqing 400715, China.

College of Science, Wuhan University of Science and Technology, Wuhan, Hubei 430081, China.

Email address: zhoujz@swu.edu.cn