

# An inverse boundary value problem for the inhomogeneous porous medium equation

Cătălin I. Cârstea\*    Tuhin Ghosh†    Gen Nakamura‡

## Abstract

In this paper we establish uniqueness in the inverse boundary value problem for the two coefficients in the inhomogeneous porous medium equation  $\epsilon \partial_t u - \nabla \cdot (\gamma \nabla u^m) = 0$ , with  $m > 1$ , in dimension 3 or higher, which is a degenerate parabolic type quasilinear PDE. Our approach relies on using a Laplace transform to turn the original equation into a coupled family of nonlinear elliptic equations, indexed by the frequency parameter ( $1/h$  in our definition) of the transform. A careful analysis of the asymptotic expansion in powers of  $h$ , as  $h \rightarrow \infty$ , of the solutions to the transformed equation, with special boundary data, allows us to obtain sufficient information to deduce the uniqueness result.

## 1 Introduction

We would like to begin by recalling the classical inverse boundary value problem of Calderón, introduced in [5]. Suppose  $\Omega \subset \mathbb{R}^n$ , with  $n \geq 3$ , is a bounded, smooth domain and  $\gamma \in C^\infty(\Omega)$  is a bounded positive function with a positive lower bound. If we consider the (forward) boundary value problem

$$\begin{cases} \nabla \cdot (\gamma(x) \nabla u(x)) = 0, & x \in \Omega \\ u|_{\partial\Omega} = g \in C^\infty(\partial\Omega), \end{cases} \quad (1)$$

---

\*School of Mathematics, Sichuan University, Chengdu, Sichuan, 610064, P.R.China; email: catalin.carstea@gmail.com

†Department of Mathematics, Universität Bielefeld, 33615 Bielefeld, Germany; email: tghosh@math.uni-bielefeld.de

‡Department of Mathematics, Hokkaido University, Sapporo 060-0808, Japan; email: gnaka@math.sci.hokudai.ac.jp

then we can introduce the Dirichlet-to-Neumann map (DN map)

$$\Lambda_\gamma(g) = \gamma \partial_\nu u|_{\partial\Omega} = \gamma \nu \cdot \nabla u|_{\partial\Omega}, \quad (2)$$

where  $u$  is the solution of (1), and  $\nu$  is the outer pointing unit normal on  $\partial\Omega$ . Calderón's problem refers to the question of determining  $\gamma$  from a knowledge of  $\Lambda_\gamma$ . The important subproblem of uniqueness is the question of showing that the correspondence  $\gamma \rightarrow \Lambda_\gamma$  is, in fact, injective. This question has been answered in the affirmative in [34]. Note that, by the linearity of the DN map, if  $\Lambda_\gamma(g)$  is given for only non-negative Dirichlet data  $g$ , then the whole  $\Lambda_\gamma$  operator, defined on complex valued functions, can be recovered. We will use this observation later in the paper.

The subject of this paper is the following boundary value problem for the (inhomogeneous) porous medium equation (PME)

$$\begin{cases} \epsilon(x) \partial_t u(t, x) - \nabla \cdot (\gamma(x) \nabla u^m(t, x)) = 0, & (t, x) \in (0, \infty) \times \Omega, \\ u(0, x) = 0, \quad u|_{[0, \infty) \times \partial\Omega} = \phi(t, x), \quad u \geq 0. \end{cases} \quad (3)$$

Here  $m > 1$ , and the coefficients  $\epsilon$  and  $\gamma$  are bounded positive functions, which also have strictly positive lower bounds. For convenience we will assume  $\epsilon, \gamma \in C^\infty(\overline{\Omega})$ . Note that consistency requires that  $\phi(0, x) = 0$ ,  $\phi \geq 0$ .

The Dirichlet-to-Neumann map associated to (3) (informally) should be

$$\Lambda_{\epsilon, \gamma}^{PM}(\phi) = \gamma \partial_\nu u^m|_{[0, \infty) \times \partial\Omega}. \quad (4)$$

In order to rigorously define this map, we must first discuss the existence of solutions to (3). We will first introduce a notion of weak solutions on arbitrary, finite time intervals. Then we will see how these lead to a notion of solutions to (3) for infinite time.

As its name suggests, the PME can be seen as a model for the flow of a gas through a porous medium, with the function  $u(t, x)$  being the density of the gas at time  $t$  and position  $x$ . The parameters  $\epsilon$  and  $\gamma$  then depend on the particular gas considered, and also on properties of the medium. From the purely mathematical point of view, the PME is interesting as a particularly simple nonlinear modification of the classical heat equation, which nevertheless can exhibit very different properties from its linear counterpart. It can be classified as a degenerate parabolic type quasilinear equation. From the point of view of the study of inverse problems, the degenerate parabolic nature of the PME makes it a particularly interesting case to study, as known results of any kind for degenerate parabolic or elliptic PDE are still very few.

A good reference for the mathematical theory of the PME is the monograph [35]. Though the particular case of the inhomogeneous equation is not

treated there, most of what follows regarding the existence of solutions to the boundary value problem is closely based on the treatment of the homogeneous case in [35].

Before giving the specific definition of weak solutions we will be working with, we must introduce a few notations. For  $0 < T \leq \infty$ , let  $Q_T := (0, T) \times \Omega$  and  $S_T := (0, T) \times \partial\Omega$ . Let

$$C_t(Q_T) = \{\varphi \in C^\infty(\overline{Q_T}) : \text{supp } \varphi \cap [\{T\} \times \Omega] = \emptyset\} \quad (5)$$

and let  $H_t^1(Q_T)$  be the completion of this space in  $H^1(Q_T)$ . We will also denote by  $H_t^{\frac{1}{2}}(S_T)$  the subspace of  $H^{\frac{1}{2}}(S_T)$  that consists of traces of  $H_t^1(Q_T)$  functions. Similarly, we also introduce

$$C_\diamond(Q_T) = \{\varphi \in C^\infty(\overline{Q_T}) : \text{supp } \varphi \cap [S_T \cup \{T\} \times \Omega] = \emptyset\}, \quad (6)$$

and its completion  $H_\diamond^1(Q_T)$  in  $H^1(Q_T)$ .

Since we will be working with weak solutions, it is useful to recall a few things about boundary traces. We will denote by  $\tau_\Omega$  the boundary trace operator for the domain  $\Omega$ . It is bounded between  $H^1(\Omega)$  and  $H^{\frac{1}{2}}(\partial\Omega)$ .

We will denote by  $\tau_{Q_T}$  the boundary trace operator for  $Q_T$  to the side boundary  $S_T$ . It is not hard to check that  $\tau_{Q_T}$  is bounded from  $L^2((0, T); H^1(\Omega))$  to  $L^2((0, T); H^{\frac{1}{2}}(\partial\Omega))$ .

For our convenience later in the paper, we would like to consider the equation (3) with an added source term

$$\begin{cases} \epsilon(x)\partial_t u(t, x) - \nabla \cdot (\gamma(x)\nabla u^m(t, x)) = f(t, x), & (t, x) \in Q_\infty, \\ u(0, x) = 0, \quad u|_{[0, \infty) \times \partial\Omega} = \phi(t, x), \quad u \geq 0. \end{cases} \quad (7)$$

**Definition 1.** *Suppose  $0 < T < \infty$ . We say that  $u \in L^\infty(Q_T)$ ,  $u \geq 0$ , is a weak solution of (7) in  $Q_T$  if it satisfies the following conditions:*

1.  $\nabla(u^m)$  exists in the sense of distributions and  $\nabla(u^m) \in L^2(Q_T)$ ;
2. for any  $\varphi \in C_\diamond(Q_T)$  (or, equivalently, any  $\varphi \in H_\diamond^1(Q_T)$ )

$$\int_{Q_T} \gamma \nabla \varphi \cdot \nabla(u^m) \, dt \, dx - \int_{Q_T} \epsilon \partial_t \varphi u \, dt \, dx = \int_{Q_T} \varphi f \, dt \, dx; \quad (8)$$

3.  $\tau_{Q_T}(u^m) = \phi^m$ .

In section 2 we will prove the following result.

**Theorem 1.** For any  $0 < T < \infty$ , if  $\phi^m \in C(\overline{S_T}) \cap C^{0,1}(S_T)$ ,  $\phi \geq 0$ , and  $\phi(0, x) = 0$  for all  $x \in \partial\Omega$ , and if  $f \in L^\infty(Q_T)$ ,  $f \geq 0$ , then there exists a unique weak solution  $u$  of (7) in  $Q_T$ . This solution satisfies the energy estimate<sup>1</sup>

$$\|u^m\|_{L^2((0,T);H^1(\Omega))} \leq C \left( \|\phi^m\|_{C^{0,1}(\Omega)} + \|\phi\|_{C(\Omega)} + \|f\|_{L^\infty(Q_T)} \right). \quad (9)$$

If  $f = 0$ , it also satisfies the maximum principle

$$0 \leq \sup_{Q_T} u \leq \sup_{S_T} \phi. \quad (10)$$

If  $\phi_1 \leq \phi_2$ ,  $f_1 \leq f_2$  are as above and give rise to weak solutions  $u_1$  and  $u_2$ , then

$$u_1 \leq u_2. \quad (11)$$

Since the length of the time interval in the above theorem is arbitrary, it follows that we also have infinite time weak solutions, in the sense specified below.

**Corollary 1.** If  $\phi^m \in C(\overline{S_\infty}) \cap C_{loc}^{0,1}(S_\infty)$ ,  $\phi \geq 0$ , and  $\phi(0, x) = 0$  for all  $x \in \partial\Omega$ , then there exists  $u : Q_\infty \rightarrow [0, \infty)$  such that, for any  $0 < T < \infty$ ,  $u$  is a weak solution of (3) in  $Q_T$  and

$$0 \leq \sup_{Q_T} u \leq \sup_{S_T} \phi. \quad (12)$$

This allows us to rigorously define the Dirichlet-to-Neumann map for all  $\phi$  satisfying the conditions in Corollary 1. If  $u$  is the corresponding weak solution and  $\psi \in H_t^1(Q_T)$ , then we can define the Neumann data  $\gamma\partial_\nu u^m|_{S_T}$  by

$$\langle \gamma\partial_\nu u^m|_{S_T}, \psi|_{S_T} \rangle = \int_{Q_T} (\gamma\nabla\psi \cdot \nabla(u^m) - \epsilon\partial_t\psi u) dt dx. \quad (13)$$

We see then that  $\Lambda_{\epsilon,\gamma}^{PM}(\phi) \in \left(H_t^{\frac{1}{2}}(S_T)\right)'$ .

Suppose now that we have two sets of coefficients,  $\epsilon^{(i)}$ ,  $\gamma^{(i)}$  and  $\epsilon^{(ii)}$ ,  $\gamma^{(ii)}$  that are as above. The main result of our paper is the following.

**Theorem 2.** If  $\Lambda_{\epsilon^{(i)},\gamma^{(i)}}^{PM}(\phi) = \Lambda_{\epsilon^{(ii)},\gamma^{(ii)}}^{PM}(\phi)$ , for all  $\phi$  such that  $\phi^m \in C(\overline{S_\infty}) \cap C_{loc}^{0,1}(S_\infty)$ ,  $\phi \geq 0$ , and  $\phi(0, x) = 0$  for all  $x \in \partial\Omega$ , then  $\gamma^{(i)} = \gamma^{(ii)}$  and  $\epsilon^{(i)} = \epsilon^{(ii)}$ .

---

<sup>1</sup>The constant  $C$  here will depend on  $T$ .

We will prove this result in section 3. Our method involves first going from an equation for the function  $u$  to an equivalent problem for the function  $v = u^m$ . We then use a Laplace transform in time to introduce functions

$$V(h, x) = \int_0^{\infty} e^{-\frac{t}{h}} v(t, x) dt. \quad (14)$$

These satisfy a family of equations, indexed by  $h$ , of the form

$$\nabla \cdot (\gamma(x) \nabla V(h, x)) = \mathcal{N}(h, x), \quad (15)$$

with the right hand side also being constructed from the solution  $u$ . If we choose boundary data of the form  $\phi^m(t, x) = tg(x)$ , we are then able to derive the first two terms in the asymptotic expansion of  $V(h, x)$  in powers of  $h$ , as  $h \rightarrow \infty$ . The corresponding expansion of the Dirichlet-to-Neumann map leads to separate integral identities that the differences  $\gamma^{(i)} - \gamma^{(ii)}$  and  $\epsilon^{(i)} - \epsilon^{(ii)}$  must satisfy. These quickly lead to the claimed uniqueness result.

The result presented in this paper fits into a series of other works on inverse boundary value problems for nonlinear equations. In fact, our method resembles the “second linearization” approach that has been used in the past for (non-degenerate) elliptic and parabolic semilinear and quasilinear problems. We list here some of the important papers in this field. For semilinear equations, examples include [13], [16], [18], [19], [22], [23], [26], [25], [32]. For quasilinear equations, not in divergence form, see [17]. For quasilinear equations in divergence form see [8], [7], [9], [10], [11], [12], [15], [20], [28], [29], [30], [31], [33], (also [6] for quasilinear time-harmonic Maxwell systems).

Results for degenerate equations, such as the PME, are quite few. Examples would include the following works for the weighted p-Laplace equation,  $\nabla \cdot (a(x)|\nabla u|^{p-2}\nabla u) = 0$ , which is a degenerate elliptic quasilinear PDE: [4], [1], [2], [14], [3], [21]. Note that uniqueness in the inverse boundary value problem for the coefficient  $a$  in this equation, without additional assumptions such as monotonicity (as in [14]), has not yet been shown. To the best of our knowledge, this paper is the first such result for a degenerate parabolic or elliptic quasilinear PDE.

## 2 The forward problem

In this section we give a proof to Theorem 1, which gives the existence of finite time weak solutions. The method used is very similar to what can be

found in [35], the difference being that in that reference the coefficients are taken to be constant.

We begin by introducing a sequence  $u_k$ ,  $k = 1, 2, \dots$ , consisting of solutions to the boundary value problem

$$\begin{cases} \epsilon \partial_t u_k - \nabla \cdot (\gamma \nabla u_k^m) = f_k, \\ u_k(0, x) = \frac{1}{k}, \quad u_k|_{[0, T) \times \partial\Omega} = \phi + \frac{1}{k}, \end{cases} \quad (16)$$

where we choose functions  $f_k \in C^\infty(Q_T)$  such that

$$f(t, x) \leq f_{k+1}(t, x) \leq f_k(t, x) \leq f(t, x) + \frac{1}{k}. \quad (17)$$

Intuitively, we expect the solutions  $u_k$  (if they exist) to satisfy the maximum principle

$$\frac{1}{k} \leq u_k(t, x) \leq \frac{1}{k} + \sup_{S_T} \phi + t \sup_{Q_T} f_k. \quad (18)$$

With this in mind, we redefine the  $u_k$  to be the solutions of

$$\begin{cases} \epsilon \partial_t u_k - \nabla \cdot (a_k(x, u_k) \nabla u_k) = f_k, \\ u_k(0, x) = \frac{1}{k}, \quad u_k|_{[0, T) \times \partial\Omega} = \phi + \frac{1}{k}, \end{cases} \quad (19)$$

where the function  $a_k(x, \lambda)$  is chosen so as to make (19) a uniform quasilinear parabolic problem (i.e. it avoids the degeneracy at  $\lambda = 0, \infty$ ), but such that if

$$\frac{1}{k} \leq \lambda \leq \sup_{S_T} \phi + T \sup_{Q_T} f + \frac{1+T}{k}, \quad (20)$$

we have

$$a_k(x, \lambda) = m\gamma(x)\lambda^{m-1}. \quad (21)$$

In this case (see [24], or [27]), the problem (19) has a solution  $u_k \in C^{1,2}(Q_T) \cap C(\overline{Q_T})$  which furthermore satisfies the maximum principle (18). Since (18) is satisfied, it follows that  $u_k$  is in fact also a solution to (16).

Note that  $u_k$  is also a solution to  $\epsilon \partial_t u_k - \nabla \cdot (a_{k+1}(x, u_k, \nabla u_k)) = f_k$ . Therefore, by the comparison principle (see [27, Theorem 9.7]), we have that

$$0 \leq u_{k+1} \leq u_k, \quad \forall k = 1, 2, \dots \quad (22)$$

We can then define the pointwise limit

$$u(t, x) = \lim_{k \rightarrow \infty} u_k(t, x). \quad (23)$$

This function will turn out to be the weak solution to (3) that we are searching for.

Since clearly  $u \leq u_k$  for all  $k$ , by (18) we have that

$$0 \leq u(t, x) \leq \sup_{S_T} \phi + t \sup_{Q_T} f. \quad (24)$$

By the monotone convergence theorem we have that

$$\lim_{k \rightarrow \infty} \int_{Q_T} (u_k - u)^2 dt dx = 0, \quad (25)$$

i.e. that  $u_k \rightarrow u$  in  $L^2(Q_T)$ . Similarly, it also holds that  $u_k^m \rightarrow u^m$  in  $L^2(Q_T)$ .

We will denote by  $\tilde{\phi} : Q_T \rightarrow [0, \infty)$  a smooth extension of the Dirichlet data, such that we still have  $\tilde{\phi}(0, x) = 0$  for all  $x \in \Omega$ . We can choose this extension in such a way that

$$\|\tilde{\phi}\|_{W^{1,\infty}(Q_T)} \leq C \|\phi\|_{C^{0,1}(S_T)}, \quad (26)$$

with a constant  $C > 0$ .

Let

$$\eta_k = u_k^m - \left( \tilde{\phi} + \frac{1}{k} \right)^m, \quad (27)$$

which is zero on  $S_T$ . We can multiply (16) by  $\eta_k$  and integrate over  $Q_T$ . Note that

$$\begin{aligned} & - \int_{Q_T} \eta_k \nabla(\gamma \nabla u_k^m) dt dx \\ &= \int_{Q_T} \gamma |\nabla u_k^m|^2 dt dx - \int_{Q_T} \gamma \nabla \left( \tilde{\phi} + \frac{1}{k} \right)^m \cdot \nabla u_k^m dt dx, \end{aligned} \quad (28)$$

and that

$$\begin{aligned} \int_{Q_T} \epsilon \partial_t u_k \eta_k dt dx &= \int_{Q_T} \epsilon \partial_t u_k u_k^m dt dx - \int_{Q_T} \epsilon \partial_t u_k \left( \tilde{\phi} + \frac{1}{k} \right)^m dt dx \\ &= \int_{\Omega} \frac{\epsilon}{m+1} u_k^{m+1}(T) dx - \int_{\Omega} \epsilon u_k(T) \left( \tilde{\phi}(T) + \frac{1}{k} \right)^m dx \\ &\quad + \int_{Q_T} \epsilon u_k \partial_t \left( \tilde{\phi} + \frac{1}{k} \right)^m dt dx. \end{aligned} \quad (29)$$

Putting these two together we have

$$\begin{aligned}
& \int_{Q_T} \gamma |\nabla u_k^m|^2 dt dx + \int_{\Omega} \frac{\epsilon}{m+1} u_k^{m+1}(T) dx \\
&= \int_{Q_T} \gamma \nabla \left( \tilde{\phi} + \frac{1}{k} \right)^m \cdot \nabla u_k^m dt dx + \int_{\Omega} \epsilon u_k(T) \left( \tilde{\phi}(T) + \frac{1}{k} \right)^m dx \\
&\quad - \int_{Q_T} \epsilon u_k \partial_t \left( \tilde{\phi} + \frac{1}{k} \right)^m dt dx + \int_{Q_T} f_k \left[ u_k^m - \left( \tilde{\phi} + \frac{1}{k} \right)^m \right]. \quad (30)
\end{aligned}$$

By straightforward estimates we can conclude that there exists a constant  $C > 0$ , depending on  $m$  and the upper and lower bounds of  $\epsilon$  and  $\gamma$ , such that

$$\begin{aligned}
& \int_{Q_T} |\nabla u_k^m|^2 dt dx + \int_{\Omega} u_k^{m+1}(T) dx \leq C \left[ \|u_k\|_{L^2(Q_T)}^2 + \|u_k^m\|_{L^2(Q_T)}^2 + \|f_k\|_{L^2(Q_T)}^2 \right. \\
&\quad \left. + \left\| \left( \tilde{\phi} + \frac{1}{k} \right)^m \right\|_{H^1(Q_T)}^2 + \left\| \left( \tilde{\phi} + \frac{1}{k} \right)^m \right\|_{L^{\frac{m+1}{m}}(\Omega)} \right]. \quad (31)
\end{aligned}$$

There is then a constant  $C' > 0$ , independent of  $k$ , such that

$$\begin{aligned}
& \int_{Q_T} |\nabla u_k^m|^2 dt dx < C' \left( \left\| \left( \phi + \frac{1}{k} \right)^m \right\|_{C^{0,1}(S_T)}^2 + \left\| \left( \phi + \frac{1}{k} \right)^m \right\|_{C(S_T)} \right. \\
&\quad \left. + \left\| \phi + \frac{1}{k} \right\|_{C(S_T)}^2 + \left\| f + \frac{1}{k} \right\|_{L^\infty(Q_T)}^2 \right). \quad (32)
\end{aligned}$$

It follows that (a subsequence of)  $\nabla u_k^m$  converges weakly in  $L^2(Q_T)$  to a limit  $U$ . It is easy to see that  $U = \nabla u^m$  in the sense of distributions.

Since we have that  $\tau_{Q_T}(u_k^m)(t) \rightarrow \phi^m$  in  $L^2((0, T); H^{\frac{1}{2}}(\partial\Omega))$  by construction, and also that  $\tau_{Q_T}(u_k^m) \rightharpoonup \tau_{Q_T}(u^m)$ , it follows that

$$\tau_{Q_T}(u^m) = \phi^m. \quad (33)$$

This shows that  $u$  is a weak solution of (3) in  $Q_T$ .

In order to prove uniqueness of weak solutions, we start by assuming  $u_1$  and  $u_2$  are both weak solutions of (3) in  $Q_T$ . We have that for all test



functions  $\varphi$

$$\int_{Q_T} \gamma(\nabla u_1^m - \nabla u_2^m) \nabla \varphi \, dt \, dx - \int_{Q_T} \epsilon(u_1 - u_2) \partial_t \varphi \, dt \, dx = 0. \quad (34)$$

Let

$$\varphi(t, x) = \int_t^T (u_1^m(s, x) - u_2^m(s, x)) \, ds. \quad (35)$$

It is not hard to check that  $\varphi \in H_\diamond^1(Q_T)$ , so we can use it in the above identity. We obtain

$$\frac{1}{2} \int_{\Omega} \gamma \left( \int_0^T (\nabla u_1^m - \nabla u_2^m) \, dt \right)^2 \, dx + \int_{Q_T} \epsilon(u_1 - u_2) (u_1^m - u_2^m) \, dt \, dx = 0. \quad (36)$$

The first term is clearly positive and, by the monotonicity of the  $z \rightarrow z^m$  function in  $\mathbb{R}_+$ , so is the second. It follows that  $u_1 = u_2$ .

Regarding the energy inequality claimed in the statement of the Theorem, it is enough to note that

$$\begin{aligned} \|\nabla u^m\|_{L^2(Q_T)} &\leq \liminf_{k \rightarrow \infty} \|\nabla u_k^m\|_{L^2(Q_T)} \\ &\leq C \left( \|\phi^m\|_{C^{0,1}(S_T)} + \|\phi\|_{C(S_T)} + \|f\|_{L^\infty(Q_T)} \right). \end{aligned} \quad (37)$$

Finally, suppose we have boundary data and sources  $\phi_1 \leq \phi_2$ ,  $f_1 \leq f_2$ , with corresponding solutions  $u_1$  and  $u_2$ . It is enough to observe that the approximating sequences defined above must satisfy

$$u_{1,k} \leq u_{2,k}, \quad (38)$$

and this property is preserved in the limit. This concludes the proof.

### 3 The inverse problem

In this section we give a proof of Theorem 2. We do this by transforming the equation so that the principal term becomes linear. Then we use a Laplace transform in time to transform the equation into a (coupled) family of nonlinear elliptic equations. Below, when we write  $\epsilon$  and  $\gamma$ , we mean any one of the pairs  $\epsilon^{(i)}$ ,  $\gamma^{(i)}$  or  $\epsilon^{(ii)}$ ,  $\gamma^{(ii)}$ .

### 3.1 The first reformulation of the problem

We first perform the change of function  $v = u^m$ . The new function satisfies the equation

$$\begin{cases} \epsilon(x)\partial_t v(t, x)^{\frac{1}{m}} - \nabla \cdot (\gamma(x)\nabla v(t, x)) = 0, \\ v(0, x) = 0, \quad v|_{[0, \infty) \times \partial\Omega} = f(t, x), \quad v \geq 0, \end{cases} \quad (39)$$

where  $f = \phi^m$ . It is clear that the notions of weak solutions we have introduced for equation (3) give corresponding notions of weak solutions to (39), whose existence is then guaranteed by Theorem 1. The associated Dirichlet-to-Neumann map is

$$\Lambda_{\epsilon, \gamma}^v(f) = \gamma \partial_\nu v|_{[0, \infty) \times \partial\Omega}, \quad (40)$$

and it contains the same information as  $\Lambda_{\epsilon, \gamma}^{PM}$ , when considering  $f \in C(\overline{S_\infty}) \cap C^\infty(S_\infty)$ ,  $f \geq 0$ , and such that  $f(0, x) = 0$  for all  $x \in \Omega$ .

### 3.2 Reformulation of the problem using the Laplace transform

The next reformulation of the problem uses the Laplace transform to turn our equation into an elliptic one. From now on we will only consider boundary data of a particular form, namely  $f(t, x) = tg(x)$ , with  $g \in C(\partial\Omega)$ . For  $h > 0$  define

$$V(h, x) = \int_0^\infty e^{-\frac{t}{h}} v(t, x) dt. \quad (41)$$

Note that since (in the sense of distributions) we have

$$\nabla V(h, x) = \int_0^\infty e^{-\frac{t}{h}} \nabla v(t, x) dt, \quad (42)$$

it then follows that

$$\begin{aligned} \|\nabla V(h, \cdot)\|_{L^2(\Omega)} &\leq \int_0^\infty e^{-\frac{t}{h}} \|\nabla v(t, \cdot)\|_{L^2(\Omega)} dt \\ &\leq \left( \int_0^\infty e^{-\frac{t}{h}} dt \right)^{\frac{1}{2}} \left( \int_{Q_\infty} e^{-\frac{t}{h}} |\nabla v(t, x)|^2 dt dx \right)^{\frac{1}{2}} < \infty, \quad (43) \end{aligned}$$

where the last inequality holds since, by Theorem 1,

$$\int_{Q_T} |\nabla v(t, x)|^2 dt dx \quad (44)$$

has an at most polynomial growth as  $T \rightarrow \infty$ , so the integral over  $Q_\infty$  converges. Therefore we have that  $V(h, \cdot) \in H^1(\Omega)$ .

If  $\varphi \in C_0^\infty(\Omega)$ , and  $\zeta \in C_0^\infty(\mathbb{R})$  is such that  $\zeta(t) = 1$  when  $|t| \leq 1$  and  $\zeta(t) = 0$  when  $|t| \geq 2$ , then

$$\begin{aligned} \langle \nabla(\gamma \nabla V(h, \cdot)), \varphi \rangle &= - \int_{\Omega} \gamma(x) \nabla V(h, x) \cdot \nabla \varphi(x) dx \\ &= - \lim_{T \rightarrow \infty} \int_{Q_\infty} \zeta\left(\frac{t}{T}\right) e^{-\frac{t}{h}} \gamma(x) \nabla v(t, x) \cdot \nabla \varphi(x) dt dx \\ &= - \lim_{T \rightarrow \infty} \int_{Q_\infty} \epsilon(x) \left[ \frac{1}{T} \zeta'\left(\frac{t}{T}\right) e^{-\frac{t}{h}} - \frac{1}{h} \zeta\left(\frac{t}{T}\right) e^{-\frac{t}{h}} \right] v^{\frac{1}{m}}(t, x) \varphi(x) dt dx \\ &= \int_{\Omega} \varphi(x) \left[ \frac{1}{h} \epsilon(x) \int_0^\infty e^{-\frac{t}{h}} v^{\frac{1}{m}}(t, x) dt \right] dx. \end{aligned} \quad (45)$$

It follows that

$$\nabla \cdot (\gamma(x) \nabla V(h, x)) = \mathcal{N}(h, x), \quad (46)$$

where

$$\mathcal{N}(h, x) = \frac{1}{h} \epsilon(x) \int_0^\infty e^{-\frac{t}{h}} v^{\frac{1}{m}}(t, x) dt. \quad (47)$$

The value  $V(h, x)$  takes on the boundary (in the sense of traces) is

$$\int_0^\infty e^{-\frac{t}{h}} t g(x) dt = h^2 g(x). \quad (48)$$

We then conclude that  $V$  satisfies the family of boundary value problems

$$\begin{cases} \nabla \cdot (\gamma(x) \nabla V(h, x)) = \mathcal{N}(h, x), \\ V|_{\partial\Omega} = h^2 g, \quad V \geq 0. \end{cases} \quad (49)$$

The associated DN map is essentially the Laplace transform of  $\Lambda_{\epsilon, \gamma}^v$ , namely

$$\Lambda_{\epsilon, \gamma}^h(g) = h^{-2} \gamma \partial_\nu V(h, \cdot)|_{\partial\Omega}. \quad (50)$$

We have added the  $h^{-2}$  factor for the sake of convenience in the computations that follow below.

Let  $m'$  be such that  $m'^{-1} + m^{-1} = 1$ . Then by Hölder's inequality

$$\begin{aligned} 0 \leq \mathcal{N}(h, x) &= \frac{1}{h} \epsilon(x) \int_0^\infty e^{-\frac{t}{m'h}} \left( e^{-\frac{t}{h}} v(t, x) \right)^{\frac{1}{m}} dt \\ &\leq h^{-\frac{1}{m}} \epsilon(x) (V(h, x))^{\frac{1}{m}}, \end{aligned} \quad (51)$$

so

$$\begin{aligned} \|\mathcal{N}(h, \cdot)\|_{L^\infty(\Omega)} &\leq h^{-\frac{1}{m}} \|\epsilon\|_{L^\infty(\Omega)} \|V(h, \cdot)\|_{L^\infty(\Omega)}^{\frac{1}{m}} \\ &\leq Ch^{-\frac{1}{m}} \|V(h, \cdot)\|_{W^{1,p}(\Omega)}^{\frac{1}{m}}, \end{aligned} \quad (52)$$

for a  $p > n$ .

By elliptic regularity estimates for (49) we have

$$\begin{aligned} \|V(h, \cdot)\|_{W^{2,p}(\Omega)} &\leq C \left( h^2 \|g\|_{W^{2-\frac{1}{p},p}(\partial\Omega)} + \|\mathcal{N}(h, \cdot)\|_{L^p(\Omega)} \right) \\ &\leq C \left( h^2 \|g\|_{W^{2-\frac{1}{p},p}(\partial\Omega)} + h^{-\frac{1}{m}} \|V(h, \cdot)\|_{W^{1,p}(\Omega)}^{\frac{1}{m}} \right). \end{aligned} \quad (53)$$

Then

$$\begin{aligned} &\max (\|V(h, \cdot)\|_{W^{2,p}(\Omega)}, 1) \\ &\leq C \left( h^2 \|g\|_{W^{2-\frac{1}{p},p}(\partial\Omega)} + 1 + h^{-\frac{1}{m}} [\max (\|V(h, \cdot)\|_{W^{1,p}(\Omega)}, 1)]^{\frac{1}{m}} \right) \\ &\leq C \left( h^2 \|g\|_{W^{2-\frac{1}{p},p}(\partial\Omega)} + 1 + h^{-\frac{1}{m}} \max (\|V(h, \cdot)\|_{W^{1,p}(\Omega)}, 1) \right). \end{aligned} \quad (54)$$

There clearly exists an  $h_0 > 0$  such that if  $h > h_0$  we have

$$\|V(h, \cdot)\|_{W^{2,p}(\Omega)} \leq h^2 C \left( \|g\|_{W^{2-\frac{1}{p},p}(\partial\Omega)} + h^{-2} \right). \quad (55)$$

### 3.3 First order asymptotic expansions and the determination of $\gamma$

We make the following Ansatz

$$V(h, x) = h^2 V_0(x) + R_1(h, x), \quad (56)$$

where  $V_0 \in W^{2,p}(\Omega)$  solves the equation

$$\begin{cases} \nabla \cdot (\gamma(x) \nabla V_0(x)) = 0, \\ V_0|_{\partial\Omega} = g. \end{cases} \quad (57)$$

It follows that  $R_1$  must satisfy

$$\begin{cases} \nabla \cdot (\gamma(x) \nabla R_1(h, x)) = \mathcal{N}(h, x), \\ R_1|_{\partial\Omega} = 0. \end{cases} \quad (58)$$

We observe that  $R_1$  is a subsolution for the elliptic operator  $\nabla \cdot \gamma \nabla$ , so we must have that  $R_1 \leq 0$ .

By elliptic estimates we have that  $R_1$  is bounded uniformly in  $h > h_0$

$$\begin{aligned} \|R_1(h, \cdot)\|_{W^{2,p}(\Omega)} &\leq C \|\mathcal{N}(h, \cdot)\|_{L^p(\Omega)} \\ &\leq Ch^{-\frac{1}{m}} \|V(h, \cdot)\|_{W^{1,p}(\Omega)}^{\frac{1}{m}} \leq Ch^{\frac{1}{m}} \|g\|_{W^{2-\frac{1}{p},p}(\partial\Omega)}^{\frac{1}{m}}. \end{aligned} \quad (59)$$

The asymptotic behavior of  $\Lambda_{\epsilon, \gamma}^h(g)$  as  $h \rightarrow \infty$  is then

$$\Lambda_{\epsilon, \gamma}^h(g) = \gamma \partial_\nu V_0(\cdot)|_{\partial\Omega} + \mathcal{O}(h^{\frac{1}{m}-2}). \quad (60)$$

Now, since we must have  $\Lambda_{\epsilon^{(i)}, \gamma^{(ii)}}^h(g) = \Lambda_{\epsilon^{(i)}, \gamma^{(ii)}}^h(g)$  for all  $g \in C(\overline{\Omega}) \cap C^\infty(\Omega)$ , with  $g \geq 0$ , it follows that corresponding orders in  $h$  of their asymptotic expansions must also be equal. From the  $h^0$  order term we then get that the DN maps for the original Calderón problem, with conductivities  $\gamma^{(i)}$  and  $\gamma^{(ii)}$ , are equal. That is, we have  $\Lambda_{\gamma^{(i)}} = \Lambda_{\gamma^{(ii)}}$ . It follows then from [34] that  $\gamma^{(i)} = \gamma^{(ii)}$ .

### 3.4 Asymptotic expansions up to the second order and the determination of $\epsilon$

In order to also prove the equality of the  $\epsilon$  coefficients, we need to look deeper into the asymptotic expansion of the DN map. Noting that

$$h^2 V_0(x) = \int_0^\infty e^{-\frac{t}{h}} t V_0(x) dt, \quad (61)$$

and also that  $V_0 \geq 0$ , let

$$\mathcal{N}_0(h, x) = \frac{1}{h} \epsilon(x) \int_0^\infty e^{-\frac{t}{h}} t^{\frac{1}{m}} V_0^{\frac{1}{m}}(x) dt = \Gamma\left(1 + \frac{1}{m}\right) \epsilon(x) h^{\frac{1}{m}} V_0^{\frac{1}{m}}(x), \quad (62)$$

where

$$\Gamma(z) = \int_0^{\infty} e^{-s} s^{z-1} ds. \quad (63)$$

We refine the Ansatz (56) to

$$V(h, x) = h^2 V_0(x) + h^{\frac{1}{m}} V_1(x) + R_2(h, x), \quad (64)$$

where  $V_1 \in W^{2,p}(\Omega)$  is the solution to

$$\begin{cases} \nabla \cdot (\gamma(x) \nabla V_1(x)) = h^{-\frac{1}{m}} \mathcal{N}_0(h, x), \\ V_1|_{\partial\Omega} = 0. \end{cases} \quad (65)$$

$V_1$  is clearly independent of  $h$ . Also, as  $V_1$  is a subsolution for the elliptic operator  $\nabla \cdot \gamma \nabla$ , it follows that  $V_1 \leq 0$ .

Since

$$h^{\frac{1}{m}} V_1(x) = \frac{1}{\Gamma\left(\frac{1}{m}\right)} \int_0^{\infty} e^{-\frac{t}{h}} t^{\frac{1}{m}-1} V_1(x) dt, \quad (66)$$

we would like to use  $V_1$  to also define

$$\mathcal{N}_1(h, x) = \frac{1}{h} \epsilon(x) \int_0^{\infty} e^{-\frac{t}{h}} \left( t V_0(x) + \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-1} V_1(x) \right)^{\frac{1}{m}} dt. \quad (67)$$

Note though that the quantity under the integral is not defined for small  $t$ . There is however a  $\sigma > 0$  such that

$$\mathcal{N}_1(h, x) = \frac{1}{h} \epsilon(x) \int_{\sigma}^{\infty} e^{-\frac{t}{h}} \left( t V_0(x) + \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-1} V_1(x) \right)^{\frac{1}{m}} dt \quad (68)$$

is well defined. We will use (68) as the definition of  $\mathcal{N}_1$ .

Returning to (64), we see that  $R_2$  must satisfy the equation

$$\begin{cases} \nabla \cdot (\gamma(x) \nabla R_2(h, x)) = (\mathcal{N}(h, x) - \mathcal{N}_0(h, x)), \\ R_2|_{\partial\Omega} = 0. \end{cases} \quad (69)$$

Note that

$$\begin{aligned} & \mathcal{N}(h, x) - \mathcal{N}_0(h, x) \\ & \leq \epsilon(x) \left[ h^{-\frac{1}{m}} V^{\frac{1}{m}}(h, x) - h^{\frac{1}{m}} \Gamma\left(1 + \frac{1}{m}\right) V_0^{\frac{1}{m}}(x) \right] \\ & \leq \epsilon(x) h^{\frac{1}{m}} \left[ (V_0(x) + h^{-2} R_1(h, x))^{\frac{1}{m}} - V_0^{\frac{1}{m}}(x) \right] \leq 0, \end{aligned} \quad (70)$$

where we have used the fact that  $0 < \Gamma\left(1 + \frac{1}{m}\right) < 1$  and our previous observation that  $R_1 \leq 0$ . It follows that  $R_2$  is a supersolution for the elliptic operator  $\nabla \cdot \gamma \nabla$ , so we have that  $R_2 \geq 0$ .

Let  $\chi : \mathbb{R} \rightarrow [0, 1]$  be a smooth, non-decreasing, cutoff function such that  $\chi(t) = 1$  if  $t \geq \sigma$  and  $\chi(t) = 0$  if  $t \leq \frac{\sigma}{2}$ . We may choose  $\sigma$  so that

$$w(t, x) = \chi^m(t) \left[ tV_0(x) + \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-1} V_1(x) \right] \quad (71)$$

is always non-negative. Then

$$\begin{aligned} \epsilon \partial_t w^{\frac{1}{m}} - \nabla \cdot (\gamma \nabla w) &= \chi' \left[ tV_0(x) + \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-1} V_1(x) \right]^{\frac{1}{m}} \\ &+ \epsilon \frac{1}{m} \chi^m w^{\frac{1}{m}-1} \left[ V_0 - \left(1 - \frac{1}{m}\right) \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-2} V_1 \right] \\ &- \epsilon \chi^m \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-1} \Gamma\left(1 + \frac{1}{m}\right) V_0^{\frac{1}{m}} \\ &\geq \frac{\epsilon \chi^m V_0}{m} \left( w^{\frac{1}{m}-1} - t^{\frac{1}{m}-1} V_0^{\frac{1}{m}-1} \right) \geq 0. \end{aligned} \quad (72)$$

Note also that  $\epsilon \partial_t w^{\frac{1}{m}} - \nabla \cdot (\gamma \nabla w) \in L^\infty(Q_T)$  for any  $0 < T < \infty$ . By the last part of Theorem 1, it follows that  $w \leq v$ . In particular, we have

$$tV_0(x) + \Gamma\left(\frac{1}{m}\right)^{-1} t^{\frac{1}{m}-1} V_1(x) \leq v(t, x), \quad \forall t \geq \sigma. \quad (73)$$

Therefore

$$\mathcal{N}_1(h, x) \leq \mathcal{N}(h, x), \quad (74)$$

so

$$\mathcal{N}_1(h, x) - \mathcal{N}_0(h, x) \leq \mathcal{N}(h, x) - \mathcal{N}_0(h, x) \leq 0. \quad (75)$$

We can estimate

$$\begin{aligned} \mathcal{N}_0(h, x) - \mathcal{N}_1(h, x) &= \frac{1}{h} \epsilon(x) \int_{\sigma}^{\infty} e^{-\frac{t}{h}} \\ &\times \left[ t^{\frac{1}{m}} V_0^{\frac{1}{m}}(x) - \left( t V_0(x) + \Gamma \left( \frac{1}{m} \right)^{-1} t^{\frac{1}{m}-1} V_1(x) \right)^{\frac{1}{m}} \right] dt \\ &+ \frac{1}{h} \epsilon(x) \int_0^{\sigma} e^{-\frac{t}{h}} t^{\frac{1}{m}} V_0^{\frac{1}{m}}(x) dt. \end{aligned} \quad (76)$$

In the sense of  $L^\infty(\Omega)$  norms we have that

$$\begin{aligned} \frac{1}{h} \epsilon(x) \int_{\sigma}^{\infty} e^{-\frac{t}{h}} \left[ t^{\frac{1}{m}} V_0^{\frac{1}{m}}(x) - \left( t V_0(x) + \Gamma \left( \frac{1}{m} \right)^{-1} t^{\frac{1}{m}-1} V_1(x) \right)^{\frac{1}{m}} \right] dt \\ = \mathcal{O}(h^{\frac{2}{m}-2}). \end{aligned} \quad (77)$$

Furthermore

$$\frac{1}{h} \int_0^{\sigma} e^{-\frac{t}{h}} t^{\frac{1}{m}} dt = \mathcal{O}(h^{-1}). \quad (78)$$

Let  $M = \min(1, 2 - \frac{2}{m}) > 0$ . It follows that

$$\|R_2(h, \cdot)\|_{W^{2,p}(\Omega)} = \mathcal{O}(h^{-M}). \quad (79)$$

This gives the following asymptotic expansion for  $\Lambda_{\epsilon, \gamma}^h$  as  $h \rightarrow \infty$

$$\Lambda_{\epsilon, \gamma}^h(g) = \gamma \partial_\nu V_0(\cdot)|_{\partial\Omega} + h^{\frac{1}{m}-2} \gamma \partial_\nu V_1(\cdot)|_{\partial\Omega} + \mathcal{O}(h^{-M-2}). \quad (80)$$

It is easy to see that for any  $m > 1$  we have

$$\frac{1}{m} - 2 > -M - 2, \quad (81)$$

so equation (80) identifies the first two terms in the the asymptotic expansion of the DN map without any ambiguity.

Let  $W$  be any smooth solution to  $\nabla \cdot (\gamma \nabla W) = 0$ . We have that

$$\begin{aligned} \langle \gamma \partial_\nu V_1|_{\partial\Omega}, W|_{\partial\Omega} \rangle &= \int_{\Omega} \gamma \nabla V_1 \cdot \nabla W \, dx + \Gamma \left( 1 + \frac{1}{m} \right) \int_{\Omega} \epsilon V_0^{\frac{1}{m}} W \, dx \\ &= \Gamma \left( 1 + \frac{1}{m} \right) \int_{\Omega} \epsilon V_0^{\frac{1}{m}} W \, dx. \end{aligned} \quad (82)$$



If  $\Lambda_{\epsilon,\gamma}^h$  is known, then so are all the integral quantities of the form that appears on the right hand side of the above equation.

We can take  $V_0$  to be of the form

$$V_0(x) = 1 + sH(x), \quad \nabla \cdot (\gamma \nabla H) = 0, \quad H > 0. \quad (83)$$

Then

$$m \frac{d}{ds} \Big|_{s=0} \int_{\Omega} \epsilon (1 + sH(x))^{\frac{1}{m}} W \, dx = \int_{\Omega} \epsilon HW \, dx \quad (84)$$

is determined by  $\Lambda_{\epsilon,\gamma}^h$ . From here it is easy to see that we must have

$$\int_{\Omega} (\epsilon^{(i)} - \epsilon^{(ii)}) UW \, dx = 0 \quad (85)$$

for all  $U, W$  such that  $\nabla \cdot (\gamma \nabla U) = \nabla \cdot (\gamma \nabla W) = 0$ . As shown in [34], the span of such products  $UV$  is dense in  $L^2(\Omega)$ , so this implies that  $\epsilon^{(i)} = \epsilon^{(ii)}$ .

**Acknowledgments:** C.C. was supported by NSF of China under grant 11931011. The research of T.G. is supported by the Collaborative Research Center, membership no. 1283, Universität Bielefeld. Work on this project began when C.C. and T.G. visited the National Center for Theoretical Sciences (NCTS), National Taiwan University. The authors acknowledge the support provided by NCTS during that visit.

## References

- [1] T. Brander. Calderón problem for the p-Laplacian: First order derivative of conductivity on the boundary. *Proceedings of the American Mathematical Society*, 144(1):177–189, 2016.
- [2] T. Brander, B. Harrach, M. Kar, and M. Salo. Monotonicity and enclosure methods for the p-Laplace equation. *SIAM Journal on Applied Mathematics*, 78(2):742–758, 2018.
- [3] T. Brander, J. Ilmavirta, and M. Kar. Superconductive and insulating inclusions for linear and non-linear conductivity equations. *Inverse Problems & Imaging*, 12(1):91–123, 2018.
- [4] T. Brander, M. Kar, and M. Salo. Enclosure method for the p-Laplace equation. *Inverse Problems*, 31(4):045001, 2015.

- [5] A. P. Calderón. On an inverse boundary value problem. In *Seminar on Numerical Analysis and its Applications to Continuum Physics (Rio de Janeiro, 1980)*, pages 65–73. Soc. Brasil. Mat., Rio de Janeiro, 1980.
- [6] C. I. Cârstea. On an inverse boundary value problem for a nonlinear time harmonic Maxwell system. *Journal of Inverse and Ill-posed Problems*, 1(ahead-of-print), 2020.
- [7] C. I. Cârstea and A. Feizmohammadi. A density property for tensor products of gradients of harmonic functions and applications. *arXiv preprint arXiv:2009.11217*, 2020.
- [8] C. I. Cârstea and A. Feizmohammadi. An inverse boundary value problem for certain anisotropic quasilinear elliptic equations. *Journal of Differential Equations*, 284:318–349, 2021.
- [9] C. I. Cârstea, A. Feizmohammadi, Y. Kian, K. Krupchyk, and G. Uhlmann. The Calderón inverse problem for isotropic quasilinear conductivities. *arXiv preprint arXiv:2103.05917*, 2021.
- [10] C. I. Cârstea and M. Kar. Recovery of coefficients for a weighted p-Laplacian perturbed by a linear second order term. *arXiv preprint arXiv:2001.01436*, 2020.
- [11] C. I. Cârstea, G. Nakamura, and M. Vashisth. Reconstruction for the coefficients of a quasilinear elliptic partial differential equation. *Applied Mathematics Letters*, 2019.
- [12] H. Egger, J.-F. Pietschmann, and M. Schlottbom. Simultaneous identification of diffusion and absorption coefficients in a quasilinear elliptic problem. *Inverse Problems*, 30(3):035009, 2014.
- [13] A. Feizmohammadi and L. Oksanen. An inverse problem for a semi-linear elliptic equation in Riemannian geometries. *Journal of Differential Equations*, 269(6):4683–4719, 2020.
- [14] C.-Y. Guo, M. Kar, and M. Salo. Inverse problems for p-Laplace type equations under monotonicity assumptions. *Rend. Istit. Mat. Univ. Trieste*, 48:79–99, 2016.
- [15] D. Hervas and Z. Sun. An inverse boundary value problem for quasilinear elliptic equations. *Communications in Partial Differential Equations*, 27(11-12):2449–2490, 2002.

- [16] V. Isakov. On uniqueness in inverse problems for semilinear parabolic equations. *Archive for Rational Mechanics and Analysis*, 124(1):1–12, 1993.
- [17] V. Isakov. Uniqueness of recovery of some quasilinear partial differential equations. *Communications in Partial Differential Equations*, 26(11-12):1947–1973, 2001.
- [18] V. Isakov and A. I. Nachman. Global uniqueness for a two-dimensional semilinear elliptic inverse problem. *Transactions of the American Mathematical Society*, 347(9):3375–3390, 1995.
- [19] V. Isakov and J. Sylvester. Global uniqueness for a semilinear elliptic inverse problem. *Communications on Pure and Applied Mathematics*, 47(10):1403–1410, 1994.
- [20] H. Kang and G. Nakamura. Identification of nonlinearity in a conductivity equation via the Dirichlet-to-Neumann map. *Inverse Problems*, 18(4):1079, 2002.
- [21] M. Kar and J.-N. Wang. Size estimates for the weighted p-Laplace equation with one measurement. *Discrete & Continuous Dynamical Systems-B*, 22(11):0, 2017.
- [22] K. Krupchyk and G. Uhlmann. Partial data inverse problems for semilinear elliptic equations with gradient nonlinearities. *arXiv preprint arXiv:1909.08122*, 2019.
- [23] Katya Krupchyk and Gunther Uhlmann. A remark on partial data inverse problems for semilinear elliptic equations. *Proceedings of the American Mathematical Society*, 148(2):681–685, 2020.
- [24] O. A. Ladyzhenskaia, V. A. Solonnikov, and N. N. Ural'tseva. *Linear and quasi-linear equations of parabolic type*, volume 23. American Mathematical Soc., 1988.
- [25] M. Lassas, T. Liimatainen, Y.-H. Lin, and M. Salo. Partial data inverse problems and simultaneous recovery of boundary and coefficients for semilinear elliptic equations. *arXiv preprint arXiv:1905.02764*, 1905.
- [26] M. Lassas, T. Liimatainen, Y.-H. Lin, and M. Salo. Inverse problems for elliptic equations with power type nonlinearities. *arXiv preprint arXiv:1903.12562*, 2019.

- [27] G. M. Lieberman. *Second Order Parabolic Differential Equations*. World Scientific, 1996.
- [28] C. Munoz and G. Uhlmann. The Calderón problem for quasilinear elliptic equations. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 2020.
- [29] R. Shankar. Recovering a quasilinear conductivity from boundary measurements. *arXiv preprint arXiv:1910.07890*, 2019.
- [30] Z. Sun. On a quasilinear inverse boundary value problem. *Mathematische Zeitschrift*, 221(1):293–305, 1996.
- [31] Z. Sun. Anisotropic inverse problems for quasilinear elliptic equations. In *Journal of Physics: Conference Series*, volume 12, page 015. IOP Publishing, 2005.
- [32] Z. Sun. An inverse boundary-value problem for semilinear elliptic equations. *Electronic Journal of Differential Equations (EJDE)[electronic only]*, 2010:Paper–No, 2010.
- [33] Z. Sun and G. Uhlmann. Inverse problems in quasilinear anisotropic media. *American Journal of Mathematics*, 119(4):771–797, 1997.
- [34] J. Sylvester and G. Uhlmann. A global uniqueness theorem for an inverse boundary value problem. *Annals of Mathematics*, pages 153–169, 1987.
- [35] J. L. Vázquez. *The porous medium equation: mathematical theory*. Oxford University Press on Demand, 2007.