On the stability of the stochastic gradient Langevin algorithm with dependent data stream^{*}

Miklós Rásonyi¹, Kinga Tikosi^{1,2}

Abstract

We prove, under mild conditions, that the stochastic gradient Langevin dynamics converges to a limiting law as time tends to infinity, even in the case where the driving data sequence is dependent.

Keywords: stochastic gradient, Langevin dynamics, dependent data

1. Stochastic gradient Langevin dynamics

Sampling from high-dimensional, possibly not even logconcave distributions is a challenging task, with far-reaching applications in optimization, in particular, in machine learning, see Raginsky et al. [1], Chau et al. [2], Barkhagen et al. [3], Brosse et al. [4].

Let $U : \mathbb{R}^d \to \mathbb{R}_+$ be a given function and consider the corresponding Langevin equation

$$d\Theta_t = -\nabla U(\Theta_t) \, dt + \sqrt{2} \, dW_t, \tag{1}$$

^{*}Both authors were supported by the "Lendület" grant 2015-6 of the Hungarian Academy of Sciences.

¹Alfréd Rényi Institute of Mathematics, Reáltandoda utca 13-15, 1053 Budapest, Hungary

²During the preparation of this paper the author attended the PhD school of Central European University, Budapest.

where W is a d-dimensional standard Brownian motion. Under suitable assumptions, the unique invariant probability μ for the diffusion process (1) has a density (with respect to the d-dimensional Lebesgue measure) that is proportional to $\exp(-U(x))$, $x \in \mathbb{R}^d$.

In practice, Euler approximations of (1) may be used for sampling from μ , i.e. a recursive scheme

$$\vartheta_{t+1}^{\lambda} = \vartheta_t^{\lambda} - \lambda \nabla U(\vartheta_t^{\lambda}) + \sqrt{2\lambda} \xi_{t+1}$$
(2)

is considered for some small $\lambda > 0$ and independent standard *d*-dimensional Gaussian sequence ξ_i , $i \ge 1$.

In some important applications, however, $U, \nabla U$ are unknown, one disposes only of unbiased estimates $H(\theta, Y_t), t \in \mathbb{N}$ of $\nabla U(\theta)$, where Y_t is some stationary data sequence. From this point on we switch to rigorous mathematics.

Let us fix integers $d, m \geq 1$ and a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. $\mathscr{B}(\mathscr{X})$ denotes the σ -algebra of the Borel-sets of a Polish space \mathscr{X} . For a random variable $X, \mathscr{L}(X)$ denotes its law. The Euclidean norm on \mathbb{R}^d or \mathbb{R}^m will be denoted by $|\cdot|$, while $||\cdot||_{TV}$ stands for the total variation distance of probability measures on $\mathcal{B}(\mathbb{R}^d)$. Let $B_r := \{\theta \in \mathbb{R}^k : |\theta| \leq r\}$ denote the ball of radius r, for $r \geq 0$, for both k = d and k = m, depending on the context. The notation Leb(\cdot) refers to the d-dimensional Lebesque-measure.

For $0 < \lambda \leq 1, t = 0, 1, ...$ and for a constant initial value $\theta_0 \in \mathbb{R}^d$ consider the recursion

$$\theta_{t+1}^{\lambda} = \theta_t^{\lambda} - \lambda H(\theta_t^{\lambda}, Y_t) + \sqrt{\lambda} \xi_{t+1}, \ t \in \mathbb{N}, \ \theta_0^{\lambda} := \theta_0, \tag{3}$$

where ξ_i , $i \ge 1$ is an i.i.d. sequence of *d*-dimensional random variables with

independent coordinates such that $\mathbb{E}[\xi_i] = 0$ and $E[|\xi_i|^2] = \sigma^2$ for some σ^2 . Furthermore, the density function f of ξ_i with respect to Leb is assumed strictly positive on every compact set. Assume that $(Y_t)_{t\in\mathbb{Z}}$ is a strict sense stationary process with values in \mathbb{R}^m an it is independent of the noise process $(\xi_t)_{t\geq 1}$. Finally, $H: \mathbb{R}^d \times \mathbb{R}^m \to \mathbb{R}^d$ is a measurable function.

A particular case of (3) is the stochastic gradient Langevin dynamics (SGLD), introduced in Welling and Teh [5], designed to learn from large datasets. See more about different versions of SGLD and their connections in Brosse et al. [4]. Note that in the present setting, unlike in SGLD, we do not assume that H is the gradient of a function and we do not assume ξ_i to be Gaussian.

A setting similar to ours was considered in Lovas and Rásonyi [6] under different assumptions. We will compare our results to those of Lovas and Rásonyi [6] at the end of Section 2 below.

The sampling error of θ_t^{λ} has been thoroughly analysed in the literature: $d(\theta_t^{\lambda}, \mu)$ has been estimated for various probability metrics d, see Chau et al. [2], Barkhagen et al. [3], Raginsky et al. [1], Brosse et al. [4]. The ergodic behaviour of θ_t^{λ} , however, has eluded attention so far. If Y_t are i.i.d. then θ_t^{λ} is a homogeneous Markov chain and standard results of Markov chain theory apply. In the more general, stationary case (considered in Barkhagen et al. [3], Chau et al. [2]), however, that machinery is not available. In the present note we study scheme (3) with stationary Y_t and establish that its law converges to a limit in total variation.

2. Main results

Assumption 1. There is a constant $\Delta > 0$ and a measurable function b: $\mathbb{R}^m \to \mathbb{R}_+$ such that, for all $\theta \in \mathbb{R}^d$ and $y \in \mathbb{R}^m$

$$\langle H(\theta, y), \theta \rangle \ge \Delta |\theta|^2 - b(y).$$
 (4)

Assumption 2. There exist constants $K_1, K_2, K_3 > 0$ and $\beta \ge 1$ such that

$$|H(\theta, y)| \le K_1 |\theta| + K_2 |y|^{\beta} + K_3.$$
(5)

Assumption 3. There exist (finite) constants $M_y, M_b > 0$ such that $\mathbb{E}[|Y_0|^{2\beta}] \le M_y$ and $\mathbb{E}[b(Y_0)] \le M_b$.

Theorem 2.1. Let Assumptions 1, 2 and 3 hold. Then, for λ small enough, the law $\mathscr{L}(\theta_t^{\lambda})$ of the iteration defined by (3) converges in total variation as $t \to \infty$ and the limit does not depend on the initialization X_0 .

In Lovas and Rásonyi [6], Δ in (4) was allowed to depend on y but b in (4) had to be constant, the process Y was assumed bounded and the process ξ Gaussian. Furthermore, in Assumption 2, β had to be 1. Under these conditions the conclusion of Theorem 2.1 was obtained, together with a rate estimate.

Theorem 2.1 above complements the results of Lovas and Rásonyi [6]: Δ must be constant in our setting but the restrictive boundedness hypothesis on Y could be removed, ξ need not be Gaussian, β in 4 can be arbitrary and b in 4 may depend on y. The examples in Section 5 demonstrate that our present results cover a wide range of relevant applications where the obtained generalizations are crucial.

3. Markov chains in random environment

The rather abstract Theorem 3.1 below, taken from Gerencsér and Rásonyi [7], is the key result we use in this paper. Let us first recall the related terminology and the assumptions.

Let \mathscr{X} and \mathscr{Y} be Polish spaces and let $(\mathscr{X}_n)_{n\in\mathbb{N}}$ (resp. $(\mathscr{Y}_n)_{n\in\mathbb{N}}$) be a nondecreasing sequence of (non-empty) Borel-sets in \mathscr{X} (resp. \mathscr{Y}). Consider a parametric family of transition kernels, i.e. a map $Q : \mathscr{Y} \times \mathscr{X} \times \mathscr{B}(\mathscr{X}) \to$ [0,1] such that for all $B \in \mathscr{B}(\mathscr{X})$ the function $(x,y) \to Q(x,y,B)$ is measurable and for every $(x,y) \in \mathscr{X} \times \mathscr{Y} Q(x,y,\cdot)$ is a probability.

An \mathscr{X} valued stochastic process $(X_t)_{t\in\mathbb{N}}$ is called a *Markov chain in a* random environment with transition kernel Q if $X_0 \in \mathscr{X}$ is deterministic (for simplicity) and

$$\mathbb{P}(X_{t+1} \in A | \mathcal{F}_t) = Q(X_t, Y_t, A), \text{ for } t \in \mathbb{N},$$
(6)

where we use the filtration $\mathcal{F}_t = \sigma(Y_k, k \in \mathbb{Z}; X_j, 0 \le j \le t).$

For a parametric family of transition kernels Q and a bounded (or non-negative) function $V : \mathscr{X} \to \mathbb{R}$ define

$$[Q(y)V](x) = \int_{\mathscr{X}} V(z)Q(x, y, dz), \text{ for } x \in \mathscr{X}.$$
(7)

Assumption 4. Let the process $(X_t)_{t\in\mathbb{N}}$ started from X_0 with be such that

$$\sup_{t\in\mathbb{N}}\mathbb{P}(X_t\notin\mathscr{X}_n)\to 0, n\to\infty.$$
(8)

Assumption 5. (Minorization condition) Let $\mathbb{P}(Y_0 \notin \mathscr{Y}_n), n \to \infty$. Assume that there exists a sequence of probability measures $(\nu_n)_{n \in \mathbb{N}}$ and a nondecreasing sequence $(\alpha_n)_{n \in \mathbb{N}}$ with $\alpha_n \in (0, 1]$ such that for all $n \in \mathbb{N}, x \in \mathbb{N}$ $\mathscr{X}_n, y \in \mathscr{Y}_n, and A \in \mathcal{B}(\mathcal{X}),$

$$Q(x, y, A) \ge \alpha_n \nu_n(A). \tag{9}$$

Theorem 3.1. (Theorem 2.11. of Gerencsér and Rásonyi [7]) Let Assumptions 4 and 5 hold. Then there exists a probability μ_* on $\mathscr{B}(\mathscr{X} \times \mathscr{Y}^{\mathbb{Z}})$ such that

$$||\mathscr{L}(X_t, (Y_{t+k})_{k\in\mathbb{Z}}) - \mu_*||_{TV} \to 0, \ as \ t \to \infty.$$

If $(X'_t)_{k \in \mathbb{N}}$ is another such Markov chain started from a different X'_0 satisfying Assumption 4 then

$$||\mathscr{L}(X_t, (Y_{t+k})_{k\in\mathbb{Z}}) - \mathscr{L}(X'_t, (Y_{t+k})_{k\in\mathbb{Z}})||_{TV} \to 0, \ as \ t \to \infty. \quad \Box$$

4. Proofs

Define the Markov chain associated to the recursive scheme (3) as

$$Q(\theta, y, A) = \mathbb{P}(\theta - \lambda H(\theta, y) + \sqrt{\lambda \xi_{n+1}} \in A),$$
(10)

for all $y \in \mathscr{Y} := \mathbb{R}^m, \, \theta \in \mathscr{X} := \mathbb{R}^d \text{ and } A \in \mathscr{B}(\mathbb{R}^d).$

Lemma 4.1. For small enough λ , under Assumptions 1 and 2, the process $(\theta_t^{\lambda})_{t \in \mathbb{N}}$ given by recursion (3) satisfies Assumption 4 with $\mathscr{X}_n := B_n$ (the ball of radius n).

PROOF. Choose $V(\theta) = |\theta|^2$. Then, since $E\xi_1 = 0$,

$$\begin{split} [Q(y)V](\theta) &= \mathbb{E}[V(\theta - \lambda H(\theta, y) + \sqrt{\lambda}\xi_1)] \\ &= |\theta|^2 + \lambda^2 |H(\theta, y)|^2 + \lambda \mathbb{E}|\xi_1|^2 - 2\lambda \langle \theta, H(\theta, y) \rangle \\ &\leq (1 - 2\lambda\Delta) |\theta|^2 + \lambda (\sigma^2 + 2b(y)) + 3\lambda^2 (K_1^2 |\theta|^2 + K_2^2 |y|^{2\beta} + K_3^2) \\ &= (1 - 2\lambda\Delta + 3\lambda^2 K_1^2) V(\theta) + \lambda (\sigma^2 + 2b(y)) + 3\lambda^2 (K_2^2 |y|^{2\beta} + K_3^2) \\ &= \gamma V(\theta) + K(y), \end{split}$$

with $K(y) = \lambda(\sigma^2 + 2b(y)) + 3\lambda^2[K_2^2|y|^{2\beta} + K_3^2]$ and $\gamma = (1 - 2\lambda\Delta + 3\lambda^2K_1^2)$. Note that for small enough $\lambda, \gamma \in (0, 1)$, independent of y.

Now using Lemma 4.2 below and setting $\theta = \theta_0$ and $y_k = Y_k$ for $k \ge 1$ we get, for each $t \ge 1$,

$$\mathbb{E}|\theta_t^{\lambda}|^2 = \mathbb{E}[Q(Y_t)Q(Y_{t-1})\dots Q(Y_1)V](\theta_0) \le \gamma^t V(\theta_0) + \sum_{i=1}^t \gamma^i \mathbb{E}K(Y_i)$$

= $\gamma^t |\theta_0|^2 + \sum_{i=1}^t \gamma^i [\lambda(\sigma^2 + 2\mathbb{E}[b(Y_i)]) + 3\lambda^2 (K_2^2 \mathbb{E}|Y_i|^{2\beta} + K_3^2)]$
 $\le |\theta_0|^2 + \frac{\gamma}{1-\gamma} [(\sigma^2 + 2M_b) + 3(K_2^2 M_y + K_3^2)] < \infty,$

by Assumption 3. Then, using Markov's inequality, we arrive at

$$\mathbb{P}(\theta_t^{\lambda} \notin \mathscr{X}_n) = \mathbb{P}(|\theta_t^{\lambda}| > n) \le \frac{\sup_t \mathbb{E}|\theta_t^{\lambda}|^2}{n^2} \to 0, \text{ as } n \to \infty. \quad \Box$$
(11)

Lemma 4.2. Assume $[Q(y)V](\theta) \leq \gamma V(\theta) + K(y)$. Then

$$[Q(y_k)Q(y_{k-1})\dots Q(y_1)V](\theta) \le \gamma^k V(\theta) + \sum_{i=1}^k \gamma^{i-1} K(y_i).$$
(12)

PROOF. We prove the statement by induction. For k = 1, it is true by assumption. Using that

$$[Q(y_2)Q(y_1)V](x) = \int_{\mathscr{X}} Q(x, y_2, dr) \int_{\mathscr{X}} V(z)Q(r, y_1, dz), \text{ for } r \in \mathscr{X}, \quad (13)$$

for k > 1 we get

$$[Q(y_k)Q(y_{k-1})\dots Q(y_1)V](\theta) = \int_{\mathscr{X}} Q(\theta, y_k, dx) [Q(y_{k-1})Q(y_{k-2})\dots Q(y_1)V](x)$$

$$\leq \int_{\mathscr{X}} \left(\gamma^{k-1}V(x) + \sum_{i=1}^{k-1} \gamma^{i-1}K(y_i) \right) Q(\theta, y_k, dx)$$

$$= \gamma^{k-1} \int_{\mathscr{X}} V(x)Q(\theta, y_k, dx) + \sum_{i=1}^{k-1} \gamma^{i-1}K(y_i)$$

$$\leq \gamma^k V(\theta) + \sum_{i=1}^k \gamma^{i-1}K(y_i). \quad \Box$$

Lemma 4.3. Define $\mathscr{X}_n = B_n$, $\mathscr{Y}_n := B_n$, $n \in \mathbb{N}$ and let Assumptions 1 and 2 hold. Then Assumption 5 is satisfied, for all λ .

Proof. For all $A \in \mathscr{B}(\mathscr{X})$,

$$\begin{split} Q(\theta, y, A) &= \mathbb{P}(\theta - \lambda H(\theta, y) + \sqrt{\lambda}\xi_1 \in A) \\ &\geq \int_{\mathbb{R}^d} \mathbb{1}_{\{\theta - \lambda H(\theta, y) + \sqrt{\lambda}\xi_1 \in A \cap B_n\}} f(w) dw \\ &= \frac{1}{\lambda^{d/2}} \int_{A \cap B_n} f\left(\frac{z - \theta + \lambda H(\theta, y)}{\sqrt{\lambda}}\right) dz \\ &\geq \frac{\operatorname{Leb}(A \cap B_n)}{\lambda^{d/2}} C(n) = \frac{\operatorname{Leb}(A \cap B_n)}{\operatorname{Leb}(B_n)} \frac{C(n) \operatorname{Leb}(B_n)}{\lambda^{d/2}} \end{split}$$

where we use that for $\theta, z \in B_n$ and $y \in B_n$ we have

$$\left|\frac{z-\theta+\lambda H(\theta,y)}{\sqrt{\lambda}}\right| \le \frac{n+n+\lambda(K_1n+K_2n^\beta+K_3)}{\sqrt{\lambda}} =: R(n),$$

therefore the integrand can be bounded from below by $C(n) := \inf_{x \in B_{R(n)}} f(x) > 0$. Then define $\nu_n(A) := \frac{\operatorname{Leb}(A \cap B_n)}{\operatorname{Leb}(B_n)}$ and $\alpha_n := \frac{C(n)\operatorname{Leb}(B_n)}{\lambda^{d/2}}$, which proves that Assumption 5 holds.

PROOF (OF THEOREM 2.1.). Follows from Lemmas 4.1, 4.3 and Theorem 3.1. $\hfill \Box$

5. Examples

5.1. Nonlinear regression

Let us consider a nonlinear regression problem which can also be seen as a one layer neural network in a supervised learning setting, where only one trainable layer connects the input and the output vectors. The training set consists of entries $Y_t = (Z_t, L_t)$ with the features $Z_t \in \mathbb{R}^{d_0}$ and the corresponding labels $L_t \in \mathbb{R}^{d_1}$ for $t \in 1, \ldots, N$. We assume that Y_t is a stationary process. Set $m := d_0 + d_1$, the dimension of Y_t .

The trainable parameters will be a matrix $W \in \mathbb{R}^{d_0 \times d_1}$ and a vector $g \in \mathbb{R}^{d_1}$, therefore the dimension of $\theta := (W, g)$ will be $d = d_0 d_1 + d_1$. The prediction function $h : \mathbb{R}^{d_0} \times \mathbb{R}^d \to \mathbb{R}^{d_1}$ is defined by $h(z, \theta) := s(Wz + g)$, where $s = (s_1, \ldots, s_{d_1})$ is a collection of nonlinear activation functions $s_i : \mathbb{R} \to \mathbb{R}$ for $i = 1, \ldots, d_1$. We will assume that each s_i and their derivatives s'_i are all bounded by some constant M_s for $i = 1, \ldots, d_1$.

Choosing the loss function to be mean-square error, one aims to minimize the empirical risk, that is

$$\min\{\mathbb{E}[|h(Z_t,\theta) - L_t|^2] + \kappa |\theta|^2\},\tag{14}$$

with some $\kappa > 0$, where the second term is added for regularization.

It is standard to solve this optimization step using gradient-based methods. For $y = (z, l) \in \mathbb{R}^{d_0} \times \mathbb{R}^{d_1}$ denote $U(\theta, y) = |h(z, \theta) - l|^2 + \kappa |\theta|^2$ and the updating function to be used in the algorithm will be

$$H(\theta, y) = \nabla U(\theta, y) = \frac{\partial}{\partial \theta} |h(z, \theta) - l|^2 + 2\kappa \theta.$$
(15)

Lemma 5.1. The function $H(\theta, y)$ defined as above satisfies Assumptions 1 and 2. *Proof.* Using the chain rule, a short calculation gives

$$\left|\frac{\partial}{\partial\theta}|h(z;\theta) - l|^2\right| = \sqrt{\sum_{i=1}^{d_0+1} \sum_{j=1}^{d_1} \left(2(h(z;\theta)_j - l_j)s_j'(\langle W_j, z \rangle + g_j)z_i\right)^2}, \quad (16)$$

where we define $z_{d_0+1} = 1$ and W_j stands for the *j*th row of *W*. Notice that by the boundedness of s' and s this is bounded in θ and at most quadratic in y. Then Assumption 2 is satisfied with $\beta = 2$.

Using the same argument about the boundedness of s and s'

$$\left| \left\langle \frac{\partial}{\partial \theta} |h(z;\theta) - l|^2, \theta \right\rangle \right| = \left| \sum_{i=1}^{d_0+1} \sum_{j=1}^{d_1} 2(h(z;\theta)_j - l_j) s_j'(W_j z + g_j) z_i \theta_{i,j} \right| \\ \leq C_0 dM_s \left(|y|^2 + 1 \right) |\theta|,$$

for some $C_0 > 0$. Using that $\left\langle \frac{\partial}{\partial \theta} \kappa |\theta|^2, \theta \right\rangle = 2\kappa |\theta|^2$, we get that $\left\langle \nabla U(\theta), \theta \right\rangle \ge c |\theta|^2 - C(|y|^4 + 1)$ with some c, C therefore Assumption 1 is satisfied with b(y) being of degree 4 in y.

5.2. A tamed algorithm for neural networks

It has been observed that in multi-layer neural networks quadratic regularization is not sufficient to guarantee dissipativity, while adding a higher order term would violate Lipschitz continuity. So the standard SGLD algorithm diverges anyway. To remedy this, certain "tamed" schemes have been suggested in Lovas et al. [8].

In contrast to the previous case now we will hidden layers between the input and output: layer 0 is the input, layer n is the output and $1, \ldots, n-1$ are the hidden layers of the neural network for some n > 1. The prediction function h will be defined as the composition of a sequence of n+1 linear transformations and activation functions, i.e. $h(z, \theta) = s_n(W_n s_{n-1}(W_{n-1} \ldots s_0(W_0 z)))$ where θ is the collection of all parameters $W_i \in \mathbb{R}^{d_{i-1}} \times \mathbb{R}^{d_i}$, $i = 1, \ldots, n$ and $s_i : \mathbb{R}^{d_i} \to \mathbb{R}^{d_i}$ is a componentwise non-linear activation function, assumed bounded together with its derivatives by some constant M_s . Therefore $h : \mathbb{R}^{d_0} \times \mathbb{R}^d \to \mathbb{R}^{d_n}$, where $d = \sum_{i=1}^n d_{i-1}d_i$ is the dimension of θ . For the case of simplicity in this case we assumed that there is no bias term g. The training set consists of entries $Y_t = (Z_t, L_t)$ with the features $Z_t \in \mathbb{R}^{d_0}$ and the corresponding labels $L_t \in \mathbb{R}^{d_n}$, the dimension of each Y_t is $m = d_0 + d_n$. We assume that Y_t is a stationary process.

As in the previous subsection, the regularized empirical risk has the form $U(\theta, y) = |h(z, \theta) - l|^2 + \frac{\eta}{2(r+1)} |\theta|^{2(r+1)}$ with some $r \ge 0, \eta > 0$. Denoting $G(\theta, y) = \nabla U(\theta, y)$, the "tamed" updating function we use will be defined as $H(\theta, y) := \frac{G(\theta, y)}{1 + \sqrt{\lambda} |\theta|^{2r}}$, for every $\theta \in \mathbb{R}^d, y \in \mathbb{R}^m$. Note that this function depends on λ !

We will use the following.

Lemma 5.2. (Proposition 4 of Lovas et al. [8])

$$\left|\frac{\partial}{\partial\theta}|h(z,\theta) - l|^2\right| \le C(1+|y|)^2 \left(1+|\theta|^{n+1}\right),\tag{17}$$

where C > 0 depends on $D = \max_{j=1,\dots,n} d_j$, n and M_s .

Lemma 5.3. For λ small enough, the conclusions of Theorem 2.1 hold for the scheme (3) with $H(\theta, y)$ defined as above, provided that $r \geq \frac{n+2}{2}$ and Assumption 3 holds.

Proof. Using Lemma 5.2, Assumption 2 can be checked as follows:

$$|H(\theta, y)| = \left| \frac{\frac{\partial}{\partial \theta} |h(z; \theta) - l|^2 + \eta \theta |\theta|^{2r}}{1 + \sqrt{\lambda} |\theta|^{2r}} \right|$$

$$\leq \left| \frac{C(1 + |y|)^2 (1 + |\theta|^{n+1})}{1 + \sqrt{\lambda} |\theta|^{2r}} \right| + \left| \frac{\eta \theta |\theta|^{2r}}{1 + \sqrt{\lambda} |\theta|^{2r}} \right| \leq K_1 |\theta| + K_2 |y|^{\beta} + K_3,$$

where $K_1 = \frac{\eta}{\sqrt{\lambda}}$, $\beta = 2$ and the constants K_2 and K_3 depend on λ, η, n and C.

Let us check Assumption 1. For the regularization term we have

$$\left\langle \frac{\eta \theta |\theta|^{2r}}{1 + \sqrt{\lambda} |\theta|^{2r}}, \theta \right\rangle = \frac{\eta |\theta|^{2r+2}}{1 + \sqrt{\lambda} |\theta|^{2r}} \ge \min\left\{ \frac{\eta}{2\sqrt{\lambda}}, \frac{\eta}{2} \right\} |\theta|^2 \ge \frac{\eta}{2} |\theta|^2 \tag{18}$$

for λ small enough.

The Cauchy inequality, Lemma 5.2 and the choice of r ensures that

$$\left| \left\langle \frac{\frac{\partial}{\partial \theta} \left(|h(z;\theta) - l|^2 \right)}{1 + \sqrt{\lambda} |\theta|^{2r}}, \theta \right\rangle \right| \le \frac{C(1 + |\theta|^{n+2})(1 + |y|^2)}{1 + \sqrt{\lambda} |\theta|^{2r}} \le K'(1 + |y|^2), \quad (19)$$

for some K' > 0. Now combining these estimates, we get

$$\langle H(\theta, y), \theta \rangle \ge \frac{\eta}{2} |\theta|^2 - K'(1+|y|^2), \tag{20}$$

therefore Assumption 1 is satisfied with $\Delta = \frac{\eta}{2}$ and b(y) is quadratic in y.

We can check that $\gamma = (1 - \eta \sqrt{\lambda} + \lambda \eta^2) < 1$ in Lemma 4.1 for λ small enough so the proof of Theorem 2.1 goes through for this choice of H.

Allowing b to be of degree 4, $\frac{n+2}{2}$ in Lemma 5.3 could be decreased to $\frac{n+1}{2}$, as easily seen.

References

- M. Raginsky, A. Rakhlin, M. Telgarsky, Non-convex learning via stochastic gradient Langevin dynamics: a nonasymptotic analysis, Proceedings of Machine Learning Research 65 (2017) 1674–1703.
- [2] N. H. Chau, Eric. Moulines, M. Rásonyi, S. Sabanis, Y. Zhang, On stochastic gradient Langevin dynamics with dependent data streams: the fully non-convex case, Preprint, arXiv:1905.13142 (2021).

- [3] M. Barkhagen, N. H. Chau, Éric. Moulines, M. Rásonyi, S. Sabanis, Y. Zhang, On stochastic gradient Langevin dynamics with dependent data streams in the logconcave case, Bernoulli 27 (2021) 1–33.
- [4] N. Brosse, A. Durmus, E. Moulines, The promises and pitfalls of stochastic gradient Langevin dynamics, in: Advances in Neural Information Processing Systems, 2018, pp. 8268–8278.
- [5] M. Welling, Y. W. Teh, Bayesian learning via stochastic gradient Langevin dynamics, in: Proceedings of the 28th international conference on machine learning (ICML-11), 2011, pp. 681–688.
- [6] A. Lovas, M. Rásonyi, Markov chains in random environment with applications in queueing theory and machine learning, To appear in Stochastic Processes and their Applications, arXiv:1911.04377 (2021).
- [7] B. Gerencsér, M. Rásonyi, Invariant measures for fractional stochastic volatility models, Preprint, arXiv:2002.04832v1 (2020).
- [8] A. Lovas, I. Lytras, M. Rásonyi, S. Sabanis, Taming neural networks with TUSLA: Non-convex learning via adaptive stochastic gradient Langevin algorithms, Preprint, arXiv:2006.14514 (2021).