FPT algorithms for packing k-safe spanning rooted sub(di)graphs

Stéphane Bessy D

LIRMM, Université de Montpellier, CNRS, Montpellier, France stephane.bessy@lirmm.fr

Florian Hörsch

Université Grenoble Alpes, Grenoble INP, CNRS, G-SCOP, Grenoble, France florian.hoersch@grenoble-inp.fr

Ana Karolinna Maia 🗅

Departamento de Computação, Universidade Federal do Ceará, Fortaleza, Brazil karolmaia@ufc.br

Dieter Rautenbach ®

Institute of Optimization and Operations Research, Ulm University, Germany dieter.rautenbach@uni-ulm.de

Ignasi Sau 👨

LIRMM, Université de Montpellier, CNRS, Montpellier, France ignasi.sau@lirmm.fr

Abstract

We study three problems introduced by Bang-Jensen and Yeo [Theor. Comput. Sci. 2015] and by Bang-Jensen, Havet, and Yeo [Discret. Appl. Math. 2016] about finding disjoint "balanced" spanning rooted substructures in graphs and digraphs, which generalize classic packing problems. Namely, given a positive integer k, a digraph D=(V,A), and a root $r\in V$, we consider the problem of finding two arc-disjoint k-safe spanning r-arborescences and the problem of finding two arc-disjoint (r,k)-flow branchings. We show that both these problems are FPT with parameter k, improving on existing XP algorithms. The latter of these results answers a question of Bang-Jensen, Havet, and Yeo [Discret. Appl. Math. 2016]. Further, given an integer k, a graph G=(V,E), and $r\in V$, we consider the problem of finding two arc-disjoint (r,k)-safe spanning trees. We show that this problem is also FPT with parameter k, again improving on a previous XP algorithm. Our main technical contribution is to prove that the existence of such spanning substructures is equivalent to the existence of substructures with size and maximum (out-)degree both bounded by a (linear or quadratic) function of k, which may be of independent interest.

2012 ACM Subject Classification Design and analysis of algorithms → Fixed parameter tractability.

Keywords and phrases Digraphs, packing problems, arborescences, branching flows, safe spanning trees, parameterized complexity, fixed-parameter tractability.

Funding Stéphane Bessy: DIGRAPHS (ANR-19-CE48-0013-02).

Ana Karolinna Maia: FUNCAP Pronem 4543945/2016 and CAPES/STIC-AmSud 88881.197438/2018-01. *Ignasi Sau*: DEMOGRAPH (ANR-16-CE40-0028), ESIGMA (ANR-17-CE23-0010), ELIT (ANR-20-CE48-0008-01), and French-German Collaboration ANR/DFG Project UTMA (ANR-20-CE92-0027).

1 Introduction

This article deals with finding certain disjoint substructures in graphs and digraphs. Throughout the article, when given a graph or a digraph, we use n for its number of vertices.

All graphs and digraphs considered in this paper are loopless, but may have multiple edges or arcs. Given a graph G = (V, E), we say that an edge e = uv is *incident* to u and v. For some $X \subseteq V$, we denote by $d_G(X)$ the number of edges that are incident to exactly

one vertex in X. For some $v \in V$, we use $N_G(v)$ for the set of vertices w such that there is an edge between v and w. A rooted graph is a graph G = (V + r, E) with a special vertex r called the root. A vertex $v \in V$ with $d_G(v) = 1$ is called a leaf of G.

Given a digraph D=(V,A) and some $X\subseteq V$, we use $\delta_D^-(X)$ (resp. $\delta_D^+(X)$) for the set of arcs entering (resp. leaving) X. We use $d_D^-(X)$ (resp. $d_D^+(X)$) for $|\delta_D^-(X)|$ (resp. $|\delta_D^+(X)|$). For a single vertex v, we abbreviate $\delta_D^+(\{v\})$ (resp. $\delta_D^-(\{v\}), d_D^+(\{v\}), d_D^-(\{v\})$) to $\delta_D^+(v)$ (resp. $\delta_D^-(v), d_D^+(v), d_D^-(v)$). We call $d_D^-(v)$ (resp. $d_D^+(v)$) the in-degree (resp. out-degree) of v. We use $N_{D}^{+}(v)$ for the set of vertices w such that there is an arc from v to w. A vertex $v \in V$ with $d^+(v) = 0$ is called a *sink* of D. Subscripts may be omitted when they are clear from the context. The underlying graph of D is obtained by replacing all arcs of A by an edge between the same two vertices. A rooted digraph is a digraph D = (V + r, A) with a special vertex r, called the *root*, whose in-degree is 0.

Packing k-safe spanning r-arborescences. The first objects we deal with are called arborescences. We remark here that the term out-branching is used in [3] to describe the same object. An r-arborescence is a directed graph X = (V + r, A) such that the underlying graph of X is a tree, the in-degree of r is 0 in X, and the in-degree of all other vertices is 1 in X. We say that r is the root of X. Observe that every $v \in V$ is the root of a unique subarborescence of X. We denote this subarborescence by B_X^v . Given a digraph D, an r-arborescence X that is a subdigraph of D is spanning in D if it has the same vertex set as D. The following is a fundamental result in digraph theory.

▶ Theorem 1 (Edmonds [6]). Let D = (V + r, A) be a rooted digraph and k a positive integer. There exists a set of k arc-disjoint spanning r-arborescences in D if and only if $d_A^-(X) \ge k \text{ for all } \emptyset \ne X \subseteq V.$

A number of alternative proofs of Theorem 1 have been found, several of which are algorithmic and yield polynomial-time algorithms to find the desired arc-disjoint spanning r-arborescences, if they exist [7,11].

This naturally raises the question whether we also can efficiently find spanning arborescences satisfying some extra properties. This consideration, as well as practical applications concerning protection against arc failures, motivated Bang-Jensen and Yeo [4] to introduce the notion of k-safe spanning arborescences. An r-arborescence X = (V + r, A) is called k-safe if $n-|V(B_X^v)| \geq k$ for all $v \in V$. Notice that it is enough that only the out-neighbours of r satisfy this latter condition for X to be k-safe. They proved the following negative result showing that in general not even the problem of finding a single k-safe spanning r-arborescence is tractable.

Theorem 2 (Bang-Jensen and Yeo [4]). Given a rooted digraph D = (V + r, A), deciding whether D has an (n-k)-safe spanning r-arborescence is NP-complete for any fixed $k \geq 3$.

In this light, a characterization in the shape of Theorem 1 clearly seems out of reach. It remains nevertheless interesting to investigate the possibility of finding arc-disjoint k-safe spanning r-arborescences for small values of k. This question has been dealt with by Bang-Jensen, Havet, and Yeo [3]. On the negative side, they implicitly proved the following result, which shows that there is little hope to algorithmically find arc-disjoint k(n)-safe spanning r-arborescences if k is a function that does not grow too slowly. While a polynomial-time algorithm for the problem they consider would not imply P = NP, it would imply the failure of the Exponential Time Hypothesis (ETH for short) of Impagliazzo and Paturi [9], stating that there is an $\varepsilon > 0$ such that there is no algorithm for solving a 3-SAT formula with ℓ variables and m clauses in time $2^{\varepsilon \ell} \cdot (\ell + m)^{O(1)}$.

▶ Theorem 3 (Bang-Jensen, Havet, and Yeo [3]). Suppose that the ETH holds, let $\varepsilon > 0$ be arbitrary and let $k : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$ be a function such that $(\log(n))^{1+\varepsilon} \leq k(n) \leq \frac{n}{2}$ for all n > 0. Further, suppose that there exists a constant C^* such that for all $c \geq C^*$ there exists an n such that k(n) = c. Then there is no algorithm running in time $n^{O(1)}$ for deciding whether a given rooted digraph D = (V + r, A) has two arc-disjoint k(n)-safe spanning r-arborescences.

On the positive side, they show that the problem becomes tractable when fixing the value of k. While several results considered in this article hold for finding an arbitrary number of disjoint objects, we focus on the case where we want to find just two of them in order to avoid technicalities. In Section 7 we discuss the generalization to more than two objects.

▶ **Theorem 4** (Bang-Jensen, Havet, and Yeo [3]). Deciding whether a given rooted digraph D = (V + r, A) contains two arc-disjoint k-safe spanning r-arborescences is XP with parameter k.

(See Subsection 2.5 for the definition of the classes XP and FPT.) Our first contribution is to improve Theorem 4 by showing that the problem is *fixed-parameter tractable* (FPT).

▶ Theorem 5. Deciding whether a given rooted digraph D = (V + r, A) contains two arc-disjoint k-safe spanning r-arborescences is FPT with parameter k. More precisely, it can be solved in time $2^{O(k^2 \cdot \log k)} \cdot n^{O(1)}$. Further, if they exist, the two arc-disjoint k-safe spanning r-arborescences can be computed within the same running time.

Packing (r,k)-flow branchings. The second structure we consider, which was introduced by Bang-Jensen and Bessy [1], builds a connection between the theory of finding arc-disjoint spanning arborescences and flow problems. A flow in a digraph D=(V,A) is a function $\mathbf{z}:A\to\mathbb{Z}_{\geq 0}$. Given a rooted digraph X=(V+r,A) and a capacity function $\mathbf{c}:A\to\mathbb{Z}_{\geq 0}$, an (r,\mathbf{c}) -branching flow is a flow $\mathbf{z}:A\to\mathbb{Z}_{\geq 0}$ such that $\mathbf{z}(a)\leq\mathbf{c}(a)$ for all $a\in A$, $\mathbf{z}(\delta^+(r))-\mathbf{z}(\delta^-(r))=n-1$ and $\mathbf{z}(\delta^-(v))-\mathbf{z}(\delta^+(v))=1$ for all $v\in V$. If X admits an (r,\mathbf{c}) -branching flow, we say that D is an (r,\mathbf{c}) -flow branching. If for some positive integer k, we have $\mathbf{c}(a)=n-k$ for all $a\in A$, we speak of an (r,k)-branching flow and an (r,k)-flow branching. Given a digraph D, an (r,k)-branching flow X that is a subdigraph of D is spanning in D if it has the same vertex set as D. If D admits a k-safe spanning r-arborescence, then it is easy to see that D is an (r,k)-flow branching, but the converse is not necessarily true, as pointed out in [3].

Bang-Jensen and Bessy [1] consider the problem of finding arc-disjoint spanning flow branchings in a rooted digraph. Among others, they show the following negative result that makes a characterization in the shape of Theorem 1 seem out of reach.

▶ Theorem 6 (Bang-Jensen and Bessy [1]). Given a rooted digraph D = (V + r, A) and a capacity function $\mathbf{c} : A \to \mathbb{Z}_{\geq 0}$, it is NP-complete to decide whether D contains two arc-disjoint spanning (r, \mathbf{c}) -flow branchings even if $\mathbf{c}(a) \in \{1, 2\}$ for all $a \in A$.

The above result has been strengthened in [3].

▶ Theorem 7 (Bang-Jensen, Havet, and Yeo [3]). Given a rooted digraph D = (V + r, A) and a fixed positive integer $k \ge 2$, it is NP-complete to decide whether D contains two arc-disjoint spanning (r, n - k)-flow branchings.

On the other hand, the problem of finding arc-disjoint spanning (r, k)-flow branchings for some small values of k turns out to be more tractable. The study of such spanning flow branchings has surprisingly many similarities with the study of k-safe spanning arborescences.

On the negative side, the following result is proven in [3]. It shows that there is little hope to algorithmically find arc-disjoint spanning (r, k(n))-flow branchings if k is a function that does not grow too slowly. It can be viewed as an analogue of Theorem 3 for (r, k)-flow branchings.

▶ Theorem 8 (Bang-Jensen, Havet, and Yeo [3]). Suppose that ETH holds, let $\varepsilon > 0$ be arbitrary and let $k: \mathbb{Z}_{>0} \to \mathbb{Z}_{>0}$ be a function such that $(\log(n))^{1+\varepsilon} \le k(n) \le \frac{n}{2}$ for all n > 0. Further, suppose that there exists a constant C^* such that for all $c \geq C^*$ there exists an n such that k(n) = c. Then there is no algorithm running in time $n^{O(1)}$ for deciding whether a given rooted digraph D = (V + r, A) has two arc-disjoint spanning (r, k(n))-flow branchings.

On the positive side, Bang-Jensen and Bessy [1] showed that the case k=1 can be solved in polynomial time. This result was again generalized by Bang-Jensen, Havet, and Yeo [3], who proved that the problem can be solved in polynomial time for every fixed value of $k \geq 1$. The following result can be viewed as an analogue of Theorem 4 for (r, k)-flow branchings.

Theorem 9 (Bang-Jensen, Havet, and Yeo [3]). Deciding whether a given rooted digraph D =(V+r,A) contains two arc-disjoint spanning (r,k)-flow branchings is XP with parameter k.

The authors of [3] ask whether the above problem is FPT. Our second contribution is an affirmative answer to this question.

- **Theorem 10.** Deciding whether a given rooted digraph D = (V + r, A) contains two arc-disjoint spanning (r, k)-flow branchings is FPT with parameter k. More precisely, it can be solved in time $2^{O(k^2 \cdot \log k)} \cdot n^{O(1)}$. Further, if they exist, the two arc-disjoint spanning (r,k)-flow branchings can be computed within the same running time.
- **Packing** (r,k)-safe spanning trees. Finally, we consider a similar problem in undirected graphs that has also been introduced in [3]. Given a graph G = (V, E), a spanning tree is a subgraph T of G that is a tree with V(T) = V. The theory of finding edge-disjoint spanning trees is also pretty rich. The most fundamental result is the following one.
- ▶ Theorem 11 (Tutte [12]). Let G = (V, E) be a graph and k a positive integer. Then G has k edge-disjoint spanning trees if and only if $\sum_{i=1}^q d_G(V_i) \ge 2k(q-1)$ for every partition $\{V_1, \ldots, V_q\} \ of \ V$.

An algorithmic proof of Theorem 11, yielding a polynomial-time algorithm, can be found in [8]. Again, we may wish to also find edge-disjoint spanning trees satisfying certain extra properties. Given a rooted tree T = (V + r, E) and some $v \in V$, we use C_v^v for the subgraph of T-v that arises from deleting the component of T-v containg r. We say that T is (r,k)-safe if for every $v \in V$, we have $|V - V(C_T^v)| \ge k$.

▶ Theorem 12 (Bang-Jensen, Havet, and Yeo [3]). Deciding whether a given rooted graph G = (V + r, E) contains two edge-disjoint (r, k)-safe spanning trees is XP with parameter k.

Again, we improve Theorem 12 as follows.

Theorem 13. Deciding whether a given rooted graph G = (V + r, E) contains two edgedisjoint (r, k)-safe spanning trees is FPT with parameter k. More precisely, it can be solved in time $2^{O(k^2 \cdot \log k)} \cdot n^{O(1)}$. Further, if they exist, the two edge-disjoint (r,k)-safe spanning trees can be computed within the same running time.

Since a hardness result in the spirit of Theorem 3 and Theorem 8 was not provided in [3], we fill this gap and prove the following theorem, whose proof is inspired by the one of [3, Theorem 5.2]. It shows that the problem is hard even if we want to find one single (r, k)-safe spanning tree.

▶ Theorem 14. Let $p \ge 1$ be a fixed positive integer. Deciding whether a given rooted graph G = (V + r, A) has p edge-disjoint (r, k)-safe spanning trees is NP-complete. Moreover, let $\varepsilon > 0$ be arbitrary and let $k : \mathbb{Z}_{\ge 0} \to \mathbb{Z}_{\ge 0}$ be a function such that $(\log(n))^{2+\varepsilon} \le k(n) \le \frac{n}{2}$ for all n > 0. Further, suppose that there exists a constant C^* such that for all $c \ge C^*$ there exists an n such that k(n) = c. Then, assuming the ETH, there is no algorithm running in time $n^{O(1)}$ for deciding whether a given rooted graph contains p edge-disjoint (r, k)-safe spanning trees.

Our techniques. In order to obtain the FPT algorithms for the three considered problems, we follow a common strategy. In a nutshell, the main ideas used in the XP algorithms of [3], and how we manage to improve them to FPT algorithms, can be summarized as follows. The algorithms of Bang-Jensen, Havet, and Yeo [3] are all based on proving the following general property for each of the considered problems, where all the substructures are rooted:

A given (di)graph contains the required substructure \mathcal{X} (i.e., a pair of disjoint spanning arborescences, flow branchings, or spanning trees) if and only if it contains another type of substructure \mathcal{X}' of size bounded by a function of k and such that, if found, it can be extended to the required substructure \mathcal{X} in polynomial time.

Once the above property is proved, an XP algorithm follows naturally: generate all candidate substructures \mathcal{X}' in time $n^{f(k)}$ and, for each of them, try to extend it to a substructure \mathcal{X} in polynomial time. Our main contribution is to prove that the above general property is still true if we replace \mathcal{X}' with another type of substructure \mathcal{X}'' having the crucial property that the candidate substructures \mathcal{X}'' can be all enumerated in time $f(k) \cdot n^{O(1)}$, hence yielding an FPT algorithm. In order to achieve this, we prove that we can restrict ourselves to objects \mathcal{X}'' whose "non-sink" vertices (i.e., those with positive (out-)degree in \mathcal{X}'') have (out-)degree, in the original (di)graph, bounded by some function of k, namely O(k) or $O(k^2)$. Intuitively, this is possible because, given a pair $\mathcal{X}' = \{X_1, X_2\}$ containing a vertex v of large (out-)degree (as a function of k) in, say, X_1 , we can safely prune the "branch" of X_1 hanging from v, with the guarantee that it will always be possible to extend the pruned substructure to another substructure of the original type. Note that since the substructures \mathcal{X}'' have size and maximum (out-)degree bounded by a function of k, we can indeed generate all candidate substructures in time $f(k) \cdot n^{O(1)}$, as required. We now make this informal explanation more concrete. For all technical definitions used in the next paragraphs, see Section 2.

Let us first focus on the problem of finding two arc-disjoint k-safe spanning r-arborescences. In this case, the substructure \mathcal{X}' is an extendable pair of arc-disjoint classic (r,k)-kernels. In Lemma 21, we restate a result from [3] that shows that the existence of this substructure is sufficient for the existence of two arc-disjoint k-safe spanning r-arborescences. We then introduce compact (r,k)-kernels. An extendable pair of arc-disjoint compact (r,k)-kernels corresponds to the substructure \mathcal{X}'' . In Lemma 28, we show that the existence of an extendable pair of arc-disjoint compact (r,k)-kernels is also sufficient for the existence of two arc-disjoint k-safe spanning r-arborescences. The proof of Lemma 28 is our main technical contribution. Having Lemma 28 at hand, the proof of Theorem 5 is easy.

As for packing (r, k)-flow branchings, the substructure \mathcal{X}' defined in [3] is an extendable pair of arc-disjoint classic (r, k)-cores. In Lemma 25, we restate a result from [3] that shows

that the existence of this substructure is sufficient for the existence of two arc-disjoint (r, k)flow branchings. We then introduce compact (r, k)-cores. An extendable pair of arc-disjoint compact (r,k)-cores corresponds to the substructure \mathcal{X}'' . In Lemma 29, we show that the existence of an extendable pair of arc-disjoint compact (r,k)-cores is also sufficient for the existence of two arc-disjoint (r, k)-flow branchings. Again, the proof of Lemma 29 is our main technical contribution and is similar to the one of Lemma 28. Again, having Lemma 29 at hand, the proof of Theorem 10 is easy.

Finally, for packing (r, k)-safe spanning trees, the substructure \mathcal{X}' defined in [3] is a completable pair of edge-disjoint classic (r,k)-certificates. In Lemma 27, we restate a result from [3] that shows that the existence of this substructure is sufficient for the existence of two arc-disjoint (r, k)-safe spanning trees. We then introduce compact (r, k)-certificates. A completable pair of edge-disjoint compact (r, k)-certificates corresponds to the substructure \mathcal{X}'' . In Lemma 30, we show that the existence of a completable pair of edge-disjoint compact (r,k)-certificates is also sufficient for the existence of two edge-disjoint (r,k)-safe spanning trees. Again, the proof of Lemma 30 is our main technical contribution and is similar to the one of Lemma 28 and Lemma 29. Again, having Lemma 30 at hand, the proof of Theorem 13 is easy.

Organization. In Section 2 we review some more technical results we need from [3], prove several preliminary results, and provide the basic definitions about parameterized complexity. In Section 3, Section 4, Section 5, and Section 6 we give the proof of Theorem 5, Theorem 10, Theorem 13, and Theorem 14, respectively. Finally, we conclude our work in Section 7.

2 **Preliminaries**

In this section we collect some more technical previous results and prove several preliminary statements. We first give some general results on graphs and digraphs and then some which are more specific to each of the particular applications. We also provide some basic definitions about parameterized complexity.

2.1 **General preliminaries**

The following is a well-known submodularity property of digraphs that can be found, for instance, in [8, Proposition 1.2.1].

▶ Proposition 15. Let D = (V, A) be a digraph and $S_1, S_2 \subseteq V$. Then $d_D^-(S_1) + d_D^-(S_2) \le V$ $d_D^-(S_1 \cup S_2) + d_D^-(S_1 \cap S_2).$

A rooted digraph D = (V+r, A) is called k-root-connected if $d_D^-(X) \ge k$ for all $X \subseteq V$. We use root-connected for 1-root-connected. In a root-connected rooted digraph D = (V + r, A) an arc $a \in A$ is called *critical* if D-a is not root-connected anymore. We now use Proposition 15 to obtain a result that will be useful when dealing with both k-safe spanning r-arborescences and spanning (r, k)-flow branchings.

▶ **Lemma 16.** Let D = (V+r, A) be a 2-root-connected rooted digraph and let D' = (V+r, A')be a root-connected rooted digraph that is obtained from D by deleting α arcs of A. Then for any $v \in V + r$, there are at most α arcs which are critical in D' and whose tail is v.

Proof. We proceed by induction on α . The statement is trivial for $\alpha = 0$. We suppose that it holds for all integers up to some α and show that it also holds for $\alpha+1$. Let $v\in V+r$ and let $\{a_1, \ldots, a_{\alpha+1}\}$ be a set of arcs in A such that $D_2 = D - \{a_1, \ldots, a_{\alpha+1}\}$ is root-connected. By the inductive hypothesis, v is the tail of at most α critical arcs in $D_1 = D - \{a_1, \dots, a_{\alpha}\}$. Suppose, for the sake of a contradiction, that there are two arcs vw_1, vw_2 which are critical in D_2 , but not in D_1 . It follows that there are sets $X_1, X_2 \subseteq V$ such that $d_{D_2}^-(X_1), d_{D_2}^-(X_2) = 1$, $d_{D_1}^-(X_1), d_{D_1}^-(X_2) = 2$, and vw_i enters X_i . Since $D_2 = D_1 - a_{\alpha+1}$, it follows that $a_{\alpha+1}$ enters $X_1 \cap X_2$, so $X_1 \cap X_2 \neq \emptyset$. As D_2 is root-connected, we have $d_{D_2}^-(X_1 \cap X_2) \geq 1$. As vw_i enters X_i , we have $v \in V + r - (X_1 \cup X_2)$ and so both vw_1 and vw_2 enter $X_1 \cup X_2$. This yields $d_{D_2}^-(X_1) + d_{D_2}^-(X_2) = 1 + 1 < 1 + 2 \leq d_{D_2}^-(X_1 \cap X_2) + d_{D_2}^-(X_1 \cup X_2)$, a contradiction to Proposition 15.

Given a 2-root-connected rooted digraph D = (V + r, A), a pair of subdigraphs $(X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2))$ of D is called *extendable* if both $D - A_1$ and $D - A_2$ are root-connected. The following is an immediate consequence of the fact that checking whether a digraph is root-connected can clearly be done in polynomial time.

▶ Lemma 17. Given a rooted digraph D = (V + r, A) and a pair of two subdigraphs $(X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2))$, we can decide in polynomial time whether (X_1, X_2) is extendable.

We now switch to some results in undirected graphs.

▶ Lemma 18. Let G = (V, E) be a graph and let T be a spanning tree of G. Let $e = uv \in E - E(T)$. Then there is some $f \in E(T)$ that is incident to u such that T - e + f is a spanning tree of G.

Proof. The graph T + f contains a unique cycle C such that $uv \in E(C)$ and the deletion of an arbitrary edge of E(C) yields a spanning tree of G. As C is is a cycle, E(C) contains an edge f different from e that is incident to u. This edge satisfies the condition.

The following result can be found in a stronger form in [8, Theorem 5.3.3].

▶ Proposition 19. Let G = (V, E) be a graph and let T_1, T_2 be spanning trees of G. Then there is a function $\sigma : E(T_1) \to E(T_2)$ such that for all $e \in E(T_1)$ both $T_1 - e + \sigma(e)$ and $T_2 - \sigma(e) + e$ are spanning trees of G.

We call a function like in Proposition 19 a tree-mapping function from T_1 to T_2 .

▶ Lemma 20. Let G = (V, E) be a graph, let T_1, T_2 be spanning trees of G, and let σ : $E(T_1) \to E(T_2)$ be a tree-mapping function from T_1 to T_2 . Further, let $e_1, e_2, e_3 \in E(T_1)$ be all incident to a common vertex v. Then $\{\sigma(e_1), \sigma(e_2), \sigma(e_3)\}$ contains at least two distinct elements.

Proof. As T_1 is a spanning tree, $T_1 - \{e_1, e_2, e_3\}$ contains three components C_1, C_2, C_3 none of which contains v such that e_i is incident to a vertex in $V(C_i)$ for i = 1, 2, 3. As $T_1 - e_i + \sigma(e_i)$ is a spanning tree, we obtain that $\sigma(e_i)$ is incident to a vertex in $V(C_i)$ for i = 1, 2, 3. As $V(C_1), V(C_2)$ and $V(C_3)$ are pairwise disjoint, the statement follows.

2.2 Preliminaries on k-safe spanning r-arborescences

Given a rooted digraph D = (V + r, A), a classic (r, k)-kernel is a subarborescence X = (V' + r, A') of D such that X is k-safe and |V'| = 2k - 2. The XP algorithm of Theorem 4 is based on the following result, which we reformulate here using our terminology.

▶ Lemma 21 (Bang-Jensen, Havet, and Yeo [3]). Let D = (V + r, A) be a rooted digraph with $|V| \ge 2k - 2$. Then D contains two arc-disjoint k-safe spanning r-arborescences if and only if D contains an extendable pair of arc-disjoint classic (r, k)-kernels. Further, the two arc-disjoint k-safe spanning r-arborescences can be constructed from the extendable pair of classic (r, k)-kernels in polynomial time.

2.3 Preliminaries on spanning (r, k)-flow branchings

We first need the following result that allows to recognize (r, k)-flow branchings.

▶ Lemma 22 (Bang-Jensen, Havet, and Yeo [3]). Given a rooted digraph D = (V + r, A) and a non-negative integer k, we can decide in polynomial time whether D is an (r, k)-flow branching.

Given a digraph D=(V,A) and two vertices $u,v\in V$, a uv-path flow is a flow \mathbf{c} such that $\mathbf{c}(a)=1$ for all arcs $a\in A(P)$ and $\mathbf{c}(a)=0$ for all $a\in A-A(P)$ for some uv-path P. Similarly, a cycle flow is a flow \mathbf{c} such that $\mathbf{c}(a)=1$ of all $a\in A(C)$ and $\mathbf{c}(a)=0$ for all $a\in A-A(C)$ for some cycle C. We need the following result on flows which is proven in a more general form in [2].

▶ Lemma 23. Let X = (V + r, A) be a flow branching and $\mathbf{z} : A \to \mathbb{Z}_{\geq 0}$ be a branching flow in X. Then there is an rv-path flow \mathbf{z}_v for all $v \in V$ and a set of cycle flows $\{\mathbf{z}_C : C \in \mathcal{C}\}$ for some set of cycles \mathcal{C} such that $\mathbf{z} = \sum_{v \in V} \mathbf{z}_v + \sum_{C \in \mathcal{C}} \mathbf{z}_C$.

We now use Lemma 23 to prove an important property of arc-minimal (r, k)-flow branchings. A rooted digraph X = (V + r, A) is called *triple-free* if it does not contain more than two arcs in the same direction between the same two vertices.

▶ Lemma 24. Let X = (V + r, A) be an arc-minimal (r, k)-flow branching with $|V| \ge 2k - 1$. Then X is triple-free.

Proof. Suppose that X contains three arcs a_1, a_2, a_3 whose tail is u and whose head is v for some $u, v \in V+r$. Further, let $\mathbf{z}: A \to \mathbb{Z}$ be an (r, k)-branching flow. By Lemma 23, we obtain that $\mathbf{z} = \sum_{v \in V} \mathbf{z}_v + \sum_{C \in \mathcal{C}} \mathbf{z}_C$ where \mathbf{z}_v is an v-path flow for all $v \in V$ and \mathbf{z}_C is a cycle flow for all $C \in \mathcal{C}$ for some set of cycles \mathcal{C} . Let $\mathbf{z}' = \sum_{v \in V} \mathbf{z}_v$. Observe that \mathbf{z}' is an (r, k)-branching flow. As all of the \mathbf{z}_v are path flows, we obtain $\mathbf{z}'(a_1) + \mathbf{z}'(a_2) + \mathbf{z}'(a_3) \leq |V| < 2(n-k)$. It follows that we can define a flow $\mathbf{z}'': A \to \mathbb{Z}$ such that $\mathbf{z}''(a_1) + \mathbf{z}''(a_2) = \mathbf{z}'(a_1) + \mathbf{z}'(a_2) + \mathbf{z}'(a_3)$, $\mathbf{z}''(a_3) = 0$ and $\mathbf{z}''(a) = \mathbf{z}'(a)$ for all $a \in A - \{a_1, a_2, a_3\}$. It is easy to see that \mathbf{z}'' is an (r, k)-branching flow, so $X - a_3$ is an (r, k)-flow branching, a contradiction to the minimality of X.

Given a rooted digraph D=(V+r,A) with $|V+r| \geq 2k$, a classic (r,k)-core is an (r,k)-flow branching X=(V'+r,A') that is a subdigraph of D with |V'|=2k-1. The XP algorithm of Theorem 9 is based on the following result, again reformulated using our terminology.

▶ Lemma 25 (Bang-Jensen, Havet, and Yeo [3]). Let D = (V + r, A) be a rooted digraph with $|V| \ge 2k - 1$. Then D contains two arc-disjoint spanning (r, k)-flow branchings if and only if D contains an extendable pair of arc-disjoint classic (r, k)-cores. Further, the two arc-disjoint spanning (r, k)-flow branchings can be constructed in polynomial time from the extendable pair of arc-disjoint classic (r, k)-cores.

2.4 Preliminaries on (r, k)-safe spanning trees

We first define an equivalent for extendability in undirected graphs. Given a rooted graph G = (V + r, E), a pair of subtrees (X_1, X_2) is called $completable^1$ if there are edge-disjoint spanning trees T_1, T_2 of G such that $E(X_i) \subseteq E(T_i)$. The following result that allows to test completability can be established using matroid theory as mentioned in [3].

▶ **Lemma 26** (Bang-Jensen, Havet, and Yeo [3]). Given a graph G = (V + r, E) and a pair of subtrees (X_1, X_2) , we can decide in polynomial time whether (X_1, X_2) is completable.

Given a rooted graph G = (V + r, E) with $|V + r| \ge 2k - 1$, a classic (r, k)-certificate is an (r, k)-safe subtree X = (V' + r, E') of G with |V'| = 2k - 2. The XP algorithm of Theorem 12 is based on the following result, again restated using our terminology.

▶ Lemma 27 (Bang-Jensen, Havet, and Yeo [3]). Let G = (V + r, E) be a rooted graph. Then G contains two edge-disjoint (r, k)-safe spanning trees if and only if G contains a completable pair of classic (r, k)-certificates. Further, given a completable pair of classic (r, k)-certificates, we can compute two edge-disjoint (r, k)-safe spanning trees in polynomial time.

2.5 Preliminaries on parameterized complexity

We refer the reader to [5] for basic background on parameterized complexity, and we recall here only some basic definitions used in this article. A parameterized problem is a decision problem whose instances are pairs $(x,k) \in \Sigma^* \times \mathbb{N}$, where k is called the parameter. A parameterized problem L is fixed-parameter tractable (FPT) if there exists an algorithm \mathcal{A} , a computable function f, and a constant c such that given an instance I = (x,k), \mathcal{A} (called an FPT algorithm) correctly decides whether $I \in L$ in time bounded by $f(k) \cdot |I|^c$. A parameterized problem L is slice-wise polynomial (XP) if there exists an algorithm \mathcal{A} and two computable functions f, g such that given an instance I = (x,k), \mathcal{A} (called an XP algorithm) correctly decides whether I is a positive instance of L in time bounded by $f(k) \cdot |I|^{g(k)}$.

A kernelization algorithm, or just $kernel^2$, for a parameterized problem L is an algorithm \mathcal{A} that, in polynomial time, generates from an instance I = (x, k) of L an equivalent instance I' = (x', k') of L such that $|x'| + k' \leq f(k)$, for some computable function $f : \mathbb{N} \to \mathbb{N}$. If f(k) is bounded from above by a polynomial of the parameter k, we say that L admits a polynomial kernel. It is well-known [5] that a parameterized problem is FPT if and only if it admits a (not necessarily polynomial) kernel.

3 An FPT algorithm for packing k-safe spanning arborescences

This section is concerned with proving Theorem 5. Given a rooted digraph D = (V + r, A) and a positive integer k, we say that a vertex $v \in V + r$ is large if $|N_D^+(v)| \ge 6k - 5$, and small otherwise. We let L_D (resp. S_D) be the set of vertices in V + r which are large (resp. small) in D.

We are now ready to introduce a new notion of (r, k)-kernels for k-safe spanning r-arborescences. A compact(r, k)-kernel is a subdigraph X = (V' + r, A') of D with $|V'| \le 2k - 2$ satisfying the following:

Note that we require a completable pair of trees to be edge-disjoint. This is in contrast with the fact that, in Subsection 2.2 and Subsection 2.3, we do not require the elements in an extendable pair to be arc-disjoint. We adopt this asymmetric choice for technical reasons arising from the proofs.

This notion should not be confused with (classic or compact) (r, k)-kernels used throughout this article.

- \blacksquare X is an r-arborescence,
- \blacksquare all vertices in $V' \cap L_D$ are sinks of X, and
- a k-safe r-arborescence Y can be obtained from X by adding a set V^* of 2k 2 |V'| new vertices and adding an arc from a vertex in $V' \cap L_D$ to v for all $v \in V^*$.

Observe that Y is not necessarily a subdigraph of D. The following result shows that compact (r, k)-kernels can be used in a similar way as classic (r, k)-kernels.

▶ Lemma 28. Let D = (V + r, A) be a 2-root-connected rooted digraph with $|V| \ge 2k - 2$. Then D contains two arc-disjoint k-safe spanning r-arborescences if and only if D contains an extendable pair of arc-disjoint compact (r, k)-kernels. Further, given an extendable pair of arc-disjoint compact (r, k)-kernels, we can find a pair of arc-disjoint k-safe spanning r-arborescences in polynomial time.

Proof. By Lemma 21, for the first part it suffices to prove that D contains a pair of arc-disjoint extendable compact (r, k)-kernels if and only if D contains a pair of arc-disjoint extendable classic (r, k)-kernels.

First let $(X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2))$ be an extendable pair of arc-disjoint classic (r, k)-kernels. Let $X_i' = (V_i' + r, A_i')$ be obtained from X_i by deleting $B_{X_i}^v - v$ for all $v \in (V_i + r) \cap L_D$. By construction, the X_i' are r-arborescences and all vertices in $(V_i' + r) \cap L_D$ are sinks in X_i' . Let Y_i be obtained from X_i' by adding the vertices in $V_i - V_i'$ and adding an arc from a vertex $v \in (V_i' + r) \cap L_D$ to a vertex $w \in V_i - V_i'$ whenever $w \in V(B_{X_i}^v) - v$. Observe that Y_i is an arborescence with $|V(Y_i)| = 2k - 1$. Further, note that $|V_i + r - V(B_{Y_i}^v)| = |V_i + r - V(B_{X_i}^v)| \ge k$ for all $v \in V_i'$ and $|V(B_{Y_i}^v)| = 0$ for all $v \in V_i - V_i'$. This yields that $|V_i + r - V(B_{Y_i}^v)| \ge |V_i + r - V(B_{X_i}^v)| \ge k$ for all $v \in V_i$ and so Y_i is a k-safe arborescence. By definition, we obtain that (X_1', X_2') is a pair of arc-disjoint compact (r, k)-kernels. Further, $D - A_i$ is a subdigraph of $D - A_i'$ that is root-connected, so $D - A_i'$ is root-connected as well. This yields that (X_1', X_2') is extendable.

Now let $(X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2))$ be an extendable pair of arc-disjoint compact (r, k)-kernels. By definition, there are k-safe arborescences Y_1, Y_2 such that Y_i is obtained from X_i by adding a set V_i^* of $2k - 2 - |V_i|$ new vertices and an arc from a vertex in $V_i \cap L_D$ to v for all $v \in V_i^*$. Let $(X'_1 = (V'_1 + r, A'_1), X'_2 = (V'_2 + r, A'_2))$ be a pair of subdigraphs of D that are vertex-maximal with the following properties:

- (i) X_i' is obtained from X_i by repeatedly adding another vertex $v \in V V_i$ and an arc of A that goes from a vertex in $V_i \cap L_D$ to v,
- (ii) $d_{X'}^+(v) \leq d_{Y_i}^+(v)$ for all $v \in V_i \cap L_D$,
- (iii) A'_1 and A'_2 are disjoint, and
- (iv) (X'_1, X'_2) is extendable.

Note that (X_1, X_2) satisfies conditions (i)-(iv), so (X'_1, X'_2) is well-defined. Also observe that if condition (ii) is satisfied with equality for all $v \in V_i \cap L_D$, then X'_i is isomorphic to Y_i , so X'_i is a k-safe arborescence and by definition also a classic (r, k)-kernel. If this is the case for both X'_1 and X'_2 , we are done by conditions (iii) and (iv).

We may therefore suppose by symmetry that there is a vertex $v \in V_1 \cap L_D$ with $d_{X_1}^+(v) < d_{Y_1}^+(v)$. For any $a = vz_a$ with $z_a \in N_D^+(v) - V_1'$, let $X_a = (V_1' \cup z_a, A_1' \cup a)$. By the maximality of X_1' , we obtain that X_a violates one of conditions (i)-(iv) for all $a \in \delta_D^+(v)$ with $z_a \in N_D^+(v) - V_1'$. By construction and the choice of v, X_a satisfies conditions (i) and (ii) for all $a \in \delta_D^+(v)$ with $z_a \in N_D^+(v) - V_1'$. If X_a violates (iii), then $z_a \in V_2'$. By (ii), we have $|V_2'| \leq |V_2 \cup V_2^*| = 2k - 2$ and so this is the case for at most 2k - 2 vertices

 $z_a \in N_D^+(v) - V_1'$. If X_a does not satisfy (iv), then a is critical in $D - A_1'$. As D is 2-root-connected and X_1' is an arborescence by construction, Lemma 16 implies that this is the case for at most $|A_1'| = |V_1'| \le |V_1 \cup V_1^*| = 2k - 2$ vertices $z_a \in N_D^+(v) - V_1'$. As $v \in L_D$ and $|V_1'| \le |V_1 \cup V_1^*| = 2k - 2$, we have $|N_D^+(v) - V_1'| \ge (6k - 5) - (2k - 2) > (2k - 2) + (2k - 2)$, so there is at least one vertex in $z \in N_D^+(v) - V_1'$ and an arc a = vz such that X_a does not violate any of conditions (i)-(iv), a contradiction.

Observe that the second part of the proof yields an algorithm for computing a pair of arc-disjoint extendable classic (r, k)-kernels. Indeed, every time we try to add an arc a = vz to X_i' , we test if (X_a, X_{3-i}') satisfies conditions (i)-(iv). Conditions (i)-(iii) can clearly be checked in polynomial time and, by Lemma 17, condition (iv) can also be checked in polynomial time. Never testing an arc that is parallel to one that we have tested already, after at most 4k-4 failed attempts, we manage to add a new vertex to V_i' . We repeat this procedure $|V_i^*| \leq 2k-2$ times. It follows that a pair of arc-disjoint extendable classic (r,k)-kernels can be computed in time $k^2 \cdot n^{O(1)} = n^{O(1)}$. By the second part of Lemma 21, we can then find the arc-disjoint k-safe spanning arborescences in D in time $n^{O(1)}$. Therefore, the overall running time of the algorithm is polynomial, as claimed.

We are now ready to proceed to the proof of Theorem 5.

Proof of Theorem 5. We may suppose that there are at most two parallel arcs from u to v for any $u, v \in V$. If |V| < 2k - 2, the problem can be solved by a brute force algorithm in time $2^{O(k^2)}$, by generating all pairs of subdigraphs of D and checking whether any of these pairs satisfies the required conditions. We may hence also suppose that $|V| \ge 2k - 2$.

We can first decide in time $n^{O(1)}$ if D is 2-root-connected. If it is not, the answer is negative, so we may suppose it is. Let $X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2)$ be two subdigraphs of D with $|V_1|, |V_2| \leq 2k - 2$. In order to test whether X_i is a compact (r, k)-kernel, we first verify if X_i is an r-arborescence such that all the vertices in $V_i \cap L_D$ are sinks in X_i . If this is the case, we add a set V_i^* of $2k - 2 - |V_i|$ new vertices to X_i . We then test all possibilities to add one arc from $(V_i + r) \cap L_D$ to v for all $v \in V_i^*$. As $|V_i + r| \leq 2k$ and $|V_i^*| \leq 2k$, there are at most $2k^{2k} = 2^{O(k \cdot \log k)}$ possibilities to check. For each of these possibilities, we can then check in polynomial time whether the obtained graph is a k-safe arborescence.

For each such pair X_1, X_2 , by the definition of classic (r, k)-kernels, we can therefore check in time $2^{O(k \cdot \log k)}$ whether both X_1 and X_2 are compact (r, k)-kernels. By Lemma 17, we can therefore decide in time $2^{O(k \cdot \log k)} + n^{O(1)}$ if (X_1, X_2) is an extendable pair of arc-disjoint compact (r, k)-kernels in D. By Lemma 28, it therefore suffices to prove that there are at most $2^{O(k^2 \cdot \log k)}$ possible candidates for the extendable pair of arc-disjoint compact (r, k)-kernels, and that these can be generated within this running time.

Let X=(V'+r,A') be a compact (r,k)-kernel in D. Observe that every vertex in V' can be reached from r by a directed path all of whose interior vertices are in S_D and whose length is at most k-1. As every vertex in S_D has at most 6k-6 out-neighbors, we obtain that the number of vertices that can be reached by such a path is at most $1+(6k-6)+(6k-6)^2+\ldots+(6k-6)^{k-1}\leq (6k)^k$. As V' contains at most 2k-2 vertices, there are at most $\binom{(6k)^k}{2k-2}\leq (6k)^{2k^2}$ possibilities to choose V'.

Now suppose that we have chosen V' of size 2k-2. As there are at most two arcs in the same direction between any two vertices, there are at most $4\binom{|V'+r|}{2} \le 16k^2$ arcs that have their head and tail in V'+r. As |A'|=2k-2, there are at most $\binom{16k^2}{2k-2} \le (16k^2)^{2k}$ possibilities to choose A'. It follows that there are at most $(6k)^{2k^2} \cdot (16k^2)^{2k}$ possibilities to choose a compact (r,k)-kernel X. As these can be computed by a brute force method, the

algorithm can finish after checking less than $f(k) = \binom{(6k)^{2k^2} \cdot (16k^2)^{2k}}{2} = 2^{O(k^2 \cdot \log k)}$ candidates for the extendable pair of arc-disjoint compact (r, k)-kernels.

If no extendable pair of arc-disjoint compact (r,k)-kernels exists, by Lemma 28, D does not contain two arc-disjoint k-safe spanning arborescences. On the other hand, once we have found an extendable pair of arc-disjoint compact (r,k)-kernels, we can compute the two arc-disjoint k-safe spanning r-arborescences in polynomial time by the second part of Lemma 28. The overall running time of the obtained algorithm is $2^{O(k^2 \cdot \log k)} \cdot n^{O(1)}$.

4 An FPT algorithm for packing spanning (r, k)-flow branchings

This section is concerned with proving Theorem 10. Slightly modifying the terminology introduced in Section 3, given a 2-root-connected digraph D = (V + r, A) and a positive integer k, we say that a vertex v is large if $|N^+(v)| \ge 20k^2 + 1$, and small otherwise. Again, we let L_D (resp. S_D) be the set of vertices in V + r which are large (resp. small) in D.

We are now ready to introduce a new notion of (r, k)-cores for spanning (r, k)-flow branchings. A *compact* (r, k)-core is a subdigraph X = (V' + r, A') of D with $|V'| \le 2k - 1$ satisfying the following:

- \blacksquare all vertices in $V' \cap L_D$ are sinks in X and
- an (r,k)-flow branching Y can be obtained from X by adding a set V^* of 2k-1-|V'| new vertices and adding an arc from a vertex in $V' \cap L_D$ to v for all $v \in V^*$.

Observe that Y is not necessarily a subdigraph of D. The following result, which is similar to Lemma 28, shows that compact (r, k)-cores can be used in a similar way as classic (r, k)-cores.

▶ Lemma 29. Let D = (V + r, A) be a 2-root-connected rooted digraph with $|V| \ge 2k - 1$. Then D has two arc-disjoint spanning (r, k)-flow branchings if and only if D contains an extendable pair of arc-disjoint compact (r, k)-cores. Further, given an extendable pair of triple-free arc-disjoint compact (r, k)-cores, we can find a pair of arc-disjoint spanning (r, k)-flow branchings in polynomial time.

Proof. By Lemma 25, for the first part it suffices to prove that D contains an extendable pair of arc-disjoint compact (r, k)-cores if and only if D contains an extendable pair of arc-disjoint classic (r, k)-cores.

First let $(X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2))$ be an extendable pair of arc-disjoint classic (r, k)-cores. Let $X_i' = (V_i' + r, A_i')$ be obtained from X_i by first deleting all arcs in A_i whose tail is a large vertex and then restricting to the subdigraph that is root-connected from r. As $A_i' \subseteq A_i$, (X_1', X_2') is extendable. It remains to show that X_1' and X_2' are compact (r, k)-cores. By construction, all vertices in $V_i' \cap L_D$ are sinks in X_i' . As X_i is an (r, k)-flow branching, there is an (r, k)-branching flow $\mathbf{z} : A_i \to \mathbb{Z}$ in X_i . Create Y_i from X_i by attaching $\mathbf{z}(\delta_{X_i}^+(v))$ arcs directed away from v to every vertex $v \in V_i' \cap L_D$. Assigning $\mathbf{z}'(a) = 1$ for all arcs leaving a large vertex in Y_i and $\mathbf{z}'(a) = \mathbf{z}(a)$ for all remaining arcs, we obtain that \mathbf{z}' is an (r, k)-branching flow in Y_i , so Y_i is an (r, k)-flow branching. Furthermore, we have $|V(Y_i) - r| = \mathbf{z}'(\delta_{Y_i}^+(r)) = \mathbf{z}(\delta_{X_i}^+(r)) = |V_i| = 2k - 1$. It follows by definition that X_i' is a compact (r, k)-core.

Now let $(X_1=(V_1+r,A_1),X_2=(V_2+r,A_2))$ be an extendable pair of arc-disjoint compact (r,k)-cores. Possibly deleting arcs, we may suppose by Lemma 24 and as $|V| \geq 2k-1$ that X_1 and X_2 are triple-free. By definition, there are (r,k)-flow branchings Y_1,Y_2 such that Y_i is obtained from X_i by adding a set V_i^* of $2k-1-|V_i|$ new vertices and adding an

arc from a vertex in $V_i \cap L_D$ to v for all $v \in V_i^*$. Let $X_1' = (V_1' + r, A_1')$ and $X_2' = (V_2' + r, A_2')$ be subdigraphs of D that are vertex-maximal with the following properties:

- (i) X_i' is obtained from X_i by repeatedly adding another vertex in $v \in V V_i$ and an arc of A that goes from a vertex in $V_i \cap L_D$ to v,
- (ii) $d_{X'}^+(v) \le d_{Y_i}^+(v)$ for all $v \in V_i \cap L_D$,
- (iii) A'_1 and A'_2 are disjoint, and
- (iv) (X'_1, X'_2) is extendable.

Note that (X_1, X_2) satisfies conditions (i)-(iv), so (X'_1, X'_2) is well-defined. Further, observe that if condition (ii) is satisfied with equality for all $v \in V_i \cap L_D$, then X'_i is isomorphic to Y_i , so X'_i is an (r, k)-flow branching, thus by definition also a classic (r, k)-core. If this is the case for both X'_1 and X'_2 , we are done by conditions (iii) and (iv).

We may therefore suppose by symmetry that there is some $v \in V_1 \cap L_D$ with $d_{X_1'}^+(v) < d_{Y_1}^+(v)$. For any $a = vz_a$ with $z_a \in N_D^+(v) - V_1'$, let $X_a = (V_1' \cup z_a, A_1' \cup a)$. By the maximality of X_1' , X_a violates one of conditions (i)-(iv) for all $a \in \delta_D^+(v)$ with $z_a \in N_D^+(v) - V_1'$. By construction and the choice of v, X_a satisfies conditions (i) and (ii) for all $a \in \delta_D^+(v)$ with $z_a \in N_D^+(v) - V_1'$. If X_a violates condition (iii), then $z_a \in V_2'$. As $|V_2'| \leq |V(Y_2)| = 2k$, this is the case for at most 2k vertices in $N_D^+(v) - V_1'$. If X_a does not satisfy (iv), then a is critical in $D - A_1'$. As X_1 is triple-free and $|V_1'| \leq 2k$ and by construction, we obtain that $|A_1'| \leq 4\binom{|V_1'|}{2} \leq 16k^2$. Now Lemma 16 implies that this is the case for at most $16k^2$ vertices in $N_D^+(v) - V_1'$. As $v \in L_D$ and $|V_1'| \leq |V_1 \cup V_1^*| = 2k - 1$, we have $|N_D^+(v) - V_1'| \geq (20k^2 + 1) - (2k - 1) > 2k + 16k^2$, so there is at least one vertex $z \in N_D^+(v) - V_1'$ and an arc a = vz such that X_a does not violate any of conditions (i)-(iv), a contradiction.

Observe that the second part of the proof yields an algorithm for computing a pair of arc-disjoint extendable classic (r,k)-cores. Indeed, every time we try to add an arc a=vz to X_i' , we test if (X_a, X_{3-i}') satisfies conditions (i)-(iv). Conditions (i)-(iii) can clearly be checked in polynomial time and, by Lemma 17, condition (iv) can also be checked in polynomial time. Never checking an arc a which is parallel to an arc we have already checked, after at most $20k^2$ failed attempts, we manage to add a new vertex to V_i' . We repeat this procedure at most $|V_i^*| \leq 2k$ times. It follows that a pair of arc-disjoint extendable classic (r,k)-cores can be computed in time $40k^3 \cdot n^{O(1)} = n^{O(1)}$. By the second part of Lemma 25, we can then find the arc-disjoint spanning (r,k)-flow branchings in D in polynomial time. The overall running time of the algorithm is polynomial, as claimed.

We are now ready to proceed to the proof of Theorem 10.

Proof of Theorem 10. First consider the case that |V| < 2k - 1. Observe that any arcminimal spanning (r,k)-flow branching has at most |V| parallel arcs between any two vertices. It follows that, for any two vertices u,v, at most $\gamma:=4k^2$ different distributions of the arcs between u and v among the two candidates for the spanning (r,k)-flow branchings have to be considered, including taking none of these arcs. Since there are $\mu:=\binom{|V|}{2}=O(k^2)$ pairs of vertices, the total number of choices for these distributions is $\gamma^\mu=2^{O(k^2\cdot\log k)}$. The problem can therefore be solved by a brute force algorithm in time $2^{O(k^2\cdot\log k)}$, by generating all $2^{O(k^2\cdot\log k)}$ pairs of candidate subdigraphs of D and checking whether any of these pairs satisfies the required conditions. We may hence suppose that $|V| \geq 2k-1$.

By Lemma 24, we may also suppose that there are at most four parallel arcs between any two vertices in D. We can first decide in polynomial time if D is 2-root-connected. If it is not, the answer is negative, so we may suppose it is. Let $X_1 = (V_1 + r, A_1), X_2 = (V_2 + r, A_2)$ be two subdigraphs of D with $|V_1|, |V_2| \leq 2k - 1$. In order to test whether X_i is a compact

(r,k)-core, we first test if all vertices in $V_i \cap L_D$ are sinks in X_i . We then add a set V_i^* of $2k-1-|V_i|$ new vertices to X_i . We then test all possibilities to add one arc from $(V_i+r)\cap L_D$ to v for all $v\in V_i^*$. As $|V_i|,|V_i^*|\leq 2k$, there are at most $2k^{2k}=2^{O(k\cdot\log k)}$ possibilities to check. By Lemma 22, we can check in time polynomial in k whether each of the resulting graphs is an (r,k)-flow branching. Thus, we can check in time $2^{O(k\cdot\log k)}$ whether both X_1 and X_2 are compact (r,k)-cores. By Lemma 17, we can therefore decide in time $2^{O(k\cdot\log k)}+n^{O(1)}$ if (X_1,X_2) is an extendable pair of arc-disjoint compact (r,k)-cores in D. By Lemma 29, it therefore suffices to prove that there are at most $2^{O(k^2\cdot\log k)}$ possible candidates for the extendable pair of arc-disjoint compact (r,k)-cores, and that these can be generated within the same running time.

Let X=(V'+r,A') be a compact (r,k)-core in D. Observe that every vertex in V' can be reached from r by a directed path all of whose internal vertices are in S_D and whose length is at most 2k-1. As every vertex in S_D has at most $20k^2$ out-neighbors, we obtain that the number of vertices that can be reached by such a path is at most $1+20k^2+(20k^2)^2+\ldots+(20k^2)^{2k-1}\leq (20k^2)^{2k}$. As V' contains at most 2k-1 vertices, there are at most $\binom{(20k^2)^{2k}}{(2k-1)}\leq (20k^2)^{4k^2}$ possibilities to choose V'. Now suppose that we have chosen V' of size at most 2k-1. As there are at most four arcs in the same direction between any two vertices, there are at most 2k-1. As there are at most 2k-1 arcs that have their head and tail in V'+r. As all arcs of A' have both ends in V'+r, there are at most 2^{32k^2} possibilities to choose A'. It follows that there are at most 2^{32k^2} possibilities to choose a compact 2^{32k^2} possibilities to choose 2^{32k^2} possibilities to choos

If no such extendable pair of arc-disjoint compact (r,k)-cores exists, by Lemma 29, D does not contain two arc-disjoint spanning (r,k)-flow branchings. On the other hand, if we find an extendable pair of arc-disjoint compact extendable (r,k)-cores, we also find such a pair (X_1, X_2) where X_1 and X_2 are arc-minimal, so by Lemma 24 triple-free. By the second part of Lemma 29, we can compute the two arc-disjoint spanning (r,k)-flow branchings in polynomial time. The overall running time of the obtained algorithm is $2^{O(k^2 \cdot \log k)} \cdot n^{O(1)}$.

5 An FPT algorithm for packing (r, k)-safe spanning trees

This section is concerned with proving Theorem 13. Again, slightly modifying the terminology introduced in Section 3 and reused in Section 4, given a rooted graph G = (V + r, E) and a positive integer k, we say that a vertex $v \in V + r$ is large if $|N_G(v)| \ge 8k - 7$, and small otherwise. And again, we let L_G (resp. (S_G)) be the set of vertices in V which are large (resp. small) in G.

We are now ready to introduce a new notion of certificates for (r, k)-safe spanning trees. A compact (r, k)-certificate is a subgraph X = (V' + r, E') of G with $|V'| \le 2k - 2$ satisfying the following:

- \blacksquare X is a tree,
- \blacksquare all vertices in $V' \cap L_G$ are leaves of X, and
- an (r,k)-safe spanning tree Y can be obtained from X by adding a set V^* of 2k-2-|V'| new vertices and adding an edge from a vertex in $V' \cap L_G$ to v for all $v \in V^*$.

Observe that Y is not necessarily a subgraph of G. The following result, which is similar to Lemma 28 and Lemma 29, shows that compact certificates can be used in a similar way as classic certificates.

▶ Lemma 30. Let G = (V + r, E) be a rooted graph with $|V| \ge 2k - 2$. Then G has two edge-disjoint (r, k)-safe spanning trees if and only if G contains a completable pair of compact (r, k)-certificates. Further, given a completable pair of compact (r, k)-certificates, we can find a pair of edge-disjoint (r, k)-safe spanning trees in polynomial time.

Proof. By Lemma 27, for the first part it suffices to prove that G contains a completable pair of compact (r,k)-certificates if and only if G contains a pair of completable classic (r,k)-certificates.

First let $(X_1 = (V_1 + r, E_1), X_2 = (V_2 + r, E_2))$ be a completable pair of classic (r, k)-certificates. Let $X_i' = (V_i' + r, E_i')$ be obtained from X_i by deleting $C_{X_i}^v$ for all $v \in (V_i + r) \cap L_G$. By construction, the X_i' are trees and all vertices in $(V_i' + r) \cap L_G$ are leaves in X_i' . Let Y_i be obtained from X_i' by adding the vertices in $V_i - V_i'$ and adding an edge from a vertex $v \in (V_i' + r) \cap L_G$ to a vertex $w \in V_i - V_i'$ whenever $w \in V(C_{X_i}^v)$. Observe that Y_i is a tree with $|V(Y_i)| = 2k - 1$. Further, we have $|V_i - V(C_{Y_i}^v)| = |V_i - V(C_{X_i}^v)| \ge k$ for all $v \in V_i'$ and $|V(C_{Y_i}^v)| = 0$ for all $v \in V_i - V_i'$. This yields that $|V_i - V(C_{Y_i}^v)| \ge |V_i - V(C_{X_i}^v)| \ge k$ for all $v \in V_i$ and so Y_i is an (r, k)-safe tree. By definition, we obtain that (X_1', X_2') is completable, we obtain that (X_1', X_2') is completable.

Now let $(X_1 = (V_1 + r, E_1), X_2 = (V_2 + r, E_2))$ be a completable pair of compact (r, k)-certificates. By definition, there are (r, k)-safe trees Y_1, Y_2 such that Y_i is obtained from X_i by adding a set V_i^* of $2k - 2 - |V_i|$ new vertices and an edge from a vertex in $V_i \cap L_G$ to v for all $v \in V_i^*$. Let $(X_1' = (V_1' + r, E_1'), X_2' = (V_2' + r, E_2'))$ be a pair of subgraphs of G that are vertex-maximal with the following properties:

- (i) X_i' is obtained from X_i by repeatedly adding another vertex $v \in V V_i$ and an edge of E that goes from a vertex in $V_i \cap L_G$ to v,
- (ii) $d_{X_i}(v) \leq d_{Y_i}(v)$ for all $v \in V_i \cap L_G$, and
- (iii) (X'_1, X'_2) is completable.

Note that (X_1, X_2) satisfies conditions (i)-(iii), so (X'_1, X'_2) is well-defined. Observe that if condition (ii) is satisfied with equality for all $v \in V_i \cap L_G$, then X'_i is isomorphic to Y_i , so X'_i is an (r, k)-safe spanning tree and by definition also a classic (r, k)-certificate. If this is the case for both X'_1 and X'_2 , we are done by condition (iii).

We may therefore suppose by symmetry that there is a vertex $v \in V_1 \cap L_G$ with $d_{X_1'}(v) < d_{Y_1}(v)$. For any $e = vz_e \in \delta_G(v)$ with $z_e \in N_G(v) - V_1'$, let $X_e = (V_1' \cup z_e, E_1' \cup e)$. By the maximality of X_1' , we obtain that X_e violates one of conditions (i)-(iii) for all $e = vz_e \in \delta_G(v)$ with $z_e \in N_G(v) - V_1'$. By construction and the choice of v, X_e satisfies conditions (i) and (ii) for all $e = vz_e \in \delta_G(v)$ with $z_e \in N_G(v) - V_1'$. It follows that (X_e, X_2') violates condition (iii) for all $e = vz_e \in \delta_G(v)$ with $z_e \in N_G(v) - V_1'$. As (X_1', X_2') is completable, there are two disjoint spanning trees T_1, T_2 of G such that $E_i' \subseteq E(T_i)$ for i = 1, 2.

▶ Claim 31. There is no $e = vz_e \in \delta_G(v) - (E(T_1) \cup E(T_2))$ with $z_e \in N_G(v) - V_1'$ such that (X_e, X_2') violates condition (iii).

Proof. Suppose otherwise. By Lemma 18, there is an edge $f \in E(T_1)$ incident to z_e such that $T_1' = T_1 - f + e$ is a spanning tree of G. As $z_e \notin V_1'$, we obtain that $f \notin E_1'$, yielding $E(X_e) \subseteq E(T_1')$. As T_1' and T_2 are edge-disjoint, we obtain that (X_z, X_2') is completable, a contradiction.

▶ Claim 32. There are at most 6k - 6 vertices $z \in N_G(v) - V_1'$ such that (X_e, X_2') violates condition (iii) for some $e = vz \in E(T_2)$.

Proof. Suppose otherwise. As $|V_2'| \leq |V(Y_2)| - 1 = 2k - 2$, we obtain that there are at least 4k - 3 vertices $z \in N_G(v) - V_1'$ such that (X_e, X_2') violates condition (iii) for some $e = vz \in E(T_2) - E_2$. Let $\sigma : E(T_2) \to E(T_1)$ be a tree-mapping function from T_2 to T_1 . By Lemma 20 and since $|E_1'| \leq 2k - 2$, there is some $z \in N_G(v) - V_1'$ and an edge $e = vz \in E(T_2) - E_2$ such that $\sigma(e) \in E(T_1) - E_1$. By definition of tree-mapping functions, $T_1' = T_1 - \sigma(e) + e$ and $T_2' = T_2 - e + \sigma(e)$ are edge-disjoint spanning trees of G. As $E(X_e) \subseteq E(T_1')$ and $E(X_2) \subseteq E(T_2')$, we obtain that (X_e, X_2') is completable, a contradiction.

As $v \in L_G$ and $|V_1'| \le |V_1 \cup V_1^*| = 2k - 2$, we have $|N_G(v) - V_1'| \ge (8k - 7) - (2k - 2) > 6k - 6$. It now follows from Claim 31 and Claim 32 that there is at least one vertex in $z \in N_G(v) - V_1'$ and an edge e = vz such that X_e does not violate any of conditions (i)-(iii), a contradiction.

Observe that the second part of the proof yields an algorithm for computing a completable pair of classic (r,k)-certificates from (X_1,X_2) . Every time we try to add an edge e to X_i' , we test if (X_e,X_{3-i}') satisfies conditions (i)-(iii). Conditions (i)-(ii) can clearly be checked in polynomial time and, by Lemma 26, condition (iii) can also be checked in polynomial time. Never checking an edge that is parallel to one we have already checked, after at most 6k-6 failed attempts, we manage to add a new vertex to V_i' . We repeat this procedure $|V_i^*| \leq 2k-2$ times. It follows that a completable pair of classic (r,k)-certificates can be computed in time $k^2 \cdot n^{O(1)} = n^{O(1)}$. By the second part of Lemma 27, we can then find two edge-disjoint (r,k)-safe spanning trees in G in polynomial time. The overall running time of the algorithm is polynomial, as claimed.

We are now ready to proceed to the proof of Theorem 13.

Proof of Theorem 13. We may suppose that there are at most two parallel edges from u to v for any $u, v \in V + r$. If |V| < 2k - 2, the problem can be solved by a brute force algorithm in time $2^{O(k^2)}$, by generating all pairs of subgraphs of G and checking whether any of these pairs satisfies the required conditions. We may hence also suppose that $|V| \ge 2k - 2$.

Let $X_1=(V_1+r,E_1), X_2=(V_2+r,E_2)$ be two subgraphs of G. In order to test whether X_i is a compact (r,k)-certificate, we first check whether X_i is a tree such that all the vertices in $X_i\cap L_G$ are leaves of X. If this is the case, we add a set V_i^* of $2k-2-|V_i|$ new vertices to X_i . We then test all possibilities to add one edge from $(V_i+r)\cap L_G$ to v for all $v\in V_i^*$. As $|V_i+r|\leq 2k$ and $|V_i^*|\leq 2k$, there are at most $2k^{2k}=2^{O(k\cdot\log k)}$ possibilities to check. For each of them, we can check in time polynomial in k if the obtained graph is an (r,k)-safe spanning tree. By the definition of compact (r,k)-certificates, we can therefore check in time $2^{O(k\cdot\log k)}$ whether both X_1 and X_2 are compact (r,k)-certificates. By Lemma 26, we can therefore decide in time $2^{O(k\cdot\log k)}+n^{O(1)}$ if (X_1,X_2) is a completable pair of compact (r,k)-certificates in G. By Lemma 30, it therefore suffices to prove that there are at most $2^{O(k^2\cdot\log k)}$ possible candidates for the completable pair of compact (r,k)-certificates, and that they can be generated within the same running time.

Let X = (V' + r, E') be a compact (r, k)-certificate in G. Observe that every vertex in V' can be reached from r by a path all of whose interior vertices are in S_G and whose length is at most k-1. As every vertex in S_G has at most 8k-8 neighbors, we obtain that the number of vertices that can be reached by such a path is at most $1+(8k-8)+(8k-8)^2+\ldots+(8k-8)^{k-1} \le (8k)^k$. As V' contains at most 2k-2 vertices, there are at most $\binom{(8k)^k}{2k-2} \le (8k)^{2k^2}$ possibilities to choose V'.

Now suppose that we have chosen V' of size 2k-2. Observe that there are at most $2\binom{|V'+r|}{2} \le 8k^2$ edges that have both ends in V'+r. As |E'|=2k-2, there are at most

 $\binom{8k^2}{2k-2} \leq (8k^2)^{2k}$ possibilities to choose A'. It follows that there are at most $(8k)^{2k^2} \cdot (8k^2)^{2k}$ possibilities to choose a compact (r,k)-certificate X. As these can be computed by a brute force method, the algorithm can finish after checking less than $f(k) = \binom{(8k)^{2k^2} \cdot (8k^2)^{2k}}{2} = 2^{O(k^2 \cdot \log k)}$ candidates for the pair of compact (r,k)-certificates.

If no completable pair of compact (r, k)-certificates exists, by Lemma 27, G does not contain two edge-disjoint (r, k)-safe spanning trees. On the other hand, once we have found a pair of completable compact (r, k)-certificates, we can compute in polynomial time the two edge-disjoint (r, k)-safe spanning trees by the second part of Lemma 27. The overall running time of the obtained algorithm is $2^{O(k^2 \cdot \log k)} \cdot n^{O(1)}$.

6 A hardness result for packing (r, k)-safe spanning trees

In this section we prove Theorem 14. It is well-known that the 3-SAT problem is NP-complete. Further, we will need the following lemma derived from the ETH using the so-called Sparsification Lemma [10].

▶ Lemma 33 (Impagliazzo et al. [10]). Assuming the ETH, there is an $\varepsilon > 0$ such that there is no algorithm for solving a 3-SAT formula with ℓ variables and m clauses in time $2^{\varepsilon m} \cdot (\ell + m)^{O(1)}$.

The proof of Theorem 14 given below is strongly inspired from the reduction given in [3, Theorem 5.2], but we provide it here entirely for the sake of completeness.

Proof of Theorem 14. Observe that, given a rooted graph G = (V + r, E) and two positive integers p and k, G contains an (r, k)-safe spanning tree if and only if the graph that is obtained from G by replacing each of its edges by p parallel copies of itself contains p edge-disjoint (r, k)-safe spanning trees. Hence, it suffices to prove the statement for p = 1. Let ϕ be an instance of 3-SAT, with variables x_1, x_2, \ldots, x_ℓ and clauses C_1, C_2, \ldots, C_m . Adding a variable that is not contained in any clause if necessary, we can assume that ℓ is even. We construct a simple rooted graph G = (V + r, E) as follows; see Figure 1 for an illustration. For $i = 1, \ldots, \ell$ let V_i be an independent set containing two vertices v_i and \bar{v}_i , and let r and t be two extra vertices. Add all possible edges between r and V_1 , between V_i and V_{i+1} for $i = 1, \ldots, \ell - 1$, and between V_ℓ and t. Next, add t vertices t is a literal of t in t




Figure 1 The rooted graph G in the proof of Theorem 14 with $C_1 = \bar{x_1} \vee x_2 \vee \bar{x_4}$.

We now prove that ϕ is satisfiable if and only if G admits an (r, k)-safe spanning tree. First assume that G contains an (r, k)-safe spanning tree T. Observe that, by construction and the definition of (r, k)-safe spanning tree, T - r has exactly two components whose vertex sets we denote by T_1 and T_2 . We may assume that T_1 contains t. Then $G[T_1]$ contains a path P from V_1 to t and T_1 contains Q. As a shortest path from V_1 to t contains $\ell + 1$ vertices, we have $|T_1| \geq q + \ell + 1$ and $n - |T_1| \leq k$. As T is (r, k)-safe, we obtain that $|T_1| = q + \ell + 1$ and T_1 consists exactly of the vertex set of the path P, which contains exactly $\ell + 1$ vertices, and the whole set Q. In particular, P intersects each V_i in exactly one vertex. Now for every variable x_i of \mathcal{F} , we set x_i to true if P contains \bar{v}_i and to false if P contains v_i . For each clause C_j , there must be a path from the corresponding vertex c_j to V_1 in T_2 . Then one of the vertices corresponding to a literal of C_j must not be contained in P and so, this literal is set to true and C_i is satisfied by it. It follows that the constructed assignment satisfies ϕ .

Conversely, assume that ϕ admits a truth assignment. Let P be the path defined so that, for every $i=1,\ldots,\ell$, it contains v_i if x_i is set to false and $\bar{v_i}$ if x_i is set to true. Further, let $T_1=V(P)\cup t\cup Q$. As ϕ is satisfied, for every $1\leq j\leq m$, there exists a path from c_j to $V_1\cup\cdots\cup V_\ell$ in $G-(T_1\cup r)$. It follows that $G-(T_1\cup r)$ is connected and we select a spanning tree of it. The union of this spanning tree with $G[T_1]$, r, and the edges incident to r is a spanning tree of G. In order to see that this spanning tree is (r,k)-safe, it suffices to observe that, as $q>k-\ell-1$, we have $|T_1|=q+\ell+1>k$ and $|V-T_1|=q+k+1+\ell-(q+\ell+1)=k$.

If we fix q = k, then the size of G is bounded by a polynomial in ℓ and m. Thus, the above reduction implies that, as 3-SAT is NP-complete, given a rooted graph G and an integer k, deciding whether G admits an (r, k)-safe spanning tree is NP-complete.

Let now ε be a positive constant and assume that k is an integer function satisfying $(\log(n))^{2+\varepsilon} \leq k(n) \leq \frac{n}{2}$ for all n > 0. Furthermore, suppose that there exists a constant C^* such that for all $c \geq C^*$ there exists an n such that k(n) = c. Finally, for the sake of a contradiction assume that there exists a polynomial-time algorithm \mathbf{A} , running in time $O(n^{c_0})$ for some $c_0 > 0$, for deciding if a given rooted graph G = (V + r, E) on n vertices contains an (r, k(n))-safe spanning tree. Then let ϕ be a 3-SAT formula with ℓ variables and m clauses. We may assume that ℓ and m are large enough so that $1 + \ell + 3\ell m/2 + m \geq C^*$. Adding trivial clauses if necessary, we may also assume that $\ell \leq m$. By hypothesis, there exists n such that $k(n) = 1 + \ell + 3\ell m/2 + m$. So, in the above reduction, we choose q to be $n - (k(n) + \ell + 1)$, in order to have $n = q + k(n) + \ell + 1$. Then, using algorithm \mathbf{A} , one could decide if ϕ is satisfiable in time $O(n^{c_0}) = O(2^{c_0 \cdot \log n}) = O(2^{c_0 \cdot k(n)^{1/(2+\varepsilon)}})$, where we have used the hypothesis that $k(n) \geq (\log(n))^{2+\varepsilon}$. Moreover, in the previous construction, we have $k(n) = 1 + \ell + 3\ell m/2 + m \leq 3m + 3m^2/2 \leq 3m^2$. So we could decide whether ϕ is satisfiable in time $O(2^{c_0 \cdot (3m^2)^{1/(2+\varepsilon)}}) = O(2^{c_0 \cdot m^{\varepsilon'}})$ with $\varepsilon' = 2/(2+\varepsilon) < 1$, a contradiction to Lemma 33 assuming the ETH.

7 Conclusion

We considered three problems on finding certain disjoint substructures in graphs and digraphs. While in our proofs we restrict to finding two of these substructures for the sake of simplicity, our results can be generalized to allow for finding an arbitrary number of them using the same proof techniques. More concretely, the following results can be established using the techniques of this article. As in [3], we omit the proofs of these generalized statements.

▶ Theorem 34. Given a rooted digraph D = (V + r, A) and an integer $p \ge 2$, deciding whether D contains p arc-disjoint k-safe spanning r-arborescences is FPT with parameter k. More precisely, the problem can be solved in time $2^{O(p \cdot k^2 \cdot \log k)} \cdot n^c$, where c is a constant

depending on p. Further, if they exist, the p arc-disjoint k-safe spanning r-arborescences can be computed within the same running time.

- ▶ **Theorem 35.** Given a rooted digraph D = (V + r, A) and an integer $p \ge 2$, deciding whether D contains p arc-disjoint (r,k)-flow branchings is FPT with parameter k. More precisely, the problem can be solved in time $2^{O(p \cdot k^2 \cdot \log k)} \cdot n^c$, where c is a constant depending on p. Further, if they exist, the p arc-disjoint (r,k)-flow branchings can be computed within the same running time.
- ▶ Theorem 36. Given a rooted graph G = (V + r, E) and an integer $p \ge 2$, deciding whether G contains p arc-disjoint (r,k)-safe spanning trees is FPT with parameter k. More precisely, the problem can be solved in time $2^{O(p \cdot k^2 \cdot \log k)} \cdot n^c$, where c is a constant depending on p. Further, if they exist, the p edge-disjoint (r,k)-safe spanning trees can be computed within the same running time.

It is natural to ask whether the dependency on k of our FPT algorithms can be improved. In the case of k-safe spanning r-arborescences (cf. Theorem 5), we can derive a lower bound from Theorem 3. Indeed, a corollary of Theorem 3 is that, assuming the ETH, for any two constants $\varepsilon>0$ and c>0, deciding whether a rooted digraph contains two arc-disjoint k-safe spanning arborescences cannot be solved in time $2^{c \cdot k^{1-\varepsilon}} \cdot n^{O(1)}$. To see this, note that if such an algorithm existed, letting $k(n) := (\log(n))^{1+\varepsilon}$ we would obtain an algorithm in time $2^{c \cdot (\log(n))^{(1+\varepsilon)(1-\varepsilon)}} \cdot n^{O(1)} = n^{O(1)}$, contradicting Theorem 3. In other words, assuming the ETH, the problem cannot be solved in time $2^{O(k^{1-\varepsilon)}} \cdot n^{O(1)}$ for any $\varepsilon>0$.

Similarly to Theorem 3, Theorem 8 implies a lower bound for packing (r,k)-flow branchings (cf. Theorem 5): assuming the ETH, deciding whether a rooted digraph contains two arc-disjoint (r,k)-flow branchings cannot be solved in time $2^{O(k^{1-\varepsilon})} \cdot n^{O(1)}$ for any $\varepsilon > 0$. Also, concerning (r,k)-safe spanning trees (cf. Theorem 13), a consequence of Theorem 14 is that, assuming the ETH, for every $p \geq 1$, deciding whether a rooted graph contains p edge-disjoint (r,k)-safe spanning trees cannot be solved in time $2^{O(k^{1/2-\varepsilon})} \cdot n^{O(1)}$ for any $\varepsilon > 0$.

There is still a significant gap between the above lower bounds, which are $2^{O(k^{1-\varepsilon})}$ or $2^{O(k^{1/2-\varepsilon})}$, and the function $2^{O(k^2 \cdot \log k)}$ in our FPT algorithms.

We did not focus on optimizing the polynomial factors in n of our algorithms, and we leave it for further research. Further, we leave as an open question whether any of the considered problems admits a polynomial kernel parameterized by k. Finally, it would be interesting to find a theorem on packing k-safe mixed arborescences in mixed graphs, hence generalizing both Theorem 34 and Theorem 36.

References

- 1 Jørgen Bang-Jensen and Stéphane Bessy. (arc-)disjoint flows in networks. *Theoretical Computer Science*, 526:28–40, 2014. doi:10.1016/j.tcs.2014.01.011.
- 2 Jørgen Bang-Jensen and Gregory Gutin. *Digraphs: Theory, Algorithms and Applications*. London, 2nd edition, 2009. URL: https://dblp.org/rec/books/daglib/0022205.bib.
- 3 Jørgen Bang-Jensen, Frédéric Havet, and Anders Yeo. The complexity of finding arc-disjoint branching flows. Discrete Applied Mathematics, 209:16–26, 2016. doi:10.1016/j.dam.2015. 10.012
- 4 Jørgen Bang-Jensen and Anders Yeo. Balanced branchings in digraphs. *Theoretical Computer Science*, 595:107–119, 2015. doi:10.1016/j.tcs.2015.06.026.
- 5 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015. doi:10.1007/978-3-319-21275-3.

- Jack Edmonds. Edge-disjoint branchings. In B. Rustin, editor, Combinatorial Algorithms, pages 91-96, New York, 1973. Academic Press.
- Jack Edmonds. Some well-solved problems in combinatorial optimization. In B. Roy, editor, Combinatorial Programming: Methods and Applications (Proceedings of the NATO Advanced Study Institute, Versailles, 1974), Reidel, Dordrecht, 1975. Academic Press. doi:10.1007/ 978-94-011-7557-9_15.
- András Frank. Connections in Combinatorial Optimization. 2011. URL: https://global.oup. $\verb|com/academic/product/connections-in-combinatorial-optimization-9780199205271|.$
- Russell Impagliazzo and Ramamohan Paturi. On the complexity of k-SAT. Journal of Computer and System Sciences, 62(2):367-375, 2001. doi:10.1006/jcss.2000.1727.
- 10 Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? Journal of Computer and System Sciences, 63(4):512-530, 2001. doi:10.1006/jcss.2001.1774.
- László Lovász. Connectivity in digraphs. Journal of Combinatorial Theory, Series B, 15(2):174-177, 1973. doi:10.1016/0095-8956(73)90018-X.
- 12 William T. Tutte. On the problem of decomposing a graph into n connected factors. Journal of the London Mathematical Society, 36(1):221-230, 1961. doi:10.1112/jlms/s1-36.1.221.