Black hole remnants are not too fast to be dark matter

Benjamin V. Lehmann^{1, 2, *} and Stefano Profumo^{1, 2, †}

¹Department of Physics, University of California Santa Cruz, 1156 High St., Santa Cruz, CA 95064, USA

²Santa Cruz Institute for Particle Physics, 1156 High St., Santa Cruz, CA 95064, USA

We comment on recent claims that recoil in the final stages of Hawking evaporation gives black hole remnants large velocities, rendering them inviable as a dark matter candidate. We point out that due to cosmic expansion, such large velocities at the final stages of evaporation are not in tension with the cold dark matter paradigm so long as they are attained at sufficiently early times. In particular, the predicted recoil velocities are robustly compatible with observations if the remnants form before the epoch of big bang nucleosynthesis, a requirement which is already imposed by the physics of nucleosynthesis itself.

Remnants of black hole evaporation have long been considered a potential dark matter candidate [1, 2]. Recently, however, Ref. [3] argued that such remnants are incompatible with cold dark matter. The reasoning is as follows: such remnants are expected to form if evaporation stalls at a mass $M_{\rm R}$ near the Planck scale $M_{\rm P}$. In the final stages of evaporation, the black hole will thus have an extremely high temperature, as $T \sim M_{\rm P}^2/M_{\rm R} \sim M_{\rm P}$. Such a black hole sheds an appreciable fraction of its mass with the emission of a small number $N_{\rm q}$ of highly energetic quanta, effectively performing a random walk in momentum due to the successive recoils. The typical momentum of the black hole after such a random walk scales as $1/\sqrt{N_{\rm q}}$, and with a few well-motivated assumptions regarding the final stages of evaporation, Ref. [3] finds that typical remnants acquire a velocity $v \sim 0.1$ as a result of the process. Ref. [3] states that these velocities are in tension with the cold dark matter paradigm. This motivated the more recent analysis of Ref. [4], which found that such bounds could be evaded in a certain class of modifications to the physics of evaporation.

Here we point that, in fact, the recoil velocity does not constrain typical models of black hole remnants as dark matter: crucially, the last step of the argument neglects the effects of cosmic expansion. These remnants may indeed be produced 'warm', with semi-relativistic speeds, but the population will inevitably cool as the Universe expands. The cold dark matter paradigm requires only that they slow to non-relativistic speeds by the time observable structures begin to form, which is much less constraining. Indeed, in standard particle dark matter scenarios, the dark matter species is relativistic at early times, being in thermal equilibrium at temperatures much greater than its mass [5]. Due to cosmic expansion, such thermal relics eventually become cold. Nor does this effect require thermal equilibrium: consider, for instance, the cosmic neutrino background, which thermally decoupled at a temperature of $\mathcal{O}(1 \text{ MeV})$, and which today is non-relativistic [6]. The momentum of each individual particle, regardless of its nature, is inversely proportional to the cosmic scale factor [7].

A species that is cold today may still have a momentum history incompatible with structure formation, since the smallest dark matter structures in the late universe formed at early times. In general, a dark matter structure of a given size starts to form when the cosmological horizon grows to encompass the length scale of the corresponding density perturbations. If dark matter is relativistic and kinetically decoupled at this time, then the dark matter particles will 'free-stream' rather than immediately forming a bound structure. Thus, the smallest dark matter structures observed today correspond to the earliest times at which dark matter is required to be non-relativistic. The situation considered here is analogous to particle dark matter models in which the dark matter species undergoes thermal decoupling while still relativistic ('warm dark matter'), which can indeed suppress the formation of structure on small scales [8–12].

The scale at which structure formation is suppressed is fixed by the comoving free streaming length $\ell_{\rm fs}$, i.e., the comoving length scale over which a typical particle travels after free streaming begins. Inhomogeneities on smaller scales are erased, and this produces an exponential cutoff in the matter power spectrum at a corresponding wavenumber $k_{\rm fs} \sim 1/\ell_{\rm fs}$. For particle dark matter that begins free streaming after kinetic decoupling, this cutoff is estimated by Ref. [11] as

$$k_{\rm fs} \simeq \left(\frac{m_{\chi}}{T_{\rm kd}}\right)^{1/2} \frac{a_{\rm eq}/a_{\rm kd}}{\log(4a_{\rm eq}/a_{\rm kd})} \frac{a_{\rm eq}}{a_0} H_{\rm eq},\tag{1}$$

where m_{χ} is the dark matter mass; $T_{\rm kd}$ is the dark matter temperature at kinetic decoupling; $a_{\rm eq}$ and $H_{\rm eq}$ denote the scale factor and the Hubble parameter at matter-radiation equality; $a_{\rm kd}$ denotes the scale factor at kinetic decoupling; and $a_0 = 1$ denotes the scale factor today. Here the ratio $m_{\chi}/T_{\rm kd}$ appears as a parameter of the phase space

^{*} benvlehmann@gmail.com

[†] profumo@ucsc.edu

distribution of dark matter in thermal equilibrium. While the black hole remnants are not in thermal equilibrium, their phase space distribution is very similar: a set of random walks in momentum space produces momenta that are approximately normally distributed, which corresponds exactly to the Maxwell–Boltzmann distribution in the non-relativistic limit. Thus, for an order of magnitude estimate, we can parametrize the mass-to-temperature ratio by the rms velocity \bar{v} , using $m_{\chi}/T \approx 3/\bar{v}^2$. Then, writing $H_{\rm eq}$ in terms of the present-day matter density $\Omega_{\rm M}$, Eq. (1) becomes

$$k_{\rm fs} \simeq \frac{\sqrt{6}H_0 \Omega_{\rm M}^{1/2} a_{\rm eq}^{1/2}}{\bar{v}_i a_i \log(4a_{\rm eq}/a_i)} \approx 0.18 \,{\rm Mpc}^{-1} \left(\frac{0.1}{\bar{v}_i}\right) \left(\frac{a_{\rm eq}}{a_i} \frac{1}{\log(4a_{\rm eq}/a_i)}\right),\tag{2}$$

where now \bar{v}_i denotes the rms velocity of the black hole remnants at the beginning of free streaming, i.e., immediately after they are accelerated by evaporation.

The smallest structures probed by observations today still correspond to scales $k \lesssim 10 \,\mathrm{Mpc}^{-1}$ in the matter power spectrum [13, 14], so to determine the latest viable formation time of the remnants, we can conservatively impose $k_{\rm fs} > 10 \,\mathrm{Mpc}^{-1}$. Taking $\bar{v}_i = 0.1$, this translates to the requirement that $a_i \lesssim 7 \times 10^{-7}$. In the most extreme case, when $N_{\rm q} = 1$ and the recoil momentum is $p \simeq M_{\rm R}$, then $E^2 = p^2 + M_{\rm R}^2 \simeq 2M_{\rm R}^2$, so $\gamma \simeq \sqrt{2}$, or $\bar{v}_i \simeq 1/\sqrt{2}$. This, in turn, corresponds to the constraint $a_i \lesssim 8 \times 10^{-8}$. A comparable bound can be derived from the effective condition that the dark matter velocity today is less than 5×10^{-7} , as stated by Refs. [15, 16] based on the results of Ref. [17]. The momentum p_0 of a remnant today is related to the momentum p_i after evaporation by $p_0 = a_i p_i$, so we find that $a_i \lesssim 5 \times 10^{-7}$ for $\bar{v}_i = 1/\sqrt{2}$.¹ These upper bounds on a_i might indeed be viewed as constraints on the evaporation history of black hole remnants

These upper bounds on a_i might indeed be viewed as constraints on the evaporation history of black hole remnants due to the recoil velocity: in order to account for dark matter, such objects must form at scale factors $a \leq 10^{-7}$. However, this is much larger than the scale factor at the epoch of big bang nucleosynthesis (BBN), $a_{\text{BBN}} \simeq 2.5 \times 10^{-10}$. BBN is highly sensitive to injected energy, so there are already stringent constraints on active evaporation during this epoch [18–21]. Typically, in phenomenologically viable models of black hole remnants, evaporation must halt when $a < a_{\text{BBN}}$ to evade these bounds. Therefore, while the recoil velocity can in principle lead to constraints on the evaporation history of black hole remnants as dark matter, the actual constraints derived from this method are subdominant to existing BBN limits.

We conclude that in any model of black hole remnants as dark matter that is observationally consistent with light element abundances, the remnants are automatically 'cold' as required by the cold dark matter paradigm. In particular, black hole remnants remain viable as a dark matter candidate.

ACKNOWLEDGMENTS

We gratefully acknowledge valuable discussions with Samuel Kováčik and Yen Chin Ong. BVL and SP are supported in part by DOE grant DE-SC0010107.

- [1] J. H. MacGibbon, Can Planck-mass relics of evaporating black holes close the universe?, Nature **329**, 308 (1987).
- [2] P. Chen and R. J. Adler, Black hole remnants and dark matter, Nucl. Phys. B Proc. Suppl. 124, 103 (2003), arXiv:gr-qc/0205106.
- [3] S. Kováčik, Hawking-Radiation Recoil of Microscopic Black Holes, (2021), arXiv:2102.06517 [gr-qc].
- [4] S. D. Gennaro and Y. C. Ong, Feasibility of Primordial Black Hole Remnants as Dark Matter in View of Hawking Radiation Recoil, (2021), arXiv:2104.08919 [gr-qc].
- [5] S. Profumo, An Introduction to Particle Dark Matter (World Scientific, 2017).
- [6] J. M. Irvine and R. Humphreys, Neutrino masses and the cosmic neutrino background, J. Phys. G 9, 847 (1983).
- [7] E. W. Kolb and M. S. Turner, The Early Universe (1990) Chap. 3.4, p. 69.
- [8] P. Bode, J. P. Ostriker, and N. Turok, Halo formation in warm dark matter models, Astrophys. J. 556, 93 (2001), arXiv:astro-ph/0010389.
- [9] V. Berezinsky, V. Dokuchaev, and Y. Eroshenko, Small-scale clumps in the galactic halo and dark matter annihilation, Phys. Rev. D 68, 103003 (2003), arXiv:astro-ph/0301551.
- [10] A. M. Green, S. Hofmann, and D. J. Schwarz, The power spectrum of SUSY-CDM on sub-galactic scales, Mon. Not. Roy. Astron. Soc. 353, L23 (2004), arXiv:astro-ph/0309621.

¹ This is a numerical coincidence, not a typographical error.

- [11] A. M. Green, S. Hofmann, and D. J. Schwarz, The first WIMPy halos, JCAP 08, 003, arXiv:astro-ph/0503387.
- [12] T. Bringmann, Particle Models and the Small-Scale Structure of Dark Matter, New J. Phys. 11, 105027 (2009), arXiv:0903.0189 [astro-ph.CO].
- [13] B. Abolfathi et al. (SDSS), The Fourteenth Data Release of the Sloan Digital Sky Survey: First Spectroscopic Data from the Extended Baryon Oscillation Spectroscopic Survey and from the Second Phase of the Apache Point Observatory Galactic Evolution Experiment, Astrophys. J. Suppl. 235, 42 (2018), arXiv:1707.09322 [astro-ph.GA].
- [14] M. A. Troxel et al. (DES), Dark Energy Survey Year 1 results: Cosmological constraints from cosmic shear, Phys. Rev. D 98, 043528 (2018), arXiv:1708.01538 [astro-ph.CO].
- [15] T. Fujita, M. Kawasaki, K. Harigaya, and R. Matsuda, Baryon asymmetry, dark matter, and density perturbation from primordial black holes, Phys. Rev. D 89, 103501 (2014), arXiv:1401.1909 [astro-ph.CO].
- [16] L. Morrison, S. Profumo, and Y. Yu, Melanopogenesis: Dark Matter of (almost) any Mass and Baryonic Matter from the Evaporation of Primordial Black Holes weighing a Ton (or less), JCAP 05, 005, arXiv:1812.10606 [astro-ph.CO].
- [17] M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese, and A. Riotto, Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with WMAP and the Lyman-alpha forest, Phys. Rev. D 71, 063534 (2005), arXiv:astro-ph/0501562.
- [18] B. V. Vainer and P. D. Naselskii, Cosmological implications of the process of primordial black hole evaporation, Soviet Ast. 22, 138 (1978).
- [19] K. Kohri and J. Yokoyama, Primordial black holes and primordial nucleosynthesis. 1. Effects of hadron injection from low mass holes, Phys. Rev. D 61, 023501 (2000), arXiv:astro-ph/9908160.
- [20] B. J. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, New cosmological constraints on primordial black holes, Phys. Rev. D 81, 104019 (2010), arXiv:0912.5297 [astro-ph.CO].
- [21] S. K. Acharya and R. Khatri, CMB and BBN constraints on evaporating primordial black holes revisited, JCAP 06, 018, arXiv:2002.00898 [astro-ph.CO].