SHARP SMOOTHING PROPERTIES OF AVERAGES OVER CURVES

HYERIM KO, SANGHYUK LEE, AND SEWOOK OH

ABSTRACT. We prove sharp smoothing properties of the averaging operator defined by convolution with a measure on a smooth nondegenerate curve γ in \mathbb{R}^d , $d \geq 3$. Despite the simple geometric structure of such curves, the sharp smoothing estimates have remained largely unknown except for those in low dimensions. Devising a novel inductive strategy, we obtain the optimal L^p Sobolev regularity estimates, which settle the conjecture raised by Beltran-Guo-Hickman-Seeger. Besides, we show the sharp local smoothing estimates for every d. As a result, we establish, for the first time, nontrivial L^p boundedness of the maximal average over dilations of γ for $d \geq 4$.

1. INTRODUCTION

The regularity property of integral transforms defined by averages over submanifolds is a fundamental subject in harmonic analysis, which has been extensively studied since the 1970s. There is an immense body of literature devoted to the subject (see, for example, [33, 21, 32, 8] and references therein). However, numerous problems remain wide open. The regularity property is typically addressed in the frameworks of L^p improving, L^p Sobolev regularity, and local smoothing estimates, to which L^p boundedness of the maximal average is also closely related. In this paper, we study the smoothing estimates for the averaging operator given by convolution with a measure supported on a curve.

Let I = [-1, 1] and γ be a smooth curve from I to \mathbb{R}^d . We define a measure \mathfrak{m}_t supported on $t\gamma$ by

$$\langle \mathfrak{m}_t, f \rangle = \int f(t\gamma(s))\psi(s)ds$$

where $\psi \in C_c^{\infty}((-1,1))$. We are concerned with $d \geq 3$ since all the problems we address in the current paper are well understood when d = 2. We consider the averaging operator

$$\mathcal{A}_t f(x) = f * \mathfrak{m}_t(x)$$

and study the above-mentioned regularity problems on \mathcal{A}_t under the assumption that γ is nondegenerate, that is to say,

(1.1)
$$\det(\gamma'(s),\ldots,\gamma^{(d)}(s)) \neq 0, \quad s \in I.$$

The L^p improving property of \mathcal{A}_t for a fixed $t \neq 0$ now has a complete characterization, see [7, 34] (also, see [36] for generalizations to variable coefficient settings). However, L^p Sobolev and local smoothing estimates for \mathcal{A}_t turned out to be more involved and are far less well understood. Recently, there has been progress in low dimensions d = 3, 4 ([24, 14, 1, 2]), but it does not seem feasible to extend the

²⁰¹⁰ Mathematics Subject Classification. 42B25 (42B20).

Key words and phrases. L^p Sobolev regularity, local smoothing, maximal function.

approaches in recent works to higher dimensions. We discuss this matter in detail near the end of the introduction. By devising an inductive strategy, we prove the optimal L^p Sobolev regularity and sharp local smoothing estimates in any dimension $d \ge 3$. As a result, we also obtain L^p boundedness of the associated maximal function which was unknown for $d \ge 4$.

 L^p Sobolev regularity. Let $2 \leq p \leq \infty$. We set $\mathcal{A}f = \mathcal{A}_1 f$ and consider the L^p Sobolev regularity estimate

(1.2)
$$\|\mathcal{A}f\|_{L^p_{\alpha}(\mathbb{R}^d)} \le C\|f\|_{L^p(\mathbb{R}^d)}.$$

When d = 2, the estimate holds if and only if $\alpha \leq 1/p$ (e.g., see [6]). In higher dimensions, however, the problem of obtaining (1.2) with the sharp smoothing order α becomes highly nontrivial except for the $L^2 \to L^2_{1/d}$ estimate which is an easy consequence of the decay of Fourier transform of \mathfrak{m}_t :

(1.3)
$$|\widehat{\mathfrak{m}}_t(\xi)| \le C(1+|t\xi|)^{-1/d}$$

It was conjectured by Beltran, Guo, Hickman, and Seeger [2, Conjecture 1] that (1.2) holds for $\alpha \leq 1/p$ if 2d - 2 . When <math>d = 3, the conjecture was verified by the conditional result of Pramanik and Seeger [24] and the decoupling inequality due to Bourgain and Demeter [4] (see [20, 35] for earlier results). The case d = 4 was recently obtained by Beltran et al [2]. Our first result proves the conjecture for every $d \geq 5$.

Theorem 1.1. Let $d \ge 3$. Suppose γ is a smooth nondegenerate curve. Then, (1.2) holds for $\alpha \le 1/p$ if p > 2(d-1).

Interpolation with the $L^2 \to L_{1/d}^2$ estimate gives (1.2) for $\alpha < (p+2)/(2dp)$ when $2 . It is also known that (1.2) fails if <math>\alpha > \alpha(p) := \min(1/p, (p+2)/(2dp))$ (see [2, Proposition 1.2]). Thus, only the estimate (1.2) with $\alpha = \alpha(p)$ remains open for 2 . Those endpoint estimates seem to be a subtleproblem. The argument in this paper provides simpler alternative proofs of theprevious results for <math>d = 3, 4. Theorem 1.1 remains valid as long as $\gamma \in C^{2d}(I)$ (see Theorem 4.1). However, we do not try to optimize the regularity assumption.

The result in Theorem 1.1 can be easily generalized to curves of different types. We say a smooth curve γ from I to \mathbb{R}^d is of finite type if there is an ℓ such that $\operatorname{span}\{\gamma^{(1)}(s),\ldots,\gamma^{(\ell)}(s)\}=\mathbb{R}^d$ for each $s \in I$. The type at s is defined to be the smallest of such ℓ and the maximal type is the supremum over $s \in I$ of the type at s. (See, e.g., [24, 12].) Using Theorem 1.1 and a rescaling argument ([24, 12]) one can obtain the following, which proves the Conjecture 2 in [2].

Corollary 1.2. Let $d \ge 3$, $\ell > d$ and $2 \le p < \infty$. Suppose γ is a curve of maximal type ℓ . Then (1.2) holds for $\alpha \le \min(\alpha(p), 1/\ell)$ if $p \ne \ell$ when $\ell \ge 2d - 2$, and if $p \in [2, 2\ell/(2d - \ell)) \cup (2d - 2, \infty)$ when $d < \ell < 2d - 2$.

By interpolation (1.2) holds for $\alpha < \min(\alpha(p), 1/\ell)$ if $p = \ell$ when $\ell \ge 2d - 2$, and if $2\ell/(2d - \ell) \le p \le 2d - 2$ when $d < \ell < 2d - 2$. These estimates are sharp. Since a finite type curve contains a nondegenerate subcurve and the $L^2 \to L^2_{1/\ell}$ estimate is optimal, (1.2) fails if $\alpha > \min(\alpha(p), 1/\ell)$. When $\ell \ge 2d - 2$, Corollary 1.2 resolves the problem of the Sobolev regularity estimate (1.2). In fact, the failure of $L^\ell \to L^\ell_{1/\ell}$ bound was shown in [2] using Christ's example [6]. By [28, Theorem 1.1], Corollary 1.2 also gives $H^1(\mathbb{R}^d) \to L^{1,\infty}(\mathbb{R}^d)$ bound on the lacunary maximal function $f \to \sup_{k \in \mathbb{Z}} |f * \mathfrak{m}_{2^k}|$ whenever γ is of finite type. Sharp local smoothing. We now consider the estimate

(1.4)
$$\|\chi(t)\mathcal{A}_t f\|_{L^p_{\alpha}(\mathbb{R}^{d+1})} \le C \|f\|_{L^p(\mathbb{R}^d)},$$

where χ is a smooth function supported in (1/2, 4). Compared with the L^p Sobolev estimate (1.2), the additional integration in t is expected to yield extra smoothing. Such a phenomenon is called *local smoothing*, which has been studied for the dispersive equations to a great extent (e.g., see [29, 9]). However, the local smoothing for the averaging operators exhibits considerably different nature.

In particular, there is no local smoothing when p = 2. Besides, a bump function example shows $\alpha \leq 1/d$. As we shall see later, the estimate (1.4) fails unless $\alpha \leq 2/p$ (Proposition 3.9). So, it seems to be plausible to conjecture that (1.4) holds for $\alpha < \min(2/p, 1/d)$ if 2 . For <math>d = 2, the conjecture follows by the recent result on Sogge's local smoothing conjecture for the wave operator ([30, 38, 16, 4]), which is due to Guth, Wang, and Zhang [11]. When d = 3, some local smoothing estimates were utilized by Pramanik and Seeger [24] and Beltran et al. [1] to prove L^p maximal bound.

Nevertheless, for $d \ge 3$, no local smoothing estimate up to the sharp order 2/p has been known previously.

Theorem 1.3. Let $d \ge 3$. Suppose γ is a smooth nondegenerate curve. Then, if $p \ge 4d - 2$, (1.4) holds true for $\alpha < 2/p$.

Theorem 1.3 remains valid as far as $\gamma \in C^{3d+1}(I)$ (see Theorem 2.2 below).

Maximal estimate. The local smoothing estimate (1.4) has been of particular interest in connection to L^p boundedness of the maximal operator

$$Mf(x) = \sup_{0 < t} |\mathcal{A}_t f(x)|$$

([19, 27, 24, 1]) and problems in geometric measure theory (see, e.g., [38] and Corollary 1.6 below). If the estimate (1.4) holds for some $\alpha > 1/p$, L^p boundedness of M follows by a standard argument relying on the Sobolev embedding ([24]).

The study of the maximal functions generated by dilations of submanifolds goes back to Stein's spherical maximal theorem [31] (see, also, [32, Ch.10] and [13]). The circular maximal theorem was later proved by Bourgain [3] (also, see [30, 19, 26, 27, 15]). Afterwards, a natural question was whether the maximal operator M under consideration in the current paper is bounded on L^p for some $p \neq \infty$ when $d \geq 3$. In view of Stein's interpolation argument based on L^2 estimate ([31]), proving L^p boundedness of M becomes more challenging as d increases since the decay of the Fourier transform of \mathfrak{m}_t gets weaker (see (1.3)). Though the question was raised as early as in the late 1980s, it remained open for any $d \geq 3$ until recently. In \mathbb{R}^3 , the first positive result was obtained by Pramanik and Seeger [24] and the range of pwas further extended to p > 4 thanks to the decoupling inequality for the cone [4]. Very recently, the authors [14] proved L^p boundedness of M on the optimal range, i.e., M is bounded on L^p if and only if p > 3. The same result was independently obtained by Beltran et al. [1].

However, no nontrival L^p bound on M has been known in higher dimensions. The following establishes existence of such maximal bound for every $d \ge 4$. **Theorem 1.4.** Let $d \ge 4$. Suppose γ is a smooth nondegenerate curve. Then, for p > 2(d-1) we have

(1.5)
$$\|Mf\|_{L^{p}(\mathbb{R}^{d})} \leq C\|f\|_{L^{p}(\mathbb{R}^{d})}.$$

The result is a consequence of Theorem 1.3. Since the estimate (1.4) holds for p = 2 and $\alpha = 1/d$, interpolation gives (1.4) for some $\alpha > 1/p$ when 2d - 2 . So, the maximal estimate (1.5) follows, as mentioned before, by a standard argument. A natural conjecture is that <math>M is bounded on L^p if and only if p > d. M can not be bounded on L^p if $p \leq d$, as can be seen by a simple adaptation of the argument in [14, Proposition 4.4]. Theorem 1.4 also extends to the finite type curves by a rescaling argument ([12, 24]). The following result is sharp when $\ell \geq 2(d-1)$.

Corollary 1.5. Let $d \ge 4$ and $\ell > d$. Suppose γ is a curve of maximal type ℓ . Then (1.5) holds if $p > \max(\ell, 2(d-1))$.

Packing of curves in \mathbb{R}^d . The sharp local smoothing estimate (1.4) in Theorem 1.3 has interesting measure theoretic consequences concerning unions of curves generated by translation and dilation of a nondegenerate curve. The following generalizes Wolff's result [38, Corollary 3], where unions of circles in \mathbb{R}^2 were considered (see also [17, 18, 37] for earlier results).

Corollary 1.6. Let γ be a smooth nondegenerate curve in \mathbb{R}^d , $d \geq 3$, and let $E \subset \mathbb{R}^{d+1}$ be a set of Hausdorff dimension greater than d-1. Suppose F is a set in \mathbb{R}^d such that $(x + t\gamma(I)) \cap F$ has positive 1-dimensional outer measure for all $(x,t) \in E$. Then F has positive outer measure.

Corollary 1.6 follows by Theorem 1.3 and the argument in [38]. The result does not hold in general without the nondegeneracy assumption on γ as one can easily see considering a curve contained in a lower dimensional affine space. The same result continues to be valid for the finite type curve. Consequently, Corollary 1.6 implies the following.

Corollary 1.6'. Let γ be a smooth finite type curve in \mathbb{R}^d , $d \geq 3$, and let E and F be compact subsets in \mathbb{R}^d . Suppose E has Hausdorff dimension greater than d-1 and for each $x \in E$ there is t(x) > 0 such that $x + t(x)\gamma(I) \subset F$. Then, F has positive measure.

Our approach. To prove L^p $(p \neq 2)$ smoothing properties of \mathcal{A}_t , we need more than the decay of $\hat{\mathfrak{m}}_t$, i.e., (1.3). When d = 2, we have rather a precise asymptotic expansion of $\hat{\mathfrak{m}}_t$, which makes it possible to relate \mathcal{A}_t to other forms of operators. In fact, one can use the estimate for the wave operator (e.g., [27, 35, 15]) to obtain local smoothing estimate. However, in higher dimensions $d \geq 3$, to compute $\hat{\mathfrak{m}}_t$ explicitly is not a simple matter. Even worse, this becomes much more complicated as dincreases since one has to take into account the derivatives $\gamma^{(k)}(s) \cdot \xi$, $k = 2, \ldots, d$. The common approach in [24, 1, 2] to get around this difficulty was to use detailed decompositions (of various scales) on the Fourier side away from the conic sets where $\hat{\mathfrak{m}}_t$ decays slowly. The consequent decompositions were then combined with the decoupling or square function estimate [20, 23, 24, 25, 1, 2]. However, this type of approach based on fine scale decomposition becomes exceedingly difficult to manage as the dimension d gets larger and, consequently, does not seem to be tractable in higher dimensions.

4

To overcome the difficulty, we develop a new strategy which allows us to dispense with such sophisticated decompositions. Before closing the introduction we briefly discuss the key ingredients of our approach.

• The main novelty of the paper lies in an induction argument which we build on the local nondegeneracy assumption:

$$\mathfrak{N}(L,B) \qquad \qquad \sum_{\ell=1}^{L} |\langle \gamma^{(\ell)}(s), \xi \rangle| \ge B^{-1} |\xi|$$

for a constant $B \geq 1$. To prove our results, we consider the operator $\mathcal{A}_t[\gamma, a]$ (see (2.2) below for its definition). Clearly, $\mathfrak{N}(d, B')$ holds for a constant B' > 0 if γ satisfies (1.1). However, instead of considering the case L = d alone, we prove the estimate for all $L = 2, \ldots, d$ under the assumption that $\mathfrak{N}(L, B)$ holds on the support of a. See Theorem 2.2 and 4.1. A trivial (yet, important) observation is that $\mathfrak{N}(L-1, B)$ implies $\mathfrak{N}(L, B)$, so we may think of $\mathcal{A}_t[\gamma, a]$ as being more degenerate as L gets larger. Thanks to this hierarchical structure, we may use an inductive strategy along the number L. See Proposition 2.3 and 4.2 below.

• We extend the rescaling [12, 14] and iteration [24] arguments. Roughly speaking, we combine the first with the induction assumption in Proposition 2.3 (or 4.2) to handle the less degenerate parts, and use the latter to deal with the remaining part. In order to generalize the arguments, we introduce a class of symbols which are naturally adjusted to a small subcurve (Definition 2.4). We also use the decoupling inequalities for the nondegenerate curves obtained by Beltran et al. [2] (Corollary 2.15). Their inequalities were deduced from those due to Bourgain, Demeter, and Guth [5]. Instead of applying the inequalities directly, we use modified forms which are adjusted to the sharp smoothing orders of the specific estimates (see (2.40) and (2.41)). This makes it possible to obtain the sharp estimates on an extended range. *Organization of the paper.* We first prove Theorem 1.3 whose proof is more involved than that of Theorem 1.1. In Section 2, we reduce the proof of Theorem 1.3 to that of Proposition 2.9, which we prove while assuming Proposition 2.10. The proof of Proposition 2.10 is given in Section 3. We prove Theorem 1.1 in Section 4.

2. Smoothing estimates with localized frequency

In this section, we consider an extension of Theorem 1.3 via microlocalization (see Theorem 2.2 below) which we can prove inductively. We then reduce the matter to proving Proposition 2.9, which we show by applying Proposition 2.10. We also obtain some preparatory results.

Let $1 \leq L \leq d$ be a positive integer and $B \geq 1$ be a large number. For quantitative control of estimates we consider the following two conditions:

(2.1)
$$\max_{0 \le j \le 3d+1} |\gamma^{(j)}(s)| \le B, \qquad s \in I,$$

$$\mathfrak{V}(L,B)$$
 $\operatorname{Vol}\left(\gamma^{(1)}(s),\ldots,\gamma^{(L)}(s)\right) \ge 1/B,$ $s \in I,$

where $Vol(v_1, \ldots, v_L)$ denotes the *L*-dimensional volume of the parallelepiped generated by v_1, \ldots, v_L . By finite decomposition, rescaling, and a change of variables, the constant *B* can be taken to be close to 1 (see Section 2.2).

Notation. For nonnegative quantities A and D, we denote $A \leq D$ if there exists an independent positive constant C such that $A \leq CD$, but the constant C may differ at each occurrence depending on the context, and $A \leq_B D$ means the inequality holds with an implicit constant depending on B. Throughout the paper, the constant C mostly depends on B. However, we do not make it explicit every time since it is clear in the context. By A = O(D) we denote $|A| \leq D$.

Definition 2.1. For $k \geq 0$, we denote $\mathbb{A}_k = \{\xi \in \mathbb{R}^d : 2^{k-1} \leq |\xi| \leq 2^{k+1}\}$. We say $a \in \mathbb{C}^{d+L+2}(\mathbb{R}^{d+2})$ is a symbol of type (k, L, B) relative to γ if supp $a \subset I \times [2^{-1}, 4] \times \mathbb{A}_k$, $\mathfrak{N}(L, B)$ holds for γ whenever $(s, t, \xi) \in \text{supp } a$ for some t, and

$$|\partial_s^j \partial_t^l \partial_{\varepsilon}^{\alpha} a(s, t, \xi)| \le B |\xi|^{-|\alpha|}$$

for $(j, l, \alpha) \in \mathcal{I}_L := \{(j, l, \alpha) : 0 \le j \le 1, 0 \le l \le 2L, |\alpha| \le d + L + 2\}.$

We define an integral operator by

(2.2)
$$\mathcal{A}_t[\gamma, a]f(x) = (2\pi)^{-d} \iint_{\mathbb{R}} e^{i(x-t\gamma(s))\cdot\xi} a(s, t, \xi) ds \,\widehat{f}(\xi) \, d\xi.$$

Note $\mathcal{A}_t f = \mathcal{A}_t[\gamma, \psi] f$. Theorem 1.3 is a consequence of the following.

Theorem 2.2. Let $\gamma \in C^{3d+1}(I)$ satisfy (2.1) and $\mathfrak{V}(L, B)$ for some $B \geq 1$. Suppose a is a symbol of type (k, L, B) relative to γ . Then, if $p \geq 4L - 2$, for any $\epsilon > 0$ there is a constant $C_{\epsilon} = C_{\epsilon}(B)$ such that

(2.3)
$$\|\mathcal{A}_t[\gamma, a]f\|_{L^p(\mathbb{R}^{d+1})} \le C_{\epsilon} 2^{(-\frac{2}{p}+\epsilon)k} \|f\|_{L^p(\mathbb{R}^d)}.$$

Theorem 2.2 is trivial when L = 1. Indeed, (2.3) follows from the estimate $|\mathcal{A}_t[\gamma, a]f(x)| \leq_B \int_I K * |f|(x - t\gamma(s)) \, ds$ where $K(x) = 2^{(d-1)k}(1 + |2^kx|)^{-d-3}$. To show this, note $|\gamma'(s) \cdot \xi| \sim 2^k$ if $(s, t, \xi) \in \text{supp } a$ for some t. By integration by parts in s, $\mathcal{A}_t[\gamma, a] = t^{-1}\mathcal{A}_t[\gamma, \tilde{a}]$ where $\tilde{a} = i(\gamma'(s) \cdot \xi \partial_s a - \gamma''(s) \cdot \xi a)/(\gamma'(s) \cdot \xi)^2$. Since $|\partial_{\xi}^{\alpha}\tilde{a}| \leq |\xi|^{-|\alpha|-1}$ for $|\alpha| \leq d+3$, routine integration by parts in ξ gives the estimate (e.g., see *Proof of Lemma 2.7*). When L = 2, Theorem 2.2 is already known by the result in [24, Theorem 4.1] and the decoupling inequality in [4].

Once we have Theorem 2.2, the proof of Theorem 1.3 is straightforward. By Littlewood-Paley decomposition it is sufficient to show (2.3) for $p \ge 4d - 2$ with $a_k(s,t,\xi) = \psi(s)\chi(t)\beta(2^{-k}|\xi|)$, where $\beta \in C_c^{\infty}((1/2,2))$. This can be made rigorous using $\iint e^{-it(\tau+\gamma(s)\cdot\xi)}\psi(s)\chi(t)dsdt = O((1+|\tau|)^{-N})$ for any N if $|\tau| \ge (1 + \max_{s \in \text{supp }\psi} |\gamma(s)|)|\xi|$. Since γ satisfies (1.1), a_k is of type (k,d,B) relative to γ for a large B. Therefore, Theorem 1.3 follows from Theorem 2.2.

Theorem 2.2 is immediate from the next proposition, which places Theorem 2.2 in an inductive framework.

Proposition 2.3. Let $2 \le N \le d$. Suppose Theorem 2.2 holds for L = N - 1. Then, Theorem 2.2 holds true with L = N.

To prove Proposition 2.3, from this section to Section 3 we fix $N \in [2, d]$, γ satisfying $\mathfrak{V}(N, B)$, and a symbol *a* of type (k, N, B) relative to γ .

One of the main ideas is that by a suitable decomposition of the symbol we can separate from $\mathcal{A}_t[\gamma, a]$ the less degenerate part which corresponds to L = N - 1. To this part we apply the assumption combined with a rescaling argument. To do this, we introduce a class of symbols which are adjusted to short subcurves of γ .

 $\mathbf{6}$

2.1. Symbols associated to subcurves. We begin with some notations. Let $N \geq 2$, and let δ and B' denote the numbers such that

$$2^{-k/N} \le \delta \le 2^{-7dN} B^{-6N}, \qquad B \le B' \le B^C$$

for a large constant $C \geq 3d + 1$. We note that $\mathfrak{V}(N-1, B')$ holds for some B'. In fact, $\mathfrak{V}(N-1, B^2)$ follows by (2.1) and $\mathfrak{V}(N, B)$.

For $s \in I$, we define a linear map $\widetilde{\mathcal{L}}_s^{\delta} : \mathbb{R}^d \mapsto \mathbb{R}^d$ as follows:

(2.4)
$$(\widetilde{\mathcal{L}}_{s}^{\delta})^{\mathsf{T}} \gamma^{(j)}(s) = \delta^{N-j} \gamma^{(j)}(s), \qquad j = 1, \dots, N-1, \\ (\widetilde{\mathcal{L}}_{s}^{\delta})^{\mathsf{T}} v = v, \qquad v \in \left(\mathbf{V}_{s}^{\gamma, N-1}\right)^{\perp},$$

where $V_s^{\gamma,\ell} = \text{span} \{ \gamma^{(j)}(s) : j = 1, \dots, \ell \}$. $\widetilde{\mathcal{L}}_s^{\delta}$ is well-defined since $\mathfrak{V}(N-1, B^2)$ holds for γ . The linear map $\widetilde{\mathcal{L}}_s^{\delta}$ naturally appears when we rescale a subcurve of length about δ (see the proofs of Lemma 2.7 and 2.8). We denote

(2.5)
$$\mathcal{L}_{s}^{\delta}(\tau,\xi) = \left(\delta^{N}\tau - \gamma(s) \cdot \widetilde{\mathcal{L}}_{s}^{\delta}\xi, \ \widetilde{\mathcal{L}}_{s}^{\delta}\xi\right), \qquad (\tau,\xi) \in \mathbb{R} \times \mathbb{R}^{d}$$

We set $G(s) = (1, \gamma(s))$ and define

$$\Lambda_k(s,\delta,B') = \bigcap_{0 \le j \le N-1} \left\{ (\tau,\xi) \in \mathbb{R} \times \mathbb{A}_k : |\langle G^{(j)}(s), (\tau,\xi) \rangle| \le B' 2^{k+5} \delta^{N-j} \right\}.$$

Definition 2.4. Let $(s_0, \delta) \in (-1, 1) \times (0, 1)$ such that $I(s_0, \delta) := [s_0 - \delta, s_0 + \delta] \subset I$. Then, by $\mathfrak{A}_k(s_0, \delta)$ we denote the set of $\mathfrak{a} \in \mathbb{C}^{d+N+2}(\mathbb{R}^{d+3})$ such that

(2.6)
$$\operatorname{supp} \mathfrak{a} \subset I(s_0, \delta) \times [2^{-1}, 2^2] \times \Lambda_k(s_0, \delta, B),$$

(2.7)
$$\left| \partial_s^j \partial_t^l \partial_{\tau,\xi}^{\alpha} \mathfrak{a}(s,t,\mathcal{L}_{s_0}^{\delta}(\tau,\xi)) \right| \le B\delta^{-j} |(\tau,\xi)|^{-|\alpha|}, \qquad (j,l,\alpha) \in \mathcal{I}_N.$$

We define $\operatorname{supp}_{\xi} \mathfrak{a} = \bigcup_{s,t,\tau} \operatorname{supp} \mathfrak{a}(s,t,\tau,\cdot)$ and $\operatorname{supp}_{s,\xi} \mathfrak{a} = \bigcup_{t,\tau} \operatorname{supp} \mathfrak{a}(\cdot,t,\tau,\cdot)$, and $\operatorname{supp}_{s} \mathfrak{a}$ and $\operatorname{supp}_{\tau,\xi} \mathfrak{a}$ are defined likewise. We note a statement $S(s,\xi)$, depending on (s,ξ) , holds on $\operatorname{supp}_{s,\xi} \mathfrak{a}$ if and only if $S(s,\xi)$ holds whenever $(s,t,\tau,\xi) \in \operatorname{supp} \mathfrak{a}$ for some t, τ .

Denote $V_s^{G,\ell} = \text{span}\{(1,0), G'(s), \ldots, G^{(\ell)}(s)\}$. We take a close look at the map \mathcal{L}_s^{δ} . By (2.4) and (2.5) we have

(2.8)
$$\begin{cases} (\mathcal{L}_{s}^{\delta})^{\mathsf{T}}G(s) = \delta^{N}(1,0), \\ (\mathcal{L}_{s}^{\delta})^{\mathsf{T}}G^{(j)}(s) = \delta^{N-j}G^{(j)}(s), & j = 1, \dots, N-1, \\ (\mathcal{L}_{s}^{\delta})^{\mathsf{T}}v = v, & v \in (\mathcal{V}_{s}^{G,N-1})^{\perp}. \end{cases}$$

The first identity is clear since $(\mathcal{L}_s^{\delta})^{\intercal}(\tau,\xi) = (\delta^N \tau, (\widetilde{\mathcal{L}}_s^{\delta})^{\intercal}\xi - \tau(\widetilde{\mathcal{L}}_s^{\delta})^{\intercal}\gamma(s))$. The second and the third follow from (2.4) since $G^{(j)} \in \{0\} \times \mathbb{R}^d$, $1 \leq j \leq N-1$, $(\mathbf{V}_s^{G,N-1})^{\perp} \subset \{0\} \times \mathbb{R}^d$, and $(\mathcal{L}_s^{\delta})^{\intercal}(0,\xi) = (0, (\widetilde{\mathcal{L}}_s^{\delta})^{\intercal}\xi)$. Furthermore, there is a constant C = C(B), independent of s and δ , such that

(2.9)
$$|\mathcal{L}_s^{\delta}(\tau,\xi)| \le C|(\tau,\xi)|.$$

Note that (2.9) is equivalent to $|(\mathcal{L}_s^{\delta})^{\intercal}(\tau,\xi)| \leq C|(\tau,\xi)|$. The inequality is clear from (2.4) because $\mathfrak{V}(N-1,B^2)$ holds and all the eigenvalues of $(\widetilde{\mathcal{L}}_s^{\delta})^{\intercal}$ are contained in the interval (0,1].

Lemma 2.5. Let $\mathcal{L}_s^{\delta}(\tau,\xi) \in \Lambda_k(s,\delta,B')$ and $\mathfrak{V}(N-1,B')$ holds for γ . Then, there exists a constant C = C(B') such that

(2.10)
$$C^{-1}|(\tau,\xi)| \le 2^k \le C|\xi|.$$

Proof. Since $\mathcal{L}^{\delta}_{s}(\tau,\xi) \in \Lambda_{k}(s,\delta,B')$, by (2.5) we have $2^{k-1} \leq |\widetilde{\mathcal{L}}^{\delta}_{s}\xi| \leq 2^{k+1}$. So, the second inequality in (2.10) is clear from (2.9) if we take $\tau = 0$.

To show the first inequality, from (2.8) we have $|\langle (1,0), (\tau,\xi) \rangle| \leq B' 2^{k+5}$ and $|\langle G^{(j)}(s), (\tau, \xi) \rangle| \leq B' 2^{k+5}, 1 \leq j \leq N-1$, because $\mathcal{L}_s^{\delta}(\tau, \xi) \in \Lambda_k(s, \delta, B')$. Also, if $v \in (\mathbf{V}_s^{G,N-1})^{\perp}$ and |v| = 1, by (2.8) we see $|\langle v, (\tau,\xi) \rangle| = |\langle v, \mathcal{L}_s^{\delta}(\tau,\xi) \rangle| \le 2^{k+1}$. Since $\mathfrak{V}(N-1, B')$ holds and $\mathbf{V}_s^{G,N-1} \oplus (\mathbf{V}_s^{G,N-1})^{\perp} = \mathbb{R}^{d+1}$, we get $|(\tau,\xi)| \le C2^k$ for some C = C(B'). \square

The following shows the matrices \mathcal{L}_s^{δ} , $\mathcal{L}_{s_0}^{\delta}$ are close to each other if so are s, s_0 .

Lemma 2.6. Let $s, s_0 \in (-1, 1)$ and γ satisfy $\mathfrak{V}(N-1, B')$. If $|s-s_0| \leq \delta$, then there exists a constant $C = C(B') \ge 1$ such that

(2.11)
$$C^{-1}|(\tau,\xi)| \le |(\mathcal{L}_{s_0}^{\delta})^{-1}\mathcal{L}_s^{\delta}(\tau,\xi)| \le C|(\tau,\xi)|.$$

Proof. It suffices to prove that (2.11) holds if $|s - s_0| \leq c\delta$ for a constant c > 0, independent of s and s_0 . Applying this finitely many times, we can remove the additional assumption. Moreover, it is enough to show

(2.12)
$$\| (\mathcal{L}_s^{\delta})^{\mathsf{T}} (\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}} - \mathbf{I} \| \lesssim_{B'} c$$

when $|s - s_0| \leq c\delta$. Here, $\|\cdot\|$ denotes a matrix norm. Taking c > 0 sufficiently small, we get (2.11).

By (2.8), $(\mathcal{L}_{s_0}^{\delta})^{\mathsf{T}}(\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}}G^{(j)}(s_0) = (\mathcal{L}_{s_0}^{\delta})^{\mathsf{T}}\delta^{-(N-j)}G^{(j)}(s_0)$ for $j = 1, \dots, N-1$. Let $s_0 = s + c'\delta$, $|c'| \leq c$. Expanding $G^{(j)}$ in Taylor series at s, by (2.1) we have

$$(\mathcal{L}_s^{\delta})^{\mathsf{T}}(\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}}G^{(j)}(s_0) = \left(\mathcal{L}_s^{\delta}\right)^{\mathsf{T}}\left(\sum_{\ell=j}^{N-1} \delta^{-(N-j)}G^{(\ell)}(s)\frac{(c'\delta)^{\ell-j}}{(\ell-j)!} + O\left(c^{N-j}B'\right)\right)$$

for $j = 1, \ldots, N - 1$. By (2.8) and the mean value theorem, we get

$$(\mathcal{L}_s^{\delta})^{\mathsf{T}}(\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}}G^{(j)}(s_0) = G^{(j)}(s_0) + O(cB'), \quad j = 1, \dots, N-1.$$

From (2.8) we also have $(\mathcal{L}_s^{\delta})^{\intercal}(\mathcal{L}_{s_0}^{\delta})^{-\intercal}(1,0) = \delta^{-N}(\mathcal{L}_s^{\delta})^{\intercal}G(s_0)$. A similar argument also shows $(\mathcal{L}_s^{\delta})^{\mathsf{T}}(\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}}(1,0) = (1,0) + O(cB').$

Let $\{v_N, \ldots, v_d\}$ denote an orthonormal basis of $(\mathcal{V}_{s_0}^{G,N-1})^{\perp}$. By $\mathfrak{V}(N-1,B')$ and (2.1) it follows that $|\gamma^{(j)}(s_0)| \ge (B')^{-1-N}, j = 1, ..., N-1$. Since $|\gamma^{(j)}(s) - \gamma^{(j)}(s)| \ge (N-1)^{-1-N}$. $\gamma^{(j)}(s_0)| \leq cB'\delta$, there is an orthonormal basis $\{v_N(s), \ldots, v_d(s)\}$ of $(\mathcal{V}_s^{G,N-1})^{\perp}$ such that $|v_j(s) - v_j| \leq_{B'} c\delta$, $j = N, \ldots, d$. So, we have $|(\mathcal{L}_s^{\delta})^{\mathsf{T}} v_j - v_j| \leq_{B'} c\delta$ by (2.9). Since $(\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}} v_j = v_j$, it follows that $|(\mathcal{L}_s^{\delta})^{\mathsf{T}} (\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}} v_j - v_j| \leq_{B'} c\delta$ by (2.9). We denote by M the matrix $[(1,0), G'(s_0), \ldots, G^{(N-1)}(s_0), v_N, \ldots, v_d]$. Then, combining all together, we have $||(\mathcal{L}_s^{\delta})^{\mathsf{T}} (\mathcal{L}_{s_0}^{\delta})^{-\mathsf{T}} M - M|| \leq_{B'} c$. Note that $\mathfrak{V}(N - 1, B')$

1, B') gives $|\mathcal{M}^{-1}v| \leq_{B'} |v|$ for $v \in \mathbb{R}^{d+1}$. Therefore, we obtain (2.12). \square

For a continuous function \mathfrak{a} supported in $I \times [1/2, 4] \times \mathbb{R} \times \mathbb{A}_k$, we set

(2.13)
$$m[\mathfrak{a}](\tau,\xi) = \iint e^{-it'(\tau+\gamma(s)\cdot\xi)}\mathfrak{a}(s,t',\tau,\xi)dsdt',$$

(2.14)
$$\mathcal{T}[\mathfrak{a}]f(x,t) = (2\pi)^{-d-1} \iint e^{i(x\cdot\xi+t\tau)}m[\mathfrak{a}](\tau,\xi)\widehat{f}(\xi)\,d\xi d\tau.$$

8

Lemma 2.7. Suppose $\mathfrak{a} \in C^{d+3}(\mathbb{R}^{d+3})$ satisfies (2.6) and (2.7) for j = l = 0 and $|\alpha| \leq d+3$. Then, there is a constant C = C(B) such that

(2.15)
$$\|\mathcal{T}[\mathfrak{a}]f\|_{L^{\infty}(\mathbb{R}^{d+1})} \leq C\delta \|f\|_{L^{\infty}(\mathbb{R}^{d})},$$

(2.16)
$$\| (1 - \tilde{\chi}) \mathcal{T}[\mathfrak{a}] f \|_{L^p(\mathbb{R}^{d+1})} \le C 2^{-k} \delta^{1-N} \| f \|_{L^p(\mathbb{R}^d)}, \quad p > 1,$$

where $\tilde{\chi} \in C_c^{\infty}((2^{-2}, 2^3))$ such that $\tilde{\chi} = 1$ on $[3^{-1}, 6]$.

Proof. We first note

(2.17)
$$\mathcal{T}[\mathfrak{a}]f(x,t) = \int K[\mathfrak{a}](s,t,\cdot) * f(x) \, ds,$$

where

(2.18)
$$K[\mathfrak{a}](s,t,x) = (2\pi)^{-d-1} \iiint e^{i(t-t',x-t'\gamma(s))\cdot(\tau,\xi)}\mathfrak{a}(s,t',\tau,\xi) \, d\xi d\tau dt'.$$

Since $\operatorname{supp}_{s} \mathfrak{a} \subset I(s_0, \delta)$, to prove (2.15) we need only to show

(2.19)
$$||K[\mathfrak{a}](s,\cdot)||_{L^{\infty}_{t}L^{1}_{x}} \leq C, \qquad s \in I(s_{0},\delta)$$

for some C = C(B) > 0. To this end, changing variables $(\tau, \xi) \to 2^k \mathcal{L}_s^{\delta}(\tau, \xi)$ in the right hand side of (2.18) and noting $|\det \mathcal{L}_s^{\delta}| = \delta^N |\det \widetilde{\mathcal{L}}_s^{\delta}| = \delta^{N(N+1)/2}$, we get

$$K[\mathfrak{a}](s,t,x) = C_* \iiint e^{i2^k(t-t',x-t\gamma(s))\cdot(\delta^N\tau,\,\widetilde{\mathcal{L}}_s^\delta\xi)}\mathfrak{a}(s,t',2^k\mathcal{L}_s^\delta(\tau,\xi))\,d\xi d\tau dt',$$

where $C_* = (2\pi)^{-d-1} \delta^{N(N+1)/2} 2^{k(d+1)}$. Since \mathfrak{a} satisfies (2.6), by (2.11) and Lemma 2.5 we have supp $\mathfrak{a}(s, t, 2^k \mathcal{L}_s^{\delta}) \subset \{(\tau, \xi) : |(\tau, \xi)| \leq_B 1\}$. Besides, by (2.7) and (2.11) it follows that $|\partial_{\tau,\xi}^{\alpha}(\mathfrak{a}(s, t, 2^k \mathcal{L}_s^{\delta}(\tau, \xi)))| \leq_B 1$ for $|\alpha| \leq d+3$. Thus, repeated integration by parts in τ, ξ yields

$$|K[\mathfrak{a}](s,t,x)| \lesssim C_* \int_{1/2}^4 \left(1 + 2^k \left| \left(\delta^N(t-t'), (\widetilde{\mathcal{L}}_s^\delta)^{\mathsf{T}}(x-t\gamma(s)) \right) \right| \right)^{-d-3} dt',$$

by which we obtain (2.19) as desired.

It is easy to show (2.16). The above estimate for $K[\mathfrak{a}]$ gives

$$\|(1-\tilde{\chi})K[\mathfrak{a}](s,t,\cdot)\|_{L^{1}_{x}} \lesssim \delta^{-N} 2^{-k} |t-1|^{-1} |1-\tilde{\chi}(t)|.$$

Since $\operatorname{supp}_s \mathfrak{a} \subset I(s_0, \delta)$, (2.16) for p > 1 follows by (2.17), Minkowski's and Young's convolution inequalities.

2.2. Rescaling. Let $\mathfrak{a} \in \mathfrak{A}_k(s_0, \delta)$. Suppose that

(2.20)
$$\sum_{j=1}^{N-1} \delta^j |\langle \gamma^{(j)}(s), \xi \rangle| \ge 2^k \delta^N / B'$$

holds on $\operatorname{supp}_{s,\xi} \mathfrak{a}$ for some B' > 0. Then, via decomposition and rescaling, we can bound the L^p norm of $\mathcal{T}[\mathfrak{a}]f$ by those of the operators given by symbols of type $(j, N - 1, \tilde{B})$ relative to a curve for some \tilde{B} and j (see Lemma 2.8 below).

To do so, we define a rescaled curve $\gamma_{s_0}^{\delta}: I \to \mathbb{R}^d$ by

(2.21)
$$\gamma_{s_0}^{\delta}(s) = \delta^{-N} (\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}} (\gamma(\delta s + s_0) - \gamma(s_0)).$$

As $\delta \to 0$, the curves $\gamma_{s_0}^{\delta}$ get close to a nondegenerate curve in N dimensional vector space, so the curves behave in a uniform way. In particular, (2.1) and $\mathfrak{V}(N, B)$ hold for some B for $\gamma_{s_0}^{\delta}$ if $\delta < \delta'$ for a constant $\delta' = \delta'(B)$ small enough.

Note $(\gamma_{s_0}^{\delta})^{(j)}(s) = \delta^{j-N}(\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\intercal}\gamma^{(j)}(\delta s + s_0), 1 \leq j \leq N-1, \text{ and } |(\gamma_{s_0}^{\delta})^{(j)}(s)| \leq B\delta,$ $N+1 \leq j \leq 3d+1$. Thus, Taylor series expansion and (2.4) give

$$(\gamma_{s_0}^{\delta})^{(j)}(s) = \sum_{k=0}^{N-j-1} \frac{\gamma^{(j+k)}(s_0)}{k!} s^k + \frac{(\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}} \gamma^{(N)}(s_0)}{(N-j)!} s^{N-j} + O(B\delta)$$

for j = 1, ..., N - 1. By (2.21), we have $(\gamma_{s_0}^{\delta})^{(N)}(s) = (\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}}\gamma^{(N)}(s_0) + O(\delta)$. We write $\gamma^{(N)}(s_0) = v + v'$ where $v \in \mathbf{V}_s^{\gamma, N-1}$ and $v' \in (\mathbf{V}_s^{\gamma, N-1})^{\perp}$. Then, $(\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}}\gamma^{(N)}(s_0) = (\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}}v + v'$. Since $|(\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}}v| \leq_B \delta$ and $|v'| \leq B$, $|(\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}}\gamma^{(N)}(s_0)| \leq B + C\delta$ for some C = C(B). Thus, $\gamma = \gamma_{s_0}^{\delta}$ satisfies (2.1) with B replaced by 3B if $\delta < \delta'$. An elementary argument (elimination) shows

$$\operatorname{Vol}\left((\gamma_{s_0}^{\delta})^{(1)}(s), \dots, (\gamma_{s_0}^{\delta})^{(N)}(s)\right) = \operatorname{Vol}\left(\gamma^{(1)}(s_0), \dots, \gamma^{(N)}(s_0)\right) + O(\delta)$$

since $(\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}} \gamma^{(N)}(s_0) = (\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}} v + v'$ and $\gamma^{(N)}(s_0) = v + v'$. Taking δ' small enough, from $\mathfrak{V}(N, B)$ for γ we see $\mathfrak{V}(N, 3B)$ hold for $\gamma = \gamma_{s_0}^{\delta}$ if $0 < \delta < \delta'$.

The next lemma (cf. [14, Lemma 2.9]) plays a crucial role in what follows.

Lemma 2.8. Let $2 \leq N \leq d$, $\mathfrak{a} \in \mathfrak{A}_k(s_0, \delta)$, and $j_* = \log(2^k \delta^N)$. Suppose (2.20) holds on $\operatorname{supp}_{s,\xi} \mathfrak{a}$. Then, there exist constants $C, \tilde{B} \geq 1$, and $\delta' > 0$ depending on B, and symbols a_1, \ldots, a_{l_*} of type $(j, N - 1, \tilde{B})$ relative to $\gamma_{s_0}^{\delta}$, such that

$$\left\|\tilde{\chi} \,\mathcal{T}[\mathfrak{a}]f\right\|_{L^{p}(\mathbb{R}^{d+1})} \leq C\delta \sum_{1 \leq l \leq C} \left\|\mathcal{A}_{t}[\gamma_{s_{0}}^{\delta}, a_{l}]\tilde{f}_{l}\right\|_{L^{p}(\mathbb{R}^{d+1})},$$

 $\|\tilde{f}_l\|_p = \|f\|_p$, and $j \in [j_* - C, j_* + C]$ as long as $0 < \delta < \delta'$.

Proof. We set $\mathfrak{a}_{\delta,s_0}(s,t,\tau,\xi) = \mathfrak{a}(\delta s + s_0,t,\tau,\xi)$. Combining (2.13) and (2.14), we write $\mathcal{T}[\mathfrak{a}]f$ as an integral (e.g., see (2.17) and (2.18)). Then, the change of variables $s \to \delta s + s_0$ and $(\tau,\xi) \to (\tau - \gamma(s_0) \cdot \xi, \xi)$ gives

$$\mathcal{T}[\mathfrak{a}]f(x,t) = (2\pi)^{-d-1}\,\delta \iint e^{i\langle x-t\gamma(s_0),\,\xi\rangle}\mathcal{J}(s,t,\xi)\widehat{f}(\xi)\,dsd\xi,$$

where

$$\mathcal{J}(s,t,\xi) = \iint e^{it\tau} e^{-it'(\tau + (\gamma(\delta s + s_0) - \gamma(s_0))\cdot\xi)} \mathfrak{a}_{\delta,s_0}(s,t',\tau - \gamma(s_0)\cdot\xi,\xi) dt'd\tau.$$

Let \tilde{f} be given by $\mathcal{F}(\tilde{f}) = |\det \delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta}|^{1-1/p} \widehat{f}(\delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta} \cdot)$ where $\mathcal{F}(\tilde{f})$ denotes the Fourier transform of \tilde{f} . Then, $\|\tilde{f}\|_p = \|f\|_p$. Changing variables $\xi \to \delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta} \xi$ gives

$$\mathcal{T}[\mathfrak{a}]f(x,t) = C_d \iint e^{i\langle x - t\gamma(s_0), \delta^{-N}\widetilde{\mathcal{L}}_{s_0}^{\delta}\xi\rangle} \mathcal{J}(s,t,\delta^{-N}\widetilde{\mathcal{L}}_{s_0}^{\delta}\xi) \mathcal{F}(\tilde{f})(\xi) \, dsd\xi,$$

where $C_d = (2\pi)^{-d-1} \, \delta |\det \delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta}|^{1/p}$. This leads us to set

(2.22)
$$\tilde{a}(s,t,\xi) = \frac{1}{2\pi} \iint e^{-it'(\tau+\gamma_{s_0}^{\delta}(s)\cdot\xi)} \tilde{\chi}(t) \mathfrak{a}_{\delta,s_0} \left(s,t+t',\delta^{-N} \mathcal{L}_{s_0}^{\delta}(\tau,\xi)\right) dt' d\tau.$$

It is easy to check $\tilde{a} \in C^{d+N+2}(\mathbb{R}^{d+2})$, since so is \mathfrak{a} and $\gamma \in C^{3d+1}$. By (2.21) and (2.5), we note $\tilde{\chi}(t)\mathcal{J}(s,t,\delta^{-N}\widetilde{\mathcal{L}}_{s_0}^{\delta}\xi) = 2\pi e^{-it\gamma_{s_0}^{\delta}(s)\cdot\xi} \tilde{a}(s,t,\xi)$. Therefore,

$$\tilde{\chi}(t)\mathcal{T}[\mathfrak{a}]f(x,t) = \delta |\det \delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta}|^{\frac{1}{p}} \mathcal{A}_t[\gamma_{s_0}^{\delta}, \tilde{a}] \tilde{f}\left(\delta^{-N} (\widetilde{\mathcal{L}}_{s_0}^{\delta})^{\mathsf{T}} (x - t\gamma(s_0))\right),$$

10

and a change of variables gives

(2.23)
$$\left\| \tilde{\chi} \mathcal{T}[\mathfrak{a}] f \right\|_{L^{p}(\mathbb{R}^{d+1})} = \delta \left\| \mathcal{A}_{t}[\gamma_{s_{0}}^{\delta}, \tilde{a}] \tilde{f} \right\|_{L^{p}(\mathbb{R}^{d+1})}$$

We shall obtain symbols of type $(j, N - 1, \tilde{B})$ from \tilde{a} via decomposition and rescaling. To this end, we first note

(2.24)
$$\operatorname{supp}_{\xi} \tilde{a} \subset \left\{ \xi \in \mathbb{R}^d : C^{-1} \delta^N 2^k \le |\xi| \le C \delta^N 2^k \right\}$$

for a constant $C = C(B) \ge 1$. This follows by Lemma 2.5 since there exists τ such that $\delta^{-N} \mathcal{L}^{\delta}_{s_0}(\tau,\xi) \in \Lambda_k(s_0,\delta,B)$ if $\xi \in \operatorname{supp}_{\xi} \tilde{a}$. We claim

(2.25)
$$|\partial_s^j \partial_t^l \partial_{\xi}^{\alpha} \tilde{a}(s,t,\xi)| \lesssim_B |\xi|^{-|\alpha|}, \qquad (j,l,\alpha) \in \mathcal{I}_{N-1}.$$

To show (2.25), let us set

$$\mathfrak{b}(s,t,t',\tau,\xi) = \tilde{\chi}(t)\mathfrak{a}_{\delta,s_0}(s,t+t',\delta^{-N}\mathcal{L}^{\delta}_{s_0}(\tau,\xi)).$$

Note $0 \le j \le 1$. Taking derivatives on both sides of (2.22), we have

$$\partial_s^j \partial_t^l \partial_{\xi}^{\alpha} \tilde{a}(s,t,\xi) = \mathcal{I}[\mathfrak{b}_1] := \frac{1}{2\pi} \iint e^{-it'(\tau + \gamma_{s_0}^{\delta}(s) \cdot \xi)} \mathfrak{b}_1(s,t,t',\tau,\xi) \, dt' d\tau$$

where

$$\mathfrak{b}_1 = \sum_{\substack{u_1+u_2=j,\\\alpha_1+\alpha_2+\alpha_3=\alpha}} C_{\alpha,u} \left(t'\gamma_{s_0}^{\delta}{}'\cdot\xi\right)^{u_1-|\alpha_1|} (t'\gamma_{s_0}^{\delta}{}')^{\alpha_1} (t'\gamma_{s_0}^{\delta})^{\alpha_2} \,\partial_s^{u_2} \,\partial_t^l \partial_{\xi}^{\alpha_3} \mathfrak{b}$$

with $0 \le u_1 \le 1$, $0 \le |\alpha_1| \le u_1$, and constants $C_{\alpha,u}$ satisfying $|C_{\alpha,u}| = 1$. Integration by parts $u_1 + |\alpha_2|$ times in τ gives $\partial_s^j \partial_t^l \partial_{\xi}^{\alpha} \tilde{a} = \mathcal{I}[\mathfrak{b}_2]$, where

$$\mathfrak{b}_2 = \sum_{\substack{u_1+u_2=j,\\\alpha_1+\alpha_2+\alpha_3=\alpha}} C'_{\alpha,u} \left(\gamma_{s_0}^{\delta}\,'\cdot\xi\right)^{u_1-|\alpha_1|} (\gamma_{s_0}^{\delta}\,')^{\alpha_1} (\gamma_{s_0}^{\delta})^{\alpha_2}\,\partial_{\tau}^{u_1+|\alpha_2|}\partial_s^{u_2}\partial_t^l\partial_{\xi}^{\alpha_3}\mathfrak{k}^{\alpha_4}$$

with constants $C'_{\alpha,u}$ satisfying $|C'_{\alpha,u}| = 1$. We decompose $\mathcal{I}[\mathfrak{b}_2] = \mathcal{I}[\chi_E \mathfrak{b}_2] + \mathcal{I}[\chi_{E^c}\mathfrak{b}_2]$ where $E = \{(\tau,\xi) : |\tau + \gamma_{s_0}^{\delta}(s) \cdot \xi| \leq 1\}$. Then, integrating by parts in t' for $\mathcal{I}[\chi_{E^c}\mathfrak{b}_2]$, we obtain

$$|\mathcal{I}[\mathfrak{b}_2]| \lesssim \iint \chi_E|\mathfrak{b}_2| + \frac{\chi_{E^c}|\partial_{t'}^2\mathfrak{b}_2|}{|\tau + \gamma_{s_0}^{\delta}(s) \cdot \xi|^2} \, dt' d\tau.$$

Since $\mathfrak{a} \in \mathfrak{A}_k(s_0, \delta)$, $|\partial_s^{j'} \partial_t^{l'} \partial_{\tau,\xi}^{\alpha'} \mathfrak{b}| \lesssim_B |\xi|^{-|\alpha'|}$ for $(j', l', \alpha') \in \mathcal{I}_N$. It is also clear that $|\gamma_{s_0}^{\delta'}(s)| \lesssim 1$ if $\delta < \delta'$. Thus, $|\mathfrak{b}_2| = O(|\xi|^{-|\alpha|})$ and $|\partial_{t'}^2 \mathfrak{b}_2| = O(|\xi|^{-|\alpha|})$ if $l \leq 2(N-1)$. Since $\partial_s^j \partial_t^l \partial_{\xi}^{\alpha} \tilde{a} = \mathcal{I}[\mathfrak{b}_2]$, we get (2.25).

Now, we decompose \tilde{a} . Let $\tilde{\chi}_1, \tilde{\chi}_2$, and $\tilde{\chi}_3 \in C_c^{\infty}(\mathbb{R})$ such that $\tilde{\chi}_1 + \tilde{\chi}_2 + \tilde{\chi}_3 = 1$ on supp $\tilde{\chi}$ and supp $\tilde{\chi}_\ell \subset [2^{\ell-3}, 2^\ell]$. Also, let $\beta \in C_c^{\infty}((2^{-1}, 2))$ such that $\sum \beta(2^{-k} \cdot) = 1$ on \mathbb{R}_+ . Then, we set

$$a_{\ell,j}(s,t,\xi) = \tilde{\chi}_{\ell}(t)\beta(2^{-j}|\xi|)\tilde{a}(s,t,\xi),$$

so $\sum_{\ell,j} a_{\ell,j} = \tilde{a}$. By (2.24), $a_{\ell,j} = 0$ if $|j - j_*| > C$ for some C > 0.

Denoting $(a)_{\rho}(s,t,\xi) = a(s,\rho t,\rho^{-1}\xi)$, via rescaling we observe $\mathcal{A}_{\rho t}[\gamma_{s_0}^{\delta},a]g(x) = \mathcal{A}_t[\gamma_{s_0}^{\delta},(a)_{\rho}]g(\rho \cdot)(x/\rho)$. Thus, changes of variables yield

$$\|\mathcal{A}_t[\gamma_{s_0}^{\delta}, a_{\ell,j}]\tilde{f}\|_{L^p(\mathbb{R}^{d+1})} = 2^{(\ell-2)/p} \|\mathcal{A}_t[\gamma_{s_0}^{\delta}, (a_{\ell,j})_{2^{\ell-2}}]\tilde{f}_{\ell}\|_{L^p(\mathbb{R}^{d+1})},$$

where $\tilde{f}_{\ell} = 2^{(\ell-2)d/p} \tilde{f}(2^{\ell-2} \cdot)$. Since $\mathcal{A}_t[\gamma_{s_0}^{\delta}, \tilde{a}] = \sum_{\ell,j} \mathcal{A}_t[\gamma_{s_0}^{\delta}, a_{\ell,j}]$, by (2.23) we get

$$\left\|\tilde{\chi} \,\mathcal{T}[\mathfrak{a}]f\right\|_{L^{p}(\mathbb{R}^{d+1})} \lesssim \delta \sum_{\ell,j} \left\|\mathcal{A}_{t}[\gamma_{s_{0}}^{\delta},(a_{\ell,j})_{2^{\ell-2}}]f_{\ell}\right\|_{L^{p}(\mathbb{R}^{d+1})}.$$

To complete the proof, we only have to relabel $(a_{\ell,j})_{2^{\ell-2}}$, $\ell = 1, 2, 3, j_* - C \leq j \leq j_* + C$. Indeed, since $\tilde{a} \in C^{d+N+2}$, $(a_{\ell,j})_{2^{\ell-2}} \in C^{d+N+2}$, which is supported in $I \times [2^{-1}, 4] \times \mathbb{A}_{j+\ell-2}$. Obviously, (2.25) holds for $\tilde{a} = (a_{\ell,j})_{2^{\ell-2}}$ because $\ell = 1, 2, 3$. Changing variables $s \to \delta s + s_0$ and $\xi \to \delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta} \xi$ in (2.20), by (2.21) we see that (2.20) on $\operatorname{supp}_{s,\xi} \mathfrak{a}$ is equivalent to $\sum_{j=1}^{N-1} |\langle (\gamma_{s_0}^{\delta})^{(j)}(s), \xi \rangle| \geq 2^k \delta^N / B'$ for $(s,\xi) \in \operatorname{supp}_{s,\xi} \mathfrak{a}_{\delta,s_0}(\cdot, \delta^{-N} \mathcal{L}_{s_0}^{\delta} \cdot)$. Note $\operatorname{supp}_{s,\xi} \mathfrak{a}_{\delta,s_0}(\cdot, \delta^{-N} \mathcal{L}_{s_0}^{\delta} \cdot) \supset \operatorname{supp}_{s,\xi} \widetilde{a}$. So, the same holds on $\operatorname{supp}_{s,\xi} \widetilde{a}$ and hence on $\operatorname{supp}_{s,\xi} (a_{\ell,j})_{2^{\ell-2}}$ if B' replaced by 2B'. Therefore, $C^{-1}(a_{\ell,j})_{2^{\ell-2}}$ is of type $(j + \ell - 2, N - 1, \widetilde{B})$ relative to $\gamma_{s_0}^{\delta}$ for a large constant C = C(B).

2.3. **Preliminary decomposition and reduction.** For the proof of Proposition 2.3, we make some reductions by decomposing the symbol *a*. We fix a sufficiently small positive constant

$$\delta_* \le \min\{\delta', (2^{7d}B^6)^{-N}\},\$$

which is to be specified in what follows. Here δ' is the number given in Lemma 2.8.

We recall that γ satisfies (2.1), $\mathfrak{N}(N, B)$, $\mathfrak{V}(N, B)$, and a is of type (k, N, B) relative to γ . We set

(2.26)
$$\eta_N(s,\xi) = \prod_{1 \le j \le N-1} \beta_0 \Big(B 2^{-k-1} \delta_*^{j-N} \langle \gamma^{(j)}(s), \xi \rangle \Big),$$

where $\beta_0 \in C_c^{\infty}((-1,1))$ such that $\beta_0 = 1$ on [-1/2, 1/2]. It is easy to see $|\partial_s^j \partial_t^l \partial_{\xi}^\alpha(a\eta_N)| \leq C |\xi|^{-|\alpha|}$ for $(j,l,\alpha) \in \mathcal{I}_N$, and the same holds for $a(1-\eta_N)$.

Note $\sum_{j=1}^{N-1} |\gamma^{(j)}(s) \cdot \xi| \ge (2B)^{-1} \delta_*^N |\xi|$ on $\operatorname{supp}_{s,\xi}(a(1-\eta_N))$. So, we see $a(1-\eta_N)$ is a symbol of type (k, N-1, B') for $B' = CB^2 \delta_*^{-C}$ with a large C. Applying the assumption (Theorem 2.2 with L = N - 1 and B = B'), we obtain

$$\|\mathcal{A}_t[\gamma, a(1-\eta_N)]f\|_{L^p(\mathbb{R}^{d+1})} \le C2^{(-\frac{2}{p}+\epsilon)k} \|f\|_{L^p(\mathbb{R}^d)}, \quad p \ge 4N-6$$

Thus, it suffices to consider $\mathcal{A}_t[\gamma, a\eta_N]$. Since $\mathfrak{N}(N, B)$ holds on $\operatorname{supp}_{s,\xi} \mathfrak{a}$,

(2.27)
$$|\gamma^{(N)}(s) \cdot \xi| \ge (2B)^{-1}|\xi|$$

whenever $(s, t, \xi) \in \operatorname{supp} a\eta_N$ for some t.

Basic assumption Before we continue to prove the estimate for $\mathcal{A}_t[\gamma, a\eta_N]$, we make several assumptions which are clearly permissible by elementary decompositions.

Decomposing a we may assume that $\operatorname{supp}_{\xi} a$ is contained in a narrow conic neighborhood and $\operatorname{supp}_{s} a \subset I(s_0, \delta_*)$ for some s_0 . Let us set

$$\Gamma_k = \left\{ \xi \in \mathbb{A}_k : \operatorname{dist} \left(|\xi|^{-1} \xi, |\xi'|^{-1} \xi' \right) < \delta_* \text{ for some } \xi' \in \operatorname{supp}_{\xi}(a\eta_N) \right\}.$$

We may also assume $\gamma^{(N-1)}(s') \cdot \xi' = 0$ for some $(s', \xi') \in I(s_0, \delta_*) \times \Gamma_k$. Otherwise, $|\gamma^{(N-1)}(s) \cdot \xi| \gtrsim |\xi|$ on $\operatorname{supp}_{s,\xi} a\eta_N$ and hence $a\eta_N = 0$ if we take *B* large enough. By (2.27) and the implicit function theorem, there exists σ such that

(2.28)
$$\gamma^{(N-1)}(\sigma(\xi)) \cdot \xi = 0$$

in a narrow conic neighborhood of ξ' where $\gamma \in C^{2d+2}$ since $\gamma \in C^{3d+1}(I)$. So, decomposing a further, we may assume $\sigma \in C^{2d+2}(\Gamma_k)$ and $\sigma(\xi) \in I(s_0, \delta_*)$ for $\xi \in \Gamma_k$. Furthermore, since σ is homogeneous of degree zero,

(2.29)
$$|\partial_{\xi}^{\alpha}\sigma(\xi)| \le C|\xi|^{-|\alpha|}, \qquad \xi \in \Gamma_k$$

for a constant C = C(B) if $|\alpha| \le 2d+2$. Any symbol which appears in what follows is to be given by decomposing the symbol *a* with appropriate cutoff functions. So, the s, ξ -supports of the symbols are assumed to be contained in $I(s_0, \delta_*) \times \Gamma_k$.

We break a to have further localization on the Fourier side. Let

$$\mathfrak{a}_1(s,t,\tau,\xi) = a\eta_N \,\beta_0 \left(2^{-2k} \delta_*^{-2N} |\tau + \langle \gamma(s),\xi \rangle|^2 \right)$$

and $\mathfrak{a}_0 = a\eta_N - \mathfrak{a}_1$. Then, by Fourier inversion

$$\mathcal{A}_t[\gamma, a\eta_N]f = \mathcal{T}[\mathfrak{a}_1]f + \mathcal{T}[\mathfrak{a}_0]f.$$

It is easy to show $\|\mathcal{T}[\mathfrak{a}_0]f\|_p \lesssim_B 2^{-2k} \|f\|_p$ for $1 \leq p \leq \infty$. Indeed, consider $\tilde{\mathfrak{a}}_0 = -(\tau + \gamma(s) \cdot \xi)^{-2} \partial_t^2 \mathfrak{a}_0$. By (2.13) and integration by parts in t', $m[\mathfrak{a}_0] = m[\tilde{\mathfrak{a}}_0]$ and hence $\mathcal{T}[\mathfrak{a}_0] = \mathcal{T}[\tilde{\mathfrak{a}}_0]$. Thanks to (2.17), it is sufficient to show

$$|K[\tilde{\mathfrak{a}}_0](s,t,x)| \le C \, 2^{k(d-1)} \int \left(1 + 2^k |t - t'| + 2^k |x - t'\gamma(s)|\right)^{-d-3} dt'$$

for a constant $C = C(B, \delta_*)$. Note $|\tau + \langle \gamma(s), \xi \rangle| \gtrsim 2^k$ on supp $\tilde{\mathfrak{a}}_0$, and recall (2.18). Rescaling and integration by parts in τ, ξ , as in the proof of Lemma 2.7, show the estimate.

The difficult part is to estimate $\mathcal{T}[\mathfrak{a}_1]$. Since δ_* is a fixed constant, it is obvious that $C^{-1}\mathfrak{a}_1 \in \mathfrak{A}_k(s_0, \delta_*)$ for some $C = C(B, \delta_*)$. So, the desired estimate for $\mathcal{T}[\mathfrak{a}_1]$ follows once we have the next proposition.

Proposition 2.9. Let $\mathfrak{a} \in \mathfrak{A}_k(s_0, \delta_*)$ with $\operatorname{supp}_{\xi} \mathfrak{a} \subset \Gamma_k$. Suppose Theorem 2.2 holds for L = N - 1. Then, if $p \ge 4N - 2$, for $\epsilon > 0$ we have

(2.30)
$$\left\|\mathcal{T}[\mathfrak{a}]f\right\|_{L^p(\mathbb{R}^{d+1})} \le C_{\epsilon} 2^{-\frac{2}{p}k+\epsilon k} \|f\|_{L^p(\mathbb{R}^d)}.$$

Therefore, the proof of Proposition 2.3 is completed if we prove Proposition 2.9. For the purpose, we use Proposition 2.10 below, which allows us to decompose $\mathcal{T}[\mathfrak{a}]$ into operators given by symbols with smaller *s*-supports while the consequent minor parts have acceptable bounds. This type of argument was used in [24] when L = 2.

Let δ_0 and δ_1 be positive numbers such that

(2.31)
$$2^{7d}B^6\delta_0^{(N+1)/N} \le \delta_1 \le \delta_0 \le \delta_*, \quad 2^{-k/N} \le \delta_1.$$

Then, it is clear that

(

2.32)
$$B^{6N}\delta_0^{j+1} \le 2^{-7dN}\delta_1^j, \qquad j = 1, \dots, N.$$

For $n \ge 0$, we denote $\mathfrak{J}_n^{\mu} = \{ \nu \in \mathbb{Z} : |2^n \delta_1 \nu - \delta_0 \mu| \le \delta_0 \}.$

Proposition 2.10. For μ such that $\delta_0 \mu \in I(s_0, \delta_*) \cap \delta_0 \mathbb{Z}$, let $\mathfrak{a}^{\mu} \in \mathfrak{A}_k(\delta_0 \mu, \delta_0)$ with $\sup_{s,\xi} \mathfrak{a}^{\mu} \subset I(s_0, \delta_*) \times \Gamma_k$. Suppose Theorem 2.2 holds for L = N - 1. Then, if $p \geq 4N - 2$, for $\epsilon > 0$ there exist a constant $C_{\epsilon} = C_{\epsilon}(B) \geq 2$ and symbols $\mathfrak{a}_{\nu} \in \mathfrak{A}_k(\delta_1 \nu, \delta_1)$ with $\operatorname{supp}_{s,\xi} \mathfrak{a}_{\nu} \subset I(s_0, \delta_*) \times \Gamma_k$, $\nu \in \bigcup_{\mu} \mathfrak{J}_0^{\mu}$, such that

$$\left(\sum_{\mu} \|\mathcal{T}[\mathfrak{a}^{\mu}]f\|_{p}^{p}\right)^{\frac{1}{p}} \leq C_{\epsilon} \left(\delta_{1}/\delta_{0}\right)^{\frac{2N}{p}-1-\epsilon} \left(\sum_{\nu} \|\mathcal{T}[\mathfrak{a}_{\nu}]f\|_{p}^{p}\right)^{\frac{1}{p}} + C_{\epsilon} \delta_{0}^{-\frac{2N}{p}+1+\epsilon} 2^{-\frac{2}{p}k+2\epsilon k} \|f\|_{p}$$

Assuming Proposition 2.10, we prove Proposition 2.9.

2.4. **Proof of Proposition 2.9.** Let $\mathfrak{a} \in \mathfrak{A}_k(s_0, \delta_*)$. We may assume $s_0 = \delta_* \mu$ for some $\mu \in \mathbb{Z}$. To apply Proposition 2.10 iteratively, we need to choose an appropriate decreasing sequence of positive numbers since the decomposition is subject to the condition (2.31).

Let $\delta_0 = \delta_*$, so $(2^{7d}B^6)^N \delta_0 < 1$. Let J be the largest integer such that

$$(2^{7d}B^6)^{N(\frac{N+1}{N})^{J-1}-N}\delta_0^{(\frac{N+1}{N})^{J-1}} > 2^{-\frac{k}{N}}.$$

So, $J \leq C_1 \log k$ for a constant $C_1 \geq 1$. We set

(2.33)
$$\delta_J = 2^{-\frac{k}{N}}, \qquad \delta_j = (2^{7d} B^6)^{N(\frac{N+1}{N})^j - N} \delta_0^{(\frac{N+1}{N})^j}$$

for $j = 1, \ldots, J - 1$. Thus, it follows that

(2.34)
$$2^{7d}B^6\delta_j^{(N+1)/N} \le \delta_{j+1} < \delta_j, \qquad j = 0, \dots, J-1.$$

For a given $\epsilon > 0$, let $\tilde{\epsilon} = \epsilon/4$. Since $\mathfrak{a} \in \mathfrak{A}_k(\delta_0 \mu, \delta_0)$ and (2.31) holds for δ_0 and δ_1 , applying Proposition 2.10 to $\mathcal{T}[\mathfrak{a}]$, we have

$$\|\mathcal{T}[\mathfrak{a}]f\|_{p} \leq C_{\tilde{\epsilon}} (\delta_{1}/\delta_{0})^{\frac{2N}{p}-1-\tilde{\epsilon}} (\sum_{\nu_{1}} \|\mathcal{T}[\mathfrak{a}_{\nu_{1}}]f\|_{p}^{p})^{\frac{1}{p}} + C_{\tilde{\epsilon}} \delta_{0}^{-\frac{2N}{p}+1+\tilde{\epsilon}} 2^{-\frac{2}{p}k+2\tilde{\epsilon}k} \|f\|_{p},$$

where $\mathfrak{a}_{\nu_1} \in \mathfrak{A}_k(\delta_1\nu_1, \delta_1), \nu_1 \in \mathfrak{J}_0^{\mu}$. Thanks to (2.34) we may apply again Proposition 2.10 to $\mathcal{T}[\mathfrak{a}_{\nu_1}]$ while δ_0, δ_1 replaced by δ_1, δ_2 , respectively. Repeating this procedure up to *J*-th step yields symbols $\mathfrak{a}_{\nu} \in \mathfrak{A}_k(\delta_J\nu, \delta_J), \delta_J\nu \in \delta_J\mathbb{Z} \cap I(\delta_0\mu, \delta_0)$, such that

$$\|\mathcal{T}[\mathfrak{a}]f\|_{p} \leq C_{\tilde{\epsilon}}^{J} \delta_{J}^{\frac{2N}{p}-1-\tilde{\epsilon}} \Big(\sum_{\nu} \|\mathcal{T}[\mathfrak{a}_{\nu}]f\|_{p}^{p}\Big)^{\frac{1}{p}} + \sum_{0 \leq j \leq J-1} C_{\tilde{\epsilon}}^{j+1} \delta_{0}^{-\frac{2N}{p}+1+\tilde{\epsilon}} 2^{-\frac{2}{p}k+2\tilde{\epsilon}k} \|f\|_{p}$$

for $p \ge 4N - 2$. Now, assuming

(2.35)
$$\left(\sum_{\nu} \|\mathcal{T}[\mathfrak{a}_{\nu}]f\|_{p}^{p}\right)^{1/p} \lesssim_{B} 2^{-k/N} \|f\|_{p}, \qquad 2 \le p \le \infty$$

for the moment, we can finish the proof of Proposition 2.9. Since $C_{\tilde{\epsilon}} \geq 2$, combining the above inequalities, we get

$$\|\mathcal{T}[\mathfrak{a}]f\|_p \lesssim_B C_{\hat{\epsilon}}^{J+1} \left(2^{-\frac{2}{p}k + \frac{\tilde{\epsilon}}{N}k} + 2^{-\frac{2}{p}k + 2\tilde{\epsilon}k}\right) \|f\|_p.$$

Note $J \leq C_1 \log k$, so $C_{\tilde{\epsilon}}^{J+1} \leq C' 2^{\epsilon k/2}$ for some C' if k is sufficiently large. Thus, the right hand side is bounded by $C 2^{-2k/p+\epsilon k} ||f||_p$.

It remains to show (2.35) for $2 \leq p \leq \infty$. By interpolation, it is enough to obtain (2.35) for $p = \infty$ and p = 2. The case $p = \infty$ follows by (2.15) since $\mathfrak{a}_{\nu} \in \mathfrak{A}_k(\delta_J \nu, \delta_J)$. So, we need only to prove (2.35) for p = 2. To do this, we first observe the following, which shows $\operatorname{supp}_{\ell} \mathfrak{a}_{\nu}$ are finitely overlapping.

Lemma 2.11. For $b \ge 1$, $s \in I(s_0, \delta_*)$, and $0 < \delta \le \delta_*$, let us set

(2.36)
$$\Lambda'_k(s,\delta,b) = \bigcap_{1 \le j \le N-1} \left\{ \xi \in \Gamma_k : |\langle \gamma^{(j)}(s), \xi \rangle| \le b 2^k \delta^{N-j} \right\}.$$

If $\Lambda'_k(s_1, \delta, b) \cap \Lambda'_k(s_2, \delta, b) \neq \emptyset$ for some $s_1, s_2 \in I(s_0, \delta_*)$, then there is a constant C = C(B) such that $|s_1 - s_2| \leq Cb\delta$.

Proof. Let $\xi \in \Lambda'_k(s_1, \delta, b) \cap \Lambda'_k(s_2, \delta, b)$. Since $|\gamma^{(N-1)}(s_j) \cdot \xi| \leq b2^k \delta$, j = 1, 2, by (2.28) and (2.27) we see $|s_j - \sigma(\xi)| \leq 2^2 bB\delta$, j = 1, 2, using the mean value theorem. This implies $|s_1 - s_2| \leq 2^3 bB\delta$.

We recall (2.13). Since (2.27) holds on $\operatorname{supp}_{s,\xi} \mathfrak{a}_{\nu}$, by van der Corput's lemma (e.g., see [32, Corollary, p. 334]) we have

$$|m[\mathfrak{a}_{\nu}](\tau,\xi)| \lesssim 2^{-k/N} \left(\|\mathfrak{a}_{\nu}(\cdot,t,\tau,\xi)\|_{\infty} + \|\partial_{s}\mathfrak{a}_{\nu}(\cdot,t,\tau,\xi)\|_{1} \right) \lesssim_{B} 2^{-k/N}$$

The second inequality is clear since $\mathfrak{a}_{\nu} \in \mathfrak{A}_k(\delta_J\nu, \delta_J)$. From (2.14) note $\mathcal{F}(\mathcal{T}[\mathfrak{a}_{\nu}]f) = m[\mathfrak{a}_{\nu}]\widehat{f}$. Since $\operatorname{supp}_{\xi}\mathfrak{a}_{\nu} \subset S_{\nu} := \Lambda'_k(\delta_J\nu, \delta_J, 2^5B)$, $\operatorname{supp}_{\xi}\mathcal{F}(\mathcal{T}[\mathfrak{a}_{\nu}]f) \subset S_{\nu}$ by (2.13). By Lemma 2.11 it follows that the sets S_{ν} overlap at most C = C(B) times. Therefore, Plancherel's theorem and the estimate above yield

$$\|\sum_{\nu} \mathcal{T}[\mathfrak{a}_{\nu}]f\|_{2}^{2} \lesssim_{B} 2^{-2k/N} \sum_{\nu} \int_{S_{\nu}} \int_{\{\tau: |\tau+\gamma(\delta_{J}\nu)\cdot\xi| \le 2^{5}B\}} d\tau \, |\widehat{f}(\xi)|^{2} \, d\xi$$

since supp $\mathfrak{a}_{\nu} \subset \Lambda_k(\delta_J \nu, \delta_J, B)$. This gives (2.35) for p = 2.

2.5. Decoupling inequalities. We denote $\mathbf{r}_{\circ}^{N}(s) = (s, s^{2}/2!, \dots, s^{N}/N!)$, and consider a collection of curves from I to \mathbb{R}^{N} which are small perturbations of \mathbf{r}_{\circ}^{N} :

$$\mathfrak{C}(\epsilon_{\circ}; N) := \{ \mathbf{r} \in \mathbf{C}^{2N+1}(I) : \| \mathbf{r} - \mathbf{r}_{\circ}^{N} \|_{\mathbf{C}^{2N+1}(I)} < \epsilon_{\circ} \}.$$

For $\mathbf{r} \in \mathfrak{C}(\epsilon_{\circ}; N)$ and $s \in I$, we define

$$\mathcal{N}_{\mathbf{r}}(s,\delta) = \Big\{ \mathbf{r}(s) + \sum_{1 \le j \le N} u_j \mathbf{r}^{(j)}(s) : |u_j| \le \delta^j, \ j = 1, \dots, N \Big\}.$$

Let $s_1, \ldots, s_l \in I$ be δ -separated points, i.e., $|s_n - s_j| \geq \delta$ if $n \neq j$, such that $\bigcup_{i=1}^l (s_j - \delta, s_j + \delta) \supset I$. Then, we set

$$\theta_j = \mathcal{N}_{\mathbf{r}}(s_j, \delta), \qquad 1 \le j \le l.$$

The following is due to Bourgain, Demeter, and Guth [5] (also see [10]).

Theorem 2.12. Let $0 < \delta \ll 1$. Suppose $\mathbf{r} \in \mathfrak{C}(\epsilon_{\circ}; N)$ for a small enough $\epsilon_{\circ} > 0$. Then, if $2 \le p \le N(N+1)$, for $\epsilon > 0$ we have

(2.37)
$$\| \sum_{1 \le j \le l} f_j \|_{L^p(\mathbb{R}^N)} \le C_{\epsilon} \delta^{-\epsilon} \Big(\sum_{1 \le j \le l} \| f_j \|_{L^p(\mathbb{R}^N)}^2 \Big)^{1/2}$$

whenever supp $\widehat{f}_j \subset \theta_j$ for $1 \leq j \leq l$.

The constant C_{ϵ} can be taken to be independent of particular choices of the δ -separated points s_1, \ldots, s_l . One can obtain a conical extension of the inequality (2.37) by modifying the argument in [4] which deduces the decoupling inequality for the cone from that for the paraboloid (see [2, Proposition 7.7]). Let us consider conical sets

$$\bar{\theta}_j = \{(\eta, \rho) \in \mathbb{R}^N \times [1, 2] : \eta/\rho \in \theta_j\}, \quad 1 \le j \le l.$$

Corollary 2.13. Let $0 < \delta \leq 1$ and let $\mathbf{r} \in \mathfrak{C}(\epsilon_{\circ}; N)$ with a small enough $\epsilon_{\circ} > 0$. Then, if $2 \leq p \leq N(N+1)$, for $\epsilon > 0$ we have

(2.38)
$$\| \sum_{1 \le j \le l} F_j \|_{L^p(\mathbb{R}^{N+1})} \le C_\epsilon \delta^{-\epsilon} \Big(\sum_{1 \le j \le l} \|F_j\|_{L^p(\mathbb{R}^{N+1})}^2 \Big)^{1/2}$$

whenever supp $\widehat{F}_j \subset \overline{\theta}_j$ for $1 \leq j \leq l$.

The decoupling inequality (2.38) does not fit the symbols which appear later when we decompose \mathfrak{a} (see Section 3.1 and Section 4.2). As to be seen later, those symbols are related to the slabs of the following form.

Definition 2.14. Let $N \geq 2$ and $\tilde{\mathbf{r}} \in \mathfrak{C}(\epsilon_{\circ}; N + 1)$. For $s \in I$, we denote by $\mathbf{s}(s, \delta, \rho; \tilde{\mathbf{r}})$ the set of $(\tau, \eta) \in \mathbb{R} \times \mathbb{R}^{N}$ which satisfies

$$\rho^{-1} \le |\langle \tilde{\mathbf{r}}^{(N+1)}(s), (\tau, \eta) \rangle| \le 2\rho,$$
$$|\langle \tilde{\mathbf{r}}^{(j)}(s), (\tau, \eta) \rangle| \le \delta^{N+1-j}, \qquad j = N, \dots, 1,$$

The same form of decoupling inequality continues to be valid for the slabs $\mathbf{s}(s_1, \delta, 1; \tilde{\mathbf{r}}), \ldots, \mathbf{s}(s_l, \delta, 1; \tilde{\mathbf{r}})$. Beltran et al. [2, Theroem 4.4] showed, using the Frenet–Serret formulas, that those slabs can be generated by conical extensions of the slabs given by a nondegenerate curve in \mathbb{R}^N . The following is a consequence of Corollary 2.13 and a simple manipulation using decomposition and rescaling.

Corollary 2.15. Let $0 < \delta \leq 1$, $\rho \geq 1$, and $\tilde{\mathbf{r}} \in \mathfrak{C}(\epsilon_{\circ}; N+1)$ for a small enough $\epsilon_{\circ} > 0$. Denote $\mathbf{s}_{j} = \mathbf{s}(s_{j}, \delta, \rho; \tilde{\mathbf{r}})$ for $1 \leq j \leq l$. Then, if $2 \leq p \leq N(N+1)$, for $\epsilon > 0$ there is a constant $C_{\epsilon} = C_{\epsilon}(\rho)$ such that

(2.39)
$$\| \sum_{1 \le j \le l} F_j \|_{L^p(\mathbb{R}^{N+1})} \le C_\epsilon \delta^{-\epsilon} \Big(\sum_{1 \le j \le l} \|F_j\|_{L^p(\mathbb{R}^{N+1})}^2 \Big)^{1/2}$$

whenever $\operatorname{supp} \widehat{F_j} \subset \mathbf{s}_j$ for $1 \leq j \leq l$.

For our purpose of proving Proposition 2.10, we use a modified form. If $p_* \in [2, N(N+1)]$, then we have

$$\Big\| \sum_{1 \le j \le l} F_j \Big\|_{L^p(\mathbb{R}^{N+1})} \le C_\epsilon \delta^{-1 + \frac{2+p_*}{2p} - \epsilon} \Big(\sum_{1 \le j \le l} \|F_j\|_{L^p(\mathbb{R}^{N+1})}^p \Big)^{1/p}$$

for $p \ge p_*$. The case $p = p_*$ follows by (2.39) and Hölder's inequality. Interpolation with the trivial $\ell^{\infty}L^{\infty}-L^{\infty}$ estimate gives the estimate for $p > p_*$. One may choose different p_* for the particular purposes. In fact, for the local smoothing estimate we take $p_* = 4N - 2$ to have

(2.40)
$$\left\| \sum_{1 \le j \le l} F_j \right\|_{L^p(\mathbb{R}^{N+1})} \le C_{\epsilon} \delta^{-1 + \frac{2N}{p} - \epsilon} \Big(\sum_{1 \le j \le l} \|F_j\|_{L^p(\mathbb{R}^{N+1})}^p \Big)^{1/p} \right\|_{L^p(\mathbb{R}^{N+1})}^{1/p}$$

for $p \ge 4N-2$ (see Section 3.2). For the L^p Sobolev regularity estimate, we observe

(2.41)
$$\|\sum_{1 \le j \le l} F_j\|_{L^p(\mathbb{R}^{N+1})} \le C_{\epsilon_0} \delta^{-1+\frac{N+1}{p}+\epsilon_0} \Big(\sum_{1 \le j \le l} \|F_j\|_{L^p(\mathbb{R}^{N+1})}^p\Big)^{1/p}$$

holds for some $\epsilon_0 = \epsilon_0(p) > 0$ if $2N . Indeed, we need only to take <math>p_* > 2N$ close enough to 2N. The presence of ϵ_0 in (2.41) is crucial for proving the optimal Sobolev regularity estimate (see Proposition 4.5).

The inequalities (2.40) and (2.41) obviously extend to cylindrical forms via the Minkowski inequality. For example, set $\tilde{\mathbf{s}}_j = \{(\xi, \eta) \in \mathbb{R}^{N+1} \times \mathbb{R}^M : \xi \in \mathbf{s}_j\}$ for $1 \leq j \leq l$. Then, using (2.40), we have

(2.42)
$$\|\sum_{1 \le j \le l} G_j\|_{L^p(\mathbb{R}^{N+M+1})} \le C_\epsilon \delta^{-1+\frac{2N}{p}-\epsilon} \Big(\sum_{1 \le j \le l} \|G_j\|_{L^p(\mathbb{R}^{N+M+1})}^2\Big)^{1/2}$$

whenever \widehat{G}_j is supported in $\widetilde{\mathbf{s}}_j$. Clearly, we also have a similar extension of (2.41).

3. Decomposition of the symbols

In this section, we prove Proposition 2.10 by applying the decoupling inequality. Meanwhile, the induction assumption (Theorem 2.2 with L = N - 1) plays an important role. We decompose a given symbol $\mathfrak{a}^{\mu} \in \mathfrak{A}_k(\delta_0\mu, \delta_0)$ into the symbols with their *s*-supports contained in intervals of length about δ_1 while the consequent minor contribution is controlled within an acceptable bound. To achieve it up to δ_1 satisfying (2.31), we approximate $\langle G(s), (\tau, \xi) \rangle$ in a local coordinate system near the set $\{(s, \xi) : \langle \gamma^{(N-1)}(s), \xi \rangle = 0\}$.

3.1. Decomposition of the symbol \mathfrak{a}^{μ} . We begin by introducing some notations. Fixing $\mu \in \mathbb{Z}$ such that $\delta_0 \mu \in I(s_0, \delta_*)$, we consider linear maps

$$y^j_{\mu}(\tau,\xi) = \langle G^{(j)}(\delta_0\mu), (\tau,\xi) \rangle, \qquad j = 0, 1, \dots, N.$$

In particular, $y^j_{\mu}(\tau,\xi) = \langle \gamma^{(j)}(\delta_0 \mu), \xi \rangle$ if $1 \le j \le N$. By (2.27) it follows that

(3.1)
$$|y^N_{\mu}(\tau,\xi)| \ge (2B)^{-1}|\xi|.$$

We denote

$$\omega_{\mu}(\xi) = rac{y_{\mu}^{N-1}(au,\xi)}{y_{\mu}^{N}(au,\xi)},$$

which is close to $\delta_0 \mu - \sigma(\xi)$ (see (3.5)). Then, we define $\mathfrak{g}^N_\mu, \mathfrak{g}^{N-1}_\mu, \ldots, \mathfrak{g}^0_\mu$ recursively, by setting $\mathfrak{g}^N_\mu = y^N_\mu$, and

(3.2)
$$\mathfrak{g}_{\mu}^{j}(\tau,\xi) = y_{\mu}^{j}(\tau,\xi) - \sum_{\ell=j+1}^{N} \frac{\mathfrak{g}_{\mu}^{\ell}(\tau,\xi)}{(\ell-j)!} (\omega_{\mu}(\xi))^{\ell-j}, \quad j = N-1,\ldots,0.$$

Note that $\mathfrak{g}_{\mu}^{N-1} = 0$ and (3.2) can be rewritten as follows:

(3.3)
$$y_{\mu}^{m}(\tau,\xi) = \sum_{\ell=m}^{N} \frac{\mathfrak{g}_{\mu}^{\ell}(\tau,\xi)}{(\ell-m)!} (\omega_{\mu}(\xi))^{\ell-m}, \quad m = 0, \dots, N.$$

The identity continues to hold for m = N since $\mathfrak{g}^N_\mu = y^N_\mu$. Apparently, $\mathfrak{g}^1_\mu, \ldots, \mathfrak{g}^N_\mu$ are independent of τ since so are y^1_μ, \ldots, y^N_μ .

For $j = 1, \ldots, N$, set

(3.4)
$$\mathcal{E}_{j}(\xi) := (y_{\mu}^{N}(\tau,\xi))^{-1} \int_{\sigma(\xi)}^{\delta_{0}\mu} \frac{\langle \gamma^{(N+1)}(r), \xi \rangle}{j!} (\sigma(\xi) - r)^{j} dr.$$

By (3.4) with j = 1 and integration by parts, we have

(3.5)
$$\mathcal{E}_1(\xi) = \sigma(\xi) - \delta_0 \mu + \omega_\mu(\xi).$$

Lemma 3.1. For $0 \le j \le N - 1$, we have

(3.6)
$$\langle G^{(j)}(\sigma(\xi)), (\tau,\xi) \rangle = \sum_{\ell=j}^{N} \frac{\mathfrak{g}_{\mu}^{\ell} (\mathcal{E}_{1})^{\ell-j}}{(\ell-j)!} - y_{\mu}^{N} \mathcal{E}_{N-j}.$$

Proof. When j = N - 1, (3.6) is clear. To show (3.6) for j = 0, 1, ..., N - 2, by Taylor's theorem with integral remainder we have

$$\langle G^{(j)}(\sigma(\xi)), (\tau, \xi) \rangle = \sum_{m=j}^{N} y^m_\mu(\tau, \xi) \frac{(\sigma(\xi) - \delta_0 \mu)^{m-j}}{(m-j)!} - y^N_\mu(\tau, \xi) \mathcal{E}_{N-j}(\xi).$$

Using (3.3) and then changing the order of the sums, we see

$$\langle G^{(j)}(\sigma(\xi)), (\tau, \xi) \rangle = \sum_{\ell=j}^{N} \mathfrak{g}^{\ell}_{\mu} \Big(\sum_{m=j}^{\ell} \frac{(\sigma(\xi) - \delta_0 \mu)^{m-j}}{(\ell-m)!(m-j)!} (\omega_{\mu})^{\ell-m} \Big) - y^N_{\mu} \mathcal{E}_{N-j}.$$

The sum over *m* equals $(\sigma(\xi) - \delta_0 \mu + \omega_\mu)^{\ell-j} / (\ell-j)!$. So, (3.6) follows by (3.5). \Box

We now decompose the symbol $\mathfrak{a}^{\mu} \in \mathfrak{A}_k(\delta_0\mu, \delta_0)$ by making use of \mathfrak{g}^j_{μ} , $j = 0, \ldots, N-2$. We define

(3.7)
$$\mathfrak{G}_{N}^{\mu}(s,\tau,\xi) = \sum_{j=0}^{N-2} \left(2^{-k} \mathfrak{g}_{\mu}^{j}(\tau,\xi) \right)^{\frac{2N!}{N-j}} + (s - \sigma(\xi))^{2N!}.$$

Let $\beta_N = \beta_0 - \beta_0(2^{2N!} \cdot)$, so $\sum_{\ell \in \mathbb{Z}} \beta_N(2^{2N!\ell} \cdot) = 1$. We also take $\zeta \in C_c^{\infty}((-1, 1))$ such that $\sum_{\nu \in \mathbb{Z}} \zeta(\cdot - \nu) = 1$. For $n \ge 0$ and $\nu \in \mathfrak{J}_n^{\mu}$, we set

$$\mathfrak{a}_{\nu}^{\mu,n} = \mathfrak{a}^{\mu} \times \begin{cases} \beta_0 \left(\delta_1^{-2N!} \mathfrak{G}_N^{\mu} \right) \zeta(\delta_1^{-1} s - \nu), & n = 0, \\ \beta_N \left((2^n \delta_1)^{-2N!} \mathfrak{G}_N^{\mu} \right) \zeta(2^{-n} \delta_1^{-1} s - \nu), & n \ge 1. \end{cases}$$

Then, it follows that

(3.8)
$$\mathfrak{a}^{\mu} = \sum_{n \ge 0} \sum_{\nu \in \mathfrak{J}_n^{\mu}} \mathfrak{a}_{\nu}^{\mu, n}$$

Lemma 3.2. There is a constant C = C(B) such that $C^{-1}\mathfrak{a}_{\nu}^{\mu,n} \in \mathfrak{A}_{k}(2^{n}\delta_{1}\nu, 2^{n}\delta_{1})$ for $n \geq 0$, μ , and ν .

The proof of Lemma 3.2 is elementary though it is somewhat involved. We postpone the proof until Section 3.3.

We collect some elementary facts regarding $\mathfrak{a}_{\nu}^{\mu,n}$. First, we may assume

$$(3.9) 2n \delta_1 \lesssim_B \delta_0$$

since, otherwise, $\mathfrak{a}_{\nu}^{\mu,n} = 0$. Note $|\langle \gamma^{(N-1)}(\delta_0 \mu), \xi \rangle| \leq B 2^{k+5} \delta_0$ if $\xi \in \operatorname{supp}_{\xi} \mathfrak{a}_{\mu}$. Then, (2.27), (2.28), and the mean value theorem show

$$(3.10) \qquad \qquad |\sigma(\xi) - \delta_0 \mu| \le B^2 2^7 \delta_0$$

for $\xi \in \operatorname{supp}_{\xi} \mathfrak{a}_{\mu}$. If $(\tau, \xi) \in \operatorname{supp}_{\tau,\xi} \mathfrak{a}_{\mu} \subset \Lambda_{k}(\delta_{0}\mu, \delta_{0}, B), |y_{\mu}^{j}(\tau, \xi)| \leq B2^{k+5}\delta_{0}^{N-j}$ for $0 \leq j \leq N-1$. Since $|\omega_{\mu}| \leq B^{2}\delta_{0}$, (3.2) gives $|\mathfrak{g}_{\mu}^{j}(\tau, \xi)| \leq_{B} 2^{k+5}\delta_{0}^{N-j}$ for $0 \leq j \leq N-2$. Therefore, $\mathfrak{G}_{N}^{\mu} \leq_{B} \delta_{0}^{2N!}$ on the support of \mathfrak{a}_{μ} . This gives (3.9). Since $\mathfrak{G}_{N}^{\mu} \leq (2^{n}\delta_{1})^{2N!}$ on $\operatorname{supp} \mathfrak{a}_{\nu}^{\mu,n}$, the following hold on the support of $\mathfrak{a}_{\nu}^{\mu,n}$:

$$(3.11) |s - \sigma(\xi)| \le 2^n \delta_1,$$

(3.12)
$$2^{-k}|\mathfrak{g}^{j}_{\mu}(\tau,\xi)| \le (2^{n}\delta_{1})^{N-j}, \qquad 0 \le j \le N-1.$$

The inequality (3.12) holds true for j = N - 1 since $\mathfrak{g}_{\mu}^{N-1} = 0$. We also have

(3.13)
$$|\mathcal{E}_{j}(\xi)| \leq B^{2} (B^{2} 2^{7} \delta_{0})^{j+1},$$

$$(3.14) \qquad \qquad |\sigma(\xi) - 2^n \delta_1 \nu| \le 2^{n+1} \delta_1.$$

on $\operatorname{supp}_{\xi} \mathfrak{a}_{\nu}^{\mu,n}$. By using (3.4), (3.10), and (3.1), it is easy to show (3.13). Since $|s - 2^n \delta_1 \nu| \leq 2^n \delta_1$ on $\operatorname{supp}_s \mathfrak{a}_{\nu}^{\mu,n}$, (3.14) follows by (3.11).

3.2. Proof of Proposition 2.10. By (3.8) and the Minkowski inequality we have

(3.15)
$$\left(\sum_{\mu} \|\mathcal{T}[\mathfrak{a}^{\mu}]f\|_{p}^{p}\right)^{1/p} \leq \sum_{n\geq 0} \left(\sum_{\mu} \|\sum_{\nu\in\mathfrak{J}_{n}^{\mu}} \mathcal{T}[\mathfrak{a}_{\nu}^{\mu,n}]f\|_{p}^{p}\right)^{1/p}$$

We apply the inequality (2.40) to $\sum_{\nu \in \mathfrak{J}_n^{\mu}} \mathcal{T}[\mathfrak{a}_{\nu}^{\mu,n}]f$ after a suitable linear change of variables. The symbols $\mathfrak{a}_{\nu}^{\mu,0}$ are to constitute the set $\{\mathfrak{a}_{\nu}\}$ while the operators associated to $\mathfrak{a}_{\nu}^{\mu,n}$, $n \geq 1$ are to be handled similarly as in Section 2.

Applying the decoupling inequality. To prove Proposition 2.10, we first show

(3.16)
$$\left\| \sum_{\nu \in \mathfrak{J}_n^{\mu}} \mathcal{T}[\mathfrak{a}_{\nu}^{\mu,n}] f \right\|_p \le C_{\epsilon} \left(2^n \delta_1 / \delta_0 \right)^{\frac{2N}{p} - 1 - \epsilon} \left(\sum_{\nu \in \mathfrak{J}_n^{\mu}} \left\| \mathcal{T}[\mathfrak{a}_{\nu}^{\mu,n}] f \right\|_p^p \right)^{1/p} \right)^{1/p}$$

for $p \ge 4N - 2$. To apply the inequality (2.40), we consider $\operatorname{supp}_{\tau,\xi} \mathfrak{a}_{\nu}^{\mu,n}$, which contains the Fourier support of $\mathcal{T}[\mathfrak{a}_{\nu}^{\mu,n}]f$ as is clear from (2.13) and (2.14).

We set

$$\mathbf{y}_{\mu}(\tau,\xi) = \left(y_{\mu}^{0}(\tau,\xi), \dots, y_{\mu}^{N}(\tau,\xi)\right)$$

Lemma 3.3. Let $\mathbf{r} = \mathbf{r}_{\circ}^{N+1}$ and \mathcal{D}_{δ} denote the matrix $(\delta^{-N}e_1, \delta^{1-N}e_2, \dots, \delta^0e_{N+1})$ where e_j denotes the *j*-th standard unit vector in \mathbb{R}^{N+1} . On $\operatorname{supp}_{\tau,\xi} \mathfrak{a}_{\nu}^{\mu,n}$, we have

(3.17)
$$\left| \left\langle \mathcal{D}_{\delta_0} \mathbf{y}_{\mu}(\tau,\xi), \mathbf{r}^{(j)} \left(\frac{2^n \delta_1}{\delta_0} \nu - \mu \right) \right\rangle \right| \lesssim 2^k \left(\frac{2^n \delta_1}{\delta_0} \right)^{N+1-j}, \quad 1 \le j \le N,$$

(3.18)
$$(2B)^{-1}2^{k-1} \le |\langle \mathbf{y}_{\mu}(\tau,\xi), \mathbf{r}^{(N+1)} \rangle| \le B2^{k+1}$$

Proof. We write $\mathbf{r} = (\mathbf{r}_1, \dots, \mathbf{r}_{N+1})$. Note $\mathbf{r}_m^{(j)}(s) = \frac{s^{m-j}}{(m-j)!}$ for $m \ge j$. By (3.3) we have

$$y_{\mu}^{m-1}\mathbf{r}_{m}^{(j)}(2^{n}\delta_{1}\nu - \delta_{0}\mu) = \sum_{\ell=m-1}^{N} \mathfrak{g}_{\mu}^{\ell} \frac{(2^{n}\delta_{1}\nu - \delta_{0}\mu)^{m-j}}{(\ell+1-m)!(m-j)!} \,\omega_{\mu}^{\ell+1-m}$$

for $m \ge j$. Since $\mathbf{r}_m^{(j)}(s) = 0$ for j > m, taking sum over m gives

$$\langle \mathbf{y}_{\mu}, \mathbf{r}^{(j)}(2^{n}\delta_{1}\nu - \delta_{0}\mu) \rangle = \sum_{\ell=j-1}^{N} \mathfrak{g}_{\mu}^{\ell} \frac{(2^{n}\delta_{1}\nu - \delta_{0}\mu + \omega_{\mu})^{\ell+1-j}}{(\ell+1-j)!}.$$

From (3.5) note $2^n \delta_1 \nu - \delta_0 \mu + \omega_\mu = 2^n \delta_1 \nu - \sigma(\xi) + \mathcal{E}_1$. Thus, (3.14), (3.13) with j = 1, and (2.32) with j = 1 show $|2^n \delta_1 \nu - \delta_0 \mu + \omega_\mu| \leq 2^n \delta_1$. Using (3.12), we obtain

$$\left|\left\langle \mathbf{y}_{\mu}(\tau,\xi), \mathbf{r}^{(j)}(2^{n}\delta_{1}\nu - \delta_{0}\mu)\right\rangle\right| \lesssim 2^{k}(2^{n}\delta_{1})^{N+1-j}, \qquad 1 \le j \le N.$$

By homogeneity it follows that $\langle \eta, \mathbf{r}^{(j)}(\delta_0 s) \rangle = \delta_0^{N+1-j} \langle \mathcal{D}_{\delta_0} \eta, \mathbf{r}^{(j)}(s) \rangle$ for $\eta \in \mathbb{R}^{N+1}$. Therefore, we get (3.17). For (3.18) note $\mathbf{r}^{(N+1)} = (0, \dots, 0, 1)$, so $\langle \mathbf{y}_{\mu}, \mathbf{r}^{(N+1)} \rangle = y_{\mu}^N$ and (3.18) follows by (3.1).

Let $V = \operatorname{span}\{\gamma'(\delta_0\mu), \ldots, \gamma^{(N)}(\delta_0\mu)\}$ and $\{v_{N+1}, \ldots, v_d\}$ be an orthonormal basis of V^{\perp} . Since γ satisfies $\mathfrak{V}(N, B)$, for each $\xi \in \mathbb{R}^d$ we can write

(3.19)
$$\xi = \overline{\xi} + \sum_{N+1 \le j \le d} y_j(\xi) v_j,$$

where $\overline{\xi} \in \mathcal{V}$ and $y_j(\xi) \in \mathbb{R}$, $N+1 \leq j \leq d$. We define a linear map $Y^{\delta_0}_{\mu}$ by $Y^{\delta_0}_{\mu}(\tau,\xi) = \left(2^{-k}\mathcal{D}_{\delta_0}\mathbf{y}_{\mu}(\tau,\xi), y_{N+1}(\xi), \dots, y_d(\xi)\right).$ Then, by (3.17) and (3.18) we see

(3.20)
$$Y^{\delta_0}_{\mu}(\operatorname{supp}_{\tau,\xi} \mathfrak{a}^{\mu,n}_{\nu}) \subset \mathbf{s}\left(\frac{2^n \delta_1}{\delta_0}\nu - \mu, C\frac{2^n \delta_1}{\delta_0}, 2^2 B; \mathbf{r}^{N+1}_{\circ}\right) \times \mathbb{R}^{d-N}$$

for some C > 1. Thus, we have the inequality (2.40) for $\delta = C2^n \delta_1/\delta_0$, the collection of slabs $\mathbf{s}(2^n \delta_1 \nu / \delta_0 - \mu, C2^n \delta_1 / \delta_0, CB; \mathbf{r}_{\circ}^{N+1}), \nu \in \mathfrak{J}_n^{\mu}$. Therefore, via cylindrical extension in y_{N+1}, \ldots, y_d (see (2.42)) and the change of variables $(\tau, \xi) \to \mathbf{Y}_{\mu}^{\delta_0}(\tau, \xi)$ we obtain (3.16) since the decoupling inequality is not affected by affine change of variables in the Fourier side.

Combining (3.15) and (3.16), we obtain

$$ig(\sum_{\mu} \|\mathcal{T}[\mathfrak{a}^{\mu}]f\|_p^pig)^{1/p} \leq \sum_{n\geq 0} \mathbf{E}_n$$

for $p \ge 4N - 2$, where

$$\mathbf{E}_n = C_\epsilon \left(2^n \delta_1 / \delta_0 \right)^{\frac{2N}{p} - 1 - \epsilon} \left(\sum_{\mu} \sum_{\nu \in \mathfrak{J}_n^{\mu}} \| \mathcal{T}[\mathfrak{a}_{\nu}^{\mu, n}] f \|_p^p \right)^{1/p}$$

Since the intervals $I(\delta_0\mu, \delta_0)$ overlap, there are at most three nonzero $\mathfrak{a}_{\nu}^{\mu,0}$ for each ν . We take $\mathfrak{a}_{\nu} = \mathfrak{a}_{\nu}^{\mu,0}$ which maximizes $\|\mathcal{T}[\mathfrak{a}_{\nu}^{\mu,0}]f\|_p$. Then, it is clear that $\mathbf{E}_0 \leq 3^{1/p}C_{\epsilon}(\delta_1/\delta_0)^{\frac{2N}{p}-1-\epsilon}(\sum_{\nu}\|\mathcal{T}[\mathfrak{a}_{\nu}]f\|_p^p)^{1/p}$. By Lemma 3.2, $C^{-1}\mathfrak{a}_{\nu} \in \mathfrak{A}_k(\delta_1\nu, \delta_1)$ for a constant C. Thus, the proof of Proposition 2.10 is now reduced to showing

(3.21)
$$\sum_{n\geq 1} \mathbf{E}_n \lesssim_B \delta_0^{-\frac{2N}{p}+1+\epsilon} 2^{-\frac{2}{p}k+2\epsilon k} \|f\|_p, \quad p\geq 4N-2.$$

Estimates for \mathbf{E}_n when $n \ge 1$. To show (3.21) we decompose $\mathfrak{a}_{\nu}^{\mu,n}$ so that (3.26) or (3.27) (see Lemma 3.5 below) holds on the s, ξ -supports of the resulting symbols. If (3.26) holds, we use the assumption after rescaling, whereas we handle the other case using estimates for the kernels of the operators.

Let

(3.22)
$$\bar{\mathfrak{G}}_{N}^{\mu}(s,\xi) = \sum_{1 \le j \le N-2} \left(2^{-k} \mathfrak{g}_{\mu}^{j} \right)^{\frac{2N!}{N-j}} + \left(s - \sigma(\xi) \right)^{2N!}.$$

The right hand side is independent of τ since so are \mathfrak{g}^{j}_{μ} , $1 \leq j \leq N-2$. Let $C_{0} = 2^{2d}B$. We set

(3.23)
$$\mathfrak{a}_{\nu,1}^{\mu,n} = \mathfrak{a}_{\nu}^{\mu,n} \beta_0 \Big((2^{-k} \mathfrak{g}_{\mu}^0)^{2(N-1)!} / (C_0^{2N!} \bar{\mathfrak{G}}_N^{\mu}) \Big), \quad n \ge 1.$$

and $\mathfrak{a}_{\nu,2}^{\mu,n} = \mathfrak{a}_{\nu}^{\mu,n} - \mathfrak{a}_{\nu,1}^{\mu,n}$, so $\mathfrak{a}_{\nu}^{\mu,n} = \mathfrak{a}_{\nu,1}^{\mu,n} + \mathfrak{a}_{\nu,2}^{\mu,n}$. Similarly as before, we have the following, which we prove in Section 3.4.

Lemma 3.4. There exists a constant C = C(B) such that $C^{-1}\mathfrak{a}_{\nu,1}^{\mu,n}$, and $C^{-1}\mathfrak{a}_{\nu,2}^{\mu,n}$ are contained in $\mathfrak{A}_k(2^n\delta_1\nu, 2^n\delta_1)$ for $n \ge 1$.

The estimate (3.21) follows if we show

(3.24)
$$\left(\sum_{\mu}\sum_{\nu\in\mathfrak{J}_{n}^{\mu}} \|\mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f\|_{p}^{p}\right)^{1/p} \leq C_{\epsilon}2^{-\frac{2}{p}k+\epsilon k}(2^{n}\delta_{1})^{-\frac{2N}{p}+1+\epsilon}\|f\|_{p}, \quad p \geq 4N-6,$$

for any $\epsilon > 0$, and

(3.25)
$$(\sum_{\mu} \sum_{\nu \in \mathfrak{J}_n^{\mu}} \|\mathcal{T}[\mathfrak{a}_{\nu,2}^{\mu,n}]f\|_p^p)^{1/p} \lesssim_B 2^{-\frac{(N+2)k}{2N}} (2^n \delta_1)^{-\frac{N}{2}} \|f\|_p, \quad 2 \le p \le \infty$$

when $n \ge 1$. Thanks to (3.9), those estimates give

$$\sum_{n\geq 1} \mathbf{E}_n \leq C_{\epsilon} \delta_0^{-\frac{2N}{p}+1+\epsilon} \sum_{1\leq n\leq \log_2(C\delta_0/\delta_1)} \left(2^{-\frac{2}{p}k+\epsilon k} + 2^{-\frac{(N+2)k}{2N}} (2^n \delta_1)^{\frac{2N}{p}-\frac{N+2}{2}-\epsilon}\right) \|f\|_p$$

for $p \ge 4N-2$. Note $\log_2(\delta_0/\delta_1) \le Ck$ from (2.31). So, (3.21) follows since 4N-2 > 4N/(N+2) and $\delta_1 \ge 2^{-k/N}$.

In order to prove the estimates (3.24) and (3.25), we start with the next lemma.

Lemma 3.5. Let $n \ge 1$. For a constant C = C(B) > 0, we have the following:

(3.26)
$$\sum_{1 \le j \le N-1} (2^n \delta_1)^{-(N-j)} |\langle \gamma^{(j)}(s), \xi \rangle| \ge C2^k, \qquad (s,\xi) \in \operatorname{supp}_{s,\xi} \mathfrak{a}_{\nu,1}^{\mu,n},$$

(3.27)
$$(2^n \delta_1)^{-N} |\tau + \langle \gamma(s), \xi \rangle| \ge C2^k, \qquad (s,\xi) \in \operatorname{supp}_{s,\xi} \mathfrak{a}_{\nu,2}^{\mu,n}.$$

Proof. We first prove (3.26). Since $\mathfrak{G}_N^{\mu} \geq 2^{-2N!-1} (2^n \delta_1)^{2N!}$ on $\operatorname{supp}_{s,\xi} \mathfrak{a}_{\nu}^{\mu,n}$, one of the following holds on $\operatorname{supp} \mathfrak{a}_{\nu,1}^{\mu,n}$:

(3.28)
$$|s - \sigma(\xi)| \ge (2^3 C_0 B)^{-1} 2^n \delta_1,$$

(3.29)
$$2^{-k}|\mathfrak{g}_{\mu}^{j}(\tau,\xi)| \ge (2^{2}C_{0})^{-(N-j)}(2^{n}\delta_{1})^{N-j}$$

for some $1 \leq j \leq N-2$, where $C_0 = 2^{2d}B$ (see (3.23)). If (3.28) holds, by (2.27) and (2.28) it follows that $(2^n\delta_1)^{-1}|\langle \gamma^{(N-1)}(s), \xi \rangle| \gtrsim 2^k$. Thus, to show (3.26) we may assume (3.28) fails, i.e., (3.29) holds for some $1 \leq j \leq N-2$. So, there is an integer $\ell \in [0, N-2]$ such that (3.29) fails for $\ell + 1 \leq j \leq N-2$, whereas (3.29) holds for $j = \ell$. By (3.6) and (3.13), we have

$$(3.30) |\langle G^{(\ell)}(\sigma(\xi)), (\tau, \xi) \rangle| \ge |\mathfrak{g}_{\mu}^{\ell}| - \sum_{j=\ell+1}^{N} |\mathfrak{g}_{\mu}^{j}| \frac{(B^{6}2^{14}\delta_{0}^{2})^{j-\ell}}{(j-\ell)!} - 2B^{3}(B^{2}2^{7}\delta_{0})^{N+1-\ell} |\xi|.$$

Thus, by (2.32), $|\langle G^{(\ell)}(\sigma(\xi)), (\tau, \xi) \rangle| \geq (2^3 C_0)^{-(N-\ell)} 2^k (2^n \delta_1)^{N-\ell}$. Also, (3.6) and our choice of ℓ give $|\langle G^{(j)}(\sigma(\xi)), (\tau, \xi) \rangle| \leq (2C_0)^{-(N-j)} 2^k (2^n \delta_1)^{N-j}$ for $\ell + 1 \leq j \leq N-2$. Combining this with $|s - \sigma(\xi)| < (2^3 C_0 B)^{-1} 2^n \delta_1$ and expanding $G^{(\ell)}$ in Taylor series at $\sigma(\xi)$, we see $|\langle G^{(\ell)}(s), (\tau, \xi) \rangle| \geq C 2^k (2^n \delta_1)^{N-\ell}$ for some C = C(B) > 0. This proves (3.26).

We now show (3.27), which is easier. On $\sup \mathfrak{a}_{\nu,2}^{\mu,n}$, $2^{-k}|\mathfrak{g}_{\mu}^{0}| \geq 2^{-N-1}(2^{n}\delta_{1})^{N}$ and $2^{-k}|\mathfrak{g}_{\mu}^{j}| \leq 2C_{0}^{-(N-j)}(2^{n}\delta_{1})^{N-j}$ for $j = 1, \ldots, N-2$. Using (3.30) with $\ell = 0$, by (2.32) and (2.31) we get $(2^{n}\delta_{1})^{-N}|\tau + \langle \gamma(\sigma(\xi)), \xi \rangle| \geq 2^{-N-2}2^{k}$. We also note that $|s-\sigma(\xi)| \leq 2C_{0}^{-1}2^{n}\delta_{1}$ and $|\langle G^{(j)}(\sigma(\xi)), (\tau, \xi) \rangle| \leq C_{0}^{-1}2^{k}(2^{n}\delta_{1})^{N-j}$ for $1 \leq j \leq N-2$ on $\sup \mathfrak{a}_{\nu,2}^{\mu,n}$. Since $|\langle G^{(N)}(s), (\tau, \xi) \rangle| \leq B2^{k+1}$, using Taylor series expansion at $\sigma(\xi)$ as above, we see (3.27) holds true for some C = C(B) > 0.

Additionally, we make use of disjointness of $\operatorname{supp}_{\xi} \mathfrak{a}_{\nu}^{\mu,n}$ by combining Lemma 2.11 and the next one.

Lemma 3.6. There is a positive constant C = C(B) such that

$$(3.31) \qquad \qquad |(\widetilde{\mathcal{L}}_s^{\delta})^{-1}\xi| \le Cb2^k$$

whenever $\xi \in \Lambda'_k(s, \delta, b)$ (see (2.36)). If $\xi \in \Gamma_k$ and (3.31) holds with C = 1, then $\xi \in \Lambda'_k(s, \delta, C_1b)$ for some $C_1 = C_1(B) > 0$.

Proof. Let $\eta \in \mathbb{R}^d$ and $\{v_N, \ldots, v_d\}$ be an orthonormal basis of $(\operatorname{span}\{\gamma^{(j)}(s) : 1 \leq j \leq N-1\})^{\perp}$. We write $\eta = \sum_{j=1}^{N-1} \mathbf{c}_j \gamma^{(j)}(s) + \sum_{j=N}^d \mathbf{c}_j v_j$. Since $\mathfrak{V}(N, B)$ holds for γ , $|\eta| \sim |(\mathbf{c}_1, \cdots, \mathbf{c}_d)|$. Let $\xi \in \Lambda'_k(s, \delta, b)$. Then, (2.4) gives

$$\langle \eta, (\widetilde{\mathcal{L}}_s^{\delta})^{-1} \xi \rangle = \langle (\widetilde{\mathcal{L}}_s^{\delta})^{-\mathsf{T}} \eta, \xi \rangle = \sum_{j=1}^{N-1} \delta^{j-N} \mathbf{c}_j \langle \gamma^{(j)}(s), \xi \rangle + \sum_{j=N}^d \mathbf{c}_j \langle v_j, \xi \rangle.$$

Thus, by (2.36) we get $|\langle \eta, (\widetilde{\mathcal{L}}_s^{\delta})^{-1}\xi \rangle| \leq Cb|\eta|2^k$, which shows (3.31). By (2.4), $\langle \gamma^{(j)}(s), \xi \rangle = \delta^{N-j} \langle \gamma^{(j)}(s), (\widetilde{\mathcal{L}}_s^{\delta})^{-1}\xi \rangle$ for $1 \leq j \leq N-1$. Therefore, (3.31) with C = 1 gives $|\langle \gamma^{(j)}(s), \xi \rangle| \leq C_1 b \delta^{N-j} 2^k$ for a constant $C_1 > 0$ when $1 \leq j \leq N - 1$. This proves the second statement.

Now, we are ready to prove the estimates (3.24) and (3.25). We first show (3.24).

Proof of (3.24). By Lemma 3.4, $C^{-1}\mathfrak{a}_{\nu,1}^{\mu,n} \in \mathfrak{A}_k(2^n\delta_1\nu, 2^n\delta_1)$ for some C > 0, and (3.26) holds on $\operatorname{supp}_{s,\underline{\xi}}\mathfrak{a}_{\nu,1}^{\mu,n}$. Thus, taking $\delta = 2^n\delta_1$ and $s_0 = 2^n\delta_1\nu$, we may use Lemma 2.8 for $\tilde{\chi} \mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f$ to get

$$\left\| \tilde{\chi} \mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}] f \right\|_{L^p(\mathbb{R}^{d+1})} \le C \sum_{1 \le l \le C} \delta \left\| \mathcal{A}_t[\gamma_{s_0}^{\delta}, a_l] \tilde{f}_l \right\|_{L^p(\mathbb{R}^{d+1})}$$

where $\|\tilde{f}_l\|_p = \|f\|_p$, a_l are of type (j, N-1, B') relative to $\gamma_{s_0}^{\delta}$ for some B' > 0, and $2^j \sim (2^n \delta_1)^N 2^k$. As seen before, $\gamma = \gamma_{s_0}^{\delta}$ satisfies $\mathfrak{V}(N, 3B)$ and (2.1) with Breplaced by 3B for $\delta \leq \delta_*$. So, $\gamma = \gamma_{s_0}^{\delta}$ satisfies $\mathfrak{V}(N-1, B')$ for a large B'. Therefore, we may apply the assumption (Theorem 2.2 with L = N - 1) to

 $\mathcal{A}_t[\gamma_{s_0}^{\delta}, a_l]$, which gives $\|\mathcal{A}_t[\gamma_{s_0}^{\delta}, a_l]f\|_p \leq C_{\epsilon} (2^k (2^n \delta_1)^N)^{-\frac{2}{p}+\epsilon} \|f\|_p$ for a constant $C_{\epsilon} = C_{\epsilon}(B')$. Consequently, we obtain

$$\|\tilde{\chi}\mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f\|_p \le C_{\epsilon} 2^{-\frac{2}{p}k+\epsilon k} (2^n \delta_1)^{1-\frac{2N}{p}+\epsilon} \|f\|_p$$

for $p \geq 4(N-1)-2$. Besides, since $C^{-1}\mathfrak{a}_{\nu,1}^{\mu,n} \in \mathfrak{A}_k(2^n\delta_1\nu, 2^n\delta_1)$, by (2.16) we have $\|(1-\tilde{\chi})\mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f\|_{L^p(\mathbb{R}^{d+1})} \lesssim_B 2^{-k}(2^n\delta_1)^{1-N}\|f\|_{L^p(\mathbb{R}^d)}$ for p > 1. Note $2^n\delta_1 \gtrsim$ $2^{-k/N}$. Combining those two estimates yields

(3.32)
$$\|\mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f\|_{p} \le C_{\epsilon} 2^{-\frac{2}{p}k+\epsilon k} (2^{n}\delta_{1})^{1-\frac{2N}{p}+\epsilon} \|f\|_{p}.$$

To exploit disjointness of $\operatorname{supp}_{\xi} \mathfrak{a}_{\nu,1}^{\mu,n}$, we define a multiplier operator by

$$\mathcal{F}(P_s^{\delta}f)(\xi) = \beta_0 \left(|(\widetilde{\mathcal{L}}_s^{\delta})^{-1}\xi| / (C_0 2^k) \right) \widehat{f}(\xi)$$

for a constant $C_0 > 0$. Since $\operatorname{supp}_{\xi} \mathfrak{a}_{\nu,1}^{\mu,n} \subset \Lambda'_k(2^n \delta_1 \nu, 2^n \delta_1, 2^5 B)$, by Lemma 3.6 we may choose C_0 large enough so that $\beta_0(|(\widetilde{\mathcal{L}}_{2^n\delta_1\nu}^{2^n\delta_1})^{-1}\cdot|/(C_02^k)) = 1$ on $\operatorname{supp}_{\xi}\mathfrak{a}_{\nu,1}^{\mu,n}$. Thus, $\mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f = \mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]P_{2^n\delta_1\nu}^{2^n\delta_1}f$. Combining this and (3.32), we obtain

$$\left(\sum_{\mu} \sum_{\nu \in \mathfrak{J}_{n}^{\mu}} \|\mathcal{T}[\mathfrak{a}_{\nu,1}^{\mu,n}]f\|_{p}^{p}\right)^{1/p} \leq C_{\epsilon} 2^{-\frac{2}{p}k+\epsilon k} (2^{n}\delta_{1})^{1-\frac{2N}{p}+\epsilon} \left(\sum_{\mu} \sum_{\nu \in \mathfrak{J}_{n}^{\mu}} \|P_{2^{n}\delta_{1}\nu}^{2^{n}\delta_{1}}f\|_{p}^{p}\right)^{1/p}$$

for a constant $C_{\epsilon} = C_{\epsilon}(B)$ if $p \ge 4N - 6$. Therefore, (3.24) follows if we show

(3.33)
$$(\sum_{\mu} \sum_{\nu \in \mathfrak{J}_n^{\mu}} \|P_{2^n \delta_1 \nu}^{2^n \delta_1} f\|_p^p)^{1/p} \lesssim_B \|f\|_p, \qquad 2 \le p \le \infty$$

By interpolation it suffices to obtain (3.33) for $p = 2, \infty$. The case $p = \infty$ is trivial since $\|P_{2n\delta_1\nu}^{2^n\delta_1}f\|_{\infty} \lesssim \|f\|_{\infty}$. For p=2, (3.33) follows by Plancherel's theorem since $\operatorname{supp} \beta_0 \left(|(\widetilde{\mathcal{L}}_{2^n \delta_1 \nu}^{2^n \delta_1})^{-1} \cdot |/(C_0 2^k) \right) \widehat{f}, \nu \in \mathfrak{J}_n^{\mu}$ are finitely overlapping. Indeed, by Lemma 3.6 we have $\operatorname{supp} \beta_0 \left(|(\widetilde{\mathcal{L}}_{2^n \delta_1 \nu}^{2^n \delta_1})^{-1} \cdot |/(C_0 2^k) \right) \widehat{f} \subset \Lambda'_k(2^n \delta_1 \nu, 2^n \delta_1, C_1 B)$ for a constant C_1 . It is clear from lemma 2.11 that $\Lambda'_k(2^n\delta_1\nu, 2^n\delta_1, C_1B), \nu \in \mathfrak{J}_n^{\mu}$ overlap at most C = C(B) times.

The proof of (3.25) is much easier since we have a favorable estimate for the kernel of $\mathcal{T}[\mathfrak{a}_{\nu,2}^{\mu,n}]$ thanks to the lower bound (3.27).

Proof of (3.25). Let

$$\mathfrak{b}(s,t,\tau,\xi) = i^{-1}(\tau + \langle \gamma(s),\xi \rangle)^{-1} \partial_t \mathfrak{a}_{\nu,2}^{\mu,n}(s,t,\tau,\xi).$$

Then, integration by parts in t shows $m[\mathfrak{a}_{\nu,2}^{\mu,n}] = m[\mathfrak{b}]$. Note (3.27) holds and $C^{-1}\mathfrak{a}_{\nu,2}^{\mu,n} \in \mathfrak{A}_k(2^n\delta_1\nu,2^n\delta_1)$ for a constant $C \geq 1$. Thus, $\mathfrak{a} := C^{-1}2^k(2^n\delta_1)^N\mathfrak{b}$ satisfies, with $\delta = 2^n \delta_1$ and $s_0 = 2^n \delta_1 \nu$, (2.6) and (2.7) for $0 \le j \le 1, 0 \le l \le 2N-1$, $|\alpha| \le d + N + 2$. Applying (2.15), we obtain $\|\mathcal{T}[\mathfrak{a}_{\nu,2}^{\mu,n}]f\|_{\infty} \lesssim_B 2^{-k} (2^n \delta_1)^{1-N} \|f\|_{\infty}$. Since $\delta_1 \geq 2^{-k/N}$, this gives

(3.34)
$$\|\mathcal{T}[\mathfrak{a}_{\nu,2}^{\mu,n}]f\|_{\infty} \lesssim_{B} 2^{-\frac{(N+2)k}{2N}} (2^{n}\delta_{1})^{-\frac{N}{2}} \|f\|_{\infty}.$$

By interpolation it is sufficient to show (3.25) for p = 2. Note $||b(\cdot, t, \tau, \xi)||_{\infty} +$ $\|\partial_s b(\cdot, t, \tau, \xi)\|_1 \lesssim 2^{-k} (2^n \delta_1)^{-N}$. Thus, (2.27) and van der Corput's lemma in s give $|m[\mathfrak{a}_{\nu,2}^{\mu,n}](\tau,\xi)| \lesssim 2^{-k(1+N)/N} (2^n \delta_1)^{-N}$. Since $\operatorname{supp}_{\xi} \mathfrak{a}_{\nu,2}^{\mu,n} \subset \Lambda'_k(2^n \delta_1 \nu, 2^n \delta_1, 2^5 B)$, as before, we have $\mathcal{T}[\mathfrak{a}_{2,\nu}^{\mu,n}]f = \mathcal{T}[\mathfrak{a}_{2,\nu}^{\mu,n}]P_{2^n\delta_1\nu}^{2^n\delta_1}f$ with $C_0 > 0$ large enough. Thus, by Plancherel's theorem

$$\begin{aligned} \|\mathcal{T}[\mathfrak{a}_{\nu,2}^{\mu,n}]f\|_{L^2}^2 &\lesssim_B 2^{-\frac{2(1+N)}{N}k} (2^n \delta_1)^{-2N} \iint_{\{\tau:|\mathfrak{g}_{\mu}^0(\tau,\xi)| \leq 2^{k+1}(2^n \delta_1)^N\}} d\tau \, |\mathcal{F}(P_{2^n \delta_1 \nu}^{2^n \delta_1} f)(\xi)|^2 \, d\xi. \end{aligned}$$
Combining this and (3.33) yields (3.25) for $p=2.$

Combining this and (3.33) yields (3.25) for p = 2.

3.3. Proof of Lemma 3.2. To simplify notations, we denote

$$\delta_{\star} = 2^n \delta_1, \qquad s_{\star} = 2^n \delta_1 \nu$$

for the rest of this section. To prove Lemma 3.2, we verify (2.6) and (2.7) with $\mathfrak{a} = \mathfrak{a}_{\nu}^{\mu,n}, \ \delta = \delta_{\star}, \ \text{and} \ s_0 = s_{\star}.$ The first is easy. In fact, since $\mathfrak{a}^{\mu} \in \mathfrak{A}_k(\delta_0 \mu, \delta_0)$ and $\operatorname{supp}_{s} \mathfrak{a}_{\mu}^{\mu,n} \subset I(s_{\star},\delta_{\star})$, we only need to show

(3.35)
$$|\langle G^{(j)}(s_{\star}), (\tau, \xi) \rangle| \le B2^{k+5} \delta_{\star}^{N-j}, \quad j = 0, \dots, N-1$$

on $\operatorname{supp}_{\tau,\xi} \mathfrak{a}_{\nu}^{\mu,n}$. Using (3.6) and (3.12) together with (2.32) and (3.13), one can easily obtain

(3.36)
$$|\langle G^{(j)}(\sigma(\xi)), (\tau, \xi) \rangle| \le 2^{k+1} \delta^{N-j}_{\star}, \quad j = 0, \dots, N-1$$

on $\operatorname{supp}_{\tau,\xi} \mathfrak{a}_{\nu}^{\mu,n}$. Expanding $\langle G^{(j)}(s), (\tau,\xi) \rangle$ in Taylor's series at $\sigma(\xi)$ gives (3.35) since (3.14) holds.

We now proceed to show (2.7) with $\mathfrak{a} = \mathfrak{a}_{\nu}^{\mu,n}$, $\delta = \delta_{\star}$, and $s_0 = s_{\star}$. Since $\mathfrak{a}_{\nu}^{\mu,n}$ consists of three factors \mathfrak{a}^{μ} , $\beta_N(\delta_{\star}^{-2N!}\mathfrak{G}_N^{\mu})$, and $\zeta(\delta_{\star}^{-1}s-\nu)$, by Leibniz's rule it is sufficient to consider the derivatives of each of them. The bounds on the derivatives $\zeta(\delta_{-}^{-1}s - \nu)$ are clear. So, it suffices to show (2.7) for

$$\mathfrak{a} = \mathfrak{a}^{\mu}, \ \beta_N(\delta_{\star}^{-2N!}\mathfrak{G}_N^{\mu})$$

with $\delta = \delta_{\star}$ and $s_0 = s_{\star}$ whenever $(\tau, \xi) \in \operatorname{supp} \mathfrak{a}_{\nu}^{\mu, n}(s, t, \mathcal{L}_{s_{\star}}^{\delta_{\star}})$.

We handle \mathfrak{a}^{μ} first. That is to say, we show

$$(3.37) \qquad \left|\partial_s^j \partial_t^l \partial_{\tau,\xi}^{\alpha} \left(\mathfrak{a}^{\mu}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}}(\tau,\xi))\right)\right| \lesssim_B \delta_{\star}^{-j} |(\tau,\xi)|^{-|\alpha|}, \qquad (j,l,\alpha) \in \mathcal{I}_N,$$

for $(\tau,\xi) \in \operatorname{supp} \mathfrak{a}_{\nu}^{\mu,n}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}}\cdot)$. Since $\mathfrak{a}^{\mu} \in \mathfrak{A}_{k}(\delta_{0}\mu,\delta_{0})$ and $|s_{\star} - \delta_{0}\mu| \leq \delta_{0}$, we have

$$(3.38) \qquad |\partial_s^j \partial_t^l \partial_{\tau,\xi}^{\alpha} \left(\mathfrak{a}^{\mu}(s,t,\mathcal{L}^{\delta_0}_{s_\star}(\tau,\xi)) \right)| \lesssim_B \delta_0^{-j} |(\tau,\xi)|^{-|\alpha|}, \qquad (j,l,\alpha) \in \mathcal{I}_N$$

One can show this using (2.11). We consider $\mathcal{U} := (\mathcal{L}_{s_{\star}}^{\delta_0})^{-1} \mathcal{L}_{s_{\star}}^{\delta_{\star}}$. By (2.8) we have $|\mathcal{U}^{\intercal}z| \lesssim_B |z|$ because $|\delta_0^{-1}2^n \delta_1| \lesssim_B 1$. Thus, (3.38) gives

$$\left|\partial_s^j \partial_t^l \partial_{\tau,\xi}^{\alpha} \left(\mathfrak{a}^{\mu}(s,t,\mathcal{L}_{s_{\star}}^{\delta_0} \mathcal{U}(\tau,\xi)) \right) \right| \lesssim_B \delta_0^{-j} |\mathcal{U}(\tau,\xi)|^{-|\alpha|}$$

for $(\tau,\xi) \in \operatorname{supp} \mathfrak{a}_{\nu}^{\mu,n}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}}).$ Let $(\tau,\xi) \in \operatorname{supp} \mathfrak{a}_{\nu}^{\mu,n}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}}).$ Then, $\mathcal{L}_{s_{\star}}^{\delta_{0}}\mathcal{U}(\tau,\xi) = \mathcal{L}_{s_{\star}}^{\delta_{\star}}(\tau,\xi) \in \Lambda_{k}(s_{\star},\delta_{\star},B)$, so $|\widetilde{\mathcal{L}}_{s}^{\delta_{\star}}\xi| \sim |(\tau,\xi)|$ by Lemma 2.5. This and (2.9) give

$$|(\tau,\xi)| \sim |\widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}} \xi| \le |\mathcal{L}_{s_{\star}}^{\delta_{\star}}(\tau,\xi)| \le |\mathcal{U}(\tau,\xi)|$$

for $(\tau,\xi) \in \operatorname{supp} \mathfrak{a}_{\nu}^{\mu,n}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}})$. So, we obtain (3.37) since $\delta_{\star} \leq \delta_{0}$.

We continue to show (2.7) for $\mathfrak{a} = \beta_N(\delta_{\star}^{-2N!}\mathfrak{G}_N^{\mu})$. Note $(2^n\delta_1)^{-2N!}\mathfrak{G}_N^{\mu}$ is a sum of $(\delta_{\star}^{-1}(s-\sigma(\xi)))^{2N!}$ and $(\delta_{\star}^{-(N-j)}2^{-k}\mathfrak{g}_{\mu}^{j})^{2N!/(N-j)}$, $0 \leq j \leq N-2$. Since the exponents 2N!/(N-j) are even integers, for the desired bounds on $\partial_{\tau,\ell}^{\alpha}(\beta_N(\delta_{\star}^{-2N!}\mathfrak{G}_N^{\mu}))$ it suffices to show the same bounds on the derivatives of

$$\delta_{\star}^{-1}(s-\sigma(\xi)), \qquad \delta_{\star}^{-(N-j)}2^{-k}\mathfrak{g}_{\mu}^{j}, \quad 0 \le j \le N-2.$$

The bound on $\partial_{\xi}^{\alpha} \delta_{\star}^{-1}(s-\sigma)$ is a consequence of (2.10) and the following lemma. To simplify notations, we denote

$$\Xi = \mathcal{L}_{s_{\star}}^{\delta_{\star}}(\tau,\xi), \qquad \tilde{\Xi} = \widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}}\xi$$

Lemma 3.7. If $\Xi \in \operatorname{supp}_{\tau, \mathcal{E}} \mathfrak{a}_{\nu}^{\mu, n}$, then we have

(3.39)
$$|\delta_{\star}^{-1}\partial_{\xi}^{\alpha}(\sigma(\tilde{\Xi}))| \lesssim_{B} |\xi|^{-|\alpha|}, \qquad 1 \le |\alpha| \le 2d+2.$$

Proof. By (2.28), $\gamma^{(N-1)}(\sigma(\tilde{\Xi})) \cdot \tilde{\Xi} = 0$. Differentiation gives

(3.40)
$$\gamma^{(N)}(\sigma(\tilde{\Xi})) \cdot \tilde{\Xi} \nabla_{\xi}(\sigma(\tilde{\Xi})) + (\widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}})^{\mathsf{T}} \gamma^{(N-1)}(\sigma(\tilde{\Xi})) = 0.$$

Denote $s = \sigma(\tilde{\Xi})$. By (2.4), $(\widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}})^{\intercal}\gamma^{(N-1)}(s) = \delta_{\star}(\widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}})^{\intercal}(\widetilde{\mathcal{L}}_{s}^{\delta_{\star}})^{-\intercal}\gamma^{(N-1)}(s)$. Since $|s_{\star}-s| \leq \delta_{\star}$, i.e., (3.11), by Lemma 2.6 we have $|(\widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}})^{\mathsf{T}}\gamma^{(N-1)}(\sigma(\widetilde{\Xi}))| \lesssim_{B} \delta_{\star}$. Besides, $|\gamma^{(N)}(\sigma(\tilde{\Xi})) \cdot \tilde{\Xi}| \gtrsim |\tilde{\Xi}| \sim 2^k$ (see (2.27)). Thus, (3.40) and (2.10) give

$$|\nabla_{\xi}(\sigma(\Xi))| \lesssim_B \delta_{\star} |\xi|^{-1},$$

which proves (3.39) with $|\alpha| = 1$.

We show the bounds on the derivatives of higher order by induction. Assume that (3.39) holds true for $|\alpha| \leq L$. Let α' be a multi-index such that $|\alpha'| = L + 1$. Then, differentiating (3.40) and using the induction assumption, one can easily see $\gamma^{(N)}(\sigma(\tilde{\Xi})) \cdot \tilde{\Xi} \ \partial_{\xi}^{\alpha'}(\sigma(\tilde{\Xi})) = O(\delta_{\star}|\xi|^{-L})$, by which we get (3.39) for $|\alpha| = L+1$. Since $\sigma \in C^{2d+2}$, one can continue this as far as $L \leq 2d+1$.

The proof of Lemma 3.2 is now completed if we show

(3.41)
$$\left|2^{-k}\partial_{\tau,\xi}^{\alpha}\left(\mathfrak{g}_{\mu}^{\ell}(\Xi)\right)\right| \lesssim_{B} \delta_{\star}^{N-\ell} 2^{-k|\alpha|}, \qquad |\alpha| \le d+N+2$$

for $0 \leq \ell \leq N-2$ whenever $\Xi \in \operatorname{supp} \mathfrak{a}^{\mu,n}_{\mu}(s,t,\cdot)$. To this end, we use the following.

Lemma 3.8. For j = 0, ..., N, we set

$$A_j = \delta_{\star}^{-(N-j)} 2^{-k} \langle G^{(j)}(\sigma(\tilde{\Xi})), \Xi \rangle$$

If $(\tau,\xi) \in \operatorname{supp} \mathfrak{a}_{\nu}^{\mu,n}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}})$, then for $j = 0, \ldots, N$ we have

$$(3.42) \qquad \qquad |\partial_{\tau,\xi}^{\alpha} A_j| \lesssim_B |(\tau,\xi)|^{-|\alpha|}, \qquad 1 \le |\alpha| \le 2d+2.$$

Proof. When j = N, the estimate (3.42) follows by Lemma 3.7 and (2.10). So, we may assume $j \leq N - 1$. Differentiating A_j , we have

$$\nabla_{\tau,\xi} A_j = B_j + D_j,$$

where

$$B_j = \delta_{\star}^{-1} \big(0, \nabla_{\xi}(\sigma(\tilde{\Xi})) \big) A_{j+1}, \qquad D_j = \delta_{\star}^{-(N-j)} 2^{-k} (\mathcal{L}_{s_{\star}}^{\delta_{\star}})^{\mathsf{T}} G^{(j)}(\sigma(\tilde{\Xi})).$$

Note $(\mathcal{L}_{s_{\star}}^{\delta_{\star}})^{\intercal}G^{(j)}(s_{\star}) = \delta_{\star}^{N-j}G^{(j)}(s_{\star})$ for $0 \leq j \leq N-1$. Since $|s_{\star} - \sigma(\tilde{\Xi})| \lesssim \delta_{\star}$, similarly as before, Lemma 2.6 and (2.8) give

(3.43)
$$|(\mathcal{L}_{s_{\star}}^{\delta_{\star}})^{\mathsf{T}}G^{(j)}(\sigma(\tilde{\Xi}))| \lesssim_{B} \delta_{\star}^{N-j}, \qquad 0 \le j \le N-1$$

By Lemma 3.7 and (3.36), $|B_j| \leq |\xi|^{-1}$. Thus, for $\Xi \in \Lambda_k(s_\star, \delta_\star, B)$, we have

$$|\nabla_{\tau,\xi} A_j| \lesssim_B |\xi|^{-1} + 2^{-k} \lesssim_B |(\tau,\xi)|^{-1}, \quad j = 0, \dots, N-1,$$

For the second inequality we use (2.10). This gives (3.42) when $|\alpha| = 1$.

To show (3.42) for $2 \leq |\alpha| \leq 2d + 2$, we use backward induction. By (2.28) we note $A_{N-1} = 0$, so (3.42) trivially holds when j = N - 1. We now assume that (3.42) holds true if $j_0 + 1 \leq j \leq N - 1$ for some j_0 . Lemma 3.7, (2.10), and the induction assumption show $\partial_{\tau,\xi}^{\alpha'} B_{j_0} = O(|(\tau,\xi)|^{-1-|\alpha'|})$ for $1 \leq |\alpha'| \leq 2d + 1$. Concerning D_{j_0} , observe that $\partial_{\xi}^{\alpha'} (G^{(j_0)}(\sigma(\tilde{\Xi})))$ is given by a sum of the terms

$$G^{(j)}(\sigma(\tilde{\Xi}))\prod_{n=1}^{j-j_0}\partial_{\xi}^{\alpha'_n}(\sigma(\tilde{\Xi})),$$

where $j \geq j_0$ and $\alpha'_1 + \cdots + \alpha'_{j-j_0} = \alpha'$. Hence, Lemma 3.7, (3.43), and (2.10) give $\partial_{\xi}^{\alpha'} D_{j_0} = O(|(\tau,\xi)|^{-1-|\alpha'|})$ for $1 \leq |\alpha'| \leq 2d+1$. Therefore, combining the estimates for B_{j_0} and D_{j_0} , we get $\partial_{\tau,\xi}^{\alpha'} \nabla_{\tau,\xi} A_{j_0} = O(|(\tau,\xi)|^{-1-|\alpha'|})$. This proves (3.42) for $j = j_0$.

Before proving (3.41), we first note

(3.44)
$$|\partial_{\xi}^{\alpha}(\mathcal{E}_{j}(\tilde{\Xi}))| \lesssim_{B} \delta_{\star}^{j} |\xi|^{-|\alpha|}, \quad |\alpha| \le 2d+2$$

for j = 1, ..., N. This can be shown by a routine computation. Indeed, differentiating (3.4), and using Lemma 3.7 and (2.32), one can easily see (3.44) since $|\sigma(\tilde{\Xi}) - \delta_0 \mu| \leq \delta_0$.

To show (3.41) for $0 \le \ell \le N - 2$, we again use backward induction. Observe that (3.41) holds for $\ell = N, N - 1$, and assume that (3.41) holds for $j + 1 \le \ell \le N$ for some $j \le N - 2$. By (3.6) we have

$$2^{-k}\mathfrak{g}_{\mu}^{j} = \delta_{*}^{N-j}A_{j} - \sum_{j+1 \le \ell \le N} (2^{-k}\mathfrak{g}_{\mu}^{\ell})(\mathcal{E}_{1})^{\ell-j}/(\ell-j)! + 2^{-k}y_{\mu}^{N}\mathcal{E}_{N-j}.$$

Thus, by Lemma 3.8 and (3.44), we get (3.41) with $\ell = j$. This completes the proof of Lemma 3.2.

3.4. **Proof of Lemma 3.4.** Lemma 3.4 can be proved in a similar way as the previous subsection. So, we shall be brief.

By Lemma 3.2 we have $C^{-1}\mathfrak{a}_{\nu}^{\mu,n} \in \mathfrak{A}_k(s_\star, \delta_\star)$ for a constant $C \geq 1$, so it suffices to show $C^{-1}\mathfrak{a}_{\nu,1}^{\mu,n} \in \mathfrak{A}_k(s_\star, \delta_\star)$ for some $C \geq 1$. The support condition (2.6) is obvious, so we need only to show (2.7) with $\mathfrak{a} = \mathfrak{a}_{\nu,1}^{\mu,n}$, $\delta = \delta_\star$, and $s_0 = s_\star$. Moreover, recalling (3.23), it is enough to consider the additional factor only, i.e., to show

$$\left|\partial_{\tau,\xi}^{\alpha} \left(\beta_0 \left(\frac{\left(\delta_{\star}^{-N} 2^{-k} \mathfrak{g}_{\mu}^0(\mathcal{L}_{s_{\star}}^{\delta_{\star}}(\tau,\xi))\right)^{2(N-1)!}}{C_0^{2N!} \delta_{\star}^{-2N!} \bar{\mathfrak{G}}_N^{\mu}(s, \widetilde{\mathcal{L}}_{s_{\star}}^{\delta_{\star}}\xi)}\right)\right)\right| \lesssim |(\tau,\xi)|^{-|\alpha|}$$

for $(\tau,\xi) \in \operatorname{supp} \mathfrak{a}_{\nu,1}^{\mu,n}(s,t,\mathcal{L}_{s_{\star}}^{\delta_{\star}}\cdot)$. Since $\delta_{\star}^{-2N!}\overline{\mathfrak{G}}_{N}^{\mu} \gtrsim 1$ on $\operatorname{supp}_{s,\xi} \mathfrak{a}_{\nu,1}^{\mu,n}$, one can obtain the estimate in the same way as in the proof of Lemma 3.2.

3.5. Sharpness of Theorem 1.3. Before closing this section, we show the optimality of the regularity exponent α in Theorem 1.3.

Proposition 3.9. Suppose (1.4) holds for $\psi(0) \neq 0$. Then $\alpha \leq 2/p$.

Proof. We write $\gamma = (\gamma_1, \ldots, \gamma_d)$. Via an affine change of variables, we may assume $\gamma_1(0) = 0$ and $\gamma'_1(s) \neq 0$ on an interval $J = [-\delta_0, \delta_0]$ for $0 < \delta_0 \ll 1$. Since $\psi(0) \neq 0$, we may also assume $\psi \ge 1$ on J.

We choose $\zeta_0 \in \mathcal{S}(\mathbb{R})$ such that $\operatorname{supp} \widehat{\zeta}_0 \subset [-1,1]$ and $\zeta_0 \geq 1$ on $[-r_1,r_1]$ where $r_1 = 1 + 2 \max\{|\gamma(s)| : s \in J\}$. Denoting $\overline{x} = (x_1, \ldots, x_{d-1})$ and $\overline{\gamma}(t) = (\gamma_1(t), \ldots, \gamma_{d-1}(t))$, we define

$$\bar{\mathcal{A}}_t h(x) = \int e^{it\lambda\gamma_d(s)} \zeta_0(x_d - t\gamma_d(s)) h(\bar{x} - t\bar{\gamma}(s)) \psi(s) \, ds.$$

Let $\zeta \in C_c^{\infty}((-2,2))$ be a positive function such that $\zeta = 1$ on [-1,1]. For a positive constant $c \ll \delta_0$, let $g_1(\bar{x}) = \sum_{\nu \in \lambda^{-1} \mathbb{Z} \cap [-c,c]} \zeta(\lambda | \bar{x} + \bar{\gamma}(\nu) |)$. We consider

$$g(\bar{x}) = e^{-i\lambda\varphi(x_1)}g_1(\bar{x}).$$

where $\varphi(s) = \gamma_d \circ (-\gamma_1)^{-1}(s)$. We claim that, if c is small enough,

 $\begin{aligned} (3.45) & |\bar{\mathcal{A}}_t g(x)| \gtrsim 1, \qquad (x,t) \in S_c, \\ \text{where } S_c &= \{(x,t) : |\bar{x}| \leq c\lambda^{-1}, |x_d| \leq c, |t-1| \leq c\lambda^{-1}\}. \text{ To show this, note} \\ \bar{\mathcal{A}}_t g(x) &= \int e^{i\lambda(t\gamma_d(s) - \varphi(x_1 - t\gamma_1(s)))} \zeta_0(x_d - t\gamma_d(s)) g_1(\bar{x} - t\bar{\gamma}(s)) \, \psi(s) \, ds. \end{aligned}$

Let
$$(x,t) \in S_c$$
. Then, $\operatorname{supp} g_1(\bar{x} - t\bar{\gamma}(\cdot)) \subset [-C_1c, C_1c]$ for some $C_1 > 0$. Since $\varphi(s) = \gamma_d \circ (-\gamma_1)^{-1}(s)$, by the mean value theorem we see $|\varphi(x_1 - t\gamma_1(s)) - \gamma_d(s)| \leq |\varphi(x_1 - t\gamma_1(s)) - \gamma_d(s)|$

 $\varphi(s) = \frac{1}{4}\varphi(-\gamma_1)^{-1}$ (s), by the mean value theorem we see $|\varphi(x_1 - v_1(s))^{-1}|_{d}(s)| \le 2r_0c\lambda^{-1}$ where $r_0 = 10r_1 \max\{|\partial_s\varphi(s)|: s \in (-\gamma_1)(J_*)\}$ and $J_* = [-(C_1+1)c, (C_1+1)c]$. Thus, we have

(3.46)
$$|t\gamma_d(s) - \varphi(x_1 - t\gamma_1(s))| \le 3r_0c\lambda^{-1}.$$

Besides, if λ is sufficiently large, $g_1(\bar{x}-t\bar{\gamma}(s)) = \sum_{\nu\in\lambda^{-1}\mathbb{Z}\cap[-c,c]}\zeta(\lambda|\bar{x}-(t-1)\bar{\gamma}(s)+\bar{\gamma}(\nu)-\bar{\gamma}(s)|) \gtrsim 1$ if $s\in[-c/2,c/2]$. Since $\operatorname{supp} g_1(\bar{x}-t\bar{\gamma}(\cdot))\subset J$ with c small enough and $\zeta_0(x_d-t\gamma_d(s))\geq 1$, we get $\int \zeta_0(x_d-t\gamma_d(s))g_1(\bar{x}-t\bar{\gamma}(s))\psi(s)\,ds\gtrsim 1$. Therefore, (3.45) follows by (3.46) if c is small enough, i.e., $c\ll 1/(3r_0)$.

We set $f(x) = e^{-i\lambda x_d}\zeta_0(x_d)g(\bar{x})$. Then, $\chi(t)\mathcal{A}_t f(x) = e^{-i\lambda x_d}\chi(t)\bar{\mathcal{A}}_t g(x)$. By our choice of ζ_0 , supp $\widehat{f} \subset \{\xi : |\xi_d + \lambda| \le 1\}$, so supp $\mathcal{F}(\chi(t)\mathcal{A}_t f) \subset \{(\tau,\xi) : |\xi_d + \lambda| \le 1\}$. This gives

(3.47)
$$\lambda^{\alpha} \| \chi(t) \mathcal{A}_t f \|_{L^p(\mathbb{R}^{d+1})} \lesssim \| \chi(t) \mathcal{A}_t f \|_{L^p_{\alpha}(\mathbb{R}^{d+1})}.$$

26

Indeed, $\lambda^{\alpha} \| \chi(t) \mathcal{A}_t f \|_{L^p(\mathbb{R}^{d+1})} \lesssim \| \chi(t) \mathcal{A}_t f \|_{L^p(\mathbb{R}_{t,\bar{x}};L^p_{\alpha}(\mathbb{R}_{x_d}))}$ by Mihlin's multiplier theorem in x_d . Similarly, one also sees $\| F \|_{L^p(\mathbb{R}_{t,\bar{x}};L^p_{\alpha}(\mathbb{R}_{x_d}))} \leq C \| F \|_{L^p_{\alpha}(\mathbb{R}^{d+1})}$ for $\alpha \geq 0$ and any F. Combining those inequalities gives (3.47).

From (3.45) we have $\|\chi(t)\mathcal{A}_t f\|_p = \|\chi(t)\bar{\mathcal{A}}_t g\|_p \ge C\lambda^{-d/p}$. Note that $\operatorname{supp} g$ is contained in a $O(\lambda^{-1})$ -neighborhood of $-\bar{\gamma}$, so it follows that $\|f\|_p \lesssim \lambda^{-(d-2)/p}$. Therefore, by (3.47) the inequality (1.4) implies $\lambda^{\alpha}\lambda^{-d/p} \lesssim \lambda^{-(d-2)/p}$. Taking $\lambda \to \infty$ gives $\alpha \le 2/p$.

4. L^p Sobolev regularity

In this section we prove Theorem 1.1, whose proof proceeds in a similar way as that of Theorem 1.3. However, we provide some details to make it clear how the optimal bounds are achieved. Since there are no t, τ variables for the symbols, the proof is consequently simpler but some modifications are necessary.

For a large $B \ge 1$, we assume

(4.1)
$$\max_{0 \le j \le 2d} |\gamma^{(j)}(s)| \le B, \qquad s \in I.$$

Let $2 \leq L \leq d$. For γ satisfying $\mathfrak{V}(L, B)$ we say $\bar{a} \in \mathbb{C}^{d+1}(\mathbb{R}^{d+1})$ is a symbol of type (k, L, B) relative to γ if supp $\bar{a} \subset I \times \mathbb{A}_k$, $\mathfrak{N}(L, B)$ holds for γ on supp \bar{a} , and

(4.2)
$$|\partial_s^j \partial_{\xi}^{\alpha} \bar{a}(s,\xi)| \le B|\xi|^{-|\alpha|}$$

for $0 \leq j \leq 1$ and $|\alpha| \leq d+1$. As before, Theorem 1.1 is a straightforward consequence of the following. We denote $\mathcal{A}[\gamma, \bar{a}] = \mathcal{A}_1[\gamma, \bar{a}]$.

Theorem 4.1. Suppose $\gamma \in C^{2d}(I)$ satisfies (4.1) and \bar{a} is a symbol of type (k, L, B) relative to γ for some $B \ge 1$. Then, if p > 2(L-1), for a constant C = C(B)

(4.3)
$$\|\mathcal{A}[\gamma,\bar{a}]f\|_{L^{p}(\mathbb{R}^{d})} \leq C2^{-k/p}\|f\|_{L^{p}(\mathbb{R}^{d})}.$$

In order to prove Theorem 1.1, we consider $\bar{a}_k(s,\xi) := \psi(s)\beta(2^{-k}|\xi|)$, where $\beta \in C_c^{\infty}((1/2,4))$. By (1.1) \bar{a}_k is a symbol of type (k,d,B) relative to γ for some B, thus Theorem 4.1 gives (4.3) for p > 2(d-1). The estimate (4.3) for each dyadic pieces can be put together by the result in [22]. So, we get (1.2) with $\alpha = \alpha(p)$ when p > 2(d-1) (e.g., see [2]).

Interpolation with $\|\mathcal{A}[\gamma, \bar{a}_k]f\|_2 \leq 2^{-k/d} \|f\|_2$ which follows from (1.3) gives $\|\mathcal{A}[\gamma, \bar{a}_k]f\|_p \leq_B 2^{-\alpha k} \|f\|_p$ for $\alpha \leq \alpha(p)$ with strict inequality when $p \in (2, 2(d-1)]$. Using those estimates, we can prove Corollary 1.2. Indeed, if γ is a curve of maximal type $\ell > d$, a typical scaling argument gives $\|\mathcal{A}[\gamma, \bar{a}_k]f\|_p \leq_B 2^{-\min(\alpha(p), 1/\ell)k} \|f\|_p$ for $p \neq \ell$ when $\ell \geq 2d-2$, and for $p \in [2, 2\ell/(2d-\ell)) \cup (2d-2, \infty)$ when $d < \ell < 2d-2$. As above, one can combine the estimates ([22]) to get (1.2).

4.1. **Proof of Theorem 4.1.** The case L = 2 is easy. Since \bar{a} is a symbol of type (k, 2, B) relative to γ , van der Corput's lemma and Plancherel's theorem give (4.3) for p = 2. Interpolation with L^{∞} estimate shows (4.3) for $p \ge 2$. When $L \ge 3$, we have the following, which immediately yields Theorem 4.1.

Proposition 4.2. Let $3 \le N \le d$. Suppose Theorem 4.1 holds for L = N - 1. Then Theorem 4.1 holds true with L = N.

To prove the proposition, we fix $N \in [3, d]$ and γ satisfying $\mathfrak{V}(N, B)$, and \bar{a} of type (k, N, B) relative to γ .

For s_0 and $\delta > 0$ such that $I(s_0, \delta) \subset I$, let

$$\bar{\Lambda}_k(s_0,\delta,B) = \bigcap_{1 \le j \le N-1} \left\{ \xi \in \mathbb{A}_k : |\langle \gamma^{(j)}(s_0),\xi \rangle| \le B2^{k+5} \delta^{N-j} \right\}.$$

By $\bar{\mathfrak{A}}_k(s_0, \delta)$ we denote a collection of $\bar{\mathfrak{a}} \in \mathbf{C}^{d+1}(\mathbb{R}^{d+1})$ such that $\operatorname{supp} \bar{\mathfrak{a}} \subset I(s_0, \delta) \times \bar{\Lambda}_k(s_0, \delta, B)$ and $|\partial_s^j \partial_{\xi}^{\alpha} \bar{\mathfrak{a}}(s, \widetilde{\mathcal{L}}_{s_0}^{\delta} \xi)| \leq B \delta^{-j} 2^{-k|\alpha|}$ for $0 \leq j \leq 1$ and $|\alpha| \leq d+1$.

The next lemma which plays the same role as Lemma 2.8 can be shown through routine adaptation of the proof of Lemma 2.8.

Lemma 4.3. Let $\bar{\mathfrak{a}} \in \bar{\mathfrak{A}}_k(s_0, \delta)$ and $j_* = \log(2^k \delta^N)$. Suppose (2.20) holds on supp $\bar{\mathfrak{a}}$. Then, there exist constants $C, \tilde{B} \ge 1$, and $\delta' > 0$ depending on B, and symbols $\bar{\mathfrak{a}}_1, \ldots, \bar{\mathfrak{a}}_{l_*}$ of type $(j, N - 1, \tilde{B})$ relative to $\gamma_{s_0}^{\delta}$, such that

$$\|\mathcal{A}[\gamma,\bar{\mathfrak{a}}]f\|_{L^{p}(\mathbb{R}^{d})} \leq C\delta \sum_{1 \leq l \leq C} \|\mathcal{A}[\gamma_{s_{0}}^{\delta},\bar{\mathfrak{a}}_{l}]\tilde{f}_{l}\|_{L^{p}(\mathbb{R}^{d})},$$

 $\|\tilde{f}_l\|_p = \|f\|_p$, and $j \in [j_* - C, j_* + C]$ as long as $0 < \delta < \delta'$.

The order of necessary regularity on γ is reduced since $\bar{\mathfrak{a}}$ is independent of τ, t . Actually, we may take $\tilde{a}(s,\xi) = \bar{\mathfrak{a}}(\delta s + s_0, \delta^{-N} \widetilde{\mathcal{L}}_{s_0}^{\delta} \xi)$ while following the *Proof of Lemma 2.8* since validity of (4.2) is clear for $\bar{a} = \tilde{a}$.

Using η_N (see (2.26)), we break

$$\mathcal{A}[\gamma, \bar{a}] = \mathcal{A}[\gamma, \bar{a}\eta_N] + \mathcal{A}[\gamma, \bar{a}(1-\eta_N)].$$

Note that $C^{-1}\bar{a}(1-\eta_N)$ is of type (k, N-1, B') relative to γ for some large constants B' and C, so we may apply the assumption to $\mathcal{A}[\gamma, \bar{a}(1-\eta_N)]f$. Consequently, we have the estimate (4.3) for $\bar{a} = \bar{a}(1-\eta_N)$ if p > 2N-4.

To obtain the estimate for $\mathcal{A}[\gamma, \bar{a}\eta_N]$, as before, we may assume $\sup \bar{a}\eta_N \subset I(s_0, \delta_*) \times \bar{\Gamma}_k$ for some s_0 and a small δ_* . Here, $\bar{\Gamma}_k$ is defined in the same way as Γ_k by replacing $a\eta_N$ by $\bar{a}\eta_N$. Since (2.27) holds on $\supp(\bar{a}\eta_N)$, we may work under the same *Basic assumption* as in Section 2.3. That is to say, we have σ on $\bar{\Gamma}_k$ satisfying (2.28) and $\sigma(\xi) \in I(s_0, \delta_*)$ for $\xi \in \bar{\Gamma}_k$. Furthermore, $\sigma \in C^{d+1}$ since $\gamma \in C^{2d}(I)$, and (2.29) holds for $\xi \in \bar{\Gamma}_k$ and $|\alpha| \leq d + 1$. Thus, (4.2) remains valid for the symbols given subsequently by decomposing \bar{a} with cutoff functions associated with σ , and $\bar{\mathfrak{G}}_N^{\mu}$.

Apparently, $C^{-1}\bar{a}\eta_N \in \bar{\mathfrak{A}}_k(s_0, \delta_*)$ for a constant $C = C(B, \delta_*)$, therefore the proof of Proposition 4.2 is completed if we show the following.

Proposition 4.4. Let $3 \leq N \leq d$ and $\bar{\mathfrak{a}} \in \bar{\mathfrak{A}}_k(s_0, \delta_*)$ with $\operatorname{supp}_{\xi} \bar{\mathfrak{a}} \subset \bar{\Gamma}_k$. Suppose Theorem 4.1 holds for L = N - 1. Then, if p > 2(N - 1), we have (4.3).

We prove Proposition 4.4 using the next, which corresponds to Proposition 2.10. In what follows, we denote $\mathcal{A}[\bar{a}] = \mathcal{A}[\gamma, \bar{a}]$.

Proposition 4.5. Let δ_0 and δ_1 satisfy (2.31). For μ such that $\delta_0 \mu \in I(s_0, \delta_*) \cap \delta_0 \mathbb{Z}$, let $\bar{\mathfrak{a}}^{\mu} \in \bar{\mathfrak{A}}_k(\delta_0 \mu, \delta_0)$ with $\operatorname{supp} \bar{\mathfrak{a}}^{\mu} \subset I(s_0, \delta_*) \times \bar{\Gamma}_k$. Suppose Theorem 4.1 holds for L = N - 1. Then, if $p \in (2N - 2, \infty)$, there are constants $\epsilon_0 > 0$, $C_0 = C_0(\epsilon_0, B) \geq 2$, and symbols $\bar{\mathfrak{a}}_{\nu} \in \bar{\mathfrak{A}}_k(\delta_1 \nu, \delta_1)$ with $\operatorname{supp} \bar{\mathfrak{a}}_{\nu} \subset I(s_0, \delta_*) \times \bar{\Gamma}_k$, $\nu \in \cup_{\mu} \mathfrak{J}_0^{\mu}$, such that

$$\left(\sum_{\mu} \|\mathcal{A}[\bar{\mathfrak{a}}^{\mu}]f\|_{p}^{p}\right)^{\frac{1}{p}} \leq C_{0}\left(\delta_{1}/\delta_{0}\right)^{\frac{N}{p}-1+\epsilon_{0}}\left(\sum_{\nu} \|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}]f\|_{p}^{p}\right)^{\frac{1}{p}} + C_{0}\delta_{0}^{-\frac{N}{p}+1}2^{-\frac{k}{p}}\|f\|_{p}.$$

28

Let δ' be given as in Lemma 4.3, and let $\delta_{\circ} > 0$ be a positive constant such that

(4.4)
$$\delta_{\circ} \le \min\{\delta', (2^{7d}B^6)^{-N}C_0^{-2N/\epsilon_0}\}$$

Proof of Proposition 4.4. Set $\delta_0 = \delta_0$, and let $\delta_1, \ldots, \delta_J$ be given by (2.33). Then, applying Proposition 4.5 iteratively up to *J*-th step (see Section 2.4), we have symbols $\bar{\mathfrak{a}}_{\nu} \in \mathfrak{A}_k(\delta_J \nu, \delta_J), \, \delta_J \nu \in I(s_0, \delta_0)$, such that

$$\left\|\mathcal{A}[\bar{\mathfrak{a}}]f\right\|_{p} \leq C_{0}^{J}\delta_{J}^{\frac{N}{p}-1+\epsilon_{0}} \left(\sum_{\nu} \|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}]f\|_{p}^{p}\right)^{1/p} + 2^{-\frac{k}{p}}\delta_{0}^{-\frac{N}{p}+1-\epsilon_{0}} \sum_{0 \leq j \leq J-1} C_{0}^{j+1}\delta_{j}^{\epsilon_{0}}\|f\|_{p}.$$

By (4.4) and (2.33), $\delta_j \leq C_0^{-2((N+1)/N)^j N/\epsilon_0}$, $0 \leq j \leq J-1$. So, $\sum_{j=0}^{J-1} C_0^{j+1} \delta_j^{\epsilon_0} \leq C_1$ for a constant C_1 , and $C_0^J \delta_J^{\epsilon_0} \leq C_1$. Thus, the matter is now reduced to showing

$$\left(\sum_{\nu} \|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}]f\|_{L^{p}(\mathbb{R}^{d})}^{p}\right)^{1/p} \lesssim_{B} 2^{-\frac{k}{N}} \|f\|_{L^{p}(\mathbb{R}^{d})}, \qquad 2 \le p \le \infty,$$

which corresponds to (2.35). The case $p = \infty$ follows from the estimate $\|\mathcal{A}[\bar{\mathfrak{a}}]f\|_{L^{\infty}}$ $\leq C\delta \|f\|_{L^{\infty}}$ when $\bar{\mathfrak{a}} \in \bar{\mathfrak{A}}_k(s_0, \delta)$ for some s_0, δ (cf. (2.15)). One can obtain this in the same manner as in the proof of Lemma 2.7. The case p = 2 can be handled similarly as before, using Plancherel's theorem and van der Corput's lemma combined with Lemma 2.11 and (2.27).

The proof of Proposition 4.5 is similar to that of Proposition 2.10. Instead of (2.40) we use the estimate (2.41), in which the exponent is adjusted to the sharp Sobolev regularity estimate. However, a similar approach breaks down if one tries to obtain the local smoothing estimate (1.4) with the optimal regularity $\alpha = 2/p$. To do so, we need the inequality (2.39) for 4N - 2 . However, there is no such estimate available when <math>N = 2.

4.2. Proof of Proposition 4.5. Let $\bar{\mathfrak{a}}^{\mu} \in \bar{\mathfrak{A}}_k(\delta_0\mu, \delta_0)$. For $\nu \in \mathfrak{J}_n^{\mu}$, set

$$\bar{\mathfrak{a}}_{\nu}^{\mu,n} = \bar{\mathfrak{a}}^{\mu} \times \begin{cases} \beta_0 \left(\delta_1^{-2N!} \,\bar{\mathfrak{G}}_N^{\mu} \right) \zeta(\delta_1^{-1} s - \nu), & n = 0, \\ \beta_N \left((2^n \delta_1)^{-2N!} \,\bar{\mathfrak{G}}_N^{\mu} \right) \zeta(2^{-n} \delta_1^{-1} s - \nu), & n \ge 1, \end{cases}$$

(see (3.22)). Let $\bar{\mathbf{y}}_{\mu} = (y_{\mu}^{1}, \dots, y_{\mu}^{N})$, and let $\bar{\mathcal{D}}_{\delta}$ denote the $N \times N$ matrix $(\delta^{1-N}\bar{e}_{1}, \delta^{2-N}\bar{e}_{2}, \dots, \delta^{0}\bar{e}_{N})$ where \bar{e}_{j} is the *j*-th standard unit vector in \mathbb{R}^{N} . Recalling (3.19), we consider a linear map

$$\bar{\mathbf{Y}}^{\delta_0}_{\mu}(\xi) = \left(2^{-k}\bar{\mathcal{D}}_{\delta_0}\bar{\mathbf{y}}_{\mu}, y_{N+1}, \dots, y_d\right).$$

Let **r** denote the curve \mathbf{r}_{\circ}^{N} . Note that (3.11) and (3.12) hold on $\operatorname{supp} \bar{\mathfrak{a}}_{\nu}^{\mu,n}$. Similarly as in *Proof of Lemma 3.3*, we see $|\langle \bar{\mathbf{y}}_{\mu}, \mathbf{r}^{(j)}((2^{n}\delta_{1}/\delta_{0})\nu-\mu)\rangle| \lesssim 2^{k}(2^{n}\delta_{1}/\delta_{0})^{N-j}$ for $1 \leq j \leq N-1$ and $2^{k-2}/B \leq |\langle \bar{\mathbf{y}}_{\mu}, \mathbf{r}^{(N)} \rangle| \leq CB2^{k}$ on $\operatorname{supp}_{\xi} \bar{\mathfrak{a}}_{\nu}^{\mu,n}$. Thus, as before (cf. (3.20)), we have

$$\bar{\mathbf{Y}}^{\delta_0}_{\mu}(\operatorname{supp}_{\xi} \bar{\mathfrak{a}}^{\mu,n}_{\nu}) \subset \mathbf{s}\Big(\frac{2^n \delta_1}{\delta_0} \nu - \mu, \, C\frac{2^n \delta_1}{\delta_0}, \, CB; \, \mathbf{r}^N_{\circ}\Big) \times \mathbb{R}^{d-N}$$

for some C > 0. Note $\operatorname{supp} \mathcal{F}(\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f) \subset \operatorname{supp}_{\xi} \bar{\mathfrak{a}}_{\nu}^{\mu,n}$. Therefore, changing variables, by (2.41) with N replaced by N-1 and its cylindrical extension (e.g.,(2.42)), we get

(4.5)
$$\left\|\sum_{\nu\in\mathfrak{J}_n^{\mu}}\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f\right\|_p \leq C_0 \left(2^n \delta_1/\delta_0\right)^{\frac{N}{p}-1+\epsilon_0} \left(\sum_{\nu\in\mathfrak{J}_n^{\mu}}\left\|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f\right\|_p^p\right)^{1/p}\right\|_{\infty}$$

for $2N-2 (cf. (3.16)). Since <math>\mathcal{A}[\bar{\mathfrak{a}}^{\mu}]f = \sum_{\nu \in \mathfrak{J}_n^{\mu}} \mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f$, by Minkowski's inequality and (4.5), we have $(\sum_{\mu} \|\mathcal{A}[\bar{\mathfrak{a}}^{\mu}]f\|_{p}^{p})^{1/p}$ bounded by

$$\sum_{n\geq 0} \bar{\mathbf{E}}_n := C_0 \sum_{n\geq 0} \left(2^n \delta_1 / \delta_0 \right)^{\frac{N}{p} - 1 + \epsilon_0} \left(\sum_{\mu} \sum_{\nu \in \mathfrak{J}_n^{\mu}} \|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu, n}] f\|_p^p \right)^{1/p}.$$

The proof of Lemma 3.2 also shows $C^{-1}\bar{\mathfrak{a}}_{\nu}^{\mu,n} \in \bar{\mathfrak{A}}_{k}(2^{n}\delta_{1}\nu,2^{n}\delta_{1})$ for a positive constant C. Therefore, the matter is reduced to obtaining

(4.6)
$$\left(\sum_{\mu}\sum_{\nu\in\mathfrak{J}_{n}^{\mu}}\|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f\|_{L^{p}(\mathbb{R}^{d})}^{p}\right)^{1/p} \lesssim_{B} (2^{n}\delta_{1})^{1-\frac{N}{p}}2^{-\frac{k}{p}}\|f\|_{L^{p}(\mathbb{R}^{d})}, \qquad n \ge 1$$

for p > 2(N-2). This gives $\sum_{n\geq 1} \bar{\mathbf{E}}_n \lesssim_B \delta_0^{-N/p+1} 2^{-k/p} ||f||_p$ since $2^n \delta_1 \leq C \delta_0$. The proof of (4.6) is similar with that of (3.24). Since $C^{-1} \bar{\mathfrak{a}}_{\nu}^{\mu,n} \in \bar{\mathfrak{A}}_k(2^n \delta_1 \nu, 2^n \delta_1)$, we have $\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f = \mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]P_{2^n\delta_1\nu}^{2^n\delta_1}f$. Besides, (3.28) or (3.29) for some $0 \leq j \leq N-2$ holds on supp $\bar{\mathfrak{a}}_{\mu,n}^{\mu,n}$. Thus, (2.20) holds with $\delta = 2^n \delta_1$ for some B' on supp $\bar{\mathfrak{a}}_{\mu,n}^{\mu,n}$ for $n \geq 1$ (see Proof of Lemma 3.5). Therefore, applying Lemma 4.3 to $\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f$ and then the assumption (Theorem 4.1 with L = N - 1), we obtain

$$\|\mathcal{A}[\bar{\mathfrak{a}}_{\nu}^{\mu,n}]f\|_{L^{p}(\mathbb{R}^{d})} \lesssim_{B} (2^{n}\delta_{1})^{1-\frac{N}{p}} 2^{-\frac{k}{p}} \|P_{2^{n}\delta_{1}\nu}^{2^{n}\delta_{1}}f\|_{p}$$

This combined with (3.33) gives (4.6) as desired.

Acknowledgement. H. Ko was supported by NRF2019R1A6A3A01092525. S. Lee and S. Oh were supported by NRF2021R1A2B5B02001786. The authors would like to thank N. Bez, A. Seeger, and J. Wright for their valuable comments.

References

- [1] D. Beltran, S. Guo, J. Hickman, A. Seeger, Sharp L^p bounds for the helical maximal function, arXiv:2102.08272.
- _____, Sobolev improving for averages over curves in \mathbb{R}^4 , Adv. Math. **393** (2021) 108089.
- [3] J. Bourgain, Averages in the plane over convex curves and maximal operators, J. Analyse. Math. 47 (1986), 69–85.
- [4] J. Bourgain, C. Demeter, The proof of the ℓ^2 decoupling conjecture, Ann. of Math. 182 (2015), 351 - 389.
- [5] J. Bourgain, C. Demeter, L. Guth, Proof of the main conjecture in Vinogradov's mean value theorem for degrees higher than three, Ann. of Math. 184 (2016), 633-682.
- [6] M. Christ, Failure of an endpoint estimate for integrals along curves, Fourier analysis and partial differential equations (Miraflores de la Sierra, 1992), Stud. Adv. Math., CRC, Boca Raton, FL, 1995, pp.163-168.
- [7]____, Convolution, curvature, and combinatorics. A case study, Int. Math. Res. Not. 19 (1998), 1033-1048.
- [8] M. Christ, A. Nagel, E. M. Stein, S. Wainger, Singular and maximal Radon transforms: analysis and geometry, Ann. of Math. (2) 150 (1999), 489-577.
- P. Constantin, J.-C. Saut, Local smoothing properties of dispersive equations, J. Amer. Math. Soc. 1 (1988), 413–439.
- [10] S. Guo, Z. Li, P.-L. Yung, P. Zorin-Kranich, A short proof of ℓ^2 decoupling for the moment curve, Amer. J. Math. 143 (2021), 1983-1998.
- [11] L. Guth, H. Wang, R. Zhang, A sharp square function estimate for the cone in \mathbb{R}^3 , Ann. of Math. 192 (2020), 551–581.
- [12] S. Ham, S. Lee, Restriction estimates for space curves with respect to general measure, Adv. Math. 254 (2014), 251-279.
- I. Ikromov, M. Kempe, D. Müller, Estimates for maximal functions associated with hyper-[13]surfaces in \mathbb{R}^3 and related problems of harmonic analysis, Acta Math. **204** (2010), 151–271.
- [14] H. Ko, S. Lee, S. Oh, Maximal estimate for average over space curve, Invent. math. (2022), advance online publication, https://doi.org/10.1007/s00222-021-01094-6.

- [15] S. Lee, Endpoint estimates for the circular maximal function, Proc. Amer. Math. Soc. 131 (2003), 1433–1442.
- [16] S. Lee, A. Vargas, On the cone multiplier in \mathbb{R}^3 , J. Funct. Anal. **263** (2012), 925–940.
- [17] J. M. Marstrand, Packing circles in the plane, Proc. London Math. Soc. 55 (1987), 37–58.
- [18] T. Mitsis, On a problem related to sphere and circle packing, J. London Math. Soc. (2) 60 (1999), 501–516.
- [19] G. Mockenhaupt, A. Seeger, C. D. Sogge, Wave front sets, local smoothing and Bourgain's circular maximal theorem, Ann. of Math. 136 (1992), 207–218.
- [20] D. Oberlin, H. Smith, C. D. Sogge, Averages over curves with torsion, Math. Res. Lett. (1998), 535–539.
- [21] D. H. Phong, E. M. Stein, Radon transforms and torsion, Internat. Math. Res. Notices 1991, no. 4, 49–60.
- [22] M. Pramanik, K.M. Rogers, A. Seeger, A Calderón-Zygmund estimate with applications to generalized Radon transforms and Fourier integral operators, Stud. Math. 202 (2011) 1–15.
- [23] M. Pramanik, A. Seeger, L^p Sobolev regularity of a restricted X-ray transform in \mathbb{R}^3 , Harmonic analysis and its applications, 47–64, Yokohama Publ., Yokohama 2006.
- [24] _____, L^p regularity of averages over curves and bounds for associated maximal operators, Amer. J. Math. **129** (2007), 61–103.
- [25] _____, L^p Sobolev estimates for a class of integral operators with folding canonical relations,
 J. Geom. Anal. (2020), https://doi.org/10.1007/s12220-020-00388-0.
- [26] W. Schlag, A generalization of Bourgain's circular maximal theorem, J. Amer. Math. Soc. 10 (1997), 103–122.
- [27] W. Schlag, C. D. Sogge, Local smoothing estimates related to the circular maximal theorem, Math. Res. Let. 4 (1997), 1–15.
- [28] A. Seeger, J. Wright, Problems on averages and lacunary maximal functions. Marcinkiewicz Centenary Volume, pp. 235–250, Banach Center Publ., vol. 95, Polish Acad. Sci. Inst. Math., Warsaw, 2011.
- [29] P. Sjölin, Regularity of solutions to the Schrödinger equation, Duke Math. J. 55 (1987), 699–715.
- [30] C. D. Sogge, Propagation of singularities and maximal functions in the plane, Invent. Math. 104 (1991), 349–376.
- [31] E. M. Stein, Maximal functions: spherical means, Proc. Nat. Acad. Sci. USA 73 (1976), 2174–2175.
- [32] _____, Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals, Princeton Univ. Press, Princeton, NJ, 1993.
- [33] E. M. Stein, S. Wainger, Problems in harmonic analysis related to curvature, Bull. Amer. Math. Soc. 84 (1978), 1239–1295.
- [34] B. Stovall, Endpoint bounds for a generalized Radon transform, J. Lond. Math. Soc. 80 (2009), 357–374.
- [35] T. Tao, A. Vargas, A bilinear approach to cone multipliers. II. Applications, Geom. Funct. Anal. 10 (2000), 216–258.
- [36] T. Tao, J. Wright, L^p improving bounds for averages along curves, J. Amer. Math. Soc. 16 (2003), 605–638.
- [37] T. Wolff, A Kakeya type problem for circles, Amer. J. Math. 119 (1997), 985–1026.
- [38] _____, Local smoothing type estimates on L^p for large p, Geom. Funct. Anal. 10 (2000), 1237–1288.

DEPARTMENT OF MATHEMATICAL SCIENCES AND RIM, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, REPUBLIC OF KOREA

Email address: kohr@snu.ac.kr

Email address: shklee@snu.ac.kr

Email address: dhtpdnr0220@snu.ac.kr