

The Science Case for LIGO-India

**M. Saleem^{1,7}, Javed Rana⁹, V. Gayathri^{5,6}, Aditya Vijaykumar²,
Srashti Goyal², Surabhi Sachdev⁹, Jishnu Suresh¹¹, S.
Sudhagar⁴, Arunava Mukherjee⁸, Gurudatt Gaur¹², Bangalore
Sathyaprakash⁹, Archana Pai⁵, Rana X Adhikari^{2,3}, P. Ajith²,
Sukanta Bose^{4,10}**

¹ Chennai Mathematical Institute, Siruseri 603103, Tamilnadu, India

² International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bangalore 560089, India

³ LIGO Laboratory, California Institute of Technology, USA

⁴ Inter-University Centre for Astronomy and Astrophysics (IUCAA), Post Bag 4, Ganeshkhind, Pune 411 007, India

⁵ Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India

⁶ Department of Physics, University of Florida, PO Box 118440, Gainesville, FL 32611-8440, USA

⁷ School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA

⁸ Saha Institute of Nuclear Physics, HBNI, 1/AF Bidhannagar, Kolkata-700064, India

⁹ Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

¹⁰ Department of Physics and Astronomy, Washington State University, 1245 Webster, Pullman, WA 99164-2814, USA

¹¹ Institute for Cosmic Ray Research (ICRR), The University of Tokyo, Kashiwa City, Chiba 277-8582, Japan

¹² Institute of Advanced Research, Gandhinagar 382 426, Gujarat, India

E-mail: sukanta@iucaa.in

10 January 2022

Abstract. The global network of gravitational-wave detectors has completed three observing runs with ~ 50 detections of merging compact binaries. A third LIGO detector, with comparable astrophysical reach, is to be built in India (LIGO-Aundha) and expected to be operational during the latter part of this decade. Multiple detectors operating at different parts of the globe will provide several pairs of interferometers with longer baselines and an increased network SNR. This will improve the sky localisation of GW events. Multiple detectors simultaneously in operation will also increase the baseline duty factor, thereby, leading to an improvement in the detection rates and, hence, the completeness of surveys. In this paper, we quantify the improvements due to the expansion of the LIGO Global Network (LGN) in the precision with which source properties will be measured. We also present examples of how this expansion will give a boost to tests of fundamental physics.

Contents

1	Introduction	3
1.1	Detectors	3
1.2	Simulations	3
2	CBC Detection rates	5
2.1	Detection criteria	5
2.2	Improvement in the effective duty-factor of a network	6
2.3	Detection rates	7
3	Parameter estimation	7
3.1	Improvement in errors for binary black hole events	8
3.1.1	Sky-localization:	9
3.1.2	Luminosity distance and inclination angle:	9
3.1.3	Source masses	10
3.2	Enhancement in matter effects: source classification and BNS properties .	10
4	Sky localisation and early warning	13
4.1	Early warning of binary neutron star mergers	15
5	Tests of GR	19
5.1	Improved constraints on deviations from GR	19
5.2	Constraints the nature of GW polarisations	20
6	Conclusions and summary	21

1. Introduction

The global network of gravitational-wave detectors (comprising the two LIGO interferometers [1] and the Virgo interferometer [2]) have completed three observing runs with ~ 50 detections of merging compact binaries [3]. A fourth detector in Japan [4] is now being commissioned and is expected to join the global network in 2022. A third LIGO detector with comparable astrophysical reach is being built in India [5] and is to be operational during the latter part of this decade. Multiple detectors operating at different parts of the globe will provide several pairs of interferometers with longer baselines and higher network SNRs than we have today. This will tremendously help improve the sky localization of GW events. Multiple detectors operating simultaneously, will also improve the duty factor of the network leading to improvements in the detection rates.

In this paper we quantify the improvements due to the addition of LIGO-Aundha to the LIGO Global Network (LGN). We quantitatively describe how the improvements can give better astrophysical insights about the source properties and how that improves our knowledge of physics and cosmology. We find that the addition of a new detector in India has substantial benefits to the scientific capabilities of the LGN.

1.1. Detectors

The LGN will consist of 3 interferometers in the upgraded configuration of Advanced LIGO (so-called A+) [6], with the third detector in Aundha, in the Hingoli district, in the eastern part of the state of Maharashtra, India. It is expected that the two LIGO detectors in the U.S. will be upgraded into this configuration in ~ 2026 and that the detector in Aundha will come online soon after. Following the existing naming convention $\ddagger$ the detector in India will be referred to as LIGO–Aundha (A). The LIGO Global Network with and without LIGO-Aundha will be denoted as AHL and HL, respectively.

1.2. Simulations

The compact binary coalescences (CBCs) observed in the GW window range in total mass from $3–150 M_{\odot}$. While most of the binary systems harbor primary objects with mass $< 45 M_{\odot}$, a few systems have the primary heavier than $45 M_{\odot}$. The recently published gravitational-wave transient catalog (GWTC-2) [3] considers several population mass distribution models to obtain the merger rates. We use a suitable mass model for the simulations [7] with the parameters inferred from the observed binary black-hole mergers. In this model, the primary mass follows a power-law distribution with some spectral index up to a certain maximum mass and a uniform Gaussian component with a finite width to account for high masses, together with a smoothing function at low masses to avoid a hard cut-off. The mass ratio follows a smoothed

$\ddagger$ where the detectors are named after the nearby town; LIGO–Hanford (H) and LIGO–Livingston (L)




Figure 1: Strain noise spectral density of the LIGO Interferometers. The Advanced LIGO/Virgo O3 noise, the Advanced LIGO design sensitivity, and the A+ sensitivity (for the LIGO detectors: Aundha, Hanford, and Livingston)

power-law distribution. We choose the median values of the hyper-parameters of these distributions inferred in [7] for simulations.

Astrophysical models suggest black holes in a binary system with isotropic spins in a dense environment such as globular clusters or galactic centers. At the same time, we expect black-hole spins to get aligned with the orbital angular momentum for isolated binaries [8, 9]. The black hole spin distribution uses a model that is a mixture of both these possibilities [7]. We use this model for drawing the spins of both the compact objects in a binary for our simulations. Besides, the binary sources are oriented uniformly and distributed uniformly over the sky and placed uniformly in co-moving volume up to a red-shift of 1.5 using the Planck 2015 cosmology [10].

The binary black hole (BBH) simulations described above are used in various studies below on quantifying the improvement in the performance of the network arising from its expansion to include LIGO-Aundha. These include a discussion of BBH detection rates in Section 2 and quantifying the improvement in the estimation of binary parameters in Section 3. In Sec. Section 4, we discuss the possibility of sending early warning alerts to electromagnetic and particle observatories before the epoch of binary coalescence. We make projections in Section 5 on how a detected BBH population can be used to place observational bounds on deviations from General Relativity (GR).

2. CBC Detection rates

Coalescing compact binaries involving neutron stars and black holes are, so far, the only GW sources detected, in the past GW observing runs [11, 3]. The inclusion of LIGO-Aundha in the LGN will boost the rate at which we detect such binaries. This enhancement will arise owing to improved sky-coverage, distance reach, and baseline duty factor, which is the effective observation period of a detector network. In this section, we quantitatively assess the improvement in the CBC detection rate (R_{det}) for AHL *vis à vis* the HL network.

We focus here on the stellar-mass BBHs, which are the main contributor to the menagerie of signals observed by LIGO-Virgo so far. Our analysis can be straightforwardly extended to classes of CBC sources that involve neutron stars.

For an astrophysical population of BBHs with a comoving constant merger-rate density r_{merg} in units of $\text{Gpc}^{-3}\text{yr}^{-1}$, the detection rates (per year) is given by $R_{\text{det}} = r_{\text{merg}} \times \langle VT \rangle$, where $\langle VT \rangle$ is the population-marginalized detection volume averaged over the period of observation for any given detector network (for more details, see [3, 12] and the references therein). The assumption that r_{merg} is non-evolving w.r.t redshift is a simplified assumption and hence could affect the rates we reported in this paper, however it has negligible impact on the rates comparison between two networks which is the goal of this study. The factor $\langle VT \rangle$ crucially depends on the number of detectors, their sensitivity as well as the search methodologies and their ability to treat the non-Gaussian noisy transients in the multi-detector data.

2.1. Detection criteria

We assume that the noise in any detector is Gaussian, with a vanishing mean, and is uncorrelated with the noise in any other detector. Then the network coherent SNR-squared is the sum of the SNR-squared of signals in the individual detectors [13]. A source is considered as detected by a network when the following conditions are met (a similar set of criteria is considered in [14]):

- (i) **Coherent network SNR criterion:** For an N-detector network, this criterion is $\sum_{k=1}^N \rho_k^2 \geq 12$ where ρ_k is the SNR at the k^{th} detector. The threshold of 12 is an arbitrary choice but reasonable because the false-alarm probability associated with this value will be low enough to make a confident detection case. This choice is conservative in the sense that there have been detections reported in past observing runs with network SNR below 12.
- (ii) **Multi-detector coincidence criterion:** $\sum_{k=1}^N \rho_k^2 \geq 12$ and $\rho_k > 4$ for any two of the N detectors.

Arguably, the simplest way to identify interesting detection candidates is to apply the first criterion. Its biggest advantage is that it allows for picking up sources that are loud enough in one detector but weak in the others, e.g., if located in their blind-spots.




Figure 2: [Left] Network duty factors of the HL and AHL networks as functions of the single-detector duty factor. [Right] The distribution of the detection rates of stellar-mass binary black-hole detection rates for the same two networks using the GWTC-2 population models.

This can happen since none of the detectors has identical orientation. Nevertheless, this criterion has a few limitations: For instance, a loud noise-transient in a single detector can give rise to a trigger that satisfies this network criterion and, therefore, gets misclassified as a detection candidate. On the other hand, if one requires that at least two detectors record a high enough SNR, such as what the second criterion above employs, then the false-alarm rate reduces significantly, albeit by sacrificing some degree of sky coverage.

2.2. Improvement in the effective duty-factor of a network

Duty factor of a detector (network) is defined as the fraction of clock time for which the detector (network) acquires science quality data. Assuming that each detector in the network has a duty factor of d_f , one can analytically compute the effective duty factor $d_{f\text{eff}}$ for each multi-detector network. For the multi-detector coincidence criterion, $d_{f\text{eff}}$ is the fraction of the observation period during which at least two of the detectors are simultaneously collecting science-quality data while for the network SNR criterion, it is the fraction of observing period when at least one of the detectors is observing in science mode. For an N -detector network, the effective duty factor is

$$d_{f\text{eff}} = \sum_{k=N_{\min}}^N {}^N C_k d_f^k (1 - d_f)^{N-k}, \quad (1)$$

where $N_{\min}$ is the minimum number of detectors required by the coincidence criterion. Figure 2 shows how the effective duty factor improves with the addition of LIGO-Aundha. Assuming 90% single-detector duty-factor, the AHL duty-factor gets boosted by a factor of ~ 1.2 compared to the HL network if one follows the criterion-(ii), while the improvement is only a one per-cent under criterion-(i), see e.g. Figure 2.

Network	Criterion (i)	Criterion (ii)
HL	$550.0^{+30.0}_{-29.0}$	$502.0^{+29.0}_{-27.0}$
AHL	$775.0^{+35.0}_{-35.0}$	$754.0^{+35.0}_{-33.0}$

Table 1: Detection rates (in yr^{-1}) of stellar-mass binary black holes in HL and AHL networks, assuming A+ sensitivity and a duty factor of 90% for all the detectors.

2.3. Detection rates

We perform extensive simulations to estimate the detection rates of merging compact binaries for the HL and AHL configurations. The published detections of binary black hole mergers provide an up to date median BBH merger rate density of $r_{\text{merg}} = 23 \text{ Gpc}^{-3} \text{ yr}^{-1}$ [7]. With this rate density and a uniform source distribution in the comoving volume, we populate ~ 8684 sources up to a redshift of 1.5. § We perform 8000 batches of simulations, with each batch containing 8684 sources with the mass and spin distributions following the ones detailed in Section 1.2. We further distribute the sources uniformly over the sky with the binary orientation distributed uniformly. For all the sources, we apply the detection criteria (i) and (ii) and obtain the detection rates (R_{det}). The right panel of Figure 2 provides the distribution of the estimated (R_{det}).

Table 1 provides the detection rate estimates of BBH. Compared to HL, the detection rate in AHL increases by 41% and 50% for criteria (i) and (ii), respectively. Besides the duty factor, the sky coverage of the two networks determines their detection rates. Since the coincidence criterion-(ii) exhibits a preference for shortlisting highly significant events, one can expect that with LIGO-Aundha one will see a perceptible increase in such events under that criterion.

3. Parameter estimation

With the expansion of the LIGO global network and the consequent enhancement in the signal-to-noise ratios of the CBC detections and mitigation of parameter degeneracies, one would anticipate improvements in the astrophysical parameter estimation. In this section, we employ CBC signal simulations to obtain quantitative support for this expectation.

For a BBH system in a circular orbit, the gravitational-wave signal is characterized by component masses (m_1, m_2), component spins ($\vec{S}_1, \vec{S}_2$), the luminosity distance (D_L), orbital inclination angle (ι), polarisation angle (ψ), sky-position angles (α, δ) and the coalescence time and phase (t_c, ϕ_c). For a binary neutron star (BNS) system, we

§ It is somewhat an arbitrary choice that we truncate the population at a maximum redshift of 1.5. However, this is motivated by the fact that at higher redshifts, the actual comoving rate density could significantly be different from the merger rate density at $z = 0$ (the one we assumed in this study) due to the star formation rate as well as the distribution of delay time between the formation and the coalescence of the binary.

require at least two additional parameters in the form of component tidal deformability parameters (Λ_1, Λ_2).

The CBC signal’s multi-dimensional parameter space harbors correlations and degeneracies among different parameter pairs, contributing to the uncertainties in the measurements of the individual parameters. For several of these parameters, the error-bar scales inversely with the signal-to-noise ratio (for loud signals) [15]. While this holds particularly well for the intrinsic binary parameters, such as component masses and spins, the aforementioned degeneracies among some pairs, e.g., (i) the sky-location angles α and δ and (ii) d_L and ι , can not often be removed despite high SNR. The expansion of LGN with LIGO-Aundha, in addition to increasing the SNR, will enhance parameter estimation accuracy by providing an independent observation of the source that can significantly reduce the degeneracies among some of the parameters.

In Sec. 3.1 we focus on general parameter estimation for select BBH events, and in Sec. 3.2 we present the primary results of masses and tidal effects in BNS systems.

3.1. Improvement in errors for binary black hole events

For this study, we simulated binary black hole signals modeled after two of the observed binary black holes, namely, (i) the loudest BBH, GW150914 [16] and (ii) the most massive BBH, GW190521 [17]. In fact, GW150914 is the first binary black hole merger observed by two LIGO detectors and the loudest event so far, with a coherent SNR of 24. The observed component masses were $36 M_\odot$ and $29 M_\odot$ with a remnant BH of $62 M_\odot$, and the event was located at a luminosity distance of 450 Mpc. It was localized in a huge sky-patch, spanning 590 sq. degs.

GW190521 is the most massive and among the farthest (5 Gpc) binary black hole mergers observed so far. Its component masses are $85 M_\odot$ and $66 M_\odot$. The remnant was estimated to have a mass of $142 M_\odot$. This is the first intermediate-mass BH candidate observed in the gravitational-wave window.

For our two simulations, the injected values of the key parameters are listed in Table 2 where we choose the masses and spins to be identical to those inferred for GW150914 and GW190521. From our 8000 batches of BBH simulations described in Sec. 2, it was found that the population-averaged ratio of SNR at AHL to the SNR at HL lies in the range of 1.3–1.4. We choose the injected sky positions in such a way that the SNR at AHL is ~ 1.4 times the SNR at the HL so that it resembles the average behaviour of SNR improvement.

The run-of-the-mill Bayesian parameter estimation approach assumes stationary Gaussian detector noise and a reliable, faithful Einstein’s GR signal model for the GW signal from the compact binary merger. An up to date suite of models for complete CBC waveforms constructed by combining various approaches include phenomenological models, such as IMRPhenom models [18], the effective one-body EOBNR waveforms that use inputs from numerical relativity [19, 20, 21], and the NRSurrogates waveforms derived from numerical relativity simulations [22, 23, 24]. In our analysis, we use

the *IMRPhenomPv2* [25, 26, 27, 28, 29] waveform model for both injections as well as recovery. We use the Bilby [30] software package, with its in-built sampler dynesty, to perform the parameter estimation. We perform this analysis with *zero-noise* injections and the likelihood computed using the A+ PSD.

We tabulate the results in terms of improvement in the 90% credible intervals on various astrophysical parameters in Table-2 and present pictorially in Fig. 3 the posterior probability contours (at 90%, and 68% credible levels). In that figure, the left and right panels depict the results for the GW150914- and GW190521-like injections, respectively.

Parameter	GW150914-like		GW190521-like	
	Injected	Improvement	Injected	Improvement
Chirpmass ($M_\odot$)	28.1	33%	64.6	39%
Total mass ($M_\odot$)	65.0	33%	149.6	40%
D_L in Gpc	2.5	35%	5.3	36%
θ_j in deg.	45	27%	45	70%
Sky localization in deg ² .		92%		96%

Table 2: Parameter estimation improvement in 90% credible intervals in expanding the network from HL to AHL: We use BBH signals modelled after GW150914 and GW190521 and estimate the improvement in sky-localization, luminosity distance, binary inclination, masses and spins.

3.1.1. Sky-localization: The detector pair comprising LIGO-Aundha and LIGO-Livingston provides the longest baseline amongst all pairs of existing/in-construction detectors. This amounts to an increase in the accuracy of source localization. For the GW150914-like injection, the 90% credible 2-D localization area is $\sim 114 \text{ deg}^2$ which improves to $\sim 9 \text{ deg}^2$ with AHL. This amounts to 92% reduction in the localization uncertainty. For GW190521-like injection, we find a $\sim 96\%$ reduction, with the respective localization area for HL and AHL configurations being $\sim 971 \text{ deg}^2$ and $\sim 35 \text{ deg}^2$. More discussion on the sky-localization can be found in Sec. 4 and the reader may also refer to earlier studies on localization, e.g., Ref. [31] and the references therein.

3.1.2. Luminosity distance and inclination angle: The three-detector configuration plays a crucial role in breaking the degeneracy between the luminosity distance D_L and the inclination angle ι . For the GW150914-like system, the errors in D_L and the inclination improve by 35% and 27%, respectively, for AHL relative to HL. Similarly, for the GW190521-like injection the error improvement in the same parameters is 36% and 70%, respectively. The improved distance measurements result from the reduced 3D localization volume of the source, primarily due to the reduced 2D sky-localization of the source by the AHL network. This will have direct implications in the measurements

of cosmological parameters [32, 33]. We also find significant improvement in the inclination angle measurement of the binary (θ_{jn}) which could be partly due to the breaking the distance-inclination degeneracy. Though this analysis has been performed on binary black hole mergers, similar improvements are expected in the inclination angles of binary neutron stars and neutron star-black hole mergers as well [34] which are favourite candidates to have associated EM counterparts. Accurate knowledge of the inclination angle is key in doing multimessenger astronomy, for making predictions on the possible EM counterparts and in understanding the physical process that drives the EM counterparts [35, 36, 37]. Further, improved precision in the binary inclination helps to probe the gravitational-wave polarization of the signal. This improvement directly impacts probing alternative theories of gravity with gravitational wave signals. In Sec. 5.2 we discuss how polarization measurements benefit from the expansion of LGN.

3.1.3. Source masses :

Improvement in the luminosity distance reflects in the improvement in the source masses. For GW150914-like injection, the errors on both the source-frame chirpmass and total mass improve by $\sim 33\%$. Similarly, for GW190521-like injection, these improvements are 39% and 40% respectively. See Fig. 3 for the $m_1 - m_2$ contour plots for both the events. Accurate knowledge of the intrinsic source parameters helps in the population synthesis studies of compact binary mergers and obtains constraints on the merger rate density [7].

3.2. Enhancement in matter effects: source classification and BNS properties

Neutron stars are characterized by their component masses (m_1, m_2) and the corresponding tidal deformability parameters (Λ_1, Λ_2) [38] of each component. The presence of matter is predominantly captured by the effective tidal deformability parameter ($\tilde{\Lambda}$) which is defined by a suitable combination of m_1, m_2, Λ_1 and Λ_2 [39]. Black holes in general relativity are predicted to have zero tidal deformability, i.e., $\Lambda_1 = \Lambda_2 = 0$. For a binary black hole (BBH) system, this leads to $\tilde{\Lambda} = 0$ irrespective of their component masses and spins. Moreover, precise estimation of the tidal deformability parameters can constrain the theoretically proposed equations of state of the neutron stars and thus shed light on the nature of internal composition of the star [40, 41, 42].

Here we illustrate how the addition of LIO-detector at Aundh-site can potentially impact our ability to constrain the effective tidal deformability parameter ($\tilde{\Lambda}$) as well as discriminate it from the $\tilde{\Lambda} = 0$ case corresponding to the BBH event in the fully Bayesian statistical framework [43].

We consider a set of candidate binary neutron star (BNS) events with source properties consistent with the first BNS event GW170817 [40, 43]. Although the chirp-mass (M_c) was very well determined to be $1.188 M_\odot$, the component masses have broader uncertainties due to lack of precision in the mass-ratio (e.g., symmetric mass-




Figure 3: Posterior distributions of certain parameters for GW150914-like (left) and GW190521-like (right) simulated signals in the HL and AHL networks: Top, middle and bottom panels correspond to the parameters $m_1^{src} - m_2^{src}$, $D_L - \iota$ and $RA - Dec$, respectively. The true values are shown by a black star. The 95% and 65% confidence intervals are shown by solid and dash-dotted lines, respectively.

ratio, η) estimation. Moreover, although GW170817 could successfully rule out the stiffest equations of state, it still has sufficiently broader uncertainty in estimated tidal deformability parameters which makes it consistent with a wide variety of neutron star EOSs.

Source parameters			Network	Measurement accuracies			Improvements in %		
m_1, m_2 ($M_\odot$)	Λ_1, Λ_2 [$\tilde{\Lambda}$]	D_L (Mpc)		ΔM_c ($M_\odot$)	$\Delta\eta$	$\Delta\tilde{\Lambda}$	ΔM_c (in %)	$\Delta\eta$ (in %)	$\Delta\tilde{\Lambda}$ (in %)
1.35,1.35	400,400 [400]	40	LH	9.7e-5	6.8e-3	131.9	23.7	14.7	24.5
			AHL	7.4e-5	5.8e-3	99.6			
1.35,1.35	857,857 [857]	40	LH	9.8e-5	6.8e-3	156.9	15.3	14.7	26.6
			AHL	8.3e-5	5.8e-3	115.2			
1.60,1.17	120,980 [551.5]	40	LH	1.3e-4	9.6e-3	147.9	15.4	13.5	25.5
			AHL	1.1e-4	8.3e-3	110.9			
1.35,1.35	400,400 [400]	100	LH	1.6e-4	7.6e-3	269.0	18.8	16.3	19.1
			AHL	1.3e-4	5.6e-3	217.7			
1.35,1.35	857,857 [857]	100	LH	1.7e-4	7.5e-3	418.7	17.6	10.6	47.8
			AHL	1.4e-4	6.7e-3	218.6			
1.60,1.17	120,980 [551.5]	100	LH	1.9e-4	1.0e-2	384.2	15.8	10.0	44.1
			AHL	1.6e-4	9.0e-3	215.0			
1.35,1.35	400,400 [400]	250	LH	4.7e-4	8.5e-3	1141.5	25.5	10.6	40.8
			AHL	3.5e-4	7.6e-3	675.4			
1.35,1.35	857,857 [857]	250	LH	3.5e-4	6.9e-3	1298.6	20.0	17.4	54.7
			AHL	2.8e-4	5.7e-3	588.5			
1.60,1.17	120,980 [551.5]	250	LH	3.2e-4	1.0e-2	2264.7	9.4	10.0	69.9
			AHL	2.9e-4	9.0e-3	638.51			

Table 3: This table summarizes comparisons of key properties of binary neutron star mergers events with high and relatively low-SNR events. Each of the BNS events is simulated with two neutron star EOSs, one softer (SLy4) and one stiffer (BHBA ϕ). The uncertainties in mass parameters, namely in chirp-mass (ΔM_c) and symmetric mass-ratio ($\Delta\eta$) as well as uncertainties in the effective tidal deformability parameter ($\Delta\tilde{\Lambda}$) of the BNS systems are quoted with 90% credible level (see subsection 3.2).

We perform a systematic injection-study of Bayesian parameter estimation for a set of simulated signals from BNS events covering the extreme corners of the parameter space component masses and tidal deformability parameters consistent with GW170817. Initially, we consider all the sources to be located at a luminosity distance (D_L) of 40 Mpc (similar to the GW170817 event). We consider one equal mass ($m_1 = m_2 = 1.35M_\odot$) BNS and an unequal mass ($m_1 = 1.60M_\odot, m_2 = 1.17M_\odot$) BNS with soft EOS, namely SLy4 [44], consistent with GW170817 observation. For the equal mass case, we also consider the possibility of the BNS events made up of a stiff EOS, namely BHBA ϕ [45].

Given the current estimation of BNS merger event rates (see 2), it is statistically unlikely to observe such an event at a distance $D_L \sim 40$ Mpc in the near future. We, thus, perform simulations for Bayesian parameter estimation considering the entire set of events to occur in addition, at a luminosity distance $D_L = 100$ Mpc and $D_L = 250$ Mpc, too. In our simulations, we use IMRPHENOMDNRTIDAL waveform model [46] for

the coalescing BNS systems with slow ($|s_1|, |s_2| \leq 0.05$), in-plane component spinning configurations for simplicity since astronomical distributions demonstrate that more rapidly spinning BNS systems are rare as well as this configuration captures the key aspects reasonably well. We summarize the measurement accuracies in mass and tidal deformability parameters with 90% Bayesian uncertainty intervals in table 3.2.

This study demonstrates that for the very high-SNR events (with comparable single detector SNR $\sim 110 - 130$ in each of H, L, and A detectors) the improvement of precision in M_c is in the range of 15 – 25% and in η is of about 10 – 17% for AHL-network of detectors as compared to the HL-network. The improvement in $\tilde{\Lambda}$ estimation in favor of the AHL-network relative to two US-based detectors is also nominal, about 25%. As the source distance increases resulting in a decrease in SNR, the improvement in precession for D_L and η does not change much for comparable SNRs in the three detectors. However, we find that for lower SNR events the precision in $\tilde{\Lambda}$ improves significantly. For the set of BNS observation at $D_L = 100$ Mpc, we find that improvements can be in the range of 20% to 45%. For the more distant sources, e.g., at $D_L = 250$ the improvements are generally more than 40%, and can be as high as 70% in favor of AHL relative to the HL-network. Moreover, for such distant sources, the lack of precision in $\tilde{\Lambda}$ can render it difficult to rule out the BBH-case corresponding to $\tilde{\Lambda} = 0$, particularly for the soft (SLy4-like) EOS. (As a comparison to the range of $\tilde{\Lambda}$ parameter for different theoretically motivated neutron star EOS models please refer to [47].) Thus, for the events with relatively weak signals – which will be originated at more distance away and thus relatively abundant in numbers – the source classification (i.e., BBH Vs BNS/NSBH) will get significantly enhanced. This will be important for generating alerts for the subsequent follow-up with astronomical observations across the electromagnetic spectrum.

4. Sky localisation and early warning

One of the main advantages of expanding the HL network to include LIGO-India is that it substantially improves the localization of CBCs in the sky [48, 49]. BNSs and a fraction of NSBHs have long been expected to produce prompt counterparts and afterglows in all electromagnetic (EM) bands. For BNS mergers in particular, it has been hypothesized that the post-merger central engine can launch short gamma-ray bursts (sGRBs) [50, 51], kilonovae [52, 53], and radio waves and X-rays before and after merger [54, 55, 56, 57].

These emissions carry information about both the progenitors – e.g., the equation of state of neutron stars – and the circum-merger environment. The prompt, and often, transient emission on the one hand and the late-time afterglow on the other hand complement each other in conveying that information, as was demonstrated amply by the multi-messenger observations of the binary neutron star event, GW170817 [58]. The joint observation of GWs followed by the sGRB, GRB 170817A, and the kilonova AT 2017gfo, [58] confirmed the several-decade-old hypothesis that compact object

mergers were progenitors of these exotic transients. However, GW170817 is so far the only gravitational-wave event to be observed in other channels. Improvements in GW detectors and expansion of the GW network is therefore required to realize more multi-messenger observations and expand our knowledge about the physical processes that occur in these systems.

The chances of telescopes spotting that EM emission improve if the localization area in the sky associated with the GW signal is small. This is particularly true for tracking down optical counterparts since the fields of view (FOV), or beam sizes, of these telescopes are small (sub-arcminutes) compared to the typical GW sky-localization area. The small localization with the rapid search strategies [59, 60, 61] can enhance the probability of finding the optical counterpart of the GW source. For prompt and transient emission, a narrow sky-area implies a small number of telescope slews and a quicker locking on to the target before it fades [62]. In the case of kilonovae and prolonged afterglows, narrow sky-areas can help since they allow observation of the EM-candidate over multiple epochs. The radio follow-up is the complimentary discovery for day-time and dust-obscured events, where the hunt in optical is difficult. In that case, a small volume in 3D localization is important for the galaxy targeted radio observation to get arcsecond localization [63].

Moreover, if the sky-localization is sharper, then spectroscopy becomes possible, which can provide not only clues on the progenitor composition but also the redshift of the event. The first discovery of the optical-counterpart of the event GW170817 was after ~ 11 hours of the GW trigger. Detection of the EM-counterparts must be much quicker if their prompt emissions are the desired target.

To demonstrate the benefit of including LIGO-Aundha in the GW network, we simulate a population of binary neutron stars and compare the distribution of GW localization in the two detector networks: HL and AHL. We generate a population of 9,308,544 simulated BNS signals using the TaylorF2 [64, 65, 66, 67] waveform model. Both source-frame component masses are drawn from a Gaussian distribution between $1.0 M_{\odot} < m_1, m_2 < 2.0 M_{\odot}$ with mean mass of $1.33 M_{\odot}$ and standard deviation of $0.09 M_{\odot}$, modeled after observations of galactic BNSs [68] (note, however [69]). The component spins are aligned or anti-aligned with respect to the orbital angular momentum with the dimensionless spin amplitude on the neutron stars restricted to 0.05, motivated by the low spins of BNSs expected to merge within a Hubble time [70, 71]. The signals are distributed uniformly in sky, orientation, and comoving volume up to a redshift of $z = 0.4$. We simulate the GW signal and calculate the expected SNR in Gaussian noise considering the three LIGO detectors at A+ sensitivity for each BNS. We mimic the results from a matched-filter GW search pipeline (current low-latency matched-filter searches running on LIGO-Virgo data include GstLAL [72, 73], PyCBCLive [74], MBTAOnline [75], SPIIR [76]) by considering the signals that pass a network SNR threshold of 12.0 to be ‘detected’. We then calculate the sky posteriors for the detected candidates using a rapid Bayesian localization tool, BAYESTAR [77]. We use the most recent BNS local merger rate from [69] of $320^{+410}_{-240} \text{Gpc}^{-3} \text{yr}^{-1}$ to estimate the

number of events detected per year in the detector network.

In Figure 4, we show the distributions (left: cumulative, right: density) of the sky localizations (90% credible interval) of the BNSs that pass the fiducial SNR threshold of 12 for the two detector networks: HL in green and AHL in cyan. The shaded regions show the uncertainty in the number of detections due to the uncertainty in the current local BNS merger rate of $320 \text{ Gpc}^{-3} \text{ yr}^{-1}$ [69]. The improvement due to addition of LIGO-India to the network is clearly visible in this figure; the AHL network detects about twice (17 – 175) as many signals as the HL (8 – 84) network. Further the peak of distribution for HL is around 800 deg^2 , about twice that for the AHL network.

Figure 5 shows the shape and the areal projection of a few localization on the sky. The HL localization are shown in green contours. They are long arc-shaped and larger in area. Most of HL localization are extended over both the hemisphere. When we include LIGO-India, the degeneracy breaks and the localization shrinks to either of the hemisphere. In this plot, a few red and blue contours have no corresponding green contours. These are the events detected by AHL but not by HL. The orientation of the HL localization are concentric. On the other hand AHL localization are randomly oriented.

This study compares the improvement in gravitational-wave localization for HL and AHL for similar sources at different parts of the sky. To do that, we divide the entire sky into equal-area pixels in HEALPix $\parallel$ format of NSIDE 16 [78]. We inject one BNS source in each pixel and calculate the localization area of the source using BAYESTAR for HL and AHL. All the injected sources are of $m_1, m_2 = 1.4 M_\odot$ at a distance of 100 Mpc. All the injected BNS sources have an orbital inclination of 5 deg. In Figure 6, the density of color code represents the area of the 90% credible region of the localization in deg^2 . The pixel value is the 90% probable area of the localization of the source injected in that pixel. In the map, the darker red pixels represent larger in localization area. For the HL network, the minimum localization area is $\sim 45 \text{ deg}^2$, and the largest area is $\sim 1732 \text{ deg}^2$. On the other hand, the smallest and biggest localization areas for the AHL are $\sim 1 \text{ deg}^2$ and $\sim 21 \text{ deg}^2$.

4.1. Early warning of binary neutron star mergers

August 17, 2017 saw the beginning of a new era in multi-messenger astronomy. The joint detection of GWs by the LIGO and Virgo interferometers and the sGRB by the Fermi-GBM and INTEGRAL satellite from the BNS coalescence, GW170817 [79, 58] confirmed the long-standing hypothesis that compact object mergers were progenitors of short GRBs. Apart from the gamma-ray burst, which was observed ~ 2 s after the merger event, the first manual follow-up observations took place ~ 8 hours after the epoch of merger [58]. This delay was caused by the delay in sending out GW information: the GW alert was sent out ~ 40 minutes [80], and the sky localization ~ 4.5

$\parallel$ A HEALPix map of NSIDE is a representation the full-sky filled by $N_{pix} = 12 \times NSIDE^2$ equal area pixels. The value of a pixel is the probability of finding the GW source in that pixel area in the sky.




Figure 4: Distributions (left: cumulative, right: density) of the sky localizations (90% credible interval) of the BNSs that pass the fiducial SNR threshold of 12 for the two detector networks: HL (green) and AHL (cyan). Using the latest median BNS merger rate from [69] of $320 \text{ Gpc}^{-3} \text{ yr}^{-1}$, we find that the HL (AHL) network is expected to detect ~ 33 (69) events per year. The shaded regions represent the uncertainty in the BNS merger rate estimate.




Figure 5: The contours on the sky are 90% probability contour of the localizations. The blue and purple contours are corresponding to the network AHL and HL, respectively. These are some selected sources from the simulation distributed over the sky to show the improvement by AHL. AHL also detects every signal which is detected by HL. The isolated blue contours correspond to the sources detected by AHL but not detected by HL.




Figure 6: Comparing the improvement in gravitational-wave localization for HL and AHL for similar sources at different sky locations. The sky map is divided into equal-area pixels in the HEALPix format. Each pixel has one injected BNS source in it and the value of the pixel is the 90% probable area of the localization of the source calculated by BAYESTAR. All of the BNS are injected at 100 Mpc and of $m_1 = m_2 = 1.4 M_\odot$ with a 5 deg. inclination angle.

hours [81] after the signal arrived on earth. By the time EM telescopes participating in the follow-up program received the alerts the source was below the horizon for them.

For a fraction of BNS events it will be possible to issue alerts up to $\delta t \sim 60$ s before the epoch of merger [82, 83, 84]. Pre-merger or *early warning* detections will facilitate electromagnetic observations of the prompt emission, which encodes the initial conditions of the outflow and the state of the merger remnant. Early optical and ultraviolet observations will be key to our understanding of *r*-process nucleosynthesis [85] and shock-heated ejecta [86], while prompt X-ray emission would reveal the final state of the remnant [87, 88, 89]. Early observations made in the radio band could indicate pre-merger magnetosphere interactions [90], and might be able to test models that predict BNS mergers as a possible precursor of fast radio bursts [91, 92, 93]. Early-warning GW alerts have recently been implemented [94, 95] and also demonstrated [96] recently.

Here we will compare the prospects of early-warning detection of GWs from BNSs for the two detector networks: HL and AHL. We follow the framework layed out in [94], which implemented an early warning GW pipeline using the GstLAL matched-filtering software suite [97]. In particular, we consider 6 different discrete frequency cut-offs 29 Hz, 32 Hz, 38 Hz, 49 Hz, 56 Hz, and 1024 Hz to analyze signal recovery at (approximately) 58 s, 44 s, 28 s, 14 s, 10 s, and 0 s before merger. We use the population of BNSs described earlier and the same criteria for ‘detected’ signal, the signals that

pass an SNR threshold of 12.0 in each frequency configuration are detected with the corresponding pre-merger latency. We then use BAYESTAR to localize all the detected signals for each frequency configuration.

Our results are shown in Figure 7. For both networks (left: HL, right: AHL), we show the cumulative distributions of the 90% credible intervals of the sky localizations for each pre-merger time considered in our simulation. The y-axis is translated to number of detections per year assuming the current median BNS merger rate estimate and 100% duty cycle of the networks. We note that at each frequency and pre-merger time configuration, the AHL network is able to detect about twice the number of events as compared to the HL network. In particular, the number of events per year that could be detected at least 10 s before merger are 7 (15) for the HL (AHL) network. Adding LIGO-India to the network will also greatly reduce the area in the sky to which these events can be localized, thereby vastly improving prospects of observing EM emissions before and/or at merger. The HL network is expected to detect one event every two years before merger that is also localized to 1000 deg^2 , while the AHL network is expected to detect ~ 5 events every year before merger that are also localized to 1000 deg^2 .




Figure 7: Cumulative distributions of the sky localizations (90% credible interval) of the BNSs that pass the fiducial SNR threshold of 12 for the two detector networks: HL (left) and AHL (right). The different colors show the different frequency bandwidths or pre-merger times considered in our work. Using the latest median BNS merger rate from [69] of $320 \text{ Gpc}^{-3} \text{ yr}^{-1}$, we find that the HL (AHL) network is expected to detect ~ 7 (15) mergers per year at least 10 s before merger. Out of these, the HL network is expected to detect one event every two years before merger that is also localized to 1000 deg^2 , while the AHL network is expected to detect ~ 5 events every year before merger that are also localized to 1000 deg^2 .

These localizations are quite large for optical telescopes, which generally have very small FOVs in comparison. Some of the largest field telescopes, such as the BlackGEM array ($0.65 \text{ m}/2.7 \text{ deg}^2$ per telescope) with 3 telescopes planned in the first phase of operation eventually expanding to 15 telescopes [98], the Zwicky Transient

Facility (1.2m/47deg²) [99], the Dark Energy Camera (4m/3.8deg²) [100], the Rubin Observatory (8.4m/9.6deg²) [101], the Swope Telescope (1m/7deg²) [102], the Subaru Telescope (8.2m/1.7deg²) [103], etc. all have FOVs only a small fraction of the GW localizations. Therefore, adding LIGO-India to the GW will be very crucial in order to observe any pre-merger and prompt emissions.

5. Tests of GR

5.1. Improved constraints on deviations from GR

Most of the tests of general relativity performed in [104, 105, 106] are null tests in the sense that they look for deviations around the expectation from GR. These deviation parameters can be measured from each detected event and, assuming certain properties, can be combined across multiple events to get tighter constraints. Measurement of deviations in the post-Newtonian parameters [107, 108, 109, 110], inspiral-merger-ringdown consistency test [111, 112, 113], and the upper bound on the mass of the graviton [114], etc., fall under this category.

We illustrate the potential of getting tighter bounds on the GR deviation parameters with the AHL network using the measurement of the mass of the graviton. In GR, GWs are non-dispersive. Hence the corresponding force carrier, graviton, should have zero rest mass. But there are alternative theories to GR that permit a non-trivial dispersion relation. We assume the following phenomenological form [115, 114, 116, 117, 118, 119, 120, 121],

$$E^2 = p^2 c^2 + A_\alpha p^\alpha c^\alpha, \quad (2)$$

where E and p are the energy and momentum of the GW, and A_α and α are phenomenological parameters. The phenomenological parameters are related to the graviton's mass by $m_g = \sqrt{A_0}/c^2$, with the condition $A_0 > 0$.

The 90% upper bound on the mass of the graviton measured from GW150914 is 9.9×10^{-23} eV/c² [105]. If the graviton is indeed massless, then this constraint should get tighter with detections louder than GW150914, as well as with more number of events. We assume that the upper bound scales inversely with the SNR of the detection. So, for a given event i with SNR ρ_i , the 90% upper bound on the graviton mass σ_i is given by $\sigma_i = \sigma_0 \rho_0 / \rho_i$. Here, σ_0 and ρ_0 are the 90% upper bound on graviton mass and the SNR obtained from GW150914. If N events are detected, we can combine the constraints to obtain the following combined constraint:

$$\sigma_{\text{comb}} = \frac{1}{N} \sqrt{\sum_{i=1}^N \sigma_i^2} = \frac{\sigma_0 \rho_0}{N} \sqrt{\sum_{i=1}^N \rho_i^{-2}}. \quad (3)$$

We simulate sources in the HL and AHL networks using the models specified in Sec. 1.2, and consider only those events detected by the multi-detector coincidence criterion (criterion (ii) in Sec. 2). We find that there are, on average, 502 events per year in the HL network and 754 events per year in AHL network that satisfy this criterion.

We apply the above prescription to obtain the constraints and find $\sigma_{\text{comb}}^{\text{HLA}}/\sigma_{\text{comb}}^{\text{HL}} \approx 0.8$, i.e. the constraints obtained with the AHL network are 20% tighter than the ones obtained with the HL network.

5.2. Constraints the nature of GW polarisations

In GR, GWs have only two independent polarisation states — i.e., two transverse quadrupole (or tensor) modes. In comparison, a general metric theory of gravity can admit up to six polarisation modes. In this sense, GW polarisations offer an interesting test of GR. GW polarisations can be constrained from observations of spinning neutron stars [122, 123] and stochastic background [124, 125, 126], as well as from observations of compact binary mergers [127, 128, 129, 130, 131, 132, 133]. While the detectability of spinning neutron stars or stochastic background is uncertain, we are expecting to detect hundreds to thousands of compact binary mergers in the next few years using ground-based GW detectors. Note that each GW detector observes only *one* linear combination of these polarisations. Due to the near co-alignment of the LIGO-Hanford and -Livingston detectors, they measure essentially the *same* linear combination of polarisations in a binary merger signal. Hence, currently the LIGO detectors alone are practically incapable of resolving even the two polarisation states predicted by GR. Additional detectors around the globe, including LIGO-India, will enable observing those two states and potentially constrain the additional non-GR polarisation states.

Given the data from a network of GW detectors, we can compare the posterior probabilities of different hypotheses, for example, one hypothesis stating that the binary phase evolution is exactly as predicted by GR, while the alternative hypothesis accommodating the presence of additional modes [127]. Motivated by the limited number of linearly independent detectors to observe the polarisation modes, the current probes of the nature of GW polarisations have employed highly simplified hypotheses as alternatives to GR. That is, the alternative hypothesis assumes that the polarisations contain only scalar modes (h_b and h_l) or only vector modes (h_x and h_y) or only tensor modes (h_+ and $h_\times$) [130, 134, 129, 132].

We perform a simulation study that compares the ability of the 3-detector network involving LIGO India to distinguish different polarisation models, as compared to the 2-detector network involving only LIGO Hanford and Livingston. For each polarisation hypothesis — tensor H_t , vector H_v , and scalar H_s , the model waveforms are generated using the corresponding antenna patterns, but always assuming that the time evolution of the polarisations follow that of the GR modes. That is, $h_b(t) = h_x(t) = h_+(t)$ and $h_l(t) = h_y(t) = h_\times(t)$. Since each detector I has different antenna pattern function F_I^A for each GW polarisation A , the strain measured at detector is a different linear combination of the polarisations modes: $h_I(t) = F_I^A(\alpha, \delta, \psi, t)h_A(t)$, where α, δ denotes the sky location of the source and ψ the polarisation angle. For simplicity, no noise is added to the simulated signals. Further, we considered GW signals from non-spinning binary black holes as our signal model.

We use the standard Gaussian likelihood model for estimating the posteriors of the parameters under different polarisation hypotheses (see, e.g., [135]), using the BILBY software package [30]. Posteriors are computed over the following parameters $m_1, m_2, \alpha, \delta, d_L, \iota, \psi, \phi_0, t_0$. We use uniform priors in redshifted component masses of the binary ($m_1, m_2 \in [3, 500]M_\odot$), isotropic priors in sky location (uniform in $\alpha, \sin\delta$) and orientation (uniform in $\cos\iota, \phi_0$), uniform in polarisation angle ψ , and a volumetric prior $\propto d_L^2$ on luminosity distance. The Bayesian evidence of each polarisation model is obtained as part of the parameter estimation.

We simulate ~ 200 GR (tensor) double coincident injections for each one of HL and AHL networks, and do parameter estimation and compute evidences $P(d|H_p)$ for each of the polarisation hypothesis $H_p \in \{H_t, H_v, H_s\}$. From those evidences, the likelihood ratio (Bayes factor) for tensor v/s vector (B_v^t) and tensor v/s scalar (B_s^t) hypothesis are calculated for various combination of detectors. The distribution of the Bayes factors $B_s^t := P(d|H_t)/P(d|H_s)$ and $B_v^t = P(d|H_t)/P(d|H_v)$ are plotted in Fig. 8. We can see that the 3-detector AHL network has a much better ability (larger Bayes factors) to distinguish the polarisation models as compared to the 2-detector HL network.

6. Conclusions and summary

We have analyzed the performance of the LIGO Global Network, in particular focusing on the improvement that comes about from the addition of a third detector in India. We focused on compact binary coalescences, involving black holes and neutron stars, as our sources. We find, overall, a significant improvement in the precision with which various binary parameters can be measured. This is especially significant for the sky localisation of compact binaries as well as the related ability to issue early warning for BNS mergers. Precise localisation is crucial in spotting and associating kilonovae with such mergers. Correct associations are necessary for understanding the influence of progenitor properties on kilonova properties, such as their spectra. They also have a bearing on constraining neutron star equation of state and the measurement of the Hubble parameter without invoking the cosmic-distance ladder.

Reducing the alert time for these mergers by several to a few tens of seconds can impact the ability of astronomers to slew their telescopes in time to capture prompt afterglow emissions, not to mention pre-merger EM signals. Prompt afterglows were missed in the observation of GW170817 and can provide important clues about the short GRB engine. This is one of the next frontiers in GRB physics that the GW network can contribute to.

The increased detection rate of CBCs with the addition of the third LIGO detector in India will also allow stronger constraints to be placed on possible deviations from GR. By assuming that constraints from a given signal would scale inversely with the SNR, and combining constraints across events, we find that the AHL network will offer a 20% improvement over the HL network on the graviton mass upper limit. We also showed how an additional detector in the network aids in discriminating among




Figure 8: The distribution of the Bayes factors that show our ability to distinguish different polarisation models. The simulated signals always follow tensor polarisations (as predicted by GR) and Bayesian evidence of three different polarisation models are computed. We can see that the 3-detector AHL network has a much better ability (larger Bayes factors) to distinguish the polarisation models as compared to the 2-detector HL network.

different polarisation models: Here we limited ourselves to models where GWs have only tensor, only vector or only scalar polarisation modes – taking all modes to have the same phase evolution; some of these assumptions can be relaxed in the future by using the null stream reconstruction [136, 131, 133].

While our primary focus here has been CBCs, the LIGO Global Network will also impact science pursuable with other signals. One such signal is an astrophysical stochastic gravitational-wave background (SGWB) arising from the superposition of inspiral signals from populations of binaries of black holes and neutron stars [137, 138]. By combining the detections of dozens of individual binaries, on the one hand, with upper-limits on the power spectra of SGWB, on the other hand, past studies [139, 140] have constrained the rate of evolution of CBCs over redshift. The spread in this rate will shrink by 20% with the third detector added to this network. The addition of a third detector will also help to understand the correlated terrestrial noise sources, which in turn will play a crucial role in confidently claiming any SGWB detection. Moreover, if the astrophysical SGWB has significant anisotropies, probing

them requires better sky coverage. LIGO-India's inclusion in the existing detector network will help to resolve these finer angular structures using the existing mapping techniques [141, 142, 143, 144].

In the future, it will be interesting to study other types of sources (and not just compact binaries), the effects of realistic interferometer noise, and the presence of other detectors in the network.

Acknowledgements

We would like to thank our colleagues in the LIGO-India Scientific Collaboration and the LIGO-India Project for valuable inputs. We appreciate the several discussions we had with members of the various working groups in the LIGO-Virgo-KAGRA collaborations. In particular, we thank K. G. Arun, Bala Iyer, Shivaraj Kandhasamy, Jose Matthew, Fred Raab, Rory Smith, and Tarun Souradeep for valuable discussions and inputs. This work makes use of NUMPY [145], SCIPY [146], MATPLOTLIB [147], ASTROPY [148, 149], JUPYTER [150], DYNESTY [151], BILBY [30] and PESUMMARY [152] software packages. Thanks are also due to the Department of Science and Technology (DST) and the Department of Atomic Energy (DAE) of India. Specifically, MS acknowledges the support from the Infosys Foundation, the Swarnajayanti fellowship grant DST/SJF/PSA-01/2017-18, and the support from the National Science Foundation with grants PHY-1806630 and PHY-2010970, AP acknowledges support from the DST-SERB Matrics grant MTR/2019/001096, AM acknowledges support from the DST-SERB Start-up Research Grant SRG/2020/001290, and PA, AV, and SG acknowledge support from DAE under project no. RTI4001. PA's research was also supported by the Max Planck Society through a Max Planck Partner Group at ICTS-TIFR and by the Canadian Institute for Advanced Research through the CIFAR Azrieli Global Scholars program. SS is supported by an Eberly postdoctoral fellowship at Pennsylvania State University and BSS is supported by NSF grant PHYS-1836779. Thanks are due to computational support provided by the Alice (ICTS-TIFR) and Sarathi (IUCAA) clusters and computing resources in SINP. In addition, the authors are also grateful for the computational resources provided by LIGO Laboratory and Leonard E Parker Center for Gravitation, Cosmology and Astrophysics at the University of Wisconsin-Milwaukee and supported by National Science Foundation Grants PHY-0757058, PHY-0823459, PHY-1626190 and PHY-1700765. This paper has been assigned the internal LIGO preprint number P2100073.

- [1] Aasi J *et al.* (LIGO Scientific Collaboration) 2015 *Class. Quantum Grav.* **32** 074001 (*Preprint* [1411.4547](#))
- [2] Acernese F *et al.* (Virgo Collaboration) 2015 *Class. Quantum Grav.* **32** 024001 (*Preprint* [1408.3978](#))
- [3] Abbott R *et al.* (LIGO Scientific, Virgo) 2020 (*Preprint* [2010.14527](#))
- [4] KAGRA Collaboration, Akutsu T, Ando M, Arai K, Arai Y, Araki S, Araya A, Aritomi N, Asada H, Aso Y, Bae S, Bae Y, Baiotti L, Bajpai R, Barton M A, Cannon K, Cao Z, Capocasa E, Chan M, Chen C, Chen K, Chen Y, Chiang C Y, Chu H, Chu Y K, Eguchi S, Enomoto Y, Flaminio R, Fujii Y, Fujikawa F, Fukunaga M, Fukushima M, Gao D, Ge G, Ha S, Hagiwara A, Haino S, Han W B, Hasegawa K, Hattori K, Hayakawa H, Hayama K, Himemoto Y, Hiranuma Y, Hirata N, Hirose E, Hong Z, Hsieh B H, Huang C Z, Huang H Y, Huang P, Huang Y C, Huang Y, Hui D C Y, Ide S, Ikenoue B, Imam S, Inayoshi K, Inoue Y, Ioka K, Ito K, Itoh Y, Izumi K, Jeon C, Jin H B, Jung K, Jung P, Kaihotsu K, Kajita T, Kakizaki M, Kamiizumi M, Kanda N, Kang G, Kashiyama K, Kawaguchi K, Kawai N, Kawasaki T, Kim C, Kim J, Kim J C, Kim W S, Kim Y M, Kimura N, Kita N, Kitazawa H, Kojima Y, Kokeyama K, Komori K, Kong A K H, Kotake K, Kozakai C, Kozu R, Kumar R, Kume J, Kuo C, Kuo H S, Kuromiya Y, Kuroyanagi S, Kusayanagi K, Kwak K, Lee H K, Lee H W, Lee R, Leonardi M, Li T G F, Li K L, Lin L C C, Lin C Y, Lin F K, Lin F L, Lin H L, Liu G C, Luo L W, Majorana E, Marchio M, Michimura Y, Mio N, Miyakawa O, Miyamoto A, Miyazaki Y, Miyo K, Miyoki S, Mori Y, Morisaki S, Moriwaki Y, Nagano K, Nagano S, Nakamura K, Nakano H, Nakano M, Nakashima R, Nakayama Y, Narikawa T, Naticchioni L, Negishi R, Nguyen Quynh L, Ni W T, Nishizawa A, Nozaki S, Obuchi Y, Ogaki W, Oh J J, Oh K, Oh S H, Ohashi M, Ohishi N, Ohkawa M, Ohta H, Okutani Y, Okutomi K, Oohara K, Ooi C P, Oshino S, Otabe S, Pan K, Pang H, Parisi A, Park J, Pena Arellano F E, Pinto I, Sago N, Saito S, Saito Y, Sakai K, Sakai Y, Sakuno Y, Sato S, Sato T, Sawada T, Sekiguchi T, Sekiguchi Y, Shao L, Shibagaki S, Shimizu R, Shimoda T, Shimode K, Shinkai H, Shishido T, Shoda A, Somiya K, Son E J, Sotani H, Sugimoto R, Suresh J, Suzuki T, Suzuki T, Tagoshi H, Takahashi H, Takahashi R, Takamori A, Takano S, Takeda H, Takeda M, Tanaka H, Tanaka K, Tanaka K, Tanaka T, Tanaka T, Tanioka S, Tapia San Martin E N, Telada S, Tomaru T, Tomigami Y, Tomura T, Travasso F, Trozzo L, Tsang T, Tsao J S, Tsubono K, Tsuchida S, Tsuna D, Tsutsui T, Tsuzuki T, Tuyenbayev D, Uchikata N, Uchiyama T, Ueda A, Uehara T, Ueno K, Ueshima G, Uraguchi F, Ushiba T, van Putten M H P M, Vocca H, Wang J, Washimi T, Wu C, Wu H, Wu S, Xu W R, Yamada T, Yamamoto K, Yamamoto K, Yamamoto T, Yamashita K, Yamazaki R, Yang Y, Yokogawa K, Yokoyama J, Yokozawa T, Yoshioka T, Yuzurihara H, Zeidler S, Zhan M, Zhang H, Zhao Y and Zhu Z H 2020 *arXiv e-prints* arXiv:2008.02921 (*Preprint* [2008.02921](#))
- [5] Iyer B *et al.* 2011 LIGO-India, proposal of the consortium for Indian initiative in gravitational-wave observations (IndIGO) URL <https://dcc.ligo.org/LIGO-M1100296/public>
- [6] Miller J, Barsotti L, Vitale S, Fritschel P, Evans M and Sigg D 2015 *Phys. Rev. D* **91**(6) 062005 URL <https://link.aps.org/doi/10.1103/PhysRevD.91.062005>
- [7] Abbott R *et al.* (LIGO Scientific, Virgo) 2020 (*Preprint* [2010.14533](#))
- [8] Kalogera V 2000 *Astrophys. J.* **541** 319–328 (*Preprint* [astro-ph/9911417](#))
- [9] Rodriguez C L, Zevin M, Pankow C, Kalogera V and Rasio F A 2016 *Astrophys. J. Lett.* **832** L2 (*Preprint* [1609.05916](#))
- [10] Ade P A R *et al.* (Planck) 2016 *Astron. Astrophys.* **594** A13 (*Preprint* [1502.01589](#))
- [11] Abbott B P *et al.* (LIGO Scientific, Virgo) 2019 *Phys. Rev. X* **9** 031040 (*Preprint* [1811.12907](#))
- [12] Abbott B *et al.* (LIGO Scientific, Virgo) 2019 *Astrophys. J. Lett.* **882** L24 (*Preprint* [1811.12940](#))
- [13] Pai A, Dhurandhar S and Bose S 2001 *Phys. Rev.* **D64**(4) 042004 (*Preprint* [0009078](#))
- [14] Abbott B *et al.* (KAGRA, LIGO Scientific, VIRGO) 2018 *Living Rev. Rel.* **21** 3 (*Preprint* [1304.0670](#))
- [15] Cutler C and Flanagan E E 1994 *Phys. Rev. D* **49** 2658–2697 (*Preprint* [gr-qc/9402014](#))
- [16] Abbott B P *et al.* (LIGO Scientific Collaboration and Virgo Collaboration) 2016 *Phys. Rev. Lett.* **116** 061102 (*Preprint* [1602.03837](#))
- [17] Abbott R *et al.* (LIGO Scientific, Virgo) 2020 *Phys. Rev. Lett.* **125** 101102 (*Preprint* [2009.01075](#))
- [18] Ajith P, Hannam M, Husa S, Chen Y, Brügmann B, Dorband N, Müller D, Ohme F, Pollney D,

- Reisswig C *et al.* 2011 *Physical review letters* **106** 241101
- [19] Buonanno A and Damour T 1999 *Phys. Rev. D* **59** 084006 (*Preprint* [gr-qc/9811091](#))
- [20] Buonanno A and Damour T 2000 *Phys. Rev. D* **62** 064015 (*Preprint* [gr-qc/0001013](#))
- [21] Barausse E and Buonanno A 2010 *Phys. Rev. D* **81** 084024 (*Preprint* [0912.3517](#))
- [22] Field S E, Galley C R, Hesthaven J S, Kaye J and Tiglio M 2014 *Phys. Rev. X* **4** 031006 (*Preprint* [1308.3565](#))
- [23] Blackman J, Field S E, Galley C R, Szilágyi B, Scheel M A, Tiglio M and Hemberger D A 2015 *Phys. Rev. Lett.* **115** 121102 (*Preprint* [1502.07758](#))
- [24] Varma V, Field S E, Scheel M A, Blackman J, Kidder L E and Pfeiffer H P 2019 *Phys. Rev. D* **99** 064045 (*Preprint* [1812.07865](#))
- [25] Hannam M, Schmidt P, Bohé A, Haegel L, Husa S *et al.* 2014 *Phys.Rev.Lett.* **113** 151101 (*Preprint* [1308.3271](#))
- [26] Schmidt P, Ohme F and Hannam M 2015 *Phys. Rev.* **D91** 024043 (*Preprint* [1408.1810](#))
- [27] Husa S, Khan S, Hannam M, Pürrer M, Ohme F, Jiménez Forteza X and Bohé A 2016 *Phys. Rev.* **D93** 044006 (*Preprint* [1508.07250](#))
- [28] Khan S, Husa S, Hannam M, Ohme F, Pürrer M, Jiménez Forteza X and Bohé A 2016 *Phys. Rev.* **D93** 044007 (*Preprint* [1508.07253](#))
- [29] Bohé A, Hannam M, Husa S, Ohme F, Puerrer M and Schmidt P 2016 Phenompv2 - technical notes for lal implementation Tech. Rep. LIGO-T1500602 LIGO Project URL <https://dcc.ligo.org/LIGO-T1500602>
- [30] Ashton G, Hübner M, Lasky P D, Talbot C, Ackley K, Biscoveanu S, Chu Q, Divakarla A, Easter P J, Goncharov B, Hernandez Vivanco F, Harms J, Lower M E, Meadors G D, Melchor D, Payne E, Pitkin M D, Powell J, Sarin N, Smith R J E and Thrane E 2019 *Apjs* **241** 27 (*Preprint* [1811.02042](#)) URL <https://ui.adsabs.harvard.edu/abs/2019ApJS..241...27A>
- [31] Pankow C, Rizzo M, Rao K, Berry C P L and Kalogera V 2020 *Astrophys. J.* **902** 71 (*Preprint* [1909.12961](#))
- [32] Abbott B P *et al.* (LIGO Scientific, Virgo) 2021 *Astrophys. J.* **909** 218 (*Preprint* [1908.06060](#))
- [33] Abbott B P *et al.* (LIGO Scientific, Virgo, 1M2H, Dark Energy Camera GW-E, DES, DLT40, Las Cumbres Observatory, VINROUGE, MASTER) 2017 *Nature* **551** 85–88 (*Preprint* [1710.05835](#))
- [34] Rodriguez C L, Farr B, Raymond V, Farr W M, Littenberg T B, Fazi D and Kalogera V 2014 *Astrophys. J.* **784** 119 (*Preprint* [1309.3273](#))
- [35] Arun K G, Tagoshi H, Mishra C K and Pai A 2014 *Phys. Rev. D* **90** 024060 (*Preprint* [1403.6917](#))
- [36] Saleem M 2020 *Mon. Not. Roy. Astron. Soc.* **493** 1633–1639 (*Preprint* [1905.00314](#))
- [37] Saleem M, Resmi L, Arun K and Mohan S 2020 *The Astrophysical Journal* **891** 130
- [38] Hinderer T 2008 *The Astrophysical Journal* **677** 1216–1220 (*Preprint* [0711.2420](#))
- [39] Wade L, Creighton J D E, Ochsner E, Lackey B D, Farr B F, Littenberg T B and Raymond V 2014 *Physical Review D* **89** 103012 (*Preprint* [1402.5156](#))
- [40] Abbott B P, Abbott R, Abbott T, Abernathy M, Acernese F, Ackley K, Adams C, Adams T, Addesso P, Adhikari R *et al.* 2017 *Physical review letters* **119** 161101 (*Preprint* [1710.05832](#))
- [41] Flanagan É É and Hinderer T 2008 *Physical Review D* **77** 021502 (*Preprint* [0709.1915](#))
- [42] Read J S, Lackey B D, Owen B J and Friedman J L 2009 *Physical Review D* **79** 124032 (*Preprint* [0812.2163](#))
- [43] Abbott B P, Abbott R, Abbott T, Abernathy M, Acernese F, Ackley K, Adams C, Adams T, Addesso P, Adhikari R *et al.* 2019 *Physical Review X* **9** 011001 (*Preprint* [1805.11579](#))
- [44] Douchin F and Haensel P 2001 *Astronomy & Astrophysics* **380** 151–167 (*Preprint* [astro-ph/0111092](#))
- [45] Banik S, Hempel M and Bandyopadhyay D 2014 *The Astrophysical Journal Supplement* **214** 22 (*Preprint* [1404.6173](#))
- [46] Dietrich T, Khan S, Dudi R, Kapadia S J, Kumar P, Nagar A, Ohme F, Pannarale F, Samajdar A, Bernuzzi S, Carullo G, Del Pozzo W, Haney M, Markakis C, Pürrer M, Riemenschneider G, Setyawati Y E, Tsang K W and Van Den Broeck C 2019 *Phys. Rev. D* **99** 024029 (*Preprint* [1805.11579](#))

- 1804.02235)
- [47] Abbott B P, Abbott R, Abbott T D, Abraham S, Acernese F, Ackley K, Adams C, Adya V B, Affeldt *et al.* 2020 *Classical and Quantum Gravity* **37** 045006 (Preprint 1908.01012)
 - [48] Fairhurst S 2014 *Journal of Physics: Conference Series* **484** 012007 (Preprint 1205.6611)
 - [49] Pankow C, Rizzo M, Rao K, Berry C P L and Kalogera V 2020 *The Astrophysical Journal* **902** 71 (Preprint 1909.12961)
 - [50] Lattimer J and Schramm D 1976 *Astrophys. J.* **210** 549
 - [51] Lee W H and Ramirez-Ruiz E 2007 *New J. Phys.* **9** 17 (Preprint astro-ph/0701874)
 - [52] Li L X and Paczynski B 1998 *Astrophys. J. Lett.* **507** L59 (Preprint astro-ph/9807272)
 - [53] Metzger B, Martinez-Pinedo G, Darbha S, Quataert E, Arcones A, Kasen D, Thomas R, Nugent P, Panov I and Zinner N 2010 *Mon. Not. Roy. Astron. Soc.* **406** 2650 (Preprint 1001.5029)
 - [54] Nakar E and Piran T 2011 *Nature* **478** 82–84 (Preprint 1102.1020)
 - [55] Metzger B D and Berger E 2012 *The Astrophysical Journal* **746** 48
 - [56] Metzger B D and Zivancev C 2016 *Mon. Not. Roy. Astron. Soc.* **461** 4435–4440 (Preprint 1605.01060)
 - [57] Hallinan G *et al.* 2017 *Science* **358** 1579–1583 (Preprint 1710.05435)
 - [58] Abbott B P *et al.* (LIGO Scientific, Virgo, Fermi GBM, INTEGRAL, IceCube, AstroSat Cadmium Zinc Telluride Imager Team, IPN, Insight-Hxmt, ANTARES, Swift, AGILE Team, 1M2H Team, Dark Energy Camera GW-EM, DES, DLT40, GRAWITA, Fermi-LAT, ATCA, ASKAP, Las Cumbres Observatory Group, OzGrav, DWF (Deeper Wider Faster Program), AST3, CAASTRO, VINROUGE, MASTER, J-GEM, GROWTH, JAGWAR, CaltechNRAO, TTU-NRAO, NuSTAR, Pan-STARRS, MAXI Team, TZAC Consortium, KU, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS, BOOTES, MWA, CALET, IKI-GW Follow-up, H.E.S.S., LOFAR, LWA, HAWC, Pierre Auger, ALMA, Euro VLBI Team, Pi of Sky, Chandra Team at McGill University, DFN, ATLAS Telescopes, High Time Resolution Universe Survey, RIMAS, RATIR, SKA South Africa/MeerKAT) 2017 *Astrophys. J.* **848** L12 (Preprint 1710.05833)
 - [59] Rana J, Singhal A, Gadre B, Bhalerao V and Bose S 2017 *The Astrophysical Journal* **838** 108 (Preprint 1603.01689)
 - [60] Coughlin M W *et al.* 2018 *MNRS* **478** 692–702 (Preprint 1803.02255)
 - [61] Ghosh S, Bloemen S, Nelemans G, Groot P J and Price L R 2016 *Astronomy and Astrophysics* **592** A82 (Preprint 1511.02673)
 - [62] Rana J, Anand S and Bose S 2019 *The Astrophysical Journal* **876** 104 (Preprint 1902.08378)
 - [63] Rana J and Mooley K P 2019 *arXiv e-prints* arXiv:1904.07335 (Preprint 1904.07335)
 - [64] Sathyaprakash B S and Dhurandhar S V 1991 *Phys. Rev.* **D44**(12) 3819–3834
 - [65] Blanchet L, Damour T, Iyer B R, Will C M and Wiseman A G 1995 *Phys. Rev. Lett.* **74** 3515–3518 (Preprint gr-qc/9501027)
 - [66] Blanchet L, Damour T, Esposito-Farese G and Iyer B R 2005 *Phys. Rev.* **D71** 124004 (Preprint gr-qc/0503044)
 - [67] Buonanno A, Iyer B R, Ochsner E, Pan Y and Sathyaprakash B S 2009 *Phys. Rev.* **D80**(8) 084043 (Preprint 0907.0700)
 - [68] Özel F and Freire P 2016 *Ann. Rev. Astron. Astrophys.* **54** 401–440 (Preprint 1603.02698)
 - [69] Abbott B P, Abbott R, Abbott T D, Abraham S, Acernese F, Ackley K, Adams C, Adhikari R X, Adya V B, Affeldt C and *et al* 2020 *The Astrophysical Journal* **892** L3 ISSN 2041-8213 URL <http://dx.doi.org/10.3847/2041-8213/ab75f5>
 - [70] Burgay M *et al.* 2003 *Nature* **426** 531–533 (Preprint astro-ph/0312071)
 - [71] Zhu X, Thrane E, Osłowski S, Levin Y and Lasky P D 2018 *Phys. Rev.* **D98** 043002 (Preprint 1711.09226)
 - [72] Sachdev S *et al.* 2019 (Preprint 1901.08580)
 - [73] Hanna C *et al.* 2020 *Phys. Rev.* **D101** 022003 (Preprint 1901.02227)
 - [74] Nitz A H, Dal Canton T, Davis D and Reyes S 2018 *Phys. Rev.* **D98** 024050 (Preprint 1805.11174)

- [75] Adams T, Buskulic D, Germain V, Guidi G M, Marion F, Montani M, Mours B, Piergiovanni F and Wang G 2016 *Class. Quant. Grav.* **33** 175012 (*Preprint* [1512.02864](#))
- [76] Chu Q 2017 *Low-latency detection and localization of gravitational waves from compact binary coalescences* Ph.D. thesis The University of Western Australia
- [77] Singer L P and Price L R 2016 *Phys. Rev. D* **93**(2) 024013 URL <https://link.aps.org/doi/10.1103/PhysRevD.93.024013>
- [78] Gorski K M *et al.* 2005 *The Astrophysical Journal* **622** 759–771 (*Preprint* [astro-ph/0409513](#))
- [79] Abbott B P, Abbott R, Abbott T, Acernese F, Ackley K, Adams C, Adams T, Addesso P, Adhikari R, Adya V *et al.* 2017 *Physical Review Letters* **119** 161101
- [80] LIGO Scientific Collaboration V C 2017 *GCN* **21505** URL <https://gcn.gsfc.nasa.gov/gcn3/21505.gcn3>
- [81] LIGO Scientific Collaboration V C 2017 *GCN* **21513** URL <https://gcn.gsfc.nasa.gov/gcn3/21513.gcn3>
- [82] Cannon K *et al.* 2012 *Astrophys. J.* **748** 136 (*Preprint* [1107.2665](#))
- [83] Chu Q, Howell E J, Rowlinson A, Gao H, Zhang B, Tingay S J, Boër M and Wen L 2016 *Monthly Notices of the Royal Astronomical Society* **459** 121–139 ISSN 0035-8711 (*Preprint* <https://academic.oup.com/mnras/article-pdf/459/1/121/8113983/stw576.pdf>) URL <https://doi.org/10.1093/mnras/stw576>
- [84] Akcay S 2019 *Annalen Phys.* **531** 1800365 (*Preprint* [1808.10057](#))
- [85] Nicholl M *et al.* 2017 *Astrophys. J.* **848** L18 (*Preprint* [1710.05456](#))
- [86] Metzger B D 2017 *Preprint: arXiv* [1710.05931](#) (*Preprint* [1710.05931](#))
- [87] Metzger B D and Piro A L 2014 *Mon. Not. Roy. Astron. Soc.* **439** 3916–3930 (*Preprint* [1311.1519](#))
- [88] Ciolfi R and Siegel D M 2015 *Astrophys. J.* **798** L36 (*Preprint* [1411.2015](#))
- [89] Siegel D M and Ciolfi R 2016 *Astrophys. J.* **819** 15 (*Preprint* [1508.07939](#))
- [90] Most E R and Philippov A A 2020 *The Astrophysical Journal* **893** L6 ISSN 2041-8213 URL <http://dx.doi.org/10.3847/2041-8213/ab8196>
- [91] Totani T 2013 *Pub. Astron. Soc. Jpn.* **65** L12 (*Preprint* [1307.4985](#))
- [92] Wang J S, Yang Y P, Wu X F, Dai Z G and Wang F Y 2016 *Astrophys. J.* **822** L7 (*Preprint* [1603.02014](#))
- [93] Dokuchaev V I and Eroshenko Yu N 2017 (*Preprint* [1701.02492](#))
- [94] Sachdev S, Magee R, Hanna C, Cannon K, Singer L, SK J R, Mukherjee D, Caudill S, Chan C, Creighton J D E and *et al* 2020 *The Astrophysical Journal* **905** L25 ISSN 2041-8213 URL <http://dx.doi.org/10.3847/2041-8213/abc753>
- [95] Nitz A H, Schäfer M and Dal Canton T 2020 *Astrophys. J. Lett.* **902** L29 (*Preprint* [2009.04439](#))
- [96] Magee R, Chatterjee D, Singer L P, Sachdev S, Kovalam M, Mo G, Anderson S, Brady P, Brockill P, Cannon K, Canton T D, Chu Q, Clearwater P, Codoreanu A, Drago M, Godwin P, Ghosh S, Greco G, Hanna C, Kapadia S J, Katsavounidis E, Oloworaran V, Pace A E, Panther F, Patwary A, Pietri R D, Piotrkowski B, Prestegard T, Rei L, Sreekumar A K, Szczepańczyk M J, Valsan V, Viets A and Wade M 2021 First demonstration of early warning gravitational wave alerts (*Preprint* [2102.04555](#))
- [97] Cannon K, Caudill S, Chan C, Cousins B, Creighton J D E, Ewing B, Fong H, Godwin P, Hanna C, Hooper S, Huxford R, Magee R, Meacher D, Messick C, Morisaki S, Mukherjee D, Ohta H, Pace A, Privitera S, de Ruiter I, Sachdev S, Singer L, Singh D, Tapia R, Tsukada L, Tsuna D, Tsutsui T, Ueno K, Viets A, Wade L and Wade M 2020 Gstlal: A software framework for gravitational wave discovery (*Preprint* [2010.05082](#))
- [98] BlackGEM 2020 The blackgem telescope array "<https://astro.ru.nl/blackgem/>" accessed 2020-07-17
- [99] ZTF 2020 The zwicky transient facility "<https://www.ztf.caltech.edu/>" accessed 2020-07-17
- [100] Flaugher B, Diehl H T, Honscheid K, Abbott T M C, Alvarez O, Angstadt R, Annis J T, Antonik M, Ballester O, Beaufore L and *et al* 2015 *The Astronomical Journal* **150** 150 ISSN 1538-3881 URL <http://dx.doi.org/10.1088/0004-6256/150/5/150>

- [101] LSST 2020 The large synoptic survey telescope "<https://www.lsst.org/lsst/>" accessed 2020-07-17
- [102] SWOPE 2020 The swope telescope "<https://obs.carnegiescience.edu/swope>" accessed 2020-07-17
- [103] Subaru 2020 The subaru telescope "<https://subarutelescope.org/en/>" accessed 2020-08-03
- [104] Abbott B P *et al.* (LIGO Scientific Collaboration and Virgo Collaboration) 2016 *Phys. Rev. Lett.* **116** 221101 (*Preprint* [1602.03841](#))
- [105] Abbott B *et al.* (LIGO Scientific, Virgo) 2019 *Phys. Rev. D* **100** 104036 (*Preprint* [1903.04467](#))
- [106] Abbott R *et al.* (LIGO Scientific, Virgo) 2020 (*Preprint* [2010.14529](#))
- [107] Yunes N and Pretorius F 2009 *Phys. Rev. D* **80** 122003 (*Preprint* [0909.3328](#))
- [108] Li T, Del Pozzo W, Vitale S, Van Den Broeck C, Agathos M, Veitch J, Grover K, Sidery T, Sturani R and Vecchio A 2012 *Physical Review D* **85** 082003
- [109] Li T, Del Pozzo W, Vitale S, Van Den Broeck C, Agathos M, Veitch J, Grover K, Sidery T, Sturani R and Vecchio A 2012 *Journal of Physics: Conference Series* vol 363 (IOP Publishing) p 012028
- [110] Agathos M, Del Pozzo W, Li T G F, Van Den Broeck C, Veitch J and Vitale S 2014 *Phys. Rev. D* **89** 082001 (*Preprint* [1311.0420](#))
- [111] Hughes S A and Menou K 2005 *Astrophys. J.* **623** 689 (*Preprint* [astro-ph/0410148](#))
- [112] Ghosh A, Del Pozzo W and Ajith P 2016 *Phys. Rev. D* **94** 104070 (*Preprint* [1505.05607](#))
- [113] Ghosh A *et al.* 2016 *Phys. Rev. D* **94** 021101(R) (*Preprint* [1602.02453](#))
- [114] Mirshekari S, Yunes N and Will C M 2012 *Phys. Rev. D* **85** 024041 (*Preprint* [1110.2720](#))
- [115] Yunes N, Yagi K and Pretorius F 2016 *Phys. Rev. D* **94** 084002 (*Preprint* [1603.08955](#))
- [116] Will C M 1998 *Phys. Rev. D* **57** 2061 (*Preprint* [gr-qc/9709011](#))
- [117] Calcagni G 2010 *Phys. Rev. Lett.* **104** 251301 (*Preprint* [0912.3142](#))
- [118] Amelino-Camelia G 2002 *Nature* **418** 34–35 (*Preprint* [gr-qc/0207049](#))
- [119] Horava P 2009 *Phys. Rev. D* **79** 084008 (*Preprint* [0901.3775](#))
- [120] Sefiedgar A, Nozari K and Sepangi H 2011 *Phys. Lett. B* **696** 119–123 (*Preprint* [1012.1406](#))
- [121] Kostelecký V and Mewes M 2016 *Phys. Lett. B* **757** 510–514 (*Preprint* [1602.04782](#))
- [122] Isi M, Pitkin M and Weinstein A J 2017 *Phys. Rev. D* **96** 042001 (*Preprint* [1703.07530](#))
- [123] Abbott B P, Abbott R, Abbott T D, Acernese F, Ackley K, Adams C, Adams T, Addesso P, Adhikari R X and *et al* (LIGO Scientific Collaboration and Virgo Collaboration) 2017 *Phys. Rev. D* **96**(12) 122006 URL <https://link.aps.org/doi/10.1103/PhysRevD.96.122006>
- [124] Callister T, Biscoveanu A, Christensen N, Isi M, Matas A, Minazzoli O, Regimbau T, Sakellariadou M, Tasson J and Thrane E 2017 *Phys. Rev. X* **7** 041058 (*Preprint* [1704.08373](#))
- [125] Abbott B P *et al.* (LIGO Scientific, Virgo) 2018 *Phys. Rev. Lett.* **120** 201102 (*Preprint* [1802.10194](#))
- [126] Nishizawa A, Taruya A, Hayama K, Kawamura S and Sakagami M a 2009 *Phys. Rev. D* **79** 082002 (*Preprint* [0903.0528](#))
- [127] Isi M and Weinstein A J 2017 (*Preprint* [1710.03794](#))
- [128] Pang P T, Lo R K, Wong I C, Li T G and Van Den Broeck C 2020 *Phys. Rev. D* **101** 104055 (*Preprint* [2003.07375](#))
- [129] Abbott B, Abbott R, Abbott T, Acernese F, Ackley K, Adams C, Adams T, Addesso P, Adhikari R, Adya V and *et al* 2019 *Physical Review Letters* **123** ISSN 1079-7114 URL <http://dx.doi.org/10.1103/PhysRevLett.123.011102>
- [130] Abbott B P *et al.* (LIGO Scientific Collaboration and Virgo Collaboration) 2017 *Phys. Rev. Lett.* **119**(14) 141101 URL <https://link.aps.org/doi/10.1103/PhysRevLett.119.141101>
- [131] Chatziioannou K, Yunes N and Cornish N 2012 *Physical Review D* **86** ISSN 1550-2368 URL <http://dx.doi.org/10.1103/PhysRevD.86.022004>
- [132] Takeda H, Morisaki S and Nishizawa A 2021 *Physical Review D* **103** ISSN 2470-0029 URL <http://dx.doi.org/10.1103/PhysRevD.103.064037>
- [133] Hagihara Y, Era N, Iikawa D, Nishizawa A and Asada H 2019 *Physical Review D* **100** ISSN 2470-0029 URL <http://dx.doi.org/10.1103/PhysRevD.100.064010>
- [134] Abbott B *et al.* (LIGO Scientific, Virgo) 2019 *Phys. Rev. D* **100** 104036 (*Preprint* [1903.04467](#))

- [135] Veitch J *et al.* 2015 *Phys. Rev.* **D91** 042003 (*Preprint* 1409.7215)
- [136] Hayama K and Nishizawa A 2013 *Phys. Rev. D* **87**(6) 062003 URL <https://link.aps.org/doi/10.1103/PhysRevD.87.062003>
- [137] Regimbau T and Mandic V 2008 *Classical and Quantum Gravity* **25** 184018 URL <https://doi.org/10.1088%2F0264-9381%2F25%2F18%2F184018>
- [138] Regimbau T 2011 *Research in Astronomy and Astrophysics* **11** 369–390 URL <https://doi.org/10.1088%2F1674-4527%2F11%2F4%2F001>
- [139] Abbott R *et al.* (LIGO Scientific, Virgo, KAGRA) 2021 (*Preprint* 2101.12130)
- [140] Callister T, Fishbach M, Holz D and Farr W 2020 *Astrophys. J. Lett.* **896** L32 (*Preprint* 2003.12152)
- [141] Mitra S, Dhurandhar S, Souradeep T, Lazzarini A, Mandic V, Bose S and Ballmer S 2008 *Phys. Rev. D* **77**(4) 042002 URL <https://link.aps.org/doi/10.1103/PhysRevD.77.042002>
- [142] Talukder D, Mitra S and Bose S 2011 *Phys. Rev. D* **83**(6) 063002 URL <https://link.aps.org/doi/10.1103/PhysRevD.83.063002>
- [143] Ain A, Suresh J and Mitra S 2018 *Phys. Rev. D* **98** 024001 (*Preprint* 1803.08285)
- [144] Suresh J, Ain A and Mitra S 2020 (*Preprint* 2011.05969)
- [145] van der Walt S, Colbert S C and Varoquaux G 2011 *Comput. Sci. Eng.* **13** 22–30 (*Preprint* 1102.1523)
- [146] Virtanen P *et al.* 2020 *Nature Meth.* (*Preprint* 1907.10121)
- [147] Hunter J D 2007 *Computing in Science & Engineering* **9** 90–95
- [148] Robitaille T P *et al.* (Astropy) 2013 *Astron. Astrophys.* **558** A33 (*Preprint* 1307.6212)
- [149] Price-Whelan A *et al.* 2018 *Astron. J.* **156** 123 (*Preprint* 1801.02634)
- [150] Kluyver T, Ragan-Kelley B, Pérez F, Granger B, Bussonnier M, Frederic J, Kelley K, Hamrick J, Grout J, Corlay S, Ivanov P, Avila D, Abdalla S, Willing C and development team J 2016 *Positioning and Power in Academic Publishing: Players, Agents and Agendas* ed Loizides F and Schmidt B (Netherlands: IOS Press) pp 87–90 URL <https://eprints.soton.ac.uk/403913/>
- [151] Higson E, Handley W, Hobson M and Lasenby A 2019 *Statistics and Computing* **29** 891–913 (*Preprint* 1704.03459)
- [152] Hoy C and Raymond V 2020 (*Preprint* 2006.06639)