

CONCENTRATED SOLUTIONS TO FRACTIONAL SCHRÖDINGER EQUATIONS WITH PRESCRIBED L^2 -NORM

QING GUO, PENG LUO, CHUNHUA WANG AND JING YANG

ABSTRACT. In this work, we investigate the existence and local uniqueness of normalized k -peak solutions for the fractional Schrödinger equations with attractive interactions with a class of degenerated trapping potential with non-isolated critical points.

Precisely, applying the finite dimensional reduction method, we first obtain the existence of k -peak concentrated solutions and especially describe the relationship between the chemical potential μ and the attractive interaction a . Second, after precise analysis of the concentrated points and the Lagrange multiplier, we prove the local uniqueness of the k -peak solutions with prescribed L^2 -norm, by use of the local Pohozaev identities, the blow-up analysis and the maximum principle associated to the nonlocal operator $(-\Delta)^s$.

To our best knowledge, there is few results on the excited normalized solutions of the fractional Schrödinger equations before this present work. The main difficulty lies in the non-local property of the operator $(-\Delta)^s$. First, it makes the standard comparison argument in the ODE theory invalid to use in our analysis. Second, because of the algebraic decay involving the approximate solutions, the estimates, on the Lagrange multiplier for example, would become more subtle. Moreover, when studying the corresponding harmonic extension problem, several local Pohozaev identities are constructed and we have to estimate several kinds of integrals that never appear in the classic local Schrödinger problems. In addition, throughout our discussion, we need to distinct the different cases of $p - 1 < \frac{4s}{N}$, $p - 1 = \frac{4s}{N}$, and $p - 1 > \frac{4s}{N}$, which are called respectively that the mass-subcritical, the mass-critical, and the mass-supercritical case, due to the mass-constraint condition. Another difficulty comes from the influence of the different degenerate rates along different directions at the critical points of the potential $V(x)$.

1. INTRODUCTION AND MAIN RESULTS

In this paper, we consider the following fractional Schrödinger equation

$$\begin{cases} (-\Delta)^s u + V(x)u = au^p + \mu u, & x \in \mathbb{R}^N, \\ u \in H^s(\mathbb{R}^N), \end{cases} \quad (1.1)$$

under the mass constraint

$$\int_{\mathbb{R}^N} u^2(x) dx = 1, \quad (1.2)$$

where $s \in (0, 1)$, $p \in (1, 2_s^* - 1)$ with $2_s^* = \frac{2N}{N-2s}$ is the fractional critical Sobolev exponent.

The fractional Laplacian operator, appearing in many areas including biological modeling, physics and mathematical finances, can be regarded as the infinitesimal generator of a stable

Date: January 7, 2022.

Key words and phrases. Normalized solutions, Multi-peak solutions, Fractional Schrödinger equations, Degenerated trapping potential.

Levy process [1]. For any $s \in (0, 1)$, $(-\Delta)^s$ is the nonlocal operator defined as

$$(-\Delta)^s u = c(N, s) P.V. \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy,$$

where $P.V.$ is the principal value and $c(N, s) = \pi^{2s + \frac{N}{2}} \Gamma(s + \frac{N}{2}) / \Gamma(-s)$. One could refer to [12, 22] for more details on the fractional Laplacian operator. Particularly, this nonlocal operator $(-\Delta)^s$ in \mathbb{R}^N can be expressed as a generalized Dirichlet-to-Neumann map for a certain elliptic boundary value problem with local differential operators defined on the upper half-space $\mathbb{R}_+^{N+1} = \{(x, t) : x \in \mathbb{R}^N, t > 0\}$. Precisely, by [7], for any $u \in \dot{H}^s(\mathbb{R}^N)$, set

$$\tilde{u}(x, t) = \mathcal{P}_s[u] = \int_{\mathbb{R}^N} \mathcal{P}_s(x - z, t) u(z) dz, \quad (x, t) \in \mathbb{R}_+^{N+1},$$

where

$$\mathcal{P}_s(x, t) = \beta(N, s) \frac{t^{2s}}{(|x|^2 + t^2)^{\frac{N+2s}{2}}}$$

with a constant $\beta(N, s)$ such that $\int_{\mathbb{R}^N} \mathcal{P}_s(x, 1) dx = 1$. Then $\tilde{u} \in L^2(t^{1-2s}, K)$ for any compact set K in $\overline{\mathbb{R}_+^{N+1}}$, $\nabla \tilde{u} \in L^2(t^{1-2s}, \mathbb{R}_+^{N+1})$ and $\tilde{u} \in C^\infty(\mathbb{R}_+^{N+1})$. Moreover, \tilde{u} satisfies

$$\begin{cases} \operatorname{div}(t^{1-2s} \nabla \tilde{u}) = 0, & x \in \mathbb{R}_+^{N+1}, \\ -\lim_{t \rightarrow 0} t^{1-2s} \partial_t \tilde{u}(x, t) = \omega_s (-\Delta u)^s u(x), & x \in \mathbb{R}^N, \end{cases}$$

in the distribution sense, where $\omega_s = 2^{1-2s} \Gamma(1-s) / \Gamma(s)$. Moreover, it holds that

$$\|\tilde{u}\|_{L^2(t^{1-2s}, \mathbb{R}_+^{N+1})} = \omega_s \|u\|_{\dot{H}^s}.$$

Without loss of generality, we may assume $\omega_s = 1$.

Problems with fractional Laplacian have been extensively studied recently, see for example [2–7, 17–21, 23, 24, 31–36, 38] and the references therein. In particular, the existence of multiple spike solution involving the stable critical points of the potential $V(x)$ was considered in [15], where the critical points of $V(x)$ seem to be isolated.

We first recall the well-known results about the ground state of the following equation

$$(-\Delta)^s u + u = u^p, \quad u > 0, \quad x \in \mathbb{R}^N, \quad u(0) = \max_{x \in \mathbb{R}^N} u(x). \quad (1.3)$$

Let $N \geq 1, s \in (0, 1)$ and $1 < p < 2_s^* - 1$. Then the following hold (c.f.[18, 19]).

- (i)(Uniqueness) The ground state solution $U \in H^s(\mathbb{R}^N)$ of (1.3) is unique.
- (ii)(Symmetry, regularity, and decay) $U(x)$ is radial, positive, and strictly decreasing in $|x|$. Moreover, $U \in H^{2s+1}(\mathbb{R}^N) \cap C^\infty(\mathbb{R}^N)$ and satisfies

$$\frac{C_1}{1 + |x|^{N+2s}} \leq U(x) \leq \frac{C_2}{1 + |x|^{N+2s}}, \quad \text{for } x \in \mathbb{R}^N,$$

with some constants $C_2 \geq C_1 > 0$.

- (iii)(Non-degeneracy) The linearized operator $L_0 = (-\Delta)^s + 1 - pU^{p-1}$ is non-degenerate, i.e. its kernel is given by

$$\ker L_0 = \operatorname{span}\{\partial_{x_1} U, \partial_{x_2} U, \dots, \partial_{x_N} U\}.$$

Moreover, By Lemma C.2 of [19], for $j = 1, \dots, N$, $\partial_{x_j} U$ has the decay estimate

$$|\partial_{x_j} U| \leq \frac{C}{1 + |x|^{N+2s}}.$$

Throughout this paper, **we assume**

$$N \geq \max\{2s, 2 - 2s\}$$

and set

$$a_* = \int_{\mathbb{R}^N} U^2, \quad a_0 \equiv \begin{cases} +\infty, & \text{if } p - 1 < \frac{4s}{N}, \\ (ka_*)^{\frac{p-1}{2}}, & \text{if } p - 1 = \frac{4s}{N}, \\ 0, & \text{if } p - 1 > \frac{4s}{N}. \end{cases} \quad (1.4)$$

Our first result is as follows.

Theorem 1.1. *Suppose that u_a satisfies (1.1)-(1.2) concentrated at some points as $a \rightarrow a_0$. Then it holds that $\mu_a \rightarrow -\infty$ as $a \rightarrow a_0$.*

Moreover, if $p - 1 \leq \frac{4s}{N}$, there holds that

$$u_a = \left(\frac{-\mu_a}{a} \right)^{\frac{1}{p-1}} \left(\sum_{i=1}^k U((-\mu_a)^{\frac{1}{2s}}(x - x_{a,i})) + \omega_a \right), \quad (1.5)$$

with $\int_{\mathbb{R}^N} \left(-\frac{1}{\mu_a} |(-\Delta)^{\frac{s}{2}} \omega_a|^2 + \omega_a^2 \right) = o\left((-\mu_a)^{-\frac{N}{2s}} \right)$.

We call u_a a k -peak solution of (1.1)-(1.2) if u_a satisfies (1.5). The problem has been extensively investigated if the critical points of $V(x)$ are isolated, while few is known otherwise. In this paper, we assume that $V(x)$ obtains its local minimum or local maximum at a closed $N - 1$ dimensional hyper-surface $\Gamma_i (i = 1, \dots, k)$, satisfying $\Gamma_i \cap \Gamma_j = \emptyset$ if $i \neq j$. Precisely, we assume that

(V). There exist some $\delta > 0$ and some C^2 compact hyper surfaces $\Gamma_i (i = 1, \dots, k)$ without boundary, satisfying

$$V(x) = V_i, \quad \frac{\partial V(x)}{\partial \nu_i} = 0, \quad \frac{\partial^2 V(x)}{\partial \nu_i^2} \neq 0, \quad \text{for any } x \in \Gamma_i \text{ and } i = 1, \dots, k,$$

where $V_i \in \mathbb{R}$, ν_i is the unit outward normal of Γ_i at $x \in \Gamma_i$. Moreover, $V(x) \in C^4(\cup_{i=1}^k W_{\delta,i})$ with $W_{\delta,i} := \{x \in \mathbb{R}^N : \text{dist}(x, \Gamma_i) < \delta\}$.

Remark 1.2. *We point out that the assumption (V) was first introduced in [30], where they studied the Bose-Einstein condensates. The assumption (V) implies that the potential $V(x)$ obtains its local minimum or local maximum on the hypersurface Γ_i for $i = 1, \dots, k$. If $\delta > 0$ is small, we set $\Gamma_{t,i} = \{x : V(x) = t\} \cap W_{\delta,i}$, which consists of two compact hypersurfaces in \mathbb{R}^N without boundary for $t \in [V_0, V_0 + \sigma)$ or $t \in (V_0 - \sigma, V_0]$ provided $\sigma > 0$ is small enough. Moreover, the outward unit normal vector $\nu_{t,i}(x)$ and the j -th principal tangential unit vector $\tau_{t,i,j}(x), j = 1, \dots, N - 1$ of $\Gamma_{t,i}$ at x are Lip-continuous in $W_{\delta,i}$.*

We can apply the Pohozaev identities to show that a k -peak solution of (1.1)-(1.2) must concentrate at some critical points of $V(x)$. The following result explains where the concentrated points locate on $\Gamma = \cup_{i=1}^k \Gamma_i$.

Theorem 1.3. *Under the condition (V), if u_a is a k -peak solution of (1.1)-(1.2), concentrating at $\{b_1, b_2, \dots, b_k\}$ with $b_i \in \Gamma$, $b_i \neq b_j$ if $i \neq j$, and $|x_{a,i} - \Gamma_i| = O((-\mu_a)^{-\frac{1}{2s}})$ as $a \rightarrow a_0$, then*

$$(D_{\tau_{i,j}} \Delta V)(b_i) = 0, \quad \text{with } i = 1, \dots, k \text{ and } j = 1, \dots, N-1, \quad (1.6)$$

where $\tau_{i,j}$ is the j -th principal tangential unit vector of Γ at b_i .

To study the converse of Theorem 1.3, we need another non-degenerate condition on the critical point of $V(x)$. In fact, we define that $x_0 \in \Gamma_i$ is non-degenerate on Γ_i if there holds that

$$\frac{\partial^2 V(x_0)}{\partial \nu_i^2} \neq 0, \quad \text{and } \det \left(\left(\frac{\partial^2 \Delta V(x_0)}{\partial \tau_{i,l} \partial \tau_{i,j}} \right)_{1 \leq l, j \leq N-1} \right) \neq 0.$$

Theorem 1.4. *Assume that the condition (V) holds. If $b_i \in \Gamma$ are non-degenerate critical points of $V(x)$ on Γ for $i = 1, \dots, k$ satisfying (1.6) and $b_i \neq b_j$ for $i \neq j$, then (1.1)-(1.2) has a k -peak solution u_a concentrating at b_1, \dots, b_k as $a \rightarrow a_0$.*

On the other hand, if we assume the function $\Delta V(x)|_{x \in \Gamma_{i_0}}$ has an isolated maximum point $b \in \Gamma_{i_0}$ with some $i_0 \in \{1, \dots, k\}$, that is $\Delta V(x) < \Delta V(b)$ for all $x \in \Gamma_{i_0} \cap (B_\delta(b) \setminus b)$, then we can obtain a k -peak solution concentrating at one point.

Theorem 1.5. *Assume (V) and $\frac{\partial^2 V(x)}{\partial \nu_{i_0}^2} \neq 0$ for any $x \in \Gamma_{i_0}$ with some $i_0 \in \{1, \dots, k\}$. If $b \in \Gamma_{i_0}$ is an isolated maximum point of $\Delta V(x)|_{x \in \Gamma_{i_0}}$ on Γ_{i_0} , then for any integer $k > 0$, problem (1.1)-(1.2) has a k -peak solution u_a concentrating at b .*

Another main result of this paper is the following local uniqueness result.

Theorem 1.6. *Suppose (V), and if further $N \geq 2s+4$. Let $u_a^{(1)}(x)$ and $u_a^{(2)}(x)$ be two k -peak solutions of (1.1)-(1.2) concentrating at b_1, \dots, b_k with $b_i \in \Gamma$, and $b_i \neq b_j$ if $i \neq j$. If b_i is non-degenerate, $i = 1, \dots, k$, $\sum_{i=1}^k \Delta V(b_i) \neq 0$ when $p-1 = \frac{4s}{N}$, and*

$$\left(\frac{\partial^2 \Delta V(b_i)}{\partial \tau_{i,l} \partial \tau_{i,j}} \right)_{1 \leq l, j \leq N-1} + \frac{\partial \Delta V(b_i)}{\partial \nu_i} \text{diag}(\kappa_{i,1}, \dots, \kappa_{i,N-1}), \quad \text{for } i = 1, \dots, k$$

is non-singular, where $\kappa_{i,j}$ is the j -th principal curvature of Γ at b_i for $j = 1, \dots, N-1$, then there exists a small positive number σ , such that

$$u_a^{(1)}(x) \equiv u_a^{(2)}(x)$$

for all a with $0 < |a - (ka_*)^{\frac{p-1}{2}}| \leq \sigma$ if $p-1 = \frac{4s}{N}$, or $0 < a \leq \sigma$ if $p-1 > \frac{4s}{N}$, or $a \geq \frac{1}{\sigma}$ if $p-1 < \frac{4s}{N}$.

Remark 1.7. *In fact, the result of Theorem 1.6 can hold for more general p and N . Here, in order to avoiding writing too dispersively, we assume that $N \geq 2s+4$.*

As far as we know, there are very few results on the local uniqueness for the fractional Schrödinger equations, especially with the subcritical nonlinearities. In particular, since the Lagrange multiplier μ_a in (1.1) depends on the solution u_a , the corresponding linearized operator has changed, which brings more nontrivial analysis, as mentioned in [30]. For the local uniqueness results of peak (or bubbling) solutions, the classical moving plane method is

not available. If x_0 is a non-degenerate critical point of $V(x)$, that is, (D^2V) is non-singular at x_0 , one can prove the local uniqueness of the peak solution concentrating at x_0 either by counting the local degree of the corresponding reduced finite dimensional problem as in [8, 10, 27], or by using Pohozaev type identities as in [9, 16, 26, 28, 29]. One of the advantage in applying the Pohozaev identities can be found in dealing with the degenerate case (see [9, 16, 26]). In these results, the rate of degeneracy along each direction is the same, although the critical point x_0 is degenerate. Following [30], under the condition (V), the function $V(x)$ is non-degenerate along the normal direction ν_i of Γ_i . But along each tangential direction of Γ_i , $V(x)$ is degenerate. Such non-uniform degeneracy makes the estimates more sophisticated.

At the end of this section, we outline the main idea of the proof and discuss the main difficulties. For the existence result, we first consider the following problem without constraint,

$$\begin{cases} (-\Delta)^s w + (\lambda + V(x))w = w^p, & x \in \mathbb{R}^N, \\ w \in H^s(\mathbb{R}^N), \end{cases} \quad (1.7)$$

where $\lambda > 0$ is a large parameter. For large $\lambda > 0$, by the standard reduction argument, we could construct various positive solutions concentrating at some stable points of $V(x)$. Particularly, we can construct positive k -peak solutions for (1.7) of the form

$$w_\lambda(x) = \lambda^{\frac{1}{p-1}} \left(\sum_{j=1}^k U(\lambda^{\frac{1}{2s}}(x - x_{\lambda,i})) + \omega_\lambda \right),$$

with $\int_{\mathbb{R}^N} \left(\frac{1}{\lambda} |(-\Delta)^{\frac{s}{2}} \omega_\lambda|^2 + \omega_\lambda^2 \right) = o\left(\lambda^{-\frac{N}{2s}}\right)$. Let $u_\lambda = \frac{w_\lambda}{\left(\int_{\mathbb{R}^N} w_\lambda^2\right)^{\frac{1}{2}}}$. Then $\int_{\mathbb{R}^N} u_\lambda^2 = 1$, and

$$\begin{cases} (-\Delta)^s u_\lambda + (\lambda + V(x))u_\lambda = a_\lambda u_\lambda^p, & x \in \mathbb{R}^N, \\ u_\lambda \in H^s(\mathbb{R}^N), \end{cases}$$

with

$$a_\lambda = \left(\int_{\mathbb{R}^N} w_\lambda^2 \right)^{\frac{p-1}{2}} = \left(k \lambda^{\frac{2}{p-1} - \frac{N}{2s}} (a_* + o(1)) \right)^{\frac{p-1}{2}}.$$

We note that $a_\lambda > 0$. Moreover, as $\lambda \rightarrow +\infty$, $a_\lambda \rightarrow a_0$, which is defined by (1.4). Therefore, we obtain a concentrated solution with k -peaks for (1.7) with normalized L^2 -norm, where $\mu = -\lambda$ and some suitable a_λ . Hence, we are sufficed to answer a converse question that for any $a > 0$ close to a_0 , whether one can choose a suitable large $\lambda = \lambda_a > 0$, such that (1.1)–(1.2) hold with $\mu = -\lambda_a$, $u_a = \frac{w_{\lambda_a}}{\left(\int_{\mathbb{R}^N} w_{\lambda_a}^2\right)^{\frac{1}{2}}}$, finally concluding Theorem 1.4. In fact,

we solve the problem by discussing the relationship between a and μ .

Recently, for the classical BECs problem ($s = 1, p = 3, N = 2, 3$), a complete description of the solutions u_a concentrating at a_* related to problem was given in [30]. Specifically, in this case if u_a is a solution of problem (1.1)–(1.2) with $s = 1, p = 3$ and $N = 2, 3$, which is concentrated at some points as $a \rightarrow a_0$ with $a_0 \in \mathbb{R}$, then it holds

$$a_0 = k a_* > 0 \quad \text{and} \quad \mu = \mu_a \rightarrow -\infty \quad \text{as} \quad a \rightarrow a_0,$$

where k is the number of peaks of the concentrated solution u_a . Moreover, the existence and local uniqueness of this kind concentrated solutions have been proved in [30]. Unlike the argument used in [30], a non-existence result, which is of great importance to prove the main

result, could not be obtained by a standard comparison argument in ODE theory. We would apply a totally different method—the sliding method—corresponding to the non-local operator to show a counterpart result (Theorem 1.1).

On the other hand, we point out that, due to the non-local property of the operator $(-\Delta)^s$, we cannot construct the local Pohozaev identities directly, which, however, is inevitable when we carry out the blow-up analysis especially in the proof of the local uniqueness. To this end, we are supposed to study the corresponding harmonic extension problem (see section 4 and section 5), and have to estimate several kind of integrals which never appear in the classic local Schrödinger problems. Similar arguments can be found in [25] etc..

Last but not least, throughout our discussion, we need to distinct the different cases of $p-1 < \frac{4s}{N}$, $p-1 = \frac{4s}{N}$, and $p-1 > \frac{4s}{N}$, which are called respectively that the mass-subcritical, the mass-critical, and the mass-supercritical case. Especially compared with the problem without any constraint, in the proof of the local uniqueness result, we have to adopt one more different local Pohozaev identity, associated with the mass-constraint condition.

This paper is organized as follows. In section 2, we prove Theorem 1.1 using the sliding method. Then in section 3, we estimate the Lagrange multiplier μ_a in terms of a . The results for the location of the peaks and for the existence of peak solutions are proved in section 4, and the local uniqueness of peak solutions are investigated in section 5.

2. PROOF OF THEOREM 1.1 BY A NON-EXISTENCE RESULT

Lemma 2.1. *Assume that $P(x)$ satisfies $P(x) > 1$ in $B_R(0) \setminus B_r(0)$ for some fixed $r > 0$ and large $R > 0$. Then the problem*

$$(-\Delta)^s u = P(x)u, \quad u > 0, \quad \text{in } \mathbb{R}^N$$

has no solution.

Proof. We take λ_1 as the first eigenvalue of $(-\Delta)^s$ and ϕ as the associated eigenfunction satisfying

$$\begin{cases} (-\Delta)^s \phi = \lambda_1 \phi, & x \in B, \\ \phi = 0, & x \in B^c, \end{cases}$$

where $B \subset B_R(0) \setminus B_r(0)$ is a large ball. Since $R > 0$ is large enough, it may hold that

$$\lambda_1 \leq 1 < P(x) \text{ in } B.$$

Now, we set $w(x) := \max_{x \in B} \frac{\phi(x)}{u(x)} \cdot u(x)$. Obviously, $w(x) \geq \phi(x)$ and $w(x_0) = \phi(x_0)$ with $x_0 \in B$ the maximal point of $\frac{\phi(x)}{u(x)}$. To get a contradiction, we are sufficed to prove that $w(x) > \phi(x)$ by use of the maximum principle.

In fact, if we denote $\gamma = \max_{x \in B} \frac{\phi(x)}{u(x)}$, then there holds that

$$(-\Delta)^s w = \gamma(-\Delta)^s u = \gamma P(x)u \geq \gamma \lambda_1 u = \lambda_1 \max_{x \in B} \frac{\phi(x)}{u(x)} u \geq \lambda_1 \phi = (-\Delta)^s \phi,$$

which gives that

$$(-\Delta)^s w \geq (-\Delta)^s \phi \text{ in } B.$$

On the other hand, $w \geq \phi$ in B^c obviously. Moreover, since here $w \not\equiv \phi$, the strong maximum principle implies that $w > \phi$ concluding the Lemma by contradiction. \square

Remark 2.2. *For the classical BECs problem when $s = 1, p = 3$ and $N = 2, 3$ in [30], the standard comparison method from the ODE theory is sufficed to show such a non-existence result, which, however, is invalid in the fractional case. We take the sliding method to overcome this obstacle, which, developed by Chen, Li and Zhu [11], would be typically available in the study of some other similar problems when dealing with the fractional operator.*

Proof of Theorem 1.1. We divide the proofs into following three steps.

Step 1. We first show $\mu_a \rightarrow -\infty$. By contradiction, we first suppose $|\mu_a| \leq M$. Since $\int_{\mathbb{R}^N} u_a^2 = 1$, we could use the argument in [21] to get u_a is uniformly bounded, which means that u_a does not blow up. Then if $\mu_a \rightarrow +\infty$, we set $P(x) = \mu_a - V(x) + au_a^{p-1}$. Since u_a concentrates at some points, we may assume

$$au_a^{p-1} > -1 \text{ in } \mathbb{R}^N \setminus B_r(0) \text{ for some } r > 0.$$

Hence, for any $R > 0$, we have

$$P(x) > 1 \text{ for } x \in B_R(0) \setminus B_r(0),$$

which gives a contradiction by Lemma 2.1.

Step 2. It holds that $a > 0$. If not, we could obtain then

$$(-\Delta)^s u_a \leq 0.$$

By the maximum principle corresponding to the fractional operator [21], we find $u_a \leq 0$, which is a contradiction.

Step 3. In the case of $p - 1 \leq \frac{4s}{N}$. Denote $v_a(x) = (-\mu_a)^{-\frac{1}{p-1}} u_a((-\mu_a)^{-\frac{1}{2s}} x)$. Then

$$(-\Delta)^s v_a + \left(1 + \frac{1}{-\mu_a} V((-\mu_a)^{-\frac{1}{2s}} x)\right) v_a = av_a^p$$

and $\int_{\mathbb{R}^N} v_a^2 = (-\mu_a)^{\frac{N}{2s} - \frac{2}{p-1}}$. Therefore, we apply the Moser iteration to get v_a is bounded uniformly in a . Let y_a be a maximum point of v_a . Then we could show that a is bounded from below. Applying the blow-up argument established by [21], we obtain that there exists some integer $k > 0$ such that

$$v_a = \sum_{i=1}^k U_{a_0}(x - y_{a,i}) + \bar{\omega}_a,$$

for some $y_{a,i} \in \mathbb{R}^N$ with $|y_{a,i} - y_{a,j}| \rightarrow +\infty$ for $i \neq j$, where U_{a_0} is the unique positive solution of

$$\begin{cases} (-\Delta)^s U_{a_0} + U_{a_0} = a_0 U_{a_0}^p, \\ U_{a_0}(0) = \max_{x \in \mathbb{R}^N} U_{a_0}(x), \end{cases}$$

and

$$\int_{\mathbb{R}^N} (|(-\Delta)^{\frac{s}{2}} \bar{\omega}_a|^2 + \bar{\omega}_a^2) = o(1)$$

Since $\int_{\mathbb{R}^N} u_a^2 = 1$, we have

$$\int_{\mathbb{R}^N} v_a^2 = (-\mu_a)^{\frac{N}{2s} - \frac{2}{p-1}}.$$

In view of $U_a = a_0^{-\frac{1}{p-1}} U$, there holds that

$$(-\mu_a)^{\frac{N}{2s} - \frac{2}{p-1}} = k a_0^{-\frac{2}{p-1}} a_* + o(1),$$

implying the conclusion. \square

3. SOME ESTIMATES BY POHOZAEV IDENTITIES

We use Pohozaev identities to estimate μ_a with respect to a . Let $\varepsilon = (-\mu_a)^{-\frac{1}{2s}}$ and $v_a = (-\frac{\mu_a}{a})^{-\frac{1}{p-1}} u_a$. Then (1.1) turns to

$$\varepsilon^{2s} (-\Delta)^s v_a + (1 + \varepsilon^{2s} V(x)) v_a = v_a^p, \quad v_a \in H^s(\mathbb{R}^N). \quad (3.1)$$

For any $a > 0$, we define the norm $\|v\|_a = \left(\int_{\mathbb{R}^N} \varepsilon^{2s} |(-\Delta)^{\frac{s}{2}} v|^2 + v^2 \right)^{\frac{1}{2}}$.

Hence, a k -peak solution of (3.1) has the form of

$$v_a(x) = \sum_{i=1}^k U_{\varepsilon, x_{a,i}}(x) + \varphi_a(x),$$

where

$$U_{\varepsilon, x_{a,i}}(x) = (1 + \varepsilon^{2s} V_i)^{\frac{1}{p-1}} U \left(\frac{(1 + \varepsilon^{2s} V_i)^{\frac{1}{2s}} (x - x_{a,i})}{\varepsilon} \right), \quad \|\varphi_a\|_a = o(\varepsilon^{\frac{N}{2}}).$$

We write then

$$\varepsilon^{2s} (-\Delta)^s \varphi_a + \left((1 + \varepsilon^{2s} V(x)) - p \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^{p-1} \right) \varphi_a = N_a(\varphi_a) + l_a(x),$$

where

$$l_a(x) = -\varepsilon^{2s} \sum_{i=1}^k (V(x) - V_i) U_{\varepsilon, x_{a,i}} + \left(\sum_{i=1}^k U_{\varepsilon, x_{a,i}} \right)^p - \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^p,$$

and

$$N_a(\varphi_a) = \left(\sum_{i=1}^k U_{\varepsilon, x_{a,i}} + \varphi_a \right)^p - \left(\sum_{i=1}^k U_{\varepsilon, x_{a,i}} \right)^p - p \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^{p-1} \varphi_a. \quad (3.2)$$

Set

$$E_{a, x_{a,i}} := \left\{ u \in H^s(\mathbb{R}^N) : \left\langle u, \frac{\partial U_{\varepsilon, x_{a,i}}}{\partial x_j} \right\rangle_a = 0, j = 1, \dots, N \right\}.$$

One may move $x_{a,i}$ such that the perturbing term $\varphi_a \in \bigcap_{i=1}^k E_{a, x_{a,i}}$.

Let L_a be the bounded linear operator from $H^s(\mathbb{R}^N)$ to itself, with the form of

$$\langle L_a u, v \rangle_a = \int_{\mathbb{R}^N} \varepsilon^{2s} (-\Delta)^{\frac{s}{2}} u (-\Delta)^{\frac{s}{2}} v + (1 + \varepsilon^{2s} V(x)) uv - p \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^{p-1} uv. \quad (3.3)$$

It is standard to prove the following result.

Lemma 3.1. *There exists some constant $c > 0$, such that for all a around a_0 , it holds then*

$$\|L_a u\|_a \geq c \|u\|_a, \quad \text{for all } u \in \bigcap_{i=1}^k E_{a, x_{a,i}}.$$

Lemma 3.2. *A k -peak solution v_a of (3.1) concentrating at b_1, \dots, b_k has the following form*

$$v_a(x) = \sum_{i=1}^k U_{\varepsilon, x_{a,i}}(x) + \varphi_a(x),$$

with $\varphi_a \in \bigcap_{i=1}^k E_{a, x_{a,i}}$ and

$$\begin{aligned} \|\varphi_a\|_a = & O\left(\left| \sum_{i=1}^k (V(x_{a,i}) - V_i) \right| \varepsilon^{\frac{N}{2}+2s} + \left| \sum_{i=1}^k \nabla V(x_{a,i}) \right| \varepsilon^{\frac{N}{2}+2s+1} + \varepsilon^{\frac{N}{2}+2s+2} \right) \\ & + \begin{cases} O\left(\varepsilon^{\frac{N}{2} + \frac{p}{2}(N+2s)} \right), & \text{if } 1 < p \leq 2, \\ O\left(\varepsilon^{\frac{N}{2} + (N+2s)} \right), & \text{if } p > 2. \end{cases} \end{aligned} \quad (3.4)$$

Proof. By calculation, we have for any $v \in H^s(\mathbb{R}^N)$,

$$\begin{aligned} \langle l_a(x), v \rangle_a &= -\varepsilon^{2s} \sum_{i=1}^k \int_{\mathbb{R}^N} (V(x) - V_i) U_{\varepsilon, x_{a,i}} v + \int_{\mathbb{R}^N} \left(\left(\sum_{i=1}^k U_{\varepsilon, x_{a,i}} \right)^p - \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^p \right) v \\ &= -\varepsilon^{2s} \sum_{i=1}^k \int_{\mathbb{R}^N} (V(x) - V(x_{a,i})) U_{\varepsilon, x_{a,i}} v - \varepsilon^{2s} \sum_{i=1}^k \int_{\mathbb{R}^N} (V(x_{a,i}) - V_i) U_{\varepsilon, x_{a,i}} v \\ &\quad + \int_{\mathbb{R}^N} \left(\left(\sum_{i=1}^k U_{\varepsilon, x_{a,i}} \right)^p - \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^p \right) v := -A_1 - A_2 + A_3. \end{aligned}$$

We estimate A_1 as follows

$$\begin{aligned} A_1 &= \varepsilon^{2s} \sum_{i=1}^k \int_{\mathbb{R}^N} (V(x) - V(x_{a,i})) U_{\varepsilon, x_{a,i}} v \\ &\leq \varepsilon^{2s} \sum_{i=1}^k \int_{B_\delta(x_{a,i})} (\nabla V(x_{a,i}) \cdot (x - x_{a,i}) + O(|x - x_{a,i}|^2)) U_{\varepsilon, x_{a,i}} v \\ &\quad + C \varepsilon^{2s} \left(\sum_{i=1}^k \int_{B_\delta^c(x_{a,i})} U_{\varepsilon, x_{a,i}}^2 \right)^{\frac{1}{2}} \|v\|_{L^2} \\ &\leq C \varepsilon^{2s + \frac{N}{2} + 1} \left| \sum_{i=1}^k \nabla V(x_{a,i}) \right| \cdot \|v\|_a + \varepsilon^{2s + \frac{N}{2} + 2} \|v\|_a. \end{aligned}$$

Similarly,

$$A_2 = \varepsilon^{2s} \sum_{i=1}^k \int_{\mathbb{R}^N} (V(x_{a,i}) - V_i) U_{\varepsilon, x_{a,i}} v \leq C \varepsilon^{2s + \frac{N}{2}} \sum_{i=1}^k |V(x_{a,i}) - V_i| \cdot \|v\|_a.$$

Finally, since

$$|A_3| = \left| \int_{\mathbb{R}^N} \left(\left(\sum_{i=1}^k U_{\varepsilon, x_{a,i}} \right)^p - \sum_{i=1}^k U_{\varepsilon, x_{a,i}}^p \right) v \right| = \begin{cases} o\left(\int_{\mathbb{R}^N} \sum_{i \neq j} U_{\varepsilon, x_{a,i}}^{p-1} U_{\varepsilon, x_{a,j}} v \right), & \text{if } p > 2, \\ o\left(\int_{\mathbb{R}^N} \sum_{i \neq j} U_{\varepsilon, x_{a,i}}^{\frac{p}{2}} U_{\varepsilon, x_{a,j}}^{\frac{p}{2}} v \right), & \text{if } 1 < p \leq 2, \end{cases}$$

then by Lemma A.1, we compute directly that if $p > 2$

$$\begin{aligned} \left| \int_{\mathbb{R}^N} \sum_{i \neq j} U_{\varepsilon, x_{a,i}}^{p-1} U_{\varepsilon, x_{a,j}} v \right| &\leq C \left(\int_{\mathbb{R}^N} \sum_{i \neq j} U_{\varepsilon, x_{a,i}}^{2(p-1)} U_{\varepsilon, x_{a,j}}^2 \right)^{\frac{1}{2}} \|v\|_a \\ &\leq C \varepsilon^{\frac{N}{2}} \sum_{i \neq j} \left(\int_{\mathbb{R}^N} \frac{1}{(1 + |y - \frac{x_{a,j}}{\varepsilon}|)^{2(N+2s)(p-1)}} \frac{1}{(1 + |y - \frac{x_{a,i}}{\varepsilon}|)^{2(N+2s)}} \right)^{\frac{1}{2}} \|v\|_a \\ &\leq C \varepsilon^{\frac{N}{2}} \sum_{i \neq j} \frac{1}{|\frac{x_{a,j} - x_{a,i}}{\varepsilon}|^{N+2s}} \left(\int_{\mathbb{R}^N} \frac{1}{(1 + |y - \frac{x_{a,j}}{\varepsilon}|)^{2(N+2s)(p-1)}} + \frac{1}{(1 + |y - \frac{x_{a,i}}{\varepsilon}|)^{2(N+2s)(p-1)}} \right)^{\frac{1}{2}} \|v\|_a \\ &\leq C \varepsilon^{\frac{N}{2} + N + 2s} \|v\|_a; \end{aligned}$$

while for $1 < p \leq 2$,

$$\begin{aligned} \left| \int_{\mathbb{R}^N} \sum_{i \neq j} U_{\varepsilon, x_{a,i}}^{\frac{p}{2}} U_{\varepsilon, x_{a,j}}^{\frac{p}{2}} v \right| &\leq C \left(\int_{\mathbb{R}^N} \sum_{i \neq j} U_{\varepsilon, x_{a,i}}^p U_{\varepsilon, x_{a,j}}^p \right)^{\frac{1}{2}} \|v\|_a \\ &\leq C \varepsilon^{\frac{N}{2}} \sum_{i \neq j} \left(\int_{\mathbb{R}^N} \frac{1}{(1 + |y - \frac{x_{a,j}}{\varepsilon}|)^{p(N+2s)}} \frac{1}{(1 + |y - \frac{x_{a,i}}{\varepsilon}|)^{p(N+2s)}} \right)^{\frac{1}{2}} \|v\|_a \\ &\leq C \varepsilon^{\frac{N}{2}} \sum_{i \neq j} \frac{1}{|\frac{x_{a,j} - x_{a,i}}{\varepsilon}|^{\frac{p}{2}(N+2s)}} \|v\|_a \leq C \varepsilon^{\frac{N}{2} + \frac{p}{2}(N+2s)} \|v\|_a. \end{aligned}$$

To sum up, the result has been proved. □

Lemma 3.3. *It holds that*

$$\int_{\mathbb{R}^N} U^2 = \left(\frac{N}{s(p+1)} - \frac{N}{2s} + 1 \right) \int_{\mathbb{R}^N} U^{p+1}.$$

Proof. It is directly obtained by the following Pohozaev identities:

$$\frac{N-2s}{2} \int_{\mathbb{R}_+^{N+1}} t^{1-2s} |\nabla \tilde{U}|^2 = -\frac{N}{2} \int_{\mathbb{R}^N} U^2 + \frac{N}{p+1} \int_{\mathbb{R}^N} U^{p+1}$$

and

$$\int_{\mathbb{R}_+^{N+1}} t^{1-2s} |\nabla \tilde{U}|^2 = - \int_{\mathbb{R}^N} U^2 + \int_{\mathbb{R}^N} U^{p+1}.$$

□

Proposition 3.4. *There holds that*

$$\begin{aligned}
 a^{\frac{2}{p-1}} &= \sum_{i=1}^k (V_i - \mu)^{\frac{2}{p-1} - \frac{N}{2s}} \int_{\mathbb{R}^N} U^2 + O\left(\left|\sum_{i=1}^k (V(x_{a,i}) - V_i)\right| (-\mu)^{\frac{2}{p-1} - 1 - \frac{N}{2s}} \right. \\
 &\quad \left. + \left|\sum_{i=1}^k \nabla V(x_{a,i})\right| (-\mu)^{\frac{2}{p-1} - 1 - \frac{N}{2s} - \frac{1}{2s}} + (-\mu)^{\frac{2}{p-1} - 1 - \frac{N}{2s} - \frac{1}{s}}\right) \\
 &\quad + O\left(\sum_{i \neq j} (-\mu)^{\frac{2}{p-1} - \frac{N}{s} - 1} |x_{a,j} - x_{a,i}|^{-(N+2s)}\right) + \begin{cases} O\left((- \mu)^{\frac{2}{p-1} - \frac{N}{2s} - \frac{pN}{4s} - \frac{p}{2}}\right), & \text{if } 1 < p \leq 2, \\ O\left((- \mu)^{\frac{2}{p-1} - \frac{N}{s} - 1}\right), & \text{if } p > 2. \end{cases}
 \end{aligned}$$

Proof. Let u_a be a solution of (1.1)-(1.2). It holds that

$$\begin{aligned}
 1 &= \int_{\mathbb{R}^N} |u_a(x)|^2 = \int_{\mathbb{R}^N} \left(\sum_{i=1}^k \frac{(V_i - \mu)^{\frac{1}{p-1}}}{a} U((V_i - \mu)^{\frac{1}{2s}}(x - x_{a,i})) + \frac{(-\mu)^{\frac{1}{p-1}}}{a} \varphi_a \right)^2 \\
 &= \sum_{i=1}^k \frac{(V_i - \mu)^{\frac{2}{p-1} - \frac{N}{2s}}}{a^{\frac{2}{p-1}}} \int_{\mathbb{R}^N} U^2 + \sum_{i=1}^k \frac{(-\mu)^{\frac{1}{p-1}}}{a^{\frac{2}{p-1}}} (V_i - \mu)^{\frac{1}{p-1} - \frac{N}{2s}} \int_{\mathbb{R}^N} U(x) \varphi_a((V_i - \mu)^{-\frac{1}{2s}}x + x_{a,i}) \\
 &\quad + \sum_{i \neq j} \frac{(V_i - \mu)^{\frac{1}{p-1} - \frac{N}{2s}}}{a^{\frac{2}{p-1}}} (V_j - \mu)^{\frac{1}{p-1}} \int_{\mathbb{R}^N} U(x) U\left(\frac{V_j - \mu}{V_i - \mu} \frac{1}{2s}x - (V_j - \mu)^{\frac{1}{2s}}(x_{a,j} - x_{a,i})\right) \\
 &\quad + \frac{(-\mu)^{\frac{2}{p-1}}}{a^{\frac{2}{p-1}}} \int_{\mathbb{R}^N} \varphi_a^2 \\
 &= \sum_{i=1}^k \frac{(V_i - \mu)^{\frac{2}{p-1} - \frac{N}{2s}}}{a^{\frac{2}{p-1}}} \int_{\mathbb{R}^N} U^2 + \sum_{i=1}^k \frac{(-\mu)^{\frac{1}{p-1}}}{a^{\frac{2}{p-1}}} (V_i - \mu)^{\frac{1}{p-1} - \frac{N}{4s}} O(\|\varphi_a\|_{L^2}) \\
 &\quad + O\left(\sum_{i \neq j} \frac{(V_i - \mu)^{\frac{1}{p-1} - \frac{N}{2s}} (V_j - \mu)^{\frac{1}{p-1} - \frac{N}{2s} - 1}}{a^{\frac{2}{p-1}}} |x_{a,j} - x_{a,i}|^{-(N+2s)}\right) + \frac{(-\mu)^{\frac{2}{p-1}}}{a^{\frac{2}{p-1}}} \int_{\mathbb{R}^N} \varphi_a^2 \\
 &= \sum_{i=1}^k \frac{(V_i - \mu)^{\frac{2}{p-1} - \frac{N}{2s}}}{a^{\frac{2}{p-1}}} \int_{\mathbb{R}^N} U^2 + \frac{(-\mu)^{\frac{2}{p-1} - \frac{N}{4s}}}{a^{\frac{2}{p-1}}} O\left(\left|\sum_{i=1}^k (V(x_{a,i}) - V_i)\right| \varepsilon^{\frac{N}{2} + 2s} \right. \\
 &\quad \left. + \left|\sum_{i=1}^k \nabla V(x_{a,i})\right| \varepsilon^{\frac{N}{2} + 2s + 1} + \varepsilon^{\frac{N}{2} + 2s + 2}\right) + \frac{(-\mu)^{\frac{2}{p-1} - \frac{N}{4s}}}{a^{\frac{2}{p-1}}} \begin{cases} O\left(\varepsilon^{\frac{N}{2} + \frac{p}{2}(N+2s)}\right), & \text{if } 1 < p \leq 2, \\ O\left(\varepsilon^{\frac{N}{2} + (N+2s)}\right), & \text{if } p > 2, \end{cases} \\
 &\quad + O\left(\sum_{i \neq j} \frac{(-\mu)^{\frac{2}{p-1} - \frac{N}{s} - 1}}{a^{\frac{2}{p-1}}} |x_{a,j} - x_{a,i}|^{-(N+2s)}\right),
 \end{aligned}$$

which gives the result. \square

4. EXISTENCE OF THE PEAK SOLUTIONS

4.1. Locating of the peaks.

Let v_a be a peak solution of (3.1). Following the argument in section 5 of [15], we obtain that the concentrating solution v_a to (3.1) must satisfy that

$$\sup_{x \in \mathbb{R}^N} \left(\sum_{i=1}^k \frac{1}{\left(1 + \left|\frac{x-x_{a,i}}{\varepsilon}\right|^2\right)^{\frac{N+2s-\theta}{2}}}\right)^{-1} |v_a(x)| < +\infty, \quad (4.1)$$

where θ is any constant such that $\theta \in (0, \frac{N}{2} + 2s)$. Now we quote the extension of v_a and its equation:

$$\begin{cases} \operatorname{div}(t^{1-2s} \nabla \tilde{v}_a) = 0, & x \in \mathbb{R}_+^{N+1}, \\ -\varepsilon^{2s} \lim_{t \rightarrow 0} t^{1-2s} \partial_t \tilde{v}_a(y, t) = v_a^p - (1 + \varepsilon^{2s} V(x)) v_a, & x \in \mathbb{R}^N. \end{cases} \quad (4.2)$$

Denote

$$\begin{aligned} B_\rho(x_0) &= \{y \in \mathbb{R}^N : |y - x_0| \leq \rho\} \subseteq \mathbb{R}^N, \\ \mathbf{B}_\rho^+(x_0) &= \{Y = (y, t) : |Y - (x_0, 0)| \leq \rho, t > 0\} \subseteq \mathbb{R}_+^{N+1}, \\ \partial' \mathbf{B}_\rho^+(x_0) &= \{Y = (y, t) : |Y - (x_0, 0)| \leq \rho, t = 0\} \subseteq \mathbb{R}^N, \\ \partial'' \mathbf{B}_\rho^+(x_0) &= \{Y = (y, t) : |Y - (x_0, 0)| = \rho, t > 0\} \subseteq \mathbb{R}_+^{N+1}, \\ \partial \mathbf{B}_\rho^+(x_0) &= \partial' \mathbf{B}_\rho^+(x_0) \cup \partial'' \mathbf{B}_\rho^+(x_0). \end{aligned}$$

Multiplying (4.2) by $\frac{\partial \tilde{v}_a}{\partial x_j}$ and integrating by parts on $B_\rho(x_{a,i})$ (or $\mathbf{B}_\rho^+(x_{a,i})$) with some $\rho > 0$, we have the following Pohozaev identities:

$$\begin{aligned} \varepsilon^{2s} \int_{B_\rho(x_{a,i})} \frac{\partial V(x)}{\partial x_j} v_a^2 &= -2\varepsilon^{2s} \int_{\partial'' \mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} \frac{\partial \tilde{v}_a}{\partial \nu} \frac{\partial \tilde{v}_a}{\partial y^j} + \varepsilon^{2s} \int_{\partial'' \mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} |\nabla \tilde{v}_a|^2 \nu_j \\ &\quad + \int_{\partial B_\rho(x_{a,i})} (1 + \varepsilon^{2s} V(x)) v_a^2 \nu_j - \frac{2}{p+1} \int_{\partial B_\rho(x_{a,i})} v_a^{p+1} \nu_j. \end{aligned} \quad (4.3)$$

Using Lemma A.2 and Remark A.3, we estimate that for $(y-x)^2 + t^2 = \rho^2, t > 0, \theta \in (0, \frac{N}{2} + 2s)$,

$$\begin{aligned} |\tilde{v}_a(y, t)| &\leq C \sum_{i=1}^k \int_{\mathbb{R}^N} \frac{t^{2s}}{(|y-\xi|+t)^{N+2s}} \frac{1}{\left(1 + \left|\frac{\xi-x_{a,i}}{\varepsilon}\right|\right)^{N+2s-\theta}} d\xi \\ &\leq C \begin{cases} \varepsilon^{N+2s-\theta} \sum_{i=1}^k \frac{1}{(1 + |y-x_{a,i}|)^{N+2s-\theta}}, & \theta > 2s, \\ \varepsilon^N \sum_{i=1}^k \frac{1}{(1 + |y-x_{a,i}|)^{N+2s-\theta}}, & \theta < 2s. \end{cases} \end{aligned}$$

Similarly, we could also prove that

$$\left| \frac{\partial}{\partial y^j} \tilde{v}_a(y, t) \right|, \left| \frac{\partial}{\partial \nu} \tilde{v}_a(y, t) \right| \leq C \begin{cases} \varepsilon^{N+2s-\theta} \sum_{i=1}^k \frac{1}{(1 + |y-x_{a,i}|)^{N+2s-\theta}}, & \theta > 2s, \\ \varepsilon^N \sum_{i=1}^k \frac{1}{(1 + |y-x_{a,i}|)^{N+2s-\theta}}, & \theta < 2s. \end{cases}$$

Therefore, we estimate the boundary terms as follows: if $\theta > 2s$

$$\begin{aligned} & -2\varepsilon^{2s} \int_{\partial''\mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} \frac{\partial \tilde{v}_a}{\partial \nu} \frac{\partial \tilde{v}_a}{\partial y^j} \\ & \leq C\varepsilon^{2s} \varepsilon^{2(N+2s-\theta)} \int_{\partial''\mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} \left(\sum_{i=1}^k \frac{1}{(1+|y-x_{a,i}|)^{N+2s-\theta}} \right)^2 \leq C\varepsilon^{2(N+3s-\theta)}, \end{aligned}$$

and similarly,

$$\begin{aligned} & \varepsilon^{2s} \int_{\partial''\mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} |\nabla \tilde{v}_a|^2 \nu_j \leq C\varepsilon^{2(N+3s-\theta)}; \\ & \int_{\partial B_\rho(x_{a,i})} (1 + \varepsilon^{2s} V(x)) v_a^2 \nu_j \leq C\varepsilon^{2(N+2s-\theta)}; \\ & \frac{2}{p+1} \int_{\partial B_\rho(x_{a,i})} v_a^{p+1} \nu_j \leq C\varepsilon^{(p+1)(N+2s-\theta)}. \end{aligned}$$

While if $\theta < 2s$,

$$\begin{aligned} & -2\varepsilon^{2s} \int_{\partial''\mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} \frac{\partial \tilde{v}_a}{\partial \nu} \frac{\partial \tilde{v}_a}{\partial y^j} \leq C\varepsilon^{2s} \varepsilon^{2N} \int_{\partial''\mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} \left(\sum_{i=1}^k \frac{1}{(1+|y-x_{a,i}|)^{N+2s-\theta}} \right)^2 \\ & \leq C\varepsilon^{2(N+s)}; \\ & \varepsilon^{2s} \int_{\partial''\mathbf{B}_\rho^+(x_{a,i})} t^{1-2s} |\nabla \tilde{v}_a|^2 \nu_j \leq C\varepsilon^{2(N+s)}; \\ & \int_{\partial B_\rho(x_{a,i})} (1 + \varepsilon^{2s} V(x)) v_a^2 \nu_j \leq C\varepsilon^{2(N+2s-\theta)}; \end{aligned}$$

and

$$\int_{\partial B_\rho(x_{a,i})} v_a^{p+1} \nu_j \leq C\varepsilon^{(p+1)(N+2s-\theta)}.$$

Hence (4.3) gives then for $\theta \in (0, \frac{N}{2} + 2s)$,

$$\int_{B_\rho(x_{a,i})} \frac{\partial V(x)}{\partial x_j} v_a^2 = \begin{cases} O(\varepsilon^{2(N+s-\theta)}), & \theta > 2s, \\ O(\varepsilon^{\min\{2N, 2(N+s-\theta)\}}), & \theta < 2s, \end{cases} \quad (4.4)$$

which implies the first necessary condition for the concentrated points b_i :

$$\nabla V(b_i) = 0, \quad \text{for } i = 1, \dots, k.$$

Remark 4.1. In our discussion, since we are sufficed to consider certain small $\theta \in (0, \frac{N}{2} + 2s)$ fixed, then for simplicity, we may as well suppose

$$\theta < s.$$

Hence (4.4) turns to be

$$\int_{B_\rho(x_{a,i})} \frac{\partial V(x)}{\partial x_j} v_a^2 = O(\varepsilon^{2N}). \quad (4.5)$$

Proof of Theorem 1.3. In view of $x_{a,i} \rightarrow b_i \in \Gamma_i$, if Γ_i is a local minimum set of $V(x)$, there exists some $t_a \in [V_i, V_i + \theta]$; while if Γ_i is a local maximum set of $V(x)$, there exists some $t_a \in [V_i - \theta, V_i]$, such that $x_{a,i} \in \Gamma_{t_a,i}$.

Let $H(x) = \langle \nabla V(x), \tau_{a,i} \rangle$ and $\tau_{a,i}$ be the unit tangential vector of $\Gamma_{t_a,i}$ at $x_{a,i}$. Then

$$H(x_{a,i}) = 0.$$

On the one hand, we have the expansion for $x \in B_\rho(x_{a,i})$,

$$H(x) = \langle \nabla H(x_{a,i}), x - x_{a,i} \rangle + \frac{1}{2} \langle \langle \nabla^2 H(x_{a,i}), x - x_{a,i} \rangle, x - x_{a,i} \rangle + o(|x - x_{a,i}|^2),$$

and then,

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} H(x) U_{\varepsilon, x_{a,i}}^2(x) \\ &= -2 \int_{B_\rho(x_{a,i})} H(x) U_{\varepsilon, x_{a,i}}(x) \varphi_a - \int_{B_\rho(x_{a,i})} H(x) \varphi_a^2 + O(\varepsilon^{2(N+s-\theta)}) \\ &= O\left(\left(\varepsilon^{\frac{N}{2}+1} |\nabla H(x_{a,i})| + \varepsilon^{\frac{N}{2}+2}\right) \|\varphi_a\|_a + \varepsilon |\nabla H(x_{a,i})| \cdot \|\varphi_a\|_a^2\right) + O\left(\varepsilon^{2(N+s-\theta)}\right) \\ &= O\left(\varepsilon^{N+2s+2}\right), \end{aligned} \quad (4.6)$$

where we used the fact that, by assumption,

$$|V(x_{a,i}) - V_i| = O(\varepsilon) \text{ and } \|\varphi_a\|_a = O\left(\varepsilon^{\frac{N}{2}+2s+1}\right).$$

On the other hand, since $H(x_{a,i}) = 0$,

$$\int_{B_\rho(x_{a,i})} H(x) U_{\varepsilon, x_{a,i}}^2(x) = \frac{1}{2N} \varepsilon^{N+2} \Delta H(x_{a,i}) \int_{\mathbb{R}^N} |x|^2 U^2 + O(\varepsilon^{N+4})$$

which, combined with (4.6), gives that

$$\Delta H(x_{a,i}) = O(\varepsilon^{2s}).$$

Hence from the condition (V), we get (1.6) concluding Theorem 1.3. \square

4.2. Existence of the normalized peak solutions.

For the existence result of problem (1.7) with large $\lambda > 0$, we set $\eta = \lambda^{-\frac{1}{2s}}$ and $w(x) \mapsto \lambda^{\frac{1}{p-1}} w(x)$, and (1.7) turns to

$$\begin{cases} \eta^{2s} (-\Delta)^s w + (1 + \eta^{2s} V(x)) w = w^p, & x \in \mathbb{R}^N, \\ w \in H^s(\mathbb{R}^N), \end{cases} \quad (4.7)$$

which is equivalent to

$$\begin{cases} \operatorname{div}(t^{1-2s}\nabla\tilde{w}) = 0, & x \in \mathbb{R}_+^{N+1}, \\ -\eta^{2s}\lim_{t \rightarrow 0} t^{1-2s}\partial_t\tilde{w}(y, t) = w^p - (1 + \eta^{2s}V(x))w, & x \in \mathbb{R}^N. \end{cases}$$

We denote $\langle u, v \rangle_\eta = \int_{\mathbb{R}^N} (\eta^{2s}(-\Delta)^{\frac{s}{2}}u(-\Delta)^{\frac{s}{2}}v + uv)$, and $\|u\|_\eta = \langle u, u \rangle_\eta^{\frac{1}{2}}$. In the following, for $\eta > 0$ small, we construct a k -peak solution u_η to (4.7) concentrating at b_1, \dots, b_k .

Proposition 4.2. *There exists an $\eta_0 > 0$, such that for any $\eta \in (0, \eta_0]$, and $d(z_i, \Gamma_i) = O(\eta)$, there exist $\omega_{\eta, z}$ with $z = (z_1, \dots, z_k)$, such that*

$$\int_{\mathbb{R}^N} (\eta^{2s}(-\Delta)^{\frac{s}{2}}w_\eta(-\Delta)^{\frac{s}{2}}\psi + (1 + \eta^{2s}V(x))w_\eta\psi) = \int_{\mathbb{R}^N} w_\eta^p\psi, \quad \text{for all } \psi \in F_{\eta, z},$$

where $w_\eta(x) = \sum_{i=1}^k U_{\eta, z} + \omega_{\eta, z}(x)$, and

$$F_{\eta, z} = \left\{ \psi(x) \in H^s(\mathbb{R}^N) : \langle v, \frac{\partial U_{\eta, z}}{\partial x_j} \rangle_\eta = 0, j = 1, \dots, N, i = 1, \dots, k \right\}.$$

Moreover, for $N \geq \max\{2 - 2s, 2s\}$

$$\begin{aligned} \|\omega_{\eta, z}\|_\eta = O & \left(\sum_{i=1}^k |V(z_i) - V_i| \eta^{\frac{N}{2}+2s} + \sum_{i=1}^k |\nabla V(z_i)| \eta^{\frac{N}{2}+2s+1} + \eta^{\frac{N}{2}+2s+2} \right) \\ & + \begin{cases} O(\eta^{\frac{N}{2}+\frac{p}{2}(N+2s)}), & \text{if } 1 < p \leq 2 \\ O(\eta^{\frac{N}{2}+(N+2s)}), & \text{if } p > 2 \end{cases} = O(\eta^{\frac{N}{2}+2s+\gamma}) \end{aligned} \quad (4.8)$$

with $\gamma = \max\{0, 1 - 2s\}$.

Remark 4.3. *The proof of the existence result is standard, and we omit the details. The last inequality in (4.8) holds because that $N \geq \max\{2 - 2s, 2s\}$ implies $\frac{p}{2}(N + 2s) > 2s + \gamma$.*

It is standard to know that, to obtain a true solution of (4.7), we need to choose z such that

$$\begin{aligned} & -\eta^{2s} \int_{\partial'' B_\rho^+(x_{a,i})} t^{1-2s} \frac{\partial \tilde{\omega}_{\eta, z}}{\partial \nu} \frac{\partial \tilde{\omega}_{\eta, z}}{\partial y^j} + \frac{1}{2} \eta^{2s} \int_{\partial'' B_\rho^+(x_{a,i})} t^{1-2s} |\nabla \tilde{\omega}_{\eta, z}|^2 \nu_j \\ & = - \int_{\partial B_\rho(x_{a,i})} (1 + \eta^{2s}V(x)) \omega_{\eta, z} \frac{\partial \omega_{\eta, z}}{\partial y^j} + \int_{\partial B_\rho(x_{a,i})} \omega_{\eta, z}^p \frac{\partial \omega_{\eta, z}}{\partial y^j}, \end{aligned}$$

which, similar to the proof of (4.5), is equivalent to

$$\int_{B_\rho(x_{a,i})} \frac{\partial V(x)}{\partial x_j} w_{\eta, z}^2 = O(\eta^{2N}).$$

We consider $z = (z_1, \dots, z_k)$, z_i close to b_i , $z_i \in \Gamma_{t,i}$, for some t close to V_i . Denote ν_i as the unit normal vector of $\Gamma_{t,i}$ at z_i , while $\tau_{i,j}$ as the principal directions of $\Gamma_{t,i}$, with $i = 1, \dots, k$, $j = 1, \dots, N - 1$. Then, at z_i , there holds that

$$D_{\tau_{i,j}} V(z_i) = 0, \quad \text{for } j = 1, \dots, N - 1, \quad \text{and } |\nabla V(z_i)| = |D_{\nu_i} V(z_i)|.$$

Moreover, we have the following results.

Lemma 4.4. *Under the assumption (V), then*

$$\int_{B_\rho(x_{a,i})} D_{\nu_i} V(x) w_{\eta,z}^2 = O(\eta^{2N})$$

is equivalent to

$$D_{\nu_i} V(z_i) = O(\eta^{2s+\gamma}).$$

Proof. On the one hand,

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} D_{\nu_i} V(x) U_{\eta,z_i}^2(x) \\ &= -2 \int_{B_\rho(x_{a,i})} D_{\nu_i} V(x) U_{\eta,z_i}(x) \omega_{\eta,z} - \int_{B_\rho(x_{a,i})} D_{\nu_i} V(x) \omega_{\eta,z}^2 + O(\eta^{2N}) \\ &= O(\eta^{N+2s+\gamma}). \end{aligned} \tag{4.9}$$

On the other hand,

$$\int_{B_\rho(x_{a,i})} D_{\nu_i} V(x) U_{\eta,z_i}^2(x) = a_* \eta^N D_{\nu_i} V(z_i) + O(\eta^{N+2})$$

which, combined with (4.9), gives that

$$D_{\nu_i} V(z_i) = O(\eta^{2s+\gamma}).$$

□

Lemma 4.5. *Under the assumption (V), then*

$$\int_{B_\rho(x_{a,i})} D_{\tau_i} V(x) w_{\eta,z}^2 = O(\eta^{2N})$$

is equivalent to

$$(D_{\tau_i} \Delta) V(z_i) = O\left(\sum_{i=1}^k |V(z_i) - V_i| \eta^{2s-1} + \eta^{2s}\right).$$

Proof. Let $H(x) = \langle \nabla V(x), \tau_i \rangle$. On the one hand,

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} H(x) U_{\eta,z_i}^2(x) \\ &= -2 \int_{B_\rho(x_{a,i})} H(x) U_{\eta,z_i}(x) \omega_{\eta,z} - \int_{B_\rho(x_{a,i})} H(x) \omega_{\eta,z}^2 + O(\eta^{2N}) \\ &= O\left(\left(\eta^{\frac{N}{2}+1} |\nabla H(z_i)| + \eta^{\frac{N}{2}+2}\right) \|\omega_{\eta,z}\|_\eta + |\nabla H(z_i)| \|\omega_{\eta,z}\|_\eta^2\right) + O(\eta^{2N}). \end{aligned} \tag{4.10}$$

On the other hand, since $H(x_{a,i}) = 0$, we set

$$B = \frac{1}{N} \int_{\mathbb{R}^N} |x|^2 U^2, \tag{4.11}$$

and then

$$\int_{B_\rho(x_{a,i})} H(x) U_{\eta,z_i}^2(x) = \frac{B}{2} \eta^{N+2} \Delta H(z_i) + O(\eta^{N+4})$$

which, combined with (4.10), gives the result.

□

Theorem 4.6. *For $\lambda > 0$ large, the problem (1.7) has a solution w_λ of the form*

$$w_\lambda(x) = \sum_{i=1}^k \lambda^{\frac{1}{p-1}} U\left(\lambda^{\frac{1}{2s}}(x - x_{\lambda,i})\right) + \omega_\lambda,$$

where $x_{\lambda,i} \rightarrow b_i$ and $\int_{\mathbb{R}^N} (|(-\Delta)^{\frac{s}{2}} \omega_\lambda|^2 + \omega_\lambda^2) \rightarrow 0$ as $\lambda \rightarrow +\infty$.

Proof. We are sufficed to solve (4.7). From Lemma 4.4 and Lemma 4.5, we know that solving (4.7) is equivalent to solving $d(z_i, \Gamma_i) = O(\eta)$,

$$D_{\nu_i} V(z_i) = O(\eta^{2s+\gamma}), \quad (D_{\tau_i} \Delta) V(z_i) = O\left(\sum_{i=1}^k |V(z_i) - V_i| \eta^{2s-1} + \eta^{2s}\right).$$

Now we take $\bar{z}_i \in \Gamma_i$ to be the point such that $z_i - \bar{z}_i = \alpha_i \nu_i$ for some $\alpha_i \in \mathbb{R}$. Then, we have $D_{\nu_i} V(\bar{z}_i) = 0$. Then,

$$D_{\nu_i} V(z_i) = D_{\nu_i} V(z_i) - D_{\nu_i} V(\bar{z}_i) = D_{\nu_i \nu_i}^2 V(\bar{z}_i) \langle z_i - \bar{z}_i, \nu_i \rangle + O(|z_i - \bar{z}_i|^2).$$

By the non-degenerate assumption, it holds that

$$D_{\nu_i} V(z_i) = O(\eta^{2s+\gamma}) \Leftrightarrow \langle z_i - \bar{z}_i, \nu_i \rangle = O(\eta^{2s+\gamma} + |z_i - \bar{z}_i|^2),$$

which means that

$$|z_i - \bar{z}_i| = O(\eta^{2s+\gamma}) \Rightarrow d(z_i, \Gamma_i) = |z_i - \bar{z}_i| = O(\eta). \quad (4.12)$$

Let $\bar{\tau}_{i,j}$ be the j -th tangential unit vector of Γ_i at \bar{z}_i . By the assumption (V), we have

$$(D_{\tau_{i,j}} \Delta) V(z_i) = (D_{\bar{\tau}_{i,j}} \Delta) V(\bar{z}_i) + O(|z_i - \bar{z}_i|) = (D_{\bar{\tau}_{i,j}} \Delta) V(\bar{z}_i) + O(\eta^{2s+\gamma}),$$

and

$$\begin{aligned} (D_{\bar{\tau}_{i,j}} \Delta) V(\bar{z}_i) &= (D_{\bar{\tau}_{i,j}} \Delta) V(\bar{z}_i) - (D_{\tau_{i,j,0}} \Delta) V(b_i) \\ &= \langle (\nabla_T D_{\tau_{i,j,0}} \Delta V)(b_i), \bar{z}_i - b_i \rangle + O(|\bar{z}_i - b_i|^2), \end{aligned}$$

where ∇_T is the tangential gradient on Γ_i at $b_i \in \Gamma_i$, and $\tau_{i,j,0}$ is the j -th tangential unit vector of Γ_i at b_i . Therefore, we have

$$(D_{\tau_{i,j}} \Delta) V(z_i) = O\left(\sum_{i=1}^k |V(z_i) - V_i| \eta^{2s-1} + \eta^{2s}\right),$$

which implies that

$$\langle (\nabla_T D_{\tau_{i,j,0}} \Delta V)(b_i), \bar{z}_i - b_i \rangle = O\left(\sum_{i=1}^k |V(z_i) - V_i| \eta^{2s-1} + \eta^{2s}\right) = O(\eta^{2s}). \quad (4.13)$$

Hence, we can solve (4.12) and (4.13) to obtain $z_i = x_{\eta,i}$ with $x_{\eta,i} \rightarrow b_i$. □

Now we are in a position to prove Theorem 1.4.

Proof of Theorem 1.4. Let w_λ be a peak solution obtained in Theorem 4.6. Define $u_\lambda = w_\lambda / \left(\int_{\mathbb{R}^N} w_\lambda^2 \right)^{\frac{1}{2}}$. Then $\int_{\mathbb{R}^N} u_\lambda^2 = 1$ and

$$(-\Delta)^s u_\lambda + V(x)u_\lambda = a_\lambda u_\lambda^p - \lambda u_\lambda, \quad a_\lambda = \int_{\mathbb{R}^N} w_\lambda^2. \quad (4.14)$$

Similar to the proof of Proposition 3.4, there holds that

$$\lambda^{\frac{N}{2s} - \frac{2}{p-1}} \left(\int_{\mathbb{R}^N} w_\lambda^2 \right)^{\frac{2}{p-1}} = ka_* + o(1), \quad \text{as } a \rightarrow a_0$$

with $a_0 = 0$ if $\frac{N}{2s} - \frac{2}{p-1} > 0$, $a_0 = (ka_*)^{\frac{p-1}{2}}$ if $\frac{N}{2s} - \frac{2}{p-1} = 0$, while $a_0 = +\infty$ if $\frac{N}{2s} - \frac{2}{p-1} < 0$.

Take $\lambda_0 > 0$ large and $b_0 = \int_{\mathbb{R}^N} w_{\lambda_0}^2$. Then, if $\frac{N}{2s} - \frac{2}{p-1} > 0$, for any $a \in (0, b_0)$, by the mean value theorem, there exists some large λ_a , such that the solution u_a to (1.1) with $\lambda = \lambda_a$ satisfies $\int_{\mathbb{R}^N} w_\lambda^2 = a$, and for such a , we obtain a k -peak solution for (1.7), with $\mu_a = -\lambda_a$.

Similarly, if $\frac{N}{2s} - \frac{2}{p-1} = 0$, for any a between b_0 and $(ka_*)^{\frac{p-1}{2}}$, while if $\frac{N}{2s} - \frac{2}{p-1} < 0$, for any $a > b_0$ large enough, there exists λ_a , such that the solution u_a to (1.1) with $\lambda = \lambda_a$ satisfies $\int_{\mathbb{R}^N} w_\lambda^2 = a$, and for such a , we obtain a k -peak solution for (1.7), with $\mu_a = -\lambda_a$. □

4.3. Clustering peak solutions.

In the end of this section, we are concerned with the clustering multi-peak solutions to problem (1.1)–(1.2), and sketch the proof of Theorem 1.5.

The function $\Delta V(x)|_{x \in \Gamma_i}$ has a minimum point and a maximum point. Let us assume that the function $\Delta V(x)|_{x \in \Gamma_{i_0}}$ has isolated maximum point $b \in \Gamma_{i_0}$, that is $\Delta V(x) < \Delta V(b)$ for all $x \in \Gamma_{i_0} \cap (B_\delta(b) \setminus b)$. Let $x_{\eta,j} \rightarrow b, j = 1, \dots, k, \frac{|x_{\eta,i} - x_{\eta,j}|}{\eta} \rightarrow +\infty, i \neq j$ as $\eta \rightarrow 0$. We take $\sum_{j=1}^k U_{\eta, x_{\eta,j}}$ as the approximate solution of (4.7) and sufficed to show the following result.

Proposition 4.7. *Assume (V) and $\frac{\partial^2 V(x)}{\partial \nu_{i_0}^2} \neq 0$ for any $x \in \Gamma_{i_0}$ with some $i_0 \in \{1, \dots, k\}$. If $b \in \Gamma_{i_0}$ is an isolated maximum point of $\Delta V(x)|_{x \in \Gamma_{i_0}}$ on Γ_{i_0} , then for any integer $k > 0$, there exists an $\eta_0 > 0$, such that for any $\eta \in (0, \eta_0]$, problem (4.7) has a solution w_η of the form*

$$w_\eta(x) = \sum_{j=1}^k U_{\eta, x_{\eta,j}} + \omega_\eta,$$

where $x_{\eta,j} \rightarrow b, j = 1, \dots, k, \frac{|x_{\eta,i} - x_{\eta,j}|}{\eta} \rightarrow +\infty, i \neq j$, and $\int_{\mathbb{R}^N} (\eta^{2s} |(-\Delta)^{\frac{s}{2}} \omega_\eta|^2 + \omega_\eta^2) = o(\eta^{\frac{N}{2}})$ as $\eta \rightarrow 0$.

Proof. Define the energy functional

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^N} \left(\eta^{2s} |(-\Delta)^{\frac{s}{2}} u|^2 + (1 + \eta^{2s} V(x)) u^2 \right) - \frac{1}{p+1} \int_{\mathbb{R}^N} u^{p+1}.$$

We obtain the energy expansion:

$$\begin{aligned}
 & I\left(\sum_{j=1}^k U_{\eta, x_{\eta, j}}\right) \\
 &= (1 + \eta^{2s} V_{i_0})^{-\frac{N}{2s}} \left(k E_1 \eta^N + E_2 \eta^{N+2s} \sum_{j=1}^k \frac{\partial^2 V(\bar{x}_{\eta, j})}{\partial \nu_{i, j}^2} r_{\eta, j}^2 + E_3 \eta^{N+2s+2} \sum_{j=1}^k \Delta V(\bar{x}_{\eta, j}) \right. \\
 & \quad - \sum_{j>m} (c_0 + o(1)) \eta^N \left(\frac{\eta}{|x_{\eta, m} - x_{\eta, j}|} \right)^{N+2s} \\
 & \quad \left. + O\left(\eta^{N+2s+3} + \eta^N \sum_{j>m} \left(\frac{\eta}{|x_{\eta, m} - x_{\eta, j}|} \right)^{N+2s+\tau} + \eta^{N+2s} r_{\eta, j}^3 \right) \right), \tag{4.15}
 \end{aligned}$$

where $E_l, c_0 > 0, l = 2, 3$,

$$E_1 = \left(\frac{1}{2} - \frac{1}{p+1} \right) \int_{\mathbb{R}^N} U^{p+1} > 0 \text{ and } r_{\eta, j} = |x_{\eta, j} - \bar{x}_{\eta, j}|,$$

with $\bar{x}_{\eta, j} \in \Gamma_{i_0}$ the point such that $|x_{\eta, j} - \bar{x}_{\eta, j}| = d(x_{\eta, j}, \Gamma_{i_0})$.

In fact, it is easy to get

$$\begin{aligned}
 & I\left(\sum_{j=1}^k U_{\eta, x_{\eta, j}}\right) \\
 &= (1 + \eta^{2s} V_{i_0})^{-\frac{N}{2s}} \left(k \left(\frac{1}{2} - \frac{1}{p+1} \right) \int_{\mathbb{R}^N} U^{p+1} + E'_2 \eta^{N+2s} \sum_{j=1}^k (V(x_{\eta, j}) - V(\bar{x}_{\eta, j})) \right. \\
 & \quad + E'_3 \eta^{N+2s+2} \sum_{j=1}^k \Delta V(x_{\eta, j}) - \sum_{j>m} (c_0 + o(1)) \eta^N \left(\frac{\eta}{|x_{\eta, m} - x_{\eta, j}|} \right)^{N+2s} \\
 & \quad \left. + O\left(\eta^{N+2s+3} + \eta^N \sum_{j>m} \left(\frac{\eta}{|x_{\eta, m} - x_{\eta, j}|} \right)^{N+2s+\tau} \right) \right), \tag{4.16}
 \end{aligned}$$

and

$$V(x_{\eta, j}) = 1 + \frac{\partial^2 V(\bar{x}_{\eta, j})}{\partial \nu_{i, j}^2} r_{\eta, j}^2 + O(r_{\eta, j}^3) \text{ and } \Delta V(x_{\eta, j}) = \Delta V(\bar{x}_{\eta, j}) + O(r_{\eta, j}). \tag{4.17}$$

So, (4.15) follows from (4.16) and (4.17).

To obtain a solution w_η of the form $\sum_{j=1}^k U_{\eta, x_{\eta, j}} + \omega_\eta$, we can first carry out the reduction argument as in Proposition 4.2 to obtain ω_η , satisfying

$$\|\omega_\eta\|_\eta = O\left(\eta^{\frac{N}{2}+2s+1} + \eta^{\frac{N}{2}} \sum_{i \neq j} \left(\frac{\eta}{|x_{\eta, i} - x_{\eta, j}|} \right)^{\min\{\frac{p}{2}, 1\}(N+2s)} \right). \tag{4.18}$$

Define

$$K(x_{\eta, 1}, \dots, x_{\eta, k}) = I(U_{\eta, x_{\eta, j}} + \omega_\eta).$$

From (4.18), we get the same expansion (4.15) for $K(x_{\eta,1}, \dots, x_{\eta,k})$. Now we set

$$\mathcal{B} = \{(r, \bar{x}) : r \in (-\delta\eta^{\frac{1}{2}}, \delta\eta^{\frac{1}{2}}), \bar{x} \in B_\delta(b_i) \cap \Gamma_i\}.$$

If Γ_{i_0} is a local maximum set of $V(x)$ and $\frac{\partial^2 V(\bar{x})}{\partial \nu_{i_0}^2} < 0$ for any $\bar{x} \in \Gamma_{i_0}$, then it is easy to prove that $K(x_{\eta,1}, \dots, x_{\eta,k})$ has a critical point, which is a maximum point of K in

$$S_{\eta,k} := \{(x_{\eta,1}, \dots, x_{\eta,k}) : x_{\eta,j} \in \mathcal{B}, \frac{|x_{\eta,j} - x_{\eta,m}|}{\eta} > \eta^{\sigma-1}, m \neq j\},$$

where $\sigma \in (\frac{1}{2}, \frac{N-1}{N+2s})$ is some constant such that both $\sigma > \frac{1}{2}$ and $(1-\sigma)(N+2s) > 2s+1$ hold.

If Γ_{i_0} is a local minimum set of $V(x)$ and $\frac{\partial^2 V(\bar{x})}{\partial \nu_{i_0}^2} > 0$ for any $\bar{x} \in \Gamma_{i_0}$, then

$$E_2 \frac{\partial^2 V(\bar{x})}{\partial \nu_{i_0}^2} r^2 + E_3 \eta^2 \Delta V(\bar{x})$$

has a saddle point $(0, b)$ in \mathcal{B} . We can use a topological argument as in [13, 14] to prove that $K(x_{\eta,1}, \dots, x_{\eta,k})$ has a critical point in $S_{\eta,k}$. \square

5. LOCAL UNIQUENESS

5.1. Precise estimates by Pohozaev identities.

From Lemma 3.2, we know that a k -peak solution v_a to equation (3.1) can be written as

$$v_a(x) = \sum_{i=1}^k U_{\varepsilon, x_{a,i}}(x) + \varphi_a(x),$$

where $|x_{a,i} - b_i| = o(1)$, $d(x_{a,i}, \Gamma_i) = O(\varepsilon)$, $\varepsilon = (-\mu_a)^{-\frac{1}{2s}}$,

$$U_{\varepsilon, x_{a,i}}(x) = (1 + \varepsilon^{2s} V_i)^{\frac{1}{p-1}} U\left(\frac{(1 + \varepsilon^{2s} V_i)^{\frac{1}{2s}}(x - x_{a,i})}{\varepsilon}\right),$$

$\varphi_a \in \bigcap_{i=1}^k E_{a, x_{a,i}}$. For $N \geq 2s+2$, it holds that $\frac{p}{2}(N+2s) > 2s+1$, which implies then

$$\|\varphi_a\|_a = \|\varphi_a\|_\varepsilon = O\left(\left|\sum_{i=1}^k (V(x_{a,i}) - V_i)\right| \varepsilon^{\frac{N}{2}+2s} + \left|\sum_{i=1}^k \nabla V(x_{a,i})\right| \varepsilon^{\frac{N}{2}+2s+1} + \varepsilon^{\frac{N}{2}+2s+1}\right). \quad (5.1)$$

Moreover, $x_{a,i} \in \Gamma_{t_a, i}$ with some $t_a \rightarrow V_i$. Denote $\nu_{a,i}$ and $\tau_{a,i,j}$ as the unit normal vector and the principal direction of $\Gamma_{t_a, i}$ at $x_{a,i}$ respectively. Then at $x_{a,i}$, there hold that

$$D_{\tau_{a,i,j}} V(x_{a,i}) = 0, \quad |\nabla V(x_{a,i})| = |D_{\nu_{a,i}} V(x_{a,i})|.$$

Lemma 5.1. *Under the assumption (V), it holds that*

$$D_{\nu_{a,i}} V(x_{a,i}) = O\left(\varepsilon^{\min\{2, 2s+1\}}\right).$$

Proof. Using (4.5), we obtain that

$$\int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) v_a^2 = O(\varepsilon^{2N}).$$

On the one hand,

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) U_{\varepsilon, x_{a,i}}^2(x) \\ &= -2 \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) U_{\varepsilon, x_{a,i}}(x) \varphi_a - \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) \varphi_a^2 + O(\varepsilon^{2N}) \\ &= O\left(\varepsilon^{\frac{N}{2}} |D_{\nu_{a,i}} V(x_{a,i})| \|\varphi_a\|_a + \varepsilon^{\frac{N}{2}+1} \|\varphi_a\|_a + \|\varphi_a\|_a^2\right) + O(\varepsilon^{2N}) \\ &= O\left(\sum_{i=1}^k |V(x_{a,i}) - V_i| \varepsilon^{N+2s} + \left| \sum_{i=1}^k \nabla V(x_{a,i}) \right| \varepsilon^{N+2s+1} + \varepsilon^{N+2s+2}\right) = O(\varepsilon^{N+2s+1}). \end{aligned} \quad (5.2)$$

On the other hand, using the definition (4.11)

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) U_{\varepsilon, x_{a,i}}^2(x) \\ &= a_* \varepsilon^N (1 + \varepsilon^{2s} V_i)^{\frac{2}{p-1}} D_{\nu_{a,i}} V(x_{a,i}) + \frac{B}{2} \varepsilon^{N+2} \Delta D_{\nu_{a,i}} V(x_{a,i}) + O(\varepsilon^{N+4}), \end{aligned} \quad (5.3)$$

which, combined with (5.2), gives that

$$D_{\nu_{a,i}} V(x_{a,i}) = O(\varepsilon^{\min\{2, 2s+1\}}).$$

□

Now we take $\bar{x}_{a,i} \in \Gamma_i$ be the point such that $x_{a,i} - \bar{x}_{a,i} = \alpha_{a,i} \nu_{a,i}$ for some $\alpha_{a,i} \in \mathbb{R}$.

Lemma 5.2. *Under the condition (V), we have that*

$$\begin{cases} \bar{x}_{a,i} - b_i = L_i \varepsilon^2 + O(\varepsilon^{2s+2}), \\ x_{a,i} - \bar{x}_{a,i} = -\frac{B}{2a_*} \Delta D_{\nu_{a,i}} V(b_i) \left(\frac{\partial^2 V(b_i)}{\partial \nu_i^2} \right)^{-1} \varepsilon^2 + O(\varepsilon^{2s+2}), \end{cases} \quad (5.4)$$

where B is as in (4.11), L_i is a vector depending on b_i and $i = 1, \dots, k$.

Proof. From (5.3), and

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) U_{\varepsilon, x_{a,i}}^2(x) \\ &= -2 \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) U_{\varepsilon, x_{a,i}}(x) \varphi_a - \int_{B_\rho(x_{a,i})} D_{\nu_{a,i}} V(x) \varphi_a^2 + O(\varepsilon^{2N}) \\ &= O\left(\varepsilon^{\frac{N}{2}} |D_{\nu_{a,i}} V(x_{a,i})| \cdot \|\varphi_a\|_a + \varepsilon^{\frac{N}{2}+1} \|\varphi_a\|_a + \|\varphi_a\|_a^2\right) + O(\varepsilon^{2N}) \\ &= O(\varepsilon^{N+2s+2}), \end{aligned}$$

we obtain that

$$\begin{aligned}
& (a_* + O(\varepsilon^{2s}))D_{\nu_{a,i}}V(x_{a,i}) + \frac{B}{2}\varepsilon^2\Delta D_{\nu_{a,i}}V(x_{a,i}) \\
&= O\left(\varepsilon^{2s+2} + \varepsilon^{2s+1}\sum_{i=1}^k|V(x_{a,i}) - V_i|\right) \\
&= O\left(\varepsilon^{2s+2} + \varepsilon^{2s+1}\sum_{i=1}^k|x_{a,i} - \bar{x}_{a,i}|^2\right).
\end{aligned}$$

Since $\frac{\partial^2 V(b_i)}{\partial \nu_i^2} \neq 0$, the outward vector $\nu_{a,i}(x)$ and the tangential unit vector $\tau_{a,i}(x)$ of $\Gamma_{a,i}$ at $x_{a,i}$ are Lip-continuous in $W_{\delta,i}$, we obtain from the above that

$$x_{a,i} - \bar{x}_{a,i} = -\frac{B}{2a_*}\Delta D_{\nu_{a,i}}V(b_i)\left(\frac{\partial^2 V(b_i)}{\partial \nu_i^2}\right)^{-1}\varepsilon^2 + O\left(\varepsilon^{2s+2} + \varepsilon^{2s+1}\sum_{i=1}^k|x_{a,i} - \bar{x}_{a,i}|^2\right). \quad (5.5)$$

From (5.1) and (5.5), for $\frac{4s+4}{N+2s} < p < \frac{N+2s}{N-2s}$ (which can be satisfied by $N \geq 2s + 4$),

$$\|\varphi_a\|_a = O\left(\sum_{i=1}^k|x_{a,i} - \bar{x}_{a,i}|\varepsilon^{\frac{N}{2}+2s} + \varepsilon^{\frac{N}{2}+4s+2} + \varepsilon^{\frac{N}{2}+2s+2}\right) = O(\varepsilon^{\frac{N}{2}+2s+2}). \quad (5.6)$$

Let $H(x) = \langle \nabla V(x), \tau_{a,i} \rangle$. Then for $N \geq 2s + 4$,

$$\begin{aligned}
& \int_{B_\rho(x_{a,i})} H(x)U_{\varepsilon,x_{a,i}}^2(x) \\
&= -2 \int_{B_\rho(x_{a,i})} H(x)U_{\varepsilon,x_{a,i}}(x)\varphi_a - \int_{B_\rho(x_{a,i})} H(x)\varphi_a^2 \\
&= -2 \int_{B_\rho(x_{a,i})} H(x)U_{\varepsilon,x_{a,i}}(x)\varphi_a + O(\|\varphi_a\|_a^2) + O(\varepsilon^{2N}) \\
&= -2 \int_{B_\rho(x_{a,i})} \langle \nabla H(x_{a,i}), x - x_{a,i} \rangle U_{\varepsilon,x_{a,i}}(x)\varphi_a + O(\varepsilon^{N+2s+4})
\end{aligned} \quad (5.7)$$

On the one hand, since $\nabla V = 0$, for $x \in \Gamma_i$,

$$\nabla H(x_{a,i}) = \langle \nabla^2 V(x_{a,i}), \tau_{a,i} \rangle = \langle \nabla^2 V(\bar{x}_{a,i}), \bar{\tau}_{a,i} \rangle + O(|x_{a,i} - \bar{x}_{a,i}|) = O(|x_{a,i} - \bar{x}_{a,i}|), \quad (5.8)$$

where $\bar{x}_{a,i} \in \Gamma_i$ is the point such that $x_{a,i} - \bar{x}_{a,i} = \beta_{a,i}\nu_{a,i}$ for some $\beta_{a,i} \in \mathbb{R}$ and $\bar{\tau}_{a,i}$ is the tangential vector of Γ_i at $\bar{x}_{a,i} \in \Gamma_i$. Hence,

$$\begin{aligned}
& \int_{B_\rho(x_{a,i})} \langle \nabla H(x_{a,i}), x - x_{a,i} \rangle U_{\varepsilon,x_{a,i}}(x)\varphi_a \\
&= O(\varepsilon^{\frac{N}{2}+1}|\nabla H(x_{a,i})| \cdot \|\varphi_a\|_a) = O(\varepsilon^{N+2s+3}|x_{a,i} - \bar{x}_{a,i}|) = O(\varepsilon^{N+2s+5}).
\end{aligned}$$

Then by (5.7) and (5.8), it follows

$$\int_{B_\rho(x_{a,i})} H(x)U_{\varepsilon,x_{a,i}}^2(x) = O(\varepsilon^{N+2s+4}). \quad (5.9)$$

On the other hand, applying the Taylor's expansion, we obtain that

$$\begin{aligned} \int_{B_\rho(x_{a,i})} H(x) U_{\varepsilon, x_{a,i}}^2(x) &= \frac{B \varepsilon^{N+2}}{2} (1 + V_i \varepsilon^{2s}) (D_{\tau_{a,i}} \Delta V)(x_{a,i}) \\ &+ \frac{H_{\tau_i} \varepsilon^{N+4}}{24} + O(\varepsilon^{N+2s+4}), \end{aligned} \quad (5.10)$$

where

$$H_{\tau_i} = \sum_{l=1}^2 \sum_{m=1}^2 \frac{\partial^4 H(b_i)}{\partial x_l^2 \partial x_m^2} \int_{\mathbb{R}^N} x_l^2 x_m^2 U^2.$$

Combining (5.9) and (5.10), we obtain that

$$(D_{\tau_{a,i}} \Delta V)(x_{a,i}) = -\frac{H_{\tau_i} \varepsilon^2}{12B} + O(\varepsilon^{2s+2}). \quad (5.11)$$

By (5.5), we get

$$\begin{aligned} (D_{\tau_{a,i}} \Delta) V(x_{a,i}) &= (D_{\bar{\tau}_{a,i}} \Delta) V(\bar{x}_{a,i}) + \langle A_{\tau_i}, x_{a,i} - \bar{x}_{a,i} \rangle + O(|x_{a,i} - \bar{x}_{a,i}|^2) \\ &= (D_{\bar{\tau}_{a,i}} \Delta) V(\bar{x}_{a,i}) + B_{\tau_i} \varepsilon^2 + O(\varepsilon^{2s+2}), \end{aligned} \quad (5.12)$$

where A_{τ_i} is a vector depending on b_i and B_i is a constant depending on b_i . Moreover,

$$(D_{\bar{\tau}_{a,i}} \Delta) V(\bar{x}_{a,i}) = ((D_{\tau_{a,i}}^2 \Delta) V(b_i))(\bar{x}_{a,i} - b_i) + O(|\bar{x}_{a,i} - b_i|^2). \quad (5.13)$$

Therefore, by (5.11), (5.12) and (5.13), there hold that

$$((D_{\tau_{a,i}}^2 \Delta) V(b_i))(x_{a,i}) (\bar{x}_{a,i} - b_i) = -\left(\frac{H_{\tau_i}}{12B} + B_{\tau_{a,i}}\right) \varepsilon^2 + O(\varepsilon^{2s+2}) + O(|\bar{x}_{a,i} - b_i|^2).$$

Since $(D_{\tau_{a,i}}^2 \Delta) V(b_i)$ is non-singular, we conclude the proof. \square

5.2. Local uniqueness.

We assume $\frac{4s+4}{N+2s} < p < \frac{N+2s}{N-2s}$, which can be satisfied by $N \geq 2s + 4$. Set

$$\delta_a = \begin{cases} \left(\frac{a^{\frac{2}{p-1}}}{ka_*}\right)^{\frac{1}{N-\frac{4s}{p-1}}}, & p-1 \neq \frac{4s}{N}, \\ \left(\frac{2s}{(s+2)B_1} |ka_* - a^{\frac{2}{p-1}}|\right)^{\frac{1}{2(s+1)}}, & p-1 = \frac{4s}{N}, \end{cases}$$

where $B_1 = \frac{1}{N} \sum_{i=1}^k \Delta V(b_i) \int_{\mathbb{R}^N} |x|^2 U^2(x)$.

Proposition 5.3. *Under the assumption (V), there hold that*

$$-\mu_a \delta_a^{2s} = 1 + \gamma_1 \delta_a^{2s} + O(\delta_a^{2s+2}), \quad (5.14)$$

and

$$x_{a,i} - b_i = \bar{L}_i \delta_a^2 + O(\delta_a^{2s+2}), \quad i = 1, \dots, k, \quad (5.15)$$

where γ_1 and the vector \bar{L}_i are constants.

Proof. (a) Firstly, if $p-1 \neq \frac{4s}{N}$, we follow the proof of Proposition 3.4 to find

$$\begin{aligned} \frac{a^{\frac{2}{p-1}}}{(-\mu_a)^{\frac{2}{p-1}-\frac{N}{2s}}} &= \sum_{i=1}^k \left(1 + \frac{V_i}{-\mu_a}\right)^{\frac{2}{p-1}-\frac{N}{2s}} \int_{\mathbb{R}^N} U^2 + O\left(\left|\sum_{i=1}^k (V(x_{a,i}) - V_i)\right|(-\mu_a)^{-1}\right. \\ &\quad \left.+ \left|\sum_{i=1}^k \nabla V(x_{a,i})\right|(-\mu_a)^{-1-\frac{1}{2s}} + (-\mu_a)^{-1-\frac{1}{s}}\right) + \begin{cases} O((-\mu_a)^{-\frac{pN}{4s}-\frac{p}{2}}), & \text{if } 1 < p \leq 2, \\ O((-\mu_a)^{-\frac{N}{2s}-1}), & \text{if } p > 2, \end{cases} \\ &\quad + O\left((-\mu_a)^{-\frac{N}{2s}-1} |x_{a,j} - x_{a,i}|^{-(N+2s)}\right), \end{aligned}$$

which implies that

$$\begin{aligned} \frac{a^{\frac{2}{p-1}}}{ka_*(-\mu_a)^{\frac{2}{p-1}-\frac{N}{2s}}} &= 1 + \sum_{i=1}^k \left(\frac{2}{p-1} - \frac{N}{2s}\right) \frac{V_i}{-\mu_a} + O\left(\left|\sum_{i=1}^k (V(x_{a,i}) - V_i)\right|(-\mu_a)^{-1}\right. \\ &\quad \left.+ \left|\sum_{i=1}^k \nabla V(x_{a,i})\right|(-\mu_a)^{-1-\frac{1}{2s}} + (-\mu_a)^{-1-\frac{1}{s}}\right) + \begin{cases} O((-\mu_a)^{-\frac{pN}{4s}-\frac{p}{2}}), & \text{if } 1 < p \leq 2, \\ O((-\mu_a)^{-\frac{N}{2s}-1}), & \text{if } p > 2. \end{cases} \\ &\quad + O\left((-\mu_a)^{-\frac{N}{2s}-1} |x_{a,j} - x_{a,i}|^{-(N+2s)}\right). \end{aligned}$$

Set $\delta_a = \left(\frac{a^{\frac{2}{p-1}}}{ka_*}\right)^{\frac{1}{N-\frac{4s}{p-1}}}$. Then we obtain (5.14). Finally, (5.15) can be obtained by (5.14) and Lemma 5.2.

(b) Secondly, we turn to deal with the case of $p-1 = \frac{4s}{N}$. From (5.4), it holds that

$$x_{a,i} - b_i = \bar{L}_i(-\mu_a)^{-\frac{1}{s}} + O((-\mu_a)^{-\frac{1}{s}-1}),$$

with some vector \bar{L}_i . Moreover, in view of (5.6), $\|\varphi_a\|_a = O(\varepsilon^{\frac{N}{2}+2s+2})$.

We know that u_a is the k -peak solutions of (1.1)-(1.2), and we have the following Pohozaev identity:

$$\begin{aligned} & - \int_{\partial'' B_\rho(x_{a,i})} t^{1-2s} \frac{\partial \tilde{u}_a}{\partial \nu} \langle Y - X_{a,i}, \nabla \tilde{u}_a \rangle + \frac{1}{2} \int_{\partial'' B_\rho(x_{a,i})} t^{1-2s} |\nabla \tilde{u}_a|^2 \langle Y - X_{a,i}, \nu \rangle \\ & - \frac{N-2s}{2} \int_{\partial'' B_\rho(x_{a,i})} t^{1-2s} \frac{\partial \tilde{u}_a}{\partial \nu} \tilde{u}_a \\ & - \int_{\partial B_\rho(x_{a,i})} \left(\frac{(u_a)^{p+1}}{p+1} - \frac{-\mu_a + V(x)}{2} (u_a)^2 \right) \langle y - x_{a,i}, \nu \rangle \\ & = \int_{B_\rho(x_{a,i})} \left(\frac{N-2s}{2} - \frac{N}{p+1} \right) |u_a|^{p+1} + s(-\mu_a + V(x))(u_a)^2 \\ & \quad + \frac{1}{2} \langle y - x_{a,i}, \nabla V(x) \rangle (u_a)^2, \end{aligned}$$

which can be rewritten as

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} sV(x)(u_a)^2 + \frac{1}{2} \langle y - x_{a,i}, \nabla V(x) \rangle (u_a)^2 \\ & = \int_{B_\rho(x_{a,i})} s\mu_a(u_a)^2 + a \left(\frac{N}{p+1} - \frac{N-2s}{2} \right) |u_a|^{p+1} + \int_{\partial'' B_\rho(x_{a,i})} W_1 + \int_{\partial B_\rho(x_{a,i})} W_1, \end{aligned} \tag{5.16}$$

where

$$W_1 = -t^{1-2s} \frac{\partial \tilde{u}_a}{\partial \nu} \langle Y - X_{a,i}, \nabla \tilde{u}_a \rangle + \frac{1}{2} t^{1-2s} |\nabla \tilde{u}_a|^2 \langle Y - X_{a,i}, \nu \rangle - \frac{N-2s}{2} t^{1-2s} \frac{\partial \tilde{u}_a}{\partial \nu} \tilde{u}_a,$$

$$W_2 = \left(-\frac{a(u_a)^{p+1}}{p+1} + \frac{(-\mu_a + V(x))}{2} (u_a)^2 \right) \langle y - x_{a,i}, \nu \rangle.$$

Recall that

$$u_a(x) = \sum_{i=1}^k \left(\frac{-\mu_a + V_i}{a} \right)^{\frac{1}{p-1}} \left(U \left((-\mu_a + V_i)^{\frac{1}{2s}} (x - x_{a,i}) \right) + \varphi_a \right).$$

Then,

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} sV(x)(u_a)^2 + \frac{1}{2} \langle y - x_{a,i}, \nabla V(x) \rangle (u_a)^2 \\ &= \left(\frac{-\mu_a + V_i}{a} \right)^{\frac{2}{p-1}} \int_{B_\rho(x_{a,i})} \left(s(V(x) - V_i) + \frac{1}{2} \langle y - x_{a,i}, \nabla V(x) \rangle \right) U^2 \left((-\mu_a + V_i)^{\frac{1}{2s}} (x - x_{a,i}) \right) \\ & \quad + sV_i \int_{B_\rho(x_{a,i})} u_a^2 + O \left((-\mu_a)^{-2-\frac{2}{s}} \right) \\ &= \frac{\frac{s}{2} + 1}{(-\mu_a + V_i)^{\frac{1}{s}}} \Delta V(x_{a,i}) \int_{\mathbb{R}^N} |x|^2 U^2(x) + sV_i \frac{a_*}{a^{\frac{2}{p-1}}} + o \left((-\mu_a + V_i)^{-\frac{1}{s}} \right). \end{aligned}$$

Moreover, we calculate more precisely that for some constants c_l , $l = 1, 2, 3$,

$$\Delta V(x_{a,i}) = \Delta V(b_i) + c_1 (-\mu_a)^{-\frac{1}{s}} + O \left((-\mu_a)^{-\frac{1}{s}-1} \right).$$

Hence,

$$\begin{aligned} & \int_{B_\rho(x_{a,i})} sV(x)(u_a)^2 + \frac{1}{2} \langle y - x_{a,i}, \nabla V(x) \rangle (u_a)^2 \\ &= \frac{\frac{s}{2} + 1}{(-\mu_a + V_i)^{\frac{1}{s}}} \Delta V(b_i) \int_{\mathbb{R}^N} |x|^2 U^2(x) + sV_i \frac{a_*}{a^{\frac{2}{p-1}}} + c_2 (-\mu_a)^{-\frac{1}{s}-1} + O \left((-\mu_a)^{-\frac{2}{s}-1} \right). \end{aligned}$$

Now summing (5.16) from $i = 1$ to $i = k$, since $\frac{N}{p+1} - \frac{N-2s}{2} = \frac{2s^2}{N+2s} > 0$, we have

$$\begin{aligned} & s\mu_a + \frac{2as^2}{N+2s} \int_{\mathbb{R}^N} |u_a|^{p+1} \\ &= \sum_{i=1}^k \frac{\frac{s}{2} + 1}{(-\mu_a)^{\frac{1}{s}}} \Delta V(b_i) \int_{\mathbb{R}^N} |x|^2 U^2(x) + \sum_{i=1}^k sV_i \frac{a_*}{a^{\frac{2}{p-1}}} + c_2 (-\mu_a)^{-\frac{1}{s}-1} + O \left((-\mu_a)^{-\frac{2}{s}-1} \right). \end{aligned} \tag{5.17}$$

On the other hand, by the orthogonality $\int_{\mathbb{R}^N} U^p((-\mu_a + V_i)^{\frac{1}{2s}}(x - x_{a,i}))\varphi_a = 0$ for $i = 1, \dots, k$, there holds that

$$\begin{aligned} \int_{\mathbb{R}^N} u_a^{p+1} &= \sum_{i=1}^k \left(\frac{-\mu_a + V_i}{a} \right)^{\frac{p+1}{p-1}} \left((-\mu_a + V_i)^{-\frac{N}{2s}} \int_{\mathbb{R}^N} U^{p+1} \right. \\ &\quad \left. + (p+1) \int_{\mathbb{R}^N} (U^p((-\mu_a + V_i)^{\frac{1}{2s}}(x - x_{a,i}))\varphi_a + O(|\varphi_a|^{p+1})) \right) \\ &= \frac{N+2s}{2s} \frac{a_*}{a^{\frac{p+1}{p-1}}} \sum_{i=1}^k (-\mu_a + V_i)^{\frac{p+1}{p-1} - \frac{N}{2s}} + O(\mu_a^{\frac{p+1}{p-1}} \|\varphi_a\|_a^2) \\ &= \frac{N+2s}{2s} \frac{a_*}{a^{\frac{p+1}{p-1}}} \sum_{i=1}^k (-\mu_a + V_i)^{\frac{p+1}{p-1} - \frac{N}{2s}} + O((-\mu_a)^{-1 - \frac{2}{s}}) \\ &= \frac{N+2s}{2s} \frac{a_*}{a^{\frac{p+1}{p-1}}} \sum_{i=1}^k (-\mu_a + V_i) + O((-\mu_a)^{-1 - \frac{2}{s}}), \end{aligned}$$

which, combined with (5.17), gives then

$$s(-\mu_a)^{1+\frac{1}{s}}(ka_* - a^{\frac{2}{p-1}}) = \left(\frac{s}{2} + 1\right) \sum_{i=1}^k \Delta V(b_i) \int_{\mathbb{R}^N} |x|^2 U^2(x) + c_3(-\mu_a)^{-1} + O((-\mu_a)^{-\frac{1}{s}-1}).$$

If we set $\delta = \left(\frac{2s}{(s+2)B_1} |ka_* - a^{\frac{2}{p-1}}| \right)^{\frac{1}{2(s+1)}}$, then we obtain (5.14), and (5.15) can be found by (5.14) and Lemma 5.2. \square

By a change of variable, the problem (1.1)-(1.2) can be rewritten as

$$\delta_a^{2s}(-\Delta)^s u + (-\mu_a \delta_a^{2s} + \delta_a^{2s} V(x))u = u^p, \quad u \in H^s(\mathbb{R}^N), \quad (5.18)$$

$$\int_{\mathbb{R}^N} u^2 = (a\delta_a^{2s})^{\frac{2}{p-1}}. \quad (5.19)$$

Similar to Lemma 3.2, the k -peak solution of (5.18)–(5.19) concentrating at b_1, \dots, b_k can be written as

$$u = \sum_{i=1}^k \bar{U}_{\delta_a, x_{a,i}} + \bar{\varphi}_a$$

with $|x_{a,i} - b_i| = o(1)$, $\|\bar{\varphi}_a\|_{\delta_a} = o(\delta_a^{\frac{N}{2}})$. Moreover,

$$\bar{\varphi}_a \in \bigcap_{i=1}^k \bar{E}_{a, x_{a,i}} := \left\{ \phi \in H^s(\mathbb{R}^N) : \left\langle \phi, \frac{\partial \bar{U}_{\delta_a, x_{a,i}}}{\partial x_j} \right\rangle_{\delta_a} = 0, j = 1, \dots, N \right\},$$

where

$$\bar{U}_{\delta_a, x_{a,i}}(x) := (1 + (\gamma_1 + V_i)\delta_a^{2s})^{\frac{1}{p-1}} U\left(\frac{(1 + (\gamma_1 + V_i)\delta_a^{2s})^{\frac{1}{2s}}(x - x_{a,i})}{\delta_a}\right),$$

$$\|\phi\|_{\delta_a}^2 = \int_{\mathbb{R}^N} (\delta_a^2 |(-\Delta)^{\frac{s}{2}} \phi|^2 + \phi^2).$$

Now, we write the equation (5.18) as

$$\bar{L}_a(\bar{\varphi}_a) = N_a(\bar{\varphi}_a) + \bar{l}_a(x),$$

where

$$\bar{L}_a(\bar{\varphi}_a) = \delta_a^{2s}(-\Delta)^s \bar{\varphi}_a + [1 + (-\mu_a \delta_a^{2s} - 1) + \delta_a^{2s} V] \bar{\varphi}_a - p \sum_{i=1}^k \bar{U}_{\delta_a, x_{a,i}}^{p-1} \bar{\varphi}_a,$$

N_a, L_a are defined by (3.2), (3.3), and

$$\bar{l}_a(x) = (\mu_a \delta_a^{2s} + 1 + \gamma_1 \delta_a^{2s} - \delta_a^{2s} V(x) + V_i \delta_a^{2s}) \sum_{i=1}^k \bar{U}_{\delta_a, x_{a,i}} + \left(\sum_{i=1}^k \bar{U}_{\delta_a, x_{a,i}} \right)^p - \sum_{i=1}^k \bar{U}_{\delta_a, x_{a,i}}^p.$$

Lemma 5.4. *It holds that*

$$\|\bar{\varphi}_a\|_{\delta_a} = O(\delta_a^{\frac{N}{2}+2s+2}). \quad (5.20)$$

Proof. Similar to Lemma 3.2, we obtain (5.20) only from proving

$$\|\bar{l}_a\|_{\delta_a} = O\left(\sum_{i=1}^k |V(x_{a,i}) - V_i| \delta_a^{\frac{N}{2}+2s} + \left| \sum_{i=1}^k \nabla V(x_{a,i}) \right| \delta_a^{\frac{N}{2}+2s+1} + \delta_a^{\frac{N}{2}+2s+2}\right) = O(\delta_a^{\frac{N}{2}+2s+2}).$$

□

Let $u_a^{(1)}$ and $u_a^{(2)}$ be two k -peak solutions to (5.18) and (5.19) concentrating at k points b_1, \dots, b_k , of the form

$$u_a^{(l)} = \sum_{i=1}^k \bar{U}_{\delta_a, x_{a,i}^{(l)}} + \bar{\varphi}_a^{(l)}, \quad \bar{\varphi}_a^{(l)} \in \bigcap_{i=1}^k \bar{E}_{a, x_{a,i}^{(l)}} \quad l = 1, 2. \quad (5.21)$$

Set $\xi_a(x) = \frac{u_a^{(1)}(x) - u_a^{(2)}(x)}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^N)}}$. Then $\|\xi_a\|_{L^\infty(\mathbb{R}^N)} = 1$. We get from (5.18) that

$$\delta^{2s}(-\Delta)^s \xi_a + C_a(x) \xi_a = g_a(x),$$

where

$$C_a(x) = \delta^{2s} V(x) - \delta^{2s} \mu_a^{(1)} - p \int_0^1 (t u_a^{(1)} + (1-t) u_a^{(2)})^{p-1} dt,$$

$$g_a(x) = \frac{\delta_a^{2s} (\mu_a^{(1)} - \mu_a^{(2)})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^N)}} u_a^{(2)}.$$

Following [15] and (4.1), we can prove that

$$|\xi_a(x)| + |\nabla \xi_a(x)| \leq C \sum_{i=1}^k \frac{1}{\left(1 + \left|\frac{x - x_{a,i}}{\delta_a}\right|^2\right)^{\frac{N+2s-\theta}{2}}}. \quad (5.22)$$

Applying blowing-up technique, we set

$$\bar{\xi}_{a,i}(x) = \xi_{a,i} \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right), \quad P_i = 1 + (\gamma_1 + V_i) \delta^{2s}.$$

The $\bar{\xi}_{a,i}$ satisfies that

$$(-\Delta)^s \bar{\xi}_{a,i} + \frac{1}{P_i} C_a \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) \bar{\xi}_{a,i} = \frac{1}{P_i} g_a \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right). \quad (5.23)$$

Lemma 5.5. For $x \in B_{\delta_a^{-1} P_i^{2s} \rho}(0)$, it holds that

$$\frac{1}{P_i} C_a \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) = 1 - p U^{p-1}(x) + O \left(\delta_a^{2s} + \sum_{l=1}^2 \varphi_a^{(l)} \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) \right), \quad (5.24)$$

and

$$\frac{1}{P_i} g_a \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) = -\frac{p-1}{ka_*} U(x) \sum_{j=1}^k \int_{\mathbb{R}^N} U^p(x) \bar{\xi}_{a,j}(x) + O \left(\delta_a^{2s+2} + \sum_{l=1}^2 \varphi_a^{(l)} \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) \right). \quad (5.25)$$

Proof. Since

$$\begin{aligned} C_a \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) &= \delta^{2s} V \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) - \delta^{2s} \mu_a^{(1)} \\ &\quad - p \int_0^1 (t u_a^{(1)} + (1-t) u_a^{(2)})^{p-1} \left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)} \right) dt, \end{aligned}$$

we can obtain (5.24) directly.

From (5.18) and (5.19), for $l = 1, 2$, we get that

$$\mu_a^{(l)} \delta_a^{\frac{4s}{p-1} + 2s} a^{\frac{2}{p-1}} = \delta_a^{2s} \int_{\mathbb{R}^N} \left(|(-\Delta)^{\frac{s}{2}} u_a^{(l)}|^2 + V(x) (u_a^{(l)})^2 \right) - \int_{\mathbb{R}^N} |u_a^{(l)}|^{p+1},$$

which implies that

$$\begin{aligned} & \frac{\delta_a^{\frac{4s}{p-1} + 2s} a^{\frac{2}{p-1}} (\mu_a^{(1)} - \mu_a^{(2)})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^N)}} \\ &= \delta_a^{2s} \int_{\mathbb{R}^N} \left((-\Delta)^{\frac{s}{2}} u_a^{(1)} + (-\Delta)^{\frac{s}{2}} u_a^{(2)} \right) \cdot (-\Delta)^{\frac{s}{2}} \xi_a + V(x) (u_a^{(1)} + u_a^{(2)}) \xi_a \\ &\quad - \int_{\mathbb{R}^N} \frac{(|u_a^{(1)}|^{p+1} - |u_a^{(2)}|^{p+1})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^N)}} \\ &= \delta_a^{2s} \int_{\mathbb{R}^N} \left((-\Delta)^{\frac{s}{2}} u_a^{(1)} + (-\Delta)^{\frac{s}{2}} u_a^{(2)} \right) \cdot (-\Delta)^{\frac{s}{2}} \xi_a + V(x) (u_a^{(1)} + u_a^{(2)}) \xi_a \\ &\quad - \int_{\mathbb{R}^N} \frac{|u_a^{(1)}|^{p+1} - |u_a^{(2)}|^{p+1}}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^N)}} - \mu_a^{(2)} \delta_a^{2s} \int_{\mathbb{R}^N} (u_a^{(1)} + u_a^{(2)}) \xi_a \\ &= (\mu_a^{(1)} - \mu_a^{(2)}) \delta_a^{2s} \int_{\mathbb{R}^N} u_a^{(1)} \xi_a + \int_{\mathbb{R}^N} (|u_a^{(1)}|^p + |u_a^{(2)}|^p) \xi_a - (p+1) \int_{\mathbb{R}^N} \xi_a \int_0^1 (t u_a^{(1)} + (1-t) u_a^{(2)})^p dt. \end{aligned} \quad (5.26)$$

Hence, in view of (5.14), (5.15), (5.20), (5.26) gives that

$$\begin{aligned} \frac{\delta_a^{2s}(\mu_a^{(1)} - \mu_a^{(2)})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^N)}} &= -\frac{p-1}{ka_*\delta_a^N} \left(\int_{\mathbb{R}^N} (|u_a^{(1)}|^p + |u_a^{(2)}|^p)\xi_a - (p+1)\xi_a \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p dt \right) \\ &\quad + O(\delta_a^{2s+2}), \end{aligned}$$

which implies (5.25). \square

Since $|\bar{\xi}_{a,i}| \leq 1$, we suppose that $\bar{\xi}_{a,i} \rightarrow \bar{\xi}_i(x)$ in $C_{loc}^1(\mathbb{R}^N)$. Lemma 5.5 actually shows the following result.

Lemma 5.6. *There hold that*

$$(-\Delta)^s \bar{\xi}_i + (1 - pU^{p-1})\bar{\xi}_i = -\frac{p-1}{ka_*} U(x) \sum_{j=1}^k \int_{\mathbb{R}^N} U^p(x) \bar{\xi}_j(x), \quad i = 1, \dots, k. \quad (5.27)$$

In order to show $\bar{\xi}_i = 0$, we write

$$\bar{\xi}_{a,i}(x) = \sum_{j=0}^N \gamma_{a,i,j} \Psi_j + \bar{\zeta}_{a,i}(x), \quad \text{in } H^s(\mathbb{R}^N), \quad (5.28)$$

where $\Psi_j (j = 0, 1, \dots, N)$ are functions in (A.3), $\bar{\zeta}_{a,i} \in \bar{E}$ and

$$\bar{E} = \{u \in H^s(\mathbb{R}^N), \langle u, \Psi_j \rangle = 0, \text{ for } j = 0, 1, \dots, N\}.$$

It is standard to show the following problem.

Lemma 5.7. *For any $u \in \bar{E}$, there holds that*

$$\|\bar{L}(u)\| \geq c_0 \|u\|,$$

where $c_0 > 0$ is a constant, and the linear operator is defined by

$$\bar{L}(u) = (-\Delta)^s u + (1 - pU^{p-1})u + \frac{p-1}{a_*} U(x) \int_{\mathbb{R}^N} U^p(x) u(x) dx. \quad (5.29)$$

Proposition 5.8. *The $\bar{\xi}_{a,i}(x)$ defined by (5.28) satisfies that*

$$\|\bar{\zeta}_{a,i}\| = O(\delta_a^{2s}). \quad (5.30)$$

Proof. Using (5.23)-(5.27), for any $\psi \in H^s(\mathbb{R}^N)$ we have by Lemma 5.5 and (5.20) that

$$\begin{aligned} \langle \bar{L}(\bar{\zeta}_{a,i}), \psi \rangle &= \int_{\mathbb{R}^N} \left[(1 - pU^{p-1} - \frac{1}{P_i} C_a(\frac{\delta_a}{P_i^{1/2s}} x + x_{a,i}^{(1)})) \bar{\xi}_{a,i} \right] \psi \\ &\quad + \int_{\mathbb{R}^N} \left[\frac{1}{P_i} g_a(\frac{\delta_a}{P_i^{1/2s}} x + x_{a,i}^{(1)}) + \frac{p-1}{ka_*} U(x) \sum_{j=1}^k \int_{\mathbb{R}^N} U^p(x) \bar{\xi}_{a,j}(x) \right] \psi \\ &= O(\delta_a^{2s}) \|\psi\| + O\left(\int_{\mathbb{R}^N} \sum_{l=1}^2 \varphi_a^{(l)}(\frac{\delta_a}{P_i^{1/2s}} x + x_{a,i}^{(1)}) |\psi| \right) \\ &= O(\delta_a^{2s}) \|\psi\| + O(\delta_a^{-\frac{N}{2}}) \|\varphi_a^{(l)}\|_{\delta_a} \|\psi\| = O(\delta_a^{2s}) \|\psi\|, \end{aligned}$$

which implies (5.30) by Lemma 5.7. \square

In the following we prove

$$\gamma_{a,i,j} = o(1) \text{ for } j = 0, 1, \dots, N.$$

First, we have the estimate for ξ_a .

Lemma 5.9. *It holds that*

$$\begin{aligned} & \delta_a^{2s} \sum_{i=1}^k \int_{B_\rho(x_{a,i}^{(1)})} \left(sV(x) + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \right) (u_a^{(1)} + u_a^{(2)}) \xi_a \\ &= (\mu_a^{(1)} - \mu_a^{(2)}) \delta_a^{2s} \int_{\mathbb{R}^N} u_a^{(1)} \xi_a + \int_{\mathbb{R}^N} ((u_a^{(1)})^p + (u_a^{(2)})^p) \xi_a \\ & \quad - (p+1) \int_{\mathbb{R}^N} \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p \xi_a \\ & \quad + \left(\frac{N}{p+1} - \frac{N-2s}{2} \right) (p+1) \int_{B_\rho(x_{a,i}^{(1)})} \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p \xi_a + O(\delta_a^{2N+2s}). \end{aligned} \tag{5.31}$$

Proof. Since $u_a^{(l)}$ are the k -peak solutions of (5.18)–(5.19), we have the following Pohozaev identity:

$$\begin{aligned} & -\delta_a^{2s} \int_{\partial^\nu \mathbf{B}_\rho(x_{a,i}^{(1)})} t^{1-2s} \frac{\partial \tilde{u}_a^{(l)}}{\partial \nu} \langle Y - X_{a,i}, \nabla \tilde{u}_a^{(l)} \rangle + \frac{\delta_a^{2s}}{2} \int_{\partial^\nu \mathbf{B}_\rho(x_{a,i}^{(1)})} t^{1-2s} |\nabla \tilde{u}_a^{(l)}|^2 \langle Y - X_{a,i}, \nu \rangle \\ & - \frac{N-2s}{2} \delta_a^{2s} \int_{\partial^\nu \mathbf{B}_\rho(x_{a,i}^{(1)})} t^{1-2s} \frac{\partial \tilde{u}_a^{(l)}}{\partial \nu} \tilde{u}_a^{(l)} \\ & - \int_{\partial B_\rho(x_{a,i}^{(1)})} \left(\frac{(u_a^{(l)})^{p+1}}{p+1} - \frac{\delta_a^{2s} (-\mu_a^{(l)} + V(x))}{2} (u_a^{(l)})^2 \right) \langle y - x_{a,i}, \nu \rangle \\ &= \int_{B_\rho(x_{a,i}^{(1)})} \left(\frac{N-2s}{2} - \frac{N}{p+1} \right) (u_a^{(l)})^{p+1} + s \delta_a^{2s} (-\mu_a^{(l)} + V(x)) (u_a^{(l)})^2 \\ & \quad + \frac{\delta_a^{2s}}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle (u_a^{(l)})^2, \end{aligned}$$

which can be rewritten as

$$\begin{aligned} & \delta_a^{2s} \int_{B_\rho(x_{a,i}^{(1)})} \left(sV(x) (u_a^{(l)})^2 + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle (u_a^{(l)})^2 \right) \\ &= \int_{B_\rho(x_{a,i}^{(1)})} \delta_a^{2s} s \mu_a^{(l)} (u_a^{(l)})^2 + \left(\frac{N}{p+1} - \frac{N-2s}{2} \right) (u_a^{(l)})^{p+1} \\ & \quad + \delta_a^{2s} \int_{\partial^\nu \mathbf{B}_\rho(x_{a,i}^{(1)})} W_1^{(l)} + \delta_a^{2s} \int_{\partial B_\rho(x_{a,i}^{(1)})} W_2^{(l)}, \end{aligned}$$

where

$$\begin{aligned} W_1^{(l)} &= -t^{1-2s} \frac{\partial \tilde{u}_a^{(l)}}{\partial \nu} \langle Y - X_{a,i}, \nabla \tilde{u}_a^{(l)} \rangle + \frac{1}{2} t^{1-2s} |\nabla \tilde{u}_a^{(l)}|^2 \langle Y - X_{a,i}, \nu \rangle - \frac{N-2s}{2} t^{1-2s} \frac{\partial \tilde{u}_a^{(l)}}{\partial \nu} \tilde{u}_a^{(l)}, \\ W_2^{(l)} &= \left(-\frac{(u_a^{(l)})^{p+1}}{\delta_a^{2s}(p+1)} + \frac{(-\mu_a^{(l)} + V(x))}{2} (u_a^{(l)})^2 \right) \langle y - x_{a,i}, \nu \rangle. \end{aligned}$$

Then, the above local Pohozaev identity implies that

$$\begin{aligned}
 & \delta_a^{2s} \int_{B_\rho(x_{a,i}^{(1)})} \left(sV(x) + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \right) (u_a^{(l)} + u_a^{(2)}) \xi_a \\
 &= \frac{(\mu_a^{(1)} - \mu_a^{(2)}) \delta_a^{2s}}{\|u_a^{(l)} - u_a^{(2)}\|_{L^\infty}} \int_{B_\rho(x_{a,i}^{(1)})} (u_a^{(1)})^2 + \mu_a^{(2)} \delta_a^{2s} \int_{B_\rho(x_{a,i}^{(1)})} (u_a^{(1)} + u_a^{(2)}) \xi_a \\
 &+ \left(\frac{N}{p+1} - \frac{N-2s}{2} \right) (p+1) \int_{B_\rho(x_{a,i}^{(1)})} \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p \xi_a + J_{a,i,1} + J_{a,i,2},
 \end{aligned} \tag{5.32}$$

where

$$\begin{aligned}
 J_{a,i,1} &= \int_{\partial'' \mathbf{B}_\rho(\mathbf{x}_{a,i}^{(1)})} \frac{\delta_a^{2s} (W_1^{(1)} - W_1^{(2)})}{\|u_a^{(l)} - u_a^{(2)}\|_{L^\infty}} \\
 &= -\delta_a^{2s} \int_{\partial'' \mathbf{B}_\rho(\mathbf{x}_{a,i}^{(1)})} t^{1-2s} \frac{\partial \tilde{u}_a^{(1)}}{\partial \nu} \langle Y - X_{a,i}, \nabla \tilde{\xi}_a \rangle - \delta_a^{2s} \int_{\partial'' \mathbf{B}_\rho(\mathbf{x}_{a,i}^{(1)})} t^{1-2s} \frac{\partial \tilde{\xi}_a}{\partial \nu} \langle Y - X_{a,i}, \nabla \tilde{u}_a^{(2)} \rangle \\
 &+ \frac{\delta_a^{2s}}{2} \int_{\partial'' \mathbf{B}_\rho(\mathbf{x}_{a,i}^{(1)})} t^{1-2s} \nabla(\tilde{u}_a^{(1)} + \tilde{u}_a^{(2)}) \cdot \nabla \tilde{\xi}_a \langle Y - X_{a,i}, \nu \rangle \\
 &- \frac{N-2s}{2} \delta_a^{2s} \int_{\partial'' \mathbf{B}_\rho(\mathbf{x}_{a,i}^{(1)})} t^{1-2s} \left(\frac{\partial \tilde{\xi}_a}{\partial \nu} \tilde{u}_a^{(1)} - \frac{\partial \tilde{u}_a^{(2)}}{\partial \nu} \tilde{\xi}_a \right) = O(\delta_a^{2s+2N}),
 \end{aligned}$$

and since $p+1 > 2$

$$J_{a,i,2} = \int_{\partial B_\rho(x_{a,i}^{(1)})} \frac{\delta_a^{2s} (W_2^{(1)} - W_2^{(2)})}{\|u_a^{(l)} - u_a^{(2)}\|_{L^\infty}} = O(\delta_a^{2s+2(N+2s-\theta)} + \delta_a^{(p+1)N+2ps}) = O(\delta_a^{2s+2(N+2s-\theta)}).$$

Summing (5.32) from $i = 1$ to $i = k$ and considering the decay (4.1) and (5.22),

$$\begin{aligned}
 & \sum_{i=1}^k \delta_a^{2s} \int_{B_\rho(x_{a,i}^{(1)})} \left(sV(x) + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \right) (u_a^{(l)} + u_a^{(2)}) \xi_a \\
 &= \frac{(\mu_a^{(1)} - \mu_a^{(2)}) \delta_a^{2s}}{\|u_a^{(l)} - u_a^{(2)}\|_{L^\infty}} \int_{\mathbb{R}^N} (u_a^{(1)})^2 + \mu_a^{(2)} \delta_a^{2s} \int_{\mathbb{R}^N} (u_a^{(1)} + u_a^{(2)}) \xi_a \\
 &+ \left(\frac{N}{p+1} - \frac{N-2s}{2} \right) (p+1) \int_{B_\rho(x_{a,i}^{(1)})} \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p \xi_a + O(\delta_a^{2N+2s}),
 \end{aligned}$$

which, combined with (5.26), gives (5.31), concluding the proof. \square

In the following, we first show $\gamma_{a,i,0} = o(1)$ for $i = 1, \dots, k$.

Lemma 5.10. *It holds that*

$$\gamma_{a,i,0} = o(1), \quad i = 1, \dots, k.$$

Proof. From (5.15), (5.20), (5.21) and

$$u_a^{(2)} \left(\frac{\delta_a}{P_i^{\frac{1}{2s}}} x + x_{a,i}^{(1)} \right) = P_i U(x) + O\left(\frac{P_i^{\frac{1}{2s}} |x_{a,i}^{(1)} - x_{a,i}^{(2)}|}{\delta_a} \right) + \bar{\varphi}_a \left(\frac{\delta_a}{P_i^{\frac{1}{2s}}} x + x_{a,i}^{(1)} \right),$$

we obtain that

$$\begin{aligned}
& \int_{B_\rho(x_{a,i}^{(1)})} \left(sV(x) + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \right) (u_a^{(l)} + u_a^{(2)}) \xi_a \\
&= \int_{B_\rho(x_{a,i}^{(1)})} \left(s(V(x) - V_i) + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \right) \left(\sum_{i=1}^k (\bar{U}_{\delta_a, x_{a,i}^{(1)}} + \bar{U}_{\delta_a, x_{a,i}^{(2)}}) + \bar{\varphi}_a^{(1)} + \bar{\varphi}_a^{(2)} \right) \xi_a \\
&\quad + \int_{B_\rho(x_{a,i}^{(1)})} sV_i (u_a^{(l)} + u_a^{(2)}) \xi_a \\
&= 2 \int_{B_\rho(x_{a,i}^{(1)})} \left(s(V(x) - V_i) + \frac{1}{2} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \right) \bar{U}_{\delta_a, x_{a,i}^{(1)}} \xi_a \\
&\quad + \int_{B_\rho(x_{a,i}^{(1)})} sV_i (u_a^{(l)} + u_a^{(2)}) \xi_a + O(\delta_a^{N+2s+2}).
\end{aligned}$$

From (5.4) (5.15), (5.28) and (5.30), we obtain that

$$\begin{aligned}
& \int_{B_\rho(x_{a,i}^{(1)})} (V(x) - V_i) \bar{U}_{\delta_a, x_{a,i}^{(1)}} \xi_a \\
&= \delta_a^N \int_{\mathbb{R}^N} \left(V\left(\frac{\delta_a}{P_i^{2s}} x + x_{a,i}^{(1)}\right) - V_i \right) U(x) \left(\sum_{j=0}^N \gamma_{a,i,j} \Psi_j \right) + O(\delta_a^{N+2s+2}) \\
&= -\frac{B}{2} \Delta V(b_i) \gamma_{a,i,0} \delta_a^{N+2} + O(\delta_a^{N+2s+2} + \delta_a^{N+3}),
\end{aligned}$$

where B is the constant as defined in (4.11).

Similarly,

$$\int_{B_\rho(x_{a,i}^{(1)})} \langle y - x_{a,i}^{(1)}, \nabla V(x) \rangle \bar{U}_{\delta_a, x_{a,i}^{(1)}} \xi_a = -\frac{B}{2} \Delta V(b_i) \gamma_{a,i,0} \delta_a^{N+2} + O(\delta_a^{N+2s+2} + \delta_a^{N+3}).$$

Therefore, it holds

$$\begin{aligned}
LHS \text{ of (5.31)} &= -\frac{2s+1}{4} B \Delta V(b_i) \gamma_{a,i,0} \delta_a^{N+2+2s} + O(\delta_a^{N+6s} + \delta_a^{N+3+2s}) \\
&\quad + s \delta_a^{2s} V_i \int_{B_\rho(x_{a,i}^{(1)})} (u_a^{(l)} + u_a^{(2)}) \xi_a.
\end{aligned} \tag{5.33}$$

Moreover, we have

$$\begin{aligned}
& \int_{B_\rho(x_{a,i}^{(1)})} (u_a^{(l)} + u_a^{(2)}) \xi_a \\
&= 2 \gamma_{a,i,0} \delta_a^N \int_{\mathbb{R}^N} U \left(U + \frac{p-1}{2s} x \cdot \nabla U \right) + O(|x_{a,i}^{(1)} - x_{a,i}^{(1)}| \delta_a^{N-1} + \delta_a^{\frac{N}{2}} \|\bar{\varphi}_a^{(2)}\|_{\delta_a}) \\
&= 2 \gamma_{a,i,0} \delta_a^N \left(1 - \frac{N(p-1)}{4s} \right) \int_{\mathbb{R}^N} U^2 + O(\delta_a^{N+2s+2} + \delta_a^{N+3}),
\end{aligned}$$

which, combined with (5.33), gives that

$$\begin{aligned} \text{LHS of (5.31)} &= -\frac{2s+1}{4}B\Delta V(b_i)\gamma_{a,i,0}\delta_a^{N+2+2s} + O(\delta_a^{N+6s} + \delta_a^{N+3+2s}) \\ &\quad + sV_i a_* \gamma_{a,i,0} \left(1 - \frac{N(p-1)}{4s}\right) \delta_a^{N+2s}. \end{aligned}$$

On the other hand, for the RHS of (5.31),

$$\begin{aligned} &\int_{\mathbb{R}^N} ((u_a^{(1)})^p + (u_a^{(2)})^p) \xi_a - (p+1) \int_{\mathbb{R}^N} \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p \xi_a \\ &\quad + \left(\frac{N}{p+1} - \frac{N-2s}{2}\right)(p+1) \int_{B_\rho(x_{a,i}^{(1)})} \int_0^1 (tu_a^{(1)} + (1-t)u_a^{(2)})^p \xi_a \\ &= \left(N - (p-1) - \frac{(N-2s)(p+1)}{2}\right) \left(1 - \frac{N}{p+1}\right) \gamma_{a,i,0} \int_{\mathbb{R}^N} U^{p+1} + O(\delta_a^{N+2+4s}) \\ &= \left(N - (p-1) - \frac{(N-2s)(p+1)}{2}\right) \left(1 - \frac{N(p-1)}{2s(p+1)}\right) \frac{1}{\left(\frac{N}{s(p+1)} - \frac{N}{2s} + 1\right)} a_* \gamma_{a,i,0} \\ &= -(\tilde{\gamma} + o(1)) a_* \gamma_{a,i,0} \delta_a^N \end{aligned}$$

with some constant $\tilde{\gamma} > 0$ since $N \geq 2s + 2$ and $p < \frac{N+2s}{N-2s}$. Moreover, by (5.14), (5.15) and (5.20), it holds that

$$(\mu_a^{(1)} - \mu_a^{(2)}) \delta_a^{2s} \int_{\mathbb{R}^N} u_a^{(1)} \xi_a = O(\delta_a^{N+6s}).$$

To sum up, we finally obtain that

$$\begin{aligned} &-\frac{2s+1}{4}B\Delta V(b_i)\gamma_{a,i,0}\delta_a^{N+2+2s} + O(\delta_a^{N+6s} + \delta_a^{N+3+2s} + \delta_a^{2N+2s}) \\ &\quad + sV_i a_* \gamma_{a,i,0} \left(1 - \frac{N(p-1)}{4s}\right) \delta_a^{N+2s} = -(\tilde{\gamma} + o(1)) a_* \gamma_{a,i,0} \delta_a^N, \end{aligned}$$

which implies $\gamma_{a,i,0} = o(1)$. □

Lemma 5.11. *It holds that $\gamma_{a,i,j} = o(1)$, for $i = 1, \dots, k$ and $j = 1, \dots, N$.*

Proof. This Lemma can be proved just following that in [30], so we only sketch the main steps as follows.

Step 1: To prove $\gamma_{a,i,N} = O(\delta_a)$ for $i = 1, \dots, k$.

On one hand, using (4.5), we deduce

$$\int_{B_\rho(x_{a,i}^{(1)})} \frac{\partial V(x)}{\partial \nu_{a,i}} A_a(x) \xi_a = O(\delta_a^{2N}), \quad (5.34)$$

where $\nu_{a,i}$ is the outward unit vector of $\partial B_\rho(x_{a,i}^{(1)})$ at x , $A_a(x) = \sum_{l=1}^2 u_a^{(l)}(x)$.

On the other hand, by (5.15), and the Taylor expansions

$$\begin{aligned}
& \int_{B_\rho(x_{a,i}^{(1)})} \frac{\partial V(x)}{\partial \nu_{a,i}} A_a(x) \xi_a \\
&= \int_{\mathbb{R}^N} \frac{\partial V(x_{a,i}^{(1)})}{\partial \nu_{a,i}} A_a(x) \xi_a + \int_{\mathbb{R}^N} \left\langle \nabla \frac{\partial V(x_{a,i}^{(1)})}{\partial \nu_{a,i}}, x - x_{a,i}^{(1)} \right\rangle A_a(x) \xi_a + O(\delta_a^{N+2}) \quad (5.35) \\
&= -\frac{\partial^2 V(x_{a,i}^{(1)})}{\partial \nu_{a,i}^2} a_* \gamma_{a,i,N} \delta_a^{N+1} + O(\delta_a^{N+2}).
\end{aligned}$$

Then (5.34) and (5.35) imply $\gamma_{a,i,N} = O(\delta_a)$.

Step 2: To prove $\gamma_{a,i,j} = o(1)$ for $i = 1, \dots, k$ and $j = 1, \dots, N-1$.

Similar to (5.34), we first have

$$\int_{B_\rho(x_{a,i}^{(1)})} \frac{\partial V(x)}{\partial \tau_{a,i,j}} A_a(x) \xi_a = O(\delta_a^{2N}), \text{ for } i = 1, \dots, k, j = 1, \dots, N-1. \quad (5.36)$$

Using suitable rotation, we assume that $\tau_{a,i,1} = (1, 0, \dots, 0), \dots, \tau_{a,i,N-1} = (0, \dots, 0, 1, 0)$ and $\nu_{a,i} = (0, \dots, 0, 1)$. Under the condition (V), we know

$$\begin{aligned}
\frac{\partial V(\delta_a x + x_{a,i}^{(1)})}{\partial \tau_{a,i,j}} &= \delta_a \sum_{l=1}^N \frac{\partial^2 V(x_{a,i}^{(1)})}{\partial x_l \partial \tau_{a,i,j}} x_l + \frac{\delta_a^2}{2} \sum_{m=1}^N \sum_{l=1}^N \frac{\partial^3 V(x_{a,i}^{(1)})}{\partial x_m \partial x_l \partial \tau_{a,i,j}} x_m x_l \\
&\quad + \frac{\delta_a^3}{6} \sum_{s=1}^N \sum_{m=1}^N \sum_{l=1}^N \frac{\partial^4 V(x_{a,i}^{(1)})}{\partial x_s \partial x_l \partial x_m \partial \tau_{a,i,j}} x_s x_l x_m + o(\delta_a^3 |x|^3), \text{ in } B_{d\delta_a^{-1}}(0). \quad (5.37)
\end{aligned}$$

By (1.6), (5.4), (5.20), (5.28), and the symmetry of $\varphi_j(x)$, we find, for $j = 1, \dots, N-1$,

$$\begin{aligned}
& \sum_{m=1}^N \sum_{l=1}^N \frac{\partial^3 V(x_{a,i}^{(1)})}{\partial x_m \partial x_l \partial \tau_{a,i,j}} \int_{B_{d\delta_a^{-1}}(0)} A_a(\delta_a x + x_{a,i}^{(1)}) \bar{\xi}_{a,i} x_m x_l \\
&= B \gamma_{a,i,0} \frac{\partial \Delta V(x_{a,i}^{(1)})}{\partial \tau_{a,i,j}} + O(\delta_a^{2s}) = O(|x_{a,i}^{(1)} - b_i|) + O(\delta_a^{2s}) = O(\delta_a^{2s}).
\end{aligned}$$

Also from Step 1, we get

$$\begin{aligned}
& \sum_{s=1}^N \sum_{m=1}^N \sum_{l=1}^N \frac{\partial^4 V(x_{a,i}^{(1)})}{\partial x_s \partial x_l \partial x_m \partial \tau_{a,i,j}} \int_{B_{d\delta_a^{-1}}(0)} A_a(\delta_a x + x_{a,i}^{(1)}) \bar{\xi}_{a,i} x_s x_l x_m \\
&= -3B \left(\sum_{q=1}^{N-1} \frac{\partial^2 \Delta V(b_i)}{\partial \tau_{a,i,q} \partial \tau_{a,i,j}} \gamma_{a,i,q} \right) + o(1).
\end{aligned}$$

By (A.2), we estimate

$$\frac{\partial^2 V(x_{a,i}^{(1)})}{\partial x_l \partial \tau_{a,i,j}} = -\frac{\partial V(x_{a,i}^{(1)})}{\partial \nu_{a,i}} \kappa_{i,l}(x_{a,i}^{(1)}) \delta_{lj}, \quad l, j = 1, \dots, N-1.$$

Since $\frac{\partial V(\bar{x}_{a,i}^{(1)})}{\partial \nu_{a,i}} = 0$, from (5.4), we find

$$\frac{\partial^2 V(x_{a,i}^{(1)})}{\partial x_l \partial \tau_{a,i,j}} = \frac{B}{2a_*} \frac{\partial \Delta V(b_i)}{\partial \nu_i} \delta_a^2 \kappa_{i,l}(b_i) \delta_{lj} + o(\delta_a^2). \quad (5.38)$$

Therefore from (5.28), (5.20) and (5.38), we get

$$\sum_{l=1}^N \frac{\partial^2 V(x_{a,i}^{(1)})}{\partial x_l \partial \tau_{a,i,j}} \int_{B_{\delta a^{-1}}(0)} A_a(\delta_a x + x_{a,i}^{(1)}) \bar{\xi}_{a,i} x_l = -\frac{B}{2} \frac{\partial \Delta V(b_i)}{\partial \nu_i} \delta_a^2 \kappa_{i,j}(b_i) \gamma_{a,i,j} + o(\delta_a^2). \quad (5.39)$$

Combining (5.37)–(5.39), we obtain

$$\begin{aligned} & \int_{B_\rho(x_{a,i}^{(1)})} \frac{\partial V(x)}{\partial \tau_{a,i,j}} A_a(x) \xi_a \\ &= -\frac{B}{2} \frac{\partial \Delta V(b_i)}{\partial \nu_i} \delta_a^{N+3} \kappa_{i,j}(b_i) \gamma_{a,i,j} - \frac{B}{2} \left(\sum_{l=1}^{N-1} \frac{\partial^2 \Delta V(b_i)}{\partial \tau_{i,l} \partial \tau_{i,j}} \gamma_{a,i,l} \right) \delta_a^{N+3} + o(\delta_a^{N+3}). \end{aligned} \quad (5.40)$$

From (5.36) and (5.40), we find

$$\frac{\partial \Delta V(b_i)}{\partial \nu_i} \kappa_{i,j}(b_i) \gamma_{a,i,j} + \left(\sum_{l=1}^{N-1} \frac{\partial^2 \Delta V(b_i)}{\partial \tau_{i,l} \partial \tau_{i,j}} \gamma_{a,i,l} \right) = o(1),$$

which implies $\gamma_{a,i,j} = o(1)$ for $i = 1, \dots, k$ and $j = 1, \dots, N-1$. \square

Now we are ready to prove Theorem 1.6.

Proof of Theorem 1.6: First, for large fixed R , (5.24) and (5.25) give

$$C_a(x) \geq \frac{1}{2}, \quad |g_a(x)| \leq C \sum_{i=1}^k \frac{\delta_a^{N+2s}}{|x - x_{a,i}^{(1)}|^{N+2s}}, \quad x \in \mathbb{R}^N \setminus \bigcup_{i=1}^k B_{R\delta_a}(x_{a,i}^{(1)}).$$

Using the comparison principle associated to the nonlocal operator $(-\Delta)^s$, we get

$$\xi_a(x) = o(1), \quad \text{in } \mathbb{R}^N \setminus \bigcup_{i=1}^k B_{R\delta_a}(x_{a,i}^{(1)}).$$

On the other hand, it follows from (5.28), (5.30), Lemmas 5.10 and 5.11 that

$$\xi_a(x) = o(1), \quad \text{in } \bigcup_{i=1}^k B_{R\delta_a}(x_{a,i}^{(1)}).$$

This is in contradiction with $\|\xi_a\|_{L^\infty(\mathbb{R}^N)} = 1$. So $u_a^{(1)}(x) \equiv u_a^{(2)}(x)$ for $a \rightarrow a_0$. \square

APPENDIX

A. BASIC ESTIMATES

Lemma A.1 (c.f. [37]). *Let $\alpha, \beta \geq 1$ be two constants. For any $0 < \delta \leq \min\{\alpha, \beta\}$, and $x_1 \neq x_2$, it holds that*

$$\frac{1}{(1 + |y - x_1|)^\alpha (1 + |y - x_2|)^\beta} \leq \frac{C}{|x_1 - x_2|^\delta} \left(\frac{1}{(1 + |y - x_1|)^{\alpha+\beta-\delta}} + \frac{1}{(1 + |y - x_2|)^{\alpha+\beta-\delta}} \right).$$

Lemma A.2 (c.f. [25]). *Let $\rho > \theta > 0$ be two constants. Suppose $(y-x)^2 + t^2 \geq \rho^2, t > 0$ and $\alpha > N$. Then, when $0 < \beta < N$, it holds that*

$$\int_{\mathbb{R}^N} \frac{1}{(t+|z|)^\alpha |y-z-x|^\beta} \leq C \left(\frac{1}{(1+|y-x|)^\beta} \frac{1}{t^{\alpha-N}} + \frac{1}{(1+|y-x|)^{\alpha+\beta-N}} \right), \quad (\text{A.1})$$

and

$$\int_{\mathbb{R}^N \setminus B_\theta(0)} \frac{1}{(t+|z|)^\alpha |y-z-x|^\beta} \leq \frac{C}{(1+|y-x|)^\beta} \frac{1}{\theta^{\alpha-N}};$$

when $N < \beta$, we have that

$$\int_{\mathbb{R}^N \setminus B_\theta(y-x)} \frac{1}{(t+|z|)^\alpha |y-z-x|^\beta} \leq C \left(\frac{1}{(1+|y-x|)^\beta} \frac{1}{t^{\alpha-N}} + \frac{1}{(1+|y-x|)^\beta} \frac{1}{\theta^{\beta-N}} \right),$$

where $C > 0$ is a constant independent of θ .

Remark A.3. *For some $\epsilon \rightarrow 0$, in the case of $\beta < N$, (A.1) implies directly that*

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{1}{(t+|z|)^\alpha \left(1 + \frac{|y-z-x|}{\epsilon}\right)^\beta} &\leq C\epsilon^\beta \int_{\mathbb{R}^N} \frac{1}{(t+|z|)^\alpha |y-z-x|^\beta} \\ &\leq C\epsilon^\beta \left(\frac{1}{(1+|y-x|)^\beta} \frac{1}{t^{\alpha-N}} + \frac{1}{(1+|y-x|)^\alpha} \right). \end{aligned}$$

While in the case of $\beta > N$, following the proof in [25], we can get indeed that,

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{1}{(t+|z|)^\alpha \left(1 + \frac{|y-z-x|}{\epsilon}\right)^\beta} &\leq C \left(\frac{\epsilon^\beta}{(1+|y-x|)^\beta} \frac{1}{t^{\alpha-N}} + \frac{\epsilon^N}{(1+|y-x|)^\alpha} \right) \\ &\leq C\epsilon^N \left(\frac{1}{(1+|y-x|)^\beta} \frac{1}{t^{\alpha-N}} + \frac{1}{(1+|y-x|)^\alpha} \right). \end{aligned}$$

Now let $\Gamma \in C^2$ be a closed hypersurface in \mathbb{R}^N . For $y \in \Gamma$, let $\nu(y)$ and $T(y)$ denote respectively the outward unit normal to Γ at y and the tangent hyperplane to Γ at y . The curvatures of Γ at a fixed point $y_0 \in \Gamma$ are determined as follows. By a rotation of coordinates, we can assume that $y_0 = 0$ and $\nu(0)$ is the x_N -direction, and x_j -direction is the j -th principal direction. In some neighborhood $\mathcal{N} = \mathcal{N}(0)$ of 0, we have

$$\Gamma = \{x : x_N = \varphi(x')\},$$

where $x' = (x_1, \dots, x_{N-1})$,

$$\varphi(x') = \frac{1}{2} \sum_{j=1}^{N-1} \kappa_j x_j^2 + O(|x'|^3),$$

where κ_j , is the j -th principal curvature of Γ at 0. The Hessian matrix $[D^2\varphi(0)]$ is given by

$$[D^2\varphi(0)] = \text{diag}[\kappa_1, \dots, \kappa_{N-1}].$$

Suppose that W is a smooth function, such that $W(x) = a$ for all $x \in \Gamma$.

Lemma A.4 (c.f. [30]). *We have*

$$\left. \frac{\partial W(x)}{\partial x_l} \right|_{x=0} = 0, \quad l = 1, \dots, N-1,$$

$$\left. \frac{\partial^2 W(x)}{\partial x_m \partial x_l} \right|_{x=0} = - \left. \frac{\partial W(x)}{\partial x_N} \right|_{x=0} \kappa_i \delta_{ml}, \text{ for } m, l = 1, \dots, N-1, \quad (\text{A.2})$$

where $\kappa_1, \dots, \kappa_{N-1}$, are the principal curvatures of Γ at 0.

B. LINEARIZATION

Lemma B.1. *Let $\xi = (\xi_1, \xi_2, \dots, \xi_k)$ be the solution of the following problem*

$$(-\Delta)^s \xi_i + (1 - pU^{p-1})\xi_i = -\frac{p-1}{ka_*} U(x) \sum_{j=1}^k \int_{\mathbb{R}^N} U^p(x) \xi_j(x), \quad i = 1, \dots, k. \quad (\text{A.1})$$

Then, there holds that

$$\xi_i(x) = \sum_{j=0}^N \gamma_{i,j} \Psi_j, \quad (\text{A.2})$$

where $\gamma_{i,j}$ are constants, and

$$\Psi_0 = U + \frac{p-1}{2s} x \cdot \nabla U, \quad \Psi_j = \frac{\partial U}{\partial x_j}, \quad j = 1, \dots, N. \quad (\text{A.3})$$

Moreover, $\gamma_{i,0} = \gamma_{l,0}$ for $i, l = 1, \dots, k$.

Proof. Setting

$$\bar{L}(u) = (-\Delta)^s u + (1 - pU^{p-1})u \text{ and } \Psi^j = (\Psi_j, \dots, \Psi_j),$$

then it is easy to find that $\bar{L}(\Psi^j) = 0$ and Ψ^j solve (A.1) for $j = 1, \dots, N$.

On the other hand, we denote

$$\Psi = (U + \frac{p-1}{2s} x \cdot \nabla U, \dots, U + \frac{p-1}{2s} x \cdot \nabla U),$$

and then by checking the extension equation associated with (A.1), we could obtain that Ψ solves (A.1) too. Noting that Ψ, Ψ^j are linearly independent, we obtain (A.2).

Moreover, putting (A.2) into (A.1), we obtain that $\gamma_{i,0} = \gamma_{l,0}$ for any $i, l = 1, \dots, k$. \square

Acknowledgments The authors would like to thank Professor Wenxiong Chen for the helpful discussion with him. Guo was supported by NSFC grants (No.11771469). Luo and Wang were supported by the Fundamental Research Funds for the Central Universities(No.KJ02072020-0319). Luo was supported by NSFC grants (No.11701204, No.11831009). Wang was supported by NSFC grants (No.12071169). Yang was supported by NSFC grants (No.11601194).

REFERENCES

- [1] D. Applebaum, Levy processes and stochastic calculus, Second edition, Cambridge Studies in Advanced Mathematics,116, Cambridge University Press, Cambridge, 2009.
- [2] B. Barrios, E. Colorado, A. de Pablo, U. Snchez, On some critical problems for the fracitonal Laplacian operator, J. Differential Equations, 252(2012), 6133–6162.
- [3] C. Brndle, E. Colorado, A. de Pablo, U. Snchez, A concave-convex elliptic problem involving the fractional Laplacian, Proc. Roy. Soc. Edinburgh Sect. A 143(2013),39–71.
- [4] B. Barrios, E. Colorado, R. Servadei, F. Soria, A critical fractional equation with concave-convex power nonlinearities, Ann. Inst. H. Poincar Anal. Non Linaire, 32(2015), 875–900.
- [5] X. Cabr, Y. Sire, Nonlinear equations for fractional Laplacians I: regularit, maximum principles, and Hamiltonian estimates, Ann. Inst. H. Poincar Anal. Non Linaire, 31(2014), 23–53.

- [6] X. Cabré, J. Tan, Positive solutions of nonlinear problems involving the square root of the Laplacian, *Adv. Math.* 224(2010), 2052–2093.
- [7] Caffarelli, L., Silvestre, L., An extension problem related to the fractional Laplacian. *Commun. Partial Diff. Equ.* 32 (2007), 1245–1260.
- [8] D. Cao, H. Heinz, *Uniqueness of positive multi-lump bound states of nonlinear Schrödinger equations.* *Math. Z.* 243 (2003), 599–642.
- [9] D. Cao, S. Li, P. Luo, *Uniqueness of positive bound states with multi-bump for nonlinear Schrödinger equations.* *Calc. Var. Partial Differential Equations* 54 (2015), 4037–4063.
- [10] D. Cao, E. Noussair, S. Yan, *Existence and uniqueness results on single-peaked solutions of a semilinear problem.* *Ann. Inst. H. Poincaré Anal. Non Linéaire* 15 (1998), 73–111.
- [11] W. Chen, C. Li, J. Zhu, Fractional equations with indefinite nonlinearities, *Disc. Cont. Dyna. Sys.*, 39(2019), 1257–1268.
- [12] W. Chen, Y. Li, P. Ma, *The fractional Laplacian*, World Scientific Publishing Co Pte Ltd, Singapore, 2019.
- [13] E. Dancer, S. Yan, *Interior and boundary peak solutions for a mixed boundary value problem.* *Indiana Univ. Math. J.* 48 (1999), 1177–1212.
- [14] E. Dancer, S. Yan, *On the existence of multipeak solutions for nonlinear field equations on \mathbb{R}^N .* *Discrete Contin. Dynam. Systems* 6 (2000), 39–50.
- [15] J. Dávila, M. del Pino, J. Wei, *Concentrating standing waves for the fractional nonlinear Schrödinger equation.* *J. Differential Equations*, 256(2014), 858–892.
- [16] Y. Deng, C. Lin, S. Yan, *On the prescribed scalar curvature problem in \mathbb{R}^N , local uniqueness and periodicity.* *J. Math. Pures Appl.* 104 (2015), 1013–1044.
- [17] S. Dipierro, G. Palatucci, E. Valdinoci, Existence and symmetry results for a Schrödinger type problem involving the fractional Laplacian, *Matematiche (Catania)*, 68(2013), 201–216.
- [18] R. L. Frank, E. Lenzmann, Uniqueness of non-linear ground states for fractional Laplacians in \mathbb{R} , *Acta Math.* 210(2013), 261–318.
- [19] R. L. Frank, E. Lenzmann, L. Silvestre, Uniqueness of radial solutions for the fractional Laplacian, *Commun. Pure Appl. Math.* 69(2016), 1671–1726.
- [20] P. Felmer, A. Quaas, J. Tan, Positive solutions of the nonlinear Schrödinger equation with the fractional Laplacian, *Proc. Roy. Soc. Edinburgh Sect. A* 142(2012), 1237–1262.
- [21] J. Jin, Y. Li, J. Xiong, On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions, *J. Eur. Math. Soc.* 16(2014), 1111–1171.
- [22] E. Di Nezza, G. Palatucci, E. Valdinoci, Hitchhiker’s guide to the fractional Sobolev spaces, *Bull. Sci. Math.*, 136 (2012), 521–573.
- [23] Y. Guo, T. Liu, J. Nie, Solutions for fractional operator problem via local Pohozaev identities, *Adv. Nonlinear Stud.* 20 (2020), 185–211.
- [24] Y. Guo, J. Nie, Infinitely many non-radial solutions for the prescribed curvature problem of fractional operator, *Discrete and Continuous Dynamical Systems*, 36(12)(2016), 6873–6898.
- [25] Y. Guo, J. Nie, M. Niu, Z. Tang, Local uniqueness and periodicity for the prescribed scalar curvature problem of fractional operator in \mathbb{R}^N , *Calc. Var. Partial Differential Equations* 56 (2017), Paper No. 118, 41 pp.
- [26] Y.X. Guo, S. Peng, S. Yan, *Local uniqueness and periodicity induced by concentration.* *Proc. Lond. Math. Soc.* 114 (2017), 1005–1043.
- [27] L. Ghangetas, *Uniqueness of positive solutions of a nonlinear elliptic equation involving the critical exponent.* *Nonlinear Anal.* 20 (1993), 571–603.
- [28] M. Grossi, *On the number of single-peak solutions of the nonlinear Schrödinger equations.* *Ann. Inst. H. Poincaré Anal. Non Linéaire* 19 (2002), 261–280.
- [29] Y. Guo, C. Lin, J. Wei, *Local uniqueness and refined spike profiles of ground states for two-dimensional attractive Bose-Einstein condensates.* *SIAM J. Math. Anal.* 49 (2017), 3671–3715.
- [30] P. Luo, S. Peng, S. Yan, *Excited states on Bose-Einstein condensates with attractive interactions.* arXiv:1909.08828.
- [31] M. Niu, Z. Tang, L. Wang, Solutions for conformally invariant fractional Laplacian equations with multi-bumps centered in lattices, *J. Differential Equations*, 266(2019), 1756–1831.
- [32] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, *Commun. Pure Appl. Math.* 60(1)(2007), 67–112.

- [33] Y. Sire, E. Valdinoci, Fractional Laplacian phase transitions and boundary reactions: a geometric inequality and a symmetry result, *J. Funct. Anal.* 256(6)(2009), 1842–1864.
- [34] J. Tan, The Brezis-Nirenberg type problem involving the square root of the Laplacian, *Calc. Var. Partial Differential Equations*, 42(2011), 21–41.
- [35] J. Tan, J. Xiong, A Harnack inequality for fractional Laplace equations with lower order terms. *Discrete Contin. Dyn. Syst.* 31(2011), 975–983.
- [36] M. Weinstein, Solitary waves of nonlinear dispersive evolution equations with critical power nonlinearities, *J. Diff. Equ.* 69(1987), 192–203.
- [37] J. Wei, S. Yan, Infinitely many solutions for the prescribed scalar curvature problem on S^N , *J. Funct. Anal.* 258(2010), 3048–3081.
- [38] S. Yan, J. Yang, X. Yu, Equations involving fractional Laplacian operator: compactness and application. *J. Funct. Anal.* 269(2015), 47–79.

(Qing Guo) COLLEGE OF SCIENCE, MINZU UNIVERSITY OF CHINA, BEIJING 100081, CHINA
Email address: guoqing0117@163.com

(Peng Luo) SCHOOL OF MATHEMATICS AND STATISTICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN 430079, CHINA
Email address: pluo@mail.ccnu.edu.cn

(Chunhua Wang) SCHOOL OF MATHEMATICS AND STATISTICS AND HUBEI KEY LABORATORY OF MATHEMATICAL SCIENCES, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN 430079, CHINA
Email address: chunhuawang@mail.ccnu.edu.cn

(Jing Yang) SCHOOL OF SCIENCE, JIANGSU UNIVERSITY OF SCIENCE AND TECHNOLOGY, ZHENJIANG 212003, CHINA
Email address: yyangecho@163.com