

Classification of irreducible integrable modules for extended affine Lie algebras with
center acting trivially

Santanu Tantubay¹, Punita Batra²

^{1,2} Harish-Chandra Research Institute, HBNI, Prayagraj (Allahabad) 211019, India

¹santanutantubay@hri.res.in

²batra@hri.res.in

*

*

ABSTRACT. We classify the irreducible integrable modules for the twisted toroidal extended affine Lie algebras with finite dimensional weight spaces where the finite dimensional center acts trivially. We have proved that the entire central extension part acts trivially on the modules and our modules turn out to be highest weight modules.

Msc: 17B67, 17B66.

Keywords: twisted toroidal extended affine Lie algebras

1. INTRODUCTION

Extended affine Lie algebras (EALAs) form a category of important Lie algebras consisting of finite dimensional semisimple Lie algebras, affine Lie algebras and some other classes of Lie algebras. Twisted toroidal extended affine Lie algebras are examples of EALAs. The structure theory of EALAs have been developed by several mathematicians like Allison, Azam, Berman, Gao, Neher, Pianzola and Yoshii (see [1], [16], [17] and references therein). With an extended affine Lie algebra, we associate a graded ideal, the core and its central quotient, the centerless core, both are Lie tori. So it is important to study the representation theory of EALAs.

Let g be a finite dimensional simple Lie algebra over \mathbb{C} and A_n be a Laurent polynomial ring in $n \geq 2$ commuting variables t_1, \dots, t_n . Let $L(g) = g \otimes A_n$ be a loop algebra. Let $L(g) \oplus \Omega_{A_n}/dA_n$ be the universal central extension of $L(g)$. This is called toroidal Lie algebras (TLA). Let $Der(A_n)$ be the Lie algebra of derivations of A_n . Then $L(g) \oplus \Omega_{A_n}/dA_n \oplus Der(A_n)$ is called the full toroidal Lie algebra (FTLA). TLA and FTLA are not EALA, as they do not admit any non-degenerate symmetric invariant bilinear form. So instead of $Der(A_n)$, one takes S_n the subalgebra of $Der(A_n)$, consisting of divergence zero vector fields. The Lie algebra $L(g) \oplus \Omega_{A_n}/dA_n \oplus S_n$ is called a toroidal extended affine Lie algebra which admits a non-degenerate symmetric bilinear form, and hence is an example of EALAs.

Here we consider more general EALAs. We take $\sigma_1, \dots, \sigma_n$ to be finite order commuting automorphisms of g and consider the multi-loop algebra $\bigoplus_{k \in \mathbb{Z}^n} g(\bar{k}) \otimes t^k$. The representations of multiloop algebras have been studied by Lau in [18]. We assume that this multi-loop algebra is a Lie torus. Now we consider the universal central extension of multi-loop algebra and add S_n , the Lie algebra consisting of divergence zero vector fields on A_n . This Lie algebra is called twisted toroidal extended affine Lie algebra, we denote it by τ . In [2], S.E. Rao, S. Sharma, P. Batra classified irreducible integrable modules for τ with finite dimensional weight spaces, where the zero degree central operators act nontrivially. In [15], the classification of irreducible integrable modules with finite dimensional weight spaces have been done for $n = 2$, both zero and non zero case. In this paper we will classify irreducible integrable modules with finite dimensional weight spaces for τ , where the zero degree central operators act trivially for any $n \geq 3$. We make use of some important results from [11] and [12] in order to classify our modules.

The paper has been organized as follows. In Section 2, we define the twisted toroidal extended affine Lie algebra τ as above. This Lie algebra has a natural triangular decomposition given by $\tau = \tau^- \oplus \tau^0 \oplus \tau^+$ (see Section 3). In Section 3, we prove the existence of highest weight vector with respect to this triangular decomposition. We show that the highest weight space is irreducible module for τ^0 as well as it is Γ -graded. In Section 4, we prove that if zero degree center $\mathbb{C}\text{-span}\{K_1, \dots, K_n\}$ acts trivially on V , then the whole central extension part $Z = \Omega_{A_n(m)}/d_{A_n(m)}$ acts trivially on V . In Section 5, to classify the modules we prove an important result Proposition 5.1(2). We have used a new method to prove this. In Section 6, we have proved our main theorem, Theorem 6.6 using a result of [11].

2. NOTATION AND PRELIMINARIES

Let $A_n = \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_n^{\pm 1}]$ be a Laurent polynomial ring in n variables. Let $L(g) = g \otimes A_n$ be the corresponding loop algebra, where g is a finite dimensional simple Lie algebra over \mathbb{C} with a Cartan subalgebra h . Let Ω_{A_n} be a vector space spanned by the symbols $t^k K_i, 1 \leq i \leq n, k \in \mathbb{Z}^n$. Let dA_n be the subspace spanned by $\sum_{i=1}^n k_i t^k K_i$. It is well known that $\tilde{L}(g) = L(g) \oplus \Omega_{A_n}/dA_n$ is the universal central extension of $L(g)$ with the following brackets:

$$[x(p), y(q)] = [x, y](p + q) + (x|y) \sum_{i=1}^n p_i t^{p+q} K_i,$$

where $x(p) = x \otimes t^p$. The basis for the Lie algebra $Der(A_n)$ is $\{d_i, t^r d_i | 1 \leq i \leq n, 0 \neq r \in \mathbb{Z}^n\}$. Now $Der(A_n)$ acts on Ω_{A_n}/dA_n by

$$t^p d_a(t^q K_b) = q_a t^{p+q} K_b + \delta_{ab} \sum_{c=1}^n p_c t^{p+q} K_c.$$

There are two non-trivial 2-cocycle of $Der(A_n)$ with values in Ω_{A_n}/dA_n :

$$\phi_1(t^p d_a, t^q d_b) = -q_a p_b \sum_{i=1}^n p_i t^{p+q} K_i$$

$$\phi_2(t^p d_a, t^q d_b) = -p_a q_b \sum_{i=1}^n p_i t^{p+q} K_i.$$

Let ϕ be any linear combination of ϕ_1 and ϕ_2 . Then $\tau = L(g) \oplus \Omega_{A_n}/dA_n \oplus Der(A_n)$ is a Lie algebra with the following brackets and is called a full toroidal Lie algebra:

$$[t^p d_a, X(q)] = q_a X(p + q),$$

$$[t^p d_a, t^q K_b] = q_a t^{p+q} K_b + \delta_{ab} \sum_{i=1}^n p_i t^{p+q} K_i,$$

$$[t^p d_a, t^q d_b] = q_a t^{p+q} d_b - p_b t^{p+q} d_a + \phi(t^p d_a, t^q d_b).$$

Now consider the subalgebra of divergence zero vector fields S_n of $Der(A_n)$. One can define $S_n = \{D(u, r) | (u|r) = 0, u \in \mathbb{C}^n, r \in \mathbb{Z}^n\}$. Now consider the subalgebra $\tau_{div} = L(g) \oplus \Omega_{A_n}/dA_n \oplus S_n$ of τ . It is well known that unlike τ , τ_{div} possesses a non-degenerate symmetric, invariant bilinear form and is called as toroidal extended affine Lie algebra. The form is defined as follows:

$$(X(r)|Y(s)) = \delta_{r,-s}(X|Y),$$

$$\forall X, Y \in \overset{\circ}{g}, r, s \in \mathbb{Z}^n;$$

$$(\sum_{i=1}^n a_i t^r d_i | t^s K_d) = \delta_{r,-s} a_d.$$

All other brackets of bilinear form are zero.

Now Let g_1 be any arbitrary finite dimensional simple Lie algebra over \mathbb{C} with a cartan subalgebra h_1 . Let $\Delta(g_1, h_1) = \text{supp}_{h_1}(g_1)$. Then $\Delta_1^\times = \Delta^\times(g_1, h_1) = \Delta(g_1, h_1) - \{0\}$ is an irreducible reduced finite root system with atmost two root lengths. Define

$$\Delta_{1,en}^\times = \Delta_1^\times \cup 2\Delta_{1,sh}^\times \quad \text{if } \Delta_1^\times = B_l \text{ types}$$

$$\Delta_1^\times \quad \text{otherwise.}$$

Definition 2.1. A finite dimensional g_1 -module V is said to satisfy condition (M) if V is irreducible with dimension greater than 1 and weights of V relative to h_1 are contained in $\Delta_{1,en}^\times$.

Now recall that g is a finite dimensional simple Lie algebra with a Cartan subalgebra h and let $\sigma_1, \sigma_2, \dots, \sigma_n$ be the commuting automorphisms of g of order m_1, m_2, \dots, m_n respectively. Let $m = (m_1, m_2, \dots, m_n) \in \mathbb{Z}^n$. Define $\Gamma = m_1 \mathbb{Z} \oplus \dots \oplus m_n \mathbb{Z}$ and $\Lambda := \mathbb{Z}^n / \Gamma$. Then we have $g = \bigoplus_{\bar{k} \in \Lambda} g(\bar{k})$, where $\bar{k} \in \Lambda$, and $g(\bar{k}) = \{X \in g | \sigma_i(X) = \zeta_i^{k_i} X, 1 \leq i \leq n\}$, where ζ_i are m_i -th primitive root of unity for $i = 0, \dots, n$.

Definition 2.2. A multiloop algebra $\bigoplus_{\bar{k} \in \mathbb{Z}^n} g(\bar{k}) \otimes t^k$ is called a Lie torus LT if

- (1) $g(\bar{0})$ is a finite dimensional simple Lie algebra.

- (2) For $\bar{k} \neq 0$ and $g(\bar{k}) \neq 0$, $g(\bar{k}) \cong U(\bar{k}) \oplus W(\bar{k})$, where $U(\bar{k})$ is trivial as $g(\bar{0})$ -module and either $W(\bar{k})$ is zero or satisfy condition (M).
- (3) The order of the group generated by $\sigma_i, 1 \leq i \leq n$ is equal to the product of orders of each σ_i , for $1 \leq i \leq n$.

Let $h(\bar{0})$ denote a Cartan subalgebra of $g(\bar{0})$. Then by [13] Lemma 3.1.3, $h(\bar{0})$ is ad-diagonalizable on g and $\Delta^\times = \Delta^\times(g, h(\bar{0}))$ is an irreducible finite root system in $h(\bar{0})$ (Proposition 3.3.5, [13]). Let $\Delta_0 := \Delta(g(\bar{0}), h(\bar{0}))$. One of the main properties of Lie tori is that $\Delta := \Delta_{0,en}$ (Proposition 3.2.5, [1]). Let $A_n(m) = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ and $S_n(m) = \{D(u, r) | (u|r) = 0, u \in \mathbb{C}^n, r \in \Gamma\}$. Now take $\tau = LT \oplus \Omega_{A_n(m)}/d_{A_n(m)} \oplus S_n(m)$. One can easily check that if $x \in g(\bar{k}), y \in g(\bar{l})$ with $(x|y) \neq 0$, then $\bar{k} + \bar{l} \in \Gamma$. Therefore we can check that τ is a well defined Lie algebra called as twisted toroidal extended affine Lie algebra. The aim of this paper is to classify the irreducible integrable modules of twisted toroidal EALAs where K_i 's are acting trivially $\forall i \in \{1, \dots, n\}$.

3. EXISTENCE OF HIGHEST WEIGHT SPACE

In this section we will give a root space decomposition of τ . Let $H = h(\bar{0}) \oplus \sum_{i=1}^n \mathbb{C}K_i \oplus \sum_{i=0}^n \mathbb{C}d_i$ be our Cartan subalgebra for the root space decomposition of τ . Define $\delta_i, w_i \in H^*$ by setting

$$\begin{aligned} \delta_i(h(\bar{0})) &= 0, \delta_i(K_j) = 0 \text{ and } \delta_i(d_j) = \delta_{ij}. \text{ and} \\ w_i(h(\bar{0})) &= 0, w_i(K_j) = \delta_{ij} \text{ and } w_i(d_j) = 0. \end{aligned}$$

Take $\delta_\beta = \sum_{i=1}^n \beta_i \delta_i$ for $\beta \in \mathbb{C}^n$. For $k \in \mathbb{Z}^n$, we shall refer to the vector $\delta_{k+\gamma}$ as the translate of δ_k by the vector δ_γ where $\gamma \in \mathbb{C}^n$. Define $g(\bar{k}, \alpha) := \{x \in g(\bar{k}) | [h, x] = \alpha(h)x, \forall h \in h(\bar{0})\}$. Then we have $\tau = \bigoplus_{\beta \in \Delta} \tau_\beta$, where $\Delta \subseteq \{\alpha + \delta_k | \alpha \in \Delta_{0,en}, k \in \mathbb{Z}^n\}$. We have $\tau_{\alpha+\delta_k} = g(\bar{k}, \alpha) \otimes t^k$ for $\alpha \neq 0$, and $\tau_{\delta_k} = g(\bar{k}, 0) \otimes t^k \oplus \bigoplus_{i=1}^n \mathbb{C}t^k K_i \oplus \bigoplus_{i=1}^n \mathbb{C}t^k d_i$ for $0 \neq k \in \mathbb{Z}^n$ and $\tau_0 = H$.

In order to get a non-degenerate form on H^* we extend $\alpha \in h(\bar{0})^*$ to H by defining $\alpha(K_i) = \alpha(d_i) = 0, \forall 1 \leq i \leq n$. Then $(h(\bar{0})|K_i) = 0, (\delta_k|\delta_l) = (w_i|w_j) = 0$ and $(\delta_i|w_j) = \delta_{ij}$ and form on $h(\bar{0})$ is the restriction of the form of g . One can easily check that this form is non-degenerate on H^* . A root $\beta = \alpha + \delta_k$ is called a real root if $\alpha \neq 0$. Let Δ^{re} denote the set of all real roots and $\beta^\vee = \alpha^\vee + \frac{2}{(\alpha|\alpha)} \sum_{i=1}^n k_i K_i$ is a co-root of β , where α^\vee is the co-root of $\alpha \in \Delta_{0,en}$. For $\gamma \in \Delta^{re}$ define $r_\gamma(\lambda) = \lambda - \lambda(\gamma^\vee)\gamma$ for $\lambda \in H^*$. Let W be the Weyl group of τ generated by $r_\gamma, \forall \gamma \in \Delta^{re}$.

Definition 3.1. A τ -module V is called integrable if

- (1) $V = \bigoplus_{\lambda \in H^*} V_\lambda$, where $V_\lambda = \{v \in V | h.v = \lambda(h)v \forall h \in H\}$ and $\dim(V_\lambda) < \infty$
- (2) All the real root vectors act locally nilpotently on V , i.e., $g(\bar{k}, \alpha) \otimes t^k$ acts locally nilpotently on V for all $0 \neq \alpha \in \Delta_{0,en}$.

Now we have the standard proposition

Proposition 3.2. Let V be an irreducible integrable module for τ . Then

- (1) $P(V) = \{\gamma \in H^* | V_\gamma \neq 0\}$ is W -invariant.
- (2) $\dim(V_\gamma) = \dim(V_{w\gamma}), \forall w \in W$.
- (3) If $\lambda \in P(V)$ and $\gamma \in \Delta^{re}$, then $\lambda(\gamma^\vee) \in \mathbb{Z}$.
- (4) If $\lambda \in P(V)$ and $\gamma \in \Delta^{re}$, and $\lambda(\gamma^\vee) > 0$, then $\lambda - \gamma \in P(V)$.

Now consider the natural triangular decomposition of τ

$$\tau^+ = \bigoplus_{\alpha > 0, k \in \mathbb{Z}^n} g(\bar{k}, \alpha) \otimes t^k$$

$$\tau^- = \bigoplus_{\alpha < 0, k \in \mathbb{Z}^n} g(\bar{k}, \alpha) \otimes t^k$$

$$\tau^0 = \bigoplus_{k \in \mathbb{Z}^n} g(\bar{k}, 0) \otimes t^k \oplus Z \oplus S_n(m)$$

Where $Z = \Omega_{A_n}/d_{A_n}$.

We have V is an integrable module for τ , so it will be integrable module for $\widetilde{LT} = LT \oplus \Omega_{A_n}/d_{A_n} \oplus D$, where D is the \mathbb{C} linear span of $d_i, 1 \leq i \leq n$.

Lemma 3.1. *Let V be an integrable \widetilde{LT} -module . Then there exists some $\lambda \in P(V)$ such that $\lambda + \eta \notin P(V)$*

This is the lemma 3.6 of [4].

Theorem 3.3. *The highest weight space $V_+ = \{v \in V | \tau^+.v = 0\}$ is non-zero.*

Proof. \widetilde{LT} is a Lie subalgebra of τ . So an integrable module for τ with finite dimensional weight spaces will be an integrable module for \widetilde{LT} with finite dimensional spaces. Now take $\lambda \in P(V)$ as in lemma 3.1. Using Proposition 3.2, it can be proved that $\lambda|_{h(\bar{0})}$ is a dominant integral weight of $g(\bar{0})$. Now using the same idea as [14] (Theorem 2.4(ii)), we can find $\mu \in P(V)$ such that $V_{\mu+\alpha+\delta_k} = 0, \forall \alpha \in \Delta_0^+, k \in \mathbb{Z}^n$. Again using Proposition 3.2 we will get $V_{\mu+2\alpha+\delta_k} = 0, \forall \alpha \in \Delta_0^+, k \in \mathbb{Z}^n$. Now our theorem is a direct consequence of $\Delta = \Delta_{0,en}$. \square

We can see that V_+ is a τ^0 module.

Lemma 3.2. (1) V_+ is irreducible module over τ^0 .

(2) $V = U(\tau^-)V_+$.

Proof. Using PBW theorem and some weight arguments we can prove (1). Using (1) and PBW theorem we can prove (2) \square

Lemma 3.3. (1) *There exists unigue $\bar{\lambda} \in h(\bar{0})^*$ and a $\beta \in \mathbb{C}^n$, here β need not be unique such that the weights of V_+ will be of the form $\bar{\lambda} + \delta_{r+\beta}$ where $r \in \mathbb{Z}^n$.*
 (2) $\bar{\lambda}$ is a dominant integral.

Proof. (1) We can see that $h(\bar{0})$ commutes with τ^0 . Now using Lemma 3.2 we will get $h(\bar{0})$ acts by scalars on V_+ and hence it will act by a single linear functional on V_+ . Let us denote it by $\bar{\lambda}$. If $D(u, r) \in S_n(m)$, then $D(u, r).V_\mu \subseteq V_{\mu+\delta_r}$ for $\mu \in P(V)$.
 (2) Direct consequence of Proposition 3.2 (4). \square

If $\bar{\lambda} = 0$ in Lemma 3.3, then using Proposition 3.2, we can check that only possible weights of V are $\delta_\beta, \beta \in \mathbb{C}^n$. Now one can prove that $LT + \Omega_{A_n(m)}/d_{A_n(m)}$ acts trivially on V . Then V will be irreducible module for $S_n(m)$ with finite dimensional weight spaces with respect to the cartan $D = \bigoplus_{i=1}^n \mathbb{C}d_i$. It is still unknown about the irreducible modules for $S_n(m)$ with finite-dimensional weight spaces. So we will take $\bar{\lambda} \neq 0$ in Lemma 3.3.

Take $\lambda \in P(V)$ from Lemma 3.1. Then by our assumption there exists $\alpha \in \Delta_0$ such that $\lambda(h_\alpha) \neq 0$. Then using Lemma 3.2 we can get

$$V_+ = \oplus_{r \in \mathbb{Z}^n} V_+(r),$$

where $V_+(r) = \{v \in V_+ | d_i.v = (\lambda(d_i) + m_i)v, 1 \leq i \leq n\}$. So V_+ is \mathbb{Z}^n graded. Now for $r \in \Gamma$ we define $V'(r) = \oplus_{r_i \leq k_i < m_i + r_i} V_k$. Then $V_+ = \oplus_{r \in \Gamma} V'_+(r)$ is Γ graded.

4. ACTION OF CENTER

In this section we will prove that $\Omega_{A_n(m)}/d_{A_n(m)}$ acts trivially on V . The following two lemmas are proved in [2] as well as in [7]

Proposition 4.1. *There are only finitely many $h(\bar{0}) \otimes A_n(m) \oplus \Omega_{A_n(m)}/d_{A_n(m)}$ -submodules of V_+*

Proposition 4.2. *The weight spaces of V_+ are uniformly bounded.*

Proposition 4.3. *$\Omega_{A_n(m)}/d_{A_n(m)}$ acts trivially on V .*

Proof. From Proposition 4.1, take a minimal $h(\bar{0}) \otimes A_n(m) \oplus \Omega_{A_n(m)}/d_{A_n(m)}$ -submodule of V_+ , assume W . Now clearly W will be irreducible. Take $W' = U(g(\bar{0}) \otimes A_n(m) \oplus \Omega_{A_n(m)}/d_{A_n(m)} \oplus D)W$. We can see that W goes injectively to the irreducible quotient of W' . Now using [5] we can see that $\Omega_{A_n(m)}/d_{A_n(m)}$ acts trivially on W . One can easily check that $S = \{v \in V | t^r K_i.v = 0, 1 \leq i \leq n\}$ is a non-zero τ -submodule of V . Hence we are done by irreducibility of V . \square

5. ACTION OF $h_\alpha \otimes t^k$ ON HIGHEST WIEGHT SPACE

Lemma 5.1. *Let $h \otimes t^k \in \tau^0$, where $k \in \Gamma \setminus \{0\}$ and $h \in h(\bar{0})$. If there exists a nonzero vector of V_+ on which $h \otimes t^k$ acts trivially, then $h \otimes t^k$ is locally nilpotent on V_+ .*

Proof. Consider the set $S = \{v \in V_+ | \text{there is a } n \in \mathbb{N} \text{ s.t. } (h \otimes t^k)^n.v = 0\}$. By assumption clearly this set is nonzero. We know that $h \otimes t^k$ commutes with $\bigoplus_{k \in \mathbb{Z}^n} g(\bar{k}, 0) \otimes t^k$. So $(\bigoplus_{k \in \mathbb{Z}^n} g(\bar{k}, 0) \otimes t^k)S \subseteq S$. Now using the equation

$$(h \otimes t^k)^n.D(u, r) = D(u, r).(h \otimes t^k)^n - n(u.k)(h \otimes t^k)^{n-1}.h \otimes t^{k+r}$$

we can prove that $S_n(m).S \subseteq S$. Therefore S is a nonzero τ^0 -submodule of V_+ . Hence by irreducibility of V_+ we can say that $S = V_+$. So the result. \square

The idea of the proof of following two lemma is from [12], but the calculations will be tottaly different. So we will give the proofs.

Lemma 5.2. *Let $h \otimes t^k$ acts locally nilpotently on V_+ , for every $k \in \Gamma \setminus \{0\}$, where $h \in h(\bar{0})$. Then $h \otimes t^k$ acts trivially on V_+ , for every $k \in \Gamma \setminus \{0\}$.*

Proof. Let $p \in \Gamma \setminus \{0\}$. Then by above discussion we will get $(h \otimes t^{-p}.h \otimes t^p)^N = 0$. Therefore $(h \otimes t^{-p})^N.(h \otimes t^p)^N = 0$.

Let $r_1 \in \Gamma \setminus \{0\}$ be such that $r_1 \notin \mathbb{Q}.p$. Then there exists $u \in \mathbb{C}^n$ such that $(u, p) \neq 0$ and $(u, r_1) = 0$. Therefore we have the following

$$D(u, r_1).(h \otimes t^{-p})^N.(h \otimes t^p)^N = 0$$

This equation will give us

$N(u, p)h \otimes t^{r_1+p} \cdot (h \otimes t^{-p})^N \cdot (h \otimes t^p)^{N-1} - N(u, p)h \otimes t^{r_1-p} \cdot (h \otimes t^{-p})^{N-1} \cdot (h \otimes t^p)^{N-1} = 0$
Now applying $h \otimes t^p$ on the above equation we will get

$$h \otimes t^{r_1-p} \cdot (h \otimes t^{-p})^{N-1} \cdot (h \otimes t^p)^{N+1} = 0.$$

Using induction on j , we want to prove, $h \otimes t^{r_1-p} \dots h \otimes t^{r_j-p} \cdot (h \otimes t^{-p})^{N-j} \cdot (h \otimes t^p)^{N+j} = 0$.

Where $1 \leq j \leq N$ and $r_1, \dots, r_j \in \Gamma$ with $\sum_{i=1}^j \epsilon_i r_i \notin \mathbb{Q}.p$ and $\epsilon_i \in \{0, 1\}$.

For $j = 1$, we already prove this. Now assume the result holds for $1 \leq j < N$.

Let $r \in \Gamma$ with $r + \sum_{i=1}^j \epsilon_i r_i \notin \mathbb{Q}.p$. So we can find $u \in \mathbb{C}^n$ such that $(u, p) \neq 0$ and $(u, r) = 0$. Therefore we have

$$D(u, r) \cdot h \otimes t^{r_1-p} \dots h \otimes t^{r_j-p} \cdot (h \otimes t^{-p})^{N-j} \cdot (h \otimes t^p)^{N+j} = 0.$$

From this equation we will get,

$$(N+j)(u, p)h \otimes t^{r_1-p} \dots h \otimes t^{r_j-p} \cdot h \otimes t^{r+p} \cdot (h \otimes t^{-p})^{N-j} \cdot (h \otimes t^p)^{N+j-1} - (N-j)(u, p)h \otimes t^{r_1-p} \dots h \otimes t^{r_j-p} \cdot h \otimes t^{r-p} \cdot (h \otimes t^{-p})^{N-j-1} \cdot (h \otimes t^p)^{N+j} = 0.$$

Applying $h \otimes t^p$, we have

$$h \otimes t^{r_1-p} \dots h \otimes t^{r_j-p} \cdot h \otimes t^{r-p} \cdot (h \otimes t^{-p})^{N-j} \cdot (h \otimes t^p)^{N+j} = 0.$$

Now taking $j = N$,

$$h \otimes t^{r_1-p} \dots h \otimes t^{r_N-p} \cdot (h \otimes t^p)^{2N} = 0$$

From this equation, using induction again, we can prove

$$h \otimes t^{r_1-p} \dots h \otimes t^{r_{N+j}-p} \cdot (h \otimes t^p)^{2N-j} = 0,$$

for all $0 \leq j \leq 2N$ with $\sum_{i=1}^{N+j} \epsilon_i r_i \notin \mathbb{Q}.p$ and $\epsilon_i \in \{0, 1\}$.

In this case instead of taking $D(u, r)$ we take $D(u, r - 2p)$, it is easy to find such $u \in \mathbb{C}^n$. Hence we will have

$$h \otimes t^{r_1-p} \dots h \otimes t^{r_{3N}-p} = 0,$$

with all $r_i \in \Gamma$ and $\sum_{i=1}^{3N} \epsilon_i r_i \notin \mathbb{Q}.p$.

Let $s_1, \dots, s_{3N} \in \Gamma \setminus \{0\}$ be arbitrary. Take $p \in \Gamma \setminus \bigcup_{\epsilon_i \in \{0,1\}} \mathbb{Q} \cdot (\sum_{i=1}^{3N} \epsilon_i s_i)$. Now taking $r_i = s_i + p$ in the above equation we can get

$$h \otimes t^{s_1} \dots h \otimes t^{s_{3N}} = 0 \dots (*) .$$

Take $W = \{v \in V_+ / h_\alpha \otimes t^p \cdot v = 0, \forall p \in \Gamma \setminus \{0\}\}$. By (*), W is nonzero. We can easily check that W is a τ^0 -submodule of V_+ . Hence by the irreducibility of V_+ , we have the lemma. \square

Lemma 5.3. $h \otimes t^k$ acts injectively or trivially on V_+ , for all $k \in \Gamma \setminus \{0\}$.

Proof. Let $h \otimes t^p$ acts injectively on V_+ , for some $p \in \Gamma \setminus \{0\}$.

Claim:(1) For any $q \in \Gamma \setminus \mathbb{Q}.p$, if $h \otimes t^q$ acts locally nilpotently, then it will act nilpotently.

Assume $h \otimes t^q$ acts locally nilpotently. Since $\dim(V_+(r)) \leq N$ for all $r \in \Gamma$, we will have $(h \otimes t^{-q})^N \cdot (h \otimes t^q)^N = 0$.

Suppose there exists $1 \leq k \leq N$ and N_k , such that $(h \otimes t^{-q})^k \cdot (h \otimes t^q)^{N_k} = 0$.

We assume $q \notin \mathbb{Q}.p$. So there exists $u \in \mathbb{C}^n$ such that $(u, p+q) = 0$ and $(u, q) \neq 0$.

Now applying $D(u, p+q)$, to the above equation we will have

$$N_k(u, q)h \otimes t^{p+q} \cdot (h \otimes t^{-p})^k (h \otimes t^p)^{N_k-1} - k(u, q)h \otimes t^p \cdot (h \otimes t^{-p})^{k-1} \cdot (h \otimes t^p)^{N_k} = 0.$$

Now applying $h \otimes t^q$, we have $h \otimes t^p \cdot (h \otimes t^{-p})^{k-1} \cdot (h \otimes t^p)^{N_k+1} = 0$. Since $h \otimes t^p$ acts injectively on V_+ , so $(h \otimes t^{-p})^{k-1} \cdot (h \otimes t^p)^{N_k+1} = 0$. Now repeating the process we can find $N_0 \in \mathbb{N}$ such that $(h_\alpha \otimes t^q)^{N_0} = 0$.

Claim:(2) If $q \in \Gamma \setminus \{0\} \cap \mathbb{Q}.p$, then $h \otimes t^q$ acts injectively on V_+ .

First we will prove that $h \otimes t^{\frac{a}{b}p}$ acts injectively, if $a \in \mathbb{Z}_-, b \in \mathbb{Z}_+$.

If for some such $a, b \in \mathbb{Z}$, $h \otimes t^{\frac{a}{b}p}$ is not acting injectively then by lemma 5.1, it will act locally nilpotently. Therefore $(h \otimes t^p)^{-a} \cdot (h \otimes t^{\frac{a}{b}p})^b$ acts nilpotently on each homogeneous spaces. Hence $(h \otimes t^p)^{-aN} \cdot (h \otimes t^{\frac{a}{b}p})^{bN} = 0$. Therefore $(h \otimes t^{\frac{a}{b}p})^{bN} = 0$, since $h \otimes t^p$ acts injectively on V_+ . Assume $N_0 \in \mathbb{N}$ be smallest such that $(h \otimes t^{\frac{a}{b}p})^{N_0} = 0$. Let $q \in \Gamma \setminus \mathbb{Q}.p$, then we can find $u \in \mathbb{C}^n$ such that, $(u, q - \frac{a}{b}p) = 0$ and $(u, p) \neq 0$. Now take

$$0 = D(u, q - \frac{a}{b}p) \cdot (h \otimes t^{\frac{a}{b}p})^{N_0} = N_0(u, \frac{a}{b}p) (h \otimes t^{\frac{a}{b}p})^{N_0-1} \cdot h \otimes t^q \cdot V_+$$

From the minimality of N_0 , we can say from the above equation and lemma 5.1 that $h \otimes t^q$ acts locally nilpotently. Therefore by claim (1) $h \otimes t^q$ acts nilpotently on V_+ . Choose N' be the smallest s.t, $(h \otimes t^q)^{N'} = 0$. Choose $u \in \mathbb{C}^n$ s.t $(u, p - q) = 0$ and $(u, q) \neq 0$. Then $0 = D(u, p - q) (h \otimes t^q)^{N'} = N'(u, q) (h \otimes t^q)^{N'-1} \cdot h \otimes t^p$.

Now by minimality of N' , we can see that $h \otimes t^p$ is not acting injectively, which is a contradiction. So our claim (2) is true for $a \in \mathbb{Z}_-, b \in \mathbb{Z}_+$. Now let both a, b are positive. We know by above $h \otimes t^{-ap}$ acts injectively. By our assumption $h \otimes t^{-ap} \cdot (h \otimes t^{\frac{a}{b}p})^b$ acts nilpotently on each homogeneous spaces. Now repeating the same process as above we will get the contradiction, $h \otimes t^p$ not acting injectively on V_+ . So our claim (2) is true. We are now going to prove that $h \otimes t^q$ acts injectively for all $q \in \Gamma \setminus \{0\}$. We already prove it for $q \in \mathbb{Q}.p$. Let $q \notin \mathbb{Q}.p$ and $h \otimes t^q$ does not acts injectively on V_+ . Therefore by lemma 5.1 and claim(1), we can say that $h \otimes t^q$ acts nilpotently on V_+ . Using the minimal power of $h \otimes t^q$, for which it acts trivially, we will arrive at $h \otimes t^p$ not acting injectively on V_+ , a contradiction to our assumption. Hence using lemma 5.1 and 5.2 we have the lemma. \square

Now we have choosen $\alpha \in \Delta_0$ such that $\bar{\lambda}(h_\alpha) \neq 0$. Now if there is a $k \in \Gamma \setminus \{0\}$ such that $h_\alpha \otimes t^k$ is not injective on V_+ , then by Lemma 5.3 we can say that $h_\alpha \otimes t^m$ acts trivially on V_+ . Then using [6] we can say that h_α acts trivially on V_+ , which is a contradiction to our assumption.

We know $V_+ = \oplus_{r \in \Gamma} V_+(r)$. Let $\{v_0, \dots, v_k\}$ be a basis of $V_+(0)$. Therefore by above discussion, $\{h_\alpha \otimes t^p \cdot v_0, \dots, h_\alpha \otimes t^p \cdot v_k\}$ will be a basis of $V_+(p)$ for all $p \in \Gamma$. Now assume, $h_\alpha \otimes t^p \cdot h_\alpha \otimes t^q \cdot (v_1, \dots, v_k) = h_\alpha \otimes t^{p+q} \cdot (v_1, \dots, v_k) A_{p,q}$, where $A_{p,q} \in M_k(\mathbb{C})$. Since $h_\alpha \otimes t^p$ are commuting family, we can easily see that $\{A_{p,q} | p, q \in \Gamma\}$ is a commuting family. Therefore we can find an invertible matrix $S \in M_k(\mathbb{C})$ such that $\{B_{p,q} = S^{-1} A_{p,q} S | p, q \in \Gamma\}$ is upper triangular family. Set $(w_1, \dots, w_k) = (v_1, \dots, v_k) \cdot S$, then we have $h_\alpha \otimes t^p \cdot h_\alpha \otimes t^q \cdot (w_1, \dots, w_k) = h_\alpha \otimes t^{p+q} \cdot (w_1, \dots, w_k) B_{p,q}$. So without loss of generality we can assume that $A_{p,q}$ is upper triangular for all $p, q \in \Gamma$.

Lemma 5.4. Let $p, q \in \Gamma$, then there exists $\lambda_{p,q}$ and $v_0 \in V_+(0)$ such that $(h_\alpha \otimes t^p \cdot h_\alpha \otimes t^q - \lambda_{p,q} h_\alpha \otimes t^{p+q}) v_0 = T_{p,q} v_0 = 0$.

Proof. Since for $\lambda \in \mathbb{C}$, $(h_\alpha \otimes t^p \cdot h_\alpha \otimes t^q - \lambda h_\alpha \otimes t^{p+q}) \cdot (v_1, \dots, v_k) = h_\alpha \otimes t^{p+q} \cdot (v_1, \dots, v_k) (A_{p,q} - \lambda I_k)$. Now the Lemma follows from $A_{p,q}$ has eigenvalues. \square

Lemma 5.5. *For $p, q \in \Gamma$, $\lambda \in \mathbb{C}$, if there exists $v \in V_+(0) \setminus \{0\}$ such that $T_\lambda \cdot v = 0$, then T_λ acts locally nilpotently on V_+ .*

Proof. The proof will be same as Lemma 5.1. \square

Now using Lemma 5.4 and Lemma 5.5, we can say that for every $p, q \in \Gamma$, $A_{p,q}$ has exactly one eigenvalue (say, $\lambda_{p,q}$).

Proposition 5.1. (1) $h_\alpha \otimes t^k$ acts injectively on V_+ for every $k \in \Gamma$.

(2) $h_\alpha \otimes t^r \cdot h_\alpha \otimes t^s = \lambda_{r,s} h_\alpha \otimes t^{r+s}$ on V_+ , where $\lambda_{r,s} = \lambda$ for all $r \neq 0, s \neq 0, r+s \neq 0$, $\lambda_{r,-r} = \mu$ for all $r \neq 0$ and $\lambda_{0,r} = \bar{\lambda}(h_\alpha)$ for all $r \in \Gamma$. Further we have $\mu \lambda_{0,r} = \lambda^2 \neq 0$.

(3) $\dim(V'_+(r)) = \dim(V'_+(s))$ for all $r, s \in \Gamma$.

Proof. (1) is a direct consequence of above discussion. (3) also follows from (1). Now we will prove (2). Set $V_0 = \sum_{i=0}^{k-1} \mathbb{C} v_i$, where $v_0 = 0$. Since $A_{p,q}$ is upper triangular, we have $T_{p,q} \cdot v_l \in \sum_{i=0}^{l-1} h_\alpha \otimes t^{p+q} \cdot v_i$. We can easily see that $T_{p,q} \cdot V_+(-p-q) \subseteq V_0$. Now for any $p, q, r \in \Gamma$, with $u \in \mathbb{C}^n$ such that $(u, r) = 0$, define $S_{p,q}^r = (u, p) h_\alpha \otimes t^{p+r} \cdot h_\alpha \otimes t^q + (u, q) h_\alpha \otimes t^p \cdot h_\alpha \otimes t^{q+r} - \lambda_{p,q}(u, p+q) h_\alpha \otimes t^{p+q+r}$. Therefore $h_\alpha \otimes t^{-(p+q+r)} \cdot S_{p,q}^r \cdot v_k = \lambda_{-(p+q+r), p+q+r}((u, p) \lambda_{p+r, q} + (u, q) \lambda_{p, q+r} - (u, p+q) \lambda_{p,q}) \cdot v_k + w'$, where $w' \in V_0$. One can also check that $h_\alpha \otimes t^{-(p+q+r)} \cdot S_{p,q}^r \cdot V_0 \subseteq V_0$. Now we can see that $[D(u, r), T_{p,q}] = S_{p,q}^r$. Given any $p, q, r \in \Gamma$, and $u \in \mathbb{C}^n$ with $(u, r) = 0$ we have

$$0 = (h_\alpha \otimes t^{-(p+q+r)})^k \cdot (D(u, r))^k \cdot (T_{p,q})^k \cdot v_k = T_{p,q} \cdot w + k! (h_\alpha \otimes t^{-(p+q+r)})^k \cdot (S_{p,q}^r)^k \cdot v_k,$$

where $w \in V_+(-p-q)$.

From the above equation, we will have

$$(u, p) \lambda_{p+r, q} + (u, q) \lambda_{p, q+r} - (u, p+q) \lambda_{p,q} = 0, \dots (**),$$

for any $p, q, r \in \Gamma$.

Take $W = \{v \in V_+ | T_{p,q} \cdot v = 0\}$, then $v_1 \in W$. Hence using the above equation we can check that W is a nonzero τ^0 -submodule of V_+ , therefore by irreducibility of V_+ , $W = V_+$.

Now taking $p = jq$, with $j \in \mathbb{Q} \setminus \{0, -1\}$ in (**), we will get

$$j(u, q) \lambda_{jq+r, q} + (u, q) \lambda_{j, q+r} - (j+1)(u, q) \lambda_{p,q} = 0.$$

Taking, $r \in \Gamma \setminus \mathbb{Q} \cdot q$, we can see that

$$j \lambda_{jq+r, q} + \lambda_{j, q+r} - (j+1) \lambda_{q,q} = 0.$$

Putting $j = 1$, we have $\lambda_{q, q+r} = \lambda_{q,q}$ and hence $\lambda_{jq+r, q} = \lambda_{j, q+r} = \lambda_{q+r, q+r} = \lambda_{q,q}$ for all $r \in \Gamma \setminus \mathbb{Q} \cdot q$. Therefore $\lambda_{p,q} = \lambda_{q,q} (= \lambda, \text{ say})$, if $p \neq 0, q \neq 0, p+q \neq 0$.

Taking $r = -p-q$ with $p \in \Gamma \setminus \mathbb{Q} \cdot q$ in (**), we have $\lambda_{p, -p} = \lambda_{q, -q} (= \mu, \text{ say})$ and hence for every $p, q \in \Gamma$. It is easy to see that $\lambda_{r,0} = \bar{\lambda}(h_\alpha)$.

Let us assume $p, q \in \Gamma$ be such that $p \neq 0, q \neq 0, p+q \neq 0$. then for any $v \in V_+$, we have

$$(h_\alpha \otimes t^p \cdot h_\alpha \otimes t^{-p}) \cdot (h_\alpha \otimes t^q \cdot h_\alpha \otimes t^{-q}) \cdot v = (\mu)^2 \cdot (\bar{\lambda}(h_\alpha))^2 v = \lambda_{p,q} \lambda_{-p, -q} \lambda_{p+q, -p-q} \bar{\lambda}(h_\alpha) \cdot v.$$

hence we have the identity of (2). \square

6. CLASSIFICATION

Now recall that our Lie algebra reduces to $\tau = LT \oplus S_n(m)$ with $\tau^0 = \oplus_{k \in \mathbb{Z}^n} g(\bar{k}, 0) \otimes t^k \oplus S_n(m)$. Take $g' = \{x \in g \mid [h, x] = 0, \forall h \in h(\bar{0})\}$. Now since g' is invariant under σ_i 's, therefore g' is Λ graded. It is easy to see that $L(g', \sigma) = LT^0$. Let us take $S'_n(m) = \text{span}\{D(u, r) - D(u, 0) \mid u \in \mathbb{C}^n, r \in \Gamma, (u, r) = 0\}$. We can easily check that $S'_n(m)$ is a Lie subalgebra of $S_n(m)$. Furthermore let us set $L = S'_n(m) \ltimes L(g', \sigma)$ and $W = \text{span}\{h_\alpha \otimes t^r \cdot v - v \mid r \in \Gamma, v \in V_+\}$. We can see that W is a L module.

Lemma 6.1. (1) W is a proper L -submodule of V_+ .

(2) $\tilde{V} = V_+/W$ is a finite dimensional L -module.

Proof. The proof is parallel to Lemma 7.2 of [7]. Let $z_i = h_\alpha \otimes t_i^m$ for each $i = 1, \dots, n$. Without loss of generality we can assume that $\lambda_{r,s} = 1$ for $r \neq 0, s \neq 0, r + s \neq 0$. Therefore we can say that $W = \text{span}\{z_i \cdot v - v \mid v \in V_+\}$. Now the proof will be same as [7] \square

Let $\beta \in \mathbb{C}^n$ be as in Lemma 3.3. Then for any L module V' we can give a τ^0 module structure on $L(V')$ by

$$x \otimes t^k \cdot (v_1 \otimes t^s) = ((x \otimes t^k) \cdot v_1) \otimes t^{k+s}.$$

$$D(u, r) \cdot (v_1 \otimes t^s) = ((D(u, r) - D(u, 0)) \cdot v_1) \otimes t^{r+s} + (u, s + \beta)(v_1 \otimes t^{r+s})$$

for all $v_1 \in V', x \in g(\bar{k}, 0)$ and $D(u, r) \in S_n(m)$.

For $v \in V_+$, let \bar{v} be the image of v in \tilde{V} . Now define

$$\phi : V_+ \rightarrow L(\tilde{V})$$

$$\text{by } v \mapsto \bar{v} \otimes t^k \text{ for } v \in V_+(k).$$

This map is clearly a nonzero τ^0 -module homomorphism. Hence by irreducibility of V_+ it follows that $V_+ \cong \phi(V_+)$ is a τ^0 submodule of $L(\tilde{V})$.

Clearly L is naturally Λ graded. Now since V_+ and W is \mathbb{Z}^n graded, therefore they are naturally Λ graded and hence so \tilde{V} . Therefore $\tilde{V} = \oplus_{\bar{p} \in \Lambda} \tilde{V}(\bar{p})$.

Now for $\bar{p} \in \Lambda$ we set

$$L(\tilde{V})(\bar{p}) = \{v \otimes t^{k+r+p} \mid v \in \tilde{V}(\bar{k}), r \in \Gamma, k \in \mathbb{Z}^n\}$$

It can be easily verified that $L(\tilde{V})(\bar{p})$ is a τ^0 submodule of $L(\tilde{V})$. The following result can be deduced similarly as in [3] (Same result is also proved in [3]).

Proposition 6.1. (1) $V_+ \cong L(\tilde{V})(\bar{0})$ as τ^0 -modules.

(2) \tilde{V} is Λ -graded-irreducible module over L .

(3) \tilde{V} is completely reducible module over L and all its irreducible components are mutually isomorphic as $S'_n(m) \ltimes h(\bar{0}) \otimes A(m)$ -modules.

Now we will concentrate on irreducible representation of L . Let (W, π) be a finite dimensional representation of L . Let $\pi(L(g', \sigma)) = g^1$, then $\pi(L) = g^1 \oplus g^2$, where g^2 is the unique complement of g^1 in $gl(W)$ (Proposition 19.1(b) of [8]). So W will be an irreducible module for $g^1 \oplus g^2$. Therefore $W \cong W_1 \otimes W_2$, where W_1 and W_2 are irreducible modules for g^1 and g^2 respectively ([9]). Let $g' = g'_{ss} \oplus R$, where g' and R are Levi and radical part of g' . Then as $\sigma_i(g') = g'$ and $\sigma_i(R) = R$ for $1 \leq i \leq n$, we have $L(g', \sigma) = L(g'_{ss}, \sigma) \oplus L(R, \sigma)$. Now W_1 is irreducible module for $L(g', \sigma)$. As R is solvable ideal, it follows that $\pi(L(R, \sigma))$ lies in the center of $\pi(L)$, which is atmost one dimensional. Hence $L(R, \sigma)$ acts as a scalar on W . So W_1 will be a irreducible module for $L(g'_{ss}, \sigma)$.

Fix a positive integer l . For each i , let $a_i = (a_{i,1}, \dots, a_{i,l})$ such that $a_{i,j}^{m_i} \neq a_{i,t}^{m_i}$ for $j \neq t$ (*). Let g be a finite dimensional semisimple Lie algebra. Let $\sigma_1, \dots, \sigma_n$ be finite order automorphisms on g of order m_1, \dots, m_n respectively. Let $L(g, \sigma)$ be the corresponding multiloop algebra. Let $I = \{(i_1, i_2, \dots, i_n) | 1 \leq i_j \leq l\}$. Now for $S = (i_1, i_2, \dots, i_n) \in I$ and $r = (r_1, r_2, \dots, r_n) \in \mathbb{Z}^n$, $a_S^r = a_{1,i_1}^{r_1} a_{2,i_2}^{r_2} \dots a_{n,i_n}^{r_n}$. Now consider the evaluation map $\phi : g \otimes A \rightarrow \bigoplus g$ (l^n copies), $\phi(X \otimes t^r) = (a_{I_1}^r, a_{I_2}^r, \dots, a_{I_{l^n}}^r)$, where I_1, I_2, \dots, I_{l^n} is some order on I . Now consider the restriction of ϕ to $L(g, \sigma)$.

Theorem 6.2. *Let W' be a finite dimensional irreducible representation of $L(g, \sigma)$. Then the representation factors through $\bigoplus g$ (l^n copies).*

The proof of this result comes from [10].

Proposition 6.3. *Let W_1 be irreducible module for $L(g'_{ss}, \sigma)$ as above. Then the representation of $L(g'_{ss}, \sigma)$ factors through only one copy of $\bigoplus g'_{ss}$. So $g_{ss}^1 \cong g'_{ss}$.*

Proof. We know by Theorem 6 that the representation factors through l^n copies, for some positive integer l . we will prove here that $l = 1$. The proof is same as [2]. Choose the i -th piece of g'_{ss} and choose the projection of the map π , say π_i onto it. Doing the same calculation as in [2] we will get $\pi_i(S'_n(m)) = 0$ and $a_{I_i}^r = 1$ for all $r \in \Gamma$. Now suppose there is atleast two pieces, say i -th and j -th piece is there. Therefore I_i and I_j are two different element of I with $a_{I_i}^r = 1 = a_{I_j}^r$ for all $r \in \Gamma$. Let $I_i = (i_1, i_2, \dots, i_n)$ and $I_j = (j_1, j_2, \dots, j_n)$. Therefore there is k with $1 \leq k \leq n$ such that $i_k \neq j_k$. Now if we take $r = (0, \dots, m_k, \dots, 0)$ then $a_{I_i}^r = 1 = a_{I_j}^r$ will give $a_{k,i_k}^{m_k} = a_{k,j_k}^{m_k}$, a contradiction to (*). So there is atleast one piece. \square

Now we know $\pi_i(S'_n(m)) = 0$, therefore $g^2 \subseteq \pi(S'_n(m))$. Now our aim is to understand finite dimensional irreducible modules for $S'_n(m)$. There is a relation between finite dimensional S'_n modules and $S_n \ltimes A$ -modules with finite dimensional weight spaces.

Theorem 6.4. *Suppose W is a finite dimensional irreducible sl_n -module (extend W to gl_n by letting I act trivially). Let $\alpha, \beta \in \mathbb{C}^n$. Take $L(W) = W \otimes A$ and the action $D(u, r)(w \otimes t^k) = (u \cdot k + \beta)w \otimes t^{k+r} + \sum_{i,j} u_i r_j E_{ji} w \otimes t^{k+r}$, for $r \neq 0$ and $D(u, 0)(w \otimes t^k) = (u \cdot \alpha + k)w \otimes t^k$ and $t^r(w \otimes t^k) = w \otimes t^{k+r}$. And all irreducible representations of $S_n \ltimes A$ with finite dimensional weight spaces occur in this way.*

This result is proved in [11]

Theorem 6.5. *Let W be finite dimensional irreducible module for sl_n and extend it trivially as before. Let E_{ij} be generators of gl_n . Then W can be made into S'_n -module by the action: $(D(u, r) - D(u, 0)) \cdot w = \sum_{i,j} E_{ji} w + (u \cdot \zeta)w$, where $\zeta \in \mathbb{C}^n$. In this case W will be irreducible module for S'_n . Every irreducible finite dimensional S'_n module occur in this way.*

Proof. See Theorem 4.5 of [2] and discussion thereafter. \square

We know \tilde{V} is completely reducible L module. Therefore $\tilde{V} = \bigoplus_{i=1}^K \tilde{V}_i$ for some $K \in \mathbb{N}$. Then by the previous discussion each $\tilde{V}_i \cong W_1^i \otimes W_2^i$ as $g'_{ss} \oplus sl_n$ module, where W_1^i, W_2^i are irreducible modules for g'_{ss} and sl_n respectively. since each component \tilde{V}_i are isomorphic as $S'_n(m) \ltimes (h(\bar{0}) \otimes A(m))$ modules, we can take $W_2^i \cong W_2^1 (= W_2)$, say as sl_n -modules for each $i \in \{1, \dots, K\}$. Now consider $W_1 = \sum_{i=1}^K W_1^i$, which is

a $L(g'_{ss}, \sigma)$ -module, in particular g'_{ss} module. Since each W_1^i are irreducible, without loss of generality we can assume that the above sum is direct. It is easy to see that L is Λ -graded with zero-th component $S'_n(m) \times (h(\bar{0}) \otimes A(m))$ and since \tilde{V} is Λ graded irreducible module (6.1), we can take W_1 as Λ -graded irreducible $L(g'_{ss}, \sigma)$ module and W_2 is zero graded as $S'_n(m)$ lies inside the zero-th graded component of L .

We now define a τ^0 -module structure on $W_1 \otimes W_2 \otimes A_n$ by

$$\begin{aligned} X \otimes t^k(w_1 \otimes w_2 \otimes t^l) &= Xw_1 \otimes w_2 \otimes t^{k+l}, \text{ for } k, l \in \mathbb{Z}^n \text{ and } X \in g'_{ss}(\bar{k}). \\ D(u, r)(w_1 \otimes w_2 \otimes t^k) &= (u.k + \beta)w_1 \otimes w_2 \otimes t^{k+r} + w_1 \otimes ((\sum_{i,j} u_i r_j E_{ji} w_2) \otimes t^{k+r} \text{ for } \\ &\quad r \neq 0 \text{ and } D(u, r) \in S_n(m). \\ D(u, 0)(w_1 \otimes w_2 \otimes t^k) &= (u.k + \alpha)w_1 \otimes w_2 \otimes t^k. \end{aligned}$$

Now take any one dimensional representation of $L(R, \sigma)$ say ψ . Then for $y \in R(\bar{k})$ we take $y \otimes t^k(w_1 \otimes w_2 \otimes t^l) = \psi(y)(w_1 \otimes w_2 \otimes t^{k+l})$. Since W_1 is Λ graded which is compatible with Λ -gradation of g'_{ss} , so the submodule $V' = \bigoplus_{k \in \mathbb{Z}^n} W_{1,k} \otimes W_2 \otimes t^k$ will be irreducible module for τ^0 . One can easily check that $L(\tilde{V})(\bar{0}) \cong V'$ as τ^0 -module. Now as in [7] consider the triangular decomposition of LT given by

$$LT = \tau^- \oplus L(g'_{ss}, \sigma) \oplus L(R, \sigma) \oplus \tau^+.$$

Let V_1 be the unique Λ -graded irreducible quotient of the induced module for W_1 and let (V_1^i, ρ_i) denote the irreducible quotient of the induced representation of W_1^i , for each $1 \leq i \leq K$. Now by [7] we can say that $V_1 = \bigoplus_{i=1}^K V_1^i$ is Λ -graded irreducible module for LT .

Define τ -module structure on $V_1 \otimes W_2 \otimes A$ by

$$\begin{aligned} (x \otimes t^l) \cdot (\sum_{i=1}^K v_1^i \otimes v_2 \otimes t^k) &= \sum_{i=1}^K (\rho_i(x \otimes t^l) v_1^i) \otimes v_2 \otimes t^{k+l}, \\ D(u, r) \cdot (\sum_{i=1}^K v_1^i \otimes v_2 \otimes t^k) &= \\ (u.k + \beta) (\sum_{i=1}^K v_1^i \otimes v_2 \otimes t^{k+r}) &+ \sum_{i=1}^K \sum_{l,j} v_1 \otimes (u_l r_j m_j E_{jl}) \cdot v_2 \otimes t^{k+r}. \end{aligned}$$

Let $V_1 = \bigoplus_{\bar{k} \in \Lambda} V_{1,\bar{k}}$. Take $\bigoplus_{k \in \mathbb{Z}^n} V_{1,\bar{k}} \otimes W_2 \otimes t^k$, which is a τ -submodule of $V_1 \otimes W_2 \otimes A$. One can prove that $\bigoplus_{k \in \mathbb{Z}^n} V_{1,\bar{k}} \otimes W_2 \otimes t^k$ is irreducible τ module.

Theorem 6.6. *Let V be an irreducible integrable τ module with finite dimensional weight spaces, with all K_i acting trivially for $1 \leq i \leq n$. Then $V \cong \bigoplus_{k \in \mathbb{Z}^n} V_{1,\bar{k}} \otimes W_2 \otimes t^k$.*

REFERENCES

- [1] Bruce Allison, Stephen Berman, John Faulkner, Arturo Pianzola, *Multiloop realization of extended affine Lie algebras and Lie tori*, C Trans. Am. Math. Soc.361(9) (2009) 4807-4842
- [2] S. Eswara Rao, Sachin S. Sharma, Punita Batra, *Integrable modules for twisted toroidal extended affine Lie algebras*, Journal of Algebra 556 (2020) 1057-1072
- [3] Punita Batra, Senapathi Eswara Rao, *On integrable modules for the twisted full toroidal Lie algebra*, J. Lie Theory 28 (1)(2018) 79-105
- [4] Souvik Pal, *Integrable modules for graded Lie tori with finite-dimensional weight spaces*, J. Pure and Applied Algebra.285(9)(2021)
- [5] S. Eswara Rao, *Classification of irreducible integrable modules for toroidal Lie algebras with finite dimensional weight spaces*, J. of Algebra 277(2004)318-348
- [6] Vyjayanthi Chari, Andrew Pressley, *Weyl modules for classical and quantum affine algebras*, Represent. Theory 5(2001) 191-223.
- [7] Souvik Pal, S. Eswara Rao, *Classification of level zero irreducible integrable modules for twisted full toroidal Lie algebra*, Journal of Algebra
- [8] James E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Graduate Texts in Mathematics, vol. 9, Springer- Verlag, New York-Berlin, 1978, second printing, revised.

- [9] Haisheng Li, On certain categories of modules for affine Lie algebras, *Math. Z.* 248 (3) (2004) 635-664.
- [10] Erhard Neher, Alistair Savage, Prasad Senesi, Irreducible finite-dimensional representation of equivariant map algebras, *Trans. Am. Math. soc.* 364(5)(2012) 2619-2646.
- [11] Yuly Billig, John Talboom, Classification of category \mathcal{J} modules for divergence zero vector fields on torus, *J. Algebra* 500 (2018) 498-516.
- [12] Xiangqian Guo, Genqiang Liu, Jet modules for the centerless Virasoro-like algebra, *J. Algebra Appl.* 18(1) (2019) 1950002.
- [13] Katsuyuki Naoi, Multiloop Lie algebras and the construction of extended affine Lie algebras. *J. Algebra*, 323(8):2103-2129, 2010
- [14] V. Chari, Integrable representations of affine Lie algebras, *Invent. Math.* 85(1986), no. 2, 317-335.
- [15] Chen, Fulin; Li, Zhiqiang; Tan, Shaobin; Classification of integrable representations for toroidal extended affine Lie algebras, *J. Algebra* 514(2021), 1-37.
- [16] Erhard Neher, Extended affine Lie algebras, *C. R. Math. Acad. Sci. Soc. R. Can.* 26 (3) (2004) 90-96.
- [17] Yoji Yoshii, Lie tori-a simple characterization of extended affine Lie algebras, *Publ. Res. Inst. Math. Sci.* 42 (3) (2006) 739-762.
- [18] M. Lau, Representations of multiloop algebras, *Pacific J. Math.* 245 (2010), no. 1, 167-184.