

COHOMOLOGICAL OBSTRUCTIONS AND WEAK CROSSED PRODUCTS OVER WEAK HOPF ALGEBRAS

RAMÓN GONZÁLEZ RODRÍGUEZ AND ANA BELÉN RODRÍGUEZ RAPOSO

ABSTRACT. Let H be a cocommutative weak Hopf algebra and let (B, φ_B) a weak left H -module algebra. In this paper, for a twisted convolution invertible morphism $\sigma : H \otimes H \rightarrow B$ we define its obstruction θ_σ as a degree three Sweedler 3-cocycle with values in the center of B . We obtain that the class of this obstruction vanish in third Sweedler cohomology group $\mathcal{H}_{\varphi_{Z(B)}}^3(H, Z(B))$ if, and only if, there exists a twisted convolution invertible 2-cocycle $\alpha : H \otimes H \rightarrow B$ such that $H \otimes B$ can be endowed with a weak crossed product structure with α keeping a cohomological-like relation with σ . Then, as a consequence, the class of the obstruction of σ vanish if, and only if, there exists a cleft extension of B by H .

KEYWORDS: Weak Hopf algebra, Sweedler cohomology, weak crossed products, cleft extension, obstruction.

MSC2020: 18M05, 16T05, 20J06.

INTRODUCTION

Crossed products of a Hopf algebra by an algebra have been widely studied in relation to extensions of algebras, generalizing classical results of semidirect products and extensions of groups, together with a generalization of group cohomology to the Hopf algebra setting. In [17] Sweedler defines the so-called Sweedler's cohomology for a cocommutative Hopf algebra H and a commutative H -module algebra B . In this paper he also shows that any cleft H -extension of algebras $B \hookrightarrow A$ (that is, roughly speaking, a split extension) can be realized as a crossed product given in terms of the action of H on B and a 2-cocycle $\sigma : H \otimes H \rightarrow B$. Moreover, cleft extensions of B are classified by the second cohomology group $\mathcal{H}^2(H, B)$. Several generalizations of these results were carried out by Doi and Takeuchi [7], Blattner, Cohen and Montgomery [4] and Blattner and Montgomery [5] by dropping out the conditions of cocommutativity and commutativity, and the associativity of the action $\varphi_B : H \otimes B \rightarrow B$ and thus, the use of Sweedler's cohomology. However some of its formalism is preserved: for an arbitrary Hopf algebra H and an arbitrary algebra B , a crossed product is given in terms of a measuring $\varphi_B : H \otimes B \rightarrow B$ and a formal 2-cocycle $\sigma : H \otimes H \rightarrow B$ that must also satisfy the twisted condition needed to substitute the associativity of φ_B . Moreover, two such crossed products are equivalent if the cocycles satisfy a cohomological-like equivalence. This last result was interpreted in an actual cohomological setting by Doi in [8], where he shows that cleft extensions of an algebra B by a cocommutative Hopf algebra H with the same action are classified by $\mathcal{H}^2(H, Z(B))$, where $Z(B)$ denotes the center of B . All these

results can be interpreted in a symmetric monoidal category with base object K (see for example [1] and [11] for cleft extensions in a monoidal setting).

The next objective became to decide when an algebra B admits a cleft extension by H . Following some classical results of obstructions to extensions of groups (see, for example, [13]), Schauenburg finds in [16] a relation between the third Sweedler's cohomology group $\mathcal{H}^3(H, Z(B))$ and cleft extensions, where $Z(B)$ denotes the center of B . For a measuring φ_B and a twisted morphism σ , he generalizes the notion of obstruction as Sweedler three cocycle θ_σ on H with values on the center of B and shows that the class $[\theta_\sigma] \in \mathcal{H}^3(H, Z(B))$ vanish if, and only if, φ_B and σ give rise to a crossed product on $H \otimes B$ and, at last, to a cleft extension.

With the apparition of weak Hopf algebras as generalizations of groupoid algebras (see [6]) all the theory of cleft extensions, Sweedler's cohomology and crossed products needed a deep review. Recall that the main point of an algebra-coalgebra H to be weak is that its unit does not need to be coassociative, nor its counit associative. These apparently innocent generalizations conceptually imply the existence of two objects, different from the base object K in the ground monoidal category when H is actually weak, that somehow will play the role of K . From a practical point of view, this lack of (co)associativity of the (co)unit forces to a change in the definition of regular morphisms, and thus to a change in the tackling of cleft extensions, cohomological interpretations of crossed products and a rethinking of cohomology and crossed products themselves. For the cocommutative case these problems were successfully solved in [2] and [3], where the authors explore the meaning of cleft extension and weak crossed product for a cocommutative weak Hopf algebra H weakly acting on an algebra B , and define Sweedler's cohomology in weak contexts. In order to achieve these objectives, they consider the unit in $\text{Reg}(H, B)$ as $\varphi_B \circ (H \otimes \eta_B)$ (and thus, regular morphisms depend on φ_B and we denote the set by $\text{Reg}_{\varphi_B}(H, B)$), where φ_B is the weak action of H on B , and η_B is the unit of B . Moreover, to study weak crossed products they consider a preunit instead of a unit, so they obtain an algebra as a subobject of $H \otimes B$, whose product is given in terms of φ_B and a twisted formal 2-cocycle $\sigma : H \otimes H \rightarrow B$. In such terms, they are able to define a cohomology theory for a cocommutative weak Hopf algebra H and a commutative H -module algebra B . Moreover, they identify cleft H -extensions of a weak H -module algebra B with products with convolution invertible 2-cocycle and classify them by $\mathcal{H}_{\varphi_B}^2(H, Z(B))$, this is, the second cohomology group. The relation of weak crossed products and cleft extensions for the non-cocommutative case was also studied in [12] by Guccione, Guccione and Valqui.

So once we have the proper concepts of cleft extensions, weak crossed products and Sweedler's cohomology for the weak setting, we just need to find out the role of obstructions in relation to cleft extensions and their cohomological meaning, and these are the main objectives of the present paper. In order to attain such objectives, we first make a wide review of weak crossed products, and we find that we just need a measuring $\varphi_B : H \otimes B \rightarrow B$ together with a twisted morphism $\sigma : H \otimes H \rightarrow B$ that does not need to be convolution invertible but a formal 2-cocycle to define a weak crossed product on $H \otimes B$. Moreover we obtain necessary and sufficient conditions for weak crossed products to be equivalent that, in particular, are given in terms of morphisms in $\text{Reg}_{\varphi_B}(H, B)$. We finally use these results in the particular case of a cocommutative weak Hopf algebra H and a weak H -module algebra (B, φ_B) .

We consider a twisted convolution invertible morphism $\sigma : H \otimes H \rightarrow B$ and define its cohomological obstruction θ_σ through the center of B . We obtain that this obstruction vanish in $\mathcal{H}_{\varphi_{Z(B)}}^3(H, Z(B))$ if, and only if, there exists a twisted convolution invertible 2-cocycle $\alpha : H \otimes H \rightarrow B$ such that $H \otimes B$ can be endowed with a weak crossed product structure with α keeping a cohomological-like relation with σ . This result means, in terms of cleft extensions, that if (B, φ_B) is a weak H -module algebra with $\sigma : H \otimes H \rightarrow B$ twisted and convolution invertible then its obstruction vanish if, and only if, there exists a cleft extension of B by H .

Throughout this paper \mathbf{C} denotes a strict symmetric monoidal category with tensor product \otimes , unit object K and symmetry isomorphism c . There is no loss of generality in assuming that \mathbf{C} is strict because every monoidal category is monoidally equivalent to a strict one. Then, we may work as if the constraints were all identities. We also assume that in \mathbf{C} every idempotent morphism splits, i.e., for any morphism $q : M \rightarrow M$ such that $q \circ q = q$ there exists an object N , called the image of q , and morphisms $i : N \rightarrow M$, $p : M \rightarrow N$ such that $q = i \circ p$ and $p \circ i = id_N$ (id_N denotes the identity morphism for N). The morphisms p and i will be called a factorization of q . Note that N , p and i are unique up to isomorphism. The categories satisfying this property constitute a broad class that includes, among others, the categories with epi-monic decomposition for morphisms and categories with (co)equalizers. Finally, given objects M , N , P and a morphism $f : N \rightarrow P$, we write $M \otimes f$ for $id_M \otimes f$ and $f \otimes M$ for $f \otimes id_M$.

An algebra in \mathbf{C} is a triple $A = (A, \eta_A, \mu_A)$, where A is an object in \mathbf{C} and $\eta_A : K \rightarrow A$ (unit), $\mu_A : A \otimes A \rightarrow A$ (product) are morphisms in \mathbf{C} such that $\mu_A \circ (A \otimes \eta_A) = id_A = \mu_A \circ (\eta_A \otimes A)$, $\mu_A \circ (A \otimes \mu_A) = \mu_A \circ (\mu_A \otimes A)$. We say that an algebra A is commutative if $\mu_A = \mu_A \circ c_{A,A}$.

Given two algebras $A = (A, \eta_A, \mu_A)$ and $B = (B, \eta_B, \mu_B)$, a morphism $f : A \rightarrow B$ in \mathbf{C} is an algebra morphism if $\mu_B \circ (f \otimes f) = f \circ \mu_A$ and $f \circ \eta_A = \eta_B$.

Also, if A , B are algebras in \mathbf{C} , the object $A \otimes B$ is an algebra in \mathbf{C} , where $\eta_{A \otimes B} = \eta_A \otimes \eta_B$ and $\mu_{A \otimes B} = (\mu_A \otimes \mu_B) \circ (A \otimes c_{B,A} \otimes B)$. Note that, if A and B are commutative algebras, so is $A \otimes B$.

A coalgebra in \mathbf{C} is a triple $D = (D, \varepsilon_D, \delta_D)$, where D is an object in \mathbf{C} and $\varepsilon_D : D \rightarrow K$ (counit), $\delta_D : D \rightarrow D \otimes D$ (coproduct) are morphisms in \mathbf{C} such that $(\varepsilon_D \otimes D) \circ \delta_D = id_D = (D \otimes \varepsilon_D) \circ \delta_D$, $(\delta_D \otimes D) \circ \delta_D = (D \otimes \delta_D) \circ \delta_D$. We say that a coalgebra D is cocommutative if $\delta_D = c_{D,D} \circ \delta_D$.

If $D = (D, \varepsilon_D, \delta_D)$ and $E = (E, \varepsilon_E, \delta_E)$ are coalgebras, a morphism $f : D \rightarrow E$ in \mathbf{C} is a coalgebra morphism if $(f \otimes f) \circ \delta_D = \delta_E \circ f$ and $\varepsilon_E \circ f = \varepsilon_D$.

If D , E are coalgebras in \mathbf{C} , the tensor product $D \otimes E$ is a coalgebra in \mathbf{C} , where $\varepsilon_{D \otimes E} = \varepsilon_D \otimes \varepsilon_E$ and $\delta_{D \otimes E} = (D \otimes c_{D,E} \otimes E) \circ (\delta_D \otimes \delta_E)$. Note that, if D and E are cocommutative coalgebras, so is $D \otimes E$.

Finally, if A is an algebra, C is a coalgebra and $f : C \rightarrow A$, $g : C \rightarrow A$ are morphisms in \mathbf{C} , we define the convolution product by $f * g = \mu_A \circ (f \otimes g) \circ \delta_C$.

1. GENERALITIES ON MEASURINGS AND CROSSED PRODUCTS IN A WEAK SETTING

In this section we resume some basic facts about the general theory of weak crossed products in \mathbf{C} , introduced in [9], particularized for measurings over a weak Hopf algebra H . Firstly, we recall the notion

of weak Hopf algebra, introduced in [6], and summarize some basic properties of these algebraic objects in a monoidal setting.

Definition 1.1. A weak bialgebra H is an object in \mathcal{C} with an algebra structure (H, η_H, μ_H) and a coalgebra structure $(H, \varepsilon_H, \delta_H)$ such that the following axioms hold:

- (a1) $\delta_H \circ \mu_H = (\mu_H \otimes \mu_H) \circ \delta_{H \otimes H}$,
- (a2) $\varepsilon_H \circ \mu_H \circ (\mu_H \otimes H) = (\varepsilon_H \otimes \varepsilon_H) \circ (\mu_H \otimes \mu_H) \circ (H \otimes \delta_H \otimes H)$
 $= (\varepsilon_H \otimes \varepsilon_H) \circ (\mu_H \otimes \mu_H) \circ (H \otimes (c_{H,H} \circ \delta_H) \otimes H)$,
- (a3) $(\delta_H \otimes H) \circ \delta_H \circ \eta_H = (H \otimes \mu_H \otimes H) \circ (\delta_H \otimes \delta_H) \circ (\eta_H \otimes \eta_H)$
 $= (H \otimes (\mu_H \circ c_{H,H}) \otimes H) \circ (\delta_H \otimes \delta_H) \circ (\eta_H \otimes \eta_H)$.

Moreover, if there exists a morphism $\lambda_H : H \rightarrow H$ in \mathcal{C} (called the antipode of H) satisfying

- (a4) $id_H * \lambda_H = ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H)$,
- (a5) $\lambda_H * id_H = (H \otimes (\varepsilon_H \circ \mu_H)) \circ (c_{H,H} \otimes H) \circ (H \otimes (\delta_H \circ \eta_H))$,
- (a6) $\lambda_H * id_H * \lambda_H = \lambda_H$,

we will say that the weak bialgebra is a weak Hopf algebra.

We say that H is commutative, if it is commutative as algebra and we say that H is cocommutative if it is cocommutative as coalgebra.

1.2. Let H be a weak biagebra. For $n \geq 1$, we denote by $H^{\otimes n}$ the n -fold tensor power $H \otimes \cdots \otimes H$. By $H^{\otimes 0}$ we denote the unit object of \mathcal{C} , i.e., $H^{\otimes 0} = K$. If $n \geq 2$, $m_H^{\otimes n}$ denotes the morphism $m_H^{\otimes n} : H^{\otimes n} \rightarrow H$ defined by $m_H^{\otimes 2} = \mu_H$ and by $m_H^{\otimes 3} = m_H^{\otimes 2} \circ (H \otimes \mu_H)$, \dots , $m_H^{\otimes k} = m_H^{\otimes (k-1)} \circ (H^{\otimes k-2} \otimes \mu_H)$ for $k > 2$. On the other hand, with $\delta_{H^{\otimes n}}$ we denote the coproduct defined in the coalgebra $H^{\otimes n}$. Then by the coassociativity of δ_H and the naturality of c , for $k = 1, \dots, n-1$,

$$\delta_{H^{\otimes n}} = \delta_{H^{\otimes (n-k)} \otimes H^{\otimes k}}$$

holds. By [2, Proposition 2.10] we have that

$$\delta_H \circ m_H^{\otimes n} = (m_H^{\otimes n} \otimes m_H^{\otimes n}) \circ \delta_{H^{\otimes n}}.$$

Finally, note that, if H is cocommutative, then so is $H^{\otimes n}$.

1.3. For any weak bialgebra, if we define the morphisms Π_H^L (target), Π_H^R (source), $\overline{\Pi}_H^L$ and $\overline{\Pi}_H^R$ by

$$\begin{aligned} \Pi_H^L &= ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H), \\ \Pi_H^R &= (H \otimes (\varepsilon_H \circ \mu_H)) \circ (c_{H,H} \otimes H) \circ (H \otimes (\delta_H \circ \eta_H)), \\ \overline{\Pi}_H^L &= (H \otimes (\varepsilon_H \circ \mu_H)) \circ ((\delta_H \circ \eta_H) \otimes H), \\ \overline{\Pi}_H^R &= ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes (\delta_H \circ \eta_H)), \end{aligned}$$

it is straightforward to show that they are idempotent and the equalities

$$(1) \quad \Pi_H^L \circ \overline{\Pi}_H^L = \Pi_H^L, \quad \Pi_H^L \circ \overline{\Pi}_H^R = \overline{\Pi}_H^R, \quad \Pi_H^R \circ \overline{\Pi}_H^L = \overline{\Pi}_H^L, \quad \Pi_H^R \circ \overline{\Pi}_H^R = \Pi_H^R,$$

$$(2) \quad \overline{\Pi}_H^L \circ \Pi_H^L = \overline{\Pi}_H^L, \quad \overline{\Pi}_H^L \circ \Pi_H^R = \Pi_H^R, \quad \overline{\Pi}_H^R \circ \Pi_H^L = \Pi_H^L, \quad \overline{\Pi}_H^R \circ \Pi_H^R = \overline{\Pi}_H^R,$$

hold.

On the other hand, denote by H_L the image of the target morphism Π_H^L and let $p_H^L : H \rightarrow H_L$, $i_H^L : H_L \rightarrow H$ be the morphisms such that $i_H^L \circ p_H^L = \Pi_H^L$ and $p_H^L \circ i_H^L = id_{H_L}$. Then

$$(H_L, \eta_{H_L} = p_H^L \circ \eta_H, \mu_{H_L} = p_H^L \circ \mu_H \circ (i_H^L \otimes i_H^L))$$

is an algebra and

$$(H_L, \varepsilon_{H_L} = \varepsilon_H \circ i_H^L, \delta_{H_L} = (p_H^L \otimes p_H^L) \circ \delta_H \circ i_H^L)$$

is a coalgebra. The morphisms η_{H_L} , μ_{H_L} , ε_{H_L} and δ_{H_L} are the unique morphisms satisfying

$$(3) \quad i_H^L \circ \eta_{H_L} = \eta_H, \quad i_H^L \circ \mu_{H_L} = \mu_H \circ (i_H^L \otimes i_H^L),$$

$$(4) \quad \varepsilon_{H_L} \circ p_H^L = \varepsilon_H, \quad \delta_{H_L} \circ p_H^L = (p_H^L \otimes p_H^L) \circ \delta_H$$

respectively.

For the morphisms target and source we have the following identities:

$$(5) \quad \Pi_H^L \circ \mu_H \circ (H \otimes \Pi_H^L) = \Pi_H^L \circ \mu_H, \quad \Pi_H^R \circ \mu_H \circ (\Pi_H^R \otimes H) = \Pi_H^R \circ \mu_H,$$

$$(6) \quad \overline{\Pi}_H^L \circ \mu_H \circ (H \otimes \overline{\Pi}_H^L) = \overline{\Pi}_H^L \circ \mu_H, \quad \overline{\Pi}_H^R \circ \mu_H \circ (\overline{\Pi}_H^R \otimes H) = \overline{\Pi}_H^R \circ \mu_H,$$

$$(7) \quad (H \otimes \Pi_H^L) \circ \delta_H \circ \Pi_H^L = \delta_H \circ \Pi_H^L, \quad (\Pi_H^R \otimes H) \circ \delta_H \circ \Pi_H^R = \delta_H \circ \Pi_H^R,$$

$$(8) \quad (H \otimes \overline{\Pi}_H^R) \circ \delta_H \circ \overline{\Pi}_H^R = \delta_H \circ \overline{\Pi}_H^R, \quad (\overline{\Pi}_H^L \otimes H) \circ \delta_H \circ \overline{\Pi}_H^L = \delta_H \circ \overline{\Pi}_H^L,$$

$$(9) \quad \mu_H \circ (H \otimes \Pi_H^L) = ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes c_{H,H}) \circ (\delta_H \otimes H),$$

$$(10) \quad (H \otimes \Pi_H^L) \circ \delta_H = (\mu_H \otimes H) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H),$$

$$(11) \quad \mu_H \circ (\Pi_H^R \otimes H) = (H \otimes (\varepsilon_H \circ \mu_H)) \circ (c_{H,H} \otimes H) \circ (H \otimes \delta_H)$$

$$(12) \quad (\Pi_H^R \otimes H) \circ \delta_H = (H \otimes \mu_H) \circ (c_{H,H} \otimes H) \circ (H \otimes (\delta_H \circ \eta_H))$$

$$(13) \quad \mu_H \circ (\overline{\Pi}_H^R \otimes H) = ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes \delta_H),$$

$$(14) \quad \mu_H \circ (H \otimes \overline{\Pi}_H^L) = (H \otimes (\varepsilon_H \circ \mu_H)) \circ (\delta_H \otimes H),$$

$$(15) \quad (\overline{\Pi}_H^L \otimes H) \circ \delta_H = (H \otimes \mu_H) \circ ((\delta_H \circ \eta_H) \otimes H),$$

$$(16) \quad (H \otimes \overline{\Pi}_H^R) \circ \delta_H = (\mu_H \otimes H) \circ (H \otimes (\delta_H \circ \eta_H)),$$

$$(17) \quad \begin{aligned} \delta_H \circ \eta_H &= (\Pi_H^R \otimes H) \circ \delta_H \circ \eta_H = (H \otimes \Pi_H^L) \circ \delta_H \circ \eta_H = (H \otimes \overline{\Pi}_H^R) \circ \delta_H \circ \eta_H \\ &= (\overline{\Pi}_H^L \otimes H) \circ \delta_H \circ \eta_H, \end{aligned}$$

$$(18) \quad \begin{aligned} \varepsilon_H \circ \mu_H &= \varepsilon_H \circ \mu_H \circ (\Pi_H^R \otimes H) = \varepsilon_H \circ \mu_H \circ (H \otimes \Pi_H^L) = \varepsilon_H \circ \mu_H \circ (\overline{\Pi}_H^R \otimes H) \\ &= \varepsilon_H \circ \mu_H \circ (H \otimes \overline{\Pi}_H^L). \end{aligned}$$

If H is a weak Hopf algebra in \mathbb{C} , the antipode λ_H is unique, antimultiplicative, anticomultiplicative and leaves the unit and the counit invariant, i.e.:

$$(19) \quad \lambda_H \circ \mu_H = \mu_H \circ (\lambda_H \otimes \lambda_H) \circ c_{H,H}, \quad \delta_H \circ \lambda_H = c_{H,H} \circ (\lambda_H \otimes \lambda_H) \circ \delta_H,$$

$$(20) \quad \lambda_H \circ \eta_H = \eta_H, \quad \varepsilon_H \circ \lambda_H = \varepsilon_H.$$

Also, it is straightforward to show that Π_H^L, Π_H^R satisfy the equalities

$$(21) \quad \Pi_H^L = id_H * \lambda_H, \quad \Pi_H^R = \lambda_H * id_H, \quad \Pi_H^L * id_H = id_H = id_H * \Pi_H^R,$$

$$\Pi_H^R * \lambda_H = \lambda_H = \lambda_H * \Pi_H^L,$$

$$(22) \quad \Pi_H^L * \Pi_H^L = \Pi_H^L, \quad \Pi_H^R * \Pi_H^R = \Pi_H^R$$

and

$$(23) \quad \Pi_H^L = \lambda_H \circ \bar{\Pi}_H^L = \bar{\Pi}_H^R \circ \lambda_H, \quad \Pi_H^R = \bar{\Pi}_H^L \circ \lambda_H = \lambda_H \circ \bar{\Pi}_H^R.$$

Finally we also have

$$(24) \quad \mu_H \circ c_{H,H} \circ (H \otimes \bar{\Pi}_H^L) \circ \delta_H = id_H = \mu_H \circ c_{H,H} \circ (\bar{\Pi}_H^R \otimes H) \circ \delta_H.$$

Now we recall the notions of measuring, left weak H -module algebra, and left H -module algebra.

Definition 1.4. Let H be a weak Hopf algebra and let B be an algebra. We say that the morphism $\varphi_B : H \otimes B \rightarrow B$ is a measuring if

$$(b1) \quad \varphi_B \circ (H \otimes \mu_B) = \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes B \otimes B).$$

Set $u_1^{\varphi_B} = \varphi_B \circ (H \otimes \eta_B)$. If φ_B is a measuring satisfying

$$(b2) \quad \varphi_B \circ (\eta_H \otimes B) = id_B,$$

$$(b3) \quad u_1^{\varphi_B} \circ \mu_H = \varphi_B \circ (H \otimes u_1^{\varphi_B}),$$

we will say that (B, φ_B) is a left weak H -module algebra. If we replace (b3) by

$$\varphi_B \circ (\mu_H \otimes B) = \varphi_B \circ (H \otimes \varphi_B)$$

we will say that (B, φ_B) is a left H -module algebra.

If (B, φ_B) is a left weak H -module algebra the following equivalent conditions are satisfied:

$$(25) \quad \varphi_B \circ (\Pi_H^L \otimes B) = \mu_B \circ (u_1^{\varphi_B} \otimes B),$$

$$(26) \quad \varphi_B \circ (\bar{\Pi}_H^L \otimes B) = \mu_B \circ c_{B,B} \circ (u_1^{\varphi_B} \otimes B),$$

$$(27) \quad u_1^{\varphi_B} \circ \Pi_H^L = u_1^{\varphi_B},$$

$$(28) \quad u_1^{\varphi_B} \circ \bar{\Pi}_H^L = u_1^{\varphi_B},$$

$$(29) \quad u_1^{\varphi_B} \circ \mu_H = u_1^{\varphi_B} \circ \mu_H \circ (H \otimes \bar{\Pi}_H^L),$$

$$(30) \quad u_1^{\varphi_B} \circ \mu_H = u_1^{\varphi_B} \circ \mu_H \circ (H \otimes \Pi_H^L).$$

Let φ_B be a measuring and $n \geq 1$. With $\varphi_B^{\otimes n}$ we will denote the morphism

$$\varphi_B^{\otimes n} : H^{\otimes n} \otimes B \rightarrow B$$

defined as $\varphi_B^{\otimes 1} = \varphi_B$ and $\varphi_B^{\otimes n} = \varphi_B \circ (H \otimes \varphi_B^{\otimes(n-1)})$.

Note that, by (b1) of Definition 1.4, proceeding by induction, it is easy to show that

$$(31) \quad \varphi_B^{\otimes n} \circ (H^{\otimes n} \otimes \mu_B) = \mu_B \circ (\varphi_B^{\otimes n} \otimes \varphi_B^{\otimes n}) \circ (H^{\otimes n} \otimes c_{H^{\otimes n}, B} \otimes B) \circ (\delta_{H^{\otimes n}} \otimes B \otimes B)$$

holds.

For $n \geq 2$ we define the morphism

$$(32) \quad u_n^{\varphi_B} = \varphi_B \circ (m_H^{\otimes n} \otimes \eta_B) = \varphi_B^{\otimes(n-1)} \circ (H^{\otimes(n-1)} \otimes u_1^{\varphi_B}).$$

Note that

$$(33) \quad u_n^{\varphi_B} = u_1^{\varphi_B} \circ m_H^{\otimes n}$$

and, by [2, Proposition 2.11], we have

$$(34) \quad u_n^{\varphi_B} * u_n^{\varphi_B} = u_n^{\varphi_B}.$$

Definition 1.5. Let H be a weak Hopf algebra and let B be an algebra. For a measuring φ_B and any morphism $\sigma : H \otimes H \rightarrow B$, we define the morphisms

$$P_{\varphi_B} : H \otimes B \rightarrow B \otimes H, \quad F_\sigma : H \otimes H \rightarrow B \otimes H, \quad G_\sigma : H \otimes H \rightarrow H \otimes B,$$

by

$$(35) \quad P_{\varphi_B} = (\varphi_B \otimes H) \circ (H \otimes c_{H, B}) \circ (\delta_H \otimes B),$$

$$(36) \quad F_\sigma = (\sigma \otimes \mu_H) \circ \delta_{H^{\otimes 2}}$$

and

$$(37) \quad G_\sigma = (\mu_H \otimes \sigma) \circ \delta_{H^{\otimes 2}}.$$

By [2, Proposition 3.3] and some easy computations we have the following result.

Proposition 1.6. *Let H be a weak Hopf algebra and let $\varphi_B : H \otimes B \rightarrow B$ be a measuring. The morphism P_{φ_B} defined in (35) satisfies*

$$(38) \quad (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (P_{\varphi_B} \otimes B) = P_{\varphi_B} \circ (H \otimes \mu_B).$$

The morphisms $\nabla_{B \otimes H}^{\varphi_B} : B \otimes H \rightarrow B \otimes H$ and $\nabla_{H \otimes B}^{\varphi_B} : H \otimes B \rightarrow H \otimes B$ defined by

$$\nabla_{B \otimes H}^{\varphi_B} = (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (B \otimes H \otimes \eta_B)$$

and

$$\nabla_{H \otimes B}^{\varphi_B} = (H \otimes \mu_B) \circ (((H \otimes \varphi_A) \circ (\delta_H \otimes \eta_A)) \otimes A)$$

are idempotent. Also, we have the following identities:

$$(39) \quad P_{\varphi_B} \circ (H \otimes \eta_B) = (u_1^{\varphi_B} \otimes H) \circ \delta_H,$$

$$(40) \quad (B \otimes \varepsilon_H) \circ P_{\varphi_B} = \varphi_B,$$

$$(41) \quad \nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B} = P_{\varphi_B},$$

$$(42) \quad \nabla_{B \otimes H}^{\varphi_B} = (\mu_B \otimes H) \circ (B \otimes ((u_1^{\varphi_B} \otimes H) \circ \delta_H)),$$

$$(43) \quad \nabla_{H \otimes B}^{\varphi_B} = (H \otimes \mu_B) \circ (((H \otimes u_1^{\varphi_B}) \circ \delta_H) \otimes B),$$

$$(44) \quad \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H) = (u_1^{\varphi_B} \otimes H) \circ \delta_H,$$

$$(45) \quad \nabla_{H \otimes B}^{\varphi_B} \circ (H \otimes \eta_B) = (H \otimes u_1^{\varphi_B}) \circ \delta_H,$$

$$(46) \quad (\mu_B \otimes H) \circ (B \otimes \nabla_{B \otimes H}^{\varphi_B}) = (B \otimes \nabla_{B \otimes H}^{\varphi_B}) \circ (\mu_B \otimes H),$$

$$(47) \quad (H \otimes \mu_B) \circ (\nabla_{H \otimes B}^{\varphi_B} \otimes B) = (\nabla_{H \otimes B}^{\varphi_B} \otimes B) \circ (H \otimes \mu_B),$$

$$(48) \quad (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} = \mu_B \circ (B \otimes u_1^{\varphi_B}),$$

$$(49) \quad (\varepsilon_H \otimes B) \circ \nabla_{H \otimes B}^{\varphi_B} = \mu_B \circ (u_1^{\varphi_B} \otimes B),$$

$$(50) \quad (B \otimes \delta_H) \circ \nabla_{B \otimes H}^{\varphi_B} = (\nabla_{B \otimes H}^{\varphi_B} \otimes H) \circ (B \otimes \delta_H),$$

$$(51) \quad (\delta_H \otimes B) \circ \nabla_{H \otimes B}^{\varphi_B} = (H \otimes \nabla_{H \otimes B}^{\varphi_B}) \circ (\delta_H \otimes B),$$

$$(52) \quad \mu_B \circ (u_1^{\varphi_B} \otimes \varphi_B) \circ (\delta_H \otimes B) = \varphi_B,$$

$$(53) \quad (\mu_B \otimes H) \circ (u_1^{\varphi_B} \otimes P_{\varphi_B}) \circ (\delta_H \otimes B) = P_{\varphi_B},$$

$$(54) \quad (B \otimes \varepsilon_H) \circ P_{\varphi_B} \circ (H \otimes \eta_B) = u_1^{\varphi_B},$$

$$(55) \quad (\mu_B \otimes H) \circ (u_1^{\varphi_B} \otimes c_{H,B}) \circ (\delta_H \otimes B) = (\mu_B \otimes H) \circ (H \otimes c_{B,B}) \circ ((P_{\varphi_B} \circ (H \otimes \eta_B)) \otimes B).$$

On the other hand, by a similar proof to the one used in [2, Proposition 3.4] we have the following proposition.

Proposition 1.7. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring and let $\sigma : H \otimes H \rightarrow B$ be a morphism. The morphisms F_σ and G_σ defined in (36) and (37) satisfy the identities:*

$$(56) \quad (B \otimes \delta_H) \circ F_\sigma = (F_\sigma \otimes \mu_H) \circ \delta_{H \otimes 2},$$

$$(57) \quad (\delta_H \otimes B) \circ G_\sigma = (\mu_H \otimes G_\sigma) \circ \delta_{H \otimes 2}.$$

Moreover, we also have the proposition:

Proposition 1.8. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring. The equality*

$$(58) \quad \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ P_{\varphi_B} = \varphi_B,$$

holds.

*Let $\sigma : H \otimes H \rightarrow B$ be a morphism and $u_2^{\varphi_B}$ be the morphism defined in (32). If $\sigma * u_2^{\varphi_B} = \sigma$, the equality*

$$(59) \quad \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ F_\sigma = \sigma,$$

holds and, as a consequence, we have the following identities:

$$(60) \quad \nabla_{B \otimes H}^{\varphi_B} \circ F_\sigma = F_\sigma,$$

$$(61) \quad (B \otimes \varepsilon_H) \circ F_\sigma = \sigma.$$

*Moreover, if $\sigma : H \otimes H \rightarrow B$ is a morphism satisfying (60) and (61), we have that $\sigma * u_2^{\varphi_B} = \sigma$.*

Proof. Note that, (58) holds because

$$\mu_B \circ (B \otimes u_1^{\varphi_B}) \circ P_{\varphi_B} \stackrel{(48)}{=} (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B} \stackrel{(41)}{=} (B \otimes \varepsilon_H) \circ P_{\varphi_B} \stackrel{(40)}{=} \varphi_B.$$

Trivially,

$$(62) \quad \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ F_\sigma = \sigma * u_2^{\varphi_B}$$

and then we obtain (59). On the other hand,

$$\nabla_{B \otimes H}^{\varphi_B} \circ F_\sigma \stackrel{(56)}{=} ((\mu_B \circ (B \otimes u_1^{\varphi_B}) \circ F_\sigma) \otimes \mu_H) \circ \delta_{H \otimes 2} \stackrel{(59)}{=} F_\sigma.$$

Therefore,

$$(B \otimes \varepsilon_H) \circ F_\sigma \stackrel{(60)}{=} (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ F_\sigma \stackrel{(48)}{=} \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ F_\sigma \stackrel{(59)}{=} \sigma.$$

Finally, let $\sigma : H \otimes H \rightarrow B$ be a morphism satisfying (60) and (61). Then,

$$\sigma \stackrel{(61)}{=} (B \otimes \varepsilon_H) \circ F_\sigma \stackrel{(60)}{=} (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ F_\sigma \stackrel{(48)}{=} \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ F_\sigma \stackrel{(62)}{=} \sigma * u_2^{\varphi_B}.$$

□

1.9. Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring and let $\sigma : H \otimes H \rightarrow B$ be a morphism. In a similar way to what was proven in the previous proposition, we can ensure that the equality

$$(63) \quad \mu_B \circ (u_1^{\varphi_B} \otimes B) \circ G_\sigma = u_2^{\varphi_B} * \sigma$$

holds. As a consequence we can obtain the following result:

Proposition 1.10. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring and let $\sigma : H \otimes H \rightarrow B$ be a morphism. Let $u_2^{\varphi_B}$ be the morphism defined in (32). If $u_2^{\varphi_B} * \sigma = \sigma$, the equality*

$$(64) \quad \mu_B \circ (u_1^{\varphi_B} \otimes B) \circ G_\sigma = \sigma,$$

holds and, as a consequence, we have the following identities:

$$(65) \quad \nabla_{H \otimes B}^{\varphi_B} \circ G_\sigma = G_\sigma,$$

$$(66) \quad (\varepsilon_H \otimes B) \circ G_\sigma = \sigma.$$

*Moreover, if $\sigma : H \otimes H \rightarrow B$ is a morphism satisfying (65) and (66), we have that $u_2^{\varphi_B} * \sigma = \sigma$.*

Remark 1.11. By the previous propositions, [12, Propositions 2.7 and 2.8] and (1), (2), if $\sigma : H \otimes H \rightarrow B$ satisfies that $\sigma * u_2^{\varphi_B} = \sigma$, we have

$$(67) \quad \sigma \circ (\mu_H \otimes H) \circ (H \otimes \Pi_H^R \otimes H) = \sigma \circ (H \otimes \mu_H) \circ (H \otimes \Pi_H^R \otimes H),$$

$$(68) \quad \sigma \circ (\mu_H \otimes H) \circ (H \otimes \bar{\Pi}_H^L \otimes H) = \sigma \circ (H \otimes \mu_H) \circ (H \otimes \bar{\Pi}_H^L \otimes H)$$

and, if $u_2^{\varphi_B} * \sigma = \sigma$, the equalities

$$(69) \quad \sigma \circ (\mu_H \otimes H) \circ (H \otimes \Pi_H^L \otimes H) = \sigma \circ (H \otimes \mu_H) \circ (H \otimes \Pi_H^L \otimes H),$$

$$(70) \quad \sigma \circ (\mu_H \otimes H) \circ (H \otimes \bar{\Pi}_H^R \otimes H) = \sigma \circ (H \otimes \mu_H) \circ (H \otimes \bar{\Pi}_H^R \otimes H)$$

hold.

1.12. Let H be a weak Hopf algebra and let B be an algebra. By the previous results, if $\varphi_B : H \otimes B \rightarrow B$ is a measuring, and $\sigma : H \otimes H \rightarrow B$ is a morphism such that $\sigma * u_2^{\varphi_B} = \sigma$, we have a quadruple

$$\mathbb{B}_H = (B, H, \psi_H^B = P_{\varphi_B}, \sigma_H^B = F_\sigma)$$

as the ones introduced in [9] to define the notion of weak crossed product. For the quadruple \mathbb{B}_H there exists a product in $B \otimes H$ defined by

$$\mu_{B \otimes_{\varphi_B} H} = (\mu_B \otimes H) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H)$$

and let $\mu_{A \times_{\varphi_B} H}$ be the product

$$\mu_{B \times_{\varphi_B} H} = p_{B \otimes H}^{\varphi_B} \circ \mu_{B \otimes H} \circ (i_{B \otimes H}^{\varphi_B} \otimes i_{B \otimes H}^{\varphi_B}),$$

where $B \times_{\varphi_B} H$, $i_{B \otimes H}^{\varphi_B} : B \times_{\varphi_B} H \rightarrow B \otimes H$ and $p_{B \otimes H}^{\varphi_B} : B \otimes H \rightarrow B \times_{\varphi_B} H$ denote the image, the injection, and the projection associated to the factorization of $\nabla_{B \otimes H}^{\varphi_B}$.

Following [9] we say that \mathbb{B}_H satisfies the twisted condition if

$$(71) \quad (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (F_\sigma \otimes B) = (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B})$$

and the cocycle condition holds if

$$(72) \quad (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (F_\sigma \otimes H) = (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes F_\sigma).$$

Note that, if \mathbb{B}_H satisfies the twisted condition, by [9, Proposition 3.4], and (60) we obtain

$$(73) \quad (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes \nabla_{B \otimes H}^{\varphi_B}) = (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H),$$

$$(74) \quad (\mu_B \otimes H) \circ (A \otimes F_\sigma) \circ (\nabla_{B \otimes H}^{\varphi_B} \otimes H) = (\mu_B \otimes H) \circ (B \otimes F_\sigma).$$

Then, composing with $B \otimes \varepsilon_H$ in (74), and applying (61) we have

$$(75) \quad \mu_B \circ (B \otimes \sigma) \circ (\nabla_{B \otimes H}^{\varphi_B} \otimes H) = \mu_B \circ (B \otimes \sigma).$$

Therefore, if \mathbb{B}_H satisfies the twisted condition, the following equality

$$(76) \quad \mu_B \circ (B \otimes \sigma) \circ ((\nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H)) \otimes H) = \sigma.$$

holds.

Theorem 1.13. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring, and let $\sigma : H \otimes H \rightarrow B$ be a morphism such that $\sigma * u_2^{\varphi_B} = \sigma$. Let \mathbb{B}_H be the associated quadruple. Then, the following assertions hold.*

(i) *The quadruple \mathbb{B}_H satisfies the twisted condition (71) iff σ satisfies the twisted condition*

$$(77) \quad \mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B}) = \mu_B \circ (B \otimes \varphi_B) \circ (F_\sigma \otimes B).$$

(ii) *The quadruple \mathbb{B}_H satisfies the cocycle condition (72) iff σ satisfies the cocycle condition*

$$(78) \quad \mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes F_\sigma) = \mu_B \circ (B \otimes \sigma) \circ (F_\sigma \otimes B).$$

Proof. The proof follows as in [2, Theorems 3.12 and 3.13] because the cocommutativity condition for H is not necessary and (61) holds. □

If the twisted and the cocycle conditions hold, the product $\mu_{B \otimes_{\varphi_B}^{\sigma} H}$ is associative and normalized with respect to $\nabla_{B \otimes H}^{\varphi_B}$, i.e.,

$$(79) \quad \nabla_{B \otimes H}^{\varphi_B} \circ \mu_{B \otimes_{\varphi_B}^{\sigma} H} = \mu_{B \otimes_{\varphi_B}^{\sigma} H} = \mu_{B \otimes_{\varphi_B}^{\sigma} H} \circ (\nabla_{B \otimes H}^{\varphi_B} \otimes \nabla_{B \otimes H}^{\varphi_B})$$

and we have

$$(80) \quad \mu_{B \otimes_{\varphi_B}^{\sigma} H} \circ (\nabla_{B \otimes H}^{\varphi_B} \otimes B \otimes H) = \mu_{B \otimes_{\varphi_B}^{\sigma} H} = \mu_{B \otimes_{\varphi_B}^{\sigma} H} \circ (B \otimes H \otimes \nabla_{B \otimes H}^{\varphi_B}).$$

Then, $\mu_{B \times_{\varphi_B}^{\sigma} H}$ is associative as well (see [9, Proposition 3.7]). Hence we define:

Definition 1.14. If \mathbb{B}_H satisfies (71) and (72) we say that $(B \otimes H, \mu_{B \otimes_{\varphi_B}^{\sigma} H})$ is a weak crossed product.

The next natural question that arises is if it is possible to endow $\mu_{B \times_{\varphi_B}^{\sigma} H}$ with a unit, and hence with an algebra structure. As we recall in [9], we need to use the notion of preunit to obtain this unit. In our setting, if $\mu_{B \otimes_{\varphi_B}^{\sigma} H}$ is an associative product, by [9, Remark 2.4], $\nu : K \rightarrow B \otimes H$ is a preunit if

$$(81) \quad \mu_{B \otimes_{\varphi_B}^{\sigma} H} \circ (B \otimes H \otimes \nu) = \mu_{B \otimes_{\varphi_B}^{\sigma} H} \circ (\nu \otimes B \otimes H), \quad \nu = \mu_{B \otimes_{\varphi_B}^{\sigma} H} \circ (\nu \otimes \nu).$$

By [9, Corollary 3.12], we know that, if ν is a preunit for $(B \otimes H, \mu_{B \otimes_{\varphi_B}^{\sigma} H})$, the object $B \times_{\varphi_B}^{\sigma} H$ is an algebra with product $\mu_{B \times_{\varphi_B}^{\sigma} H}$ and unit $\eta_{B \times_{\varphi_B}^{\sigma} H} = p_{B \otimes H}^{\varphi_B} \circ \nu$.

The following proposition is a tool to establish the conditions under which the morphism $\nu = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$ is a preunit for $\mu_{B \otimes_{\varphi_B} H}$.

Proposition 1.15. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring, and let $\sigma : H \otimes H \rightarrow B$ be a morphism such that $\sigma * u_2^{\varphi_B} = \sigma$. Then, the following equalities hold.*

$$(82) \quad \sigma \circ (\eta_H \otimes H) = \sigma \circ c_{H,H} \circ (H \otimes \overline{\Pi}_H^L) \circ \delta_H,$$

$$(83) \quad \sigma \circ (H \otimes \eta_H) = \sigma \circ (H \otimes \Pi_H^R) \circ \delta_H.$$

Proof. The first equality follows from equalities (68) and (24), and the second one is consequence of (67) and (21). \square

Definition 1.16. Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring, and let $\sigma : H \otimes H \rightarrow B$ be a morphism. We say that σ satisfies the normal condition if

$$(84) \quad \sigma \circ (\eta_H \otimes H) = \sigma \circ (H \otimes \eta_H) = u_1^{\varphi_B}.$$

Therefore, if $\sigma * u_2^{\varphi_B} = \sigma$, by Proposition 1.15, σ is normal if and only if

$$\sigma \circ c_{H,H} \circ (H \otimes \overline{\Pi}_H^L) \circ \delta_H = \sigma \circ (H \otimes \Pi_H^R) \circ \delta_H = u_1^{\varphi_B}.$$

Theorem 1.17. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring such that*

$$(85) \quad \nabla_{B \otimes H}^{\varphi_B} \circ (B \otimes \eta_H) = P_{\varphi_B} \circ (\eta_H \otimes B),$$

*and let $\sigma : H \otimes H \rightarrow B$ be a morphism such that $\sigma * u_2^{\varphi_B} = \sigma$. Let \mathbb{B}_H be the associated quadruple and assume that \mathbb{B}_H satisfies the twisted and the cocycle conditions (71) and (72). Then, $\nu = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$ is a preunit for the weak crossed product associated to \mathbb{B}_H if and only if*

$$(86) \quad F_\sigma \circ (\eta_H \otimes H) = F_\sigma \circ (H \otimes \eta_H) = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H).$$

Proof. By [9, Theorem 3.11] a morphism ν is a preunit for the associated weak crossed if and only if

$$(87) \quad (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes \nu) = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H),$$

$$(88) \quad (\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (\nu \otimes H) = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H)$$

and

$$(89) \quad (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (\nu \otimes B) = (\mu_B \otimes H) \circ (B \otimes \nu)$$

hold. Then the theorem follows because, on the one hand

$$\begin{aligned} & (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (\nu \otimes B) \\ &= (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ ((P_{\varphi_B} \circ (\eta_H \otimes \eta_B)) \otimes B) \text{ (by the unit properties)} \\ &= P_{\varphi_B} \circ (\eta_H \otimes B) \text{ (by (38) and the unit properties)} \\ &= \nabla_{B \otimes H}^{\varphi_B} \circ (B \otimes \eta_H) \text{ (by (85))} \\ &= (\mu_B \otimes H) \circ (B \otimes \nu) \text{ (by the unit properties).} \end{aligned}$$

and, on the other hand, by the unit properties and (71), we have

$$(\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes \nu) = F_\sigma \circ (H \otimes \eta_H).$$

Finally, by (74), the equality

$$(\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (\nu \otimes B) = F_\sigma \circ (\eta_H \otimes H)$$

holds. \square

Remark 1.18. Note that if ν is a preunit for the weak crossed product associated to \mathbb{B}_H , by [9, Theorem 3.11], the equality (89) holds for ν . Then,

$$(90) \quad \nabla_{B \otimes H}^{\varphi_B} \circ \nu = \nu$$

holds. Therefore the preunit of a weak crossed product, if it exists, is unique because if $(B \otimes H, \mu_{B \otimes \varphi_B H})$ admits two preunits ν_1, ν_2 , we have $\eta_{B \otimes \varphi_B H} = p_{B \otimes H}^{\varphi_B} \circ \nu_1 = p_{B \otimes H}^{\varphi_B} \circ \nu_2$ and then

$$\nu_1 = \nabla_{B \otimes H}^{\varphi_B} \circ \nu_1 = \nabla_{B \otimes H}^{\varphi_B} \circ \nu_2 = \nu_2.$$

As a consequence of Theorem 1.17, and by Proposition 1.15 we have:

Corollary 1.19. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring, let $\sigma : H \otimes H \rightarrow B$ be a morphism and let \mathbb{B}_H be the associated quadruple such that the assumptions of Theorem 1.17 hold. Then, $\nu = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$ is a preunit for the weak crossed product associated to \mathbb{B}_H if and only if σ satisfies the normal condition (84).*

Proof. Considering (44), the proof follows from the equalities

$$F_\sigma \circ (\eta_H \otimes H) = ((\sigma \circ c_{H,H} \circ (H \otimes \overline{\Pi}_H^L) \circ \delta_H) \otimes H) \circ \delta_H,$$

and

$$F_\sigma \circ (H \otimes \eta_H) = ((\sigma \circ (H \otimes \Pi_H^R) \circ \delta_H) \otimes H) \circ \delta_H,$$

which hold by (15), (12) and the naturality of c . \square

Therefore, as a consequence of the previous results, we obtain the complete characterization of weak crossed products associated to a measuring.

Corollary 1.20. *Let H be a weak Hopf algebra, let $\varphi_B : H \otimes B \rightarrow B$ be a measuring, let $\sigma : H \otimes H \rightarrow B$ be a morphism and let \mathbb{B}_H be the associated quadruple such that the assumptions of Theorem 1.17 hold. Then the following statements are equivalent:*

- (i) *The product $\mu_{B \otimes \varphi_B H}$ is associative with preunit $\nu = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$ and normalized with respect to $\nabla_{B \otimes H}^{\varphi_B}$.*
- (ii) *The morphism σ satisfies the twisted condition (77), the cocycle condition (78) and the normal condition (84).*

Remark 1.21. Let H be a weak Hopf algebra. If (B, φ_B) is a left weak H -module algebra the equality (85) holds because:

$$\nabla_{B \otimes H}^{\varphi_B} \circ (B \otimes \eta_H)$$

$$\begin{aligned}
&= (\mu_B \otimes H) \circ (B \otimes ((u_1^{\varphi_B} \otimes H) \circ \delta_H \circ \eta_H)) \text{ (by (42))} \\
&= ((\varphi_B \circ c_{B,H}) \otimes H) \circ (B \otimes ((\overline{\Pi}_H^L \otimes H) \circ \delta_H \circ \eta_H)) \text{ (by (26))} \\
&= ((\varphi_B \circ c_{B,H}) \otimes H) \circ (B \otimes (\delta_H \circ \eta_H)) \text{ (by (17))} \\
&= P_{\varphi_B} \circ (\eta_H \otimes B) \text{ (by naturality of } c \text{).}
\end{aligned}$$

Then, if we work with a left weak H -module algebra (B, φ_B) , Corollary 1.20 holds without assuming (85).

2. EQUIVALENT WEAK CROSSED PRODUCTS

The general theory of equivalent weak crossed products was presented in [10]. In this section we remember the criterion obtained in [10] that characterises the equivalence between two weak crossed products and we give the translation of this criterion to the particular setting of weak crossed products induced by measurings.

We shall start by introducing the notion of equivalence of weak crossed products induced by measurings.

Definition 2.1. Let H be a weak Hopf algebra, let $\varphi_B, \phi_B : H \otimes B \rightarrow B$ be measurings, and let $\sigma, \tau : H \otimes H \rightarrow B$ be morphisms such that $\sigma * u_2^{\varphi_B} = \sigma, \tau * u_2^{\phi_B} = \tau$. Assume that σ, τ satisfy the twisted condition (77) and the 2-cocycle condition (78), and suppose that ν is a preunit for $\mu_{B \otimes_{\varphi_B} H}$, and u is a preunit for $\mu_{B \otimes_{\phi_B} H}$. We say that $(B \otimes H, \mu_{B \otimes_{\varphi_B} H})$ and $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$ are equivalent weak crossed products if there is an isomorphism

$$\Upsilon : B \times_{\varphi_B}^{\sigma} H \rightarrow B \times_{\phi_B}^{\tau} H$$

of algebras, left B -modules and right H -comodules, where the left actions are defined by $\varphi_{B \times_{\varphi_B}^{\sigma} H} = p_{B \otimes H}^{\varphi_B} \circ (\mu_B \otimes H) \circ (B \otimes i_{B \otimes H}^{\varphi_B})$, $\varphi_{B \times_{\phi_B}^{\tau} H} = p_{B \otimes H}^{\phi_B} \circ (\mu_B \otimes H) \circ (B \otimes i_{B \otimes H}^{\phi_B})$, and the right coactions are $\rho_{B \times_{\varphi_B}^{\sigma} H} = (p_{B \otimes H}^{\varphi_B} \otimes H) \circ (B \otimes \delta_H) \circ i_{B \otimes H}^{\varphi_B}$, $\rho_{B \times_{\phi_B}^{\tau} H} = (p_{B \otimes H}^{\phi_B} \otimes H) \circ (B \otimes \delta_H) \circ i_{B \otimes H}^{\phi_B}$.

In our setting the general criterion [10, Theorem 1.7] that characterizes equivalent weak crossed products admits the following formulation:

Theorem 2.2. Let H be a weak Hopf algebra, let $\varphi_B, \phi_B : H \otimes B \rightarrow B$ be measurings and let $\sigma, \tau : H \otimes H \rightarrow B$ be morphisms such that $\sigma * u_2^{\varphi_B} = \sigma, \tau * u_2^{\phi_B} = \tau$. Assume that σ, τ satisfy the twisted condition (77), the 2-cocycle condition (78) and suppose that ν is a preunit for $\mu_{B \otimes_{\varphi_B} H}$ and u is a preunit for $\mu_{B \otimes_{\phi_B} H}$. Let $(B, H, P_{\varphi_B}, F_{\sigma})$ be the quadruple associated to φ_B , and σ and let $(B, H, P_{\phi_B}, F_{\tau})$ be the one associated to ϕ_B and τ . The following assertions are equivalent:

- (i) The weak crossed products $(B \otimes H, \mu_{B \otimes_{\varphi_B} H})$ and $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$ are equivalent.
- (ii) There exist two morphisms $T, S : B \otimes H \rightarrow B \otimes H$, of left B -modules for the trivial action $\varphi_{B \otimes H} = \mu_B \otimes H$, and right H -modules for the trivial coaction $\rho_{B \otimes H} = B \otimes \delta_H$, satisfying the conditions

$$(91) \quad T \circ \nu = u,$$

$$(92) \quad T \circ \mu_{B \otimes_{\varphi_B} H} = \mu_{B \otimes_{\phi_B} H} \circ (T \otimes T),$$

$$(93) \quad S \circ T = \nabla_{B \otimes H}^{\varphi_B}, \quad T \circ S = \nabla_{B \otimes H}^{\phi_B}.$$

(iii) *There exist two morphisms $\theta, \gamma : H \rightarrow B \otimes H$ of right H -modules for the trivial coaction satisfying the conditions*

$$(94) \quad \theta = \nabla_{B \otimes H}^{\varphi_B} \circ \theta,$$

$$(95) \quad (\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H),$$

$$(96) \quad P_{\phi_B} = (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B),$$

$$(97) \quad F_\tau = (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \mu_{B \otimes \varphi_B H} \circ (\theta \otimes \theta),$$

$$(98) \quad u = (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nu.$$

Proof. The proof of this theorem is the one developed in [10, Theorem 1.7]. It is not difficult to check the right H -comodule condition for the morphisms S, T, θ and γ . We leave the details of the proof to the reader. \square

Proposition 2.3. *Let H be a weak Hopf algebra, let $\varphi_B, \phi_B : H \otimes B \rightarrow B$ be measurings and let $\sigma, \tau : H \otimes H \rightarrow B$ be morphisms such that $\sigma * u_2^{\varphi_B} = \sigma, \tau * u_2^{\phi_B} = \tau$. Assume that σ, τ satisfy the twisted condition (77), the 2-cocycle condition (78) and suppose that $\nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$ is a preunit for $\mu_{B \otimes \varphi_B H}$ and $\nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H)$ is a preunit for $\mu_{B \otimes \tau_{\phi_B} H}$. If $(B \otimes H, \mu_{B \otimes \varphi_B H})$ and $(B \otimes H, \mu_{B \otimes \tau_{\phi_B} H})$ are equivalent weak crossed products, there exists morphisms $T, S : B \otimes H \rightarrow B \otimes H$ of left B -modules for the trivial action and right H -comodules for the trivial coaction such that*

$$(99) \quad \nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H) = T \circ (\eta_B \otimes \eta_H), \quad \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H) = S \circ (\eta_B \otimes \eta_H).$$

Proof. If $(B \otimes H, \mu_{B \otimes \varphi_B H})$ and $(B \otimes H, \mu_{B \otimes \tau_{\phi_B} H})$ are equivalent weak crossed products, there exists an isomorphism of algebras, left B -modules and right H -comodules $\Upsilon : B \times_{\varphi_B}^{\sigma} H \rightarrow B \times_{\phi_B}^{\tau} H$. By (i) \Rightarrow (ii) of the previous theorem there exists two morphisms of left B -modules and right H -comodules $T, S : B \otimes H \rightarrow B \otimes H$, defined by

$$T = i_{B \otimes H}^{\phi_B} \circ \Upsilon \circ p_{B \otimes H}^{\varphi_B}, \quad S = i_{B \otimes H}^{\varphi_B} \circ \Upsilon^{-1} \circ p_{B \otimes H}^{\phi_B}$$

and satisfying the conditions (93),

$$(100) \quad T \circ S \circ T = T$$

and

$$(101) \quad S \circ T \circ S = S.$$

Then,

$$\nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H) \stackrel{(91)}{=} T \circ \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H) \stackrel{(93)}{=} T \circ S \circ T \circ (\eta_B \otimes \eta_H) \stackrel{(100)}{=} T \circ (\eta_B \otimes \eta_H).$$

On the other hand, if ν and u are the preunits of $(B \otimes H, \mu_{B \otimes_{\varphi_B} H})$ and $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$, by (91) we have that $S \circ T \circ \nu = S \circ u$. Then, by (93) we have that $\nabla_{B \otimes H}^{\varphi_B} \circ \nu = S \circ u$ and applying (90) we obtain that

$$(102) \quad S \circ u = \nu$$

holds. Therefore, in our particular case, we have

$$\nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H) \stackrel{(102)}{=} S \circ \nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H) \stackrel{(93)}{=} S \circ T \circ S \circ (\eta_B \otimes \eta_H) \stackrel{(101)}{=} S \circ (\eta_B \otimes \eta_H).$$

□

Theorem 2.4. *Let H be a weak Hopf algebra, let $\varphi_B, \phi_B : H \otimes B \rightarrow B$ be measurings, and let $\sigma, \tau : H \otimes H \rightarrow B$ be morphisms such that $\sigma * u_2^{\varphi_B} = \sigma$, $\tau * u_2^{\phi_B} = \tau$. Assume that σ, τ satisfy the twisted condition (77), the 2-cocycle condition (78) and suppose that ν is a preunit for $\mu_{B \otimes_{\varphi_B} H}$, and u is a preunit for $\mu_{B \otimes_{\phi_B} H}$. The following assertions are equivalent:*

- (i) *The weak crossed products $(B \otimes H, \mu_{B \otimes_{\varphi_B} H})$ and $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$ are equivalent.*
- (ii) *There exists two morphisms $h, h^{-1} : H \rightarrow B$ such that*

$$(103) \quad h^{-1} * h = u_1^{\varphi_B},$$

$$(104) \quad h * h^{-1} * h = h, \quad h^{-1} * h * h^{-1} = h^{-1},$$

$$(105) \quad \phi_B = \mu_B \circ (\mu_B \otimes h^{-1}) \circ (h \otimes P_{\varphi_B}) \circ (\delta_H \otimes B),$$

$$(106) \quad \tau = \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H)).$$

$$(107) \quad u = ((\mu_B \circ (B \otimes h^{-1})) \otimes H) \circ (B \otimes \delta_H) \circ \nu,$$

Proof. First we will prove that (i) \Rightarrow (ii). By Theorem 2.2 there exists two morphisms $T, S : B \otimes H \rightarrow B \otimes H$ of left B -modules for the trivial action and right H -comodules for trivial coaction defined as in the proof of the previous proposition and satisfying the conditions (91), (92), (93), (100) and (101). Also, S preserves the preunit, i.e., (102) holds, and S is multiplicative, i.e., $S \circ \mu_{B \otimes_{\phi_B} H} = \mu_{B \otimes_{\varphi_B} H} \circ (S \otimes S)$ ([10, (37)]). Also, by (ii) \Rightarrow (iii) of Theorem 2.2, there exists two morphisms $\theta, \gamma : H \rightarrow B \otimes H$ of right H -comodules defined by

$$(108) \quad \theta = S \circ (\eta_B \otimes H), \quad \gamma = T \circ (\eta_B \otimes H).$$

Then,

$$(109) \quad S = (\mu_B \otimes H) \circ (B \otimes \theta), \quad T = (\mu_B \otimes H) \circ (B \otimes \gamma).$$

For θ and γ the equalities (94), (95), (96), (97), and (98) hold. Moreover,

$$(110) \quad \gamma = \nabla_{B \otimes H}^{\phi_B} \circ \gamma,$$

$$(111) \quad (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \theta = \nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes H),$$

$$(112) \quad P_{\varphi_B} = (\mu_B \otimes H) \circ (\mu_B \otimes \theta) \circ (B \otimes P_{\phi_B}) \circ (\gamma \otimes B),$$

$$(113) \quad F_\sigma = (\mu_B \otimes H) \circ (B \otimes \theta) \circ \mu_{B \otimes \varphi_B^* H} \circ (\gamma \otimes \gamma),$$

$$(114) \quad \nu = (\mu_B \otimes H) \circ (B \otimes \theta) \circ u,$$

also hold. Define

$$(115) \quad h = (B \otimes \varepsilon_H) \circ \theta, \quad h^{-1} = (B \otimes \varepsilon_H) \circ \gamma.$$

Then, by the condition of right H -comodule morphism for θ and γ , we have

$$(116) \quad \theta = (h \otimes H) \circ \delta_H, \quad \gamma = (h^{-1} \otimes H) \circ \delta_H.$$

The equality (103) holds because

$$\begin{aligned} & h^{-1} * h \\ &= (B \otimes \varepsilon_H) \circ (\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma \text{ (by the comodule morphism condition for } \gamma \text{ and counit properties)} \\ &= (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H) \text{ (by (95))} \\ &= u_1^{\varphi_B} \text{ (by counit properties)}. \end{aligned}$$

Also,

$$\begin{aligned} & h * h^{-1} * h \\ &= \mu_B \circ (H \otimes \varepsilon_H \otimes \varphi_B) \circ (\theta \otimes \eta_B) \text{ (by the comodule morphism condition for } \theta \text{ and (103))} \\ &= (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ \theta \text{ (by counit properties and naturality of } c) \\ &= (B \otimes \varepsilon_H) \circ \theta \text{ (by (94))} \\ &= h \text{ (by counit properties)} \end{aligned}$$

and

$$\begin{aligned} & h^{-1} * h * h^{-1} \\ &= (\mu_B \otimes \varepsilon_H) \circ (B \otimes \gamma) \circ P_{\varphi_B} \circ (H \otimes \eta_B) \text{ (by naturality of } c) \\ &= (\mu_B \otimes \varepsilon_H) \circ (B \otimes \gamma) \circ \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H) \text{ (by properties of } \eta_B) \\ &= (\mu_B \otimes \varepsilon_H) \circ (B \otimes \gamma) \circ (\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma \text{ (by (95))} \\ &= (\mu_B \otimes \varepsilon_H) \circ (B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H))) \circ \gamma \text{ (by the associativity of } \mu_B \text{ and (111))} \\ &= (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ \gamma \text{ (by (46) and properties of } \eta_B) \\ &= h^{-1} \text{ (by (110))}. \end{aligned}$$

The equality (107) follows directly from (98) because γ is a morphism of right H -comodules. Moreover, composing in (96) with $B \otimes \varepsilon_H$, by (116) we prove (105). Finally, (106) holds because

$$\begin{aligned} & \tau \\ &= (B \otimes \varepsilon_H) \circ F_\tau \text{ (by (61) for } \tau) \\ &= (\mu_B \otimes \varepsilon_H) \circ (B \otimes \gamma) \circ \mu_{B \otimes \varphi_B^* H} \circ (\theta \otimes \theta) \text{ (by (97))} \\ &= (\mu_B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^* H} \circ ((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H) \text{ (by (116))}. \end{aligned}$$

Conversely, to prove (ii) \Rightarrow (i), define

$$\theta = (h \otimes H) \circ \delta_H, \quad \gamma = (h^{-1} \otimes H) \circ \delta_H.$$

Then θ and γ are morphisms of right H -comodules, $h = (B \otimes \varepsilon_H) \circ \theta$ and $h^{-1} = (B \otimes \varepsilon_H) \circ \gamma$. To prove the equivalence between $(B \otimes H, \mu_{B \otimes \varphi_B H})$ and $(B \otimes H, \mu_{B \otimes \phi_B H})$, we must show that (94), (95), (96), (97) and (98) hold. First note that, (98) follows from (107). Also, (94) holds because:

$$\begin{aligned} & \nabla_{B \otimes H}^{\varphi_B} \circ \theta \\ &= ((h * u_1^{\varphi_B}) \otimes H) \circ \delta_H \text{ (by the coassociativity of } \delta_H \text{ and (42))} \\ &= ((h * h^{-1} * h) \otimes H) \circ \delta_H \text{ (by (103))} \\ &= \theta \text{ (by (104)).} \end{aligned}$$

On the other hand, (95) follows by

$$\begin{aligned} & (\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma \\ &= ((h^{-1} * h) \otimes H) \circ \delta_H \text{ (by coassociativity of } \delta_H) \\ &= (u_1^{\varphi_B} \otimes H) \circ \delta_H \text{ (by (103))} \\ &= \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H) \text{ (by (44)).} \end{aligned}$$

and (96) follows by

$$\begin{aligned} & P_{\phi_B} \\ &= ((\mu_B \circ (\mu_B \otimes h^{-1})) \otimes H) \circ (h \otimes ((P_{\varphi_B} \otimes H) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B))) \circ (\delta_H \otimes B) \text{ (by (105) and} \\ & \quad \text{coassociativity of } \delta_H) \\ &= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B) \text{ (by coassociativity of } \delta_H \text{ and the naturality of } c). \end{aligned}$$

Finally, (97) holds because

$$\begin{aligned} & F_{\tau} \\ &= ((\mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B H}) \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H))) \otimes \mu_H \circ \delta_{H \otimes 2} \text{ (by (106))} \\ &= ((\mu_B \circ (\mu_B \otimes h^{-1})) \otimes H) \circ (\mu_B \otimes ((F_{\sigma} \otimes \mu_H) \circ \delta_{H \otimes 2})) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \text{ (by coassociativity} \\ & \quad \text{of } \delta_H \text{ and the naturality of } c) \\ &= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (\mu_B \otimes F_{\sigma}) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \text{ (by (56))} \\ &= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \mu_{B \otimes \varphi_B H} \circ (\theta \otimes \theta) \text{ (by the definition of } \mu_{B \otimes \varphi_B H}). \end{aligned}$$

□

Remark 2.5. Note that, in the conditions of (ii) of Theorem 2.4, composing with $H \otimes \eta_B$ in (105), we obtain the identity

$$(117) \quad h * h^{-1} = u_1^{\phi_B}.$$

Definition 2.6. Let H be a weak Hopf algebra and let $\varphi_B : H \otimes B \rightarrow B$ be a measuring. We will say that the pair of morphisms $h, h^{-1} : H \rightarrow B$ is a gauge transformation for φ_B if they satisfy (103) and (104).

By the previous Theorem 2.4 we know that, under suitable conditions, equivalent weak crossed products are related by gauge transformations. After the next discussion, we should be able to secure that the converse is also true.

2.7. Let H be a weak Hopf algebra and let $\varphi_B : H \otimes B \rightarrow B$ be a measuring. Let (h, h^{-1}) be a gauge transformation for φ_B and let $\sigma : H \otimes H \rightarrow B \otimes H$ be a morphism satisfying the identity $\sigma * u_2^{\varphi_B} = \sigma$, the

twisted condition (77) and the 2-cocycle condition (78). Suppose that ν is a preunit for the associated weak crosse product $\mu_{B \otimes_{\varphi_B} H}$.

Define θ and γ as in (116), i.e., $\theta = (h \otimes H) \circ \delta_H$ and $\gamma = (h^{-1} \otimes H) \circ \delta_H$. Then θ and γ are morphisms of right H -comodules. Also, by (42), the coassociativity of δ_H and the condition of gauge transformation, we have that $\nabla_{B \otimes H}^{\varphi_B} \circ \theta = \theta$ and then (94) holds. By similar arguments and the associativity of μ_B we obtain the equality

$$(118) \quad (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nabla_{B \otimes H}^{\varphi_B} = (\mu_B \otimes H) \circ (B \otimes \gamma).$$

Moreover, by the coassociativity of δ_H and the condition of gauge transformation we have

$$(\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma = (u_1^{\varphi_B} \otimes H) \circ \delta_H = \nabla_{B \otimes H}(\eta_B \otimes H)$$

and then (95) holds. As a consequence, we obtain

$$(119) \quad (\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) = \nabla_{B \otimes H}^{\varphi_B}.$$

Define $\varphi_B^h : H \otimes B \rightarrow B$ by

$$(120) \quad \varphi_B^h = \mu_B \circ (\mu_B \otimes h^{-1}) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B)$$

and $\sigma^h : H \otimes H \rightarrow B$ by

$$(121) \quad \sigma^h = \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B} H} \circ (\theta \otimes \theta).$$

Then, φ_B^h is a measuring because

$$\begin{aligned} & \mu_B \circ (\varphi_B^h \otimes \varphi_B^h) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes B \otimes B) \\ &= \mu_B \circ (B \otimes \mu_B) \circ ((\mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B)) \otimes (\mu_B \circ (u_1^{\varphi_B} \otimes \varphi_B) \circ (\delta_H \otimes B)) \otimes h^{-1}) \circ (H \otimes c_{H,B} \otimes c_{H,B}) \\ & \quad \circ (H \otimes H \otimes c_{H,B} \otimes B) \circ (((H \otimes \delta_H) \circ \delta_H) \otimes B \otimes B) \text{ (by naturality of } c, \text{ coassociativity of } \delta_H, \text{ associativity} \\ & \quad \text{of } \mu_B, \text{ and (103))} \\ &= \mu_B \circ (B \otimes \mu_B) \circ (h \otimes (\mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes B \otimes B)) \otimes h^{-1}) \circ (H \otimes H \otimes B \otimes c_{H,B}) \\ & \quad \circ (H \otimes H \otimes c_{H,B} \otimes B) \circ (((H \otimes \delta_H) \circ \delta_H) \otimes B \otimes B) \text{ (by (52), coassociativity of } \delta_H \text{ and associativity of } \mu_B \\ & \quad \mu_B) \\ &= \varphi_B^h \circ (H \otimes \mu_B) \text{ (by and (b1) of Definition 1.4 and naturality of } c) \end{aligned}$$

and trivially, by the coassociativity of δ_H and the naturality of c , we have that

$$P_{\varphi_B^h} = (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B)$$

Therefore (96) holds. Also, by the associativity of μ_B , (42) and the condition of gauge transformation

$$(122) \quad u_1^{\varphi_B^h} = h * h^{-1}$$

holds. Then, as a consequence of the previous identity, we have that

$$(123) \quad (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \theta = (u_1^{\varphi_B^h} \otimes H) \circ \delta_H$$

and

$$(124) \quad (\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \gamma) \circ \theta)) = \nabla_{B \otimes H}^{\varphi_B^h}.$$

hold.

On the other hand for F_{σ^h} we have

$$\begin{aligned}
& F_{\sigma^h} \\
&= ((\mu_B \circ (\mu_H \otimes h^{-1})) \otimes H) \circ (\mu_B \otimes ((F_\sigma \otimes \mu_H) \circ \delta_{H^{\otimes 2}})) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \text{ (by the naturality} \\
&\quad \text{of } c \text{ and the coassociativity of } \delta_H) \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\theta \otimes \theta) \text{ (by (56))}
\end{aligned}$$

and (97) also holds. Moreover, $\sigma^h * u_2^{\varphi_B^h} = \sigma^h$, because

$$\begin{aligned}
& \sigma^h * u_2^{\varphi_B^h} \\
&= \mu_B \circ ((\mu_B \otimes (\mu_B \otimes h^{-1})) \otimes (\varphi_B^h \circ (H \otimes \eta_B))) \circ (\mu_B \otimes ((F_\sigma \otimes \mu_H) \circ \delta_{H^{\otimes 2}})) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \\
&\quad \text{(by the naturality of } c, \text{ the coassociativity of } \delta_H \text{ and the condition of morphism of right } H\text{-comodules for } \theta) \\
&= \mu_B \circ ((\mu_B \otimes (\mu_B \otimes h^{-1})) \otimes (\varphi_B^h \circ (H \otimes \eta_B))) \circ (\mu_B \otimes ((B \otimes \delta_H) \circ F_\sigma)) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \\
&\quad \text{(by (56))} \\
&= \mu_B \circ (\mu_B \otimes (h^{-1} * u_1^{\varphi_B^h})) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \text{ (by the associativity of } \mu_B) \\
&= \mu_B \circ (\mu_B \otimes (h^{-1} * h * h^{-1})) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta) \text{ (by (122))} \\
&= \sigma^h \text{ (by (104) and associativity of } \mu_B).
\end{aligned}$$

By (i) of Theorem 1.13 to obtain that $(B, H, P_{\varphi_B^h}, F_{\sigma^h})$ satisfies the twisted condition is enough to prove that (77) holds. Indeed,

$$\begin{aligned}
& \mu_B \circ (B \otimes \sigma^h) \circ (P_{\varphi_B^h} \otimes H) \circ (H \otimes P_{\varphi_B^h}) \\
&= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B}^\sigma H} \circ (((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma))) \otimes \theta) \circ (\mu_B \otimes H \otimes H) \\
&\quad \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes ((\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B))) \text{ (by the associativity of } \mu_B) \\
&= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\nabla_{B \otimes H}^{\varphi_B} \otimes \theta) \circ (\mu_B \otimes H \otimes H) \circ (B \otimes P_{\varphi_B} \otimes H) \\
&\quad \circ (\theta \otimes ((\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B))) \text{ (by (119))} \\
&= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\mu_B \otimes H \otimes \theta) \circ (B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B}) \otimes H) \\
&\quad \circ (\theta \otimes ((\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B))) \text{ (by the associativity of } \mu_B) \\
&= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\mu_B \otimes H \otimes \theta) \circ (B \otimes P_{\varphi_B} \otimes H) \\
&\quad \circ (\theta \otimes ((\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B))) \text{ (by (41))} \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (P_{\varphi_B} \otimes B))) \otimes H \\
&\quad \circ (\theta \otimes ((\mu_B \otimes \theta) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\theta \otimes B))) \text{ (by the definition of } \mu_{B \otimes_{\varphi_B}^\sigma H}) \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \\
&\quad \circ (\theta \otimes ((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ P_{\varphi_B})) \circ (\theta \otimes B) \text{ (by (38)} \\
&\quad \text{and the associativity of } \mu_B) \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes ((\mu_B \otimes H) \circ (B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B})) \circ (\theta \otimes B))) \\
&\quad \text{(by (119))} \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (\mu_B \otimes ((\mu_B \otimes H) \circ (B \otimes F_\sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B}))) \circ (B \otimes P_{\varphi_B} \otimes H \otimes B) \circ (\theta \otimes \theta \otimes B) \\
&\quad \text{(by (41), (38) and the associativity of } \mu_B) \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (B \otimes P_{\varphi_B}) \circ ((\mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\theta \otimes \theta)) \otimes B) \text{ (by (71))} \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (B \otimes P_{\varphi_B}) \circ ((\nabla_{B \otimes H}^{\varphi_B} \circ \mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\theta \otimes \theta)) \otimes B) \text{ (by (79))} \\
&= \mu_B \circ (\mu_B \otimes h^{-1}) \circ ((\mu_B \circ (B \otimes (h^{-1} * h))) \otimes P_{\varphi_B}) \circ (B \otimes \delta_H \otimes B) \circ ((\mu_{B \otimes_{\varphi_B}^\sigma H} \circ (\theta \otimes \theta)) \otimes B) \\
&\quad \text{(by (42) and the condition of gauge transformation)}
\end{aligned}$$

$$= \mu_B \circ (B \otimes P_{\varphi_B^h}) \circ (F_{\sigma^h} \otimes B) \text{ (by the associativity of } \mu_B \text{ and the coassociativity of } \delta_H \text{)}.$$

Also, by (ii) of Theorem 1.13, to obtain that $(B, H, P_{\varphi_B^h}, F_{\sigma^h})$ satisfies the cocycle condition is enough to prove that (78) holds. Indeed,

$$\begin{aligned} & \mu_B \circ (B \otimes \sigma^h) \circ (P_{\varphi_B^h} \otimes H) \circ (H \otimes F_{\sigma^h}) \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ (\mu_B \otimes H \otimes \theta) \circ (B \otimes \mu_B \otimes H \otimes H) \\ & \quad \circ (B \otimes B \otimes ((\mu_B \otimes H) \circ (B \otimes (\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ P_{\varphi_B}) \otimes H) \circ (B \otimes P_{\varphi_B} \otimes \gamma) \\ & \quad \circ (\theta \otimes (\mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta))) \text{ (by the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ (\mu_B \otimes H \otimes \theta) \circ (\mu_B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B}) \otimes H) \circ (B \otimes P_{\varphi_B} \otimes \gamma) \\ & \quad \circ (\theta \otimes (\mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta))) \text{ (by (119) and the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ (\mu_B \otimes H \otimes \theta) \circ (\mu_B \otimes P_{\varphi_B} \otimes H) \circ (B \otimes P_{\varphi_B} \otimes \gamma) \\ & \quad \circ (\theta \otimes (\mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta))) \text{ (by (41))} \\ &= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (\mu_B \otimes F_{\sigma}) \circ (\mu_B \otimes P_{\varphi_B} \otimes H) \\ & \quad \circ (\mu_B \otimes P_{\varphi_B} \otimes ((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ F_{\sigma})) \\ & \quad \circ (B \otimes P_{\varphi_B} \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta \otimes \theta) \text{ (by (38), the definition of } \mu_{B \otimes \varphi_B^h} \text{ and the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (\mu_B \otimes ((\mu_B \otimes H) \circ (B \otimes F_{\sigma}) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ F_{\sigma})))) \\ & \quad \circ (\mu_B \otimes P_{\varphi_B} \otimes H \otimes H) \circ (B \otimes P_{\varphi_B} \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta \otimes \theta) \text{ (by (119) and the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (B \otimes \mu_B \otimes H) \circ (B \otimes B \otimes ((\mu_B \otimes H) \circ (B \otimes F_{\sigma}) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes F_{\sigma}))) \\ & \quad \circ (\mu_B \otimes P_{\varphi_B} \otimes H \otimes H) \circ (B \otimes P_{\varphi_B} \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes \theta \otimes \theta) \text{ (by (60) and the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (B \otimes \mu_B \otimes H) \circ (\mu_B \otimes B \otimes F_{\sigma}) \\ & \quad \circ (B \otimes B \otimes ((\mu_B \otimes H) \circ (B \otimes F_{\sigma}) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B}))) \otimes H) \circ (B \otimes P_{\varphi_B} \otimes H \otimes \theta) \circ (\theta \otimes \theta \otimes H) \\ & \quad \text{(by (72) and the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ ((\mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta)) \otimes \theta) \text{ (by (71) and the associativity of } \mu_B \text{)} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ ((\nabla_{B \otimes H}^{\varphi_B} \circ \mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta)) \otimes \theta) \text{ (by (79))} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ (((\mu_B \otimes H) \circ (B \otimes ((u_1^{\varphi_B} \otimes H) \circ \delta_H))) \circ \mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta)) \otimes \theta \\ & \quad \text{(by (42))} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes \varphi_B^h} \circ (((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ \mu_{B \otimes \varphi_B^h} \circ (\theta \otimes \theta)) \otimes \theta) \\ & \quad \text{(by (119) and the condition of gauge transformation)} \\ &= \mu_B \circ (B \otimes \sigma^h) \circ (F_{\sigma^h} \otimes B) \text{ (by the associativity of } \mu_B \text{)}. \end{aligned}$$

If we define ν^h by $\nu^h = (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nu$ we have that (98) holds trivially. Moreover ν^h is a preunit for $(B \otimes H, \mu_{B \otimes \varphi_B^h})$ because (87), (88) and (89) hold. Indeed, (87) follows by

$$\begin{aligned} & \mu_B \circ (B \otimes F_{\sigma^h}) \circ (P_{\varphi_B^h} \otimes H) \circ (H \otimes \nu^h) \\ &= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \mu_{B \otimes \varphi_B^h} \circ (\mu_B \otimes H \otimes \theta) \\ & \quad \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ P_{\varphi_B}) \otimes H) \circ (\theta \otimes ((\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nu)) \\ & \quad \text{(by the associativity of } \mu_B \text{)} \\ &= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (\mu_B \otimes F_{\sigma}) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}))) \otimes H) \circ (B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B}) \otimes \theta) \\ & \quad \circ (\theta \otimes ((\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nu)) \text{ (by (119), the definition of } \mu_{B \otimes \varphi_B^h} \text{ and the associativity of } \mu_B \text{)} \\ &= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (\mu_B \otimes F_{\sigma}) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (P_{\varphi_B} \otimes B))) \otimes \theta \\ & \quad \circ (\theta \otimes ((\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nu)) \text{ (by (41))} \end{aligned}$$

$$\begin{aligned}
&= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes ((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ \nu))) \\
&\quad \text{(by (38) and the associativity of } \mu_B \text{)} \\
&= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (\mu_B \otimes F_\sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (\theta \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ \nu)) \quad \text{(by (119))} \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nabla_{B \otimes H}^{\varphi_B} \circ \mu_{B \otimes \varphi_B}^{\sigma} H \circ ((\nabla_{B \otimes H}^{\varphi_B} \circ \theta) \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ \nu)) \quad \text{(by (94) and (118))} \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nabla_{B \otimes H}^{\varphi_B} \circ \theta \quad \text{(by the condition of unit for } p_{B \otimes H}^{\varphi_B} \circ \nu \text{)} \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \theta \quad \text{(by (94))} \\
&= (u_1^{\varphi_B^h} \otimes H) \circ \delta_H \quad \text{(by (123))} \\
&= \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H) \quad \text{(by (123) and the properties of } \eta_B \text{).}
\end{aligned}$$

Also, (88) holds because

$$\begin{aligned}
&\mu_B \circ (B \otimes F_{\sigma^h}) \circ (\nu^h \otimes H) \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \mu_{B \otimes \varphi_B}^{\sigma} H \circ (((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ \nu) \otimes \theta) \\
&\quad \text{(by the associativity of } \mu_B \text{)} \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nabla_{B \otimes H}^{\varphi_B} \circ \mu_{B \otimes \varphi_B}^{\sigma} H \circ ((\nabla_{B \otimes H}^{\varphi_B} \circ \theta) \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ \nu)) \quad \text{(by (94) and (118))} \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \nabla_{B \otimes H}^{\varphi_B} \circ \theta \quad \text{(by the condition of unit for } p_{B \otimes H}^{\varphi_B} \circ \nu \text{)} \\
&= (\mu_B \otimes H) \circ (B \otimes \gamma) \circ \theta \quad \text{(by (94))} \\
&= (u_1^{\varphi_B^h} \otimes H) \circ \delta_H \quad \text{(by (123))} \\
&= \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H) \quad \text{(by (123) and the properties of } \eta_B \text{).}
\end{aligned}$$

Finally,

$$\begin{aligned}
&\mu_B \circ (B \otimes P_{\varphi_B^h}) \circ (\nu^h \otimes H) \\
&= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (((\mu_B \otimes H) \circ (B \otimes ((\mu_B \otimes H) \circ (B \otimes \theta) \circ \gamma)) \circ \nu) \otimes B) \\
&\quad \text{(by the associativity of } \mu_B \text{)} \\
&= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ ((\nabla_{B \otimes H}^{\varphi_B} \circ \nu) \otimes B) \quad \text{(by (119))} \\
&= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes P_{\varphi_B}) \circ (\nu \otimes B) \quad \text{(by (90))} \\
&= (\mu_B \otimes H) \circ (\mu_B \otimes \gamma) \circ (B \otimes \nu) \quad \text{(by (89))} \\
&= (\mu_B \otimes H) \circ (B \otimes \nu^h) \quad \text{(by the associativity of } \mu_B \text{).}
\end{aligned}$$

and (89) holds.

Therefore, as a consequence of the previous facts, we have a theorem that generalizes to the monoidal setting [15, Theorem 5.4].

Theorem 2.8. *Let H be a weak Hopf algebra, let $\varphi_B, \phi_B : H \otimes B \rightarrow B$ be measurings, and let $\sigma, \tau : H \otimes H \rightarrow B$ be morphisms such that $\sigma * u_2^{\varphi_B} = \sigma, \tau * u_2^{\phi_B} = \tau$. Assume that σ, τ satisfy the twisted condition (77), the 2-cocycle condition (78) and suppose that ν is a preunit for $\mu_{B \otimes \varphi_B}^{\sigma} H$, and u is a preunit for $\mu_{B \otimes \phi_B}^{\tau} H$. The weak crossed products $(B \otimes H, \mu_{B \otimes \varphi_B}^{\sigma} H)$ and $(B \otimes H, \mu_{B \otimes \phi_B}^{\tau} H)$ are equivalent if and only if there exists a gauge transformation (h, h^{-1}) for φ_B such that $\phi_B = \varphi_B^h, \tau = \sigma^h$ and $u = \nu^h$.*

3. REGULAR MORPHISMS

Definition 3.1. Let H be a weak Hopf algebra and φ_B be a measuring. With $\text{Reg}_{\varphi_B}(H, B)$ we will denote the set of regular morphisms between H and B , i.e., a morphism $h : H \rightarrow B$ is a regular

morphism if there exists a morphism $h^{-1} : H \rightarrow B$, called the convolution inverse of h , such that the pair (h, h^{-1}) is a gauge transformation for φ_B and

$$(125) \quad h * h^{-1} = u_1^{\varphi_B},$$

holds. Then, $Reg_{\varphi_B}(H, B)$, with the convolution as a product, is a group with unit $u_1^{\varphi_B}$.

3.2. Let \mathcal{P}_{φ_B} be the set of all pairs (ϕ_B, τ) , where:

- (i) The morphism $\phi_B : H \otimes B \rightarrow B$ is a measuring satisfying $u_1^{\phi_B} = u_1^{\varphi_B}$.
- (ii) The morphism $\tau : H \otimes H \rightarrow B$ is such that $\tau = \tau * u_2^{\phi_B}$ and the associated quadruple $(B, H, P_{\phi_B}, F_{\tau})$ satisfies the twisted condition and the cocycle condition.
- (iii) The associated weak crossed product $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$ admits a preunit v .

By the results proved in the previous section we know that $Reg_{\varphi_B}(H, B)$ acts on \mathcal{P}_{φ_B} . The action

$$R : Reg_{\varphi_B}(H, B) \times \mathcal{P}_{\varphi_B} \rightarrow \mathcal{P}_{\varphi_B},$$

is defined by

$$(126) \quad R(h, (\phi_B, \tau)) = (\phi_B^h, \tau^h).$$

Proposition 3.3. *Let (B, φ_B) , (B, ϕ_B) be a left weak H -module algebra and let $h : H \rightarrow B$ be a morphism such that $h * u_1^{\varphi_B} = h = u_1^{\phi_B} * h$. Then, the following assertions are equivalent:*

- (i) $h \circ \eta_H = \eta_B$.
- (ii) $h \circ \Pi_H^L = u_1^{\phi_B}$.
- (iii) $h \circ \overline{\Pi}_H^L = u_1^{\varphi_B}$.

Moreover, if $h * \eta_H = \eta_B$, the identity

$$(127) \quad (\mu_B \otimes H) \circ (B \otimes ((h \otimes H) \circ \delta_H \circ \eta_H)) = \nabla_{B \otimes H}^{\varphi_B} \circ (B \otimes \eta_H),$$

holds.

Also, if $g : H \rightarrow B$ is a morphism such that $g * u_1^{\phi_B} = g = u_1^{\varphi_B} * g$, the following assertions are equivalent:

- (iv) $g \circ \eta_H = \eta_B$.
- (v) $g \circ \Pi_H^L = u_1^{\varphi_B}$.
- (vi) $g \circ \overline{\Pi}_H^L = u_1^{\phi_B}$.

Then, if $g * \eta_H = \eta_B$, the identity

$$(128) \quad (\mu_B \otimes H) \circ (B \otimes ((g \otimes H) \circ \delta_H \circ \eta_H)) = \nabla_{B \otimes H}^{\phi_B} \circ (B \otimes \eta_H)$$

holds.

Moreover, if there exists $h^{-1} : H \rightarrow B$ such that (h, h^{-1}) is a gauge transformation for φ_B and $h * h^{-1} = u_1^{\phi_B}$ holds, we have $h * \eta_H = \eta_B$ iff $h^{-1} * \eta_H = \eta_B$.

Proof. By the properties of η_H and (b2) of Definition 1.4, we obtain that (ii) \Rightarrow (i), and (iii) \Rightarrow (i). Also, (i) \Rightarrow (ii) holds because,

$$h \circ \Pi_H^L$$

$$\begin{aligned}
&= (u_1^{\phi_B} * h) \circ \Pi_H^L \text{ (by } h = u_1^{\phi_B} * h) \\
&= \mu_B \circ (u_1^{\phi_B} \otimes h) \circ (((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes c_{H,H}) \circ (\delta_H \otimes H)) \circ H \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H) \\
&\quad \text{(by naturality of } c \text{ and coassociativity of } \delta_H) \\
&= \mu_B \circ ((u_1^{\phi_B} \circ \mu_H \circ (H \otimes \Pi_H^L)) \otimes h) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H) \text{ (by (9))} \\
&= \mu_B \circ ((\phi_B \circ (H \otimes u_1^{\phi_B})) \otimes h) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H) \text{ (by (b3) of Definition 1.4 and (30))} \\
&= \mu_B \circ ((\phi_B \circ (\overline{\Pi}_H^L \otimes B)) \otimes h) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes u_1^{\phi_B}) \text{ (by the naturality of } c \text{ and (17))} \\
&= \mu_B \circ ((\mu_B \circ c_{B,B} \circ (u_1^{\phi_B} \otimes B)) \otimes h) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes u_1^{\phi_B}) \text{ (by (26))} \\
&= \mu_B \circ c_{B,B} \circ (((u_1^{\phi_B} * h) \circ \eta_H) \otimes u_1^{\phi_B}) \text{ (by naturality of } c \text{ and associativity of } \mu_B) \\
&= \mu_B \circ c_{B,B} \circ ((h \circ \eta_H) \otimes u_1^{\phi_B}) \text{ (by } h = u_1^{\phi_B} * h) \\
&= u_1^{\phi_B} \text{ (by (i), naturality of } c, \text{ and properties of } \eta_B).
\end{aligned}$$

On the other hand, (i) \Rightarrow (iii) because

$$\begin{aligned}
&h \circ \overline{\Pi}_H^L \\
&= (h * u_1^{\varphi_B}) \circ \overline{\Pi}_H^L \text{ (by } h = h * u_1^{\varphi_B}) \\
&= \mu_B \circ (h \otimes u_1^{\varphi_B}) \circ (H \otimes ((H \otimes (\varepsilon_H \circ \mu_H)) \circ (\delta_H \otimes H))) \circ ((\delta_H \circ \eta_H) \otimes H) \text{ (by the coassociativity of } \delta_H) \\
&= \mu_B \circ (h \otimes (u_1^{\varphi_B} \circ \mu_H \circ (H \otimes \overline{\Pi}_H^L))) \circ ((\delta_H \circ \eta_H) \otimes H) \text{ (by (14))} \\
&= \mu_B \circ (h \otimes \varphi_B) \circ ((\delta_H \circ \eta_H) \otimes u_1^{\varphi_B}) \text{ (by (b3) of Definition 1.4 and (29))} \\
&= \mu_B \circ (h \otimes (\varphi_B \circ (\Pi_H^L \otimes B))) \circ ((\delta_H \circ \eta_H) \otimes u_1^{\varphi_B}) \text{ (by (17))} \\
&= \mu_B \circ (h \otimes (\mu_B \circ (u_1^{\varphi_B} \otimes B))) \circ ((\delta_H \circ \eta_H) \otimes u_1^{\varphi_B}) \text{ (by (25))} \\
&= \mu_B \circ (((h * u_1^{\varphi_B}) \circ \eta_H) \otimes u_1^{\varphi_B}) \text{ (by associativity of } \mu_B) \\
&= \mu_B \circ ((h \circ \eta_H) \otimes u_1^{\varphi_B}) \text{ (by } h = h * u_1^{\varphi_B}) \\
&= u_1^{\varphi_B} \text{ (by (i), and properties of } \eta_B).
\end{aligned}$$

As a consequence of these equivalences, we obtain (127) because

$$\begin{aligned}
&(\mu_B \otimes H) \circ (B \otimes ((h \otimes H) \circ \delta_H \circ \eta_H)) \\
&= (\mu_B \otimes H) \circ (B \otimes (((h \circ \overline{\Pi}_H^L) \otimes H) \circ \delta_H \circ \eta_H)) \text{ (by (17))} \\
&= \nabla_{B \otimes H}^{\varphi_B} \circ (B \otimes \eta_H) \text{ (by (iii) and (42))}
\end{aligned}$$

The proof for the equivalences associated to g and (128) are similar and we leave the details to the reader.

Finally, assume that there exists $h^{-1} : H \rightarrow B$ such that (h, h^{-1}) is a gauge transformation for φ_B and $h * h^{-1} = u_1^{\phi_B}$ holds. If $h \circ \eta_H = \eta_B$, we have

$$\begin{aligned}
&h^{-1} \circ \eta_H \\
&= (u_1^{\varphi_B} * h^{-1}) \circ \eta_H \text{ (by } u_1^{\varphi_B} * h^{-1} = h^{-1}) \\
&= ((h \circ \overline{\Pi}_H^L) * h^{-1}) \circ \eta_H \text{ (by (ii))} \\
&= u_1^{\phi_B} \circ \eta_H \text{ (by (17) and } u_1^{\phi_B} = h * h^{-1}) \\
&= \eta_B \text{ (by (b2) of Definition 1.4).}
\end{aligned}$$

Conversely, if $h^{-1} \circ \eta_H = \eta_B$, by similar arguments (in this case $g = h$),

$$h \circ \eta_H = (h * u_1^{\varphi_B}) \circ \eta_H = (h * (h^{-1} \circ \Pi_H^L)) \circ \eta_H = u_1^{\phi_B} \circ \eta_H = \eta_B.$$

□

As a particular instance of the previous proposition we have the following corollary.

Corollary 3.4. *Let (B, φ_B) be a left weak H -module algebra and let $h : H \rightarrow B$ be a morphism such that $h * u_1^{\varphi_B} = h = u_1^{\varphi_B} * h$. Then, the following assertions are equivalent:*

- (i) $h \circ \eta_H = \eta_B$.
- (ii) $h \circ \Pi_H^L = u_1^{\varphi_B}$.
- (iii) $h \circ \overline{\Pi}_H^L = u_1^{\varphi_B}$.

Moreover, if $h \in \text{Reg}_{\varphi_B}(H, B)$ with convolution inverse $h^{-1} : H \rightarrow B$, we have $h * \eta_H = \eta_B$ iff $h^{-1} * \eta_H = \eta_B$. Then, under these conditions, if $h * \eta_H = \eta_B$, the following assertions hold and are equivalent:

- (iv) $h^{-1} \circ \eta_H = \eta_B$.
- (v) $h^{-1} \circ \Pi_H^L = u_1^{\varphi_B}$.
- (vi) $h^{-1} \circ \overline{\Pi}_H^L = u_1^{\varphi_B}$.

Definition 3.5. Let H be a weak Hopf algebra and $\varphi_B : H \otimes B \rightarrow B$ be a measuring. With

$$\text{Reg}_{\varphi_B}^t(H, B)$$

we will denote the set of morphisms $h : H \rightarrow B$ in $\text{Reg}_{\varphi_B}(H, B)$ such that

$$(129) \quad h \circ \eta_H = \eta_B.$$

Then, if (B, φ_B) is a left weak H -module algebra, by Corollary 3.4, $\text{Reg}_{\varphi_B}^t(H, B)$ is a subgroup of $\text{Reg}_{\varphi_B}(H, B)$.

Remark 3.6. The set $\text{Reg}_{\varphi_B}^t(H, B)$ also acts on \mathcal{P}_{φ_B} , i.e., we have a map

$$R' : \text{Reg}_{\varphi_B}^t(H, B) \times \mathcal{P}_{\varphi_B} \rightarrow \mathcal{P}_{\varphi_B},$$

defined by

$$R'(h, (\phi_B, \tau)) = R(h, (\phi_B, \tau)),$$

where R is the action defined in (126).

Note that, if (B, φ_B) is a left weak H -module algebra, the measuring φ_B^h defined in (120) satisfies (b2) of Definition 1.4 because we have

$$\begin{aligned} & \varphi_B^h \circ (\eta_H \otimes B) \\ &= \mu_B \circ ((\mu_B \circ ((h \circ \overline{\Pi}_H^L) \otimes B)) \otimes h^{-1}) \circ (H \otimes P_{\varphi_B}) \circ ((\delta_H \circ \eta_H) \otimes B) \text{ (by (17))} \\ &= \mu_B \circ ((\mu_B \circ (u_1^{\varphi_B} \otimes B)) \otimes h^{-1}) \circ (H \otimes P_{\varphi_B}) \circ ((\delta_H \circ \eta_H) \otimes B) \text{ (by (i) } \Rightarrow \text{ (iii) of Corollary 3.4)} \\ &= \mu_B \circ (\mu_B \otimes h^{-1}) \circ (B \otimes P_{\varphi_B}) \circ ((P_{\varphi_B} \circ (\eta_H \otimes \eta_B)) \otimes B) \text{ (by (39))} \\ &= \mu_B \circ (B \otimes h^{-1}) \circ P_{\varphi_B} \circ (\eta_H \otimes B) \text{ (by (38) and properties of } \eta_B) \\ &= \mu_B \circ (\varphi_B \otimes B) \circ (H \otimes c_{B, B}) \circ (((H \otimes (h^{-1} \circ \Pi_H^L)) \circ \delta_H \circ \eta_H) \otimes B) \text{ (by (17) and naturality of } c) \\ &= \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ P_{\varphi_B} \circ (\eta_H \otimes B) \text{ (by (v) of Corollary 3.4 and the naturality of } c) \\ &= \varphi_B \circ (\eta_H \otimes B) \text{ (by (58) for } \phi_B) \\ &= id_B \text{ (by (b2) of Definition 1.4).} \end{aligned}$$

Theorem 3.7. *Let H be a weak Hopf algebra, let $(B, \varphi_B), (B, \phi_B)$ be left weak H -module algebras and let $\sigma, \tau : H \otimes H \rightarrow B$ be morphisms such that $\sigma * u_2^{\varphi_B} = \sigma$, $\tau * u_2^{\phi_B} = \tau$. Assume that σ, τ satisfy the twisted condition (77), the 2-cocycle condition (78) and the normal condition (84). The following assertions are equivalent:*

- (i) *The weak crossed products $(B \otimes H, \mu_{B \otimes_{\varphi_B} H})$ and $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$ are equivalent.*
- (ii) *There exists a gauge transformation (h, h^{-1}) for φ_B such that (117),*

$$(130) \quad h \circ \eta_H = \eta_B,$$

$$(131) \quad \mu_B \circ (B \otimes h) \circ P_{\phi_B} = \mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B),$$

$$(132) \quad \mu_B \circ (B \otimes h) \circ F_\tau = (\mu_B \otimes H) \circ (\mu_B \otimes \sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H)),$$

hold.

Proof. We first prove (i) \Rightarrow (ii). By Corollary 1.20, we know that $(B \otimes H, \mu_{B \otimes_{\varphi_B} H})$ and $(B \otimes H, \mu_{B \otimes_{\phi_B} H})$ are weak crossed products with preunits $\nu = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$, $u = \nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H)$, respectively. As in the proof of Theorem 2.4 define θ, γ by (108) and h, h^{-1} by (115). Then, using that T, S are morphisms of left B -modules and (116) we have the following identities:

$$(133) \quad (B \otimes \varepsilon_H) \circ T = \mu_B \circ (B \otimes h^{-1}), \quad (B \otimes \varepsilon_H) \circ S = \mu_B \circ (B \otimes h).$$

By (i) \Rightarrow (ii) of Theorem 2.4, the pair (h, h^{-1}) is a gauge transformation for φ_B and the identities (105), (106) and (107) hold. Therefore, by Remark (2.5) we obtain that (117) holds.

On the other hand, we obtain (130) because

$$\begin{aligned} & h \circ \eta_H \\ &= (B \otimes \varepsilon_H) \circ S \circ (\eta_B \otimes \eta_H) \text{ (by (133))} \\ &= (B \otimes \varepsilon_H) \circ \nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H) \text{ (by (99))} \\ &= u_1^{\phi_B} \circ \eta_H \text{ (by naturality of } c \text{ and counit properties)} \\ &= \eta_B \text{ (by (b2) of Definition 1.4).} \end{aligned}$$

Now, by the proof of (i) \Rightarrow (ii) of Theorem 2.4 we know that S is multiplicative, i.e.,

$$(134) \quad S \circ \mu_{B \otimes_{\phi_B} H} = \mu_{B \otimes_{\varphi_B} H} \circ (S \otimes S).$$

Then composing with $\eta_B \otimes H \otimes B \otimes \eta_H$ in the previous identity we have

$$\begin{aligned} & S \circ \mu_{B \otimes_{\phi_B} H} \circ (\eta_B \otimes H \otimes B \otimes \eta_H) \\ &= (\mu_B \otimes H) \circ (\mu_B \otimes \theta) \circ (B \otimes (F_\tau \circ (H \otimes \eta_H))) \circ P_{\phi_B} \text{ (by (109) and properties of } \eta_B) \\ &= (\mu_B \otimes H) \circ (\mu_B \otimes \theta) \circ (B \otimes (\nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes H))) \circ P_{\phi_B} \text{ (by (86))} \\ &= (\mu_B \otimes H) \circ (B \otimes \theta) \circ \nabla_{B \otimes H}^{\phi_B} \circ P_{\phi_B} \text{ (by the left } B\text{-linearity of } \nabla_{B \otimes H}^{\phi_B} \text{ and properties of } \eta_B) \\ &= (\mu_B \otimes H) \circ (B \otimes \theta) \circ P_{\phi_B} \text{ (by (41).)} \end{aligned}$$

and

$$\begin{aligned} & \mu_{B \otimes_{\varphi_B} H} \circ (S \otimes S) \circ (\eta_B \otimes H \otimes B \otimes \eta_H) \\ &= \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes (((\mu_B \otimes H) \circ (B \otimes ((h \otimes H) \circ \delta_H \circ \eta_H)))) \text{ (by (109), (116), and properties} \end{aligned}$$

$$\begin{aligned}
& \text{of } \eta_B) \\
&= \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ (B \otimes \eta_H))) \text{ (by (127))} \\
&= \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes B \otimes \eta_H) \text{ (by (80))} \\
&= (\mu_B \otimes H) \circ (\mu_B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes H))) \circ (B \otimes P_{\varphi_B}) \circ (((h \otimes H) \circ \delta_H) \otimes B) \text{ (by (86))} \\
&= (\mu_B \otimes H) \circ (B \otimes (\nabla_{B \otimes H}^{\varphi_B} \circ P_{\varphi_B})) \circ (((h \otimes H) \circ \delta_H) \otimes B) \text{ (by the left } B\text{-linearity of } \nabla_{B \otimes H}^{\varphi_B} \text{ and properties of } \eta_B) \\
&= (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (((h \otimes H) \circ \delta_H) \otimes B) \text{ (by the left } B\text{-linearity of } \nabla_{B \otimes H}^{\varphi_B} \text{ and (41))}.
\end{aligned}$$

Therefore,

$$(135) \quad (\mu_B \otimes H) \circ (B \otimes \theta) \circ P_{\phi_B} = (\mu_B \otimes H) \circ (B \otimes P_{\varphi_B}) \circ (((h \otimes H) \circ \delta_H) \otimes B)$$

holds and, composing with $B \otimes \varepsilon_H$ in (135), we obtain (131) by the naturality of c and (40).

Finally, composing with $\eta_B \otimes H \otimes \eta_B \otimes H$ in (134) we have

$$\begin{aligned}
& S \circ \mu_{B \otimes_{\phi_B} H} \circ (\eta_B \otimes H \otimes \eta_B \otimes H) \\
&= (\mu_B \otimes H) \circ (B \otimes ((h \otimes H) \circ \delta_H)) \circ F_{\tau} \text{ (by (74), (109), (116) and the properties of } \eta_B)
\end{aligned}$$

and

$$\begin{aligned}
& \mu_{B \otimes_{\varphi_B} H} \circ (S \otimes S) \circ (\eta_B \otimes H \otimes \eta_B \otimes H) \\
&= \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H)) \text{ (by (109), properties of } \eta_B, \text{ and (116))}.
\end{aligned}$$

As a consequence,

$$(136) \quad (\mu_B \otimes H) \circ (B \otimes ((h \otimes H) \circ \delta_H)) \circ F_{\tau} = \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H))$$

holds and, composing with $B \otimes \varepsilon_H$, we obtain (132) by the counit properties and (61).

Conversely, consider that (ii) holds. To prove that (ii) \Rightarrow (i), following (ii) \Rightarrow (i) of Theorem 2.4, we only need to obtain the equalities (105), (106) and (107). Indeed, note that by Proposition 3.3, $h^{-1} \circ \eta_H = \eta_B$ because (130) holds. Then, (105) holds because

$$\begin{aligned}
& \mu_B \circ ((\mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B)) \otimes h^{-1}) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \\
& \mu_B \circ (B \otimes (\mu_B \circ (h \otimes h^{-1}))) \circ (P_{\phi_B} \otimes H) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \text{ (by (131) and the associativity of } \mu_B) \\
&= \mu_B \circ (B \otimes (h * h^{-1})) \circ P_{\phi_B} \text{ (by the naturality of } c \text{ and coassociativity of } \delta_H) \\
&= \mu_B \circ (B \otimes u_1^{\phi_B}) \circ P_{\phi_B} \text{ (by (117))} \\
&= \phi_B \text{ (by (58) for } \phi_B).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \mu_B \circ (B \otimes h^{-1}) \circ \mu_{B \otimes_{\varphi_B} H} \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H)) \\
&= \mu_B \circ (((\mu_B \otimes H) \circ (\mu_B \otimes \sigma)) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H))) \otimes (h^{-1} \circ \mu_H) \\
& \quad \circ \delta_{H \otimes 2} \text{ (by naturality of } c \text{ and coassociativity of } \delta_H) \\
&= \mu_B \circ (B \otimes (\mu_B \circ (h \otimes h^{-1}))) \circ (F_{\tau} \otimes \mu_H) \circ \delta_{H \otimes 2} \text{ (by (132) and associativity of } \mu_B) \\
&= \mu_B \circ (B \otimes (h * h^{-1})) \circ F_{\tau} \text{ (by (56))} \\
&= \mu_B \circ (B \otimes u_1^{\phi_B}) \circ F_{\tau} \text{ (by (117))} \\
&= \tau \text{ (by (59))}
\end{aligned}$$

and then (106) holds. Finally, we obtain (107) because

$$\begin{aligned}
& ((\mu_B \circ (B \otimes h^{-1})) \otimes H) \circ (B \otimes \delta_H) \circ \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H) \\
&= ((u_1^{\varphi_B} * h^{-1}) \otimes H) \circ \delta_H \circ \eta_H \text{ (by (44) and by the coassociativity of } \delta_H) \\
&= (h^{-1} \otimes H) \circ \delta_H \circ \eta_H \text{ (by the gauge transformation condition)} \\
&= ((h^{-1} \circ \overline{\Pi}_H^L) \otimes H) \circ \delta_H \circ \eta_H \text{ (by (17))} \\
&= (u_1^{\phi_B} \otimes H) \circ \delta_H \circ \eta_H \text{ (by (vi) of Proposition 3.3)} \\
&= \nabla_{B \otimes H}^{\phi_B} \circ (\eta_B \otimes \eta_H) \text{ (by (44))}.
\end{aligned}$$

□

3.8. As a consequence of the previous theorem, it is possible to define a groupoid, denoted by \mathcal{G}_H^B whose objects are pairs

$$(\varphi_B, \sigma),$$

where (B, φ_B) is a left weak H -module algebra, $\sigma : H \otimes H \rightarrow B$ is a morphism such that

$$u_2^{\varphi_B} * \sigma = \sigma = \sigma * u_2^{\varphi_B}$$

and the associated quadruple \mathbb{B}_H satisfies the twisted, cocycle and normal conditions.

A morphism between two objects $(\varphi_B, \sigma), (\phi_B, \tau)$ of \mathcal{G}_H^B is defined by a morphism $h : H \rightarrow B$ for which there exists a morphism $h^{-1} : H \rightarrow B$ such that (h, h^{-1}) is a gauge transformation for φ_B satisfying the conditions (ii) of Theorem 3.7. If $h : (\varphi_B, \sigma) \rightarrow (\phi_B, \tau)$, $g : (\phi_B, \tau) \rightarrow (\chi_B, \omega)$ are morphisms in \mathcal{G}_H^B , the composition, denoted by $g \odot h$, is defined by

$$g \odot h = g * h.$$

The previous composition is well defined because, if $l = g * h$ and $l^{-1} = h^{-1} * g^{-1}$, it is easy to show that (l, l^{-1}) is a gauge transformation for φ_B and $l * l^{-1} = u_1^{\chi_B}$. Also,

$$\begin{aligned}
& l \circ \eta_H \\
&= (g * (h \circ \Pi_H^L)) \circ \eta_H \text{ (by (17))} \\
&= (g * u_1^{\phi_B}) \circ \eta_H \text{ (by (ii) of Proposition 3.3)} \\
&= g \circ \eta_H \text{ (by the condition of gauge transformation)} \\
&= \eta_B \text{ (by (130))}
\end{aligned}$$

and then (130) holds for l . The equality (131) for l follows by

$$\begin{aligned}
& \mu_B \circ (B \otimes l) \circ P_{\chi_B} \\
&= \mu_B \circ ((\mu_B \circ (B \otimes g) \circ P_{\chi_B}) \otimes h) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \text{ (by the associativity of } \mu_B, \text{ the coassociativity of } \delta_H \text{ and the naturality of } c) \\
&= \mu_B \circ ((\mu_B \circ (g \otimes \phi_B) \circ (\delta_H \otimes B)) \otimes h) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \text{ (by (131))} \\
&= \mu_B \circ (g \otimes ((\mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B))) \circ (\delta_H \otimes B) \text{ (by the coassociativity of } \delta_H) \\
&= \mu_B \circ (g \otimes (\mu_B \circ (B \otimes h) \circ P_{\phi_B})) \circ (\delta_H \otimes B) \text{ (by (131))} \\
&= (\mu_B \circ (l \otimes \phi_B) \circ (\delta_H \otimes B)) \text{ (by the associativity of } \mu_B \text{ and the coassociativity of } \delta_H)
\end{aligned}$$

and (132) holds because

$$\begin{aligned}
& \mu_B \circ (B \otimes l) \circ F_\omega \\
&= \mu_B \circ ((\mu_B \circ (B \otimes g) \circ F_\omega) \otimes (h \circ \mu_H)) \circ \delta_{H \otimes 2} \text{ (by the associativity of } \mu_B, \text{ the coassociativity of } \delta_{H \otimes 2})
\end{aligned}$$

$$\begin{aligned}
&= \mu_B \circ ((\mu_B \circ (\mu_B \otimes \tau)) \circ (B \otimes P_{\phi_B} \otimes H) \circ (((g \otimes H) \circ \delta_H) \otimes ((g \otimes H) \circ \delta_H))) \otimes (h \circ \mu_H) \circ \delta_{H^{\otimes 2}} \\
&\quad \text{(by (132))} \\
&= \mu_B \circ ((\mu_B \circ (B \otimes \phi_B)) \otimes (\mu_B \circ (B \otimes h) \circ F_\tau)) \circ (B \otimes H \otimes c_{H,B} \otimes H) \circ (((g \otimes \delta_H) \circ \delta_H) \otimes ((g \otimes H) \circ \delta_H)) \\
&\quad \text{(by the associativity of } \mu_B, \text{ the coassociativity of } \delta_H \text{ and the naturality of } c) \\
&= \mu_B \circ ((\mu_B \circ (B \otimes \phi_B)) \otimes (\mu_B \circ (\mu_B \otimes \sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (((h \otimes H) \circ \delta_H) \otimes ((h \otimes H) \circ \delta_H)))) \\
&\quad \circ (B \otimes H \otimes c_{H,B} \otimes H) \circ (((g \otimes \delta_H) \circ \delta_H) \otimes ((g \otimes H) \circ \delta_H)) \text{ (by the associativity of } \mu_B \text{ and (132))} \\
&= \mu_B \circ ((\mu_B \circ (B \otimes (\mu_B \circ (B \otimes h) \circ P_{\phi_B}))) \otimes (\mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H))) \\
&\quad \circ (((g \otimes \delta_H) \circ \delta_H) \otimes (((g \otimes h) \circ \delta_H) \otimes H) \circ \delta_H) \text{ (by the associativity of } \mu_B, \text{ the coassociativity of } \delta_H \text{ and} \\
&\quad \text{the naturality of } c) \\
&= \mu_B \circ ((\mu_B \circ (B \otimes (\mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B)))) \otimes (\mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H))) \\
&\quad \circ (((g \otimes \delta_H) \circ \delta_H) \otimes (((g \otimes h) \circ \delta_H) \otimes H) \circ \delta_H) \text{ (by the associativity of } \mu_B \text{ and (131))} \\
&= \mu_B \circ (B \otimes (\mu_B \circ ((\mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes B \otimes B)) \otimes \sigma))) \\
&\quad \circ (B \otimes H \otimes ((B \otimes c_{H,B}) \circ (c_{H,B} \otimes B)) \otimes H) \circ (((g \otimes \delta_H) \circ \delta_H) \otimes (((g \otimes h) \circ \delta_H) \otimes H) \circ \delta_H) \\
&\quad \text{(by the associativity of } \mu_B, \text{ the coassociativity of } \delta_H \text{ and the naturality of } c) \\
&= \mu_B \circ (\mu_B \otimes \sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (((l \otimes H) \circ \delta_H) \otimes ((l \otimes H) \circ \delta_H)) \text{ (by the associativity of } \mu_B \text{ and} \\
&\quad \text{(b1) of Definition 1.4).}
\end{aligned}$$

The identity of (φ_B, σ) is $id_{(\varphi_B, \sigma)} = u_1^{\varphi_B}$ because $h * u_1^{\varphi_B} = h$ and $(u_1^{\varphi_B}, u_1^{\varphi_B})$ is a gauge transformation for φ_B satisfying (125). The equality (130) follows from (b2) of Definition 1.4, (131) is a consequence of the naturality of c and (b1) of Definition 1.4 and (132) holds because:

$$\begin{aligned}
&(\mu_B \otimes H) \circ (\mu_B \otimes \sigma) \circ (B \otimes P_{\varphi_B} \otimes H) \circ (((u_1^{\varphi_B} \otimes H) \circ \delta_H) \otimes ((u_1^{\varphi_B} \otimes H) \circ \delta_H)) \\
&= \mu_B \circ ((\mu_B \circ (u_1^{\varphi_B} \otimes \varphi_B) \circ (\delta_H \otimes u_1^{\varphi_B})) \otimes \sigma) \circ \delta_{H^{\otimes 2}} \text{ (by naturality of } c \text{ and coassociativity of } \delta_H) \\
&= \mu_B \circ ((\varphi_B \circ (H \otimes u_1^{\varphi_B})) \otimes \sigma) \circ \delta_H^{\otimes 2} \text{ (by (52))} \\
&= u_2^{\varphi_B} * \sigma \text{ (by (b3) of Definition 1.4)} \\
&= \sigma \text{ (by (78))} \\
&= \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ F_\sigma \text{ (by (59)).}
\end{aligned}$$

As a consequence, h is an isomorphism with inverse h^{-1} and $(g \odot h)^{-1} = h^{-1} \odot g^{-1}$. Therefore, \mathcal{G}_H^B is a groupoid.

3.9. Let H be a weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma : H \otimes H \rightarrow B$ be a morphism such that $\sigma * u_2^{\varphi_B} = \sigma$. Assume that σ satisfies the twisted condition (77), the 2-cocycle condition (78) and the normal condition (84). Let h be a morphism in $Reg_{\varphi_B}^t(H, B)$. Then (h, h^{-1}) is a gauge transformation for φ_B such that (125) and (130) hold. Define φ_B^h and σ^h as in (120) and (121) respectively. Then, by (2.7), φ_B^h is a measuring such that (122) holds. Therefore $u_1^{\varphi_B} = u_1^{\varphi_B^h}$ and then

$$(137) \quad \nabla_{B \otimes H}^{\varphi_B^h} = \nabla_{B \otimes H}^{\varphi_B}.$$

Moreover, by Remark 3.6, we know that φ_B^h satisfies (b2) of Definition 1.4. On the other hand, σ^h is such that $\sigma^h * u_2^{\varphi_B^h} = \sigma^h$ and satisfies the twisted condition (77), the cocycle condition (78) and

$$\nu^h = (\mu_B \otimes H) \circ (B \otimes ((h^{-1} \otimes H) \circ \delta_H)) \circ \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$$

is a preunit for the associated weak crossed product $(B \otimes H, \mu_{B \otimes_{\varphi_B^h} H})$. Note that

$$\begin{aligned}
& \nu^h \\
&= ((u_1^{\varphi_B} * h^{-1}) \otimes H) \circ \delta_H \circ \eta_H \text{ (by the coassociativity of } \delta_H) \\
&= (h^{-1} \otimes H) \circ \delta_H \circ \eta_H \text{ (by the condition of gauge transformation)} \\
&= ((h^{-1} \circ \overline{\Pi}_H^L) \otimes H) \circ \delta_H \circ \eta_H \text{ (by (17))} \\
&= (u_1^{\varphi_B} \otimes H) \circ \delta_H \circ \eta_H \text{ (by (vi) of Corollary (3.4))} \\
&= \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H) \text{ (by (44))} \\
&= \nabla_{B \otimes H}^{\varphi_B^h} \circ (\eta_B \otimes \eta_H) \text{ (by (137)).}
\end{aligned}$$

Therefore,

$$\nu^h = \nabla_{B \otimes H}^{\varphi_B^h} \circ (\eta_B \otimes \eta_H) = \nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H) = \nu.$$

Also, φ_B^h satisfies (26) because

$$\begin{aligned}
& \varphi_B^h \circ (\overline{\Pi}_H^L \otimes B) \\
&= \mu_B \otimes (\mu_B \otimes h^{-1}) \circ (B \otimes P_{\varphi_B}) \circ (((h \circ \overline{\Pi}_H^L) \otimes H) \circ \delta_H \circ \overline{\Pi}_H^L \otimes B) \text{ (by (8))} \\
&= \mu_B \otimes (\mu_B \otimes h^{-1}) \circ (u_1^{\varphi_B} \otimes P_{\varphi_B}) \circ ((\delta_H \circ \overline{\Pi}_H^L) \otimes B) \text{ (by (ii) of Corollary 3.4)} \\
&= \mu_B \otimes (B \otimes h^{-1}) \circ P_{\varphi_B} \circ (\overline{\Pi}_H^L \otimes B) \text{ (by (53))} \\
&= \mu_B \circ ((\varphi_B \circ (\overline{\Pi}_H^L \otimes B)) \otimes h^{-1}) \circ (H \otimes c_{H,B}) \circ ((\delta_H \circ \overline{\Pi}_H^L) \otimes B) \text{ (by (8))} \\
&= \mu_B \circ ((\mu_B \circ c_{B,B} \circ (u_1^{\varphi_B})) \otimes h^{-1}) \circ (H \otimes c_{H,B}) \circ ((\delta_H \circ \overline{\Pi}_H^L) \otimes B) \text{ (by (26) for } \varphi_B) \\
&= \mu_B \circ (B \otimes (u_1^{\varphi_B} * h^{-1})) \circ c_{H,B} \circ (\overline{\Pi}_H^L \otimes B) \text{ (by the associativity of } \mu_B \text{ and the naturality of } c) \\
&= \mu_B \circ c_{B,B} \circ ((h^{-1} \circ \overline{\Pi}_H^L) \otimes B) \text{ (by the naturality of } c \text{ and the condition of gauge transformation)} \\
&= \mu_B \circ c_{B,B} \circ (u_1^{\varphi_B} \otimes B) \text{ (by (vi) of Corollary 3.4)} \\
&= \mu_B \circ c_{B,B} \circ (u_1^{\varphi_B^h} \otimes B) \text{ (by } u_1^{\varphi_B} = u_1^{\varphi_B^h})
\end{aligned}$$

As a consequence, using the same proof than in Remark 1.21, we obtain that (85) holds for φ_B^h . Therefore, by Corollary 1.19, we have that σ^h satisfies the normal condition (84), i.e.,

$$\sigma^h \circ (\eta_H \otimes H) = u_1^{\varphi_B^h} = \sigma^h \circ (H \otimes \eta_H).$$

Finally, if B is commutative and H is coconmutative, the equality

$$(138) \quad \mu_B \circ (B \otimes h) \circ P_{\varphi_B} = \mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B)$$

holds and then

$$\begin{aligned}
& \varphi_B^h \\
&= \mu_B \circ ((\mu_B \circ (h \otimes \varphi_B) \circ (\delta_H \otimes B)) \otimes h^{-1}) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \text{ (by the coassociativity of } \delta_H) \\
&= \mu_B \circ ((\mu_B \circ (B \otimes h) \circ P_{\varphi_B}) \otimes h^{-1}) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \text{ (by (138))} \\
&= \mu_B \circ (B \otimes (h * h^{-1})) \circ P_{\varphi_B} \text{ (by the coassociativity of } \delta_H, \text{ the associativity of } \mu_B \text{ and the naturality of } c) \\
&= \mu_B \circ (B \otimes u_1^{\varphi_B}) \circ P_{\varphi_B} \text{ (by (125))} \\
&= \varphi_B \text{ (by (58)).}
\end{aligned}$$

4. HOM-PRODUCTS, INVERTIBLE MORPHISMS AND CENTERS

In this subsection, for a weak Hopf algebra H and an algebra B , we will explore a product in $\text{Hom}_C(H^{\otimes n} \otimes B, B)$ that will permit us to extend some results about the factorization through the center of B , given in [14] for Hopf algebras, to the weak Hopf algebra setting.

Definition 4.1. Let H be a weak Hopf algebra and let $\varphi_B : H \otimes B \rightarrow B$ be a measuring. Let φ and $\psi \in \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B)$. We define the product

$$\wedge : \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B) \times \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B) \rightarrow \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B)$$

between φ and ψ as

$$\varphi \wedge \psi = \varphi \circ (H^{\otimes n} \otimes \psi) \circ (\delta_{H^{\otimes n}} \otimes B).$$

Obviously, \wedge is an associative product because $\delta_{H^{\otimes n}}$ is coassociative.

We say that a morphism $\varphi \in \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B)$ is φ_B -invertible if there exists a morphism $\varphi^\dagger \in \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B)$ such that

$$(139) \quad \varphi \wedge \varphi^\dagger = \mu_B \circ (u_n^{\varphi_B} \otimes B).$$

Proposition 4.2. Let H be a weak Hopf algebra and let $\varphi_B : H \otimes B \rightarrow B$ be a measuring. For $\omega : H^{\otimes n} \rightarrow B$ define

$$\overline{\omega} = \mu_B \circ (\omega \otimes B), \quad \overline{\omega}^{op} = \mu_B \circ c_{B,B} \circ (\omega \otimes B).$$

Then, if $\omega, \theta \in \text{Hom}_{\mathcal{C}}(H^{\otimes n}, B)$ and $\gamma \in \text{Hom}_{\mathcal{C}}(H^{\otimes n} \otimes B, B)$ the following equalities hold:

- (i) $\overline{\omega} \wedge \overline{\theta} = \overline{\omega * \theta}$.
- (ii) If H is cocommutative, $\overline{\omega}^{op} \wedge \gamma = \mu_B \circ (\gamma \otimes \omega) \circ (H^{\otimes n} \otimes c_{H^{\otimes n}, B}) \circ (\delta_{H^{\otimes n}} \otimes B)$.
- (iii) If H is cocommutative, $\overline{\omega}^{op} \wedge \overline{\theta}^{op} = \overline{\theta * \omega}^{op}$.
- (iv) If H is cocommutative, $\overline{\omega}^{op} \wedge \overline{\theta} = \overline{\theta} \wedge \overline{\omega}^{op}$.
- (v) If (B, φ_B) is a left weak H -module algebra, $\overline{u_n^{\varphi_B}} \wedge \varphi_B^{\otimes n} = \varphi_B^{\otimes n}$.
- (vi) If H is cocommutative and (B, φ_B) is a left weak H -module algebra, $\overline{u_n^{\varphi_B}}^{op} \wedge \varphi_B^{\otimes n} = \varphi_B^{\otimes n}$.
- (vii) $\overline{\varphi_B \circ (H \otimes \omega)} \wedge (\varphi_B \circ (H \otimes \gamma)) = \varphi_B \circ (H \otimes (\overline{\omega} \wedge \gamma))$.
- (viii) If H is cocommutative, $\overline{\varphi_B \circ (H \otimes \omega)}^{op} \wedge (\varphi_B \circ (H \otimes \gamma)) = \varphi_B \circ (H \otimes (\overline{\omega}^{op} \wedge \gamma))$.

Proof. The proof of (i) follows directly from the associativity of μ_B . If H is cocommutative, so is $H^{\otimes n}$ and, by the naturality of c , we obtain (ii). By similar reasoning and using the associativity of μ_B we obtain (iii) and (iv). On the other hand,

$$\overline{u_n^{\varphi_B}} \wedge \varphi_B^{\otimes n}$$

$$= \mu_B \circ (\varphi_B^{\otimes n} \otimes \varphi_B^{\otimes n}) \circ (H^{\otimes n} \otimes c_{H^{\otimes n}, B} \otimes B) \circ (\delta_{H^{\otimes n}} \otimes \eta_B \otimes B) \stackrel{(31)}{=} \varphi_B^{\otimes n} \circ (H^{\otimes n} \otimes (\mu_B \circ (\eta_B \otimes B))) = \varphi_B^{\otimes n},$$

and then (v) holds. Similarly, using that $H^{\otimes n}$ is cocommutative, the naturality of c and (31) we prove (vi).

The identity, (vii) follows from

$$\begin{aligned} & \overline{\varphi_B \circ (H \otimes \omega)} \wedge (\varphi_B \circ (H \otimes \gamma)) \\ &= \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes ((\omega \otimes \theta) \circ (\delta_{H^{\otimes n}} \otimes B))) \text{ (by the naturality of } c) \\ &= \varphi_B \circ (H \otimes (\overline{\omega} \wedge \gamma)) \text{ (by (31))} \end{aligned}$$

and, similarly, using that H is cocommutative, we obtain (viii). \square

Remark 4.3. The equivalence of measurings (or, in particular, of weak actions) in terms of gauge transformations acquires a new meaning in terms of this product. Actually, if H is cocommutative, the action described in (126) on a measuring ϕ_B can be seen as a conjugation by gauge transformations in the following way:

$$\phi_B^h = \overline{h} \wedge \overline{h^{-1}}^{op} \wedge \phi_B.$$

Moreover observe that for a cocommutative weak Hopf algebra H and measurings φ_B and ϕ_B satisfying conditions of Theorem 2.4, we can re-write equality (105) using the Hom-product as

$$\phi_B = \overline{h} \wedge \overline{h^{-1}}^{op} \wedge \varphi_B.$$

Also in this way, equality (131) of Theorem 3.7 can be interpreted as $\overline{h}^{op} \wedge \phi_B = \overline{h} \wedge \varphi_B$, in coherence with the action of gauge transformations as a conjugation given above.

Proposition 4.4. *Let H be a cocommutative weak Hopf algebra and let (B, φ_B) be a left weak H -module algebra. A morphism $\sigma : H^{\otimes 2} \rightarrow B$ satisfies the twisted condition (77) if and only if*

$$(140) \quad \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} = \overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B))$$

holds.

Proof. The proof follows from the following facts. First, note that by definition of F_σ , we have that

$$\mu_B \circ (B \otimes \varphi_B) \circ (F_\sigma \otimes B) = \overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B)).$$

On the other hand, if H is cocommutative, by the naturality of c , we have

$$\mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B}) = \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2}.$$

□

Definition 4.5. Let H be a weak Hopf algebra and (B, φ_B) be a left weak H -module algebra. For $n \geq 1$, with $Reg_{\varphi_B}(H^{\otimes n}, B)$ we will denote the set of morphisms $\sigma : H^{\otimes n} \rightarrow B$ such that there exists a morphism $\sigma^{-1} : H^{\otimes n} \rightarrow B$ (the convolution inverse of σ) satisfying the following equalities:

$$(141) \quad \sigma * \sigma^{-1} = \sigma^{-1} * \sigma = u_n^{\varphi_B},$$

$$(142) \quad \sigma * \sigma^{-1} * \sigma = \sigma, \quad \sigma^{-1} * \sigma * \sigma^{-1} = \sigma^{-1}.$$

Note that, for $n = 1$, we recover the group $Reg_{\varphi_B}(H, B)$ introduced in Definition 3.1. For any n , $Reg_{\varphi_B}(H^n, B)$ is a group with unit element $u_n^{\varphi_B}$ because by (34) we know that $u_n^{\varphi_B} * u_n^{\varphi_B} = u_n^{\varphi_B}$. Also, if B is commutative and H is cocommutative, we have that $Reg_{\varphi_B}(H^{\otimes n}, B)$ is an abelian group.

We denote by $Reg_{\varphi_B}(H_L, B)$ the set of morphisms $g : H_L \rightarrow B$ such that there exists a morphism $g^{-1} : H_L \rightarrow B$ (the convolution inverse of g) satisfying

$$g * g^{-1} = g^{-1} * g = u_0^{\varphi_B}, \quad g * g^{-1} * g = g, \quad g^{-1} * g * g^{-1} = g^{-1},$$

where $u_0^{\varphi_B} = u_1^{\varphi_B} \circ i_H^L$. Then by (27) we have $u_1^{\varphi_B} = u_0^{\varphi_B} \circ p_H^L$.

Definition 4.6. For an algebra B we define the center of B as a subobject $Z(B)$ of B with a monomorphism $z_B : Z(B) \rightarrow B$ satisfying the identity

$$(143) \quad \mu_B \circ (B \otimes z_B) = \mu_B \circ c_{B,B} \circ (B \otimes z_B)$$

and such that, if $f : A \rightarrow B$ is a morphism for which $\mu_B \circ (B \otimes f) = \mu_B \circ c_{B,B} \circ (B \otimes f)$ holds, there exists a unique morphism $f' : A \rightarrow Z(B)$ satisfying $z_B \circ f' = f$. As a consequence, we obtain that $Z(B)$ is a commutative algebra, where $\eta_{Z(B)}$ is the unique morphism such that

$$(144) \quad z_B \circ \eta_{Z(B)} = \eta_B$$

and $\mu_{Z(B)}$ is the unique morphism such that

$$(145) \quad z_B \circ \mu_{Z(B)} = \mu_B \circ (z_B \otimes z_B).$$

For example, if \mathbf{C} is a closed category with equalizers and α_B and β_B are the unit and the counit, respectively, of the \mathbf{C} -adjunction $B \otimes - \dashv [B, -] : \mathbf{C} \rightarrow \mathbf{C}$, the center of B can be obtained by the following equalizer diagram:

$$\begin{array}{ccccc} Z(B) & \xrightarrow{z_B} & B & \xrightleftharpoons[\theta_B]{\vartheta_B} & [B, B] \end{array}$$

where $\vartheta_B = [B, \mu_B] \circ \alpha_B(B)$ and $\theta_B = [B, \mu_B \circ c_{B,B}] \circ \alpha_B(B)$. Then in the category of modules over a commutative ring the center is an equalizer object.

Finally, note that by (143), composing with the symmetry isomorphism we obtain

$$\mu_B \circ (z_B \otimes B) = \mu_B \circ c_{B,B} \circ (z_B \otimes B).$$

Example 4.7. Let H be cocommutative weak Hopf algebra and let (B, φ_B) be a left weak H -module algebra. Then, $\Pi_H^L = \overline{\Pi}_H^L$ and by (25) and (26) we have that $\mu_B \circ c_{B,B} \circ (u_1^{\varphi_B} \otimes B) = \mu_B \circ (u_1^{\varphi_B} \otimes B)$. Then, $u_1^{\varphi_B}$ factors through $Z(B)$. Therefore, there exists a unique morphism $v_1^{\varphi_B} : H \rightarrow Z(B)$ such that

$$z_B \circ v_1^{\varphi_B} = u_1^{\varphi_B}.$$

Then, taking into account the equality (33), we obtain

$$\mu_B \circ c_{B,B} \circ (u_n^{\varphi_B} \otimes B) = \mu_B \circ c_{B,B} \circ ((u_1^{\varphi_B} \circ m_H^{\otimes n}) \otimes B) = \mu_B \circ ((u_1^{\varphi_B} \circ m_H^{\otimes n}) \otimes B) = \mu_B \circ (u_n^{\varphi_B} \otimes B)$$

and, as a consequence, $u_n^{\varphi_B}$ factors through $Z(B)$. Therefore, there exists a unique morphism $v_n^{\varphi_B} : H \otimes H \rightarrow Z(B)$ such that

$$z_B \circ v_n^{\varphi_B} = u_n^{\varphi_B}.$$

Remark 4.8. Let H be a weak Hopf algebra. Let $\omega : H^{\otimes n} \rightarrow B$ be a morphism. Then, ω factors through the center of B if and only if $\overline{\omega} = \overline{\omega}^{op}$. Therefore, if H is cocommutative and (B, φ_B) is a left weak H -module algebra, $\overline{u_n^{\varphi_B}} = \overline{u_n^{\varphi_B}}^{op}$ for all $n \geq 1$.

Also, if ω factors through the center of B , $\omega * \tau = \tau * \omega$ for all morphism $\tau : H^{\otimes n} \rightarrow B$.

Proposition 4.9. Let H be a cocommutative weak Hopf algebra. Let (B, φ_B) be a left weak H -module algebra and $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Then, φ_B is φ_B -invertible.

Proof. Let h_σ and $h_{\sigma^{-1}}$ be the morphisms defined by

$$h_\sigma = \sigma \circ (H \otimes \lambda_H) \circ \delta_H, \quad h_{\sigma^{-1}} = \sigma^{-1} \circ (H \otimes \lambda_H) \circ \delta_H.$$

Then, $h_\sigma \in \text{Reg}_{\varphi_B}(H, B)$ and $h_\sigma^{-1} = h_{\sigma^{-1}}$. Indeed, first note that

$$\begin{aligned} & h_\sigma * h_{\sigma^{-1}} \\ &= \mu_B \circ (\sigma \otimes \sigma^{-1}) \circ (H \otimes c_{H,H} \otimes H) \circ (\delta_H \otimes ((\lambda_H \otimes \lambda_H) \circ \delta_H)) \circ \delta_H \quad (\text{by the coassociativity and} \\ & \quad \text{the cocommutativity of } \delta_H \text{ and the naturality of } c) \\ &= (\sigma * \sigma^{-1}) \circ (H \otimes \lambda_H) \circ \delta_H \quad (\text{the cocommutativity of } \delta_H \text{ and (19)}) \\ &= u_1^{\varphi_B} \circ \mu_H \circ (H \otimes \lambda_H) \circ \delta_H \quad (\text{by } \sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)) \\ &= u_1^{\varphi_B} \circ \Pi_H^L \quad (\text{by the definition of } \Pi_H^L) \\ &= u_1^{\varphi_B} \quad (\text{by (27)}) \end{aligned}$$

and similarly, $h_{\sigma^{-1}} * h_\sigma = u_1^{\varphi_B}$. Also, by the coassociativity and the cocommutativity of δ_H , the naturality of c , (19) and $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ we have that

$$h_\sigma * h_{\sigma^{-1}} * h_\sigma = (\sigma * \sigma^{-1} * \sigma) \circ (H \otimes \lambda_H) \circ \delta_H = h_\sigma$$

and $h_{\sigma^{-1}} * h_\sigma * h_{\sigma^{-1}} = h_{\sigma^{-1}}$ hold.

Now, let φ_σ be the morphism defined by

$$\varphi_\sigma = \mu_B \circ (\mu_B \otimes B) \circ (h_\sigma \otimes B \otimes h_{\sigma^{-1}}) \circ (H \otimes c_{H,H}) \circ (\delta_H \otimes H).$$

Then, φ_σ is φ_B -invertible with inverse defined by

$$\varphi_\sigma^\dagger = \mu_B \circ (\mu_B \otimes B) \circ (h_{\sigma^{-1}} \otimes B \otimes h_\sigma) \circ (H \otimes c_{H,H}) \circ (\delta_H \otimes H).$$

Indeed:

$$\begin{aligned} & \varphi_\sigma \wedge \varphi_\sigma^\dagger \\ &= \mu_B \circ (B \otimes (\mu_B \circ c_{B,B})) \otimes (((h_\sigma * h_{\sigma^{-1}}) \otimes (h_\sigma * h_{\sigma^{-1}}) \circ \delta_H) \otimes B) \quad (\text{by the coassociativity and the} \\ & \quad \text{cocommutativity of } \delta_H, \text{ the naturality of } c \text{ and the associativity of } \mu_B) \\ &= \mu_B \circ (u_1^{\varphi_B} \otimes (\mu_B \circ c_{B,B} \circ (u_1^{\varphi_B} \otimes B))) \circ (\delta_H \otimes B) \quad (\text{by } h_\sigma \in \text{Reg}_{\varphi_B}(H, B)) \\ &= \mu_B \circ ((u_1^{\varphi_B} * u_1^{\varphi_B}) \otimes B) \quad (\text{by the factorization of } u_1^{\varphi_B} \text{ through the center of } B) \\ &= \mu_B \circ (u_1^{\varphi_B} \otimes B) \quad (\text{by (34)}). \end{aligned}$$

On the other hand, let be the morphism $\varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B))$. For this morphism we have the following

$$\begin{aligned} & \overline{h_\sigma}^{op} \wedge (\varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B))) \\ &= \mu_B \circ (\varphi_B \otimes \sigma) \circ (H \otimes c_{H,B} \otimes H) \circ (\delta_H \otimes (c_{H,B} \circ (H \otimes \varphi_B) \circ ((\delta_H \circ \lambda_H) \otimes B))) \circ (\delta_H \otimes B) \quad (\text{by the} \\ & \quad \text{coassociativity and the cocommutativity of } \delta_H, \text{ the naturality of } c \text{ and (19)}) \\ &= \mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B}) \circ (((H \otimes \lambda_H) \circ \delta_H) \otimes B) \quad (\text{by the cocommutativity of } \delta_H \text{ and} \\ & \quad \text{the naturality of } c) \\ &= \mu_B \circ (B \otimes \varphi_B) \circ (F_\sigma \otimes B) \circ (((H \otimes \lambda_H) \circ \delta_H) \otimes B) \quad (\text{by (77)}) \\ &= \mu_B \circ (B \otimes \varphi_B) \circ (((h_\sigma \otimes \Pi_H^L) \circ \delta_H) \otimes B) \quad (\text{by the cocommutativity of } \delta_H, \text{ the naturality of } c \text{ and (19)}) \\ &= \mu_B \circ (B \otimes \mu_B) \circ (((h_\sigma \otimes u_1^{\varphi_B}) \circ \delta_H) \otimes B) \quad (\text{by (25)}) \\ &= \mu_B \circ ((h_\sigma * u_1^{\varphi_B}) \otimes B) \quad (\text{by the associativity of } \mu_B) \end{aligned}$$

$$= \overline{h_\sigma} \text{ (by } h_\sigma \in \text{Reg}_{\varphi_B}(H, B))$$

and, as a consequence,

$$(146) \quad \varphi_\sigma = \varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B))$$

holds because, on the one hand,

$$\begin{aligned} & \overline{h_{\sigma^{-1}}}^{op} \wedge (\overline{h_\sigma}^{op} \wedge (\varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B)))) \\ &= \overline{h_\sigma * h_{\sigma^{-1}}}^{op} \wedge (\varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B))) \text{ (by associativity of } \wedge \text{ and (iii) of Proposition 4.2)} \\ &= \overline{u_1^{\varphi_B}}^{op} \wedge (\varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B))) \text{ (by } h_\sigma \in \text{Reg}_{\varphi_B}(H, B)) \\ &= (\overline{u_1^{\varphi_B}}^{op} \wedge \varphi_B) \wedge (\varphi_B \circ (\lambda_H \otimes B)) \text{ (by associativity of } \wedge) \\ &= \varphi_B \wedge (\varphi_B \circ (\lambda_H \otimes B)) \text{ (by (vi) of Proposition 4.2)} \end{aligned}$$

and, on the other hand, by the cocommutativity of δ_H , the naturality of c and the associativity of μ_B

$$\overline{h_{\sigma^{-1}}}^{op} \wedge \overline{h_\sigma} = \varphi_\sigma.$$

Finally, define the morphism φ_B^\dagger by

$$\varphi_B^\dagger = (\varphi_B \circ (\lambda_H \otimes B)) \wedge \varphi_{\sigma^{-1}}.$$

Then, by (146) we have

$$\varphi_B \wedge \varphi_B^\dagger = \varphi_\sigma \wedge \varphi_\sigma^\dagger = \mu_B \circ (u_1^{\varphi_B} \otimes B)$$

and therefore φ_B is φ_B -invertible with inverse φ_B^\dagger . \square

Proposition 4.10. *Let H be a cocommutative weak Hopf algebra. Let (B, φ_B) be a left weak H -module algebra and suppose that φ_B is φ_B -invertible. Then, $\varphi_B^{\otimes n}$ is $\varphi_B^{\otimes n}$ -invertible.*

Proof. By assumption the assertion is true for $n = 1$. We will proceed by induction assuming that it is true for $n - 1$, i.e., $\varphi_B^{\otimes(n-1)}$ is $\varphi_B^{\otimes(n-1)}$ -invertible with inverse $\varphi_B^{\otimes(n-1)\dagger}$, and we will prove it for n . Indeed: Define $\varphi_B^{\otimes n\dagger}$ by

$$\varphi_B^{\otimes n\dagger} = \varphi_B^{\otimes(n-1)\dagger} \circ (H^{\otimes(n-1)} \otimes \varphi_B^\dagger) \circ (c_{H, H^{\otimes(n-1)}} \otimes B).$$

Then,

$$\begin{aligned} & \varphi_B^{\otimes n} \wedge \varphi_B^{\otimes n\dagger} \\ &= \varphi_B \circ (H \otimes (\varphi_B^{\otimes(n-1)} \wedge \varphi_B^{\otimes(n-1)\dagger})) \circ (H \otimes H^{\otimes(n-1)} \otimes \varphi_B^\dagger) \circ (H \otimes c_{H, H^{\otimes(n-1)}} \otimes B) \circ (\delta_H \otimes H^{\otimes(n-1)} \otimes B) \\ & \quad \text{(by naturality of } c) \\ &= \varphi_B \circ (H \otimes (\mu_B \circ (u_{n-1}^{\varphi_B} \otimes B))) \circ (H \otimes H^{\otimes(n-1)} \otimes \varphi_B^\dagger) \circ (H \otimes c_{H, H^{\otimes(n-1)}} \otimes B) \circ (\delta_H \otimes H^{\otimes(n-1)} \otimes B) \\ & \quad \text{(by the induction hypothesis)} \\ &= \varphi_B \circ (H \otimes \mu_B) \circ (H \otimes \varphi_B^\dagger \otimes B) \circ (\delta_H \otimes (c_{B, B} \circ (u_{n-1}^{\varphi_B} \otimes B))) \text{ (by the factorization of } u_{n-1}^{\varphi_B} \text{ through} \\ & \quad \text{the center of } B \text{ and the naturality of } c) \\ &= \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H, B} \otimes B) \circ (\delta_H \otimes \varphi_B^\dagger \otimes B) \circ (\delta_H \otimes (c_{B, B} \circ (u_{n-1}^{\varphi_B} \otimes B))) \text{ (by (b1) of} \\ & \quad \text{Definition 1.4)} \\ &= \mu_B \circ ((\varphi_B \wedge \varphi_B^\dagger) \otimes \varphi_B) \circ (H \otimes c_{H, B} \otimes B) \circ (\delta_H \otimes (c_{B, B} \circ (u_{n-1}^{\varphi_B} \otimes B))) \text{ (by naturality of } c \text{ and the} \\ & \quad \text{coassociativity and cocommutativity of } \delta_H) \end{aligned}$$

$$\begin{aligned}
&= \mu_B \circ ((\mu_B \circ (u_1^{\varphi_B} \otimes B)) \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes (c_{B,B} \circ (u_{n-1}^{\varphi_B} \otimes B))) \text{ (by the } \varphi_B\text{-invertibility of } \varphi_B) \\
&= \mu_B \circ (B \otimes (\mu_B \circ (u_1^{\varphi_B} \otimes \varphi_B) \circ (\delta_H \otimes B))) \circ (c_{H,B} \otimes B) \circ (H \otimes (c_{B,B} \circ (u_{n-1}^{\varphi_B} \otimes B))) \text{ (by the naturality of } c, \text{ the associativity of } \mu_B \text{ and the factorization of } u_1^{\varphi_B} \text{ through the center of } B) \\
&= \mu_B \circ (B \otimes \varphi_B) \circ (c_{H,B} \otimes B) \circ (H \otimes (c_{B,B} \circ (u_{n-1}^{\varphi_B} \otimes B))) \text{ (by (52))} \\
&= \mu_B \circ c_{B,B} \circ (u_n^{\varphi_B} \otimes B) \text{ (by the naturality of } c) \\
&= \mu_B \circ (u_n^{\varphi_B} \otimes B) \text{ (by the factorization of } u_n^{\varphi_B} \text{ through the center of } B)
\end{aligned}$$

and, therefore $\varphi_B^{\otimes n}$ is $\varphi_B^{\otimes n}$ -invertible. □

Proposition 4.11. *Let H be a cocommutative weak Hopf algebra. Let (B, φ_B) be a left weak H -module algebra and suppose that φ_B is φ_B -invertible. Then, $\omega \in \text{Reg}_{\varphi_B}(H^{\otimes n}, B)$ satisfies*

$$(147) \quad \overline{\omega} \wedge \varphi_B^{\otimes n} = \overline{\omega}^{op} \wedge \varphi_B^{\otimes n}$$

if and only if it factors through the center of B .

Proof. Assume that (147) holds. Then, by the associativity of μ_B and $\omega \in \text{Reg}_{\varphi_B}(H^{\otimes n}, B)$, we have

$$\overline{\omega} \wedge \varphi_B^{\otimes n} \wedge \varphi_B^{\otimes n \dagger} = \overline{\omega} \wedge (\mu_B \circ (u_n^{\varphi_B} \otimes B)) = \mu_B \circ ((\omega * u_n^{\varphi_B}) \otimes B) = \overline{\omega}.$$

On the other hand,

$$\begin{aligned}
&\overline{\omega}^{op} \wedge \varphi_B^{\otimes n} \wedge \varphi_B^{\otimes n \dagger} \\
&= \overline{\omega}^{op} \wedge (\mu_B \circ (u_n^{\varphi_B} \otimes B)) \text{ (by the } \varphi_B^{\otimes n}\text{-invertibility)} \\
&= \mu_B \circ c_{B,B} \circ (\omega \otimes (\mu_B \circ c_{B,B} \circ (u_n^{\varphi_B} \otimes B))) \circ (\delta_{H^{\otimes n}} \otimes B) \text{ (by the factorization of } u_n^{\varphi_B} \text{ through the center of } B) \\
&= \mu_B \circ (B \otimes (\mu_B \circ c_{B,B})) \circ (c_{B,B} \otimes B) \circ (B \otimes c_{B,B}) \circ (((\omega \otimes u_n^{\varphi_B}) \circ \delta_{H^{\otimes n}}) \otimes B) \text{ (by the naturality of } c \text{ and the associativity of } \mu_B) \\
&= \mu_B \circ c_{B,B} \circ ((u_n^{\varphi_B} * \omega) \otimes B) \text{ (by the naturality of } c \text{ and the cocommutativity of } \delta_H) \\
&= \overline{\omega}^{op} \text{ (by } \omega \in \text{Reg}_{\varphi_B}(H^{\otimes n}, B)).
\end{aligned}$$

Therefore, $\overline{\omega} = \overline{\omega}^{op}$ and, as a consequence, ω factors through the center of B .

Conversely, if ω factors through the center of B , by Remark 4.8, we have that $\overline{\omega} = \overline{\omega}^{op}$ and then (147) holds trivially. □

Proposition 4.12. *Let H be a cocommutative weak Hopf algebra. Let (B, φ_B) be a left weak H -module algebra and suppose that φ_B is φ_B -invertible. Then, if $\omega \in \text{Reg}_{\varphi_B}(H^{\otimes n}, B)$ satisfies (147), ω^{-1} also satisfies (147). Then, as a consequence, ω^{-1} factors through the center of B .*

Proof. By the equalities of Proposition 4.2, Remark 4.8 and Proposition 4.11 we have the following:

$$\overline{\omega} \wedge \overline{\omega^{-1}} = \overline{\omega * \omega^{-1}} = \overline{u_n^{\varphi_B}} = \overline{u_n^{\varphi_B}}^{op} = \overline{\omega^{-1} * \omega}^{op} = \overline{\omega}^{op} \wedge \overline{\omega^{-1}}^{op} = \overline{\omega} \wedge \overline{\omega^{-1}}^{op}.$$

Then, as a consequence, we have that

$$\overline{\omega^{-1}} \wedge \varphi_B^{\otimes n} = \overline{\omega^{-1}} \wedge \overline{u_n^{\varphi_B}} \wedge \varphi_B^{\otimes n} = \overline{\omega^{-1}} \wedge \overline{u_n^{\varphi_B}}^{op} \wedge \varphi_B^{\otimes n} = \overline{u_n^{\varphi_B}}^{op} \wedge \overline{\omega^{-1}} \wedge \varphi_B^{\otimes n} = \overline{u_n^{\varphi_B}} \wedge \overline{\omega^{-1}} \wedge \varphi_B^{\otimes n}$$

$$\begin{aligned}
&= \overline{\omega^{-1}} \wedge \overline{\omega} \wedge \overline{\omega^{-1}} \wedge \varphi_B^{\otimes n} = \overline{\omega^{-1}} \wedge \overline{\omega} \wedge \overline{\omega^{-1}}^{op} \wedge \varphi_B^{\otimes n} = \overline{u_n^{\varphi_B}} \wedge \overline{\omega^{-1}}^{op} \wedge \varphi_B^{\otimes n} = \overline{\omega^{-1}}^{op} \wedge \overline{u_n^{\varphi_B}} \wedge \varphi_B^{\otimes n} \\
&= \overline{\omega^{-1}}^{op} \wedge \varphi_B^{\otimes n}.
\end{aligned}$$

Therefore, ω^{-1} satisfies (147) and, by the previous proposition, ω^{-1} factors through the center of B . \square

Proposition 4.13. *Let H be a cocommutative weak Hopf algebra. Let (B, φ_B) be a left weak H -module algebra and $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Then, φ_B induces a left H -module algebra structure on the center of B , where the action $\varphi_{Z(B)} : H \otimes Z(B) \rightarrow Z(B)$ is the factorization of $\varphi_B \circ (H \otimes z_B) : H \otimes Z(B) \rightarrow B$ through the center of B .*

Proof. First note that, by (143), (b1) of Definition 1.4 and the naturality of c , we obtain that the identity

$$(148) \quad \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_B \otimes z_B \otimes B) = \mu_B \circ c_{B,B} \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_B \otimes z_B \otimes B)$$

holds. Then, on the one hand,

$$\begin{aligned}
&\mu_B \circ (\varphi_B \otimes (\mu_B \circ (u_1^{\varphi_B} \otimes B))) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_B \otimes z_B \otimes B) \\
&= \mu_B \circ (\mu_B \otimes B) \circ (\varphi_B \otimes c_{B,B}) \circ (H \otimes c_{B,B} \otimes B) \circ (((H \otimes u_1^{\varphi_B}) \circ \delta_H) \otimes z_B \otimes B) \text{ (by (143) and the} \\
&\quad \text{naturality of } c) \\
&= \mu_B \circ (\mu_B \otimes B) \circ (\varphi_B \otimes c_{B,B}) \circ (H \otimes c_{B,B} \otimes B) \circ (((c_{B,H} \circ (u_1^{\varphi_B} \otimes H)) \circ \delta_H) \otimes z_B \otimes B) \text{ (by the} \\
&\quad \text{cocommutativity of } \delta_H \text{ and the naturality of } c) \\
&= \mu_B \circ (B \otimes (\mu_B \circ c_{B,B} \circ (u_1^{\varphi_B} \otimes B))) \circ ((c_{H,B} \circ (H \otimes \varphi_B)) \circ (\delta_H \otimes z_B)) \otimes B \text{ (by the naturality of } c \\
&\quad \text{and the associativity of } \mu_B) \\
&= \mu_B \circ ((\mu_B \circ c_{B,B} \circ (u_1^{\varphi_B} \otimes \varphi_B)) \circ (\delta_H \otimes z_B)) \otimes B \text{ (by the naturality of } c, \text{ the associativity of } \mu_B \text{ and (143))} \\
&= \mu_B \circ ((\mu_B \circ (u_1^{\varphi_B} \otimes \varphi_B)) \circ (\delta_H \otimes z_B)) \otimes B \text{ (by (143))} \\
&= \mu_B \circ ((\varphi_B \circ (H \otimes z_B)) \otimes B) \text{ (by (52))}
\end{aligned}$$

and, on the other hand,

$$\begin{aligned}
&\mu_B \circ (\varphi_B \otimes (\mu_B \circ (u_1^{\varphi_B} \otimes B))) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes z_B \otimes B) \\
&= \mu_B \circ (\varphi_B \otimes (\varphi_B \wedge \varphi_B^\dagger)) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes z_B \otimes B) \text{ (by the } \varphi_B\text{-invertibility of } \varphi_B) \\
&= \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes \varphi_B^\dagger) \circ (\delta_H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes z_B \otimes B) \text{ (by the coassociativity of } \delta_H \\
&\quad \text{and the naturality of } c) \\
&= \mu_B \circ c_{B,B} \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes \varphi_B^\dagger) \circ (\delta_H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes z_B \otimes B) \text{ (by (148))} \\
&= \mu_B \circ c_{B,B} \circ (\varphi_B \otimes (\varphi_B \wedge \varphi_B^\dagger)) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes z_B \otimes B) \text{ (by the naturality of } c \text{ and the} \\
&\quad \text{coassociativity of } \delta_H) \\
&= \mu_B \circ (\mu_B \otimes B) \circ (B \otimes c_{B,B}) \circ ((c_{B,B} \circ (\varphi_B \otimes u_1^{\varphi_B})) \circ (H \otimes c_{H,B})) \circ (\delta_H \otimes z_B) \otimes B \text{ (by the} \\
&\quad \varphi_B\text{-invertibility of } \varphi_B \text{ and the naturality of } c) \\
&= \mu_B \circ (\mu_B \otimes B) \circ (B \otimes c_{B,B}) \circ (((u_1^{\varphi_B} \otimes \varphi_B) \circ (\delta_H \otimes z_B)) \otimes B) \text{ (by the naturality of } c \text{ and the} \\
&\quad \text{cocommutativity of } \delta_H) \\
&= \mu_B \circ (B \otimes \mu_B) \circ (c_{B,B} \otimes B) \circ (B \otimes c_{B,B}) \circ (((u_1^{\varphi_B} \otimes \varphi_B) \circ (\delta_H \otimes z_B)) \otimes B) \text{ (by (143) and the} \\
&\quad \text{associativity of } \mu_B) \\
&= \mu_B \circ c_{B,B} \circ ((\mu_B \circ (u_1^{\varphi_B} \otimes \varphi_B)) \circ (\delta_H \otimes z_B)) \otimes B \text{ (by the naturality of } c) \\
&= \mu_B \circ c_{B,B} \circ ((\varphi_B \circ (H \otimes z_B)) \otimes B) \text{ (by (52))}.
\end{aligned}$$

Therefore, as a consequence of the previous equalities, we have that

$$\mu_B \circ ((\varphi_B \circ (H \otimes z_B)) \otimes B) = \mu_B \circ c_{B,B} \circ ((\varphi_B \circ (H \otimes z_B)) \otimes B)$$

and this implies that there exists a unique morphism $\varphi_{Z(B)} : H \otimes Z(B) \rightarrow Z(B)$ such that

$$(149) \quad z_B \circ \varphi_{Z(B)} = \varphi_B \circ (H \otimes z_B).$$

The pair $(Z(B), \varphi_{Z(B)})$ is a left H -module algebra. Indeed, using that z_B is a monomorphism we have that $\varphi_{Z(B)} \circ (\eta_H \otimes Z(B)) = id_{Z(B)}$ because by (149) $z_B \circ \varphi_{Z(B)} \circ (\eta_H \otimes Z(B)) = z_B$ holds. Also, we have the identity

$$(150) \quad \mu_B \circ ((\varphi_B \circ (H \otimes \varphi_B)) \otimes B) \circ (H \otimes H \otimes (c_{B,B} \circ (\sigma \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) = \mu_B \circ (B \otimes \varphi_B) \circ (G_\sigma \otimes z_B)$$

since

$$\begin{aligned} & \mu_B \circ ((\varphi_B \circ (H \otimes \varphi_B)) \otimes B) \circ (H \otimes H \otimes (c_{B,B} \circ (\sigma \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \\ &= \mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes P_{\varphi_B}) \circ (H \otimes H \otimes z_B) \text{ (by the naturality of } c) \\ &= \mu_B \circ (B \otimes \varphi_B) \circ (F_\sigma \otimes z_B) \text{ (by (77))} \\ &= \mu_B \circ c_{B,B} \circ (B \otimes (z_B \circ \varphi_{Z(B)})) \circ (F_\sigma \otimes Z(B)) \text{ (by (143) and (149))} \\ &= \mu_B \circ (\varphi_B \otimes B) \circ (H \otimes c_{B,B}) \otimes (((\mu_H \otimes \sigma) \circ c_{H^{\otimes 2}, H^{\otimes 2}} \circ \delta_{H^{\otimes 2}}) \otimes z_B) \text{ (by the naturality of } c) \\ &= \mu_B \circ (\varphi_B \otimes B) \circ (H \otimes c_{B,B}) \otimes (G_\sigma \otimes z_B) \text{ (by the cocommutativity of } \delta_{H^{\otimes 2}}). \end{aligned}$$

Then,

$$\begin{aligned} & \mu_B \circ ((\mu_B \circ ((\varphi_B \circ (H \otimes \varphi_B)) \otimes B) \circ (H \otimes H \otimes (c_{B,B} \circ (\sigma \otimes B)))) \circ (\delta_{H^{\otimes 2}} \otimes B)) \otimes B \\ & \circ (H \otimes H \otimes (c_{B,B} \circ (\sigma^{-1} \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \\ &= \mu_B \circ ((\varphi_B \circ (H \otimes \varphi_B)) \otimes B) \circ (H \otimes H \otimes (c_{B,B} \circ ((\sigma * \sigma^{-1}) \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \text{ (by the coassociativity} \\ & \text{ of } \delta_{H^{\otimes 2}}) \\ &= \mu_B \circ ((\varphi_B \circ (H \otimes \varphi_B)) \otimes B) \circ (H \otimes H \otimes (c_{B,B} \circ (u_2^{\varphi_B} \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \text{ (by } \sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)) \\ &= \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes ((\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ (\delta_H \otimes z_B \otimes \eta_B))) \text{ (by} \\ & \text{ the naturality of } c) \\ &= \varphi_B \circ (H \otimes (\varphi_B \circ (H \otimes (\mu_B \circ (z_B \otimes \eta_B))))) \text{ (by (b1) of Definition 1.4)} \\ &= \varphi_B \circ (H \otimes (\varphi_B \circ (H \otimes z_B))) \text{ (by the properties of } \eta_B) \\ &= z_B \circ \varphi_{Z(B)} \circ (H \otimes \varphi_{Z(B)}) \text{ (by (149))} \end{aligned}$$

and, on the other hand,

$$\begin{aligned} & \mu_B \circ ((\mu_B \circ (B \otimes \varphi_B) \circ (G_\sigma \otimes B)) \otimes B) \circ (H \otimes H \otimes (c_{B,B} \circ (\sigma^{-1} \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \\ &= \mu_B \circ (\varphi_B \otimes B) \circ (\mu_H \otimes (c_{B,B} \circ ((\sigma * \sigma^{-1}) \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \text{ (by the coassociativity of } \delta_{H^{\otimes 2}}) \\ &= \mu_B \circ (\varphi_B \otimes B) \circ (\mu_H \otimes (c_{B,B} \circ (u_2^{\varphi_B} \otimes B))) \circ (\delta_{H^{\otimes 2}} \otimes z_B) \text{ (by } \sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)) \\ &= \mu_B \circ (\varphi_B \otimes u_1^{\varphi_B}) \circ (H \otimes c_{H,B}) \circ (((\mu_H \otimes \mu_H) \circ \delta_{H^{\otimes 2}}) \otimes z_B) \text{ (by the naturality of } c) \\ &= \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H,B} \otimes B) \circ ((\delta_H \circ \mu_H) \otimes z_B \otimes \eta_B) \text{ (by (a1) of Definition 1.1)} \\ &= \varphi_B \circ (\mu_H \otimes (\mu_B \circ (z_B \otimes \eta_B))) \text{ (by (b1) of Definition 1.4)} \\ &= \varphi_B \circ (\mu_H \otimes z_B) \text{ (by the properties of } \eta_B) \\ &= z_B \circ \varphi_{Z(B)} \circ (\mu_H \otimes Z(B)) \text{ (by (149)).} \end{aligned}$$

Therefore, as a consequence of (150), we have that $\varphi_{Z(B)} \circ (H \otimes \varphi_{Z(B)}) = \varphi_{Z(B)} \circ (\mu_H \otimes Z(B))$ and this implies that $(Z(B), \varphi_{Z(B)})$ is a left H -module. Finally, it is a left H -module algebra because composing with the monomorphism z_B we have

$$\begin{aligned} & z_B \circ \mu_{Z(B)} \circ (\varphi_{Z(B)} \otimes \varphi_{Z(B)}) \circ (H \otimes c_{H, Z(B)} \otimes Z(B)) \circ (\delta_H \otimes Z(B) \otimes Z(B)) \\ &= \mu_B \circ (\varphi_B \otimes \varphi_B) \circ (H \otimes c_{H, B} \otimes B) \circ (\delta_H \otimes z_B \otimes z_B) \text{ (by (145), (149) and the naturality of } c) \\ &= \varphi_B \circ (H \otimes (\mu_B \circ (z_B \otimes \eta_B))) \text{ (by (b1) of Definition 1.4)} \\ &= \varphi_B \circ (H \otimes (z_B \circ \mu_{Z(B)})) \text{ (by (145))} \\ &= z_B \circ \varphi_{Z(B)} \circ (H \otimes \mu_{Z(B)}) \text{ (by (149))} \end{aligned}$$

and

$$\begin{aligned} & z_B \circ \varphi_{Z(B)} \circ (H \otimes u_1^{\varphi_{Z(B)}}) \\ &= \varphi_B \circ (H \otimes u_1^{\varphi_B}) \text{ (by (144), (149))} \\ &= u_1^{\varphi_B} \circ \mu_H \text{ (by (b3) of Definition 1.4)} \\ &= z_B \circ u_1^{\varphi_{Z(B)}} \circ \mu_H \text{ (by (144), (149)).} \end{aligned}$$

□

Remark 4.14. Note that, under the conditions of the previous proposition, the equality

$$(151) \quad z_B \circ u_1^{\varphi_{Z(B)}} = u_1^{\varphi_B}$$

holds.

4.15. Let H be a cocommutative weak Hopf algebra. By [3, Theorem 3.1] we know that, if (A, φ_A) is a left weak H -module algebra and $\sigma \in \text{Reg}_{\varphi_A}(H^{\otimes 2}, A)$ satisfies the twisted condition (77), (A, φ_A) is a left H -module algebra if and only if the morphism σ factorizes through the center of A . Moreover, by [3, Corollary 3.1], (A, φ_A) is a left H -module algebra if and only if the morphism $u_2^{\varphi_A}$ satisfies the twisted condition (77).

Proposition 4.16. Let H be a cocommutative weak Hopf algebra and let (B, φ_B) be a left weak H -module algebra. Let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Then, $\alpha \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfies the twisted condition (77) if, and only if, there exists $\tau \in \text{Reg}_{\varphi_{Z(B)}}(H^{\otimes 2}, Z(B))$ such that

$$(152) \quad \alpha = (z_B \circ \tau) * \sigma.$$

Proof. Suppose that α satisfies the twisted condition (77). We will see that $\sigma * \alpha^{-1}$ factors through the center of B . Following Proposition 4.11, to prove it we will see that $\overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2} = \overline{\sigma * \alpha^{-1}}^{op} \wedge \varphi_B^{\otimes 2}$. First, note that

$$\varphi_B \circ (\mu_H \otimes B) = \overline{\sigma^{-1}} \wedge \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2}$$

because

$$\begin{aligned} & \varphi_B \circ (\mu_H \otimes B) \\ &= \overline{u_2^{\varphi_B}} \wedge (\varphi_B \circ (\mu_H \otimes B)) \text{ (by (a1) of Definition 1.1 and (52))} \\ &= \overline{\sigma^{-1}} \wedge \overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B)) \text{ (by } \sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B) \text{ and (i) of Proposition 4.2)} \\ &= \overline{\sigma} * \overline{\alpha^{-1}}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (140)).} \end{aligned}$$

Thus, for α we have the same identity and then

$$(153) \quad \overline{\sigma^{-1}} \wedge \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} = \overline{\alpha^{-1}} \wedge \overline{\alpha}^{op} \wedge \varphi_B^{\otimes 2}$$

holds. As a consequence,

$$(154) \quad \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} = \overline{\alpha}^{op} \wedge \overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2}$$

also holds since

$$\begin{aligned} & \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} \\ &= \overline{\sigma}^{op} \wedge \overline{u_2^{\varphi_B}} \wedge \varphi_B^{\otimes 2} \text{ (by (v) of Proposition 4.2)} \\ &= \overline{u_2^{\varphi_B}} \wedge \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (iv) of Proposition 4.2)} \\ &= \overline{\sigma} \wedge \overline{\sigma^{-1}} \wedge \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by } \sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B) \text{ and (i) of Proposition 4.2)} \\ &= \overline{\sigma} \wedge \overline{\alpha^{-1}} \wedge \overline{\alpha}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (153))} \\ &= \overline{\alpha}^{op} \wedge \overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2} \text{ (by (iv) and (i) of Proposition 4.2).} \end{aligned}$$

Therefore,

$$\begin{aligned} & \overline{\sigma * \alpha^{-1}}^{op} \wedge \varphi_B^{\otimes 2} \\ &= \overline{\alpha^{-1}}^{op} \wedge \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (iii) of Proposition 4.2)} \\ &= \overline{\alpha^{-1}}^{op} \wedge \overline{\alpha}^{op} \wedge \overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2} \text{ (by 154)} \\ &= \overline{u_2^{\varphi_B}}^{op} \wedge \overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2} \text{ (by } \alpha \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B) \text{ and (iii) of Proposition 4.2)} \\ &= \overline{u_2^{\varphi_B}} \wedge \overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2} \text{ (by Remark 4.8)} \\ &= \overline{\sigma * \alpha^{-1}} \wedge \overline{u_2^{\varphi_B}} \wedge \varphi_B^{\otimes 2} \text{ (by the factorization of } u_2^{\varphi_B} \text{ through the center of } B) \\ &= \overline{\sigma * \alpha^{-1}} \wedge \varphi_B^{\otimes 2} \text{ (by (v) of Proposition 4.2)} \end{aligned}$$

and this implies that $\sigma * \alpha^{-1}$ factors through the center of B . Then, by Proposition 4.12, the morphism $(\sigma * \alpha^{-1})^{-1} = \alpha * \sigma^{-1}$ also factors through the center of B . If τ is the factorization, we have that $z_B \circ \tau = \alpha * \sigma^{-1}$. Then, (152) holds.

Conversely, if (152) holds for $\tau \in \text{Reg}_{\varphi_{Z(B)}}(H^{\otimes 2}, Z(B))$, we have that

$$\begin{aligned} & \overline{\alpha}^{op} \wedge \varphi_B^{\otimes 2} \\ &= (\overline{z_B \circ \tau}) * \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (152))} \\ &= \overline{\sigma}^{op} \wedge \overline{z_B \circ \tau}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (iii) of Proposition 4.2)} \\ &= \overline{\sigma}^{op} \wedge \overline{z_B \circ \tau} \wedge \varphi_B^{\otimes 2} \text{ (by the factorization through the center of } B) \\ &= \overline{z_B \circ \tau} \wedge \overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2} \text{ (by (iv) of Proposition 4.2)} \\ &= \overline{z_B \circ \tau} \wedge \overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B)) \text{ (by (140))} \\ &= (\overline{z_B \circ \tau}) \circ \overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B)) \text{ (by (i) of Proposition 4.2)} \end{aligned}$$

and, therefore, α satisfies the twisted condition. \square

5. COHOMOLOGICAL OBSTRUCTIONS IN A WEAK SETTING

In the beginning of this section we review the basic facts about the Sweedler cohomology in a weak setting. This cohomology was introduced in [2] as a generalization of the classical Sweedler cohomology for Hopf algebras [17]. The groups $\text{Reg}_{\varphi_B}(H_L, B)$ and $\text{Reg}_{\varphi_B}(H^{\otimes n}, B)$, introduced in the previous section, will be the objects of the corresponding cosimplicial complex.

5.1. Let H be a cocommutative weak Hopf algebra and let (B, φ_B) be a left weak H -module algebra. Following [2] we define the coface operators as the group morphisms

$$\begin{aligned} \partial_{0,i} : \text{Reg}_{\varphi_B}(H_L, B) &\rightarrow \text{Reg}_{\varphi_B}(H, B), \quad i \in \{0, 1\} \\ \partial_{0,0}(g) &= \varphi_B \circ (H \otimes (g \circ p_L \circ \Pi_H^R)) \circ \delta_H, \quad \partial_{0,1}(g) = g \circ p_L, \\ \partial_{k,i} : \text{Reg}_{\varphi_B}(H^{\otimes k}, B) &\rightarrow \text{Reg}_{\varphi_B}(H^{\otimes(k+1)}, B), \quad k \geq 1, \quad i \in \{0, 1, \dots, k+1\} \\ \partial_{k,i}(\sigma) &= \begin{cases} \varphi_B \circ (H \otimes \sigma), & i = 0 \\ \sigma \circ (H^{i-1} \otimes \mu_H \otimes H^{k-i}), & i \in \{1, \dots, k\} \\ \sigma \circ (H^{\otimes(k-1)} \otimes (\mu_H \circ (H \otimes \Pi_H^L))), & i = k+1, \end{cases} \end{aligned}$$

On the other hand, we define the codegeneracy operators by $s_{1,0} : \text{Reg}_{\varphi_B}(H, B) \rightarrow \text{Reg}_{\varphi_B}(H_L, B)$,

$$s_{1,0}(h) = h \circ i_L,$$

and $s_{k+1,i} : \text{Reg}_{\varphi_B}(H^{\otimes(k+1)}, B) \rightarrow \text{Reg}_{\varphi_B}(H^{\otimes k}, B)$, $k \geq 1$, $i \in \{0, 1, \dots, k\}$,

$$s_{k+1,i}(\sigma) = \sigma \circ (H^{\otimes i} \otimes \eta_H \otimes H^{\otimes(k-i)}).$$

Taking into account the codegeneracy operators, we define the groups

$$\text{Reg}_{\varphi_B}^+(H^{\otimes(k+1)}, B) = \bigcap_{i=0}^k \text{Ker}(s_{k+1,i}),$$

$$\text{Reg}_{\varphi_B}^+(H_L, B) = \{g \in \text{Reg}_{\varphi_B}(H_L, B) ; g \circ \eta_{H_L} = \eta_B\}.$$

Note that $\text{Reg}_{\varphi_B}^+(H, B)$ is the group $\text{Reg}_{\varphi_B}^t(H, B)$ introduced in Definition 3.5 because

$$\begin{aligned} \text{Reg}_{\varphi_B}^+(H, B) &= \text{Ker}(s_{1,0}) = \{h \in \text{Reg}_{\varphi_B}(H, B) ; h \circ i_H^L = u_0^{\varphi_B}\} \\ &= \{h \in \text{Reg}_{\varphi_A}(H, B) ; h \circ \Pi_H^L = u_1^{\varphi_B}\} = \{h \in \text{Reg}_{\varphi_B}(H, B) ; h \circ \eta_H = \eta_B\} = \text{Reg}_{\varphi_B}^t(H, B). \end{aligned}$$

Also, $\text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ is the subgroup of $\text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ formed by the elements satisfying the normal condition (84) because

$$\begin{aligned} \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B) &= \text{Ker}(s_{2,0}) \cap \text{Ker}(s_{2,1}) \\ &= \{\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B) ; \sigma \circ (\eta_H \otimes H) = \sigma \circ (H \otimes \eta_H) = u_1^{\varphi_B}\} \end{aligned}$$

and finally

$$\begin{aligned} \text{Reg}_{\varphi_B}^+(H^{\otimes 3}, B) &= \text{Ker}(s_{3,0}) \cap \text{Ker}(s_{3,1}) \cap \text{Ker}(s_{3,2}) \\ &= \{\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 3}, B) ; \sigma \circ (\eta_H \otimes H \otimes H) = \sigma \circ (H \otimes \eta_H \otimes H) = \sigma \circ (H \otimes H \otimes \eta_H) = u_2^{\varphi_B}\}. \end{aligned}$$

5.2. Let H be a cocommutative weak Hopf algebra. If (A, φ_A) is a left H -module algebra, by [2] the groups $Reg_{\varphi_A}(H_L, A)$ and $Reg_{\varphi_A}(H^{\otimes n}, A)$, $n \geq 1$ are the objects of a cosimplicial complex of groups with the previous coface and codegeneracy operators. In this case,

$$D_{\varphi_A}^k = \partial_{k,0} * \partial_{k,1}^{-1} * \cdots * \partial_{k,k+1}^{(-1)^{k+1}}$$

denote the coboundary morphisms of the cochain complex

$$\begin{aligned} Reg_{\varphi_A}(H_L, A) &\xrightarrow{D_{\varphi_A}^0} Reg_{\varphi_A}(H, A) \xrightarrow{D_{\varphi_A}^1} Reg_{\varphi_A}(H^{\otimes 2}, A) \xrightarrow{D_{\varphi_A}^2} \cdots \\ \cdots &\xrightarrow{D_{\varphi_A}^{k-1}} Reg_{\varphi_A}(H^{\otimes k}, A) \xrightarrow{D_{\varphi_A}^k} Reg_{\varphi_A}(H^{\otimes(k+1)}, A) \xrightarrow{D_{\varphi_A}^{k+1}} \cdots \end{aligned}$$

associated to the cosimplicial complex $Reg_{\varphi_A}(H^{\otimes \bullet}, A)$.

5.3. Let H be a weak Hopf algebra. If (B, φ_B) is a left weak H -module algebra and if $\sigma \in Reg_{\varphi_B}(H^{\otimes 2}, B)$, by [12, Proposition 5.5], the morphism $E(\sigma) : H^{\otimes 3} \rightarrow B$ defined by $E(\sigma) = \sigma \otimes \varepsilon_H$ satisfies the following identities:

$$(155) \quad E(\sigma) * u_3^{\varphi_B} = u_3^{\varphi_B} * E(\sigma) = \partial_{2,3}(\sigma).$$

Then, using that $\partial_{2,3}$ is a group morphism, we have

$$u_3^{\varphi_B} = \partial_{2,3}(\sigma)^{-1} * u_3^{\varphi_B} * E(\sigma) = \partial_{2,3}(\sigma)^{-1} * E(\sigma).$$

Therefore,

$$(156) \quad \partial_{2,3}(\sigma) = E(\sigma).$$

Similarly,

$$(157) \quad \partial_{2,3}(\sigma^{-1}) = E(\sigma^{-1}) * u_3^{\varphi_B} = u_3^{\varphi_B} * E(\sigma) = E(\sigma^{-1}),$$

where $E(\sigma^{-1}) = \sigma^{-1} \otimes \varepsilon_H$.

Then, if (B, φ_B) is a left H -module algebra and H is cocommutative, by (155), the second coboundary morphism of the cosimplicial complex $Reg_{\varphi_B}(H^{\otimes \bullet}, B)$ admits the following form:

$$(158) \quad D_{\varphi_B}^2(\sigma) = \partial_{2,0}(\sigma) * \partial_{2,1}(\sigma^{-1}) * \partial_{2,2}(\sigma) * E(\sigma^{-1}).$$

5.4. Let H be a cocommutative weak Hopf algebra. If (A, φ_A) is a commutative left H -module algebra, $(Reg_{\varphi_A}(H^{\otimes \bullet}, A), D_{\varphi_A}^{\bullet})$ gives the Sweedler cohomology of H in (A, φ_A) . Therefore, the k th group will be defined by

$$\mathcal{H}_{\varphi_A}^k(H, A) = \frac{Ker(D_{\varphi_A}^k)}{Im(D_{\varphi_A}^{k-1})}$$

for $k \geq 1$.

The normalized cochain subcomplex of A , denoted by $(Reg_{\varphi_A}^+(H^{\otimes \bullet}, B), D_{\varphi_A}^{\bullet+})$, is defined by the groups $Reg_{\varphi_A}^+(H^{\otimes(k+1)}, A)$, $Reg_{\varphi_A}^+(H_L, A)$ with $D_{\varphi_A}^{k+}$ the restriction of $D_{\varphi_A}^k$ to $Reg_{\varphi_A}^+(H^{\otimes \bullet}, A)$.

We have that $(Reg_{\varphi_A}^+(H^{\otimes \bullet}, A), D_{\varphi_A}^{\bullet+})$, is a subcomplex of $(Reg_{\varphi_A}(H^{\otimes \bullet}, A), D_{\varphi_A}^{\bullet})$ and the injection map between $(Reg_{\varphi_A}^+(H^{\otimes \bullet}, A), D_{\varphi_A}^{\bullet+})$ and $(Reg_{\varphi_A}(H^{\otimes \bullet}, A), D_{\varphi_A}^{\bullet})$ induces an isomorphism of cohomology

$$I_{\varphi_A}^k : \mathcal{H}_{\varphi_A}^k(H, A) \rightarrow \mathcal{H}_{\varphi_A}^{k+}(H, A).$$

5.5. Assume that H is a weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma, \tau \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfying the twisted condition (77) and the 2-cocycle condition (78). Then by Theorem 3.7, $(B \otimes H, \mu_{B \otimes \varphi_B H})$ and $(B \otimes H, \mu_{B \otimes \tau_{\varphi_B H}})$ are equivalent if, and only if, there exists $h \in \text{Reg}_{\varphi_B}^+(H, B)$ satisfying (138) and (132). Then, if H is cocommutative, by [2, Corollary 4.8, Theorem 4.9], $(B \otimes H, \mu_{B \otimes \varphi_B H})$ and $(B \otimes H, \mu_{B \otimes \tau_{\varphi_B H}})$ are equivalent if, and only if, there exists $h \in \text{Reg}_{\varphi_B}^+(H, B)$ such that the equalities (138) and

$$(159) \quad \sigma * \partial_{1,1}(h) = \partial_{1,0}(h) * \partial_{1,2}(h) * \tau,$$

hold. Note that the equality (138) is always true if B is commutative and H is cocommutative. Then, under these conditions, if (B, φ_B) is a left H -module algebra, the equivalence between two weak crossed products $(B \otimes H, \mu_{B \otimes \varphi_B H})$ and $(B \otimes H, \mu_{B \otimes \tau_{\varphi_B H}})$ is determined by the existence of h in $\text{Reg}_{\varphi_B}^+(H, B)$ satisfying the equality (159). In this case, if σ and $\tau \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$, (159) is equivalent to say that

$$\sigma * \tau^{-1} \in \text{Im}(D_{\varphi_B}^{1+}),$$

i.e., $[\sigma] = [\tau]$ in $\mathcal{H}_{\varphi_B}^{2+}(H, B)$.

5.6. Let H be a weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$. Then, using the coface operators, it is an easy exercise to prove that σ satisfy the cocycle condition (78) if and only if

$$(160) \quad \partial_{2,0}(\sigma) * \partial_{2,2}(\sigma) = \partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)$$

holds. Then, by (155), we have that σ satisfy the cocycle condition (78) if and only if σ satisfies the equality

$$(161) \quad \partial_{2,0}(\sigma) * \partial_{2,2}(\sigma) = E(\sigma) * \partial_{2,1}(\sigma).$$

Definition 5.7. Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$. We define the pre-obstruction of σ as the morphism $w_\sigma : H^{\otimes 3} \rightarrow B$, where

$$w_\sigma = \partial_{2,0}(\sigma) * \partial_{2,2}(\sigma) * \partial_{2,1}(\sigma)^{-1} * \partial_{2,3}(\sigma)^{-1}.$$

The using that $\partial_{2,1}(\sigma)^{-1} = \partial_{2,1}(\sigma^{-1})$ and $\partial_{2,3}(\sigma)^{-1} = \partial_{2,3}(\sigma^{-1})$, by the previous considerations, we have that σ satisfies the cocycle condition (78) if and only if $w_\sigma = u_3^{\varphi_B}$. Also, note that by (157), we have

$$w_\sigma = \partial_{2,0}(\sigma) * \partial_{2,2}(\sigma) * \partial_{2,1}(\sigma^{-1}) * E(\sigma^{-1}).$$

Note that $\omega_\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 3}, B)$ and it is easy to show that

$$(162) \quad \omega_\sigma = S_\sigma * R_\sigma,$$

where

$$S_\sigma = \partial_{2,0}(\sigma) * \partial_{2,2}(\sigma) = \mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes F_\sigma)$$

and

$$R_\sigma = \partial_{2,1}(\sigma^{-1}) * E(\sigma^{-1}) = \mu_B \circ (\sigma^{-1} \otimes B) \circ (H \otimes c_{B,H}) \circ (G_{\sigma^{-1}} \otimes H),$$

are morphisms in $\text{Reg}_{\varphi_B}(H^{\otimes 3}, B)$, where

$$S_\sigma^{-1} = \mu_B \circ (\sigma^{-1} \otimes \varphi_B) \circ (H \otimes c_{H,H} \otimes B) \circ (\delta_H \otimes G_{\sigma^{-1}})$$

and

$$R_\sigma^{-1} = \mu_B \circ (B \otimes \sigma) \circ (F_\sigma \otimes H).$$

Proposition 5.8. *Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$. Let ω_σ be the pre-obstruction of σ . Then*

$$(163) \quad \partial_{3,4}(\omega_\sigma) = \omega_\sigma \otimes \varepsilon_H.$$

Proof. Note that by (162) we know that $\omega_\sigma = S_\sigma * R_\sigma$. Then, if we prove that

$$\partial_{3,4}(S_\sigma) = S_\sigma \otimes \varepsilon_H, \quad \partial_{3,4}(R_\sigma) = R_\sigma \otimes \varepsilon_H$$

hold, we obtain (163) since

$$\partial_{3,4}(\omega_\sigma) = \partial_{3,4}(S_\sigma * R_\sigma) = \partial_{3,4}(S_\sigma) * \partial_{3,4}(R_\sigma) = (S_\sigma \otimes \varepsilon_H) * (R_\sigma \otimes \varepsilon_H) = (S_\sigma * R_\sigma) \otimes \varepsilon_H = \omega_\sigma \otimes \varepsilon_H.$$

First note that

$$\begin{aligned} & \partial_{3,4}(S_\sigma) \\ &= \mu_B \circ (B \otimes \sigma) \circ (P_{\varphi_B} \otimes H) \circ (H \otimes ((\sigma \otimes \mu_H) \circ (H \otimes c_{H,H} \otimes H) \circ (\delta_H \otimes ((\mu_H \circ (H \otimes \Pi_H^L)) \otimes (\mu_H \circ (H \otimes \Pi_H^L))) \circ \delta_{H^{\otimes 2}})))) \text{ (by (i) of [2, Proposition 2.6], (a1) of Definition 1.1 and the naturality of } c) \\ &= \mu_B \circ (B \otimes \partial_{3,2}(\partial_{2,3}(\sigma))) \circ ((P_{\varphi_B} \circ (H \otimes \partial_{2,3}(\sigma))) \otimes H \otimes H \otimes H) \circ (H \otimes \delta_{H^{\otimes 3}}) \text{ (by the naturality of } c \text{ and the associativity of } \mu_H) \\ &= S_\sigma \otimes \varepsilon_H \text{ (by (156) and the naturality of } c). \end{aligned}$$

Finally, $\partial_{3,4}(R_\sigma) = R_\sigma \otimes \varepsilon_H$ holds because:

$$\begin{aligned} & \partial_{3,4}(R_\sigma) \\ &= \mu_B \circ (\partial_{2,3}(\sigma^{-1}) \otimes B) \circ (H \otimes H \otimes c_{B,H}) \circ (H \otimes c_{B,H} \otimes H) \circ (G_{\sigma^{-1}} \otimes H \otimes H) \text{ (by the naturality of } c) \\ &= R_\sigma \otimes \varepsilon_H \text{ (by the naturality of } c \text{ and (156) for } \sigma^{-1}). \end{aligned}$$

□

Proposition 5.9. *Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Then, the pre-obstruction of σ factors through the center of B .*

Proof. We will use Proposition 4.11 to obtain that ω_σ factors through the center of B . To prove that

$$\overline{\omega_\sigma} \wedge \varphi_B^{\otimes 3} = \overline{\omega_\sigma}^{op} \wedge \varphi_B^{\otimes 3}$$

we first see

$$(164) \quad \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} = \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B))$$

and

$$(165) \quad \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} = \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)).$$

Indeed:

$$\begin{aligned}
& \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} \\
&= \overline{\partial_{2,2}(\sigma)}^{op} \wedge \overline{\partial_{2,0}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} \text{ (by (iii) of Proposition 4.2)} \\
&= \overline{\partial_{2,2}(\sigma)}^{op} \wedge (\varphi_B \circ (H \otimes (\overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2}))) \text{ (by the naturality of } c, \text{ (b1) of Definition 1.4 and the cocommutativity of } \delta_H) \\
&= \overline{\partial_{2,2}(\sigma)}^{op} \wedge (\varphi_B \circ (H \otimes (\overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B)))) \text{ (by (140))} \\
&= \overline{\partial_{2,2}(\sigma)}^{op} \wedge \overline{\partial_{2,0}(\sigma)} \wedge (\varphi_B \circ (H \otimes (\varphi_B \circ (\mu_H \otimes B)))) \text{ (by (vii) of Proposition 4.2)} \\
&= \overline{\partial_{2,0}(\sigma)} \wedge \overline{\partial_{2,2}(\sigma)}^{op} \wedge (\varphi_B^{\otimes 2} \circ (H \otimes \mu_H \otimes B)) \text{ (by (iv) of Proposition 4.2)} \\
&= \overline{\partial_{2,0}(\sigma)} \wedge \overline{\partial_{2,2}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by (a1) of Definition 1.1, the naturality of } c \text{ and (140))} \\
&= \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by the naturality of } c)
\end{aligned}$$

and

$$\begin{aligned}
& \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} \\
&= \overline{\partial_{2,1}(\sigma)}^{op} \wedge \overline{\partial_{2,3}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} \text{ (by (iii) of Proposition 4.2)} \\
&= \overline{\partial_{2,1}(\sigma)}^{op} \wedge ((\overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2}) \circ (H \otimes H \otimes \varphi_B)) \text{ (by the naturality of } c, \text{ the counit properties and (156))} \\
&= \overline{\partial_{2,1}(\sigma)}^{op} \wedge ((\overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B))) \circ (H \otimes H \otimes \varphi_B)) \text{ (by (140))} \\
&= \overline{\partial_{2,1}(\sigma)}^{op} \wedge \overline{\partial_{2,3}(\sigma)} \wedge (\varphi_B \circ (\mu_H \otimes \varphi_B)) \text{ (by the naturality of } c \text{ and the counit properties)} \\
&= \overline{\partial_{2,3}(\sigma)} \wedge \overline{\partial_{2,1}(\sigma)}^{op} \wedge (\varphi_B \circ (\mu_H \otimes \varphi_B)) \text{ (by (iv) of Proposition 4.2)} \\
&= \overline{\partial_{2,3}(\sigma)} \wedge ((\overline{\sigma}^{op} \wedge \varphi_B^{\otimes 2}) \circ (\mu_H \otimes H \otimes B)) \text{ (by the naturality of } c \text{ and (a1) of Definition 1.1)} \\
&= \overline{\partial_{2,3}(\sigma)} \wedge ((\overline{\sigma} \wedge (\varphi_B \circ (\mu_H \otimes B))) \circ (\mu_H \otimes H \otimes B)) \text{ (by (140))} \\
&= \overline{\partial_{2,3}(\sigma)} \wedge \overline{\partial_{2,1}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by the naturality of } c \text{ and (a1) of Definition 1.1)} \\
&= \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by (i) of Proposition 4.2)}.
\end{aligned}$$

Also,

$$\begin{aligned}
& \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)} \wedge \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \\
&= \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by (iv) of Proposition 4.2)} \\
&= \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} \text{ (by 165)} \\
&= \overline{\partial_{2,3}(\sigma) * \partial_{2,1}(\sigma) * \partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge \varphi_B^{\otimes 3} \text{ (by (iii) of Proposition 4.2)} \\
&= \overline{u_3^{\varphi_B} \circ \varphi_B^{\otimes 3}} \wedge \varphi_B^{\otimes 3} \text{ (by the property of group morphism for } \partial_{2,1} \text{ and } \partial_{2,3}) \\
&= \varphi_B^{\otimes 3} \text{ (by (v) of Proposition 4.2)}
\end{aligned}$$

and, as a consequence, the following identity holds:

$$(166) \quad \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) = \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})} \wedge \varphi_B^{\otimes 3}.$$

Therefore,

$$\begin{aligned}
& \overline{\omega_\sigma}^{op} \wedge \varphi_B^{\otimes 3} \\
&= \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)}^{op} \wedge \varphi_B^{\otimes 3} \text{ (by (iii) of Proposition 4.2)} \\
&= \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by (164))} \\
&= \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)} \wedge \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})}^{op} \wedge (\varphi_B \circ (m_H^{\otimes 3} \otimes B)) \text{ (by (iv) of Proposition 4.2)} \\
&= \overline{\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)} \wedge \overline{\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})} \wedge \varphi_B^{\otimes 3} \text{ (by (166))} \\
&= \overline{\omega_\sigma} \wedge \varphi_B^{\otimes 3} \text{ (by (i) of Proposition 4.2)}
\end{aligned}$$

□

Definition 5.10. Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). The obstruction of σ is defined as the unique morphism $\theta_\sigma : H^{\otimes 3} \rightarrow Z(B)$ such that $z_B \circ \theta_\sigma = \omega_\sigma$, where ω_σ is the pre-obstruction of σ .

Note that, by the previous proposition, we can assure that θ_σ exists. Also, $\theta_\sigma \in \text{Reg}_{\varphi_{Z(B)}}(H^{\otimes 3}, Z(B))$.

Theorem 5.11. Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Then, the pre-obstruction of σ is a 3-cocycle, i.e., the following equality holds:

$$(167) \quad \partial_{3,0}(\omega_\sigma) * \partial_{3,2}(\omega_\sigma) * \partial_{3,4}(\omega_\sigma) = \partial_{3,1}(\omega_\sigma) * \partial_{3,3}(\omega_\sigma).$$

Proof. In order to prove the theorem we will see some equalities. First of all observe that by the definition of the pre-obstruction ω_σ we have:

$$(168) \quad \partial_{2,0}(\sigma) * \partial_{2,2}(\sigma) = \omega_\sigma * \partial_{2,3}(\sigma) * \partial_{2,1}(\sigma).$$

Now using that $\partial_{3,2}$ is group morphism we have that

$$(169) \quad \partial_{3,2}(\partial_{2,0}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) = \partial_{3,2}(\omega_\sigma) * \partial_{3,2}(\partial_{2,3}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)).$$

But observe that, by the associativity of μ_H and (156), we have

$$(170) \quad \partial_{3,2}(\partial_{2,3}(\sigma)) = \partial_{3,4}(\partial_{2,2}(\sigma)),$$

$$(171) \quad \partial_{3,3}(\partial_{2,2}(\sigma)) = \partial_{3,2}(\partial_{2,2}(\sigma)),$$

and, trivially,

$$(172) \quad \partial_{3,2}(\partial_{2,0}(\sigma)) = \partial_{3,0}(\partial_{2,1}(\sigma)).$$

Then, as a consequence of (169), (170) and (172) we obtain

$$(173) \quad \partial_{3,0}(\partial_{2,1}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) = \partial_{3,2}(\omega_\sigma) * \partial_{3,4}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)).$$

On the other hand, by (156), we have

$$(174) \quad \partial_{3,0}(\partial_{2,3}(\sigma)) = \partial_{3,4}(\partial_{2,0}(\sigma)),$$

and, trivially,

$$(175) \quad \partial_{3,3}(\partial_{2,0}(\sigma)) = \partial_{3,0}(\partial_{2,2}(\sigma))$$

holds.

Also,

$$(176) \quad \partial_{3,3}(\partial_{2,1}(\sigma^{-1})) * \partial_{3,3}(\partial_{2,3}(\sigma^{-1})) = \partial_{3,1}(\partial_{2,2}(\sigma^{-1})) * \partial_{3,4}(\partial_{2,3}(\sigma^{-1}))$$

holds, because

$$\begin{aligned} & \partial_{3,3}(\partial_{2,1}(\sigma^{-1})) * \partial_{3,3}(\partial_{2,3}(\sigma^{-1})) \\ &= \partial_{3,3}(\partial_{2,1}(\sigma^{-1}) * \partial_{2,3}(\sigma^{-1})) \text{ (by the condition of group morphism for } \partial_{3,3}) \\ &= \mu_B \circ (\sigma^{-1} \otimes B) \circ (H \otimes c_{B,H}) \circ (G_{\sigma^{-1}} \otimes \mu_H) \text{ (by (157), the counit properties and the naturality of } c) \end{aligned}$$

$$= \partial_{3,1}(\partial_{2,2}(\sigma^{-1})) * \partial_{3,4}(\partial_{2,3}(\sigma^{-1})) \text{ (by (a1) of Definition 1.1, naturality of } c \text{ and (18))}.$$

Then, as a consequence of (176), we have the identity

$$(177) \quad \partial_{3,3}(\partial_{2,3}(\sigma)) * \partial_{3,3}(\partial_{2,1}(\sigma)) = \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,2}(\sigma))$$

and using that $\partial_{3,3}$ is a group morphism, (175) and (171) we can assure that

$$(178) \quad \partial_{3,0}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) = \partial_{3,3}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,2}(\sigma))$$

holds.

Moreover,

$$(179) \quad \partial_{3,0}(\partial_{2,0}(\sigma)) * \partial_{3,4}(\partial_{2,3}(\sigma)) = \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,0}(\sigma))$$

holds because

$$\begin{aligned} & \partial_{3,0}(\partial_{2,0}(\sigma)) * \partial_{3,4}(\partial_{2,3}(\sigma)) \\ &= \mu_B \circ ((\varphi_B \circ (H \otimes \varphi_B)) \otimes \sigma) \circ (H \otimes H \otimes c_{H,B} \otimes H) \circ (H \otimes H \otimes H \otimes c_{H,B}) \circ (\delta_{H^{\otimes 2}} \otimes ((B \otimes \varepsilon_H) \circ F_\sigma)) \\ & \quad \text{(by (18) and the naturality of } c) \\ &= \mu_B \circ (B \otimes \sigma) \otimes (P_{\varphi_B} \otimes H) \circ (H \otimes (P_{\varphi_B} \circ (H \otimes \sigma))) \text{ (by the naturality of } c \text{ and (61))} \\ &= \mu_B \circ (B \otimes \varphi_B) \circ (F_\sigma \otimes \sigma) \text{ (by (77))} \\ &= \mu_B \circ (B \otimes \varphi_B) \circ (F_\sigma \otimes ((\varepsilon_H \otimes B) \circ G_\sigma)) \text{ (by (66))} \\ &= \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,0}(\sigma)) \text{ (by (156), the naturality of } c \text{ and (18))} \end{aligned}$$

and by (156) and the associativity of μ_H we obtain the equalities

$$(180) \quad \partial_{3,1}(\partial_{2,3}(\sigma)) = \partial_{3,4}(\partial_{2,1}(\sigma)),$$

$$(181) \quad \partial_{3,1}(\partial_{2,1}(\sigma)) = \partial_{3,2}(\partial_{2,1}(\sigma)).$$

Finally, observe that, as ω_σ factors through the center of B , for all $i \in \{1, 2, 3, 4\}$ and $\tau \in \text{Reg}_{\varphi_B}(H^{\otimes 4}, B)$, we have

$$(182) \quad \tau * \partial_{3,i}(\omega_\sigma) = \partial_{3,i}(\omega_\sigma) * \tau.$$

Therefore, we conclude the proof by cancellation because in one hand

$$\begin{aligned} & \partial_{3,0}(\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) \\ &= \partial_{3,0}(\partial_{2,0}(\sigma)) * \partial_{3,0}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) \text{ (by the condition of group morphism for } \partial_{3,0}) \\ &= \partial_{3,0}(\omega_\sigma) * \partial_{3,0}(\partial_{2,3}(\sigma)) * \partial_{3,0}(\partial_{2,1}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) \text{ (by the condition of group morphism for } \partial_{3,0} \text{ and (168))} \\ &= \partial_{3,0}(\omega_\sigma) * \partial_{3,0}(\partial_{2,3}(\sigma)) * \partial_{3,2}(\omega_\sigma) * \partial_{3,4}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)) \text{ (by (173))} \\ &= \partial_{3,0}(\omega_\sigma) * \partial_{3,2}(\omega_\sigma) * \partial_{3,0}(\partial_{2,3}(\sigma)) * \partial_{3,4}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)) \text{ (by (182))} \\ &= \partial_{3,0}(\omega_\sigma) * \partial_{3,2}(\omega_\sigma) * \partial_{3,4}(\partial_{2,0}(\sigma)) * \partial_{3,4}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)) \text{ (by (174))} \\ &= \partial_{3,0}(\omega_\sigma) * \partial_{3,2}(\omega_\sigma) * \partial_{3,4}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,4}(\partial_{2,1}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)) \text{ (by the condition of group morphism for } \partial_{3,4} \text{ and (168))}, \end{aligned}$$

and on the other hand

$$\partial_{3,0}(\partial_{2,0}(\sigma) * \partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma))$$

$$\begin{aligned}
&= \partial_{3,0}(\partial_{2,0}(\sigma)) * \partial_{3,0}(\partial_{2,2}(\sigma)) * \partial_{3,2}(\partial_{2,2}(\sigma)) \text{ (by the condition of group morphism for } \partial_{3,0}) \\
&= \partial_{3,0}(\partial_{2,0}(\sigma)) * \partial_{3,3}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,2}(\sigma)) \text{ (by (178))} \\
&= \partial_{3,3}(\omega_\sigma) * \partial_{3,0}(\partial_{2,0}(\sigma)) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,2}(\sigma)) \text{ (by (182))} \\
&= \partial_{3,3}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,0}(\sigma)) * \partial_{3,1}(\partial_{2,2}(\sigma)) \text{ (by (179))} \\
&= \partial_{3,3}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\omega_\sigma) * \partial_{3,1}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,1}(\sigma)) \text{ (by the condition of group morphism} \\
&\quad \text{for } \partial_{3,1} \text{ and (168))} \\
&= \partial_{3,1}(\omega_\sigma) * \partial_{3,3}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,3}(\sigma)) * \partial_{3,1}(\partial_{2,1}(\sigma)) \text{ (by (182))}. \\
&= \partial_{3,1}(\omega_\sigma) * \partial_{3,3}(\omega_\sigma) * \partial_{3,4}(\partial_{2,3}(\sigma)) * \partial_{3,4}(\partial_{2,1}(\sigma)) * \partial_{3,2}(\partial_{2,1}(\sigma)) \text{ (by (180) and (181))}.
\end{aligned}$$

□

Theorem 5.12. *Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Let θ_σ the obstruction of σ . Then, $\theta_\sigma \in \text{Im}(D_{\varphi_{Z(B)}}^2)$ if, and only if, there exists $\alpha \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ that satisfies the twisted condition (77) and the cocycle condition (78).*

Proof. If $\theta_\sigma \in \text{Im}(D_{\varphi_{Z(B)}}^2)$, there exists $\tau \in \text{Reg}_{\varphi_{Z(B)}}(H^{\otimes 2}, Z(B))$ such that $D_{\varphi_{Z(B)}}^2(\tau) = \theta_\sigma$. Then,

$$(183) \quad z_B \circ D_{\varphi_{Z(B)}}^2(\tau) = \omega_\sigma.$$

By Proposition 4.16, the morphism $\alpha = (z_B \circ \tau^{-1}) * \sigma$ satisfies the twisted condition (77) and belongs to $\text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$. On the other hand,

$$\begin{aligned}
&\omega_\alpha \\
&= D_{\varphi_B}^2(z_B \circ \tau^{-1}) * \omega_\sigma \text{ (by the properties of } \partial_{i,j} \text{ and } \tau) \\
&= z_B \circ D_{\varphi_{Z(B)}}^2(\tau^{-1}) * \omega_\sigma \text{ (by the properties of } D_{\varphi_B}^2) \\
&= \omega_\sigma^{-1} * \omega_\sigma \text{ (by (183))} \\
&= u_3^{\varphi_B} \text{ (by (141) for } \omega_\sigma)
\end{aligned}$$

and, as a consequence, α satisfies the cocycle condition (78).

Conversely, assume that there exists $\alpha \in \text{Reg}_{\varphi_B}(H^{\otimes 2}, B)$ that satisfies the twisted condition (77) and the cocycle condition (78). Then, by Proposition 4.16, there there exists $\tau \in \text{Reg}_{\varphi_{Z(B)}}(H^{\otimes 2}, Z(B))$ such that (152) holds, i.e., $\alpha = (z_B \circ \tau) * \sigma$. As a consequence, $\sigma = (z_B \circ \tau)^{-1} * \alpha$ and $\theta_\sigma \in \text{Im}(D_{\varphi_{Z(B)}}^2)$ since

$$\omega_\sigma = D_{\varphi_B}^2(z_B \circ \tau^{-1}) * \partial_{2,0}(\alpha) * \partial_{2,2}(\alpha) * \partial_{2,1}(\alpha^{-1}) * \partial_{2,3}(\alpha^{-1}) \stackrel{(160)}{=} D_{\varphi_B}^2(z_B \circ \tau^{-1}) = z_B \circ D_{\varphi_{Z(B)}}^2(\tau^{-1}).$$

□

Proposition 5.13. *Let H be a cocommutative weak Hopf algebra and let (B, φ_B) be a left weak H -module algebra. Let $\sigma \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Then, $\alpha \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfies the twisted condition (77) if, and only if, there exists $\tau \in \text{Reg}_{\varphi_{Z(B)}}^+(H^{\otimes 2}, Z(B))$ satisfying (152).*

Proof. First note that, if H is cocommutative, (D, φ_D) is a left weak H -module algebra and $\beta \in \text{Reg}_{\varphi_D}(H^{\otimes 2}, D)$, using that $\overline{\Pi}_H^L = \Pi_H^L$ and (82) we obtain that

$$(184) \quad \beta \circ (\eta_H \otimes H) = \beta \circ (\Pi_H^L \otimes H) \circ \delta_H$$

holds. Also, (83), holds for β and therefore β satisfies the normal condition (84), i.e., $\beta \in \text{Reg}_{\varphi_D}^+(H^{\otimes 2}, D)$ if and only if

$$(185) \quad \beta \circ (\Pi_H^L \otimes H) \circ \delta_H = \beta \circ (H \otimes \Pi_H^R) \circ \delta_H = u_1^{\varphi_D}.$$

Let $\alpha \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfying the twisted condition (77). By Proposition 4.16 there exists $\tau \in \text{Reg}_{\varphi_{Z(B)}}(H^{\otimes 2}, Z(B))$ satisfying (152). Then, $z_B \circ \tau = \alpha * \sigma^{-1}$ and τ satisfies the normal condition (84) because, in one hand,

$$\begin{aligned} & z_B \circ \tau \circ (\Pi_H^L \otimes H) \circ \delta_H \\ &= \mu_B \circ ((\alpha \circ (\Pi_H^L \otimes H)) \circ (\sigma^{-1} \circ (\Pi_H^L \otimes H))) \circ \delta_{H^{\otimes 2}} \circ \delta_H \text{ (by the naturality of } c \text{ and (i) of} \\ & \quad \text{[2, Proposition 2.6])} \\ &= (\alpha \circ (\Pi_H^L \otimes H) \circ \delta_H) * (\sigma^{-1} \circ (\Pi_H^L \otimes H) \circ \delta_H) \text{ (by the cocommutativity of } \delta_H) \\ &= u_2^{\varphi_B} * u_2^{\varphi_B} \text{ (by (185))} \\ &= z_B \circ u_2^{\varphi_{Z(B)}} \text{ (by (151))} \end{aligned}$$

and, on the other hand, using the same arguments we have $z_B \circ \tau \circ (H \otimes \Pi_H^R) \circ \delta_H = z_B \circ u_2^{\varphi_{Z(B)}}$.

Conversely, if there exists $\tau \in \text{Reg}_{\varphi_{Z(B)}}^+(H^{\otimes 2}, Z(B))$ satisfying (152), by the previous arguments, we obtain that

$$\alpha \circ (\Pi_H^L \otimes H) \circ \delta_H = (z_B \circ \tau \circ (\Pi_H^L \otimes H) \circ \delta_H) * (\sigma^{-1} \circ (\Pi_H^L \otimes H) \circ \delta_H) = (z_B \circ u_2^{\varphi_{Z(B)}}) * u_2^{\varphi_B} = u_2^{\varphi_B}$$

and similarly $\alpha \circ (H \otimes \Pi_H^R) \circ \delta_H = u_2^{\varphi_B}$. Therefore, $\alpha \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$. \square

Remark 5.14. Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma, \beta \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Let $\theta_\sigma, \theta_\beta$ the corresponding obstructions of σ and β . Then, by the previous proposition, it is easy to show that $[\theta_\sigma] = [\theta_\beta]$ in $\mathcal{H}_{\varphi_{Z(B)}}^{3+}(H, Z(B))$, i.e., θ_σ and θ_β are cohomologous.

Corollary 5.15. *Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Let θ_σ the obstruction of σ . Then, $\theta_\sigma \in \text{Im}(D_{\varphi_{Z(B)}}^{2+})$ if, and only if, there exists $\alpha \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ that satisfies the twisted condition (77) and the cocycle condition (78).*

Proof. The result is a direct consequence of Theorem 5.12 and Proposition 5.13. \square

Corollary 5.16. *Let H be a cocommutative weak Hopf algebra, let (B, φ_B) be a left weak H -module algebra and let $\sigma \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ satisfying the twisted condition (77). Let θ_σ the obstruction of σ . Then, $[\theta_\sigma] = 0$ in $\mathcal{H}_{\varphi_{Z(B)}}^{3+}(H, Z(B))$ if and only if there exists a morphism $\alpha \in \text{Reg}_{\varphi_B}^+(H^{\otimes 2}, B)$ that satisfies the twisted condition (77), the cocycle condition (78) and the normal condition (84).*

Proof. The proof follows by the previous corollary and Corollary 1.20 \square

As a consequence of this corollary we can assure that the obstruction vanishes if and only if there exists a weak crossed product with preunit $\nabla_{B \otimes H}^{\varphi_B} \circ (\eta_B \otimes \eta_H)$ and normalized with respect to $\nabla_{B \otimes H}^{\varphi_B}$. Equivalently, by [12, Theorem 6.17, Corollary 6.18], this is equivalent to say that B admits a H -cleft extension (see also [3, Proposition 3.5]).

FUNDING

The authors were supported by Ministerio de Ciencia e Innovación of Spain. Agencia Estatal de Investigación. Unión Europea - Fondo Europeo de Desarrollo Regional. Grant: Homología, homotopía e invariantes categóricos en grupos y álgebras no asociativas.

REFERENCES

- [1] J. N. Alonso Álvarez, J. M. Fernández Vilaboa, Cleft extensions in braided categories, *Commun. Algebra* 28 (2000), 3185-3196.
- [2] J. N. Alonso Álvarez, J. M. Fernández Vilaboa, R. González Rodríguez, Cohomology of algebras over weak Hopf algebras, *Homol. Homotopy Appl.* 16 (2014), 341-369.
- [3] J. N. Alonso Álvarez, J. M. Fernández Vilaboa, R. González Rodríguez, Crossed products over weak Hopf algebras related to cleft extensions and cohomology, *Chinese Ann. Math. B.* 35 (2014), 161-190.
- [4] R. Blattner, M. Cohen, S. Montgomery, Crossed products and inner actions of Hopf algebras, *T. Am. Math. Soc.* 298 (1986), 671-711.
- [5] R. Blattner, S. Montgomery, Crossed products and Galois extensions of Hopf algebras, *Pac. J. Math.* 137 (1989) 37-54.
- [6] G. Böhm, F. Nill, K. Szlachányi, Weak Hopf algebras, I. Integral theory and C^* -structure, *J. Algebra* 221 (1999), 385-438.
- [7] Y. Doi, M. Takeuchi, Cleft comodule algebras for a bialgebra, *Comm. Algebra* 14 (1986) 801-817.
- [8] Y. Doi, Equivalent crossed products for a Hopf algebra, *Comm. Algebra* 17 (1989), 3053-3085.
- [9] J. M. Fernández Vilaboa, R. González Rodríguez, A. B. Rodríguez Raposo, Preunits and weak crossed products, *J. Pure Appl. Algebra* 213 (2009), 2244-2261.
- [10] J. M. Fernández Vilaboa, R. González Rodríguez, A. B. Rodríguez Raposo, Equivalences for weak crossed products, *Commun. Algebra* 44 (2016), 4519-4545.
- [11] J. M. Fernández Vilaboa, E. Villanueva Nóvoa, A characterization of the cleft comodule triples, *Commun. Algebra* 16 (1998), 613-622.
- [12] Jorge A. Guccione, Juan J. Guccione, C. Valqui, Cleft extensions of weak Hopf algebras, *J. Algebra* 547 (2020), 668-710.
- [13] S. MacLane, *Homology. Classics in Mathematics.* Springer-Verlag, Berlin. 1975.
- [14] S. Montgomery, Binvertible actions of Hopf algebras, *Israel J. Math.* 83 (1993), 45-71.
- [15] A. B. Rodríguez Raposo, Crossed products for weak Hopf algebras, *Comm. Algebra* 37 (2009), 2274-2289.
- [16] P. Schauenburg, Cohomological obstructions to cleft extensions over cocommutative Hopf algebras, *K-Theory* 24 (2001), 227-242.
- [17] M. Sweedler, Cohomology of algebras over Hopf algebras, *T. Am. Math. Soc.* 133 (1968), 205-239.

(Ramón González Rodríguez) UNIVERSIDADE DE VIGO. DEPARTAMENTO DE MATEMÁTICA APLICADA II, E.E. TELECOMUNICACIÓN, CAMPUS UNIVERSITARIO LAGOAS-MARCOSENDE, E-36280 VIGO, SPAIN.

Email address, R. González: rgon@dma.uvigo.es

(Ana Belén Rodríguez Raposo) UNIVERSIDADE DE SANTIAGO DE COMPOSTELA. DEPARTAMENTO DE DIDÁCTICAS APLICADAS, FACULTADE DE CIENCIAS DA EDUCACIÓN, CAMPUS NORTE, E-15771 SANTIAGO DE COMPOSTELA, SPAIN.

Email address, M.B. Rodríguez: anabelen.rodriguez.raposo@usc.es