

On the tangential cone condition for electrical impedance tomography

Stefan Kindermann*

December 8, 2021

Abstract

We state some sufficient criteria for the tangential cone conditions to hold for the electrical impedance tomography problem. The results are based on an estimate for the first-order Taylor residual for the forward operator and some convexity results in the Löwner order. As a consequence, for conductivities satisfying certain monotonicity properties, the tangential cone condition is verified.

1 Introduction

The electrical impedance tomography problem is a classical inverse problem, where the aim is to extract information about the conductivity from boundary measurements of current/voltage pairs. Starting with the definition of the problem in the seminal paper of Calderon [3], it has been investigated in various directions and now serves as a paradigmatic instance of a parameter identification problem from boundary measurements.

The common mathematical formulation is to consider solutions of the boundary value problem on a Lipschitz domain Ω ,

$$\operatorname{div}(\gamma \nabla u) = 0 \quad \text{in } \Omega, \quad u = f \quad \text{on } \Omega. \quad (1)$$

The data for the inverse problem are multiple or infinitely many pairs of Cauchy data $(f, \gamma \frac{\partial}{\partial n} u|_{\partial \Omega})$ on the boundary, and the interest is to recover the conductivity $\gamma(x)$ in the interior Ω . As it is typical for such identification problems with only boundary data, this leads under usual circumstances to a nonlinear severely ill-posed problem, and without strong restrictions on the conductivity, one can at best only expect conditional logarithmic stability [10]. Various classical uniqueness and stability results are collected, e.g., in [2] or [10].

In the following, we assume that the unknown conductivity can be written as a perturbation $\delta\gamma$ of a known background, which we take without loss of generality as 1. Thus, we assume throughout that

$$\gamma(x) = 1 + \delta\gamma(x), \quad \underline{\alpha} \leq \gamma(x) \leq \bar{\alpha}, \quad \text{a.e. in } \Omega, \quad (2)$$

*Industrial Mathematics Institute, Johannes Kepler University Linz, Austria. kindermann@indmath.uni-linz.ac.at.

with positive constants $\underline{\alpha}, \bar{\alpha}$ to ensure ellipticity and stability of the partial differential equation.

In order to solve the problem, it is convenient to frame it into operator-theoretic language. The above-mentioned Cauchy-data (in the case of complete data), are equivalent to the knowledge of the Dirichlet-to-Neumann operator. Thus, we introduce the parameter-to-data map:

$$F(\gamma) := \Lambda_\gamma - \Lambda_1, \quad (3)$$

where Λ_γ is the Dirichlet-to-Neumann map for (1). Solving the inverse problem is then equivalent to inverting F . We give a precise definition of the associate spaces X, Y in $F : X \rightarrow Y$ in the next section.

The main theme of this article concerns not the solution of this problem but the investigation of the nonlinearity of F . This is highly relevant when applying regularization methods, in particular, iterative ones. Indeed, the convergence theory of iterative regularization methods such as Landweber's method requires some restrictions on the nonlinearity that quantify the deviation of the problem from a linear one. In this work, we focus on the well-known tangential cone conditions and of its variants (cf. [14, 6, 12]).

For a general inverse problem with a differentiable parameter-to-data map F between Hilbert spaces, the so-called strong tangential cone condition [14, 6] is satisfied if, with F' denoting the Fréchet-derivative of F , there is an η , $1 > \eta > 0$, such that

$$\|F(\tilde{x}) - F(x) - F'(x)(\tilde{x} - x)\| \leq \eta \|F(\tilde{x}) - F(x)\| \quad (\text{stc})$$

holds for all \tilde{x}, x in a neighborhood of some x_0 . A weaker version, the weak-tangential cone conditions [14] holds if an η , $1 > \eta > 0$, exists such that

$$(F(\tilde{x}) - F(x) - F'(x)(\tilde{x} - x), F(\tilde{x}) - F(x))_Y \leq \eta \|F(\tilde{x}) - F(x)\|^2. \quad (\text{wtc})$$

Moreover, the weak tangential cone condition with $\eta = 1$ reads as

$$(F'(x)(\tilde{x} - x), F(\tilde{x}) - F(x))_Y \geq 0, \quad (\text{qcon})$$

which yields a weaker condition than (wtc) that has been proposed in [12]. Note that by the parallelogram identity, the inequalities (wtc) and (qcon) may be equivalently rewritten as

$$\begin{aligned} & \|F(\tilde{x}) - F(x) - F'(x)(\tilde{x} - x)\|^2 \\ & \leq (2\eta - 1) \|F(\tilde{x}) - F(x)\|^2 + \|F'(x)(\tilde{x} - x)\|^2. \end{aligned} \quad (4)$$

These inequalities are central to the convergence theory of the nonlinear Landweber method and many other iterative regularization methods. They are a replacement to coercivity estimates, which cannot exist in the ill-posed case. It is a classical result that (under some standard additional assumptions) the strong tangential cone condition with $\eta \leq \frac{1}{2}$ implies strong convergence of the Landweber method [6, 11]. Similarly, the weak-tangential cone condition [14] imply nonexpansivity of the iteration, and in particular, weak (subsequential) convergence of the iterates. (Of course, all this in connection with parameter choice rules.) The condition (qcon) imply that the iterates stay in a neighborhood of the solution, which also yields weak (subsequential) convergence [12].

It is surprising that in view of its importance, the tangential cone conditions for the impedance tomography problem could only be verified in a few special cases. For instance, Lechleitner and Rieder [13] have proven (stc) in a semidiscrete case, essentially by using a stability result for the discrete problem. Interestingly, de Hoop, Liu and Scherzer [4] have proven (stc) for a class of piecewise constant conductivities being constant on finitely many regions. The proof is based on a Lipschitz stability result of Alessandrini and Vessella [1]. In both cases, the stability constants might get quite large, thus, the cone conditions can in practice only be theoretically verified in a very narrow neighborhood.

Except from these few cases, the validity of the above tangential cone conditions is completely open, which is extremely puzzling given the fact that the Landweber method has successfully been applied to the impedance tomography problem in many situations, and only inveterate sceptics would doubt its convergence.

Our article aims to gain further understanding of this puzzle (though without completely resolving it) by analyzing and establishing sufficient conditions for the tangential cone conditions. The main contribution is that a condition of the form

$$\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)^2\| \leq C\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|$$

suffices; see (27). This is established by a useful bound for the remainder in a first-order Taylor expansion in (24).

One result that is probably most relevant in practice is that the tangential cone conditions are satisfied for conductivities that satisfy certain monotonicity properties (e.g., a purely positive perturbation of the background conductivity). A central tool in this paper, which might be of more general interest, is a convexity result of the parameter-to-solution map in the Löwner ordering given in Theorems 1 and 2.

2 Problem setup and operator estimates

We formulate some standard assumption and specify the notation. We make use of the space of L^2 -vector fields:

$$L^2(\Omega)^n := \{v : \Omega \rightarrow \mathbb{R}^n \mid \int_{\Omega} |v(x)|^2 < \infty\}.$$

The Dirichlet data f are canonically chosen in $H^{\frac{1}{2}}(\partial\Omega)$, and thus, the associated boundary value problem

$$\Delta u_{1,f} = 0 \text{ in } \Omega, \quad u_{1,f} = f \text{ on } \partial\Omega$$

has a unique solution in $H^1(\Omega)$. Moreover, by the Poincaré inequality, the mapping $f \rightarrow \nabla u_{1,f}$ is an isomorphism from $H^{\frac{1}{2}}(\partial\Omega) \rightarrow L^2(\Omega)^n$, such that we may define the $H^{\frac{1}{2}}(\partial\Omega)$ -norm in this paper as

$$\|f\|_{H^{\frac{1}{2}}(\partial\Omega)} := \|\nabla u_{1,f}\|_{L^2(\Omega)^n}.$$

Moreover, we define $u_{\gamma,f}$ as the solution in $H^1(\Omega)$ of the problem (1)

$$\operatorname{div}(\gamma \nabla u_{\gamma,f}) = 0 \text{ in } \Omega, \quad u_{\gamma,f} = f \text{ on } \partial\Omega.$$

For a conductivity γ satisfying (2), the Dirichlet-to-Neumann operator Λ_γ maps a $f \in H^{\frac{1}{2}}(\partial\Omega)$ to $\gamma \frac{\partial}{\partial n} u_{\gamma,f} \in H^{-\frac{1}{2}}(\partial\Omega)$ continuously. It is well-known that the Dirichlet-to-Neumann operators can be rewritten in terms of energy integrals: Defining F by (3), we have for γ_1, γ_2 satisfying (2) that (cf. [10])

$$F(\gamma_1) - F(\gamma_2) = \Lambda_{\gamma_1} - \Lambda_{\gamma_2} \quad \text{with} \quad (5)$$

$$\langle \Lambda_{\gamma_1} - \Lambda_{\gamma_2} f, g \rangle = \int_{\Omega} (\gamma_1 - \gamma_2) \nabla u_{\gamma_1, f} \cdot \nabla u_{\gamma_2, g} dx. \quad (6)$$

Here and in the following we denote by $\langle \cdot, \cdot \rangle$ the duality pairing in $H^{-\frac{1}{2}} \times H^{\frac{1}{2}}$:

$$\langle g, f \rangle = \langle g, f \rangle_{H^{-\frac{1}{2}}(\partial\Omega), H^{\frac{1}{2}}(\partial\Omega)}.$$

Moreover, the mapping $F(\gamma)$ is Fréchet-differentiable with respect to the L^∞ -norm, and the derivative can be expressed as

$$F'[\gamma_1]w = \Lambda'_{\gamma_1}(w) \quad (7)$$

$$\langle \Lambda'_{\gamma_1}(w)f, g \rangle = \int_{\Omega} w \nabla u_{\gamma_1, f} \cdot \nabla u_{\gamma_1, g} dx, \quad w \in L^\infty(\Omega). \quad (8)$$

Our analysis is based on the following assumptions, which we assume to hold for the rest of the article.

Assumption 1.

We assume given a (finite or countably infinite) sequence of linear independent and orthogonal Dirichlet data $(f_i)_{i \in I}$, $f_i \in H^{\frac{1}{2}}(\partial\Omega)$.

We set as domain of definition of the parameter-to-data map

$$D(F) := \{\gamma \in L^\infty(\Omega) \mid \underline{\alpha} \leq \gamma \leq \bar{\alpha}, \text{ and (9) holds}\},$$

with

$$\sum_{i, j \in I} |\langle [\Lambda_\gamma - \Lambda_1]f_i, f_j \rangle|^2 < \infty. \quad (9)$$

The first assumption is not much of a restriction, as the Dirichlet data are part of the experimental design and can be chosen orthogonal. Also the second one is not severe, as (9) holds if we only have finitely many Dirichlet data, or in case of infinitely many f_i if the deviations from the background conductivity $\delta\gamma$ have a common compact support inside of Ω . This is also a usual assumption in the impedance tomography problem. We note that we do not a priori require that the f_i form a complete basis in $H^{\frac{1}{2}}(\partial\Omega)$. Thus, much of our analysis is also valid in case of finitely many measurements or measurements on only a part of the boundary.

Associated to the set of Dirichlet data, we set

$$V_D := \text{span}(f_i)_{i \in I} \subset H^{\frac{1}{2}}(\partial\Omega).$$

The quadratic form $\langle (\Lambda_\gamma - \Lambda_1)f, g \rangle$ associated to the Dirichlet-to-Neumann mapping defines a linear operator $\Lambda_\gamma - \Lambda_1 : V_D \rightarrow V'_D$. Hence, the parameter to solution map can be defined as a mapping in the following spaces

$$F : D(F) \subset X := L^\infty(\Omega) \rightarrow Y := L(V_D, V'_D) \subset L(H^{\frac{1}{2}}(\partial\Omega), H^{-\frac{1}{2}}(\partial\Omega)) \\ \gamma \rightarrow \Lambda_\gamma - \Lambda_1$$

with the norm

$$\|F(\gamma)\|_Y^2 := \sum_{i,j \in I} |\langle [\Lambda_\gamma - \Lambda_1]f_i, f_j \rangle|^2.$$

Introducing the Riesz isomorphism $\mathcal{I} : H^{-\frac{1}{2}} \rightarrow H^{\frac{1}{2}}$, we may write the norm

$$\|F(\gamma)\|_Y^2 = \sum_{i,j \in I} \left| (\mathcal{I}[\Lambda_\gamma - \Lambda_1]f_i, f_j)_{H^{\frac{1}{2}}, H^{\frac{1}{2}}} \right|^2 = \|\mathcal{I}[\Lambda_\gamma - \Lambda_1]\|_{HS(V_D)}^2$$

as Hilbert-Schmidt norm of $\mathcal{I}(\Lambda_\gamma - \Lambda_1)$ for the operator mapping between the space V_D . Thus, the image space is equipped with a Hilbert space structure. Note that for selfadjoint compact operators, the Hilbert-Schmidt norm is the sum of squares of the eigenvalues. We will denote by $\|\cdot\|_{HS}$ the Hilbert-Schmidt norm, omitting the underlying space where the operators map, and by $\|\cdot\|_{2,L(X,Y)}$ the operator norm for a linear operator from $X \rightarrow Y$. Moreover, for functions in L^∞ , we set $\|\cdot\|_\infty = \|\cdot\|_{L^\infty}$.

For later use we also recall the Löwner ordering for selfadjoint operators on a Hilbert space H : We have

$$A \leq_L B \Leftrightarrow (Ax, x)_H \leq (Bx, x)_H \quad \forall x \in H.$$

A useful property of this ordering is that for Hilbert-Schmidt operators,

$$0 \leq A \leq_L B \Rightarrow \|A\|_{HS} \leq \|B\|_{HS}. \quad (10)$$

This follows from Weyl's inequality for the eigenvalues and since the Löwner ordering implies a corresponding eigenvalue inequality. Moreover, the following result with an arbitrary bounded operator T and T^* its adjoint will be used frequently:

$$A \leq_L B \Rightarrow T^*AT \leq_L T^*BT. \quad (11)$$

We define the operator $L(\gamma)^{-1}$ as the solution operator for (1) with homogeneous Dirichlet condition, i.e., for γ satisfying (2), we set

$$\begin{aligned} L(\gamma)^{-1} : H^{-1}(\Omega) &\rightarrow H_0^1(\Omega) \\ h &\rightarrow v, \end{aligned}$$

where v is the solution of

$$\operatorname{div}(\gamma \nabla v) = h \quad \text{in } \Omega \quad v = 0 \quad \text{on } \partial\Omega.$$

Furthermore, for a function $\kappa \in L^\infty(\Omega)$, we define the multiplication operator

$$H_\kappa : L^2(\Omega)^n \rightarrow L^2(\Omega)^n \quad \vec{f}(x) \rightarrow \kappa(x)\vec{f}(x) \quad (12)$$

and the bounded operator

$$T_\gamma : L^2(\Omega)^n \rightarrow L^2(\Omega)^n \quad T_\gamma := \nabla L(\gamma)^{-1} \operatorname{div}.$$

With this notation we state the following useful lemma:

Lemma 1. For any γ_1, γ_2 with $\|\frac{\gamma_1 - \gamma_2}{\gamma_2}\|_\infty < 1$ satisfying (2) and any $f \in H^{\frac{1}{2}}(\partial\Omega)$, we have

$$\nabla u_{\gamma_1, f} = (I - T_{\gamma_2} H_{\gamma_2 - \gamma_1})^{-1} \nabla u_{\gamma_2, f},$$

Proof. By definition of the inhomogeneous Dirichlet problem, we have that $u_{\gamma_2, f} = u_{\gamma_1, f} + w$, where w satisfies the homogenous problem

$$\operatorname{div}(\gamma_2 \nabla w) = -\operatorname{div}(\gamma_2 u_{\gamma_1, f}) = -\operatorname{div}((\gamma_2 - \gamma_1) u_{\gamma_1, f}),$$

and where we used that $u_{\gamma_2, f}$ solves the problem (1) with $\gamma = \gamma_2$. Thus,

$$u_{\gamma_2, f} = [I - L(\gamma_2)]^{-1} \operatorname{div}((\gamma_2 - \gamma_1) \nabla) u_{\gamma_1, f}.$$

Applying the gradient yields

$$\nabla u_{\gamma_2, f} = [I - \nabla L(\gamma_2)]^{-1} \operatorname{div}[(\gamma_2 - \gamma_1) \nabla] u_{\gamma_1, f}.$$

Under the stated assumptions, the operator on the left has a bounded inverse, as can also be seen from the next proposition. \square

Note that the multiplication operator satisfies $H_{\kappa_1} H_{\kappa_2} = H_{\kappa_1 \kappa_2}$, these operators commute, and $\|H_\kappa\|_2 \leq \|\kappa\|_\infty$.

The operators T_γ play an important role as they can be expressed as projection operator.

Proposition 1. The operator T_γ can be written as

$$T_\gamma = H_{\gamma^{\frac{1}{2}}}^{-1} R_\gamma H_{\gamma^{\frac{1}{2}}},$$

where R_γ is an orthogonal projection operator on $L^2(\Omega)^n$. Moreover, the range of R_γ is orthogonal to the space

$$\{z \in L^2(\Omega)^n \mid \operatorname{div}(\gamma^{\frac{1}{2}} z) = 0\}. \quad (13)$$

Proof. Consider $R_\gamma = H_{\gamma^{\frac{1}{2}}} T_\gamma H_{\gamma^{\frac{1}{2}}}$. We show that $R_\gamma^2 = R_\gamma$. Let $y \in L^2(\Omega)^n$. Then by definition $R_\gamma y = \gamma^{\frac{1}{2}} \nabla v_y$, where $v_y \in H_0^1$ satisfies the variational formulation

$$\int_\Omega \gamma \nabla v_y \cdot \nabla w \, dx = \int_\Omega \gamma^{\frac{1}{2}} y \cdot \nabla w \, dx$$

for arbitrary $w \in H_0^1$. Taking as $y = R_\gamma z = \gamma^{\frac{1}{2}} \nabla v_z$, where v_z satisfies the same equation as v_y with y replaced by z , we arrive at

$$\int_\Omega \gamma \nabla v_y \cdot \nabla w \, dx = \int_\Omega \gamma^{\frac{1}{2}} y \cdot \nabla w \, dx = \int_\Omega \gamma \nabla v_z \cdot \nabla w \, dx = \int_\Omega \gamma^{\frac{1}{2}} z \cdot \nabla w \, dx,$$

thus $R_\gamma^2 y = R_\gamma y$.

Next, we show that R_γ is selfadjoint. Indeed, let $y, z \in L^2(\Omega)^n$ be arbitrary and define v_y, v_z as before. Then

$$(R_\gamma y, z)_{L^2(\Omega)^n} = \int_\Omega \gamma^{\frac{1}{2}} \nabla v_y \cdot z \, dx = \int_\Omega \gamma \nabla v_y \cdot \nabla v_z \, dx,$$

which is symmetric in y, z , thus $R_\gamma y = R_\gamma^* y$, hence it is selfadjoint. As a consequence, R_γ is an orthogonal projector. Finally we show that R_γ annihilates the space in (13). Let $z \in L^2(\Omega)^n$ be in the space in (13). Then

$$R_\gamma z = \gamma^{\frac{1}{2}} \nabla L(\gamma^\dagger)^{-1} \operatorname{div}(\gamma^{\frac{1}{2}} z) = 0$$

since $\operatorname{div}(\gamma^{\frac{1}{2}} z) = 0$ by definition. \square

A useful observation is the following monotonicity property:

Lemma 2. *If $w(x) \geq 0$ a.e., $w \in L^\infty(\Omega)$, then $F'[\gamma]w \geq_L 0$. In particular, for $w_1, w_2 \in L^\infty(\Omega)$ with $0 \leq w_1 \leq w_2$ a.e. in Ω , and any $\gamma \in D(F)$, we have the estimate*

$$\|F'[\gamma](w_1)\|_Y \leq \|F'[\gamma](w_2)\|_Y. \quad (14)$$

Proof. For $f \in H^{\frac{1}{2}}(\partial\Omega)$ we have by (7) that

$$\langle (F'[\gamma]w)f, f \rangle = \langle \Lambda'_\gamma(w), f, f \rangle = \int_\Omega w(x) |\nabla u_{\gamma, f}|^2 dx \geq 0.$$

The second result follows now from (10) since the assumption implies

$$0 \leq_L \Lambda'_\gamma(w_1) \leq_L \Lambda'_\gamma(w_2).$$

\square

We can now state the important structural convexity property of the forward map.

Theorem 1. *For any $\gamma, \gamma^\dagger \in D(F)$ that satisfy $\|\frac{\gamma - \gamma^\dagger}{\gamma^\dagger}\|_\infty < 1$, we have*

$$0 \leq_L \Lambda(\gamma) - \Lambda(\gamma^\dagger) - \Lambda'_\gamma(\gamma - \gamma^\dagger) \leq_L \Lambda'_\gamma\left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger}\right). \quad (15)$$

Proof. Define

$$B(\gamma, \gamma^\dagger) := \Lambda(\gamma) - \Lambda(\gamma^\dagger) - \Lambda'_\gamma(\gamma - \gamma^\dagger). \quad (16)$$

Then, from (5) and (7), it follows that

$$\langle B(\gamma, \gamma^\dagger)f, f \rangle = \int_\Omega (\gamma - \gamma^\dagger) \nabla u_{\gamma, f} \cdot (\nabla u_{\gamma^\dagger, f} - \nabla u_{\gamma, f}) dx.$$

Using Lemma 1 with $\gamma_2 = \gamma^\dagger$ and $\gamma_1 = \gamma$ and (12), we write this as

$$\begin{aligned} \langle B(\gamma, \gamma^\dagger)f, f \rangle &= (H_{\gamma - \gamma^\dagger} \nabla u_{\gamma, f}, [(I - T_{\gamma^\dagger} H_{\gamma^\dagger - \gamma}) - I] \nabla u_{\gamma, f})_{L^2(\Omega)^n} \\ &= (H_{\gamma - \gamma^\dagger} \nabla u_{\gamma, f}, T_{\gamma^\dagger} H_{\gamma - \gamma^\dagger} \nabla u_{\gamma, f})_{L^2(\Omega)^n} \end{aligned} \quad (17)$$

Since T_{γ^\dagger} is positive definite by Proposition 1, the last term is positive, which proves that $B(\gamma, \gamma^\dagger) \geq_L 0$.

For the upper bound, we define the shortcut

$$\Delta\gamma := \frac{\gamma - \gamma^\dagger}{(\gamma^\dagger)^{\frac{1}{2}}}.$$

Using Proposition 1, the fact that an orthogonal projector satisfies $P \leq_L I$, and (11), we find that

$$\begin{aligned} \langle B(\gamma, \gamma^\dagger) f, f \rangle &= (H_{\Delta\gamma} \nabla u_{\gamma, f}, R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f})_{L^2(\Omega)^n} \\ &\leq (H_{\Delta\gamma} \nabla u_{\gamma, f}, H_{\Delta\gamma} \nabla u_{\gamma, f})_{L^2(\Omega)^n} \\ &= \int_{\Omega} \frac{(\gamma - \gamma^\dagger)^2}{(\gamma^\dagger)} |\nabla u_{\gamma, f}|^2 dx = \langle \Lambda'_\gamma ((\Delta\gamma)^2) f, f \rangle. \end{aligned} \quad (18)$$

This verifies the upper bound. \square

The upper bound can be strengthened by a more detailed analysis. The following result is an improvement.

Theorem 2. *With the same assumption on γ, γ^\dagger as in Theorem 1, let $B(\gamma, \gamma^\dagger)$ be as in (16), i.e.,*

$$B(\gamma, \gamma^\dagger) = \Lambda_\gamma - \Lambda_{\gamma^\dagger} - \Lambda'_\gamma (\gamma - \gamma^\dagger)$$

Then

$$\langle B(\gamma, \gamma^\dagger) f, f \rangle \leq \inf_{w \in L^2(\Omega)^n : \operatorname{div}((\gamma^\dagger)^{\frac{1}{2}} w) = 0} \|H_{\Delta\gamma} \nabla u_{\gamma, f} - w\|^2.$$

Moreover, the following estimates holds:

$$B(\gamma, \gamma^\dagger) \leq_L \Lambda'_\gamma \left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger} \right) - [\Lambda_\gamma - \Lambda_{\gamma^\dagger}] \Lambda_{\gamma^\dagger}^{-1} [\Lambda_\gamma - \Lambda_{\gamma^\dagger}], \quad (19)$$

$$B(\gamma, \gamma^\dagger) \leq_L \Lambda'_\gamma \left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger} \right) - \Lambda'_\gamma \left((\gamma - \gamma^\dagger) \frac{\gamma}{\gamma^\dagger} \right) \left[\Lambda'_\gamma \left(\frac{\gamma^2}{\gamma^\dagger} \right) \right]^{-1} \Lambda'_\gamma \left((\gamma - \gamma^\dagger) \frac{\gamma}{\gamma^\dagger} \right). \quad (20)$$

Proof. With (18) we find that

$$\langle B(\gamma, \gamma^\dagger) f, f \rangle = \|R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f}\|^2 = \inf_{w \in R_{\gamma^\dagger}^\perp} \|H_{\Delta\gamma} \nabla u_{\gamma, f} - w\|^2,$$

where $R_{\gamma^\dagger}^\perp$ is the orthogonal complement of the range of R_{γ^\dagger} . By Proposition 1, the set of w with $\operatorname{div}((\gamma^\dagger)^{\frac{1}{2}} w)$ is subset of R_{γ^\dagger} , which yields the first inequality.

Now we take $w = w_c$ as

$$w_c = \sum_{i=1}^{\infty} c_i (\gamma^\dagger)^{\frac{1}{2}} \nabla u_{\gamma^\dagger, f_i},$$

where $c_i \in \ell^2$ are coefficients to be specified below. Note that $\operatorname{div}((\gamma^\dagger)^{\frac{1}{2}} w_c) = 0$ since $\nabla u_{\gamma^\dagger, f_i}$ solves (1) with $\gamma = \gamma^\dagger$.

Expanding the square we find that

$$\begin{aligned} \|H_{\Delta\gamma} \nabla u_{\gamma, f} - w_c\|^2 &= \int H_{\Delta\gamma}^2 |\nabla u_{\gamma, f}|^2 dx - 2 \sum_{i=1}^{\infty} c_i \int_{\Omega} (\gamma - \gamma^\dagger) \nabla u_{\gamma, f} \cdot \nabla u_{\gamma^\dagger, f_i} dx \\ &\quad + \sum_{i, j=1}^{\infty} c_i c_j \int \gamma^\dagger \nabla u_{\gamma^\dagger, f_i} \cdot \nabla u_{\gamma^\dagger, f_j} dx \\ &= \langle \Lambda'_\gamma ((\Delta\gamma)^2) f, f \rangle - 2 \sum_{i=1}^{\infty} c_i \langle \Lambda_\gamma - \Lambda_{\gamma^\dagger} f, f_i \rangle \\ &\quad + \sum_{i, j=1}^{\infty} c_i c_j \langle \Lambda_{\gamma^\dagger} f_i, f_j \rangle. \end{aligned}$$

Note that f_i is an orthogonal basis such that $f_c = \sum c_i f_i$ represents an element in $H^{\frac{1}{2}}$ and hence, the last line can be written in terms of f, f_c and the Dirichlet-to-Neumann operators. Now minimizing over f_c yields that

$$\inf_c \|H_{\Delta\gamma} \nabla u_{\gamma, f} - w\|^2 = \langle \Lambda'_\gamma ((\Delta\gamma)^2) f, f \rangle - \langle \Lambda_{\gamma^\dagger}^{-1} (\Lambda_\gamma - \Lambda_{\gamma^\dagger}) f, (\Lambda_\gamma - \Lambda_{\gamma^\dagger}) f \rangle,$$

which proves the first result.

For the second one, we take

$$w_c = \sum_{i=1}^{\infty} c_i \frac{\gamma}{(\gamma^\dagger)^{\frac{1}{2}}} \nabla u_{\gamma, f_i},$$

which is again in R_{γ^\dagger} . As above, we find an analogous result with

$$\begin{aligned} & \|H_{\Delta\gamma} \nabla u_{\gamma, f} - w_c\|^2 \\ &= \int H_{\Delta\gamma}^2 |\nabla u_{\gamma, f}^2| dx - 2 \sum_{i=1}^{\infty} c_i \int_{\Omega} (\gamma - \gamma^\dagger) \frac{\gamma}{\gamma^\dagger} \nabla u_{\gamma, f} \cdot \nabla u_{\gamma, f_i} dx \\ & \quad + \sum_{i, j=1}^{\infty} c_i c_j \int \frac{\gamma^2}{\gamma^\dagger} \nabla u_{\gamma, f_i} \cdot \nabla u_{\gamma, f_j} dx \\ &= \langle \Lambda'_\gamma ((\Delta\gamma)^2) f, f \rangle - 2 \langle \Lambda'_\gamma \left((\gamma - \gamma^\dagger) \frac{\gamma}{\gamma^\dagger} \right) f, f_c \rangle + \langle \Lambda'_\gamma \left(\frac{\gamma^2}{\gamma^\dagger} \right) f_c, f_c \rangle. \end{aligned}$$

Proceeding as above yields the second result. \square

Next we derive some useful ordering and norm estimates: By rearranging terms and switching γ and γ^\dagger in (15) using the lower bound, and then the upper bound in (15), we obtain the following two inequalities.

Corollary 1. *Under the same assumptions as in Theorem 1, we have*

$$\Lambda'_\gamma (\gamma - \gamma^\dagger) \leq_L \Lambda(\gamma) - \Lambda(\gamma^\dagger) \leq_L \Lambda'_{\gamma^\dagger} (\gamma - \gamma^\dagger) \quad (21)$$

as well as

$$\Lambda'_{\gamma^\dagger} \left(\frac{\gamma^\dagger}{\gamma} (\gamma - \gamma^\dagger) \right) \leq_L \Lambda(\gamma) - \Lambda(\gamma^\dagger) \leq_L \Lambda'_\gamma \left(\frac{\gamma}{\gamma^\dagger} (\gamma - \gamma^\dagger) \right). \quad (22)$$

Note that the lower bound in (15) and consequently also in (21) was already been established by von Harrach and Seo [8] and used in various uniqueness and stability estimates [7, 9]. Our upper bound in (15) seem to be new to the knowledge of the author.

We may observe by a Taylor expansion, that the middle term in the inequality (15) is

$$-\frac{1}{2} F''[\gamma] (\gamma^\dagger - \gamma, \gamma^\dagger - \gamma) + o(|\gamma^\dagger - \gamma|^2) = -\frac{1}{2} F''[\gamma^\dagger] (\gamma^\dagger - \gamma, \gamma^\dagger - \gamma),$$

while the left-hand side is $F'[\gamma^\dagger] \left(\frac{|\gamma^\dagger - \gamma|^2}{\gamma^\dagger} \right) + o(|\gamma^\dagger - \gamma|^2)$. By setting $\gamma = \gamma^\dagger + \epsilon w$ for arbitrary $w \in L^\infty$, and taking the limit, we obtain

$$0 \leq -F''[\gamma^\dagger](w, w) \leq 2F'[\gamma^\dagger] \left(\frac{|w|^2}{\gamma^\dagger} \right).$$

Hence the second derivative of F is always negative definite. A similar definite result can be obtained for all even derivatives.

For later use, we also establish a related inequality:

Lemma 3. *Let $\xi_\dagger := \|\frac{\gamma-\gamma^\dagger}{\gamma^\dagger}\|_\infty < 1$ Then we have the estimates*

$$0 \leq_L \Lambda'_{\gamma^\dagger}(\gamma - \gamma^\dagger) - \Lambda'_\gamma(\gamma - \gamma^\dagger) \leq_L (2 + \xi_\dagger) \Lambda'_\gamma \left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger} \right) \quad (23)$$

Proof. Define

$$\mathcal{A} := \Lambda'_{\gamma^\dagger}(\gamma - \gamma^\dagger) - \Lambda'_\gamma(\gamma - \gamma^\dagger).$$

Then using Lemma 1 with $\gamma_1 = \gamma$, $\gamma_2 = \gamma^\dagger$, we find

$$\begin{aligned} \langle \mathcal{A}f, f \rangle &= \int (\gamma - \gamma^\dagger) (|\nabla u_{\gamma^\dagger, f}|^2 - |\nabla u_{\gamma, f}|^2) dx \\ &= \left(H_{\gamma - \gamma^\dagger} (I + H_{(\gamma^\dagger)^{-\frac{1}{2}}} R_{\gamma^\dagger} H_{\Delta\gamma}) \nabla u_{\gamma, f}, (I + H_{(\gamma^\dagger)^{-\frac{1}{2}}} R_{\gamma^\dagger} H_{\Delta\gamma}) \nabla u_{\gamma, f} \right) \\ &\quad - (H_{\gamma - \gamma^\dagger} \nabla u_{\gamma, f}, \nabla u_{\gamma, f}) \\ &= \left(\left[2H_{\Delta\gamma} R_{\gamma^\dagger} H_{\Delta\gamma} + H_{\Delta\gamma} R_{\gamma^\dagger} H_{\frac{\Delta\gamma}{(\gamma^\dagger)^{\frac{1}{2}}}} R_{\gamma^\dagger} H_{\Delta\gamma} \right] \nabla u_{\gamma, f}, \nabla u_{\gamma, f} \right) \\ &= \left(\left[2I + H_{\frac{\Delta\gamma}{(\gamma^\dagger)^{\frac{1}{2}}}} \right] R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f}, R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f} \right), \end{aligned}$$

where we used that $R_{\gamma^\dagger}^2 = R_{\gamma^\dagger}$ and all operators are selfadjoint. Since $\xi_\dagger = \left\| \frac{\Delta\gamma}{(\gamma^\dagger)^{\frac{1}{2}}} \right\|$, we find that

$$0 \leq_L (2 - \xi_\dagger)I \leq_L 2I + H_{\frac{\Delta\gamma}{(\gamma^\dagger)^{\frac{1}{2}}}} \leq_L (2 + \xi_\dagger)I,$$

and the lower bound follows as well as the first upper bound since

$$\begin{aligned} (R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f}, R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f}) &= \|R_{\gamma^\dagger} H_{\Delta\gamma} \nabla u_{\gamma, f}\|^2 \leq \|H_{\Delta\gamma} \nabla u_{\gamma, f}\|^2 \\ &= \langle \Lambda'_\gamma(\Delta\gamma)^2 f, f \rangle. \end{aligned}$$

□

As a consequence of Theorem 1 and 2 we find the following upper bounds

Theorem 3. *Under the same assumptions as in Theorem 1, we have*

$$\|F(\gamma) - F(\gamma^\dagger) - F'[\gamma](\gamma - \gamma^\dagger)\|_Y^2 \leq \|F'[\gamma] \left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger} \right)\|_Y^2. \quad (24)$$

Moreover, there exists a constant C depending only on $\underline{\alpha}, \bar{\alpha}, \Omega$ such that

$$\|F(\gamma) - F(\gamma^\dagger)\|_Y^2 \leq C \sum_{i \in I} \left\langle \Lambda'_\gamma \left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger} \right) f_i, f_i \right\rangle \quad (25)$$

$$\|F'[\gamma] \left(\frac{(\gamma - \gamma^\dagger)\gamma}{\gamma^\dagger} \right)\|_Y^2 \leq C \sum_{i \in I} \left\langle \Lambda'_\gamma \left(\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger} \right) f_i, f_i \right\rangle. \quad (26)$$

The last two upper bound can be interpreted as the trace norm of the operator $\mathcal{I}\Lambda'_\gamma(\frac{|\gamma-\gamma^\dagger|^2}{\gamma^\dagger})$, i.e., the ℓ^1 -norm of the eigenvalues.

Proof. The estimate (24) is a direct consequence of (15). In view of (25), we find as a consequence of (19) and the fact that $\Lambda_{\gamma^\dagger}^{-1}$ is positive definite that

$$0 \leq_L [\Lambda_\gamma - \Lambda_{\gamma^\dagger}] \Lambda_{\gamma^\dagger}^{-1} [\Lambda_\gamma - \Lambda_{\gamma^\dagger}] \leq_L \Lambda_\gamma ((\Delta\gamma)^2).$$

Standard elliptic estimates yield

$$\langle \Lambda_{\gamma^\dagger} f, f \rangle \sim \|f\|_{H^{\frac{1}{2}}}^2 = \langle \mathcal{I}f, f \rangle.$$

where \mathcal{I} is the Riesz isomorphism. With $\Delta F := [\Lambda_\gamma - \Lambda_{\gamma^\dagger}]$ we find that

$$\|\mathcal{I}(\Delta F)f\|_{H^{\frac{1}{2}}}^2 = \langle (\Delta F), \mathcal{I}(\Delta F)f \rangle \leq C \langle \Lambda'_\gamma ((\Delta\gamma)^2) f, f \rangle.$$

An equivalent alternative definition of the Hilbert-Schmidt norm is the sum of the left-hand side over an orthogonal system f_i , which yields the desired inequality (25). In the same way we obtain (26) from (20). \square

Remark 1. We note that the norm estimates (14), (24), (25), and (26) remain valid, wenn the Y -norm is replaced by the operator norm $\|\cdot\|_{L(H^{\frac{1}{2}}, H^{-\frac{1}{2}})}$. This follows from the fact that the inequalities are derived from the Löwner ordering and hold in particular for the largest eigenvalues, which agree with the operator norm.

3 Tangential cone conditions

We can now state the first sufficient conditions for the tangential cone conditions. An immediate corollary of the upper bound (24) leads to the first condition:

Theorem 4. *Let $\gamma, \gamma^\dagger \in D(F)$ with $\|\frac{\gamma-\gamma^\dagger}{\gamma^\dagger}\|_\infty < 1$. If*

$$\|F'[\gamma](\frac{|\gamma-\gamma^\dagger|^2}{\gamma^\dagger})\|_Y \leq \|F'[\gamma](\gamma - \gamma^\dagger)\|_Y,$$

then the weak tangential cone condition with $\eta \geq \frac{1}{2}$ is satisfied.

If for some $\zeta < 1$, it holds that

$$\|F'[\gamma](\frac{|\gamma-\gamma^\dagger|^2}{\gamma^\dagger})\|_Y \leq \zeta \|F'[\gamma](\gamma - \gamma^\dagger)\|_Y,$$

then the strong tangential cone condition with $\eta = \frac{\zeta}{1-\zeta}$ is satisfied.

Proof. In the second case, we have that

$$\begin{aligned} \|F'[\gamma](\gamma - \gamma^\dagger)\|_Y &\leq \|F'[\gamma](\gamma - \gamma^\dagger) - (F(\gamma) - F(\gamma^\dagger))\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y \\ &\leq \|F'[\gamma](\frac{|\gamma-\gamma^\dagger|^2}{\gamma^\dagger})\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y \\ &\leq \zeta \|F'[\gamma](\gamma - \gamma^\dagger)\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y. \end{aligned}$$

From this, we conclude that

$$\|F'[\gamma](\gamma - \gamma^\dagger)\|_Y \leq \frac{1}{1-\zeta} \|F(\gamma) - F(\gamma^\dagger)\|_Y$$

Thus,

$$\begin{aligned} \|(F(\gamma) - F(\gamma^\dagger) - F'[\gamma](\gamma - \gamma^\dagger))\|_Y &\leq \|F'[\gamma](\frac{|\gamma - \gamma^\dagger|^2}{\gamma^\dagger})\|_Y \\ &\leq \zeta \|F'[\gamma](\gamma - \gamma^\dagger)\|_Y \leq \frac{\zeta}{1 - \zeta} \|F(\gamma) - F(\gamma^\dagger)\|_Y. \end{aligned}$$

The weak cone condition for $\eta \geq \frac{1}{2}$ follows easily from (4) and the upper estimate (15). \square

Since the above conditions involve der Fréchet-derivative at γ , it is of interest to derive localized conditions that only require $F'[\gamma^\dagger]$. This is established in the next theorem.

Theorem 5. *Let $\gamma, \gamma^\dagger \in D(F)$ with $\|\frac{\gamma - \gamma^\dagger}{\gamma^\dagger}\|_\infty < 1$ and $\|\frac{\gamma - \gamma^\dagger}{\gamma}\|_\infty < 1$. If it holds that*

$$\|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|^2)\|_Y \leq \theta_\eta \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y, \quad (27)$$

with

$$\theta_\eta \leq \frac{\alpha \eta}{4 + \eta},$$

then the strong tangential cone condition is satisfied with this η . Moreover, the estimate (27) holds if there is a constant C such that

$$\|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|)\|_Y \leq C \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y, \quad (28)$$

and

$$\|\gamma - \gamma^\dagger\|_\infty \leq C \theta_\eta.$$

Proof. We have

$$\begin{aligned} \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y &\leq \|F'[\gamma^\dagger](\gamma - \gamma^\dagger) + (F(\gamma^\dagger) - F(\gamma))\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y \\ &\leq \|F'[\gamma^\dagger](\frac{|\gamma - \gamma^\dagger|^2}{\gamma})\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y \\ &\leq \frac{1}{\gamma} \|\infty\| \|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|^2)\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y \\ &\leq \frac{1}{\alpha} \theta_\eta \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y + \|F(\gamma) - F(\gamma^\dagger)\|_Y. \end{aligned}$$

Thus,

$$(1 - \frac{\theta_\eta}{\alpha}) \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y \leq \|F(\gamma) - F(\gamma^\dagger)\|_Y$$

Using (23), with γ, γ^\dagger swapped and the constant $C = 2 + \xi_\dagger = 2 + \|\frac{\gamma - \gamma^\dagger}{\gamma}\|_\infty$ we find,

$$\begin{aligned} &\|F(\gamma) - F(\gamma^\dagger) - F'[\gamma](\gamma^\dagger - \gamma)\|_Y \\ &\leq \|F(\gamma) - F(\gamma^\dagger) - F'[\gamma^\dagger](\gamma^\dagger - \gamma)\|_Y + \|F'[\gamma^\dagger](\gamma^\dagger - \gamma) - F'[\gamma](\gamma^\dagger - \gamma)\|_Y \\ &\leq \|F'[\gamma^\dagger](\frac{|\gamma - \gamma^\dagger|^2}{\gamma})\|_Y + C \|F'[\gamma^\dagger](\frac{|\gamma - \gamma^\dagger|^2}{\gamma})\|_Y \\ &\leq (1 + C) \|\frac{1}{\gamma}\|_\infty \|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|^2)\|_Y \leq (3 + \xi) \frac{1}{\alpha} \theta_\eta \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y \\ &\leq (3 + \xi) \frac{\frac{\theta_\eta}{\alpha}}{1 - \frac{\theta_\eta}{\alpha}} \|F(\gamma) - F(\gamma^\dagger)\|_Y. \end{aligned}$$

Thus if

$$(3 + \xi) \frac{\frac{\theta_\eta}{\underline{\alpha}}}{1 - \frac{\theta_\eta}{\underline{\alpha}}\|\infty} \leq \eta \Leftrightarrow \frac{\theta_\eta}{\underline{\alpha}} \leq \frac{\eta}{3 + \xi + \eta} \Leftarrow \frac{\theta_\eta}{\underline{\alpha}} \leq \frac{\eta}{3 + 1 + \eta},$$

then the strong tangential cone condition is verified. The sufficiency of (28) follows from this bound by using (14) with $w_1 = (\gamma - \gamma^\dagger)^2$ and $w_2 = \|\gamma - \gamma^\dagger\|_\infty |\gamma - \gamma^\dagger|$, and thus

$$\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)^2\|_Y \leq \|\gamma - \gamma^\dagger\|_\infty \|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|)\|_Y.$$

□

An immediate corollary is obtained for the in case that γ is below or above the true conductivity since such a monotonicity yields (27).

Corollary 2. *Let $\gamma, \gamma^\dagger \in D(F)$, with $\|\gamma - \gamma^\dagger\|_{L^\infty} \leq \underline{\alpha}$. Assume that either*

$$\gamma(x) \leq \gamma^\dagger(x) \quad \text{or} \quad \gamma(x) \geq \gamma^\dagger(x) \quad \forall x \in \Omega \text{ a.e.}$$

Then for

$$\|\gamma - \gamma^\dagger\|_\infty \leq \theta_\eta,$$

with θ_η as in Theorem 5, the strong tangential cone condition is satisfied with some given η .

Proof. Because of monotonicity we have that $|\gamma(x) - \gamma^\dagger(x)| = \pm(\gamma(x) - \gamma^\dagger(x))$, such that the result follows from (28) with $C = 1$. □

Another corollary of this theorem is that the cone conditions are satisfied if a source condition or a conditionals stability estimate holds.

Corollary 3. *Let X be a Hilbert space that is continuously embedded into L^∞ . Assume that a source condition*

$$\gamma - \gamma^\dagger = (F'[\gamma^\dagger]^* F'[\gamma^\dagger])^\mu \omega$$

holds with $\mu > \frac{1}{2}$, where $F'[\gamma^\dagger]^*$ is the adjoint in X . Then, for $\|\gamma - \gamma^\dagger\|_\infty$ sufficiently small, the strong tangential cone condition holds for a given $\eta \leq \frac{1}{2}$.

Proof. The left-hand side of (27) is bounded by

$$\|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|^2)\|_Y \leq L \|\gamma - \gamma^\dagger\|_\infty^2 \leq L \|\gamma - \gamma^\dagger\|_X^2, \quad L = \|F'[\gamma^\dagger]\|_{2,L(L^\infty,Y)}$$

The source condition implies a stability estimate (see, e.g., [5, p. 59])

$$\|\gamma - \gamma^\dagger\|_X \leq \|\omega\|^{\frac{1}{1+2\mu}} \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y^{\frac{2\mu}{2\mu+1}}.$$

Combining the inequalities yields

$$\begin{aligned} \|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|^2)\|_Y &\leq L \|\omega\|^{\frac{2}{1+2\mu}} \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y^{\frac{4\mu}{2\mu+1}-1} \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y \\ &\leq L \|\omega\|^{\frac{2}{1+2\mu}} (L \|\gamma - \gamma^\dagger\|_\infty)^{\frac{2\mu-1}{2\mu+1}} \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y. \end{aligned}$$

If $\mu < \frac{1}{2}$ then the assumptions in Theorem 5 are verified for

$$\|\gamma - \gamma^\dagger\|_\infty \leq \left(\frac{\theta_\eta}{L \|\omega\|^{\frac{2}{1+2\mu}}} \right)^{\frac{2\mu+1}{2\mu-1}}, \quad \|\gamma - \gamma^\dagger\|_\infty < \underline{\alpha},$$

with θ_η as given in this theorem. □

In particular, it follows that the tangential cone condition locally holds true in a dense subset of $N(F'[\gamma^\dagger])^\perp$.

An interesting observation is that the left-hand side in (28) generates a (semi)-norm

$$\|w\|_* := \|F'[\gamma^\dagger](|w|)\|_Y.$$

Indeed, take z_1, z_2 in L^∞ arbitrary, then $0 \leq |z_1(x) + z_2(x)| \leq |z_1(x)| + |z_2(x)|$. Then, the triangle inequality for $\|w\|_*$ follows from (14) with $w_1 = |z_1 + z_2|$ and $w_2 = |z_1| + |z_2|$ and the triangle inequality for $\|\cdot\|_Y$. Thus, (28) can be interpreted as an equivalence condition between two (semi)-norms. (Note that the reverse inequality is easy to obtain). The well-known norm equivalence in finite-dimensional spaces leads to the following result:

Corollary 4. *Let $\gamma, \gamma^\dagger \in D(F)$ with $\gamma - \gamma^\dagger$ being in a finite-dimensional space X_n , and let $F'[\gamma^\dagger]$ be injective on X_n . Then there is a dimension-dependent constant C_n such that for all*

$$\|\gamma - \gamma^\dagger\|_\infty \leq C_n \theta_\eta,$$

the strong tangential cone condition is satisfied for some given η .

We may slightly generalize the monotonicity result. We denote the positive and negative part of $\gamma - \gamma^\dagger$ by

$$(\gamma - \gamma^\dagger)^+(x) = \max\{\gamma(x) - \gamma^\dagger(x), 0\} \quad (\gamma - \gamma^\dagger)^- := -\min\{\gamma(x) - \gamma^\dagger(x), 0\}.$$

such that

$$\gamma - \gamma^\dagger = (\gamma - \gamma^\dagger)^+ - (\gamma - \gamma^\dagger)^-.$$

Theorem 6. *Assume that there exists a constant C or a constant $\nu < 1$ such that*

$$\|F'[\gamma^\dagger](\gamma - \gamma)^\dagger(x)\|_Y \leq C \|F'[\gamma^\dagger](\gamma - \gamma)\|_Y \quad \text{or} \quad (29)$$

$$\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)^+(x)\|_Y \leq \nu \|F'[\gamma^\dagger](\gamma - \gamma)^\dagger(x)\|_Y \quad (30)$$

(or the respective inequalities with $()^+$ and $()^-$ swapped) hold.

In case (29), if

$$\|\gamma - \gamma^\dagger\|_\infty \leq \theta_\eta \frac{1}{(2C + 1)}$$

holds, and in case (30), if

$$\|\gamma - \gamma^\dagger\|_\infty \leq \frac{\theta_\eta}{3}(1 - \nu)$$

holds, with θ_η from Theorem 5, then (27) is satisfied. In particular, under one of these conditions the tangential cone condition holds with any given η if additionally $\|\gamma - \gamma^\dagger\|$ is sufficiently small.

Proof. Set $p(x) = (\gamma - \gamma^\dagger)^+(x)$ and $n(x) = (\gamma - \gamma^\dagger)^+(-)$, then $\gamma - \gamma^\dagger = p - n$ and $(\gamma - \gamma^\dagger)^2 = p^2 + n^2$. We have

$$\begin{aligned} \|F'[\gamma^\dagger](|\gamma - \gamma^\dagger|)\|_Y &\leq \|F'[\gamma^\dagger](p + n)\|_Y \leq \|F'[\gamma^\dagger]p\|_Y + \|F'[\gamma^\dagger]n\|_Y \\ &\leq \|F'[\gamma^\dagger]p\|_Y + \|F'[\gamma^\dagger](n - p)\|_Y + \|F'[\gamma^\dagger]p\|_Y \\ &\leq 2C \|F'[\gamma^\dagger](p - n)\|_Y + \|F'[\gamma^\dagger](n - p)\|_Y \leq (2C + 1) \|F'[\gamma^\dagger](\gamma - \gamma)^\dagger\|_Y. \end{aligned}$$

Thus,

$$\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)^2\|_Y \leq \|\gamma - \gamma^\dagger\|_\infty (2C + 1) \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y$$

and (27) holds. In case of (30), we estimate

$$\begin{aligned} \|F'[\gamma^\dagger](\gamma - \gamma^\dagger)\|_Y &= \|F'[\gamma^\dagger]p - F'[\gamma^\dagger]n\|_Y \geq \left| \|F'[\gamma^\dagger]p\|_Y - \|F'[\gamma^\dagger]n\|_Y \right| \\ &\geq (1 - \nu) \|F'[\gamma^\dagger]p\|_Y. \end{aligned}$$

Hence (29) holds with $C = \frac{1}{1-\nu}$. The result then follows from the first part with $\nu \geq 0$. \square

Finally, as the most constructive result, we establish the local tangential cone conditions for C^2 -conductivities with “unbalanced” positive and negative part.

Theorem 7. *Assume that $\|\gamma - \gamma^\dagger\|_{C^2(\Omega)} \leq C_1$ and $\|(\gamma - \gamma^\dagger)\|_\infty \leq C_2 < \frac{\theta_\eta}{3}$ with θ_η from Theorem 5 given some $\eta \leq 1$. There is a nonnegative nondecreasing function ψ such that if*

$$\|(\gamma - \gamma^\dagger)^-\|_\infty \leq \psi(\|(\gamma - \gamma^\dagger)\|_\infty) \quad (31)$$

holds (or with the roles of + and - swapped), then the strong tangential cone condition is satisfied with this η .

Proof. Without loss of generality we assume that $\|(\gamma - \gamma^\dagger)^+\|_\infty > \|(\gamma - \gamma^\dagger)^-\|_\infty$ and hence that $\|(\gamma - \gamma^\dagger)\|_\infty = \|(\gamma - \gamma^\dagger)^+\|_\infty$. The case that that negative and positive part have equal norm is ruled out by the assumptions of the theorem. Let x_0 be a point in Ω , where the maximum M of $\delta\gamma := (\gamma - \gamma^\dagger)^+$, is attained. Then $\delta\gamma'(x_0) = 0$, and with the C^2 -bound we may find an estimate

$$(\gamma - \gamma^\dagger)^+(x) \geq M - \frac{C_1}{2} \|x - x_0\|^2$$

Thus for $\|x - x_0\|^2 \leq \frac{M}{C_1}$, we have that $(\gamma - \gamma^\dagger)^+(x) \geq \frac{M}{2}$. By the monotonicity result in (14) it follows that

$$\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)^+(x)\|_Y \geq \frac{\|(\gamma - \gamma^\dagger)^+\|_\infty}{2} \|F'[\gamma^\dagger]\chi_{B_{\frac{M}{C_1}}(x_0)}\|_Y$$

where $B_r(x_0)$ is the unit ball with center x_0 and radius r and χ denotes the characteristic function. Define

$$\kappa(m) := \inf_{B_m(x_0) \subset \Omega} \|F'[\gamma^\dagger]\chi_{B_m}(x_0)\|.$$

This defines a nonnegative and nondecreasing function. Thus,

$$\|F'[\gamma^\dagger](\gamma - \gamma^\dagger)^+(x)\|_Y \geq \frac{M}{2} \phi\left(\frac{M}{C_1}\right).$$

Define $\nu := 1 - \frac{3C_2}{\theta_\eta}$. By assumption in the theorem, $0 < \nu < 1$. We let

$$\psi(M) := \frac{1}{2L} M \kappa\left(\frac{M}{C_1}\right) \nu.$$

If (31) is satisfied with this ψ , then with $M = \|(\gamma - \gamma^\dagger)^+\|_\infty$, we have

$$\begin{aligned} \|F'[\gamma^\dagger](\gamma - \gamma)^\ominus(x)\|_Y &\leq L\|(\gamma - \gamma)^\ominus\|_\infty \leq \psi(\|(\gamma - \gamma)^\ominus\|_\infty) \\ &\leq \nu \frac{M}{2} \kappa\left(\frac{M}{C_1}\right) \leq \nu \|F'[\gamma^\dagger](\gamma - \gamma)^\ominus(x)\|_Y. \end{aligned}$$

Thus (30) holds and since

$$\frac{\theta_\eta}{3}(1 - \nu) = C_2 \geq \|(\gamma - \gamma)^\ominus\|_\infty,$$

the result follows from Theorem 6. \square

The relevance of this result is that we have the tangential cone condition satisfied in a L^∞ -ball if the positive (or negative) part of the difference of conductivities dominates the negative (resp. positive) part.

Remark 2. The last theorem might explain to some extent the convergence behaviour of the Landweber iteration for the impedance tomography problem. Fact is that the cone conditions are only required for the iterates γ_k and the true conductivity γ^\dagger . In many numerical experiments, the initial value for the iteration is often chosen as being strictly below the true conductivity, for instance, if γ^\dagger correspond to inclusions that have higher conductivities than the background one, and naturally the initial values of the iteration are taken as that background. Then by the last theorem, respectively, Corollary 2, the cone condition is satisfied for the initial iterate, and the iteration will stay at least bounded. This will also be true for a certain number of the following iterations. However, it is not guaranteed that this will hold for all iterations. In a lucky case, a monotonicity property will hold up to a stopping index, and then the iteration appear as convergent. However, in an unlucky case the cone condition might get violated with the effect that the iterates can diverge even though the stopping criteria is not yet met. This effect, may mistakenly be regarded as semiiteration, i.e., divergence by data error, although this has nothing to do with noisy data.

For a fair investigation of the convergence of the Landweber method, it would be interesting to start with a γ that has values below and above of γ^\dagger on a high number of regions, e.g., $\gamma_0 = \gamma^\dagger +$ highly oscillatory. An interesting question is whether the iterates of the Landweber method would still remain bounded in this case.

References

- [1] G. ALESSANDRINI AND S. VESSELLA, *Lipschitz stability for the inverse conductivity problem*, Adv. in Appl. Math., 35 (2005), pp. 207–241.
- [2] L. BORCEA, *Electrical impedance tomography*, Inverse Problems, 18 (2002), pp. R99–R136.
- [3] A.-P. CALDERÓN, *On an inverse boundary value problem*, in Seminar on Numerical Analysis and its Applications to Continuum Physics (Rio de Janeiro, 1980), Soc. Brasil. Mat., Rio de Janeiro, 1980, pp. 65–73.

- [4] M. V. DE HOOP, L. QIU, AND O. SCHERZER, *Local analysis of inverse problems: Hölder stability and iterative reconstruction*, Inverse Problems, 28 (2012), pp. 045001, 16.
- [5] H. W. ENGL, M. HANKE, AND A. NEUBAUER, *Regularization of inverse problems*, vol. 375 of Mathematics and its Applications, Kluwer Academic Publishers Group, Dordrecht, 1996.
- [6] M. HANKE, A. NEUBAUER, AND O. SCHERZER, *A convergence analysis of the Landweber iteration for nonlinear ill-posed problems*, Numer. Math., 72 (1995), pp. 21–37.
- [7] B. HARRACH, *Uniqueness and Lipschitz stability in electrical impedance tomography with finitely many electrodes*, Inverse Problems, 35 (2019), pp. 024005, 19.
- [8] B. HARRACH AND J. K. SEO, *Exact shape-reconstruction by one-step linearization in electrical impedance tomography*, SIAM J. Math. Anal., 42 (2010), pp. 1505–1518.
- [9] B. HARRACH AND M. ULLRICH, *Monotonicity-based shape reconstruction in electrical impedance tomography*, SIAM J. Math. Anal., 45 (2013), pp. 3382–3403.
- [10] V. ISAKOV, *Inverse problems for partial differential equations*, vol. 127 of Applied Mathematical Sciences, Springer-Verlag, New York, 1998.
- [11] B. KALTENBACHER, A. NEUBAUER, AND O. SCHERZER, *Iterative regularization methods for nonlinear ill-posed problems*, vol. 6 of Radon Series on Computational and Applied Mathematics, Walter de Gruyter GmbH & Co. KG, Berlin, 2008.
- [12] S. KINDERMANN, *Convergence of the gradient method for ill-posed problems*, Inverse Probl. Imaging, 11 (2017), pp. 703–720.
- [13] A. LECHLEITER AND A. RIEDER, *Newton regularizations for impedance tomography: convergence by local injectivity*, Inverse Problems, 24 (2008), pp. 065009, 18.
- [14] O. SCHERZER, *Convergence criteria of iterative methods based on Landweber iteration for solving nonlinear problems*, J. Math. Anal. Appl., 194 (1995), pp. 911–933.