On the Majorana solution to the Thomas-Fermi equation

Francisco M. Fernández^{*} and Javier Garcia NIFTA, DQT, Sucursal 4, C. C. 16, 1900 La Plata, Argentina

Abstract

We analyse the solution to the Thomas-Fermi equation discovered by Majorana. We show that the series for the slope at origin enables one to obtain results of accuracy far beyond those provided by available methods. We also estimate the radius of convergence of this series and conjecture that the singularity closest to origin is a square-root branch point.

1 Introduction

There has been great interest in the accurate calculation of the solution to the nonlinear differential equation that comes from the Thomas-Fermi model for neutral atoms [1–13]. Several approaches have been applied for this purpose; for example: Padé Hankel method [1–3, 5], fractional order of rational Euler functions [6], fractional order of rational Bessel functions collocation method [7], fractional order of rational Jacobi functions [8], rational Chebyshev functions [4], fractional order of rational Chebyshev functions of the second kind [10], a hybrid approach based on the collocation and Newton-Kantorovich methods plus fractional order of rational Legendre functions [11], Newton iteration with spectral

^{*}fernande@quimica.unlp.edu.ar

algorithms based on fractional order of rational Gegenbauer functions [12] and rational Chebyshev series accelerated through coordinate transformations [13]. We have just mentioned the most accurate results. Other authors have already obtained less accurate ones and more often than not reported many wrong digits as shown in the tables of some of the papers just mentioned [6–11].

The purpose of this paper is the discussion of a semi-analytical solution to the Thomas-Fermi equation discovered by Majorana in 1928 that remained unpublished for a long time as revealed in an enlightening pedagogical article by Esposito [14]. Although the Majorana solution was mentioned in some of the papers just quoted [4–8,8–10] none of those authors used it for testing their calculations. They did not realize that this approach enables one to obtain the slope at origin with any desired accuracy and in fact 100 significant digits have been reported in Wikipedia (https://en.wikipedia.org/wiki/Thomas%E2%80%93Fermi_equation#cite_note-11). As far as we know, this is the most accurate value of the slope at origin available nowadays.

One may reasonably argue that theoretical results of such an accuracy have no physical meaning. However, the Thomas-Fermi equation is commonly chosen as a benchmark for testing algorithms for solving nonlinear differential equations numerically and, for this reason, accurate results may be useful. The purpose of this paper is the analysis of the remarkable accuracy of the Majorana approach to the Thomas-Fermi equation. In section 2 we summarize the main equations shown by Esposito [14], in section 3 we analyze the convergence properties of the Majorana series numerically and in section 4 we summarize the main results and draw conclusions.

2 The Majorana transformation

The Thomas-Fermi equation for neutral atoms can be easily reduced to the dimensionless nonlinear second-order differential equation

$$\varphi''(x) = x^{-1/2} \varphi(x)^{3/2}, \ \varphi(0) = 1, \ \varphi(\infty) = 0.$$
(1)

In order to solve this equation one has to determine the unknown slope at origin $\varphi'_0 = \varphi'(0)$ that is consistent with the boundary conditions. By means of the change of independent and dependent variables

$$t = 144^{-1/6} x^{1/2} \varphi^{1/6}, \ u = -\left(\frac{16}{3}\right)^{1/3} \varphi^{-4/3} \varphi', \tag{2}$$

Majorana derived the nonlinear first-order differential equation [14]

$$\frac{du(t)}{dt} = 8\frac{tu(t)^2 - 1}{1 - t^2u(t)}, \ u(1) = 1.$$
(3)

It can be proved that the slope at origin is given by

$$\varphi_0' = -\left(\frac{3}{16}\right)^{1/3} u(0). \tag{4}$$

The solution u(t) can be expanded as

$$u(t) = \sum_{j=0}^{\infty} a_j \tau^j, \ a_0 = 1, \ \tau = 1 - t,$$
(5)

where the coefficients a_n satisfy the recurrence relation [14].

$$a_{m} = \frac{1}{2(m+8) - (m+1)a_{1}} \left\{ \sum_{n=1}^{m-2} a_{m-n} \left[(n+1)a_{n+1} - 2(n+4)a_{n} + (n+7)a_{n-1} \right] + \left[(m+7) - 2(m+3)a_{1} \right] a_{m-1} + (m+6)a_{1}a_{m-2} \right\}$$
(6)

and a_1 is a root of $a_1^2 - 18a_1 + 8 = 0$. If $a_1 = 9 - \sqrt{73}$ then the coefficients a_m are all positive and Esposito [14] estimated $a_n/a_{n-1} \sim 4/5$ for $n \to \infty$. On the other hand, when $a_1 = 9 + \sqrt{73}$ the magnitude of the coefficients $|a_n|$ appears to increase unboundedly and the ratio $|a_n/a_{n-1}|$ oscillates. Therefore, one expects to obtain reasonable results only for the first choice that we consider from now on.

It follows from equations (4) and (5) that one can obtain approximate solutions to the slope at origin from the sequence of partial sums

$$\varphi'_{0,N} = -\left(\frac{3}{16}\right)^{1/3} S_N, \ S_N = \sum_{j=0}^N a_j, \ N = 1, 2, \dots$$
 (7)

Esposito [14] estimated $\varphi'_0 = -1.588$ (notice that there is a misprint in his paper) and this result has been cited by several authors [6–10]. However, none of them tried to obtain accurate results from equations (6) and (7), except for the accurate value of the slope at origin in Wikipedia mentioned above.

This approach exhibits two great advantages: first, the expansion coefficients a_n are positive so that $S_N < S_{N+1}$ and the sequence converges from below. Second, the expansion coefficients decrease exponentially as shown in Figure 1 and, consequently, the rate of convergence is remarkable. From the first $N \leq 5000$ partial sums we estimate

 $\varphi_{0,5000}' = -1.58807102261137531271868450942395010945274662167482 \\ 561676567741816655196115430926233203397013842866526981 \\ 402442246654652538163078118812938288132770903659214744 \\ 086806778901475626169733142185514721772803432126893905 \\ 784750244483250437472432504160153205506367895247811586 \\ 772904901159198239550290936872919960125747115454641997 \\ 589356517736943217510752047768900396683865938312577933 \\ 839690794052221882937586$

that is supposed to be accurate to the last digit as we carried out the calculation numerically with sufficient accuracy (a simple Python program for this purpose is available at https://zenodo.org/record/4681779#.YH2Zorh1Yqg). It is unlikely that any of the approaches applied to this problem [1–7, 9–11, 13] (and references therein) can provide a result of such an accuracy. Notice that the error of a calculation based on S_N is roughly of the order of a_{N+1} .

3 Analysis of the series

In this section we will try to determine some features of the Majorana series (5) numerically. To this end we resort to a method discussed by Hunter and Guerrieri [15] some time ago that we develop in what follows in a less general

and simpler way, more suitable for present needs.

The function

$$f(x) = A \left(1 - \frac{x}{x_0} \right)^{\nu}, \tag{8}$$

exhibits a singular point at $x = x_0$ for any real exponent ν , except when it is a positive integer. It is sufficient for present purposes to consider x real. This function can be expanded in a Taylor series

$$f(x) = \sum_{j=0}^{\infty} f_j x^j, \tag{9}$$

that converges for all $|x| < |x_0|$. It follows from the differential equation

$$\left(1 - \frac{x}{x_0}\right)f' = -\frac{\nu}{x_0}f,\tag{10}$$

that the expansion coefficients satisfy the recurrence relation

$$(n+1)f_{n+1} = \frac{n}{x_0}f_n - \frac{\nu}{x_0}f_n, \ n = 0, 1, \dots$$
(11)

We appreciate that the ratio f_{n+1}/f_n is a linear function of 1/(n+1)

$$\frac{f_{n+1}}{f_n} = \frac{1}{x_0} - \frac{\nu+1}{x_0(n+1)}.$$
(12)

From equation (11) we can derive two linear equations

$$x_0(n+1)f_{n+1} + \nu f_n = nf_n,$$

$$x_0nf_n + \nu f_{n-1} = (n-1)f_{n-1},$$
(13)

from which we obtain

$$x_{0} = \frac{f_{n}f_{n-1}}{(n+1)f_{n+1}f_{n-1} - nf_{n}^{2}},$$

$$\nu = \frac{(n^{2}-1)f_{n+1}f_{n-1} - n^{2}f_{n}^{2}}{(n+1)f_{n+1}f_{n-1} - nf_{n}^{2}}.$$
(14)

As stated in section 2, the coefficients a_n are all positive and decrease exponentially which suggests that u(t) may exhibit a singular point as shown in equation (8). We can therefore obtain both the location τ_0 and the exponent ν of the singular point closest to the origin from the approximate straight line

$$\frac{a_{n+1}}{a_n} \approx \frac{1}{\tau_0} - \frac{\nu+1}{\tau_0(n+1)}, \ n \gg 1,$$
(15)

as suggested by equation (12). Alternatively, we may resort to equation (14) and estimate these parameters from

$$\tau_{0,n} = \frac{a_n a_{n-1}}{(n+1) a_{n+1} a_{n-1} - n a_n^2}, \ \tau_0 = \lim_{n \to \infty} \tau_{0,n},$$
$$\nu_n = \frac{(n^2 - 1) a_{n+1} a_{n-1} - n^2 a_n^2}{(n+1) a_{n+1} a_{n-1} - n a_n^2}, \ \nu = \lim_{n \to \infty} \nu_n.$$
(16)

The second and third columns of Table 1 show values of $\tau_{0,n}$ and ν_n , respectively, obtained from equation (16) for sufficiently large values of n. The convergence is rather slow but it seems that ν_n converges towards $\nu = 1/2$ from below when $n \to \infty$. The convergence of these sequences can be improved by means of of Aitken extrapolation [16] and results obtained from the last 15 entries in the third column of table 1 confirms this conjecture. The fourth column shows results for

$$\tau_{0,n} \left(\nu = 1/2\right) = \frac{n-1/2}{n+1} \frac{a_n}{a_{n+1}}.$$
(17)

It seems that $\tau_{0,n}(\nu_n)$ and $\tau_{0,n}(\nu = 1/2)$ appear to be monotonously decreasing and increasing, respectively. On assuming that this behaviour already applies to all *n* we conjecture that 1.20168605688 < τ_0 < 1.2016860577. Straightforward application of Aitken extrapolation to the last 15 entries in both columns enables us to sharpen those bounds to 1.2016860571 < τ_0 < 1.2016860575. The same analysis on the third column yields $\nu = 0.4998$.

On the other hand, from a numerical fit of the ratio a_{n+1}/a_n to a straight line we estimate $\tau_0 = 1.2016860577$ and $\nu = 0.4997$ in agreement with the results above.

4 Conclusions

In this paper we have shown that the Majorana transformation brought to light by Esposito [14] in a pedagogical paper enables one to obtain the slope at origin of the solution to the Thomas-Fermi equation with an accuracy that has not been achieved with any of the methods proposed so far. The slope at origin is necessary for the application of any approach because it is the relevant unknown in the nonlinear differential equation just mentioned. In addition to it, we estimated the parameters that determine the singular point closest to the origin of the function u(t) appearing in the Majorana transformation of the dimensionless Thomas-Fermi equation. We conjectured that the singularity is a square-root branch point and estimated its location with reasonable precision by means of lower and upper bounds.

References

- F. M. Fernández, Comment on: "Series solution to the Thomas-Fermi equation" [Phys. Lett. A 365 (2007) 111], Phys. Lett. A 372 (2008) 5258-5260.
- [2] S. Abbasbandy and C. Bervillier, Analytic continuation of Taylor series and the boundary value problems of some nonlinear ordinary differential equations, Appl. Math. Comput. 218 (2011) 2178-2199.
- [3] F. M. Fernández, Rational approximation to the Thomas-Fermi equations, Appl. Math. Comput. 207 (2011) 6433-6436.
- [4] J. P. Boyd, Rational Chebyshev Series for the Thomas-Fermi Function: Endpoint Singularities and Spectral Methods, J. Comp. Appl. Math. 244 (2013) 90-101.
- [5] P. Amore, J. P. Boyd, and F. M. Fernández, Accurate calculation of the solutions to the Thomas-Fermi equations, Appl. Math. Comput. 232 (2014) 929-943.
- [6] K. Parand, H. Yousefi, M. Delkhosh, and A. Ghaderi, A novel numerical technique to obtain an accurate solution to the Thomas-Fermi equation, Eur. Phys. J. Plus 131 (2016) 228.
- [7] K. Parand, A. Ghaderi, M. Delkhosh, and H. Yousefi, A new approach for solving nonlinear Thomas-Fermi equation based on fractional order rational Bessel functions, Elect. J. Diff. Eq. 2016 (2016) 1-18.

- [8] K. Parand, P. Mazaheri, and M. Delkhosh, Fractional order of rational Jacobi functions for solving the non-linear singular Thomas-Fermi equation, Eur. Phys. J. Plus 132 (2017) 77.
- [9] K. Parand and M. Delkhosh, Accurate solution of the Thomas-Fermi equation using the fractional order of rational Chebyshev functions, J. Comp. Appl. Math. 317 (2017) 624-642.
- [10] K. Parand, K. Raibei, and M. Delkhosh, An efficient numerical method for solving nonlinear Thomas-Fermi equation, Acta Univ. Sapientiae Mathematica 10 (2018) 134-151.
- [11] F. A. Parand, Z. Kalantari, M. Delkhosh, and F. Mirahmadian, A computationally hybrid Method for Solving a famous physical problem on an unbounded domain, Commun. Theor. Phys. 71 (2019) 9-15.
- [12] A. H. Hadian-Rasanan, N. Mehran, A. Bahramnezhad, M. M. Moayeri, and K. Parand, A comparison between pre-Newton and post-Newton approaches for solving a physical singular second-order boundary problem in the semi-infinite interval, arXiv:1909.04066 [math.NA]
- [13] X. Zhang and J. P. Boyd, Revisiting the Thomas-Fermi equation: accelerating rational Chebyshev series through coordinate transformations, Appl. Num. Math. 135 (2019) 186-205.
- [14] S. Esposito, Majorana solution of the Thomas-Fermi equation, Am. J. Phys. 70 (2002) 852-856.
- [15] C. Hunter and B. Guerrieri, Deducing the properties of singularities of functions from their Taylor series coefficients, SIAM J. Appl. Math. 39 (1980) 248-263.
- [16] G. Dahlquist and A Björck, Numerical Methods, Prentice-Hall, Englewood Cliffs, 1974.

Table 1: Parameters for the singular point

 $\tau_{0,n}(\nu_n)$ n400000 1.20168605769264029435361803588 1.20168605769043354866722488742401000 402000 1.20168605768824926820209670303 403000 1.2016860576860796865697237324404000 1.20168605768392281024420573709 405000 1.20168605768178079115740801082 406000 1.20168605767965783307100573377 407000 1.2016860576775544722516936779408000 1.20168605767546042850468283397 409000 1.20168605767338062590953642012410000 1 20168605767131927381271870206 411000 1.20168605766927692214546468536 412000 1.20168605766724329181078603638 413000 1.20168605766522800138146309189414000 1.20168605766322121156205411946415000 1.20168605766123753508189521398 416000 1.2016860576592620129791566382 417000 1.20168605765730427494019569306 418000 1.20168605765535448415903736306 1.20168605765342574745544322327419000 420000 1.20168605765150765389984597273 421000 1.20168605764960529812976599862 422000 1.201686057647710116814306249461.20168605764583596292330946038423000 424000 1.20168605764397169184576048614425000 1.20168605764212265017319322498426000 1.20168605764028028246218469624427000 1.20168605763845844575564932541 428000 1.20168605763664591943599140173 429000 1.20168605763484830282540948745430000 1.20168605763305679932369313611 431000 1.20168605763128535363451068823 432000 1.20168605762952275020995754139 433000 1.20168605762777459527635201156434000 1.2016860576260320960994033668435000 1.20168605762430916779681415112 436000 1.20168605762259470597020488796437000 1.20168605762089421604880153902 438000 1.20168605761919894526367540425 439000 1.2016860576175228147547327723440000 1 20168605761585472407406745998 441000 1.20168605761420016645747662319

 ν_n 0.499705403241764669703589077642 0.4997061387078161609832196909530.49970686850423206121412536936 0.4997075951930702647210850595880.499708319421314682402898280176 0.499709040445317571614381306288 0.4997097568209175629405145049790.4997104683319716692758525421780.499711178432852468843090560012 0.4997118854371809481420521073380 499712587885926043832104906147 0.499713285557507085244966336909 0.499713981941132448940288267412 0.499714673721611178261717764848 0.4997153642551535663759741445310.49971604848615901584789918849 0.499716731547383522068454193635 0.499717410088710305095020620403 0.4997180874991608873221026631340.4997187592004941474002086698290.499719428790372125524826408218 0.499720094468778248392634189577 0.499720759214935382469376826240.4997214181458194061123389614430.4997220751522537040502367000560.4997227283296280294345082607250.499723380683722672209827082160.499724027284788849437199852618 0.499724672088646643944149066554 0.4997253130836879635092475461510.499725953390862034718500932713 0.499726588003858223074707568887 0.4997272209148135760683241439440.499727850091775822559729268675 0.49972847868434786817072797909 0.4997291016512756204488330472750.499729722982543095775678947105 0.499730340664761125504584005280.499730957863058367311230730290.499731569488391115336882033527 0 499732179566977791786651829719 0.499732786072290704444252059124

 $\tau_{0,n}(\nu=1/2)$

1.20168605680760714618188774385 1.201686056809811459479000520751.20168605681199934431717238544 1.201686056814170963524403428431.201686056816326477946112427931.20168605681846604642319422634 1.201686056820589825867435067641.20168605682269797125288861779 1.20168605682479063564593371776 1.20168605682686797024999802803 1 20168605682893012444449010861 1.201686056830977245781374905041.20168605683300948001349255953 1.201686056835026971141774261291.201686056837029861429066602831.20168605683901829144174101888 1.201686056840992400029754386541.201686056842952324395139484651.2016860568448982000997565051 1.201686056846830161105965595941.2016860568487483397506528092 1.20168605685065286682152455643 1.20168605685254387155404180481.201686056854421481673399469431.201686056856285823370373190711.201686056858137021371835936051.201686056859975198938367447771.20168605686180047790432363325 1.20168605686361297865286240567 1.201686056865412820183749590841.201686056867200120111674723641.20168605686897499470303323157 1.20168605687073755885098422327 1.201686056872487926141350098011.20168605687422620884996548424 1.20168605687595251797812152143 1.20168605687766696322636321741 1.20168605687936965305979320755 1.20168605688106069470346942218 1.201686056882740194177039113611 2016860568844082562677809745 1.20168605688606498459441012161




Figure 1: Logarithm of the expansion coefficients