

Pair correlations of logarithms of integers

Jouni Parkkonen Frédéric Paulin

With an appendix by Étienne Fouvry

January 7, 2022

Abstract

We study the correlations of pairs of logarithms of positive integers at various scalings, either with trivial weights or with weights given by the Euler function, proving the existence of pair correlation functions. We prove that at the linear scaling, the pair correlations exhibit level repulsion, as it sometimes occurs in statistical physics. We prove total loss of mass phenomena at superlinear scalings, and Poissonian behaviour at sublinear scalings. The case of Euler weights has applications to the pair correlation of the lengths of common perpendicular geodesic arcs from the maximal Margulis cusp neighborhood to itself in the modular curve $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}_{\mathbb{R}}^2$.¹

1 Introduction

When studying the asymptotic distribution of a sequence of finite subsets of $\mathbb{R}$, finer information is sometimes given by the statistics of the spacing (or gaps) between pairs or k -tuples of elements, seen at an appropriate scaling. This problematic is largely developed in quantum chaos, including energy level spacings or clusterings, and in statistical physics, including molecular repulsion or interstitial distribution. The general setting of such a study may be described as follows. Let $\mathcal{F} = ((F_N)_{N \in \mathbb{N}}, \omega)$ be a nondecreasing sequence of finite subsets F_N of a finite dimensional Euclidean space E , endowed with a *multiplicity function* $\omega : \bigcup_{N \in \mathbb{N}} F_N \rightarrow]0, +\infty[$ (or *weight function*). Let $\psi : \mathbb{N} \rightarrow]0, +\infty[$ be a nondecreasing *scaling function*. The *pair correlation measure of $\mathcal{F}$ at time N with scaling $\psi(N)$* is the measure on E with finite support

$$\mathcal{R}_N^{\mathcal{F}, \psi} = \sum_{x, y \in F_N : x \neq y} \omega(x) \omega(y) \Delta_{\psi(N)(y-x)},$$

where Δ_z denotes the unit Dirac mass at z . When the sequence of measures $(\mathcal{R}_N^{\mathcal{F}, \psi})_{N \in \mathbb{N}}$, appropriately renormalized, weak-star converges to a measure $g \mathrm{Leb}_E$ absolutely continuous with respect to the Lebesgue measure Leb_E of E , the Radon-Nikodym derivative $g = g_{\mathcal{F}, \psi}$ is called the asymptotic *pair correlation function of $\mathcal{F}$ for the scaling ψ* . When $g_{\mathcal{F}, \psi}$ vanishes on a neighbourhood of 0 in E , we say that the pair $(\mathcal{F}, \psi)$ *exhibits level repulsion*.

¹**Keywords:** pair correlation, logarithms of integers, level repulsion, Euler function. **AMS codes:** 11K38, 11J83, 11N37, 53C22.

If the family $\mathcal{F}$ consists of subsets of the unit interval $[0, 1]$, then it is customary to use the cardinality of the finite set F_N as the scaling function. See for example [BocZ], where $F_N = \{\frac{p}{q} : p, q \in \mathbb{N}, p \leq q, (p, q) = 1, 0 < q \leq N\}$ is the set of Farey fractions of order N in $[0, 1]$ (without multiplicities, hence $\omega \equiv 1$), so that $\psi(N) = \frac{3N^2}{\pi} + O(N \ln N)$. Montgomery studied (under the Riemann hypothesis) the pair correlations of the imaginary parts of the zeros (with their multiplicity as zeros) of the Riemann zeta function ζ in the seminal paper [Mon]. The number of zeros $\frac{1}{2} + it$ of ζ with imaginary part t in the interval $[0, N]$ is asymptotic to $\frac{N \ln N}{2\pi}$ as $N \rightarrow +\infty$ and the scaling used in [Mon] is, analogously to the unit interval case of [BocZ], by the inverse of the *average gap* : $\psi(N) = (\frac{N \ln N}{2\pi})/N = \frac{\ln N}{2\pi}$.

In Sections 2 and 3, we study the pair correlations of the family of the logarithms of positive integers

$$\mathcal{L}_{\mathbb{N}} = ((L_N = \{\ln n : 0 < n \leq N\})_{N \in \mathbb{N}}, \omega \equiv 1)$$

without multiplicities. In order to simplify the statements in this introduction, we only consider power scalings $\psi : N \mapsto N^\alpha$ for $\alpha \geq 0$, and we denote these scaling functions by id^α .


Theorem 1.1 *Let $\alpha \geq 0$. As $N \rightarrow +\infty$, the normalized pair correlation measures $\frac{1}{N^{2-\alpha}} \mathcal{R}_N^{\mathcal{L}_{\mathbb{N}}, \text{id}^\alpha}$ on $\mathbb{R}$ weak-star converge to a measure $g_{\mathcal{L}_{\mathbb{N}}, \text{id}^\alpha} \text{Leb}_{\mathbb{R}}$ with pair correlation function given by*

$$g_{\mathcal{L}_{\mathbb{N}}, \text{id}^\alpha} : t \mapsto \begin{cases} \frac{1}{2} e^{-|s|} & \text{if } \alpha = 0 \\ \frac{1}{2} & \text{if } 0 < \alpha < 1 \\ \frac{1}{2t^2} [|t|] ([|t|] + 1) & \text{if } \alpha = 1 \\ 0 & \text{if } \alpha > 1 . \end{cases}$$

We refer to Theorems 2.1 and 3.1, for more complete versions of Theorem 1.1, with congruence restrictions and with more general scaling functions, as well as for error terms. These error terms, as well as the ones in Theorems 4.1 and 5.1, constitute the main technical parts of this paper.

The renormalisation by $\frac{1}{N^{2-\alpha}}$ in Theorem 1.1 is naturally chosen in order for the pair correlation function to be finite. As the finite set L_N , whose order is N , is contained in the minimal interval $[0, \ln N]$, the average gap in L_N is $\frac{\ln N}{N}$. Scaling by the inverse $\psi(N) = \frac{N}{\ln N}$ of the average gap (as in a particular case of Theorem 3.1), as well as by id^α for $0 < \alpha < 1$ (as in the above statement) gives a pair correlation function which is constant nonzero, a characteristic of a *Poissonian distribution*. As in the above result for $\alpha > 1$ and more generally by Theorem 3.1, if the scaling function ψ grows faster than linearly, then the pair correlation function vanishes : the empirical measures $\mathcal{R}_N^{\mathcal{L}_{\mathbb{N}}, \psi}$ have a total loss of mass at infinity, actually whatever the renormalisation is (the support of the measure itself converges to infinity). The transition from Poissonian to zero correlation occurs at linear scalings, where a more exotic pair correlation function appears (see for instance [RS] for Poissonian and [HK, LS] for non-Poissonian pair correlation phenomena). Since $g_{\mathcal{L}_{\mathbb{N}}, \text{id}}$ vanishes on $] -1, 1[$, the pair $(\mathcal{L}_{\mathbb{N}}, \text{id})$ exhibits a level repulsion.

The figure below gives the graph of the pair correlation function $g_{\mathcal{L}_{\mathbb{N}}, \text{id}}$ of $\mathcal{L}_{\mathbb{N}}$ at the linear scaling $\psi = \text{id} : N \mapsto N$ in the interval $[-15, 15]$ compared with the graph of the constant function $\frac{1}{2}$. The graph is similar to certain radial distribution functions in statistical physics, see for example [ZP, Sect. II], [SdH, Fig. 7], [Cha, page 199] or [Boh, page 18].



Instead of the pair correlations, one can study the gaps between consecutive elements in the subsets F_N of the real line or, most often, of the unit interval. Marklof and Strömbergsson [MaS] have computed the gap distribution of the fractional parts of the family $\mathcal{L}_{\mathbb{N}}$ (with a linear scaling and linear renormalisation) and showed that the limiting gap distribution has two jump discontinuities.

In Section 6, we prove that the pair correlation measures of the lengths of the common perpendiculars between the maximal Margulis cusp neighbourhood and itself in the modular curve $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}_{\mathbb{R}}^2$ are (up to a factor 2) the pair correlation measures of the weighted family

$$\mathcal{L}_{\mathbb{N}}^{\varphi} = ((L_N = \{\ln n : 0 < n \leq N\})_{N \in \mathbb{N}}, \omega = \varphi \circ \exp)$$

of logarithms of integers, with weights given by the Euler function $\varphi : n \mapsto \mathrm{Card}(\mathbb{Z}/n\mathbb{Z})^{\times}$, see Proposition 6.1. See [PS1, PS2] for results on the pair correlation of the lengths of closed geodesics in negatively curved manifolds.

We study the pair correlations of the arithmetically defined family $\mathcal{L}_{\mathbb{N}}^{\varphi}$ in Sections 4 and 5, where we find the pair correlation function without scaling and with linear scaling.


Theorem 1.2 (1) *As $N \rightarrow +\infty$, the pair correlation measures $\mathcal{R}_N^{\mathcal{L}_{\mathbb{N}}^{\varphi}, 1}$ on $\mathbb{R}$, renormalized to be probability measures, weak-star converge to the probability measure $g_{\mathcal{L}_{\mathbb{N}}^{\varphi}, 1} \mathrm{Leb}_{\mathbb{R}}$, with pair correlation function $g_{\mathcal{L}_{\mathbb{N}}^{\varphi}, 1} : s \mapsto e^{-2|s|}$.*

(2) *As $N \rightarrow +\infty$, the normalized pair correlation measures $\frac{1}{N^3} \mathcal{R}_N^{\mathcal{L}_{\mathbb{N}}^{\varphi}, \mathrm{id}}$ (with linear scaling) on $\mathbb{R}$ weak-star converge to the measure $g_{\mathcal{L}_{\mathbb{N}}^{\varphi}, \mathrm{id}} \mathrm{Leb}_{\mathbb{R}}$, with pair correlation function*

$$g_{\mathcal{L}_{\mathbb{N}}^{\varphi}, \mathrm{id}} : s \mapsto \frac{1}{s^4} \prod_{p \text{ prime}} \left(1 - \frac{2}{p^2}\right) \sum_{k=1}^{\lfloor |s| \rfloor} k^3 \prod_{p \text{ prime}, p \mid k} \left(1 + \frac{1}{p(p^2 - 2)}\right). \quad (1)$$

We refer to Theorems 4.1 and 5.1 for more complete versions of Theorem 1.2 with congruence restrictions, and for error terms. When the congruences are nontrivial, the proof of the second claim of Theorem 1.2 uses a generalization of Mirsky's formula (see [Mir]) that is proved in Appendix A by Étienne Fouvry.

The figure below gives the graph of the pair correlation function $g_{\mathcal{L}_{\mathbb{N}}^{\varphi}, \mathrm{id}}$ compared with the graph of the constant function with value $\frac{1}{4} \prod_{p \text{ prime}} \left(1 - \frac{2}{p^2}\right) \left(1 + \frac{1}{p^2(p^2 - 2)}\right) \simeq 0.09239$, which is the limit of the pair correlation function $g_{\mathcal{L}_{\mathbb{N}}^{\varphi}, \mathrm{id}}$ at $\pm\infty$ by Proposition 5.5.



Theorems 4.1 and 5.1 imply pair correlation results for the common perpendiculars of cusps neighborhoods in the modular curve and on quotients of the hyperbolic plane by Hecke congruence subgroups of $\mathrm{PSL}(\mathbb{Z})$, see Corollary 6.2 for precise statements.

Further directions. It would be interesting, given a discrete subgroup Γ of $\mathrm{PSL}_2(\mathbb{R})$, to study the asymptotic of the pair correlation measures of the complex translation lengths $\ell_{\mathbb{C}}(\gamma)$ with absolute value at most N of the elements $\gamma \in \Gamma$, and given a discrete subgroup Γ of a semi-simple connected real Lie group G with finite center and without compact factor, of the Cartan projections $\mu(\gamma)$ with Killing norm at most N of the elements $\gamma \in \Gamma$. See Section 6 for the problem of the asymptotic of the pair correlation measures of common perpendiculars in negative curvature, which will be studied more completely in subsequent works of the authors.

When the finite-dimensional Euclidean space E (where the family of finite sets $(F_N)_{N \in \mathbb{N}}$ sits) is replaced by a locally compact metric space (X, d) , we may also consider the positive measure on $]0, +\infty[$ with finite support $\mathcal{R}_N^{\mathcal{F}, \psi} = \sum_{x, y \in F_N : x \neq y} \omega(x) \omega(y) \Delta_{\psi(N)d(x, y)}$.

Acknowledgements: The authors thank a lot Etienne Fouvry for his proofs of Lemma 4.2, Proposition 5.5 and Theorem A.1 and for the agreeing to contribute the appendix to this paper. This research was supported by the French-Finnish CNRS IEA BARP.

Notation. We introduce here some of the notation used throughout the paper.

The pushforward of a measure μ by a mapping f is denoted by $f_*\mu$. We denote by $\mathrm{sg} : \mathbb{R} \rightarrow \mathbb{R}$ the change of sign map $t \mapsto -t$.

For every interval I in $\mathbb{R}$, we denote by Leb_I the Lebesgue measure on I and by $\mathbb{1}_I$ the characteristic function of I . We denote by $\mathrm{BV}(I)$ the vector space of measurable functions $f : I \rightarrow \mathbb{R}$ with finite total variation $\mathrm{Var}(f)$. For every $k \in \{0, 1\}$, we denote by $C_c^k(I)$ the real vector space of C^k -smooth functions $f : I \rightarrow \mathbb{C}$ with compact support in I . We denote by $\|f\|_{\infty} = \sup_{x \in I} |f(x)|$ the uniform norm of $f \in C_c^0(I)$.

In addition to the above, more or less standard, notation, we will use the following indexing sets in Sections 2, 3, 4 and 5. Let us fix throughout the paper $a, b \in \mathbb{N} - \{0\}$ with $a \leq b$. For every $N \in \mathbb{N} - \{0\}$, let

$$\begin{aligned} I_N &= I_{N, a, b} = \{(m, n) \in \mathbb{N}^2 : 0 < m, n \leq N, m \neq n, m, n \equiv a \pmod{b}\}, \\ I_N^- &= \{(m, n) \in \mathbb{N}^2 : 0 < m < n \leq N, m, n \equiv a \pmod{b}\}, \\ I_N^+ &= \{(m, n) \in \mathbb{N}^2 : 0 < n < m \leq N, m, n \equiv a \pmod{b}\}, \end{aligned}$$

so that $I_N = I_N^- \sqcup I_N^+$ is the disjoint union of I_N^- and I_N^+ .

Except in the proof of Lemma 4.2, of Proposition 5.5, and in the whole Appendix A, for every function g of a variable in $\mathbb{N} - \{0\}$, possibly depending on parameters (including a and b), we will denote by $O(g)$ (or $O_b(g)$ when we want to insist on the possible dependence on the parameter b) any function f on $\mathbb{N} - \{0\}$ such that there exists a constant C' depending only on the parameter b and a constant N_0 possibly depending on the parameters such that for every $N \geq N_0$, we have $|f(N)| \leq C' |g(N)|$.

2 Pair correlations without weights nor scaling

For every $N \in \mathbb{N} - \{0\}$, the (not normalised) *pair correlation measure* of the logarithms of integers congruent to a modulo b at time N , with trivial multiplicities and with trivial scaling function, is

$$\nu_N = \sum_{(m,n) \in I_N} \Delta_{\ln m - \ln n}.$$

If we consider the following nondecreasing sequence of finite subsets of $\mathbb{R}$ with trivial multiplicity

$$\mathcal{L}_N^{a,b} = (L_N^{a,b} = \{\ln n : 0 < n \leq N, n \equiv a \pmod{b}\})_{N \in \mathbb{N}}, \omega \equiv 1),$$

then, with the notation of the introduction, we have $\mathcal{L}_N^{1,1} = \mathcal{L}_N$ and $\nu_N = \mathcal{R}_N^{\mathcal{L}_N^{a,b},1}$.

Theorem 2.1 *As $N \rightarrow +\infty$, the measures ν_N on $\mathbb{R}$, renormalized to be probability measures, weak-star converge to the measure absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$, with Radon-Nikodym derivative the function $g_{\mathcal{L}_N^{a,b},1} : s \mapsto \frac{1}{2} e^{-|s|}$:*

$$\frac{\nu_N}{\|\nu_N\|} \xrightarrow{*} g_{\mathcal{L}_N^{a,b},1} \text{ Leb}_{\mathbb{R}}.$$

Furthermore, for every $f \in C_c^0(\mathbb{R}) \cap \text{BV}(\mathbb{R})$, we have

$$\frac{\nu_N}{\|\nu_N\|}(f) = \frac{1}{2} \int_{s \in \mathbb{R}} f(s) e^{-|s|} ds + O_b\left(\frac{\|f\|_{\infty} + \text{Var}(f)}{N}\right).$$

When $a = b = 1$, this result implies the case $\alpha = 0$ of Theorem 1.1 in the introduction, with pair correlation function $g_{\mathcal{L}_N,1} = g_{\mathcal{L}_N^{1,1},1}$.

Proof. For every $q \in \mathbb{N} - \{0\}$ with $q \equiv a \pmod{b}$, let $q' \in \mathbb{N}$ be such that $q = a + q'b$ and

$$J_q = \{p \in \mathbb{N} : 0 < p < q, p \equiv a \pmod{b}\} = \{a + kb : 0 \leq k < q'\}. \quad (2)$$

Let

$$\omega_q = \sum_{p \in J_q} \Delta_{\frac{p}{q}},$$

which is a finitely supported measure on $[0, 1]$, with total mass $\|\omega_q\| = q'$. When $q' \neq 0$, we hence have $\|\omega_q\| = \frac{q}{b} + O(1)$ and $\frac{1}{\|\omega_q\|} = \frac{b}{q} + O(\frac{1}{q^2})$. When $q' \neq 0$, we denote by $\overline{\omega}_q = \frac{\omega_q}{\|\omega_q\|}$ the renormalisation of ω_q to a probability measure on $[0, 1]$. By well known Riemann sum arguments, we have, as $q \rightarrow +\infty$,

$$\overline{\omega}_q \xrightarrow{*} \text{Leb}_{[0,1]}.$$

Let $f \in \text{BV}([0, 1])$, and note that f is bounded, with $\|f\|_\infty \leq f(0) + \text{Var}(f)$. Denoting by M_k and m_k the maximum and minimum respectively of f on $[\frac{a+kb}{q}, \frac{a+(k+1)b}{q}]$ for $0 \leq k < q'$, we have

$$\begin{aligned} & \left| \int_0^1 f(t) dt - \frac{b}{q} \omega_q(f) \right| = \left| \int_0^1 f(t) dt - \sum_{p \in J_q} \frac{b}{q} f\left(\frac{p}{q}\right) \right| \\ & \leq \left| \int_0^{\frac{a}{q}} f(t) dt \right| + \sum_{k=0}^{q'-1} \left| \int_{\frac{a+kb}{q}}^{\frac{a+(k+1)b}{q}} f(t) dt - \frac{b}{q} f\left(\frac{a+kb}{q}\right) \right| \\ & \leq \frac{b}{q} \|f\|_\infty + \sum_{k=0}^{q'-1} \frac{b}{q} (M_k - m_k) \leq (\|f\|_\infty + \text{Var}(f)) \frac{b}{q}. \end{aligned}$$

When $q' \neq 0$, since $|\omega_q(f)| \leq \|\omega_q\| \|f\|_\infty$, we hence have

$$\begin{aligned} \overline{\omega}_q(f) &= \int_0^1 f(t) dt - \int_0^1 f(t) dt + \frac{b}{q} \omega_q(f) + O\left(\frac{1}{q^2}\right) \omega_q(f) \\ &= \int_0^1 f(t) dt + O\left(\frac{\|f\|_\infty + \text{Var}(f)}{q}\right). \end{aligned}$$

For every $N \in \mathbb{N} - \{0\}$, with $N \geq a + b$, let us define

$$\mu_N^- = \sum_{(m,n) \in I_N^-} \Delta_{\frac{m}{n}} = \sum_{1 \leq q \leq N, q \equiv a \pmod{b}} \omega_q,$$

which is a finitely supported measure on $[0, 1]$. Its total mass is equal to

$$\|\mu_N^-\| = \sum_{1 \leq q \leq N, q \equiv a \pmod{b}} \|\omega_q\| = \sum_{0 \leq q' \leq \lfloor \frac{N-a}{b} \rfloor} (q' + O(1)) = \frac{N^2}{2b^2} + O(N).$$

Hence $\frac{1}{\|\mu_N^-\|} = \frac{2b^2}{N^2} + O(\frac{1}{N^3})$. For $f \in \text{BV}([0, 1])$, we have (taking $\|\omega_q\| \overline{\omega}_q(f) = 0$ if $q = a$)

$$\begin{aligned} \frac{\mu_N^-(f)}{\|\mu_N^-\|} &= \frac{1}{\|\mu_N^-\|} \sum_{1 \leq q \leq N, q \equiv a \pmod{b}} \|\omega_q\| \overline{\omega}_q(f) \\ &= \int_0^1 f(t) dt + \frac{1}{\|\mu_N^-\|} \sum_{1 \leq q \leq N, q \equiv a \pmod{b}} O(\|f\|_\infty + \text{Var}(f)) \\ &= \int_0^1 f(t) dt + O\left(\frac{\|f\|_\infty + \text{Var}(f)}{N}\right). \end{aligned}$$

Notice that $\ln$ is an increasing homeomorphism from $[0, 1]$ to $[-\infty, 0]$. For every element $N \in \mathbb{N} - \{0\}$, let us define

$$\nu_N^\pm = \sum_{(m,n) \in I_N^\pm} \Delta_{\ln \frac{m}{n}},$$

so that $\nu_N^- = \ln_* \mu_N^- = \nu_N|_{]-\infty, 0]}$, and $\|\nu_N^-\| = \|\mu_N^-\|$. We have, for every $f \in \text{BV}([-\infty, 0])$,

$$\begin{aligned} \frac{\nu_N^-(f)}{\|\nu_N^-\|} &= \frac{\mu_N^-(f \circ \ln)}{\|\mu_N^-\|} = \int_0^1 f \circ \ln(t) dt + O\left(\frac{\|f \circ \ln\|_\infty + \text{Var}(f \circ \ln)}{N}\right) \\ &= \int_{-\infty}^0 f(s) e^s ds + O\left(\frac{\|f\|_\infty + \text{Var}(f)}{N}\right). \end{aligned}$$

Since $\nu_N = \nu_N^- + \nu_N^+$, since $\nu_N^+ = \text{sg}_* \nu_N^-$, we have $\|\nu_N^\pm\| = \frac{1}{2} \|\nu_N\|$ and the result follows. $\square$

Let us give some numerical illustrations of Theorem 2.1 when $a = b = 1$. For every $N \in \mathbb{N} - \{0\}$, let

$$\mathcal{D}_N : s \mapsto \frac{\text{Card}\{(p, q) \in I_N : \ln \frac{p}{q} \leq s\}}{\text{Card } I_N},$$


which is the cumulative distribution function at time N of the differences of pairs of logarithms of integers, that is, for all $s, s' \in \mathbb{R}$ with $s < s'$, we have

$$\frac{\nu_N}{\|\nu_N\|} (]s, s']) = \mathcal{D}_N(s') - \mathcal{D}_N(s).$$

The above theorem says that the function $\mathcal{D}_N$ converges pointwise as $N \rightarrow +\infty$ to the C^1 (but not C^2) function

$$\mathcal{D} : s \mapsto \begin{cases} \frac{1}{2} e^s & \text{if } s \leq 0 \\ 1 - \frac{1}{2} e^{-s} & \text{if } s \geq 0 \end{cases}$$

(with derivative $\mathcal{D}' = g_{\mathcal{L}_{\mathbb{N}}, 1}$), which is the asymptotic cumulative distribution function of the differences of pairs of logarithms of integers. This is illustrated by the figure below, which shows $\mathcal{D}_{15}$ in green.



3 Pair correlations without weights and with scaling

In this section, we study the pair correlations of logarithms of integers at various scaling, now assumed to converge to $+\infty$. We fix two nondecreasing positive functions, respectively $\psi : \mathbb{N} - \{0\} \rightarrow]0, +\infty[$ and $\psi' : \mathbb{N} - \{0\} \rightarrow]0, +\infty[$, which will give the scaling factors on the difference of pairs of logarithms and the renormalizing factors on their distribution.

For every $N \in \mathbb{N} - \{0\}$, the (not normalised) *pair correlation measure* of the logarithms of integers congruent to a modulo b at time N with trivial multiplicities and with scaling $\psi(N)$ is the (Borel, positive) measure with finite support in $\mathbb{R}$ defined by

$$\mathcal{R}_N^{\mathcal{L}_{\mathbb{N}}, \psi} = \sum_{(m, n) \in I_N} \Delta_{\psi(N)(\ln m - \ln n)},$$

and the normalized one is $\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_{\mathbb{N}}, \psi}$.

Theorem 3.1 Assume that the nondecreasing positive function ψ satisfies $\lim_{+\infty} \psi = +\infty$ and $\lim_{N \rightarrow +\infty} \frac{\psi(N)}{N} = \lambda_\psi \in [0, +\infty]$. As $N \rightarrow +\infty$, the measures $\mathcal{R}_N^{\mathcal{L}_N, \psi}$ on $\mathbb{R}$, normalized by $\psi'(N)$ as given below, weak-star converge to a measure $g_{\mathcal{L}_N, \psi} \text{Leb}_{\mathbb{R}}$ absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$,

$$\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi} \xrightarrow{*} g_{\mathcal{L}_N, \psi} \text{Leb}_{\mathbb{R}},$$

with Radon-Nikodym derivative the function

$$g_{\mathcal{L}_N, \psi} : t \mapsto \begin{cases} 0 & \text{if } \lambda_\psi = +\infty, \text{ for any } \psi', \\ \frac{1}{2b^2} & \text{if } \lambda_\psi = 0 \text{ and } \psi'(N) = \frac{N^2}{\psi(N)}, \\ \frac{1}{2t^2} \lfloor \frac{|t|}{b\lambda_\psi} \rfloor \left(\lfloor \frac{|t|}{b\lambda_\psi} \rfloor + 1 \right) & \text{if } \lambda_\psi \neq 0, +\infty \text{ and } \psi'(N) = \psi(N). \end{cases}$$

Furthermore, if $\lambda_\psi \neq 0, +\infty$, for every $f \in C_c^1(I)$ with support contained in $[-A, A]$, we have

$$\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi}(f) = \int_{s \in \mathbb{R}} f(s) g_{\mathcal{L}_N, \psi}(s) ds + O_b \left(A \|f\|_\infty \left(|\lambda_\psi - \frac{\psi(N)}{N}| + \frac{A}{N} \right) + \|f'\|_\infty \frac{A^3}{N} \right).$$

The pair correlation function $g_{\mathcal{L}_N, \psi}$ depends on b but it is independent of a . The above result shows in particular that renormalizing to probability measures (taking $\psi'(N) = N^2 - N$) is inappropriate, as the limiting measure would always be 0.

When $\alpha = 0$, $a = b = 1$ and $\psi = \text{id}^\alpha : N \rightarrow N^\alpha$, the measure $\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi}$ corresponds to the one denoted by $\frac{1}{N^{2-\alpha}} \mathcal{R}_N^{\mathcal{L}_N, \text{id}^\alpha}$ in the introduction. The above result thus implies the cases $\alpha > 0$ of Theorem 1.1 in the introduction, as well as the comment about the scaling by the inverse of the average gap $\psi(N) = \frac{N}{\ln N}$, for which $\lambda_\psi = 0$.

The fact that $g_{\mathcal{L}_N, \psi}$ vanishes when $\lambda_\psi = +\infty$ means that the sequence of measures $(\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi})_{N \in \mathbb{N}}$ on $\mathbb{R}$ has a total loss of mass at infinity. For error terms when $\lambda_\psi = +\infty$ and $\lambda_\psi = 0$, see respectively Equation (6) and Equation (8).

Proof. Note that the change of variables $(m, n) \mapsto (n, m)$ in I_N proves that we have $\mathcal{R}_N^{\mathcal{L}_N, \psi} |_{]-\infty, 0]} = \text{sg}_* (\mathcal{R}_N^{\mathcal{L}_N, \psi} |_{[0, +\infty[})$. We will thus only study the convergence of the measures $\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi}$ on $[0, +\infty[$, and deduce the global result by the symmetry of $g_{\mathcal{L}_N, \psi}$.

For every $N \in \mathbb{N} - \{0\}$ and for every $p \in \mathbb{N}$ with $p \equiv 0 \pmod b$ and $0 < p < N$, let $N_p = \lfloor \frac{N-p-a}{b} \rfloor$, let

$$J_{p, N} = \{q \in \mathbb{N} : 1 \leq q \leq N - p, q \equiv a \pmod b\} = \{a + kb : 0 \leq k \leq N_p\}, \quad (3)$$

and let

$$\omega_{p, N} = \sum_{q \in J_{p, N}} \Delta_{\psi(N) \frac{p}{q}} \quad \text{and} \quad \mu_N^+ = \sum_{0 < p < N, p \equiv 0 \pmod b} \omega_{p, N}.$$

Then $\omega_{p, N}$ is a measure on $[0, +\infty[$, with finite support contained in $[\frac{\psi(N)}{N-p}p, \psi(N)p]$, and with total mass $\|\omega_{p, N}\| = N_p + 1$. When this total mass is not 0, which occurs if and only if $N \geq p+a$, we denote by $\overline{\omega}_{p, N} = \frac{\omega_{p, N}}{\|\omega_{p, N}\|}$ the renormalisation of $\omega_{p, N}$ to a probability measure on $]0, +\infty[$. The support of the measure μ_N^+ on $]0, +\infty[$ is contained in $[\frac{\psi(N)}{N}, \psi(N)N]$. The motivation for the definition of the measure μ_N^+ comes from the following lemma.

Lemma 3.2 *For every $A > 0$ and for every $f \in C_c^1(\mathbb{R})$ with compact support contained in $[0, A]$, we have, as $N \rightarrow +\infty$,*

$$| \mathcal{R}_N^{\mathcal{L}_N, \psi}(f) - \mu_N^+(f) | = O \left(A^3 \|f'\|_\infty \left(\frac{N}{\psi(N)} \right)^2 \right).$$

In particular, if $\frac{1}{\psi'(N)} \left(\frac{N}{\psi(N)} \right)^2$ tends to 0 as $N \rightarrow +\infty$, the measures $\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi} |_{[0, +\infty]}$ and $\frac{1}{\psi'(N)} \mu_N^+$ on $[0, +\infty]$ are asymptotic for the weak-star convergence of measures on $[0, +\infty]$, and we will study the weak-star convergence of the latter one.

Proof. By the change of variable $(p, q) \mapsto (m = p + q, n = q)$, we have

$$\mathcal{R}_N^{\mathcal{L}_N, \psi} |_{[0, +\infty]} = \sum_{(m, n) \in I_N^+} \Delta_{\psi(N) \ln \frac{m}{n}} = \sum_{\substack{0 < q \leq N-p, q \equiv a \pmod{b} \\ 0 < p < N, p \equiv 0 \pmod{b}}} \Delta_{\psi(N) \ln(1 + \frac{p}{q})}.$$

By definition, we have

$$\mu_N^+ = \sum_{\substack{0 < q \leq N-p, q \equiv a \pmod{b} \\ 0 < p < N, p \equiv 0 \pmod{b}}} \Delta_{\psi(N) \frac{p}{q}}.$$

Since the support of f is contained in $[0, A]$, if a pair (p, q) occurs in the index of the sum defining either $\mathcal{R}_N^{\mathcal{L}_N, \psi}(f)$ or $\mu_N^+(f)$ with nonzero summand, then $\psi(N) \ln(1 + \frac{p}{q}) \leq A$. This implies that $\frac{p}{q} = O\left(\frac{A}{\psi(N)}\right)$ since $\lim_{+\infty} \psi = +\infty$, and that $p = O\left(\frac{AN}{\psi(N)}\right)$ since $q \leq N$. For all $x, y \in [0, +\infty]$, we have

$$|\Delta_x(f) - \Delta_y(f)| = |f(x) - f(y)| \leq \|f'\|_\infty |x - y|.$$

Recall that $|\ln(1 + t) - t| = O(t^2)$ as $t \rightarrow 0$. Hence, by a uniform majoration of the terms of the sum below,


$$| \mathcal{R}_N^{\mathcal{L}_N, \psi}(f) - \mu_N^+(f) | \leq \sum_{\substack{1 \leq q \leq N \\ 1 \leq p \leq O\left(\frac{AN}{\psi(N)}\right)}} \|f'\|_\infty \psi(N) O\left(\left(\frac{p}{q}\right)^2\right) = O\left(A^3 \|f'\|_\infty \left(\frac{N}{\psi(N)}\right)^2\right). \quad \square$$

Let us now study the convergence properties of the (renormalized) measures $\omega_{p, N}$ and of their sums μ_N^+ as $N \rightarrow +\infty$.

Let $\iota :]0, +\infty[\rightarrow]0, +\infty[$ be the involutive diffeomorphism $t \mapsto \frac{1}{t}$. We have

$$\iota_* \omega_{p, N} = \sum_{q \in J_{p, N}} \Delta_{\frac{q}{\psi(N)p}}.$$

As q varies in $J_{p, N}$, the above Dirac masses are taken on the distribution of points given by the following picture.



As in the proof of Theorem 2.1, for every C^1 function $f :]0, +\infty[\rightarrow \mathbb{R}$ with compact support, we have

$$\begin{aligned} & \left| \int_0^{\frac{N-p}{\psi(N)p}} f(t) dt - \frac{b}{\psi(N)p} \iota_* \omega_{p, N}(f) \right| = \left| \int_0^{\frac{N-p}{\psi(N)p}} f(t) dt - \sum_{q \in J_{p, N}} \frac{b}{\psi(N)p} f\left(\frac{q}{\psi(N)p}\right) \right| \\ & \leq \left| \int_0^{\frac{a}{\psi(N)p}} f(t) dt \right| + \sum_{k=0}^{N_p-1} \left| \int_{\frac{a+kb}{\psi(N)p}}^{\frac{a+(k+1)b}{\psi(N)p}} f(t) dt - \frac{b}{\psi(N)p} f\left(\frac{a+kb}{\psi(N)p}\right) \right| + \left| \int_{\frac{a+N_p b}{\psi(N)p}}^{\frac{N-p}{\psi(N)p}} f(t) dt \right| \\ & \leq (2 \|f\|_{]0, \frac{N-p}{\psi(N)p}] } + \text{Var}(f|_{]0, \frac{N-p}{\psi(N)p}]}) \frac{b}{\psi(N)p}. \end{aligned}$$

Hence for every C^1 function $f :]0, +\infty[\rightarrow \mathbb{R}$ with compact support, since ι is a diffeomorphism, we have

$$\begin{aligned} \omega_{p, N}(f) &= \frac{\psi(N)p}{b} \int_0^{\frac{N-p}{\psi(N)p}} f \circ \iota(s) ds + O(\|f \circ \iota\|_{]0, \frac{N-p}{\psi(N)p}] } + \text{Var}(f \circ \iota|_{]0, \frac{N-p}{\psi(N)p}]}) \\ &= \frac{\psi(N)p}{b} \int_{\frac{\psi(N)p}{N-p}}^{+\infty} f(t) \frac{dt}{t^2} + O(\|f\|_{[\frac{\psi(N)p}{N-p}, +\infty[} + \text{Var}(f|_{[\frac{\psi(N)p}{N-p}, +\infty[})). \end{aligned}$$

For every $t > 0$, let

$$\theta_N(t) = \frac{1}{t^2} \sum_{\substack{0 < p < N \\ p \equiv 0 \pmod{b}}} \frac{p}{b} \mathbb{1}_{[\frac{\psi(N)p}{N-p}, +\infty[}(t).$$

Then

$$\begin{aligned} \theta_N(t) &= \frac{1}{t^2} \sum_{0 < k < N/b} k \mathbb{1}_{[\frac{\psi(N)bk}{N-bk}, +\infty[}(t) = \frac{1}{t^2} \sum_{0 < k \leq \frac{tN}{b(\psi(N)+t)}} k \\ &= \frac{1}{2t^2} \left\lfloor \frac{tN}{b(\psi(N)+t)} \right\rfloor \left(\left\lfloor \frac{tN}{b(\psi(N)+t)} \right\rfloor + 1 \right). \end{aligned} \quad (4)$$

In particular, we have $\theta_N(t) = 0$ if and only if $t \in [0, \frac{b\psi(N)}{N-b}]$.

Thus, if the support of f is contained in the interval $[0, A]$, since $\frac{\psi(N)p}{N-p} \leq A$ if and only if $p \leq \frac{AN}{\psi(N)+A}$, we have,

$$\begin{aligned} \mu_N^+(f) &= \sum_{0 < p < N, p \equiv 0 \pmod{b}} \omega_{p, N}(f) \\ &= \psi(N) \int_0^{+\infty} f(t) \left(\sum_{\substack{0 < p < N \\ p \equiv 0 \pmod{b}}} \frac{p}{b} \mathbb{1}_{[\frac{\psi(N)p}{N-p}, +\infty[}(t) \right) \frac{dt}{t^2} + O((\|f\|_\infty + \text{Var}(f)) \frac{AN}{\psi(N)}) \\ &= \psi(N) \int_0^{+\infty} f(t) \theta_N(t) dt + O((\|f\|_\infty + \text{Var}(f)) \frac{AN}{\psi(N)}). \end{aligned} \quad (5)$$

Case 1. Assume first that $\lambda_\psi = +\infty$, that is, $\lim_{N \rightarrow +\infty} \frac{N}{\psi(N)} = 0$. Then for every $A \geq 1$, if N is large enough, then for every $t \in [0, A]$, we have $\theta_N(t) = 0$. Thus, whatever the normalizing function ψ' is, we have a total loss of mass at infinity :

$$\frac{1}{\psi'(N)} \mu_N^+ \xrightarrow{*} 0.$$

More precisely, for every C^1 function $f :]0, +\infty[\rightarrow \mathbb{R}$ with compact support contained in $[0, A]$, we have

$$\frac{1}{\psi'(N)} \mu_N^+(f) = O\left((\|f\|_\infty + \text{Var}(f)) \frac{AN}{\psi(N)\psi'(N)}\right).$$

Since $\text{Var}(f) = \int_0^A |f'| \leq A \|f'\|_\infty$, we have, by Lemma 3.2, that $\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi} \xrightarrow{*} 0$ and

$$\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi}(f) = O\left((\|f\|_\infty + \|f'\|_\infty) \frac{A^3 N}{\psi(N)\psi'(N)}\right). \quad (6)$$

Case 2. Now assume that $\lambda_\psi = 0$, that is, $\lim_{N \rightarrow +\infty} \frac{\psi(N)}{N} = 0$. By Equation (4), if we have $t \geq \frac{b\psi(N)}{N-b}$, then

$$\theta_N(t) = \frac{1}{2t^2} \left(\frac{tN}{b(\psi(N) + t)} + O(1) \right)^2 = \frac{N^2}{2b^2\psi(N)^2} \left(1 + O\left(\frac{t}{\psi(N)}\right) + O\left(\frac{\psi(N)}{tN}\right) \right)^2,$$

therefore

$$\frac{\psi(N)^2}{N^2} \theta_N(t) = \frac{1}{2b^2} + O\left(\frac{t}{\psi(N)}\right) + O\left(\frac{\psi(N)}{tN}\right). \quad (7)$$

Since θ_N vanishes on $[0, \frac{b\psi(N)}{N-b}[$, this proves that $\frac{\psi(N)^2}{N^2} \theta_N$ is bounded on any compact subset of $[0, +\infty[$, and pointwise converges to the constant function $\frac{1}{2b^2}$. Hence by the Lebesgue dominated convergence theorem, we have

$$\frac{\psi(N)}{N^2} \mu_N^+ \xrightarrow{*} \frac{1}{2b^2} \text{Leb}_{[0, +\infty[}.$$

More precisely, for every $A \geq 3$, for every C^1 function $f :]0, +\infty[\rightarrow \mathbb{R}$ with compact support contained in $[0, A]$, by Equations (5) and (7) and since $\psi(N) \leq N$ for N large enough, we have

$$\begin{aligned} \frac{\psi(N)}{N^2} \mu_N^+(f) &= \frac{1}{2b^2} \int_{\frac{b\psi(N)}{N-b}}^{+\infty} f(t) dt + O\left(\frac{1}{\psi(N)} \int_{\frac{b\psi(N)}{N-b}}^A t |f(t)| dt\right) \\ &\quad + O\left(\frac{\psi(N)}{N} \int_{\frac{b\psi(N)}{N-b}}^A \frac{1}{t} |f(t)| dt\right) + O\left((\|f\|_\infty + \text{Var}(f)) \frac{A}{N}\right) \\ &= \frac{1}{2b^2} \int_0^{+\infty} f(t) dt + O\left(\frac{\psi(N)}{N} \|f\|_\infty\right) + O\left(\frac{A^2}{\psi(N)} \|f\|_\infty\right) \\ &\quad + O\left(\frac{\psi(N)}{N} \|f\|_\infty \left(\ln A - \ln \frac{b\psi(N)}{N-b}\right)\right) + O\left((\|f\|_\infty + \|f'\|_\infty) \frac{A^2}{N}\right) \\ &= \frac{1}{2b^2} \int_0^{+\infty} f(t) dt + O\left(\|f\|_\infty \left(\frac{\psi(N) \ln A}{N} + \frac{A^2}{\psi(N)} - \frac{\psi(N)}{N} \ln \frac{\psi(N)}{N}\right)\right) \\ &\quad + O\left(\|f'\|_\infty \frac{A^2}{N}\right). \end{aligned}$$

Let $\psi'(N) = \frac{N^2}{\psi(N)}$. By Lemma 3.2, we hence have

$$\frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi} \xrightarrow{*} \frac{1}{2b^2} \text{Leb}_{\mathbb{R}}$$

Futhermore, for every C^1 function $f : \mathbb{R} \rightarrow \mathbb{R}$ with compact support contained in $[-A, A]$, we have

$$\begin{aligned} \frac{1}{\psi'(N)} \mathcal{R}_N^{\mathcal{L}, \psi}(f) &= \frac{1}{2b^2} \int_{\mathbb{R}} f(t) dt + O\left(\|f\|_{\infty} \left(\frac{\psi(N) \ln A}{N} + \frac{A^2}{\psi(N)} - \frac{\psi(N)}{N} \ln \frac{\psi(N)}{N} \right) \right) \\ &\quad + O\left(\|f'\|_{\infty} \frac{A^3}{\psi(N)}\right). \end{aligned} \quad (8)$$

Case 3. Let us finally assume that $\lim_{N \rightarrow +\infty} \frac{\psi(N)}{N} = \lambda_{\psi}$ belongs to $]0, +\infty[$. Let us consider the map $\theta_{\infty} : [0, +\infty[\rightarrow \mathbb{R}$ defined by

$$t \mapsto \frac{1}{t^2} \sum_{k=1}^{\infty} k \mathbb{1}_{[b\lambda_{\psi}k, +\infty[}(t) = \frac{1}{2t^2} \left\lfloor \frac{t}{b\lambda_{\psi}} \right\rfloor \left(\left\lfloor \frac{t}{b\lambda_{\psi}} \right\rfloor + 1 \right).$$

It vanishes on $[0, b\lambda_{\psi}[$, is uniformly bounded, tends to $\frac{1}{2b^2\lambda_{\psi}^2}$ as $t \rightarrow +\infty$, and is piecewise continous, with discontinuities at $b\lambda_{\psi}\mathbb{N} - \{0\}$. See the first picture in the introduction when $a = b = \lambda_{\psi} = 1$.

By Equation (4), the sequence of uniformly bounded maps $(\theta_N)_{N \in \mathbb{N}}$ converges almost everywhere to θ_{∞} (more precisely, it converges at least at every point of $[0, +\infty[- b\lambda_{\psi}\mathbb{N}$). Hence by Equation (5) and by the Lebesgue dominated convergence theorem, we have

$$\frac{1}{\psi(N)} \mu_N^+ \xrightarrow{*} \theta_{\infty} \text{ Leb}_{[0, +\infty[}.$$

Let $A \geq 1$ and $k \in \mathbb{N}$. Note that $b\lambda_{\psi}k \leq A$ implies that $k \leq \frac{A}{b\lambda_{\psi}} \leq \frac{2A}{b\lambda_{\psi}}$. If N is large enough so that $\frac{\psi(N)}{N} \geq \frac{\lambda_{\psi}}{2}$, then $\frac{\psi(N)bk}{N-bk} \leq A$ implies that $k \leq \frac{AN}{b(\psi(N)+A)} \leq \frac{2A}{b\lambda_{\psi}}$. Hence for every $t \in [0, A]$, we have

$$|\theta_{\infty}(t) - \theta_N(t)| \leq \frac{1}{t^2} \sum_{k=1}^{\frac{2A}{b\lambda_{\psi}}} k \left| \mathbb{1}_{[b\lambda_{\psi}k, +\infty[}(t) - \mathbb{1}_{[\frac{\psi(N)bk}{N-bk}, +\infty[}(t) \right|.$$

For every continuous function $f : [0, +\infty[\rightarrow \mathbb{R}$ with compact support in $[0, A]$, we therefore have

$$\begin{aligned} \left| \int_0^{+\infty} f(\theta_{\infty} - \theta_N) dt \right| &= O\left(\|f\|_{\infty} \sum_{k=1}^{\frac{2A}{b\lambda_{\psi}}} k \left| b\lambda_{\psi}k - \frac{\psi(N)bk}{N-bk} \right| \right) \\ &= O\left(A^3 \|f\|_{\infty} \left| \lambda_{\psi} - \frac{\psi(N)}{N} \right| + O\left(\frac{A}{N}\right) \right). \end{aligned}$$

By Equation (5), for every C^1 function $f : [0, +\infty[\rightarrow \mathbb{R}$ with compact support in $[0, A]$, we thus have

$$\begin{aligned} \frac{1}{\psi(N)} \mu_N^+(f) &= \int f \theta_{\infty} d\text{Leb}_{[0, +\infty[} \\ &\quad + O\left(A^3 \|f\|_{\infty} \left(\left| \lambda_{\psi} - \frac{\psi(N)}{N} \right| + O\left(\frac{A}{N}\right) \right) \right) + O\left(\|f'\|_{\infty} \frac{A^2}{N}\right). \end{aligned}$$

With $g_{\mathcal{L}_N, \psi} : \mathbb{R} \rightarrow \mathbb{R}$ given by $t \mapsto \theta_\infty(|t|)$, by Lemma 3.2, it follows that

$$\frac{1}{\psi(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi} \xrightarrow{*} g_{\mathcal{L}_N, \psi} \text{Leb}_{\mathbb{R}} .$$


Futhermore, for every $A \geq 1$ and every C^1 function $f : \mathbb{R} \rightarrow \mathbb{R}$ with compact support contained in $[-A, A]$, we have

$$\frac{1}{\psi(N)} \mathcal{R}_N^{\mathcal{L}_N, \psi}(f) = \int_{\mathbb{R}} f(t) g_{\mathcal{L}_N, \psi}(t) dt \quad (9)$$

$$+ O\left(A^3 \|f\|_\infty \left(\left| \lambda_\psi - \frac{\psi(N)}{N} \right| + \frac{A}{N} \right)\right) + O\left(\|f'\|_\infty \frac{A^3}{N}\right) . \quad (10)$$

This concludes the proof of Theorem 3.1. $\square$

Let us give a numerical illustration of Theorem 3.1 when $a = b = 1$ and $\psi(N) = N$. The following figure shows in red an approximation of the pair correlation function $g_{\mathcal{L}_N, \psi}$ computed using $\mathcal{R}_{2000}^{\mathcal{L}_N, \psi}$, and in blue the pair correlation function $g_{\mathcal{L}_N, \psi}$ in the interval $[-4, 4]$.



4 Pair correlations with Euler weights without scaling

In this section, we study the weighted family

$$\mathcal{L}_N^{a,b,\varphi} = \left((L_N^{a,b})_{N \in \mathbb{N}}, \omega = \varphi \circ \exp \right) .$$

The (not normalised) *pair correlation measure* of the logarithms of integers congruent to a modulo b at time N with multiplicities given by the Euler function φ , for the trivial scaling function, is

$$\tilde{\nu}_N = \sum_{(m,n) \in I_N} \varphi(n) \varphi(m) \Delta_{\ln m - \ln n} .$$

With the notation of the introduction, we have $\mathcal{L}_N^{1,1,\varphi} = \mathcal{L}_N^\varphi$ and $\tilde{\nu}_N = \mathcal{R}_N^{\mathcal{L}_N^{a,b,\varphi}, 1}$.

Theorem 4.1 *As $N \rightarrow +\infty$, the measures $\tilde{\nu}_N$ on $\mathbb{R}$, renormalized to be probability measures, weak-star converge to the measure absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$, with Radon-Nikodym derivative the function $g_{\mathcal{L}_N^{\varphi,a,b},1} : s \mapsto e^{-2|s|}$:*

$$\frac{\tilde{\nu}_N}{\|\tilde{\nu}_N\|} \xrightarrow{*} g_{\mathcal{L}_N^{\varphi,a,b},1} \text{Leb}_{\mathbb{R}} .$$

Furthermore, for all $f \in C_c^1(\mathbb{R})$ and $\alpha \in [\frac{1}{2}, 1[$, we have

$$\frac{\tilde{\nu}_N}{\|\tilde{\nu}_N\|}(f) = \int_{s \in \mathbb{R}} f(s) e^{-2|s|} ds + O_b \left(\frac{\ln N}{N^{1-\alpha}} \|f\|_{\infty} + \frac{1}{N^{\alpha}} \|e^{|s|} f'(s)\|_{\infty} \right) .$$

When $a = b = 1$, the measure $\tilde{\nu}_N$ corresponds to the one denoted by $\mathcal{R}_N^{\varphi,1}$ in the introduction. The above result gives the first assertion of Theorem 1.2 in the introduction, with pair correlation function $g_{\mathcal{L}_N^{\varphi},1} = g_{\mathcal{L}_N^{\varphi,1,1},1}$.

Proof. The first assertion of Theorem 4.1 follows from the second one, by taking for instance $\alpha = \frac{1}{2}$ and by the density of C^1 -smooth functions with compact support in the space of continuous functions with compact support on $\mathbb{R}$.

For every $q \in \mathbb{N} - \{0\}$ with $q \equiv a \pmod{b}$, let $q' \in \mathbb{N}$ be such that $q = a + q'b$ and let J_q be given by Equation (2). We now define

$$\tilde{\omega}_q = \sum_{p \in J_q} \varphi(p) \Delta_{\frac{p}{q}} ,$$

which is a finitely supported measure on $[0, 1]$, and nonzero if and only if $q' \neq 0$. In order to compute its total mass, we will use the following elementary adaptation of Mertens' formula (see for example [HaW, Thm. 330]). We have not found its proof in the literature, hence we provide one, due to Fouvry.

Let $(a, b) \in \mathbb{N} - \{0\}$ be the greatest common divisor of a and b . Let

$$c_{a,b} = \frac{\varphi((a, b))}{b(a, b)} \prod_{p \text{ prime}, p|b} \left(1 - \frac{1}{p^2}\right)^{-1} ,$$

and note that $c_{a,b} > 0$ is uniformly bounded from above when a and b vary in $\mathbb{N} - \{0\}$, and tends to 0 as $b \rightarrow +\infty$. When $a = b = 1$, we have $c_{a,b} = 1$, and the following result is exactly Mertens' formula.

Lemma 4.2 *There exists $C > 0$ such that for all integers $a, b \geq 1$ and real numbers $x \geq 1$, we have*

$$\left| \sum_{1 \leq n \leq x, n \equiv a \pmod{b}} \varphi(n) - \frac{3c_{a,b}}{\pi^2} x^2 \right| \leq C x \ln(2x) .$$

Proof. (Fouvry) In this proof, for every function g of a variable in $[1, +\infty[$, possibly depending on parameters, we use the notation $O(g)$ in order to denote any function f on $[1, +\infty[$ such that there exists a constant C , independent of the variable and of all the parameters, such that $|f| \leq C|g|$. We do not need a more precise error term.

Let $S(x, a, b)$ be the above sum. We refer for instance to [HaW] for the definition of the Möbius function $\mu : \mathbb{N} - \{0\} \rightarrow \{-1, 0, 1\}$, of the Dirichlet convolution $f * g$ of two

maps $f, g : \mathbb{N} - \{0\} \rightarrow \mathbb{R}$ and for the Möbius inversion formula, which in particular gives that $\varphi = \mu * \text{id}$. Hence

$$S(x, a, b) = \sum_{\substack{1 \leq n \leq x \\ n \equiv a \pmod{b}}} \sum_{md=n} \mu(d) m = \sum_{1 \leq d \leq x} \mu(d) \sum_{\substack{1 \leq m \leq x/d \\ md \equiv a \pmod{b}}} m.$$

Let us fix $d \geq 1$. Let us consider the congruence equation $md \equiv a \pmod{b}$ with unknown m . It has no solution if the greatest common divisor (b, d) of b and d does not divide a . If (b, d) does divide a , let $a' = \frac{a}{(b, d)}$, $b' = \frac{b}{(b, d)}$ and $d' = \frac{d}{(b, d)}$, so that the congruence equation is equivalent to $md' \equiv a' \pmod{b'}$. Since d' is coprime with b' , it is invertible modulo b' , and we denote its inverse by $\overline{d'}$. The congruence equation becomes $m \equiv a' \overline{d'} \pmod{b'}$. The classical formula $\sum_{1 \leq m \leq y, m \equiv a' \overline{d'} \pmod{b'}} 1 = \frac{y}{b'} + O(1)$ gives, by a summation by parts, the equality

$$\sum_{1 \leq m \leq y, m \equiv a' \overline{d'} \pmod{b'}} m = \frac{y^2}{2b'} + O(y).$$

Therefore

$$\begin{aligned} S(x, a, b) &= \sum_{1 \leq d \leq x, (b, d) | a} \mu(d) \left(\frac{(b, d)}{2b} \left(\frac{x}{d} \right)^2 + O\left(\frac{x}{d} \right) \right) \\ &= \frac{x^2}{2b} \left(\sum_{1 \leq d \leq x, (b, d) | a} \mu(d) \frac{(b, d)}{d^2} \right) + O(x \ln(2x)). \end{aligned}$$

Using the Eulerian product formula of the zeta function, giving $\prod_{p \text{ prime}} (1 - \frac{1}{p^2}) = \frac{1}{\zeta(2)} = \frac{6}{\pi^2}$,

and the expression $\varphi(n) = n \prod_{p \text{ prime}, p | n} (1 - \frac{1}{p})$ of the Euler function in terms of the prime factors, we have, by decomposing d into prime powers and using the definition of the Möbius function,

$$\begin{aligned} \sum_{1 \leq d \leq x, (b, d) | a} \mu(d) \frac{(b, d)}{d^2} &= \sum_{d \geq 1, (b, d) | a} \mu(d) \frac{(b, d)}{d^2} + O\left(\frac{b}{x} \right) \\ &= \prod_{p \text{ prime}, p \nmid b} \left(1 - \frac{1}{p^2} \right) \prod_{p \text{ prime}, p | a, p | b} \left(1 - \frac{1}{p} \right) + O\left(\frac{b}{x} \right) \\ &= \frac{1}{\zeta(2)} \prod_{p \text{ prime}, p | b} \left(1 - \frac{1}{p^2} \right)^{-1} \frac{\varphi((a, b))}{(a, b)} + O\left(\frac{b}{x} \right) = \frac{6}{\pi^2} \frac{c_{a,b}}{c_a} + O\left(\frac{b}{x} \right). \end{aligned}$$

This proves Lemma 4.2. □

Lemma 4.2 says that if $q' \neq 0$ (that is, when $q > a$), then

$$\|\tilde{\omega}_q\| = \frac{3 c_{a,b} q^2}{\pi^2} + O(q \ln(2q)), \quad (11)$$

and in particular $\frac{1}{\|\tilde{\omega}_q\|} = \frac{\pi^2}{3 c_{a,b} q^2} (1 + O(\frac{\ln q}{q}))$.

Lemma 4.3 *We have as $q \rightarrow +\infty$,*

$$\frac{\tilde{\omega}_q}{\|\tilde{\omega}_q\|} \xrightarrow{*} 2t \, d\text{Leb}_{[0,1]}(t) .$$

More precisely, for all $f \in C^1([0, 1])$ and $\alpha \in [\frac{1}{2}, 1[$, we have

$$\frac{\tilde{\omega}_q}{\|\tilde{\omega}_q\|}(f) = \int_0^1 2t f(t) \, dt + O\left(\frac{\ln q}{q^{1-\alpha}} \|f\|_\infty + \frac{1}{q^\alpha} \|f'\|_\infty\right) .$$

Proof. The first assertion follows from the second one, by taking for instance $\alpha = \frac{1}{2}$ and by the density of C^1 -smooth functions in the space of continuous functions on $[0, 1]$.

Let $Q = \lfloor q^\alpha \rfloor \in \mathbb{N} - \{0\}$. For all $n \in \{0, \dots, Q-1\}$ and $t \in]nq^{-\alpha}, (n+1)q^{-\alpha}]$, we have by the mean value theorem

$$f(t) = f(nq^{-\alpha}) + O(q^{-\alpha} \|f'\|_\infty) .$$

Since $n \leq Q \leq q^\alpha$, we hence have

$$\begin{aligned} \int_{nq^{-\alpha}}^{(n+1)q^{-\alpha}} 2t f(t) \, dt &= (f(nq^{-\alpha}) + O(q^{-\alpha} \|f'\|_\infty)) \int_{nq^{-\alpha}}^{(n+1)q^{-\alpha}} 2t \, dt \\ &= \frac{1}{q^\alpha} ((2n+1)q^{-\alpha} f(nq^{-\alpha}) + O(nq^{-2\alpha} \|f'\|_\infty)) . \end{aligned} \quad (12)$$

Using twice Equation (11) and the formula for $\frac{1}{\|\tilde{\omega}_q\|}$ following it, we have

$$\begin{aligned} &\sum_{p \in]nq^{1-\alpha}, (n+1)q^{1-\alpha}] \cap J_q} f\left(\frac{p}{q}\right) \frac{1}{\|\tilde{\omega}_q\|} \varphi(p) \\ &= (f(nq^{-\alpha}) + O(q^{-\alpha} \|f'\|_\infty)) \frac{\pi^2}{3c_{a,b}q^2} (1 + O(\frac{\ln q}{q})) \\ &\quad \times \left(\frac{3c_{a,b}((n+1)^2 - n^2)}{\pi^2} q^{2-2\alpha} + O((n+1)q^{1-\alpha} \ln((n+1)q^{1-\alpha})) \right) \\ &= \frac{1}{q^\alpha} ((2n+1)q^{-\alpha} f(nq^{-\alpha}) + O(\frac{n}{q^{2\alpha}} \|f'\|_\infty + \frac{n \ln q}{q} \|f\|_\infty)) . \end{aligned} \quad (13)$$

Again using Equation (11), we have

$$\sum_{p \in]Qq^{1-\alpha}, q[\cap J_q} \frac{\varphi(p)}{\|\tilde{\omega}_q\|} |f(\frac{p}{q})| = O\left(\frac{q^2 - (Qq^{1-\alpha})^2}{q^2} \|f\|_\infty\right) = O(q^{-\alpha} \|f\|_\infty) . \quad (14)$$

By cutting the sum defining $\tilde{\omega}_q$ and the integral from 0 to 1, by using (12), (13) and (14),

since $Q \leq q^\alpha$ and $|1 - Q q^{-\alpha}| = O(q^{-\alpha})$, we have (using $\alpha \geq \frac{1}{2}$ for the last equality)

$$\begin{aligned}
& \left| \frac{\tilde{\omega}_q}{\|\tilde{\omega}_q\|}(f) - \int_0^1 2t f(t) dt \right| \\
&= \left| \sum_{n=0}^{Q-1} \left(\sum_{p \in]n q^{1-\alpha}, (n+1)q^{1-\alpha}] \cap J_q} \frac{\varphi(p)}{\|\tilde{\omega}_q\|} f\left(\frac{p}{q}\right) - \int_{n q^{-\alpha}}^{(n+1)q^{-\alpha}} 2t f(t) dt \right) \right| \\
&\quad + \sum_{p \in]Q q^{1-\alpha}, q[\cap J_q} \frac{\varphi(p)}{\|\tilde{\omega}_q\|} \left| f\left(\frac{p}{q}\right) \right| + \int_{Q q^{-\alpha}}^1 2t |f(t)| dt \\
&= O(q^{-\alpha} \|f\|_\infty) + \frac{1}{q^\alpha} \sum_{n=0}^{Q-1} O\left(\frac{n}{q^{2\alpha}} \|f'\|_\infty + \frac{n \ln q}{q} \|f\|_\infty + n q^{-2\alpha} \|f'\|_\infty\right) \\
&= O(q^{-\alpha} \|f\|_\infty) + \frac{1}{q^\alpha} O\left(\frac{Q^2}{q^{2\alpha}} \|f'\|_\infty + \frac{Q^2 \ln q}{q} \|f\|_\infty\right) \\
&= O\left(\frac{\ln q}{q^{1-\alpha}} \|f\|_\infty + \frac{\|f'\|_\infty}{q^\alpha}\right).
\end{aligned}$$

This proves Lemma 4.3. □

For every $N \in \mathbb{N} - \{0\}$, let us define

$$\tilde{\mu}_N^- = \sum_{(m,n) \in I_N^-} \varphi(m) \varphi(n) \Delta_{\frac{m}{n}} = \sum_{1 \leq q \leq N, q \equiv a \pmod b} \varphi(q) \tilde{\omega}_q,$$

which is a finitely supported measure on $[0, 1]$. By Lemma 4.2, its total mass is

$$\begin{aligned}
\|\tilde{\mu}_N^-\| &= \sum_{1 \leq q \leq N, q \equiv a \pmod b} \varphi(q) \|\tilde{\omega}_q\| = \sum_{(m,n) \in I_N^-} \varphi(m) \varphi(n) \\
&= \frac{1}{2} \left(\left(\sum_{\substack{1 \leq q \leq N \\ q \equiv a \pmod b}} \varphi(q) \right)^2 - \sum_{\substack{1 \leq q \leq N \\ q \equiv a \pmod b}} \varphi(q)^2 \right) = \frac{9 c_{a,b}^2 N^4}{2 \pi^4} + O(N^3 \ln N).
\end{aligned}$$

For $f \in C^1([0, 1])$, by Lemma 4.3, since $q^{1+\alpha}(\ln q) \leq N^{1+\alpha}(\ln N)$ and $q^{2-\alpha} \leq N^{2-\alpha}$ when q occurs in the summations below, we have

$$\begin{aligned}
\frac{\tilde{\mu}_N^-(f)}{\|\tilde{\mu}_N^-\|} &= \frac{1}{\|\tilde{\mu}_N^-\|} \sum_{1 \leq q \leq N, q \equiv a \pmod b} \varphi(q) \|\tilde{\omega}_q\| \frac{\tilde{\omega}_q(f)}{\|\tilde{\omega}_q\|} \\
&= \int_0^1 2t f(t) dt + \frac{1}{\|\tilde{\mu}_N^-\|} \sum_{\substack{1 \leq q \leq N \\ q \equiv a \pmod b}} \varphi(q) \|\tilde{\omega}_q\| O\left(\frac{\ln q}{q^{1-\alpha}} \|f\|_\infty + \frac{1}{q^\alpha} \|f'\|_\infty\right) \\
&= \int_0^1 2t f(t) dt + O\left(\frac{1}{N^4} \sum_{\substack{1 \leq q \leq N \\ q \equiv a \pmod b}} \varphi(q) (q^{1+\alpha}(\ln q) \|f\|_\infty + q^{2-\alpha} \|f'\|_\infty)\right) \\
&= \int_0^1 2t f(t) dt + O\left(\frac{\ln N}{N^{1-\alpha}} \|f\|_\infty + \frac{1}{N^\alpha} \|f'\|_\infty\right).
\end{aligned}$$

For every $N \in \mathbb{N} - \{0\}$, let us define

$$\tilde{\nu}_N^\pm = \sum_{(m,n) \in I_N^\pm} \varphi(m) \varphi(n) \Delta_{\ln \frac{m}{n}},$$

so that $\tilde{\nu}_N^- = \ln_* \tilde{\mu}_N^- = \tilde{\nu}_N|_{]-\infty, 0]}$, and $\|\tilde{\nu}_N^-\| = \|\tilde{\mu}_N^-\|$. We have, for every $f \in C_c^1(]-\infty, 0])$,

$$\begin{aligned} \frac{\tilde{\nu}_N^-(f)}{\|\tilde{\nu}_N^-\|} &= \frac{\tilde{\mu}_N^-(f \circ \ln)}{\|\tilde{\mu}_N^-\|} = \int_0^1 2t f(\ln t) dt + O\left(\frac{\ln N}{N^{1-\alpha}} \|f \circ \ln\|_\infty + \frac{1}{N^\alpha} \|(f \circ \ln)'\|_\infty\right) \\ &= \int_{-\infty}^0 2f(s) e^{2s} ds + O\left(\frac{\ln N}{N^{1-\alpha}} \|f\|_\infty + \frac{1}{N^\alpha} \|e^{-s} f'(s)\|_\infty\right). \end{aligned}$$

Since $\tilde{\nu}_N = \tilde{\nu}_N^- + \tilde{\nu}_N^+$, since $\tilde{\nu}_N^+ = \text{sg}_* \tilde{\nu}_N^-$, we have $\|\tilde{\nu}_N^\pm\| = \frac{1}{2} \|\tilde{\nu}_N\|$ and Theorem 4.1 follows. $\square$

Let us give some numerical illustrations of Theorem 4.1 with $a = b = 1$. For every $N \in \mathbb{N} - \{0\}$, let

$$\tilde{\mathcal{D}}_N : s \mapsto \frac{\sum_{0 \leq m \neq n \leq N : (\ln m - \ln n) \leq s} \varphi(m) \varphi(n)}{\sum_{0 \leq m \neq n \leq N} \varphi(m) \varphi(n)}.$$


This is the cumulative distribution function at time N of the unscaled differences of the logarithms of natural numbers weighted by the Euler function, that is, for all $s, s' \in \mathbb{R}$ with $s < s'$, we have

$$\frac{\mathcal{R}_N^{\mathcal{L}_N^\varphi, 1}}{\|\mathcal{R}_N^{\mathcal{L}_N^\varphi, 1}\|} (]s, s']) = \tilde{\mathcal{D}}_N(s') - \tilde{\mathcal{D}}_N(s).$$

Theorem 4.1 with $a = b = 1$ says that the function $\tilde{\mathcal{D}}_N$ pointwise converges as $N \rightarrow +\infty$ to the C^1 (but not C^2) function

$$\tilde{\mathcal{D}} : s \mapsto \begin{cases} \frac{1}{2} e^{2s} & \text{if } s \leq 0 \\ 1 - \frac{1}{2} e^{-2s} & \text{if } s \geq 0 \end{cases}$$

(with derivative $\tilde{\mathcal{D}}' = g_{\mathcal{L}_N^\varphi, 1}$). This is illustrated by the figure below, which shows $\tilde{\mathcal{D}}_{15}$ in red.



5 Pair correlations with Euler weights and linear scaling

In this section, we study the pair correlations of the family $\mathcal{L}_N^{\varphi, a, b}$ defined at the beginning of Section 4, now with a linear scaling. We leave to the reader the study of a general

scaling ψ , assumed to converge to $+\infty$. For every $N \in \mathbb{N} - \{0\}$, the (not normalised) *pair correlation measure* of the logarithms of integers, congruent to a modulo b , at time N with multiplicities given by the Euler function and with scaling N is the (Borel, positive) measure with finite support in $\mathbb{R}$ defined by

$$\tilde{\mathcal{R}}_N = \sum_{(m,n) \in I_N} \varphi(m) \varphi(n) \Delta_{N(\ln m - \ln n)}.$$

With the notation of the introduction, we have $\tilde{\mathcal{R}}_N = \mathcal{R}_N^{\mathcal{L}_N^{\varphi,a,b}, \text{id}}$.

For every $k \in \mathbb{N} - \{0\}$, we consider the arithmetic constant $c_{a,b,k}$ defined in Equation (21) of Appendix A. Note that $c_{a,b,k} > 0$ is uniformly bounded from above when a, b, k vary in $\mathbb{N} - \{0\}$, and has a positive lower bound on a and k when b is fixed, by Equation (22) of Appendix A.

Theorem 5.1 *As $N \rightarrow +\infty$, the family $(\frac{1}{N^3} \tilde{\mathcal{R}}_N)_{N \in \mathbb{N}}$ of measures on $\mathbb{R}$ weak-star converges to the measure absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$, with Radon-Nikodym derivative the function*

$$g_{\mathcal{L}_N^{\varphi,a,b}, \text{id}} : s \mapsto \frac{1}{s^4} \sum_{1 \leq k \leq |s|, k \equiv 0 \pmod b} c_{a,b,k} k^3,$$

that is, as $N \rightarrow +\infty$,

$$\frac{1}{N^3} \tilde{\mathcal{R}}_N \xrightarrow{*} g_{\mathcal{L}_N^{\varphi,a,b}, \text{id}} \text{Leb}_{\mathbb{R}}.$$

Furthermore, for all $f \in C^1(\mathbb{R})$ with compact support contained in $[-A, A]$, where $A \geq 1$, and for any $\alpha \in [\frac{1}{2}, 1[$, we have

$$\frac{1}{N^3} \tilde{\mathcal{R}}_N(f) = \int_{s \in \mathbb{R}} f(s) g_{\mathcal{L}_N^{\varphi,a,b}, \text{id}}(s) ds + O_b \left(\frac{A^7 \ln^2 N}{N^{1-\alpha}} \|f\|_{\infty} + \frac{A^3}{N^{\alpha}} \|f'\|_{\infty} \right).$$

When $a = b = 1$, the measure $\tilde{\mathcal{R}}_N$ corresponds to the one denoted by $\mathcal{R}_N^{\mathcal{L}_N^{\varphi}, \text{id}}$ in the introduction. The above result gives the second assertion of Theorem 1.2 in the introduction, with pair correlation function $g_{\mathcal{L}_N^{\varphi}, \text{id}} = g_{\mathcal{L}_N^{\varphi,1,1}, \text{id}}$, using Mirsky's value of $c_{1,1,k}$ given by Equation (23), as explained in Remark A.6 of Appendix A.

Note that, as the proof below shows, the total mass of $\tilde{\mathcal{R}}_N$ is equivalent to cN^4 as $N \rightarrow +\infty$, for some constant $c > 0$, hence renormalising $\tilde{\mathcal{R}}_N$ to be a probability measure makes it weak-star converge to the zero measure on the noncompact space $\mathbb{R}$ (a total loss of mass phenomenon).

Proof. The first assertion of Theorem 5.1 follows from the second one, by taking for instance $\alpha = \frac{1}{2}$ and by the density of C^1 -smooth functions with compact support in the space of continuous functions with compact support on $\mathbb{R}$.

The change of variables $(m, n) \mapsto (n, m)$ in I_N gives $\tilde{\mathcal{R}}_N|_{]-\infty, 0]} = \text{sg}_* (\tilde{\mathcal{R}}_N|_{[0, +\infty[})$. We will thus only study the convergence of the measures $\frac{1}{N^3} \tilde{\mathcal{R}}_N$ on $[0, +\infty[$, and deduce the global result by the symmetry of $g_{\mathcal{L}_N^{\varphi,a,b}, \text{id}}$.

For every $N \in \mathbb{N} - \{0\}$ and for every $p \in \mathbb{N}$ with $p \equiv 0 \pmod b$ and $0 < p < N$, let $J_{p,N}$ be given par Equation (3). We now define the key auxiliary measure by

$$\tilde{\omega}_{p,N} = \sum_{q \in J_{p,N}} \varphi(q) \varphi(q+p) \Delta_{\frac{q}{N^p}}.$$

Then $\tilde{\omega}_{p,N}$ is a measure on $[0, +\infty[$, with finite support contained in $[\frac{1}{Np}, \frac{N-p}{Np}]$, hence in $[0, 1]$. The measure $\tilde{\omega}_{p,N}$ is nonzero if and only if $N \geq a + p$. In order to compute its total mass, we use an adaptation with congruences of a formula by Mirsky (see [Mir, Thm. 9]) proved in the appendix by Fouvry. Theorem A.1 says that if $N \geq a + p$, then

$$\|\tilde{\omega}_{p,N}\| = \frac{c_{a,b,p}}{3} (N-p)^3 + O(N^2(p + \ln(N-p))^2), \quad (15)$$

and in particular $\frac{1}{\|\tilde{\omega}_{p,N}\|} = \frac{3}{c_{a,b,p}(N-p)^3} (1 + O(\frac{N^2(p + \ln(N-p))^2}{(N-p)^3}))$.

The next result implies that the measures $\tilde{\omega}_{p,N}$, once normalized to be probability measures, weak-star converge to the measure $d\mu(t) = 3(\frac{Np}{N-p})^3 t^2 d\text{Leb}_{[\frac{1}{Np}, \frac{N-p}{Np}]}$, which is absolutely continuous with respect to the Lebesgue measure on the interval $[\frac{1}{Np}, \frac{N-p}{Np}]$.

Lemma 5.2 *For every $p \in \mathbb{N}$ with $0 < p < N$ and $p \equiv 0 \pmod{b}$, for every $\alpha \in]0, 1[$ and for every $f \in C_c^1([0, 1])$, we have*

$$\begin{aligned} \frac{\tilde{\omega}_{p,N}}{\|\tilde{\omega}_{p,N}\|}(f) &= \int_{\frac{1}{Np}}^{\frac{N-p}{Np}} 3\left(\frac{Np}{N-p}\right)^3 t^2 f(t) dt \\ &+ O\left(\left(\frac{(Np)^3}{(N-p)^3(Np)^\alpha} + \frac{(Np)^6(p + \ln(Np))^2}{(N-p)^6(Np)^{1-\alpha}}\right) \|f\|_\infty + \frac{1}{(Np)^\alpha} \|f'\|_\infty\right). \end{aligned}$$

Proof. As in the proof of Lemma 4.3, we will estimate the difference of the main terms in the above centered formula by cutting the sum defining the renormalized measure $\tilde{\omega}_{p,N}$ and by cutting similarly the integral from $\frac{1}{Np}$ to $\frac{N-p}{Np}$.

Let $Q = \lfloor (Np)^\alpha \frac{N-p}{Np} \rfloor \in \mathbb{N}$. For all $n \in \{0, \dots, Q-1\}$, we thus define

$$S_n = \sum_{q \in]n(Np)^{1-\alpha}, (n+1)(Np)^{1-\alpha}] \cap J_{p,N}} f\left(\frac{q}{Np}\right) \frac{1}{\|\tilde{\omega}_{p,N}\|} \varphi(q) \varphi(q+p)$$

and

$$I_n = \int_{n(Np)^{-\alpha}}^{(n+1)(Np)^{-\alpha}} 3\left(\frac{Np}{N-p}\right)^3 t^2 f(t) dt.$$

Let us also define the following remaining terms

$$S_{\text{end}} = \sum_{q \in]Q(Np)^{1-\alpha}, N-p] \cap J_{p,N}} f\left(\frac{q}{Np}\right) \frac{1}{\|\tilde{\omega}_{p,N}\|} \varphi(q) \varphi(q+p)$$

and

$$I_{\text{end}} = \int_{Q(Np)^{-\alpha}}^{\frac{N-p}{Np}} 3\left(\frac{Np}{N-p}\right)^3 t^2 f(t) dt.$$

For all $t \in]n(Np)^{-\alpha}, (n+1)(Np)^{-\alpha}]$, we have by the mean value theorem

$$f(t) = f(n(Np)^{-\alpha}) + O((Np)^{-\alpha} \|f'\|_\infty).$$

Using twice Theorem A.1 and the formula for $\frac{1}{\|\tilde{\omega}_{p,N}\|}$ following Equation (15), and using the inequality $(n+1) \leq Q \leq (Np)^\alpha \frac{N-p}{Np}$, we have

$$\begin{aligned}
S_n &= \left(f(n(Np)^{-\alpha}) + O((Np)^{-\alpha} \|f'\|_\infty) \right) \left(\frac{1}{(N-p)^3} \left(1 + O\left(\frac{N^2(p + \ln(N-p))^2}{(N-p)^3} \right) \right) \right) \\
&\quad \times \left(((n+1)(Np)^{1-\alpha})^3 - (n(Np)^{1-\alpha})^3 \right. \\
&\quad \left. + O\left((p + (n+1)(Np)^{1-\alpha})^2 (p + \ln((n+1)(Np)^{1-\alpha}))^2 \right) \right) \\
&= \frac{(Np)^3}{(N-p)^3(Np)^\alpha} \left((3n^2 + 3n + 1)(Np)^{-2\alpha} f(n(Np)^{-\alpha}) + O\left(\frac{n^2}{(Np)^{3\alpha}} \|f'\|_\infty \right) \right. \\
&\quad \left. + O\left(\left(\frac{n^2 N^2 (p + \ln(N-p))^2}{(N-p)^3 (Np)^{2\alpha}} + \frac{(n + \frac{p}{(Np)^{1-\alpha}})^2 (p + \ln(Np))^2}{(Np)^{1+\alpha}} \right) \|f\|_\infty \right) \right). \quad (16)
\end{aligned}$$

We also have

$$\begin{aligned}
I_n &= \left(f(n(Np)^{-\alpha}) + O((Np)^{-\alpha} \|f'\|_\infty) \right) \int_{n(Np)^{-\alpha}}^{(n+1)(Np)^{-\alpha}} 3 \left(\frac{Np}{N-p} \right)^3 t^2 dt \\
&= \frac{(Np)^3}{(N-p)^3(Np)^\alpha} \left((3n^2 + 3n + 1)(Np)^{-2\alpha} f(n(Np)^{-\alpha}) + O\left(\frac{n^2}{(Np)^{3\alpha}} \|f'\|_\infty \right) \right). \quad (17)
\end{aligned}$$

Similarly, since $Q \geq (Np)^\alpha \frac{N-p}{Np} - 1$ and $N-p \leq Np$, we have

$$S_{\text{end}} = O\left(\frac{(N-p)^3 - (Q(Np)^{1-\alpha})^3}{(N-p)^3} \|f\|_\infty \right) = O\left(\frac{(Np)^3}{(N-p)^3(Np)^\alpha} \|f\|_\infty \right) \quad (18)$$

and

$$I_{\text{end}} = O\left(\left(\frac{Np}{N-p} \right)^3 \left(\left(\frac{N-p}{Np} \right)^3 - (Q(Np)^{-\alpha})^3 \right) \|f\|_\infty \right) = O\left(\frac{(Np)^3}{(N-p)^3(Np)^\alpha} \|f\|_\infty \right). \quad (19)$$

Note that $\sum_{n=0}^{Q-1} \left(n + \frac{p}{(Np)^{1-\alpha}} \right)^2 = O\left(\left(Q + \frac{p}{(Np)^{1-\alpha}} \right)^3 \right) = O\left(\frac{(Np)^{3\alpha}}{p^3} \right)$. Putting together Equations (16), (17), (18) and (19), we have (again using $N-p \leq Np$)

$$\begin{aligned}
&\left| \frac{\tilde{\omega}_{p,N}}{\|\tilde{\omega}_{p,N}\|}(f) - \int_{\frac{1}{Np}}^{\frac{N-p}{Np}} 3 \left(\frac{Np}{N-p} \right)^3 t^2 f(t) dt \right| = \left| \sum_{n=0}^{Q-1} (S_n - I_n) + S_{\text{end}} - I_{\text{end}} \right| \\
&\leq \sum_{n=0}^{Q-1} |S_n - I_n| + |S_{\text{end}}| + |I_{\text{end}}| \\
&= O\left(\frac{(Np)^3}{(N-p)^3(Np)^\alpha} \|f\|_\infty \right) + \frac{(Np)^3}{(N-p)^3(Np)^\alpha} \sum_{n=0}^{Q-1} O\left(\frac{n^2}{(Np)^{3\alpha}} \|f'\|_\infty + \right. \\
&\quad \left. \left(\frac{n^2 N^2 (p + \ln(N-p))^2}{(N-p)^3 (Np)^{2\alpha}} + \frac{(n + \frac{p}{(Np)^{1-\alpha}})^2 (p + \ln(Np))^2}{(Np)^{1+\alpha}} \right) \|f\|_\infty + \frac{n^2}{(Np)^{3\alpha}} \|f'\|_\infty \right) \\
&= O\left(\frac{(Np)^3}{(N-p)^3(Np)^\alpha} \|f\|_\infty + \frac{1}{(Np)^\alpha} \|f'\|_\infty + \frac{(Np)^6 (p + \ln(Np))^2}{(N-p)^6 (Np)^{1-\alpha}} \|f\|_\infty \right).
\end{aligned}$$

This proves Lemma 5.2. □

Now, let us introduce the sum

$$\tilde{\mu}_N^+ = \sum_{0 < p < N, p \equiv 0 \pmod{b}} \iota_* \tilde{\omega}_{p,N} = \sum_{\substack{1 \leq q \leq N-p, 0 < p < N \\ q \equiv a \pmod{b}, p \equiv 0 \pmod{b}}} \varphi(q) \varphi(q+p) \Delta_{N \frac{p}{q}},$$

where as previously $\iota : t \mapsto \frac{1}{t}$ (noting that the measures $\tilde{\omega}_{p,N}$ are supported in $]0, +\infty[$).

Lemma 5.3 *For every $f \in C^1([0, +\infty[)$ with compact support contained in $[0, A]$, where $A \geq 1$, we have*

$$|\tilde{\mathcal{R}}_N(f) - \tilde{\mu}_N(f)| = O(A^3 N^2 \|f'\|_\infty).$$

Proof. Using the change of variables $(p, q) \mapsto (m = p + q, n = q)$, we have

$$\tilde{\mathcal{R}}_N|_{[0, +\infty[} = \sum_{(m,n) \in I_N^+} \varphi(m) \varphi(n) \Delta_{N \ln \frac{m}{n}} = \sum_{\substack{0 < p < N, 1 \leq q \leq N-p \\ p \equiv 0 \pmod{b}, q \equiv a \pmod{b}}} \varphi(q) \varphi(q+p) \Delta_{N \ln(1 + \frac{p}{q})}.$$

As in the proof of Lemma 3.2, since the support of f is contained in $[0, A]$, if a pair (p, q) occurs in the index of the sum defining either $\tilde{\mathcal{R}}_N(f)$ or $\tilde{\mu}_N^+(f)$ with nonzero corresponding summand, then $\frac{p}{q} = O(\frac{A}{N})$ and $p = O(A)$. By the mean value theorem, we then have

$$\begin{aligned} |f(N \frac{p}{q}) - f(N \ln(1 + \frac{p}{q}))| &\leq \|f'\|_\infty |N \frac{p}{q} - N \ln(1 + \frac{p}{q})| \\ &= \|f'\|_\infty N O((\frac{p}{q})^2) = O(\frac{A^2}{N} \|f'\|_\infty). \end{aligned}$$

Thus, using Theorem A.1 and Equation (22) in Appendix A, we have

$$\begin{aligned} |\tilde{\mathcal{R}}_N(f) - \tilde{\mu}_N^+(f)| &\leq \sum_{1 \leq p \leq O(A), 1 \leq q \leq N} \varphi(q) \varphi(q+p) O(\frac{A^2}{N} \|f'\|_\infty) \\ &\leq \sum_{1 \leq p \leq O(A)} O(N^3) O(\frac{A^2}{N} \|f'\|_\infty) = O(A^3 N^2 \|f'\|_\infty). \end{aligned}$$

This proves Lemma 5.3. □

Lemma 5.4 *For all $\alpha \in [\frac{1}{2}, 1[$ and $f \in C^1([0, +\infty[)$ with compact support contained in $[0, A]$, where $A \geq 1$, we have, as $N \rightarrow +\infty$,*

$$\frac{1}{N^3} \tilde{\mu}_N(f) = \int_0^\infty f(s) g_{\mathcal{L}_N^{\varphi, a, b}, \text{id}}(s) ds + O\left(\frac{A^7 \ln^2 N}{N^{1-\alpha}} \|f\|_\infty + \frac{A^3}{N^\alpha} \|f'\|_\infty\right).$$

Proof. Let A and f be as in the statement. Since the support of $\tilde{\omega}_{p,N}$ is contained in $[\frac{1}{Np}, \frac{N-p}{Np}]$, the support of $\iota_* \tilde{\omega}_{p,N}$ is contained in $[\frac{Np}{N-p}, Np]$. In particular the measures $\tilde{\mu}_N$ and $g_{\mathcal{L}_N^{\varphi, a, b}, \text{id}}(s) ds$ both vanish on $[0, 1]$. Hence we may assume that the support of f is contained in $[1, +\infty[$, so that the support of $f \circ \iota$ is contained in $]0, 1]$.

Note that $\|f \circ \iota\|_\infty = \|f\|_\infty$ and $\|(f \circ \iota)'\|_\infty = \|t^2 f'(t)\|_\infty \leq A^2 \|f'\|_\infty$, since the support of f' is contained in $[0, A]$.

By the definition of $\tilde{\mu}_N^+$, by Equation (15) and Lemma 5.2, since $N - p \leq N$ and by the restriction on p , explained in the proof of Lemma 5.3, in the summation defining $\tilde{\mu}_N^+(f)$ due to the support of f , we have

$$\begin{aligned}
\tilde{\mu}_N^+(f) &= \sum_{0 < p < N, p \equiv 0 \pmod b} \iota_* \tilde{\omega}_{p,N}(f) = \sum_{0 < p < N, p \equiv 0 \pmod b} \|\tilde{\omega}_{p,N}\| \frac{\tilde{\omega}_{p,N}}{\|\tilde{\omega}_{p,N}\|}(f \circ \iota) \\
&= \sum_{0 < p < N, p \equiv 0 \pmod b, p \leq O(A)} \left(\frac{c_{a,b,p}}{3} (N-p)^3 + O(N^2(p + \ln(N-p))^2) \right) \\
&\quad \times \left(\int_{\frac{1}{Np}}^{\frac{N-p}{Np}} 3 \left(\frac{Np}{N-p} \right)^3 t^2 f \circ \iota(t) dt \right. \\
&\quad \left. + O \left(\left(\frac{(Np)^3}{(N-p)^3(Np)^\alpha} + \frac{(Np)^6(p + \ln(Np))^2}{(N-p)^6(Np)^{1-\alpha}} \right) \|f \circ \iota\|_\infty + \frac{1}{(Np)^\alpha} \|(f \circ \iota)'\|_\infty \right) \right) \\
&= N^3 \sum_{0 < p < N, p \equiv 0 \pmod b, p \leq O(A)} c_{a,b,p} p^3 \int_{\frac{Np}{N-p}}^{Np} \frac{1}{s^4} f(s) ds \\
&\quad + N^3 \sum_{0 < p < N, p \equiv 0 \pmod b, p \leq O(A)} O \left(\frac{p^3 N^2 (p + \ln(N-p))^2}{(N-p)^3} \|f\|_\infty \right) \\
&\quad + O \left(\left(\frac{p^{3-\alpha}}{N^\alpha} + \frac{N^3 p^{5+\alpha} (p + \ln(Np))^2}{(N-p)^3 N^{1-\alpha}} \right) \|f\|_\infty + \frac{A^2}{(Np)^\alpha} \|f'\|_\infty \right).
\end{aligned}$$


Noting that $\int_p^{\frac{Np}{N-p}} \frac{1}{s^4} f(s) ds = O \left(\frac{N^2 p + N p^2 + p^3}{(Np)^3} \|f\|_\infty \right)$, we therefore have, for N large compared to A ,

$$\begin{aligned}
\frac{1}{N^3} \tilde{\mu}_N^+(f) &= \left(\sum_{0 < p < N, p \equiv 0 \pmod b} c_{a,b,p} p^3 \int_p^{+\infty} \frac{1}{s^4} f(s) ds \right) + O \left(\frac{A^4}{N} \|f\|_\infty \right) \\
&\quad + O \left(\frac{A^4 \ln^2(N)}{N} \|f\|_\infty + \left(\frac{A^4}{N^\alpha} + \frac{A^7 \ln^2(N)}{N^{1-\alpha}} \right) \|f\|_\infty + \frac{A^3}{N^\alpha} \|f'\|_\infty \right) \\
&= \int_0^{+\infty} f(s) g_{\mathcal{L}_N^{\varphi,a,b}, \text{id}}(s) ds + O \left(\left(\frac{A^4}{N^\alpha} + \frac{A^7 \ln^2(N)}{N^{1-\alpha}} \right) \|f\|_\infty + \frac{A^3}{N^\alpha} \|f'\|_\infty \right).
\end{aligned}$$

This proves Lemma 5.4, using that $\frac{A^4}{N^\alpha} = O \left(\frac{A^7 \ln^2(N)}{N^{1-\alpha}} \right)$ if $\alpha \geq \frac{1}{2}$. $\square$

Theorem 5.1 now follows from Lemmas 5.3 and 5.4, as explained in the beginning of the proof. $\square$

We close this section with a numerical illustration of Theorem 5.1 when $a = b = 1$. The following figure shows in red an approximation of the pair correlation function $g_{\mathcal{L}_N^{\varphi}, \text{id}} = g_{\mathcal{L}_N^{\varphi,1,1}, \text{id}}$ computed using $\tilde{\mathcal{H}}_{2000}$ in the interval $[-10, 10]$, to be compared with the graph of $g_{\mathcal{L}_N^{\varphi}, \text{id}}$ in the introduction.



The fact that the graph of $g_{\mathcal{L}_N^\varphi, \text{id}}$ has a horizontal asymptote near $\pm\infty$ follows from the following result.

Proposition 5.5 *We have $\lim_{s \rightarrow \pm\infty} g_{\mathcal{L}_N^\varphi, \text{id}}(s) = \frac{1}{4} \prod_{p \text{ prime}} (1 - \frac{2}{p^2})(1 + \frac{1}{p^2(p^2-2)})$.*

Proof. (Fouvry) In this proof, we use the same convention concerning $O(\cdot)$ as in the beginning of the proof of Lemma 4.2.

We consider the multiplicative² function $f : n \mapsto \prod_{p \text{ prime}, p|n} (1 + \frac{1}{p(p^2-2)})$ and the constant $C_1 = \prod_{p \text{ prime}} (1 + \frac{1}{p^2(p^2-2)})$. Let us prove that uniformly in $x \geq 1$, we have

$$\sum_{1 \leq n \leq x} n^3 f(n) = \frac{C_1}{4} x^4 + O(x^3). \quad (20)$$

By Equation (1), this proves Proposition 5.5.

Let $g = f * \mu$ be the Dirichlet convolution of f with the Möbius function μ . Then g is multiplicative. For every prime p , we have

$$g(p) = f(p) \mu(1) + f(1) \mu(p) = \frac{1}{p(p^2-2)}$$

and $g(p^k) = f(p^k) \mu(1) + f(p^{k-1}) \mu(p) = 0$ for every $k \geq 2$. Therefore, for every $m \geq 1$, we have

$$g(m) = \mu(m)^2 \prod_{p \text{ prime}, p|m} \frac{1}{p(p^2-2)}.$$

Lemma 5.6 *For every $m \geq 1$, we have $0 \leq g(m) \leq m^{-3} \prod_{p \text{ prime}} (1 - \frac{2}{p^2})^{-1}$.*

Proof. This is immediate if $\mu(m) = 0$. Otherwise, $m = p_1 \dots p_k$ with $p_1, \dots, p_k$ pairwise distinct primes, and

$$0 \leq m^3 g(m) = \prod_{i=1}^k \frac{p_i^3}{p_i(p_i^2-2)} = \prod_{i=1}^k (1 - \frac{2}{p_i^2})^{-1} \leq \prod_{p \text{ prime}} (1 - \frac{2}{p^2})^{-1} < +\infty. \quad \square$$

²Recall that an arithmetic function f is *multiplicative* if $f(1) = 1$ and for all coprime integers m, n , we have $f(mn) = f(m)f(n)$.

Therefore, using the Möbius inversion formula $f = g * \mathbf{1}$ for the first equality, Lemma 5.6 for the fifth equality and an Eulerian product (since g is multiplicative and vanishes on integers divisible by a nontrivial square) for the sixth equality, we have, with $S(x) = \sum_{1 \leq k \leq x} f(k)$,

$$\begin{aligned} S(x) &= \sum_{\substack{m, n \geq 1 \\ mn \leq x}} g(m) = \sum_{1 \leq m \leq x} g(m) \sum_{1 \leq n \leq x/m} 1 = \sum_{1 \leq m \leq x} g(m) \left(\frac{x}{m} + O(1) \right) \\ &= x \sum_{m=1}^{\infty} \frac{g(m)}{m} + O \left(x \sum_{m \geq x} \frac{g(m)}{m} \right) + O \left(\sum_{1 \leq m \leq x} g(m) \right) = x \sum_{m=1}^{\infty} \frac{g(m)}{m} + O(1) \\ &= x \prod_{p \text{ prime}} \left(1 + \frac{1}{p^2(p^2 - 2)} \right) + O(1) = C_1 x + O(1). \end{aligned}$$

By summation by parts, we hence have

$$\begin{aligned} \sum_{1 \leq n \leq x} n^3 f(n) &= \int_1^x t^3 d[S(t)] = [t^3(C_1 t + O(1))]_1^x - 3 \int_1^x t^2(C_1 t + O(1)) dt \\ &= \frac{C_1}{4} x^4 + O(x^3). \end{aligned}$$

This proves Equation (20) and concludes the proof of Proposition 5.5. $\square$

6 Pair correlations of common perpendiculars in the modular curve $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}_{\mathbb{R}}^2$

In this section, we give a geometric motivation for the introduction of the Euler function as multiplicities in the family $\mathcal{L}_{\mathbb{N}}^{\varphi}$ of logarithms of natural numbers. We refer to [PP, BPP] for more information.

Let Y be a nonelementary geodesically complete connected proper locally $\mathrm{CAT}(-1)$ good orbispace, so that the underlying space of Y is $\Gamma \backslash \tilde{Y}$ with $\tilde{Y}$ a geodesically complete proper $\mathrm{CAT}(-1)$ space and Γ a discrete group of isometries of $\tilde{Y}$ preserving no point nor pair of points in $\tilde{Y} \cap \partial_{\infty} \tilde{Y}$. Let D^- and D^+ be connected proper nonempty properly immersed locally convex closed subsets of Y , that is, D^- and D^+ are the Γ -orbits of proper nonempty closed convex subsets $\tilde{D}^-$ and $\tilde{D}^+$ of $\tilde{Y}$. A *common perpendicular* α between D^- and D^+ is the Γ -orbit of the unique shortest arc $\tilde{\alpha}$ between $\tilde{D}^-$ and $\gamma \tilde{D}^+$ for some $\gamma \in \Gamma$ such that $d(\tilde{D}^-, \gamma \tilde{D}^+) > 0$. The *multiplicity* $\mathrm{mult}(\alpha)$ of α is the ratio A/B where

- A is the number of elements $(\gamma_-, \gamma_+) \in (\Gamma/\Gamma_{D^-}) \times (\Gamma/\Gamma_{\gamma D^+})$ such that $\tilde{\alpha}$ is the unique shortest arc between $\gamma_- \tilde{D}^-$ and $\gamma_+ \gamma \tilde{D}^+$, and
- B is the cardinality of the pointwise stabilizer of $\tilde{\alpha}$ in Γ .

The *length* $\lambda(\alpha)$ of the common perpendicular α is the length of the geodesic segment $\tilde{\alpha}$ in $\tilde{Y}$. For every ℓ in the set $\mathrm{OL}^{\natural}(D^-, D^+)$ of lengths of common perpendiculars, the *length multiplicity* of ℓ is the sum of the multiplicities of the common perpendiculars between D^- , D^+ having the length ℓ :

$$\omega(\ell) = \sum_{\substack{\alpha \text{ common perpendicular} \\ \text{between } D^- \text{ and } D^+ \text{ with } \lambda(\alpha)=\ell}} \mathrm{mult}(\alpha).$$

If $\text{Perp}(D^-, D^+)$ is the set of all common perpendiculars from D^- to D^+ with multiplicities, then $(\lambda(\alpha))_{\alpha \in \text{Perp}(D^-, D^+)}$ is the *marked ortholength spectrum* from D^- to D^+ , and the set $\text{OL}(D^-, D^+) = (\text{OL}^{\natural}(D^-, D^+), \omega)$ of the lengths of the common perpendiculars endowed with the length multiplicity ω is the *ortholength spectrum* from D^- to D^+ .

The *pair correlation measure of the common perpendiculars from D^- to D^+* is the pair correlation measure of

$$\mathcal{F}_{D^-, D^+} = ((F_N = \text{OL}^{\natural}(D^-, D^+) \cap [0, 2 \ln N])_{N \in \mathbb{N}}, \omega).$$

We will study the asymptotic properties of the pair correlation measures $\mathcal{R}_N^{\mathcal{F}_{D^-, D^+}, \psi}$ for appropriate scalings ψ in a subsequent paper, and we only consider in this paper the following example.

Let

$$\tilde{Y} = \mathbb{H}_{\mathbb{R}}^2 = (\{z \in \mathbb{C} : \text{Im } z > 0\}, ds^2 = \frac{d(\text{Re } z)d(\text{Im } z)}{(\text{Im } z)^2})$$

be the upper halfspace model of the real hyperbolic plane with constant curvature -1 . For every $b \in \mathbb{N} - \{0\}$, let $\Gamma_0[b]$ be *Hecke's congruence subgroup modulo b* of the *modular group* $\text{PSL}_2(\mathbb{Z})$, which is the preimage of the upper triangular subgroup $\text{PSL}(\mathbb{Z}/b\mathbb{Z})$ under the reduction morphism $\text{PSL}_2(\mathbb{Z}) \rightarrow \text{PSL}(\mathbb{Z}/b\mathbb{Z})$. It acts faithfully by homographies on $\mathbb{H}_{\mathbb{R}}^2$, and is a lattice in the isometry group of $\mathbb{H}_{\mathbb{R}}^2$. Let $Y^b = \Gamma_0[b] \backslash \mathbb{H}_{\mathbb{R}}^2$, which is a finite (ramified) cover of the *modular curve* $\text{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}_{\mathbb{R}}^2$. Let $\tilde{D}^- = \tilde{D}^+$ be the horoball $\mathcal{H}_{\infty} = \{z \in \mathbb{C} : \text{Im } z \geq 1\}$ in $\mathbb{H}_{\mathbb{R}}^2$, whose image $D^- = D^+$ in Y^b is a *Margulis neighbourhood* of a cusp of Y^b . If $b = 1$, then $D^- = D^+$ is actually the maximal Margulis neighbourhood of the unique cusp of Y^b . To emphasize the dependence on the integer b , we will use the notation $\mathcal{F}_{D^-, D^+}^b = \mathcal{F}_{D^-, D^+}$ for the family of lengths of common perpendiculars between D^- and D^+ .

The following result says that the pair correlation measures of the common perpendiculars from this Margulis cusp neighbourhood to itself are, up to the homothety of factor 2, the pair correlation measures of the logarithms of the natural numbers congruent to 0 modulo b , with multiplicities given by the Euler function φ .

We use in the following result the notation $\mathcal{R}_N^{\mathcal{L}_N^{\varphi, b, b}, \psi}$ of the introduction with


$$\mathcal{L}_N^{\varphi, b, b} = ((L_N = \{\ln n : 0 < n \leq N, n \equiv 0 \pmod{b}\})_{N \in \mathbb{N}}, \omega = \varphi \circ \exp).$$

Proposition 6.1 *For every scaling function ψ and every $N \in \mathbb{N}$, if $f : t \mapsto 2t$, then*

$$\mathcal{R}_N^{\mathcal{F}_{D^-, D^+}^b, \psi} = f_* (\mathcal{R}_N^{\mathcal{L}_N^{\varphi, b, b}, \psi}).$$

Proof. The orbit of $\mathcal{H}_{\infty}$ under $\Gamma_0[b]$ consists, besides $\mathcal{H}_{\infty}$ itself, of the Euclidean disks $\mathcal{H}_{\frac{p}{q}}$ of Euclidean radius $\frac{1}{2q^2}$ tangent to the horizontal line at the rational numbers $\frac{p}{q}$ with $q > 0$, $q \equiv 0 \pmod{b}$ and $(p, q) = 1$.

Every common perpendicular between D^- and D^+ has a unique representative which starts from the Euclidean segment $[i, i + 1[$ on the boundary of $\mathcal{H}_\infty$ and ends on the boundary of $\mathcal{H}_q$ with $\frac{p}{q} \in \mathbb{Q} \cap [0, 1[$ and $q \equiv 0 \pmod{b}$. Its hyperbolic length is $2 \ln q$. In particular, we have $\text{OL}^b(D^-, D^+) = \{2 \ln q : q \geq 2, q \equiv 0 \pmod{b}\}$.



Since $\text{PSL}_2(\mathbb{R})$ acts simply transitively on the unit tangent bundle of $\mathbb{H}_{\mathbb{R}}^2$, the multiplicities of the common perpendiculars are equal to 1. Hence the length multiplicity of $2 \ln q$ is exactly the number of elements $p \in \mathbb{Z}/q\mathbb{Z}$ coprime with q , that is, $\omega(2 \ln q) = \varphi(q)$. $\square$

The following results, computing the pair correlation functions at trivial or linear scaling of the lengths of the common perpendiculars from the Margulis cusp neighbourhood at infinity to itself in Hecke's modular curve $\Gamma_0[b] \backslash \mathbb{H}_{\mathbb{R}}^2$, follow immediately from Theorems 4.1 and 5.1 with $a = b$, which also give an error term, using Proposition 6.1.

Corollary 6.2 (1) For all b , as $N \rightarrow +\infty$, the pair correlation measures $\mathcal{R}_N^{b, D^-, D^+, 1}$ on $\mathbb{R}$, renormalized to be probability measures, weak-star converge to a measure absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$, with pair correlation function given by $s \mapsto \frac{1}{2} e^{-|s|}$.

(2) For all b , as $N \rightarrow +\infty$, the pair correlation measures $\frac{1}{N^3} \mathcal{R}_N^{b, D^-, D^+, \text{id}}$ on $\mathbb{R}$ weak-star converge to a measure absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$, with pair correlation function given by $s \mapsto \frac{1}{s^4} \sum_{1 \leq k \leq |s|, k \equiv 0 \pmod{b}} c_{b,b,k} k^3$, where $c_{b,b,k}$ is defined in Equation (21). $\square$

A Appendix : A Mirsky formula with congruences, by Etienne Fouvry

The aim of this appendix is to give a proof of a version with congruences of an arithmetic formula due to Mirsky [Mir], which is used in the proof of Theorem 5.1.

Let $a, b, k \in \mathbb{N}$ be fixed integers satisfying $a, b \geq 1$. Denoting by φ the Euler function, we give an asymptotic formula, as $x \geq 1$ tends to $+\infty$, for the quantity

$$S(x; a, b, k) = \sum_{\substack{1 \leq n \leq x \\ n \equiv a \pmod{b}}} \varphi(n) \varphi(n + k) .$$

Throughout this appendix, the letter p denotes as usual a varying positive prime in $\mathbb{Z}$, and we use the convention on $O(\cdot)$ of the beginning of Lemma 4.2. For all integers $\alpha, \beta \geq 1$, we denote as usual by (α, β) and $[\alpha, \beta]$ their positive gcd et lcm, respectively.

Theorem A.1 *Let*

$$c_{a,b,k} = \frac{1}{b} \prod_{\substack{p \\ (p,b) \mid k+a}} \left(1 - \frac{(p,b)}{p^2}\right) \prod_{\substack{p \\ (p,b) \mid a}} \left(1 - (p,b) \frac{\kappa_{a,b,k}(p) \kappa'_k(p)}{p^2}\right), \quad (21)$$

where

$$\kappa_{a,b,k}(p) = \begin{cases} (1 - \frac{(p,b)}{p^2})^{-1} & \text{if } (p,b) \mid a+k \\ 1 & \text{otherwise} \end{cases} \quad \text{and} \quad \kappa'_k(p) = \begin{cases} 1 - \frac{1}{p} & \text{if } p \mid k \\ 1 & \text{otherwise.} \end{cases}$$

There exists an absolute constant $K > 0$ such that for all integers $a, b \geq 1$ and $k \geq 0$ and real number $x \geq 1$, we have

$$\left| S(x; a, b, k) - c_{a,b,k} \left(\frac{x^3}{3} + \frac{kx^2}{2} \right) \right| \leq K(x(x+k)(\ln 2x) \ln(2x+k)).$$

Remark A.2 Before proving this theorem, we give some considerations on the constant $c_{a,b,k}$. We start from the obvious inequalities, for every prime p ,

$$1 \leq \kappa_{a,b,k}(p) \leq 2 \quad \text{and} \quad \frac{1}{2} \leq \kappa'_k(p) \leq 1.$$

From these inequalities and the definition (21), we deduce that $c_{a,b,k}$ satisfies the following inequalities

$$\frac{1}{b} \Lambda(a, b, k) \prod_{\substack{p \geq 3 \\ (p,b) \mid a+k}} \left(1 - \frac{(p,b)}{p^2}\right) \prod_{\substack{p \geq 3 \\ (p,b) \mid a}} \left(1 - 2 \cdot \frac{(p,b)}{p^2}\right) \leq c_{a,b,k} < \frac{1}{b}, \quad (22)$$

where the local factor $\Lambda(a, b, k)$, obtained by separating the case $p = 2$ in the Euler products, is defined by

$$\Lambda(a, b, k) = \begin{cases} 5/8 & \text{if } 2 \nmid b \text{ and } 2 \mid k, \\ 1/2 & \text{if } 2 \nmid b \text{ and } 2 \nmid k, \\ 1 & \text{if } 2 \mid b, 2 \nmid a \text{ and } 2 \mid k, \\ 1/2 & \text{if } 2 \mid b, 2 \nmid a \text{ and } 2 \nmid k, \\ 1/4 & \text{if } 2 \mid b, 2 \mid a \text{ and } 2 \mid k, \\ 1/2 & \text{if } 2 \mid b, 2 \mid a \text{ and } 2 \nmid k. \end{cases}$$

We have the positive lower bound $\Lambda(a, b, k) \geq \frac{1}{4}$ in all cases. This shows in particular that $c_{a,b,k} > 0$. A deeper look also leads to the statement that the positive product $b c_{a,b,k}$ can be arbitrarily small : it suffices to consider the case where all the integers a, b and k are all divisible by the t smallest primes, and letting the integer t tend to infinity.

Theorem A.1 was already known when $a = b = 1$, this result is due to [Mir, Thm. 9, Eq. (30)], with

$$c_{1,1,k} = \prod_p \left(1 - \frac{2}{p^2}\right) \prod_{p \mid k} \left(1 + \frac{1}{p(p^2 - 2)}\right). \quad (23)$$

For this computation, see Remark A.6 below.

Proof of Theorem A.1. For every $x \geq 1$, let us first prove that there exists a constant $c_{a,b,k} \in]0, 1]$ such that the sum

$$\tilde{S}(x) = \sum_{\substack{1 \leq n \leq x \\ n \equiv a \pmod{b}}} \frac{\varphi(n)}{n} \cdot \frac{\varphi(n+k)}{n+k}$$

satisfies the asymptotic formula, uniformly in $a, b \geq 1$, $k \geq 0$ and $x \geq 1$,

$$\tilde{S}(x) = c_{a,b,k} x + O((\ln 2x) \ln(2x+k)). \quad (24)$$

Theorem A.1 follows classically, by applying Abel's summation formula

$$\sum_{1 \leq n \leq x} a_n f(n) = \left(\sum_{1 \leq n \leq x} a_n \right) f(x) - \int_1^x \left(\sum_{1 \leq n \leq t} a_n \right) f'(t) dt$$

to the numerical sequence $(a_n = \frac{\varphi(n)}{n} \cdot \frac{\varphi(n+k)}{n+k} \delta_n)_{n \geq 1}$, where $\delta_n = 1$ if $n \equiv a \pmod{b}$ and $\delta_n = 0$ otherwise, and to the function $f : [1, +\infty[\rightarrow \mathbb{R}$ of class C^1 defined by $x \mapsto x(x+k)$. We indeed have

$$\begin{aligned} S(x; a, b, k) &= \tilde{S}(x)x(x+k) - \int_1^x \tilde{S}(t)(2t+k) dt \\ &= \frac{c_{a,b,k}}{3} x^3 + \frac{c_{a,b,k}k}{2} x^2 + \left(\frac{2}{3} + \frac{k}{2}\right)c_{a,b,k} + O(x(x+k)(\ln 2x) \ln(2x+k)), \end{aligned}$$

which gives the result since $c_{a,b,k} \leq 1$.

Let us denote by $\mathbf{1}$ the constant arithmetic function with value 1. The convolution equality $\varphi = \mu \star \text{id}$ implies by division that $\frac{\varphi}{\text{id}} = \frac{\mu}{\text{id}} \star 1$. Applying twice this formula, we have

$$\tilde{S}(x) = \sum_{1 \leq d \leq x} \frac{\mu(d)}{d} \sum_{1 \leq \delta \leq x+k} \frac{\mu(\delta)}{\delta} \sum_{\substack{1 \leq n \leq x \\ d|n, \delta|(n+k) \\ n \equiv a \pmod{b}}} 1.$$

The system of three congruences $n \equiv \begin{cases} 0 \pmod{d} \\ -k \pmod{\delta} \\ a \pmod{b} \end{cases}$ has a solution $n \leq x$ if and only if there exists an integer $m \leq x/d$ such that $n = dm$ and

$$\begin{cases} dm \equiv -k \pmod{\delta} \\ dm \equiv a \pmod{b}. \end{cases} \quad (25)$$

When $(d, \delta) \nmid k$ or when $(d, b) \nmid a$, no solution exists. We hence have

$$\tilde{S}(x) = \sum_{\substack{1 \leq d \leq x, 1 \leq \delta \leq x+k \\ (d, \delta) | k, (d, b) | a}} \frac{\mu(d)}{d} \frac{\mu(\delta)}{\delta} \text{Card}\{m \leq x/d : \begin{matrix} dm \equiv -k \pmod{\delta} \\ dm \equiv a \pmod{b} \end{matrix}\}.$$

If $(d, \delta) \mid k$ and $(d, b) \mid a$, let us denote by $\overline{\frac{d}{(d, \delta)}}$ the multiplicative inverse of the integer $\frac{d}{(d, \delta)}$ modulo $\frac{\delta}{(d, \delta)}$ and by $\overline{\frac{d}{(d, b)}}$ the multiplicative inverse of the integer $\frac{d}{(d, b)}$ modulo $\frac{b}{(d, b)}$.

The system of two congruences (25), after division of its first equation by (d, δ) and of its second equation by (d, b) , is then equivalent to the system

$$\begin{cases} m \equiv -\frac{k}{(d, \delta)} \frac{\overline{d}}{(d, \delta)} \pmod{\frac{\delta}{(d, \delta)}} \\ m \equiv \frac{a}{(d, b)} \frac{\overline{d}}{(d, b)} \pmod{\frac{b}{(d, b)}} . \end{cases}$$

This system has a solution if and only if the following divisibility condition holds

$$\begin{aligned} & \left(\frac{\delta}{(d, \delta)}, \frac{b}{(d, b)} \right) \mid \frac{k}{(d, \delta)} \frac{\overline{d}}{(d, \delta)} + \frac{a}{(d, b)} \frac{\overline{d}}{(d, b)} \\ \Leftrightarrow & \left(\frac{\delta}{(d, \delta)}, \frac{b}{(d, b)} \right) \mid \frac{k}{(d, \delta)} \frac{d}{(d, b)} + \frac{a}{(d, b)} \frac{d}{(d, \delta)} \end{aligned}$$

Since the integers $\frac{d}{(d, b)}$ and $\frac{d}{(d, \delta)}$ are coprime with the gcd $(\frac{\delta}{(d, \delta)}, \frac{b}{(d, b)})$, we deduce by multiplication that the above condition holds if and only if we have

$$(\delta(d, b), b(d, \delta)) \mid d(k + a) .$$

The following lemma, where $O(1)$ is uniformly bounded in $\alpha, \beta, \alpha_0, \beta_0$, is elementary.

Lemma A.3 *For all integers $\alpha, \beta, \alpha_0, \beta_0 \geq 1$ and real number $y \geq 1$, we have*

$$\text{Card}\{m \leq y : m \equiv \alpha_0 \pmod{\alpha} \text{ and } m \equiv \beta_0 \pmod{\beta}\} = \begin{cases} 0 & \text{if } \alpha_0 \not\equiv \beta_0 \pmod{(\alpha, \beta)} \\ \frac{y}{[\alpha, \beta]} + O(1) & \text{otherwise} . \quad \square \end{cases}$$

This lemma implies that $\text{Card}\{m \leq x/d : \begin{matrix} dm \equiv -k \pmod{\delta} \\ dm \equiv a \pmod{b} \end{matrix}\} = \frac{x}{d \left[\frac{\delta}{(d, \delta)}, \frac{b}{(d, b)} \right]} + O(1)$ under the assumption that $(d, \delta) \mid k$ and $(d, b) \mid a$, where $O(1)$ is uniformly bounded in a, b, k, d, δ . By the classical majoration of the harmonic series, we have

$$\sum_{1 \leq d \leq x, 1 \leq \delta \leq x+k} \frac{1}{d} \frac{1}{\delta} = O((\ln 2x) \ln(2x + k)) .$$

By the relation between lcm, gcd and product of two positive integers, we hence have

$$\tilde{S}(x) = x \sum_{\substack{1 \leq d \leq x, 1 \leq \delta \leq x+k \\ (d, \delta) \mid k, (d, b) \mid a \\ (\delta(d, b), b(d, \delta)) \mid d(k+a)}} \frac{\mu(d)}{d} \frac{\mu(\delta)}{\delta} \frac{(\delta(d, b), b(d, \delta))}{d\delta b} + O((\ln 2x) \ln(2x + k)) ,$$

uniformly in $a, b \geq 1, k \geq 0$ and $x \geq 1$.

Completing the sum with the indices $d > x$ and $\delta > x + k$ introduces an error of the form (uniformly in $a, b \geq 1, k \geq 0$ and $x \geq 1$)

$$O\left(\sum_{d \geq x, \delta \geq 1} \frac{(d, \delta)}{d^2 \delta^2}\right) = O\left(\sum_{t \geq 1, d' \geq x/t, \delta' \geq 1} \frac{t}{(td')^2 (t\delta')^2}\right) = O\left(\frac{1}{x}\right) .$$

This proves Formula (24) by setting

$$c_{a, b, k} = \sum_{\substack{d, \delta \geq 1 \\ (d, \delta) \mid k, (d, b) \mid a \\ (\delta(d, b), b(d, \delta)) \mid d(k+a)}} \frac{\mu(d)}{d} \frac{\mu(\delta)}{\delta} \frac{(\delta(d, b), b(d, \delta))}{d\delta b} . \quad (26)$$

Let us now prove Equation (21). By Remark A.2, this implies that $0 < c_{a,b,k} \leq 1$, hence completes the proof of Theorem A.1.

Proof of Equation (21). For every integer $d \geq 1$, let χ_d be the characteristic function of the set of integers $\delta \geq 1$ such that $(\delta, d) \mid k$. For every integer $d \geq 1$, let us define

$$\psi_d : \delta \mapsto \left(\delta, \frac{b}{(d,b)} (d, \delta) \right). \quad (27)$$

Note that the assertion $(\delta(d, b), b(d, \delta)) \mid d(k + a)$ is equivalent to the assertion

$$\psi_d(\delta) \mid \frac{d}{(d,b)} (k + a).$$

For every integer $d \geq 1$, let χ_d^* be the characteristic function of the set of integers $\delta \geq 1$ such that the above divisibility assertion is satisfied. Let us define

$$c^* : d \mapsto \sum_{\delta \geq 1} \frac{\mu(\delta)}{\delta^2} \chi_d(\delta) \chi_d^*(\delta) \psi_d(\delta) \quad (28)$$

(this arithmetic function c^* depends on the constants a, b, k). Equation (26) then becomes

$$c_{a,b,k} = \frac{1}{b} \sum_{\substack{d \geq 1 \\ (d,b) \mid a}} \frac{\mu(d)}{d^2} (d, b) c^*(d). \quad (29)$$

In order to transform the series $c^*(d)$ defined by Formula (28) into an Eulerian product and in order to analyse it, we will use the following two lemmas.

Lemma A.4 *For every integer $d \geq 1$, the arithmetic functions χ_d , χ_d^* and ψ_d are multiplicative.*

Proof. We have $\chi_d(1) = \chi_d^*(1) = \psi_d(1) = 1$. Let δ_1, δ_2 be two coprime integers.

The equality $(\delta_1 \delta_2, d) = (\delta_1, d)(\delta_2, d)$ and the fact that (δ_1, d) and (δ_2, d) are coprime imply the multiplicativity of χ_d .

In order to prove the multiplicativity of the function ψ_d , we write

$$\psi_d(\delta_1 \delta_2) = \left(\delta_1 \delta_2, \frac{b}{(d,b)} (d, \delta_1 \delta_2) \right) = \left(\delta_1, \frac{b}{(d,b)} (\delta_1, d)(\delta_2, d) \right) \left(\delta_2, \frac{b}{(d,b)} (\delta_1, d)(\delta_2, d) \right).$$

Since δ_1 is coprime to (δ_2, d) and since δ_2 is coprime to (δ_1, d) , we obtain as wanted the equality $\psi_d(\delta_1 \delta_2) = \psi_d(\delta_1) \psi_d(\delta_2)$.

Finally, the multiplicativity of the function χ_d^* is a consequence of the multiplicativity of the function ψ_d and of the fact that $\psi_d(\delta_1)$ and $\psi_d(\delta_2)$ are coprime. $\square$

Lemma A.5 *For every prime p and every integer $d \geq 1$, we have*

$$\psi_d(p) = \begin{cases} p & \text{if } p \mid d, \\ (p, b) & \text{otherwise,} \end{cases}$$

and

$$\chi_d(p) \chi_d^*(p) = 1 \Leftrightarrow \begin{cases} p \mid (d, k) & \text{and } p \mid \frac{d}{(d,b)} (k + a), \\ \text{or} \\ p \nmid d & \text{and } (p, b) \mid k + a. \end{cases}$$

Proof. The first formula follows from the definition of $\psi_d(p)$ (see Formula (27)) by considering the three cases $(p \mid d)$, $(p \nmid d \text{ and } p \mid b)$, and $(p \nmid d \text{ and } p \nmid b)$.

The second formula follows from the first one, from the definitions of $\chi_d(p)$ and $\chi_d^*(p)$, and from the fact that $\chi_d(p) \chi_d^*(p) = 1$ if and only if $\chi_d(p) = \chi_d^*(p) = 1$, by considering the two cases $(p \mid d)$ and $(p \nmid d)$. $\square$

The arithmetic function $\delta \mapsto \mu(\delta) \chi_d(\delta) \chi_d^*(\delta) \psi_d(\delta)$ being multiplicative by Lemma A.4, and vanishing on the nontrivial powers of primes, the series defining $c^*(d)$ in Formula (28) may be written as an Eulerian product

$$c^*(d) = \prod_p \left(1 - \frac{\chi_d(p) \chi_d^*(p) \psi_d(p)}{p^2} \right) = \prod_{\substack{p \\ \chi_d(p) \chi_d^*(p) = 1}} \left(1 - \frac{\psi_d(p)}{p^2} \right). \quad (30)$$

By Equations (29) and (30), and by Lemma A.5, we have

$$c_{a,b,k} = \frac{1}{b} \sum_{\substack{d \geq 1 \\ (d,b) \mid a}} \frac{\mu(d)}{d^2} (d,b) \prod_{\substack{p \nmid d \\ (p,b) \mid k+a}} \left(1 - \frac{(p,b)}{p^2} \right) \prod_{\substack{p \mid (d,k) \\ p \mid \frac{d}{(d,b)}(k+a)}} \left(1 - \frac{1}{p} \right).$$

Let us define $\Gamma_{a,b,k} = \prod_{\substack{p \\ (p,b) \mid k+a}} \left(1 - \frac{(p,b)}{p^2} \right)$, so that

$$c_{a,b,k} = \frac{\Gamma_{a,b,k}}{b} \sum_{\substack{d \geq 1 \\ (d,b) \mid a}} \frac{\mu(d)}{d^2} (d,b) \prod_{\substack{p \mid d \\ (p,b) \mid k+a}} \left(1 - \frac{(p,b)}{p^2} \right)^{-1} \prod_{\substack{p \mid (d,k) \\ p \mid \frac{d}{(d,b)}(k+a)}} \left(1 - \frac{1}{p} \right). \quad (31)$$

For every integer $d \geq 1$ without square factor such that $(d,b) \mid a$, we have

$$\begin{aligned} \prod_{\substack{p \mid (d,k) \\ p \mid \frac{d}{(d,b)}(k+a)}} \left(1 - \frac{1}{p} \right) &= \prod_{\substack{p \mid (d,k) \\ p \mid \frac{d}{(d,b)}}} \left(1 - \frac{1}{p} \right) \prod_{\substack{p \mid (d,k) \\ p \mid k+a}} \left(1 - \frac{1}{p} \right) \prod_{\substack{p \mid (d,k) \\ p \mid \frac{d}{(d,b)}(k+a)}} \left(1 - \frac{1}{p} \right)^{-1} \\ &= \prod_{p \mid \left(\frac{d}{(d,b)}, k \right)} \left(1 - \frac{1}{p} \right) \prod_{p \mid (d,a,k)} \left(1 - \frac{1}{p} \right) \prod_{p \mid \left(\frac{d}{(d,b)}, a, k \right)} \left(1 - \frac{1}{p} \right)^{-1} \\ &= \prod_{p \mid \left(\frac{d}{(d,b)}, k \right)} \left(1 - \frac{1}{p} \right) \prod_{p \mid (d,a,b,k)} \left(1 - \frac{1}{p} \right) \\ &= \prod_{p \mid \left(\frac{d}{(d,b)}, k \right)} \left(1 - \frac{1}{p} \right) \prod_{p \mid (d,b,k)} \left(1 - \frac{1}{p} \right) = \prod_{p \mid (d,k)} \left(1 - \frac{1}{p} \right). \end{aligned}$$

Thus, Equation (31) writes $c_{a,b,k}$ as a series $\frac{\Gamma_{a,b,k}}{b} \sum_{\substack{d \geq 1 \\ (d,b) \mid a}} \frac{f(d)}{d^2}$ where f is a multiplicative

function, which vanishes on the nontrivial powers of primes. By Eulerian product, we have therefore proved Equation (21). $\square$

Remark A.6 When $a = b = 1$, we indeed recover Mirsky’s result [Mir, Thm. 9, Eq. (30)]. Indeed, by Equation (21), we have

$$\begin{aligned}
c_{1,1,k} &= \prod_p \left(1 - \frac{1}{p^2}\right) \prod_{p|k} \left(1 - \frac{(1 - \frac{1}{p^2})^{-1}(1 - \frac{1}{p})}{p^2}\right) \prod_{p \nmid k} \left(1 - \frac{(1 - \frac{1}{p^2})^{-1}}{p^2}\right) \\
&= \prod_p \left(1 - \frac{1}{p^2}\right) \prod_{p|k} \left(1 - \frac{p-1}{p(p^2-1)}\right) \prod_p \left(1 - \frac{1}{p^2-1}\right) \prod_{p|k} \left(1 - \frac{1}{p^2-1}\right)^{-1} \\
&= \prod_p \left(1 - \frac{2}{p^2}\right) \prod_{p|k} \left(1 + \frac{1}{p(p^2-2)}\right).
\end{aligned}$$

References

- [BocZ] F. P. Boca and A. Zaharescu. *The correlations of Farey fractions*. J. London Math. Soc. **72** (2005) 25–39.
- [Boh] O. Bohigas. *Random matrix theories and chaotic dynamics*. In “Chaos et physique quantique” (Les Houches, 1989), 87–199, North-Holland, 1991.
- [BPP] A. Broise-Alamichel, J. Parkkonen, and F. Paulin. *Equidistribution and counting under equilibrium states in negative curvature and trees. Applications to non-Archimedean Diophantine approximation*. With an Appendix by J. Buzzi. Prog. Math. **329**, Birkhäuser, 2019.
- [Cha] D. Chandler. *Introduction to modern statistical mechanics*. Oxford Univ. Press, 1987.
- [HaW] G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. Oxford Univ. Press, sixth ed., 2008.
- [HK] R. Hofer and L. Kaltenböck. *Pair correlations of Halton and Niederreiter sequences are not Poissonian*. Monat. Math. **194** (2021) 789–809.
- [LS] G. Larcher and W. Stockinger. *Some negative results related to Poissonian pair correlation problems*. Discrete Math. **343** (2020) 111656.
- [Mir] L. Mirsky. *Summation formulae involving arithmetic functions*. Duke Math. J. **16** (1949), 261–272 .
- [MaS] J. Marklof and A. Strömbergsson. *Gaps between logs*. Bull. Lond. Math. Soc. **45** (2013) 1267–1280.
- [Mon] H. L. Montgomery. *The pair correlation of zeros of the zeta function*. In "Analytic number theory" (Proc. Sympos. Pure Math. Vol. XXIV, St. Louis Univ., 1972), pp 181–193, Amer. Math. Soc. 1973.
- [PP] J. Parkkonen and F. Paulin. *Counting common perpendicular arcs in negative curvature*. Erg. Theo. Dyn. Sys. **37** (2017) 900–938.
- [PS1] M. Pollicott and R. Sharp. *Correlations for pairs of closed geodesics*. Invent. Math. **163** (2006) 1–24.
- [PS2] M. Pollicott and R. Sharp. *Correlations of Length Spectra for Negatively Curved Manifolds*. Commun. Math. Phys. **319** (2013) 515–533 .
- [RS] Z. Rudnick and P. Sarnak. *The pair correlation function of fractional parts of polynomials*. Commun. Math. Phys. **194** (1998) 61–70.
- [SdH] A. Santos and M. Lopez de Haro. *Radial distribution function for hard spheres in fractal dimensions: A heuristic approximation*. Phys. Rev. E **93** (2016) 062126.

- [ZP] F. Zernike and J. A. Prins. *Die Beugung von Röntgenstrahlen in Flüssigkeiten als Effekt der Molekulanordnung*. Z. Physik **41** (1927) 184–194.

Department of Mathematics and Statistics, P.O. Box 35
40014 University of Jyväskylä, FINLAND.
e-mail: jouni.t.parkkonen@jyu.fi

Laboratoire de mathématique d’Orsay, UMR 8628 CNRS,
Université Paris-Saclay,
91405 ORSAY Cedex, FRANCE
e-mail: frederic.paulin@universite-paris-saclay.fr

Laboratoire de mathématique d’Orsay, UMR 8628 CNRS,
Université Paris-Saclay,
91405 ORSAY Cedex, FRANCE
e-mail: etienne.fouvry@universite-paris-saclay.fr