

**$q$ -DIFFERENCE EQUATION FOR GENERALIZED TRIVARIATE  $q$ -HAHN POLYNOMIALS**SAMA ARJIKA<sup>1</sup> AND MAHAMAN KABIR MAHAMAN<sup>2</sup>

**ABSTRACT.** In this paper, we introduce a family of trivariate  $q$ -Hahn polynomials  $\Psi_n^{(a)}(x, y, z|q)$  as a general form of Hahn polynomials  $\psi_n^{(a)}(x|q)$ ,  $\psi_n^{(a)}(x, y|q)$  and  $F_n(x, y, z; q)$ . We represent  $\Psi_n^{(a)}(x, y, z|q)$  by the homogeneous  $q$ -difference operator  $\tilde{L}(a, b; \theta_{xy})$  introduced by Srivastava *et al* [H. M. Srivastava, S. Arjika and A. Sherif Kelil, *Some homogeneous  $q$ -difference operators and the associated generalized Hahn polynomials*, Appl. Set-Valued Anal. Optim. **1** (2019), pp. 187–201.] to derive: extended generating, Rogers formula, extended Rogers formula and Srivastava-Agarwal type generating functions involving  $\Psi_n^{(a)}(x, y, z|q)$  by the  $q$ -difference equation.

**1. Introduction**

In this paper, we adopt the common conventions and notations on  $q$ -series. For the convenience of the reader, we provide a summary of the mathematical notations, basics properties and definitions to be used in the sequel. We refer to the general references (see [9]) for the definitions and notations. Throughout this paper, we assume that  $|q| < 1$ .

For complex numbers  $a$ , the  $q$ -shifted factorials are defined by:

$$(a; q)_0 := 1, \quad (a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k), \quad (a; q)_\infty := \prod_{k=0}^{\infty} (1 - aq^k) \quad (1.1)$$

and  $(a_1, a_2, \dots, a_r; q)_m = (a_1; q)_m (a_2; q)_m \cdots (a_r; q)_m$ ,  $m \in \{0, 1, 2, \dots\}$ .

The  $q$ -binomial coefficient is defined as [5]

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} = \frac{(q^{-n}; q)_k}{(q; q)_k} (-1)^k q^{nk - \binom{k}{2}}, \quad \text{for } 0 \leq k \leq n.$$

The basic or  $q$ -hypergeometric function in the variable  $z$  (see Slater [12, Chap. 3], Srivastava and Karlsson [13, p. 347, Eq. (272)] for details) is defined as:

$${}_r\Phi_s \left[ \begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} q; z \right] = \sum_{n=0}^{\infty} \frac{[(-1)^n q^{\binom{n}{2}}]^{1+s-r} (a_1, a_2, \dots, a_r; q)_n}{(b_1, b_2, \dots, b_s; q)_n (q; q)_n} z^n,$$

when  $r > s + 1$ . Note that, for  $r = s + 1$ , we have:

$${}_{r+1}\Phi_r \left[ \begin{matrix} a_1, a_2, \dots, a_{r+1}; \\ b_1, b_2, \dots, b_r; \end{matrix} q; z \right] = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1}; q)_n}{(b_1, b_2, \dots, b_r; q)_n (q; q)_n} z^n.$$

We will be mainly concerned with the Cauchy polynomials as given below [4]

$$p_n(x, y) := (x - y)(x - qy) \cdots (x - q^{n-1}y) = (y/x; q)_n x^n \quad (1.2)$$

*Key words and phrases.*  $q$ -difference equation; homogeneous  $q$ -operator; Hahn polynomials; generating functions.

<sup>1</sup>Department of Mathematics and Informatics, University of Agadez, Niger. <sup>2</sup>Department of Mathematics, ENS, University Abdou Moumouni of Niamey, B.P.: 10 896, Niamey, Niger.

Emails: rjksama2008@gmail.com (S.A.), mahamankabir@yahoo.fr (M.K.M).

2010 *Mathematics Subject Classification.* 05A30, 39A13, 33D15, 33D45.

with the Srivastava-Agarwal type generating function

$$\sum_{n=0}^{\infty} p_n(x, y) \frac{(\lambda; q)_n t^n}{(q; q)_n} = {}_2\Phi_1 \left[ \begin{matrix} \lambda, y/x; \\ 0; \end{matrix} q; xt \right]. \quad (1.3)$$

For  $\lambda = 0$ , we get the generating function [4]

$$\sum_{n=0}^{\infty} p_n(x, y) \frac{t^n}{(q; q)_n} = \frac{(yt; q)_{\infty}}{(xt; q)_{\infty}}. \quad (1.4)$$

The generating function (1.4) is also the homogeneous version of the Cauchy identity or the  $q$ -binomial theorem given by [5]

$$\sum_{k=0}^{\infty} \frac{(a; q)_k}{(q; q)_k} z^k = {}_1\Phi_0 \left[ \begin{matrix} a; \\ -; \end{matrix} q; z \right] = \frac{(az; q)_{\infty}}{(z; q)_{\infty}}, \quad |z| < 1. \quad (1.5)$$

Putting  $a = 0$ , the relation (1.5) becomes Euler's identity [5]

$$\sum_{k=0}^{\infty} \frac{z^k}{(q; q)_k} = \frac{1}{(z; q)_{\infty}} \quad |z| < 1 \quad (1.6)$$

and its inverse relation [5]

$$\sum_{k=0}^{\infty} \frac{(-1)^k q^{\binom{k}{2}} z^k}{(q; q)_k} = (z; q)_{\infty}. \quad (1.7)$$

Saad and Sukhi [11] defined the  $q$ -difference operator  $\theta_{xy}$

$$\theta_{xy}\{f(x, y)\} := \frac{f(q^{-1}x, y) - f(x, qy)}{q^{-1}x - y}, \quad (1.8)$$

which turns out to be suitable for dealing with the Cauchy polynomials. Their corresponding  $q$ -exponential operator is

$$\mathbb{E}(z\theta_{xy}) = \sum_{k=0}^{\infty} \frac{q^{\binom{k}{2}}}{(q; q)_k} (z\theta_{xy})^k. \quad (1.9)$$

Recently, Srivastava, Arjika and Kelil [14] have introduced the  $q$ -difference operator  $\widetilde{L}(a, b; \theta_{xy})$

$$\widetilde{L}(a, b; \theta_{xy}) = \sum_{k=0}^{\infty} \frac{q^{\binom{k}{2}} (a; q)_k}{(q; q)_k} (b\theta_{xy})^k, \quad (1.10)$$

to study  $q$ -polynomials and related generating functions.

In this paper, our goal is to generalize the results of Srivastava, Arjika and Kelil [14], and Mohameed [1]. We first construct the following generalized trivariate  $q$ -Hahn polynomials as

$$\Psi_n^{(a)}(x, y, z|q) = (-1)^n q^{-\binom{n}{2}} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-1)^k q^{\binom{k}{2}} (a; q)_k p_{n-k}(y, x) z^k. \quad (1.11)$$

**Remark 1.** For  $a = 0$ , the generalized trivariate  $q$ -Hahn polynomials  $\Psi_n^{(a)}(x, y, z|q)$  are the well known trivariate  $q$ -polynomials  $F_n(x, y, z; q)$  investigated by Mohameed (see [1] for more details), i.e.,

$$\Psi_n^{(0)}(x, y, z|q) = F_n(x, y, z; q). \quad (1.12)$$

If we let  $a = 0$ ,  $y = ax$  and  $z = y$ , the generalized trivariate  $q$ -Hahn polynomials  $\Psi_n^{(a)}(x, y, z|q)$  reduce to the second Hahn polynomials  $\psi_n^{(a)}(x, y|q)$  [3], i.e.,

$$\Psi_n^{(0)}(x, ax, y|q) = \psi_n^{(a)}(x, y|q). \quad (1.13)$$

Also,  $a = 0, y = ax$  and  $z = 1$ , the generalized trivariate  $q$ -Hahn  $\Psi_n^{(a)}(x, y, z|q)$  reduce to Hahn polynomials  $\psi_n^{(a)}(x|q)$  [2], i.e.,

$$\Psi_n^{(0)}(x, ax, 1|q) = \psi_n^{(a)}(x|q). \quad (1.14)$$

The polynomials (1.11) can be represented by the homogeneous  $q$ -difference operator (1.10) as follows.

**Proposition 2.**

$$\Psi_n^{(a)}(x, y, z|q) = \tilde{L}(a, z; \theta_{xy}) \left\{ (-1)^n q^{-\binom{n}{2}} p_n(y, x) \right\}. \quad (1.15)$$

*Proof.* By identity (1.10) and taking into account  $\theta_{xy} p_n(y, x) = -(1 - q^n) p_{n-1}(y, x)$ , we get the result.  $\square$

In light of  $\theta_{xy}^k [(xt; q)_\infty / (yt; q)_\infty] = (-t)^k [(xt; q)_\infty / (yt; q)_\infty]$ , we have the following identity

$$\tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xt; q)_\infty}{(yt; q)_\infty} \right\} = \frac{(xt; q)_\infty}{(yt; q)_\infty} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; zt \right]. \quad (1.16)$$

The main object of this paper is to use the  $q$ -difference equation to derive some identities such as: extended generating function, Rogers formula, extended Rogers formula and Srivastava-Agarwal type generating functions.

The paper is organized as follows. In Section 2, we state two theorems and give the proofs. We derive an extended generating function for these  $q$ -polynomials. In Section 3, we state the Rogers formula and extended Rogers formula and give the proofs by the  $q$ -difference equation. In Section 4, we obtain Srivastava-Agarwal type generating functions involving the generalized trivariate  $q$ -Hahn polynomials by the method of  $q$ -difference equation.

## 2. MAIN RESULTS AND PROOFS

In this section, we introduce another extension of  $q$ -Hahn polynomials. Then, we represent it by the homogeneous  $q$ -difference operator and derive an extended generating function.

**Theorem 3.** Let  $f(a, b, x, y, z)$  be an 5-variable analytic function at  $(a, b, x, y, z) = (0, 0, 0, 0, 0) \in \mathbb{C}^5$ . If  $f(a, b, x, y, z)$  satisfies the  $q$ -difference equation

$$(q^{-1}x - y) [f(a, b, x, y, z) - f(a, b, x, y, qz)] = z [f(a, b, q^{-1}x, y, qz) - f(a, b, x, qy, qz)] + az [f(a, b, x, qy, q^2z) - f(a, b, q^{-1}x, y, q^2z)], \quad (2.1)$$

then we have:

$$f(a, b, x, y, z) = \tilde{L}(a, z; \theta_{xy}) \{ f(a, b, x, y, 0) \}. \quad (2.2)$$

**Corollary 4.** Let  $f(b, x, y, z)$  be an 4-variable analytic function at  $(b, x, y, z) = (0, 0, 0, 0) \in \mathbb{C}^4$ . If  $f(b, x, y, z)$  satisfies the  $q$ -difference equation

$$(q^{-1}x - y) [f(b, x, y, z) - f(b, x, y, qz)] = z [f(b, q^{-1}x, y, qz) - f(b, x, qy, qz)], \quad (2.3)$$

then we have:

$$f(b, x, y, z) = \mathbb{E}(z\theta_{xy}) \{ f(b, x, y, 0) \}. \quad (2.4)$$

*Proof.* From the theory of several complex variables [10], we begin to solve the  $q$ -difference equation (2.3). First we may assume that

$$f(a, b, x, y, z) = \sum_{n=0}^{\infty} A_n(a, b, x, y) z^n. \quad (2.5)$$

Substituting (2.5) into (2.3), we get:

$$(q^{-1}x - y) \sum_{n=0}^{\infty} (1 - q^n) A_n(a, b, x, y) z^n = \sum_{n=0}^{\infty} q^n (1 - aq^n) [A_n(a, b, q^{-1}x, y) - A_n(a, b, x, qy)] z^{n+1}.$$

Comparing coefficients of  $z^n$ ,  $n \geq 1$ , we find that

$$(q^{-1}x - y)(1 - q^n) A_n(a, b, x, y) = q^{n-1} (1 - aq^{n-1}) [A_{n-1}(a, b, q^{-1}x, y) - A_{n-1}(a, b, x, qy)].$$

After simplification, we get:

$$A_n(a, b, x, y) = q^{n-1} \frac{1 - aq^{n-1}}{1 - q^n} \theta_{xy} \{A_{n-1}(a, b, x, y)\}.$$

By iteration, we gain

$$A_n(a, b, x, y) = q^{\binom{n}{2}} \frac{(a; q)_n}{(q; q)_n} \theta_{xy}^n [A_0(a, b, x, y)]. \quad (2.6)$$

Just taking  $z = 0$  in (2.5), we immediately obtain  $A_0(a, b, x, y) = f(a, b, x, y, 0)$ . Substituting (2.6) back into (2.5), we achieve (2.3).  $\square$

**Theorem 5** (Extended generating function for  $\Psi_n^{(a)}(x, y, z|q)$ ). For  $|yt| < 1$ , we have:

$$\sum_{n=0}^{\infty} \Psi_{n+k}^{(a)}(x, y, z|q) \frac{(-1)^{n+k} q^{\binom{n+k}{2}} t^n}{(q; q)_n} = t^{-k} \frac{(xt; q)_{\infty}}{(yt; q)_{\infty}} \sum_{n=0}^k \frac{(q^{-k}; q)_n q^n}{(xt, q; q)_n} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; ztq^n \right]. \quad (2.7)$$

**Corollary 6.** For  $|yt| < 1$ , we have:

$$\sum_{n=0}^{\infty} F_{n+k}(x, y, z; q) \frac{(-1)^{n+k} q^{\binom{n+k}{2}} t^n}{(q; q)_n} = t^{-k} \frac{(xt, zt; q)_{\infty}}{(yt; q)_{\infty}} {}_3\Phi_2 \left[ \begin{matrix} q^{-k}, yt, 0; \\ xt, zt; \end{matrix} q; q \right]. \quad (2.8)$$

**Remark 7.** For  $a = 0$ , (2.7) reduces (2.8). For  $a = 0$  and  $k = 0$ , (2.7) and (2.8) reduce to the generating function for  $\Psi_n^{(a)}(x, y, z|q)$

$$\sum_{n=0}^{\infty} \Psi_n^{(a)}(x, y, z|q) \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} = \frac{(xt; q)_{\infty}}{(yt; q)_{\infty}} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; zt \right], \quad |yt| < 1 \quad (2.9)$$

and [1, Theorem 2.6].

To prove the Theorem 5, the following Lemma is necessary.

**Lemma 8.**  $q$ -Chu-Vandermonde formula [5, Eq. (II.6)]

$${}_2\Phi_1 \left[ \begin{matrix} q^{-n}, a; \\ c; \end{matrix} q; q \right] = \frac{(c/a; q)_n}{(c; q)_n} a^n. \quad (2.10)$$

*Proof of Theorem 5.* Denoting the right-hand side of equation (2.7) by  $F(a, t, x, y, z)$ , we have:

$$F(a, t, x, y, z) = t^{-k} \sum_{n=0}^k \frac{(q^{-k}; q)_n q^n}{(q; q)_n} \frac{(xtq^n; q)_{\infty}}{(ytq^n; q)_{\infty}} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; ztq^n \right]. \quad (2.11)$$

Because equation (2.11) satisfies (2.3), we have:

$$\begin{aligned} F(a, t, x, y, z) &= \widetilde{L}(a, z; \theta_{xy}) \{F(a, t, x, y, 0)\} \\ &= \widetilde{L}(a, z; \theta_{xy}) \left\{ t^{-k} \sum_{n=0}^k \frac{(q^{-k}; q)_n q^n}{(q; q)_n} \frac{(xtq^n; q)_{\infty}}{(ytq^n; q)_{\infty}} \right\} \end{aligned}$$

$$\begin{aligned}
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xt; q)_\infty}{(yt; q)_\infty} t^{-k} \sum_{n=0}^k \frac{(q^{-k}, yt; q)_n q^n}{(xt, q; q)_n} \right\} \\
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xt; q)_\infty}{(yt; q)_\infty} t^{-k} {}_2\Phi_1 \left[ \begin{matrix} q^{-k}, yt; \\ q, q \end{matrix} \middle| xt; \right] \right\} \text{ by (2.10)} \\
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{p_k(y, x) (xt; q)_\infty}{(xt; q)_k (yt; q)_\infty} \right\} \text{ by (1.4)} \\
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ p_k(y, x) \sum_{n=0}^{\infty} \frac{p_n(y, xq^k) t^n}{(q; q)_n} \right\} \\
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ \sum_{n=0}^{\infty} p_{n+k}(y, x) \frac{t^n}{(q; q)_n} \right\} \\
 &= \sum_{n=0}^{\infty} \tilde{L}(a, z; \theta_{xy}) \left\{ (-1)^{n+k} q^{-\binom{n+k}{2}} p_{n+k}(y, x) \right\} (-1)^{n+k} q^{\binom{n+k}{2}} \frac{t^n}{(q; q)_n} \text{ by (1.15)} \\
 &= \sum_{n=0}^{\infty} \Psi_{n+k}^{(a)}(x, y, z|q) \frac{(-1)^{n+k} q^{\binom{n+k}{2}} t^n}{(q; q)_n},
 \end{aligned}$$

which is the left-hand side of (2.7). □

### 3. THE ROGERS FORMULA FOR $\Psi_n^{(a)}(x, y, z|q)$

In this section, we give Rogers formula and extended Rogers formula for the generalized trivariate  $q$ -Hahn polynomials  $\Psi_n^{(a)}(x, y, z|q)$  by using the homogeneous  $q$ -difference equations.

**Theorem 9** (Rogers's-type formula for  $\Psi_n^{(a)}(x, y, z|q)$ ). *We have:*

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \Psi_{n+m}^{(a)}(x, y, z|q) (-1)^{n+m} q^{\binom{n+m}{2}} \frac{t^n}{(q; q)_n} \frac{s^m}{(q; q)_m} = \frac{(xs; q)_\infty}{(t/s, ys; q)_\infty} \sum_{k=0}^{\infty} \frac{(ys; q)_k q^k}{(sq/t, xs, q; q)_k} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} \middle| q; zsq^k \right], \quad (3.1)$$

where  $\max\{|t/s|, |ys|\} < 1$ .

**Corollary 10.** *We have:*

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} F_{n+m}(x, y, z; q) (-1)^{n+m} q^{\binom{n+m}{2}} \frac{t^n}{(q; q)_n} \frac{s^m}{(q; q)_m} = \frac{(xs, zs; q)_\infty}{(t/s, ys; q)_\infty} {}_4\Phi_3 \left[ \begin{matrix} yt, 0, 0, 0; \\ sq/t, xs, zs; \end{matrix} \middle| q; q \right], \quad (3.2)$$

where  $\max\{|t/s|, |ys|\} < 1$ .

*Proof.* Denoting the right-hand side of equation (3.1) by  $G(a, s, x, y, z)$ , we have:

$$G(a, s, x, y, z) = \frac{1}{(t/s; q)_\infty} \sum_{k=0}^{\infty} \frac{q^k}{(sq/t, q; q)_k} \frac{(xsq^k; q)_\infty}{(ysq^k; q)_\infty} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} \middle| q; zsq^k \right]. \quad (3.3)$$

Because equation (3.3) satisfies (2.3), by (2.4), we have:

$$\begin{aligned}
 G(a, s, x, y, z) &= \tilde{L}(a, z; \theta_{xy}) \{G(a, s, x, y, 0)\} \\
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{1}{(t/s; q)_\infty} \sum_{k=0}^{\infty} \frac{q^k}{(sq/t, q; q)_k} \frac{(xsq^k; q)_\infty}{(ysq^k; q)_\infty} \right\} \\
 &= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} \sum_{k=0}^{\infty} \frac{(ys; q)_k q^k}{(xs, q; q)_k} \frac{1}{(t/s; q)_\infty (sq/t; q)_k} \right\}
 \end{aligned}$$

$$\begin{aligned}
&= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} \sum_{k=0}^{\infty} \frac{(ys; q)_k q^k (-t/s)^k q^{-\binom{k}{2}-k}}{(xs, q; q)_k (tq^{-k}/s; q)_\infty} \right\} \\
&= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} \sum_{k=0}^{\infty} \frac{(ys; q)_k q^k (-t/s)^k q^{-\binom{k}{2}-k}}{(xs, q; q)_k} \sum_{n=0}^{\infty} \frac{(t/s)^n q^{-nk}}{(q; q)_n} \right\} \\
&= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} \sum_{k=0}^{\infty} \frac{(ys; q)_k q^k}{(xs, q; q)_k} \sum_{n=k}^{\infty} \frac{(-t/s)^n q^{\binom{k}{2}-nk}}{(q; q)_{n-k}} \right\} \\
&= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} \sum_{n=0}^{\infty} \frac{(t/s)^n}{(q; q)_n} \sum_{k=0}^n \frac{(q^{-n}, ys; q)_k q^k}{(xs, q; q)_k} \right\} \\
&= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} \sum_{n=0}^{\infty} \frac{(t/s)^n}{(q; q)_n} {}_2\Phi_1 \left[ \begin{matrix} q^{-n}, ys; \\ xs; \end{matrix} \middle| q; q \right] \right\} \\
&= \sum_{n=0}^{\infty} \frac{t^n}{(q; q)_n} \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(xs; q)_\infty}{(ys; q)_\infty} s^{-n} {}_2\Phi_1 \left[ \begin{matrix} q^{-n}, ys; \\ xs; \end{matrix} \middle| q; q \right] \right\} \text{ by (2.10)} \\
&= \sum_{n=0}^{\infty} \frac{t^n}{(q; q)_n} \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{p_n(y, x) (xs; q)_\infty}{(xs; q)_n (ys; q)_\infty} \right\} \text{ by (2.12)} \\
&= \sum_{n=0}^{\infty} \frac{t^n}{(q; q)_n} \sum_{m=0}^{\infty} \Psi_{n+m}^{(a)}(x, y, z|q) (-1)^{n+m} q^{\binom{n+m}{2}} \frac{s^m}{(q; q)_m},
\end{aligned}$$

which is the left-hand side of (3.1).  $\square$

**Theorem 11** (Extended Roger's-type formula for  $\Psi_n^{(a)}(x, y, z|q)$ ). *We have:*

$$\begin{aligned}
&\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \Psi_{n+m+k}^{(a)}(x, y, z|q) \frac{(-1)^{n+m+k} q^{\binom{n+m+k}{2}} t^n s^m \omega^k}{(q; q)_{n+m} (q; q)_m (q; q)_k} \\
&= \frac{(x\omega; q)_\infty}{(s/t, t/\omega, y\omega; q)_\infty} \sum_{j=0}^{\infty} \frac{(y\omega; q)_j q^j}{(x\omega, q\omega/t, q; q)_j} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} \middle| q; z\omega q^j \right], \quad (3.4)
\end{aligned}$$

where  $\max\{|s/t|, |t/\omega|, |y\omega|\} < 1$ .

**Remark 12.** For  $s = 0$ , (3.4) reduces to (3.1).

*Proof of Theorem 11.* Denoting the right-hand side of equation (3.4) by  $H(a, \omega, x, y, z)$ , we have:

$$H(a, \omega, x, y, z) = \frac{1}{(s/t, t/\omega; q)_\infty} \sum_{k=0}^{\infty} \frac{q^k}{(q\omega/t, q; q)_k} \cdot \frac{(x\omega q^k; q)_\infty}{(y\omega q^k; q)_\infty} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} \middle| q; z\omega q^k \right]. \quad (3.5)$$

Because equation (3.5) satisfies (2.3), by (2.4), we have:

$$\begin{aligned}
H(a, \omega, x, y, z) &= \tilde{L}(a, z; \theta_{xy}) \{H(a, \omega, x, y, 0)\} \\
&= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{1}{(s/t, t/\omega; q)_\infty} \sum_{k=0}^{\infty} \frac{q^k}{(q\omega/t, q; q)_k} \frac{(x\omega q^k; q)_\infty}{(y\omega q^k; q)_\infty} \right\} \\
&= \frac{1}{(s/t; q)_\infty} \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{1}{(t/\omega; q)_\infty} \sum_{k=0}^{\infty} \frac{q^k}{(q\omega/t, q; q)_k} \frac{(x\omega q^k; q)_\infty}{(y\omega q^k; q)_\infty} \right\} \\
&= \frac{1}{(s/t; q)_\infty} \tilde{L}(a, z; \theta_{xy}) \{F(a, \omega, x, y, 0)\} \text{ by (2.10)}
\end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{(s/t; q)_\infty} \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \Psi_{m+k}^{(a)}(x, y, z|q) \frac{(-1)^{m+k} q^{\binom{m+k}{2}} t^m}{(q; q)_m} \frac{\omega^k}{(q; q)_k} \\
 &= \sum_{n=0}^{\infty} \frac{(s/t)^n}{(q; q)_n} \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \Psi_{m+k}^{(a)}(x, y, z|q) \frac{(-1)^{m+k} q^{\binom{m+k}{2}} t^m}{(q; q)_m} \frac{\omega^k}{(q; q)_k} \\
 &= \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \sum_{n=m}^{\infty} \Psi_{n+m}^{(a)}(x, y, z|q) \frac{(-1)^{m+k} q^{\binom{m+k}{2}} t^{m-n}}{(q; q)_m} \frac{s^n}{(q; q)_n} \frac{\omega^k}{(q; q)_k}. \tag{3.6}
 \end{aligned}$$

Replacing  $m$  by  $m + n$  in (3.6), we obtain:

$$H(a, \omega, x, y, z) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \Psi_{n+m+k}^{(a)}(x, y, z|q) \frac{(-1)^{n+m+k} q^{\binom{n+m+k}{2}} t^n s^m \omega^k}{(q; q)_m (q; q)_{n+m} (q; q)_k},$$

which is the left-hand side of (3.4). □

#### 4. SRIVASTAVA-AGARWAL TYPE GENERATING FUNCTIONS FOR GENERALIZED TRIVARIATE $q$ -HAHN POLYNOMIALS

The Hahn polynomials [7, 8] (or Al-Salam and Carlitz polynomials [2]) are defined as

$$\Phi_n^{(a)}(x|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (a; q)_k x^k. \tag{4.1}$$

Srivastava and Agarwal gave the following generating function.

**Lemma 13.** [15, Eq. (3.20)] *Suppose that  $\max\{|t|, |xt|\} < 1$ , we have:*

$$\sum_{n=0}^{\infty} \Phi_n^{(a)}(x|q) \frac{(\lambda; q)_n t^n}{(q; q)_n} = \frac{(\lambda t; q)_\infty}{(t; q)_\infty} {}_2\Phi_1 \left[ \begin{matrix} \lambda, a; \\ q, xt \end{matrix} \right]. \tag{4.2}$$

The generating function (4.2) is called Srivastava-Agarwal type generating functions for the Al-Salam-Carlitz polynomials [15].

In this section, we give Srivastava-Agarwal type generating function for the generalized trivariate  $q$ -Hahn polynomials  $\Psi_n^{(a)}(x, y, z|q)$  by the homogeneous  $q$ -difference equation.

**Theorem 14.** *For  $|yvt| < 1$ , we have:*

$$\sum_{n=0}^{\infty} \Psi_n^{(a)}(x, y, z|q) p_n(\nu, \mu) \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} = \frac{(\mu/\nu, xvt; q)_\infty}{(yvt; q)_\infty} \sum_{n=0}^{\infty} \frac{(yvt; q)_n (\mu/\nu)^n}{(xvt, q; q)_n} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; zvtq^n \right]. \tag{4.3}$$

**Corollary 15.** *For  $|yvt| < 1$ , we have:*

$$\sum_{n=0}^{\infty} F_n(x, y, z; q) p_n(\nu, \mu) \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} = \frac{(\mu/\nu, xvt, zvt; q)_\infty}{(yvt; q)_\infty} {}_3\Phi_2 \left[ \begin{matrix} yvt, 0, 0; \\ xvt, zvt; \end{matrix} q; \frac{\mu}{\nu} \right]. \tag{4.4}$$

**Corollary 16.** *For  $|yt| < 1$ , we have:*

$$\sum_{n=0}^{\infty} \Psi_n^{(a)}(x, y, z|q) (\lambda; q)_n \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} = \frac{(\lambda, xt; q)_\infty}{(yt; q)_\infty} \sum_{n=0}^{\infty} \frac{(yt; q)_n \lambda^n}{(xt, q; q)_n} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; ztq^n \right]. \tag{4.5}$$

**Remark 17.** *For  $\nu = 1$ , (4.4) reduces to (4.5).*

**Corollary 18.** For  $|axt| < 1$ , we have:

$$\sum_{n=0}^{\infty} \psi_n^{(a)}(x, y|q)(\lambda; q)_n \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} = \frac{(\lambda, xt, yt; q)_{\infty}}{(axt; q)_{\infty}} {}_3\Phi_2 \left[ \begin{matrix} axt, 0, 0; \\ xt, yt; \end{matrix} q; \lambda \right] \quad (4.6)$$

and

$$\sum_{n=0}^{\infty} \psi_n^{(a)}(x|q)(\lambda; q)_n \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} = \frac{(\lambda, xt, t; q)_{\infty}}{(axt; q)_{\infty}} {}_3\Phi_2 \left[ \begin{matrix} axt, 0, 0; \\ xt, t; \end{matrix} q; \lambda \right]. \quad (4.7)$$

**Remark 19.** Setting  $v = 1$ ,  $a = 0$ ,  $y = ax$  and  $z = y$  in (4.4), we get (4.6). For  $v = 1$ ,  $a = 0$ ,  $y = ax$  and  $z = 1$ , (4.4) reduces to (4.7). For  $y = 1$ , (4.6) reduces to (4.7).

Before we prove the Theorem 15, the following Lemma is necessary.

**Lemma 20.** [6, Eq. (III.1)] For  $\{|c|, |z|, |b|\} < 1$ , we have:

$${}_2\Phi_1 \left[ \begin{matrix} a, b; \\ c; \end{matrix} q; z \right] = \frac{(b, az; q)_{\infty}}{(c, z; q)_{\infty}} {}_2\Phi_1 \left[ \begin{matrix} c/b, z; \\ az; \end{matrix} q; b \right]. \quad (4.8)$$

*Proof of Theorems 15.* Denoting the right-hand side of equation (4.4) by  $H'(a, t, x, y, z)$ , we have:

$$H'(a, t, x, y, z) = (\mu/v; q)_{\infty} \sum_{n=0}^{\infty} \frac{(xvtq^n; q)_{\infty} (\mu/v)^n}{(yvtq^n; q)_{\infty} (q; q)_n} {}_1\Phi_1 \left[ \begin{matrix} a; \\ 0; \end{matrix} q; zvtq^n \right]. \quad (4.9)$$

Because equation (4.9) satisfies (2.3), by (2.4), we have:

$$\begin{aligned} H'(a, t, x, y, z) &= \tilde{L}(a, z; \theta_{xy}) \{H'(a, t, x, y, 0)\} \\ &= \tilde{L}(a, z; \theta_{xy}) \left\{ (\mu/v; q)_{\infty} \sum_{n=0}^{\infty} \frac{(xvtq^n; q)_{\infty} (\mu/v)^n}{(yvtq^n; q)_{\infty} (q; q)_n} \right\} \\ &= \tilde{L}(a, z; \theta_{xy}) \left\{ \frac{(\mu/v, xvt; q)_{\infty}}{(yvt; q)_{\infty}} {}_2\Phi_1 \left[ \begin{matrix} yvt, 0; \\ xvt; \end{matrix} q; \frac{\mu}{v} \right] \right\}. \end{aligned}$$

By using (4.8) and (1.3), the last relation becomes

$$\begin{aligned} H'(a, t, x, y, z) &= \tilde{L}(a, z; \theta_{xy}) \left\{ \sum_{n=0}^{\infty} p_n(y, x) \frac{p_n(v, \mu) t^n}{(q; q)_n} \right\} \\ &= \sum_{n=0}^{\infty} \tilde{L}(a, z; \theta_{xy}) \left\{ (-1)^n q^{-\binom{n}{2}} p_n(y, x) \right\} p_n(v, \mu) \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n} \text{ by (1.15)} \\ &= \sum_{n=0}^{\infty} \Psi_n^{(a)}(x, y, z|q) p_n(v, \mu) \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q; q)_n}, \end{aligned}$$

which is the left-hand side of (4.4). This achieves the proof.  $\square$

#### ACKNOWLEDGMENTS

The author would like to thank the referees and editors for their many valuable comments and suggestions. This work was supported by the Zhejiang Provincial Natural Science Foundation of China (No. LY21A010019).

## REFERENCES

- [1] M. A. Abdhusein, *Two operator representations for the trivariate  $q$ -polynomials and Hahn polynomials*, Ramanujan J. **40** (2016), 491–509.
- [2] A. Al-Salam and L. Carlitz, *Some orthogonal  $q$ -polynomials*, Math. Nachr. **30** (1965), 47–61.
- [3] J. Cao, *On Carlitz's trilinear generating functions*, Appl. Math. Comput. **218** (2012), 9839–9847.
- [4] W. Y. C. Chen, A. M. Fu and B. Zhang, *The homogeneous  $q$ -difference operator*, Adv. App. Math. **31** (2003), 659–668.
- [5] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd edn. Cambridge University Press, Cambridge, 2004.
- [6] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Cambridge University Press, Cambridge, 1990.
- [7] W. Hahn, *Über Orthogonalpolynome, die  $q$ -Differenzgleichungen*, Math. Nachr. **2** (1949), 434.
- [8] W. Hahn, *Beitrage zur Theorie der Heineschen Reihen; Die 24 Integrale der hypergeometrischen  $q$ -Differenzgleichung; Das  $q$ -Analogon der Laplace-Transformation*, Math. Nachr. **2** (1949), 340–379.
- [9] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its  $q$ -analogue report*, Delft University of Technology, 1998.
- [10] R. M. Range, *Complex analysis: A brief tour into higher dimensions*, Amer. Math. Mon. **110** (2003), 89–108.
- [11] H. L. Saad and A. A. Sukhi, *Another homogeneous  $q$ -difference operator*, Appl. Math. Comput. **215** (2010), 4332–4339.
- [12] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge Univ. Press, Cambridge/London/New York, 1966.
- [13] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted (Ellis Horwood, Chichester); Wiley, New York, 1985.
- [14] H. M. Srivastava, S. Arjika and A. Sherif Kelil, *Some homogeneous  $q$ -difference operators and the associated generalized Hahn polynomials*, Appl. Set-Valued Anal. Optim. **1**(2019), 187–201.
- [15] H. M. Srivastava and A.K. Agarwal, *Generating functions for a class of  $q$ -polynomials*, Ann. Mat. Pure Appl. (Ser. 4) **154** (1989), 99–109.