

QUADRATIC AND SYMPLECTIC STRUCTURES ON 3-(HOM)- ρ -LIE ALGEBRAS

ZAHRA BAGHERI ¹ AND ESMAEIL PEYGHAN ^{2,*}

^{1,2} DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, ARAK UNIVERSITY,
ARAK, 38156-8-8349, IRAN

*correspondence : e – peyghan@araku.ac.ir

ABSTRACT. Our purpose in this paper is the generalization of the notions of quadratic and symplectic structures to the case of 3-(Hom)- ρ -Lie algebras. We describe some properties of them by expressing the related lemmas and theorems. Also, we introduce the concept of 3-pre-(Hom)- ρ -Lie algebras and define their representation.

1. INTRODUCTION

At the first, Hartwig, Larsson and Silvestrov in [24] generalized Lie algebras and constructed a new class of Lie algebras in the name of Hom-Lie algebras. These algebras very soon were attracted considerable attentions and many algebraic and geometric structures were introduced on them. In 1994, the other concept was emerged as " ρ -Lie algebra" or "Lie color algebra" by P.J. Bongaarts [16] and then in 1998, Scheunert and Zhang introduced a new notion which is called a cohomology theory of Lie color algebras [32]. Also, the notion of Hom-Lie color algebras, as an extension of Hom-Lie superalgebras to G -graded algebras, was introduced by Yuan [38]. These Lie algebras were applied by the researchers and in 2015, Abdaoui, Ammar and Makhlof defined representations and a cohomology of Hom-Lie color algebras in [1]. After two years, in 2017, B. Sun et al. studied some concepts such as T^* -extension and abelian extension on Hom-Lie color algebras [35].

The notion of pre-Lie algebra was given by Gerstenhaber in [22] in the study of deformations and the cohomology theory of associative algebras. These algebras in the other research articles appear with titles such as left-symmetric algebras or right-symmetric algebras. Also, the notion of Hom-pre-Lie algebra was introduced in [28]. This kind of algebras in the construction of Hom-Lie 2-algebras has an important role (see [33]). Representation of a Hom-pre-Lie algebra was constructed in [34], in the study of Hom-pre-Lie bialgebras. The geometrization and the bialgebra theory of these algebras was studied in [34, 40]. For the commutator bracket $[\cdot, \cdot]_c$, every Hom-pre-Lie algebra (A, \cdot, α) gives rise to a Hom-Lie algebra $(A, [\cdot, \cdot]_c, \alpha)$. This Lie algebra is called the subadjacent Hom-Lie algebra and denoted by A^c . Actually, we can say the Hom-pre-Lie algebras have a interconnected with the Hom-Lie algebras.

The concept of n -Lie algebra was introduced by Filippov in 1985. These Lie algebras are known by different names such as Filippov algebras, Nambu-Lie algebras, Lie n -algebras. The special case of these algebras are 3-Lie algebras and in the Hom case 3-Hom-Lie algebras, which extracted from 3-Lie algebras base on Hom-Lie algebras. These algebras were defined and checked by H. Ataguema et al. in [7]. Y. Liu et al. took into consideration these algebras and studied the representations and module-extensions on them in [27]. Later, the new notions such as 3-Lie color algebra, 3-Hom-Lie color algebra and 3-pre-Lie

2010 *Mathematics Subject Classification.* 15A63, 17B10, 16W25, 17B40, 17B70, 17B75, 37J10.

Key words and phrases. 3- ρ -Lie algebra, 3-pre- ρ -Lie algebra, quadratic structure, representation, symplectic structure.

algebra were constructed and studied in [12, 31, 39]. In 1996, Y. Daletskii and V. Kushnirenich introduced the notion of n -Lie superalgebra, as a natural generalization of n -Lie algebra ([18]). When $n = 3$, the 3-Lie superalgebras are a particular case of 3- ρ -Lie algebras when the abelian group is \mathbb{Z}_2 (see for more details [2, 3, 23]).

Ternary Lie algebras have vast applications in various realms of modern physics and mechanics such as Nambu, field and Hamiltonian theories. For example, in Nambu consideration for a case that the algebra of functions in a manifold M is ternary, one can construct a bracket defined by a trivector on M that is famous as Nambu bracket in literature. In this way, the dynamics of M is controlled by two Hamiltonians in return of classical case. Indeed, 3-ary bracket in ternary Lie algebra is a suitable alternative for Poisson alternative in Hamiltonian mechanics. To study how Ternary Lie algebras can model this frameworks, see [21, 29, 36]. Nahm equations describing some facts in string and superstring theories can be formulized using ternary Lie algebras also. In [13] it is described the benefits of replacing ternary Lie algebras instead of using simple Lie algebras in lifted Nahm equations and in [9] the authors for the lifted Nahm equations replaced the Lie algebra in the Nahm equation by a 3-Lie algebra. In the framework of Bagger-Lambert Gustavsson model of multiple $M2$ branes, it is possible to revisit the super-symmetric alternative of world-volume theory of M -theory membrane in special cases such $N = 2$. See [9, 10, 25, 30] for more details in the way.

In 2006, quadratic Lie algebras were discussed by I. Bajo et al. in [11]. A quadratic Lie algebra comes equipped with a non-degenerate, symmetric and invariant bilinear form. Quadratic Lie algebras are important in Mathematics and Physics specially in conformal field theory [20]. S. Benayadi and A. Makhlouf, in 2010 studied the quadratic Hom-Lie algebras [15] as a generalization of the classical case. This process continued until a generalization to the case of quadratic (even quadratic) color Lie algebras was obtained in [37]. Also, quadratic color Hom-Lie algebras were introduced and studied by F. Ammar et al. in [5]. Also, the quadratic-algebraic structures have been studied on the class of Γ -graded Lie algebras specially in the case where $\Gamma = \mathbb{Z}_2$. These algebras are called homogeneous (even or odd) quadratic Lie superalgebras (see [4, 8, 14, 32]).

In this paper, we discuss the quadratic and symplectic structures on 3-(Hom)- ρ -Lie algebras and study the representations of them. The other aim is to introduce the notion of 3-pre-(Hom)- ρ -Lie algebras and check their phase spaces. Also, we study the relationship between the 3-pre-(Hom)- ρ -Lie algebras and 3-(Hom)- ρ -Lie algebras.

This paper is arranged as follows: In Section 2, we give a brief introduction to 3- ρ -Lie algebras and study the quadratic and symplectic structures on 3- ρ -Lie algebras. Also, we study the metrics and derivations on 3- ρ -Lie algebras and we explore the relationship between the symplectic structure and the invertible derivation which is antisymmetric with respect to a metric. Section 3 is devoted to study of representation theory on 3- ρ -Lie algebras and some classification results will be given. In Section 4, we introduce the notion of 3-pre- ρ -Lie algebras and discuss some basic properties of 3-pre- ρ -Lie algebras including representation theory and some classification results. Also, we define ρ - \mathcal{O} -operator associated to the representation (V, μ) of a 3- ρ -Lie algebra and construct a 3- ρ -pre-Lie algebra structure on the representation V by a ρ - \mathcal{O} -operator and bilinear map μ . This section also contains the notion of a phase space of a 3- ρ -Lie algebra and it will be shown that a 3- ρ -Lie algebra has a phase space if and only if it is sub-adjacent to a 3- ρ -pre-Lie algebra. In Section 5, we study the constructions of Section 2, such as quadratic, symplectic structures, 3-associative Lie algebras in the case of 3-Hom- ρ -lie algebras. The

representation theory of 3-Hom- ρ -lie algebras is also included in this section. Section 6 is devoted to the 3-pre-Hom- ρ -Lie algebras, which are studied similar to the classical case in Section 4. Actually, we give the similar results to Section 4 in the case of 3-pre-Hom- ρ -Lie algebras.

2. QUADRATIC 3- ρ -LIE ALGEBRAS

We begin this section by describing the class of ρ -Lie algebras, which includes the special multiplication that is known as bracket and a two-cycle. This Lie algebras is called 3- ρ -Lie algebra. We summarize some definitions and introduce the notions such as quadratic structure, symplectic structure ω , metric φ , derivation D and φ -antisymmetric derivation D on 3- ρ -Lie algebras and show that a symplectic structure ω may be defined on a 3- ρ -Lie algebra if and only if there exists a φ -antisymmetric invertible derivation D . Also, We give a brief exposition of 3-associative algebras and give the constructions of direct product 3- ρ -Lie algebra structures from 3-associative algebras.

Let us consider $(G, +)$ as an abelian group over a field \mathbb{K} ($\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$). A map $\rho : G \times G \longrightarrow \mathbb{K}^*$ with the properties

$$\rho(a, b) = \rho(b, a)^{-1}, \quad a, b \in G, \quad (2.1)$$

$$\rho(a + b, c) = \rho(a, c)\rho(b, c), \quad a, b, c \in G, \quad (2.2)$$

is called a two-cycle. From two above properties we can deduce $\rho(a, b) \neq 0$, $\rho(0, b) = 1$ and $\rho(c, c) = \pm 1$ for all $a, b, c \in G$, $c \neq 0$. Note that if \mathcal{B} is a G -graded vector space ($\mathcal{B} = \bigoplus_{a \in G} \mathcal{B}_a$), an element $f \in \mathcal{B}_a$ is called homogeneous of G -degree a . We denote by $Hg(\mathcal{B})$ the set of homogeneous elements in \mathcal{B} and by $|f|$ the G -degree of $f \in Hg(\mathcal{B})$

A ρ -Lie algebra is a triple $(\mathcal{B}, [\cdot, \cdot]_{\mathcal{B}}, \rho)$, where

- i. \mathcal{B} is a G -graded vector space,
- ii. ρ is a two-cycle,
- iii. $[\cdot, \cdot]_{\mathcal{B}}$ is a bilinear map on \mathcal{B} such that

- $|[f, g]_{\rho}| = |f| + |g|$,
- $[f, g]_{\rho} = -\rho(|f|, |g|)[g, f]_{\rho}$,
- $\rho(|h|, |f|)[f, [g, h]_{\rho}]_{\rho} + \rho(|g|, |h|)[h, [f, g]_{\rho}]_{\rho} + \rho(|f|, |g|)[g, [h, f]_{\rho}]_{\rho} = 0$.

The third condition is equivalent to

$$[f, [g, h]_{\rho}]_{\rho} = [[f, g]_{\rho}, h]_{\rho} + \rho(|f|, |g|)[g, [f, h]_{\rho}]_{\rho}.$$

In the next, for simplicity we use $\rho(f, g)$ instead of $\rho(|f|, |g|)$.

Definition 2.1. A triple $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho)$ is called a 3- ρ -Lie algebra if \mathcal{B} is a G -graded vector space, $[\cdot, \cdot, \cdot]_{\mathcal{B}} : \mathcal{B} \times \mathcal{B} \times \mathcal{B} \longrightarrow \mathcal{B}$ is a 3-linear map, ρ is a two-cycle and moreover the following conditions are satisfied

- (1) $|[f_1, f_2, f_3]_{\mathcal{B}}| = |f_1| + |f_2| + |f_3|$,
- (2) $[f_1, f_2, [g_1, g_2, g_3]_{\mathcal{B}}]_{\mathcal{B}} = [[f_1, f_2, g_1]_{\mathcal{B}}, g_2, g_3]_{\mathcal{B}} + \rho(f_1 + f_2, g_1)[g_1, [f_1, f_2, g_2]_{\mathcal{B}}, g_3]_{\mathcal{B}} + \rho(f_1 + f_2, g_1 + g_2)[g_1, g_2, [f_1, f_2, g_3]_{\mathcal{B}}]_{\mathcal{B}}$, (ρ -fundamental identity)
- (3) $[f_1, f_2, f_3]_{\mathcal{B}} = -\rho(f_1, f_2)[f_2, f_1, f_3]_{\mathcal{B}} = -\rho(f_2, f_3)[f_1, f_3, f_2]_{\mathcal{B}}$. (ρ -skew-symmetry property)

We denote the 3- ρ -Lie algebra $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho)$ by \mathcal{B}_{ρ} .

Example 2.2. One of the most important case of 3- ρ -Lie algebras are 3-ary Lie superalgebras when abelian group G is \mathbb{Z}_2 .

In the following, we extend to 3- ρ -Lie algebras the study of quadratic color Hom-Lie algebras introduced in [5] and in the next of this paper we investigate Hom version. We apply a non-degenerate bilinear form on 3- ρ -Lie algebra \mathcal{B}_{ρ} to define a quadratic structure.

Definition 2.3. Let $\chi : \mathcal{B}_{\rho} \times \mathcal{B}_{\rho} \longrightarrow \mathbb{K}$ be a non-degenerate bilinear form on \mathcal{B}_{ρ} . Then χ is called a quadratic structure of \mathcal{B}_{ρ} if, for any $f_1, f_2, f_3, f_4 \in \text{Hg}(\mathcal{B}_{\rho})$, the following conditions are satisfied

- i. $\chi(f_1, f_2) = \rho(f_1, f_2)\chi(f_2, f_1)$ (ρ -symmetric),
- ii. $\chi([f_1, f_2, f_3]_{\mathcal{B}}, f_4) = \chi(f_1, [f_2, f_3, f_4]_{\mathcal{B}})$ (invariant).

Also $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \chi)$ is said to be a quadratic 3- ρ -Lie algebra if χ is a quadratic structure of \mathcal{B} . We use the symbol $\mathcal{B}_{\rho, \chi}$ to represent any quadratic 3- ρ -Lie algebra $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \chi)$.

Definition 2.4. Let $\text{End}(\mathcal{B}_{\rho})$ be the endomorphism algebra of \mathcal{B}_{ρ} . A vector subspace $\text{Cent}(\mathcal{B}_{\rho})$ of $\text{End}(\mathcal{B}_{\rho})$ is defined by

$$\text{Cent}(\mathcal{B}_{\rho}) = \{\psi \in \text{End}(\mathcal{B}_{\rho}) : \psi([f, g, h]_{\mathcal{B}}) = [\psi(f), g, h]_{\mathcal{B}} = \rho(\psi, f)[f, \psi(g), h]_{\mathcal{B}} = \rho(\psi, f + g)[f, g, \psi(h)]_{\mathcal{B}}\}. \quad (2.3)$$

$\text{Cent}(\mathcal{B}_{\rho})$ is a subalgebra of $\text{End}(\mathcal{B}_{\rho})$ and it is called the centroid of \mathcal{B}_{ρ} .

In the following proposition, we construct 3- ρ -Lie algebras from a 3- ρ -Lie algebra and an even element in its centroid.

Proposition 2.5. Let $\psi \in \text{Cent}(\mathcal{B}_{\rho})$ be an even element. If we define two multiplications $[\cdot, \cdot, \cdot]^{\psi}$ and $[\cdot, \cdot, \cdot]_{\psi}^{\psi}$ on \mathcal{B} by

$$[f, g, h]^{\psi} = [\psi(f), g, h]_{\mathcal{B}}, \quad [f, g, h]_{\psi}^{\psi} = [\psi(f), \psi(g), \psi(h)]_{\mathcal{B}}, \quad \forall f, g, h \in \mathcal{B}_{\rho},$$

then we have two 3- ρ -Lie algebras $(\mathcal{B}, [\cdot, \cdot, \cdot]^{\psi}, \rho)$ and $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\psi}^{\psi}, \rho)$.

Proof. The definition of the bracket $[\cdot, \cdot, \cdot]^{\psi}$ and this fact that ψ is an even element in $\text{Cent}(\mathcal{B}_{\rho})$, lead to

- (a) $[f_1, f_2, [g_1, g_2, g_3]^{\psi}]^{\psi} = \psi^2[f_1, f_2, [g_1, g_2, g_3]_{\mathcal{B}}]_{\mathcal{B}}$,
- (b) $[[f_1, f_2, g_1]^{\psi}, g_2, g_3]^{\psi} = \psi^2[[f_1, f_2, g_1]_{\mathcal{B}}, g_2, g_3]_{\mathcal{B}}$,
- (c) $\rho(f_1 + f_2, g_1)[g_1, [f_1, f_2, g_2]^{\psi}, g_3]^{\psi} = \rho(f_1 + f_2, g_1)\psi^2[g_1, [f_1, f_2, g_2]_{\mathcal{B}}, g_3]_{\mathcal{B}}$,
- (d) $\rho(f_1 + f_2, g_1 + g_2)[g_1, g_2, [f_1, f_2, g_3]^{\psi}]^{\psi} = \rho(f_1 + f_2, g_1 + g_2)\psi^2[g_1, g_2, [f_1, f_2, g_3]_{\mathcal{B}}]_{\mathcal{B}}$.

According to the above equalities and this fact that \mathcal{B}_{ρ} is a 3- ρ -Lie algebra, it is easy to show that $[\cdot, \cdot, \cdot]^{\psi}$ is a 3- ρ -Lie algebra structure. Also, by the similar way, we can show that $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\psi}^{\psi}, \rho)$ is a 3- ρ -Lie algebra. \square

Proposition 2.6. Consider $\mathcal{B}_{\rho, \chi}$ and an even invertible χ -symmetric element $\psi \in \text{Cent}(\mathcal{B}_{\rho})$. If we define the even bilinear form χ_{ψ} by $\chi_{\psi}(f, g) = \chi(\psi(f), g)$, then we have two quadratic 3- ρ -Lie algebras $(\mathcal{B}, [\cdot, \cdot, \cdot]^{\psi}, \rho, \chi_{\psi})$ and $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\psi}^{\psi}, \rho, \chi_{\psi})$.

Proof. χ_ψ is a non-degenerate and ρ -symmetric bilinear form, since ψ is invertible and χ is a non-degenerate, ρ -symmetric bilinear form on \mathcal{B} . Then, it is enough to show that χ_ψ is invariant. For this, we have

$$\begin{aligned}\chi_\psi([f_1, f_2, f_3]^\psi, f_4) &= \chi(\psi[\psi(f_1), f_2, f_3]_A, f_4) = \chi([\psi(f_1), f_2, f_3], \psi(f_4)) \\ &= \chi(\psi(f_1), [f_2, f_3, \psi(f_4)]_A) = \chi_\psi(f_1, [f_2, f_3, f_4]^\psi).\end{aligned}$$

Also, we can show that $\chi_\psi([f_1, f_2, f_3]^\psi, f_4) = \chi_\psi(f_1, [f_2, f_3, f_4]^\psi)$, where f_1, f_2, f_3, f_4 are homogeneous elements of \mathcal{B} . \square

Definition 2.7. A 3-associative algebra is a pair (g, μ) consisting of a G -graded vector space g and an even 3-linear map $\mu : g \times g \times g \longrightarrow g$ (i.e. $\mu(g_a, g_b, g_c) \subset g_{a+b+c}$), such that for all $a, b, c, d, e \in g$

$$\mu(a, b, \mu(c, d, e)) = \mu(\mu(a, b, c), d, e).$$

If ρ is a two-cycle on G and the linear map μ is the ρ -symmetry with respect to the displacement of every two elements, the 3-associative algebra (g, μ) is called commutative and denoted by $g_{\rho, C}$.

Definition 2.8. A quadratic 3-associative algebra is a 3-associative algebra (g, μ) equipped with a ρ -symmetric, invariant and non-degenerate bilinear form χ_g . We denote a quadratic 3-associative algebra (g, μ, χ_g) and a commutative quadratic 3-associative algebra respectively by g_{χ_g} and g_{ρ, C, χ_g} .

Note that if A and A' are two G -graded vector spaces, then $A \otimes A'$ is also a G -graded vector space such that

$$A \otimes A' = \bigoplus_{\gamma \in G} (A \otimes A')_\gamma,$$

where $(A \otimes A')_\gamma = A_a \otimes A'_{a'}$ and so $f \otimes g \in (A \otimes A')_\gamma$ has the degree $|f \otimes g| = \gamma = a + a'$.

According to the above point, we have the following theorem, which gives us a quadratic 3- ρ -Lie algebra structure starting from a quadratic 3- ρ -Lie algebra and a quadratic commutative 3-associative algebra.

Theorem 2.9. Consider $\mathcal{B}_{\rho, \chi}$ and g_{ρ, C, χ_g} . The tensor product $(E = \mathcal{B} \otimes g, [\cdot, \cdot, \cdot]_E, \rho, \chi_E)$ is a quadratic 3- ρ -Lie algebra, such that $[\cdot, \cdot, \cdot]_E$ and χ_E are given by

$$[f \otimes a, g \otimes b, h \otimes c]_E = \rho(a, g + h)\rho(b, h)[f, g, h]_{\mathcal{B}} \otimes \mu(a, b, c), \quad (2.4)$$

$$\chi_E(f \otimes a, g \otimes b) = \rho(a, g)\chi(f, g)\chi_g(a, b), \quad (2.5)$$

for all $f, g, h \in Hg(\mathcal{B}_{\rho, \chi})$ and $a, b, c \in Hg(g_{\rho, \chi})$.

Proof. Using (2.4) and the commutative property of product μ , we get

$$\begin{aligned}
[f \otimes a, g \otimes b, [h_1 \otimes c, h_2 \otimes d, h_3 \otimes e]_E]_E &= \rho(a+b, h_1+h_2+h_3)\rho(c, h_2+h_3)\rho(d, h_3)\rho(a, g) \\
&\quad [f, g, [h_1, h_2, h_3]_{\mathcal{B}}]_{\mathcal{B}} \otimes \mu(a, b, \mu(c, d, e)), \\
[[f \otimes a, g \otimes b, h_1 \otimes c]_E, h_2 \otimes d, h_3 \otimes e]_E &= \rho(a+b, h_1+h_2+h_3)\rho(c, h_2+h_3)\rho(d, h_3)\rho(a, g) \\
&\quad [[f, g, h_1]_{\mathcal{B}}, h_2, h_3]_{\mathcal{B}} \otimes \mu(a, b, \mu(c, d, e)), \\
\rho(f+a+g+b, h_1+c)[h_1 \otimes c, [f \otimes a, g \otimes b, h_2 \otimes d]_E, h_3 \otimes e]_E &= \rho(a+b, h_1+h_2+h_3)\rho(c, h_2+h_3) \\
&\quad \rho(d, h_3)\rho(a, g)\rho(f+g, h_1)[h_1, [f, g, h_2]_{\mathcal{B}}, h_3]_{\mathcal{B}} \otimes \mu(a, b, \mu(c, d, e)), \\
\rho(f+a+g+b, h_1+c+h_2+d)[h_1 \otimes c, h_2 \otimes d, [f \otimes a, g \otimes b, h_3 \otimes e]_E]_E &= \rho(a+b, h_1+h_2+h_3) \\
&\quad \rho(c, h_2+h_3)\rho(d, h_3)\rho(a, g)\rho(f+g, h_1+h_2)[h_1, h_2, [f, g, h_3]_{\mathcal{B}}]_{\mathcal{B}} \otimes \mu(a, b, \mu(c, d, e)),
\end{aligned}$$

where $f, g, h_1, h_2, h_3 \in Hg(\mathcal{B}_{\rho, \chi})$ and $a, b, c, d, e \in Hg(g_{\rho, \chi, C})$. So $[\cdot, \cdot, \cdot]_E$ is a 3- ρ -Lie algebra structure on E . Also, we have

$$\chi_E(f \otimes a, g \otimes b) = \rho(f+g, g+b)\chi_E(g \otimes b, f \otimes a),$$

and

$$\begin{aligned}
\chi_E([f \otimes a, g \otimes b, h \otimes c]_E, h_1 \otimes d) &= \rho(a, g+h)\rho(b, h)([f, g, h]_A \otimes \mu(a, b, c), h_1 \otimes d) \\
&= \rho(a, g+h)\rho(b, h)\rho(a+b+c, h_2)\chi([f, g, h]_A, h_1)\chi_g(\mu(a, b, c), d) \\
&= \rho(a, g+h)\rho(b, h)\rho(a+b+c, h_2)\chi(f, [g, h, h_1]_{\mathcal{B}})\chi_g(a, \mu(b, c, d)) \\
&= \rho(b, h)\rho(b+c, h_1)\chi_E(f \otimes a, [g, h, h_1] \otimes \mu(b, c, d)) \\
&= \chi_E(f \otimes a, [g \otimes b, h \otimes c, h_1 \otimes d]_E),
\end{aligned}$$

which imply χ_E is a quadratic structure. \square

2.1. Ideal of 3- ρ -Lie algebras. In this part, we introduce the definitions of subalgebra and ideal of \mathcal{B}_{ρ} and give some properties related to ideals of \mathcal{B}_{ρ} .

Definition 2.10. A subalgebra of \mathcal{B}_{ρ} is a sub-vector space $I \subseteq \mathcal{B}_{\rho}$ such that $[I, I, I]_{\mathcal{B}} \subseteq I$. I is also called an ideal of \mathcal{B}_{ρ} if $[I, \mathcal{B}_{\rho}, \mathcal{B}_{\rho}]_{\mathcal{B}} \subseteq I$.

Definition 2.11. Let I be an ideal of $\mathcal{B}_{\rho, \chi}$.

- (i) I is said to be non-degenerate if $\chi|_{I \times I}$ is non-degenerate.
- (ii) The orthogonal I^{\perp} of I , with respect to χ , is defined by

$$I^{\perp} = \{f \in \mathcal{B}_{\rho}, \chi(f, g) = 0 \ \forall g \in I\}. \quad (2.6)$$

Lemma 2.12. Let $Z(\mathcal{B}_{\rho})$ be the center of \mathcal{B}_{ρ} and defined by

$$Z(\mathcal{B}_{\rho}) = \{f \in \mathcal{B}_{\rho} \mid [f, g, h]_{\mathcal{B}} = 0 \ \forall g, h \in \mathcal{B}_{\rho}\}. \quad (2.7)$$

Then $Z(\mathcal{B}_{\rho})$ is an ideal of \mathcal{B}_{ρ} .

Proof. It is clear that $[Z(\mathcal{B}_{\rho}), \mathcal{B}_{\rho}, \mathcal{B}_{\rho}] = \{0\} \subseteq Z(\mathcal{B}_{\rho})$. \square

Lemma 2.13. Let I be an ideal of $\mathcal{B}_{\rho, \chi}$. Then I^{\perp} with respect to χ is an ideal of $\mathcal{B}_{\rho, \chi}$.

Proof. At first, we show that $[I^\perp, \mathcal{B}_{\rho, \chi}, \mathcal{B}_{\rho, \chi}] \subseteq I^\perp$. For $f \in I^\perp$, $g, h \in Hg(\mathcal{B}_\rho)$ and $s \in I$ we have

$$\chi([f, g, h]_{\mathcal{B}}, s) = \chi(f, [g, h, s]_{\mathcal{B}}) = \rho(g + h, s)\chi(f, [s, g, h]) = 0,$$

since $f \in I^\perp$ and I is an ideal of $\mathcal{B}_{\rho, \chi}$. \square

Symplectic Structure: In the following, we introduce the notions like symplectic structure ω , metric φ , derivation D and φ -antisymmetric derivation D on \mathcal{B}_ρ and show that a symplectic structure ω may be defined on \mathcal{B}_ρ if and only if there exists a φ -antisymmetric invertible derivation D of \mathcal{B}_ρ .

Definition 2.14. A non-degenerate and ρ -skew-symmetric bilinear form $\omega \in \wedge^2 \mathcal{B}^*$ is called a symplectic structure on \mathcal{B}_ρ if

$$\begin{aligned} &\omega([f_1, f_2, f_3]_{\mathcal{B}}, f_4) - \rho(f_1, f_2 + f_3 + f_4)\omega([f_2, f_3, f_4]_{\mathcal{B}}, f_1) \\ &+ \rho(f_1 + f_2, f_3 + f_4)\omega([f_3, f_4, f_1]_{\mathcal{B}}, f_2) - \rho(f_1 + f_2 + f_3, f_4)\omega([f_4, f_1, f_2]_{\mathcal{B}}, f_3) = 0. \end{aligned}$$

We will say that $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \chi, \omega)$ is a quadratic symplectic 3- ρ -Lie algebra if (\mathcal{B}, χ) is quadratic and (\mathcal{B}, ω) is symplectic.

In the next example, we are going to check the 4-dimensional 3- ρ -Lie algebra, which its classical case is given in [19].

Example 2.15. Let us consider \mathcal{B} as a $G = \mathbb{Z}_2^3$ -graded vector space with the basis $\{l_1, l_2, l_3, l_4\}$. Then $\mathcal{B} = \mathcal{B}_{(1,0,0)} \oplus \mathcal{B}_{(0,1,0)} \oplus \mathcal{B}_{(0,0,1)} \oplus \mathcal{B}_{(1,1,1)}$ with

$$l_1 \in \mathcal{B}_{(1,0,0)}, \quad l_2 \in \mathcal{B}_{(0,1,0)}, \quad l_3 \in \mathcal{B}_{(0,0,1)}, \quad l_4 \in \mathcal{B}_{(1,1,1)}.$$

Also, $\rho : G \times G \longrightarrow \mathbb{C}^*$ is defined by the matrix

$$[\rho(a, b)] = \begin{bmatrix} 1 & -1 & -1 & 1 \\ -1 & 1 & -1 & 1 \\ -1 & -1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix},$$

where $(a, b) \in \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1)\} \times \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1)\}$. The 3- ρ -Lie algebra bracket associate to the basis $\{l_1, l_2, l_3, l_4\}$ is given by

$$\begin{aligned} [l_1, l_2, l_3]_{\mathcal{B}} &= l_4, & [l_1, l_2, l_4]_{\mathcal{B}} &= l_3, \\ [l_2, l_3, l_4]_{\mathcal{B}} &= l_1, & [l_1, l_3, l_4]_{\mathcal{B}} &= l_2. \end{aligned}$$

It is clear that any non-degenerate ρ -skew symmetric bilinear form is a symplectic structure on \mathcal{B} . For example

$$\begin{aligned} \omega_1 &= l_3^* \wedge l_1^* + l_4^* \wedge l_2^*, & \omega_2 &= l_2^* \wedge l_1^* + l_4^* \wedge l_3^*, & \omega_3 &= l_2^* \wedge l_1^* + l_3^* \wedge l_4^*, \\ \omega_4 &= l_1^* \wedge l_2^* + l_4^* \wedge l_3^*, & \omega_5 &= l_1^* \wedge l_2^* + l_3^* \wedge l_4^*, & \omega_6 &= l_1^* \wedge l_3^* + l_2^* \wedge l_4^*, \end{aligned}$$

where $\{l_1^*, l_2^*, l_3^*, l_4^*\}$ is the dual basis of $\{l_1, l_2, l_3, l_4\}$.

Definition 2.16. A derivation on \mathcal{B}_ρ is a linear map $D : \mathcal{B}_\rho \longrightarrow \mathcal{B}_\rho$ such that

$$D[f, g, h]_{\mathcal{B}} = [D(f), g, h] + \rho(D, f)[f, D(g), h] + \rho(D, f + g)[f, g, D(h)],$$

for any $f, g, h \in Hg(\mathcal{B}_\rho)$. Let us denote by $Der(\mathcal{B}_\rho)$ the space of all derivations of \mathcal{B}_ρ .

Definition 2.17. A metric on ρ -Lie algebra \mathcal{B} is a non-degenerate ρ -symmetric bilinear form $\varphi : \mathcal{B} \times \mathcal{B} \rightarrow \mathbb{K}$ such that

$$\varphi([f, g], h) = -\rho(f, g)\varphi(g, [f, h]).$$

Also, non-degenerate ρ -symmetric bilinear form φ is said to be a metric on \mathcal{B}_ρ if

$$\varphi([f, g, h], z) + \rho(f + g, h)\varphi(h, [f, g, z]) = 0, \quad \forall f, g, h, z \in Hg(\mathcal{B}_\rho). \quad (2.8)$$

In this case, $(\mathcal{B}_\rho, \varphi)$ is called a metric 3- ρ -Lie algebra and we denote it by $\mathcal{B}_{\rho, \varphi}$.

Definition 2.18. Consider $\mathcal{B}_{\rho, \chi}$ ($(\mathcal{B}_{\rho, \varphi})$) and let D be an endomorphism or a derivation of \mathcal{B}_ρ . D is called χ -antisymmetric (φ -antisymmetric) if

$$\chi(D(f), g) = -\rho(D, f)\chi(f, D(g)) \quad (\varphi(D(f), g) = -\rho(D, f)\varphi(f, D(g))),$$

for all $f, g \in Hg(\mathcal{B}_\rho)$. Denote by $\text{End}_\chi(\mathcal{B}_\rho)$ ($\text{End}_\varphi(\mathcal{B}_\rho)$) or $\text{Der}_\chi(\mathcal{B}_\rho)$ ($\text{Der}_\varphi(\mathcal{B}_\rho)$) the space of χ -antisymmetric (φ -antisymmetric) endomorphism or derivation of \mathcal{B}_ρ .

Proposition 2.19. Consider the 3- ρ -Lie algebra \mathcal{B}_ρ . The following assertions are equivalent:

- i. The existence of a symplectic structure on \mathcal{B}_ρ ,
- ii. The existence of an invertible derivation $D \in \text{Der}_\varphi(\mathcal{B}_\rho)$.

Proof. At first, we assume that there is an invertible derivation D on \mathcal{B}_ρ such that $\varphi(D(f), g) = -\rho(D, f)\varphi(f, D(g))$. For $f, g \in \mathcal{B}_\rho$, define $\omega(f, g) = \varphi(D(f), g)$. Thus ω is a non-degenerate and ρ -skew symmetric bilinear form. Now, we show that it is a symplectic structure. For homogeneous elements $f, g, h, z \in \mathcal{B}_\rho$, we deduce

$$\begin{aligned} & \omega([f_1, f_2, f_3]_{\mathcal{B}}, f_4) - \rho(f_1, f_2 + f_3 + f_4)\omega([f_2, f_3, f_4]_{\mathcal{B}}, f_1) + \rho(f_1 + f_2, f_3 + f_4)\omega([f_3, f_4, f_1]_{\mathcal{B}}, f_2) \\ & \quad - \rho(f_1 + f_2 + f_3, f_4)\omega([f_4, f_1, f_2]_{\mathcal{B}}, f_3) \\ & = \varphi(D[f_1, f_2, f_3]_{\mathcal{B}}, f_4) - \rho(f_1, f_2 + f_3 + f_4)\varphi(D[f_2, f_3, f_4]_{\mathcal{B}}, f_1) \\ & \quad + \rho(f_1 + f_2, f_3 + f_4)\varphi(D[f_3, f_4, f_1]_{\mathcal{B}}, f_2) - \rho(f_1 + f_2 + f_3, f_4)\varphi(D[f_4, f_1, f_2]_{\mathcal{B}}, f_3) \\ & = \varphi(D[f_1, f_2, f_3]_{\mathcal{B}}, f_4) - \varphi(D[f_1, f_2, f_3]_{\mathcal{B}}, f_4) = 0. \end{aligned}$$

Conversely is similar. □

Consider the quadratic 3- ρ -Lie algebra $\mathcal{B}_{\rho, \chi}$. For any symmetric bilinear form χ' on \mathcal{B}_ρ , there is an associated map $D : \mathcal{B}_\rho \rightarrow \mathcal{B}_\rho$ satisfying

$$\chi'(f, g) = \chi(D(f), g), \quad \forall f, g \in \mathcal{B}_\rho. \quad (2.9)$$

Since χ and χ' are symmetric, then D is symmetric with respect to χ , i.e.,

$$\chi(D(f), g) = \rho(D, f)\chi(f, D(g)), \quad \forall f, g \in Hg(\mathcal{B}_\rho).$$

Lemma 2.20. Let χ' be defined by (2.9) on \mathcal{B}_ρ . Then

- i. χ' is invariant if and only if D satisfies

$$D([f, g, h]_{\mathcal{B}}) = [D(f), g, h]_{\mathcal{B}} = \rho(D, f)[f, D(g), h]_{\mathcal{B}} = \rho(D, f + g)[f, g, D(h)]_{\mathcal{B}}. \quad (2.10)$$

- ii. χ' is non-degenerate if and only if D is invertible.

Proof. (i) Assuming $f, g, h, s \in \mathcal{B}_\rho$, we have

$$\chi'([f, g, h]_\mathcal{B}, s) = \chi(D([f, g, h]_\mathcal{B}), s) \quad \text{and} \quad \chi'(f, [g, h, s]_\mathcal{B}) = \chi(D(f), [g, h, s]_\mathcal{B}).$$

Since χ is invariant, so $\chi'(f, [g, h, s]_\mathcal{B}) = \chi([D(f), g, h]_\mathcal{B}, s)$. On the other hand, since χ is non-degenerate, therefore χ' is invariant if and only if $D([f, g, h]_\mathcal{B}) = [D(f), g, h]_\mathcal{B}$. Also, since 3-Lie algebra structure $[\cdot, \cdot, \cdot]_\mathcal{B}$ is anticommutative with respect to displacement of every two elements, then $D([f, g, h]_\mathcal{B}) = \rho(D, f)[f, D(g), h]_\mathcal{B} = \rho(D, f + g)[f, g, D(h)]_\mathcal{B}$.

(ii) We assume that χ' is non-degenerate and $f \in \mathcal{B}_\rho$ such that $D(f) = 0$. Then $\chi(D(f), \mathcal{B}_\rho) = 0$ and so $\chi'(f, \mathcal{B}_\rho) = 0$. Therefore $f = 0$ and D is invertible. Conversely, assume that D is invertible and $\chi'(f, \mathcal{B}_\rho) = 0$. So $\chi(D(f), \mathcal{B}_\rho) = 0$. Since χ is non-degenerate, thus $D(f) = 0$. D is invertible, therefore $f = 0$ and χ' is non-degenerate. \square

Definition 2.21. A ρ -symmetric map $D : \mathcal{B}_\rho \longrightarrow \mathcal{B}_\rho$ satisfying (2.10) is called a centromorphism of \mathcal{B}_ρ . The space of centromorphisms of \mathcal{B}_ρ is denote by $\mathcal{C}(\mathcal{B}_\rho)$.

Proposition 2.22. Let $\delta \in \text{Der}_\chi(\mathcal{B}_\rho)$ and $D \in \mathcal{C}(\mathcal{B}_\rho)$ such that $D \circ \delta = \delta \circ D$. Then $D \circ \delta$ is also a χ -antisymmetric derivation of \mathcal{B}_ρ .

Proof. We have

$$\begin{aligned} D \circ \delta[f, g, h] &= D[\delta(f), g, h]_\mathcal{B} + \rho(\delta, f)D[f, \delta(g), h]_\mathcal{B} + \rho(\delta, f + g)[f, g, \delta(h)]_\mathcal{B} \\ &= [D \circ \delta(f), g, h]_\mathcal{B} + \rho(D + \delta, f)D[f, D \circ \delta(g), h]_\mathcal{B} + \rho(\delta, f + g)[f, g, D \circ \delta(h)]_\mathcal{B}, \end{aligned}$$

and $B(D \circ \delta(f), g) = \rho(D + \delta, f)B(f, \delta \circ D(g)) = -\rho(D + \delta, f)B(f, D \circ \delta(g))$ for all $f, g \in Hg(\mathcal{B}_\rho)$. So we deduce that $D \circ \delta \in \text{Der}_\chi(\mathcal{B}_\rho)$. \square

3. REPRESENTATION OF 3- ρ -LIE ALGEBRAS

In this section we extend the representation theory of 3-Lie algebras introduced in [12] and [26] to color Lie version. We introduce a representation of 3- ρ -Lie algebras and discuss the cases of adjoint and coadjoint representations for 3- ρ -Lie algebras.

Consider $\mathcal{L} = \wedge^2 \mathcal{B}_\rho$ and call it the fundamental set. Defining

$$[(f_1, f_2), (g_1, g_2)]_\mathcal{L} = ([f_1, f_2, g_1], g_2) + \rho(f_1 + f_2, g_1)(g_1, [f_1, f_2, g_2]), \quad (3.1)$$

we construct a ρ -Lie algebra structure on \mathcal{L} .

Example 3.1. Let $(\mathcal{B}, [\cdot, \cdot, \cdot]_\mathcal{B}, \rho, \varphi)$ be a metric 3- ρ -Lie algebra. Then $(\mathcal{L}, [\cdot, \cdot]_\mathcal{L}, \varphi_\mathcal{L})$ is a metric ρ -Lie algebra, where

$$\varphi_\mathcal{L}((f_1, f_2), (g_1, g_2)) = \varphi([f_1, f_2, g_1]_\mathcal{B}, g_2).$$

Definition 3.2. The pair (V, μ) is called a representation of \mathcal{B}_ρ if the following conditions are satisfied

$$\mu[(f_1, f_2), (g_1, g_2)]_{\mathcal{L}} = \mu(f_1, f_2)\mu(g_1, g_2) - \rho(f_1 + f_2, g_1 + g_2)\mu(g_1, g_2)\mu(f_1, f_2), \quad (3.2)$$

$$\begin{aligned} \mu([g_1, g_2, g_3]_{\mathcal{B}}, f) &= \mu(g_1, g_2)\mu(g_3, f) + \rho(g_1, g_2 + g_3)\mu(g_2, g_3)\mu(g_1, f) \\ &\quad + \rho(g_1 + g_2, g_3)\mu(g_3, g_1)\mu(g_2, f), \end{aligned} \quad (3.3)$$

$$\begin{aligned} \mu(g, [f_1, f_2, f_3]_{\mathcal{B}}) &= \rho(g, f_1 + f_2)\mu(f_1, f_2)\mu(g, f_3) + \rho(g, f_2 + f_3)\rho(f_1, f_2 + f_3)\mu(f_2, f_3)\mu(g, f_1) \\ &\quad + \rho(g, f_1 + f_3)\rho(f_1 + f_2, f_3)\mu(f_3, f_1)\mu(g, f_2), \end{aligned} \quad (3.4)$$

where V is a G -graded vector space and μ is a linear map from $\mathcal{L} = \wedge^2 \mathcal{B}_\rho$ to $\text{gl}(V)$.

One of the important examples of the 3- ρ -Lie algebra representation is the adjoint representation, which is a bilinear map $ad : \mathcal{B}_\rho \times \mathcal{B}_\rho \longrightarrow \text{gl}(\mathcal{B}_\rho)$ defined by $ad(f_1, f_2)(f_3) = [f_1, f_2, f_3]_{\mathcal{B}}$, that is a representation of \mathcal{B}_ρ on itself.

Lemma 3.3. For the representation (V, μ) of \mathcal{B}_ρ , we have

$$\begin{aligned} 0 &= \rho(f_1 + f_2, g_1)\mu(g_1, [f_1, f_2, g_2]_{\mathcal{B}}) + \mu([f_1, f_2, g_1]_{\mathcal{B}}, g_2) \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\rho(g_1 + g_2, f_1)\mu(f_1, [g_1, g_2, f_2]_{\mathcal{B}}) + \rho(f_1 + f_2, g_1 + g_2)\mu([g_1, g_2, f_1]_{\mathcal{B}}, f_2). \end{aligned}$$

Proof. Using Definition 3.2, we obtain

$$\begin{aligned} &\rho(f_1 + f_2, g_1)\mu(g_1, [f_1, f_2, g_2]_{\mathcal{B}}) + \mu([f_1, f_2, g_1]_{\mathcal{B}}, g_2) \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\mu([g_1, g_2, f_1]_{\mathcal{B}}, f_2) \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\rho(g_1 + g_2, f_1)\mu(f_1, [g_1, g_2, f_2]_{\mathcal{B}}) \\ &= \mu(f_1, f_2)\mu(g_1, g_2) - \rho(f_1 + f_2, g_1 + g_2)\mu(g_1, g_2)\mu(f_1, f_2) \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\mu(g_1, g_2)\mu(f_1, f_2) - \mu(f_1, f_2)\mu(g_1, g_2) = 0. \end{aligned}$$

□

Consider the pair (\mathcal{B}_ρ, V) , where V is a G -graded vector space. The direct sum $\mathcal{B} \oplus V$ is also graded and $(\mathcal{B} \oplus V)_a = \mathcal{B}_a \oplus V_a$. Furthermore, a homogeneous element of $\mathcal{B} \oplus V$ has the form $f + v$ such that $f \in \mathcal{B}_\rho$ and $v \in V$ and $|f + v| = |f| = |v|$.

In the following, we discuss some properties of 3- ρ -Lie algebras representations. Actually, the following theorem examines the relationship between the 3- ρ -Lie algebra representation and the existence of 3- ρ -Lie algebra structure on the graded vector space $\mathcal{B} \oplus V$:

Proposition 3.4. The pair (V, μ) is a representation of \mathcal{B}_ρ if and only if $(\mathcal{B} \oplus V, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus V})$ is a 3- ρ -Lie algebra, where the structure $[\cdot, \cdot, \cdot]_{\mathcal{B} \oplus V}$ is given by

$$[f_1 + v_1, f_2 + v_2, f_3 + v_3]_{\mathcal{B} \oplus V}^\mu = [f_1, f_2, f_3]_{\mathcal{B}} + \mu(f_1, f_2)v_3 + \rho(f_1, f_2 + f_3)\mu(f_2, f_3)v_1 + \rho(f_1 + f_2, f_3)\mu(f_3, f_1)v_2,$$

for homogeneous elements $f_1, f_2, f_3 \in \mathcal{B}_\rho$ and $v_1, v_2, v_3 \in V$.

Proof. The proof of this proposition follows by a direct computation and the definition of the representation. □

In the following, we describe the dual representations and coadjoint representations of 3- ρ -Lie algebras such as the adjoint representations. At first, we recall the dual space of a graded vector space and explore the graded space of direct sum of a graded vector space and its dual.

Let $\mathcal{B} = \oplus_{a \in G} \mathcal{B}_a$ be a G -graded space. Then \mathcal{B}^* , the dual space of \mathcal{B} , is also G -graded space and

$$\mathcal{B}_a^* = \{\alpha \in \mathcal{B}^* \mid \alpha(f) = 0, \forall f : |f| \neq -a\}.$$

Furthermore, the direct sum $\mathcal{B} \oplus \mathcal{B}^*$ is G -graded, where

$$\mathcal{B} \oplus \mathcal{B}^* = \oplus_{a \in G} (\mathcal{B} \oplus \mathcal{B}^*)_a = \oplus_{a \in G} (\mathcal{B}_a \oplus \mathcal{B}_a^*),$$

A homogeneous element of $\mathcal{B} \oplus \mathcal{B}^*$ has the form $f + \alpha$ such that $f \in \mathcal{B}$ and $\alpha \in \mathcal{B}^*$ and $|f + \alpha| = |f| = |\alpha|$.

Proposition 3.5. *Let (V, μ) be a ρ -skew-symmetric representation of \mathcal{B}_ρ and let V^* be the dual vector space of V . Defining the linear map $\mu^* : \mathcal{B}_\rho \times \mathcal{B}_\rho \longrightarrow \text{gl}(V^*)$ by $\mu^*(f_1, f_2)(\varrho) = -\rho(f_1 + f_2, \varrho)\varrho \circ \mu(f_1, f_2)$, where $f_1, f_2 \in \mathcal{B}$, $\varrho \in V^*$, (V^*, μ^*) defines a representation of \mathcal{B}_ρ , which is called the dual representation.*

Proof. By invoking Definition 3.2 and Lemma 3.3, we can prove this proposition by direct computations. \square

Remark 3.6. *An example of the dual representation is the linear map $\text{ad}^* : \mathcal{B}_\rho \times \mathcal{B}_\rho \longrightarrow \text{gl}(\mathcal{B}_\rho^*)$ defined by*

$$\text{ad}^*(f_1, f_2)(\varrho)(f_3) = -\rho(f_1 + f_2, \varrho)\varrho[f_1, f_2, f_3]_{\mathcal{B}} = -\rho(f_1 + f_2, \varrho)\text{ad}(f_1, f_2)(f_3),$$

for $f_1, f_2, f_3 \in \text{Hg}(\mathcal{B}_\rho)$ and $\varrho \in \mathcal{B}_\rho^*$, which is a representation of \mathcal{B} on \mathcal{B}_ρ^* (ad^* is called coadjoint representation). Since ad^* is a representation of \mathcal{B}_ρ , then by Proposition 3.4, $\mathcal{B} \oplus \mathcal{B}^*$ is a 3- ρ -Lie algebra by the following structure

$$\begin{aligned} [f_1 + \alpha_1, f_2 + \alpha_2, f_3 + \alpha_3]_{\mathcal{B} \oplus \mathcal{B}^*}^{\text{ad}^*} &= [f_1, f_2, f_3]_{\mathcal{B}} + \text{ad}^*(f_1, f_2)\alpha_3 \\ &\quad + \rho(f_1, f_2 + f_3)\text{ad}^*(f_2, f_3)\alpha_1 + \rho(f_1 + f_2, f_3)\text{ad}^*(f_3, f_1)\alpha_2. \end{aligned} \quad (3.5)$$

Considering $\varphi : \mathcal{B} \oplus \mathcal{B}^* \times \mathcal{B} \oplus \mathcal{B}^* \longrightarrow \mathbb{K}$ by $\varphi(f + \alpha, g + \beta) = \alpha(g) + \rho(f, g)\beta(f)$, we have a metric 3- ρ -Lie algebra $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}^{\text{ad}^*}, \rho, \varphi)$.

4. 3-PRE- ρ -LIE ALGEBRA

In [12] the notion of 3-pre-Lie algebras was introduced as a generalization of pre-Lie algebras. In this section, we introduce the notion of 3-pre- ρ -Lie algebras and investigate their properties. Also, the 3-pre- ρ -Lie algebras representations and the phase spaces of 3-pre- ρ -Lie algebras will be examined.

Definition 4.1. *A triple $(\mathcal{B}, \{\cdot, \cdot, \cdot\}, \rho)$ is called a 3-pre- ρ -Lie algebra if the following equalities hold*

$$\begin{aligned} \{f_1, f_2, f_3\} &= |f_1| + |f_2| + |f_3|, \\ \{f_1, f_2, f_3\} &= -\rho(f_1, f_2)\{f_2, f_1, f_3\}, \end{aligned} \quad (4.1)$$

$$\begin{aligned} \{f_1, f_2, \{g_1, g_2, g_3\}\} &= \{[f_1, f_2, g_1]_c, g_2, g_3\} + \rho(f_1 + f_2, g_1)\{g_1, [f_1, f_2, g_2]_c, g_3\} \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\{g_1, g_2, \{f_1, f_2, g_3\}\}, \end{aligned} \quad (4.2)$$

$$\begin{aligned} \{[f_1, f_2, f_3]_c, g_1, g_2\} &= \{f_1, f_2, \{f_3, g_1, g_2\}\} + \rho(f_1, f_2 + f_3)\{f_2, f_3, \{f_1, g_1, g_2\}\} \\ &\quad + \rho(f_1 + f_2, f_3)\{f_3, f_1, \{f_2, g_1, g_2\}\}, \end{aligned} \quad (4.3)$$

where \mathcal{B} is a G -graded vector space, $\{., ., .\} : \otimes^3 \mathcal{B} \longrightarrow \mathcal{B}$ is a trilinear map, ρ is a two-cycle and

$$[f_1, f_2, f_3]_c = \{f_1, f_2, f_3\} + \rho(f_1, f_2 + f_3)\{f_2, f_3, f_1\} + \rho(f_1 + f_2, f_3)\{f_3, f_1, f_2\}.$$

To simplify notation, we write $\mathcal{B}_{\{\}} instead of $(\mathcal{B}, \{., ., .\}, \rho)$.$

Proposition 4.2. $([., ., .]_c, \rho)$ give a 3- ρ -Lie algebra structure, when $(\mathcal{B}, \{., ., .\}, \rho)$ is a 3-pre- ρ -Lie algebra. $(\mathcal{B}, [., ., .]_c, \rho)$ is called the sub-adjacent 3- ρ -Lie algebra of \mathcal{B} and is denoted by \mathcal{B}_ρ^c .

Proof. Using the definition of $[., ., .]_c$, (4.2) and (4.3), we have

$$\begin{aligned} & [f_1, f_2, [f_3, f_4, f_5]_c]_c - [[f_1, f_2, f_3]_c, f_4, f_5]_c - \rho(f_1 + f_2, f_3)[f_3, [f_1, f_2, f_4]_c, f_5]_c \\ & - \rho(f_1 + f_2, g_1 + g_2)[f_3, f_4, [f_1, f_2, f_5]_c]_c = 0. \end{aligned}$$

□

Note that, $\mathcal{B}_{\{\}}$ is called the compatible 3- ρ -pre-Lie algebra structure on the 3- ρ -Lie algebra \mathcal{B}_ρ^c . The following lemma gives a representation of \mathcal{B}_ρ^c by the left multiplication:

Lemma 4.3. Defining the left multiplication $L : \wedge^2 \mathcal{B}_{\{\}} \longrightarrow gl(\mathcal{B}_{\{\}})$ by $L(f, g)h = \{f, g, h\}$, (\mathcal{B}, L) is a representation of \mathcal{B}_ρ^c .

Proof. By Definition 3.2, we have

$$\begin{aligned} L([f_1, f_2], (g_1, g_2))_{\mathcal{L}}(h) &= L([f_1, f_2, g_1]_c, g_2)(h) + \rho(f_1 + f_2, g_1)L(g_1, [f_1, f_2, g_2]_c)(h) \\ &= \{[f_1, f_2, g_1]_c, g_2, h\} + \rho(f_1 + f_2, g_1)\{g_1, [f_1, f_2, g_2]_c, h\} \\ &= \{f_1, f_2, [g_1, g_2, h]_c\} - \rho(f_1 + f_2, g_1 + g_2)\{g_1, g_2, \{f_1, f_2, h\}\} \\ &= L(f_1, f_2)L(g_1, g_2)(h) - \rho(f_1 + f_2, g_1 + g_2)L(g_1, g_2)L(f_1, f_2)(h). \end{aligned}$$

The other ones prove in a similar way.

□

Representation of 3- ρ -pre-Lie algebras: Here, we are going to represent the representation theory of 3- ρ -pre-Lie algebras. Also, the dual representation will be given.

Definition 4.4. A pair $(\mu, \tilde{\mu})$ is called a representation of $\mathcal{B}_{\{\}} on G -graded vector space V if for all homogeneous elements $f_1, f_2, f_3, f_4 \in Hg(\mathcal{B})$, the following equalities hold$

$$\begin{aligned} \tilde{\mu}(f_1, \{f_2, f_3, f_4\}) &= \rho(f_1, f_2 + f_3)\mu(f_2, f_3)\tilde{\mu}(f_1, f_4) + \rho(f_1, f_3 + f_4)\rho(f_2, f_3 + f_4)\tilde{\mu}(f_3, f_4)\tilde{\mu}(f_1, f_2) \\ &\quad - \rho(f_1, f_2 + f_4)\rho(f_3, f_4)\tilde{\mu}(f_2, f_4)\tilde{\mu}(f_1, f_3) + \rho(f_1 + f_2, f_3 + f_4)\tilde{\mu}(f_3, f_4)\mu(f_1, f_2) \\ &\quad + \rho(f_1, f_2 + f_3 + f_4)\rho(f_3, f_4)\tilde{\mu}(f_2, f_4)\tilde{\mu}(f_3, f_1) \\ &\quad - \rho(f_1, f_2 + f_4)\rho(f_3, f_4)\tilde{\mu}(f_2, f_4)\mu(f_1, f_3) \\ &\quad - \rho(f_2, f_3 + f_4)\rho(f_1, f_2 + f_3 + f_4)\tilde{\mu}(f_3, f_4)\tilde{\mu}(f_2, f_1), \end{aligned}$$

$$\begin{aligned} \mu(f_1, f_2)\tilde{\mu}(f_3, f_4) &= \rho(f_1 + f_2, f_3 + f_4)\tilde{\mu}(f_3, f_4)\mu(f_1, f_2) - \rho(f_1 + f_2, f_3 + f_4)\rho(f_1, f_2)\tilde{\mu}(f_3, f_4)\tilde{\mu}(f_2, f_1) \\ &\quad + \rho(f_1 + f_2, f_3 + f_4)\tilde{\mu}(f_3, f_4)\tilde{\mu}(f_1, f_2) + \tilde{\mu}([f_1, f_2, f_3]_c, f_4) \\ &\quad + \rho(f_1 + f_2, f_3)\tilde{\mu}(f_3, \{f_1, f_2, f_4\}), \end{aligned}$$

$$\tilde{\mu}([f_1, f_2, f_3]_c, f_4) = \mu(f_1, f_2)\tilde{\mu}(f_3, f_4) + \rho(f_1, f_2 + f_3)\mu(f_2, f_3)\tilde{\mu}(f_1, f_4) + \rho(f_1 + f_2, f_3)\mu(f_3, f_1)\tilde{\mu}(f_2, f_4),$$

$$\begin{aligned} \tilde{\mu}(f_3, f_4)\mu(f_1, f_2) &= \rho(f_1, f_2)\tilde{\mu}(f_3, f_4)\tilde{\mu}(f_2, f_1) - \tilde{\mu}(f_3, f_4)\tilde{\mu}(f_1, f_2) + \rho(f_3 + f_4, f_1 + f_2)\mu(f_1, f_2)\tilde{\mu}(f_3, f_4) \\ &\quad - \rho(f_3 + f_4, f_1 + f_2)\rho(f_1, f_2)\tilde{\mu}(f_2, \{f_1, f_3, f_4\}) + \rho(f_3 + f_4, f_1 + f_2)\tilde{\mu}(f_1, \{f_2, f_3, f_4\}), \end{aligned}$$

where $\mu : \wedge^2 \mathcal{B}_\rho^c \rightarrow gl(V)$ is a representation of the 3- ρ -Lie algebra \mathcal{B}_ρ^c and $\tilde{\mu} : \otimes^2 \mathcal{B} \rightarrow gl(V)$ is a bilinear map.

In the following lemma, we construct a representation of $\mathcal{B}_{\{\}}$ by the left and right multiplication. The left and right multiplication are two bilinear maps of the form $L, R : \otimes^2 \mathcal{B}_{\{\}} \rightarrow gl(\mathcal{B}_{\{\}})$ which are defined by $L(f, g)h = \{f, g, h\}$ and $R(f, g)h = \rho(f + g, h)\{h, f, g\}$, respectively.

Lemma 4.5. Let $\mathcal{B}_{\{\}}$ be a 3-pre- ρ -Lie algebra. Then the pair (L, R) is a representation of $\mathcal{B}_{\{\}}$ on itself (this representation is called the regular representation).

Proof. For sample, let us examine one of the properties of the representation. For this, we have

$$\begin{aligned} &L(f_1, f_2)R(f_3, f_4)f_5 + \rho(f_1, f_2 + f_3)L(f_2, f_3)R(f_1, f_4)f_5 + \rho(f_1 + f_2, f_3)L(f_3, f_1)R(f_2, f_4)f_5 \\ &= \rho(f_3 + f_4, f_5)\{f_1, f_2, \{f_5, f_3, f_4\}\} + \rho(f_1, f_2 + f_3)\rho(f_1 + f_4, f_5)\{f_2, f_3, \{f_5, f_1, f_4\}\} \\ &\quad + \rho(f_1 + f_2, f_3)\rho(f_2 + f_4, f_5)\{f_3, f_1, \{f_5, f_2, f_4\}\} \\ &= -\rho(f_4, f_5)\{[f_1, f_2, f_3]_c, f_5, f_4\} = \rho(f_1 + f_2 + f_3 + f_4, f_5)\{f_5, [f_1, f_2, f_3]_c, f_4\} \\ &= R([f_1, f_2, f_3]_c, f_4)f_5. \end{aligned}$$

□

Consider $(\mathcal{B}, \{., ., .\}, \rho)$ and (V, μ) as a 3-pre- ρ -Lie algebra and a representation of \mathcal{B}_ρ^c , respectively. Then $(V, \mu, 0)$ is a representation of $\mathcal{B}_{\{\}}$.

Proposition 4.6. *Let $(V, \mu, \tilde{\mu})$ be a representation of $\mathcal{B}_{\{\cdot\}}$. Then $(\mathcal{B} \oplus V, \{\cdot, \cdot, \cdot\}_{\mu, \tilde{\mu}}, \rho)$ is a 3-pre- ρ -Lie algebra, where the operation $\{\cdot, \cdot, \cdot\}_{\mu, \tilde{\mu}} : \otimes^3(\mathcal{B} \oplus V) \longrightarrow \mathcal{B} \oplus V$ is defined by*

$$\{f_1 + v_1, f_2 + v_2, f_3 + v_3\}_{\mu, \tilde{\mu}} = \{f_1, f_2, f_3\} + \mu(f_1, f_2)v_3 + \rho(f_1, f_2 + f_3)\tilde{\mu}(f_2, f_3)v_1 - \rho(f_2, f_3)\tilde{\mu}(f_1, f_3)v_2, \quad (4.4)$$

and $f_1, f_2, f_3 \in Hg(\mathcal{B})$ and $v_1, v_2, v_3 \in Hg(V)$,

Proof. Using Definitions 4.1 and 4.4 and a direct computation, we get the result. \square

Let V be a graded vector space. Define the operator $\zeta : \otimes^2 V \longrightarrow \otimes^2 V$ by $\zeta(f_1 \otimes f_2) = f_2 \otimes f_1$, where $f_1 \otimes f_2 \in \otimes^2 V$.

Proposition 4.7. *Let $(V, \mu, \tilde{\mu})$ be a representation of $\mathcal{B}_{\{\cdot\}}$. If we define the bilinear map $\nu : \wedge^2 \mathcal{B} \longrightarrow gl(V)$ by $\nu(f_1, f_2) = (\mu - \rho(f_1, f_2)\tilde{\mu}\zeta + \tilde{\mu})(f_1, f_2)$ for all $f_1, f_2 \in Hg(\mathcal{B})$, then (V, ν) is a representation of \mathcal{B}_ρ^c .*

Proof. Proposition 4.6 implies that we have the 3-pre- ρ -Lie algebra $(\mathcal{B} \oplus V, \{\cdot, \cdot, \cdot\}_{\mu, \tilde{\mu}}, \rho)$. Also, Proposition 4.2 implies that $(\mathcal{B}^c \oplus V, [\cdot, \cdot, \cdot]_c, \rho)$ is the sub-adjacent 3- ρ -Lie algebra, where

$$\begin{aligned} [f_1 + v_1, f_2 + v_2, f_3 + v_3]_c^{\mu, \tilde{\mu}} &= \{f_1 + v_1, f_2 + v_2, f_3 + v_3\}_{\mu, \tilde{\mu}} + \rho(f_1, f_2 + f_3)\{f_2 + v_2, f_3 + v_3, f_1 + v_1\}_{\mu, \tilde{\mu}} \\ &\quad + \rho(f_1 + f_2, f_3)\{f_3 + v_1, f_1 + v_1, f_2 + v_2\}_{\mu, \tilde{\mu}}. \end{aligned}$$

By using (4.4), we get

$$\begin{aligned} [f_1 + v_1, f_2 + v_2, f_3 + v_3]_c^{\mu, \tilde{\mu}} &= [f_1, f_2, f_3]_c + (\mu - \rho(f_1, f_2)\tilde{\mu}\zeta + \tilde{\mu})(f_1, f_2)v_3 \\ &\quad + \rho(f_1, f_2 + f_3)(\mu - \rho(f_2, f_3)\tilde{\mu}\zeta + \tilde{\mu})(f_2, f_3)v_1 \\ &\quad + \rho(f_1 + f_2, f_3)(\mu - \rho(f_3, f_1)\tilde{\mu}\zeta + \tilde{\mu})(f_3, f_1)v_2 \\ &= [f_1, f_2, f_3]_c + \nu(f_1, f_2)v_3 + \rho(f_1, f_2 + f_3)\nu(f_2, f_3)v_1 \\ &\quad + \rho(f_1 + f_2, f_3)\nu(f_3, f_1)v_2. \end{aligned} \quad (4.5)$$

Finally, by Proposition 3.4, we deduce that (V, ν) is a representation of \mathcal{B}_ρ^c . \square

According to the above proposition, we can deduce that if $(\mathcal{B}, \{\cdot, \cdot, \cdot\}, \rho)$ is a 3-pre- ρ -Lie algebra with the representation $(V, \mu, \tilde{\mu})$, then the 3-pre- ρ -Lie algebras $(\mathcal{B} \oplus V, \{\cdot, \cdot, \cdot\}_{\mu, \tilde{\mu}}, \rho)$ and $(\mathcal{B} \oplus V, \{\cdot, \cdot, \cdot\}_{\nu, 0}, \rho)$ have the same sub-adjacent 3- ρ -Lie algebra given by (4.5).

In the next proposition, we find the dual of the representation $(V, \mu, \tilde{\mu})$ of 3-pre- ρ -Lie algebra $(\mathcal{B}, \{\cdot, \cdot, \cdot\}, \rho)$. Let us define $\nu^*(f_1, f_2) = \mu^* - \rho(f_1, f_2)\tilde{\mu}^*\zeta + \tilde{\mu}^*(f_1, f_2)$.

Proposition 4.8. *Consider $\mathcal{B}_{\{\cdot\}}$ equipped with a ρ -skew-symmetric representation $(V, \mu, \tilde{\mu})$. Then $(V^*, \nu^*, -\tilde{\mu}^*)$ is a representation of $\mathcal{B}_{\{\cdot\}}$, which is called the dual representation of the representation $(V, \mu, \tilde{\mu})$.*

Proof. By Propositions 3.5, 4.7 and Definition 4.4, we get the result. \square

If $\mathcal{B}_{\{\cdot\}}$ is a 3-pre- ρ -Lie algebra with the ρ -skew-symmetric representation $(V, \mu, \tilde{\mu})$, then the 3-pre- ρ -Lie algebras $(\mathcal{B} \oplus V^*, \{\cdot, \cdot, \cdot\}_{\mu^*, 0}, \rho)$ and $(\mathcal{B} \oplus V^*, \{\cdot, \cdot, \cdot\}_{\nu^*, -\tilde{\mu}^*}, \rho)$ have the same sub-adjacent 3- ρ -Lie algebra $(\mathcal{B}^c \oplus V^*, [\cdot, \cdot, \cdot]_c^{\mu^*}, \rho)$.

ρ - \mathcal{O} -operator. In the following, we define the notion of ρ - \mathcal{O} -operator associated to the representation (V, μ) of \mathcal{B}_ρ , then we construct a 3- ρ -pre-Lie algebra structure on the representation V by a ρ - \mathcal{O} -operator T and bilinear map μ . In the next, we investigate the relationship between compatible 3- ρ -pre-Lie algebra structure and an invertible ρ - \mathcal{O} -operator of \mathcal{B}_ρ (for the classical case refer to [12]).

Definition 4.9. Let \mathcal{B} be a 3- ρ -Lie algebra with the representation (V, μ) . An even linear operator $T : V \longrightarrow \mathcal{B}$ is called a ρ - \mathcal{O} -operator associated to (V, μ) if T satisfies

$$[Tx, Ty, Tz]_{\mathcal{B}} = T(\mu(Tx, Ty)z + \rho(x, y + z)(\mu(Ty, Tz)x + \rho(x + y, z)\mu(Tz, Tx)y), \quad \forall x, y, z \in V.$$

Lemma 4.10. Consider the following three assumptions

- i. \mathcal{B}_ρ is a 3- ρ -Lie algebra,
- ii. (V, μ) is a ρ -skew-symmetric representation of \mathcal{B}_ρ ,
- iii. T is a ρ - \mathcal{O} -operator associated to (V, μ) .

If we define a new multiplication on V by

$$\{x, y, z\}_V = \mu(Tx, Ty)z, \quad \forall x, y, z \in V, \quad (4.6)$$

then $(V, \{\cdot, \cdot, \cdot\}_V)$ is a 3-pre- ρ -Lie algebra.

Proof. Suppose that $x, y, z \in Hg(V)$. It is easy to see that

$$\{x, y, z\} = -\rho(x, y)\{y, x, z\}, \quad T[x, y, z]_c = [Tx, Ty, Tz].$$

Also, for $x_1, x_2, x_3, x_4, x_5 \in Hg(V)$, we have

$$\{x_1, x_2, \{x_3, x_4, x_5\}\} = \mu(Tx_1, Tx_2)\mu(Tx_3, Tx_4)x_5, \quad (4.7)$$

$$\{[x_1, x_2, x_3]_c, x_4, x_5\} = \mu([Tx_1, Tx_2, Tx_3], Tx_4)x_5, \quad (4.8)$$

$$\rho(x_1 + x_2, x_3)\{x_3, [x_1, x_2, x_4]_c, x_5\} = \rho(x_1 + x_2, x_3)\mu(Tx_3, [Tx_1, Tx_2, Tx_4])x_5, \quad (4.9)$$

$$\rho(x_1 + x_2, x_3 + x_4)\{x_3, x_4, \{x_1, x_2, x_5\}\} = \rho(x_1 + x_2, x_3 + x_4)\mu(Tx_3, Tx_4)\mu(Tx_1, Tx_2)x_5. \quad (4.10)$$

By Definition 3.2 and (4.7)–(4.10), we get the result. \square

Corollary 4.11. By the above proposition, we deduce that $(V, [\cdot, \cdot, \cdot]_C)$ is a 3- ρ -Lie algebra as the sub-adjacent of the 3- ρ -pre-Lie algebra $(V, \{\cdot, \cdot, \cdot\})$. Furthermore, $T(V) = \{Tv | v \in V\} \subset A$ is a subalgebra of \mathcal{B}_ρ and there is an induced 3- ρ -pre-Lie algebra structure on $T(V)$ given by

$$\{Tu, Tv, Tw\}_{T(V)} := T\{u, v, w\}, \quad \forall u, v, w \in V. \quad (4.11)$$

Lemma 4.12. Consider 3- ρ -Lie algebra \mathcal{B}_ρ . Then the following assertions are equivalent

- i. existence of a compatible 3-pre- ρ -Lie algebra structure,
- ii. existence of an invertible ρ - \mathcal{O} -operator.

Proof. Let T be an invertible ρ - \mathcal{O} -operator of \mathcal{B}_ρ associated to (V, μ) . BY (4.6), (4.11) and this fact that T is a ρ - \mathcal{O} -operator, we can easily deduce that $(A = T(V), \{\cdot, \cdot, \cdot\}_{T(V)})$ is a compatible 3- ρ -pre-Lie algebra. Conversely, the identity map id is an ρ - \mathcal{O} -operator of the sub-adjacent 3- ρ -Lie algebra of a 3- ρ -pre-Lie algebra associated to the representation (\mathcal{B}, L) . \square

4.1. Phase spaces of 3- ρ -Lie algebras. In this section, the notion of a phase space of a 3- ρ -Lie algebra will be introduced and we show that a 3- ρ -Lie algebra has a phase space if and only if it is sub-adjacent to a 3- ρ -pre-Lie algebra.

Proposition 4.13. *Consider the symplectic 3- ρ -Lie algebra $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \omega)$. Then the following structure defines a compatible 3-pre- ρ -Lie algebra structure on \mathcal{B}*

$$\omega(\{f, g, h\}, s) = -\rho(f + g, h)\omega(h, [f, g, s]_{\mathcal{B}}), \quad \forall f, g, h, s \in \mathcal{B}. \quad (4.12)$$

Proof. Let us define a linear map $T : \mathcal{B}^* \longrightarrow \mathcal{B}$ by $\omega(f, g) = (T^{-1}(f))(g)$, or equivalently, $\omega(T\alpha, g) = \alpha(g)$ for all $f, g \in Hg(\mathcal{B})$ and $\alpha \in \mathcal{B}^*$. Since ω is a symplectic structure, we deduce that T is an invertible ρ - \mathcal{O} -operator associated to (\mathcal{B}^*, Ad^*) . Also, by (4.6) and (4.11), there exists a compatible 3- ρ -pre-Lie algebra on \mathcal{B} given by $\{f, g, h\} = T(ad^*(f, g)T^{-1}(h))$. So, (4.12) holds. \square

Definition 4.14. $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \omega)$ is called a phase space of \mathcal{B}_ρ , if

- i. $[\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}$ is a 3- ρ -Lie algebra structure on the vector space $\mathcal{B} \oplus \mathcal{B}^*$,
- ii. ω is a natural non-degenerate ρ -skew-symmetric bilinear form on $\mathcal{B} \oplus \mathcal{B}^*$ given by

$$\omega(f + \alpha, g + \beta) = \alpha(g) - \rho(f, g)\beta(f), \quad (4.13)$$

- such that $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \omega)$ is a symplectic 3- ρ -Lie algebra,
- iii. $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho)$ and $(\mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B}^*}, \rho)$ are 3- ρ -Lie subalgebras of $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho)$.

Theorem 4.15. *A 3- ρ -Lie algebra has a phase space if and only if it is sub-adjacent to a 3-pre- ρ -Lie algebra.*

Proof. Let $\mathcal{B}_{\{\}} be a 3-pre- ρ -Lie algebra. By Propositions 3.4, 3.5, 4.13 and Lemma 4.3 we have the 3- ρ -Lie algebra $(\mathcal{B}^c \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B}^c \oplus \mathcal{B}^*}^{L^*}, \rho)$. Moreover, for all $f_1, f_2, f_3, f_4 \in Hg(\mathcal{B})$ and $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in Hg(\mathcal{B}^*)$, by$

(3.5) with respect to L^* , we have

$$\begin{aligned} & \omega([f_1 + \alpha_1, f_2 + \alpha_2, f_3 + \alpha_3]_{\mathcal{B}^c \oplus \mathcal{B}^*}^{L^*}, f_4 + \alpha_4) \\ &= -\rho(f_1 + f_2 + f_3, f_4)\alpha_4\{f_1, f_2, f_3\} - \rho(f_1, f_2 + f_3)\rho(f_1 + f_2 + f_3, f_4)\alpha_4\{f_2, f_3, f_1\} \\ & \quad - \rho(f_1 + f_2, f_3)\rho(f_1 + f_2 + f_3, f_4)\alpha_4\{f_3, f_1, f_2\} - \rho(f_1 + f_2, f_3)\alpha_3\{f_1, f_2, f_4\} \\ & \quad - \alpha_1\{f_2, f_3, f_4\} - \rho(f_1 + f_2, f_3)\rho(f_3 + f_1, f_2)\alpha_2\{f_3, f_1, f_4\}, \end{aligned}$$

$$\begin{aligned} & \rho(f_1, f_2 + f_3 + f_4)\omega([f_2 + \alpha_2, f_3 + \alpha_3, f_4 + \alpha_4]_{\mathcal{B}^c \oplus \mathcal{B}^*}^{L^*}, f_1 + \alpha_1) \\ &= -\alpha_1\{f_2, f_3, f_4\} - \rho(f_2, f_3 + f_4)\alpha_1\{f_3, f_4, f_2\} - \rho(f_2 + f_3, f_4)\alpha_1\{f_4, f_2, f_3\} \\ & \quad - \rho(f_1, f_2 + f_3 + f_4)\rho(f_2 + f_3, f_4)\alpha_4\{f_2, f_3, f_1\} - \rho(f_1, f_2 + f_3 + f_4)\alpha_2\{f_3, f_4, f_1\} \\ & \quad - \rho(f_1, f_2 + f_3 + f_4)\rho(f_2 + f_3, f_4)\rho(f_4 + f_2, f_3)\alpha_3\{f_4, f_2, f_1\}, \end{aligned}$$

$$\begin{aligned} & \rho(f_1 + f_2, f_3 + f_4)\omega([f_3 + \alpha_3, f_4 + \alpha_4, f_1 + \alpha_1]_{\mathcal{B}^c \oplus \mathcal{B}^*}^{L^*}, f_2 + \alpha_2) \\ &= -\rho(f_1, f_3 + f_4 + f_2)\alpha_2\{f_3, f_4, f_1\} \\ & \quad - \rho(f_1, f_4 + f_1)\rho(f_3, f_4)\alpha_2\{f_4, f_1, f_3\} \\ & \quad - \rho(f_2, f_3 + f_4)\rho(f_1 + f_3 + f_4, f_2)\alpha_2\{f_1, f_3, f_4\} \\ & \quad - \rho(f_2, f_3 + f_4)\alpha_1\{f_3, f_4, f_2\} - \rho(f_1 + f_2, f_3 + f_4)\alpha_3\{f_4, f_1, f_2\} \\ & \quad - \rho(f_2, f_3 + f_4)\rho(f_3 + f_1, f_4)\alpha_4\{f_1, f_3, f_2\}, \end{aligned}$$

$$\begin{aligned} & \rho(f_1 + f_2 + f_3, f_4)\omega([f_4 + \alpha_4, f_1 + \alpha_1, f_2 + \alpha_2]_{\mathcal{B}^c \oplus \mathcal{B}^*}^{L^*}, f_3 + \alpha_3) \\ &= -\rho(f_1 + f_2, f_3 + f_4)\alpha_3\{f_4, f_1, f_2\} - \rho(f_1 + f_2, f_3)\alpha_3\{f_1, f_2, f_4\} \\ & \quad - \rho(f_1 + f_2, f_3 + f_4)\rho(f_4 + f_1, f_2)\alpha_3\{f_2, f_4, f_1\} - \rho(f_1 + f_3, f_4)\rho(f_1, f_2)\alpha_2\{f_4, f_1, f_3\} \\ & \quad - \rho(f_3, f_4)\rho(f_1 + f_2, f_4)\alpha_4\{f_1, f_2, f_3\} - \rho(f_3, f_4)\alpha_1\{f_2, f_4, f_3\}. \end{aligned}$$

By the above relations, we deduce that ω is a symplectic structure on $\mathcal{B}^c \oplus \mathcal{B}^*$. Moreover, \mathcal{B}^c and \mathcal{B}^* are the subalgebra and the abelian subalgebra of $\mathcal{B}^c \oplus \mathcal{B}^*$, respectively. Thus, the symplectic 3- ρ -Lie algebra $(\mathcal{B}^c \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B}^c \oplus \mathcal{B}^*}^{L^*}, \rho, \omega)$ is a phase space of \mathcal{B}_ρ^c . Conversely, let $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \omega)$ be a phase space of \mathcal{B}_ρ . By Proposition 4.13, there is a compatible 3-pre- ρ -Lie algebra structure $\{\cdot, \cdot, \cdot\}$ on $\mathcal{B} \oplus \mathcal{B}^*$. Thus

$$\omega(\{f, g, h\}, z) = -\rho(f + g, h)\omega(h, [f, g, z]_{\mathcal{B} \oplus \mathcal{B}^*}) = -\rho(f + g, h)\omega(h, [f, g, z]_{\mathcal{B}}) = 0 \quad \forall f, g, h, z \in Hg(\mathcal{B}).$$

The last equality holds, since \mathcal{B}_ρ is a subalgebra of $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho)$. Therefore, $\{f, g, h\} \in \mathcal{B}_\rho$ that says $(\mathcal{B}, \{\cdot, \cdot, \cdot\}_{|\mathcal{B}}, \rho)$ is a subalgebra of the 3-pre- ρ -Lie algebra $(\mathcal{B} \oplus \mathcal{B}^*, \{\cdot, \cdot, \cdot\}, \rho)$ and so its sub-adjacent 3- ρ -Lie algebra \mathcal{B}_ρ^c is the 3- ρ -Lie algebra \mathcal{B}_ρ . \square

Corollary 4.16. *Let $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \omega)$ be a phase space of \mathcal{B}_ρ and $(\mathcal{B} \oplus \mathcal{B}^*, \{\cdot, \cdot, \cdot\}, \rho)$ be the associated 3-pre- ρ -Lie algebra. Then $(\mathcal{B}, \{\cdot, \cdot, \cdot\}_{|\mathcal{B}}, \rho)$ and $(\mathcal{B}^*, \{\cdot, \cdot, \cdot\}_{|\mathcal{B}^*}, \rho)$ are subalgebras of the 3-pre- ρ -Lie algebra $(\mathcal{B} \oplus \mathcal{B}^*, \{\cdot, \cdot, \cdot\}, \rho)$.*

Lemma 4.17. *Let (V, μ) be a representation of \mathcal{B}_ρ and $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}^{\mu^*}, \rho, \omega)$ be a phase space of it. Then there exists a 3-pre- ρ -Lie algebra structure on \mathcal{B} given by*

$$\{f, g, h\} = \mu(f, g)h, \quad \forall f, g, h \in \mathcal{B}.$$

Proof. Assume that $f, g, h \in Hg(\mathcal{B})$ and $\alpha \in Hg(\mathcal{B}^*)$. Then, we have

$$\begin{aligned} \rho(f + g + h, \alpha)\alpha(\{f, g, h\}) &= -\omega(\{f, g, h\}, \alpha) = \rho(f + g, h)\omega(h, [f, g, \alpha]_{\mathcal{B} \oplus \mathcal{B}^*}) = \rho(f + g, h)\omega(h, \mu^*(f, g)\alpha) \\ &= -\rho(f + g, h)\rho(h, f + g + \alpha)\mu^*(f, g)\alpha(h) = \rho(f + g + h, \alpha)\alpha\mu(f, g)h. \end{aligned}$$

□

5. QUADRATIC 3-HOM- ρ -LIE ALGEBRAS

In this section, we study 3-Hom- ρ -lie algebras, which 3- ρ -lie algebras are special case of these Lie algebras with respect to the map Id . The constructions of Section 2, such as quadratic, symplectic structures, 3-associative Lie algebras and etc, will be investigated in this section to color Hom-Lie case. In this section, the representation theory of 3-Hom- ρ -lie algebras is also included.

Definition 5.1. *Multiple $(\mathcal{B}, [\cdot, \cdot]_{\mathcal{B}}, \rho, \phi)$ is called a 2-Hom- ρ -Lie algebra (or simply a Hom- ρ -Lie algebra) if*

- i. \mathcal{B} is a G -graded vector space, where G is an abelian group,
- ii. ρ is a two-cycle,
- iii. $\phi : \mathcal{B} \longrightarrow \mathcal{B}$ is an even linear map,
- iv. $[\cdot, \cdot]_{\mathcal{B}}$ is a bilinear map on \mathcal{B} , satisfying the following conditions

- $|[f, g]_{\mathcal{B}}| = |f| + |g|$,
- $[f, g]_{\mathcal{B}} = -\rho(f, g)[g, f]_{\mathcal{B}}$,
- $\rho(h, f)[\phi(f), [g, h]_{\mathcal{B}}]_{\mathcal{B}} + \rho(g, h)[\phi(h), [f, g]_{\mathcal{B}}]_{\mathcal{B}} + \rho(f, g)[\phi(g), [h, f]_{\mathcal{B}}]_{\mathcal{B}} = 0 \quad \forall f, g, h \in \mathcal{B}.$

The third condition is equivalent to

$$[\phi(f), [g, h]_{\mathcal{B}}]_{\mathcal{B}} = [[f, g]_{\mathcal{B}}, \phi(h)]_{\mathcal{B}} + \rho(f, g)[\phi(g), [f, h]_{\mathcal{B}}]_{\mathcal{B}}.$$

Definition 5.2. *Let*

- i. \mathcal{B} be a G -graded vector space,
- ii. $\phi : \mathcal{B} \longrightarrow \mathcal{B}$ be an even linear map,
- iii. ρ be a two-cycle,
- iv. $[\cdot, \cdot, \cdot]_{\mathcal{B}}$ be a trilinear map on \mathcal{B} satisfying

$$|[f_1, f_2, f_3]_{\mathcal{B}}| = |f_1| + |f_2| + |f_3|,$$

$$\begin{aligned} [\phi(f_1), \phi(f_2), [g_1, g_2, g_3]_{\mathcal{B}}]_{\mathcal{B}} &= [[f_1, f_2, g_1]_{\mathcal{B}}, \phi(g_2), \phi(g_3)]_{\mathcal{B}} + \rho(f_1 + f_2, g_1)[\phi(g_1), [f_1, f_2, g_2]_{\mathcal{B}}, \phi(g_3)]_{\mathcal{B}} \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)[\phi(g_1), \phi(g_2), [f_1, f_2, g_3]_{\mathcal{B}}]_{\mathcal{B}}, \end{aligned}$$

$$[f_1, f_2, f_3]_{\mathcal{B}} = -\rho(f_1, f_2)[f_2, f_1, f_3]_{\mathcal{B}} = -\rho(f_2, f_3)[f_1, f_3, f_2]_{\mathcal{B}},$$

for any $f_1, f_2, f_3, g_1, g_2 \in Hg(\mathcal{B})$. Then $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi)$ is called a 3- ρ -Lie algebra. Let us denote the 3-Hom- ρ -Lie algebra $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi)$ by $\mathcal{B}_{\rho, \phi}$.

Definition 5.3. Let $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \phi)$ and $(\mathcal{A}, [\cdot, \cdot, \cdot]_{\mathcal{A}}, \psi)$ be two 3-Hom- ρ -Lie algebras. A linear map $\alpha : \mathcal{B} \longrightarrow \mathcal{A}$ is said to be a morphism of 3-Hom- ρ -Lie algebras if for all $f, g, h \in \mathcal{B}$

$$\alpha[f, g, h]_{\mathcal{B}} = [\alpha(f), \alpha(g), \alpha(h)]_{\mathcal{A}},$$

and

$$\alpha \circ \phi = \psi \circ \alpha.$$

Definition 5.4. $\mathcal{B}_{\rho, \phi}$ is said to be

- (1) multiplicative if ϕ is a Lie algebra morphism, i.e. for any $f, g, h \in \mathcal{B}$, $\phi[f, g, h]_{\mathcal{B}} = [\phi(f), \phi(g), \phi(h)]_{\mathcal{B}}$.
- (2) regular if ϕ is an automorphism for $[\cdot, \cdot, \cdot]_{\mathcal{B}}$,
- (3) involutive if $\phi^2 = Id_{\mathcal{B}}$.

In the following theorem, a new 3-Hom- ρ -Lie algebra structure will be constructed by combining a 3-Hom- ρ -Lie algebra structure and an even 3- ρ -Lie algebra endomorphism.

Theorem 5.5. Consider $\mathcal{B}_{\rho, \phi}$ and let $\beta : \mathcal{B}_{\rho, \phi} \longrightarrow \mathcal{B}_{\rho, \phi}$ be an even 3- ρ -Lie algebra endomorphism. Then $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\beta} = \beta \circ [\cdot, \cdot, \cdot]_{\mathcal{B}}, \beta \circ \phi)$ is a 3-Hom- ρ -Lie algebra.

Proof. It is easy to see that $[\cdot, \cdot, \cdot]_{\beta}$ is ρ -skew symmetric with respect to the displacement of every two elements. Now, we check the ρ -fundamental identity:

- (a) $[\beta \circ \phi(f_1), \beta \circ \phi(f_2), [f_3, f_4, f_5]_{\beta}]_{\beta} = [\beta \circ \phi(f_1), \beta \circ \phi(f_2), \beta \circ [f_3, f_4, f_5]_{\mathcal{B}}]_{\beta}$
 $= \beta^2[\phi(f_1), \phi(f_2), [f_3, f_4, f_5]_{\mathcal{B}}]_{\mathcal{B}},$
- (b) $[[f_1, f_2, f_3]_{\beta}, \beta \circ \phi(f_4), \beta \circ \phi(f_5)]_{\beta} = \beta^2[[f_1, f_2, f_3]_{\mathcal{B}}, \phi(f_4), \phi(f_5)]_{\mathcal{B}},$
- (c) $\rho(f_1 + f_2, f_3)[\beta \circ \phi(f_3), [f_1, f_2, f_4]_{\beta}, \beta \circ \phi(f_5)]_{\beta} = \rho(f_1 + f_2, f_3)\beta^2[\phi(f_3), [f_1, f_2, f_4]_{\mathcal{B}}, \phi(f_5)]_{\mathcal{B}}$
- (d) $\rho(f_1 + f_2, f_3 + f_4)[\beta \circ \phi(f_3), \beta \circ \phi(f_4), [f_1, f_2, f_5]_{\beta}]_{\beta} = \rho(f_1 + f_2, f_3 + f_4)\beta^2[\phi(f_3), \phi(f_4), [f_1, f_2, f_5]_{\mathcal{B}}]_{\mathcal{B}}.$

According to the above four items and this fact that \mathcal{B} is a 3-Hom- ρ -Lie algebra, the proof is complete. \square

Example 5.6. Consider $\mathcal{B}_{\rho, \phi}$ and let ϕ be a 3- ρ -Lie algebra morphism. Then $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\phi} = \phi \circ [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi)$ is a multiplicative 3-Hom- ρ -Lie algebra.

Definition 5.7. A non-degenerate bilinear form χ on $\mathcal{B}_{\rho, \phi}$ satisfying

- i. $\chi(f_1, f_2) = \rho(f_1, f_2)\chi(f_2, f_1) \quad \forall f_1, f_2 \in Hg(\mathcal{B})$ (ρ -symmetric),
- ii. $\chi([f_1, f_2, f_3]_{\mathcal{B}}, f_4) = \chi(f_1, [f_2, f_3, f_4]_{\mathcal{B}}) \quad \forall f_1, f_2, f_3, f_4 \in \mathcal{B}$ (invariant),
- iii. $\chi(\phi(f_1), f_2) = \chi(f_1, \phi(f_2)) \quad (\phi \text{ is } \chi\text{-symmetric}).$

is called a quadratic structure on $\mathcal{B}_{\rho, \phi}$. A quadratic 3-Hom- ρ -Lie algebra is a 3-Hom- ρ -Lie algebra equipped with a quadratic structure. We denote a quadratic 3-Hom- ρ -Lie algebra by $\mathcal{B}_{\rho, \phi, \chi}$. Also, We recover quadratic 3- ρ -Lie algebras when $\phi = id_{\mathcal{B}}$.

Consider the subalgebra $\text{Cent}(\mathcal{B}_{\rho})$ of $\text{End}(\mathcal{B}_{\rho})$ defined by 2.3, we have the following:

Proposition 5.8. Let ψ be an even element in $\text{Cent}(\mathcal{B}_{\rho, \phi})$. For any $f, g, h \in Hg(\mathcal{B})$, we define two following structures

$$[f, g, h]^{\psi} = [\psi(f), g, h]_{\mathcal{B}}, \quad [f, g, h]_{\psi}^{\psi} = [\psi(f), \psi(g), \psi(h)]_{\mathcal{B}}.$$

Then, we have the following 3-Hom- ρ -Lie algebras

$$\begin{aligned} &(\mathcal{B}, [\cdot, \cdot, \cdot], \rho, \psi \circ \phi), \quad (\mathcal{B}, [\cdot, \cdot, \cdot]^\psi, \rho, \phi), \quad (\mathcal{B}, [\cdot, \cdot, \cdot]^\psi, \rho, \psi \circ \phi), \\ &(\mathcal{B}, [\cdot, \cdot, \cdot]^\psi, \rho, \phi), \quad (\mathcal{B}, [\cdot, \cdot, \cdot]^\psi, \rho, \psi \circ \phi). \end{aligned}$$

Proof. This proposition will be proved in the similar way of Proposition 2.5. \square

Proposition 5.9. *Let ψ be an even invertible and χ -symmetric element of the centroid of $\mathcal{B}_{\rho, \phi, \chi}$ such that $\psi \circ \phi = \phi \circ \psi$. Then \mathcal{B} together with two structures $[\cdot, \cdot, \cdot]^\psi, [\cdot, \cdot, \cdot]^\psi$ and the even bilinear form χ_ψ defined by $\chi_\psi(f, g) = \chi(\psi(f), g)$ makes two quadratic 3-Hom- ρ -Lie algebras.*

Proof. According to the proof of Proposition 2.6, it is enough to check that $\chi_\psi(\phi(f), g) = \chi_\psi(f, \phi(g))$. For this, we have

$$\chi_\psi(\phi(f), g) = \chi(\psi \circ \phi(f), g) = \chi(\phi \circ \psi(f), g) = \chi(\psi(f), \phi(g)) = \chi_\psi(f, \phi(g)).$$

\square

Definition 5.10. *Let g be a G -graded vector space, $\mu : g \times g \times g \longrightarrow g$ be an even 3-linear map (i.e. $\mu(g_a, g_b, g_c) \subset g_{a+b+c}$) and $\alpha : g \longrightarrow g$ be an even homomorphism such that for elements $a, b, c, d, e \in g$*

$$\mu(\alpha(a), \alpha(b), \mu(c, d, e)) = \mu(\mu(a, b, c), \alpha(d), \alpha(e)),$$

then (g, μ, α) is called a 3-Hom-associative algebra. If the bilinear map μ is the ρ -symmetry with respect to the displacement of every two elements, the 3-Hom-associative algebra (g, μ, α) is called commutative. We denote this algebra by $g_{\rho, \phi, C}$.

Definition 5.11. *A quadratic structure on 3-Hom-associative algebra g is a ρ -symmetric, invariant and non-degenerate bilinear form χ_g on g such that α is χ_g -symmetric. Let us denoted by $g_{\rho, \chi}$ a quadratic 3-Hom-associative algebra.*

Theorem 5.12. *Consider $g_{\rho, \phi, \chi, C}$ and $\mathcal{B}_{\rho, \phi, \chi}$. Then the tensor product $E = \mathcal{B} \otimes g$ equipped with the structures*

$$\begin{aligned} [f \otimes a, g \otimes b, h \otimes c]_E &= \rho(a, g + h)\rho(b, h)[f, g, h]_{\mathcal{B}} \otimes \mu(a, b, c), \\ \chi_E(f \otimes a, g \otimes b) &= \rho(a, g)\chi_A(f, g)\chi_g(a, b), \\ \phi_E(f \otimes a, g \otimes b) &= \phi(f) \otimes \alpha(a), \quad \forall f, g, h \in Hg(\mathcal{B}), a, b, c \in Hg(g), \end{aligned}$$

is a quadratic 3-Hom- ρ -Lie algebra.

Proof. By the proof of Theorem 2.9, we can easily show that $(E, [\cdot, \cdot, \cdot]_E, \rho, \phi_E)$ is a 3-Hom- ρ -Lie algebra and χ_E is ρ -symmetric and invariant. It is enough to check that ϕ_E is χ_E -symmetric, which completes the proof. For this, we have

$$\begin{aligned} \chi_E(\phi_E(f \otimes a), g \otimes b) &= \chi_E(\phi(f) \otimes \alpha(a), g \otimes b) = \rho(a, g)\chi(\phi(f), g)\chi_g(\alpha(a), b) \\ &= \rho(a, g)\chi(f, \phi(g))\chi_g(a, \alpha(b)) = \chi_E(f \otimes a, \phi_E(g \otimes b)). \end{aligned}$$

\square

Ideals of 3-Hom- ρ -Lie algebras: In this part, we introduce the definitions of subalgebra and ideal of $\mathcal{B}_{\rho, \phi}$ and give some properties related to ideals of $\mathcal{B}_{\rho, \phi}$.

Definition 5.13. A Hom-subalgebra of $\mathcal{B}_{\rho,\phi}$ is defined as a sub-vector space $I \subseteq \mathcal{B}_{\rho,\phi}$ with the property $[I, I, I]_{\mathcal{B}} \subseteq I$ and $\phi(I) \subseteq I$. I also is called a Hom ideal of \mathcal{B} if $\phi(I) \subseteq I$ and $[I, \mathcal{B}_{\rho,\phi}, \mathcal{B}_{\rho,\phi}]_{\mathcal{B}} \subseteq I$.

Lemma 5.14. $Z(\mathcal{B}_{\rho,\phi})$, the center of multiplicative quadratic 3-Hom- ρ -Lie algebra $\mathcal{B}_{\rho,\phi,\chi}$, is an Hom-ideal of $\mathcal{B}_{\rho,\phi}$.

Proof. It is easy to see that $[Z(\mathcal{B}_{\rho,\phi}), \mathcal{B}_{\rho,\phi}, \mathcal{B}_{\rho,\phi}] \subseteq \mathcal{B}_{\rho,\phi}$. We assume that $f \in Z(\mathcal{B}_{\rho,\phi})$ and $g, h, m \in \mathcal{B}$, so

$$\begin{aligned} \chi([\phi(f), g, h]_{\mathcal{B}}, m) &= \chi(\phi(f), [g, h, m]_{\mathcal{B}}) = \chi(f, \phi[g, h, m]_{\mathcal{B}}) \\ &= \chi(f, [\phi(g), \phi(h), \phi(m)]_{\mathcal{B}}) = \chi([f, \phi(g), \phi(h)]_{\mathcal{B}}, \phi(m)) = 0. \end{aligned}$$

The last equality holds, since $f \in Z(\mathcal{B}_{\rho,\phi})$. On the other hand, since χ is non-degenerate, thus the above relation gives $[\phi(f), g, h]_{\mathcal{B}} = 0$ for any $g, h \in \mathcal{B}$. This implies that $\phi(f) \in Z(\mathcal{B}_{\rho,\phi})$ and therefore $Z(\mathcal{B}_{\rho,\phi})$ is an Hom-ideal of $\mathcal{B}_{\rho,\phi}$. \square

Lemma 5.15. The orthogonal I^\perp of I (I is a Hom-ideal of $\mathcal{B}_{\rho,\phi,\chi}$) with respect to χ is a Hom-ideal of $\mathcal{B}_{\rho,\phi}$.

Proof. It is clear that $[I^\perp, \mathcal{B}_{\rho,\phi}, \mathcal{B}_{\rho,\phi}] \subseteq I^\perp$. Also, for $f \in I^\perp$, $g \in I$, we have $B(\phi(f), g) = B(f, \phi(g)) = 0$, since $\phi(I) \subseteq I$ and $f \in I^\perp$. \square

Symplectic structure of 3-Hom- ρ -Lie algebras: This part is devoted to study of symplectic structure, metrics and their properties. Also, we show that a symplectic structure ω may be defined on $\mathcal{B}_{\rho,\phi}$ if and only if there exists an invertible derivation of $\mathcal{B}_{\rho,\phi}$ that is antisymmetric with respect to metric.

Definition 5.16. If $\omega \in \wedge^2 \mathcal{B}^*$ is a non-degenerate and ρ -skew-symmetric bilinear form such that

$$\begin{aligned} \omega([f_1, f_2, f_3]_{\mathcal{B}}, \phi(f_4)) - \rho(f_1, f_2 + f_3 + f_4)\omega([f_2, f_3, f_4]_{\mathcal{B}}, \phi(f_1)) \\ + \rho(f_1 + f_2, f_3 + f_4)\omega([f_3, f_4, f_1]_{\mathcal{B}}, \phi(f_2)) - \rho(f_1 + f_2 + f_3, f_4)\omega([f_4, f_1, f_2]_{\mathcal{B}}, \phi(f_3)) = 0, \end{aligned}$$

then ω is called a symplectic structure and $(\mathcal{B}, [., ., .]_{\mathcal{B}}, \rho, \phi, \omega)$ is said to be a symplectic 3-Hom- ρ -Lie algebra.

We will say that $(\mathcal{B}, [., ., .]_{\mathcal{B}}, \rho, \phi, \chi, \omega)$ is a quadratic symplectic 3-Hom- ρ -Lie algebra if (\mathcal{B}, χ) is quadratic and (\mathcal{B}, ω) is symplectic.

Example 5.17. Let $(\mathcal{B}, [., ., .]_{\mathcal{B}}, \rho, \omega)$ be the 4-dimensional symplectic 3- ρ -Lie algebra provided in Example 2.15. We consider the multiplex $(\mathcal{B}, [., ., .]_{\phi}, \rho, \phi)$ defining a 3-Hom- ρ -Lie algebra, according to Theorem 5.5, such that the bracket $[., ., .]_{\phi}$ and the even linear map ϕ are defined by

$$\begin{aligned} [l_1, l_2, l_3]_{\phi} &= l_4, & \phi(l_4) &= l_4, \\ [l_1, l_2, l_4]_{\phi} &= l_3, & \phi(l_3) &= -l_3, \\ [l_1, l_3, l_4]_{\phi} &= l_2, & \phi(l_2) &= -l_2, \\ [l_2, l_3, l_4]_{\phi} &= l_1, & \phi(l_1) &= -l_1. \end{aligned}$$

Moreover, $(\mathcal{B}, [., ., .]_{\phi}, \rho, \phi)$ together with the bilinear map ω is given in Example 2.15 is a 4-dimensional symplectic 3-Hom- ρ -Lie algebra.

In the following, we define a derivation of $\mathcal{B}_{\rho,\phi}$ and examine the relationship between symplectic structures and φ -symmetric invertible derivations.

Definition 5.18. A linear map D on $\mathcal{B}_{\rho,\phi}$ satisfying

$$D \circ \phi = \phi \circ D,$$

$$D[f, g, h]_{\mathcal{B}} = [D(f), g, h] + \rho(D, f)[f, D(g), h] + \rho(D, f + g)[f, g, D(h)],$$

is called a derivation of $\mathcal{B}_{\rho,\phi}$. We denote by $\text{Der}(\mathcal{B}_{\rho,\phi})$ the space of all derivations of $\mathcal{B}_{\rho,\phi}$.

Definition 5.19. $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi, \varphi, \mathbb{K})$ is called a metric 3-Hom- ρ -Lie algebra, if

- i. \mathbb{K} is a field (\mathbb{C} or \mathbb{R}),
- ii. $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi)$ is a 3-Hom- ρ -Lie algebra,
- iii. $\varphi : \mathcal{B} \times \mathcal{B} \longrightarrow \mathbb{K}$ is a non-degenerate ρ -symmetric bilinear form on \mathcal{B} satisfying the following condition

$$\varphi([f, g, h]_{\mathcal{B}}, \phi(z)) + \rho(f + g, h)\varphi(\phi(h), [f, g, z]_{\mathcal{B}}) = 0, \quad \forall f, g, h, z \in Hg(\mathcal{B}).$$

Let us denote by $\mathcal{B}_{\rho,\phi,\varphi}$ the metric 3-Hom- ρ -Lie algebra $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi, \varphi, \mathbb{K})$.

Proposition 5.20. There exists a symplectic structure on $\mathcal{B}_{\rho,\phi}$ if and only if there exists an invertible derivation $D \in \text{Der}_{\varphi}(\mathcal{B}_{\rho,\phi})$.

Proof. It is enough to define $\omega(f, g) = \varphi(D(f), g)$. □

For any symmetric bilinear form χ' on $\mathcal{B}_{\rho,\phi,\chi}$, there is an associated map $D : \mathcal{B} \longrightarrow \mathcal{B}$ satisfying

$$\chi'(f, g) = \chi(D(f), g), \quad \forall f, g \in Hg(\mathcal{B}). \quad (5.1)$$

Since χ and χ' are symmetric, then D is symmetric with respect to χ , i.e.,

$$\chi(D(f), g) = \rho(D, f)\chi(f, D(g)), \quad \forall f, g \in Hg(\mathcal{B}).$$

Lemma 5.21. Let $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi, \chi)$ be a quadratic 3-Hom- ρ -Lie algebra and χ' defined by (5.1). Then χ' is invariant if and only if

$$D([f, g, h]_{\mathcal{B}}) = [D(f), g, h]_{\mathcal{B}} = \rho(D, f)[f, D(g), h]_{\mathcal{B}} = \rho(D, f + g)[f, g, D(h)]_{\mathcal{B}}. \quad (5.2)$$

Also, χ' is non-degenerate if and only if D is invertible and ϕ is χ' -symmetric if and only if $D \circ \phi = \phi \circ D$.

Proof. By the proof of Lemma 2.20, it is enough to show that $\chi'(\phi(f), g) = \chi'(f, \phi(g))$. For this, we have

$$\chi'(\phi(f), g) = \chi(D \circ \phi(f), g) = \chi(\phi \circ D(f), g) = \chi(D(f), \phi(g)) = \chi'(f, \phi(g)).$$

□

Definition 5.22. A ρ -symmetric map $D : \mathcal{B}_{\rho,\phi} \longrightarrow \mathcal{B}_{\rho,\phi}$ satisfying (2.10) is called a centromorphism of $\mathcal{B}_{\{\cdot, \cdot, \cdot\}, \phi}$. We denote by $\mathcal{C}(\mathcal{B}_{\rho,\phi})$ the space of all centromorphisms of $\mathcal{B}_{\rho,\phi}$.

Proposition 5.23. Let $\delta \in \text{Der}_{\chi}(\mathcal{B}_{\rho,\phi})$ and $D \in \mathcal{C}(\mathcal{B}_{\rho,\phi})$ such that $D \circ \delta = \delta \circ D$ and $D \circ \phi = \phi \circ D$. Then $D \circ \delta$ is also a χ -antisymmetric derivation of $\mathcal{B}_{\rho,\phi}$.

Proof. By the proof of Proposition 2.22, it is enough to show that $(D \circ \delta) \circ \phi = \phi \circ (D \circ \delta)$. Since $\delta \circ \phi = \phi \circ \delta$, we have

$$(D \circ \delta) \circ \phi = D \circ (\delta \circ \phi) = D \circ (\phi \circ \delta) = (D \circ \phi) \circ \delta = (\phi \circ D) \circ \delta = \phi \circ (D \circ \delta).$$

□

5.1. Representation of 3-Hom- ρ -Lie algebras. The representation theory of 3-Hom- ρ -Lie algebras is studied in [31]. In this part, we recall main definition of representation and some properties, then we add a brief details which are not discussed in that article.

Consider the multiplicative 3-Hom- ρ -Lie algebra $\mathcal{B}_{\rho,\phi}$. The fundamental set $\mathcal{L} = \wedge^2 \mathcal{B}_{\rho,\phi}$ together with operation

$$[(f_1, f_2), (g_1, g_2)]_{\mathcal{L}} = ([f_1, f_2, g_1], \phi(g_2)) + \rho(f_1 + f_2, g_1)(\phi(g_1), [f_1, f_2, g_2]), \quad (5.3)$$

and the even linear map $\phi_1 : \mathcal{L} \rightarrow \mathcal{L}$ defined by $\phi_1(f_1, f_2) = (\phi(f_1), \phi(f_2))$ is the multiplicative Hom- ρ -Lie algebra.

Definition 5.24. [31] *A bilinear map $\mu : \mathcal{L} \rightarrow \mathfrak{gl}(V)$ is called a representation of $\mathcal{B}_{\rho,\phi}$ on vector space V with respect to $\beta \in \mathfrak{gl}(V)$ if the following equalities are satisfied*

$$\mu([(f_1, f_2), (g_1, g_2)]_{\mathcal{L}}) \circ \beta = \mu(\phi_1(f_1, f_2))\mu(g_1, g_2) - \rho(f_1 + f_2, g_1 + g_2)\mu(\phi_1(g_1, g_2))\mu(f_1, f_2), \quad (5.4)$$

$$\begin{aligned} \mu([g_1, g_2, g_3], \phi(f)) \circ \beta &= \mu(\phi_1(g_1, g_2))\mu(g_3, f) + \rho(g_1, g_2 + g_3)\mu(\phi_1(g_2, g_3))\mu(g_1, f) \\ &\quad + \rho(g_1 + g_2, g_3)\mu(\phi_1(g_3, g_1))\mu(g_2, f), \end{aligned} \quad (5.5)$$

$$\begin{aligned} \mu(\phi(g), [f_1, f_2, f_3]) \circ \beta &= \rho(g, f_1 + f_2)\mu(\phi_1(f_1, f_2))\mu(g, f_3) + \rho(g, f_2 + f_3)\rho(f_1, f_2 + f_3)\mu(\phi_1(f_2, f_3))\mu(g, f_1) \\ &\quad + \rho(g, f_1 + f_3)\rho(f_1 + f_2, f_3)\mu(\phi_1(f_3, f_1))\mu(g, f_2). \end{aligned} \quad (5.6)$$

If $\mathcal{B}_{\rho,\phi}$ is multiplicative, in addition to the above conditions, the following condition must be hold too

$$\mu(\phi(f), \phi(g)) \circ \beta = \beta \circ \mu(f, g).$$

Lemma 5.25. *Let (V, μ, β) be a representation of $\mathcal{B}_{\rho,\phi}$. Then, we have*

$$\begin{aligned} 0 &= \rho(f_1 + f_2, g_1)\mu(\phi(g_1), [f_1, f_2, g_2]_{\mathcal{B}}) \circ \beta + \mu([f_1, f_2, g_1]_{\mathcal{B}}, \phi(g_2)) \circ \beta \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\rho(g_1 + g_2, f_1)\mu(\phi(f_1), [g_1, g_2, f_2]_{\mathcal{B}}) \circ \beta \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\mu([g_1, g_2, f_1]_{\mathcal{B}}, \phi(f_2)) \circ \beta. \end{aligned}$$

Proof. For the proof refer to Lemma 3.3. □

If $V = \mathcal{B}_{\rho,\phi}$ and $\phi = \beta \in \mathfrak{gl}(\mathcal{B}_{\rho,\phi})$, then the bilinear map $ad : \mathcal{B} \times \mathcal{B} \rightarrow \mathfrak{gl}(\mathcal{B})$ defined by $ad(f_1, f_2)(f_3) = [f_1, f_2, f_3]_{\mathcal{B}}$ is a representation of $\mathcal{B}_{\rho,\phi}$ with respect to $\beta = \phi$.

Proposition 5.26. *Consider the 3-Hom- ρ -Lie algebra $\mathcal{B}_{\rho,\phi}$. Let V be a graded vector space, $\beta \in \mathfrak{gl}(V)$ and $\mu : \wedge^2 \mathcal{B} \rightarrow \mathfrak{gl}(V)$ be a bilinear map. Then (V, μ, β) is a representation of $\mathcal{B}_{\rho,\phi}$ if and only if $\mathcal{B} \oplus V$ is a 3-Hom- ρ -Lie algebra with the following structures*

$$\begin{aligned} [f_1 + v_1, f_2 + v_2, f_3 + v_3]_{\mathcal{B} \oplus V}^{\mu} &= [f_1, f_2, f_3]_{\mathcal{B}} + \mu(f_1, f_2)v_3 \\ &\quad + \rho(f_1, f_2 + f_3)\mu(f_2, f_3)v_1 + \rho(f_1 + f_2, f_3)\mu(f_3, f_1)v_2, \\ \psi(f_1 + v_1) &= \phi(f_1) + \beta(v_1), \end{aligned}$$

where $f_1, f_2, f_3 \in Hg(\mathcal{B})$ and $v_1, v_2, v_3 \in Hg(V)$.

Consider the representation (V, μ, β) of $\mathcal{B}_{\rho, \phi}$ and V^* as the dual of vector space V . We define bilinear map $\tilde{\mu} : \mathcal{B} \times \mathcal{B} \longrightarrow \text{End}(V^*)$ by $\tilde{\mu}(f_1, f_2)(\varrho) = -\rho(f_1 + f_2, \varrho)\varrho \circ \mu(f_1, f_2)$, where $f_1, f_2 \in Hg(\mathcal{B})$, $\varrho \in V^*$ and set $\tilde{\beta}(\varrho) = \varrho \circ \beta$.

Note that by Definition 3.5 and Lemma 5.25, we can deduce that if (V, μ, β) is a ρ -skew symmetric representation, then $(V^*, \tilde{\mu}, \tilde{\beta})$ is a representation of $\mathcal{B}_{\rho, \phi}$.

By abuse the above notation, (\mathcal{B}^*, ad^*) with respect to $\tilde{\beta}(\varrho) = \varrho \circ \phi$ is a representation of $\mathcal{B}_{\rho, \phi}$. So by Proposition 5.26, $\mathcal{B} \oplus \mathcal{B}^*$ is a 3-Hom- ρ -Lie algebra together with the following structure

$$\begin{aligned} [f_1 + \alpha_1, f_2 + \alpha_2, f_3 + \alpha_3]_{\mathcal{B} \oplus \mathcal{B}^*}^{ad^*} &= [f_1, f_2, f_3]_{\mathcal{B}} + ad^*(f_1, f_2)\alpha_3 \\ &\quad + \rho(f_1, f_2 + f_3)ad^*(f_2, f_3)\alpha_1 + \rho(f_1 + f_2, f_3)ad^*(f_3, f_1)\alpha_2, \\ (\phi + \phi^*)(f + \alpha) &= \phi(f) + \alpha \circ \phi. \end{aligned}$$

Now, consider the bilinear map $\varphi(f + \alpha, g + \beta) = \alpha(g) + \rho(f, g)\beta(f)$. Then $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}^{ad^*}, \rho, \phi + \phi^*, \varphi)$ is a metric 3-Hom- ρ -Lie algebra.

6. 3-PRE-HOM- ρ -LIE ALGEBRAS

This part is devoted to the 3-pre-Hom- ρ -Lie algebras, which are studied similar to the classical case in Section 4. Actually, we give the similar results to Section 4 in the case of 3-pre-Hom- ρ -Lie algebras.

Definition 6.1. $(\mathcal{B}, \{\cdot, \cdot, \cdot\}, \rho, \phi)$ is called a 3-pre-Hom- ρ -Lie algebra if

- i. \mathcal{B} is a G -graded vector space,
- ii. $\phi : \mathcal{B} \longrightarrow \mathcal{B}$ is an even linear map,
- iii. $\{\cdot, \cdot, \cdot\} : \otimes^3 \mathcal{B} \longrightarrow \mathcal{B}$ is a trilinear map satisfying the following relations

$$\begin{aligned} |\{f_1, f_2, f_3\}| &= |f_1| + |f_2| + |f_3|, \\ \{f_1, f_2, f_3\} &= -\rho(f_1, f_2)\{f_2, f_1, f_3\}, \\ \{\phi(f_1), \phi(f_2), \{g_1, g_2, g_3\}\} &= \{\{f_1, f_2, g_1\}_c, \phi(g_2), \phi(g_3)\} + \rho(f_1 + f_2, g_1)\{\phi(g_1), [f_1, f_2, g_2]_c, \phi(g_3)\} \\ &\quad + \rho(f_1 + f_2, g_1 + g_2)\{\phi(g_1), \phi(g_2), \{f_1, f_2, g_3\}\}, \\ \{\{f_1, f_2, f_3\}_c, \phi(g_1), \phi(g_2)\} &= \{\phi(f_1), \phi(f_2), \{f_3, g_1, g_2\}\} + \rho(f_1, f_2 + f_3)\{\phi(f_2), \phi(f_3), \{f_1, g_1, g_2\}\} \\ &\quad + \rho(f_1 + f_2, f_3)\{\phi(f_3), \phi(f_1), \{f_2, g_1, g_2\}\}, \end{aligned}$$

where

$$[f_1, f_2, f_3]_c = \{f_1, f_2, f_3\} + \rho(f_1, f_2 + f_3)\{f_2, f_3, f_1\} + \rho(f_1 + f_2, f_3)\{f_3, f_1, f_2\}.$$

We denoted by $\mathcal{B}_{\{\cdot, \cdot, \cdot\}, \rho, \phi}$ the 3-pre-Hom- ρ -Lie algebra $(\mathcal{B}, \{\cdot, \cdot, \cdot\}, \rho, \phi)$.

For a 3-pre-Hom- ρ -Lie algebra $(\mathcal{B}, \{\cdot, \cdot, \cdot\}, \rho, \phi)$, it is easy to see that $(\mathcal{B}, [\cdot, \cdot, \cdot]_c, \rho, \phi)$ is a 3-Hom- ρ -Lie algebra. We denote it by $\mathcal{B}_{\rho, \phi}^c$.

Let us define two multiplications $L : \wedge^2 \mathcal{B} \longrightarrow gl(\mathcal{B})$ and $R : \otimes^2 \mathcal{B} \longrightarrow gl(\mathcal{B})$ on $\mathcal{B}_{\{\cdot, \cdot, \cdot\}, \rho, \phi}$, which is called respectively the left and right multiplications, by $L(f, g)h = \{f, g, h\}$ and $R(f, g)h = \rho(f + g, h)\{h, f, g\}$. Then (\mathcal{B}, L) is a representation of $\mathcal{B}_{\rho, \phi}^c$ with respect to $\beta = \phi$.

Definition 6.2. The triple $(V, \mu, \tilde{\mu})$ consisting of a G -graded vector space V , a representation μ of $\mathcal{B}_{\rho, \phi}^c$ and a bilinear map $\tilde{\mu} : \otimes^2 \mathcal{B} \longrightarrow gl(V)$ is called a representation of $\mathcal{B}_{\{\cdot, \cdot, \cdot\}, \phi}$ with respect to $\beta \in gl(V)$ if for all homogeneous elements $f_1, f_2, f_3, f_4 \in Hg(\mathcal{B})$, the following equalities hold

$$\begin{aligned} \tilde{\mu}(\phi(f_1), \{f_2, f_3, f_4\}) \circ \beta &= \rho(f_1, f_2 + f_3) \mu(\phi(f_2), \phi(f_3)) \tilde{\mu}(f_1, f_4) \\ &\quad + \rho(f_1, f_3 + f_4) \rho(f_2, f_3 + f_4) \tilde{\mu}(\phi(f_3), \phi(f_4)) \tilde{\mu}(f_1, f_2) \\ &\quad - \rho(f_1, f_2 + f_4) \rho(f_3, f_4) \tilde{\mu}(\phi(f_2), \phi(f_4)) \tilde{\mu}(f_1, f_3) \\ &\quad + \rho(f_1 + f_2, f_3 + f_4) \tilde{\mu}(\phi(f_3), \phi(f_4)) \mu(f_1, f_2) \\ &\quad + \rho(f_1, f_2 + f_3 + f_4) \rho(f_3, f_4) \tilde{\mu}(\phi(f_2), \phi(f_4)) \tilde{\mu}(f_3, f_1) \\ &\quad - \rho(f_1, f_2 + f_4) \rho(f_3, f_4) \tilde{\mu}(\phi(f_2), \phi(f_4)) \mu(f_1, f_3) \\ &\quad - \rho(f_2, f_3 + f_4) \rho(f_1, f_2 + f_3 + f_4) \tilde{\mu}(\phi(f_3), \phi(f_4)) \tilde{\mu}(f_2, f_1), \end{aligned}$$

$$\begin{aligned} \mu(\phi(f_1), \phi(f_2)) \tilde{\mu}(f_3, f_4) &= \rho(f_1 + f_2, f_3 + f_4) \tilde{\mu}(\phi(f_3), \phi(f_4)) \mu(f_1, f_2) \\ &\quad - \rho(f_1 + f_2, f_3 + f_4) \rho(f_1, f_2) \tilde{\mu}(\phi(f_3), \phi(f_4)) \tilde{\mu}(f_2, f_1) \\ &\quad + \rho(f_1 + f_2, f_3 + f_4) \tilde{\mu}(\phi(f_3), \phi(f_4)) \tilde{\mu}(f_1, f_2) + \tilde{\mu}([f_1, f_2, f_3]_{\mathcal{B}}, \phi(f_4)) \circ \beta \\ &\quad + \rho(f_1 + f_2, f_3) \tilde{\mu}(\phi(f_3), \{f_1, f_2, f_4\}), \end{aligned}$$

$$\begin{aligned} \tilde{\mu}([f_1, f_2, f_3]_{\mathcal{B}}, \phi(f_4)) \circ \beta &= \mu(\phi(f_1), \phi(f_2)) \tilde{\mu}(f_3, f_4) + \rho(f_1, f_2 + f_3) \mu(\phi(f_2), \phi(f_3)) \tilde{\mu}(f_1, f_4) \\ &\quad + \rho(f_1 + f_2, f_3) \mu(\phi(f_3), \phi(f_1)) \tilde{\mu}(f_2, f_4), \end{aligned}$$

$$\begin{aligned} \tilde{\mu}(\phi(f_3), \phi(f_4)) \mu(f_1, f_2) &= \rho(f_1, f_2) \mu(\phi(f_3), \phi(f_4)) \tilde{\mu}(f_2, f_1) - \tilde{\mu}(\phi(f_3), \phi(f_4)) \tilde{\mu}(f_1, f_2) \\ &\quad + \rho(f_3 + f_4, f_1 + f_2) \mu(\phi(f_1), \phi(f_2)) \tilde{\mu}(f_3, f_4) \\ &\quad - \rho(f_3 + f_4, f_1 + f_2) \rho(f_1, f_2) \tilde{\mu}(\phi(f_2), \{f_1, f_3, f_4\}) \circ \beta \\ &\quad + \rho(f_3 + f_4, f_1 + f_2) \tilde{\mu}(\phi(f_1), \{f_2, f_3, f_4\}) \circ \beta. \end{aligned}$$

For example $(\mathcal{B}, L, R, \phi)$ is a representation of $\mathcal{B}_{\{\cdot, \cdot, \cdot\}, \phi}$. Let us to define the operation $\{., ., .\}_{\mu, \tilde{\mu}} : \otimes^2(\mathcal{B} \oplus V) \longrightarrow \mathcal{B} \oplus V$ and the linear map $\psi : \mathcal{B} \oplus V \longrightarrow \mathcal{B} \oplus V$ by

$$\begin{aligned} \{f_1 + v_1, f_2 + v_2, f_3 + v_3\}_{\mu, \tilde{\mu}} &= \{f_1, f_2, f_3\} + \mu(f_1, f_2) v_3 + \rho(f_1, f_2 + f_3) \tilde{\mu}(f_2, f_3) v_1 - \rho(f_2, f_3) \tilde{\mu}(f_1, f_3) v_2, \\ \psi(f_1 + v_1) &= \phi(f_1) + \beta(v_1), \end{aligned}$$

for all $f_1, f_2, f_3 \in Hg(\mathcal{B})$ and $v_1, v_2, v_3 \in V$. Then $(\mathcal{B} \oplus V, \{., ., .\}_{\mu, \tilde{\mu}}, \rho, \psi)$ is a 3-pre-Hom- ρ -Lie algebra.

Consider the operator $\zeta : \otimes^2 V \longrightarrow \otimes^2 V$ given by $\zeta(f_1 \otimes f_2) = f_2 \otimes f_1$, where V is a graded vector space and $f_1 \otimes f_2 \in \otimes^2 V$.

Using the above notation, in the following proposition, we give a representation of $\mathcal{B}_{\rho, \phi}^c$:

Proposition 6.3. Let $(V, \mu, \tilde{\mu}, \beta)$ be a representation of $\mathcal{B}_{\{\cdot, \cdot, \cdot\}, \phi}$. Then there exists a representation $\nu : \wedge^2 \mathcal{B} \longrightarrow gl(V)$ of $\mathcal{B}_{\rho, \phi}^c$ on V that is given by $\nu(f_1, f_2) = (\mu - \rho(f_1, f_2) \tilde{\mu} \xi + \tilde{\mu})(f_1, f_2)$ for all $f_1, f_2 \in Hg(\mathcal{B})$.

The above proposition conclude that if $(V, \mu, \tilde{\mu})$ is a representation of $\mathcal{B}_{\{\}, \phi}$, then the 3-pre-Hom- ρ -Lie algebras $(\mathcal{B} \oplus V, \{., ., .\}_{\mu, \tilde{\mu}}, \rho, \psi)$ and $(\mathcal{B} \oplus V, \{., ., .\}_{\nu, 0}, \rho, \psi)$ have the same sub-adjacent 3-Hom- ρ -Lie algebra given by (4.5).

Let us define the dual of the representation (V, ν) by $\nu^*(f_1, f_2) = \mu^* - \rho(f_1, f_2)\tilde{\mu}^*\xi + \tilde{\mu}^*(f_1, f_2)$. In the following proposition, we will talk about the dual of the representation $(V, \mu, \tilde{\mu})$:

Proposition 6.4. *Consider $\mathcal{B}_{\{\}, \phi}$ equipped with the ρ -skew-symmetric representation $(V, \mu, \tilde{\mu}, \beta)$. Then $(V^*, \nu^*, -\tilde{\mu}^*, \beta)$ is a representation of $\mathcal{B}_{\{\}, \phi}$, which is called the dual representation of $(V, \mu, \tilde{\mu})$.*

Remark 6.5. *Consider $\mathcal{B}_{\{\}, \phi}$ as a 3-pre-Hom- ρ -Lie algebra with the ρ -skew-symmetric representation $(V, \mu, \tilde{\mu}, \beta)$. Then the 3-pre-Hom- ρ -Lie algebras $(\mathcal{B} \oplus V^*, \{., ., .\}_{\mu^*, 0}, \rho, \psi)$ and $(\mathcal{B} \oplus V^*, \{., ., .\}_{\nu^*, -\tilde{\mu}^*}, \rho, \psi)$ have the same sub-adjacent 3-Hom- ρ -Lie algebra $(\mathcal{B}^c \oplus V^*, [., ., .]_{\mathcal{B}^c}^{\mu^*}, \rho, \phi)$ with the representation $(V^*, \mu^*, \tilde{\beta})$.*

In the following definition, we define the ρ -Hom- \mathcal{O} -operator associated to representation (V, μ, β) . The importance of defining this operator is that it can be used to construct a 3-pre-Hom- ρ -Lie algebra structure on representation V .

Definition 6.6. *Let (V, μ, β) be a representation of $\mathcal{B}_{\rho, \phi}$. If an even linear operator $T : V \longrightarrow \mathcal{B}_{\rho, \phi}$ satisfies*

$$T \circ \beta = \phi \circ T,$$

$$[Tx, Ty, Tz]_{\mathcal{B}} = T(\mu(Tx, Ty)z + \rho(x, y + z)(\mu(Ty, Tz)x + \rho(x + y, z)\mu(Tz, Tx)y), \quad \forall x, y, z \in V,$$

it is called a ρ -Hom- \mathcal{O} -operator associated to representation (V, μ, β) .

Lemma 6.7. *Let (V, μ, β) be a ρ -skew-symmetric representation of $\mathcal{B}_{\rho, \phi}$ and $T : V \longrightarrow \mathcal{B}_{\rho, \phi}$ is a ρ -Hom- \mathcal{O} -operator associated to (V, μ, β) . Then $(V, \{., ., .\}, \rho, \psi)$ is a 3-pre-Hom- ρ -Lie algebra, where*

$$\psi = \beta, \tag{6.1}$$

$$\{x, y, z\} = \mu(Tx, Ty)z, \quad \forall x, y, z \in V. \tag{6.2}$$

Proof. Suppose that $x, y, z \in Hg(V)$. It is easy to see that

$$\{x, y, z\} = -\rho(x, y)\{y, x, z\}, \quad T[x, y, z]_c = [Tx, Ty, Tz].$$

Also, for $x_1, x_2, x_3, x_4, x_5 \in Hg(V)$, we have

$$\begin{aligned} \{\psi(x_1), \psi(x_2), \{x_3, x_4, x_5\}\} &= \mu(T \circ \psi(x_1), T \circ \psi(x_2))\mu(Tx_3, Tx_4)x_5 \\ &= \mu(\phi \circ T(x_1), \phi \circ T(x_2))\mu(Tx_3, Tx_4)x_5, \end{aligned} \tag{6.3}$$

$$\begin{aligned} \{[x_1, x_2, x_3]_c, \psi(x_4), \psi(x_5)\} &= \mu([Tx_1, Tx_2, Tx_3], T \circ \psi(x_4))x_5 \\ &= \mu([Tx_1, Tx_2, Tx_3], \phi \circ T(x_4))x_5, \end{aligned} \tag{6.4}$$

$$\begin{aligned} \rho(x_1 + x_2, x_3)\{\psi(x_3), [x_1, x_2, x_4]_c, \psi(x_5)\} &= \rho(x_1 + x_2, x_3)\mu(T \circ \psi(x_3), [Tx_1, Tx_2, Tx_4])x_5 \\ &= \rho(x_1 + x_2, x_3)\mu(\phi \circ T(x_3), [Tx_1, Tx_2, Tx_4])x_5, \end{aligned} \tag{6.5}$$

$$\begin{aligned} \rho(x_1 + x_2, x_3 + x_4)\{\psi(x_3), \psi(x_4), \{x_1, x_2, x_5\}\} &= \rho(x_1 + x_2, x_3 + x_4)\mu(T \circ \psi(x_3), T \circ \psi(x_4))\mu(Tx_1, Tx_2)x_5 \\ &= \rho(x_1 + x_2, x_3 + x_4)\mu(\phi \circ T(x_3), \phi \circ T(x_4))\mu(Tx_1, Tx_2)x_5. \end{aligned} \tag{6.6}$$

By Definition 5.24 and the equalities (6.3)–(6.6), we get the result. \square

Lemma 6.8. *Consider $\mathcal{B}_{\rho,\phi}$ as a 3-Hom- ρ -Lie algebra. Then the following assertions are equivalent*

- i. *existence of a compatible 3-pre-Hom- ρ -Lie algebra structure,*
- ii. *existence of an invertible ρ - \mathcal{O} -operator.*

Proof. Refer to the proof of Lemma 4.12. □

Phase Space: In the following, we extend the study of phase space of a 3- ρ -Lie algebra to 3-Hom- ρ -Lie algebras and show that a 3-Hom- ρ -Lie algebra has a phase space if and only if it is sub-adjacent to a 3-pre-Hom- ρ -Lie algebra.

Proposition 6.9. *Consider the symplectic 3-Hom- ρ -Lie algebra $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi, \omega)$. Then the following structure defines a compatible 3-pre-Hom- ρ -Lie algebra structure on \mathcal{B}*

$$\omega(\{f, g, h\}, \phi(s)) = -\rho(f + g, h)\omega(\phi(h), [f, g, s]_{\mathcal{B}}), \quad \forall f, g, h, s \in Hg(\mathcal{B}).$$

Proof. Let us define a linear map $T : \mathcal{B}^* \longrightarrow \mathcal{B}$ by $\omega(\phi(f), g) = \omega(f, \phi(g)) = (T^{-1}(f))(g)$, or equivalently, $\omega(T\alpha, \phi(g)) = \alpha(g)$ for all $f, g \in Hg(\mathcal{B})$ and $\alpha \in \mathcal{B}^*$. Since ω is a symplectic structure, we deduce that T is an invertible ρ - \mathcal{O} -operator associated to (\mathcal{B}^*, Ad^*) . Also, by (4.6) and (6.2), there exists a compatible 3- ρ -pre-Lie algebra on B given by $\{f, g, h\} = T(ad^*(f, g)T^{-1}(h))$. So

$$\begin{aligned} \omega(\{f, g, h\}, \phi(s)) &= \omega(T(ad^*(f, g)T^{-1}(h)), \phi(s)) = ad^*(f, g)T^{-1}(h)(s) \\ &= -\rho(f + g, h)T^{-1}(h)[f, g, s]_{\mathcal{B}} = -\rho(f + g, h)\omega(\phi(h), [f, g, s]_{\mathcal{B}}). \end{aligned}$$

□

Definition 6.10. $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \phi + \phi^*, \omega)$ is called a phase space of the 3-Hom- ρ -Lie algebra \mathcal{B} , if

- i. $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \phi + \phi^*)$ is a 3-Hom- ρ -Lie algebra,
- ii. ω is a natural non-degenerate skew-symmetric bilinear form on $\mathcal{B} \oplus \mathcal{B}^*$ given by

$$\omega(f + \alpha, g + \beta) = \alpha(g) - \rho(f, g)\beta(f), \tag{6.7}$$

such that $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \phi + \phi^*, \omega)$ is a symplectic 3-Hom- ρ -Lie algebra,

- iii. $(\mathcal{B}, [\cdot, \cdot, \cdot]_{\mathcal{B}}, \rho, \phi)$ and $(\mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B}^*}, \rho, \phi^*)$ are 3-Hom- ρ -Lie subalgebras of $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}, \rho, \phi + \phi^*)$.

Theorem 6.11. *A 3-Hom- ρ -Lie algebra has a phase space if and only if it is sub-adjacent to a 3-pre-Hom- ρ -Lie algebra.*

Proof. By Theorem 4.15, it is easy to get the result. □

Lemma 6.12. *Let (V, μ, β) be a representation of $\mathcal{B}_{\rho,\phi}$ and $(\mathcal{B} \oplus \mathcal{B}^*, [\cdot, \cdot, \cdot]_{\mathcal{B} \oplus \mathcal{B}^*}^{\mu^*}, \rho, \phi + \phi^*, \omega)$ be a phase space of it. Then the relation $\{f, g, h\} = \mu(f, g)h$ constructs a 3-pre-Hom- ρ -Lie algebra structure on \mathcal{B} if $\beta = \phi$.*

Proof. Assume that $f, g, h \in Hg(\mathcal{B})$ and $\alpha \in Hg(\mathcal{B}^*)$. Then, we have

$$\begin{aligned}
 \rho(f + g + h, \alpha) \alpha \circ \phi(\{f, g, h\}) &= -\omega(\{f, g, h\}, \alpha \circ \phi) = \rho(f + g, h) \omega(\phi(h), [f, g, \alpha]_{\mathcal{B} \oplus \mathcal{B}^*}) \\
 &= \rho(f + g, h) \omega(\phi(h), \mu^*(f, g) \alpha) \\
 &= -\rho(f + g, h) \rho(h, f + g + \alpha) \mu^*(f, g) \alpha(\phi(h)) \\
 &= \rho(f + g + h, \alpha) \alpha \circ \phi \mu(f, g) h.
 \end{aligned}$$

□

Data Availability Statements. The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

1. K. Abdaoui, F. Ammar and A. Makhlouf, *Construction and cohomology of color Hom-Lie algebras*, J. Commu. Alg., **43**(11) (2015), 4581–4612.
2. V. Abramov, *Matrix 3-Lie superalgebras and BRST supersymmetry*, Int. J. Geom. Meth. Mod. Phys., **14**(11), 1750160, (2017).
3. V. Abramov, *Super 3-Lie algebras induced by super Lie algebras*, Adv. Appl. Clifford Alg., **27**(1) (2017), 9–16.
4. H. Albuquerque, E. Barreiro, and S. Benayadi, *Odd-quadratic Lie superalgebras*, J. Geom. Phys., **60** (2010), 230–250.
5. F. Ammara, I. Ayadib, S. Mabrouka, A. Makhlouf, *Quadratic color Hom-Lie algebras*, Ass. Non-ass. alg. app., **311** (2020), 287–312.
6. A. R. Arman, S. Silvestrov and M. R. Farhangdoost, *Extensions of hom-Lie color algebras*, Geor. Math. J., (2019).
7. H. Ataguema, A. Makhlouf and S. Silvestrov, *Generalization of n -ary Nambu algebras and beyond*, J. Math. Phys., **50**(8) (2009), 15pp.
8. I. Ayadi, H. Benamor, and S. Benayadi, *Lie superalgebras with some homogeneous structures*, J. Alg. Appl., **11**(5) (2012).
9. J. Bagger and N. Lambert, *Gauge symmetry and supersymmetry of multiple M2-branes*, Phys. Rev. D, **77**(6) (2008), 6 pp.
10. J. Bagger and N. Lambert, *Three-algebras and $N = 6$ Chern-Simons gauge theories*, Phys. Rev. D, **79**(2) (2009), 025002, 8 pp.
11. I. Bajo, S. Benayadi and A. Medina, *Symplectic structures on quadratic Lie algebras*, J. Alg, **316** (2006), 174–188.
12. C. Bai, L. Guo and Y. Sheng, *Bialgebras, the classical yang-baxter equation and manin triples for 3-Lie algebras*, Adv. Theo. Math. Phys., **23** (2019), 27–74.
13. A. Basu and J.A. Harvey, *The M2-M5 brane system and a generalized Nahm's equation*, Nucl. Phys. B, **713**(136) (2005), 136–150.
14. H. Benamor and S. Benayadi, *Double extension of quadratic Lie superalgebras*, Comm. Alg., **27** (1999), 67–88.
15. S. Benayadi and A. Makhlouf, *Hom-Lie algebras with symmetric invariant nondegenerate bilinear forms*, J. Geom. Phys., **76** (2014), 38–60.
16. P. J. Bongaarts and H. G. J. Pijls, *Almost commutative algebra and differential calculus on the quantum hyperplane*, J. Math. Phys. **35**(2) (1994), 959–970.
17. N. Cantarini and V. Kac, *Classification of Simple Linearly Compact n -Lie Superalgebras*, Comm. Math. Phys. **298** (3) (2010), 833–853.
18. Y. Daletskii and V. Kuznirenich, *Inclusion of Nambu-Takhtajan algebra in formal differential geometry structure*, Dop. NAN Ukr., **4** (1996), 12–18.
19. V. T. Filippov, *n -Lie algebras*, Siberian. Math. J., **26** (1985), 879–891.
20. J. M. Figueroa-O'Farrill and S. Stanciu, *On the structure of symmetric self-dual Lie algebras*, J. Math. Phys. **37**(2) (1996), 451–464.
21. P. Gautheron, *Some remarks concerning Nambu mechanics*, Lett. Math. Phys., **37** (1996), 103–116.

22. M. Gerstenhaber, *The cohomology structure of an associative ring*, Ann. Math., **78** (1963), 267–288.
23. B. Guan, L. Chen and B. Sun, *3-ary Hom-Lie Superalgebras induced by Hom-Lie superalgebras*, Adv. Appl. Clifford Algebras, **27** (2017), 3063–3082.
24. J. T. Hartwig, D. Larsson and S. D. Silvestrov, *Deformations of Lie algebras using σ -derivations*, J. Alg., **295** (2006), 314–361.
25. P. Ho, R. Hou and Y. Matsuo, *Lie 3-algebra and multiple M2-branes*. J. High. Energy Phys., **6** (2008), 30 pp.
26. Sh. M. Kasymov, *On a theory of n -Lie algebras*, Algebra Logika **26**(3) (1987), 277–297.
27. Y. Liu, L. Chen and Y. Ma, *Representations and module-extensions of 3-hom-Lie algebras*, J. Geom. Phys., **98** (2015), 376–383.
28. A. Makhlouf and S. D. Silvestrov, *Hom-algebra structures*, J. Gen. Lie Theory. Appl., **2** (2008), 51–64.
29. Y. Nambu, *Generalized Hamiltonian dynamics*, Phys. Rev. D, **7** (1973), 2405–2412.
30. G. Papadopoulos, *M2-branes, 3-Lie algebras and Plucker relations*, J. High Energy Phys., **5** (2008), 9pp.
31. E. Peyghan, Z. Bagheri, I. Gultekin and A. Gezer, *Representations and deformations of 3-Hom- ρ -Lie algebras*, arXiv:2104.08816v2, (2021).
32. M. Scheunert and R. B. Zhang, *Cohomology of Lie superalgebras and their generalizations*, J. Math. Phys., **39** (1998), 5024–5061.
33. Y. Sheng and D. Chen, *Hom-Lie 2 algebras*, J. Alg., **376** (2013), 174–195.
34. Q. Sun and H. Li, *On parakähler hom-Lie algebras and hom-left-symmetric bialgebras*. Comm. Algebra, **45**(1) (2017), 105–120.
35. B. Sun, L. Chen and Y. Liu, *T^* -extensions and abelian extensions of Hom-Lie color algebras*, Revista De La Unión Matemática Argentina., **59**(1) (2018), 123–142.
36. L. Takhtajan, *On foundation of the generalized Nambu mechanics*, Commun. Math. Phys., (1994), 295–316.
37. S. Wang, L. Zhu, and Y. Su, *Non-degenerate invariant bilinear forms on Lie color algebras*, Algebra Colloquium, **17** (2010), 365–374.
38. L. Yuan, *Hom-Lie color algebra structures*, J. Commu. Alg. **40**(2) (2012), 575–592.
39. T. Zhang, *Cohomology and deformations of 3-Lie colour algebras*, Lin. Multi. Alg., **63**(4) (2015), 651–671.
40. Q. Zhang, H. Yu and C. Wang, *Hom-Lie algebroids and hom-left-symmetric algebroids*. J. Geom. Phys., **116** (2017), 187–203.