

Drinfeld Twists on Skew Braces

Aryan Ghobadi

Queen Mary University of London
School of Mathematics, Mile End Road
London E1 4NS, UK
Email: a.ghobadi@qmul.ac.uk

Abstract

We introduce the notion of Drinfeld twists for both set-theoretical YBE solutions and skew braces. We give examples of such twists and show that all twists between skew braces come from families of isomorphisms between their additive groups. We then describe the relation between these definitions and co-twists on FRT-type Hopf algebras in the category SupLat , and prove that any co-twist on a co-quasitriangular Hopf algebra in SupLat induces a Drinfeld twist on its remnant skew brace. We go on to classify co-twists on bicrossproduct Hopf algebras coming from groups with unique factorisation, and the twists which they induce on skew braces.

2020 *Mathematics Subject Classification*: 16T25, 18M15, 16T99, 17B37

Keywords: Drinfeld twist, Hopf algebra, join-semilattice, matched pairs of groups, skew braces, Yang-Baxter equation

1 Introduction

Set-theoretical solutions of the Yang-Baxter equation (YBE) and their classification have attracted a lot of attention in recent years [2, 4, 3, 8, 16]. A prominent role in this theory is played by *skew braces*, which are sets with two compatible group structures [8]. Each set-theoretical YBE solution has a corresponding universal skew brace, while any skew brace comes equipped with a set-theoretical YBE solution. In fact, any skew brace is equivalent to a group with a compatible YBE solution on it, called a braiding operator. In the recent work [7], we showed that skew braces are in fact remainders of co-quasitriangular Hopf algebras (CQHA) in the category SupLat , which has complete lattices as objects and join-preserving maps as morphisms. Given any CQHA in this category we can construct a group called the remnant of the Hopf algebra, which comes equipped with a braiding operator. Conversely, any group with a braiding operator appears as the remnant of a CQHA in SupLat , but not necessarily of a unique one. The first fruits of this theory already appeared in [7], where the correspondence between skew braces and groups with braiding operators was described as the shadow of the theory of transmutation for CQHAs [14]. In addition to explaining aspects of the theory of skew braces, this machinery enables us to apply standard methods in the theory of Hopf algebras to produce new skew braces and set-theoretical YBE solutions. In this work, we will introduce the theory of Drinfeld twists for skew braces using this machinery.

Drinfeld twists were originally introduced in [5, 6] in the study of quasi-Hopf algebras. Given a Drinfeld twist on an ordinary Hopf algebra, we can produce a new Hopf algebra structure on the same space by conjugating the coalgebra structure by the twist element, see [15]. Moreover, if the Hopf algebra is equipped with a quasitriangular structure, the twisted Hopf algebra also obtains a quasitriangular structure. The dual of this theory also works and is utilised under the name of Drinfeld co-twists [13]. By applying this theory in the category of SupLat , we see that any co-twist on a CQHA in SupLat

produces a new skew brace structure on its remnant. The study of co-twist on the FRT-type Hopf algebras constructed in [7], then reveals how we can formulate a novel notion of twists directly on skew braces.

In Section 2, we introduce the notion of Drinfeld twists for skew braces, Definitions 2.2, and show directly how such a twist induces new skew brace structure on the same set, Theorem 2.5. The analogous definition and result for twists on set-theoretical YBE solutions also appear in Definition 2.1 and Theorem 2.1. In Example 2.1, we show that any involutive YBE solution can be obtained by a twist on the permutation solution, while in Example 2.5 we show that any skew brace (G, \cdot, \star) can be obtained by a twist on the trivial skew brace, (G, \star, \star) , defined by its additive group (G, \star) . We also comment on the difference between Drinfeld twists on the universal skew brace of a YBE solution and twists on the solution itself in Example 2.4. Beyond examples, in Theorem 2.2, we show how to compose and invert twists, and how one can view a twist on a YBE solution as a morphism going from the original solution to the new solution obtained by twisting. The same techniques follow for twists on skew braces, Theorem 2.6, and we obtain a groupoid which has skew brace as its objects and Drinfeld twists as morphisms. This result together with Example 2.5 allow us to translate the problem of whether two skew braces are related via Drinfeld twists fully in terms of their additive groups, Theorem 2.7. Finally, in Corollary 2.8 we conclude that twists between a pair of skew braces (G, \cdot, \star) and (G, \cdot', \star') correspond to families of group isomorphisms $\{f_g : (G, \star) \rightarrow (G, \star')\}_{g \in G}$ satisfying $f_g(g) = g$.

While the theory of Drinfeld twists on skew braces can be understood independently, it is inspired by the Hopf algebraic point of view in SupLat. Our main results extending the work in [7] appear in Section 3. As mentioned earlier, studying co-twists on the FRT-type algebras constructed in [7], reveal the definition of twists on YBE solutions and skew braces directly, Theorem 3.3. More generally, we show that a Drinfeld co-twist on any arbitrary CQHA in SupLat induces a Drinfeld twist on its remnant skew brace, Theorem 3.1. In Section 4, we classify co-twists on dualizable CQHAs in SupLat, which correspond to matched pairs of groups [17], and thereby describe the induced twists coming from these structures, Theorem 4.2. Since any skew brace gives rise to such a pair of groups in a trivial way, we obtain an explicit family of twists for any given skew brace in Corollary 4.4. We conclude the manuscript by presenting a number of open problems regarding the theory of Drinfeld twists on skew braces in Section 5.

Lastly, let us comment on the similarities of our work with other sources which study linear YBE solutions. In [9], the authors present the notion of twist on linear YBE solutions, by studying Drinfeld twists on the FRT bialgebras. Our definition of Drinfeld twists on set-theoretical YBE solutions and its relation to the SupLat-FRT algebras can be seen as the set-theoretical version of their work. In particular, the proof of Theorem 2.1 is practically the same as Proposition 1 in [9]. However, the introduction of twists on skew braces is the truly novel portion of our work. A discussion relating Drinfeld twists and braces also appears [4, 3], where the authors study Drinfeld twists on linear FRT-type algebras coming from the linearisation of involutive YBE solutions. We are able to rephrase part of the work done in [3] in terms of our notion of Drinfeld twists in Example 2.1.

Notation. Throughout this work, X will always denote a set and indices on a map such as $r_{ij} : X^3 \rightarrow X^3$ will denote the application of r to the i and j -th components of X^3 e.g. $r_{23} = \text{id}_X \times r$. We use the standard notation $r(x, y) = (\sigma_x(y), \gamma_y(x))$ for maps $\sigma_x, \gamma_y : X \rightarrow X$. The flip map on the set X^2 which sends a pair (x, y) to (y, x) will always be denoted by fl_X . For any set S , id_S denotes the identity map. By a groupoid we mean a category where all morphisms are invertible. The triple (G, m, e) will always denote a group, where $e \in G$ is the identity element and m the multiplication operation. We will use m and \cdot interchangeably in calculations. The term skew brace will always refer to skew left brace as defined in [8]. In Sections 3 and 4 we adapt the notation of [7] and [11], respectively.

Acknowledgements. The author would like to thank Shahn Majid for his helpful comments on an

earlier draft of this work.

2 Twists on Skew Braces

In this section, we introduce the notion of Drinfeld twists on set-theoretical YBE solutions and skew braces. While twists on YBE solutions first appeared in [9], the definition of twists on skew braces, Definition 2.2, is new and its relation with Drinfeld twists on Hopf algebras will be explored in Section 3. In this section, we show how these twists can be viewed as morphisms between YBE solutions and skew braces, respectively, and determine when two skew braces can be related via such twists. Note that we only consider the braid form of the Yang-Baxter equation and by a set-theoretical YBE solution we mean a set X and an *invertible* map $r : X^2 \rightarrow X^2$ satisfying $r_{23}r_{12}r_{23} = r_{12}r_{23}r_{12}$.

Definition 2.1. A *Drinfeld twist* on a set-theoretical YBE solution (X, r) consists of a triple of invertible maps $F : X^2 \rightarrow X^2$ and $\Phi, \Psi : X^3 \rightarrow X^3$ satisfying

$$(F \times \text{id}_X)\Psi = (\text{id}_X \times F)\Phi \quad (\text{T1})$$

$$\Phi(\text{id}_X \times r) = (\text{id}_X \times r)\Phi \quad (\text{T2})$$

$$\Psi(r \times \text{id}_X) = (r \times \text{id}_X)\Psi \quad (\text{T3})$$

The notion of twists on YBE solutions has previously appeared in [9], but is written in terms of $(F^{-1}, \Phi^{-1}, \Psi^{-1})$. As observed in [9], the triple (F, Φ, Ψ) is determined uniquely by the pair (F, G) , where $G = F_{12}\Psi = F_{23}\Phi$ and we could re-write the conditions of Definition 2.1 for this pair, but in application this will only lengthen our proofs.

Theorem 2.1. Given a Drinfeld twist (F, Φ, Ψ) on a set-theoretical YBE solution (X, r) , we obtain a new YBE solution, (X, FrF^{-1}) .

Proof. For this proof we observe the simple fact that (T1) implies $F_{23}^{-1}F_{12} = \Phi\Psi^{-1}$. We now show that $R_{23}R_{12}R_{23} = R_{12}R_{23}R_{12}$ holds for $R = FrF^{-1}$ by simplifying both sides:

$$\begin{aligned} R_{23}R_{12}R_{23} &= F_{23}r_{23}F_{23}^{-1}F_{12}r_{12}F_{12}^{-1}F_{23}r_{23}F_{23}^{-1} = F_{23}r_{23}\Phi\Psi^{-1}r_{12}F_{12}^{-1}F_{23}r_{23}F_{23}^{-1} \\ &= F_{23}\Phi r_{23}r_{12}\Psi^{-1}F_{12}^{-1}F_{23}r_{23}F_{23}^{-1} = F_{23}\Phi r_{23}r_{12}\Phi^{-1}r_{23}F_{23}^{-1} = F_{23}\Phi r_{23}r_{12}r_{23}\Phi^{-1}F_{23}^{-1} \\ R_{12}R_{23}R_{12} &= F_{12}r_{12}F_{12}^{-1}F_{23}r_{23}F_{23}^{-1}F_{12}r_{12}F_{12}^{-1} = F_{12}r_{12}F_{12}^{-1}F_{23}r_{23}\Phi\Psi^{-1}r_{12}F_{12}^{-1} \\ &= F_{12}r_{12}F_{12}^{-1}F_{23}\Phi r_{23}r_{12}\Psi^{-1}F_{12}^{-1} = F_{12}r_{12}\Psi r_{23}r_{12}\Psi^{-1}F_{12}^{-1} = F_{12}\Psi r_{12}r_{23}r_{12}\Psi^{-1}F_{12}^{-1} \end{aligned}$$

Since r satisfy the YBE and (T1) is equivalent to $F_{12}\Psi = F_{23}\Phi$, the two sides are equal. \square

Observe that FrF^{-1} is involutive i.e. satisfies $(FrF^{-1})^2 = \text{id}_{X^2}$ if and only if r is involutive. In fact, the statements $r^n = \text{id}_{X^2}$ and $(FrF^{-1})^n = \text{id}_{X^2}$ are equivalent for any natural number $n \in \mathbb{N}$. Recall, that a YBE solution r is said to be non-degenerate if σ_x, γ_y are bijections for all $x, y \in X$. For FrF^{-1} to also be non-degenerate we need f_x, f'_y to be bijections, where $F(x, y) = (f_y(x), f'_x(y))$.

We can view a Drinfeld twist (F, Φ, Ψ) on a solution (X, r) as an “arrow” from the solution (X, r) to (X, FrF^{-1}) . From this perspective, we obtain a category which has set-theoretical YBE solution as objects and Drinfeld twists as morphisms:

Theorem 2.2. The class of set-theoretical YBE solutions, together with Drinfeld twists viewed as morphisms between solution form a groupoid.

Proof. Each YBE solution has an associated trivial twist $(\text{id}_{X^2}, \text{id}_{X^3}, \text{id}_{X^3})$ which acts as an identity morphism. For composition of twists, let (F, Φ, Ψ) be a twist on (X, r) and (G, ϕ, ψ) be a twist on (X, FrF^{-1}) . We define the composition of these twists to be the triple

$$(GF, F_{23}^{-1}\phi F_{23}\Phi, F_{12}^{-1}\psi F_{12}\Psi)$$

It is straightforward to check that this triple defines a twist on r :

$$\begin{aligned} \text{(T1): } & (GF)_{23}F_{23}^{-1}\phi F_{23}\Phi = G_{23}F_{23}F_{23}^{-1}\phi F_{12}\Psi = G_{12}\psi F_{12}\Psi = (GF)_{12}F_{12}^{-1}\psi F_{12}\Psi \\ \text{(T2): } & r_{23}F_{23}^{-1}\phi F_{23}\Phi = F_{23}^{-1}(F_{23}r_{23}F_{23}^{-1})\phi F_{23}\Phi = F_{23}^{-1}\phi(F_{23}r_{23}F_{23}^{-1})F_{23}\Phi = F_{23}^{-1}\phi F_{23}\Phi r_{23} \end{aligned}$$

A symmetric argument demonstrates (T3). Finally, we observe that Drinfeld twists are invertible. If (X, R) is obtained by a twist (F, Ψ, Φ) on (X, r) , then the triple $(F^{-1}, F_{23}\Phi^{-1}F_{23}^{-1}, F_{12}\Psi^{-1}F_{12}^{-1})$ form a twist on R , which recovers $r = F^{-1}RF$ by Theorem 2.1. One can easily check that the latter triple becomes the inverse of (F, Ψ, Φ) , with respect to the composition which we have defined. \square

By Theorem 2.2, we can say two YBE solutions are *related* by twists, since if one is obtained as a twist of the other, then the converse statement also holds true. Note that Drinfeld twists only send a YBE solution on a set X to another solution on the same underlying set X . The important conclusion of Theorem 2.2 is that any solution in a connected component of the described groupoid can be obtained as a twist of another solution in the same component. In other words, if solutions r_1 and r_2 on a set X are related to a solution (X, r_3) via twists, then r_1 and r_2 are also related via Drinfeld twists.

Example 2.1. In [3], it is stated that any involutive YBE solution can be viewed as a *quasi-admissible twist* of the permutation solution. Although [3] is discussing linear Hopf algebras built on these solutions, the proof of Proposition 3.13 of [3] boils down to the following observation: if $r(x, y) = (\sigma_x(y), \gamma_y(x))$ satisfies the YBE we can introduce $F(x, y) = (x, \sigma_x(y))$, $\Phi(x, y, z) = (x, \sigma_x(y), \sigma_{\gamma_y(x)}(z))$ and $\Psi(x, y, z) = (x, y, \sigma_x(\sigma_y(z)))$, so that the triple (F, Φ, Ψ) define a Drinfeld twist on (X, r) . Conditions (T1), (T2) and (T3), all follow directly from r being a YBE solution and the identities $\sigma_{\sigma_x(y)}(\sigma_{\gamma_x(y)}(z)) = \sigma_x(\sigma_y(z))$ and $\sigma_{\gamma_{\sigma_y(z)}(x)}(\gamma_z(y)) = \gamma_{\sigma_x(y)}(\sigma_{\gamma_y(x)}(z))$. Consequently, by Theorem 2.1 we obtain a new YBE solution defined by $FrF^{-1}(x, y) = (y, \sigma_y(\gamma_{\sigma_x^{-1}(y)}(x)))$. Observe that $\sigma_y(\gamma_{\sigma_x^{-1}(y)}(x))$ appears in the first component of $r^2(x, \sigma_x^{-1}(y))$. Going back to the involutive setting considered in [3], we have that $FrF^{-1}(x, y) = (y, x)$, since $r^2 = \text{id}_{X^2}$. Therefore, the permutation solution (X, fl_X) is obtained from a twist on (X, r) , and by Theorem 2.2, (X, r) can be obtained by the inverse twist on (X, fl_X) .

By the above example, any involutive solution on X is obtained as a twist of the permutation solution and by Theorem 2.2, we can conclude the following result.

Corollary 2.3. *If X is a set and r_1 and r_2 are involutive YBE solutions in X , then there exists a twist (F, Φ, Ψ) on r_1 such that $r_2 = Fr_1F^{-1}$.*

Example 2.2. Let us consider the non-involutive YBE solution from Example 3.4 of [16]. We take $X = \{1, 2, 3, 4\}$ and $r(x, y) = (\sigma(y), \gamma(x))$, where $\sigma = (12)$ and $\gamma = (34)$, where $(12), (34) \in S_4$ are 2-cycle permutations with standard notation. Consider $F(x, y) = (\sigma(x), \gamma(y))$ with $\Phi(x, y, z) = (\gamma(\sigma(x)), \sigma(y), \sigma(z))$ and $\Psi(x, y, z) = (\gamma(x), \gamma(y), \gamma(\sigma(z)))$. The twisted YBE solution we obtain is $FrF^{-1}(x, y) = (\gamma(y), \sigma(x))$. This can be generalised to any YBE solution of the form $r(x, y) = (\sigma(x), \gamma(y))$, where bijections σ and γ commute and $\sigma^2 = \gamma^2 = \text{id}_X$.

Example 2.3. For any YBE solution $r(x, y) = (\sigma(y), \gamma(x))$ we can introduce a twist in the style of Example 2.1, $F(x, y) = (x, \kappa(y))$, where κ is a bijection on X which commutes with σ and γ and we define $\Phi(x, y, z) = (x, \kappa(y), \kappa(z))$ and $\Psi(x, y, z) = (x, y, \kappa(\kappa(z)))$. In the case of (X, r) of Example 2.2, κ can be either $\sigma = (12)$ or $\gamma = (34)$, giving the twisted YBE solutions $(x, y) \mapsto (y, \sigma(\gamma(x)))$ and $(x, y) \mapsto (\sigma(\gamma(y)), x)$, respectively.

Now we briefly recall the definition of a group with braiding operator from [10]. A *braiding operator* on a group (G, m, e) is a map $r : G^2 \rightarrow G^2$ satisfying

$$r(e, g) = (g, e), \quad r(g, e) = (e, g) \quad (1)$$

$$rm_{12} = m_{23}r_{12}r_{23} \quad (2)$$

$$rm_{23} = m_{12}r_{23}r_{12} \quad (3)$$

$$mr = m \quad (4)$$

It follows by definition that r is a non-degenerate solution of the YBE, Corollary 3 of [10].

Definition 2.2. A *Drinfeld twist* on a group (G, m, e) with a braiding operator r consists of a Drinfeld twist (F, Φ, Ψ) on the underlying YBE solution (G, r) satisfying the additional conditions

$$\Psi(x, y, e) = (x, y, e), \quad \Phi(e, x, y) = (e, x, y) \quad (G1)$$

$$F(e, x) = (e, x), \quad F(x, e) = (x, e) \quad (G2)$$

$$m_{23}\Phi(x, y, z) = F(x, yz) \quad (G3)$$

$$m_{12}\Psi(x, y, z) = F(xy, z) \quad (G4)$$

where $x, y, z \in G$.

Although this definition does not appear to be related to Drinfeld twists on Hopf algebras, the relation between the two will be clarified in Section 3, by the study of Hopf algebras in SupLat.

Lemma 2.4. *If (G, m, e) is a group with a braiding operator r , and (F, Φ, Ψ) is as in Definition 2.2, it follows that*

$$\Phi(x, y, e) = (F(x, y), e), \quad \Phi(x, e, y) = F_{13}(x, e, y) \quad (L1)$$

$$\Psi(e, x, y) = (e, F(x, y)), \quad \Psi(x, e, y) = F_{13}(x, e, y) \quad (L2)$$

hold for any $x, y \in G$.

Proof. The second equation in (L1) follows from the first equation together with (T2) and (1) and the first equation in (L1) is a consequence of (T1), (G1) and (G2):

$$\Phi(x, y, e) = F_{23}^{-1}F_{23}\Phi(x, y, e) = F_{23}^{-1}F_{12}\Psi(x, y, e) = F_{23}^{-1}(F(x, y), e) = (F(x, y), e)$$

$$\Phi(x, e, y) = r_{23}^{-1}\Phi r_{23}(x, e, y) = r_{23}^{-1}\Phi(x, y, e) = r_{23}^{-1}(F(x, y), e) = F_{13}(x, e, y)$$

The proof of (L2) is completely analogous. □

Observe that by Lemma 2.4, F can be recovered from Φ or Ψ , and twists as defined in Definition 2.2 are completely determined by the pair (Φ, Ψ) . Additionally, (T1) implies that Ψ is determined by F and Φ . Hence, Definition 2.2 can then be re-written in terms of a single map Φ : a twist on a group (G, m, e)

with a braiding operator r consists of a bijection $\Phi : X^3 \rightarrow X^3$ such that $\Phi(x, y, e) = (\overline{\Phi}(x, y), e)$ for a bijection $\overline{\Phi} : X^2 \rightarrow X^2$, and Φ satisfies (T2) and

$$\Phi(e, x, y) = (e, x, y), \quad \overline{\Phi}(x, e) = (x, e) \quad (5)$$

$$m_{23}\Phi = \overline{\Phi}m_{23} \quad (6)$$

$$m_{12}\overline{\Phi}_{12}^{-1}\overline{\Phi}_{23}\Phi = \overline{\Phi}m_{12} \quad (7)$$

$$r_{12}\overline{\Phi}_{12}^{-1}\overline{\Phi}_{23}\Phi = \overline{\Phi}_{12}^{-1}\overline{\Phi}_{23}\Phi r_{12} \quad (8)$$

for $x, y \in G$. Notice that Ψ is recovered from Φ as $\overline{\Phi}_{12}^{-1}\overline{\Phi}_{23}\Phi$, so that (T1) holds and (7) and (8) are translations of (G4) and (G3), respectively. Similarly, (6) replaces (G3). The remaining conditions in (G1) and (G2) follow by (5) and (6), since $\Phi(e, x, e) = (\overline{\Phi}(e, x), e)$ and $\overline{\Phi}_{12}^{-1}\overline{\Phi}_{23}\Phi(x, y, e) = \overline{\Phi}_{12}^{-1}\overline{\Phi}_{23}(\overline{\Phi}(x, y), e) = \overline{\Phi}_{12}^{-1}\overline{\Phi}_{12}(x, y, e)$. Although, the data for a twist can be captured in a single map Φ , we will continue to use triples (F, Φ, Ψ) , since they are much more intuitive to work with and make our proofs less complicated.

Theorem 2.5. *Let (F, Φ, Ψ) be a Drinfeld twist on a group (G, m, e) with a braiding operator r . Then (G, mF^{-1}, e) defines a new group structure on the set G with a braiding operator FrF^{-1} .*

Proof. We must first demonstrate that (G, mF^{-1}, e) is a group. First, we observe that mF^{-1} is an associative operation by (T1) and (G3), (G4):

$$\begin{aligned} (mF^{-1})(mF^{-1})_{12} &= mF^{-1}m_{12}F_{12}^{-1} = mm_{12}\Psi^{-1}F_{12}^{-1} = mm_{23}\Phi^{-1}F_{23}^{-1} \\ &= mF^{-1}m_{23}F_{23}^{-1} = (mF^{-1})(mF^{-1})_{23} \end{aligned}$$

By definition (G2) implies that $e \in G$ acts as the identity element for mF^{-1} . What remains is the existence of inverses with respect to mF^{-1} . Let $x \in G$, then for any element $y \in G$, we first observe that $\Phi(y, x, x^{-1}) = (y, z, z^{-1})$, for some $z \in G$, since

$$m_{23}\Phi(y, x, x^{-1}) = Fm_{23}(y, x, x^{-1}) = F(y, e) = (y, e)$$

A similar argument shows that $\Psi(x, x^{-1}, y) = (l, l^{-1}, y)$, for some $l \in G$. Now we observe that $F_{23}\Phi(x, x^{-1}, x) = (x, x_F, x^{-1})$, for some x_F . By the previous argument, there exists a pair $z, l \in G$ such that

$$\begin{aligned} (x, F(z, z^{-1})) &= F_{23}\Phi(x, x^{-1}, x) = F_{12}\Psi(x, x^{-1}, x) = (F(l, l^{-1}), x) \\ \Rightarrow mF^{-1}(x_F, x) &= m(z, z^{-1}) = e, \quad mF^{-1}(x, x_F) = m(l, l^{-1}) = e \end{aligned}$$

Consequently, x_F is the inverse of x with respect to mF^{-1} and $(G, m, 1)$ forms a group.

In the second part of the proof we must show that FrF^{-1} is a braiding operator on the group. Condition (4) follows by definition since $(mF^{-1})(FrF^{-1}) = mrF^{-1} = mF^{-1}$ and (1) follows by (G2) and r satisfying (1) e.g. $FrF^{-1}(1, x) = Fr(1, x) = F(x, 1) = (x, 1)$. To prove (2), we utilise the equivalent form of (T1), $F_{23}^{-1}F_{12} = \Phi\Psi^{-1}$, and (T2), (G3), (T1) and (G4), respectively:

$$\begin{aligned} (mF^{-1})_{23}(FrF^{-1})_{12}(FrF^{-1})_{23} &= m_{23}F_{23}^{-1}F_{12}r_{12}F_{12}^{-1}F_{23}r_{23}F_{23}^{-1} = m_{23}\Phi\Psi^{-1}r_{12}\Psi\Phi^{-1}r_{23}F_{23}^{-1} \\ &= \Phi m_{23}r_{12}r_{23}\Phi^{-1}F_{23}^{-1} = Frm_{12}\Psi^{-1}F_{12}^{-1} = FrF^{-1}m_{12}F_{12}^{-1} = (FrF^{-1})(mF^{-1})_{12} \end{aligned}$$

Proving (3) is analogous to the argument above and is left to the reader. \square

The above results have been stated in terms of groups with braiding operators, which are equivalent structures to skew braces. The latter objects consist of a set B with two compatible group structures (B, \cdot) and (B, \star) . In particular, any group (G, \cdot) with braiding operator r gives rise to a skew brace (G, \cdot, \star) , where $x \star y := x \cdot \sigma_x^{-1}(y)$, while the braiding operator r can be recovered from the group operations of a skew brace (G, \cdot, \star) . We refer to Section 3 of [16] for additional details, where the products \cdot and \star are denoted by \circ and \cdot , respectively. Although r can be determined in terms of the skew brace operations, \cdot and \star , the author has not been able to find a way to re-write the axioms of Definition 2.2 solely in terms of \cdot and \star . However, by the end of this section we will fully classify all twists between skew braces and describe them in terms of the skew brace operations. From this point onwards, we will use the terms group with braiding operator and skew brace interchangeably and refer to (G, \star) as the *additive group* of the skew brace (G, \cdot, \star) , as done in [16].

In the next example we will reflect on the difference between twists on a YBE solution (X, r) and twists on the solution's universal skew brace $G(X, r) = \langle x \in X \mid x \cdot y = \sigma_x(y) \cdot \gamma_y(x), \forall x, y \in X \rangle$.

Example 2.4. Let (X, r) be as in Example 2.2. As discussed in Example 3.4 of [16], the universal skew brace of (X, r) , denoted by G , will be the free abelian group on two generators, a and b , with the flip map fl_G as its braiding operator. The natural map $i : X \rightarrow G$ sends 1, 2 to generator a and 3, 4 to generator b . First, let us consider the problem of extending a twist (f, ϕ, ψ) on (X, r) to a twist of skew braces (F, Φ, Ψ) on (G, fl_G) , so that the triples commute with i . The issue here is that i is not injective. For example the triple $f = (13) \times (13)$, $\phi = (13) \times \text{id}_{X^2}$, $\psi = \text{id}_{X^2} \times (13)$ define a twist on (X, r) , while there cannot exist a map F such that $F(i \times i) = (i \times i)f$ holds true since $i(1) = i(2) \neq i(3)$. Even if we could define the triple on the generators, there is no unique way to extend maps such as ψ and ϕ to multiplications of the generators e.g. $\Phi(a^3, a^2, a^7)$ is not uniquely determined by (F, Φ, Ψ) being defined on the generators of the group. The reader should also note that the twists in Examples 2.2 and 2.3 all lift to the trivial twist where F, Φ and Ψ are all identity morphisms. In the opposite direction, it is also non-trivial to check whether a twist on G extends a canonical twist on X . The issue here is that a generator in G represents several elements of X . Consider the group automorphisms $\chi_{a^n b^m} : G \rightarrow G$ corresponding to group elements $a^n b^m \in G$ defined by

$$\chi_{a^n b^m}(a^l b^k) = \begin{cases} a^l b^k & \text{if } n + m \text{ is even} \\ b^l a^k & \text{if } n + m \text{ is odd} \end{cases}$$

We obtain a twist on (G, fl_G) defined by $F(g, h) = (g, \chi_g(h))$ and $\Phi(g, h, k) = (g, \chi_g(h), \chi_g(k))$ and $\Psi = (g, h, \chi_{gh}(k))$, which recovers the universal skew brace appearing in Example 3.12 of [2]. In this case, we can define $f(x, y) = (\varpi(x), \varphi(y))$ with any of the choices $\varpi = \text{id}_X, (12), (34), (12)(34)$ and $\varphi = (13)(24), (14)(23)$ and $F(i \times i) = (i \times i)f$ will follow regardless of our choice. Consequently, there is no natural choice for a triple (f, ϕ, ψ) so that (F, Φ, Ψ) is a lift of the former. For example, the triples $f = \varpi \times (14)(23)$, $\phi = \varpi \times (13)(24) \times (14)(23)$ and $\psi = \varpi \times \varpi \times \text{id}_X$ form twists lifting to (F, Φ, Ψ) for both choices $\varpi = (12)$ and $\varpi = (34)$.

Theorem 2.6. *The category which has skew braces as objects and Drinfeld twists between them as morphisms forms a subgroupoid of the groupoid of YBE solutions of Theorem 2.2.*

Proof. We only need to show that Drinfeld twists on skew braces respect the composition and inverse operation defined in Theorem 2.2. Recall from the proof of Theorem 2.2, the inverse of a triple (F, Φ, Ψ) is given by $(F^{-1}, F_{23}\Phi^{-1}F_{23}^{-1}, F_{12}\Psi^{-1}F_{12}^{-1})$ and we must demonstrate that it is indeed a twist on (G, mF^{-1}, e) with braiding operator FrF^{-1} . If the first triple satisfies (G1) and (G2), it follows directly that the inverse triple also satisfies these axioms, since the unit element of the twisted skew brace

remains the same. Now we demonstrate (G3):

$$(mF^{-1})_{23}F_{23}\Phi^{-1}F_{23}^{-1} = m_{23}F_{23}^{-1}F_{23}\Phi^{-1}F_{23}^{-1} = m_{23}\Phi^{-1}F_{23}^{-1} = F^{-1}m_{23}F_{23}^{-1} = F^{-1}(mF^{-1})_{23}$$

Condition (G4) holds in a symmetric manner.

We must also check that composing twists as defined in Theorem 2.2 respect the additional axioms in Definition 2.2. Let (F, Φ, Ψ) be a twist on (G, m, e, r) and (\mathbf{F}, ϕ, ψ) be a twist on $(G, mF^{-1}, e, FrF^{-1})$. Then the composite twist $(\mathbf{F}F, F_{23}^{-1}\phi F_{23}\Phi, F_{12}^{-1}\psi F_{12}\Psi)$ satisfies (G1) and (G2) by triviality. Conditions (G3) holds since since (F, Φ) and (\mathbf{F}, ϕ) satisfy (G3):

$$m_{23}F_{23}^{-1}\phi F_{23}\Phi = (mF^{-1})_{23}\phi F_{23}\Phi = \mathbf{F}(mF^{-1})_{23}F_{23}\Phi = \mathbf{F}m_{23}\Phi = \mathbf{F}Fm_{23}$$

Condition (G4) holds in a symmetric manner and is left to the reader. \square

Note that Drinfeld twists on YBE solutions and skew braces can be defined upto isomorphism. Explicitly, if (G, m, \cdot) with operator r and (H, m', e') with r' are isomorphic via a bijection $f : G \rightarrow H$ satisfying $m'(f \times f) = fm$ and $(f \times f)r = r'(f \times f)$, then twists (F, Φ, Ψ) on (G, m, e, r) are in bijection with twists (F', Φ', Ψ') on (H, m', e', r') , where the latter pairs are obtained by conjugation by $f \times f$ and $f \times f \times f$ e.g. $F' := (f \times f)F(f^{-1} \times f^{-1})$. Additionally, the twisted skew braces $(G, mF^{-1}, e, FrF^{-1})$ and $(G, m(F')^{-1}, e, F'r'(F')^{-1})$ obtained from corresponding twists between a pair of isomorphic skew braces will be isomorphic via the same map $f : X \rightarrow Y$, since $(f \times f)FrF^{-1} = F'r'(F')^{-1}(f \times f)$ and $fmF^{-1} = m'(F')^{-1}(f \times f)$. Hence, we could refine the groupoids of Theorems 2.2 and 2.6 to have isomorphism classes of YBE solutions and skew braces as objects, respectively.

Example 2.5. We can realise the twists defined in Example 2.1 for any skew brace (G, \cdot, r) . Define $F(x, y) = (x, \sigma_x(y))$, $\Phi(x, y, z) = (x, \sigma_x(y), \sigma_{\gamma_y(x)}(z))$ and $\Psi(x, y, z) = (x, y, \sigma_x(\sigma_y(z)))$, then (F, Φ, Ψ) form a twist on the underlying solution (G, r) by Example 2.1 and (G1) and (G2) hold trivially by (1), while (G3) and (G4) follow from (3) and (2), respectively. By Theorem 2.5 we obtain a new group structure on G defined by $mF^{-1}(x, y) = x \cdot \sigma_x^{-1}(y)$ which agrees with the additive group structure of the skew brace, \star , and a corresponding braiding operator $FrF^{-1}(x, y) = (y, \sigma_y(\gamma_{\sigma_x^{-1}(y)}(x)))$. Notice that $\sigma_y(\gamma_{\sigma_x^{-1}(y)}(x)) = y \star x \star y$, where $y \star$ denotes the inverse of $y \in G$ with respect to the operation \star . In the terminology of [8], $F(x, y) = (x, x^{-1} \cdot (x \star y))$ and this twist sends a skew brace (G, \cdot, \star) to the trivial skew brace (G, \star, \star) . The latter is the skew brace obtained by a single group (G, \star) as described in Example 1.3 of [8].

By Example 2.5, it follows that the any skew brace (G, \cdot, \star) is related to the trivial skew brace (G, \star, \star) by a twist. Since Drinfeld twists can be inverted and composed, by Theorem 2.6, we conclude that any two skew braces with the same additive groups are related via Drinfeld twists. Two observations follow from this result. Firstly, it is well known that there exist non-isomorphic skew braces with isomorphic additive groups. Hence, the notion of a twist relating two skew braces is much weaker than isomorphism of skew braces. Secondly, the problem of whether two skew braces (G, \cdot, \star) and (G, \cdot', \star') are related by Drinfeld twists fully translates into a problem about their underlying additive groups and whether there exists Drinfeld twists between (G, \star, \star) and (G, \star', \star') , which we can answer.

Theorem 2.7. *If two skew braces (G, \circ, \cdot) and (G, \cdot', \star') are related via a Drinfeld twist, then their respective additive groups (G, \cdot) and (G, \star) are isomorphic.*

Proof. One direction of the argument has already been proved in the discussion before Theorem 2.7. For the other direction, let us assume that there exists a Drinfeld (G, \circ, \cdot) relating (G, \cdot', \star') . Then by

Theorem 2.6 and Example 2.5, we can obtain a Drinfeld twist relating (G, \cdot, \cdot) and (G, \star, \star) . Hence, let (F, Φ, Ψ) be a twist on (G, \cdot, \cdot) such that $\star = \cdot F^{-1}$ and for any $x, y \in G$ we have $r'(x, y) := FrF^{-1}(x, y) = (y, y^\star \star x \star y)$, where $r(x, y) = (y, y^{-1}xy)$. Since $\star F = \cdot$, we can write $F(x, y) = (f_{xy}(x), (f_{xy}(x))^\star \star (x \cdot y))$ for a family of maps $f_g : G \rightarrow G$ corresponding to $g \in G$. Since F is a bijection, then the maps f_g are also bijective. Consequently, we have that

$$Fr(x, y) = (f_{xy}(y), (f_{xy}(y))^\star \star (xy)) = ((f_{xy}(x))^\star \star (x \cdot y), (xy)^\star \star f_{xy}(x) \star (xy)) = r'F(x, y)$$

which implies that $f_{xy}(x) \star f_{xy}(y) = xy$ for any pair $x, y \in G$ and $F(x, y) = (f_{xy}(x), f_{xy}(y))$.

By (G3) it follows that $\Phi(x, y, z) = (f_{xyz}(x), f_{xyz}(yz)(\alpha_{x,yz}(z))^{-1}, \alpha_{x,yz}(z))$ for some maps $\alpha_{x,c} : G \rightarrow G$ corresponding to $x, c \in G$. Expanding equation (T2) gives

$$(f_{xyz}(x), f_{xyz}(yz)(\alpha_{x,yz}(z^{-1}yz))^{-1}, \alpha_{x,yz}(z^{-1}yz)) = (f_{xyz}(x), \alpha_{x,yz}(z), (\alpha_{x,yz}(z))^{-1}f_{xyz}(yz))$$

which implies that $f_{xc}(c) = \alpha_{x,c}(z)\alpha_{x,c}(z^{-1}c)$ for any $x, c, z \in G$, where we have replaced the term yz by c since y is a free variable. We can also re-write $\Phi(x, y, z) = (f_{xyz}(x), \alpha_{x,yz}(y), \alpha_{x,yz}(z))$. A symmetric argument shows that $\Psi(x, y, z) = (\beta_{xy,z}(x), \beta_{xy,z}(y), f_{xyz}(z))$ for some maps $\beta_{c,z} : G \rightarrow G$ satisfying $\beta_{xy,z}(x) \cdot \beta_{xy,z}(y) = f_{xyz}(xy)$.

Now we compare the third component of equation (T1). The third component of $F_{12}\Psi(x, y, z)$ is $f_{xyz}(z)$, while the third component of $F_{23}\Phi(x, y, z)$ is $f_{f_{xyz}(yz)}(\alpha_{x,yz}(z))$. Equating the two values tells us that $\alpha_{x,c} = f_{f_{xc}(c)}^{-1}f_{xc}$ and the identity $f_{xc}(c) = \alpha_{x,c}(z) \cdot \alpha_{x,c}(z^{-1}c)$ can be re-written as

$$f_{xc}(c) = f_{f_{xc}(c)}^{-1}(f_{xc}(z)) \cdot f_{f_{xc}(c)}^{-1}(f_{xc}(z^{-1}c))$$

where $x, c, z \in G$ are all independent variables. By $f_{xy}(x) \star f_{xy}(y) = xy$, we have that

$$f_{xc}(z) \star f_{xc}(z^{-1}c) = f_{f_{xc}(c)} \left(f_{f_{xc}(c)}^{-1}(f_{xc}(z)) \right) \star f_{f_{xc}(c)} \left(f_{f_{xc}(c)}^{-1}(f_{xc}(z^{-1}c)) \right) = f_{xc}(c)$$

Thereby, we can change variables by $x = pc^{-1}$ and $c = zz'$ and observe that $f_p(z) \star f_p(z') = f_p(z \cdot z')$ for all $p, z, z' \in G$. Hence, $f_p : (G, \cdot) \rightarrow (G, \star)$ are groups isomorphisms for all $p \in G$. \square

We could continue the proof of Theorem 2.7 by comparing the first component of equation (T1), which implies that $\beta_{x,y} = f_{f_{xy}(x)}^{-1}f_{xy}$ and the identity on $\beta_{xy,z}$ follows by f_p being group morphisms. With this, (T1) holds automatically and (G1) and (G2) follow by $f_g(1) = 1$. Hence, this result fully classifies all possible twists from (G, \cdot, \cdot) to (G, \star, \star) , for isomorphic (G, \cdot) and (G, \star) . Given any family of group isomorphisms $\{f_x : (G, \cdot) \rightarrow (G, \star)\}_{x \in G}$ satisfying $f_x(x) = x$, we obtain a non-trivial twist on (G, \cdot, \cdot) by setting $F(x, y) = (f_{xy}(x), f_{xy}(y))$ and Φ and Ψ as defined in the proof of Theorem 2.7, with $\alpha_{x,y} = f_{f_{xy}(y)}^{-1}f_{xy}$ and $\beta_{x,y} = f_{f_{xy}(x)}^{-1}f_{xy}$. In the groupoid of skew braces and twists of Theorem 2.6, these twists are all possible morphisms from (G, \cdot, \cdot) to (G, \star, \star) . In particular, all endomorphisms of (G, \cdot, \cdot) in this groupoid must also come from a family of group automorphisms $\{f_x\}_{x \in G}$ satisfying $f_x(x) = x$.

Recall that in an arbitrary groupoid, if we are given morphisms $a : o_1 \rightarrow o_2$, $a_i : o_i \rightarrow o'_i$ for $i = 1, 2$ between some objects o_1, o_2, o'_1, o'_2 , then there must exist a morphism $b : o'_1 \rightarrow o'_2$ defined by $b = a_2 a a_1^{-1}$ such that $a = a_2^{-1} b a_1$. Hence, we observe that any twist $(F, \Phi, \Psi) : (G, \cdot, \star) \rightarrow (G, \cdot', \star')$ can be decomposed as the composition of twists coming from Example 2.5 and a twist from (G, \star, \star) to (G, \star', \star') which by the above result has to come from a family of group isomorphisms.

Corollary 2.8. Any twist $(F, \Phi, \Psi) : (G, \cdot, \star) \rightarrow (G, \circ, \star)$ can be decomposed as the composition of the twists in Example 2.5 and those coming from Theorem 2.7. In particular, there exists a family of group isomorphisms $\{f_x : (G, \star) \rightarrow (G, \star')\}_{x \in G}$ satisfying $f_x(x) = x$ so that

$$F(x, y) = \left(f_{x.y}(x), \underline{\sigma}_{f_{x.y}(x)}^{-1}(f_{x.y}(\sigma_x(y))) \right) = (f_{x.y}(x), f_{x.y}(x)^\circ \circ (x.y)) \quad (9)$$

where σ and $\underline{\sigma}$ denote the relevant maps for the braiding operators of (G, \cdot, \star) and (G, \circ, \star) , respectively.

3 Co-twists on Hopf Algebras in SupLat

In this section we adapt the notation of [7] where we described skew braces as remnants of CQHAs in SupLat and study Drinfeld co-twists on these Hopf algebras. In particular, we look at co-twists on the SupLat-FRT algebras, which were constructed in Section 5 of [7]. The reader should firstly note that the Hopf algebras we will describe are not linear and are Hopf algebra objects in the category SupLat and secondly that there are two FRT-type algebras which we are concerned with: H_ω denotes the SupLat-FRT algebra of a non-degenerate YBE solution, which recovers the universal skew brace of the solution as its remnant, while H_{ω_m} denotes the FRT-type algebra defined for a given skew brace, which recovers the skew brace itself as its remnant. From this point onwards, we will assume that all YBE solutions of concern are non-degenerate.

A Drinfeld co-twist on a Hopf algebra \mathcal{H} in SupLat consists of a morphism $\mathcal{F} : \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{P}(1)$, for which there exists another morphism $\mathcal{F}^{-1} : \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{P}(1)$ so that

$$\mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \cdot \mathcal{F}(a_{(2)}, b_{(2)}) = \epsilon(a) \cdot \epsilon(b) = \mathcal{F}(a_{(1)}, b_{(1)}) \cdot \mathcal{F}^{-1}(a_{(2)}, b_{(2)}) \quad (\text{cT1})$$

$$\mathcal{F}(a_{(1)} \cdot b_{(1)}, c) \cdot \mathcal{F}(a_{(2)}, b_{(2)}) = \mathcal{F}(a, b_{(1)} \cdot c_{(1)}) \cdot \mathcal{F}(b_{(2)}, c_{(2)}) \quad (\text{cT2})$$

$$\mathcal{F}(a, 1) = \epsilon(a) = F(1, a) \quad (\text{cT3})$$

hold for $a, b \in \mathcal{H}$. The first condition (cT1) says that \mathcal{F} is *convolution invertible* and (cT2) and (cT3) are sometimes called the cocycle conditions, see Section 2.3 of [15] for more details. Given a co-twist on a co-quasitriangular Hopf algebras $(\mathcal{H}, \mathcal{R})$, we obtain a new CQHA $(\mathcal{H}^\mathcal{F}, \mathcal{R}^\mathcal{F})$, where the multiplication and co-quasitriangular structure are twisted:

$$m^\mathcal{F}(a, b) = \mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \cdot a_{(2)} \cdot b_{(2)} \cdot \mathcal{F}(a_{(1)}, b_{(1)}) \quad (10)$$

$$\mathcal{R}^\mathcal{F}(a, b) = \mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \mathcal{R}(a_{(2)}, b_{(2)}) \cdot \mathcal{F}(b_{(3)}, a_{(3)}) \quad (11)$$

where $a, b \in \mathcal{H}$. We refer the reader to Section 2.3 of [15], where the proof of the dual of this statement is proved. As mentioned in Chapter 9 of [15], these proofs can be re-written in diagrammatic manner and dualised.

Recall from Section 3 and 4 of [7] that given a Hopf algebra \mathcal{H} in SupLat, we can construct a group called its remnant, $R(\mathcal{H})$, and any co-quasitriangular structure on \mathcal{H} induces a braiding operator on $R(\mathcal{H})$. The underlying set of $R(\mathcal{H})$ is defined solely by the counit of the Hopf algebra \mathcal{H} . Hence, given a twist \mathcal{F} on a CQHA, $(\mathcal{H}, \mathcal{R})$, the new skew brace, $R(\mathcal{H}^\mathcal{F})$, which we obtain will have the same underlying set as $R(\mathcal{H})$. Next we will show that \mathcal{F} in fact induces a Drinfeld twist on $R(\mathcal{H})$ in the sense of Section 2.

Before proving this statement, let us briefly emphasize the notation used in [7]. Given a CQHA $(\mathcal{H}, \mathcal{R})$, there is a natural quotient Hopf algebra \mathcal{Q} with a Hopf algebra projection $\pi : \mathcal{H} \rightarrow \mathcal{Q}$. The inclusion morphism $\iota : \mathcal{Q} \rightarrow \mathcal{H}$ sends an element $\pi(a) = \bar{a} \in \mathcal{Q}$ to $a \vee D$, where $D = \vee \epsilon^{-1}(\emptyset)$. Lastly, the remnant group of \mathcal{H} is identified as the basis of \mathcal{Q} i.e. $\mathcal{Q} = \mathcal{P}(R(\mathcal{H}))$. For more details we refer the reader to Section 3 of [7].

Theorem 3.1. *Any co-twist \mathcal{F} on a co-quasitriangular Hopf algebra $(\mathcal{H}, \mathcal{R})$ induces a twist on its remnant skew brace $R(\mathcal{H})$.*

Proof. Given a co-twist \mathcal{F} , we will define the triple (F, Φ, Ψ) more generally for \mathcal{Q} and then by demonstrating that they are invertible, it follows that they must restrict to maps on the basis elements. For $\bar{a}, \bar{b}, \bar{c} \in \mathcal{Q}$, define the triple as follows:

$$\begin{aligned} F(\bar{a}, \bar{b}) &= \mathcal{F}(a_{(1)}, b_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)})) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)}) \\ \Phi(\bar{a}, \bar{b}, \bar{c}) &= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)}), \pi(c_{(2)})) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)} \cdot c_{(3)}) \\ \Psi(\bar{a}, \bar{b}, \bar{c}) &= \mathcal{F}(a_{(1)} \cdot b_{(1)}, c_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)}), \pi(c_{(2)})) \cdot \mathcal{F}^{-1}(a_{(3)} \cdot b_{(3)}, c_{(3)}) \end{aligned}$$

Note that a more natural definition would have been of the form $F = (\pi \otimes \pi)F'(\iota \otimes \iota)$ for some $F' : \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$. However, we can follow an argument made in the beginning of the proof of Theorem 4.2 in [7], since (cT1) implies that F' commutes with $\epsilon \otimes \epsilon$, and thereby composing with ι gives the same result.

Now we define the inverse of Φ by

$$\Phi^{-1}(\bar{a}, \bar{b}, \bar{c}) = \mathcal{F}^{-1}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)}), \pi(c_{(2)})) \cdot \mathcal{F}(a_{(3)}, b_{(3)} \cdot c_{(3)})$$

The inverse of F and Ψ are obtained exactly in the same way, where \mathcal{F}^1 and \mathcal{F} are swapped in their respective definitions. It then follows easily from (cT1), that Φ^{-1} and Φ are inverses

$$\begin{aligned} \Phi^{-1}\Phi(\bar{a}, \bar{b}, \bar{c}) &= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \mathcal{F}^{-1}(a_{(2)(1)}, b_{(2)(1)} \cdot c_{(2)(1)}) \cdot (\pi(a_{(2)(2)}), \pi(b_{(2)(2)}), \pi(c_{(2)(2)})) \\ &\quad \cdot \mathcal{F}(a_{(2)(3)}, b_{(2)(3)} \cdot c_{(2)(3)}) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)} \cdot c_{(3)}) \\ &= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \mathcal{F}^{-1}(a_{(2)}, b_{(2)} \cdot c_{(2)}) \cdot (\pi(a_{(3)}), \pi(b_{(3)}), \pi(c_{(3)})) \cdot \mathcal{F}(a_{(4)}, b_{(4)} \cdot c_{(4)}) \\ &\quad \cdot \mathcal{F}^{-1}(a_{(5)}, b_{(5)} \cdot c_{(5)}) \\ &= \epsilon(a_{(1)}) \cdot \epsilon(b_{(1)} \cdot c_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)}), \pi(c_{(2)})) \epsilon(a_{(3)}) \cdot \epsilon(b_{(3)} \cdot c_{(3)}) = (\bar{a}, \bar{b}, \bar{c}) \end{aligned}$$

Similar proofs hold for F and Ψ . Hence, (F, Φ, Ψ) restrict to automorphisms on the subsets of basis elements of $\mathcal{Q}^{\otimes 2}$ and $\mathcal{Q}^{\otimes 3}$ i.e. $R(\mathcal{H})^2$ and $R(\mathcal{H})^3$. Next we demonstrate that the axioms of Definition 2.2 hold.

First, we recall that the unit of the group $R(\mathcal{H})$ is $\bar{1}$ and by (cT3), it follows directly that (G1) and (G2) hold. Additionally, recall that the multiplication on $R(\mathcal{H})$ is defined as the projection of multiplication of \mathcal{H} . Hence, if we denote the multiplication of $R(\mathcal{H})$ by m , then

$$\begin{aligned} F(\bar{a}, \bar{b} \cdot \bar{c}) &= F(\bar{a}, \overline{b \cdot c}) = \mathcal{F}(a_{(1)}, (b \cdot c)_{(1)}) \cdot (\pi(a_{(2)}), \pi((b \cdot c)_{(2)})) \cdot \mathcal{F}^{-1}(a_{(3)}, (b \cdot c)_{(3)}) \\ &= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)} \cdot c_{(2)})) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)} \cdot c_{(3)}) \\ &= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)}) \cdot \pi(c_{(2)})) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)} \cdot c_{(3)}) = m_{23}\Phi(\bar{a}, \bar{b}, \bar{c}) \end{aligned}$$

holds. A symmetric argument show that (G4) also holds.

It still remains to show that the axioms of Definition 2.1 hold i.e. we must demonstrate that (F, Φ, Ψ) defines a twist on the underlying YBE solution $(R(\mathcal{H}), r)$ as well. Observe that combining the fact that \mathcal{F} is convolution invertible, (cT1), and (cT2) provides a symmetric relation for \mathcal{F}^{-1} :

$$\mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \cdot \mathcal{F}^{-1}(a_{(2)} \cdot b_{(2)}, c) = \mathcal{F}^{-1}(b_{(1)}, c_{(1)}) \cdot \mathcal{F}^{-1}(a, b_{(2)} \cdot c_{(2)}) \quad (\text{cT2}')$$

Condition (T1) follows directly from (cT2) and (cT2'):

$$F_{23}\Phi(\bar{a}, \bar{b}, \bar{c}) = \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot \mathcal{F}(b_{(2)(1)}, c_{(2)(1)}) \cdot (\pi(a_{(2)}), \pi(b_{(2)(2)}), \pi(c_{(2)(2)}))$$

$$\begin{aligned}
& \cdot \mathcal{F}^{-1}(b_{(2)(3)}, c_{(2)(3)}) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)} \cdot c_{(3)}) \\
&= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot \mathcal{F}(b_{(2)}, c_{(2)}) \cdot (\pi(a_{(2)}), \pi(b_{(3)}), \pi(c_{(3)})) \cdot \mathcal{F}^{-1}(b_{(4)}, c_{(4)}) \\
& \quad \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(5)} \cdot c_{(5)}) \\
&= \mathcal{F}(a_{(1)} \cdot b_{(1)}, c_{(1)}) \cdot \mathcal{F}(a_{(2)}, b_{(2)}) (\pi(a_{(3)}), \pi(b_{(3)}), \pi(c_{(2)})) \cdot \mathcal{F}^{-1}(a_{(4)}, b_{(4)}) \\
& \quad \cdot \mathcal{F}^{-1}(a_{(5)} \cdot b_{(5)} \cdot c_{(3)}) = F_{12} \Psi(\bar{a}, \bar{b}, \bar{c})
\end{aligned}$$

Recall from Section 4 of [7] that r is defined by

$$r(\bar{a}, \bar{b}) = \mathcal{R}(a_{(1)}, b_{(1)}) \cdot (\pi(b_{(2)}), \pi(a_{(2)})) \cdot \mathcal{R}^{-1}(a_{(3)}, b_{(3)}) \quad (12)$$

Moreover, \mathcal{R} satisfies $\mathcal{R}(b_{(1)}, a_{(1)})a_{(2)} \cdot b_{(2)} = b_{(1)} \cdot a_{(1)} \mathcal{R}(b_{(2)}, a_{(2)})$ and as for \mathcal{F} and (cT2''), we can obtain a symmetric relation for \mathcal{R}^{-1} , since \mathcal{R} is convolution invertible. Using these relations we observe that (T2) holds:

$$\begin{aligned}
r_{23} \Phi(\bar{a}, \bar{b}, \bar{c}) &= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot \mathcal{R}(b_{(2)(1)}, c_{(2)(1)}) \cdot (\pi(a_{(2)}), \pi(c_{(2)(2)}), \pi(b_{(2)(2)})) \\
& \quad \cdot \mathcal{R}^{-1}(b_{(2)(3)}, c_{(2)(3)}) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)} \cdot c_{(3)}) \\
&= \mathcal{F}(a_{(1)}, b_{(1)} \cdot c_{(1)}) \cdot \mathcal{R}(b_{(2)}, c_{(2)}) \cdot (\pi(a_{(2)}), \pi(c_{(3)}), \pi(b_{(3)})) \cdot \mathcal{R}^{-1}(b_{(4)}, c_{(4)}) \\
& \quad \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(5)} \cdot c_{(5)}) \\
&= \mathcal{F}(a_{(1)}, b_{(1)(1)} \cdot c_{(1)(1)}) \cdot \mathcal{R}(b_{(1)(2)}, c_{(1)(2)}) \cdot (\pi(a_{(2)}), \pi(c_{(2)}), \pi(b_{(2)})) \\
& \quad \cdot \mathcal{R}^{-1}(b_{(3)(1)}, c_{(3)(1)}) \cdot \mathcal{F}^{-1}(a_{(3)}, b_{(3)(2)} \cdot c_{(3)(2)}) \\
&= \mathcal{R}(b_{(1)(1)}, c_{(1)(1)}) \cdot \mathcal{F}(a_{(1)}, c_{(1)(2)} \cdot b_{(1)(2)}) \cdot (\pi(a_{(2)}), \pi(c_{(2)}), \pi(b_{(2)})) \\
& \quad \cdot \mathcal{F}^{-1}(a_{(3)}, c_{(3)(1)} \cdot b_{(3)(1)}) \cdot \mathcal{R}^{-1}(b_{(3)(2)}, c_{(3)(2)}) \\
&= \mathcal{R}(b_{(1)}, c_{(1)}) \cdot \mathcal{F}(a_{(1)}, c_{(2)(1)} \cdot b_{(2)(1)}) \cdot (\pi(a_{(2)}), \pi(c_{(2)(2)}), \pi(b_{(2)(2)})) \\
& \quad \cdot \mathcal{F}^{-1}(a_{(3)}, c_{(2)(3)} \cdot b_{(2)(3)}) \cdot \mathcal{R}^{-1}(b_{(3)}, c_{(3)}) = \Phi r_{23}(\bar{a}, \bar{b}, \bar{c})
\end{aligned}$$

A symmetric argument shows that (T3) holds. \square

Corollary 3.2. *Let \mathcal{F} be a co-twist on $(\mathcal{H}, \mathcal{R})$, and (F, Φ, Ψ) the induced twist on $R(\mathcal{H})$, by Theorem 3.1. The induced skew brace structure induced on $R(\mathcal{H}^{\mathcal{F}})$ agrees with that induced by the twist (F, Φ, Ψ) .*

Proof. First recall that the underlying set of the remnant is solely dependent on the counit and thereby $G = R(H_{\omega_m}) = R(H^{\mathcal{F}})$ as sets. Let us denote the twisted product on $R(\mathcal{H}^{\mathcal{F}})$ by $\cdot^{\mathcal{F}}$, then by (10) we have that

$$\begin{aligned}
\bar{a} \cdot^{\mathcal{F}} \bar{b} &= \pi m^{\mathcal{F}}(a \vee D, b \vee D) = \mathcal{F}^{-1}(a_{(1)} \vee D_{(1)}, b_{(1)} \vee D_{(1)}) \cdot \pi m(a_{(2)} \vee D_{(2)}, b_{(2)} \vee D_{(2)}) \\
& \quad \cdot \mathcal{F}(a_{(3)} \vee D_{(3)}, b_{(3)} \vee D_{(3)}) \\
&= \mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \cdot \pi m(a_{(2)} \vee D, b_{(2)} \vee D) \cdot \mathcal{F}(a_{(3)}, b_{(3)}) = \pi m(\iota \otimes \iota) F^{-1}(\bar{a}, \bar{b})
\end{aligned}$$

What remains to be shown is that $r^{\mathcal{F}} = F r F^{-1}$, where we denote the induced braiding operator on $R(H^{\mathcal{F}})$ by $r^{\mathcal{F}}$, which follows by (11) and (12):

$$\begin{aligned}
r^{\mathcal{F}}(\bar{a}, \bar{b}) &= \mathcal{F}^{-1}(a_{(1)(1)}, b_{(1)(1)}) \mathcal{R}(a_{(1)(2)}, b_{(1)(2)}) \cdot \mathcal{F}(b_{(1)(3)}, a_{(1)(3)}) \cdot (\pi(b_{(2)}), \pi(a_{(2)})) \\
& \quad \cdot \mathcal{F}^{-1}(b_{(3)(1)}, a_{(3)(1)}) \mathcal{R}^{-1}(a_{(3)(2)}, b_{(3)(2)}) \cdot \mathcal{F}(a_{(3)(3)}, b_{(3)(3)}) \\
&= \mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \mathcal{R}(a_{(2)}, b_{(2)}) \cdot \mathcal{F}(b_{(3)(1)}, a_{(3)(1)}) \cdot (\pi(b_{(3)(2)}), \pi(a_{(3)(2)}))
\end{aligned}$$

$$\begin{aligned}
& \mathcal{F}^{-1}(b_{(3)(3)}, a_{(3)(3)}) \mathcal{R}^{-1}(a_{(4)}, b_{(4)}) \cdot \mathcal{F}(a_{(5)}, b_{(5)}) \\
&= \mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \mathcal{R}(a_{(2)}, b_{(2)}) \cdot F(\pi(b_{(3)}), \pi(a_{(3)})) \cdot \mathcal{R}^{-1}(a_{(4)}, b_{(4)}) \cdot \mathcal{F}(a_{(5)}, b_{(5)}) \\
&= \mathcal{F}^{-1}(a_{(1)}, b_{(1)}) \cdot Fr(\pi(a_{(2)}), \pi(b_{(2)})) \cdot \mathcal{F}(b_{(3)}, a_{(3)}) = FrF^{-1}(\bar{a}, \bar{b}) \quad \square
\end{aligned}$$

Each skew brace can arise as the remnant of non-isomorphic CQHAs in SupLat. It is not clear whether every twist on a skew brace can be extended to co-twists on some CQHA which recovers it as its remnant. In what follows we will discuss co-twists on the two families of Hopf algebras in SupLat which were studied in [7] and comment on the obstruction of extending a twist on the remnant to a co-twist on these algebras.

In Section 5.1 of [7], we constructed the SupLat-FRT algebra H_ω corresponding to a YBE solution (X, r) . As an algebra in SupLat, H_ω is generated by elements $(x, y)_1$ and $(x, y)_2$, corresponding to pairs $x, y \in X$, and has relations

$$\vee_{a \in X} \{(x, a)_1 \cdot (x, a)_2\} = 1 = \vee_{a \in X} \{(a, x)_2 \cdot (a, x)_1\} \quad (13)$$

$$(x, a)_1 \cdot (y, a)_2 = \emptyset = (a, x)_2 \cdot (a, y)_1, \text{ when } x \neq y \quad (14)$$

$$(x, y)_1 \cdot (a, b)_1 = (\sigma_x(a), \sigma_y(b))_1 \cdot (\gamma_a(x), \gamma_b(y))_1 \quad (15)$$

for $a, b, x, y \in X$. We refer the reader to Section 5.1 of [7] for additional details and the definitions of the structural maps of the Hopf algebra. It was also shown that the remnant of the SupLat-FRT algebra of a solution recovers the universal skew brace of the solution. Consequently, by Theorem 3.1 co-twists on the SupLat-FRT algebra H_ω of a solution induce twists on its universal skew brace. However, in [7] we noted that H_ω carries much more information about the solution compared to the universal skew brace. This is confirm in the study of twists since any co-twist on SupLat-FRT algebra of a solution also induces a twist on the YBE solution itself.

Theorem 3.3. *If H_ω denotes the SupLat-FRT algebra of a set-theoretical YBE solution (X, r) , then a Drinfeld co-twists on H_ω induces a Drinfeld twists on (X, r) , as in Definition 2.1.*

Proof. Let $\mathcal{F} : H_\omega \otimes H_\omega \rightarrow \mathcal{P}(1)$ denote a co-twist on H_ω . We define the triple (F, Φ, Ψ) as follows:

$$\begin{aligned}
F(x, y) &= (a, b) \text{ if and only if } \mathcal{F}(\{(x, a)_1, (y, b)_1\}) = 1 \\
\Phi(x, y, z) &= (a, b, c) \text{ if and only if } \mathcal{F}(\{(x, a)_1, (y, b)_1, (z, c)_1\}) = 1 \\
\Psi(x, y, z) &= (a, b, c) \text{ if and only if } \mathcal{F}(\{(x, a)_1, (y, b)_1, (z, c)_1\}) = 1
\end{aligned}$$

We must first demonstrate that F, Φ and Ψ are indeed maps, rather than relations between X^2 and itself. By (cT1), we have that for a fixed $a, b, a', b' \in X$:

$$\begin{aligned}
\bigvee_{m, n \in X} \mathcal{F}(\{(a, m)_1, (b, n)_1\}) \cdot \mathcal{F}^{-1}(\{(m, a')_1, (n, b')_1\}) &= \delta_{a, a'} \cdot \delta_{b, b'} \\
&= \bigvee_{m, n \in X} \mathcal{F}^{-1}(\{(a, m)_1, (b, n)_1\}) \cdot \mathcal{F}(\{(m, a')_1, (n, b')_1\})
\end{aligned}$$

Consequently, if we let $F^{-1}(x, y) := (a, b)$ if and only if $\mathcal{F}^{-1}(\{(x, a)_1, (y, b)_1\}) = 1$, then by the above equation $FF^{-1} = F^{-1}F = \text{id}_{X^2}$. Hence, both F and F^{-1} must be bijective maps, since a relation $R \subset A \times B$ between two sets is invertible by composition of relations, if and only if it represents a bijective map. Similarly, (cT1) implies that Φ and Ψ as we defined them are both well-defined and bijective maps from X^3 to itself.

Now we substitute $a = (a, a')_1$, $b = (b, b')_1$ and $c = (c, c')_1$ in equation (cT2), for arbitrary $a, a', b, b', c, c' \in X$ and observe that the left hand side

$$\bigvee_{m, n \in X} \mathcal{F}(\{(a, m)_1, (b, n)_1, (c, c')_1\}) \cdot \mathcal{F}(\{(m, a')_1, (n, b')_1\})$$

is equal to 1 if and only if $(a', b', c') = F_{12}\Psi(a, b, c)$, while the right hand side

$$\bigvee_{m, n \in X} \mathcal{F}(\{(a, a')_1, (b, m)_1, (c, n)_1\}) \cdot \mathcal{F}(\{(m, b')_1, (n, c')_1\})$$

is equal to 1 if and only if $(a', b', c') = F_{23}\Phi(a, b, c)$. Hence, (T1) holds. One the other hand, by the relations imposed in H_ω , we have that

$$\mathcal{F}(\{(x, a)_1, (y, b)_1, (z, c)_1\}) = \mathcal{F}(\{\sigma_x(y), \sigma_a(b))_1, (\gamma_y(x), \gamma_b(a))_1, (z, c)_1\})$$

This directly implies that $r_{12}\Psi = \Psi r_{12}$ and that (T3) holds. A symmetric argument implies that (T2) holds. \square

Note that the converse of the above statement is not true i.e. not every twist on a YBE solution can be realised as a co-twist on H_ω . Firstly, co-twists on H_ω are built from twists on the underlying YBE solution which lift to its universal skew brace and we have already seen counterexamples for such situations in Example 2.4. Additionally, for a twist (F, Φ, Ψ) to extend to a co-twist on H_ω , we require a lot more data. If we call the multiplication of n generators of H_ω , a word of order n , then the triple (F, Φ, Ψ) only determine the action of \mathcal{F} on pairs of words of orders $(1, 1)$, $(1, 2)$ and $(2, 1)$, respectively. For a co-twist \mathcal{F} on H_ω , we need the additional datum of its action on pairs of words of any order, which will have to be compatible with the triple via (cT2). Section III of [9] provides a full discussion of this in the linear setting, while [3] also addresses a weaker notion of twists on FRT-type algebras coming from the linearisation of involutive set-theoretical YBE solutions. The new ingredient for extending a twist to the SupLat-FRT construction, is how \mathcal{F} is defined on generators of the form $(x, y)_2$ in H_ω . These actions will be uniquely determined by the action of \mathcal{F} on generators $(x, y)_1$, but will impose a *non-degeneracy* condition on F , as discussed in our comments after Theorem 2.1.

In Section 5.2 of [7], it was also shown that any group (G, m, e) with a braiding operator r appears as the remnant of a CQHA in SupLat. The relevant Hopf algebra was denoted by H_{ω_m} and is the quotient of the SupLat-FRT algebra of the underlying YBE solution (G, r) , with additional relations $(e, e)_1 = 1 = (e, e)_2$ and

$$(a.b, c)_1 = \bigvee_{d, f \in G \text{ with } d.f=c} \{(a, d)_1, (b, f)_1\} \quad \text{and} \quad (a, b.c)_2 = \bigvee_{d, f \in G \text{ with } d.f=a} \{(f, c)_2, (d, b)_2\} \quad (16)$$

for any $a, b, c \in G$. By Theorem 3.1, any co-twist on H_{ω_m} induces a twist on the skew brace G . Alternatively, since H_{ω_m} is defined in a similar fashion to H_ω , one could define the triple (F, Φ, Ψ) as done in Theorem 3.3. Nevertheless, it's easy to observe that by definition both methods would provide the same triple. As in the proof of Theorem 3.3, the latter method provides a straightforward way to derive the definitions in Section 2, directly by studying the SupLat-FRT algebras. As we will see in the next section, this is the key advantage which the SupLat-FRT algebras posses over the classically considered groups with unique factorisation, when it comes to the study of set-theoretical YBE solutions.

4 Co-twists and Groups with Unique Factorisation

In this section, we will classify co-twist on dualizable Hopf algebras in SupLat and thereby describe the induced Drinfeld twists on their remnants. As described in Section 4.1 of [7], dualizable Hopf algebras in SupLat are all of the form $\mathcal{P}(G)$, where G is a group with unique factorization $G = G_+ \cdot G_-$ and the Hopf algebra structure of $\mathcal{P}(G)$ is given as the bicrossproduct of the group algebra $\mathcal{P}(G_-)$ and the function algebra $\mathcal{P}(G_+)$. This theory was developed first in [12] and quasitriangular structures on these Hopf algebras were classified in [11]. In this section, we will adapt the notation of [11] on groups with unique factorisation, which are also called matched pairs of groups. We refer the reader to Section 4.1 of [7] for additional details on the re-phrasing of the results of [12, 11] in the setting of SupLat.

Throughout this section we will assume $G = G_+ \cdot G_-$ is a group with unique factorisation unless stated otherwise and $\mathcal{P}(G)$ denotes the relevant Hopf algebra in SupLat.

Lemma 4.1. *Convolution invertible $\mathcal{F} : \mathcal{P}(G) \otimes \mathcal{P}(G) \rightarrow \mathcal{P}(1)$ i.e. \mathcal{F} which satisfy (cT1) are in bijection with maps $\Theta : G_- \times G_- \rightarrow G_+ \times G_+$ such that*

$$F_\Theta : (g_-, h_-) \rightarrow \left(\Theta_1(g_-, h_-)g_-, \Theta_2(g_-, h_-)h_- \right) \quad (17)$$

defines a bijection from $G_- \times G_-$ to itself, where we denote $\Theta(g_-, h_-) = (\Theta_1(g_-, h_-), \Theta_2(g_-, h_-))$.

Proof. Given a convolution invertible \mathcal{F} , we can define $\Theta(g_-, h_-) = (a_+, b_+)$ for elements where $\mathcal{F}^{-1}(a_+g_-, b_+h_-) = 1$. First, we must show that Θ is well-defined. By (cT1), for any pair $g_-, h_- \in G_-$ there exists at least one pair $a_+, b_+ \in G_+$ such that

$$\mathcal{F}^{-1}(a_+g_-, b_+h_-) = 1 = \mathcal{F}\left(a_+^{-1}(a_+g_-), b_+^{-1}(b_+h_-)\right)$$

and if $\mathcal{F}(k_+(a_+g_-), l_+(b_+h_-)) = 1$, then $k_+ = a_+^{-1}$ and $l_+ = b_+^{-1}$. A symmetric statement holds for \mathcal{F} and for any $g_-, h_- \in G_-$, there exist at least one pair $m_+, n_+ \in G_+$ such that $\mathcal{F}(m_+g_-, n_+h_-) = 1 = \mathcal{F}^{-1}(m_+^{-1}(m_+g_-), n_+^{-1}(n_+h_-))$. By the latter statement, for the pair a_+g_-, b_+h_- there exists at least one pair $k_+, l_+ \in G_+$ satisfying $\mathcal{F}(k_+(a_+g_-), l_+(b_+h_-)) = 1$ and by the first statement we have that $k_+ = a_+^{-1}$ and $l_+ = b_+^{-1}$. Hence, if $\mathcal{F}^{-1}(a_+g_-, b_+h_-) = 1$, then $\mathcal{F}(a_+^{-1}(a_+g_-), b_+^{-1}(b_+h_-)) = 1$ and the symmetric statement holds for \mathcal{F} .

Now we show that for any $g_-, h_- \in G_-$, the pair a_+, b_+ satisfying $\mathcal{F}^{-1}(a_+g_-, b_+h_-) = 1$ is unique. Assume that the pair k_+, l_+ also satisfies $\mathcal{F}^{-1}(k_+g_-, l_+h_-) = 1$, then $\mathcal{F}(k_+^{-1}(k_+g_-), l_+^{-1}(l_+h_-)) = 1$ and thereby

$$\mathcal{F}^{-1}(a_+g_-, b_+h_-) \cdot \mathcal{F}\left(k_+^{-1}(k_+g_-), l_+^{-1}(l_+h_-)\right) = 1 = \epsilon(a_+k_+^{-1}g_-) \cdot \epsilon(b_+l_+^{-1}h_-)$$

Hence, $k_+ = a_+$ and $l_+ = b_+$ and thereby Θ is a well-defined map. In a symmetric way, we can define $\Theta' : G_- \times G_- \rightarrow G_+ \times G_+$ by $\Theta'(g_-, h_-) = (a_+, b_+)$ for $\mathcal{F}(a_+g_-, b_+h_-) = 1$. By (cT1), it then follows that the maps

$$F_\Theta(g_-, h_-) = \left(\Theta_1(g_-, h_-)g_-, \Theta_2(g_-, h_-)h_- \right), \quad F_{\Theta'}(g_-, h_-) = \left(\Theta'_1(g_-, h_-)g_-, \Theta'_2(g_-, h_-)h_- \right)$$

are inverses. \square

Before we resume, recall from Section 4.1 of [7] that the remnant of $\mathcal{P}(G)$ is the group G_- and co-quasitriangular structures on $\mathcal{P}(G)$ are in bijection with pairs of maps $\eta, \xi : G_- \rightarrow G_+$ satisfying

additional conditions. Now we calculate the induced twist data on G_- by Theorem 3.1. Note that co-twist on a Hopf algebra are defined independent of any co-quasitriangular structure and should not take the braiding operator into consideration. While the SupLat-FRT algebras depended directly on the braiding, this is not the case for $\mathcal{P}(G)$. In this case, the twists induced on G_- by Theorem 3.1 will automatically be compatible with any of the possible braiding operators which arise from co-quasitriangular structures on $\mathcal{P}(G)$. We will denote the units of G_- and G_+ by e_- and e_+ , respectively.

Theorem 4.2. *Co-twists on $\mathcal{P}(G)$ are in bijection with maps $\Theta : G_- \times G_- \rightarrow G_+ \times G_+$ such that F_Θ , as defined in Lemma 4.1, is invertible and Θ satisfies $\Theta_2(e_-, a_-) = e_+ = \Theta_1(a_-, e_-)$ and the following conditions:*

$$\Theta_1 \left(\Theta_1(ab, c)a, \Theta_1(ab, c)^a b \right) \cdot \Theta_1(ab, c) = \Theta_1(a, bc) \quad (18)$$

$$\Theta_2 \left(\Theta_1(ab, c)a, \Theta_1(ab, c)^a b \right) \cdot \Theta_1(ab, c)^a = \Theta_1 \left(\Theta_2(a, bc)b, \Theta_2(a, bc)^b c \right) \cdot \Theta_2(a, bc) \quad (19)$$

$$\Theta_2(ab, c) = \Theta_2 \left(\Theta_2(a, bc)b, \Theta_2(a, bc)^b c \right) \cdot \Theta_2(a, bc)^b \quad (20)$$

where $a, b, c \in G_-$ and the equations above hold in G_+ .

Proof. First we recall that the unit element in $\mathcal{P}(G)$ is $\bigvee_{g_+ \in G_+} g_+$ and thereby (cT3) would translate to $\bigvee_{g_+ \in G_+} \mathcal{F}^{-1}(g_+, a_+ a_-) = 1$ if and only if $a_+ = e_+ \in G_+$. Equivalently, $\Theta_2(e_-, a_-) = e_+$. A symmetric argument show that $\Theta_1(a_-, e_-) = e_+$. Since Θ is formulated in terms of \mathcal{F}^{-1} , we will look at the equivalent condition to (cT2) in terms of \mathcal{F}^{-1} , (cT2'). Let $a, b, c \in G$, then the left hand side of equation (cT2') expands as follows:

$$\begin{aligned} & \bigvee_{g, h \in G_+} \mathcal{F}^{-1} \left(a_+ g^{-1}({}^g a_-), b_+ h^{-1}({}^h b_-) \right) \cdot \mathcal{F}^{-1}(ga_- hb_-, c_+ c_-) \\ &= \bigvee_{g \in G_+} \mathcal{F}^{-1} \left(a_+ g^{-1}({}^g a_-), b_+ (g^{a_-})^{-1}({}^{g^{a_-}} b_-) \right) \cdot \mathcal{F}^{-1}(ga_- b_-, c_+ c_-) \end{aligned}$$

The equation is equal to 1 if and only if $a_+ = \Theta_1 \left({}^g a_-, {}^{g^{a_-}} b_- \right) \cdot g$, $b_+ = \Theta_2 \left({}^g a_-, {}^{g^{a_-}} b_- \right) \cdot g^{a_-}$, $c_+ = \Theta_2(a_- b_-, c_-)$ and $g = \Theta_1(a_- b_-, c_-)$. While the right hand side of (cT2') expands as

$$\begin{aligned} & \bigvee_{g, h \in G_+} \mathcal{F}^{-1} \left(b_+ g^{-1}({}^g b_-), c_+ h^{-1}({}^h c_-) \right) \cdot \mathcal{F}^{-1}(a_+ a_-, gb_- hc_-) \\ &= \bigvee_{g \in G_+} \mathcal{F}^{-1} \left(b_+ g^{-1}({}^g b_-), c_+ (g^{b_-})^{-1}({}^{g^{b_-}} c_-) \right) \cdot \mathcal{F}^{-1}(a_+ a_-, gb_- c_-) \end{aligned}$$

which will be equal to 1 if and only if $b_+ = \Theta_1 \left({}^g b_-, {}^{g^{b_-}} c_- \right) \cdot g$, $c_+ = \Theta_2 \left({}^g b_-, {}^{g^{b_-}} c_- \right) \cdot g^{b_-}$, $a_+ = \Theta_1(a_-, b_- c_-)$ and $g = \Theta_2(a_-, b_- c_-)$. The equalities in Theorem then simply arise as equating the value of a_+ , b_+ and c_+ from the left and right hand side of (cT2'), respectively. \square

The works [12, 11] focus on linear Hopf algebras with *positive bases* and show that all such Hopf algebras come from finite groups with unique factorisation, and classify all *positive* quasitriangular structures on them. The proofs in these works consist of two parts. The first part is finding an isomorphism so that the structural maps don't have any scalar factor, which basically translates the problem into a problem about Hopf algebras on free lattices. The second part of their proofs classify such structures and fully translate into SupLat, see Section 4.1 of [7]. Hence, we expect that the above classification can be re-phrased as classifying all *positive* Drinfeld co-twists on linear Hopf algebras with positive bases.

Corollary 4.3. *Given a co-twist on $\mathcal{P}(G)$ defined by a map Θ , we obtain a twist $(F_\Theta, \Phi_\Theta, \Psi_\Theta)$ on the group G_- , where F_Θ is as defined in Lemma 4.1 and the other two maps are defined by*

$$\Phi_\Theta(a, b, c) = \left(\Theta_1(a, bc)_a, \Theta_2(a, bc)_b, \Theta_2(a, bc)^b_c \right) \quad (21)$$

$$\Psi_\Theta(a, b, c) = \left(\Theta_1(ab, c)_a, \Theta_1(ab, c)^a_b, \Theta_2(ab, c)_b \right) \quad (22)$$

for $a, b, c \in G_-$. In particular, the twist on G_- is independent of η and ξ and the braiding operator on G_- .

Proof. Recall from Section 4.1 of [7] that $\pi : G \rightarrow G_-$ is defined by $\pi(a_+ a_-) = \delta_{a_+, e_+} a_-$ for $a_+ \in G_+$ and $a_- \in G_-$. Hence, for $a, b \in G_-$ we have

$$\begin{aligned} F(a, b) &= \bigvee_{g, h, k, l \in G_+} \mathcal{F} \left(g^{-1}(g a), h^{-1}(h b) \right) \cdot (\pi(g k^{-1}(k a)), \pi(h l^{-1}(l b))) \cdot \mathcal{F}^{-1}(k a, l b) \\ &= \bigvee_{g, h \in G_+} \mathcal{F} \left(g^{-1}(g a), h^{-1}(h b) \right) \cdot (\pi(g a), \pi(h b)) \cdot \mathcal{F}^{-1}(g a, h b) = (\Theta_1(a, b)_a, \Theta_1(a, b)_b) = F_\Theta(a, b) \end{aligned}$$

A similar calculation follows for Φ_Θ and Ψ_Θ , using the fact that $(^h b) (^{h^b} c) = {}^h(bc)$ for $b, c \in G_-$ and $h \in G_+$ [Equation (1) in [11]]:

$$\begin{aligned} \Phi(a, b, c) &= \bigvee_{g, h, f, k, l, m \in G_+} \mathcal{F} \left(g^{-1}(g a), h^{-1}(h b) \cdot f^{-1}(f c) \right) \cdot (\pi(g k^{-1}(k a)), \pi(h l^{-1}(l b)), \pi(f m^{-1}(m b))) \\ &\quad \cdot \mathcal{F}^{-1}(k a, l b, m c) \\ &= \bigvee_{g, h, f \in G_+} \mathcal{F} \left(g^{-1}(g a), h^{-1}(h b) \cdot f^{-1}(f c) \right) \cdot (\pi(g a), \pi(h b), \pi(f b)) \cdot \mathcal{F}^{-1}(g a, h b, f c) \\ &= \bigvee_{g, h \in G_+} \mathcal{F} \left(g^{-1}(g a), h^{-1}(h b) (^{h^b} c) \right) \cdot (\pi(g a), \pi(h b), \pi(^{h^b} c)) \cdot \mathcal{F}^{-1}(g a, h b c) \\ &= \left(\Theta_1(a, bc)_a, \Theta_2(a, bc)_b, \Theta_2(a, bc)^b_c \right) \\ \Psi(a, b, c) &= \bigvee_{g, h, f, k, l, m \in G_+} \mathcal{F} \left(g^{-1}(g a) \cdot h^{-1}(h b), f^{-1}(f c) \right) \cdot (\pi(g k^{-1}(k a)), \pi(h l^{-1}(l b)), \pi(f m^{-1}(m b))) \\ &\quad \cdot \mathcal{F}^{-1}(k a, l b, m c) \\ &= \bigvee_{g, h, f \in G_+} \mathcal{F} \left(g^{-1}(g a) \cdot h^{-1}(h b), f^{-1}(f c) \right) \cdot (\pi(g a), \pi(h b), \pi(f b)) \cdot \mathcal{F}^{-1}(g a, h b, f c) \\ &= \bigvee_{g, f \in G_+} \mathcal{F} \left(g^{-1}(g a) (^{g^a} b), f^{-1}(f c) \right) \cdot (\pi(g a), \pi(^{g^a} b), \pi(f b)) \cdot \mathcal{F}^{-1}(g a b, f c) \\ &= \left(\Theta_1(ab, c)_a, \Theta_1(ab, c)^a_b, \Theta_2(ab, c)_c \right) \end{aligned}$$

□

As mentioned before the proof of this result co-quasitriangular structures on $\mathcal{P}(G)$ are classified by maps $\eta, \xi : G_- \rightarrow G_+$ satisfying conditions of Proposition 1 in [11]. Given such a pair we obtain an induced braiding operator on G_- defined by $r : (g, h) \mapsto ({}^{\eta(g)} h, g^{\xi(h)})$. Theorem 3.1 tells us that the

triples defined above produce a twist on (G_-, r) regardless of the choice of η and ξ . This can be verified directly as well and follows from the axioms which η and ξ satisfy.

A skew brace (G_-, r) coming from a matched pair of groups can also arise as the remnant of a SupLat-FRT algebra. However, given a twist on (G_-, r) the obstruction of extending the twist to a co-twist on the two CQHAs is very different. In the FRT case, we were missing additional information, whereas here the sole obstruction to extending a twist, F , to a co-twist on $\mathcal{P}(G)$ is whether it can be written as F_Θ for some Θ , which in return involves the group G_+ and its action on G_- .

The simplest case of groups with unique factorisation are provided by groups with braiding operators themselves. If (G, r) is of the latter form, then we can define $G_- = G_+ = G$ with actions

$$g_+ g_- = \sigma_{g_+}(g_-), \quad g_+^{g_-} = \gamma_{g_-}(g_+), \quad g_- g_+ = \tau_{g_+}(g_-), \quad g_-^{g_+} = \rho_{g_+}(g_-)$$

where $r^{-1}(x, y) = (\tau_x(y), \rho_y(x))$. Thereby, we obtain a group with unique factorisation G_+, G_- on the underlying set $G \times G$. The braiding operator on G_- is then recovered by $\eta = \xi = \text{id}_G$. Hence, we can re-write Theorems 4.2 and Corollary 4.3 solely in terms of (G, r) .

Corollary 4.4. *Let (G, m, e) be a group with a braiding operator. If $\Theta : G^2 \rightarrow G^2$ is a map such that $F_\Theta : G^2 \rightarrow G^2$ defined by $F_\Theta(g, h) = (\sigma_{\Theta_1(g, h)}(h), \sigma_{\Theta_2(g, h)}(g))$ is a bijection and satisfies $\Theta_2(e_-, a_-) = e_+ = \Theta_1(a_-, e_-)$ and*

$$\Theta_1(\sigma_{\Theta_1(ab, c)}(a), \sigma_{\gamma_a(\Theta_1(ab, c))}(b)) \cdot \Theta_1(ab, c) = \Theta_1(a, bc) \quad (23)$$

$$\Theta_2(ab, c) = \Theta_2(\sigma_{\Theta_2(a, bc)}(b), \sigma_{\gamma_b(\Theta_2(a, bc))}(c)) \cdot \gamma_b(\Theta_2(a, bc)) \quad (24)$$

$$\Theta_2(\sigma_{\Theta_1(ab, c)}(a), \sigma_{\gamma_a(\Theta_1(ab, c))}(b)) \cdot \gamma_a(\Theta_1(ab, c)) = \Theta_1(\sigma_{\Theta_2(a, bc)}(b), \sigma_{\gamma_b(\Theta_2(a, bc))}(c)) \cdot \Theta_2(a, bc) \quad (25)$$

for $a, b, c \in G$, then we obtain a new group (G, mF_Θ^{-1}, e) with a braiding operator $F_\Theta r F_\Theta^{-1}$.

We have already seen examples of twists which can be written of the above form in Section 2. If we define $\Theta(x, y) = (e, x)$, then $F_\Theta(x, y) = (x, \sigma_x(y))$ is bijective by definition, while (23) and (24) hold in a trivial manner and (24) reduces to $ab = \sigma_a(b)\gamma_b(a)$. With this choice of Θ , we recover the triple $F_\Theta(x, y) = (x, \sigma_x(y))$, $\Phi_\Theta(x, y, z) = (x, \sigma_x(y), \sigma_{\gamma_y(x)}(z))$ and $\Psi_\Theta(x, y, z) = (x, y, \sigma_{xy}(z))$ of Example 2.5. The reader should also note that the inverse twists to this family of twists cannot be written in the form of Corollary 4.4. By Theorem 2.6, any twist of the mentioned form has an inverse twist (F', Φ', Ψ') with $F' = F^{-1}$ on the group (G, \star) with the trivial braiding operator $(x, y) \mapsto (y, y \star x \star y)$. Due to the trivial braiding operator on (G, \star, \star) , the twists coming from Corollary 4.4 on this skew brace are independent of the choice of Θ and define the same twist given by $F_\Theta(x, y) = (y, x)$. Given this limitation, we can conclude that the general theory Drinfeld twists described in Section 2, truly goes beyond the study of twist on Hopf algebras coming from matched pairs of groups.

5 Concluding Remarks and Outlook

While in [7], we provided the theory which relates skew braces with Hopf algebra objects in a suitable category, SupLat, the introduction of Drinfeld twists in this work is the first step in advancing this theory by translating Hopf algebraic constructions, such as Drinfeld twists, into constructions which are applicable to skew braces in a compatible way. There are a number of other constructions on CQHAs which should have fruitful applications to the study of skew braces, such as the theory of co-double bosonisation [1]. The studies in Sections 3 and 4, also fully demonstrate the advantage of applying Hopf algebraic techniques to general Hopf algebras in SupLat and observing the effect on their remnants compared to studying these technique on Hopf algebras coming from matched pairs of groups.

There are several questions which remain open with regards to the discussion of Drinfeld twists on skew brace. The first natural question is whether every twist on a skew brace can be extended to a co-twist on a CQHA in SupLat, which has this skew brace as its remnant. The second part of this question is whether every twist can be extended canonically to a co-twist on the SupLat-FRT algebra H_{ω_m} related to the skew brace. As we mentioned in Section 3, more data is needed to define a co-twist on H_{ω_m} , but there might be a natural choice for selecting this data. The computational aspects of this theory might also be of interest. By Example 2.5, any skew brace can be obtained as a twist from the trivial skew brace of its additive group. Hence, one could write an algorithm for to classify twists on such objects and thereby all skew braces with the same additive group. It is not however clear whether such an algorithm or variations of it could have a computational advantage to the algorithm presented in Section 5 of [8].

References

- [1] Ryan Aziz and Shahn Majid. Co-double bosonisation and dual bases of $c_q[SL2]$ and $c_q[SL3]$. *Journal of Algebra*, 518:75–118, 2019.
- [2] David Bachiller. Solutions of the Yang-Baxter equation associated to skew left braces, with applications to racks. *Journal of Knot Theory and Its Ramifications*, 27(08):1850055, 2018.
- [3] Anastasia Doikou. Set theoretic Yang-Baxter equation, braces and Drinfeld twists. *arXiv preprint arXiv:2102.13591*, 2021.
- [4] Anastasia Doikou and Agata Smoktunowicz. Set-theoretic Yang-Baxter & reflection equations and quantum group symmetries. *arXiv preprint arXiv:2003.08317*, 2020.
- [5] Vladimir Gershonovich Drinfeld. Quasi-Hopf algebras. *Algebra i Analiz*, 1(6):114–148, 1989.
- [6] Vladimir Guerchonovitch Drinfeld. Quasi-Hopf algebras and Knizhnik-Zamolodchikov equations. In *Problems of modern quantum field theory*, pages 1–13. Springer, 1989.
- [7] Aryan Ghobadi. Skew braces as remnants of co-quasitriangular Hopf algebras in SupLat. *arXiv preprint arXiv:2009.12815*, 2020.
- [8] Leandro Guarnieri and Leandro Vendramin. Skew braces and the Yang-Baxter equation. *Mathematics of Computation*, 86(307):2519–2534, 2017.
- [9] PP Kulish and AI Mudrov. On twisting solutions to the Yang-Baxter equation. *Czechoslovak Journal of Physics*, 50(1):115–122, 2000.
- [10] Jiang-Hua Lu, Min Yan, Yong-Chang Zhu, et al. On the set-theoretical Yang-Baxter equation. *Duke Mathematical Journal*, 104(1):1–18, 2000.
- [11] Jiang-Hua Lu, Min Yan, and Yongchang Zhu. Quasi-triangular structures on Hopf algebras with positive. In *New Trends in Hopf Algebra Theory: Proceedings of the Colloquium on Quantum Groups and Hopf Algebras, La Falda, Sierras de Córdoba, Argentina, August 9-13, 1999*, volume 267, page 339. American Mathematical Soc., 2000.
- [12] Jiang-Hua Lu, Min Yan, and Yongchang Zhu. On Hopf algebras with positive bases. *Journal of Algebra*, 237(2):421–445, 2001.

- [13] Shahn Majid. Tannaka-Krein theorem for quasi-Hopf algebras and other results. In *Deformation theory and quantum groups with applications to mathematical physics*, volume 134, pages 219–232. Contemp. Maths, 1992.
- [14] Shahn Majid. Transmutation theory and rank for quantum braided groups. In *Mathematical proceedings of the Cambridge Philosophical Society*, volume 113, pages 45–70. Cambridge University Press, 1993.
- [15] Shahn Majid. *Foundations of quantum group theory*. Cambridge university press, 2000.
- [16] Agata Smoktunowicz and Leandro Vendramin. On skew braces (with an appendix by N. Byott and L. Vendramin). *Journal of Combinatorial Algebra*, 2(1):47–86, 2018.
- [17] Mitsuhiro Takeuchi. Matched pairs of groups and bismash products of Hopf algebras. *Communications in Algebra*, 9(8):841–882, 1981.