A new proof for the existence of rotationally symmetric gradient Ricci solitons

Shu-Yu Hsu
Department of Mathematics
National Chung Cheng University
168 University Road, Min-Hsiung
Chia-Yi 621, Taiwan, R.O.C.
e-mail: shuyu.sy@gmail.com

April 6, 2024

Abstract

We give a new proof for the existence of rotationally symmetric steady and expanding gradient Ricci solitons in dimension n+1, $2 \le n \le 4$, with metric $g = \frac{da^2}{h(a^2)} + a^2 d \sigma$ for some function h where $d\sigma$ is the standard metric on the unit sphere S^n in \mathbb{R}^n . More precisely for any $\lambda \ge 0$, $2 \le n \le 4$ and $\mu_1 \in \mathbb{R}$, we prove the existence of unique solution $h \in C^2((0,\infty)) \cap C^1([0,\infty))$ for the equation $2r^2h(r)h_{rr}(r) = (n-1)h(r)(h(r)-1) + rh_r(r)(rh_r(r) - \lambda r - (n-1))$, h(r) > 0, in $(0,\infty)$ satisfying h(0) = 1, $h_r(0) = \mu_1$. We also prove the existence of unique analytic solution of the about equation on $[0,\infty)$ for any $\lambda \ge 0$, $n \ge 2$ and $\mu_1 \in \mathbb{R}$. Moreover we will prove the asymptotic behaviour of the solution h for any $n \ge 2$, $\lambda \ge 0$ and $\mu_1 \in \mathbb{R} \setminus \{0\}$.

Keywords: singular elliptic equation, rotationally symmetric, Ricci steady solitons, Ricci expanding solitons

AMS 2020 Mathematics Subject Classification: Primary 35J70, 35J75 Secondary 53C21

1 Introduction

Recently there is a lot of study on Ricci solitons by S. Brendle, R.L. Bryant, H.D. Cao, D. Zhou, M. Feldman, T. Ilmanen and D. Knopf, Y. Li and B. Wang, O. Munteanu and N. Sesum, P. Petersen and W. Wylie, [B1], [Br], [C], [CZ], [FIK], [LW], [MS], [PW], etc. Ricci solitons are used extensively in the study of Ricci flow. For example in [B2] S. Brendle used singular rotationally symmetric steady solitons to construct barrier functions which

are used to prove a conjecture of Perelman on 3-dimensional ancient κ solution to the Ricci flow.

A Riemannian metric $g = (g_{ij})$ on a Riemannian manifold M is said to be a gradient Ricci soliton if there exists a smooth function f on M and a constant $\lambda \in \mathbb{R}$ such that the Ricci tensor R_{ij} of the metric g satisfies

$$R_{ij} = \nabla_i \nabla_j f - \lambda g_{ij} \quad \text{on } M. \tag{1.1}$$

The gradient soliton is called an expanding gradient Ricci soliton if $\lambda > 0$. It is called a steady gradient Ricci soliton if $\lambda = 0$ and it is called a shrinking gradient Ricci soliton if $\lambda < 0$.

By an argument similar to R.L. Bryant [Br] for any $n \ge 2$, if (M, g) is a (n+1)-dimensional rotational symmetric gradient Ricci soliton which satisfies (1.1) for some smooth function f and constant $\lambda \in \mathbb{R}$, then

$$g = dt^2 + a(t)^2 d\sigma (1.2)$$

where $d\sigma$ is the standard metric on the unit sphere S^n in \mathbb{R}^n , a(t) satisfies

$$a(t)^{2}a_{t}(t)a_{ttt}(t) = a(t)^{2}a_{tt}(t)^{2} - (n-1)a(t)a_{tt}(t) - \lambda a(t)^{3}a_{tt}(t) + a(t)a_{t}(t)^{2}a_{tt}(t) - (n-1)a_{t}(t)^{2} + (n-1)a_{t}(t)^{4},$$
(1.3)

and f satisfies

$$\begin{cases} a(t)f_{tt}(t) = \lambda a(t) - na_{tt}(t)^2 \\ a(t)a_{tt}(t) = (n-1)(1 - a_t(t)^2) - a(t)a_t(t)f_t(t) + \lambda a(t)^2. \end{cases}$$

As in [Br], by (1.3) if a(0) = 0, one can choose $a_t(0) = 1$. If one writes

$$g = \frac{da^2}{h(a^2)} + a^2 d\sigma \tag{1.4}$$

where h(r), $r = a^2$, and a(t) satisfies

$$a_t(t) = \sqrt{h(a(t)^2)},\tag{1.5}$$

then by (1.3) h satisfies the singular elliptic equation

$$2r^{2}h(r)h_{rr}(r) = (n-1)h(r)(h(r)-1) + rh_{r}(r)(rh_{r}(r)-\lambda r - (n-1)), \quad h(r) > 0.$$
 (1.6)

Note that the radial sectional curvature and the orbital sectional curvature of g is $-h_r(r)$ and (1 - h(r))/r respectively. Hence the existence of gradient Ricci soliton is equivalent to the existence of solution of (1.6) and the positivity of the sectional curvatures of g is equivalent to the condition that $h_r(r) < 0$ and 0 < h(r) < 1 holds for any r > 0.

In the paper [Br] R.L. Bryant by using power series method on (1.6) and phase plane analysis on the functions

$$x = a(t)^2$$
, $y = a_t(t)^2$, $z = a(t)a_{tt}(t)$, (1.7)

gave an incomplete proof of the existence of 3-dimensional rotationally symmetric steady and expanding gradient Ricci solitons. In this paper we will use fixed point argument to prove the existence of unique solution $h \in C^2((0,\infty)) \cap C^1([0,\infty))$ of

$$\begin{cases}
2r^{2}h(r)h_{rr}(r) = (n-1)h(r)(h(r)-1) + rh_{r}(r)(rh_{r}(r)-\lambda r - (n-1)), & h(r) > 0, \quad \forall r > 0 \\
h(0) = 1, & h_{r}(0) = \mu_{1}
\end{cases}$$
(1.8)

for any constant $\lambda \ge 0$, $\mu_1 \in \mathbb{R}$ and integer $2 \le n \le 4$. We will also use a modification of the power series method in [Br] to proof the existence of unique analytic solutions of (1.8) for any $n \ge 2$, $\lambda \ge 0$ and $\mu_1 \in \mathbb{R}$.

In [Br] R.L. Bryant by constructing explicit supersolutions of (1.6) proved that when $\lambda > 0$, $0 < \mu_1 < 3\lambda/7$ and n = 2, any solution h of (1.6) which satisfies the initial condition,

$$h(0) = 1$$
 and $h_r(0) = \mu_1$, (1.9)

satisfies $h_r(r) > 0$ for any $r \ge 0$ and $\lim_{r \to \infty} h(r) \in (1, \infty)$ exists. However it is difficult to extend the method of [Br] to the case $n \ge 2$ and $0 < \mu_1 < \lambda/n$. In this paper we will extend this result of [Br] and prove the asymptotic behaviour of the solution h of (1.8) for any $n \ge 2$, $\lambda \ge 0$ and $\mu_1 \in \mathbb{R} \setminus \{0\}$. The main results we obtain in this paper are the following.

Theorem 1.1. Let $\lambda \geq 0$, $\mu_1 \in \mathbb{R}$ and $2 \leq n \leq 4$. There exists a unique solution $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ of (1.8).

Theorem 1.2. Let $\lambda \geq 0$, $\mu_1 \in \mathbb{R}$, $n \geq 2$ and $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ be a solution of (1.8). Then

$$\begin{cases}
h(r) > 1 & and & h_r(r) > 0 & \forall r > 0 & if \mu_1 > 0 \\
0 < h(r) < 1 & and & h_r(r) < 0 & \forall r > 0 & if \mu_1 < 0
\end{cases}$$
(1.10)

Theorem 1.3. Let $\lambda < 0$, $2 \le n \le 4$ and $\mu_1 \in \mathbb{R}$. Then there exists a constant $L \ge -(n-1)/\lambda$ and a unique solution $h \in C^2((0,L)) \cap C^1([0,L))$ of (1.6) in (0, L) which satisfies (1.9).

Theorem 1.4. Let $\lambda \geq 0$, $\mu_1 \in \mathbb{R}$ and $n \geq 2$. Then there exists a unique analytic solution h of (1.8) on $[0, \infty)$.

Theorem 1.5. Let $\mu_1 < 0$ and $n \ge 2$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of

$$\begin{cases}
2r^2h(r)h_{rr}(r) = (n-1)h(r)(h(r)-1) + rh_r(r)(rh_r(r)-(n-1)), & h(r) > 0, \quad \forall r > 0 \\
h(0) = 1, & h_r(0) = \mu_1.
\end{cases}$$
(1.11)

Then

$$\lim_{r \to \infty} h(r) = 0. \tag{1.12}$$

Theorem 1.6. Let $\mu_1 < 0$ and $n \ge 2$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11). Then

$$\lim_{r \to \infty} \frac{rh_r(r)}{h(r)} = -1. \tag{1.13}$$

Theorem 1.7. Let $\mu_1 < 0$ and $n \ge 2$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11). Then

$$b_1 := \lim_{r \to \infty} rh(r) \in [0, \infty) \quad exists \tag{1.14}$$

with $b_1 > 0$ when $n \ge 4$. Moreover

$$\lim_{r \to \infty} r \left(\frac{r h_r(r)}{h(r)} + 1 \right) = \left(\frac{n-4}{n-1} \right) b_1. \tag{1.15}$$

Theorem 1.8. Let $\lambda > 0$, $\mu_1 \in \mathbb{R} \setminus \{0\}$ and $n \geq 2$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.8). Then the following holds.

(i) If $\mu_1 = \lambda/n$, then

$$h(r) = 1 + \frac{\lambda}{n}r \quad \forall r \ge 0. \tag{1.16}$$

is the explicit analytic solution of (1.8).

(ii) If $0 < \mu_1 < \lambda/n$, then $\lim_{r \to \infty} h_r(r) = 0$ and $\lim_{r \to \infty} h(r) \in (1, \infty)$ exists and

$$1 < h(r) \le 1 + \mu_1 r \quad \forall r > 0. \tag{1.17}$$

(iii) If $\mu_1 > \lambda/n$, then

$$\lim_{r \to \infty} h_r(r) = \lim_{r \to \infty} h(r) = \infty \tag{1.18}$$

and

$$h(r) \ge 1 + \mu_1 r \quad \forall r > 0.$$
 (1.19)

(iv) If $\mu_1 < 0$, then $\lim_{r \to \infty} h_r(r) = 0$ and $\lim_{r \to \infty} h(r) \in (0, 1)$.

Theorem 1.9. Let $n \ge 2$ and $\mu_1 > 0$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11). Then h satisfies (1.18) and (1.19).

The plan of the paper is as follows. In section two we will use fixed point technique to prove the existence of unique solution of (1.8). In section three we will use a modification of the power series method of [Br] to prove Theorem 1.4. In section four we will use integral representation of h and construction of appropriate auxilliary functions to prove the asymptotic behaviour of the solution of (1.11). In section five we will prove Theorem 1.8 and Theorem 1.9.

2 Existence of rotationally symmetric gradient Ricci soliton

In this section we will use fixed point argument to prove the existence of solution of (1.8). We first start with the local existence of solution of (1.6), (1.9), near the origin.

Lemma 2.1. Let λ , $\mu_1 \in \mathbb{R}$ and $2 \le n \le 4$. Then there exists a constant $\varepsilon > 0$ such that (1.6) has a unique solution $h \in C^2((0, \varepsilon)) \cap C^1([0, \varepsilon))$ in $(0, \varepsilon)$ which satisfies (1.9).

Proof: We first observe that if $h \in C^2((0, \varepsilon)) \cap C^1([0, \varepsilon))$ is a solution of (1.6) in (0, ε) for some constant $\varepsilon > 0$ which satisfies (1.9), then

$$2r^{2}h(r)h_{rr}(r) = (n-1)(h(r)-1)^{2} + r^{2}\left\{h_{r}(r)^{2} - \lambda h_{r}(r) + \frac{n-1}{r}\left(\frac{h(r)-1}{r} - h_{r}(r)\right)\right\}$$

$$\Leftrightarrow h_{rr}(r) = \frac{1}{2h(r)}\left\{\frac{n-1}{r^{2}}\left(\int_{0}^{r}h_{r}(\rho)\,d\rho\right)^{2} + h_{r}(r)^{2} - \lambda h_{r}(r) + \frac{n-1}{r^{2}}\int_{0}^{r}(h_{r}(\rho) - h_{r}(r))\,d\rho\right\}$$

$$\Leftrightarrow h_{r}(r) = \int_{0}^{r}\frac{1}{2h(s)}\left\{\frac{n-1}{s^{2}}\left(\int_{0}^{s}h_{r}(\rho)\,d\rho\right)^{2} + \frac{n-1}{s^{2}}\int_{0}^{s}(h_{r}(\rho) - h_{r}(s))\,d\rho + h_{r}(s)^{2} - \lambda h_{r}(s)\right\}\,ds$$

$$+ \mu_{1}$$

for any $0 < r < \varepsilon$. This suggests one to use fixed point argument to prove the local existence of solution of (1.8).

For any $\varepsilon > 0$ we now define the Banach space

$$\mathcal{X}_{\varepsilon} := \left\{ (h, w) : h \in C^{0,1}([0, \varepsilon]; \mathbb{R}), w \in C^{0,1}([0, \varepsilon]; \mathbb{R}) \right\}$$

with a norm given by

$$||(h, w)||_{X_{\varepsilon}} = \max(||h||_{C^{0,1}([0,\varepsilon])}, ||w||_{C^{0,1}([0,\varepsilon])})$$

where

$$||w||_{C^{0,1}([0,\varepsilon])} = \max(||w||_{L^{\infty}([0,\varepsilon])}, ||w||_{1,[0,\varepsilon]}) \quad \text{ and } \quad ||w||_{1,[0,\varepsilon]} = \sup_{s,s' \in [0,\varepsilon], s \neq s'} \frac{|w(s) - w(s')|}{|s - s'|}.$$

For any $(h, w) \in X_{\varepsilon}$, we define

$$\Phi(h, w) := (\Phi_1(h, w), \Phi_2(h, w)),$$

where

$$\begin{cases}
\Phi_{1}(h, w)(r) = 1 + \int_{0}^{r} w(\rho) d\rho, \\
\Phi_{2}(h, w)(r) = \int_{0}^{r} \frac{1}{2h(s)} \left\{ (n-1) \left(\frac{\left(\int_{0}^{s} w(\rho) d\rho \right)^{2}}{s^{2}} + \frac{\int_{0}^{s} (w(\rho) - w(s)) d\rho}{s^{2}} \right) + w(s)^{2} - \lambda w(s) \right\} ds \\
+ \mu_{1}
\end{cases}$$
(2.2)

for any $0 < r \le \varepsilon$. Let $\mathcal{D}_{\varepsilon}$ be the family of all $(h, w) \in \mathcal{X}_{\varepsilon}$ satisfying

$$\begin{cases}
||h - 1||_{L^{\infty}([0,\varepsilon])} \leq (|\mu_{1}| + 1)\varepsilon \\
||h||_{1,[0,\varepsilon]} \leq |\mu_{1}| + 1 \\
||w - \mu_{1}||_{L^{\infty}([0,\varepsilon])} \leq 3\left(n(|\mu_{1}| + 1)^{2} + |\lambda|(|\mu_{1}| + 1)\right)\varepsilon \\
||w||_{1,[0,\varepsilon]} \leq 3\left(n(|\mu_{1}| + 1)^{2} + |\lambda|(|\mu_{1}| + 1)\right) \\
h(0) = 1, \quad w(0) = \mu_{1}.
\end{cases} (2.3)$$

Note that $\mathcal{D}_{\varepsilon}$ is a closed subspace of X_{ε} . Since $(1, \mu_1) \in \mathcal{D}_{\varepsilon}$, $\mathcal{D}_{\varepsilon} \neq \phi$. Let

$$\varepsilon_1 = \frac{1}{100 \left(n(|\mu_1| + 1)^2 + |\lambda|(|\mu_1| + 1) \right)}$$

We will assume that $\varepsilon \in (0, \varepsilon_1)$ for the rest of the proof. We will show that if $\varepsilon \in (0, \varepsilon_1)$ is sufficiently small, the map $(h, w) \mapsto \Phi(h, w)$ will have a unique fixed point in $\mathcal{D}_{\varepsilon}$. We first prove that $\Phi(\mathcal{D}_{\varepsilon}) \subset \mathcal{D}_{\varepsilon}$ for any $\varepsilon \in (0, \varepsilon_1)$. In fact for any $\varepsilon \in (0, \varepsilon_1)$ and $(h, w) \in \mathcal{D}_{\varepsilon}$, by (2.3) we have

$$\frac{99}{100} \le h(r) \le \frac{101}{100} \quad \forall 0 \le r \le \varepsilon \quad \text{and} \quad ||w||_{L^{\infty}([0,\varepsilon])} \le |\mu_1| + \frac{3}{100} \le |\mu_1| + 1. \tag{2.4}$$

Hence by (2.2), (2.3) and (2.4), we have

$$|\Phi_{1}(h, w)(r_{2}) - \Phi_{1}(h, w)(r_{1})| \leq ||w||_{L^{\infty}([0, \varepsilon])}|r_{2} - r_{1}| \quad \forall 0 \leq r_{1} < r_{2} \leq \varepsilon$$

$$\Rightarrow \quad ||\Phi_{1}(h, w)||_{1, [0, \varepsilon]} \leq ||w||_{L^{\infty}([0, \varepsilon])} \leq |\mu_{1}| + 1 \tag{2.5}$$

and

$$\|\Phi_1(h, w) - 1\|_{L^{\infty}([0,\varepsilon])} \le \|w\|_{L^{\infty}([0,\varepsilon))} \varepsilon \le (|\mu_1| + 1)\varepsilon \tag{2.6}$$

and

$$\begin{split} &|\Phi_{2}(h,w)(r_{2}) - \Phi_{2}(h,w)(r_{1})|\\ \leq &\frac{100}{2 \cdot 99} \int_{r_{1}}^{r_{2}} \left(n||w||_{L^{\infty}([0,\varepsilon])}^{2} + |\lambda|||w||_{L^{\infty}([0,\varepsilon])} + \frac{(n-1)||w||_{1,[0,\varepsilon]}}{s^{2}} \int_{0}^{s} (s-\rho) \, d\rho \right) ds\\ \leq &\frac{50}{99} \left(n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1) + \frac{3(n-1)}{2} \left(n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1) \right) \right) |r_{2} - r_{1}|\\ \leq &\frac{25}{9} \left(n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1) \right) |r_{2} - r_{1}|\\ \leq &3 \left(n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1) \right) |r_{2} - r_{1}| \quad \forall 0 \leq r_{1} < r_{2} \leq \varepsilon \end{split}$$

Thus

$$\|\Phi_2(h, w)\|_{1, [0, \varepsilon]} \le 3\left(n(|\mu_1| + 1)^2 + |\lambda|(|\mu_1| + 1)\right) \tag{2.7}$$

and

$$|\Phi_2(h, w)(r) - \mu_1| \le 3\left(n(|\mu_1| + 1)^2 + |\lambda|(|\mu_1| + 1)\right)\varepsilon \quad \forall 0 \le r \le \varepsilon.$$
 (2.8)

By (2.5), (2.6), (2.7) and (2.8) we get that $\Phi(\mathcal{D}_{\varepsilon}) \subset \mathcal{D}_{\varepsilon}$ for any $\varepsilon \in (0, \varepsilon_1)$. Let $(h_1, w_1), (h_2, w_2) \in \mathcal{D}_{\varepsilon}, 0 < \varepsilon < \varepsilon_1, \delta_1 = \|(h_1, w_1) - (h_2, w_2)\|_{X_{\varepsilon}}$ and

$$E(w,s) = \frac{n-1}{s^2} \left(\int_0^s w(\rho) \, d\rho \right)^2 + \frac{n-1}{s^2} \int_0^s (w(\rho) - w(s)) \, d\rho + w(s)^2 - \lambda w(s)$$
 (2.9)

for any $w \in C^{0,1}([0, \varepsilon]; \mathbb{R}), 0 < s < \varepsilon$. Then by (2.3),

$$\frac{99}{100} \le h_i(r) \le \frac{101}{100} \quad \text{and} \quad ||w_i||_{L^{\infty}([0,\varepsilon])} \le |\mu_1| + 1 \quad \forall 0 \le r \le \varepsilon, i = 1, 2.$$
 (2.10)

and since $w_1(0) = w_2(0) = \mu_1$,

$$\begin{aligned}
&|E(w_{1},s) - E(w_{2},s)| \\
&\leq \frac{n-1}{s^{2}} \left(\int_{0}^{s} |w_{1}(\rho) - w_{2}(\rho)| d\rho \right) \left(\int_{0}^{s} |w_{1}(\rho) + w_{2}(\rho)| d\rho \right) + |w_{1}(s) - w_{2}(s)| |w_{1}(s) + w_{2}(s)| \\
&+ \frac{n-1}{s^{2}} \int_{0}^{s} |(w_{1} - w_{2})(\rho) - (w_{1} - w_{2})(s)| d\rho + |\lambda| |w_{1}(s) - w_{2}(s)| \\
&\leq \left\{ \left(\frac{n-1}{s} \int_{0}^{s} \rho d\rho + s \right) (||w_{1}||_{L^{\infty}([0,\varepsilon])} + ||w_{2}||_{L^{\infty}([0,\varepsilon])}) + \frac{n-1}{s^{2}} \int_{0}^{s} (s-\rho) d\rho + |\lambda| s \right\} ||w_{1} - w_{2}||_{1,[0,\varepsilon]} \\
&\leq \left(((n+1)(|\mu_{1}| + 1) + |\lambda|)s + \frac{n-1}{2} \right) ||w_{1} - w_{2}||_{1,[0,\varepsilon]} \quad \forall 0 < s \leq \varepsilon.
\end{aligned} \tag{2.11}$$

Similarly,

$$|E(w_{2},s)| \leq n||w_{2}||_{L^{\infty}([0,\varepsilon])}^{2} + |\lambda|||w_{2}||_{L^{\infty}([0,\varepsilon])} + \frac{n-1}{2}||w_{2}||_{1,[0,\varepsilon]}$$

$$\leq n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1) + \frac{3(n-1)}{2}\left(n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1)\right)$$

$$\leq \frac{11}{2}\left(n(|\mu_{1}|+1)^{2} + |\lambda|(|\mu_{1}|+1)\right) =: C_{1} \quad (\text{say}) \quad \forall 0 < s \leq \varepsilon. \tag{2.12}$$

Hence by (2.2), (2.3), (2.10), (2.11) and (2.12),

$$\begin{split} &|(\Phi_{2}(h_{1},w_{1})-\Phi_{2}(h_{2},w_{2}))(r_{2})-(\Phi_{2}(h_{1},w_{1})-\Phi_{2}(h_{2},w_{2}))(r_{1})|\\ &=\frac{1}{2}\left|\int_{r_{1}}^{r_{2}}\frac{E(w_{1},s)}{h_{1}(s)}\,ds-\int_{r_{1}}^{r_{2}}\frac{E(w_{2},s)}{h_{2}(s)}\,ds\right|\\ &\leq\int_{r_{1}}^{r_{2}}\frac{|E(w_{1},s)-E(w_{2},s)|}{2h_{1}(s)}\,ds+\int_{r_{1}}^{r_{2}}\frac{|E(w_{2},s)||h_{2}(s)-h_{1}(s)|}{2h_{1}(s)h_{2}(s)}\,ds\\ &\leq\frac{100}{2\cdot 99}\left(\int_{r_{1}}^{r_{2}}\left(((n+1)(|\mu_{1}|+1)+|\lambda|)s+\frac{n-1}{2}\right)ds\right)||w_{1}-w_{2}||_{1,[0,\varepsilon]}\\ &+\frac{1}{2}\left(\frac{100}{99}\right)^{2}C_{1}\left(\int_{r_{1}}^{r_{2}}s\,ds\right)||h_{1}-h_{2}||_{1,[0,\varepsilon]}\\ &\leq\left\{\frac{25}{99}\left[((n+1)(|\mu_{1}|+1)+|\lambda|)(r_{2}+r_{1})+n-1\right]+\frac{2500C_{1}}{99^{2}}(r_{2}+r_{1})\right\}|r_{2}-r_{1}|\delta_{1}\\ &\leq\left(C_{2}\varepsilon+\frac{25}{33}\right)|r_{2}-r_{1}|\delta_{1}\quad\forall 0\leq r_{1}< r_{2}\leq\varepsilon\end{aligned}$$

where

$$C_2 = (n+1)(|\mu_1|+1) + |\lambda| + \frac{5000}{99^2}C_1.$$

Thus

$$\|(\Phi_2(h_1, w_1) - \Phi_2(h_2, w_2))\|_{1,[0,\varepsilon]} \le \left(C_2\varepsilon + \frac{25}{33}\right)\delta_1. \tag{2.13}$$

Similarly,

$$\|(\Phi_2(h_1, w_1) - \Phi_2(h_2, w_2))\|_{L^{\infty}([0,\varepsilon])} \le \left(C_2\varepsilon + \frac{25}{33}\right)\varepsilon\delta_1.$$
 (2.14)

Let $\varepsilon_2 = \min\left(\varepsilon_1, \frac{1}{33C_2}\right)$ and $0 < \varepsilon < \varepsilon_2$ for the rest of the proof. Then by (2.13) and (2.14),

$$\|(\Phi_2(h_1, w_1) - \Phi_2(h_2, w_2))\|_{C^{0,1}([0,\varepsilon])} \le \left(C_2\varepsilon + \frac{25}{33}\right)\delta_1 \le \frac{26}{33}\delta_1. \tag{2.15}$$

On the other hand,

$$\begin{split} &|(\Phi_1(h_1,w_1)-\Phi_1(h_2,w_2))(r_2)-(\Phi_2(h_1,w_1)-\Phi_2(h_2,w_2))(r_1)|\\ \leq &\int_{r_1}^{r_2}|w_1(s)-w_2(s)|\,ds\\ \leq &||w_1-w_2||_{1,[0,\varepsilon]}\int_{r_1}^{r_2}s\,ds\\ \leq &\varepsilon\delta_1|r_2-r_1|\quad\forall 0\leq r_1< r_2\leq \varepsilon. \end{split}$$

Hence

$$\|\Phi_1(h_1, w_1) - \Phi_1(h_2, w_2)\|_{1, [0, \varepsilon]} \le \varepsilon \delta_1. \tag{2.16}$$

Similarly,

$$|\Phi_{1}(h_{1}, w_{1})(r) - \Phi_{1}(h_{2}, w_{2})(r)| \leq \varepsilon \delta_{1} r \leq \varepsilon^{2} \delta_{1} \quad \forall 0 \leq r \leq \varepsilon$$

$$\Rightarrow \|\Phi_{1}(h_{1}, w_{1}) - \Phi_{1}(h_{2}, w_{2})\|_{L^{\infty}([0, \varepsilon])} \leq \varepsilon \delta_{1}. \tag{2.17}$$

By (2.16) and (2.17),

$$\|\Phi_1(h_1, w_1) - \Phi_1(h_2, w_2)\|_{C^{0,1}([0,\varepsilon])} \le \varepsilon \delta_1. \tag{2.18}$$

Hence for any $0 < \varepsilon < \varepsilon_2$, by (2.15) and (2.18),

$$\|(\Phi(h_1, w_1) - \Phi(h_2, w_2))\|_{X_{\varepsilon}} \le \frac{26}{33}\delta_1.$$

Thus Φ is a contraction map on $\mathcal{D}_{\varepsilon}$. Hence by the contraction map theorem there exists a unique fixed point $(h, w) = \Phi(h, w)$ in $\mathcal{D}_{\varepsilon}$. Thus

$$\begin{cases} h(r) = 1 + \int_{0}^{r} w(\rho) d\rho, \\ w(r) = \int_{0}^{r} \frac{1}{2h(s)} \left\{ \frac{n-1}{s^{2}} \left(\int_{0}^{s} w(\rho) d\rho \right)^{2} + \frac{n-1}{s^{2}} \int_{0}^{s} (w(\rho) - w(s)) d\rho + w(s)^{2} - \lambda w(s) \right\} ds \\ + \mu_{1} \end{cases}$$
(2.19)

for any $0 < r \le \varepsilon$ with h(0) = 1, $w(0) = \mu_1$, and both h and w are differentiable on $[0, \varepsilon]$ with

$$h_r(r) = w(r) \tag{2.20}$$

and

$$w_r(r) = \frac{1}{2h(r)} \left\{ \frac{n-1}{r^2} \left(\int_0^r w(\rho) \, d\rho \right)^2 + \frac{n-1}{r^2} \int_0^r (w(\rho) - w(r)) \, d\rho + w(r)^2 - \lambda w(s) \right\}. \tag{2.21}$$

Putting (2.20) in (2.21) we get (2.1). Hence for any $0 < \varepsilon < \varepsilon_2$, (1.6) has a unique solution $h \in C^2((0, \varepsilon)) \cap C^1([0, \varepsilon))$ in $(0, \varepsilon)$ which satisfies (1.9) and the lemma follows.

We next observe that by a similar argument as the proof of Lemma 1 of [Br] we have the following result.

Lemma 2.2. (cf. Lemma 1 of [Br]) Let $\lambda \in \mathbb{R}$, $0 \le a < b$ and $n \ge 2$. Let $h \in C^2((a, b))$ be a solution of (1.6) in (a, b). Then the following holds.

- (i) Suppose there exists $x_0 \in (a, b)$ such that $h_r(x_0) \ge 0$ and $h(x_0) > 1$. Then $h_r(x) > 0$ on (x_0, b) .
- (ii) Suppose there exists $x_0 \in (a,b)$ such that $h_r(x_0) \le 0$ and $h(x_0) < 1$. Then $h_r(x) < 0$ on (x_0,b) .

Lemma 2.3. Let $\lambda \in \mathbb{R}$ and $n \ge 2$. Suppose $h \in C^2((0,L))$ satisfies (1.6) in (0, L) for some constant $L \in (0,\infty)$ such that $L < -\frac{(n-1)}{\lambda}$ if $\lambda < 0$. Then there exists a constant $c_1 > 0$ such that

$$h(r) \ge c_1 \quad \forall L/2 \le r \le L. \tag{2.22}$$

Proof: Suppose (2.22) does not hold. Then there exists a sequence $\{r_i\}_{i=1}^{\infty} \subset (L/2, L), r_i < r_{i+1}$ for all $i \in \mathbb{Z}^+$, $r_i \to L$ as $i \to \infty$, such that

$$h(r_i) \to 0 \quad \text{as } i \to \infty.$$
 (2.23)

Without loss of generality we may assume that

$$0 < h(r_{i+1}) < h(r_i) < 1 \quad \forall i \in \mathbb{Z}^+. \tag{2.24}$$

By (1.6) or Lemma 3.1 of [H],

$$(h^{-1/2}h_r)_r = (n-1)\frac{h-1}{2r^2h^{1/2}} - \frac{n-1+\lambda r}{2r} \cdot \frac{h_r}{h^{3/2}} \quad \forall 0 < r < L$$

$$\Rightarrow h_r(r) = \sqrt{h(r)} \left\{ \frac{h_r(r_1)}{\sqrt{h(r_1)}} + (n-1) \int_{r_1}^r \frac{h(\rho)-1}{2\rho^2 \sqrt{h(\rho)}} d\rho - \int_{r_1}^r \frac{((n-1)\rho^{-1}+\lambda)h_r(\rho)}{2h(\rho)^{3/2}} d\rho \right\}$$

$$= \sqrt{h(r)} \left\{ \frac{h_r(r_1)}{\sqrt{h(r_1)}} + (n-1) \int_{r_1}^r \frac{h(\rho)-1}{2\rho^2 \sqrt{h(\rho)}} d\rho + \left(\frac{n-1}{r} + \lambda\right) \frac{1}{\sqrt{h(r)}} - \left(\frac{n-1}{r_1} + \lambda\right) \frac{1}{\sqrt{h(r_1)}} + (n-1) \int_{r_1}^r \frac{d\rho}{\rho^2 \sqrt{h(\rho)}} \right\} \quad \forall r_1 < r < L.$$

Hence

$$h_{r}(r) = \frac{n-1}{r} + \lambda + \sqrt{\frac{h(r)}{h(r_{1})}} \left(h_{r}(r_{1}) - \frac{n-1}{r_{1}} - \lambda \right) + (n-1) \frac{\sqrt{h(r)}}{2} \int_{r_{1}}^{r} \frac{h(\rho) + 1}{\rho^{2} \sqrt{h(\rho)}} d\rho \qquad (2.25)$$

$$> \frac{n-1}{r} + \lambda + \sqrt{\frac{h(r)}{h(r_{1})}} \left(h_{r}(r_{1}) - \frac{n-1}{r_{1}} - \lambda \right) \quad \forall r_{1} < r < L. \qquad (2.26)$$

Putting $r = r_i$ in (2.26) and letting $i \to \infty$,

$$\liminf_{i \to \infty} h_r(r_i) \ge \frac{n-1}{L} + \lambda > 0$$

$$\Rightarrow \exists i_0 \in \mathbb{Z}^+ \text{ such that } h_r(r_i) > 0 \quad \forall i \ge i_0. \tag{2.27}$$

By (2.24) and the mean value thoerem for any $i \in \mathbb{Z}^+$ there exists $\xi_i \in (r_i, r_{I+1})$ such that

$$h_r(\xi_i) = \frac{h(r_{i+1}) - h(r_i)}{r_{i+1} - r_i} < 0.$$

This together with (2.27) implies that for any $i \ge i_0$, there exists a maximal interval (r_i, b_i) , $b_i \in (r_i, \xi_i)$, such that

$$h_r(r) > 0 \quad \forall r_i \le r < b_i \quad \text{and} \quad h_r(b_i) = 0.$$
 (2.28)

We now divide the proof into three cases.

Case 1: $h(b_{i_0}) = 1$.

By (2.28) and uniqueness solution of ODE $h(r) \equiv 1$ on (0, L). This contradicts (2.23). Hence case 1 does not hold.

Case 2: $h(b_{i_0}) > 1$.

Then by (2.28) and Lemma 2.2,

$$h_r(r) > 0 \quad \forall b_{i_0} < r < L \quad \Rightarrow \quad h(r) > h(b_{i_0}) \quad \forall b_{i_0} < r < L$$

which contradicts (2.23). Hence case 2 does not hold.

Case 3: $h(b_{i_0}) < 1$.

Then by (2.28) and Lemma 2.2,

$$h_r(r) < 0 \quad \forall b_{i_0} \le r < L.$$

This contradicts (2.27). Hence case 3 does not hold. By case 1, case 2 and case 3 we get a contradiction. Hence no such sequence $\{r_i\}_{i=1}^{\infty}$ exists. Thus there exists a constant $c_1 > 0$ such that (2.22) holds and the lemma follows.

Lemma 2.4. Let $\lambda \in \mathbb{R}$ and $n \ge 2$. Suppose $h \in C^2((0, L))$ satisfies (1.6) in (0, L) for some constant $L \in (0, \infty)$ such that $L < -(n-1)/\lambda$ if $\lambda < 0$. Then there exists a constant $c_2 > 0$ such that

$$h(r) \le c_2 \quad \forall L/2 \le r \le L. \tag{2.29}$$

Proof: Suppose (2.29) does not hold. Then there exists a sequence $\{r_i\}_{i=1}^{\infty} \subset (L/2, L), r_i < r_{i+1}$ for all $i \in \mathbb{Z}^+$, $r_i \to L$ as $i \to \infty$, such that

$$h(r_i) \to \infty \quad \text{as } i \to \infty.$$
 (2.30)

Without loss of generality we may assume that

$$h(r_{i+1}) > h(r_i) + 1 \quad \forall i \in \mathbb{Z}^+.$$
 (2.31)

By (2.31) and the mean value theorem for any $i \in \mathbb{Z}^+$ there exists $\xi_i \in (r_i, r_{i+1})$ such that

$$h_r(\xi_i) = \frac{h(r_{i+1}) - h(r_{r_i})}{r_{i+1} - r_i} > \frac{1}{r_{i+1} - r_i} > 0 \quad \forall i \in \mathbb{Z}^+$$

$$\Rightarrow h_r(\xi_i) \to \infty \quad \text{as } i \to \infty. \tag{2.32}$$

Hence by (2.32) there exists $i_0 \in \mathbb{Z}^+$ such that

$$h_r(\xi_{i_0}) > \frac{2(n-1)}{L} + \lambda \quad \Rightarrow \quad h_r(\xi_{i_0}) > \frac{n-1}{\xi_{i_0}} + \lambda.$$
 (2.33)

By replacing r_1 by ξ_{i_0} in (2.26) we get

$$h_r(r) > \frac{n-1}{r} + \lambda + \sqrt{\frac{h(r)}{h(\xi_{i_0})}} \left(h_r(\xi_{i_0}) - \frac{n-1}{\xi_{i_0}} - \lambda \right) > \frac{n-1}{r} + \lambda > 0 \quad \forall \xi_{i_0} \le r < L. \quad (2.34)$$

By (2.30) and (2.34),

$$h(r) \to \infty \quad \text{as } r \to L.$$
 (2.35)

Since $L < -(n-1)/\lambda$ if $\lambda < 0$,

$$n - 1 + \lambda r > 0 \quad \forall 0 < r < L. \tag{2.36}$$

By (1.6), (2.34) and (2.36),

$$\left(\frac{h_r}{h}\right)_r = \frac{h_{rr}}{h} - \frac{h_r^2}{h^2} = \frac{1}{2h} \left((n-1)\frac{h-1}{r^2} - \frac{(n-1+\lambda r)h_r}{rh} + \frac{h_r^2}{h}\right) - \frac{h_r^2}{h^2} \le (n-1)\frac{h-1}{2r^2h} \le \frac{2(n-1)}{L^2} \tag{2.37}$$

holds for any $\xi_{i_0} \le r < L$. Integrating (2.37) over $(\xi_{i_0}, r), r \in (\xi_{i_0}, L)$, by (2.34) we have

$$\frac{h_r(r)}{h(r)} \le \frac{h_r(\xi_{i_0})}{h(\xi_{i_0})} + \frac{2(n-1)}{L} =: C_3 \text{ (say)} \quad \forall \xi_{i_0} \le r < L$$

$$\Rightarrow \log\left(\frac{h(r)}{h(\xi_{i_0})}\right) \le C_3 L \quad \forall \xi_{i_0} \le r < L$$

$$\Rightarrow h(r) \le h(\xi_{i_0}) e^{C_3 L} \quad \forall \xi_{i_0} \le r < L.$$

which contradicts (2.35). Hence no such sequence $\{r_i\}_{i=1}^{\infty}$ exists. Thus there exists a constant $c_2 > 0$ such that (2.29) holds and the lemma follows.

Lemma 2.5. Let $\lambda \in \mathbb{R}$ and $n \ge 2$. Suppose $h \in C^2((0,L))$ satisfies (1.6) in (0, L) for some constant $L \in (0,\infty)$ such that $L < -(n-1)/\lambda$ if $\lambda < 0$. Then there exists a constant $c_3 > 0$ such that

$$h_r(r) \le c_3 \quad \forall L/2 \le r \le L.$$
 (2.38)

Proof: Let $c_2 > 0$ be as given by Lemma 2.4. Suppose (2.38) does not hold. Then there exists a sequence $\{r_i\}_{i=1}^{\infty} \subset (L/2, L), r_i < r_{i+1} \text{ for all } i \in \mathbb{Z}^+, r_i \to L \text{ as } i \to \infty$, such that

$$h_r(r_i) \to \infty \quad \text{as } i \to \infty.$$
 (2.39)

Let

$$C_4 = \frac{16}{L} \max \left(c_2 n, \sqrt{c_2 n}, n - 1 + |\lambda| L \right). \tag{2.40}$$

By (2.39) there exists $i_0 \in \mathbb{Z}^+$ such that

$$h_r(r_i) > C_4 \quad \forall i \ge i_0. \tag{2.41}$$

We now rewrite (1.6) as

$$h_{rr}(r) = \frac{h_r(r)^2}{2h(r)} + (n-1)\frac{h(r)-1}{2r^2} - \frac{n-1+\lambda r}{2r} \cdot \frac{h_r(r)}{h(r)}, \quad h(r) > 0.$$
 (2.42)

Then by (2.29), (2.41) and (2.42),

$$h_{rr}(r_{i_0}) = \frac{h_r(r_{i_0})^2}{4h(r_{i_0})} + \frac{1}{2} \left(\frac{h_r(r_{i_0})^2}{4h(r_{i_0})} + (n-1) \frac{h(r_{i_0}) - 1}{r_{i_0}^2} \right) + \frac{1}{2h(r_{i_0})} \left(\frac{h_r(r_{i_0})^2}{4} - \frac{n-1+\lambda r_{i_0}}{r_{i_0}} h_r(r_{i_0}) \right)$$

$$\geq \frac{h_r(r_{i_0})^2}{4h(r_{i_0})} + \frac{1}{2} \left(\frac{h_r(r_{i_0})^2}{4c_2} - \frac{4(n-1)}{L^2} \right) + \frac{1}{2h(r_{i_0})} \left(\frac{h_r(r_{i_0})^2}{4} - \frac{h_r(r_{i_0})^2}{16} - 16 \left(\frac{(n-1+|\lambda|L)}{L} \right)^2 \right)$$

$$\geq \frac{h_r(r_{i_0})^2}{4h(r_{i_0})}.$$
(2.43)

Hence by (2.43) and continuity there exists $0 < \delta < L - r_{i_0}$ such that

$$h_{rr}(r) > \frac{h_r(r)^2}{4h(r)} \quad \forall r_{i_0} \le r < r_{i_0} + \delta.$$
 (2.44)

Let

$$b_0 = \sup \left\{ b \in (r_{i_0}, L) : h_{rr}(r) > \frac{h_r(r)^2}{4h(r)} \quad \forall r_{i_0} \le r < b \right\}.$$

Then $b_0 \ge r_{i_0} + \delta$. Suppose $b_0 < L$. Then by (2.41) and the definition of b_0 ,

$$h_{rr}(r) > \frac{h_r(r)^2}{4h(r)} \quad \forall r_{i_0} \le r < b_0 \quad \text{and} \quad h_{rr}(b_0) = \frac{h_r(b_0)^2}{4h(b_0)}$$
 (2.45)

$$\Rightarrow h_r(b_0) > h_r(r_{i_0}) > C_4.$$
 (2.46)

Then by repeating the above argument but with b_0 and (2.46) replacing r_{i_0} and (2.41) in the proof we get

$$h_{rr}(b_0) > \frac{h_r(b_0)^2}{4h(b_0)}$$

which contradicts (2.45). Hence $b_0 = L$. By (2.41),

$$h_{rr}(r) > \frac{h_r(r)^2}{4h(r)} \quad \forall r_{i_0} \le r < L$$

 $\Rightarrow h_r(r) > h_r(r_{i_0}) > C_4 \quad \forall r_{i_0} \le r < L.$ (2.47)

By (2.29), (2.36), (2.42) and (2.47),

$$\begin{aligned} h_{rr}(r) &\leq \frac{1}{2h(r)} \left(h_r(r)^2 + (n-1) \frac{h(r)^2}{r^2} \right) \leq \frac{1}{2h(r)} \left(h_r(r)^2 + \frac{4(n-1)c_2^2}{L^2} \right) \leq \frac{h_r(r)^2}{h(r)} \quad \forall r_{i_0} \leq r < L \\ \Rightarrow \quad \frac{h_{rr}(r)}{h_r(r)} \leq \frac{h_r(r)}{h(r)} \quad \forall r_{i_0} \leq r < L \\ \Rightarrow \quad h_r(r) \leq \frac{h_r(r_{i_0})}{h(r_{i_0})} h(r) \leq c_2 \frac{h_r(r_{i_0})}{h(r_{i_0})} \quad \forall r_{i_0} \leq r < L \end{aligned}$$

which contradicts (2.39). Hence no such sequence $\{r_i\}_{i=1}^{\infty}$ exists. Thus there exists a constant $c_3 > 0$ such that (2.38) holds and the lemma follows.

Lemma 2.6. Let $\lambda \in \mathbb{R}$ and $n \ge 2$. Suppose $h \in C^2((0, L))$ satisfies (1.6) in (0, L) for some constant $L \in (0, \infty)$ such that $L < -(n-1)/\lambda$ if $\lambda < 0$. Then there exists a constant $c_4 \in \mathbb{R}$ such that

$$h_r(r) \ge c_4 \quad \forall L/2 \le r \le L.$$
 (2.48)

Proof: Let $c_1 > 0$, $c_2 > 0$, $c_3 > 0$, be as given by Lemma 2.3, Lemma 2.4 and Lemma 2.5 respectively. Suppose (2.48) does not hold. Then there exists a sequence $\{r_i\}_{i=1}^{\infty} \subset (L/2, L)$, $r_i < r_{i+1}$ for all $i \in \mathbb{Z}^+$, $r_i \to L$ as $i \to \infty$, such that

$$h_r(r_i) < 0 \quad \forall i \in \mathbb{Z}^+ \quad \text{and} \quad h_r(r_i) \to -\infty \quad \text{as } i \to \infty.$$
 (2.49)

Let C_4 be given by (2.40). By (2.49) there exists $i_0 \in \mathbb{Z}^+$ such that

$$h_r(r_i) < -C_4 \quad \forall i \ge i_0. \tag{2.50}$$

Then by (2.50) and an argument similar to the proof of Lemma 2.5 we get that (2.43) holds. Hence by (2.43) and continuity there exists $0 < \delta < L - r_{i_0}$ such that (2.44) holds. Then by (2.44),

$$h_{rr}(r) > 0 \quad \forall r_{i_0} \le r < r_{i_0} + \delta \quad \Rightarrow \quad h_r(r) > h_r(r_{i_0}) \quad \forall r_{i_0} < r < r_{i_0} + \delta.$$

Let

$$b_2 = \sup \{b \in (r_{i_0}, L) : h_r(r) > h_r(r_{i_0}) \quad \forall r_{i_0} \le r < b\}.$$

Then $b_2 \ge r_{i_0} + \delta$. Suppose $b_2 < L$. Then

$$h_r(r) > h_r(r_{i_0}) \quad \forall r_{i_0} \le r < b \quad \text{and} \quad h_r(b_2) = h_r(r_{i_0}) < -C_4$$
 (2.51)

$$\Rightarrow h_{rr}(b_2) \le 0. \tag{2.52}$$

On the other hand by (2.51) and an argument similar to the proof of Lemma 2.5 we have

$$h_{rr}(b_2) > \frac{h_r(b_2)^2}{4h(b_2)}$$

which contradicts (2.52). Hence $b_2 = L$. Thus

$$h_r(r) > h_r(r_{i_0}) \quad \forall r_{i_0} \le r < L$$

which contradicts (2.49). Hence no such sequence $\{r_i\}_{i=1}^{\infty}$ exists. Thus there exists a constant $c_4 > 0$ such that (2.48) holds and the lemma follows.

We next observe that by standard ODE theory we have the following result.

Lemma 2.7. Let $\lambda \geq 0$, $\mu_1 \in \mathbb{R}$, L > 0, $n \geq 2$, $b_1 \in (c_1, c_2)$, $b_2 \in (c_4, c_3)$ for some constants $c_2 > c_1 > 0$ and $c_3 > c_4$. Then there exists a constant $0 < \delta_1 < L/4$ depending only on c_1, c_2, c_3, c_4 such that for any $r_0 \in (L/2, L)$ (1.6) has a unique solution $\widetilde{h} \in C^2((r_0 - \delta_1, r_0 + \delta_1))$ in $(r_0 - \delta_1, r_0 + \delta_1)$ which satisfies

$$\widetilde{h}(r_0) = b_1$$
 and $\widetilde{h}_r(r_0) = b_2$. (2.53)

We are now ready for the proof of Theorem 1.1.

Proof of Theorem 1.1: Uniqueness of solution of (1.8) follows from the local uniqueness of solution of (1.6) in (0, ε) for some small $\varepsilon > 0$ which satisfies (1.9) and from the uniqueness of solution of ODE of (1.6) on $[\varepsilon/2, \infty)$ with given values of $h(\varepsilon/2)$ and $h_r(\varepsilon/2)$.

We next observe that by Lemma 2.1 there exists a constant $\varepsilon > 0$ such that (1.6) has a unique solution $h \in C^2((0,\varepsilon)) \cap C^1([0,\varepsilon))$ in $(0,\varepsilon)$ which satisfies (1.9). Let [0,L) be the maximal interval of existence of solution $h \in C^2((0,L)) \cap C^1([0,L))$ of (1.6) in (0,L) which satisfies (1.9). Suppose $L < \infty$. Then by Lemma 2.3, Lemma 2.4, Lemma 2.5 and Lemma 2.6 there exists constants $c_2 > c_1 > 0$ and $c_3 > c_4$ such that (2.22), (2.29), (2.38) and (2.48) hold.

Then by Lemma 2.7 there exists a constant $0 < \delta_1 < L$ depending only on c_1, c_2, c_3, c_4 such that for any $r_0 \in (L/2, L)$ (1.6) has a unique solution $\widetilde{h} \in C^2((r_0 - \delta_1, r_0 + \delta_1))$ in $(r_0 - \delta_1, r_0 + \delta_1)$ which satisfies (2.53) with $b_1 = h(r_0)$ and $b_1 = h_r(r_0)$. We now set $r_0 = L - (\delta_1/2)$ and extend h to a function on $[0, L + (\delta_1/2))$ by setting $h(r) = \widetilde{h}(r)$ for any $r \in [L, L + (\delta_1/2))$. Then $h \in C^2((0, L + (\delta_1/2))) \cap C^1([0, L + (\delta_1/2)))$ is a solution of (1.6) in $(0, L + \delta_1)$ which satisfies (1.9). This contradicts the choice of L. Hence $L = \infty$ and there exists a unique solution $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ of (1.8).

By Theorem 1.1 and Lemma 2.2 we get Theorem 1.2. By Lemma 2.3, Lemma 2.4, Lemma 2.5 and Lemma 2.6 and a proof similar to the proof of Theorem 1.1 we get Theorem 1.3. By scaling and the uniqueness result of Theorem 1.1 we have the following corollary.

Corollary 2.8. Let $2 \le n \le 4$. For any $\mu_1 \in \mathbb{R}$ let $h(r; \mu_1) \in C^2((0, \infty)) \cap C^1([0, \infty))$ be the unique solution of (1.11). Then

$$h(\mu r; \mu_1) = h(r; \mu \mu_1) \quad \forall \mu > 0, \mu_1 \in \mathbb{R}.$$

3 Existence of analytic solution

In this section we will use a modification of the power series method of [Br] to proof the existence of unique analytic solutions of (1.8) for any $n \ge 2$, $\lambda \ge 0$ and $\mu_1 \in \mathbb{R}$.

Lemma 3.1. Let λ , $\mu_1 \in \mathbb{R}$ and $n \geq 2$. Then there exists a constant $\varepsilon > 0$ such that (1.6) has a unique analytic solution h in $[0, \varepsilon)$ which satisfies (1.9).

Proof: Since local analytic solution of (1.6), (1.9), about r = 0 is unique, we only need to proof existence of local analytic solution of (1.6), (1.9), about r = 0. We will use a modification of the proof of Proposition 1 of [Br] to proof this lemma. Since the proof is similar to the proof of Proposition 1 of [Br] we will only sketch the argument here. Suppose the solution h of (1.6), (1.9), near the origin is of the form,

$$h(r) = \sum_{i=0}^{\infty} c_i r^i \tag{3.1}$$

for some constants c_1, c_2, \ldots Substituting (3.1) into (1.6), (1.9), and equating the coefficients of r^i , $i \in \mathbb{Z}^+$, we get that (3.1) satisfies (1.6) and (1.9) if and only if

$$c_0 = 1, \quad c_1 = \mu_1, \tag{3.2}$$

and

$$(k-1)(2k+n-1)c_k + \lambda(k-1)c_{k-1} + \sum_{1 \le j \le k-1} (3j^2 - (2+k)j - (n-1))c_j c_{k-j} = 0 \quad \forall k \ge 2. \quad (3.3)$$

Let c_k , $k \ge 2$, be given uniquely in terms of c_1, \ldots, c_{k-1} , by the recurrence relation (3.3). Note that

$$|3j^{2} - (2+k)j - (n-1)| \le (3j-2)j + (2k+n-1)j \qquad \forall 1 \le j \le k-1, k \ge 2$$

$$\le (3(k-1)-2)(k-1) + (2k+n-1)(k-1) \qquad \forall 1 \le j \le k-1, k \ge 2$$

$$\le \frac{3}{2}(2k+n-1)(k-1) + (2k+n-1)(k-1) \qquad \forall 1 \le j \le k-1, k \ge 2$$

$$\le \frac{5}{2}(2k+n-1)(k-1) \qquad \forall 1 \le j \le k-1, k \ge 2.$$

$$(3.4)$$

Hence by (3.3) and (3.4),

$$(k-1)(2k+n-1)|c_k| \le |\lambda|(k-1)|c_{k-1}| + \frac{5}{2}(2k+n-1)(k-1)\sum_{1 \le j \le k-1} |c_j||c_{k-j}| \quad \forall k \ge 2$$

$$\Rightarrow |c_k| \le |\lambda| |c_{k-1}| + \frac{5}{2} \sum_{1 \le j \le k-1} |c_j| |c_{k-j}| \quad \forall k \ge 2.$$
 (3.5)

Let

$$b(r) = \sum_{i=1}^{\infty} c_i' r^i \tag{3.6}$$

where

$$c'_1 = |\mu_1|$$
 and $c'_k = |\lambda|c'_{k-1} + \frac{5}{2} \sum_{1 \le j \le k-1} c'_j c'_{k-j} \quad \forall k \ge 2.$ (3.7)

By (3.2), (3.5), (3.7) and an induction argument on the sequences $\{c_k\}_{k=1}^{\infty}$, $\{c_k'\}_{k=1}^{\infty}$, we get,

$$|c_k| \le c_k' \quad \forall k \ge 1. \tag{3.8}$$

Hence the series (3.1) will be convergent if the series (3.6) is convergent. Now by (3.7) b(r) satisfies

$$b(r) = |\lambda| r b(r) + \frac{5}{2} b(r)^2 + c_1' r$$

$$\Leftrightarrow \frac{5}{2} b(r)^2 - (1 - |\lambda| r) b(r) + c_1' r = 0.$$
(3.9)

Note that (3.9) has a explicit solution of the form,

$$b(r) = \frac{1 - |\lambda|r - \sqrt{(1 - |\lambda|r)^2 - 10c_1'r}}{5} \quad \forall |r| \le \varepsilon$$
 (3.10)

where

$$\varepsilon = \min\left(\frac{1}{10(|\lambda|+1)}, \frac{1}{200(c_1'+1)}\right)$$

and the function given by (3.10) has a convergent Taylor series expansion on $[0, \varepsilon)$ with radius of convergence about r = 0 greater than or equal to ε . By (3.8) the series (3.1) for h also has radius of convergence about r = 0 greater than or equal to ε . Hence the lemma follows.

By Lemma 3.1 and an argument similar to the proof of Theorem 1.1 Theorem 1.4 follows. By scaling and the uniqueness result of Theorem 1.4 we have the following corollary.

Corollary 3.2. Let $n \ge 2$. For any $\mu_1 \in \mathbb{R}$ let $h(r; \mu_1)$ be the unique analytic solution of (1.11) on $[0, \infty)$. Then

$$h(\mu r; \mu_1) = h(r; \mu \mu_1) \quad \forall \mu > 0, \mu_1 \in \mathbb{R}.$$

4 Asymptotic behaviour of rotationally symmetric steady gradient Ricci soliton

In this section we will prove the asymptotic behaviour of the solution h of (1.11) as $r \to \infty$. We will assume that $\lambda = 0$, $\mu_1 < 0$, $n \ge 2$ and $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11) in this section. By Theorem 1.2 (1.10) holds. Hence

$$c_0 = \lim_{r \to \infty} h(r) \in [0, 1)$$
 exists (4.1)

and

$$c_0 < h(r) < 1 \quad \forall r > 0.$$
 (4.2)

We will prove that $c_0 = 0$. We now let

$$q(r) = \frac{rh_r(r)}{h(r)} \quad \forall r \ge 0. \tag{4.3}$$

Then by (1.10),

$$q(r) < 0 \quad \forall r > 0. \tag{4.4}$$

Note that by (1.11) and a direct computation q satisfies

$$q_r(r) = \frac{q(r)}{r} \left(1 - \frac{n-1}{2h(r)} \right) - \left(\frac{q(r)^2}{2r} + (n-1) \frac{1 - h(r)}{2rh(r)} \right) = -\frac{H(q(r), r)}{2r} \quad \forall r > 0$$
 (4.5)

where

$$H(b,r) = \frac{-4nh(r)^2 + 8(n-1)h(r) - (n-1)^2}{4h(r)^2} + \left(b + \frac{n-1-2h(r)}{2h(r)}\right)^2 \quad \forall r > 0, b \in \mathbb{R}.$$
 (4.6)

Lemma 4.1. Let $\mu_1 < 0$ and $n \ge 2$. Let $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ be a solution of (1.11) and c_0 , q, be given by (4.1) and (4.3) respectively. If there exist constants $r_0 > 0$ and C > 0 such that

$$q(r) \le -C \quad \forall r \ge r_0, \tag{4.7}$$

then $c_0 = 0$.

Proof: By (4.7),

$$\frac{rh_r(r)}{h(r)} \le -C \quad \forall r \ge r_0 \quad \Rightarrow \quad \log(h(r)/h(r_0)) \le -C\log(r/r_0) \quad \forall r \ge r_0$$

$$\Rightarrow \quad 0 \le c_0 < h(r) \le \frac{r_0^C h(r_0)}{r^C} \quad \forall r \ge r_0$$

$$\Rightarrow \quad 0 \le c_0 < h(r) \to 0 \quad \text{as } r \to \infty$$

and the lemma follows.

We are now ready to prove Theorem 1.5.

Proof of Theorem 1.5: Suppose

$$0 < c_0 < 1. (4.8)$$

Then by (4.1),

$$\lim_{\substack{r \to \infty \\ b \to 0}} H(b,r) = \frac{-4nc_0^2 + 8(n-1)c_0 - (n-1)^2}{4c_0^2} + \frac{(n-1-2c_0)^2}{4c_0^2} = \frac{(n-1)c_0(1-c_0)}{c_0^2} =: C_5 > 0 \text{ (say)}.$$
(4.9)

By (4.9) there exists $r'_0 > 0$ and $b_0 > 0$ such that

$$C_5/2 < H(b,r) < 3C_5/2 \quad \forall r \ge r'_0, |b| \le b_0.$$
 (4.10)

We now claim that there exist constants C > 0 and $r_0 > 0$ such that (4.7) holds. Suppose the claim does not hold. Then there exists a sequence $\{r_i\}_{i=1}^{\infty} \subset (r'_0, \infty)$, $r_i < r_{i+1}$ for any $i \in \mathbb{Z}^+$ and $r_i \to \infty$ as $i \to \infty$, such that

$$q(r_i) \to 0$$
 as $i \to \infty$.

Without loss of generality we may assume that

$$0 > q(r_i) > -b_0 \quad \forall i \in \mathbb{Z}^+. \tag{4.11}$$

Suppose there exists $i_0 \in \mathbb{Z}^+$ and $s_0 \in (r_{i_0}, r_{i_0+1})$ such that $q(s_0) < -b_0$. Let

$$s_1 = \sup\{s' > s_0 : q(s) < -b_0 \quad \forall s_0 \le s < s'\}.$$
 (4.12)

By (4.11) and (4.12), $s_0 < s_1 < r_{i_0+1}$,

$$q(s) < -b_0 \quad \forall s_0 \le s < s_1 \quad \text{and} \quad q(s_1) = -b_0$$
 (4.13)

$$\Rightarrow q_r(s_1) \ge 0. \tag{4.14}$$

On the other hand by (4.5), (4.10) and (4.13),

$$q_r(s_1) = -\frac{H(q_r(s_1), s_1)}{2s_1} \le -\frac{C_5}{4s_1} < 0$$

which contradicts (4.14). Thus no such constants i_0 and s_0 exist and

$$0 > q(r) \ge -b_0 \quad \forall r \ge r_1. \tag{4.15}$$

Then by (4.5), (4.10) and (4.15),

$$q_r(r) = -\frac{H(q_r(r), r)}{2r} \le -\frac{C_5}{4r} < 0 \quad \forall r \ge r_1$$

$$\Rightarrow q(r) \le q(r_1) \quad \forall r \ge r_1.$$

This contradicts the assumption that there do not exist any constants C > 0 and $r_0 > 0$ such that (4.7) holds. Hence there exist constants C > 0 and $r_0 > 0$ such that (4.7) holds. Then by Lemma 4.1 $c_0 = 0$ and the theorem follows.

We now let

$$u(r) = rh(r). (4.16)$$

and

$$p(r) = rh_r(r) \quad \forall r \ge 0. \tag{4.17}$$

Then by (1.10),

$$p(r) < 0 \quad \forall r > 0 \quad \Rightarrow \quad u_r(r) = rh_r(r) + h(r) = p(r) + h(r) < h(r) < 1 \quad \forall r > 0.$$
 (4.18)

By (1.11) and a direct computation p satisfies

$$p_r(r) = \frac{(p(r) + h(r))[p(r) - (n-1)(1 - h(r))] - (n-2)h(r)p(r)}{2u(r)} \quad \forall r > 0.$$
 (4.19)

By (4.18), (4.19) and a direct computation u satisfies

$$u_{rr}(r) = \frac{-u_r(r)(n-1-2h(r)-u_r(r)) + (n-4)h(r)^2}{2rh(r)} \quad \forall r > 0.$$
 (4.20)

Lemma 4.2. Let $\mu_1 < 0$ and $n \ge 2$. Let $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ be a solution of (1.11) and u be given by (4.16). Then for any $n \ge 4$,

$$u_r(r) > 0 \quad \forall r > 0. \tag{4.21}$$

When $2 \le n < 4$, then either (4.21) holds or there exists a constant $s_0 > 0$ such that

$$\begin{cases} u_r(r) > 0 & \forall 0 < r < s_0 \\ u_r(s_0) = 0 \\ u_r(r) < 0 & \forall r > s_0 \end{cases}$$
 (4.22)

holds. Moreover for any $n \ge 2$ when (4.21) holds,

$$\lim_{r \to \infty} u_r(r) = 0. \tag{4.23}$$

Proof: Since $u_r(0) = h(0) = 1$, there exists a constant $\delta_1 > 0$ such that $u_r(r) > 0$ for any $0 \le r < \delta_1$. Let

$$s_0 = \sup\{s' > 0 : u_r(r) > 0 \quad \forall 0 \le r < s'\}.$$

Then $s_0 \ge \delta_1$. If $s_0 = \infty$, then (4.21) holds. Suppose $s_0 < \infty$. Then

$$u_r(r) > 0 \quad \forall 0 < r < s_0 \quad \text{and} \quad u_r(s_0) = 0$$

 $\Rightarrow u_{rr}(s_0) \le 0.$ (4.24)

We now divide the proof into three cases.

Case 1: n > 4.

By (4.20),

$$u_{rr}(s_0) = \frac{(n-4)h(s_0)}{2s_0} > 0. {(4.25)}$$

which contradicts (4.24). Hence $s_0 = \infty$ and (4.21) holds.

Case 2: n = 4.

By (4.20) both the functions u_r and

$$\widetilde{w}(r) = 0 \quad \forall r \ge 0$$

satisfy

$$\widetilde{w}_r(r) = -\frac{\widetilde{w}(r)(n-1-2h(r)-\widetilde{w}(r))}{2rh(r)} \quad \forall r > 0.$$

and $\widetilde{w}(s_0) = 0 = u_r(s_0)$. Thus by uniqueness of ODE,

$$u_r(r) = \widetilde{w}(r) = 0 \quad \forall r > 0 \quad \Rightarrow \quad 1 = u_r(0) = \lim_{r \to 0} u_r(r) = 0$$

and contradiction arises. Hence $s_0 = \infty$ and (4.21) holds.

Case 3: $2 \le n < 4$.

By (4.20),

$$u_{rr}(s_0) = \frac{(n-4)h(s_0)}{2s_0} < 0.$$

Then there exists a constant $s_1 > s_0$ such that

$$u_{rr}(r) < 0 \quad \forall s_0 \le r < s_1$$

 $\Rightarrow u_r(r) < u_r(s_0) = 0 \quad \forall s_0 < r < s_1.$

Let

$$s_2 = \sup\{s' > 0 : u_r(r) < 0 \quad \forall s_0 < r < s'\}.$$

Then $s_2 \ge s_1$. If $s_2 < \infty$, then

$$u_r(r) < 0 \quad \forall s_0 < r < s_2 \quad \text{and} \quad u_r(s_2) = 0$$
 (4.26)

$$\Rightarrow u_{rr}(s_2) \ge 0. \tag{4.27}$$

By (4.20) and (4.26),

$$u_{rr}(s_2) = \frac{(n-4)h(s_2)}{2s_2} < 0 \quad \forall 2 \le n < 4$$

which contradicts (4.27). Hence $s_2 = \infty$ and (4.22) holds with $s_0 \in (0, \infty)$. Finally for any $n \ge 2$ when (4.21) holds, by (4.18), (4.21) and Theorem 1.5 we get (4.23) and the lemma follows.

By (1.10), (4.18) and Lemma 4.2 we have the following lemma.

Lemma 4.3. Let $\mu_1 < 0$ and $n \ge 2$. Suppose $h \in C^2((0,\infty)) \cap C^1([0,\infty))$ is a solution of (1.11). Let q be given by (4.3). Suppose (4.21) holds. Then

$$-1 < q(r) < 0 \quad \forall r > 0.$$

Note that by (1.12) there exists a constant $r_0 > 1$ such that

$$0 < h(r) < 1/500 \quad \forall r > r_0. \tag{4.28}$$

Lemma 4.4. Let $\mu_1 < 0$ and $2 \le n \le 4$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11). Then (1.14) holds.

Proof: Let u be given by (4.16). By Lemma 4.2 we can divide the proof into two cases. **Case 1**: (4.21) holds.

By (4.18), (4.20), (4.21) and (4.28),

$$u_{rr}(r) \leq -\frac{u_{r}(r)(n - (101/100) - u_{r}(r))}{2u(r)} \qquad \forall r > r_{0}$$

$$\Rightarrow \frac{u_{rr}}{n - (101/100) - u_{r}(r)} \leq -\frac{u_{r}(r)}{2u(r)} \qquad \forall r > r_{0}$$

$$\Rightarrow \log\left(\frac{n - (101/100) - u_{r}(r_{0})}{n - (101/100) - u_{r}(r)}\right) \leq -\frac{1}{2}\log\left(\frac{u(r)}{u(r_{0})}\right) \qquad \forall r > r_{0}$$

$$\Rightarrow u(r) \leq u(r_{0})\left(\frac{n - (101/100) - u_{r}(r)}{n - (101/100) - u_{r}(r_{0})}\right)^{2} \leq u(r_{0})\left(\frac{n - (101/100)}{n - (203/200)}\right)^{2} \qquad \forall r > r_{0}. \tag{4.29}$$

By (4.21) and (4.29) u(r) is monotone increasing and uniformly bounded in $r > r_0$. Hence $0 < b_1 = \lim_{r \to \infty} u(r) \in [0, \infty)$ exists and (1.14) follows.

Case 2: There exists a constant $s_0 > 0$ such that (4.22) holds.

By (4.22) u(r) > 0 is monotone decreasing in $r > s_0$. Hence (1.14) follows.

Lemma 4.5. Let $\mu_1 < 0$ and $2 \le n < 4$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11). Let u be given by (4.16). Suppose there exists a constant $s_0 > 0$ such that (4.22) holds. Then (4.23) holds.

Proof: Let $2 \le n < 4$. Suppose (4.23) does not hold. Then there exists a constant $0 < \delta < 1$ and a sequence $\{\overline{s}_i\}_{i=1}^{\infty}, \overline{s}_i \to \infty$ as $i \to \infty$, such that

$$u_r(\overline{s}_i) < -\delta \quad \forall i \in \mathbb{Z}^+.$$
 (4.30)

By (1.12) there exists a constant $s_2 > s_0$ such that

$$0 < h(r) \le \delta/300 \quad \forall r \ge s_2. \tag{4.31}$$

By Lemma 4.4 (1.14) holds. Then by (1.14) and (4.22), there exists a constant $s_3 > \max(s_2, 1/\delta)$ such that

$$0 < u(s_3) - u(2s_3) < 1. (4.32)$$

Hence by (4.32) and the mean value theorem there exists a constant $s_4 \in (s_3, 2s_3)$ such that

$$u_r(s_4) = \frac{u(2s_3) - u(s_3)}{s_3} > -\frac{1}{s_3} \ge -\delta.$$
 (4.33)

Then by (4.30) there exists $i_0 \in \mathbb{Z}^+$ such that $s_5 := \overline{s}_{i_0} > s_4$ and

$$u_r(s_5) < -\delta. \tag{4.34}$$

Let

$$s_6 = \inf\{s' > 0 : u_r(r) < -\delta \quad \forall s' < r \le s_5\}.$$

Then by (4.33) $s_6 \in (s_4, s_5)$ and

$$u_r(r) < -\delta \quad \forall s_6 < r \le s_5 \quad \text{and} \quad u_r(s_6) = -\delta$$
 (4.35)

$$\Rightarrow u_{rr}(s_6) \le 0. \tag{4.36}$$

By (4.31) and (4.35),

$$u_r(r)^2 \ge \delta^2 \ge 100h(r)^2 > 2(4-n)h(r)^2 \quad \forall s_6 \le r \le s_5, 2 \le n < 4.$$
 (4.37)

Then by (4.20), (4.22), (4.31) and (4.37),

$$u_{rr}(s_6) \ge \frac{-u_r(s_6)(n - (101/100) - u_r(s_6)) - (4 - n)h(s_6)^2}{2u(s_6)}$$

$$\ge \frac{u_r(s_6)^2 - (4 - n)h(s_6)^2}{2u(s_6)}$$

$$\ge \frac{u_r(s_6)^2}{4u(s_6)}$$

$$= \frac{\delta^2}{4u(s_6)} > 0 \quad \forall 2 \le n < 4$$

which contradicts (4.36). Hence (4.23) holds and the lemma follows.

Lemma 4.6. Let $\mu_1 < 0$ and $2 \le n < 4$. Suppose $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ is a solution of (1.11). Let q be given by (4.3). Suppose there exists a constant $s_0 > 0$ such that (4.22) holds. Then there exists a constant C > 0 such that

$$-C < q(r) < 0 \quad \forall r > 0. \tag{4.38}$$

Proof: Let $2 \le n < 4$ and $r_0 > 1$ be as in (4.28). By Lemma 4.5 there exists a constant $s_1 > \max(s_0, r_0)$ such that

$$-1/10 < u_r(r) < 0 \quad \forall r > s_1. \tag{4.39}$$

Suppose (4.38) does not hold for any C > 0. Then there exists a constant $s_2 > s_1$ such that $q(s_2) < -32$. Let

$$s_3 = \sup\{s' > s_2 : q(r) < -32 \quad \forall s_2 \le r < s'\}.$$

Then

$$q(r) < -32 \quad \forall s_2 \le r < s_3.$$
 (4.40)

By (4.18), (4.28) and (4.39),

$$0 > rh_r(r) > -h(r) - \frac{1}{10} \ge -\frac{11}{100} \quad \forall r > s_1. \tag{4.41}$$

By (4.5), (4.28) and (4.41),

$$rq_{r}(r) = -q(r) \left(\frac{n-1+rh_{r}(r)}{2h(r)} - 1 \right) - (n-1) \frac{1-h(r)}{2h(r)}$$

$$\geq -q(r) \left(\frac{n-(111/100)}{2h(r)} - \frac{1}{100h(r)} \right) - \frac{n-1}{2h(r)}$$

$$\geq -\frac{C_{1}(q(r)+C_{2})}{2h(r)} \quad \forall s_{2} \leq r < s_{3}, 2 \leq n < 4.$$

$$(4.42)$$

where

$$C_1 := n - \frac{113}{100} > 0$$
 and $1 < C_2 := \frac{n-1}{n - \frac{113}{100}} < 2 \quad \forall 2 \le n < 4.$ (4.43)

By (4.40) and (4.43),

$$q(r) + C_2 < 0 \quad \forall s_2 \le r < s_3, 2 \le n < 4.$$
 (4.44)

Suppose $s_3 = \infty$. Then by (4.42) and (4.44),

$$\frac{q_{r}(r)}{q(r) + C_{2}} \leq -\frac{C_{1}}{2rh(r)} \quad \forall r \geq s_{2}, 2 \leq n < 4$$

$$\Rightarrow \log\left(\frac{q(r) + C_{2}}{q(s_{2}) + C_{2}}\right) \leq -\int_{s_{2}}^{r} \frac{C_{1}}{2\rho h(\rho)} d\rho \quad \forall r \geq s_{2}, 2 \leq n < 4$$

$$\Rightarrow q(r) + C_{2} \geq (q(s_{2}) + C_{2})e^{-\int_{s_{2}}^{r} \frac{C_{1}}{2\rho h(\rho)} d\rho} \quad \forall r \geq s_{2}, 2 \leq n < 4. \tag{4.45}$$

Letting $r \to \infty$ in (4.45), by (4.28) and (4.43),

$$\liminf_{r\to\infty}q(r)\geq -C_2>-2.$$

which contradicts (4.40) and the assumption that $s_3 = \infty$. Hence $s_3 < \infty$ and $q(s_3) = -32$. By (4.42) and (4.43),

$$q_r(s_3) \ge -\frac{C_1(q(s_3) + C_2)}{2s_3h(s_3)} \ge \frac{15C_1}{s_3h(s_3)} > 0.$$
 (4.46)

By (4.46) there exists a constant $s_4 > s_3$ such that

$$q_r(r) > 0 \qquad \forall s_3 \le r \le s_4$$

$$\Rightarrow q(r) > q(s_3) = -32 \quad \forall s_3 \le r \le s_4. \tag{4.47}$$

Since (4.38) does not hold for any C > 0, there exists a constant $s_5 > s_4$ such that $q(s_5) < -32$. Let

$$s_6 = \inf\{s' < s_5 : q(r) < -32 \quad \forall s' < r \le s_5\}.$$

By (4.47) $s_6 \in (s_4, s_5)$. Then

$$q(r) < -32 \quad \forall s_6 < r \le s_5 \quad \text{and} \quad q(s_6) = -32$$
 (4.48)

$$\Rightarrow q_r(s_6) \le 0. \tag{4.49}$$

By (4.48) and an argument similar to the proof of (4.46),

$$q_r(s_6) \ge -\frac{C_1(q(s_6) + C_2)}{2s_6h(s_6)} \ge \frac{15C_1}{s_6h(s_6)} > 0$$
 (4.50)

which contradicts (4.49). Thus no such constant s_5 exists and

$$q(r) \ge -32 \quad \forall r \ge s_4$$

 $\Rightarrow q(r) > -C_3 \quad \forall r > 0$ (4.51)

where $C_3 = \min(32, 1 + \max_{0 \le r \le s_5} |q(r)|)$. This contradicts our assumption that (4.38) does not hold for any C > 0. Hence no such constant s_2 exists. Thus there exists a constant C > 0 such that (4.38) holds and the lemma follows.

We are now ready to prove Theorem 1.6.

Proof of Theorem 1.6: Let $r_0 > 1$ be as in (4.28). Let

$$F(r) = \exp\left(\frac{n-1}{2} \int_{r_0}^r \frac{d\rho}{\rho h(\rho)}\right) \quad \forall r \ge r_0. \tag{4.52}$$

Multiplying (4.5) by F(r)/r and integrating over $(r_0, r), r > r_0$,

$$\frac{F(r)q(r)}{r} = r_0^{-1}q(r_0) - \frac{1}{2} \int_{r_0}^r \left(q(\rho)^2 + \frac{(n-1)(1-h(\rho))}{h(\rho)} \right) \frac{F(\rho)}{\rho^2} d\rho$$

$$\Rightarrow q(r) = \frac{r_0^{-1}q(r_0)}{r^{-1}F(r)} - \frac{1}{2r^{-1}F(r)} \int_{r_0}^r \left(q(\rho)^2 + \frac{(n-1)(1-h(\rho))}{h(\rho)} \right) \frac{F(\rho)}{\rho^2} d\rho \quad \forall r > r_0. \tag{4.53}$$

Let $\{r_i\}_{i=1}^{\infty} \subset (r_0, \infty)$ be a sequence such that $r_i \to \infty$ as $i \to \infty$. By Lemma 4.2, Lemma 4.3 and Lemma 4.6 there exists a constant C > 0 such that (4.38) holds. Hence the sequence $\{r_i\}_{i=1}^{\infty}$ has a subsequence which we may assume without loss of generality to be the sequence itself that converges to some constant $q_1 \in [-C, 0]$ as $i \to \infty$. By (4.28),

$$F(r) \ge \exp\left(\int_{r_0}^r \frac{d\rho}{2\rho h(\rho)}\right) \ge \exp\left(50 \int_{r_0}^r \frac{d\rho}{\rho}\right) = r_0^{-50} r^{50} \quad \forall r > r_0$$

$$\Rightarrow \quad r^{-2} F(r) \ge r_0^{-50} r^{48} \to \infty \quad \text{as } r \to \infty$$

$$\Rightarrow \quad \int_{r_0}^r \left(q(\rho)^2 + \frac{(n-1)(1-h(\rho))}{h(\rho)}\right) \frac{F(\rho)}{\rho^2} d\rho \ge \int_{r_0}^r \rho^{-2} F(\rho) d\rho \to \infty \quad \text{as } r \to \infty.$$

$$(4.54)$$

By (1.12), (4.53), (4.54), (4.55) and the l'Hospital rule,

$$q_{1} = \lim_{i \to \infty} q(r_{i}) = -\lim_{i \to \infty} \frac{1}{2r_{i}^{-1}F(r_{i})} \int_{r_{0}}^{r_{i}} \left(q(\rho)^{2} + \frac{(n-1)(1-h(\rho))}{h(\rho)} \right) \frac{F(\rho)}{\rho^{2}} d\rho$$

$$= -\frac{1}{2} \lim_{i \to \infty} \frac{\left(q(r_{i})^{2} + \frac{(n-1)(1-h(r_{i}))}{h(r_{i})} \right) r_{i}^{-2}F(r_{i})}{-r_{i}^{-2}F(r_{i}) + r_{i}^{-1}F(r_{i})(n-1)(2r_{i}h(r_{i}))^{-1}}$$

$$= \lim_{i \to \infty} \frac{q(r_{i})^{2}h(r_{i}) + (n-1)(1-h(r_{i}))}{2h(r_{i}) - (n-1)}$$

$$= -1.$$

Since the sequence $\{r_i\}_{i=1}^{\infty}$ is arbitrary, (1.13) holds.

Lemma 4.7. Let $\mu_1 < 0$ and $n \ge 2$. Suppose $h \in C^2((0,\infty)) \cap C^1([0,\infty))$ is a solution of (1.11). Let

$$w(r) = \frac{u_r(r)}{h(r)^2} \quad \forall r > 0.$$
 (4.56)

Then

$$\lim_{r \to \infty} w(r) = \frac{n-4}{n-1}.$$
 (4.57)

Proof: Let q and u be given by (4.3) and (4.16) respectively. By (1.12), Theorem 1.6, Lemma 4.2 and Lemma 4.5 there exists a constant $r_0 > 1$ such that (4.28) and

$$-3/2 < q(r) < -1/2 \quad \forall r \ge r_0 \tag{4.58}$$

and

$$|u_r| \le 1/500 \quad \forall r \ge r_0 \tag{4.59}$$

hold. By (4.20) and a direct computation,

$$w_r(r) + \frac{n - 1 - 2h(r) + 4q(r)h(r) - u_r(r)}{2rh(r)}w(r) = \frac{n - 4}{2rh(r)} \quad \forall r > 0.$$
 (4.60)

Let

$$F_{1}(r) = \exp\left(\int_{r_{0}}^{r} \frac{n - 1 - 2h(\rho) + 4q(\rho)h(\rho) - u_{r}(\rho)}{2\rho h(\rho)} d\rho\right) \quad \forall r \ge r_{0}. \tag{4.61}$$

Multiplying (4.60) by F_1 and integrating over (r_0, r) , $r > r_0$,

$$w(r)F_{1}(r) = w(r_{0}) + \frac{n-4}{2} \int_{r_{0}}^{r} \frac{F_{1}(\rho)}{\rho h(\rho)} d\rho$$

$$\Rightarrow w(r) = \frac{w(r_{0})}{F_{1}(r)} + \frac{n-4}{2F_{1}(r)} \int_{r_{0}}^{r} \frac{F_{1}(\rho)}{\rho h(\rho)} d\rho \quad \forall r \geq r_{0}. \tag{4.62}$$

By (4.18), (4.28), (4.58) and (4.59),

$$\frac{n-1-2h(\rho)+4q(\rho)h(\rho)-u_{r}(\rho)}{2h(\rho)} \ge \frac{1-\frac{2}{500}-\frac{4}{500}\cdot\frac{3}{2}-\frac{1}{500}}{\frac{2}{500}} > 245$$

$$\Rightarrow F_{1}(r) \ge \exp\left(245\int_{r_{0}}^{r}\frac{d\rho}{\rho}\right) \ge (r/r_{0})^{245} \ \forall r \ge r_{0}$$

$$\to \infty \quad \text{as } r \to \infty. \tag{4.63}$$

Then by (4.28) and (4.63),

$$\int_{r_0}^{r} \frac{F_1(\rho)}{\rho h(\rho)} d\rho \ge \frac{500}{245r_0^{245}} (r^{245} - r_0^{245}) \quad \forall r \ge r_0$$

$$\to \infty \quad \text{as } r \to \infty. \tag{4.64}$$

By Lemma 4.2 and Lemma 4.5 (4.23) holds. By (1.12), (4.23), (4.62), (4.63), (4.64), the l'Hosiptal rule and Theorem 1.6, we get

$$\lim_{r \to \infty} w(r) = \lim_{r \to \infty} \frac{w(r_0)}{F_1(r)} = 0 \quad \text{if } n = 4$$

and

$$\lim_{r \to \infty} w(r) = \frac{n-4}{2} \lim_{r \to \infty} \frac{F_1(r)(rh(r))^{-1}}{F_1(r)(n-1-2h(r)+4g(r)h(r)-u_r(r))(2rh(r))^{-1}} = \frac{n-4}{n-1} \quad \forall n \ge 2, n \ne 4$$

and (4.57) follows.

Lemma 4.8. Let $\mu_1 < 0$, $n \ge 4$ and $h \in C^2((0, \infty)) \cap C^1([0, \infty))$ be a solution of (1.11). Then (1.14) holds with $b_1 > 0$.

Proof: Let *u* be given by (4.16). By Lemma 4.7 there exists a constant $s_1 > 1$ such that

$$\frac{u_{r}(r)}{h(r)^{2}} < \frac{n - (3/2)}{n - 1} =: C_{3} < 1 \quad \forall r \ge s_{1}$$

$$\Rightarrow u(r)^{-2}u_{r}(r) < C_{3}r^{-2} \quad \forall r \ge s_{1}$$

$$\Rightarrow \frac{1}{u(s_{1})} - \frac{1}{u(r)} < \frac{C_{3}}{s_{1}} - \frac{C_{3}}{r} \quad \forall r \ge s_{1}$$

$$\Rightarrow \frac{1}{u(r)} > \frac{1}{s_{1}h(s_{1})} - \frac{C_{3}}{s_{1}} + \frac{C_{3}}{r} \ge \frac{1 - C_{3}}{s_{1}} \quad \forall r \ge s_{1}$$

$$\Rightarrow u(r) < \frac{s_{1}}{1 - C_{3}} \quad \forall r \ge s_{1}.$$
(4.65)

By (4.21) and (4.65) u(r) > 0 is monotone increasing and uniformly bounded in $r > s_1$. Hence $0 < b_1 = \lim_{r \to \infty} u(r)$ exists and (1.14) follows.

Proof of Theorem 1.7:

(1.14) is proved in Lemma 4.4 and Lemma 4.8. Hence it remains to prove (1.15). Let

$$v(r) = r(q(r) + 1) \quad \forall r \ge 0.$$

Then by (4.5) and a direct computation,

$$v_r(r) + \left(\frac{n-1}{h(r)} - 5 + q(r)\right) \frac{v(r)}{2r} = \frac{n-4}{2} \quad \forall r > 0.$$
 (4.66)

By (1.12) and Theorem 1.6 there exists a constant $r_0 > 1$ such that (4.28) and (4.58) holds. Let

$$H(r) = \exp\left(\frac{1}{2} \int_{r_0}^r \left(\frac{n-1}{h(\rho)} + q(r)\right) \frac{d\rho}{\rho}\right) \quad \forall r \ge r_0.$$
 (4.67)

Multiplying (4.66) by $r^{-5/2}H(r)$ and integrating over (r_0, r) , $r > r_0$,

$$r^{-5/2}H(r)v(r) = r_0^{-5/2}v(r_0) + \frac{n-4}{2} \int_{r_0}^{r} \rho^{-5/2}H(\rho) d\rho \quad \forall r > r_0$$

$$\Rightarrow v(r) = \frac{r_0^{-5/2}v(r_0)}{r^{-5/2}H(r)} + \frac{n-4}{2r^{-5/2}H(r)} \int_{r_0}^{r} \rho^{-5/2}H(\rho) d\rho \quad \forall r > r_0. \tag{4.68}$$

By (4.28) and (4.58),

$$H(r) \ge \exp\left(\frac{1}{2} \int_{r_0}^r (100(n-1) - (3/2)) \frac{d\rho}{\rho}\right) \ge \exp\left(49 \int_{r_0}^r \frac{d\rho}{\rho}\right) = (r/r_0)^{49} \quad \forall r > r_0$$

$$\Rightarrow \quad r^{-5/2}H(r) \ge r_0^{-49} r^{93/2} \to \infty \quad \text{as } r \to \infty$$

$$\Rightarrow \quad \int_r^r \rho^{-5/2}H(\rho) d\rho \to \infty \quad \text{as } r \to \infty.$$

$$(4.69)$$

Hence for n = 4 by (4.68) and (4.69),

$$\lim_{r\to\infty}v(r)=0$$

For $n \ge 2$ and $n \ne 4$ by (1.14), (4.68), (4.69), (4.70), the l'Hospital rule and Theorem 1.6,

$$\begin{split} \lim_{r \to \infty} v(r) &= \frac{n-4}{2} \lim_{r \to \infty} \frac{1}{r^{-5/2} H(r)} \int_{r_0}^r \rho^{-5/2} H(\rho) \, d\rho \\ &= \frac{n-4}{2} \lim_{r \to \infty} \frac{r^{-5/2} H(r)}{-(7/2) r^{-7/2} H(r) + r^{-5/2} H(r) [(n-1)(2rh(r))^{-1} + q(r)(2r)^{-1}]} \\ &= \left(\frac{n-4}{n-1}\right) \lim_{r \to \infty} rh(r) = \left(\frac{n-4}{n-1}\right) b_1 \quad \forall n \ge 2, n \ne 4 \end{split}$$

where b_1 is given by (1.14) and (1.15) follows.

Alternate proof of Theorem 1.7: Let w be given by (4.56). Then by Theorem 1.6 and Lemma $\overline{4.7}$,

$$v(r) = w(r)rh(r) \quad \forall r > 0$$

$$\Rightarrow \lim_{r \to \infty} v(r) = \lim_{r \to \infty} w(r) \cdot \lim_{r \to \infty} rh(r) = \left(\frac{n-4}{n-1}\right)b_1 \quad \forall n \ge 2$$

and (1.15) follows.

Remark 4.9. By (1.15) of Theorem 1.7,

$$\frac{h_r(r)}{h(r)} \approx -\frac{1}{r} - \frac{c_1 b_1}{r^2} \quad as \ r \to \infty$$

$$\Rightarrow \log\left(\frac{h(r)}{h(r_0)}\right) \approx \log\left(\frac{r_0}{r}\right) + c_1 b_1 \left(\frac{1}{r} - \frac{1}{r_0}\right) \quad \forall 0 < r_0 < r \quad as \ r > r_0 \to \infty$$

$$\Rightarrow h(r) \approx \frac{C}{r} e^{\frac{c_1 b_1}{r}} \quad as \ r \to \infty$$

$$\Rightarrow \frac{1}{\sqrt{h(r^2)}} \approx \frac{r}{\sqrt{C}} e^{-\frac{c_1 b_1}{r^2}} \approx \frac{r}{\sqrt{C}} \left(1 - \frac{c_1 b_1}{r^2}\right) \quad as \ r \to \infty$$
(4.71)

for some constant C > 0 depending on r_0 where b_1 is as given by (1.14) and $c_1 = \frac{n-4}{n-1}$. Hence by (1.5) and (4.71),

$$t = \int_0^{a(t)} \frac{d\rho}{\sqrt{h(\rho^2)}} \approx \frac{1}{\sqrt{C}} \int_{r_0}^{a(t)} \left(\rho - \frac{c_1 b_1}{\rho}\right) d\rho \approx \frac{1}{2\sqrt{C}} \left(a(t)^2 - c_1 b_1 \log a(t)^2\right) \quad as \ a(t) \to \infty$$

where $r_0 > 0$ is a sufficiently large constant. Thus $t \to \infty$ as $a(t) \to \infty$ and the metric g given by (1.2) or equivalent by (1.4) with a(t) and h(a(t)) related by (1.5) where h is a solution of (1.11) with $\mu_1 < 0$ is a complete metric.

5 Asymptotic behaviour of rotationally symmetric expanding gradient Ricci soliton

In this section we will prove the asymptotic behaviour of the solution h of (1.8) for the case $n \ge 2$ and $\lambda > 0$ as $r \to \infty$. We will assume that $\lambda > 0$, $\mu_1 \in \mathbb{R} \setminus \{0\}$, $n \ge 2$ and $h \in C^2((0,\infty)) \cap C^1([0,\infty))$ is a solution of (1.8) in this section.

Proof of Theorem 1.8:

We first observe that by (2.25) of Lemma 2.3,

$$h_r(r) = \frac{n-1}{r} + \lambda + \sqrt{\frac{h(r)}{h(r_1)}} H_1(r_1, r) \le \frac{n-1}{r} + \lambda + \sqrt{\frac{h(r)}{h(r_1)}} H_2(r_1) \quad \forall r > r_1 > 0$$
 (5.1)

where

$$H_1(r_1, r) = h_r(r_1) - \frac{n-1}{r_1} + \frac{(n-1)\sqrt{h(r_1)}}{2} \int_{r_1}^r \frac{h(\rho) + 1}{\rho^2 \sqrt{h(\rho)}} d\rho - \lambda \quad \forall r > r_1 > 0$$
 (5.2)

and

$$H_2(r_1) = h_r(r_1) + \frac{(n-1)\sqrt{h(r_1)}}{2} \int_{r_1}^{\infty} \frac{h(\rho) + 1}{\rho^2 \sqrt{h(\rho)}} d\rho - \lambda \quad \forall r_1 > 0.$$
 (5.3)

On the other hand by Theorem 1.2 (1.10) holds. Hence by (1.10),

$$h_{\infty} := \lim_{r \to \infty} h(r) \in [0, \infty]$$
 exists.

Proof of (i): By direct computation (1.16) is the explicit analytic solution of (1.8).

Proof of (ii): Suppose $0 < \mu_1 < \lambda/n$. Let $\mu_1 < \delta < \lambda/n$. By (1.8) there exists a constant $s_0 > 0$ such that

$$h_r(r) < \delta \quad \forall 0 \le r < s_0.$$

Let

$$R_0 = \sup\{r_1 > 0 : h_r(r) < \delta \quad \forall 0 \le r < r_1\}.$$

Then $R_0 \ge s_0$. Suppose $R_0 < \infty$. Then

$$h_r(r) < \delta \quad \forall 0 \le r < R_0 \quad \text{and} \quad h_r(R_0) = \delta$$
 (5.4)

$$\Rightarrow 1 < h(r) < 1 + \delta r \quad \forall 0 < r \le R_0 \quad \text{and} \quad h_{rr}(R_0) \ge 0.$$
 (5.5)

By (1.8), (5.4) and (5.5),

$$2R_0^2h(R_0)h_{rr}(R_0) < (n-1)(1+\delta R_0)\delta R_0 + \delta R_0(\delta R_0 - \lambda R_0 - (n-1)) = \delta R_0^2(n\delta - \lambda) < 0$$

$$\Rightarrow h_{rr}(R_0) < 0$$

which contradicts (5.5). Hence $R_0 = \infty$. Thus by letting $\delta \searrow \mu_1$ in (5.4) and (5.5),

$$0 < h_r(r) \le \mu_1$$
 and $1 < h(r) \le 1 + \mu_1 r \quad \forall r > 0.$ (5.6)

Thus $h_{\infty} > 1$ and (1.17) follows. Suppose

$$h_{\infty} = \infty. \tag{5.7}$$

By (5.6) we can divide the proof of (ii) into two cases.

Case 1: $\lim_{r\to\infty} h_r(r) = 0$.

Let $0 < \delta_0 < \frac{\lambda}{2n}$. Then there exists a constant $r_0 > 1$ such that

$$h_r(r) < \delta_0 \quad \forall r \ge r_0 \tag{5.8}$$

$$\Rightarrow 1 < h(r) < h(r_0) + \delta_0 r \quad \forall r \ge r_0. \tag{5.9}$$

Hence by (5.9) for any $r_1 > r_0$,

$$\frac{\sqrt{h(r_1)}}{2} \int_{r_1}^{\infty} \frac{h(\rho) + 1}{\rho^2 \sqrt{h(\rho)}} d\rho \le (h(r_0) + \delta_0 r_1)^{1/2} \int_{r_1}^{\infty} \frac{\sqrt{h(\rho)}}{\rho^2} d\rho$$

$$\le (h(r_0) + \delta_0 r_1)^{1/2} (h(r_0) r_1^{-1} + \delta_0)^{1/2} \int_{r_1}^{\infty} \rho^{-3/2} d\rho$$

$$\le 2(h(r_0) + \delta_0 r_1)^{1/2} (h(r_0) r_1^{-1} + \delta_0)^{1/2} r_1^{-1/2}.$$
(5.10)

By (5.3) and (5.10),

$$\lim_{r_1 \to \infty} \sup H_2(r_1) \le 2(n-1)\delta_0 - \lambda < 0. \tag{5.11}$$

By (5.11) there exists a constant $r_1 > r_0$ such that

$$H_2(r_1) < 0. (5.12)$$

By (5.1), (5.7) and (5.12),

$$\limsup_{r\to\infty}h_r(r)=-\infty$$

which contradicts (1.10). Hence case 1 does not hold.

<u>Case 2</u>: There exists a constant $\delta_1 > 0$ and a sequence $\{s_i\}_{i=1}^{\infty} \subset (1, \infty)$, $s_i \to \infty$ as $i \to \infty$, such that

$$h_r(s_i) \ge \delta_1 \quad \forall i \in \mathbb{Z}^+.$$
 (5.13)

By (5.6) and (5.13) the sequence $\{s_i\}_{i=1}^{\infty}$ has a subsequence which we may assume without loss of generality to be the sequence itself such that

$$h_r(s_i) \to \delta_2 \quad \text{as } i \to \infty$$
 (5.14)

for some constant $\delta_2 \in (0, \mu_1]$. By (5.6),

$$\int_{s_{i}}^{\infty} \frac{h(\rho) + 1}{\rho^{2} \sqrt{h(\rho)}} d\rho \leq 2 \int_{s_{i}}^{\infty} \frac{\sqrt{h(\rho)}}{\rho^{2}} d\rho \leq 2(1 + \mu_{1})^{1/2} \int_{s_{i}}^{\infty} \rho^{-3/2} d\rho \leq 4(1 + \mu_{1})^{1/2} s_{i}^{-1/2} \quad \forall i \in \mathbb{Z}^{+}$$

$$\to 0 \quad \text{as } i \to \infty.$$
(5.15)

By (5.7), (5.14), (5.15) and the l'Hospital rule,

$$\lim_{i \to \infty} \frac{\sqrt{h(s_{i})}}{2} \int_{s_{i}}^{\infty} \frac{h(\rho) + 1}{\rho^{2} \sqrt{h(\rho)}} d\rho = \lim_{i \to \infty} \frac{\int_{s_{i}}^{\infty} \frac{h(\rho) + 1}{2\rho^{2} \sqrt{h(\rho)}} d\rho}{h(s_{i})^{-1/2}} = \lim_{i \to \infty} \frac{\frac{h(s_{i}) + 1}{s_{i}^{2} \sqrt{h(s_{i})}}}{h(s_{i})^{-3/2} h_{r}(s_{i})}$$

$$= \delta_{2}^{-1} \lim_{i \to \infty} \frac{h(s_{i})}{s_{i}} \cdot \lim_{i \to \infty} \frac{h(s_{i}) + 1}{s_{i}}$$

$$= \delta_{2}^{-1} \lim_{i \to \infty} h_{r}(s_{i}) \cdot \lim_{i \to \infty} h_{r}(s_{i})$$

$$= \delta_{2}. \tag{5.16}$$

Putting $r_1 = s_i$, $r = \infty$, in (5.2) and letting $i \to \infty$, by (5.14) and (5.16),

$$\lim_{i\to\infty} H_1(s_i,\infty) = \delta_2 + (n-1)\delta_2 - \lambda = n\delta_2 - \lambda \le n\mu_1 - \lambda < 0.$$

Hence there exists $i_0 \in \mathbb{Z}^+$ such that

$$H_1(s_{i_0}, \infty) < 0.$$
 (5.17)

Putting $r_1 = s_{i_0}$ in (5.1) and letting $r \to \infty$, by (5.1), (5.7) and (5.17),

$$\limsup_{r\to\infty}h_r(r)=-\infty$$

which contradicts (1.10). Hence case 2 does not hold. Thus (5.7) does not hold and $h_{\infty} < \infty$. Putting $r_1 = 1$ and letting $r \to \infty$ in (2.25), by (5.6),

$$\lim_{r \to \infty} h_r(r) = \lambda + \sqrt{\frac{h_\infty}{h(1)}} \left(h_r(1) - (n-1) - \lambda \right) + (n-1) \frac{\sqrt{h_\infty}}{2} \int_1^\infty \frac{h(\rho) + 1}{\rho^2 \sqrt{h(\rho)}} d\rho \in [0, \mu_1]. \quad (5.18)$$

Since $h_{\infty} < \infty$, by (5.18),

$$\lim_{r\to\infty}h_r(r)=0$$

and (ii) follows.

Proof of (iii): Suppose $\mu_1 > \lambda/n$. Let $\mu_1 > \delta > \lambda/n$. By (1.8) there exists a constant $s_0' > 0$ such that

$$h_r(r) > \delta \quad \forall 0 \le r < s'_0.$$

Let

$$R_0 = \sup\{r_1 > 0 : h_r(r) > \delta \quad \forall 0 \le r < r_1\}.$$

Then $R_0 \ge s_0'$. Suppose $R_0 < \infty$. Then

$$h_r(r) > \delta \quad \forall 0 \le r < R_0 \quad \text{and} \quad h_r(R_0) = \delta$$
 (5.19)

$$\Rightarrow h(r) > 1 + \delta r \quad \forall 0 < r \le R_0 \quad \text{and} \quad h_{rr}(R_0) \le 0. \tag{5.20}$$

By (1.8), (5.19) and (5.20),

$$2R_0^2 h(R_0) h_{rr}(R_0) > (n-1) (1 + \delta R_0) \delta R_0 + \delta R_0 (\delta R_0 - \lambda R_0 - (n-1)) = \delta R_0^2 (n\delta - \lambda) > 0$$

$$\Rightarrow h_{rr}(R_0) > 0$$

which contradicts (5.20). Hence $R_0 = \infty$. Thus by letting $\delta \nearrow \mu_1$ in (5.19) and (5.20),

$$h_r(r) \ge \mu_1$$
 and $h(r) \ge 1 + \mu_1 r$ $\forall r > 0$ (5.21)
 $\Rightarrow h_\infty = \lim_{r \to \infty} h(r) = \infty.$

Hence (1.19) holds. By (5.21),

$$\sqrt{h(r_1)} \int_{r_1}^{\infty} \frac{h(\rho) + 1}{2\rho^2 \sqrt{h(\rho)}} d\rho \ge \frac{(1 + \mu_1 r_1)^{1/2} \sqrt{\mu_1}}{2} \int_{r_1}^{\infty} \rho^{-3/2} d\rho \ge \mu_1 \quad \forall r_1 > 0.$$
 (5.22)

By (5.2), (5.21) and (5.22),

$$\liminf_{r_1 \to \infty} H_1(r_1, \infty) \ge \mu_1 + (n-1)\mu_1 - \lambda = n\mu_1 - \lambda > 0.$$

Hence there exists a constant $r_1 > 0$ such that

$$H_1(r_1,\infty)>0.$$

Thus there exists a constant $r'_0 > r_1$ such that

$$H_1(r_1, r) \ge H_1(r_1, r'_0) > 0 \quad \forall r \ge r'_0.$$
 (5.23)

Letting $r \to \infty$ in (5.1), by (5.23),

$$\lim_{r\to\infty}h_r(r)=\infty$$

and (iii) follows.

Proof of (iv): Suppose $h_{\infty} = 0$. Then by (5.1),

$$h_r(r) \ge \frac{n-1}{r} + \lambda + \sqrt{\frac{h(r)}{h(r_1)}} \left(h_r(r_1) - \frac{n-1}{r_1} - \lambda \right) \quad \forall r > r_1 > 0$$

$$\Rightarrow \quad \liminf_{r \to \infty} h_r(r) \ge \lambda > 0$$

which contradicts (1.10). Hence by (1.10) $h_{\infty} \in (0, 1)$. Thus,

$$\int_{1}^{\infty} \frac{h(\rho) + 1}{\rho^2 \sqrt{h(\rho)}} d\rho < \infty. \tag{5.24}$$

Putting $r_1 = 1$ and letting $r \to \infty$ in (2.25), by (1.10) and (5.24),

$$h_r^{\infty} := \lim_{r \to \infty} h_r(r) = \lambda + \sqrt{\frac{h_{\infty}}{h(1)}} \left(h_r(1) - (n-1) - \lambda \right) + (n-1) \frac{\sqrt{h_{\infty}}}{2} \int_1^{\infty} \frac{h(\rho) + 1}{\rho^2 \sqrt{h(\rho)}} d\rho \in (-\infty, 0].$$

If h_r^{∞} < 0, then there exists a constant r_0 > 0 such that

$$h_r(r) \le h_r^{\infty}/2 < 0 \quad \forall r \ge r_0$$

$$\Rightarrow h(r) \le h(r_0) + (r - r_0)(h_r^{\infty}/2) \le 1 + (r - r_0)(h_r^{\infty}/2) < 0 \quad \forall r > r_0 + (2/|h_r^{\infty}|)$$

which contradicts (1.10). Hence

$$h_r^{\infty} := \lim_{r \to \infty} h_r(r) = 0$$

and (iv) follows.

Finally by an argument similar to the proof of (iii) of Theorem 1.8 we get that Theorem 1.9 holds.

References

- [B1] S. Brendle, *Rotational symmetry of self-similar solutions to the Ricci flow*, Inventiones Mathematicae 194 (2013), 731—764.
- [B2] S. Brendle, Ancient solutions to the Ricci flow in dimension three, Acta Math. 225 (2020), 1—102.
- [Br] R.L. Bryant, *Ricci flow solitons in dimension three with SO*(3)-symmetries, available at: www.math.duke.edu/~bryant/3DRotSymRicciSolitons.pdf
- [C] H.D. Cao, *Recent progress on Ricci solitons*, Recent advances in geometric analysis", Y.I. Lee, C.S. Lin, M.P. Tsui, editors, Adv. Lect. Math. 11, Int. Press, Somerville, MA (2010), 1–38.
- [CZ] H.D. Cao and D. Zhou, *On complete gradient shrinking Ricci solitons* J. Differential Geom. 85 (2010), no. 2, 175–186.
- [FIK] M. Feldman, T. Ilmanen and D. Knopf, *Rotationally symmetric shrinking and expanding gradient Kähler Ricci solitons*, J. Differ. Geom. 65 (2003), 169–209.
- [H] K.M. Hui, Existence of singular rotationally symmetric gradient Ricci solitons in higher dimensions, arXiv:2107.13685.
- [LW] Y. Li and B. Wang, *Heat kernel on Ricci shrinkers*, Calc. Var. and PDE 59 (2020), article no. 194.
- [MS] O. Munteanu and N. Sesum, *On Gradient Ricci Solitons*, J. Geom. Anal. 23 (2013), 539–561.
- [PW] P. Petersen and W. Wylie, On the classification of gradient Ricci solitons, Geometry and topology (2010), 2277–2300.