ASYMPTOTIC STABILITY FOR A CLASS OF VISCOELASTIC EQUATIONS WITH GENERAL RELAXATION FUNCTIONS AND THE TIME DELAY

MENGLAN LIAO, ZHONG TAN*

ABSTRACT. The goal of the present paper is to study the viscoelastic wave equation with the time delay

 $|u_t|^{\rho} u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s)\Delta u(s)ds + \mu_1 u_t(x,t) + \mu_2 u_t(x,t-\tau) = b|u|^{p-2}u_t(x,t-\tau) = b|u|^{p-2}u_t(x,t-\tau)$

under initial boundary value conditions, where ρ , b, μ_1 are positive constants, μ_2 is a real number, $\tau > 0$ represents the time delay. By using the multiplier method together with some properties of the convex functions, the explicit and general stability results of energy are proved under the general assumption on the relaxation function g. This work generalizes and improves earlier results on the stability of the viscoelastic equations with the time delay in the literature.

1. INTRODUCTION

This paper concerns the following initial boundary value problem with the time delay

$$\begin{cases} |u_t|^{\rho} u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s)\Delta u(s)ds \\ +\mu_1 u_t(x,t) + \mu_2 u_t(x,t-\tau) = b|u|^{p-2}u \quad (x,t) \in \Omega \times (0,\infty), \\ u_t(x,t-\tau) = f_0(x,t-\tau) \quad (x,t) \in \Omega \times (0,\tau), \\ u(x,0) = u_0(x), \ u_t(x,0) = u_1(x) \quad x \in \Omega, \\ u(x,t) = 0 \quad (x,t) \in \partial\Omega \times [0,\infty), \end{cases}$$
(1.1)

where $\Omega \subset \mathbb{R}^N (N \geq 1)$ is a bounded domain with a smooth boundary $\partial\Omega$, the unknown u := u(x,t) is a real valued function defined on $\Omega \times (0,\infty)$, ρ , b, μ_1 are positive constants, μ_2 is a real number, $\tau > 0$ represents the time delay, g is the kernel of the memory term, and the initial data (u_0, u_1, f_0) are given functions belonging to suitable spaces. In addition, the following assumptions are given throughout this paper:

 $(\mathbf{H_1})$ The relaxation function $g: [0, \infty) \to (0, \infty)$ is a differentiable function satisfying

$$1 - \int_0^\infty g(s)ds = l > 0, \tag{1.2}$$

²⁰¹⁰ Mathematics Subject Classification. 35B40, 26A51, 93D20.

Key words and phrases. Viscoelasticity; delay term; source term; energy decay.

Supported by the National Natural Science Foundation of China (Grant Nos. 11926316, 11531010, 12071391).

^{*} Corresponding author: Zhong Tan.

and there exists a C^1 function $G: (0, \infty) \to (0, \infty)$ which is strictly increasing and strictly convex C^2 function on (0, r], $r \leq g(0)$, with G(0) = G'(0) = 0 such that

$$g'(t) \le -\zeta(t)G(g(t)) \quad \text{for } t \ge 0, \tag{1.3}$$

here $\zeta(t)$ is a positive non-increasing differentiable function.

 $(\mathbf{H_2}) \ \rho \ \text{and} \ p \ \text{satisfy}$

$$0 < \rho \le \frac{2}{N-2} \text{ for } N \ge 3 \text{ and } \rho > 0 \text{ for } N = 1, 2;$$

$$2 2 \text{ for } N = 1, 2.$$

It is well known that time delay effects which often appear in many practical applications may induce some instabilities. Some results on the local existence and blow-up of solutions to a class of equations with delay have been obtained, the interested readers can refer to [4, 12, 13, 14, 23] and the reference therein. Nicaise and Pignotti [20] considered the wave equation with a delay term in the boundary condition as well as the wave equation with a delayed velocity term and mixed Dirichlet-Neumann boundary condition in a bounded and smooth domain, respectively. By introducing suitable energies and by using some observability inequalities, they proved an exponential stability of the solution in both cases under suitable assumptions. Kirane and Said-Houari [11] studied the following viscoelastic wave equation with a delay term in internal feedback

$$u_{tt} - \Delta u + \int_0^t g(t-s)\Delta u(s)ds + \mu_1 u_t(x,t) + \mu_2 u_t(x,t-\tau) = 0$$
(1.4)

here μ_1 , μ_2 are positive constants, under the initial boundary conditions of problem (1.1). They proved the existence of a unique weak solution for $\mu_2 \leq \mu_1$ by using the Faedo-Galerkin approximations together with some energy estimates. Provided that $g: \mathbb{R}^+ \to \mathbb{R}^+$ is a C^1 function satisfying g(0) > 0 and (1.2), and there exists a positive non-increasing differentiable function $\zeta(t)$ such that

$$g'(t) \le -\zeta(t)g(t) \text{ for } t \ge 0 \text{ and } \int_0^{+\infty} \zeta(t)dt = +\infty,$$
 (1.5)

by establishing suitable Lyapunov functionals, they also obtained the corresponding exponential stability for $\mu_2 < \mu_1$ and for $\mu_2 = \mu_1$, respectively. Subsequently, Dai and Yang [7] proved an existence result of problem (1.4) without restrictions of μ_1 , $\mu_2 > 0$ and $\mu_2 \leq \mu_1$. Making use of the viscoelasticity term controls the delay term, they also proved an energy decay result for problem (1.4) in the case $\mu_1 = 0$ provided that $g : \mathbb{R}^+ \to \mathbb{R}^+$ is a C^1 function satisfying g(0) > 0 and (1.2), and there exists a positive constant ζ such that

$$g(t) \le -\zeta g(t) \text{ for } t > 0. \tag{1.6}$$

Liu [15] generalized the results obtained by Kirane and Said-Houari [11]. That is, by the similar method in [11], they established a general energy decay result for problem (1.4) with $\tau(t)$ instead of τ . In the absence of the source term $b|u|^{p-2}u$ in problem (1.1), Wu [21] proved an energy decay by the similar method in [11], and generalized the results to the time-varying delay in [22]. There are many papers concerning with the stability of viscoelastic equations with time delay, the interested readers may refer to [3, 8, 16] and the reference therein. However, the relaxation function q are mainly limited to satisfying among the three conditions, which are (1.5), (1.6) and that $q: \mathbb{R}^+ \to \mathbb{R}^+$ is a differentiable function satisfying q(0) > 0 and (1.2), and there exists a positive function $G \in C^1(\mathbb{R}^+)$ and G is linear or strictly increasing and strictly convex C^2 function on (0, r], r < 1, with G(0) = G'(0) = 0, such that

$$g'(t) \leq -G(g(t))$$
 for $t > 0$

Until recently, Chellaoua and Boukhatem [5] generalized the previous conditions that the relaxation function g satisfied, specifically investigated the following secondorder abstract viscoelastic equation in Hilbert spaces

$$\begin{cases} u_{tt} + Au - \int_0^\infty g(s)Bu(t-s)ds + \mu_1 u_t(t) + \mu_2 u_t(t-\tau) = 0 & t > 0, \\ u_t(t-\tau) = f_0(t-\tau) & t \in (0,\tau), \\ u(-t) = u_0(t), \ u_t(0) = u_1 & t \ge 0, \end{cases}$$

where $A: D(A) \to H$ and $B: D(B) \to H$ are a self-adjoint linear positive operator with domains $D(A) \subset D(B) \subset H$ such that the embeddings are dense and compact. They established an explicit and general decay results of the energy solution by introducing a suitable Lyapunov functional and some properties of the convex functions under the condition (\mathbf{H}_1) . Chellaoua and Boukhatem also addressed the stability results for the following second-order abstract viscoelastic equation in Hilbert spaces with time-varying delay in [6]

$$\begin{cases} u_{tt} + Au - \int_0^t g(t-s)Bu(s)ds + \mu_1 u_t(t) + \mu_2 u_t(t-\tau(t)) = 0 & t > 0, \\ u_t(t-\tau(0)) = f_0(t-\tau(0)) & t \in (0,\tau(0)), \\ u(0) = u_0, \ u_t(0) = u_1 & t \ge 0, \end{cases}$$

at

under the condition (\mathbf{H}_1) . It is worth pointing out that Mustafa [17] first proposed the condition (\mathbf{H}_1) to study the decay rates for the following initial boundary value problem

$$\begin{cases} u_{tt} - \Delta u + \int_0^t g(t-s)\Delta u(s)ds = 0 & \text{in } \Omega \times (0,\infty), \\ u = 0 & \text{on } \partial\Omega \times (0,\infty), \\ u(x,0) = u_0(x), \ u_t(x,0) = u_1(x) & x \in \Omega. \end{cases}$$

After that, many authors popularized the method used by Mustafa in [17]. The readers may see the references [2, 9, 10, 18, 19] to get more details.

Motivated by the above works, we are committed to considering the stability of problem (1.1) when the relaxation function q satisfies the condition (\mathbf{H}_1) . To the best of our knowledge, there is no decay result for problem (1.1) where the relaxation functions satisfy (\mathbf{H}_1) , although Wu [23] has investigated problem (1.1) and proved the blow-up result with nonpositive and positive initial energy. With minimal conditions on the relaxation function g, we establish a general and optimal energy decay rates of problem (1.1) in Theorem 3.3. Our proof is based on the multiplier method and the similar arguments in [5, 17] but it is different from before since the presence of Δu_{tt} , the time delay and the force source $b|u|^{p-2}u$. The outline of this paper is as follows: In Section 2, we give some preliminary results. Section 3 is used to present the energy decay and its proof. In Section 4, we give the possible generalizations.

2. Preliminaries

Throughout this paper, we denote by $\|\cdot\|_p$ and $\|\nabla\cdot\|_2$ the norm on $L^p(\Omega)$ with $1 \leq p \leq \infty$ and $H_0^1(\Omega)$, respectively. Let λ_1 be the first eigenvalue of the following boundary value problem

$$\begin{cases} -\Delta \psi = \lambda \psi & \text{ for } x \in \Omega, \\ \psi = 0 & \text{ for } x \in \partial \Omega. \end{cases}$$

The symbol c_s is the optimal embedding constant of $H^1_0(\Omega) \hookrightarrow L^p(\Omega)$.

In order to the completeness of results, in what follows, we state the known results in [23]. Let us introduce the new variable

$$z(x,\kappa,t) = u_t(x,t-\tau\kappa) \quad \text{for } x \in \Omega, \ \kappa \in (0,1),$$

then problem (1.1) is equivalent to

$$\begin{cases} |u_t|^{\rho} u_{tt} - \Delta u - \Delta u_{tt} + \int_0^\tau g(t-s)\Delta u(s)ds \\ + \mu_1 u_t(x,t) + \mu_2 z(x,1,t) = b|u|^{p-2}u & (x,t) \in \Omega \times (0,\infty), \\ \tau z_t(x,\kappa,t) + z_\kappa(x,\kappa,t) = 0 & (x,t) \in \Omega \times (0,\infty), \\ z(x,0,t) = u_t(x,t) & (x,t) \in \Omega \times (0,\infty), \\ z(x,\kappa,0) = f_0(x,-\tau\kappa) & x \in \Omega, \\ u_t(x,t-\tau) = f_0(x,t-\tau) & (x,t) \in \Omega \times (0,\tau), \\ u(x,0) = u_0(x), \ u_t(x,0) = u_1(x) & x \in \Omega, \\ u(x,t) = 0 & (x,t) \in \partial\Omega \times [0,\infty). \end{cases}$$
(2.1)

Theorem 2.1 (Theorem 2.3 in [23]). Suppose that $|\mu_2| \leq \mu_1$, (**H**₁) and (**H**₂) hold. Assume that u_0 , $u_1 \in H_0^1(\Omega)$ and $f_0 \in L^2(\Omega \times (0,1))$, then there exists a unique solution (u, z) of problem (2.1) satisfying

$$u, u_t \in C([0,T); H^1_0(\Omega)), \quad z \in C([0,T); L^2(\Omega \times (0,1))),$$

for T > 0.

Define the energy functional of problem (2.1) as follows

$$E(t) = \frac{1}{\rho+2} \|u_t\|_{\rho+2}^{\rho+2} + \frac{1}{2} \left(1 - \int_0^t g(s)ds\right) \|\nabla u\|_2^2 + \frac{1}{2} (g \circ \nabla u)(t) + \frac{1}{2} \|\nabla u_t\|_2^2 + \frac{\xi}{2} \int_\Omega \int_0^1 z^2(x,\kappa,t)d\kappa dx - \frac{b}{p} \|u\|_p^p,$$
(2.2)

here ξ is a positive constant so that

$$\tau|\mu_2| \le \xi \le \tau(2\mu_1 - |\mu_2|) \tag{2.3}$$

and

$$(g \circ u)(t) = \int_0^t g(t-s) \|u(s) - u(t)\|_2^2 ds.$$

Lemma 2.2 (Lemma 3.1 in [23]). E(t) is a non-increasing function and

$$E'(t) \leq -\omega(\|u_t\|_2^2 + \|z(x,1,t)\|_2^2) + \frac{1}{2}(g' \circ \nabla u)(t) - \frac{1}{2}g(t)\|\nabla u\|_2^2$$

$$\leq -\omega(\|u_t\|_2^2 + \|z(x,1,t)\|_2^2) \leq 0 \quad \text{for all } t \geq 0,$$
(2.4)

where $\omega = \min \left\{ \mu_1 - \frac{\xi}{2\tau} - \frac{|\mu_2|}{2}, \frac{\xi}{2\tau} - \frac{|\mu_2|}{2} \right\} \ge 0.$

Lemma 2.3. If u is a solution for problem (2.1) and

$$E(0) < E_1 = \frac{p-2}{2p}\sigma_1^2, \quad l \|\nabla u_0\|_2^2 < \sigma_1^2,$$

here $\sigma_1 = b^{-\frac{1}{p-2}} B_1^{-\frac{p}{p-2}}$, $B_1 = \frac{c_s^p}{l^{\frac{p}{2}}}$, then there exists a positive constant σ_2 satisfying $0 < \sigma_2 < \sigma_1$ such that

$$l \|\nabla u\|_2^2 + (g \circ \nabla u)(t) \le \sigma_2^2 \quad \text{for all } t \ge 0.$$

$$(2.5)$$

Proof. Taking the combination of (2.2) and (1.2) with the embedding $H_0^1(\Omega) \hookrightarrow L^p(\Omega)$, one has

$$E(t) \geq \frac{l}{2} \|\nabla u\|_{2}^{2} + \frac{1}{2} (g \circ \nabla u)(t) - \frac{bB_{1}^{p}}{p} \left(l^{\frac{1}{2}} \|\nabla u\|_{2} \right)^{p} \\ \geq F\left(\sqrt{l} \|\nabla u\|_{2}^{2} + (g \circ \nabla u)(t) \right),$$
(2.6)

where $F(x) = \frac{1}{2}x^2 - \frac{bB_1^p}{p}x^p$ for x > 0. We know that F is strictly increasing in $(0, \sigma_1)$, strictly decreasing in (σ_1, ∞) , and F has a maximum at σ_1 with the maximum value E_1 . Since $E(0) < E_1$, there exists a $\sigma_2 < \sigma_1$ such that $F(\sigma_2) = E(0)$. Set $\sigma_0 := \sqrt{l \|\nabla u_0\|_2^2}$, recall (2.6), then $F(\sigma_0) \leq E(0) = F(\sigma_2)$, which implies $\sigma_0 \leq \sigma_2$ since the given condition $\sigma_0^2 < \sigma_1^2$. In order to complete the proof of (2.5), we suppose by contradiction that for some $t^0 > 0$,

$$\sigma(t^0) = \sqrt{l \|\nabla u(t_0)\|_2^2} + (g \circ \nabla u)(t_0) > \sigma_2.$$

The continuity of $\sqrt{l \|\nabla u\|_2^2 + (g \circ \nabla u)(t)}$ illustrates that we may choose t^0 such that $\sigma_1 > \sigma(t^0) > \sigma_2$, then we have

$$E(0) = F(\sigma_2) < F(\sigma(t^0)) \le E(t^0).$$

This is a contradiction since Lemma 2.2.

Lemma 2.4. Under all the conditions of Lemma 2.3, there exists a positive constant \mathcal{D} such that for all $t \geq 0$,

$$||u||_p^p \le \mathcal{D}E(t) \le \mathcal{D}E(0), \tag{2.7}$$

$$\frac{1}{\rho+2} \|u_t\|_{\rho+2}^{\rho+2} + \frac{1}{2} \left(1 - \int_0^t g(s) ds\right) \|\nabla u\|_2^2 + \frac{1}{2} (g \circ \nabla u)(t) + \frac{1}{2} \|\nabla u_t\|_2^2 + \frac{\xi}{2} \int_\Omega \int_0^1 z^2(x, \kappa, t) d\kappa dx \le \mathcal{D}E(t) \le \mathcal{D}E(0).$$
(2.8)

Proof. Using the embedding $H_0^1(\Omega) \hookrightarrow L^p(\Omega)$, (2.2) and (2.5), we have

$$\begin{split} \frac{b}{p} \|u\|_{p}^{p} &\leq \frac{bB_{1}^{p}}{p} \left(l^{\frac{1}{2}} \|\nabla u\|_{2}\right)^{p} \leq \frac{bB_{1}^{p}}{p} \left(l\|\nabla u\|_{2}^{2} + (g \circ \nabla u)(t)\right)^{\frac{p-2}{2}} [l\|\nabla u\|_{2}^{2} + (g \circ \nabla u)(t)] \\ &\leq \frac{2bB_{1}^{p}}{p} \left(l\|\nabla u\|_{2}^{2} + (g \circ \nabla u)(t)\right)^{\frac{p-2}{2}} \left(E(t) + \frac{b}{p}\|u\|_{p}^{p}\right) \\ &\leq \frac{2bB_{1}^{p}}{p} \sigma_{2}^{p-2} \left(E(t) + \frac{b}{p}\|u\|_{p}^{p}\right), \end{split}$$

which yields (2.7) with

$$\mathcal{D} = \frac{2pB_1^p \sigma_2^{p-2}}{p - 2bB_1^p \sigma_2^{p-2}} > 0.$$
(2.9)

One has (2.8) by combining (2.7) with (2.2),

Lemma 2.5 (Lemma 4.1 in [2]). For $u \in H_0^1(\Omega)$, we have for all $t \ge 0$,

$$\int_{\Omega} \left(\int_{0}^{t} g(t-s)(\nabla u(s) - \nabla u(t)) ds \right)^{2} dx \le C_{\alpha}(h_{\alpha} \circ \nabla u)(t)$$
(2.10)

where, for any $0 < \alpha < 1$,

$$C_{\alpha} = \int_{0}^{\infty} \frac{g^{2}(s)}{\alpha g(s) - g'(s)} ds \text{ and } h_{\alpha}(t) = \alpha g(t) - g'(t).$$
(2.11)

Let us follow from the proof of Lemma 4.1 in [2], we have in fact

$$\int_{\Omega} \left(\int_0^t g(t-s)(u(s)-u(t))ds \right)^2 dx \le C_{\alpha}(h_{\alpha} \circ u)(t).$$
(2.12)

Lemma 2.6 (Lemma 2.2 in [6]). There exist positive constants γ and t_1 such that ()

$$g'(t) \le -\gamma g(t) \text{ for } t \in [0, t_1].$$
 (2.13)

Lemma 2.7. Let u be a solution of problem (2.1), then the functional

$$I_1(t) = \frac{1}{\rho+1} \int_{\Omega} |u_t|^{\rho} u_t u dx + \int_{\Omega} \nabla u_t \nabla u dx, \qquad (2.14)$$

satisfies, for $\varepsilon > 0$ and for all $t \ge 0$,

$$I_{1}'(t) \leq \frac{1}{\rho+1} \|u_{t}\|_{\rho+2}^{\rho+2} - \left[l - \left(1 + \frac{\mu_{1}}{\lambda_{1}} + \frac{\mu_{2}}{\lambda_{1}}\right)\varepsilon\right] \|\nabla u\|_{2}^{2} + \frac{1}{4\varepsilon} C_{\alpha}(h_{\alpha} \circ \nabla u)(t) + \frac{|\mu_{2}|}{4\varepsilon} \|z(x, 1, t)\|_{2}^{2} + b\|u\|_{p}^{p} + \|\nabla u_{t}\|_{2}^{2} + \frac{\mu_{1}}{4\varepsilon} \|u_{t}\|_{2}^{2}.$$

$$(2.15)$$

Proof. Multiplying the first identity in problem (2.1) by u, integrating on x over $\Omega,$ and then using integration by parts, we give

$$\int_{\Omega} |u_t|^{\rho} u_{tt} u dx + \int_{\Omega} \nabla u_{tt} \nabla u dx$$

= $-\left(1 - \int_0^t g(s) ds\right) \|\nabla u\|_2^2 + \int_{\Omega} \nabla u(t) \int_0^t g(t-s) \nabla (u(s) - u(t)) ds$ (2.16)
 $- \int_{\Omega} \mu_1 u_t(x,t) u dx - \int_{\Omega} \mu_2 z(x,1,t) u dx + b \|u\|_p^p.$

Differentiating (2.14) on t, and using (2.16), one has

$$I_{1}'(t) = \frac{1}{\rho+1} \|u_{t}\|_{\rho+2}^{\rho+2} + \int_{\Omega} |u_{t}|^{\rho} u_{tt} u dx + \int_{\Omega} \nabla u_{tt} \nabla u dx + \|\nabla u_{t}\|_{2}^{2}$$

$$= \frac{1}{\rho+1} \|u_{t}\|_{\rho+2}^{\rho+2} - \left(1 - \int_{0}^{t} g(s) ds\right) \|\nabla u\|_{2}^{2}$$

$$+ \int_{\Omega} \nabla u(t) \int_{0}^{t} g(t-s) \nabla (u(s) - u(t)) ds$$

$$- \int_{\Omega} \mu_{1} u_{t}(x, t) u dx - \int_{\Omega} \mu_{2} z(x, 1, t) u dx + b \|u\|_{p}^{p} + \|\nabla u_{t}\|_{2}^{2}.$$
(2.17)

Applying Cauchy's inequality with $\varepsilon > 0$ and $\lambda_1 ||u||_2^2 \le ||\nabla u||_2^2$, it follows that

$$-\int_{\Omega} \mu_1 u_t(x,t) u dx \le \frac{\mu_1}{4\varepsilon} \|u_t\|_2^2 + \mu_1 \varepsilon \|u\|_2^2 \le \frac{\mu_1}{4\varepsilon} \|u_t\|_2^2 + \frac{\mu_1 \varepsilon}{\lambda_1} \|\nabla u\|_2^2, \qquad (2.18)$$

6

$$-\int_{\Omega} \mu_2 z(x,1,t) u dx \le \frac{|\mu_2|}{4\varepsilon} \|z(x,1,t)\|_2^2 + \frac{|\mu_2|\varepsilon}{\lambda_1} \|\nabla u\|_2^2.$$
(2.19)

It follows from Cauchy's inequality with $\varepsilon>0$ and (2.10) that

$$\int_{\Omega} \nabla u(t) \int_{0}^{t} g(t-s)(\nabla u(s) - \nabla u(t)) ds \leq \varepsilon \|\nabla u\|_{2}^{2} + \frac{1}{4\varepsilon} C_{\alpha}(h \circ \nabla u)(t). \quad (2.20)$$

erting (2.18)-(2.20) into (2.17), we obtain (2.15).

Inserting (2.18)-(2.20) into (2.17), we obtain (2.15).

Lemma 2.8. Under all the conditions of Lemma 2.3, let u be a solution of problem (2.1), then the functional

$$I_{2}(t) = \int_{\Omega} \left(\Delta u_{t} - \frac{1}{\rho+1} |u_{t}|^{\rho} u_{t} \right) \int_{0}^{t} g(t-s)(u(t) - u(s)) ds dx,$$
(2.21)

satisfies, for $\delta > 0$ and for all $t \ge 0$,

$$I_{2}'(t) \leq B_{1} \|\nabla u\|_{2}^{2} + B_{2}(h_{\alpha} \circ \nabla u)(t) + \left[B_{3} - \int_{0}^{t} g(s)ds\right] \|\nabla u_{t}\|_{2}^{2} + \delta \|z(x, 1, t)\|_{2}^{2} - \int_{0}^{t} g(s)ds \cdot \frac{1}{\rho + 1} \|u_{t}\|_{\rho + 2}^{\rho + 2},$$

$$(2.22)$$

here B_1 , B_2 and B_3 are positive constants depending on δ shown in (2.31).

Proof. Differentiating (2.21) on t, and using the first identity in problem (2.1) and integration by parts, we have

$$\begin{split} I_{2}'(t) &= \int_{\Omega} \left(\Delta u_{tt} - |u_{t}|^{\rho} u_{tt} \right) \int_{0}^{t} g(t-s)(u(t) - u(s)) ds dx \\ &+ \int_{\Omega} \left(\Delta u_{t} - \frac{1}{\rho+1} |u_{t}|^{\rho} u_{t} \right) \int_{0}^{t} g_{t}(t-s)(u(t) - u(s)) ds dx \\ &+ \int_{\Omega} \left(\Delta u_{t} - \frac{1}{\rho+1} |u_{t}|^{\rho} u_{t} \right) \int_{0}^{t} g(t-s) u_{t}(t) ds dx \\ &= \int_{\Omega} \nabla u(t) \int_{0}^{t} g(t-s) \nabla (u(t) - u(s)) ds \\ &- \int_{\Omega} \int_{0}^{t} g(t-s) \nabla u(s) ds \int_{0}^{t} g(t-s) \nabla (u(t) - u(s)) ds dx \\ &+ \int_{\Omega} \mu_{1} u_{t}(x,t) \int_{0}^{t} g(t-s)(u(t) - u(s)) ds dx \\ &+ \int_{\Omega} \mu_{2} z(x,1,t) \int_{0}^{t} g(t-s)(u(t) - u(s)) ds dx \\ &- b \int_{\Omega} |u|^{p-2} u \int_{0}^{t} g(t-s) \nabla (u(t) - u(s)) ds dx \\ &- \int_{\Omega} \nabla u_{t} \int_{0}^{t} g_{t}(t-s) \nabla (u(t) - u(s)) ds dx \\ &- \int_{\Omega} \frac{1}{\rho+1} |u_{t}|^{\rho} u_{t} \int_{0}^{t} g_{t}(t-s)(u(t) - u(s)) ds dx \\ &- \int_{0}^{t} g(s) ds \cdot ||\nabla u_{t}||_{2}^{2} - \int_{0}^{t} g(s) ds \cdot \frac{1}{\rho+1} ||u_{t}||_{\rho+2}^{\rho+2} \\ &= J_{1} + J_{2} + \dots + J_{8} + J_{9}. \end{split}$$

 $\overline{7}$

It is direct from Cauchy's inequality with $\delta > 0$ and (2.10) that

$$J_{1} \leq \delta \|\nabla u\|_{2}^{2} + \frac{1}{4\delta} C_{\alpha}(h_{\alpha} \circ \nabla u)(t), \qquad (2.24)$$

$$J_{2} \leq \delta \int_{\Omega} \left(\int_{0}^{t} g(t-s) |\nabla u(s) - \nabla u(t)| + |\nabla u(t)| ds \right)^{2} dx$$

$$+ \frac{1}{4\delta} \int_{\Omega} \left(\int_{0}^{t} g(t-s) |\nabla u(s) - \nabla u(t)| ds \right)^{2} dx$$

$$\leq 2\delta \int_{\Omega} \left(\int_{0}^{t} g(t-s) |\nabla u(s) - \nabla u(t)| ds \right)^{2} dx + 2\delta(1-l)^{2} \|\nabla u(t)\|_{2}^{2} \qquad (2.25)$$

$$+ \frac{1}{4\delta} \int_{\Omega} \left(\int_{0}^{t} g(t-s) |\nabla u(s) - \nabla u(t)| ds \right)^{2} dx$$

$$\leq \left(2\delta + \frac{1}{4\delta} \right) C_{\alpha}(h_{\alpha} \circ \nabla u)(t) + 2\delta(1-l)^{2} \|\nabla u\|_{2}^{2}.$$

Cauchy's inequality with $\delta > 0$ and (2.12) yield

$$J_3 \leq \delta \|u_t\|_2^2 + \frac{\mu_1^2}{4\delta} C_\alpha(h_\alpha \circ u)(t) \leq \frac{\delta}{\lambda_1} \|\nabla u_t\|_2^2 + \frac{\mu_1^2}{4\delta\lambda_1} C_\alpha(h_\alpha \circ \nabla u)(t), \qquad (2.26)$$

$$J_4 \le \delta \|z(x, 1, t)\|_2^2 + \frac{\mu_2^2}{4\delta\lambda_1} C_\alpha(h_\alpha \circ \nabla u)(t).$$
(2.27)

It follows from Cauchy's inequality with $\delta > 0$, (2.12), the embedding $H_0^1(\Omega) \hookrightarrow L^{2(p-1)}(\Omega)$ and (2.8) that

$$J_{5} \leq b\delta \|u\|_{2(p-1)}^{2(p-1)} + \frac{b}{4\delta} \int_{\Omega} \left(\int_{0}^{t} g(t-s)(u(t)-u(s))ds \right)^{2} dx$$

$$\leq b\delta \|u\|_{2(p-1)}^{2(p-1)} + \frac{b}{4\delta} C_{\alpha}(h_{\alpha} \circ u)(t)$$

$$\leq b\delta c_{s}^{2(p-1)} \left(\frac{2\mathcal{D}}{l} E(0)\right)^{p-2} \|\nabla u\|_{2}^{2} + \frac{b}{4\delta\lambda_{1}} C_{\alpha}(h_{\alpha} \circ \nabla u)(t).$$

(2.28)

Recalling the definition of g'(t) in (2.11), and using Cauchy's inequality with $\delta > 0$, (2.10) and Hölder's inequality, one obtains

$$J_{6} = -\int_{\Omega} \nabla u_{t} \int_{0}^{t} \alpha g(t-s) \nabla (u(t) - u(s)) ds dx + \int_{\Omega} \nabla u_{t} \int_{0}^{t} h_{\alpha}(t-s) \nabla (u(t) - u(s)) ds dx \leq \delta \|\nabla u_{t}\|_{2}^{2} + \frac{\alpha^{2}}{4\delta} \int_{\Omega} \left(\int_{0}^{t} g(t-s) \nabla (u(t) - u(s)) ds \right)^{2} dx + \delta \|\nabla u_{t}\|_{2}^{2} + \frac{1}{4\delta} \int_{\Omega} \left(\int_{0}^{t} h_{\alpha}(t-s) \nabla (u(t) - u(s)) ds \right)^{2} dx$$
(2.29)
$$\leq 2\delta \|\nabla u_{t}\|_{2}^{2} + \frac{\alpha^{2}}{4\delta} C_{\alpha} (h_{\alpha} \circ \nabla u)(t) + \frac{1}{4\delta} \int_{0}^{t} h_{\alpha}(s) ds \int_{0}^{t} h_{\alpha}(t-s) \|\nabla (u(t) - u(s))\|_{2}^{2} ds \leq 2\delta \|\nabla u_{t}\|_{2}^{2} + \left(\frac{\alpha^{2}}{4\delta} C_{\alpha} + \frac{\alpha(1-l) + g(0)}{4\delta}\right) (h_{\alpha} \circ \nabla u)(t).$$

Similarly, we get

$$J_{7} = -\int_{\Omega} \frac{1}{\rho+1} |u_{t}|^{\rho} u_{t} \int_{0}^{t} \alpha g(t-s)(u(t)-u(s)) ds dx + \int_{\Omega} \frac{1}{\rho+1} |u_{t}|^{\rho} u_{t} \int_{0}^{t} h_{\alpha}(t-s)(u(t)-u(s)) ds dx \leq \frac{\delta}{\rho+1} ||u_{t}||^{2(\rho+1)}_{2(\rho+1)} + \frac{\alpha^{2}}{4(\rho+1)\delta} \int_{\Omega} \left(\int_{0}^{t} g(t-s)(u(t)-u(s)) ds \right)^{2} dx + \frac{\delta}{\rho+1} ||u_{t}||^{2(\rho+1)}_{2(\rho+1)} + \frac{1}{4(\rho+1)\delta} \int_{\Omega} \left(\int_{0}^{t} h_{\alpha}(t-s)(u(t)-u(s)) ds \right)^{2} dx \leq \frac{2\delta}{\rho+1} ||u_{t}||^{2(\rho+1)}_{2(\rho+1)} + \frac{\alpha^{2}}{4(\rho+1)\delta} C_{\alpha}(h_{\alpha} \circ u)(t) + \frac{1}{4(\rho+1)\delta} \int_{0}^{t} h_{\alpha}(s) ds \int_{0}^{t} h_{\alpha}(t-s) ||u(t)-u(s)||^{2}_{2} ds \leq \frac{2\delta}{\rho+1} c_{s}^{2(\rho+1)} \left(2\mathcal{D}E(0) \right)^{\frac{\rho}{2}} ||\nabla u_{t}||^{2}_{2} + \left(\frac{\alpha^{2}}{4(\rho+1)\delta} C_{\alpha} + \frac{\alpha(1-l)+g(0)}{4(\rho+1)\delta} \right) \frac{1}{\lambda_{1}} (h_{\alpha} \circ \nabla u)(t).$$

$$(2.30)$$

Inserting (2.25)-(2.30) into (2.23), one has

$$I_{2}'(t) \leq B_{1} \|\nabla u\|_{2}^{2} + B_{2}(h_{\alpha} \circ \nabla u)(t) + \left[B_{3} - \int_{0}^{t} g(s)ds\right] \|\nabla u_{t}\|_{2}^{2} + \delta \|z(x, 1, t)\|_{2}^{2} - \int_{0}^{t} g(s)ds \cdot \frac{1}{\rho + 1} \|u_{t}\|_{\rho + 2}^{\rho + 2},$$

with

$$\begin{cases} B_1 = \delta + 2\delta(1-l)^2 + b\delta c_s^{2(p-1)} \left(\frac{2\mathcal{D}}{l}E(0)\right)^{p-2}; \\ B_2 = \left[\frac{1}{2\delta} + 2\delta + \frac{\mu_1^2}{4\delta\lambda_1} + \frac{\mu_2^2}{4\delta\lambda_1} + \frac{b}{4\delta\lambda_1} + \frac{\alpha^2}{4\delta} + \frac{\alpha^2}{4(\rho+1)\delta\lambda_1}\right] C_{\alpha} \\ + \frac{\alpha(1-l)+g(0)}{4\delta} + \frac{\alpha(1-l)+g(0)}{4(\rho+1)\delta\lambda_1}; \\ B_3 = \frac{\delta}{\lambda_1} + 2\delta + \frac{2\delta}{\rho+1} c_s^{2(\rho+1)} \left(2\mathcal{D}E(0)\right)^{\frac{\rho}{2}}. \end{cases}$$
(2.31)

Lemma 2.9. The functional

$$I_{3}(t) = \int_{0}^{1} e^{-2\tau\kappa} \|z(x,\kappa,t)\|_{2}^{2} d\kappa$$
(2.32)

satisfies for all $t \ge 0$,

$$I'_{3}(t) = -2I_{3}(t) + \frac{1}{\tau} \|u_{t}\|_{2}^{2} - \frac{e^{-2\tau}}{\tau} \|z(x, 1, t)\|_{2}^{2}.$$
 (2.33)

Proof. Differentiating (2.32) on t, and using the second identity in (2.1), it is direct that

$$I'_{3}(t) = \int_{0}^{1} 2e^{-2\tau\kappa} \int_{\Omega} z(x,\kappa,t)z_{t}(x,\kappa,t)dxd\kappa$$

$$= -\int_{0}^{1} \frac{2}{\tau}e^{-2\tau\kappa} \int_{\Omega} z(x,\kappa,t)z_{k}(x,\kappa,t)dxd\kappa$$

$$= -\frac{1}{\tau} \int_{\Omega} \int_{0}^{1} \left(\frac{d}{d\kappa}e^{-2\tau\kappa}z^{2}(x,\kappa,t) + 2\tau e^{-2\tau\kappa}z^{2}(x,\kappa,t)\right)d\kappa dx$$

$$= -2I_{3}(t) + \frac{1}{\tau} ||u_{t}||_{2}^{2} - \frac{e^{-2\tau}}{\tau} ||z(x,1,t)||_{2}^{2}.$$

$$(2.34)$$

Lemma 2.10 (Lemma 3.4 in [17]). The functional

$$I_4(t) = \int_0^t f(t-s) \|\nabla u(s)\|_2^2 ds$$
(2.35)

satisfies for all $t \ge 0$,

$$I'_{4}(t) \le 3(1-l) \|\nabla u\|_{2}^{2} - \frac{1}{2}(g \circ \nabla u)(t), \qquad (2.36)$$

where $f(t) = \int_t^\infty g(s) ds$.

3. Stability results

In this section, we will present and prove the decay results of the energy functional E(t) based on the lemmas in Section 2. To begin with, we define a functional

$$L(t) = ME(t) + \sum_{i=1}^{3} N_i I_i(t), \qquad (3.1)$$

where M, N_1 , N_2 , N_3 are positive constants. The following lemma is shown to illustrate that L(t) is equivalent to E(t).

Lemma 3.1. Under all the conditions of Lemma 2.3, assume that M is enough large, then there exist two positive constants β_1 and β_2 such that

$$\beta_1 E(t) \le L(t) \le \beta_2 E(t). \tag{3.2}$$

Proof. Recalling the definition of $I_1(t)$ in (2.14), using Young's inequality and Cauchy's inequality, and then applying the embedding $H_0^1(\Omega) \hookrightarrow L^{\rho+2}(\Omega)$ and (2.8), it is not hard to give

$$|I_{1}(t)| \leq \frac{1}{\rho+2} \|u_{t}\|_{\rho+2}^{\rho+2} + \frac{1}{(\rho+1)(\rho+2)} \|u\|_{\rho+2}^{\rho+2} + \frac{1}{2} \|\nabla u_{t}\|_{2}^{2} + \frac{1}{2} \|\nabla u\|_{2}^{2}$$

$$\leq \frac{1}{\rho+2} \|u_{t}\|_{\rho+2}^{\rho+2} + \left[\frac{c_{s}^{\rho+2}}{(\rho+1)(\rho+2)} \left(\frac{2\mathcal{D}}{l}E(0)\right)^{\frac{\rho}{2}} + \frac{1}{2}\right] \|\nabla u\|_{2}^{2} + \frac{1}{2} \|\nabla u_{t}\|_{2}^{2}.$$

$$(3.3)$$

Recalling the definition of $I_2(t)$ in (2.21), using integration by parts, Cauchy's inequality, Young's inequality and Hölder's inequality, we give

$$\begin{aligned} |I_{2}(t)| &\leq \frac{1}{2} \|\nabla u_{t}\|_{2}^{2} + \frac{1}{2} \int_{\Omega} \left(\int_{0}^{t} g(t-s)(\nabla u(t) - \nabla u(s))ds \right)^{2} dx \\ &+ \frac{1}{\rho+2} \|u_{t}\|_{\rho+2}^{\rho+2} + \frac{1}{(\rho+1)(\rho+2)} \int_{\Omega} \left(\int_{0}^{t} g(t-s)(u(t) - u(s))ds \right)^{\rho+2} dx \\ &\leq \frac{1}{2} \|\nabla u_{t}\|_{2}^{2} + \frac{1-l}{2} (g \circ \nabla u)(t) + \frac{1}{\rho+2} \|u_{t}\|_{\rho+2}^{\rho+2} \\ &+ \frac{1}{(\rho+1)(\rho+2)} (1-l)^{\rho+1} c_{s}^{\rho+2} \left(\frac{2\mathcal{D}}{l} E(0) \right)^{\frac{\rho}{2}} (g \circ \nabla u)(t), \end{aligned}$$
(3.4)

here we have used

$$\int_{\Omega} \left(\int_{0}^{t} g(t-s)(u(t)-u(s))ds \right)^{\rho+2} dx \\
\leq \int_{\Omega} \left(\int_{0}^{t} (g(t-s))^{\frac{\rho+1}{\rho+2}} (g(t-s))^{\frac{1}{\rho+2}} (u(t)-u(s))ds \right)^{\rho+2} dx \\
\leq \left(\int_{0}^{t} g(s)ds \right)^{\rho+1} \int_{0}^{t} g(t-s) \|u(t)-u(s)\|_{\rho+2}^{\rho+2} ds \\
\leq (1-l)^{\rho+1} c_{s}^{\rho+2} \int_{0}^{t} g(t-s) \|\nabla u(t)-\nabla u(s)\|_{2}^{\rho+2} ds \\
\leq (1-l)^{\rho+1} c_{s}^{\rho+2} \left(\frac{2\mathcal{D}}{l} E(0) \right)^{\frac{\rho}{2}} (g \circ \nabla u)(t).$$
(3.5)

Therefore, it follows from (2.8) that

$$|L(t) - ME(t)| = |\sum_{i=1}^{3} N_i I_i(t)| \le CE(t),$$

here C is some positive constant.

Lemma 3.2. Under all the conditions of Lemma 2.3, for $|\mu_2| < \mu_1$, the functional L(t) defined in (3.1) satisfies, for $t \ge t_1$

$$L'(t) \leq -C_1 \|u_t\|_{\rho+2}^{\rho+2} - C_2 \|\nabla u_t\|_2^2 - 4(1-l) \|\nabla u\|_2^2 + N_1 b \|u\|_p^p + \frac{1}{4} (g \circ \nabla u)(t) - 2N_3 \int_0^1 \|z(x,\kappa,t)\|_2^2 d\kappa.$$
(3.6)

where C_1 , C_2 are positive constants given in (3.8).

Proof. Taking the combination of (2.4), (2.15) and (2.22) with (2.33), recalling (2.11), and applying $g_1 = \int_0^{t_1} g(s) ds \leq \int_0^t g(s) ds$ for $t \geq t_1$, one has

$$\begin{split} L'(t) &\leq -M\omega(\|u_t\|_2^2 + \|z(x,1,t)\|_2^2) + \frac{M}{2}(g' \circ \nabla u)(t) - \frac{M}{2}g(t)\|\nabla u\|_2^2 \\ &+ \frac{N_1}{\rho+1}\|u_t\|_{\rho+2}^{\rho+2} - N_1\Big[l - \Big(1 + \frac{\mu_1}{\lambda_1} + \frac{\mu_2}{\lambda_1}\Big)\varepsilon\Big]\|\nabla u\|_2^2 + \frac{N_1}{4\varepsilon}C_{\alpha}(h_{\alpha} \circ \nabla u)(t) \\ &+ \frac{N_1|\mu_2|}{4\varepsilon}\|z(x,1,t)\|_2^2 + N_1b\|u\|_p^{\rho} + N_1\|\nabla u_t\|_2^2 + N_1\frac{\mu_1}{4\varepsilon}\|u_t\|_2^2 \\ &+ N_2B_1\|\nabla u\|_2^2 + N_2B_2(h_{\alpha} \circ \nabla u)(t) + N_2\Big[B_3 - \int_0^t g(s)ds\Big]\|\nabla u_t\|_2^2 \\ &+ N_2\delta\|z(x,1,t)\|_2^2 - N_2\int_0^t g(s)ds \cdot \frac{1}{\rho+1}\|u_t\|_{\rho+2}^{\rho+2} \\ &- 2N_3I_3(t) + \frac{N_3}{\tau}\|u_t\|_2^2 - \frac{N_3e^{-2\tau}}{\tau}\|z(x,1,t)\|_2^2 \\ &\leq -C_1\|u_t\|_{\rho+2}^{\rho+2} - C_2\|\nabla u_t\|_2^2 - C_3\|\nabla u\|_2^2 - C_4\|z(x,1,t)\|_2^2 - C_5(h_{\alpha} \circ \nabla u)(t) \\ &+ N_1b\|u\|_p^p + \frac{\alpha M}{2}(g \circ \nabla u)(t) - C_6\|u_t\|_2^2 - 2N_3\int_0^1\|z(x,\kappa,t)\|_2^2d\kappa \end{split}$$

with

$$\begin{cases} C_{1} = N_{2}g_{1} \cdot \frac{1}{\rho+1} - \frac{N_{1}}{\rho+1}; \\ C_{2} = N_{2} \Big[g_{1} - \Big(\frac{\delta}{\lambda_{1}} + 2\delta + \frac{2\delta}{\rho+1}c_{s}^{2(\rho+1)} \Big(2\mathcal{D}E(0) \Big)^{\frac{\rho}{2}} \Big) \Big] - N_{1}; \\ C_{3} = N_{1} \Big[l - \Big(1 + \frac{\mu_{1}}{\lambda_{1}} + \frac{|\mu_{2}|}{\lambda_{1}} \Big) \varepsilon \Big] - N_{2} \Big[\delta + 2\delta(1-l)^{2} + b\delta c_{s}^{2(\rho-1)} \Big(\frac{2\mathcal{D}}{l}E(0) \Big)^{p-2} \Big]; \\ C_{4} = \omega M + N_{3} \frac{e^{-2\tau}}{\tau} - N_{1} \frac{|\mu_{2}|}{4\varepsilon} - N_{2}\delta; \\ C_{5} = \frac{M}{2} - N_{1} \frac{1}{4\varepsilon}C_{\alpha} - N_{2} \Big[\Big(\frac{1}{2\delta} + 2\delta + \frac{\mu_{1}^{2}}{4\delta\lambda_{1}} + \frac{\mu_{2}^{2}}{4\delta\lambda_{1}} + \frac{b}{4\delta\lambda_{1}} + \frac{\alpha^{2}}{4\delta} + \frac{\alpha^{2}}{4(\rho+1)\delta\lambda_{1}} \Big) C_{\alpha}; \\ + \frac{\alpha(1-l)+g(0)}{4\delta} + \frac{\alpha(1-l)+g(0)}{4(\rho+1)\delta\lambda_{1}} \Big]; \\ C_{6} = \omega M - \frac{N_{3}}{\tau} - N_{1} \frac{\mu_{1}}{4\varepsilon}, \end{cases}$$

$$(3.8)$$

where we have used the values of B_1, B_2 and B_3 defined in (2.31). Next, we choose δ such that

$$\delta < \left\{ \frac{lg_1}{16 \left[1 + 2(1-l)^2 + bc_s^{2(p-1)} \left(\frac{2\mathcal{D}}{l} E(0) \right)^{p-2} \right]}, \frac{lg_1}{1024(1-l)^2}, \frac{5g_1}{8 \left(\frac{1}{\lambda_1} + 2 + \frac{2}{\rho+1} c_s^{2(\rho+1)} \left(2\mathcal{D}E(0) \right)^{\frac{\rho}{2}} \right)} \right\}.$$

Let us choose $N_1 = \frac{3}{8}g_1N_2$, then

$$C_1 = N_2 g_1 \cdot \frac{1}{\rho+1} - \frac{3}{8} g_1 N_2 \frac{1}{\rho+1} = \frac{5}{8} g_1 N_2 \frac{1}{\rho+1} > 0, \quad C_2 > 0.$$

Let us fix

$$\varepsilon = \frac{3l}{4} \frac{1}{1 + \frac{\mu_1}{\lambda_1} + \frac{|\mu_2|}{\lambda_1}},$$

then

$$C_3 = \frac{N_1 l}{4} - N_2 \left[\delta + 2\delta (1-l)^2 + b\delta c_s^{2(p-1)} \left(\frac{2\mathcal{D}}{l} E(0) \right)^{p-2} \right] > \frac{l}{32} g_1 N_2 > 0.$$

By taking $N_2 = \frac{1}{8\delta(1-l)}$, we get

$$C_3 > \frac{l}{32}g_1N_2 = \frac{lg_1}{256\delta(1-l)} > 4(1-l).$$

Since $g'(s) \leq 0$, one has $\frac{\alpha g^2(s)}{\alpha g(s) - g'(s)} \leq g(s)$, further we get

$$\lim_{\alpha \to 0^+} \alpha C_{\alpha} = \lim_{\alpha \to 0^+} \int_0^\infty \frac{\alpha g^2(s)}{\alpha g(s) - g'(s)} ds = 0.$$

Thus, there exists $0 < \alpha_0 < 1$ so that if $\alpha < \alpha_0$, then

$$\alpha C_{\alpha} < \frac{1}{8\left[N_2\left(\frac{1}{2\delta} + 2\delta + \frac{\mu_1^2}{4\delta\lambda_1} + \frac{\mu_2^2}{4\delta\lambda_1} + \frac{b}{4\delta\lambda_1} + \frac{\alpha^2}{4\delta} + \frac{\alpha^2}{4(\rho+1)\delta\lambda_1}\right) + N_1\frac{1}{4\varepsilon}\right]}.$$

Let us choose M sufficiently large such that for $\alpha = \frac{1}{2M}$,

$$\begin{split} C_5 &= \frac{M}{4} - N_2 \Big[\frac{\alpha (1-l) + g(0)}{4\delta} + \frac{\alpha (1-l) + g(0)}{4(\rho+1)\delta\lambda_1} \Big] > 0 \\ C_4 &= \omega M + N_3 \frac{e^{-2\tau}}{\tau} - N_1 \frac{|\mu_2|}{4\varepsilon} - N_2 \delta > 0, \\ C_6 &= \omega M - \frac{N_3}{\tau} - N_1 \frac{\mu_1}{4\varepsilon} > 0, \end{split}$$

here we have used $\omega > 0$ given in Lemma 2.2 since $|\mu_2| < \mu_1$. Based on the above discussion, one has from (3.7)

$$L'(t) \leq -C_1 \|u_t\|_{\rho+2}^{\rho+2} - C_2 \|\nabla u_t\|_2^2 - 4(1-l) \|\nabla u\|_2^2 + N_1 b \|u\|_p^p + \frac{1}{4} (g \circ \nabla u)(t) - 2N_3 \int_0^1 \|z(x,\kappa,t)\|_2^2 d\kappa.$$

Now, we give the following stability results.

Theorem 3.3. For $|\mu_2| < \mu_1$, suppose that $(\mathbf{H_1})$, $(\mathbf{H_2})$ hold and

$$E(0) < E_1, \quad l \|\nabla u_0\|_2^2 < \sigma_1^2,$$

then there exist positive constants k_1, k_2, k_3 and k_4 such that the solution of problem (1.1) satisfies for all $t \ge t_1$,

$$E(t) \leq \begin{cases} k_1 e^{-k_2 \int_{t_1}^t \zeta(s) ds} & \text{for } G \text{ is linear;} \\ k_4 G_1^{-1} \left(k_3 \int_{t_1}^t \zeta(s) ds \right) & \text{for } G \text{ is nonlinear,} \end{cases}$$

where E_1 and σ_1 are shown in Lemma 2.3, $G_1(t) = \int_t^r \frac{1}{sG'(s)} ds$ is strictly decreasing and convex in (0, r] with $\lim_{t\to 0} G_1(t) = +\infty$.

Proof. Using (2.13) and (2.4), one has, for $t \ge t_1$,

$$\int_{0}^{t_{1}} g(s) \|\nabla u(t) - \nabla u(t-s)\|_{2}^{2} ds
\leq -\frac{1}{\gamma} \int_{0}^{t_{1}} g'(s) \|\nabla u(t) - \nabla u(t-s)\|_{2}^{2} ds \leq -cE'(t)$$
(3.9)

here c is used to denote a generic positive constant throughout this proof. Define a functional F(t) that is obviously equivalent to E(t) as follows

$$F(t) = L(t) + cE(t),$$

then based on (3.6), (2.2), (3.9), for some m > 0 and for any $t \ge t_1$, we have

$$F'(t) \leq -C_1 \|u_t\|_{p+2}^{p+2} - C_2 \|\nabla u_t\|_2^2 - 4(1-l) \|\nabla u\|_2^2 + N_1 b \|u\|_p^p + \frac{1}{4} (g \circ \nabla u)(t) - 2N_3 \int_0^1 \|z(x,\kappa,t)\|_2^2 d\kappa + cE'(t) \leq -mE(t) - \left(\frac{bc}{p} - N_1 b\right) \|u\|_p^p + c(g \circ \nabla u)(t) + cE'(t) \leq -mE(t) + c \int_{t_1}^t g(s) \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds,$$
(3.10)

here we have chosen N_1 so small that $\frac{bc}{p} - N_1 b > 0$. In what follows, we will discuss in two cases.

CASE 1: G IS LINEAR Multiplying (3.10) by $\zeta(t)$, using (H₁) and (2.4), one gives

$$\begin{split} \zeta(t)F'(t) &\leq -m\zeta(t)E(t) + c\zeta(t) \int_{t_1}^t g(s) \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \\ &\leq -m\zeta(t)E(t) - c \int_{t_1}^t g'(s) \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \\ &\leq -m\zeta(t)E(t) - cE'(t), \end{split}$$

which implies

$$(\zeta(t)F(t) + cE(t))' \leq -m\zeta(t)E(t)$$
 for $t \geq t_1$

Integrating the above inequality over (t_1, t) , and using the fact that $\zeta(t)F(t) + cE(t)$ is equivalent to E(t), one has

$$E(t) \le k_1 e^{-k_2 \int_{t_1}^t \zeta(s) ds} \quad \text{for } t \ge t_1,$$

here k_1 and k_2 are constants.

CASE 2: G is nonlinear Define a functional

$$H(t) = L(t) + I_4(t).$$

Taking the combination of Lemma 3.1 and the non-negativity of E(t) obtained by Lemma 2.4 with the definition of $I_4(t)$ in (2.35), it is not difficult to get the non-negativity of H(t). It follows from (3.6) and (2.36) that for some $m_1 > 0$ and $t \geq t_1$,

$$\begin{aligned} H'(t) &= L'(t) + I'_4(t) \\ &\leq -C_1 \|u_t\|_{\rho+2}^{\rho+2} - C_2 \|\nabla u_t\|_2^2 - (1-l) \|\nabla u\|_2^2 \\ &+ N_1 b \|u\|_p^p - \frac{1}{4} (g \circ \nabla u)(t) - 2N_3 \int_0^1 \|z(x,\kappa,t)\|_2^2 d\kappa \leq -m_1 E(t). \end{aligned}$$

Integrating the above inequality over (t_1, t) yields

$$m_1 \int_{t_1}^t E(s) ds \le H(t_1) - H(t) \le H(t_1),$$

which implies

$$\int_0^\infty E(s)ds < +\infty. \tag{3.11}$$

Define

$$\lambda(t) = p \int_{t_1}^t \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds,$$

by using (2.8), then we give

$$\begin{split} \lambda(t) &\leq 2p \int_0^t \left(\|\nabla u(t)\|_2^2 + \|\nabla u(t-s)\|_2^2 \right) ds \\ &\leq \frac{8p\mathcal{D}}{l} \int_0^t \left(E(t) + E(t-s) \right) ds \leq \frac{16p\mathcal{D}}{l} \int_0^t E(t-s) ds \\ &= \frac{16p\mathcal{D}}{l} \int_0^t E(s) ds \leq \int_0^\infty E(s) ds < +\infty. \end{split}$$

Thus, we can choose p so small that for $t \geq t_1$,

$$\lambda(t) < 1. \tag{3.12}$$

It is direct that

$$G(\theta z) \le \theta G(z) \quad \text{for } 0 \le \theta \le 1 \text{ and } z \in (0, r],$$
(3.13)

since G is strictly convex on (0, r] and G(0) = 0. Based on (1.3), (3.12), (3.13) and Jensen's inequality, one gives

$$\begin{split} I(t) &= \frac{1}{p\lambda(t)} \int_{t_1}^t \lambda(t)(-g'(s))p \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \\ &\geq \frac{1}{p\lambda(t)} \int_{t_1}^t \lambda(t)\zeta(s)G(g(s))p \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \\ &\geq \frac{\zeta(t)}{p\lambda(t)} \int_{t_1}^t \overline{G}(\lambda(t)g(s))p \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \\ &\geq \frac{\zeta(t)}{p} \overline{G}\Big(p \int_{t_1}^t g(s) \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds\Big), \end{split}$$

which yields

$$\int_{t_1}^t g(s) \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \le \frac{1}{p} \overline{G}^{-1} \Big(\frac{pI(t)}{\zeta(t)}\Big), \tag{3.14}$$

where G has an extension \overline{G} which is a strictly increasing and strictly convex C^2 function on $(0, +\infty)$ as in Remark 2.1 [6]. Therefore, (3.10) becomes

$$F'(t) \le -mE(t) + \frac{c}{p}\overline{G}^{-1}\left(\frac{pI(t)}{\zeta(t)}\right).$$
(3.15)

Let us define the functional

$$F_1(t) = \overline{G}'\left(\frac{r_1 E(t)}{E(0)}\right)F(t) + E(t)$$

with $0 < r_1 < r$, then F_1 is equivalent to E and

$$F_{1}'(t) = \frac{r_{1}E'(t)}{E(0)}\overline{G}''\left(\frac{r_{1}E(t)}{E(0)}\right)F(t) + \overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right)F'(t) + E'(t)$$

$$\leq -mE(t)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + \frac{c}{p}\overline{G}^{-1}\left(\frac{pI(t)}{\zeta(t)}\right)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + E'(t)$$
(3.16)

by using (3.15), (2.4), G' > 0 and G'' > 0. Let \overline{G}^* be the convex conjugate of G in the sense of Young in [1], which is given by

$$\overline{G}^*(s) = s(\overline{G}')^{-1}(s) - \overline{G}\left[(\overline{G}')^{-1}(s)\right]$$
(3.17)

and it satisfies the following Young's inequality

$$AB \le \overline{G}^*(A) + \overline{G}(B). \tag{3.18}$$

Choosing

$$A = \overline{G}'\left(\frac{r_1 E(t)}{E(0)}\right) \text{ and } B = \overline{G}^{-1}\left(\frac{pI(t)}{\zeta(t)}\right),$$

then using (3.18), (3.17) and the non-negativity of \overline{G} , (3.16) becomes

$$F_{1}'(t) \leq -mE(t)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + \frac{c}{p}\overline{G}^{-1}\left(\frac{pI(t)}{\zeta(t)}\right)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + E'(t)$$

$$\leq -mE(t)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + \frac{c}{p}\overline{G}^{*}\left(\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right)\right) + c\frac{I(t)}{\zeta(t)} + E'(t)$$

$$\leq -mE(t)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + \frac{c}{p}\frac{r_{1}E(t)}{E(0)}\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) \qquad (3.19)$$

$$-\overline{G}\left(\frac{r_{1}E(t)}{E(0)}\right) + c\frac{I(t)}{\zeta(t)} + E'(t)$$

$$\leq -mE(t)\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + \frac{c}{p}\frac{r_{1}E(t)}{E(0)}\overline{G}'\left(\frac{r_{1}E(t)}{E(0)}\right) + c\frac{I(t)}{\zeta(t)} + E'(t).$$

Note that (2.4) implies

$$I(t) \le \int_{t_1}^t -g'(s) \|\nabla u(t) - \nabla u(t-s)\|_2^2 ds \le -2E'(t),$$

then one has

$$\zeta(t)F_1'(t) \le -m\zeta(t)E(t)G'\left(\frac{r_1E(t)}{E(0)}\right) + \frac{c}{p}\zeta(t)\frac{r_1E(t)}{E(0)}G'\left(\frac{r_1E(t)}{E(0)}\right) - cE'(t) \quad (3.20)$$

by multiplying (3.19) by $\zeta(t)$ and by using the fact

$$\overline{G}'\left(\frac{r_1E(t)}{E(0)}\right) = G'\left(\frac{r_1E(t)}{E(0)}\right).$$

Define the functional $F_2(t) = \zeta(t)F_1(t) + cE(t)$ which is equivalent to E(t), which means

$$\gamma_1 F_2(t) \le E(t) \le \gamma_2 F_2(t) \tag{3.21}$$

for some γ_1 and γ_2 . Under a suitable choice of r_1 and for a positive constant k, we have

$$F_{2}'(t) \leq -k\zeta(t)\frac{E(t)}{E(0)}G'\left(\frac{r_{1}E(t)}{E(0)}\right) = -k\zeta(t)G_{2}\left(\frac{E(t)}{E(0)}\right)$$
(3.22)

with $G_2(t) = tG'(r_1t)$. Obviously, G_2 and G'_2 are positive in (0, 1] since

$$G'_2(t) = G'(r_1t) + r_1t^2G''(r_1t)$$

and the convexity of G in (0, r]. (3.22) and (3.21) imply

$$\left(\frac{\gamma_1 F_2(t)}{E(0)}\right)' \le -k\zeta(t)\frac{\gamma_1}{E(0)}G_2\left(\frac{E(t)}{E(0)}\right) \le -k_3\zeta(t)G_2\left(\frac{\gamma_1 F_2(t)}{E(0)}\right)$$
(3.23)

with $k_3 = k \frac{\gamma_1}{E(0)}$. Setting $R(t) = \frac{\gamma_1 F_2(t)}{E(0)}$, and then integrating (3.23) over (t_1, t) , one has

$$\int_{t_1}^t -\frac{R'(s)}{G_2(R(s))} ds \ge \int_{t_1}^t k_3 \zeta(s) ds.$$

Since $r_1 R(t_1) < r$, we have

$$G_1(r_1R(t)) = \int_{r_1R(t)}^{r_1R(t_1)} \frac{1}{sG'(s)} ds \ge k_3 \int_{t_1}^t \zeta(s) ds.$$

It is noted that G_1 is strictly decreasing function on (0, r] and $\lim_{t\to 0} G_1(t) = +\infty$ in Theorem 3.3, then

$$R(t) \le \frac{1}{r_1} G_1^{-1} \Big(k_3 \int_{t_1}^t \zeta(s) ds \Big).$$

Since R(t) is equivalent to E(t), further one obtains

$$E(t) \le k_4 G_1^{-1} \left(k_3 \int_{t_1}^t \zeta(s) ds \right)$$

with $k_4 = \frac{1}{r_1}$. This completes the proof of this theorem.

4. Possible generalizations

Our results can be generalized to the following initial boundary value problem with the time-varying delay $\tau(t)$

$$\begin{aligned} |u_t|^{\rho} u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s)\Delta u(s)ds \\ + \mu_1 u_t(x,t) + \mu_2 u_t(x,t-\tau(t)) &= b|u|^{p-2}u \quad (x,t) \in \Omega \times (0,\infty), \\ u_t(x,t-\tau(0)) &= f_0(x,t-\tau(0)) \qquad (x,t) \in \Omega \times (0,\tau(0)), \\ u(x,0) &= u_0(x), \ u_t(x,0) &= u_1(x) \qquad x \in \Omega, \\ u(x,t) &= 0 \qquad (x,t) \in \partial\Omega \times [0,\infty), \end{aligned}$$

equipped with the following assumptions in addition to $(\mathbf{H_1})$ and $(\mathbf{H_2}):$

 (A_1) the function $\tau \in W^{2,\infty}([0,T])$, for all T > 0 such that

$$0 < \tau_0 \le \tau(t) \le \tau_1$$
 for all $t > 0$,

$$\tau'(t) \le d < 1$$
 for all $t > 0$,

where τ_0 and τ_1 are positive numbers.

 (A_2) the coefficients of delay and dissipation satisfy

$$|\mu_2| \le \frac{2(1-d)}{2-d}\mu_1.$$

17

The above assumptions in fact are given in [6]. To be more precise, we need to define the new energy functional

$$\begin{split} E(t) &= \frac{1}{\rho+2} \|u_t\|_{\rho+2}^{\rho+2} + \frac{1}{2} \Big(1 - \int_0^t g(s) ds \Big) \|\nabla u\|_2^2 + \frac{1}{2} (g \circ \nabla u)(t) \\ &+ \frac{1}{2} \|\nabla u_t\|_2^2 + \frac{\xi}{2} \tau(t) \int_\Omega \int_0^1 z^2(x,\kappa,t) d\kappa dx - \frac{b}{p} \|u\|_p^p, \end{split}$$

where ξ satisfies

$$\frac{|\mu_2|}{1-d} \le \xi \le 2\mu_1 - |\mu_2|.$$

We also need to replace $I_3(t)$ in Lemma 2.9 by

$$I_3(t) = \int_0^1 e^{-2\tau(t)\kappa} \|z(x,\kappa,t)\|_2^2 d\kappa$$

further, one has

$$I_3'(t) \le -2I_3(t) - \frac{(1-d)e^{-2\tau_1}}{\tau_1} \|z(x,1,t)\|_2^2 + \frac{1}{\tau_0} \|u_t\|_2^2$$

as shown in Lemma 2.8 of [6]. Based on the above changes, we believe that it is possible to get the similar results in Theorem 3.3 by following the steps in this paper.

In addition, we have a conjecture that our results can also be extended to the following initial boundary value problem:

$$\begin{cases} |u_t|^{\rho} u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s)\Delta u(s)ds \\ + \mu_1 |u_t(x,t)|^{m-2} u_t(x,t) \\ + \mu_2 |u_t(x,t-\tau)|^{m-2} u_t(x,t-\tau) = b|u|^{p-2} u \quad (x,t) \in \Omega \times (0,\infty), \\ u_t(x,t-\tau) = f_0(x,t-\tau) & (x,t) \in \Omega \times (0,\tau), \\ u(x,0) = u_0(x), \ u_t(x,0) = u_1(x) & x \in \Omega, \\ u(x,t) = 0 & (x,t) \in \partial\Omega \times [0,\infty). \end{cases}$$

Acknowledgements. The author would like to express her sincere gratitude to Professor Wenjie Gao and Professor Bin Guo in Jilin University for their support and constant encouragement.

References

- [1] V. Arnold, Mathematical methods of classical mechanics. Springer, New York (1989)
- [2] F. Belhannache, M.M. Algharabli, S.A. Messaoudi, Asymptotic stability for a viscoelastic equation with nonlinear damping and very general type of relaxation functions, J. Dyn. Control Syst. 2020, 26: 45–67.
- [3] N. Boumaza, M. Saker, B. Gheraibia, Asymptotic behavior for a viscoelastic Kirchhoff-type equation with delay and source terms, *Acta Appl. Math.* 2021, 171: 18.
- [4] H. Chellaoua, Y. Boukhatem, Blow-up result for an abstract evolution problem with infinite memory and time-varying delay, Appl. Anal. 2020, 5: 1–24.
- [5] H. Chellaoua, Y. Boukhatem, Optimal decay for second-order abstract viscoelastic equation in Hilbert spaces with infinite memory and time delay, *Math. Meth. Appl. Sci.* 2021, 44: 2071–2095.
- [6] H. Chellaoua, Y. Boukhatem, Stability results for second-order abstract viscoelastic equation in Hilbert spaces with time-varying delay, Z. Angew. Math. Phys. 2021 72:46
- [7] Q.Y. Dai, Z.F. Yang, Global existence and exponential decay of the solution for a viscoelastic wave equation with a delay, Z. Angew. Math. Phys. 2014, 65: 885–903.

- [8] B.W. Feng, Well-posedness and exponential stability for a plate equation with time-varying delay and past history, Z. Angew. Math. Phys. 2017, 68:6.
- [9] B.W. Feng, H.Y. Li, Decay rates for a coupled viscoelastic Lamé system with strong damping, Math. Model. Anal. 2020, 25: 226–240.
- [10] J.H. Hassan, S.A. Messaoudi, General decay rate for a class of weakly dissipative second-order systems with memory, *Math. Meth. Appl. Sci.* 2019, 42: 2842–2853.
- [11] M. Kirane, B. Said-Houari, Existence and asymptotic stability of a viscoelastic wave equation with a delay, Z. Angew. Math. Phys. 2011, 62: 1065–1082.
- [12] M. Kafini, S.A. Messaoudi, A blow-up result in a nonlinear wave equation with delay, Mediterr. J. Math. 2016, 13: 237–247.
- [13] M. Kafini, S.A. Messaoudi,Local existence and blow up of solutions to a logarithmic nonlinear wave equation with delay, Appl. Anal. 2018, 1: 1–18.
- [14] J.R. Kang, Global nonexistence of solutions for viscoelastic wave equation with delay, Math. Meth. Appl. Sci. 2018, 41: 6834–6841.
- [15] W.J. Liu, General decay of the solution for a viscoelastic wave equation with a time-varying delay term in the internal feedback, J. Math. Phys. 2013, 54: 043504.
- [16] M.I. Mustafa, M. Kafini, Energy decay for viscoelastic plates with distributed delay and source term, Z. Angew. Math. Phys. 2016, 67: 36.
- [17] M.I. Mustafa, Optimal decay rates for the viscoelastic wave equation, Math. Meth. Appl. Sci. 2018, 41: 192–204.
- [18] M.I. Mustafa, General decay result for nonlinear viscoelastic equations, J. Math. Anal. Appl. 2018, 457: 134–152.
- [19] M.I. Mustafa, Optimal energy decay result for nonlinear abstract viscoelastic dissipative systems, Z. Angew. Math. Phys. 2021, 72:67.
- [20] S. Nicaise, C. Pignotti, Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, SIAM J. Control Optim. 2006, 45(5): 1561–1585.
- [21] S.T. Wu, Asymptotic behavior for a viscoelastic wave equation with a delay term, 2013, *Taiwanese J. Math.* 17: 765–784.
- [22] S.T. Wu, Asymptotic behavior for a viscoelastic wave equation with a time-varying delay term, J. Partial Differ. Equ. 2016, 29: 22–35.
- [23] S.T. Wu, Blow-up of solution for a viscoelastic wave equation with delay, Acta Math. Sci. Ser. B (Engl. Ed.) 2019, 39B(1): 329–338.

Menglan Liao

School of Mathematical Sciences, Xiamen University, Xiamen, Fujian, 361005, People's Republic of China

Email address: liaoml14@mails.jlu.edu.cn

Zhong Tan

School of Mathematical Sciences, Xiamen University, Xiamen, Fujian, 361005, People's Republic of China

Email address: tan850xmu.edu.cn