# Explicit universal estimate for *p*-polyharmonic equations via Morse index

Abdellaziz Harrabi<sup>1</sup>

Mathematics Department, Northern Border University, Arar, Saudi Arabia. Institut Supérieur des Mathématiques Appliquées et de l'Informatique, Universitè de Kairouan, Tunisia Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.

### Abstract

We investigate explicit universal estimate of finite Morse index solutions to

$$(-\Delta)^r u = f(x, u), \text{ in } \Omega,$$

where  $r \in \mathbb{N}$ ,  $r \ge 2$  and  $\Omega$  is a proper open subset of  $\mathbb{R}^n$ . Differently to [8, 11, 12, 18, 25, 29], we do not use a blow-up procedure which requires an available Liouville-type theorem in the whole space and where one imposes that f(x, u) has an asymptotical behavior at infinity like  $|u|^{q-1}u$ . We propose here a direct proof under large superlinear and subcritical growth conditions on f. To do so, we introduce a new interpolation inequality and we make use of a variant of the Pohozaev identity [24] to derive a local  $L^q$  estimate on a ball. Therefore, we employ a delicate boot strap argument from local  $L^q - W^{2r,q}$ -regularity. Particularly, we show that the universal constant evolves as a polynomial function of the Morse index.

Consider the following *p*-polyharmonic equation

$$\Delta_p^r u = |u|^{q-1} u$$
 in  $\mathbb{R}^n$  where  $p \ge 2$  and  $q > 1$ .

Thanks to our interpolation inequality, we improve the integral estimate and we extend the nonexistence results obtained in [18, 19] related to stable at infinity weak solutions. At last, we revised previous local  $L^p - W^{2r,p}$  estimate stated in [14, 26].

*Keywords:* Interpolation inequality, Universal estimate, *p*-polyharmonic equation, Morse index, Liouville-type theorem, Pohozaev identity, Boot strap argument. *PACS:* Primary 35G20, 35G30, Secondary 35B05, 35B09, 35B53.

#### 1. Introduction

Let  $\Omega$  be a domain of  $\mathbb{R}^n$ ,  $n \ge 2$ . For  $r \in \mathbb{N}$ ,  $r \ge 2$  and  $p \ge 2$ , let  $u \in W_{loc}^{r,p}(\Omega)$  and  $j = (j_1, j_2, ..., j_n)$  be a multi-index of order  $|j| = j_1 + j_2 + ... + j_n \le r$ . The weak j<sup>th</sup> partial derivative  $D^j u = \frac{\partial^j u}{\partial x_1^{j_1} ... \partial x_n^{j_n}}$  is defined *a.e* in  $\Omega$ . For  $1 \le s \le r$ , the magnitude of the s<sup>th</sup> gradient is defined *a.e* in  $\Omega$  by

$$|\nabla^{s} u| = \left(\sum_{|j|=s} |D^{j} u|^{p}\right)^{\frac{1}{p}}.$$
(1.1)

Email address: abdellaziz.harrabi@yahoo.fr (Abdellaziz Harrabi)

#### 1.1. Explicit universal estimate

Consider the following polyharmonic problem:

$$(-\Delta)^r u = f(x, u), \text{ in } \Omega, \tag{1.2}$$

where  $u \in C^{2r}(\Omega)$ , f and  $f' = \frac{\partial f}{\partial s}$  belong to  $C(\Omega \times \mathbb{R})$ . The associated quadratic form is defined by

$$Q_u(h) = \int_{\Omega} |D^r h|^2 - \int_{\Omega} f'(u)h^2, \ \forall h \in C_c^r(\Omega).$$

$$(1.3)$$

The Morse index of u, denoted by i(u) is defined as the maximal dimension of all subspaces V of  $C_c^r(\Omega)$  such that  $Q_{\mu}(h) < 0, \forall h \in V \setminus \{0\}.$ 

From blow-up technique one can derive universal estimate; or  $L^{\infty}$ -bounds provided that Liouville-type theorems classifying finite Morse index solutions are available. Note that even in the autonomous case f(x, u) = f(u), the blow-up procedure requires that f has the following asymptotical behavior at infinity:

 $(h_0)$ : There exist p > 1 and a positive constant  $c_0$  such that  $\lim_{|s|\to\infty} \frac{f'(s)}{|s|^{p-1}} = c_0$ . The reader may consult [3, 8, 11, 18, 29] and also [12, 25, 26, 27, 28] for positive solutions. However, this approach

fails to exhibit how the universal constant depends on i(u). We investigate here **explicit** universal estimate where we also success to relax assumption  $(h_0)$  into large superlinear and subcritical growth conditions. Precisely, we assume that there exist,  $s_0 > 0$ ,  $c_1 > 1$  and  $1 < p_1 \le p_2 < \frac{n+2r}{n-2r}$  such that for all  $(x, s) \in \Omega \times \mathbb{R} \setminus [-s_0, s_0]$ , we have

- (*h*<sub>1</sub>) (Super-linearity)  $f'(x, s)s^2 \ge p_1 f(x, s)s;$
- (*h*<sub>2</sub>) (Subcritical growth)  $(p_2 + 1)F(x, s) \ge f(x, s)s$  where  $F(x, s) = \int_0^s f(x, t)dt$ ;
- (*h*<sub>3</sub>)  $|(\nabla_{\gamma}F)(x, s)| \leq c_1(F(x, s) + 1)$ , for all  $(x, s) \in \Omega \times \mathbb{R}$ ;

 $(h_4) |f(x,s)| \le c_1$ , for all  $(x,s) \in \Omega \times [-s_0, s_0]$  and  $\pm f(x, \pm s_0) \ge \frac{1}{c_1}$ , for all  $x \in \Omega$ .

In the autonomous case, the above assumptions are reduced to  $(h_1)-(h_2)$  with  $\pm f(\pm s_0) > 0$ , which are weaker than  $(h_0)$  when  $1 . However, <math>f(s) = s_+^{p_2} - s_-^{p_1}$  (where  $s_+ = \max(s, 0)$ ,  $s_- = \max(-s, 0)$  and  $1 < p_1 \le p_2 < \frac{n+2r}{n-2r}$  satisfies  $(h_1)$ - $(h_2)$  and  $\pm f(\pm s_0) > 0$  but not  $(h_0)$ . Let  $K \in C^1(\Omega)$  be a positive function such that K,  $|\nabla K| \in L^{\infty}(\Omega)$ , then  $f(x, s) = K(x)(s_+^{p_2} - s_-^{p_1})$  satisfies  $(h_1)$ - $(h_4)$ . Let  $\Omega$  be a proper domain of  $\mathbb{R}^n$ . For  $\alpha \in (0, 1)$  and  $y \in \Omega$ , denote by  $\delta_y = \operatorname{dist}(y, \partial \Omega)$ ,  $d_y = \inf(\alpha, \delta_y)$  and set

 $\Omega_{\alpha} = \{y \in \Omega; \delta_y \ge \alpha\}$ . We use here (1.1) with p = 2. We have

**Theorem 1.1.** Assume that f satisfies  $(h_1)$ - $(h_4)$ . Then, there exist  $\alpha_0 \in (0,1)$  and a positive constant C = $C(n, r, p_1, p_2, s_0, c_1)$  independent of  $\Omega$  such that for any finite Morse index solution u of (1.2) and  $\alpha \in (0, \alpha_0)$ , we have

$$\sum_{j=0}^{2r-1} d_{y}^{j} |\nabla^{j} u(y)| \le C(1+i(u))^{\gamma_{2}} d_{y}^{\frac{2r\gamma_{1}}{p_{1}-1}}, \ \forall \ y \in \Omega,$$
(1.4)

where  $\gamma_1 = \frac{2r(p_1+1)p_2}{2r(p_2+1) - n(p_2-1)}$  and  $\gamma_2 = \frac{2r}{p_1-1}\gamma_1 + \frac{2r(p_2+1)}{2r(p_2+1) - n(p_2-1)}$ . Moreover, for any  $\alpha \in (0, \alpha_0)$ , we have

$$\|u\|_{C^{2r-1}(\Omega_{\alpha})} \le C(1+i(u))^{\gamma_2 + \frac{2\gamma_1}{p_1 - 1}} \text{ where } C = C(\alpha, n, r, p_1, p_2, s_0, c_1) > 0.$$
(1.5)

3

To prove Theorem 1.1, we introduce a new interpolation inequality (see Lemma 2.2 in section 2) to provide a first integral estimate on a ring and by virtue of a variant of Pohozaev identity, we extend this estimate to a ball. Precisely we prove

$$d_{y}^{-n} \int_{B(y,\frac{d_{y}}{2})} |f(x,u)|^{\frac{p_{2}+1}{p_{2}}} \le C(1+i(u)) \left(\frac{1+i(u)}{d_{y}}\right)^{\frac{2(p_{1}+1)r}{p_{1}-1}}, \ \forall \ y \in \Omega.$$
(1.6)

To conclude (1.4), we employ a delicate boot strap argument from a local  $L^{p}-W^{2r,p}$ -regularity. If in assumptions  $(h_{1})-(h_{2})$  we assume that  $\frac{n+2r}{n-2r} < p_{1} \le p_{2}$ , we may also derive (1.6), but the boot strap procedure fails to derive explicit universal estimate in the supercritical range. Furthermore, we can provide similar estimate of of (1.6) involving the p-polyharmonic equation. However, we do not dispose to any  $L^q$ -regularity result to star the boot strap procedure except. Particularly, explicit  $L^{\infty}$ -bounds has been obtained dealing with *p*-laplacian Dirichelet boundary problem on a bounded smooth domain [17]. This extends the result of [30] related to the second order Dirichlet boundary problem where it is shown that the  $L^{\infty}$  norm evolves less rapidity than a polynomial growth on i(u). In [20] the authors examined the influence of the type boundary condition in the biharmonic and triharmonic problems to provide similar polynomial growth estimate. The general higher order polyharmonic problem is still an open question where some local estimates near the boundary are no longer available for  $r \ge 4$  under any type boundary conditions. Observe that  $\gamma_1 \to \infty$  as  $p_2 \to \frac{n+2r}{n-2r}$ . This seems coherent with the fact that the  $L^{\infty}$  norm

has an exponential growth on i(u) when the nonlinearity f is close to the critical power as  $f(s) = \frac{|s|^{\frac{4}{n-2}}s}{\ln(s^2+2)}$  [15].

#### 1.2. Liouville type theorem

Consider the following problem

$$\Delta_p^r u = c_1 u_+^{q_1} - c_2 u_-^{q_2} \text{ in } \mathbb{R}^n, \tag{1.7}$$

where  $p \ge 2$ ,  $q_1$ ,  $q_2 > p - 1$ ,  $c_1$ ,  $c_2 > 0$  and  $u_+ = \max(0, u)$ ,  $u_- = \max(0, -u)$ . The p-polyharmonic operator  $\Delta_p^r$ is defined by

$$\Delta_p^r u = -\operatorname{div}\left\{\Delta^{j-1}(|\nabla\Delta^{j-1}u|^{p-2}\nabla\Delta^{j-1}u)\right\} \text{ if } r = 2j-1 \text{ and } \Delta^j(|\Delta^j u|^{p-2}\Delta^j u) \text{ if } r = 2j.$$

Define the main r-order differential operator by  $D_r u = \nabla \Delta^{j-1} u$  if r = 2j - 1 and  $\Delta^j u$  if r = 2j. We denote

$$D_{r}u \cdot D_{r}v = \begin{cases} \nabla \Delta^{j-1}u \cdot \nabla \Delta^{j-1}v & \text{if } r = 2j-1; \\ \Delta^{j}u\Delta^{j}v & \text{if } r = 2j; \\ |D_{r}u|^{2} = D_{r}u \cdot D_{r}u. \end{cases}$$
(1.8)

The Sobolev critical exponent of  $W^{r,p}(\mathbb{R}^n)$  is  $p^* = \frac{pn}{n-pr}$  if n > pr and  $\infty$  if  $n \le pr$ . Let us first recall some basic definitions and make the meaning of a weak stable solution. We designate by  $B_{\lambda}$  the ball of radius  $\lambda > 0$  centered at the origin and  $\overline{q} = \max(q_1, q_2)$ . Consider the Banach space  $E_{\lambda} =: W_0^{r,m}(B_{\lambda})$  if  $p - 1 < \overline{q} \leq 1$  $p^* - 1$  (respectively  $E_{\lambda} =: W_0^{r,p}(B_{\lambda}) \cap L^{\overline{q}+1}(B_{\lambda})$  if  $\overline{q} > p^* - 1$ ), endowed with the norm  $||D_r v||_{L^p(B_{\lambda})}$  (respectively by  $||D_r v||_{L^p(B_{\lambda})} + ||v||_{L^{\overline{q}+1}(B_{\lambda})}$ ). Define the energy functional by

$$I(v) := \frac{1}{p} \int_{B_{\lambda}} |D_{r}v|^{p} - \int_{B_{\lambda}} (\frac{c_{1}v_{+}^{q_{1}+1}}{q_{1}+1} + \frac{c_{2}v_{-}^{q_{2}+1}}{q_{2}+1}), \quad \forall v \in E_{\lambda}.$$

As  $p \ge 2$ , then  $I \in C^2(E_\lambda)^{-1}$ . We say that  $u \in W^{r,p}_{loc}(\mathbb{R}^n) \cap L^{q+1}_{loc}(\mathbb{R}^n)$  is a weak solution of (1.7) if for any  $h \in E_\lambda$ and for any  $\lambda > 0$ , we have

$$\int_{B_{\lambda}} |D_r u|^{p-2} D_r u \cdot D_r h = \int_{B_{\lambda}} (c_1 u_+^{q_1} - c_2 u_-^{q_2}) h.$$
(1.9)

<sup>&</sup>lt;sup>1</sup>If  $1 , the energy functional I is only <math>C^1$  functional, so it is not clear which definition of stability would be the natural one (see further comments in [10] for the p-laplacian equation).

We deal here with weak solutions since no regularity result is known for the p-polyharmonic operator for  $r \ge 2$ .

The linearized operator of (1.7) at u is given by

$$L_{u}(g,h) := \int_{B_{\lambda}} \left[ |D_{r}u|^{p-2} D_{r}g \cdot D_{r}h + (p-2)|D_{r}u|^{p-4} (D_{r}u \cdot D_{r}g)(D_{r}u \cdot D_{r}h) - (c_{1}q_{1}u_{+}^{q_{1}-1} + c_{2}q_{2}u_{-}^{q_{2}-1})gh \right], \ \forall (g,h) \in E_{\lambda}^{2}.$$

$$(1.10)$$

So, the associated quadratic form is defined by  $Q_u(h) := L_u(h, h)$  and as

$$\int_{B_{\lambda}} |D_r u|^{p-4} (D_r u \cdot D_r h)^2 \leq \int_{B_{\lambda}} |D_r u|^{p-2} |D_r h|^2, \ \forall h \in E_{\lambda},$$

then

$$Q_{u}(h) \leq (p-1) \int_{B_{\lambda}} |D_{r}u|^{p-2} |D_{r}h|^{2} - \int_{B_{\lambda}} (c_{1}q_{1}u_{+}^{q_{1}-1} + c_{2}q_{2}u_{-}^{q_{2}-1})h^{2}.$$
(1.11)

**Definition 1.1.** • We say that u is stable if for all  $\lambda > 0$  we have  $Q_u(h) \ge 0$ , for all  $h \in E_{\lambda}$ .

• For  $R_0 > 0$ , u is said to be stable outside the ball  $B_{R_0}$  if  $\forall \lambda > R_0$ ,  $\forall h \in E_{\lambda}$  with supp $(h) \subset \mathbb{R}^n \setminus B_{R_0}$ , we have  $Q_u(h) \ge 0$ . u is also called stable at infinity solution.

For  $c_1 = c_2$ ,  $q_1 = q_2$  and p = 2, there are an extensive classification results of stable at infinity solutions in lower order  $r \le 3$  (the reader may consult [3, 7, 11] for r = 1, [5, 29, 16, 9] for r = 2 and [21] for r = 3). Precisely, sharp nonexistence results have been obtained in [9, 11, 21], up to the so called Joseph-Lundgren exponent  $p_{II}(n,r)$ which is larger than the Sobolev critical exponent  $2^*$ . The approach to get nonexistence results in the supercritical range for r = 2, 3 makes use of powerful monotonicity formulas introduced respectively in [9, 4] and a delicate blow-down analysis. Very recently, classification results dealing with the nonhomogeneous polyharmonic problem have been discussed in [18] where one only requires that the term source has a subcritical growth at 0 to provide the nonexistence of nontrivial stable solutions. If  $c_1 > 0$ ,  $c_2 = 0$  and r = 1, it is shown that any finite morse index solution is spherically symmetric about some point of  $\mathbb{R}^n$  [22]. However, the situation is extremely unclear in the higher order case since the decomposition  $u = u^+ - u^-$  is no longer available for  $r \ge 2$ . When p > 2 the nonlinear degenerate operator *p*-polyharmonic does not yield any monotonicity formula, which provokes an obstruction to examine the supercritical growth case. In contrast, for r = 1, sharp classification results of stable solutions and radial stable at infinity solutions were obtained in [10] where the key ingredient is the Moser iteration argument. However, only the subcritical case related to the *p*-biharmonic problem was examined in [23] where the main integral estimate is so hard to achieve. This result has been extended in [19] to problem (1.7) with  $c_1 = c_2$  and  $q_1 = q_2$ . In this paper we classify stable and stable at infinity solutions of (1.7) for all  $r \ge 2$  and  $p \ge 2$  and  $p - 1 < q_1$ ,  $q_2 \le p^* - 1$ . For this end, we establish an interpolation inequality (see Lemma 2.3) to improve the integral estimate obtained in [18, 19]. Therefore we remove the exponential growth condition imposed on unbounded solutions. The main integral estimate reads as follows:

**Proposition 1.1.** Let  $r \ge 2$ ,  $p \ge 2$ ,  $q_1$ ,  $q_2 > p - 1$  and  $R_0 > 0$ . Let  $u \in W_{loc}^{r,p}(\mathbb{R}^n) \cap L_{loc}^{\overline{q}+1}(\mathbb{R}^n)$  be a weak solution of (1.7) which is stable outside the ball  $B_{R_0}$ . Then, there exist two positive constants  $C_0 = C_0(u, R_0, n, r, q_1, q_2, p)$  and  $C = C(n, r, q_1, q_2, p)$  such that

$$\int_{B_R} |\nabla^r u|^p + \int_{B_R} (c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1}) \le C_0 + CR^{n - \frac{pr(\overline{q}+1)}{\overline{q}+1-p}}, \ \forall R > \max(1, R_0).$$
(1.12)

Moreover, if u is a stable solution, then (1.12) holds with  $C_0 = 0$  for all R > 1. We also have

$$\int_{B_R} (c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1}) \le \int_{A_R} |\nabla^r u|^p + C R^{-pr} \int_{A_R} |u|^p.$$
(1.13)

Inequality (1.13) will be only used to prove the nonexistence of nontrivial weak stable solution when  $\overline{q} = p^* - 1$  and n > 2r.

We now give our Liouville type theorem. Set  $q = \min(q_1, q_2)$ , we have

**Theorem 1.2.** *1.* Let  $u \in W_{loc}^{r,p}(\mathbb{R}^n)$  be a weak stable solution of (1.7). If  $p - 1 < q_1, q_2 \le p^* - 1$ , then  $u \equiv 0^2$ .

2. Let  $u \in W_{loc}^{r+1,p}(\mathbb{R}^n)$  be a weak solution of (1.7) which is stable outside the ball  $B_{R_0}$ . If  $p - 1 < q_1$ ,  $q_2 < p^* - 1$ , then  $u \equiv 0$ .

*Moreover, if* n > pr and  $q = \overline{q} = p^* - 1$ , then

$$\int_{\mathbb{R}^n} |D_r u|^p = \int_{\mathbb{R}^n} (c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1}).$$
(1.14)

If  $p - 1 < \underline{q} < \overline{q} = p^* - 1$ , then  $u \ge 0$  if  $\overline{q} = q_1$  (respectively  $u \le 0$  if  $\overline{q} = q_2$ ) and (1.14) holds.

In section 2, we introduce our interpolation inequality and some preliminary technical Lemmas. Section 3 is devoted to the proof of the main Theorem 1.1. The proofs of Proposition 1.1 and Theorem 1.2 will be done in section 4. In the appendix we revised previous  $L^p - W^{2r,p}$  estimate stated in [14, 26].

#### 2. Interpolation inequalities and preliminary technical lemmas.

#### 2.1. Appropriate cut-off function and interpolation inequalities.

From the standard interpolation inequality on the unit ball  $B_1$  [1] and an obvious dilation argument, we have for every  $\varepsilon \in (0, 1)$  and for any  $v \in W_{loc}^{r,p}(\mathbb{R}^n)$ 

$$R^{r(q-r)} \int_{B_R} |\nabla^q v|^p \le \varepsilon \int_{B_R} |\nabla^r v|^p + C\varepsilon^{\frac{-q}{r-q}} R^{-pr} \int_{B_R} |v|^p, \quad \forall \ 1 \le q \le r-1,$$

$$(2.1)$$

where C = C(n, p, r) > 0. According to (2.1), we may easily derive the following weighted interpolation inequality (see [18, 13, 19, 26])

$$R^{p(q-r)}\Phi^p_q(v) \le \varepsilon \Phi^p_r(v) + C\varepsilon^{\frac{-q}{r-q}}R^{-pr}\int_{B_R} |v|^p dx.$$
(2.2)

Where  $\Phi_q$  is a family of weighted semi-norms defined by

$$\Phi_q(v) = \left(\sup_{0 < \alpha < 1} (1 - \alpha)^q \int_{B_{\alpha R}} |\nabla^q v|^p\right)^{\frac{1}{p}}, \ 0 \le q \le r$$

To make use of inequality (2.2) we employ the following family of cut-off functions

 $\psi = \psi_{(R,\alpha)} \in C_c^r(\mathbb{R}^n), \ 0 \le \psi \le 1, \ \operatorname{supp}(\psi) \subset B_{\alpha' R}, \ \psi \equiv 1, x \in B_{\alpha R},$ 

with  $\alpha \in (0, 1)$ , and  $\alpha' = \frac{1+\alpha}{2}$  where

$$\psi = \exp\left(\left(\frac{\frac{|x|}{R} - \alpha}{\frac{|x|}{R} - \alpha'}\right)^{r+1}\right) \text{ and } |\nabla^k \psi| \le C((1 - \alpha)R)^{-k} \text{ if } \alpha R < |x| < \alpha' R, \text{ and } 1 \le k \le r.$$

We introduce here a more general cut-off function related to two bounded open subset of  $\Omega$ ,  $\omega$  and  $\omega'$  such that  $\overline{\omega} \subset \omega' \subset \overline{\omega'} \subset \Omega$ . Set  $d = \operatorname{dist}(\omega, \Omega \setminus \omega')$ , we have

**Lemma 2.1.** There exists  $\psi \in C_c^{\infty}(\Omega)$  satisfying

$$\begin{cases} \operatorname{supp}(\psi) \subset \omega' \text{ and } 0 \leq \psi \leq 1, \\ \psi \equiv 1 \quad \text{if } x \in \omega, \\ |\nabla^k \psi(x)|^p \leq C d^{-kp}, \ \forall x \in \omega' \setminus \omega \text{ and } k \in \mathbb{N}, \end{cases}$$

$$(2.3)$$

where C is a positive constant depending only on (n, p, k).

<sup>2</sup>Precisely,  $p - 1 < q_1$ ,  $q_2 \le p^* - 1$  if n > pr and  $q_1, q_2 > p - 1$  if  $n \le pr$ ).

#### Proof of Lemma 2.1.

Set  $\omega_d = \{x \in \Omega, \text{ dist}(x, \omega) < \frac{d}{4}\}$ , we have  $\omega \subset \omega_d \subset \omega'$ . Let  $h = \chi_{\omega_d}$  be the indicator function of  $\omega_d$  and  $g \in C_c^{\infty}(\mathbb{R}^n)$  a nonnegative function such that

$$\operatorname{supp}(g) \subset B_1 \text{ and } \int_{\mathbb{R}^n} g(x) dx = 1.$$

Set

$$g_d(x) = \left(\frac{8}{d}\right)^n g\left(\frac{8x}{d}\right)$$
 and  $\psi(x) = \int_{B_\lambda} g_d(y)h(x-y)dy$ .

We have  $\operatorname{supp}(\psi) \subset \omega_d + B_\lambda \subset \omega'$ , and  $0 \le \psi \le 1$  and as  $\omega + B_\lambda \subset \omega_d$ . Thus,  $\psi(x) = \int_{B_\lambda} g_d(y) dy = 1$  if  $x \in \omega$  and as  $\psi(x) = \int_{\mathbb{R}^n} g_d(x - y)h(y) dy$  then  $\psi \in C_c^{\infty}(\mathbb{R}^n)$  and  $D^j\psi(x) = \int_{B_\lambda} D^j g_d(y)h(x - y) dy$ . Therefore,  $|D^j\psi(x)| \le \int_{B_\lambda} |D^j g_d| dy \le \lambda^{-|j|} \int_{B_\lambda} |D^j g(y)| dy \le Cd^{-|j|}$ , where C = ((n, |j|). Lemma 2.1 is proved.

Inequality (2.2) is essential in providing local  $L^{p}-W^{2r,p}$ -regularity [26] and the integral estimate from the stability property [18, 19]. However, compared to [11, 29, 9, 21], this estimate is weaker than the one obtained in lower order  $r \leq 3$ . We establish interpolation inequality composed with the cut-off function  $\psi$  defined in Lemma 2.1 which will be more relevant than (2.2) in testing; or truncating higher order PDE's against  $\psi u$ , as for classification and regularity problems. It will be also helpful to investigate **explicit** universal estimate via the Morse index. As in previous works [9, 11, 15, 16, 20, 21, 27, 29], we employ here Crandall-Rabinowitz's trick [6], that is, we use  $\psi^{m}$ , m > r as a test function. Let  $(q, k) \in \mathbb{N}^{*} \times \mathbb{N}^{*}$ , q + k = r, we have

**Lemma 2.2.** For every  $0 < \varepsilon < 1$  there exist two positive constants C = C(n, r, p, m) and  $C_{\varepsilon} = C(\varepsilon, n, p, r, m) > 0$  such that for any  $u \in W_{loc}^{r,p}(\Omega)$ , we have

$$\int_{\omega'} |\nabla^q u|^p |\nabla^k(\psi^m)|^p \le C d^{-pk} \int_{\omega'} |\nabla^q u|^p \psi^{p(m-k)} \le \varepsilon \int_{\omega'} |\nabla^r u|^p \psi^{pm} + C_\varepsilon d^{-pr} \int_{\omega'} |u|^p \psi^{p(m-r)}.$$
(2.4)

Consequently,

$$\int_{\omega'} |\nabla^r u|^p \psi^{pm} \le 2 \int_{\omega'} |\nabla^r (u\psi^m)|^p + Cd^{-pr} \int_{\omega'} |u|^p \psi^{p(m-r)},$$
(2.5)

and

$$\int_{\omega'} |\nabla^q u|^p |\nabla^k \psi^m|^p \le C d^{-pk} \int_{\omega'} |\nabla^q u|^p \psi^{p(m-k)} \le \varepsilon \int_{\omega'} |\nabla^r (u\psi^m)|^p + C_\varepsilon d^{-pr} \int_{\Omega} |u|^p \psi^{p(m-r)}.$$
(2.6)

When attempting to discuss stable at infinity weak solutions of (1.7), we use the following cut-off function  $\psi = \psi_{R,R_0} \in C_c^r(B_{2R}), R > 2R_0 > 0$  satisfying

$$\begin{cases} \psi \equiv 1 & \text{if } 2R_0 < |x| < R, \ \psi \equiv 0 & \text{if } |x| < R_0 & \text{or } |x| > 2R, \\ 0 \le \psi \le 1 & \text{and } |\nabla^k \psi| \le CR^{-k}, & \text{for all } R < |x| < 2R, & \text{and } 1 \le k \le r. \end{cases}$$
(2.7)

We have the analogue one of Lemma 2.2:

**Lemma 2.3.** Let  $\psi = \psi_{R,R_0}$  and  $u \in W_{loc}^{r,p}(\Omega)$ . Then, for every  $0 < \varepsilon < 1$  there exit two positive constants  $C_{\varepsilon} = C(\varepsilon, n, p, r, m) > 0$  and  $C_0 = C_0(u, R_0, \varepsilon, n, p, r, m)$  such that for all  $R > 2R_0$  we have

$$R^{-pk} \int_{B_{2R}} |\nabla^q u|^p |\nabla^k \psi^{p(m-k)} \le C_0 + \varepsilon \int_{B_{2R}} |\nabla^r u|^p \psi^{pm} + C_\varepsilon R^{-pr} \int_{\mathbb{R}^n} |u|^p \psi^{p(m-r)}.$$
(2.8)

As a consequence, we have

$$\int_{B_{2R}} |\nabla^r u|^p \psi^{pm} \le C_0 + 3 \int_{\mathbb{R}^n} |\nabla^r (u\psi^m)|^p + CR^{-pr} \int_{B_{2R}} |u|^p \psi^{p(m-r)},$$
(2.9)

and

$$R^{-pk} \int_{B_{2R}} |\nabla^q u|^p \psi^{p(m-k)} \le C_0 + \varepsilon \int_{B_{2R}} |\nabla^r (u\psi^m)|^p + C_\varepsilon R^{-pr} \int_{B_{2R}} |u|^p \psi^{p(m-r)}.$$
(2.10)

#### 2.2. Proofs of Lemma 2.2 and Lemma 2.3.

The proofs of these Lemmas are closely similar, we will do them in parallel. Working by induction we may verify that

$$\begin{cases} |\nabla^{k}\psi^{m}|^{p} \leq Cd^{-pk}\psi^{p(m-k)}, \quad x \in \omega' \setminus \omega, \\ |\nabla^{k}\psi^{m}_{R,R_{0}}|^{p} \leq CR^{-pk}\psi^{p(m-k)}_{R,R_{0}}, \quad R < |x| < 2R, \end{cases}$$
(2.11)

for all  $1 \le k \le r$  where *C* is a positive constant depending only on (n, p, r, m). The proof of (2.11) is done in the Appendix of [18]. Inequalities (2.6) and (2.10) are an immediate consequence of (2.4), (2.5) and respectively (2.8), (2.9). Also, inequalities (2.5) and (2.9) follow respectively from (2.4) and (2.8). Indeed, according to (1.1) we have

$$|\nabla^r u|^p \psi^{pm} = \sum_{|j|=r} |D^j u|^p \psi^{pm}.$$

We apply the following inequality <sup>3</sup>

$$b^{p} \le 2a^{p} + C|a - b|^{p}, \ \forall \ a, \ b > 0 \text{ where } C = C(p) > 0,$$
 (2.12)

with  $a = |D^{j}(u\psi^{m})|$  and  $b = |D^{j}(u)\psi^{m}|$  and using Leibnitz's formula [1], we arrive at

$$\begin{split} |\nabla^r u|^p \psi^{pm} &\leq 2 |\nabla^r (u\psi^m)|^p + C \sum_{|j|=r} |D^j (u\psi^m) - D^j (u)\psi^m|^p \\ &\leq 2 |\nabla^r (u\psi^m)|^p + C \sum_{q+k=r,\,q\neq r} |\nabla^q u|^p |\nabla^k \psi^m|^p. \end{split}$$

Integrate the above inequality over  $\Omega$  (respectively over  $B_{2R}$  if  $\psi = \psi_{R,R_0}$ ), and using (2.11), we obtain

$$\int_{\Omega} |\nabla^r u|^p \psi^{pm} \le 2 \int_{\Omega} |\nabla^r (u\psi^m)|^p + C \sum_{q+k=r, q \neq r} d^{-pk} \int_{\Omega} |\nabla^q u|^p \psi^{p(m-k)};$$

and

$$\int_{B_{2R}} |\nabla^r u|^p \psi^{pm} \le C_0 + 2 \int_{B_{2R}} |\nabla^r (u\psi^m)|^p + C \sum_{q+k=r, q \ne r} R^{-pk} \int_{\mathbb{R}^n} |\nabla^q u|^p \psi^{p(m-k)};$$

where

$$C_0 = \sum_{q+k=r, q \neq r} \int_{A_{R_0}} |\nabla^q u|^p |\nabla^k \psi^m|^p, \ A_{R_0} = \{R_0 < |x| < 2R_0\}.$$

Combine these inequalities with respectively (2.4) and (2.8) and choose  $\varepsilon = \frac{1}{rC}$ , we derive (2.5) and (2.9).

<sup>3</sup>In fact, we have  $(1 - 2^{-\frac{1}{p}})b \le |b - a|$  if  $b \ge 2^{\frac{1}{p}}a$ .

**Proof of** (2.4).

In the following, *C* denotes always generic positive constants depending on (n, p, r, m) only, which could be changed from one line to another. Set  $I_q = d^{-pk} \int_{\omega'} |\nabla^q u|_p^p \psi^{p(m-k)}$ . By virtue of (2.11), we have  $\int_{\Omega} |\nabla^q u|^p |\nabla^k (\psi^m)|^p \le CI_q$ . Consequently, to prove inequality (2.4) we have only to show that for every  $\varepsilon \in (0, 1)$ , we have

$$I_q \le C\varepsilon^{1-p^r} I_0 + \varepsilon I_r, \ \forall \ 1 \le q \le r-1.$$

We divide the proof of (2.13) in two steps.

Step 1. We shall establish the following first-order interpolation inequality:

$$I_q \le C \varepsilon^{1-p} I_{q-1} + \varepsilon I_{q+1}, \ \forall 1 \le q \le r-1.$$
 (2.14)

Denote by  $u_{|\omega'}$  the restriction of u on  $\omega'$ . As  $u \in W_{loc}^{r,p}(\Omega)$ , we have  $u_{|\omega'} \in W^{r,p}(\omega')$ . Thanks to Meyers-Serrin's density theorem [1], we may assume that  $u_{|\omega'} \in C^r(\omega') \cap W^{r,p}(\omega')^4$ . Let  $j = (j_1, j_2, ..., j_n)$  be a multi index with |j| = q and  $1 \le q \le r-1$ . As  $|j| \ne 0$ , there exists  $i_0 \in \{1, 2, ..., n\}$  such that  $j_{i_0} \ne 0$ . Set  $j_- = (j_1, ..., j_{i_0} - 1, ..., j_n), |j_-| = q - 1$  and  $j_+ = (j_1, ..., j_{i_0} + 1, ..., j_n), |j_+| = q + 1$ . Taking into account that  $u_{|\omega'} \in C^r(\omega') \cap W^{r,p}(\omega'), 1 \le |j| = q \le r-1$  and  $p \ge 2$ , we deduce

$$|D^{j}u|^{p-2}D^{j}u \in C^{1}(\omega') \text{ and } \frac{\partial(|D^{j}u|^{p-2}D^{j}u)}{\partial x_{i_{0}}} = (p-1)|D^{j}u|^{p-2}D^{j_{+}}u.$$
(2.15)

Recall that  $\psi \in C_c^{\infty}(\omega')$ ,  $|\nabla \psi| \le Cd^{-1}$  (see Lemma 2.1)), then integrate by parts with respect the variable  $x_{i_0}$ , we obtain

$$d^{-pk} \int_{\omega'} |D^{j}u|^{p} \psi^{p(m-k)} = d^{-pk} \int_{\omega'} |D^{j}u|^{p-2} D^{j}u \frac{\partial D^{j-u}}{\partial x_{i_{0}}} \psi^{p(m-k)}$$

$$= -(p-1)d^{-pk} \int_{\omega'} |D^{j}u|^{p-2} D^{j_{+}} u D^{j_{-}} u \psi^{p(m-k)}$$

$$-p(m-k)d^{-pk} \int_{\omega'} |D^{j}u|^{p-2} D^{j}u D^{j_{-}} u \psi^{p(m-k)-1} \frac{\partial \psi}{\partial x_{i_{0}}}$$

$$\leq Cd^{-pk} \int_{\omega'} |\nabla^{q-1}u| |\nabla^{q}u|^{p-2} |\nabla^{q+1}u| \psi^{p(m-k)}$$

$$+ Cd^{-(pk+1)} \int_{\omega'} |\nabla^{q-1}u| |\nabla^{q}u|^{p-1} \psi^{p(m-k)-1}, \qquad (2.16)$$

where C = C(n, p, r, m) > 0. As  $I_q = \sum_{|j|=q} d^{-pk} \int_{\omega'} |D^j u|^p \psi^{p(m-k)}$ , with k + q = r, then (2.16) implies  $I_q \le C(J_1 + J_2)$ , (2.17)

where

$$J_1 = d^{-pk} \int_{\omega'} |\nabla^{q-1}u| |\nabla^q u|^{p-2} |\nabla^{q+1}u| \psi^{p(m-k)} \text{ and } J_2 = d^{-(pk+1)} \int_{\omega'} |\nabla^{q-1}u| |\nabla^q u|^{p-1} \psi^{p(m-k)-1}$$

We need now the following Young's inequalities: for every  $0 < \varepsilon < 1$  and a, b, c > 0, we have

$$ab^{p-2}c \le \frac{1}{p}\varepsilon^{1-p}a^p + \frac{p-2}{p}\varepsilon b^p + \frac{1}{p}\varepsilon c^p,$$
(2.18)

<sup>&</sup>lt;sup>4</sup>When  $u_{|\omega'} \in W^{r,p}(\omega')$  we used standard approximation argument and Lebesgue's dominated convergence theorem to extend (2.13).

and

$$ab^{p-1} \le \frac{1}{p}\varepsilon^{1-p}a^p + \frac{p-1}{p}\varepsilon b^p.$$

$$(2.19)$$

Observe that pk = (k + 1) + k(p - 2) + (k - 1), p(m - k) = (m - (k + 1)) + (p - 2)(m - k) + (m - (k - 1)) and apply (2.18) with

$$a = d^{-(k+1)} |\nabla^{q-1}u| \psi^{m-(k+1)}, \ b^{p-2} = d^{-k(p-2)} |\nabla^{q}u|^{p-2} \psi^{(r-k)(p-2)}, \ c = d^{-(k-1)} |\nabla^{q+1}u| \psi^{m-(k-1)},$$

there holds

$$J_{1} \leq \frac{1}{p} \varepsilon^{1-p} I_{q-1} + \frac{(p-2)}{p} \varepsilon I_{q} + \frac{1}{p} \varepsilon I_{q+1}.$$
(2.20)

Also, as pk + 1 = (k + 1) + (p - 1)k and p(m - k) - 1 = (m - (k + 1)) + (p - 1)(m - k), according to (2.19) with

$$a = d^{-(k+1)} |\nabla^{q-1} u| \psi^{m-(k+1)}, \ b^{p-1} = d^{-(p-1)k} |\nabla^{q} u|^{p-1} \psi^{(m-k)(p-1)},$$

we obtain

$$J_{2} \leq \frac{1}{p} \varepsilon^{1-p} I_{q-1} + \frac{p-1}{p} \varepsilon I_{q}.$$
 (2.21)

Combine, (2.17) with (2.20) and (2.21), we arrive at

$$(1 - 2C\varepsilon)I_q \le C\varepsilon^{1-p}I_{q-1} + C\varepsilon I_{q+1}$$

We replace  $\varepsilon$  by  $\frac{\varepsilon}{4(1+C)}$ , thus the desired inequality (2.14) follows.

**Step 2.** Obviously inequality (2.13) is an immediate consequence of (2.14) if r = 2. However, the case  $r \ge 3$  needs a delicate iteration argument. We first prove that

$$I_q \le C \varepsilon^{1-p^r} I_0 + \varepsilon I_{q+1}, \ \forall 1 \le q \le r-1.$$
 (2.22)

The case q = 1 is obvious. For  $2 \le j \le q \le r - 1$  we apply (2.14) where we replace q by j - i and  $\varepsilon$  by  $\varepsilon^{p^i}$  with i = 0, 1, 2, ..., j - 1, we derive

$$C^{i}\varepsilon^{-p^{i}}I_{j-i} \leq C^{i+1}\varepsilon^{-p^{i+1}}I_{j-i-1} + C^{i}I_{j-i+1},$$

Set  $S_j = \sum_{i=2}^{J} I_i$ , we make the sum the above inequalities from i = 0 to i = j - 1, we obtain <sup>5</sup>

$$\varepsilon^{-1}I_{j} \leq C^{j}\varepsilon^{-p^{j}}I_{0} + I_{j+1} + \sum_{i=1}^{j-1}C^{i}I_{j-i+1}$$
  
$$\leq (C+1)^{r}\varepsilon^{-p^{r}}I_{0} + \varepsilon I_{j+1} + (C+1)^{r}S_{j}$$

Observe that  $S_j \leq S_q$  as  $I_q \geq 0$ , therefore we have

$$I_j \le C\varepsilon^{1-p^r} I_0 + \varepsilon I_{j+1} + C\varepsilon S_q \quad \text{if } 2 \le j \le q.$$

$$(2.23)$$

<sup>&</sup>lt;sup>5</sup>Recall that  $0 < \varepsilon < 1$ .

Thus, we make the sum of these inequalities from j = 2 to j = q, we arrive at

$$S_q \leq C \varepsilon^{1-p^r} I_0 + \varepsilon I_{q+1} + C \varepsilon S_q$$

We replace  $\varepsilon$  by  $\frac{\varepsilon}{2(C+1)}$ , we get

$$S_q \leq C\varepsilon^{1-p^r}I_0 + \varepsilon I_{q+1}$$
, for all  $1 \leq q \leq r-1$ .

Taking now j = q in (2.23) and combining it with the above inequality, then the desired (2.22) follows. As  $0 < \varepsilon < 1$ , from (2.22), we have

We make the sum of these inequalities, we deduce

$$I_q \le C\varepsilon^{1-p^r}I_0 + \varepsilon I_r.$$

Therefore inequality (2.13) is proven and the proof of Lemma 2.2 is completed.

Let us now end the proof of inequality (2.8) of Lemma 2.3 which is closely similar to the one of (2.4). We will only here explain how the constant  $C_0 = C_0(u, R_0, n, p, r, m)$  appears in (2.8). For  $1 \le q \le r - 1$  and k = r - q, let  $u \in W_{loc}^{r,p}(\mathbb{R}^n)$  and  $\psi = \psi_{R,R_0}$ . Set  $I_q = R^{-pk} \int_{B_{2R}} |\nabla^q u|_p^p \psi^{p(r-k)}$ . We first show that for every  $\varepsilon \in (0, 1)$  there exit two positive constants C = C(n, p, r, m) and  $C_0 = C(u, R_0, n, p, r, m)$  such that

$$I_q \le C_0 + C\varepsilon^{1-p}I_{q-1} + \varepsilon I_{q+1}, \ \forall 1 \le q \le r-1.$$
(2.25)

Recall that  $R > 2R_0$  and  $|\nabla \psi| \le CR^{-1}$  if  $x \in A_R = \{R < |x| < 2R\}$  (see (2.7)). As above integration by parts gives

$$\begin{split} R^{-pk} \int_{B_{2R}} |D^{j}u|^{p} \psi^{p(m-k)} &= -(p-1)R^{-pk} \int_{B_{2R}} D^{j}u|D^{j}u|^{p-2}D^{j}u\psi^{p(m-k)} \\ &- p(m-k)R^{-pk} \int_{A_{R}\cup A_{R_{0}}} D^{j}u|D^{j}u|^{p-2}D^{j}u\psi^{p(m-k)-1} \frac{\partial \psi}{\partial x_{i_{0}}} \\ &\leq C_{0} + (p-1)R^{-pk} \int_{B_{2R}} |D^{j}u||D^{j}u|^{p-2}|D^{j}u|\psi^{p(m-k)} \\ &+ p(m-k)R^{-pk-1} \int_{B_{2R}} |D^{j}u||D^{j}u|^{p-2}|D^{j}u|\psi^{p(m-k)-1}, \end{split}$$

where  $C_0 = pm(\inf(1, R_0))^{-rp} \int_{A_{R_0}} |D^{j_-}u| |D^{j}u|^{p-1} |\nabla \psi|$ . Therefore,

$$I_q \leq C_0 + C \left( R^{-pk} \int_{B_{2R}} |\nabla^{q-1}u| |\nabla^q u|^{p-2} |\nabla^{q+1}u| \psi^{p(m-k)} + R^{-pk-1} \int_{B_{2R}} |\nabla^{q-1}u| |\nabla^q u|^{p-1} \psi^{p(m-k)-1} \right),$$

where  $C_0 = pm(\inf(1, R_0))^{-rp} \int_{A_{R_0}} |\nabla^{q-1}u| |\nabla^q u|^{p-1} |\nabla \psi|$ . At this step, we follow the proof of (2.14), then we derive (2.25). Note that the additive constant  $C_0$  appeared in (2.25) does not provoke any mathematical difficulty to employ similar iteration argument as above which exhibits an other constant denoted again by  $C_0$  depending on  $(u, R_0, \varepsilon, n, p, r, m)$  such that

$$I_q \le C_0 + C\varepsilon^{1-p^r} I_0 + \varepsilon I_r, \ \forall 1 \le q \le r-1.$$

$$(2.26)$$

The proof of Lemma 2.3 is thereby completed.

#### 2.3. Preliminary technical Lemmas.

Thanks to our interpolation inequalities of Lemmas 2.2-2.3, we establish the following technical lemmas which will be essential to prove respectively Theorem 1.1 and Proposition 1.1. Let  $\psi = \psi_{(R,R_0)}$  the cut-off function defined in section 2 (see (2.7)). If  $R_0 = 0$  we denote by  $\psi = \psi_{(R)}$  the cut-off function defined in Lemma 2.1 with  $\omega = B_R$  and  $\omega' = B_{2R}$ . For m > r set  $A_R = \{R < |x| < 2R\}$ . We have

**Lemma 2.4.** 1. For every  $0 < \varepsilon < 1$ , there exist two positive constants  $C_0 = C_0(u, R_0, n, m, r, p, \varepsilon)$  and  $C_{\varepsilon} = C(n, m, r, p, \varepsilon)$  such that

$$\int_{B_{2R}} \left| |D_r u|^{p-2} \left( |D_r (u\psi^{\frac{pm}{2}})|^2 - D_r u D_r (u\psi^{pm}) \right) \right| \le C_0 + I(u, R, r, p, m, \varepsilon);$$
(2.27)

$$\int_{B_{2R}} \left| |D_r u|^p \psi^{pm} - \int_{B_{2R}} |D_r u|^{p-2} D_r u D_r (u \psi^{pm}) \right| \le C_0 + I(u, R, r, p, m, \varepsilon);$$
(2.28)

where

$$I(u, R, r, p, m, \varepsilon) = \varepsilon \int_{A_R} |\nabla^r u|^p \psi^{pm} + C_\varepsilon \sum_{q+k=r, q \neq r} R^{-pk} \int_{A_R} |\nabla^q u|^p |\psi^{p(m-k)}.$$
(2.29)

- 2. If  $\psi = \psi_R$ , inequalities (2.27) and (2.28) hold with  $C_0 = 0$ .
- 3. As a consequence of Lemma 2.3, we have

$$I(u, R, r, p, m, \varepsilon) \le \varepsilon \int_{B_{2R}} |\nabla^r(u\psi^m)|^p + C_\varepsilon R^{-pr} \int_{B_{2R}} |u|^p \psi^{p(m-r)}.$$
(2.30)

#### Proof of Lemma 2.4.

Let s > 1 and  $\beta_q \in \mathbb{R}$  with  $0 \le q \le r$ . From Young's inequality, we can easily derive that for every  $\epsilon > 0$  there exists  $C_{\epsilon} > 0$  such that

$$\left|\beta_r \sum_{q=0}^{r-1} \beta_q\right| \le \epsilon |\beta_r|^s + C_\epsilon \sum_{q=0}^{r-1} |\beta_q|^{\frac{s}{s-1}}.$$
(2.31)

For  $\eta \in C_c^r(\mathbb{R}^n)$  set  $A(\eta, u) = D_r(u\eta) - \eta D_r u$ . A simple computation and Leibnitz's formula [1], imply <sup>6</sup>

$$\begin{cases} \eta^{2} |D_{r}u|^{2} - D_{r}u \cdot D_{r}(u\eta^{2}) = -D_{r}u \cdot A(\eta^{2}, u); \\ |D_{r}(u\eta)|^{2} = \eta^{2} |D_{r}u|^{2} + 2\eta D_{r}u \cdot A(\eta, u) + |A(\eta, u)|^{2}; \\ |A(\eta, u)| \leq C_{r,n} \sum_{q=0}^{r-1} |\nabla^{r-q}\eta| |\nabla^{q}u|. \end{cases}$$

$$(2.32)$$

Therefore, we have

$$\left| |D_r u|^{p-2} \left( |D_r (u\eta)|^2 - D_r u D_r (u\eta^2) \right) \right| \le C \sum_{q+k=r, q \neq r} \left( |D_r u|^{p-1} |\nabla^q u| \left( \eta |\nabla^k \eta| + |\nabla^k (\eta^2)| \right) + |D_r u|^{p-2} |\nabla^q u|^2 |\nabla^k \eta|^2 \right) \le C \sum_{q+k=r, q \neq r} \left( |D_r u|^{p-1} |\nabla^q u| \left( \eta |\nabla^k \eta| + |\nabla^k (\eta^2)| \right) + |D_r u|^{p-2} |\nabla^q u|^2 |\nabla^k \eta|^2 \right) \le C \sum_{q+k=r, q \neq r} \left( |D_r u|^{p-1} |\nabla^q u| \left( \eta |\nabla^k \eta| + |\nabla^k (\eta^2)| \right) + |D_r u|^{p-2} |\nabla^q u|^2 |\nabla^k \eta|^2 \right) \le C \sum_{q+k=r, q \neq r} \left( |D_r u|^{p-1} |\nabla^q u| \left( \eta |\nabla^k \eta| + |\nabla^k (\eta^2)| \right) + |D_r u|^{p-2} |\nabla^q u|^2 |\nabla^k \eta|^2 \right)$$

and

$$\left| |D_r u|^{p-2} \left( |D_r u|^2 \eta^2 - D_r u D_r (u\eta^2) \right| \right) \le C_{r,n} |D_r u|^{p-1} \sum_{q+k=r, q \neq r} |\nabla^q u| |\nabla^k (\eta^2)|.$$

<sup>&</sup>lt;sup>6</sup>Both  $D_r$  and  $A(\eta, u)$  are respectively scalar operators if r is even, and n-vectorial operators if r is odd.

We choose now  $\eta = \psi^{\frac{pm}{2}}$ , as  $|D_r u| \le C |\nabla^r u|$ , by virtue of (2.11) we obtain <sup>7</sup>

$$\int_{B_{2R}} \left| |D_r u|^{p-2} \left( |D_r (u\psi^{\frac{pm}{2}})|^2 - D_r u D_r (u\psi^{pm}) \right) \right| \le C_0 + CS(u, R, p, m, r),$$
(2.33)

and

$$\int_{B_{2R}} \left| |D_r u|^{p-2} \left( |D_r u|^2 \psi^{pm} - D_r u D_r (u \psi^{pm}) \right) \right| \le C_{r,n} (C_0 + CS(u, R, p, m, r)),$$
(2.34)

where

$$S(u, R, p, m, r) = \int_{A_R} \left( |\nabla^r u|^{p-1} \sum_{q+k=r, q \neq r} R^{-k} |\nabla^q u| \psi^{pm-k} \right) + \int_{A_R} \left( |\nabla^r u|^{p-2} \sum_{q+k=r, q \neq r} R^{-2k} |\nabla^q u|^2 \psi^{pm-2k} \right),$$

and

$$C_0 = C_0(u, R_0, p, m, r) = \int_{A_{R_0}} \sum_{q+k=r, q \neq r} \left( |D_r u|^{p-1} |\nabla^q u| (\psi^{\frac{pm}{2}} |\nabla^k (\psi^{\frac{pm}{2}})| + |\nabla^k (\psi^{pm})|) + |D_r u|^{p-2} |\nabla^q u|^2 (|\nabla^k \psi^{\frac{pm}{2}}|^2) \right).$$

Observe that pm - k = (p - 1)m + (m - k) (respectively pm - 2k = (p - 2)m + 2(m - k)) and apply (2.31) in S(u, R, p, m, r) with  $s = \frac{p}{p-1}$ ,  $\beta_r = |\nabla^r u|^{p-1}\psi^{(p-1)m}$  and  $\beta_q = R^{-k}|\nabla^q u|\psi^{(m-k)}$  (respectively with  $s = \frac{p}{p-2}$  and  $\beta_r = |\nabla^r u|^{p-2}\psi^{(p-2)m}$  and  $\beta_q = R^{-2k}|\nabla^q u|^2\psi^{(2(m-k))}$ ), we obtain  $S(u, R, p, m, r) \le I(u, R, r, p, m, \varepsilon)$ . Hence, inequalities (2.27) and (2.28) follow the last inequality and respectively (2.33) and (2.34). ,We are now in a position to use our interpolation inequalities of Lemma 2.3 to prove (2.30). In fact as  $A_R \subset B_{2R}$ , so (2.30) is an immediate consequence of inequalities (2.9) and (2.10). The proof of Lemma 2.4 is completed.

For  $y \in \Omega$ , we designate by  $B(y, \lambda)$  the ball of radius  $\lambda > 0$  centered at y. Let  $\omega$  and  $\omega'$  be two bounded open subset of  $B(y, \lambda)$  such that  $\overline{\omega} \subset \omega' \subset \overline{\omega'} \subset B(y, \lambda)$  and  $d = \text{dist}(\omega, \Omega \setminus \omega')$ . Let  $\psi$  the cut-off function defined in Lemma 2.1. Similarly to the proof of Lemma 2.4, we derive the following

**Lemma 2.5.** For every  $0 < \varepsilon < 1$ , there exist two positive constants C = C(n, r)  $C_{\varepsilon} = C(n, m, r, \varepsilon)$  such that for any  $u \in W_{loc}^{r,2}(\Omega)$  we have

$$\int_{B(y,\lambda)} \left| |D_r(u\psi^m)|^2 - D_r u D_r(u\psi^{2m}) \right| \le \varepsilon \int_{B(y,\lambda)} |\nabla^r(u\psi^m)|^2 + C_\varepsilon d^{-2r} \int_{B(y,\lambda)} u^2 \psi^{2(m-r)};$$
(2.35)

and

$$\int_{B(y,\lambda)} \left| |D_r u|^2 \psi^{2m} - D_r u D_r (u \psi^{2m}) \right| \le \varepsilon \int_{B(y,\lambda)} |\nabla^r (u \psi^m)|^2 + C_\varepsilon d^{-2r} \int_{B(y,\lambda)} u^2 \psi^{2(m-r)}.$$
(2.36)

#### Proof of Lemma 2.5.

We only give here the outline of the proof since it is closely identical to the one of Lemma 2.4. Set

$$I(u, d, r, m, \varepsilon) = \varepsilon \int_{B(y, \lambda)} |\nabla^r u|^p \psi^{pm} + C_{\varepsilon} \sum_{q+k=r, q \neq r} R^{-pk} \int_{B(y, \lambda)} |\nabla^q u|^p |\psi^{p(m-k)}|.$$

we appeal to Lemma 2.2 and we insert inequalities (2.5) and (2.6) in the right hand side of I, we deduce

<sup>7</sup>Precisely we used (2.11) where one replaces respectively *m* by  $\frac{pm}{2}$  and *pm*.

$$I(u, d, r, m, \varepsilon) \le \varepsilon \int_{B(y, \lambda)} |\nabla^r(u\psi^m)|^2 + C_\varepsilon d^{-2r} \int_{B(y, \lambda)} u^2 \psi^{2(m-r)}.$$
(2.37)

Following now the proof of Lemma 2.4 where we replace p by 2, R by d and both  $B_{2R}$ ,  $A_R$  by  $B(y, \lambda)$ , we therefore derive the analogue one of inequalities (2.27) and (2.28) and combining them with (2.37), we obtain (2.35), (2.36). This ends the proof of Lemma 2.5.

According to the standard Calderon-Zygmund's inequality [13] and working by induction we derive the higher order analogue one [14]:

$$\int_{B(y,\lambda)} |\nabla^r v|^p \le C \int_{B(y,\lambda)} |D^r v|^p, \forall v \in W_0^{r,p}((B(y,\lambda)).$$
(2.38)

where C is a positive constant depending only on  $(n, r, p)^8$ .

Next, we recall the following elementary identity which will be used to provide a variant of Pohozaev identity [24] (see the proof in [18], page 1866<sup>9</sup>. Fix  $y \in \mathbb{R}^n$  and let  $u \in W_{loc}^{r+1,2}(\Omega)$ . We have

$$D_r u D_r (\nabla u \cdot (x - y)) = \frac{1}{2} \nabla (|D_r u|^2) \cdot (x - y) + r |D_r u|^2, \ a.e \text{ in } \mathbb{R}^n.$$
(2.39)

#### 3. Proof of Theorem 1.1.

In the following lemma, *C* denotes always generic positive constant depending only on the parameters  $(s_0, p_1, p_2)$ and the constant  $c_1$  of assumptions  $(h_1)$ - $(h_4)$ . Let q > 1 and set  $q_1 = \frac{p_2 + 1}{p_2}$ . We have

**Lemma 3.1.** There exists a positive constant *C* such that for all  $(x, s) \in \Omega \times \mathbb{R}$ , we have

- [1]  $f'(x, s)s^2 \ge p_1 f(x, s) C;$
- [2]  $(p_2 + 1)F(x, s) \ge f(x, s)s C;$
- [3]  $|s|^{p_1+1} \le C(|f(x,s)s|+1)$  and  $|s|^q \le C(|f(x,s)|^q+1);$
- [4]  $|f(x, s)s| \le f(x, s)s + C$  and  $|F(x, s)| \le C(|f(x, s)s| + 1);$
- $[5] |f(x,s)|^{q_1} \le C(|f(x,s)s|+1) \text{ and } |f(x,s)|^{\frac{q}{p_2}} \le C(|s|^q+1);$
- [6] For all  $\varepsilon \in (0, 1)$ ,  $0 \le a \le 1$  and b > 0 we have  $as^2b \le C + \varepsilon f(x, s)a^{\frac{p_1+1}{2}} + \varepsilon^{\frac{-2}{p_1-1}}b^{\frac{p_1+1}{p_1-1}}$ .

### Proof of Lemma 3.1

According to assumption  $(h_4)$ , we have

$$|F(x,s)|, |f(x,s)s| \le C, \ \forall \ (x,s) \in \Omega \times [-s_0, s_0].$$
(3.1)

Thus, points 1 and 2 are an immediate consequence of assumptions  $(h_1)$ - $(h_2)$  and points 3, 4 and 5 hold for all  $(x, s) \in \Omega \times [-s_0, s_0]$ . As the nonlinearity -f(x, -s) satisfies also  $(h_1)$ - $(h_4)$ , then we need only to prove these inequalities for all  $(x, s) \in \Omega \times [s_0, \infty)$ .

## Proof of point 3.

From  $(h_1)$  we have

$$f'(x,s)s \ge p_1 f(y,s), \ \forall (x,s) \in \Omega \times [s_0,\infty).$$
(3.2)

<sup>&</sup>lt;sup>8</sup>In fact, an obvious dilation and translation argument show that *C* does not depend on  $(y, \lambda)$ .

<sup>&</sup>lt;sup>9</sup>In [18], y = 0 and  $D_r$  is denoted by  $D^r$ .

It follows that  $\left(\frac{f(x,s)}{s^{p_1}}\right)' \ge 0$  and  $f(x,s) \ge \frac{(f(x,s_0)}{s^{p_1}_0}s^{p_1}$ . As  $f(x,s_0) \ge \frac{1}{c_1}$  for all  $x \in \Omega$  (see  $(h_4)$ ), then

$$f(x,s) \ge \frac{s^{p_1}}{c_1 s_0^{p_1}} \text{ and } f(x,s)s \ge \frac{s^{p_1+1}}{c_1 s_0^{p_1}}.$$
 (3.3)

Hence, we derive  $s^{p_1} \leq C|f(x,s)|$  and  $s^{p_1+1} \leq C|f(x,s)s| \ \forall \ (x,s) \in \Omega \times [s_0,\infty).$ Let q > 1, from the above inequality we have  $s^q \le s^{qp_1} + C \le C(|f(x, s)|^q + 1)$ . **Proof of point 4.** 

The first inequality follows from (3.3)). Integrating (3.2) over  $[s_0, s]$  and using  $(h_2)$ , we derive  $\frac{f(x, s)s}{n_2 + 1} \le F(x, s) \le 1$  $\frac{f(x, s)s}{p_1 + 1}$  + *C*, ∀(*x*, *s*) ∈ Ω × [*s*<sub>0</sub>, ∞). So, the second inequality follows. **Proof of point 5.** 

As  $f(x, s_0) \ge \frac{1}{c_1}$ ,  $(h_2)$  gives  $\left(\frac{F(x, s)}{s^{p_2+1}}\right)' \le 0$ . Then,  $(h_4)$  implies  $F(x, s) \le \frac{F(x, s_0)}{s_0^{p_2+1}}s^{p_2+1} \le \frac{c_1}{s_0^{p_2+1}}s^{p_2+1} \forall (x, s) \in \frac{F(x, s_0)}{s_0^{p_2+1}}$  $\Omega \times [s_0, \infty($ . Using again  $(h_2)$ , we obtain  $f(x, s)s \le C|s|^{p_2+1}$ ,  $\forall (x, s) \in \Omega \times [s_0, \infty)$ . From above, we have f(x, s)s = 0

|f(x,s)s| if  $(x,s) \in \Omega \times [s_0,\infty)$ . Then  $|f(x,s)|^{\frac{1}{p_2}} \le C|s|$ ,  $\forall (x,s) \in \Omega \times [s_0,\infty)$ . Therefore, for q > 1, we have

$$|f(x,s)|^{q_1} \le C|f(x,s)s|$$
 and  $|f(x,s)|^{\frac{q}{p_2}} \le C(|s|^q + 1), (x,s) \in \Omega \times [s_0,\infty).$ 

### Proof of point 6.

By the following Young's inequality  $t^2 v \le \varepsilon t^{p_1+1} + \varepsilon^{\frac{-2}{p_1-1}} v^{\frac{p_1+1}{p_1-1}}$ , with  $t = sa^{\frac{1}{2}}$  and v = b, we obtain  $as^2b \le \varepsilon s^{p_1+1}a^{\frac{p_1+1}{2}} + \varepsilon^{\frac{-2}{p_1-1}}b^{\frac{p_1+1}{p_1-1}}$ . As  $0 \le a \le 1$ , using point 3 we derive inequality 6. This end the proof of Lemma 3.1.

Let  $\alpha \in (0, 1)$ , recall that  $d_y = \inf(\alpha, \delta_y)$  where  $\delta_y = \operatorname{dist}(y, \partial \Omega)$  and  $y \in \Omega$ . We designate by  $B(x, d_y)$  the ball of radius  $d_y$  centered at y and  $A_a^b = \{x \in \mathbb{R}^n; a < |x - y| < b\}$ . For  $j = 1, 2, \dots, i(u) + 1$ , set

$$A_j := A_{a_j}^{b_j} \quad \text{with} \quad a_j = d_x \frac{2(j+i(u))}{4(i(u)+1)}, \quad b_j = d_x \frac{2(j+i(u))+1}{4(i(u)+1)}.$$
(3.4)

$$A'_{j} := A^{b'_{j}}_{a'_{j}} \quad \text{with} \quad a'_{j} = d_{x} \frac{2(j+i(u)) - \frac{1}{2}}{4(i(u)+1)}, \quad b'_{j} = d_{x} \frac{2(j+i(u)) + \frac{3}{2}}{4(i(u)+1)}. \tag{3.5}$$

From Lemma 2.1 with  $\omega = A_j$  and  $\omega' = A'_j$ , there exist  $\psi_j \in C_c^r(B(y, d_y))$  satisfying

- $\psi_j = 1$  for  $x \in A_j$  and  $0 \le \psi_j \le 1$  for  $x \in A'_j$ ;
- $\operatorname{supp}(\psi_j) \subset A'_j \text{ and } |\nabla^k \psi_j| \le C(\frac{1+i(u)}{d_x})^k.$

As dist $(A_j, B(y, d_y) \setminus A'_j) = \frac{d_y}{2(i(u) + 1)}$ , from inequality (2.11) and for m > r we have

$$|\nabla^{k}(\psi_{j}^{m})(x)|^{2} \leq C\psi_{j}^{2(m-k)} \left(\frac{1+i(u)}{d_{y}}\right)^{2k}, \ \forall x \in B(y, d_{y}).$$
(3.6)

We have  $supp(u\psi_i^m) \cap supp(u\psi_l^m) = \emptyset$  as  $A'_i \cap A'_l = \emptyset \forall 1 \le l \ne j \le 1 + i(u)$ . Consequently

$$Q_u\left(\sum_{1}^{1+i(u)}\lambda_j u\psi_j^m\right) = \sum_{1}^{1+i(u)}\lambda_j^2 Q_u(u\psi_j^m),$$

where  $Q_u$  is the quadratic form defined in (3.7)). According to the definition of i(u), there exists  $j_0 \in \{1, 2, ..., 1 + i(u)\}$  such that  $Q_u(u\psi_{j_0}^m) \ge 0$ , therefore point 1 of lemma 3.1 implies

$$\int_{B(y,d_y)} f(x,u)u\psi_{j_0}^{2m} - Cd_y^n \le \int_{B(y,d_y)} f'(x,u)(u\psi_{j_0}^m)^2 \le \int_{B(y,d_y)} |D_r(u\psi_{j_0}^m)|^2.$$
(3.7)

Step 1. We first prove the following estimate

$$\int_{A_{j_0}} |\nabla^r u|^2 + \int_{A_{j_0}} |f(x, u)u| \le C d_y^n \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}.$$
(3.8)

Multiplying equation (1.2) by  $(\frac{1+p_1}{2})u\psi_{j_0}^{2m}$  and integrating by parts, yields

$$\left(\frac{1+p_1}{2}\right)\int_{B(y,d_y)} D_r u D_r u(u\psi_{j_0}^{2m}) = \left(\frac{1+p_1}{2}\right)\int_{B(y,d_y)} f(x,u)u\psi_{j_0}^{2m}$$
(3.9)

So, the sum of (3.7) with (3.9), gives

$$\frac{p_1 - 1}{2} \left( \int_{B(y,d_y)} f(x,u) u \psi_{j_0}^{2m} + \int_{B(y,d_y)} |D_r(u\psi_{j_0}^m)|^2 \right) \le C d_y^n + \frac{p_1 + 1}{2} \left( \int_{B(y,d_y)} \left( |D_r(u\psi_{j_0}^m)|^2 - D_r u D_r u(u\psi_{j_0}^{2m}) \right) \right).$$
(3.10)

As  $u\psi_{j_0}^m \in W_0^{r,2}(B(y, d_y))$ , applying (2.38) in the left-hand side of (3.10) (with  $\lambda = d_y$ ). We also invoke Lemma 2.5 with  $\lambda = d_y$ ,  $\psi = \psi_{j_0}$  and  $d = \text{dist}(A_j, B(y, d_y) \setminus A'_j) = \frac{d_y}{2(i(u) + 1)}$ , precisely we insert inequality (2.35) in the right-hand side of (3.10), we derive

$$\int_{B(y,d_y)} |\nabla^r(u\psi_{j_0}^m)|^2 + \int_{B(y,d_y)} f(x,u)u\psi_{j_0}^{2m} \le Cd_y^n + \varepsilon \int_{B(y,d_y)} |\nabla^r(u\psi_{j_0}^m)|^2 + C_\varepsilon \left(\frac{1+i(u)}{d_x}\right)^{2r} \int_{B(y,d_y)} u^2 \psi_{j_0}^{2(m-r)}.$$
 (3.11)

Choose now  $m = \frac{(p_1 + 1)r}{2} > r$  so that  $\frac{(p_1 + 1)(m - r)}{p_1 - 1} = m$  and we apply point 6 of Lemma 3.1 with s = u,  $a = \psi_{j_0}^{2(m-r)}$  and  $b = C(\frac{1 + i(u)}{d_x})^{-2r}$ , there holds

$$C\left(\frac{1+i(u)}{d_y}\right)^{-2r}\int_{B(y,d_y)} u^2\psi_{j_0}^{2(m-r)} \leq Cd_y^n + \varepsilon \int_{B(y,d_y)} f(x,u)u\psi_{j_0}^{2m} + C_\varepsilon d_y^n \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}.$$

The last inequality with (3.11) together with point 4 of Lemma 3.1 imply <sup>10</sup>

$$\int_{B(y,d_y)} |\nabla^r (u\psi_{j_0}^m)|^2 + \int_{B(y,d_y)} |f(x,u)u|\psi_{j_0}^{2m} \le Cd_y^n \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}.$$

Thus, the desired inequality (3.8) follows as  $\psi_{j_0}^k(x) = 1$  in  $A_{j_0}$ . Step 2.

<sup>10</sup>Observe that 
$$d_y^n \le d_y^n \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}$$
 as  $d_y = \inf(\alpha, \delta_y) < 1$ .

15

We will establish a variant of the Pohozaev identity to extend the integral estimate (3.8) to the ball  $B(y, \frac{d_x}{2})$ . Precisely, we prove that

$$d_{y}^{-n} \int_{B(y,\frac{d_{y}}{2})} |f(x,u)|^{q_{1}} \le C(1+i(u)) \left(\frac{1+i(u)}{d_{y}}\right)^{\frac{2(p_{1}+1)r}{p_{1}-1}}.$$
(3.12)

Recall that  $A_{j_0} = \{a_{j_0} < |x - y| < b_{j_0} \text{ where } a_{j_0} \text{ and } b_{j_0} \text{ are defined in (3.5). So, Lemma 2.1 with } \omega = B(y, a_{j_0}), \omega' = B(y, b_{j_0}) \text{ and } d = \operatorname{dist}(B(y, a_{j_0}, B(y, d_y) \setminus B(y, b_{j_0})) = \frac{d_y}{2(i(u) + 1)} \text{ guaranties that there exists } \psi \in C_c^r(B(y, d_y)) \text{ satisfying}$ 

- $\psi \equiv 1$  in  $B(y, a_{j_0}), \psi(y) \equiv 0$  if  $|x y| \ge b_{j_0}$ ;
- $|\nabla^k \psi^{2m}| \le C \left(\frac{1+i(u)}{d_y}\right)^k \quad \forall y \in A_{j_0} \quad k=1,2,..,r.$

In one hand, multiplying (1.2) by  $(\nabla u \cdot (x - y)\psi^{2m}, m > r)$ . According to (2.39) with z = y and  $\lambda = d_y$ , integration by parts gives

$$\frac{2r-n}{2}\int_{B(y,d_y)}|D_r u|^2\psi^{2m} + n\int_{B(y,d_y)}F(x,u)\psi^{2m} = I - \int_{B(y,d_y)}(\nabla_x F)(x,u)\cdot(x-y)\psi^{2m},\tag{3.13}$$

where

$$I = \frac{1}{2} \int_{A_{j_0}} |D_r u|^2 (\nabla \psi^{2m} \cdot (x - y)) - \int_{A_{j_0}} F(x, u) (\nabla \psi^{2m} \cdot (x - y))$$

Observe that  $|x - y| \le d_y \le \alpha$  and  $|\nabla \psi^{2m} \cdot (x - y)| \le C(1 + i(u))$ ,  $\forall x \in A_{j_0}$ . Thus, point 4 of Lemma 3.1, assumption  $(h_3)$  and the integral estimate (3.8) of step 1 imply

$$\begin{cases} |I| \le C(1+i(u)) \left( d_y^n + \int_{A_{j_0}} \left( |f(y,u)u| + |D_r u|^2 \right) \right) \le C d_y^n (1+i(u)) \left( \frac{1+i(u)}{d_y} \right)^{\frac{2(p_1+1)r}{p_1-1}}; \\ \int_{B(y,d_y)} |(\nabla_x F)(x,u) \cdot (x-y)| \psi^{2m} \le \int_{B(y,d_y)} |F(x,u)| |x-y| \psi^{2m} \le C \left( d_y^n + \alpha \int_{B(y,d_y)} |f(x,u)u| \psi^{2m} \right). \end{cases}$$

These inequalities combined with (3.13) and point 2 of Lemma 3.1 imply

$$\left(\frac{2n}{(p_2+1)(n-2r)} - C\alpha\right) \int_{B(y,d_y)} f(x,u)u\psi^{2m} - \int_{B(y,d_y)} |D_r u|^2 \psi^{2m} \le Cd_y^n (1+i(u)) \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}.$$
(3.14)

On the other hand, multiplying equation (1.2) by  $u\psi^{2m}$ , we deduce

$$\int_{B(y,d_y)} D_r u D_r(u\psi^{2m}) = \int_{B(y,d_y)} f(x,u) u\psi^{2m}.$$

In view of inequality (2.36) of Lemma 2.5, we derive

$$\int_{B(y,d)} |D_r u|^2 \psi^{2m} - \int_{B(y,d_y)} f(x,u) u \phi^{2m} \le \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon (\frac{1+i(u)}{d_x})^{2r} \int_{B(y,d_y)} u^2 \psi^{2(m-r)} dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 dx = \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m$$

Choose now  $\alpha = \alpha_0 \in (0, 1)$  small enough so that  $\frac{2n}{(p_2 + 1)(n - 2r)} - C\alpha_0 > 1$ . Then the above inequality together with (3.14), imply <sup>11</sup>

$$\begin{split} \int_{B(y,d_y)} |D_r u|^2 \psi^{2m} + \int_{B(y,d_y)} f(x,u) u \psi^{2m} &\leq \varepsilon \int_{B(y,d_y)} |\nabla^r (u\psi^m)|^2 + C_\varepsilon \frac{(1+i(u))^{2r}}{d_y^{2r}} \int_{B(y,d)} u^2 \psi^{2(m-r)} \\ &+ C d_y^m (1+i(u)) \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}. \end{split}$$

Combine (2.35) with (2.36) and using again (2.38), we deduce that

$$C \int_{B(y,d_y)} |\nabla^r(u\psi^m)|^2 \le \int_{B(y,d_y)} |D_r(u\psi^m)|^2 \le \int_{B(y,d_y)} |D_r u|^2 \psi^{2m} \le \varepsilon \int_{B(y,d_y)} |\nabla^r(u\psi^m)|^2 + C_{\varepsilon} (\frac{1+i(u)}{d_y})^{2r} \int_{B(y,d)} u^2 \psi^{2(m-r)}.$$
(3.15)

We collect the two last inequalities and we choose  $m = \frac{(p_1 + 1)r}{2} > r$ , then points 4 and 6 of Lemma 3.1 imply

$$\int_{B(y,d_y)} |\nabla^r(u\psi^m)|^2 + \int_{B(y,d_y)} |f(x,u)u|\psi^{2m} \le Cd_y^n (1+i(u)\left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)p}{p_1-1}}.$$
(3.16)

As  $\psi \equiv 1$  on  $B(y, \frac{d_y}{2}) \subset B(y, a_{j_0})$ , it follows that

$$d_y^{-n} \int_{B(y,\frac{d_y}{2})} |f(x,u)|^{q_1} \le C(1+i(u)) \left(\frac{1+i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}$$

**Step 3.** In order to apply a boot-strap procedure, we use dilation argument to exploit local  $L^p - W^{2r,q}$  estimate on a fixed ball centered at the origin with radius 1. Precisely, set  $\lambda = \frac{d_y}{2} < 1$ ,  $u_{\lambda} = u(x + \lambda y)$  and  $g_{\lambda}(y) = f(x + \lambda y, u(x + \lambda y))$ ,  $y \in B_1$ , then  $u_{\lambda}$  satisfies

$$(-\Delta u_{\lambda})^{r} = \lambda^{2r} g_{\lambda} \text{ in } B_{1}, \qquad (3.17)$$

By virtue of (3.12), we have

$$\int_{B_1} |g_{\lambda}|^{q_1} = 2^n d_y^{-n} \int_{B(x, \frac{d_y}{2})} |f(y, u)|^{q_1} \le C \left(\frac{1 + i(u)}{d_y}\right)^{\frac{2(p_1+1)r}{p_1-1}}.$$
(3.18)

Appealing now local  $L^{p}-W^{2r,p}$  estimate (see corollary 6 in [26]; or Corollary 4.1 in the appendix) and applying Rellich-Kondrachov's theorem [14]. As  $0 < \lambda < 1$ , then for q > 1, point 3 of Lemma 3.1 implies

$$\|u_{\lambda}\|_{L^{q^*}(B_{\frac{1}{2}})} \le C\|u_{\lambda}\|_{W^{2r,q}(B_{\frac{1}{2}}))} \le C(\|g_{\lambda}\|_{L^{q}(B_{1})} + \|u_{\lambda}\|_{L^{q}(B_{1})}) \le C(\|g_{\lambda}\|_{L^{q}(B_{1})} + 1),$$
  
where  $q^{*} = \frac{qn}{n - 2rq}$  if  $2rq < n$  and for any  $q^{*} > 1$  if  $q = \frac{n}{2r}$ . (3.19)

and

$$\|u_{\lambda}\|_{C^{2r-1}(B_{\frac{1}{2}})} \le C\|u_{\lambda}\|_{W^{2r,q}(B_{\frac{1}{2}})} \le C(\|g_{\lambda}\|_{L^{q}(B_{1})} + \|u_{\lambda}\|_{L^{q}(B_{1})}) \le C(\|g_{\lambda}\|_{L^{q}(B_{1})} + 1), \text{ if } 2rq > n.$$
(3.20)

<sup>11</sup>Recall that  $\frac{2n}{(p_2+1)(n-2r)} > 1.$ 

$$\|g_{\lambda}\|_{L^{\frac{q^{*}}{p_{2}}}(B_{\frac{1}{2}})} \leq C\left(\|g_{\lambda}\|_{L^{q}(B_{1})}^{p_{2}}+1\right), \text{ if } 2rq \leq n.$$
(3.21)

We use now the boot strap argument to discuss the difficult case  $2rq_1 < n^{12}$ . Set  $q_2 = \frac{q_1^*}{p_2}$  and  $q_{k+1} = \frac{q_k^*}{p_2}$ . We claim that there exists  $k_0 \in \mathbb{N}^*$  such that

$$2rq_{k_0+1} > n \text{ and } 2rq_{k_0} < n.$$
 (3.22)

Indeed, suppose by contradiction that  $2rq_k < n$  for all  $k \in \mathbb{N}^*$ . Since  $\frac{1}{q_{k+1}} = \frac{p_2}{q_k} - \frac{2rp_2}{n}$ , a simple calculation gives

$$\frac{1}{q_{k+1}} = \frac{p_2^k}{q_1} - \frac{2rp_2}{n} \sum_{j=0}^{k-1} p_2^j = p_2^k \left(\frac{1}{q_1} - \frac{2rp_2}{n(p_2 - 1)}\right) + \frac{2rp_2}{n(p_2 - 1)}.$$
(3.23)

As  $1 < p_2 < \frac{n+2r}{n-2r}$  and  $q_1 = \frac{p_2+1}{p_2}$ , then  $\frac{1}{q_1} - \frac{2rp_2}{n(p_2-1)} < 0$  and  $\frac{1}{q_k} \to -\infty$ . So we reach a contradiction. Set

$$\beta = \frac{2rp_2}{n(p_2 - 1)} \left(\frac{2rp_2}{n(p_2 - 1)} - \frac{1}{q_1}\right)^{-1} = \frac{2r(p_2 + 1)}{2r(p_2 + 1) - n(p_2 - 1)}$$

From (3.23), we have  $p_2^{k_0} < \beta$  and  $p_2^{k_0+1} > \beta$ . Iterating now (3.21)  $k_0$  times and using (3), we obtain

$$||g_{\lambda}||_{L^{q_{k_{0}+1}}(B_{\frac{1}{2^{k_{0}+1}}})} \leq C(||g_{\lambda}||_{L^{q_{1}}(B_{1})}+1)^{p_{2}^{k_{0}}} \leq C(||g_{\lambda}||_{L^{q_{1}}(B_{1})}+1)^{\beta}.$$

Set  $\gamma_1 = \frac{(p_1 + 1)\beta}{q_1} = \frac{2r(p_1 + 1)p_2}{2r(p_2 + 1) - n(p_2 - 1)}$  and  $\gamma_2 = \beta + \frac{2r}{p_1 - 1}\gamma_1$ . As  $rq_{k_0 + 1} > n$ , the last inequality with (3.20) and (3.18) imply

$$\|u_{\lambda}\|_{C^{2r-1}(B_{\frac{1}{2^{k_0+1}}})} \leq C(1+i(u))^{\gamma_2} d_y^{\frac{2r}{p_1-1}\gamma_1}.$$

According to the definition of  $u_{\lambda}$ , we get  $\sum_{j=0}^{2r-1} d_{y}^{j} |(\nabla^{j} u)(y)| \leq C(1+i(u))^{\gamma_{2}} d_{y}^{\frac{2r}{p_{1}-1}\gamma_{1}}$ . So, the desired inequality (1.4) follows. Inequality (1.5) is an immediate consequence of (1.4). This achieves the proof of Theorem 1.1.

#### 4. Proofs of Proposition 1.1 and Theorem 1.2.

# *4.1. Proof of Proposition 1.1.* **Proof of** (1.12).

Recall that  $\overline{q} = \max(q_1, q_2)$  and  $q = \min(q_1, q_2) > p - 1$ . Let  $u \in W_{loc}^{r,p}(\mathbb{R}^n) \cap L_{loc}^{q+1}(\mathbb{R}^n)$  be a weak solution of (1.7) which is stable outside a ball  $B_{R_0}$ ,  $\overline{R}_0 > 0$ , as mentioned in section 2 we employ the cut-off function  $\psi = \psi_{(R,R_0)} \in C_c^r(\mathbb{R}^N)$ ,  $R > 2R_0$  (see (2.7) for more detail). When u is a stable solution, we use the cut-off function  $\psi = \psi_R$  defined by Lemma 2.1 related to  $\omega = B_R$  and  $\omega' = B_{2R}$ . For m > r, we have  $u\psi^{\frac{pm}{2}} \in W_0^{r,p}(B_{2R}) \cap L^{\overline{q}+1}(B_{2R})$ , then from (1.9) we have

$$\int_{B_{2R}} |D_r u|^{p-2} D_r u \cdot D_r (u\psi^{pm}) = \int_{B_{2R}} (c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1}) \psi^{pm}.$$
(4.1)

<sup>&</sup>lt;sup>12</sup>If  $2rq_1 > n$  (respectively  $2rq_1 = n$ ), the desired estimate (1.4) of Theorem **??** follows from (3.18) and (3.20) with  $q = q_1$  (respectively from (3.21) with  $q = q_1$  and (3.20) with  $q = \frac{n}{2r} + 1$ ).

Observe that  $\operatorname{supp}(u\psi^{\frac{pm}{2}}) \subset B_{2R} \setminus B_{R_0} \subset B_{2R} \setminus B_{R_0}$  and *u* is stable outside a ball  $B_{R_0}$ , then inequality (1.11) implies <sup>13</sup>

$$\int_{B_{2R}} (c_1 q_1 u_+^{q_1+1} + c_2 q_2 u_-^{q_2+1}) \psi^{pm} \le (p-1) \int_{B_{2R}} |D_r u|^{p-2} |D_r (u \psi^{\frac{pm}{2}})|^2.$$
(4.2)

We multiply (4.1) by  $-\frac{q+p-1}{2}$  and we add it to (4.2), we obtain <sup>14</sup>

$$\int_{B_{2R}} \left( c_1 \frac{2q_1 - \underline{q} - p + 1}{2} u_+^{q_1 + 1} + c_2 \frac{2q_2 - \underline{q} - p + 1}{2} u_-^{q_2 + 1} \right) \psi^{pm} + \frac{\underline{q} - p + 1}{2} \int_{B_{2R}} |D_r u|^{p-2} |D_r (u\psi^{\frac{pm}{2}})|^2 \le K(u, r, p, q_1, q_2, R),$$

$$(4.3)$$

where

$$K(u, r, p, q_1, q_2, R) = \frac{\underline{q} + p - 1}{2} \left( \int_{B_{2R}} |D_r u|^{p-2} \left( |D_r(u\psi^{\frac{pm}{2}})|^2 - D_r u D_r(u\psi^{pm}) \right) \right).$$

We are now in a position to apply Lemma 2.4, we insert inequality (2.27) in  $K(u, r, p, q_1, q_2, R)$ . Also, as  $u\psi^m \in$  $W_0^{r,p}(B_{2R})$ , we apply (2.38) (with y = 0 and  $\lambda = 2R$ ) in the left-hand side of (4.5). Therefore, we derive

$$\int_{B_{2R}} |\nabla^r (u\psi^m)|^p + \int_{\mathbb{R}^n} (u_+^{q_1+1} + u_-^{q_2+1})\psi^{pm} \le C_0 + \varepsilon \int_{A_R} |\nabla^r u|^p \psi^{pm} + C_\varepsilon \sum_{q+k=r,q\neq r} R^{-pk} \int_{A_R} |\nabla^q u|^p \psi^{p(m-k)}, \tag{4.4}$$

where  $C_0 = 0$  if u is a stable solution. Using again Lemma 2.4 with  $\varepsilon = \frac{1}{2}$  and applying inequality (2.30) in the right-hand side of (4.4), there holds

$$\int_{B_{2R}} |\nabla^r (u\psi^m)|^p + \int_{B_{2R}} (u_+^{q_1+1} + u_-^{q_2+1})\psi^{pm} \le C_0 + CR^{-pr} \int_{B_{2R}} |u|^p \psi^{p(m-r)} \le C_0 + CR^{-pr} \int_{B_{2R}} (u_+^p + u_-^p)\psi^{p(m-r)}.$$
 (4.5)

Let  $m = \frac{(\underline{q}+1)r}{q+1-p} > r$  so that  $pm = (\underline{q}+1)(m-r) \le (\overline{q}+1)(m-r)$ . We apply the following Young's inequality

$$a^{p}b \leq \frac{p}{t+1}a^{t+1} + \frac{t+1-p}{t+1}b^{\frac{t+1}{t+1-p}},$$

in the right-hand side of (4.5) with  $a = u_+ \psi^{(m-r)}$ ,  $t = q_1$  (respectively  $a = u_- \psi^{(m-r)}$ ,  $t = q_2$ ) and  $b = CR^{-pr}$ , we obtain

$$\begin{aligned} CR^{-pr} \int_{B_{2R}} (u_{+}^{p} + u_{-}^{p})\psi^{p(m-r)} &\leq \frac{p}{q_{1}+1} \int_{B_{2R}} u_{+}^{q_{1}+1}\psi^{pm} + \frac{p}{q_{2}+1} \int_{B_{2R}} u_{-}^{q_{2}+1})\psi^{pm} + C(R^{-\frac{pr(q_{1}+1)}{q_{1}+1-p}+n} + R^{-\frac{pr(q_{2}+1)}{q_{2}+1-p}+n}) \\ &\leq \frac{p}{q_{1}+1} \int_{B_{2R}} u_{+}^{q_{1}+1}\psi^{pm} + \frac{p}{q_{2}+1} \int_{B_{2R}} u_{-}^{q_{2}+1})\psi^{pm} + C(R^{-\frac{pr(q_{1}+1)}{q_{1}+1-p}+n}, \ \forall R > \max(1, R_{0}). \end{aligned}$$

Combining (4.5) with the above inequality and taking into account that  $\psi \equiv 1$  on  $\{2R_0 < |x| < R\}$  (respectively  $\psi_R = 1$ if |x| < R), then the main integral estimate (1.12) follows.

**Proof of inequality** (1.13).

We comeback to (4.4) with  $C_0 = 0$  (as *u* is a stable solution), we derive

$$\int_{\mathbb{R}^n} (u_+^{q_1+1} + u_-^{q_2+1})\psi^{pm} \leq C_0 + \varepsilon \int_{A_R} |\nabla^r u|^p \psi^{pm} + C_\varepsilon \sum_{q+k=r,q\neq r} R^{-pk} \int_{A_R} |\nabla^q u|^p \psi^{p(m-k)}.$$

<sup>&</sup>lt;sup>13</sup>See the paragraph before definition 1.1 for more detail.

<sup>&</sup>lt;sup>14</sup>Recall that  $1 \le p - 1 < \underline{q}$ . <sup>15</sup>As  $0 \le \psi \le 1$  we have  $\psi^{(\overline{q}+1)(m-r)} \le \psi^{pm}$ .

We insert (2.1) in the right-hand side of the above inequality, there holds <sup>16</sup>

$$\int_{B_R} (u_+^{q_1+1} + u_-^{q_2+1}) \le 2 \int_{A_R} |\nabla^r u|^p + CR^{-pr} \int_{A_R} |u|^p.$$

The proof of Proposition 1.1 is thereby completed.

#### 4.2. Proof of Theorem 1.2.

We recall that  $u \in W_{loc}^{r,p}(\mathbb{R}^n)$  is a stable weak solution of (1.7); or  $u \in W_{loc}^{r+1,p}(\mathbb{R}^n)$  is stable outside the ball  $B_{R_0}$  and  $p-1 < \underline{q} \leq \overline{q} \leq p^* - 1$  if n > pr (respectively  $p-1 < \underline{q} \leq \overline{q}$  if  $n \leq pr$ ). Then  $u \in W_{loc}^{r,p}(\mathbb{R}^n) \cup L_{loc}^{\overline{q}+1}(\mathbb{R}^n)$  and the main estimate (1.12) holds. In particular if u is a stable weak solution we have  $C_0 = 0$  and as  $n - \frac{pr(q+1)}{q+1-p} < 0$  if  $p-1 < \overline{q} < p^* - 1$ , then  $\int_{\mathbb{R}^n} (u_+^{q_1+1} + u_-^{q_2+1}) = 0$ , therefore  $u \equiv 0$ . If  $\overline{q} = p^* - 1$  and n > pr; or u is stable outside the ball  $B_{R_0}$  and  $p-1 < \overline{q} \le p^* - 1$ , from (1.12) we can easily see that  $u_+^{q_1+1} + u_-^{q_2+1} \in L^1(\mathbb{R}^n)$  and  $\nabla^r u \in L^p(\mathbb{R}^n)$ . Also, from Hölder's inequality, we have

$$\begin{split} R^{-pr} \int_{A_R} |u|^p &\leq CR^{-pr} \int_{B_{2R}} (u_+^p + u_-^p) \\ &\leq CR^{\frac{n(q_1+1-p)}{q_1+1} - pr} \left( \int_{A_R} u_+^{q_1+1} \right)^{\frac{p}{q_1+1}} + CR^{\frac{n(q_2+1-p)}{q_2+1} - pr} \left( \int_{A_R} u_-^{q_2+1} \right)^{\frac{p}{q_2+1}} \\ &\leq C \left( \int_{A_R} u_+^{q_1+1} \right)^{\frac{p}{q_1+1}} + C \left( \int_{A_R} u_-^{q_2+1} \right)^{\frac{p}{q_2+1}} . \end{split}$$

Therefore,

$$R^{-rp} \int_{A_R} |u|^p = o(1), \ \int_{A_R} (u_+^{q_1+1} + u_-^{q_2+1}) = o(1) \text{ and } \int_{A_R} |\nabla^r u|^p = o(1), \text{ as } R \to \infty.$$
(4.6)

Consequently, if u is a stable solution of (1.7) and n > pr, it follows from (1.12) and (4.6) that  $\int_{P}^{P} (u_{+}^{q_{1}+1} + u_{-}^{q_{2}+1}) =$ o(1), as  $R \to \infty$  and therefore  $u \equiv 0$ . This ends the proof of point 1.

#### **Proof of Point 2.**

To make use of inequality (2.28) of Lemma 2.4, we use again the cut-off function  $\psi_R$  defined in Lemma 2.1 with  $\omega = B_R$  and  $\omega' = B_{2R}$ . Set  $\phi = \psi_R^{mp}$ . In one hand, from equality (4.1), we have

$$\int_{B_{2R}} |D_r u|^{p-2} D_r u D_r (u\phi) = \int_{B_{2R}} (c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1})\phi$$
  
According to (2.28) and (1.13), we obtain <sup>17</sup>

(2.28) and (1.13),

$$\int_{B_{2R}} |D_r u|^p \phi - \int_{B_{2R}} (c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1}) \phi \le C \left( \int_{A_R} |\nabla^r u|^p + CR^{-pr} \int_{A_R} |u|^p \right).$$

It follows from (4.6) that

$$\int_{B_{2R}} |D_r u|^p \psi - \int_{B_{2R}} (u_+^{q_1+1} + u_-^{q_2+1})\psi = o(1).$$
(4.7)

<sup>&</sup>lt;sup>16</sup>In fact, inequality (1.13) holds when one replaces  $B_{\lambda}$  by  $A_R = \{r < |x| < 2R\}$ .

<sup>&</sup>lt;sup>17</sup>Inequality (2.28) holds (with  $C_0 = 0$  because we used  $\psi_R$  as cut-off function.

$$\int_{B_{2R}} |D_r u|^{p-2} D_r u D_r (\nabla u \cdot x\phi) = \int_{B_{2R}} (c_1 u_+^{q_1} - c_2 u_-^{q_2}) \nabla u \cdot x\phi.$$

Multiplying now (2.39) by  $|D_r u|^{p-2} D_r u$  (with y = 0), we derive

$$|D_r u|^{p-2} D_r u D_r (\nabla u \cdot x) = \frac{1}{p} \nabla (|D_r u|^p) \cdot x + r |D_r u|^p, \ a.e \text{ in } \mathbb{R}^n.$$

Thus, direct integrations by parts yield

$$\int_{B_{2R}} |D_r u|^{p-2} D_r u D_r (\nabla u \cdot x) \phi = \frac{rp-n}{p} \int_{B_{2R}} |D_r u|^p \phi - \frac{1}{p} \int_{B_{2R}} |D_r u|^p (\nabla \phi \cdot x)$$

and

$$\int_{B_{2R}} (c_1 u_+^{q_1} - c_2 u_-^{q_2}) \nabla u \cdot x \phi = -n \int_{B_{2R}} (\frac{c_1 u_+^{q_{1+1}}}{q_1 + 1} + \frac{c_2 u_-^{q_{2+1}}}{q_2 + 1}) \phi - \int_{B_{2R}} (\frac{c_1 u_+^{q_{1+1}}}{q_1 + 1} + \frac{c_2 u_-^{q_{2+1}}}{q_2 + 1}) (\nabla \phi \cdot x).$$

By virtue of (4.6) we have

$$\int_{B_{2R}} \left(\frac{c_1 u_+^{q_{1+1}}}{q_1+1} + \frac{c_2 u_-^{q_{2+1}}}{q_2+1}\right) (\nabla \phi \cdot x) (\nabla \phi \cdot x) = o(1) \text{ and } \int_{B_{2R}} |D_r u|^p (\nabla \phi \cdot x) = o(1), \text{ as } R \to \infty.$$

Collecting these equalities, we arrive at

$$n \int_{B_{2R}} \left(\frac{c_1 u_+^{q_{1+1}}}{q_1+1} + \frac{c_2 u_-^{q_{2+1}}}{q_2+1}\right) \phi - \frac{n-rp}{p} \int_{B_{2R}} |D_r u|^p \phi = o(1) \text{ as } R \to \infty.$$
(4.8)

Thus, if  $n \le pr$ , then (4.8) implies that  $\int_{B_{2R}} (\frac{c_1 u_+^{q_{1+1}}}{q_1+1} + \frac{c_2 u_-^{q_{2+1}}}{q_2+1})\phi = o(1)$ , so  $u \equiv 0$ . If n > pr and  $p - 1 < \underline{q} \le \overline{q} < p^* - 1$ , we combine (4.8) with (4.7) we derive

$$\left[c_1\left(\frac{np}{(n-rp)(q_1+1)}-1\right)\int_{B_{2R}}u_+^{q_1+1}+c_2\left(\frac{np}{(n-rp)(q_2+1)}-1\right)\int_{B_{2R}}u_-^{q_2+1}\right]\phi=o(1),$$

which implies that  $u \equiv 0$ . If  $\underline{q} = \overline{q} = p^* - 1$ , taking into account that  $c_1 u_+^{q_1+1} + c_2 u_-^{q_2+1} \in L^1(\mathbb{R}^n)$  and  $\nabla^r u \in L^p(\mathbb{R}^n)$ , we apply the dominated convergence theorem in (4.7) we obtain (1.14). Now if  $p - 1 < \underline{q} < \overline{q} = p^* - 1$ , as above we derive that

$$c_2\left(\frac{np}{(n-rp)(\underline{q}+1)}-1\right)\int_{B_{2R}}u_-^{\underline{q}+1}\phi=o(1)$$

Consequently, (1.14) holds with  $u \ge 0$  if  $\overline{q} = q_1$  (respectively  $u \le 0$  if  $\overline{q} = q_2$ ). This achieves the proof of Theorem 1.2.

Appendix: Local  $L^p$ - $W^{2r,p}$  estimate revisited.

Let  $\Omega$  be a domain of  $\mathbb{R}^n$ . consider the linear higher order elliptic problem of the form

$$Lu = g \quad \text{in} \quad \Omega. \tag{4.9}$$

Here

$$L = \left(-\sum_{i,k=1}^{n} a_{ik}(x) \frac{\partial^2}{\partial x_i \partial x_k}\right)^r + \sum_{|j| \le 2r-1} b_j(x) D^j$$

is a uniformly elliptic operator with coefficients  $b_j \in L^{\infty}_{loc}(\Omega)$  and  $a_{ik} \in C^{2r-2}(\Omega)$ , that is that there exists a constant  $\lambda > 0$  with  $\lambda^{-1}|\xi|^2 \leq \sum_{k=1}^{n} a_{ik}(x)\xi_i\xi_k \leq \lambda|\xi|^2$  for all  $\xi \in \mathbb{R}^n$ ,  $x \in \Omega$ .

$$\lambda > 0$$
 with  $\lambda^{-1} |\xi|^2 \le \sum_{i,k=1} a_{ik}(x)\xi_i\xi_k \le \lambda |\xi|^2$  for all  $\xi \in \mathbb{R}^n, x \in \Omega$ .

For  $p \ge 2$ , thanks to Lemma 2.2, we propose a direct proof of local analogue of the celebrated  $L^{p}-W^{2r,p}$  estimate of Agmon-Douglis-Nirenberg [2]. Let  $\omega$  and  $\omega'$  be two bounded open subset of  $\Omega$  such that  $\overline{\omega} \subset \omega'$  and  $\overline{\omega'} \subset \Omega$ , we have

**Corollary 4.1.** Let  $g \in L^p_{loc}(\Omega)$  for some p > 1. Then there exists a constant C > 0 depending only on  $||a_{ik}||_{C^{2r-2}(\omega')}$ ,  $||b_j||_{L^{\infty}(\omega')}$  and  $\lambda, \omega', d, n, p, r$  such that for any  $u \in W^{2r,p}_{loc}(\Omega)$  a weak solution of (4.9), we have

$$\|u\|_{W^{2r,p}(\omega)} \le C\left(\|g\|_{L^{p}(\omega')} + \|u\|_{L^{p}(\omega')}\right)$$

In the following *C* denotes a generic positive constant which depends on the parameters stated in Corollary 4.1 and  $d = \text{dist}(\omega, \Omega \setminus \omega')$ .

**Proof of Corollary 4.1.** 

,We distingue two cases

**Case 1,**  $p \ge 2$ .

We truncate the equation (4.9) by  $\psi^m$ , where  $\psi$  is the cut-off function defined in Lemma 2.1 and  $m \ge 2r$ . Thus, we have

$$L(u\psi^{m}) = g\psi^{m} + uL(\psi^{m}) + b_{0}u\psi^{m} + \sum_{1 \le |i| + |j| \le 2r-1} c_{i,j}D^{j}uD^{i}(\psi^{m}), \text{ where } c_{i,j} \in L^{\infty}(\Omega).$$

As  $\psi \in C_c^{\infty}(\omega')$  and  $u \in W^{2r,p}(\omega')$ , then  $u\psi^m \in W^{2r,p}(\omega') \cap W_0^{r,p}(\omega')$  with **compact support**, therefore Agmon-Douglis-Nirenberg's global estimate [2]<sup>18</sup> and (2.11) imply

$$\sum_{0 \le s \le 2r} \int_{\omega'} |\nabla^s(u\psi^m)|^p \le C \left( ||g||_{L^p(\omega')}^p + ||u||_{L^p(\omega')}^p + \sum_{1 \le s \le 2r-1} \sum_{1 \le q \le s} \int_{\omega'} |\nabla^q u|^p |\nabla^{s-q}\psi^m|^p \right), \tag{4.10}$$

Using now our interpolation inequality (2.6) (where one replaces r by s in (2.6)), we obtain

$$\sum_{1 \le q \le s-1} \int_{\omega'} |\nabla^q u|^p |\nabla^{s-q} \psi^m|^p \le \varepsilon \int_{\omega'} |\nabla^s (u\psi^m)|^p + C_{\varepsilon,d} \int_{\omega'} |u|^p \psi^{p(m-s)}$$

Also apply the second inequality of (2.6) with r = s + 1 and we replace  $\varepsilon$  by  $\frac{\varepsilon}{d}$ , yields

$$\int_{\omega'} |\nabla^s u|^p |\psi^{pm} \le \int_{\omega'} |\nabla^s u|^p \psi^{p(m-1)} \le \varepsilon \int_{\omega'} |\nabla^{s+1} (u\psi^m)|^p + C_{\varepsilon,d} \int_{\omega'} |u|^p \psi^{p(m-s-1)}.$$

Collecting the two last inequalities, we derive

$$\sum_{1 \le s \le 2r-1} \sum_{1 \le q \le s} \int_{\omega'} |\nabla^q u|^p |\nabla^{s-q} \psi^m|^p \le \varepsilon \sum_{0 \le s \le 2r} \int_{\omega'} |\nabla^s (u \psi^m)|^p + C_{\varepsilon,d} \int_{\omega'} |u|^p.$$

We insert the above inequality in the right-hand side of (4.10), and we choose  $\varepsilon = \frac{1}{2C}$ , we deduce

 $\|u\psi^{m}\|_{W^{2r,p}(\omega')}^{p} \leq C\left(\|g\|_{L^{p}(\omega')}^{p} + \|u\|_{L^{p}(\omega')}^{p}\right).$ 

<sup>&</sup>lt;sup>18</sup>See also Theorem 5 in [26]).

Since  $\psi(x) = 1$  if  $x \in \omega$ , we obtain

$$\|u\|_{W^{2r,p}(\omega)}^{p} \leq C\left(\|g\|_{L^{p}(\omega')}^{p} + \|u\|_{L^{p}(\omega')}^{p}\right).$$

**Case 2,** 1 < *p* < 2.

Let  $B(x,\rho)$  be the ball of radius  $\rho$  and centred at x. We apply Corollary 6 of [26]) with  $R = \frac{d}{2}$  and  $\sigma = \frac{1}{2}$ . Then, for all  $x \in \omega$ , we have

$$\begin{aligned} \|u\|_{W^{2r,p}(B(x,\frac{d}{4}))}^{p} &\leq C\left(\|g\|_{L^{p}(B(x,\frac{d}{2}))}^{p} + \|u\|_{L^{p}(B(x,\frac{d}{2}))}^{p}\right) \\ &\leq C\left(\|g\|_{L^{p}(\omega')}^{p} + \|u\|_{L^{p}(\omega')}^{p}\right). \end{aligned}$$

As  $\overline{\omega}$  is a compact set, we can find  $x_i \in \omega$ ,  $i = 1, 2, ..., k_0$  such that  $\overline{\omega} \subset \bigcup B(x_i, \frac{d}{4}) \subset \omega'$  where  $k_0 \in \mathbb{N}^*$  depending only on d and  $\omega$ . Therefore, we derive

$$\|u\|_{W^{2r,p}(\omega)}^{p} \leq Ck_{0}\left(\|g\|_{L^{p}(\omega')}^{p} + \|u\|_{L^{p}(\omega')}^{p}\right).$$

This achieves the proof of Corollary 4.1.

#### References

- [1] R.A. Adams, J.J.F. Fournier, Sobolev Spaces. Pure and Applied Mathematics. Academic Press, London, 2nd edn. xiii+305 pp (2003).
- [2] S. Agmon, A. Douglis, L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I. Comm. Pure Appl. Math. 12 (1959), 623–727.
- [3] A.Bahri and P.L.Lions, Solutions of superlinear elliptic equations and their Morse indices. Comm. Pure Appl. Math. 45 (1992), 1205-1215.
  [4] S. Blatt, Monotonicity formulas for extrinsic triharmonic maps and the triharmonic Lane–Emden equation. Journal of Differential Equations
- 262 (2017), 5691-5734.
  [5] C. Cowan, *Liouville theorems for stable Lane-Emden systems and biharmonic problems*, Nonlinearity, 26 (2013), 2357-2371.
- [6] M. G. Crandall and P. H. Rabinowitz, Some continuation and variation methods for positive solutions of nonlinear elliptic eigenvalue problems. Arch. Rat. Mech. Anal. 58 (1975), 207-218.
- [7] E.N. Dancer, Finite Morse index solutions of supercritical problems. J. Reine Angew Math. 620 (2008), 213-233.
- [8] J. Dávila, L. Dupaigne and A.Farina, Partial regularity of finite Morse index solutions to the Lane-Emden equation, J. Funct. Anal. 261 (2011), 218-232.
- J. Dávila, L. Dupaigne, K. Wang and J. Wei, A monotonicity formula and a Liouville-type theorem for a fourth order supercritical problem. Adv. Math., 258 (2014), 240–285.
- [10] L. Damascelli, A. Farina, B. Sciunzi and E. Valdinoci, Liouville results for m- Laplace equations of Lane-Emden Fowler type. Ann. I. H. Poincaré. 26 (2009), 1099-1119.
- [11] A. Farina, On the classification of solutions of the Lane–Emden equation on unbounded domains of  $\mathbb{R}^N$ , J. Math. Pures Appl. 87 (2007), 537–561.
- [12] B. Gidas and J. Spruck, A priori bounds for positive solutions of nonlinear elliptic equations, Comm. Partial Differential Equations. 6 (1981), 883-901.
- [13] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, 2nd edition, Grundlehren 224, Springer, Berlin-Heidelberg-New York-Tokyo (1983).
- [14] F. Gazzola, H. Christoph Grunau, G. Sweers, *Polyharmonic Boundary Value Problems*, Lecture Notes in Mathematics, vol.1991, Springer, 2010.
- [15] H. Hajlaoui, A. Harrabi and F. Mtiri, Morse indices of solutions for super-linear elliptic PDEs, Nonlinear Analysis 116 (2015), 180-192.
- [16] H. Hajlaoui, A. Harrabi and D. Ye, On stable solutions of the biharmonic problem with polynomial growth, Pacific J. Math. 270 (2014), 79-93.
- [17] M. K. Hamdani, A. Harrabi,  $L^{\infty}$ -norm Estimates of Weak Solutions via their Morse indices for the m-Laplacian Problems, Results in Mathematics. **74** (2019), article number (69).
- [18] A. Harrabi, High-order Bahri-Lions Liouville type theorems, Annali di Matematica Pura ed Applicata. 198 (2019), 1675–1692.
- [19] A. Harrabi, F. Mtiri, *Liouville-type theorems and existence results for stable at infinity solutions of higher order m-polyharmonic problems*, To appear in JMAA (2021).
- [20] A. Harrabi, F. Mtiri and D. Ye, Explicit L<sup>∞</sup>-norm estimates via Morse index, the bi-harmonic and tri-harmonic semilinear problems, Manuscripta Math. 270 (2019), 57–79.
- [21] A. Harrabi and B. Rahal, On the sixth-order Joseph-Lundgren exponent, Ann. Henri Poincaré. 18 (2017), 1055-1094.
- [22] A. Harrabi, S. Rebhi and A. Selmi, Solutions of superlinear equations and their Morse indices I, Duke. Math. J. 94 (1998), 141-157.
- [23] F. Mtiri and D. Ye, *Liouville theorems for stable at infinity solutions of Lane-Emden system*, Nonlinearity **32** (2019), 910-926.

- [24] S.I. Pohozaev, *Eigenfunctions of the equation*  $\Delta u + \lambda f(u) = 0$ . Sov. Math. Doklady **5** (1965), 1408-1411.
- [25] P. Polácik, P. Quittner and P. Souplet, Singularity and decay estimates in superlinear problems via Liouville-type theorems. I. Elliptic equations and systems, Duke Math. Journal. 139 (3) (2007), 555-579.
- [26] W. Reichel and T. weth, A priori bounds and a Liouville theorem on a half space for higher-order elliptic Dirichlet problems, Math. Z. 261 (2009), 805-827.
- [27] J. Serrin and H. Zou, *Cauchy-Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities*, Acta Math. **189** (2002), 79-142.
- [28] B. Sirakov, Existence results and a priori bounds for higher order elliptic equations and systems, J.Math.Pures.Appl 9 89 (2008), no. 2, 114-133.
- [29] J. Wei and D. Ye, Liouville Theorems for stable solutions of biharmonic problem, Math, Ann. 356 (4) (2013), 1599-1612.
- [30] X.F. Yang, Nodal Sets and Morse Indices of Solutions of Super-linear Elliptic PDE's. J. Funct. Anal. 160 (1998), 223-253.