

Spin-orbit correlations and exchange-bias control in twisted Janus dichalcogenide multilayers

D. Soriano^{1,2} and J. L. Lado³

¹*Radboud University, Institute for Molecules and Materials, NL-6525 AJ Nijmegen, the Netherlands*

²*Dipartimento di Ingegneria dell'Informazione, Università di Pisa, 56122 Pisa, Italy*

³*Department of Applied Physics, Aalto University, 00076 Aalto, Espoo, Finland*

(Dated: March 7, 2022)

Janus dichalcogenide multilayers provide a paradigmatic platform to engineer electronic phenomena dominated by spin-orbit coupling. Their unique spin-orbit effects stem from the mirror symmetry breaking in each layer, which induces a colossal Rashba spin-orbit effect in comparison with the conventional dichalcogenide counterparts. Here we put forward twisted dichalcogenide bilayers as a simple platform to realize spin-orbit correlated states. We demonstrate the emergence of flat bands featuring strong spin-momentum locking and the emergence of symmetry broken states with associated non-coplanar magnetization when interactions are included. We further show that the symmetry broken states can be controlled by means of a magnetic substrate, strongly impacting the non-coplanar magnetic texture of the moire unit cell. Our results put forward twisted Janus multilayers as a powerful platform to explore spin-orbit correlated physics, and highlighting the versatility of magnetic substrates to control unconventional moire magnetism.

I. INTRODUCTION

Dichalcogenide van der Waals materials^{1–6} are paradigmatic two dimensional compounds featuring strong spin-orbit coupling effects^{7–13}. The strong spin-momentum locking has motivated a variety of proposals for spin-orbitronic control.^{14–19} These phenomena comes from the intrinsic Ising spin-orbit coupling, stemming from the inversion symmetry breaking of the structure. Interestingly, a special family of dichalcogenides, known as Janus dichalcogenides^{20–23}, provide a unique new form of spin-orbit physics to these materials. Janus dichalcogenides feature an intrinsic mirror-symmetry breaking adding strong Rashba spin-orbit coupling effects^{24–26} to the already rich physics of conventional dichalcogenides.

Twist engineering has risen as a powerful knob to control the electronic structure of dichalcogenides in particular,^{27–32} and van der Waals materials in general.^{33–37} The emergence of flat bands and correlated states in dichalcogenide systems represent an illustrative example of the versatility of twist engineering in these systems.^{38–44} Spin-orbit coupling effects in these flat band systems^{45–51} have also been proposed to bring up unique possibilities for spin and valley control. However, the emergence of correlated states in Janus dichalcogenides has remained largely unexplored, and studies on spin-orbit effects have focused mainly on Ising spin-orbit coupling.

Here we put forward twisted Janus systems (Fig. 1a) as a paradigmatic example in which correlated states emerge in flat bands dominated by spin-momentum texture in reciprocal space. In particular, we demonstrate the emergence of flat bands dominated by Rashba spin-orbit coupling effects in strike contrast with conventional twisted dichalcogenide systems. When including electronic interactions, we show that these flat bands lead to non-coplanar magnetic states in real space, stemming




FIG. 1. (a) Sketch of a twisted Janus dichalcogenide, showing in different colors the different chalcogenide atoms. Panel (b) shows the low energy band structure of a monolayer Janus dichalcogenide as the strength of the Rashba spin-orbit coupling λ_R is increased in comparison with the Ising spin-orbit coupling λ_I . Panel (c) shows the moire bands of twisted Janus system at a rotation angle of 2° , at the bottom of the conduction band. Panel (d) shows an illustration of the low energy model, and panel (e) shows the different terms in the Hamiltonian. We took $\lambda_I = 0.02t$ in (b,c).

from the strong Rashba effects of the parent compounds. We finally show how the exchange proximity effect allows controlling these correlated states by combining Janus dichalcogenide bilayers with two-dimensional ferromagnets. Our results highlight the spin-orbit dominated




FIG. 2. (a,b) Band structure of a twisted Janus dichalcogenide at a twist angle of 9° , showing the S_x (a) and S_z (b) spin textures in reciprocal space. The strong spin-momentum texture arises from the intrinsic mirror symmetry breaking of each Janus monolayer, that is inherited by the twisted bilayer system. We took $\lambda_I = 0.1\lambda_R$ and $\lambda_I = 0.02t$ in (a,b).

physics of twisted Janus dichalcogenide systems and their versatility for controllable correlated spin-orbitronics. The manuscript is organized as follows. In Section II we show the impact of mirror symmetry breaking in twisted dichalcogenide systems. In Section III we show the emergence of correlations in the twisted Janus system, together with the emergence of non-coplanar magnetism. In Section IV we show how the correlated states can be controlled by the exchange proximity effect. Finally, in Section V we summarize our conclusions.

II. SPIN-TEXTURE IN TWISTED JANUS BILAYERS

We first address the emergence of flat bands in the Janus van der Waals heterostructure, and in particular, how the different types of spin-orbit affect the low energy electronic structure. In particular, Janus materials have coexisting types of spin-orbit coupling, namely Ising and Rashba^{24–26}, whose relative strength substantially impact the low energy electronic dispersion (Fig. 1b). Such interplay of spin-orbit effects further impacts the quasi flat bands of a twisted heterostructure (Fig. 1c), and therefore would be expected to have an important impact in potentially correlated states. To model these systems, we consider an effective real-space model for the Janus material that incorporates the effects of intrinsic Ising spin-orbit coupling and Rashba spin-orbit coupling. We will focus the discussion on the conduction band of the model, yet an analogous discussion can be performed for the valence band. The Janus dichalcogenide system is captured by an effective honeycomb lattice model (Fig.

1d) of the following form

$$\begin{aligned} \mathcal{H} = & \sum_{\langle ij \rangle, s} t c_{i,s}^\dagger c_{j,s} + m \sum_{i,s} \vartheta_{ii}^z c_{i,s}^\dagger c_{i,s} \\ & + \sum_{i,j,s} t^\perp(\mathbf{r}_i, \mathbf{r}_j) c_{i,s}^\dagger c_{j,s} \\ & + i\lambda_I \sum_{\langle\langle ij \rangle\rangle, s, s'} \nu_{ij} \sigma_{s,s'}^z c_{i,s}^\dagger c_{j,s'} \\ & + i\lambda_R \sum_{\langle ij \rangle, s, s'} \mathbf{d}_{ij} \cdot \boldsymbol{\sigma}_{s,s'} c_{i,s}^\dagger c_{j,s'} \end{aligned} \quad (1)$$

where t is a first neighbor hopping, $\langle \rangle$ denotes sum over first neighbors in a layer, m denotes the onsite energy imbalance between the two sites of the honeycomb lattice, $t^\perp(\mathbf{r}_i, \mathbf{r}_j)$ parametrize the interlayer coupling,^{52–54} ϑ_{ii}^z is the sublattice Pauli matrix, $\sigma_{s,s'}^z$ is the spin Pauli matrix, λ_I controls the Ising spin-orbit coupling $\nu_{ij} = \pm 1$ for clock-wise/anti-clockwise hopping,⁵⁵ $\langle\langle \rangle\rangle$ denotes second neighbors in a layer, λ_R controls the Rashba spin-orbit coupling⁵⁶ and $\mathbf{d}_{ij} = (\mathbf{r}_i - \mathbf{r}_j) \times \hat{z}$. An illustration of the different terms of the effective model of a Janus monolayer is shown in Fig. 1e. For a conventional dichalcogenide system, i.e., non-Janus one, $\lambda_R = 0$ due to the presence of mirror symmetry in the layer, and the Hamiltonian has $U(1)$ rotational spin symmetry. For a Janus system, $\lambda_R \neq 0$ breaks the $U(1)$ rotational spin-symmetry, creating a spin-momentum texture in reciprocal space. For the sake of simplicity, we take the same sign of λ_R for both layers, which physically means the two systems have the same mirror symmetry breaking. The values of λ_I and λ_R are material specific, and can be controlled by choosing different dichalcogenide semiconductors^{57,58}. From the electronic structure point of view, the value of λ_I controls the splitting at the K-point, whereas the value of λ_R controls the momentum offset of the two spin channels. As a reference, λ_R can reach up to 50 meV in WSeTe⁵⁸, and λ_I can range between 1 and 100 meV in different dichalcogenides⁵⁷. Such parameters can be extracted from the spin-splitting of the electronic structure computed with first principle methods including spin-orbit coupling effects^{57,58}. In the single-layer limit, the model of Eq. 1 captures the dispersion of a Janus monolayer in the presence of spin-orbit coupling. We finally note that this model captures mainly the electronic structure at low energies and that multiorbital Wannier models⁵⁹ could be used to capture the physics at higher energies.

We now move on to discuss in more detail the electronic structure of the twisted Janus system. In the presence of a twist, the moire length L_M emerges and can be controlled by the twist angle, and the electronic structure now is defined in the mini-Brillouin zone. The strong spin-momentum locking of the monolayer system is inherited by the electronic structure of the twisted Janus system, as shown in Fig. 2. In particular, we first take




FIG. 3. (a) Spin texture in momentum space of the lowest conduction band of a twisted Janus bilayer at an angle of 9° . Vortices around the Γ (b), M (c) and K (d) are clearly observed. The arrows denote the expectation value $(\langle S_x \rangle, \langle S_y \rangle)$ for that eigenstate. We took $\lambda_I = 0.1\lambda_R$ and $\lambda_I = 0.02t$.

a twisted system at a large twist angle of 9° degrees and compute the spin expectation value of the S_x operator and S_z operator in reciprocal space. As shown in Fig. 2ab, we observe a strong texture of the S_x operator in reciprocal space (Fig. 2a), with the S_z operator strongly mixed (Fig. 2b). This behavior is in stark contrast with non-Janus systems, in which the Ising spin-orbit coupling creates spin-polarized moire bands in reciprocal space.

The spin-momentum texture of the Janus system can be further explored by computing the spin projection in the lowest conduction band. In particular, we show in Fig. 3 a quiver map of the lowest conduction band, where the arrows denote the expectation value of the spin operators S_x and S_y , and the color the eigenvalue of the state. A strong spin texture in the whole reciprocal space is clearly seen (Fig. 3a). In particular, spin texture vortices are observed around the Γ (Fig. 3b), M (Fig. 3c) and K (Fig. 3d) points. This in-plane spin texture is absent in conventional twisted dichalcogenide multilayers and therefore is a unique feature of twisted Janus bilayers. This spin texture will heavily influence the correlated states in the system, as we discuss in the next section.

III. INTERACTION-INDUCED SYMMETRY BREAKING IN JANUS SYSTEMS

After having addressed the single-particle electronic structure, we now move on to consider the effect of

electron-electron interactions, focusing on a system with a smaller twist angle and flatter bands. We include non-local interactions up second neighbor sites in our atomistic model, taking the form

$$\begin{aligned} \mathcal{H}_I = & U \sum_i \left(c_{i,\uparrow}^\dagger c_{i,\uparrow} - \frac{1}{2} \right) \left(c_{i,\downarrow}^\dagger c_{i,\downarrow} - \frac{1}{2} \right) \\ & + V_1 \sum_{\langle ij \rangle} \left[\left(\sum_s c_{i,s}^\dagger c_{i,s} \right) \left(\sum_s c_{j,s}^\dagger c_{j,s} \right) \right] \\ & + V_2 \sum_{\langle\langle ij \rangle\rangle} \left[\left(\sum_s c_{i,s}^\dagger c_{i,s} \right) \left(\sum_s c_{j,s}^\dagger c_{j,s} \right) \right] \end{aligned} \quad (2)$$

where U parametrizes the local Hubbard interaction, V_1 the first neighbor interaction and V_2 the second neighbor interaction. By solving the previous Hamiltonian via a mean-field approximation, we can address the interaction-induced symmetry broken states in the Janus system. Importantly, we will exploit a fully rotational invariant mean-field formalism capable of capturing non-coplanar formalism. Our mean-field formalism includes all the Wick contractions in the mean-field ansatz, which in particular captures interaction-induced real-space non-coplanar moire spin-textures. For the sake of concreteness, we will focus on correlated states in the case with two electrons per unit cell.

We now address the impact of interactions in the twisted Janus system. We show in Fig. 4a the low energy electronic structure for a twisting angle of 3.5° , featuring four nearly flat bands with strong spin-momentum texture. Once we include electronic interactions at the mean-field level, we observe the emergence of a ferromagnetic state at half-filling of the low energy bands, as shown in Fig. 4b. The low energy bands in the absence of interactions form a triangular lattice in the moire supercell, as shown in the density of states of Fig. 4c. The real-space spin density in the correlated regime also presents a non-zero expectation value in the emergent moire triangular lattice, as shown in Fig. 4d. This phenomenology is, of course, common to most of the moire systems featuring strongly localized modes in real space. However, more interestingly, the moire system displays a strong non-coplanar magnetization in real space, forming a meron texture around each moire hot spot. It is worth to note that the local order parameter is not only non-collinear, but also that is not contained in a plane and thus we refer to it as non-coplanar. The emergence of such non-coplanar real-space spin texture stems from the intrinsic mirror symmetry breaking of each Janus monolayer, and in particular, from the intrinsic strong Rashba spin-orbit coupling.

We now comment on some observable transport signatures associated to this correlated state. The existence of a non-coplanar magnetic texture is expected to have an important impact on the electronic transport properties




FIG. 4. (a) Non-interacting and (b) selfconsistent electronic structure of the twisted Janus dichalcogenide at a twist angle of 3.5° . Before interactions, the nearly flat bands form a effective triangular lattice in real-space (c). The interaction-induced magnetization displays a strong texture in real-space as shown in panel (d), stemming from the Rashba spin-orbit coupling of the twisted dichalcogenide. L_M denotes the moire length. We took $U = 1.5t$, $V_1 = 1.2t$ and $V_2 = 0.3t$ and $\lambda_I = 0.1\lambda_R$ and $\lambda_I = 0.02t$.

of the system. Upon small doping away from the correlated state, free electrons will feel a strong non-coplanar spin-texture in real space that will lead to anomalous Hall effect⁶⁰. The emergence of this anomalous Hall state is well-known in other non-coplanar magnetic systems, with the paradigmatic case of skyrmion materials^{61–63}. In the strong coupling limit, by performing a local gauge rotation to the local spin axis, the effective equations of motion for electrons acquire an effective gauge field associated with the non-coplanar magnetization⁶¹. This new term enters as an anomalous velocity in the equations of motion, and it is known to give rise to the so-called anomalous skyrmion Hall effect⁶¹. Interestingly, beyond the anomalous Hall effect stemming from the real-space non-coplanar spin texture, the current system can potentially have additional contributions to the anomalous velocity stemming from the combination of spin-orbit coupling and magnetism⁵⁶. It is also worth to mention that, apart from the anomalous Hall velocity for electrons, magnetic excitations of the symmetry broken state can also show topological features. In particular, magnon excitations of the ferromagnetic state can also potentially display anomalous Hall effect^{64,65}. At the microscopic level, this stems from the Dzyaloshinskii–Moriya interaction of the effective spin model^{66,67}, which naturally arises from the intrinsic Rashba spin-orbit coupling in the electronic Hamiltonian⁶⁸.




FIG. 5. Twisted Janus bilayer with in-plane (a,c,e,g) and out-of-plane (b,d,f,h) magnetic encapsulation. Panels (a,b) show a sketch of the magnetically encapsulated Janus bilayer, panels (c,d) the non-interacting band structure, panels (e,f) the interacting band structure and panels (g,h) the real space interaction-induced magnetization texture. It is observed that the direction of the antiferromagnetic encapsulation impact both the band structure and real space magnetic texture distribution. We took $U = 0.8t$, $V_1 = 0.6t$ and $V_2 = 0.2t$, $\lambda_I = 0.1\lambda_R$ and $\lambda_I = 0.02t$, and twist angle of 4.3° .

IV. EXCHANGE CONTROLLED CORRELATIONS IN TWISTED JANUS DICHALCOGENIDES

We now move on to consider the impact of a magnetic encapsulation, and in particular, how the correlated state is impacted by it. Twisted Janus bilayer can be encapsulated between other two-dimensional materials. While usually boron-nitride encapsulation is used, other different insulators can be considered. In particular, the use




FIG. 6. Sketch of the ferromagnet/Janus interface with out-of-plane magnetization (a). Panel (b) shows the electronic bands, highlighting the existence of both Ising and Rashba spin-orbit coupling. Panel (c) shows a zoom at the K-point, highlighting the exchange splitting induced by the ferromagnet at the valleys.

of van der Waals ferromagnetic insulators^{69–71} such as CrCl_3 ⁷², CrBr_3 ⁷³, and CrI_3 ⁷⁴ as encapsulation^{75–79} provide an interesting possibility with regards to controlling a correlated state in the twisted Janus bilayer. We now will show how a magnetic encapsulation allows controlling the underlying electronic structure of the Janus system, and in particular, tuning the non-coplanar magnetic order of the correlated state. For this purpose, we now include a new term in the Hamiltonian that accounts for the impact of the effect of a magnetic substrate in the Janus bilayer, taking the form

$$\mathcal{H}_J = \sum_{i,s,s'} \tau_{ii} \mathbf{J} \cdot \sigma_{s,s'} c_{i,s}^\dagger c_{i,s'} \quad (3)$$

where $\tau_{ii} = \pm 1$ for the upper/lower layer, and $\mathbf{J}$ denotes the strength of the exchange field. We focus on the case in which the ferromagnetic encapsulation is aligned antiferromagnetically, as expected from the superexchange between layers mediated by the Janus bilayer. We will compare two scenarios for the magnetic encapsulation (Fig. 5ab), in-plane ferromagnetics such as the case of CrCl_3 with $\mathbf{J} = (J, 0, 0)$, out-of-plane ferromagnets such as the case of CrBr_3 and CrI_3 with $\mathbf{J} = (0, 0, J)$. Let us first discuss the non-interacting electronic structure of the magnetically encapsulated bilayer shown in Fig. 5cd. Depending on the alignment of the magnetic encapsulation, a big change in the low energy dispersion is observed, stemming from the combination of exchange proximity effect, Ising spin-orbit coupling, and Rashba spin-orbit coupling.

We now move on to consider the impact of interactions, shown in the interacting electronic structure of Fig. 5ef. It is interesting to note that the interaction-induced complex non-coplanar order does not impact the valley sector of the twisted dichalcogenide system. This can be verified by computing the expectation value of the valley

operator $\mathcal{V}_z$ ^{80–82}, defined in the real-space atomistic basis as $\mathcal{V}_z = \frac{i}{3\sqrt{3}} \sum_{\langle\langle ij \rangle\rangle, s} \vartheta_{ij}^z \nu_{ij} c_{i,s}^\dagger c_{j,s}$. We observe that the emergence of non-coplanar order does not create intervalley mixing, demonstrating that the complex spin texture coexists with the emergent valley conservation of the twisted dichalcogenide system. It is finally worth to emphasize that different electronic fillings can potentially lead to spontaneously valley mixed states, as proposed for twisted graphene multilayers^{78,83,84}.

More interestingly, the spatial distribution of the interaction-induced magnetic state is shown to depend strongly on the magnetic encapsulation, as shown in Fig. 5gh. In particular, we observe that the encapsulation creates a redistribution of the direction and magnitude of the magnetic moment in the z-direction in real space. This phenomenology stems from the non-trivial interference between Wannier states of the moire unit cell, whose dependence is inherited from the underlying electronic structure featuring strong spin-orbit coupling. It is worth emphasizing that in the absence of spin-orbit coupling effects, such redistribution would not happen, and therefore this is a genuine effect arising from the interplay of spin-orbit and strong interactions.

So far we have based our analysis on an effective low energy model, in particular including the impact of the ferromagnet as an effective exchange field. We now highlight via first-principles methods that this assumption is correct a specific heterostructure, in particular taking SWeSe as the dichalcogenide and CrBr_3 as the ferromagnetic insulator. We take a minimal heterostructure (Fig. 6a) capturing the interplay of Ising spin-orbit coupling, Rashba spin-orbit coupling and exchange proximity magnetism, namely a $\text{CrBr}_3/\text{SWeSe}$ heterostructure. Our calculations are performed with Quantum Espresso⁸⁵, including correlations in the ferromagnet by means of the DFT+U formalism and with LDA exchange-correlation functional⁸⁶, and PAW pseudopotentials⁸⁷. As shown in Fig. 6b, we observe that the electronic structure of the ferromagnet does not get mixed with the Janus system, highlighting that at low energies, the ferromagnet can be integrated out. By zooming at the valleys (Fig. 6c), we observe that the combination of spin-orbit coupling and exchange proximity effect breaks the original $+k \rightarrow -k$ degeneracy of the electronic structure, demonstrating the existence of the exchange proximity effect. It is worth to note that, although this calculation focuses on the interplay of spin-orbit effect and exchange proximity, correlated states could be studied purely from first-principles methods^{28,51,88–91}. These calculations would be nevertheless a remarkable challenge from the computational point of view due to the existence of non-coplanar magnetism, spin-orbit coupling, and a large number of atoms in the moire unit cell.

V. CONCLUSIONS

We have shown that a twisted Janus dichalcogenide bilayer displays an electronic structure dominated by Rashba spin-orbit coupling effects. In particular, we showed the emergence of spin-momentum spin-textures in the electronic structure, which are inherited by the flat band regime of the twisted Janus dichalcogenide system. Upon including electron-electron interactions, we showed that a non-coplanar real-space texture emerges in each unit cell of the moire system. We further showed that the non-coplanar symmetry broken state could be controlled by magnetically encapsulating the system, providing a simple means of engineering controllable non-coplanar

moire magnetism. Our results put forward twisted Janus dichalcogenide bilayers as a versatile platform to engineer spin-orbit correlated states in moire systems, exemplifying the critical role that relativistic effects can have in flat band materials.

Acknowledgments: D.S. thanks financial support from EU through the MSCA project Nr. 796795 SOT-2DvdW. J.L.L. acknowledges the computational resources provided by the Aalto Science-IT project, and the financial support from Academy of Finland Projects No. 331342 and No. 336243. We thank P. Liljeroth, S. Kezilebieke, T. Heikkilä, F. Aikebaier, T. Wolf, P. Törmä and A. Ramires for fruitful discussions.

-
- ¹ B. Radisavljevic, A. Radenovic, J. Brivio, V. Giacometti, and A. Kis, “Single-layer MoS2 transistors,” *Nature Nanotechnology* **6**, 147–150 (2011).
 - ² Andrea Splendiani, Liang Sun, Yuanbo Zhang, Tianshu Li, Jonghwan Kim, Chi-Yung Chim, Giulia Galli, and Feng Wang, “Emerging photoluminescence in monolayer MoS2,” *Nano Letters* **10**, 1271–1275 (2010).
 - ³ Sajede Manzeli, Dmitry Ovchinnikov, Diego Pasquier, Oleg V. Yazyev, and Andras Kis, “2d transition metal dichalcogenides,” *Nature Reviews Materials* **2** (2017), 10.1038/natrevmats.2017.33.
 - ⁴ Kin Fai Mak, Changyu Lee, James Hone, Jie Shan, and Tony F. Heinz, “Atomically thin mos2: A new direct-gap semiconductor,” *Phys. Rev. Lett.* **105**, 136805 (2010).
 - ⁵ Roscoe G. Dickinson and Linus Pauling, “THE CRYSTAL STRUCTURE OF MOLYBDENITE,” *Journal of the American Chemical Society* **45**, 1466–1471 (1923).
 - ⁶ J.A. Wilson and A.D. Yoffe, “The transition metal dichalcogenides discussion and interpretation of the observed optical, electrical and structural properties,” *Advances in Physics* **18**, 193–335 (1969).
 - ⁷ Andor Kormányos, Viktor Zólyomi, Neil D. Drummond, and Guido Burkard, “Spin-orbit coupling, quantum dots, and qubits in monolayer transition metal dichalcogenides,” *Phys. Rev. X* **4**, 011034 (2014).
 - ⁸ Kin Fai Mak, Keliang He, Jie Shan, and Tony F. Heinz, “Control of valley polarization in monolayer MoS2 by optical helicity,” *Nature Nanotechnology* **7**, 494–498 (2012).
 - ⁹ Hualing Zeng, Junfeng Dai, Wang Yao, Di Xiao, and Xiaodong Cui, “Valley polarization in MoS2 monolayers by optical pumping,” *Nature Nanotechnology* **7**, 490–493 (2012).
 - ¹⁰ Ting Cao, Gang Wang, Wenpeng Han, Huiqi Ye, Chuanrui Zhu, Junren Shi, Qian Niu, Pingheng Tan, Enge Wang, Baoli Liu, and Ji Feng, “Valley-selective circular dichroism of monolayer molybdenum disulphide,” *Nature Communications* **3** (2012), 10.1038/ncomms1882.
 - ¹¹ G. Sallen, L. Bouet, X. Marie, G. Wang, C. R. Zhu, W. P. Han, Y. Lu, P. H. Tan, T. Amand, B. L. Liu, and B. Urbaszek, “Robust optical emission polarization in mos2 monolayers through selective valley excitation,” *Phys. Rev. B* **86**, 081301 (2012).
 - ¹² Xiaodong Xu, Wang Yao, Di Xiao, and Tony F. Heinz, “Spin and pseudospins in layered transition metal dichalcogenides,” *Nature Physics* **10**, 343–350 (2014).
 - ¹³ Kin Fai Mak and Jie Shan, “Photonics and optoelectronics of 2d semiconductor transition metal dichalcogenides,” *Nature Photonics* **10**, 216–226 (2016).
 - ¹⁴ Manuel Offidani, Mirco Milletari, Roberto Raimondi, and Aires Ferreira, “Optimal charge-to-spin conversion in graphene on transition-metal dichalcogenides,” *Phys. Rev. Lett.* **119**, 196801 (2017).
 - ¹⁵ C. K. Safeer, Josep Ingla-Aynés, Franz Herling, José H. Garcia, Marc Vila, Nerea Ontoso, M. Reyes Calvo, Stephan Roche, Luis E. Hueso, and Fèlix Casanova, “Room-temperature spin hall effect in graphene/MoS2 van der waals heterostructures,” *Nano Letters* **19**, 1074–1082 (2019).
 - ¹⁶ Qiming Shao, Guoqiang Yu, Yann-Wen Lan, Yumeng Shi, Ming-Yang Li, Cheng Zheng, Xiaodan Zhu, Lain-Jong Li, Pedram Khalili Amiri, and Kang L. Wang, “Strong rashba-edelstein effect-induced spin-orbit torques in monolayer transition metal dichalcogenide/ferromagnet bilayers,” *Nano Letters* **16**, 7514–7520 (2016).
 - ¹⁷ J. B. S. Mendes, A. Aparecido-Ferreira, J. Holanda, A. Azevedo, and S. M. Rezende, “Efficient spin to charge current conversion in the 2d semiconductor MoS2 by spin pumping from yttrium iron garnet,” *Applied Physics Letters* **112**, 242407 (2018).
 - ¹⁸ Yunqiu Kelly Luo, Jinsong Xu, Tiancong Zhu, Guanzhong Wu, Elizabeth J. McCormick, Wenbo Zhan, Mahesh R. Neupane, and Roland K. Kawakami, “Opto-valleytronic spin injection in monolayer MoS2/few-layer graphene hybrid spin valves,” *Nano Letters* **17**, 3877–3883 (2017).
 - ¹⁹ Ahmet Avsar, Dmitrii Unuchek, Jiawei Liu, Oriol Lopez Sanchez, Kenji Watanabe, Takashi Taniguchi, Barbaros Özyilmaz, and Andras Kis, “Optospintronics in graphene via proximity coupling,” *ACS Nano* **11**, 11678–11686 (2017).
 - ²⁰ Ang-Yu Lu, Hanyu Zhu, Jun Xiao, Chih-Piao Chuu, Yimo Han, Ming-Hui Chiu, Chia-Chin Cheng, Chih-Wen Yang, Kung-Hwa Wei, Yiming Yang, Yuan Wang, Dimosthenis Sokaras, Dennis Nordlund, Peidong Yang, David A. Muller, Mei-Yin Chou, Xiang Zhang, and Lain-Jong Li, “Janus monolayers of transition metal dichalcogenides,” *Nature Nanotechnology* **12**, 744–749 (2017).
 - ²¹ Liang Dong, Jun Lou, and Vivek B. Shenoy, “Large in-plane and vertical piezoelectricity in janus transition metal

- dichalcogenides,” *ACS Nano* **11**, 8242–8248 (2017).
- 22 Fengping Li, Wei Wei, Pei Zhao, Baibiao Huang, and Ying Dai, “Electronic and optical properties of pristine and vertical and lateral heterostructures of janus MoSSe and WSSe,” *The Journal of Physical Chemistry Letters* **8**, 5959–5965 (2017).
 - 23 H. U. Din, M. Idrees, Arwa Albar, M. Shafiq, Iftikhar Ahmad, Chuong V. Nguyen, and Bin Amin, “Rashba spin splitting and photocatalytic properties of GeC – mSSe ($m = \text{Mo, w}$) van der waals heterostructures,” *Phys. Rev. B* **100**, 165425 (2019).
 - 24 Roberto Sant, Maxime Gay, Alain Marty, Simone Lisi, Rania Harrabi, Céline Vergnaud, Minh Tuan Dau, Xiaorong Weng, Johann Coraux, Nicolas Gauthier, Olivier Renault, Gilles Renaud, and Matthieu Jamet, “Synthesis of epitaxial monolayer janus SPtSe,” *npj 2D Materials and Applications* **4** (2020), 10.1038/s41699-020-00175-z.
 - 25 Idris Smaili, Slimane Laref, Jose H. Garcia, Udo Schwingenschlgl, Stephan Roche, and Aurelien Manchon, “Janus Monolayers of Magnetic Transition Metal Dichalcogenides as an All-in-One Platform for Spin-Orbit Torque,” arXiv e-prints, arXiv:2007.07579 (2020), arXiv:2007.07579 [cond-mat.mes-hall].
 - 26 Marko M. Petrić, Malte Kremser, Matteo Barbone, Ying Qin, Yasir Sayyad, Yuxia Shen, Sefaattin Tongay, Jonathan J. Finley, Andrés R. Botello-Méndez, and Kai Müller, “Raman spectrum of janus transition metal dichalcogenide monolayers wsse and mosse,” *Phys. Rev. B* **103**, 035414 (2021).
 - 27 Fengcheng Wu, Timothy Lovorn, Emanuel Tutuc, Ivar Martin, and A. H. MacDonald, “Topological insulators in twisted transition metal dichalcogenide homobilayers,” *Phys. Rev. Lett.* **122**, 086402 (2019).
 - 28 Mit H. Naik and Manish Jain, “Ultraflatbands and shear solitons in moiré patterns of twisted bilayer transition metal dichalcogenides,” *Phys. Rev. Lett.* **121**, 266401 (2018).
 - 29 Indrajit Maity, Mit H. Naik, Prabal K. Maiti, H. R. Krishnamurthy, and Manish Jain, “Phonons in twisted transition-metal dichalcogenide bilayers: Ultrasoft phasons and a transition from a superlubric to a pinned phase,” *Phys. Rev. Research* **2**, 013335 (2020).
 - 30 C. H. Lui, Zhipeng Ye, Chao Ji, Kuan-Chang Chiu, Cheng-Tse Chou, Trond I. Andersen, Casie Means-Shively, Heidi Anderson, Jenn-Ming Wu, Tim Kidd, Yi-Hsien Lee, and Rui He, “Observation of interlayer phonon modes in van der waals heterostructures,” *Phys. Rev. B* **91**, 165403 (2015).
 - 31 Aaron M. Jones, Hongyi Yu, Nirmal J. Ghimire, Sanfeng Wu, Grant Aivazian, Jason S. Ross, Bo Zhao, Jiaqiang Yan, David G. Mandrus, Di Xiao, Wang Yao, and Xiaodong Xu, “Optical generation of excitonic valley coherence in monolayer WSe₂,” *Nature Nanotechnology* **8**, 634–638 (2013).
 - 32 Britton W. H. Baugher, Hugh O. H. Churchill, Yafang Yang, and Pablo Jarillo-Herrero, “Optoelectronic devices based on electrically tunable p–n diodes in a monolayer dichalcogenide,” *Nature Nanotechnology* **9**, 262–267 (2014).
 - 33 J. M. B. Lopes dos Santos, N. M. R. Peres, and A. H. Castro Neto, “Continuum model of the twisted graphene bilayer,” *Phys. Rev. B* **86**, 155449 (2012).
 - 34 J. M. B. Lopes dos Santos, N. M. R. Peres, and A. H. Castro Neto, “Graphene bilayer with a twist: Electronic structure,” *Phys. Rev. Lett.* **99**, 256802 (2007).
 - 35 Xing-Ju Zhao, Yang Yang, Dong-Bo Zhang, and Su-Huai Wei, “Formation of bloch flat bands in polar twisted bilayers without magic angles,” *Phys. Rev. Lett.* **124**, 086401 (2020).
 - 36 Mingyuan Chen, Xiao Lin, Thao H. Dinh, Zhiren Zheng, Jialiang Shen, Qiong Ma, Hongsheng Chen, Pablo Jarillo-Herrero, and Siyuan Dai, “Configurable phonon polaritons in twisted α -MoO₃,” *Nature Materials* **19**, 1307–1311 (2020).
 - 37 Kasra Hejazi, Zhu-Xi Luo, and Leon Balents, “Non-collinear phases in moiré magnets,” *Proceedings of the National Academy of Sciences* **117**, 10721–10726 (2020).
 - 38 Fengcheng Wu, Timothy Lovorn, Emanuel Tutuc, and A. H. MacDonald, “Hubbard model physics in transition metal dichalcogenide moiré bands,” *Phys. Rev. Lett.* **121**, 026402 (2018).
 - 39 Jue Wang, Qianhui Shi, En-Min Shih, Lin Zhou, Wenjing Wu, Yusong Bai, Daniel Rhodes, Katayun Barmak, James Hone, Cory R. Dean, and X.-Y. Zhu, “Diffusivity reveals three distinct phases of interlayer excitons in mose₂/wse₂ heterobilayers,” *Phys. Rev. Lett.* **126**, 106804 (2021).
 - 40 Chenhao Jin, Emma C. Regan, Aiming Yan, M. Iqbal Bakti Utama, Danqing Wang, Sihan Zhao, Ying Qin, Sijie Yang, Zhiren Zheng, Shenyang Shi, Kenji Watanabe, Takashi Taniguchi, Sefaattin Tongay, Alex Zettl, and Feng Wang, “Observation of moiré excitons in WSe₂/WS₂ heterostructure superlattices,” *Nature* **567**, 76–80 (2019).
 - 41 Emma C. Regan, Danqing Wang, Chenhao Jin, M. Iqbal Bakti Utama, Beini Gao, Xin Wei, Sihan Zhao, Wenyu Zhao, Zuocheng Zhang, Kentaro Yumigeta, Mark Blei, Johan D. Carlström, Kenji Watanabe, Takashi Taniguchi, Sefaattin Tongay, Michael Crommie, Alex Zettl, and Feng Wang, “Mott and generalized wigner crystal states in WSe₂/WS₂ moiré superlattices,” *Nature* **579**, 359–363 (2020).
 - 42 Yuya Shimazaki, Ido Schwartz, Kenji Watanabe, Takashi Taniguchi, Martin Kroner, and Ataç Imamoğlu, “Strongly correlated electrons and hybrid excitons in a moiré heterostructure,” *Nature* **580**, 472–477 (2020).
 - 43 Yanhao Tang, Lizhong Li, Tingxin Li, Yang Xu, Song Liu, Katayun Barmak, Kenji Watanabe, Takashi Taniguchi, Allan H. MacDonald, Jie Shan, and Kin Fai Mak, “Simulation of hubbard model physics in WSe₂/WS₂ moiré superlattices,” *Nature* **579**, 353–358 (2020).
 - 44 Indrajit Maity, Prabal K. Maiti, H. R. Krishnamurthy, and Manish Jain, “Reconstruction of moiré lattices in twisted transition metal dichalcogenide bilayers,” *Phys. Rev. B* **103**, L121102 (2021).
 - 45 Heather M. Hill, Albert F. Rigosi, Kwang Taeg Rim, George W. Flynn, and Tony F. Heinz, “Band alignment in MoS₂/WS₂ transition metal dichalcogenide heterostructures probed by scanning tunneling microscopy and spectroscopy,” *Nano Letters* **16**, 4831–4837 (2016).
 - 46 Shiang Fang, Rodrick Kuate Defo, Sharmila N. Shirodkar, Simon Lieu, Georgios A. Tritsarlis, and Efthimios Kaxiras, “Ab initio tight-binding hamiltonian for transition metal dichalcogenides,” *Phys. Rev. B* **92**, 205108 (2015).
 - 47 Z. Y. Zhu, Y. C. Cheng, and U. Schwingenschlgl, “Giant spin-orbit-induced spin splitting in two-dimensional transition-metal dichalcogenide semiconductors,” *Phys. Rev. B* **84**, 153402 (2011).
 - 48 Di Xiao, Gui-Bin Liu, Wanxiang Feng, Xiaodong Xu, and

- Wang Yao, “Coupled spin and valley physics in monolayers of MoS_2 and other group-vi dichalcogenides,” *Phys. Rev. Lett.* **108**, 196802 (2012).
- ⁴⁹ D Soriano and J L Lado, “Exchange-bias controlled correlations in magnetically encapsulated twisted van der waals dichalcogenides,” *Journal of Physics D: Applied Physics* **53**, 474001 (2020).
- ⁵⁰ Zhen Zhan, Yipei Zhang, Pengfei Ly, Hongxia Zhong, Guodong Yu, Francisco Guinea, José Ángel Silva-Guillén, and Shengjun Yuan, “Tunability of multiple ultraflat bands and effect of spin-orbit coupling in twisted bilayer transition metal dichalcogenides,” *Phys. Rev. B* **102**, 241106 (2020).
- ⁵¹ Valerio Vitale, Kemal Atalar, Arash A. Mostofi, and Johannes Lischner, “Flat band properties of twisted transition metal dichalcogenide homo- and heterobilayers of MoS_2 , MoSe_2 , WS_2 and WSe_2 ,” arXiv e-prints, arXiv:2102.03259 (2021), arXiv:2102.03259 [cond-mat.mtrl-sci].
- ⁵² E. Suárez Morell, J. D. Correa, P. Vargas, M. Pacheco, and Z. Barticevic, “Flat bands in slightly twisted bilayer graphene: Tight-binding calculations,” *Phys. Rev. B* **82**, 121407 (2010).
- ⁵³ A. O. Sboychakov, A. L. Rakhmanov, A. V. Rozhkov, and Franco Nori, “Electronic spectrum of twisted bilayer graphene,” *Phys. Rev. B* **92**, 075402 (2015).
- ⁵⁴ Luis A. Gonzalez-Arraga, J. L. Lado, Francisco Guinea, and Pablo San-Jose, “Electrically controllable magnetism in twisted bilayer graphene,” *Phys. Rev. Lett.* **119**, 107201 (2017).
- ⁵⁵ C. L. Kane and E. J. Mele, “Quantum spin hall effect in graphene,” *Phys. Rev. Lett.* **95**, 226801 (2005).
- ⁵⁶ Zhenhua Qiao, Shengyuan A. Yang, Wanxiang Feng, Wang-Kong Tse, Jun Ding, Yugui Yao, Jian Wang, and Qian Niu, “Quantum anomalous hall effect in graphene from rashba and exchange effects,” *Phys. Rev. B* **82**, 161414 (2010).
- ⁵⁷ K. Kośmider, J. W. González, and J. Fernández-Rossier, “Large spin splitting in the conduction band of transition metal dichalcogenide monolayers,” *Phys. Rev. B* **88**, 245436 (2013).
- ⁵⁸ Qun-Fang Yao, Jia Cai, Wen-Yi Tong, Shi-Jing Gong, Ji-Qing Wang, Xiangang Wan, Chun-Gang Duan, and J. H. Chu, “Manipulation of the large rashba spin splitting in polar two-dimensional transition-metal dichalcogenides,” *Phys. Rev. B* **95**, 165401 (2017).
- ⁵⁹ Giovanni Pizzi, Valerio Vitale, Ryotaro Arita, Stefan Blügel, Frank Freimuth, Guillaume Géranton, Marco Gibertini, Dominik Gresch, Charles Johnson, Takashi Koretsune, Julen Ibañez-Azpiroz, Hyungjun Lee, Jae-Mo Lihm, Daniel Marchand, Antimo Marrazzo, Yuriy Mokrousov, Jamal I Mustafa, Yoshiro Nohara, Yusuke Nomura, Lorenzo Paulatto, Samuel Poncé, Thomas Ponweiser, Junfeng Qiao, Florian Thöle, Stepan S Tsirkin, Malgorzata Wierzbowska, Nicola Marzari, David Vanderbilt, Ivo Souza, Arash A Mostofi, and Jonathan R Yates, “Wannier90 as a community code: new features and applications,” *Journal of Physics: Condensed Matter* **32**, 165902 (2020).
- ⁶⁰ Di Xiao, Ming-Che Chang, and Qian Niu, “Berry phase effects on electronic properties,” *Rev. Mod. Phys.* **82**, 1959–2007 (2010).
- ⁶¹ Naoto Nagaosa and Yoshinori Tokura, “Topological properties and dynamics of magnetic skyrmions,” *Nature Nanotechnology* **8**, 899–911 (2013).
- ⁶² Su Do Yi, Shigeki Onoda, Naoto Nagaosa, and Jung Hoon Han, “Skyrmions and anomalous hall effect in a dzyaloshinskii-moriya spiral magnet,” *Phys. Rev. B* **80**, 054416 (2009).
- ⁶³ Takashi Kurumaji, Taro Nakajima, Max Hirschberger, Akiko Kikkawa, Yuichi Yamasaki, Hajime Sagayama, Hironori Nakao, Yasujiro Taguchi, Taka hisa Arima, and Yoshinori Tokura, “Skyrmion lattice with a giant topological hall effect in a frustrated triangular-lattice magnet,” *Science* **365**, 914–918 (2019).
- ⁶⁴ Y. Onose, T. Ideue, H. Katsura, Y. Shiomi, N. Nagaosa, and Y. Tokura, “Observation of the magnon hall effect,” *Science* **329**, 297–299 (2010).
- ⁶⁵ S. A. Owerre, “Topological honeycomb magnon hall effect: A calculation of thermal hall conductivity of magnetic spin excitations,” *Journal of Applied Physics* **120**, 043903 (2016).
- ⁶⁶ Tôru Moriya, “Anisotropic superexchange interaction and weak ferromagnetism,” *Phys. Rev.* **120**, 91–98 (1960).
- ⁶⁷ I. Dzyaloshinsky, “A thermodynamic theory of “weak” ferromagnetism of antiferromagnetics,” *Journal of Physics and Chemistry of Solids* **4**, 241–255 (1958).
- ⁶⁸ Jinghua Liang, Weiwei Wang, Haifeng Du, Ali Hallal, Karin Garcia, Mairbek Chshiev, Albert Fert, and Hongxin Yang, “Very large dzyaloshinskii-moriya interaction in two-dimensional janus manganese dichalcogenides and its application to realize skyrmion states,” *Phys. Rev. B* **101**, 184401 (2020).
- ⁶⁹ Michael McGuire, “Crystal and magnetic structures in layered, transition metal dihalides and trihalides,” *Crystals* **7**, 121 (2017).
- ⁷⁰ D. Soriano, M. I. Katsnelson, and J. Fernández-Rossier, “Magnetic two-dimensional chromium trihalides: A theoretical perspective,” *Nano Letters* **20**, 6225–6234 (2019).
- ⁷¹ M. Blei, J. L. Lado, Q. Song, D. Dey, O. Erten, V. Pardo, R. Comin, S. Tongay, and A. S. Botana, “Synthesis, engineering, and theory of 2d van der waals magnets,” *Applied Physics Reviews* **8**, 021301 (2021).
- ⁷² Zhe Wang, Marco Gibertini, Dumitru Dumcenco, Takashi Taniguchi, Kenji Watanabe, Enrico Giannini, and Alberto F. Morpurgo, “Determining the phase diagram of atomically thin layered antiferromagnet CrCl_3 ,” *Nature Nanotechnology* **14**, 1116–1122 (2019).
- ⁷³ M. Kim, P. Kumaravadivel, J. Birkbeck, W. Kuang, S. G. Xu, D. G. Hopkinson, J. Knolle, P. A. McClarty, A. I. Berdyugin, M. Ben Shalom, R. V. Gorbachev, S. J. Haigh, S. Liu, J. H. Edgar, K. S. Novoselov, I. V. Grigorieva, and A. K. Geim, “Micromagnetometry of two-dimensional ferromagnets,” *Nature Electronics* **2**, 457–463 (2019).
- ⁷⁴ Bevin Huang, Genevieve Clark, Efrén Navarro-Moratalla, Dahlia R. Klein, Ran Cheng, Kyle L. Seyler, Ding Zhong, Emma Schmidgall, Michael A. McGuire, David H. Cobden, Wang Yao, Di Xiao, Pablo Jarillo-Herrero, and Xiaodong Xu, “Layer-dependent ferromagnetism in a van der waals crystal down to the monolayer limit,” *Nature* **546**, 270–273 (2017).
- ⁷⁵ C. Cardoso, D. Soriano, N. A. García-Martínez, and J. Fernández-Rossier, “Van der waals spin valves,” *Phys. Rev. Lett.* **121**, 067701 (2018).
- ⁷⁶ Klaus Zollner, Paulo E. Faria Junior, and Jaroslav Fabian, “Proximity exchange effects in MoSe_2 and WSe_2 heterostructures with CrI_3 : Twist angle, layer, and gate dependence,” *Phys. Rev. B* **100**, 085128 (2019).

- ⁷⁷ Alexander M. Holmes, Samaneh Pakniyat, S. Ali Hassani Gangaraj, Francesco Monticone, Michael Weinert, and George W. Hanson, “Exchange splitting and exchange-induced nonreciprocal photonic behavior of graphene in CrI_3 -graphene van der Waals heterostructures,” *Phys. Rev. B* **102**, 075435 (2020).
- ⁷⁸ Tobias M. R. Wolf, Oded Zilberberg, Gianni Blatter, and Jose L. Lado, “Spontaneous valley spirals in magnetically encapsulated twisted bilayer graphene,” *Phys. Rev. Lett.* **126**, 056803 (2021).
- ⁷⁹ Guangze Chen and J. L. Lado, “Tunable moire spinons in magnetically encapsulated twisted van der Waals quantum spin-liquids,” arXiv e-prints, arXiv:2102.07484 (2021), arXiv:2102.07484 [cond-mat.str-el].
- ⁸⁰ E. Colomés and M. Franz, “Antichiral edge states in a modified haldane nanoribbon,” *Phys. Rev. Lett.* **120**, 086603 (2018).
- ⁸¹ Aline Ramires and Jose L. Lado, “Electrically tunable gauge fields in tiny-angle twisted bilayer graphene,” *Phys. Rev. Lett.* **121**, 146801 (2018).
- ⁸² Aline Ramires and Jose L. Lado, “Impurity-induced triple point fermions in twisted bilayer graphene,” *Phys. Rev. B* **99**, 245118 (2019).
- ⁸³ Yves H. Kwan, Glenn Wagner, Tomohiro Soejima, Michael P. Zaletel, Steven H. Simon, Siddharth A. Parameswaran, and Nick Bultinck, “Kekulé spiral order at all nonzero integer fillings in twisted bilayer graphene,” arXiv e-prints, arXiv:2105.05857 (2021), arXiv:2105.05857 [cond-mat.str-el].
- ⁸⁴ I. Y. Phinney, D. A. Bandurin, C. Collignon, I. A. Dmitriev, T. Taniguchi, K. Watanabe, and P. Jarillo-Herrero, “Strong interminivalley scattering in twisted bilayer graphene revealed by high-temperature magnetooscillations,” (2021), arXiv:2106.02088 [cond-mat.mes-hall].
- ⁸⁵ Paolo Giannozzi, Stefano Baroni, Nicola Bonini, Matteo Calandra, Roberto Car, Carlo Cavazzoni, Davide Ceresoli, Guido L Chiarotti, Matteo Cococcioni, Ismaila Dabo, Andrea Dal Corso, Stefano de Gironcoli, Stefano Fabris, Guido Fratesi, Ralph Gebauer, Uwe Gerstmann, Christos Gougoussis, Anton Kokalj, Michele Lazzeri, Layla Martin-Samos, Nicola Marzari, Francesco Mauri, Riccardo Mazzarello, Stefano Paolini, Alfredo Pasquarello, Lorenzo Paulatto, Carlo Sbraccia, Sandro Scandolo, Gabriele Sclauszero, Ari P Seitsonen, Alexander Smogunov, Paolo Umari, and Renata M Wentzcovitch, “QUANTUM ESPRESSO: a modular and open-source software project for quantum simulations of materials,” *Journal of Physics: Condensed Matter* **21**, 395502 (2009).
- ⁸⁶ J. P. Perdew and Alex Zunger, “Self-interaction correction to density-functional approximations for many-electron systems,” *Phys. Rev. B* **23**, 5048–5079 (1981).
- ⁸⁷ Andrea Dal Corso, “Pseudopotentials periodic table: From h to pu,” *Computational Materials Science* **95**, 337–350 (2014).
- ⁸⁸ Zhen Zhan, Yipei Zhang, Guodong Yu, Francisco Guinea Jose Angel Silva-Guillen, and Shengjun Yuan, “Multi-ultraflatbands tunability and effect of spin-orbit coupling in twisted bilayer transition metal dichalcogenides,” arXiv e-prints, arXiv:2005.13868 (2020), arXiv:2005.13868 [cond-mat.mes-hall].
- ⁸⁹ Yipei Zhang, Zhen Zhan, Francisco Guinea, Jose Ángel Silva-Guillén, and Shengjun Yuan, “Tuning band gaps in twisted bilayer MoS_2 ,” *Phys. Rev. B* **102**, 235418 (2020).
- ⁹⁰ Armando Pezo, Zeila Zanolli, Nils Wittemeier, Adalberto Fazzio, Stephan Roche, and Jose H. Garcia, “A Twist for Tuning the Spin-orbit Coupling in Graphene/Transition Metal Dichalcogenide Heterobilayers,” arXiv e-prints, arXiv:2011.06714 (2020), arXiv:2011.06714 [cond-mat.mes-hall].
- ⁹¹ Zheyu Lu, Stephen Carr, Daniel T. Larson, and Efthimios Kaxiras, “Lithium intercalation in moS_2 bilayers and implications for moiré flat bands,” *Phys. Rev. B* **102**, 125424 (2020).