

Calibration of mid- to far-infrared spectral lines in galaxies

Sabrina Mordini^{1,2,*}, Luigi Spinoglio^{2,**}, and Juan Antonio Fernández-Ontiveros^{2,***}

¹ Dipartimento di Fisica, Università di Roma La Sapienza, P.le A. Moro 2, I-00185 Roma, Italy

² Istituto di Astrofisica e Planetologia Spaziali (INAF-IAPS), Via Fosso del Cavaliere 100, I-00133, Roma, Italy

March 1, 2022

ABSTRACT

Context. Mid- to far-infrared (IR) lines are suited to study dust obscured regions in galaxies, because dust extinction is strongly decreasing with wavelength and therefore IR spectroscopy allows us to explore the most hidden regions of galaxies, where heavily obscured star formation as well as accretion onto supermassive black-holes at the nuclei of galaxies occur. This is mostly important for the so-called “Cosmic Noon”, i.e. at redshifts of $1 < z < 3$, when most of the baryonic mass in galaxies has been assembled.

Aims. Our goal is to provide reliable calibrations of the mid- to far-IR ionic fine structure lines, the brightest H_2 pure rotational lines and the Polycyclic Aromatic Hydrocarbons (PAHs) features, that will be used to analyse current and future observations in the mm/submm range from the ground, as well as mid-IR spectroscopy from the upcoming *James Webb Space Telescope*.

Methods. We use three samples of galaxies observed in the local Universe: star forming galaxies (196), active galactic nuclei (AGN; 90–150 for various observables) and low-metallicity dwarf galaxies (40). For each population we derive different calibrations of the observed line luminosities versus the total IR luminosities.

Results. Through the resulting calibrations, we derive spectroscopic measurements of the Star Formation Rate (SFR) and of the Black Hole Accretion Rate (BHAR) in galaxies using mid- and far-IR fine structure lines, H_2 pure rotational lines and PAH features. In particular, we derive robust star-formation tracers based on: the [CII]158 μm line; the sum of the two far-IR oxygen lines, the [OI]63 μm line and the [OIII]88 μm line; a combination of the neon and sulfur mid-IR lines; the bright PAH features at 6.2 and 11.3 μm , as well as – for the first time – the H_2 rotational lines at 9.7, 12.3 and 17 μm . We propose the [CII]158 μm line, the combination of the two neon lines ([NeII]12.8 μm and [NeIII]15.5 μm) and, for solar-like metallicity galaxies that may harbour an AGN, the PAH11.3 μm feature as the best SFR tracers. On the other hand, a reliable measure of the BHAR can be obtained using the [OIV]25.9 μm and the [NeV]14.3 and 24.3 μm lines. For the most commonly observed fine-structure lines in the far-IR we compare our calibration with the existing ALMA observations of high redshift galaxies. We find an overall good agreement for the [CII]158 μm line for both AGN and star forming galaxies, while the [OIII]88 μm line in high- z galaxies is in better agreement with the low metallicity local galaxies (dwarf galaxies sample) than with the star forming galaxies, suggesting that high- z galaxies might have strong radiation fields due to low metal abundances, as expected.

Key words. galaxies: active – galaxies: evolution – galaxies: star formation – infrared: galaxies – techniques: spectroscopic telescopes

1. Introduction

One of the major and still unsolved problems in astrophysics is related to the lack of a clear understanding of how galaxies evolve, from the time of structure formation till today, and what are the dominant processes that have influenced their evolution (Somerville & Davé 2015; Naab & Ostriker 2017; Bullock & Boylan-Kolchin 2017; Wechsler & Tinker 2018; Förster Schreiber & Wuyts 2020). From the observational scenario that has been consolidated in the last tens of years, we know that the two main energy production mechanisms along galaxy evolution are, on one side, star formation and subsequent stellar evolution and, on the other side, accretion onto the super-massive black holes (SMBHs) that form at the centre of galaxies.

The bulk of star formation and BH accretion took place at the so-called *Cosmic Noon* ($1 < z < 3$), with a steep decline toward the present epoch, in heavily obscured environments, embedded in large amounts of gas and dust, where optical and UV detected radiation corresponds to only $\sim 10\%$ of the total emitted light (Madau & Dickinson 2014) and with most of the radiated energy

being absorbed by dust and re-emitted at longer wavelengths. Both star formation and black hole accretion contribute to the dust re-emission, and therefore measuring the integrated IR continuum, through photometric observations, does not discriminate easily between both components. The spatial resolution required to isolate the bulk of the nuclear IR emission from the host galaxy ($\lesssim 100$ pc) cannot be attained in high- z galaxies, while spectral decomposition techniques (e.g., Berta et al. 2013) are ultimately dependent on templates of local active galactic nuclei (AGN), whose inner workings and dust distribution are still far from being understood (e.g. Lyu & Rieke 2020). Thus measuring star formation rate (SFR) and black hole accretion rate (BHAR) in galaxies across cosmic time, which is one of the major observational goals in galaxy evolution studies, has to be done through spectroscopic observations at wavelengths long enough to overcome dust absorption.

The mid to far-IR range is populated by a large number of atomic and molecular lines and features. In particular, the atomic and ionic fine-structure lines cover a wide range of physical parameters in terms of excitation, density and ionisation, as can be seen in fig. 4 of Spinoglio et al. (2012), showing the critical density for collisional de-excitation versus the ionisation potential of the IR fine-structure lines. These lines can easily discriminate

* E-mail: sabrina.mordini@uniroma1.it

** E-mail: luigi.spinoglio@inaf.it

*** E-mail: j.a.fernandez.ontiveros@gmail.com

among the different gas excitation conditions, from the Narrow Line Regions (NLR) excited by the active nucleus, to the HII and photo-dissociation regions (PDRs), whose origin is due to stellar excitation (see, e.g., Spinoglio & Malkan 1992; Tommasin et al. 2010; Spinoglio et al. 2015; Fernández-Ontiveros et al. 2016), offering an ideal tool to probe the highly opaque and dust-obscured regions. Most of these lines can be assessed only from space IR telescopes, while only a few transitions happen to lie in the atmospheric windows, e.g. in the spectral region of 8–13 μm .

The intermediate ionisation fine-structure lines are good tracers of the star formation and in particular the sum of the fluxes of [NeII] and [NeIII] has been shown to give a measure of the Star Formation Rate (Ho & Keto 2007; Zhuang et al. 2019) in galaxies. The low ionisation [CII] line at 158 μm is one of the brightest emission lines in star forming galaxies, and can thus trace star formation activity both in the local (De Looze et al. 2014; Herrera-Camus et al. 2015) and high redshift (Leung et al. 2020) Universe.

The mid-IR spectra of star forming galaxies between 3 and 19 μm are often dominated by emission features attributed to Polycyclic Aromatic Hydrocarbons (PAHs), due to infrared (IR) fluorescence by these large molecules containing 50–100 C-atoms, pumped by single FUV photons (Allamandola et al. 1989; Smith et al. 2007; Tielens 2008). Because they trace the FUV stellar flux, they can be used to measure star formation. These features are strong when compared to fine structure lines in star forming galaxies, and are detected not only in local galaxies, but also at redshifts up to $z \sim 4$ (Kirkpatrick et al. 2015; Riechers et al. 2014; Sajina et al. 2012). Their emission accounts for ~ 10 –20% of the total IR radiation by dust, and originate from PDRs near star forming regions. PAHs can be used to trace SFR not only in star forming galaxies (Shipley et al. 2016; Xie & Ho 2019), but also in sources where an AGN contribution is present but not the dominant source of integrated light (Shipley et al. 2013). In these AGN, the equivalent width of the PAH features can be used to estimate the star formation contribution to the total IR luminosity (Armus et al. 2007; Tommasin et al. 2010). However, in more extreme AGN, strong UV and X-ray radiation fields can also suppress the PAH emission in the vicinity of the AGN by photo-dissociation, while increasing the mid-IR continuum emission (Lacy et al. 2013). Analogously, also in Low Metallicity Galaxies (LMG, i.e. below $12 + \log(\text{O}/\text{H}) \sim 8.2$), because of the reduced formation efficiency and the increased stellar radiation hardness, the strength of the PAH features is reduced and only detected at metallicities above $1/8$ – $1/10 Z_{\odot}$ (Engelbracht et al. 2008; Cormier et al. 2015; Galliano et al. 2021).

Among the molecular lines in the mid-IR, the pure rotational transitions of H_2 are particularly important as the typical physical conditions of the gas associated with these lines can be found both in AGN and star forming galaxies. Rigopoulou et al. (2002) found that, while in star forming environments the H_2 emission can originate from PDRs, with a small contribution from shocks, in AGN dominated galaxies the X-ray emission from the central AGN plays an important role in heating large amounts of gas boosting the H_2 emission.

High ionisation lines trace AGN activity: all lines whose ionisation potential is higher than the one necessary to doubly ionise helium ($> 54.4 \text{ eV}$) cannot be efficiently produced by stellar radiation in significant amounts. Therefore, the detection of the [NeV]14.3 μm , [NeV]24.3 μm or [NeVI]7.65 μm lines probes the presence of an AGN, while the [OIV]25.9 μm line, which can also be found in energetic starbursts and LMG, is however much stronger in AGN, having an equivalent width one order of magnitude larger compared to Star Forming Galaxies (SFG)

(Tommasin et al. 2010). Typical IR line ratios used to measure the strength of the active nucleus with respect to the star formation component in a galaxy are: the [NeV]14.3 μm or 24.3 μm to [NeII]12.8 μm ratio, the [OIV]25.9 μm /[NeII]12.8 μm line ratio and the [OIV]25.9 μm /[OIII]52 μm or 88.35 μm ratio (Sturm et al. 2002; Armus et al. 2007; Tommasin et al. 2010; Spinoglio et al. 2015; Fernández-Ontiveros et al. 2016).

The main goals of this work are: (i) to revise the calibration of the mid- to far-IR lines including ionic fine-structure lines, the brightest H_2 pure rotational lines and the PAH features and (ii) to provide a local calibration of spectroscopic SFR and BHAR tracers, that can also be applied to measurements at high- z . We will follow the study presented in Spinoglio et al. (2012), update the IR spectroscopic observations presented there and include a sample of LMG, to extend the calibration to these objects. These latter are included to characterise the response of the lines to the conditions of low metallicity (average $1/5 Z_{\odot}$) (Madden et al. 2013). Low-metallicity AGN are not included because they are rare in the local Universe and only a few examples are found (e.g. Circinus, Oliva et al. 1999), but the broad- and narrow-line regions in AGN show essentially little or no chemical evolution up to $z \sim 7$ (Nagao et al. 2006; Juarez et al. 2009; Onoue et al. 2020), suggesting that the quasar phase appears mostly when galaxies are already chemically mature objects. Therefore including the metallicity dependence in the calibrations for AGN is not as relevant as in the case of star forming galaxies, where the chemical evolution with redshift is well known (Sanders et al. 2020). This work extends and update a previous study (Spinoglio et al. 2021) aimed at the preparation of the SPICA mission (Roelfsema et al. 2018) where the authors give a calibration of the most important features that were used to plan spectroscopic observations with that mission.

A further motivation of this study resides in the need to exploit the great potential of extragalactic IR spectroscopy for high redshift galaxies, which is being already explored by the Atacama Large Millimeter/submillimeter Array (ALMA, Wootten & Thompson 2009; Carpenter et al. 2020), and will have a dramatic boost with the next space IR telescopes, such as the *James Webb Space Telescope* (JWST, Gardner et al. 2006) and in a more far future, possibly by the *Origins Space Telescope*¹.

The paper is organized as follows: section 2 describes the samples of galaxies, observed in the local Universe, that we have used to derive the correlations; section 3 reports our results, in particular: section 3.1 presents the new correlations between the line luminosities and the total IR luminosities, while sections 3.2 and 3.3 give simple recipes to measure the two main parameters of the SFR and BHAR. In section 4 we discuss our results: section 4.3 compares our study to previous ones, in section 4.4 we discuss the metallicity effect on the SFR tracers, section 4.5 presents how the observations at high redshift compare with the correlations we have derived and section 4.6 shows how our results can be used to interpret present and future IR/(sub)-mm observations. Our conclusions are presented in section 5.

2. The selected lines and features and the samples of galaxies

We use the most representative samples of star forming galaxies (SFG), AGN, and low metallicity galaxies (LMG) in the local Universe for which IR spectroscopy is available, mainly from *Spitzer*-IRS (Houck et al. 2004) and *Herschel*-PACS (Poglitsch et al. 2010), but also from the *Infrared Space Observatory* (ISO)

¹ <https://asd.gsfc.nasa.gov/firs>

SWS and LWS spectrometers (de Graauw et al. 1996; Clegg et al. 1996). For each of the three galaxy populations, we derive linear relations in logarithmic space between the line luminosity and the total IR luminosity. Then we derive the best tracers of the SFR and the BHAR, using the discussed IR lines and features, and compare them with what has been reported in the literature.

In this analysis we considered, in order of decreasing ionisation/excitation:

- Four high-ionisation fine structure lines, typical of AGN: [NeVI]7.65 μm , [NeV]14.32 μm , [NeV]24.32 μm and [OIV]25.89 μm ;
- Ten intermediate ionisation fine structure lines, typical of stellar/HII regions: [SIV]10.51 μm , [NeII]12.81 μm , [NeIII]15.55 μm , [SIII]18.71 μm , [SIII]33.48 μm , [OIII]51.81 μm , [NIII]57.32 μm , [OIII]88.36 μm , [NII]121.9 μm and [NII]205 μm ;
- Five low-ionisation/neutral fine structure lines, typical of PDR: [FeII]25.99 μm , [SiII]34.81 μm , [OI]63.18 μm , [OI]145.5 μm and [CII]157.7 μm ;
- Four H₂ pure rotational lines at 9.67, 12.28, 17.03 and 28.22 μm ;
- Five PAH features at 6.2, 7.7, 8.6, 11.3 and 17 μm .

The fundamental parameters for the fine-structure lines considered in this analysis are reported in Table A.1.

In this analysis we have included those lines observed with the *Spitzer*-IRS high resolution (HR) channel in the 10–35 μm range, for which good spectra are available in the literature. The only exception is the line of [NeVI]7.65 μm , an exclusive AGN line, for which we included ISO-SWS observations. Other well known fine-structure lines, such as [ArII]6.98 μm , [ArIII]8.99 μm and [NeIII]36.0 μm , which could indeed have a relevant role for future observations, especially in view of the *JWST* launch, at the present time do not have enough high quality spectra in the literature to be included in the analysis. We do not consider upper limits to derive our correlations. This is because, in general, our statistics are quantitatively appropriate, thus making the inclusion of upper limits not necessary. Moreover, where the statistics are less precise, upper limits are usually not available for the considered lines.

The definition of the various samples of galaxies chosen to compute the correlations are described in the following sections and summarised in Table 1, with the instruments used to observe the spectral lines and features, the total number of objects selected and the references for each sample.

2.1. The AGN sample

The AGN sample has been drawn from the 12 μm selected active galaxies sample (12MGS, Rush et al. 1993), which is the brightest complete and unbiased sample of Seyfert galaxies in the local Universe. For the mid-IR fine structure lines and the H₂ rotational lines, we have used the sub-sample of the 12MGS observed by *Spitzer* IRS at high spectral resolution ($R = 600$) which contains 88 AGN (Tommasin et al. 2008, 2010). For the PAH features at 6.2 μm and 11.2 μm we have used the *Spitzer*-IRS data at low spectral resolution ($R \sim 60$ –120; Buchanan et al. 2006; Wu et al. 2009) of the 12MGS (103 objects), because this setting matches better the intrinsic width of these features. We note here that Wu et al. measure the PAH features using a spline function to determine the continuum level. In order to make these measurements comparable to those obtained using automated fitting procedures (e.g. PAHFIT, Smith et al. 2007), we apply a correction factor of 1.7 and 1.9 to increase the fluxes of the

6.2 μm and 11.3 μm PAH in Wu et al., respectively, following the differences found by Smith et al. (2007).

The adoption of the complete 12MGS as the main catalog, from which we cover about 75% of the total sample, allows us to derive statistically robust calibrations.

For the [NeVI]7.65 μm line, we could not use the 12MGS, because this line was not detected by *Spitzer* at low resolution and was outside its spectral range at high-resolution. Therefore we had to use the data from Sturm et al. (2002), which contain 8 detections of the [NeVI]7.65 μm line with *ISO*-SWS at medium resolution ($R \sim 1500$), and thus not contaminated by the PAH emission at 7.7 μm .

For the far-IR spectral range, namely the 50–205 μm , we used the catalogue of AGN assembled by Fernández-Ontiveros et al. (2016), which includes all the Seyfert galaxies and the quasars of the Véron-Cetty & Véron (2010) catalogue which have far-IR spectra observed by *Herschel*-PACS. The sample of the AGN observed in the far-IR lines counts 170 galaxies and contains about 50% of objects from the 12MGS, while the others do not come from a complete sample.

2.2. The SFG sample

The SFG sample was constructed using the *The Great Observatories All-Sky LIRG Survey* (GOALS sample, Armus et al. 2009), from which we extracted 158 galaxies, with data from Inami et al. (2013), who report the fine structure lines at high resolution in the 10–36 μm interval, and Stierwalt et al. (2014), who include the detections of the H₂ molecular lines and the PAH features at low spectral resolution. For those galaxies in the GOALS sample that have a single IRAS counterpart, but more than one source detected in the emission lines, we have added together the line/feature fluxes of all components, to consistently associate the correct line/feature emission to the total IR luminosity computed from the IRAS fluxes. To cover also lower luminosity galaxies, as the GOALS sample only includes Luminous IR Galaxies (LIRGs) and Ultra-Luminous IR Galaxies (ULIRGs), we have included 38 galaxies from Bernard-Salas et al. (2009) and Goulding & Alexander (2009), to reach the total sample of 196 galaxies with IR line fluxes in the 5.5–35 μm interval in which an AGN component is not detected. For the Bernard-Salas et al. (2009), Goulding & Alexander (2009) and the GOALS samples we excluded all the composite starburst-AGN objects identified as those with a detection of [NeV] either at 14.3 or 24.3 μm . It is worth to note that the original samples from Goulding & Alexander (2009) and Bernard-Salas et al. (2009) have spectra covering solely the central region of the galaxies. To estimate the global SFR, we corrected the published line fluxes of the *Spitzer* spectra by multiplying them by the ratio of the continuum reported in the IRAS Point Source Catalogue to the continuum measured on the *Spitzer* spectra extracted from the CASSIS database (Lebouteiller et al. 2015). We assume here that the line emission scales, at first order, with the IR brightness distribution. In particular, we considered the continuum at 12 μm for the [NeII]12.8 μm and [NeIII]15.6 μm lines, and the continuum at 25 μm for the [OIV]25.9 μm , [FeII]26 μm , [SIII]33.5 μm and [SiII]34.8 μm lines. This correction was not needed for the AGN sample and the GOALS sample because of the greater average redshift of the galaxies in the 12MGS and GOALS samples. In particular, the 12MGS active galaxies sample has a mean redshift of 0.028 (Rush et al. 1993), while the GOALS sample has a mean redshift of 0.026. The galaxies presented by Bernard-Salas et al. (2009) have instead an average redshift of 0.0074, while the sample by Goulding & Alexander (2009) has an average redshift

Table 1. Characteristics of the different samples used in this analysis. For each class of objects we report, for the different lines and features, the original sample (*heterogeneous* indicates that the sources do not belong to a specific well-defined sample of galaxies), the facility and instrument that observed the sample of galaxies, the number of objects used in the analysis and the references where the original data can be found.

Type	Lines/features	Sample	Observed by	N. of objects	References
AGN	[NeV]7.7 μ m	heterogeneous	ISO-SWS	8	Sturm et al. (2002)
	MIR lines in the 10-35 μ m range including H ₂ rotational lines	12MGS	<i>Spitzer</i> -IRS HR	88	Tommasin et al. (2008, 2010)
	PAH features	12MGS	<i>Spitzer</i> -IRS LR	103	Wu et al. (2009)
	FIR lines in the 50-205 μ m range	heterogeneous	<i>Herschel</i> -PACS and SPIRE	149	Fernández-Ontiveros et al. (2016)
SFG	MIR lines in the 10-35 μ m range	GOALS sample	<i>Spitzer</i> -IRS HR	153	Inami et al. (2013)
		+ heterogeneous		10	Bernard-Salas et al. (2009)
		+ heterogeneous		28	Goulding & Alexander (2009)
		GOALS sample		153	Stierwalt et al. (2014)
	H ₂ rotational lines	+ heterogeneous	<i>Spitzer</i> -IRS HR	10	Bernard-Salas et al. (2009)
		+ heterogeneous		28	Goulding & Alexander (2009)
	PAH features	GOALS sample	<i>Spitzer</i> -IRS LR	179	Stierwalt et al. (2014)
		+ heterogeneous		12	Brandl et al. (2006)
	FIR lines in the 50-205 μ m range	GOALS sample	<i>Herschel</i> -PACS	153	Díaz-Santos et al. (2017)
		+ heterogeneous	<i>Herschel</i> -PACS and SPIRE	20	Fernández-Ontiveros et al. (2016)
LMG	MIR lines in the 10-35 μ m range	DGS	<i>Spitzer</i> -IRS HR and LR	40	Cormier et al. (2015)
	FIR lines in the 50-158 μ m range	DGS	<i>Herschel</i> -PACS	40	Cormier et al. (2015)

of 0.0044. For the other lines in the 10–36 μ m interval, Goulding & Alexander (2009) do not report a detection, and we used the data presented in Bernard-Salas et al. (2009) for a total of 15 objects. Both Bernard-Salas et al. (2009) and Goulding & Alexander (2009) report data from the high-resolution *Spitzer*-IRS spectra. Data in the 50–205 μ m interval were taken from Díaz-Santos et al. (2017) for the GOALS sample, 20 starburst galaxies from Fernández-Ontiveros et al. (2016), and 23 objects from the ISO-LWS observations of Negishi et al. (2001), resulting in a total sample of 193 objects. Lastly, the PAH features fluxes were measured from the low resolution *Spitzer*-IRS spectra by Brandl et al. (2006), including 12 objects from the sample of Bernard-Salas et al. (2009) and 179 objects from Stierwalt et al. (2014).

2.3. The LMG sample

The LMG sample was selected from Cormier et al. (2015), where the 10–36 μ m interval was observed by *Spitzer*-IRS in both high and low resolution, and the 50–158 μ m interval was observed by *Herschel*-PACS. For the *Spitzer*-IRS data, we only considered the high resolution results, for a total sample of 40 objects.

3. Results

We derive, for each line or feature in each galaxy sample, the correlation between the logarithms of the total IR luminosity in the 8–1000 μ m range – computed from the IRAS fluxes following Sanders & Mirabel (1996) – and the line luminosity, according to the equation:

$$\log L_{\text{Line}} = (a \pm \delta a) \log L_{\text{IR}} + (b \pm \delta b) \quad (1)$$

with all luminosities expressed in units of 10^{41} erg s⁻¹. We report in Table D.1 of the Appendix D the best-fit parameters obtained for each line/feature using the orthogonal distance regression fit (Boggs & Rogers 1990), the number of objects N , and the Pearson

correlation coefficient r . We use the orthogonal distance regression because the two variables are independent each other, instead of the ordinary least-square minimisation, where one variable is dependent from the other one. This is also particularly useful to derive the inverse relation between the two variables from the best-fit coefficients.

Using these correlations, we derive the tracers for the SFR (Section 3.2) and the BHAR (Section 3.3). For the SFR, we used the [CII]158 μ m luminosity, various combinations of the luminosities of the [NeII]12.8 μ m, [NeIII]15.6 μ m, [SIII]18.7 μ m and [SIV]10.5 μ m lines, the luminosity of the PAH features at 6.2 μ m and 11.3 μ m and the luminosity of the H₂ rotational lines at 9.7 μ m, 12.3 μ m and 17.3 μ m. For the BHAR, we have used the luminosities of the [OIV]25.9 μ m, [NeV]14.3 μ m and 24.3 μ m lines.

3.1. Spectral lines and features vs. total IR luminosity

Among the different correlations derived in this work and presented in Table D.1 of the Appendix D, we highlight in this section the main results, while we refer the reader to Appendix B for the whole set of figures illustrating the correlations with the total IR luminosity.

In Fig. 1a, we present the correlation between the [CII]158 μ m line luminosity and the total IR luminosity for the three galaxy populations considered. We find that the SFG and the LMG samples follow a tight relation over six orders of magnitude in L_{IR} with a consistent regression slope and offset values within the uncertainties, from low-luminosity dwarf galaxies to LIRGs and ULIRGs, confirming the results of De Looze et al. (2014). This will be further discussed in Section 3.2.2. On the other hand, AGN have higher IR luminosities for a given [CII]158 μ m line luminosity with respect to the SFG and LMG samples, likely due to the contribution from the AGN continuum emission to the total IR luminosity. This effect is also seen in low ionisation transitions where the AGN contribution to the line emission is expected to be small, such as the [NII]122 μ m line (see Fig. 1c). However, the




Fig. 1. (a: left) The [CII]158 μm line luminosity as a function of the total IR luminosity. Blue squares represent detections in AGN, red stars indicate SFG and green triangles LMG. The solid red line represents the linear relation calculated for SFG, the blue dotted line shows the relation for AGN and the green dashed line the one for LMG. **(b: centre)** The [NeII]12.8 μm line luminosity as a function of the total IR luminosity. **(c: right)** The [NeIII]15.6 μm line luminosity as a function of the total IR luminosity. In **(b)** and **(c)** we use the same notations as in **(a)**.




Fig. 2. (a: left) The [OIII]88 μm line luminosity as a function of the total IR luminosity. **(b: centre)** The [OIV]25.9 μm line luminosity as a function of the total IR luminosity. **(c: right)** The luminosity of the PAH feature at 11.3 μm as a function of the total IR luminosity. The same legend as in Fig. 1 was used.

same effect is not observed for [OI]63 μm (see Fig. B.7b). This might be related to the higher critical density of the [OI]63 μm line, which becomes an efficient coolant in X-ray dissociation regions in the presence of an AGN (Maloney et al. 1996; Dale et al. 2004). High-excitation lines where the AGN completely dominates the line emission show the opposite behaviour, that is, the sources are shifted towards higher line intensities at a given IR luminosity (e.g. [OIV]25.9 μm in Fig. 2b). This suggests that the lack of a noticeable shift between AGN and SFGs in intermediate excitation lines such as [NeII]12.8 μm or [SIII]18.7,33.5 μm (Figs. 1b, B.4b and B.6a) might be caused by a comparable AGN contribution to both the IR luminosity and the line intensity.

We note that various authors (e.g. Herrera-Camus et al. 2015; Croxall et al. 2017) have reported an observed deficit in [CII] luminosity with the increase of total L_{IR} , in particular in ULIRGs. Different mechanisms have been proposed to explain the lower [CII] emission (Sutter et al. 2019, 2021, and references therein). In particular, Sutter et al. (2019) find that the [CII] deficit is particularly evident when the emission arises from ionised gas, while the effect is negligible when the emission comes predominantly from PDR regions.

For the [NeII]12.8 μm line (Fig. 1b) there is a slight difference between SFGs and the other samples, but the differences are not statistically relevant, being consistent within 3σ of each other.

In Fig. 1c, we report the correlation obtained for the [NeIII]15.6 μm line with the total IR luminosity, showing that the AGN and the SFG have a comparable correlation, while LMG have the [NeIII]15.6 μm line more than one order of magnitude brighter, at a given IR luminosity (see also Cormier et al. 2012).

In Fig. 2a, b and c we report the correlations obtained for the [OIII]88 μm , the [OIV]25.9 μm line luminosities and the luminosity of the PAH feature at 11.3 μm with the total IR luminosity, respectively. For the [OIII]88 μm line, LMG are on average two orders of magnitude brighter, at a given IR luminosity (see also Cormier et al. 2012), compared to the other two classes of galaxies, that are almost overlapping. This could be due to the differences in the ionising spectra and conditions in the ISM of LMG with respect to the SFG and the AGN. In LMG, an increased number of photons can ionise gas at greater distances from the star forming regions, thus facilitating cooling through ionised gas emission. Additionally, the lower dust-to-gas ratios in LMG is also expected to cause a decrease of the IR luminosities, at a fixed stellar mass, in these galaxies.

AGN have about one order of magnitude brighter [OIV]25.9 μm emission compared to SFG (see also Tommasin et al. 2010), LMG have a shallower relation with IR luminosity, with a decreasing slope at higher luminosities (Fig. 2b). The flatter slope could be related to the high ionisation potential of the

[OIV]25.9 μ m line, that is, the few ionising photons beyond 54.4 eV produced by LMGs might not scale linearly with the overall luminosity. However, no firm conclusion can be drawn because of the relatively poor statistics of LMG, for which only 16 detections of the [OIV] line are available. We note here that in order to obtain a reliable correlation for the AGN sample, we considered only objects with an AGN component at the 19 μ m continuum, as defined in Tommasin et al. (2010), greater than 85%. We apply this limit only in this case, to minimise a possible contamination in this line from emission due to strong starburst activity (see, e.g. Lutz et al. 1998). This approach is motivated by the use of the [OIV]25.9 μ m line as a BHAR tracer in Sect. 3.3, where the same reduced sample is adopted.

Fig. 2c shows that the 11.3 μ m PAH feature is present both in AGN and SFG and correlates well with the total IR luminosity, while in this analysis we do not consider the PAH detections in LMG, because detections are available only in a few cases, and thus are not enough to obtain a statistically significant result. As shown by different authors (Madden 2000; Engelbracht et al. 2005; Wu et al. 2006; Smith et al. 2007; Calzetti et al. 2007), in LMG there is evidence for a deficit of PAH emission and the available measurements present significantly weaker features than SFG. Moreover, the higher ionising continuum present in LMG contribute to destroy these features (Engelbracht et al. 2008; Cormier et al. 2015).

As will be shown in Section 3.2.5, while there is a difference of ~ 0.3 dex between the emission of SFG and AGN, the slopes of the two correlations are comparable within the errors, with the difference linked to a higher L_{IR} for equal PAH emission in AGN.

We note here that, while we report in Tab. D.1 the correlation derived for the [NeVI]7.6 μ m line, this correlation was obtained for a sample of only 8 AGN, while for the majority of AGN this line is not detected. We therefore conclude that this calibration has to be taken with caution, because it may be biased toward AGN with a high ionisation parameter ($U \sim -1$), while for a lower ionisation we expect the [NeVI] line to be considerably less prominent, as discussed in Satyapal et al. (2021, in particular in their Fig. 5).

3.2. Star formation rate tracers

3.2.1. Determination of SFR

In this section, we propose different SFR tracers. In order to calibrate the proposed SFR tracers, we use two different methods. For SFG, we measure the SFR directly from the total IR luminosity L_{IR} , following Kennicutt (1998):

$$\text{SFR} = k_{\text{IR}} * L_{\text{IR}} \quad (2)$$

where $k_{\text{IR}} = 4.5 \times 10^{-44} \text{M}_{\odot} \text{yr}^{-1} \text{erg}^{-1} \text{s}$. We apply the same conversion also to AGN sources (see Sect. 3.2.5), but limiting our sample to the sources with $L_{\text{IR}} \leq 10^{45} \text{erg s}^{-1}$. We apply this limit to minimise the AGN effect on the L_{IR} and avoid overestimating the SFR because of AGN activity.

For the LMG, we adopt the SFR derived from H_{α} and corrected by the total IR luminosity, as reported in Rémy-Ruyer et al. (2015). The total IR luminosity alone does not accurately represent the total SFR in LMG due to the lower dust-to-gas ratio in their ISM. The inclusion of the optical component for computing the SFR is necessary to properly account for the emission not reprocessed by dust.

3.2.2. $L_{[\text{CII}]158\mu\text{m}}$ –SFR relation

The [CII]158 μ m line can be used as a tracer of the SFR, as proposed by different authors (see, e.g., De Looze et al. 2014). We compare the line intensity of SFG and LMG to the SFR. Applying the orthogonal distance regression fit, we find a strong correlation between the two quantities:

$$\log \left(\frac{\text{SFR}}{\text{M}_{\odot} \text{yr}^{-1}} \right) = (0.62 \pm 0.02) + (0.89 \pm 0.02) \left(\log \frac{L_{[\text{CII}]}}{10^{41} \text{erg s}^{-1}} \right) \quad (3)$$

In Fig. 3a we show the correlation obtained for 227 galaxies, of which 37 are LMG and 190 are SFG. The relation is nearly linear, with a Pearson r -value of $r = 0.92$, and covers six orders of magnitude in SFR, indicating that the [CII] line emission is an overall good tracer of SFR for local star forming galaxies independently of their metallicity. We excluded the AGN from this correlation, since we derive the SFR from the L_{IR} and AGN have an excess IR continuum emission not due to star formation.

In Fig. 3b, we compare our results with those obtained in De Looze et al. (2014) and Sargsyan et al. (2012). We note that, while in this work and in the work by Sargsyan et al. the SFR was determined following Kennicutt (1998), De Looze et al. trace the SFR using the GALEX FUV emission (Cortese et al. 2012) and the *Spitzer*-MIPS 24 μ m emission (Rieke et al. 2004). When compared to the results obtained by De Looze et al. (2014) for their total sample, we find good agreement at luminosities of $L_{[\text{CII}]} > 10^{41} \text{erg s}^{-1}$, while there is a difference of ~ 0.3 dex for lower luminosities. In De Looze et al. the dwarf sample, when considered alone, shows a flatter slope than the total sample, equal to 0.80 ± 0.5 (see Fig. 3b, the blue diamond line). The authors link the flatter slope to an underestimation of the SFR based on the far-UV emission. We observe a similar flattening of the slope when considering the LMG sample alone, equal to 0.69 ± 0.05 , consistent with the result by De Looze et al. within 3σ of each other.

De Looze et al. (2014) find that the ULIRG population presents a scatter of almost one order of magnitude in the [CII]-SFR relation, when compared to the total sample. We find the same result in our correlation, with the ULIRG population (composed of 16 objects) lying between 0.2 and 1.0 dex above the correlation derived for the total sample. Given the small number of sources, we do not derive a specific [CII]-SFR correlation for the ULIRG sample.

3.2.3. Oxygen based SFR tracer

Besides C^+ , also O and O^{2+} are two important coolants of the ISM. The [OI]63 μ m line traces the warm and/or dense PDRs, while the [OIII]88 μ m emission line originates from diffuse, highly ionised regions near young, hot stars.

The [OIII] line can be an important SFR tracer in LMG (De Looze et al. 2014), where PDRs are weak or absent in the ISM and the ionisation field is stronger. In order to trace the SFR in SFG, however, a tracer that can account for PDRs needs to be included. For this reason, combining these two lines can provide an accurate estimate of the SFR, both in SFG and LMG, probing both neutral and ionised medium.

Fig. 4 shows the correlation obtained for a sample of 151 objects, 24 LMG and 124 SFG, of which 22 ULIRGs. The correlation, with a Pearson correlation coefficient of $r = 0.94$, can




Fig. 3. (a: left) Correlation between the [CII]158 μm line luminosity and the SFR derived from the total L_{IR} . Red stars represent SFG, green triangles the LMG. Purple squares show the ULIRG population of SFG, and the shaded green area indicates the 95% confidence interval. (b: centre) Comparison of the $\log L_{\text{[CII]}} - \log(\text{SFR})$ relation obtained in this work (black solid line) with the results obtained by De Looze et al. (2014): the green dashed line represents a sample of HII/star forming galaxies, the blue diamond line shows the results for the low metallicity dwarf sample, and the pink dash-dotted line considers the whole sample. The red dotted line shows the results obtained by Sargsyan et al. (2012). (c: right) Correlation between the [NeII]12.8 μm and [NeIII]15.6 μm summed emission lines luminosity, in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black dashed line) for SFG (red star) and from the H_{α} luminosity (corrected for the IR luminosity) for LMG (green triangles). The shaded green area indicates the 95% confidence interval. The purple squares highlight the ULIRG population in the SFG sample. The blue dashed line shows the results obtained by Zhuang et al. (2019) for the same relation.




Fig. 4. Correlation between the [OI]63 μm and [OIII]88 μm summed emission line luminosities, in units of $10^{41} \text{ erg s}^{-1}$, and the SFR derived from the total IR luminosity (black dashed line) for a composite sample of SFG (red stars) and from the H_{α} luminosity (corrected for the IR luminosity) for LMG (green triangles). Purple squares indicate the sample of ULIRGs included in the SFG sample.

be expressed by:

$$\log\left(\frac{\text{SFR}}{M_{\odot} \text{ yr}^{-1}}\right) = (0.30 \pm 0.04) + (1.25 \pm 0.04) \log\left(\frac{L_{\text{[OI]}} + L_{\text{[OIII]}}}{10^{41} \text{ erg s}^{-1}}\right) \quad (4)$$

3.2.4. Neon- and sulfur-based SFR tracers

A promising SFR tracer, proposed by Ho & Keto (2007); Zhuang et al. (2019), is the sum of the [NeII]12.8 μm and the

[NeIII]15.6 μm emission line fluxes. By adding the two lines, this tracer is fairly independent from the effects related to the hardness of the radiation field, which are stronger at lower metallicities affecting single line diagnostics. For instance, while the [NeII]12.8 μm line intensity scales consistently in LMG and SFG, the [NeIII]15.6 μm line becomes remarkably brighter in LMG (Fig. 1). Although some dependency on density is still expected, the high critical density of the sulphur ($> 7 \times 10^3 \text{ cm}^{-3}$; see Table A.1) and the neon lines ($> 5 \times 10^4 \text{ cm}^{-3}$) used in this Section guarantees a minor effect on these tracers for the vast majority of the galaxy population.

In Fig. 3c we show the correlation found between the summed luminosity of the [NeII] and [NeIII] emission lines and the SFR. This relation, obtained from data of 203 local star forming galaxies, of which 182 SFG and 21 LMG, can be expressed as:

$$\log\left(\frac{\text{SFR}}{M_{\odot} \text{ yr}^{-1}}\right) = (0.88 \pm 0.03) + (0.96 \pm 0.03) \log\left(\frac{L_{\text{[NeII]}} + L_{\text{[NeIII]}}}{10^{41} \text{ erg s}^{-1}}\right) \quad (5)$$

where $\log(L_{\text{[NeII]}} + L_{\text{[NeIII]}})$ is the luminosity corresponding to the sum of the fluxes of the [NeII]12.8 μm and the [NeIII]15.6 μm lines. This relation has a Pearson coefficient of $r=0.91$. We report in Fig. 3c the comparison between the theoretical relation obtained by Zhuang et al. (2019), and our empirical one. The relation by Zhuang et al. is derived following the assumption that the neon lines trace all the ionising photons in a star forming region. Our relation, derived from observational data, shows a lower slope, and thus a larger offset, thus suggesting a lower efficiency in reprocessing the ionising photons by the neon transitions.

Analogously to the case of the two neon lines, we also take into consideration the sum of the two sulfur lines, i.e. [SIII]18.7 μm and [SIV]10.5 μm , as a tracer of the SFR. We plot in Fig. 5a the correlation of the luminosity derived by summing [SIII]18.7 μm and [SIV]10.5 μm with the SFR derived from the




Fig. 5. (a: left) Correlation between the [SIV]10.5 μm and [SIII]18.7 μm summed emission lines luminosity, expressed in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black dashed line) for a composite catalogue of SFG (red star) and from the $\text{H}\alpha$ luminosity (corrected for the IR luminosity) for LMG (green triangles). The shaded green area in the three plots indicates the 95% confidence interval. (b: centre) Correlation between the [NeII]12.8 μm and [SIV]10.5 μm summed luminosities and the SFR derived from the total IR luminosity (black dashed line) for SFG (red star) and from the $\text{H}\alpha$ luminosity (corrected for the IR luminosity) for LMG (green triangles). (c: right) Correlation between the [NeIII]15.6 μm and [SIII]18.7 μm summed emission lines and the SFR derived from the total IR luminosity (black dashed line) for SFG (red star) and from the $\text{H}\alpha$ luminosity (corrected for the IR luminosity) for LMG (green triangles). In all three figures, the purple squares highlight the ULIRG population in the SFG sample.

total IR luminosity:

$$\log \left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}} \right) = (1.07 \pm 0.06) + (1.04 \pm 0.05) \log \left(\frac{L_{[\text{SIII}]+[\text{SIV}]} }{10^{41} \text{ erg s}^{-1}} \right) \quad (6)$$

For this relation were used 52 SFG and 25 LMG, obtaining a relation with a Pearson r coefficient of $r=0.90$.

As shown in Table A.1, Ne^{2+} has a ionisation potential (IP) of 21.56 eV, while Ne^{2+} has a IP of 40.96 eV and a potential of the next stage at 63.45 eV, thus covering the $\sim 20 - 60$ eV interval in energy. This is roughly the same ionisation interval covered by the [NeII]+[SIV], with [SIV] having a IP of 34.79 eV and the next stage IP at 47.22 eV, or by [NeIII]+[SIII], with [SIII]18.7 μm that has a IP of 23.34 eV. Analyzing the former pair, we find the linear correlation between $L_{[\text{NeII}]+[\text{SIV}]}$ and SFR shown in Fig. 5b and expressed as:

$$\log \left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}} \right) = (0.85 \pm 0.05) + (0.94 \pm 0.04) \log \left(\frac{L_{[\text{NeII}]+[\text{SIV}]} }{10^{41} \text{ erg s}^{-1}} \right) \quad (7)$$

This relation has been calculated using 77 galaxies, of which 56 SFG and 21 LMG, and has a Pearson correlation coefficient $r = 0.93$. The small number of SFG used for determining this relation is due to the lack of [SIV] measurements available in literature, due to the intrinsic line faintness in star forming galaxies (see Table D.1) and the position of the line in the silicate absorption band in the 9-11 μm range.

The relation between SFR and the [NeIII]+[SIII] emission sum is shown in Fig. 5c and can be expressed as:

$$\log \left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}} \right) = (1.16 \pm 0.04) + (1.03 \pm 0.04) \log \left(\frac{L_{[\text{NeIII}]+[\text{SIII}]} }{10^{41} \text{ erg s}^{-1}} \right) \quad (8)$$

This relation has been calculated using 165 galaxies, of which 140 SFG and 25 LMG, and has a Pearson correlation coefficient $r = 0.86$.

3.2.5. PAH SFR tracer

In the mid-IR range, the emission features due to the PAH molecules arise from PDR around HII regions embedding young stars (e.g., Draine & Li 2007), and can be used as SFR tracers. We analyse the PAH features at 6.2 and 11.3 μm , considering the same sample of 155 SFG and the sample of 103 AGN galaxies described in section 2 for the PAH analysis. As discussed in Section 3.1, we have not computed the correlation between the PAH and the IR luminosity in LMG, therefore we exclude the LMG for the SFR determination with the PAH.

The use of the PAH as a measure of the SFR was originally proposed by Wu et al. (2005), who used the *Spitzer*-IRAC camera (Fazio et al. 2004) 8 μm band, whose flux density is dominated by the strongest PAH feature, i.e that at 7.7 μm . They derive a calibration of the 8 μm SFR using the radio VLA emission at 1.4GHz and the $\text{H}\alpha$ luminosity from the SFR-radio luminosity relation given by Yun et al. (2001) and the SFR- $\text{H}\alpha$ luminosity from Kennicutt (1998), respectively.

More recently, the relatively bright PAH features at 6.2, 7.7 and 11.3 μm have been used to derive a total PAH luminosity and has been correlated to the extinction corrected $\text{H}\alpha$ luminosity of a sample of 227 galaxies (Shipley et al. 2016). For star-forming galaxies (105 galaxies), the total PAH luminosity correlates linearly with the extinction-corrected $\text{H}\alpha$ luminosity.

In Fig. 6 we show the correlation of the single PAH features and the SFR: panel (a) shows the relation calculated for the 6.2 μm feature using 142 SFG and 56 AGN, while panel (b) shows the relation for the 11.3 μm feature, for which were used 142 SFG and 77 AGN. When computing these correlations, we excluded all ULIRGs from the original sample of AGN and SFG, following the approach by Pope et al. (2013) to avoid saturation effects in the PAH-to-continuum luminosities. Moreover, when considering the AGN sample, we discarded all sources with a total IR luminosity greater than $L_{\text{IR}} = 10^{45} \text{ erg s}^{-1}$, where the total IR




Fig. 6. (a: left) Correlation between the PAH emission feature at 6.2 μm , expressed in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black line) for a composite catalogue of SFG (red star) and AGN dominated galaxies (blue square). The shaded area in the three plots indicates the 95% confidence interval. **(b: centre)** Correlation between the PAH emission feature at 11.3 μm , expressed in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black line) for a composite catalogue of SFG (red star) and AGN dominated galaxies (blue square). In panels (a) and (b) we exclude ULIRGs from both AGN and SFG samples due to the known PAH deficit in these sources, plus AGN with luminosities above $10^{45} \text{ erg s}^{-1}$ that could dominate the IR continuum used to estimate the SFR. **(c: right)** Comparison between the relation between PAH total luminosity and SFR derived by Shipley et al. (2016) (black solid line) and using our sample (red dashed line). Red stars indicate SFG, purple squares indicate the ULIRG sub-sample included in the GOALS sample, and grey circles represent a sample of local galaxies used in Xie & Ho (2019).

luminosity, and therefore the derived SFR, could be severely contaminated by the AGN. Nevertheless, these are a few sources that do not affect the derived fit parameter when they are included in the fit. The two relations are described respectively by the following equations, which also report the Pearson r coefficient relative to each equation:

$$\log\left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}}\right) = (0.37 \pm 0.04) + (0.76 \pm 0.03) \log\left(\frac{L_{\text{PAH}6.2}}{10^{41} \text{ erg s}^{-1}}\right), r = 0.87 \quad (9)$$

$$\log\left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}}\right) = (0.41 \pm 0.03) + (0.73 \pm 0.02) \log\left(\frac{L_{\text{PAH}11.3}}{10^{41} \text{ erg s}^{-1}}\right), r = 0.89 \quad (10)$$

Both results show a shallow slope, significantly lower than unity. This is a consequence of including the AGN sample. Considering only the SFG sample, the slopes would increase, becoming 0.87 for the PAH feature at 6.2 μm , and of 0.90 for the 11.3 μm feature. On the other hand, while we excluded all ULIRGs, including these objects does not result in a significant change in the slope, that remains comparable within the errors. In particular, including the entire AGN and SFG samples for the PAH feature at 6.2 μm results in a slope of 0.81 ± 0.03 , while for the PAH feature at 11.3 μm the slope would be of 0.78 ± 0.03 .

We compared our results with the one proposed by Shipley et al. (2016): in order to properly determine the SFR using the H_{α} and 24 μm luminosities, as done by these authors, we selected only the SFG, excluding the AGN, leaving our sample composed of 100 sources with the 6.2, 7.7, 11.3 PAH features, the 24 μm and the H_{α} data. We find that our estimate of the SFR is higher by ~ 0.25 dex than that derived by Shipley et al. (2016) and almost

linear in slope, (see Fig. 6c). We obtain:

$$\log\left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}}\right) = (-1.28 \pm 0.23) + (1.08 \pm 0.10) \log\left(\frac{L_{\Sigma\text{PAH}}}{10^{41} \text{ erg s}^{-1}}\right) \quad (11)$$

with a Pearson r coefficient of $r=0.66$. While the slope of our result is comparable, within the error, to that obtained by Shipley et al., we attribute the difference in the intercept to the significant difference in sample characteristics. In particular, the disagreement arises when we extrapolate the results obtained by Shipley et al. for local galaxies. The sample used by Shipley et al. is in fact composed of galaxies with redshift in the $z \sim 0.2 - 0.6$ range, while our sample has a mean redshift of $z \sim 0.027$. Moreover, while our sample is primarily composed of LIRGs and ULIRGs, the sample used by these authors is mainly constituted of galaxies in the $L_{\text{IR}} \sim 10^9 - 10^{12} L_{\odot}$ interval. We additionally compared our derived relation to a sample of local SFG described by Xie & Ho (2019), for which was available in the literature the SFR derived from H_{α} , corrected by the flux at 24 μm . This sample is composed of local galaxies and, while the bulk of these galaxies present a lower total PAH luminosity, it follows the same relation derived from the GOALS sample. This suggests that the difference observed between our result and the result by Shipley et al. is indeed due to an intrinsic difference of the sample used.

3.2.6. H_2 SFR tracers

Molecular hydrogen is the most abundant molecule in the Universe and can be found in various environments. It can be excited by UV fluorescence (Black & van Dishoeck 1987), shocks (Hollenbach & McKee 1989) or X-ray illumination (Maloney 1997), thus probing various astrophysical environments. H_2 forms on the surface of dust grains, affecting the ISM chemistry, and acts as a coolant. It is particularly important in all processes that regulate star formation and galaxy evolution, where the principal




Fig. 7. (a: left) Correlation between the H₂ molecular line at 9.67 μm , expressed in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black line) for a catalogue of SFG (red star). (b: centre) Correlation between the H₂ molecular line at 12.28 μm , expressed in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black line) for a catalogue of SFG (red star). The green triangle show LMG detection, not used to derive the correlation. (c: right) Correlation between the H₂ molecular line at 17.03 μm , expressed in units of $10^{41} \text{ erg s}^{-1}$ and the SFR derived from the total IR luminosity (black line) for a catalogue of SFG (red star). In all three figures, the shaded area shows the 95% confidence interval of the relations, and the purple squares highlight the ULIRG population in the SFG sample.

mechanism of these lines is associated with the UV radiation from massive stars.

Comparing theoretical models to observations, Rigopoulou et al. (2002) found evidence that an important fraction of H₂ emission in SFG can originate in PDRs. In this section we test the use of different H₂ molecular lines as SFR tracers. In Fig. 7a we show the correlation between the H₂ (S(3)) molecular line at 9.67 μm and the SFR determined from the total L_{IR} for the SFG in the GOALS sample. We find a relation expressed by:

$$\log\left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}}\right) = (1.57 \pm 0.02) + (0.71 \pm 0.03) \log\left(\frac{L_{\text{S}(3)}}{10^{41} \text{ erg s}^{-1}}\right) \quad (12)$$

determined using 168 SFG, with a Pearson r coefficient of $r=0.87$. Fig. 7b shows the correlation between the H₂ (S(2)) line at 12.28 μm and the SFR, determined with a sample of 126 SFG, with a Pearson coefficient of $r=0.86$ and expressed by:

$$\log\left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}}\right) = (1.71 \pm 0.02) + (0.87 \pm 0.04) \log\left(\frac{L_{\text{S}(2)}}{10^{41} \text{ erg s}^{-1}}\right) \quad (13)$$

Finally, Fig. 7c shows the correlation between the H₂ (S(1)) line at 17.03 μm and the SFR, derived from a sample of 154 SFG, with $r=0.85$ and expressed by:

$$\log\left(\frac{\text{SFR}}{\text{M}_{\odot} \text{ yr}^{-1}}\right) = (1.39 \pm 0.02) + (0.92 \pm 0.04) \log\left(\frac{L_{\text{S}(1)}}{10^{41} \text{ erg s}^{-1}}\right) \quad (14)$$

In Fig. 7b and c we also report, in green, seven detections in LMG of S(2) and S(1), respectively. We did not include the LMG sample in the correlation due to the small number of sources, limiting ourselves to a comparison of the results. While a good correlation is lacking for the S(1) line, for the S(2) line we have a good agreement between the LMG population and the SFG one.

This can be of particular interest in those LMG in which the CO emission is considerably lower than what it would correspond to the estimated SFR, implying the presence of CO-dark molecular gas that is not traced by the CO emission in LMG (Togi & Smith 2016). We speculate that the sub-linear slopes obtained for the S(2) and S(3) lines might be linked to the different gas excitation temperature associated with each line. In particular, by moving to higher excitation lines, and thus to shorter wavelengths, the H₂ transitions are originated by an increasingly warmer gas in a thinner layer of the molecular gas clouds. Tracing the warmest material might cause the flattening of the slope, since colder star-forming clouds are not detected by the higher transitions. Additionally, S(2) and S(3) lines could have a larger contribution from other excitation mechanisms, as suggested by the different excitation temperatures measured for these transitions in the Boltzmann diagrams for nearby galaxies (e.g. Tommasin et al. 2010).

We have not included the AGN sample in this Section due to the different excitation mechanisms that can contribute to the H₂ rotational lines in these sources, such as shocks or X-ray illumination. For the same reason H₂ lines are neither used as BHAR tracers in Section 3.3, since they respond to the excitation temperature rather than the hardness of the ionising continuum. This makes quite difficult to establish a clear connexion between the line intensities and the SFR or the accreted mass onto the black hole when both contributions are present.

3.3. BHAR tracers

The [OIV]25.9 μm and the [NeV]24.3 μm lines can be used to trace AGN activity. From our catalogue of AGN, derived from the 12 μm sample, we compiled from the literature the 2-10 keV X-rays fluxes corrected for absorption. We then selected all objects with hydrogen column density of $N_{\text{H}} \leq 5 \times 10^{23} \text{ cm}^{-2}$, because we want to exclude Compton-thick objects for which the 2-10 keV X-rays can be substantially absorbed, thus obtaining a sub-catalogue of 42 objects. For these objects, we investigated the correlation between the [OIV]25.9 μm line luminosity and the 2-10 keV X-ray luminosity (see Fig. 8a).




Fig. 8. (a: left) Linear correlation between the [OIV]25.9 μm and the 2-10 keV X-ray luminosity. (b: centre) Linear correlation between the [NeV]24.3 μm line luminosity and the 2-10 keV X-ray luminosity. (c: right) Linear correlation between the [OIV]25.9 μm line luminosity and the 19 μm luminosity. All luminosities are expressed in units of $10^{41} \text{ erg s}^{-1}$.

We find a correlation expressed by the equation:

$$\log\left(\frac{L_X}{10^{41} \text{ erg s}^{-1}}\right) = (1.85 \pm 0.07) + (1.21 \pm 0.10) \log\left(\frac{L_{[\text{OIV}]25.9}}{10^{41} \text{ erg s}^{-1}}\right) \quad (15)$$

with a Pearson r coefficient of $r=0.87$. In order to obtain a measure of the BHAR, it is necessary to convert the luminosity in the 2-10 keV band to the bolometric luminosity of the object, and from there to the BHAR ($L_{\text{AGN}} = \eta \dot{M}_{\text{BH}} c^2$). Different studies have been carried out to determine the best bolometric correction to apply when considering the 2-10 keV luminosity, with those by Marconi et al. (2004) and Lusso et al. (2012) being the most used in the literature. The resulting bolometric luminosities obtained applying these corrections are different, giving us a range of possible values. We applied both correction to our data, and then from the resulting bolometric luminosities we calculated two linear relations linking the [OIV]25.9 μm line luminosity to the BHAR. Assuming a radiative efficiency of $\eta=0.1$, we obtained in equation 16 the linear correlation applying the bolometric correction from Lusso et al. (2012), and in equation 17 the one applying the correction from Marconi et al. (2004). The equations are reported each followed by its Pearson r coefficient:

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-1.65 \pm 0.07) + (1.04 \pm 0.09) \log\left(\frac{L_{[\text{OIV}]25.9}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.86 \quad (16)$$

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-1.66 \pm 0.09) + (1.49 \pm 0.12) \log\left(\frac{L_{[\text{OIV}]25.9}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.87 \quad (17)$$

In a similar way, we first calculated the linear correlation between the [NeV]24.3 μm line luminosity and the 2-10 keV luminosity (see right panel in Fig. 8). In this case, we have a total of 34 objects, due to a smaller number of [NeV] available data,

and we obtained a linear relation described by the equation:

$$\log\left(\frac{L_X}{10^{41} \text{ erg s}^{-1}}\right) = (2.40 \pm 0.08) + (0.95 \pm 0.11) \log\left(\frac{L_{[\text{NeV}]24.3}}{10^{41} \text{ erg s}^{-1}}\right) \quad (18)$$

with a Pearson coefficient of $r=0.84$. From this, we then applied the same bolometric corrections, obtaining linear correlations between the line luminosity and the BHAR, expressed by equations 19 and 20 for the Lusso et al. (2012) and Marconi et al. (2004) correction respectively:

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-1.11 \pm 0.09) + (1.06 \pm 0.12) \log\left(\frac{L_{[\text{NeV}]24.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.83 \quad (19)$$

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-0.89 \pm 0.11) + (1.47 \pm 0.15) \log\left(\frac{L_{[\text{NeV}]24.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.84 \quad (20)$$

As a general trend, we find that both the [NeV]24.3 μm and [OIV]25.9 μm lines correlate linearly with the 2-10 keV X-ray luminosity, thus providing a good proxy to measure the AGN activity, as shown in Fig. 8 a and b.

Following the work by Tommasin et al. (2010), we also analysed the correlation between the [NeV]24.3 and [OIV]25.9 μm lines with the luminosity at 19 μm . The luminosity at 19 μm ($L_{19\mu\text{m}}$) was used for two main reasons. On one hand, accurate *Spitzer*-IRS observations are available for the 19 μm flux density, from two different apertures, for the considered sample of galaxies. On the other hand, the 19 μm photometry data are the only available accurate photometric data closest to the emission at 12 μm . The 12 μm emission is, in turn, the best proxy for the bolometric flux of an active galaxy (Spinoglio et al. 1995). In particular, Tommasin et al. (2010) found that the 19 μm luminosity correlates with the [NeV]14.3 μm line luminosity.

The $L_{19\mu\text{m}}$ has been used by Tommasin et al. to compute the percentage of AGN and starburst components in 51 sources of

their sample. Following these results, we selected all sources with an AGN component at $19\mu\text{m}$ equal or above 85%, and for this sub-sample of 35 objects we determined the correlation between the [OIV]25.9 μm and the [NeV]24.3 μm line luminosities with the $19\mu\text{m}$ luminosity. These correlations are shown in Fig. 8c and Fig. 9a, respectively, and expressed by Eq. 21 and 22 respectively, where r indicated the Pearson correlation coefficient, and n the number of objects used to derive the correlation.

$$\log\left(\frac{L_{19\mu\text{m}}}{10^{41} \text{ erg s}^{-1}}\right) = (2.72 \pm 0.06) + (0.77 \pm 0.10) \log\left(\frac{L_{[\text{OIV}]25.9}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.81, \quad n = 32 \quad (21)$$

$$\log\left(\frac{L_{19\mu\text{m}}}{10^{41} \text{ erg s}^{-1}}\right) = (3.12 \pm 0.06) + (0.81 \pm 0.10) \log\left(\frac{L_{[\text{NeV}]24.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.85, \quad n = 30 \quad (22)$$

We then used these relations, (equations 21 and 22,) to determine the bolometric luminosity of our sources, and then the accretion rate. In order to calculate the bolometric luminosity, we used the relation, calculated by Spinoglio et al. (1995), that links the bolometric luminosity to the $12\mu\text{m}$ luminosity. The $12\mu\text{m}$ luminosity for our sample was determined using data from Deo et al. (2009): the authors report continuum measurements at 5.5 μm , 14.7 μm and 20 μm taken from *Spitzer* low-resolution spectroscopic observations. Where possible, we interpolated the continuum slope and the $12\mu\text{m}$ continuum flux using the 5.5 μm and 14.7 μm measurements, otherwise (but only in three cases) using the extrapolation from the 14.7 μm and 20 μm fluxes. We then matched the Deo et al. sample with the Tommasin et al. sample, calculated the bolometric luminosity starting from the monochromatic $12\mu\text{m}$ luminosity for those sources with a 85% AGN component. Similarly to what has been done for the 2-10 keV luminosity, we determined the BHAR starting from the bolometric luminosity derived from the $12\mu\text{m}$ luminosity, and its relation to the [NeV]24.3 μm and [OIV]25.9 μm line luminosities, obtaining the following relations:

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-1.14 \pm 0.07) + (0.67 \pm 0.15) \log\left(\frac{L_{[\text{OIV}]25.9}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.68, \quad n = 26 \quad (23)$$

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-0.77 \pm 0.10) + (0.77 \pm 0.16) \log\left(\frac{L_{[\text{NeV}]24.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.71, \quad n = 24 \quad (24)$$

The smaller number of objects used to derive these relations is due to the lack of data for the determination of the continuum at $12\mu\text{m}$.

Measuring the BHAR requires important approximations in terms of bolometric correction, which can yield significantly different results. Between the three proposed bolometric corrections, we note that the correlations obtained when applying the Marconi correction are steeper than those obtained from the Lusso and Spinoglio correction. In particular, we compare the

results in Fig. 9: for both the [OIV]25.9 μm (panel b, at the centre) and [NeV]24.3 μm (panel c, on the right) lines, we note that the results derived from the corrections of Spinoglio et al. (1995) and Lusso et al. (2012) show a similar, flatter slope, indicating, within the errors, the expected linear relation between the line tracers and the BHAR. It is important to note that the Marconi et al. and Lusso et al. bolometric corrections are based on a third-degree polynomial transformation of the X-ray luminosity. This necessarily leads to deviations from the slopes obtained in Eq. 15 and 18.

We refer to the Appendix C for a discussion on the use of the mid-ionisation lines of [NeIII]15.5 μm and [SIV]10.5 μm as alternative BHAR tracers.

4. Discussion

4.1. Applicability of our results to composite objects

The coexistence of AGN and star formation in galaxies is well known (see, e.g. Pérez-Torres et al. 2021, for a review) and therefore we expect that many galaxies are characterised by both components at work together. In general, when considering composite objects, with both a SF and an AGN component detectable, our results in Section 3.2 and Section 3.3 can still be applied.

We use the GOALS sample to determine whether the results on the SFR tracers can be applied to composite objects (Section 3.2). The GOALS sample is composed of SFG, but for part of the sample an AGN component is also present, if not always detected. For our calibrations, we have excluded all sources for which a [NeV] emission line is detected. Here we instead include all objects of the sample. In Fig. 10 we plot the ratio of the SFR obtained using the [NeII]+[NeIII] tracer to the SFR obtained using the 11.3 μm PAH feature, as a function of the Equivalent Width (EW) of the PAH feature at 6.2 μm , as determined by Stierwalt et al. (2014). The mean ratio for the entire population is -0.049, with the median value equal to -0.023 and 50% of the sample included in the [-0.18, 0.06] interval. While there is some dispersion around zero, for the majority of the sample we obtain a similar SFR whether we use the [NeII]+[NeIII] tracer or the PAH tracer. This suggests that for mixed objects these two tracers are equivalent, and independent of the presence of an AGN. Significant differences are observable for the extreme objects in the sample: the sources with the larger AGN content are located in the top left corner, for which the SFR obtained using the [NeII]+[NeIII] tracer is significantly higher than the SFR obtained using the PAH. This is plausibly due to an increase of the [NeIII] emission related to the presence of the AGN.

4.2. Comparison among the various tracers

Summarising the results presented in Section 3.2, among the various SFR tracers presented, for galaxies dominated by SF processes we suggest that the best tracers are the [CII]158 μm line (slope $\alpha = 0.89 \pm 0.02$ and correlation coefficient $r = 0.92$) or the combination of the [OI] and [OIII] lines (slope $\alpha = 1.25 \pm 0.04$ and $r = 0.94$) for high-redshift galaxies observed from the ground with submillimeter telescopes. For galaxies observed from space or airborne facilities (but also from the new generation of the very large optical ground based telescopes in the 8-13 μm atmospheric window, for local galaxies), the best SFR tracers are the combination of the [NeII] and [NeIII] lines (slope $\alpha = 0.96 \pm 0.03$ and $r = 0.91$) or the combination of the other mid-IR fine-structure lines. For galaxies containing an AGN, the PAH features can reliably be used and the PAH 11.3 μm feature is probably the best SFR




Fig. 9. (a: left) Linear correlation between the [NeV]24.3 μm line luminosity and the 19 μm luminosity, expressed in units of $10^{41} \text{ erg s}^{-1}$. (b: centre) Comparison of the three different relations between [OIV]25.9 μm line luminosity and the BHAR: the blue dashed line reports the relation obtained from the Lusso et al. bolometric correction, the red dotted line the relation from the Marconi et al. bolometric correction, and the green solid line the results from the Spinoglio et al. correction. (c: right) Same as panel (a), for the [NeV]24.3 μm line.




Fig. 10. Comparison of the ratio between the SFR obtained using the [NeII]+[NeIII] tracer and the SFR obtained using the PAH feature at 11.3 μm , against the EW of the PAH feature at 6.2 μm for the GOALS sample. The colour gradient indicates the percentage of the AGN component in each object as determined by mid-IR tracers (Díaz-Santos et al. 2017). The black dashed line shows the mean value of the SFR ratio for the population, with the shaded area indicating where the 50% of the entire sample is located. The red dotted lines represent the 1σ interval around the mean value.

tracer (slope $\alpha = 0.75 \pm 0.04$ and $r = 0.79$). As an alternative, the [NeII] and [NeIII] lines can still be used also in AGN, with the correction to the total neon flux that can be computed using one of the [NeV] lines, as suggested by Zhuang et al. (2019).

When measuring the BHAR, following the results presented in Section 3.3, the [NeV] lines at 14.3 and 24.3 μm are exclusive probes of AGN activity, because their emission is a direct signature of the hard ionising spectrum due to the accretion process. However, these lines are fainter than the [OIV]25.9 μm line, which can be detected more easily in faint objects. When using the [OIV] line as a BHAR tracer, it is important to keep into

account the possible contamination due to SF processes, because its emission can be attributed also, to some extent, to starburst activity (Lutz et al. 1998).

4.3. Comparison with previous line calibrations

We briefly compare here the results of the line calibrations obtained in this work with those of Spinoglio et al. (2012, 2014) and Gruppioni et al. (2016), while leaving all the details of the comparisons to Appendix E. We note that in this work, to derive the correlations, we used the orthogonal distance regression method, instead of the least-squares minimisation method, which was used by the other authors, because we consider the total IR luminosity and the line luminosity two independent variables.

The AGN sample used in this work and in Spinoglio et al. is the same for the lines in the 10–35 μm interval, and the differences in the correlation are only due to the different methods of analysis. We note that in this work, in order to obtain a correlation between the total L_{IR} and the [OIV]25.9 μm line luminosity that better represents the AGN population, we did not use the entire sample of AGNs by Tommasin et al. (2008, 2010), but a sub-sample of objects with an AGN component in the 19 μm luminosity of at least 85% (see Section 3.1).

For the SFG sample, while in Spinoglio et al. the sample of galaxies described by Bernard-Salas et al. (2009) was used, in this work we have expanded the same sample by including the LIRGs and ULIRGs sample of GOALS, as described in Section 2. Nonetheless, we obtain comparable results, except in the cases of the PAH feature at 11.3 μm and of the H_2 line at 17.03 μm , for which our results show line luminosities an order of magnitude higher. This is due to the presence, in our sample, of LIRGs and ULIRGs, which shift the relation toward a steeper slope.

For the lines at wavelengths in the 50–158 μm range, we refer to the Appendix E, where we discuss the differences and plot the results of the different calibrations of Spinoglio et al. with respect to our results.

Because of the different analytical method used to determine the correlations between total and line luminosities, for the comparison with Spinoglio et al., we have recomputed our calibrations using the least square fit and give the results in the Appendix E. For the comparison with Gruppioni et al., instead, we recomputed the correlations of the Gruppioni et al. sample, applying the orthogonal distance regression and then compare these with

our results. As a general trend, we find good agreement between our results and those by Gruppioni et al., with comparable slopes within 2σ of each other. For all the details and the plots of the results of the different calibrations, we refer to the Appendix E.

We do not compare our results with those obtained by Bonato et al. (2019), since the methods of analysis are widely different. In particular, while in this work we calibrate the line luminosities leaving the slope of the correlations as a free parameter, in Bonato et al. (2019) the slope of the relation was fixed to unity, thus giving rise to substantial differences in the results.

4.4. Metallicity and SFR tracers

In Section 3.2.2 we revise the [CII]-SFR relation for a wide galaxy sample, from LMG to extreme ULIRGs. In Section 3.2.4 we derive a measure of the SFR through the neon and sulfur mid-IR lines and propose new SFR tracers using different combinations of these lines. In this section we discuss the possible effects that metallicity, and the associated changes in the ISM of these galaxies, may have on these tracers.

First of all, for the sample of dwarf galaxies, we adopted SFR values derived from the observed H_α luminosity and corrected from the total IR luminosity (Rémy-Ruyer et al. 2015). This is motivated by the underestimation of the SFR by the IR luminosity at very low metallicities ($12 + \log(O/H) \lesssim 8.5$; Lee et al. 2013), due to the lower metal abundance in these galaxies compared to SFG. In principle, the lower dust to gas ratio of LMG should also have an impact on the observed intensities of the fine-structure lines. This is, however, balanced by the higher cooling rates in these transitions, as discussed by De Looze et al. (2014).

Fig. 3a shows that the [CII]158 μm emission in LMG follows the trend found in SFG with no need to perform any additional correction for metallicity in these galaxies. Similarly, the different combinations of [NeII]12.8 μm , [NeIII]15.6 μm , [SIII]18.7 μm , and [SIV]10.5 μm , shown in Figs. 3 and 5, follow the correlation of solar-like metallicity galaxies. The higher cooling rates in LMG are particularly evident when the neon and sulfur transitions are considered. While the [NeII]12.8 μm emission in LMG scales with L_{IR} similarly as for SFG (Fig. 1b), the [NeIII]15.6 μm line becomes comparatively much brighter for a given IR luminosity, more than one order of magnitude above the correlation found for SFG, as can also be seen by the value of the constant b in the best fit equation for the [NeIII]15.6 μm line in Table A.1. When considering the sulfur lines, this effect is even more pronounced (see Fig. B.4a and Table A.1). This means that mid- to high-ionisation species such as Ne^{2+} or S^{3+} trace a contribution to the star-formation that is not revealed by either the Ne^+ and S^{2+} low-ionisation gas or the IR emission. Thus, the combination of low and high-ionisation lines allows us to trace the total star formation in both low- and solar-metallicity galaxies (Ho & Keto 2007; Zhuang et al. 2019).

In the case of the [CII]158 μm line, the Fig. 3a suggests that this transition still remains a dominant coolant of the ISM at low metallicities. Given its low ionisation potential (11.3 eV, see Table A.1), this line can originate from both neutral and ionised gas, and one could expect a decreasing contribution from the neutral component as the ionisation field becomes harder at low metallicities. However, Cormier et al. (2019) demonstrated that the PDR contribution to the global [CII]158 μm emission is still dominant for the same LMG sample used in this work. This is also in line with the results of Croxall et al. (2017), suggesting that the [CII]158 μm emission linked to ionised gas is of the order of $\sim 10\%$ in LMG, and up to maximum $< 40\%$ where a high-U is required. Moreover, analytical models developed in these studies

show a decrease of [CII] emission from ionised gas with decreasing metallicity with $\sim 55 - 75\%$ of the [CII] emission arising from PDRs in LMG, reaching almost 100% when the metallicity decreases below $1/4 Z_\odot$. Additionally, the thickness of the [CII] layer increases for molecular clouds exposed to the harder radiation fields typical of LMG. This is shown by the detection of higher [CII]/CO(1-0) ratios in local LMG when compared to SFG with solar or super-solar metallicity (e.g. Madden et al. 1997; Madden 2000; Hunter et al. 2001), and it is supported by PDR models (Bolatto et al. 1999; Röllig et al. 2006).

While a detailed study of the ionised gas and PDR structure is out of the scope of the present work, the results discussed above suggest that both [CII]158 μm and the different combinations of neon and sulphur lines are robust star formation tracers, virtually independent of dust extinction, that can be applied to a wide diversity of environments with different physical conditions and metallicities. Specifically, the variations expected from the changes in the chemical abundances are mostly balanced by the increase in the cooling rates of these transitions.

4.5. Comparison with high- z data

We compare the calibrations described in Section 3.1 obtained for local galaxies to high-redshift detections of sources obtained with ALMA. In particular, we consider detections of the [OIII]88 μm , [NII]122 μm , [CII]158 μm and [NII]205 μm lines at $z \geq 3$. Sources identified as QSOs are compared to local AGN results, while sources for which a classification is not given in the literature, or are classified as starburst galaxies, are compared to local SFG.

Fig. 11a shows the comparison between local and high redshift detections of the [CII]158 μm line for QSOs galaxies, and in particular 27 detections at $z \sim 6$ reported in Venemans et al. (2020), plus the detections by Walter et al. (2018) for one source at $z \sim 6.08$ and by Hashimoto et al. (2018) for two sources at $z \sim 7.1$, for a total of 30 sources. Fig. 11b displays the comparison of local and high redshift SFG: 84 detections from the ALPINE catalogue (Faisst et al. 2020) plus other 9 detections (Inoue et al. 2016; Vishwas et al. 2018; Walter et al. 2018; De Breuck et al. 2019; Hashimoto 2019; Harikane et al. 2019; Rybak et al. 2019) for a total of 93 objects with redshift in the $4.2 \leq z \leq 7.2$ range. While some outliers are present, the bulk of the detections in both cases lies within the prediction interval, that we show in the figures at the 95% level, suggesting that the relations derived for local galaxies hold for high redshift sources.

We compare in Fig. 11c the $L_{\text{[CII]}}$ -SFR relation extrapolated for local galaxies (see Section 3.2.2) with ALPINE detections of the [CII]158 μm emission line in starburst galaxies. We find that high redshift objects follow the same $L_{\text{[CII]}}$ -SFR relation as local SFG and LMG, thus suggesting little or no evolution in the $L_{\text{[CII]}}$ -SFR relation. In particular, no deficit of [CII] emission is seen for the highest SFR values of $\sim 100 M_\odot \text{ yr}^{-1}$, suggesting that starbursts at high- z might not behave like local ULIRGs such as those shown in Fig. 3 (purple squares). An analogous result was found by Schaerer et al. (2020), who analysed a large sample of galaxies at high redshift ($z \sim 4-6$) observed by ALMA. Leung et al. (2020) explored the possibility of tracing the SFR at the Epoch of Recombination with [CII]158 μm using simulated data. Although they find a good correlation, when compared to observed data, the simulated sample shows an average [CII] luminosity lower than the one obtained from ALMA pointed observations. A similar result was obtained for simulations of




Fig. 11. (a: left) Comparison between the $L_{\text{IR}}-L_{\text{CII}158}$ relation (black dashes line) for local AGN dominated galaxies (gray squares) and high redshift detections for QSOs (blue symbols). The shaded area shows the 95% prediction interval. **(b: centre)** Same for high redshift starburst galaxies (pink symbols). **(c: right)** Comparison of the local $\log(L_{\text{CII}})-\log(\text{SFR})$ relation (black dashes line) and $L_{\text{CII}}-\text{SFR}$ values of high redshift sources (pink symbols). Grey dots represent local star forming galaxies. The shaded area shows the 95% prediction interval for the local relation.




Fig. 12. (a: left) Comparison between the $L_{\text{IR}}-L_{\text{OIII}88}$ relation (black dashes line) for local AGN dominated galaxies (gray squares) and high redshift detections for QSOs (blue symbols). **(b: centre)** Comparison between the $L_{\text{IR}}-L_{\text{OIII}88}$ relation (red dashes line) for local SFG (gray stars) and high redshift detections for starburst galaxies (pink symbols). The red shaded area shows the 95% prediction interval of this relation. For comparison, the green dashed line shows the $L_{\text{IR}}-L_{\text{OIII}88}$ relation for local LMG, with gray squares indicating local LMG and the green shaded area giving the 95% prediction interval for this relation. **(c: right)** Comparison of the local $L_{\text{IR}}-L_{\text{NII}205}$ relation (black dashes line) for high redshift detections of SFG (pink symbols). Grey stars represent local star forming galaxies. The shaded area in all the figures shows the 95% prediction interval for the local relation.

high redshift systems (Ferrara et al. 2019) showing the presence of under-luminous [CII] emission.

In Fig. 12a we show the comparison between the local $L_{\text{IR}}-L_{\text{OIII}88}$ relation for AGN and high redshift detections in QSOs (Hashimoto et al. 2018; Walter et al. 2018), while panel (b) shows the relation between local and high redshift SFG (Inoue et al. 2016; Vishwas et al. 2018; Walter et al. 2018; De Breuck et al. 2019; Harikane et al. 2019; Hashimoto 2019; Tamura et al. 2019). We note that, while the high-redshift detections of [OIII]88 μm in QSOs are comparable with the relation derived from local AGN dominated galaxies, in SFG the majority of high redshift detections appears to be one to two orders of magnitude brighter at comparable L_{IR} . These detections are better represented by the $L_{\text{IR}}-L_{\text{OIII}88}$ relation derived for local LMG (see Fig. 12b). This suggests that the [OIII]88 μm line is produced in an environment with a higher ionising spectrum, similar to the ISM of local LMG (see Section 3.1).

Finally, the high redshift detections in panel (c) show the local $L_{\text{IR}}-L_{\text{NII}205}$ relation for SFG and 17 high-redshift detections of [NII]205 μm (Cunningham et al. 2020) with the local corre-

lation that well represents also the high redshift detections. We report in Fig. B.8 the comparison between the local $L_{\text{IR}}-L_{\text{NII}122}$ relation and high redshift SFG: in this case only one object (De Breuck et al. 2019) has been detected, while three (Harikane et al. 2019) objects have upper limits. Thus, only one object is located almost an order of magnitude higher than the local relation for SFG, and therefore there is no statistical evidence for a difference in the production mechanism of [NII] in high redshift objects, especially considering the results for the [NII]205 μm line.

4.6. Application to present and future facilities

The lines and features presented in this work can be observed in galaxies by present and future IR/(sub)-mm facilities. The *JWST* will be the next NASA observatory to explore the Universe in the near- and mid-IR spectral range. In particular, the *JWST* Mid-InfraRed Instrument (MIRI, Rieke et al. 2015; Wright et al. 2015) will be able to obtain imaging and spectra with unprecedented sensitivity in the 4.9–28.9 μm spectral range. In Fig. 13a

we compare the MIRI wavelength interval to the observability of key mid-IR lines at various redshifts. In particular, from the analysis presented in Section 3.2, it will be possible to study the SFR up to redshift $z \sim 3.5$ with the PAH feature at $6.2 \mu\text{m}$, up to redshift $z \sim 1.5$ with the PAH feature at $11.3 \mu\text{m}$, or up to $z \sim 0.8$ using combinations of the neon and sulfur lines.

On the other hand, while we do not report in Fig. 13a the [NeV]24.3 and [OIV]25.9 μm lines due to the limited wavelength range available for *JWST*, we show the [NeV]14.3 μm line, for which the same analysis carried out in Section 3.3 to derive the BHAR is applied and reported in Appendix C. The [NeV]14.3 μm line will be observed by *JWST*-MIRI up to redshift $z \leq 1$, while the [NeV]24.3 μm line will be observed up to redshift $z \leq 0.15$, and the [OIV]25.9 μm line will be observed up to redshift $z \leq 0.1$.

While *JWST*-MIRI will be able to observe molecular and atomic lines up to redshift $z \sim 2$, and the PAH up to higher redshift ($z \sim 3.5$ for the $6.2 \mu\text{m}$ feature), ALMA is able to trace the far-IR lines, like [CII]158 μm , up to redshift $z \sim 8$, as shown in Fig. 13b, providing important information on the evolution of galaxies with cosmic time. An analogous study can be found in Fig. 1 of Carilli & Walter (2013), where the CO transitions and other key tracers of the ISM are shown as a function of redshift versus frequency in terms of the observability by ALMA and JVLA (Karl J. Jansky Very Large Array, Perley et al. 2011). We also report in Fig. 13(b) the redshift of currently available data of ALMA detections: while lines like [CII]158 μm or [NII]205 μm are detected over a significant redshift interval, shorter wavelength lines, such as [OI]63 μm , present very few results. This is due to the difficulty of observing in the bands at higher frequency, for which the atmospheric absorption allows good visibility only for 10% of the total observational time.

In a near future the extremely large ground-based telescopes under construction, with dedicated instruments for *N*-band spectroscopy will be able to obtain observations of the mid-IR spectral range 8–13 μm and thus will test the [SIV] $_{10.5 \mu\text{m}}$ + [NeII] $_{12.8 \mu\text{m}}$ relation, for local galaxies and AGN. The ESO Extremely Large telescope (E-ELT; Gilmozzi & Spyromilio 2007) is expected to obtain first light at the end of 2025, with the Mid-infrared ELT Imager and Spectrograph (METIS, Brandl et al. 2021) instrument covering the *N* band with imaging and low resolution spectroscopy, the Thirty Meter Telescope (TMT; Schöck et al. 2009) will have completed first light and will be ready for science at the end of 2027, and the Giant Magellan Telescope (GMT; Johns et al. 2012) will be operational in 2029.

5. Summary and conclusions

In this work we systematically revise the calibration of lines and features in the 6–205 μm spectral range. We report new line to L_{IR} calibrations for three galaxy types: active galactic nuclei (AGN), low-metallicity galaxies (LMG) and star-forming galaxies (SFG), building well defined samples with available IR spectroscopy mainly obtained from *Herschel* and *Spitzer*. The main results of this work are:

- Statistically robust calibrations to the total IR luminosity have been obtained for the bright IR lines and features, including the PAH features, the H_2 pure rotational lines and the fine-structure lines, mostly for all the three galaxy populations, excluding the PAH features for the LMG and the high-ionisation fine structure lines for the SFG and LMG.
- The [CII]158 μm line can be used as SFR tracer for both LMG and SFG, covering 6 order of magnitudes in luminosity, independently of the source metallicity. Moreover, when

compared to high redshift detections, the correlation obtained for local galaxies can still be applied.

- The sum of the two far-IR oxygen lines, the [OI]63 μm line and the [OIII]88 μm line, can be used as a tracer for the SFR.
- A combination of low- and intermediate-ionisation lines from neon and/or sulfur is also a robust proxy for the SFR, independent of the metallicity. In particular, the [SIV]10.5 μm + [NeII]12.8 μm tracer will be accessible to ELTs facilities in the near future for galaxies in the local Universe.
- The brightest PAH features at 6.2 and 11.3 μm can be reliably used for SFR tracers in both SFG and AGN, while the lack of these features prevents their use in LMG.
- We present for the first time a correlation between the SFR and the H_2 molecular line fluxes at 9.7, 12.3 and 17.0 μm .
- The [CII]158 μm line, the combination of the two neon lines ([NeII]12.8 μm and [NeIII]15.5 μm) and, for **solar-like metallicity galaxies** that may harbour an AGN, the PAH11.3 μm feature are the best SFR tracers.
- The [NeV]14.3 μm , [NeV]24.3 μm and [OIV]25.9 μm lines are good tracers of the BHAR, derived either from the 2–10 keV luminosity or from the 12 μm luminosity.

These results will assume particular relevance with the upcoming launch of the *JWST*, which will observe with the MIRI instrument many of the mid-IR features in galaxies of the Local Universe and beyond, up to redshift of ~ 1 for the brightest lines, thus allowing to measure with spectroscopy both the SFR and the BHAR.

Acknowledgements. We thank the anonymous referee for his/her comments which helped to improve the present article. LS and JAFO acknowledge financial support by the Agenzia Spaziale Italiana (ASI) under the research contract 2018-31-HH.0.

References

- Allamandola L. J., Tielens A. G. G. M., Barker J. R., 1989, *ApJS*, 71, 733
 Armus L., et al., 2007, *ApJ*, 656, 148
 Armus L., et al., 2009, *PASP*, 121, 559
 Bernard-Salas J., et al., 2009, *ApJS*, 184, 230
 Berta S., et al., 2013, *A&A*, 551, A100
 Black J. H., van Dishoeck E. F., 1987, *ApJ*, 322, 412
 Boggs P. T., Rogers J. E., 1990, Orthogonal Distance Regression. Contemporary Mathematics Vol. 112, American Mathematical Society
 Bolatto A. D., Jackson J. M., Ingalls J. G., 1999, *ApJ*, 513, 275
 Bonato M., et al., 2019, *PASA*, 36, e017
 Brandl B. R., et al., 2006, *ApJ*, 653, 1129
 Brandl B., et al., 2021, *The Messenger*, 182, 22
 Buchanan C. L., Gallimore J. F., O’Dea C. P., Baum S. A., Axon D. J., Robinson A., Elitzur M., Elvis M., 2006, *AJ*, 132, 401
 Bullock J. S., Boylan-Kolchin M., 2017, *ARA&A*, 55, 343
 Calzetti D., et al., 2007, *ApJ*, 666, 870
 Carilli C. L., Walter F., 2013, *ARA&A*, 51, 105
 Carpenter J., Iono D., Kemper F., Wootten A., 2020, arXiv e-prints, p. arXiv:2001.11076
 Clegg P. E., et al., 1996, *A&A*, 315, L38
 Cormier D., et al., 2012, *A&A*, 548, A20
 Cormier D., et al., 2015, *A&A*, 578, A53
 Cormier D., et al., 2019, *A&A*, 626, A23
 Cortese L., et al., 2012, *A&A*, 544, A101
 Croxall K. V., et al., 2017, *ApJ*, 845, 96
 Cunningham D. J. M., et al., 2020, *MNRAS*, 494, 4090
 Dale D. A., Helou G., Brauer J. R., Cutri R. M., Malhotra S., Beichman C. A., 2004, *ApJ*, 604, 565
 De Breuck C., et al., 2019, arXiv e-prints, p. arXiv:1909.12554
 De Looze I., et al., 2014, *A&A*, 568, A62
 Deo R. P., Richards G. T., Crenshaw D. M., Kraemer S. B., 2009, *ApJ*, 705, 14
 Díaz-Santos T., et al., 2017, *ApJ*, 846, 32
 Draine B. T., Li A., 2007, *ApJ*, 657, 810
 Engelbracht C. W., Gordon K. D., Rieke G. H., Werner M. W., Dale D. A., Latter W. B., 2005, *ApJ*, 628, L29




Fig. 13. (Left) Observability of key mid-IR (dashed lines) lines compared to the *JWST*-MIRI instrument (shaded area) in the 0-1.6 redshift interval. (Right) Observability of key far-IR (dashed lines) lines compared to the ALMA bands (shaded area) in the 1-8 redshift interval. Dots represent current detection for each line.

- Engelbracht C. W., Rieke G. H., Gordon K. D., Smith J. D. T., Werner M. W., Moustakas J., Willmer C. N. A., Vanzi L., 2008, *ApJ*, 678, 804
- Faisst A. L., et al., 2020, *ApJS*, 247, 61
- Farrah D., et al., 2013, *ApJ*, 776, 38
- Fazio G. G., et al., 2004, *ApJS*, 154, 10
- Fernández-Ontiveros J. A., Spinoglio L., Pereira-Santaella M., Malkan M. A., Andreani P., Dasyra K. M., 2016, *ApJS*, 226, 19
- Ferrara A., Vallini L., Pallottini A., Gallerani S., Carniani S., Kohandel M., Decataldo D., Behrens C., 2019, *MNRAS*, 489, 1
- Förster Schreiber N. M., Wuyts S., 2020, *ARA&A*, 58, 661
- Galliano F., et al., 2021, arXiv e-prints, p. arXiv:2101.00456
- Gardner J. P., et al., 2006, *Space Sci. Rev.*, 123, 485
- Gilmozzi R., Spyromilio J., 2007, *The Messenger*, 127, 11
- Goldsmith P. F., Langer W. D., Pineda J. L., Velusamy T., 2012, *ApJS*, 203, 13
- Goulding A. D., Alexander D. M., 2009, *MNRAS*, 398, 1165
- Greenhouse M. A., Feldman U., Smith H. A., Klapisch M., Bhatia A. K., Bar-Shalom A., 1993, *ApJS*, 88, 23
- Gruppioni C., et al., 2016, *MNRAS*, 458, 4297
- Harikane Y., et al., 2019, arXiv e-prints, p. arXiv:1910.10927
- Hashimoto T., 2019, in *American Astronomical Society Meeting Abstracts #233*, p. 106.05
- Hashimoto T., Inoue A. K., Tamura Y., Matsuo H., Mawatari K., Yamaguchi Y., 2018, arXiv e-prints, p. arXiv:1811.00030
- Herrera-Camus R., et al., 2015, *ApJ*, 800, 1
- Ho L. C., Keto E., 2007, *ApJ*, 658, 314
- Hollenbach D., McKee C. F., 1989, *ApJ*, 342, 306
- Houck J. R., et al., 2004, *ApJS*, 154, 18
- Hunter D. A., et al., 2001, *ApJ*, 553, 121
- Inami H., et al., 2013, *ApJ*, 777, 156
- Inoue A. K., et al., 2016, *Science*, 352, 1559
- Johns M., et al., 2012, in *Steff L. M., Gilmozzi R., Hall H. J., eds, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 8444, Ground-based and Airborne Telescopes IV*, p. 84441H, doi:10.1117/12.926716
- Juarez Y., Maiolino R., Mujica R., Pedani M., Marinoni S., Nagao T., Marconi A., Oliva E., 2009, *A&A*, 494, L25
- Kennicutt Robert C. J., 1998, *ARA&A*, 36, 189
- Kirkpatrick A., Pope A., Sajina A., Roebuck E., Yan L., Armus L., Díaz-Santos T., Stierwalt S., 2015, *ApJ*, 814, 9
- Lacy M., et al., 2013, *ApJS*, 208, 24
- Launay J. M., Roueff E., 1977, *A&A*, 56, 289
- Lebouteiller V., Barry D. J., Goes C., Sloan G. C., Spoon H. W. W., Weedman D. W., Bernard-Salas J., Houck J. R., 2015, *ApJS*, 218, 21
- Lee J. C., Hwang H. S., Ko J., 2013, *ApJ*, 774, 62
- Leung T. K. D., Olsen K. P., Somerville R. S., Davé R., Greve T. R., Hayward C. C., Narayanan D., Popping G., 2020, *ApJ*, 905, 102
- Lusso E., et al., 2012, *MNRAS*, 425, 623
- Lutz D., Genzel R., Kunze D., Spoon H. W. W., Sturm E., Sternberg A., Moorwood A. F. M., 1998, in *Yun J., Liseau L., eds, Astronomical Society of the Pacific Conference Series Vol. 132, Star Formation with the Infrared Space Observatory*, p. 89
- Lyu J., Rieke G. H., 2020, arXiv e-prints, p. arXiv:2011.07638
- Madau P., Dickinson M., 2014, *ARA&A*, 52, 415
- Madden S. C., 2000, *New A Rev.*, 44, 249
- Madden S. C., Poglitsch A., Geis N., Stacey G. J., Townes C. H., 1997, *ApJ*, 483, 200
- Madden S. C., et al., 2013, *PASP*, 125, 600
- Maloney P. R., 1997, *Ap&SS*, 248, 105
- Maloney P. R., Hollenbach D. J., Tielens A. G. G. M., 1996, *ApJ*, 466, 561
- Marconi A., Risaliti G., Gilli R., Hunt L. K., Maiolino R., Salvati M., 2004, *MNRAS*, 351, 169
- Naab T., Ostriker J. P., 2017, *ARA&A*, 55, 59
- Nagao T., Maiolino R., Marconi A., 2006, *A&A*, 447, 863
- Negishi T., Onaka T., Chan K. W., Roellig T. L., 2001, *A&A*, 375, 566
- Oliva E., Marconi A., Moorwood A. F. M., 1999, *A&A*, 342, 87
- Onoue M., et al., 2020, *ApJ*, 898, 105
- Pérez-Torres M., Mattila S., Alonso-Herrero A., Aalto S., Efstathiou A., 2021, *A&A Rev.*, 29, 2
- Perley R. A., Chandler C. J., Butler B. J., Wrobel J. M., 2011, *ApJ*, 739, L1
- Poglitsch A., et al., 2010, *A&A*, 518, L2
- Pope A., et al., 2013, *ApJ*, 772, 92
- Rémy-Ruyer A., et al., 2015, *A&A*, 582, A121
- Riechers D. A., et al., 2014, *ApJ*, 786, 31
- Rieke G. H., et al., 2004, *ApJS*, 154, 25
- Rieke G. H., et al., 2015, *PASP*, 127, 584
- Rigopoulou D., Kunze D., Lutz D., Genzel R., Moorwood A. F. M., 2002, *A&A*, 389, 374
- Roelfsema P. R., et al., 2018, *PASA*, 35, e030
- Röllig M., Ossenkopf V., Jeyakumar S., Stutzki J., Sternberg A., 2006, *A&A*, 451, 917
- Rush B., Malkan M. A., Spinoglio L., 1993, *ApJS*, 89, 1
- Rybak M., Zavala J. A., Hodge J. A., Casey C. M., van der Werf P., 2019, arXiv e-prints, p. arXiv:1912.07652
- Sajina A., Yan L., Fadda D., Dasyra K., Huynh M., 2012, *ApJ*, 757, 13
- Sanders D. B., Mirabel I. F., 1996, *ARA&A*, 34, 749
- Sanders R. L., et al., 2020, arXiv e-prints, p. arXiv:2009.07292
- Sargsyan L., et al., 2012, *ApJ*, 755, 171
- Satyapal S., Kamal L., Cann J. M., Secrest N. J., Abel N. P., 2021, *ApJ*, 906, 35
- Schaefer D., et al., 2020, *A&A*, 643, A3
- Schöck M., et al., 2009, *PASP*, 121, 384
- Shipley H. V., Papovich C., Rieke G. H., Dey A., Jannuzi B. T., Moustakas J., Weiner B., 2013, *ApJ*, 769, 75
- Shipley H. V., Papovich C., Rieke G. H., Brown M. J. I., Moustakas J., 2016, *ApJ*, 818, 60
- Smith J. D. T., et al., 2007, *ApJ*, 656, 770
- Somerville R. S., Davé R., 2015, *ARA&A*, 53, 51

- Spinoglio L., Malkan M. A., 1992, [ApJ](#), 399, 504
- Spinoglio L., Malkan M. A., Rush B., Carrasco L., Recillas-Cruz E., 1995, [ApJ](#), 453, 616
- Spinoglio L., Dasyra K. M., Franceschini A., Gruppioni C., Valiante E., Isaak K., 2012, [ApJ](#), 745, 171
- Spinoglio L., Dasyra K. M., Franceschini A., Gruppioni C., Valiante E., Isaak K., 2014, [ApJ](#), 791, 138
- Spinoglio L., Pereira-Santaella M., Dasyra K. M., Calzoletti L., Malkan M. A., Tommasin S., Busquet G., 2015, [ApJ](#), 799, 21
- Spinoglio L., et al., 2021, arXiv e-prints, p. arXiv:2103.03584
- Stierwalt S., et al., 2014, [ApJ](#), 790, 124
- Sturm E., Lutz D., Verma A., Netzer H., Sternberg A., Moorwood A. F. M., Oliva E., Genzel R., 2002, [A&A](#), 393, 821
- Sutter J., et al., 2019, [ApJ](#), 886, 60
- Sutter J., et al., 2021, arXiv e-prints, p. arXiv:2102.08865
- Tamura Y., et al., 2019, [ApJ](#), 874, 27
- Tielens A. G. G. M., 2008, [ARA&A](#), 46, 289
- Tielens A. G. G. M., Hollenbach D., 1985, [ApJ](#), 291, 722
- Togi A., Smith J. D. T., 2016, [ApJ](#), 830, 18
- Tommasin S., Spinoglio L., Malkan M. A., Smith H., González-Alfonso E., Charmandaris V., 2008, [ApJ](#), 676, 836
- Tommasin S., Spinoglio L., Malkan M. A., Fazio G., 2010, [ApJ](#), 709, 1257
- Venemans B. P., et al., 2020, [ApJ](#), 904, 130
- Véron-Cetty M. P., Véron P., 2010, [A&A](#), 518, A10
- Vishwas A., et al., 2018, [ApJ](#), 856, 174
- Walter F., et al., 2018, [ApJ](#), 869, L22
- Wechsler R. H., Tinker J. L., 2018, [ARA&A](#), 56, 435
- Wooten A., Thompson A. R., 2009, [IEEE Proceedings](#), 97, 1463
- Wright G. S., et al., 2015, [PASP](#), 127, 595
- Wu H., Cao C., Hao C.-N., Liu F.-S., Wang J.-L., Xia X.-Y., Deng Z.-G., Young C. K.-S., 2005, [ApJ](#), 632, L79
- Wu Y., Charmandaris V., Hao L., Brandl B. R., Bernard-Salas J., Spoon H. W. W., Houck J. R., 2006, [ApJ](#), 639, 157
- Wu Y., Charmandaris V., Huang J., Spinoglio L., Tommasin S., 2009, [ApJ](#), 701, 658
- Xie Y., Ho L. C., 2019, [ApJ](#), 884, 136
- Yun M. S., Reddy N. A., Condon J. J., 2001, [ApJ](#), 554, 803
- Zhuang M.-Y., Ho L. C., Shanguan J., 2019, [ApJ](#), 873, 103
- de Graauw T., et al., 1996, [A&A](#), 315, L49

Appendix A: Characteristics of IR fine-structure lines

Table A.1. Fine-structure lines in the mid- to far-IR range. The columns correspond to the central wavelength, frequency, ionisation potential, excitation temperature, critical density, spectral and spatial resolution of the data presented in this work, and the number of AGN, starburst, and dwarfs galaxies with line detections above $3\times\text{RMS}$. Critical densities and excitation temperatures from: Launay & Roueff (1977); Tielens & Hollenbach (1985); Greenhouse et al. (1993); Sturm et al. (2002); Cormier et al. (2012); Goldsmith et al. (2012); Farrah et al. (2013).

Line	λ (μm)	ν (GHz)	I.P. (eV)	E (K)	n_{cr} (cm^{-3})	Spec. Res. (km s^{-1})	Ang. Res. (arcsec)
[Ne VI] $^2\text{P}_{3/2}-^2\text{P}_{1/2}$	7.65	39188.56	126.21	1888	2.5×10^5	~ 1500	14×20
H ₂ (0,0) S(3)	9.66	31034.41	4.48	2504	$9 \times 10^{5\text{a}}$	~ 500	4.7×11.3
[S IV] $^2\text{P}_{3/2}-^2\text{P}_{1/2}$	10.51	28524.50	34.79	1369	5.39×10^4	~ 500	4.7×11.3
H ₂ (0,0) S(2)	12.28	24413.07	4.48	1682	$2 \times 10^{5\text{a}}$	~ 500	4.7×11.3
[Ne II] $^2\text{P}_{1/2}-^2\text{P}_{3/2}$	12.81	23403.00	21.56	1123	7.00×10^5	~ 500	4.7×11.3
[Ne V] $^3\text{P}_2-^3\text{P}_1$	14.32	20935.23	97.12	1892	3×10^4	~ 500	4.7×11.3
[Ne III] $^3\text{P}_1-^3\text{P}_2$	15.56	19266.87	40.96	925	2.68×10^5	~ 500	4.7×11.3
H ₂ (0,0) S(1)	17.03	17603.78	4.48	1015	$2 \times 10^{4\text{a}}$	~ 500	4.7×11.3
[S III] $^3\text{P}_2-^3\text{P}_1$	18.71	16023.11	23.34	769	2.22×10^4	~ 500	4.7×11.3
[Ne V] $^3\text{P}_1-^3\text{P}_0$	24.32	12326.99	97.12	596	5.0×10^5	~ 500	11.1×22.3
[O IV] $^2\text{P}_{3/2}-^2\text{P}_{1/2}$	25.89	11579.47	54.94	555	10^4	~ 500	11.1×22.3
[Fe II]a $^6\text{D}_{7/2}-\text{a}^6\text{D}_{9/2}$	25.98	11539.35	7.9	553	2.2×10^6	~ 500	11.1×22.3
H ₂ (0,0) S(1)	17.03	17603.78	4.48	1015	$2 \times 10^{4\text{a}}$	~ 500	4.7×11.3
[S III] $^3\text{P}_1-^3\text{P}_0$	33.48	8954.37	23.34	430	7.04×10^3	~ 500	11.1×22.3
[Si II] $^2\text{P}_{3/2}-^2\text{P}_{1/2}$	34.81	8612.25	8.15	413	$3.4 \times 10^{5\text{a}}, 10^3$	~ 500	11.1×22.3
[O III] $^3\text{P}_2-^3\text{P}_1$	51.81	5787.57	35.12	441	3.6×10^3	~ 105	9.4 ^c
[N III] $^2\text{P}_{3/2}-^2\text{P}_{1/2}$	57.32	5230.43	29.60	251	3.0×10^3	~ 105	9.4
[O I] $^3\text{P}_2-^3\text{P}_1$	63.18	4744.77	–	228	$4.7 \times 10^{5\text{a}}$	~ 86	9.4
[O III] $^3\text{P}_1-^3\text{P}_0$	88.36	3393.01	35.12	163	510	~ 124	9.4
[N II] $^3\text{P}_2-^3\text{P}_1$	121.90	2459.38	14.53	118	310	~ 290	9.4
[O I] $^3\text{P}_1-^3\text{P}_0$	145.52	2060.07	–	98	$9.5 \times 10^{4\text{a}}$	~ 256	10.3
[C II] $^2\text{P}_{3/2}-^2\text{P}_{1/2}$	157.74	1900.54	11.26	91	$20, [2.2^{\text{a}}, 4.4^{\text{b}}] \times 10^3$	~ 238	11.2
[N II] $^3\text{P}_1-^3\text{P}_0$	205.3	1460.27	14.53	70	48	~ 297	~ 16.8

^a Critical density for collisions with hydrogen atoms.

^b Critical density for collisions with H₂ molecules.

^c The beam size for *Herschel*/PACS is dominated by the spaxel size ($9''.4$) below $\sim 120 \mu\text{m}$.

Appendix B: Line correlations




Fig. B.1. (a: left) The luminosity of the PAH feature at 6.2 μm as a function of the total IR luminosity. **(b: centre)** The [NeVI]7.65 μm line luminosity as a function of the total IR luminosity. **(c: right)** The luminosity of the PAH feature at 7.7 μm as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.2. (a: left) The luminosity of the PAH feature at 8.6 μm as a function of the total IR luminosity. **(b: centre)** The luminosity of the H_2 molecular line at 9.7 μm as a function of the total IR luminosity. **(c: right)** The [SIV]10.5 μm line luminosity as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.3. (a: left) The luminosity of the H_2 molecular line at 12.28 μm as a function of the total IR luminosity. **(b: centre)** The [NeV]14.3 μm line luminosity as a function of the total IR luminosity. **(c: right)** The luminosity of the PAH feature at 17 μm as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.4. (a: left) The luminosity of the H_2 molecular line at $17.03 \mu m$ as a function of the total IR luminosity. (b: centre) The $[SIII]18.7 \mu m$ line luminosity as a function of the total IR luminosity. (c: right) The $[FeIII]22.93 \mu m$ line luminosity as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.5. (a: left) The $[NeV]24.3 \mu m$ line luminosity as a function of the total IR luminosity. (b: centre) The $[FeII]25.99 \mu m$ line luminosity as a function of the total IR luminosity. (c: right) The luminosity of the H_2 molecular line at $28.22 \mu m$ as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.6. (a: left) The $[SIII]33.5 \mu m$ line luminosity as a function of the total IR luminosity. (b: centre) The $[SiII]34.8 \mu m$ line luminosity as a function of the total IR luminosity. (c: right) The $[OIII]52 \mu m$ line luminosity as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.7. (a: left) The [NIII]57 μm line luminosity as a function of the total IR luminosity. (b: centre) The [OI]63.18 μm line luminosity as a function of the total IR luminosity. (c: right) The [NII]122 μm line luminosity as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.8. (a: left) The [OI]145 μm line luminosity as a function of the total IR luminosity. (b: right) The [NII]122 μm line luminosity as a function of the total IR luminosity. The same legend as in Fig. 1 was used.




Fig. B.9. (a: left) Comparison between the local $L_{\text{IR}}-L_{[\text{NII}]122}$ relation for local SFG (black dashed line) and high redshift detections of [NII]122 μm line. The shaded area shows the 95% prediction interval for the local relation. The pink + sign indicates a detection for the [NII]122 μm line (De Breuck et al. 2019), while symbols with arrows indicate upper limits for high redshift sources (Harikane et al. 2019). (b: centre) Comparison between the local $L_{\text{IR}}-L_{[\text{OI}]63}$ relation for local SFG (black dashed line) and high redshift detections of [OI]63 μm line (pink symbol Rybak et al. 2019). (c: right) Comparison between the local $L_{\text{IR}}-L_{[\text{OI}]145}$ relation for local SFG (black dashed line) and high redshift detections of [OI]145 μm line (pink symbol De Breuck et al. 2019). The shaded area shows the 95% prediction interval for the local relation. Grey stars show local SFG, while grey triangles show local LMG.

Appendix C: [NeV]14.3 μm as BHAR tracer and discussion on the use of the mid-ionisation lines

In this section, following what has been done in Section 3.3, we report the results for the [NeV]14.3 μm line as BHAR tracer. In particular, starting from the total sample of Tommasin et al. (2008, 2010), excluding Compton-thick objects we obtain a sample of 38 objects. The correlation between the [NeV]14.3 μm line luminosity and the 2-10 keV X-ray luminosity is shown in Fig C.1a and is expressed by the equation:

$$\log\left(\frac{L_X}{10^{41} \text{ erg s}^{-1}}\right) = (2.58 \pm 0.10) + (1.11 \pm 0.13) \log\left(\frac{L_{[\text{NeV}]14.3}}{10^{41} \text{ erg s}^{-1}}\right) \quad (\text{C.1})$$

with a Pearson r coefficient of $r = 0.79$. From this result, we then apply the bolometric corrections by Marconi et al. (2004) and Lusso et al. (2012) to the 2-10 keV X-ray luminosity, in order to obtain the bolometric luminosity of the AGN. From the bolometric luminosities we then calculate the linear correlation linking the [NeV]14.3 μm line luminosity to the BHAR, with $L_{\text{AGN}} = \eta \dot{M}_{\text{BH}} c^2$. Assuming a radiative efficiency of $\eta = 0.1$, we report in equation C.2 the linear correlation applying the bolometric correction from Lusso et al. (2012), and in equation C.3 the one applying the Marconi et al. (2004) bolometric correction, and the relative Pearson r coefficient:

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-1.0 \pm 0.10) + (1.04 \pm 0.13) \log\left(\frac{L_{[\text{NeV}]14.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.76 \quad (\text{C.2})$$

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-0.73 \pm 0.17) + (1.49 \pm 0.13) \log\left(\frac{L_{[\text{NeV}]14.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.78 \quad (\text{C.3})$$

We have then analysed the correlation between the [NeV]14.3 μm line with the luminosity at 19 μm following the work by Tommasin et al. (2010). As in Section 3.3, we select only those sources with an AGN component at 19 μm equal or above 85%, obtaining a subsample of 32 objects. The correlation between the [NeV]14.3 μm line luminosity and the 19 μm luminosity is shown in Fig. C.1b, and expressed by:

$$\log\left(\frac{L_{19}}{10^{41} \text{ erg s}^{-1}}\right) = (3.09 \pm 0.06) + (1.01 \pm 0.10) \log\left(\frac{L_{[\text{NeV}]14.3}}{10^{41} \text{ erg s}^{-1}}\right) \quad (\text{C.4})$$

with a Pearson r coefficient of $r = 0.85$. We then calculate the bolometric luminosity from the monochromatic 12 μm luminosity extracted from Deo et al. (2009) using the relation by Spinoglio et al. (1995). From the bolometric luminosity we determine the BHAR and its relation to the [NeV]14.3 μm line luminosity, obtaining:

$$\log\left(\frac{\dot{M}_{\text{BH}}}{M_{\odot} \text{ yr}^{-1}}\right) = (-0.66 \pm 0.11) + (1.05 \pm 0.19) \log\left(\frac{L_{[\text{NeV}]14.3}}{10^{41} \text{ erg s}^{-1}}\right), \quad r = 0.68 \quad (\text{C.5})$$

for a sample of 26 objects, where r is the Pearson coefficient. The small number of objects used to derive this relation is due to the lack of data for the determination of the continuum at 12 μm .

In Fig. C.1c we compare the three different relation derived to determine the BHAR from the [NeV]14.3 μm line luminosity, as in Section 3.3, also for this line we find good agreement between the results obtained with the Lusso et al. and Spinoglio et al. bolometric corrections, while the Marconi et al. bolometric correction produces a significantly steeper relationship.




Fig. C.1. (a: left) Linear correlation between the [NeV]14.3 μm and the 2-10 keV X-ray luminosity. Blue squares show the AGN sample, and the shaded area shows the 95% confidence interval for the relation. **(b: centre)** Linear correlation between the [NeV]14.3 μm line luminosity and the 19 μm luminosity. **(c: right)** Comparison between the [NeV]14.3 μm line luminosity and the BHAR derived from the bolometric correction by Spinoglio et al. (green solid line), Marconi et al. (red dotted line) and Lusso et al. (blue dashed line).

As presented in Section 3.3 and here, high ionisation lines, with an ionisation potential above the threshold of doubly ionised helium (54.4 eV) can only be efficiently produced by AGN activity, and thus are optimal BHAR tracers. Mid-ionisation lines, however, are partially affected by AGN activity, and could in theory be used as tracers for the BHAR. For this reason, we have tested two mid-ionisation lines, namely the [SIV] 10.5 μ m line, and the [NeIII]15.6 μ m line, to evaluate their use as tracers for the X-ray luminosity in the 2-10 keV interval. Starting from the sample of Tommasin et al. (2008, 2010), as in Sect. 3.3, we exclude all Compton-thick objects and obtain a sub-sample of 44 objects for the [NeIII] line, and of 39 objects for the [SIV] line. Fig. C.2a shows the correlation between the [NeIII] line luminosity and the 2-10 keV X-ray luminosity, expressed as:

$$\log\left(\frac{L_X}{10^{41} \text{ erg s}^{-1}}\right) = (2.20 \pm 0.10) + (1.33 \pm 0.13) \log\left(\frac{L_{[\text{NeIII}]15.6}}{10^{41} \text{ erg s}^{-1}}\right) \quad (\text{C.6})$$

The correlation of the [SIV]10.5 μ m line with the 2-10 keV X-ray luminosity, shown in Fig. C.2b, can be expressed as:

$$\log\left(\frac{L_X}{10^{41} \text{ erg s}^{-1}}\right) = (2.51 \pm 0.09) + (0.98 \pm 0.10) \log\left(\frac{L_{[\text{SIV}]10.5}}{10^{41} \text{ erg s}^{-1}}\right) \quad (\text{C.7})$$

For both correlations, we find a low Pearson correlation coefficient, equal to $r = 0.63$, lower than the values found for [OIV] ($r = 0.87$) and the [NeV] lines ($r = 0.79$ and $r = 0.84$, for the 14.3 μ m and 24.3 μ m lines, respectively). This suggests that, while mid-ionisation lines can correlate with the X-ray luminosity, they may suffer from contributions due to star formation activity, and thus are far less reliable as BHAR proxies. In particular, AGN sources in low-metallicity star-forming galaxies would have a significant contamination of the [SIV] and [NeIII] lines from the stellar population, as shown by the LMG in our sample (Figs. 1c and B.2c), in contrast with the [OIV] and [NeV] lines.




Fig. C.2. (a: left) Linear correlation between the [NeIII]15.6 μ m and the 2-10 keV X-ray luminosity. Blue squares show the AGN sample, and the shaded area shows the 95% confidence interval for the relation. **(b: right)** Linear correlation between the [SIV]10.5 μ m and the 2-10 keV X-ray luminosity

Appendix D: Line and PAH calibrations

Table D.1. New calibration obtained for fine-structure lines and PAH bands. For each class of objects, are reported the slope (a) and intercept (b) of the linear correlation with relative errors, the number of data from which the linear correlation was calculated (N) and the best-fit coefficients of determination (r).

Line/feature	AGN				SFG				LMG			
	$a \pm \delta a$	$b \pm \delta b$	N	r	$a \pm \delta a$	$b \pm \delta b$	N	r	$a \pm \delta a$	$b \pm \delta b$	N	r
PAH 6.2	1.04±0.07	-2.73±0.24	65	0.87	1.15±0.06	-2.99±0.24	150	0.81	–	–	–	–
[NeVI]7.65	1.22±0.24	-4.12±0.82	8	0.89	–	–	–	–	–	–	–	–
PAH 7.7	–	–	–	–	1.29±0.07	-2.98±0.27	150	0.81	–	–	–	–
PAH 8.6	–	–	–	–	1.43±0.23	-4.25±0.67	16	0.84	–	–	–	–
H ₂ 9.67	1.20±0.28	-4.97±1.17	9	0.82	1.39±0.06	-5.43±0.26	137	0.87	–	–	–	–
[SIV]10.51	1.37±0.11	-5.19±0.38	75	0.80	1.58±0.24	-7.00±0.94	51	0.59	1.03±0.08	-3.08±0.16	29	0.91
PAH 11.3	1.04±0.06	-2.82±0.20	90	0.88	1.11±0.06	-2.79±0.23	150	0.81	–	–	–	–
H ₂ 12.28	1.25 ± 0.10	-5.41 ± 0.37	31	0.91	1.18 ± 0.05	-4.79± 0.19	114	0.91	–	–	–	–
[NeII]12.81	1.19±0.07	-4.23±0.26	86	0.85	1.04±0.43	-3.48±0.17	186	0.84	1.37±0.10	-4.36±0.20	21	0.95
[NeV]14.32	1.32±0.11	-5.01±0.39	74	0.78	–	–	–	–	–	–	–	–
[NeIII]15.56	1.44 ± 0.10	-5.16±0.36	88	0.80	1.13±0.06	-4.60±0.21	182	0.80	1.13±0.06	-3.11±0.12	28	0.96
PAH 17	–	–	–	–	1.14±0.06	-3.08±0.23	131	0.84	–	–	–	–
H ₂ 17.03	1.14±0.06	-4.64±0.19	76	0.91	1.22±0.05	-4.58±0.20	135	0.89	–	–	–	–
[SIII]18.71	1.22±0.09	-4.62±0.30	70	0.83	1.26±0.11	-4.79±0.42	140	0.60	1.14±0.06	-3.42±0.12	25	0.96
[FeIII]22.93	–	–	–	–	1.33±0.13	-5.79±0.44	14	0.64	–	–	–	–
[NeV]24.32	1.27±0.11	-4.73±0.37	65	0.80	–	–	–	–	–	–	–	–
[OIV]25.89	1.27±0.14	-4.13±0.48	34	0.82	1.15±0.05	-5.00±0.20	144	0.86	0.88±0.13	-3.71±0.27	16	0.86
[FeII]25.98	–	–	–	–	0.80±0.18	-3.99±0.51	35	0.44	–	–	–	–
H ₂ 28.22	1.04±0.14	-4.44 ± 0.45	25	0.81	1.65±0.24	-6.51±0.82	11	0.91	–	–	–	–
[SIII]33.48	1.10±0.07	-3.85±0.23	75	0.87	1.13±0.05	-3.94±0.20	170	0.83	1.11±0.08	-3.24±0.17	20	0.96
[SiII]34.81	1.21±0.08	-3.99±0.28	73	0.84	1.15±0.04	-3.72 ± 0.15	167	0.91	1.05±0.08	-3.14±0.18	17	0.95
[OIII]51.81	0.92±0.13	-3.10±0.54	19	0.85	–	–	–	–	–	–	–	–
[NIII]57.32	0.98±0.06	-3.96±0.24	31	0.94	–	–	–	–	–	–	–	–
[OI]63.18	1.07±0.04	-3.56±0.17	106	0.91	1.18±0.05	-3.86±0.21	183	0.83	1.07±0.06	-3.00±0.11	31	0.96
[OIII]88.36	1.18±0.06	-4.34±0.22	81	0.91	1.20±0.06	-4.27±0.25	117	0.85	1.05±0.06	-2.53±0.10	37	0.95
[NII]121.9	1.08±0.06	-4.43±0.22	77	0.89	0.90±0.06	-3.39±0.22	75	0.86	0.96±0.09	-4.17±0.23	8	0.97
[OI]145.5	1.06±0.07	-4.53± 0.26	64	0.88	1.03±0.05	-4.14±0.16	13	0.99	1.05±0.07	-4.14± 0.18	12	0.97
[CII]157.7	1.00±0.04	-3.18± 0.14	149	0.90	0.97±0.05	-2.76±0.20	183	0.78	1.09±0.06	-2.88±0.10	40	0.95
[NII]205	0.88±0.05	-3.88± 0.18	60	0.92	1.02±0.07	-4.36±0.23	13	0.97	–	–	–	–

Appendix E: Differences with previous results

We analyse here the differences of the line calibrations obtained in this work with respect to those of Spinoglio et al. (2012, 2014) and Gruppioni et al. (2016).

When considering the results by Spinoglio et al., we only analyse the fine structure lines in the 50-160 μ m spectral interval, for which these authors consider an heterogeneous sample of both AGN and SFG, while in this work we consider three classes of galaxies. When comparing the results, we consider the calibrations obtained by Spinoglio et al. and compare them to our results for AGN and SFG. For this comparison, we apply to our samples the ordinary least square method, which was used in Spinoglio et al. to derive their correlations. Fig E.1 shows the comparison for the [OIII]52 μ m, [NIII]57 μ m and [OI]63 μ m lines, Fig E.2 shows the comparison for the [OIII]88 μ m, [NII]122 μ m and [OI]145 μ m lines, and Fig E.3 shows the comparison for the [CII]158 μ m line.

When compared to our AGN sample, the results are consistent within the errors, except for the [NII]57 μ m and [OIII]88 μ m lines, where the slopes are comparable within 2σ of each other. If compared to the SFG results, we obtain results comparable within the errors except for [OIII]88 μ m and [NII]122 μ m lines, for which the slopes are comparable within 2σ of each other.




Fig. E.1. (a: left) The [OIII]52 μ m line luminosity versus the total IR luminosity. **(b: centre)** The [NIII]57 μ m line luminosity versus the total IR luminosity. **(c: right)** The [OI]63 μ m line luminosity versus the total IR luminosity. In the figures, Blue squares represent AGN while red stars show the SFG sample. The dashed blue line represents the linear fit, the dot-dashed red line the relation for SFG, and the black solid line the relation obtained by Spinoglio et al. (2012, 2014).




Fig. E.2. (a: left) The [OIII]88 μ m line luminosity versus the total IR luminosity. **(b: centre)** The [NII]122 μ m line luminosity versus the total IR luminosity. **(c: right)** The [OI]145 μ m line luminosity versus the total IR luminosity. The same legend as in Fig. E.1 was used.

When considering the results presented by Gruppioni et al. (2016), we take advantage of the published catalogue and apply the orthogonal distance regression to the sample presented by the authors. Gruppioni et al. present a sample of 76 AGN, divided in two sub-samples depending on the fraction of 5-40 μ m luminosity produced by the active nucleus [$f_{\text{AGN}}(5-40\mu\text{m})$]. In particular, there are 30 objects with $f_{\text{AGN}}(5-40\mu\text{m}) \leq 0.4$ and 43 objects with $f_{\text{AGN}}(5-40\mu\text{m}) > 0.4$.

Figs. E.4-E.7 show the comparison for the relations calculated for the $f_{\text{AGN}}(5-40\mu\text{m}) > 0.4$ sub-sample, and Figs. E.8-E.11 show the comparison for the relations calculated for the $f_{\text{AGN}}(5-40\mu\text{m}) \leq 0.4$ sub-sample.

As a general trend, our sample of AGN and the sample by Gruppioni et al. occupy the same region in the $L_{\text{IR}}-L_{\text{line}}$ space. We note here that, while the data to calculate the correlations were available in the literature, Gruppioni et al. only present the correlations




Fig. E.3. (a): The [CII]158 μ m line luminosity versus the total IR luminosity. The same legend as in Fig. E.1 was used

for [NeV]14.3, 24.3 μ m, [NeIII]15.6 μ m and [OIV]25.9 μ m for the $f_{\text{AGN}}(5-40\mu\text{m}) > 0.4$ sub-sample. For these relations, we find that our results show steeper slopes, but consistent within 3σ of each other.

For the $f_{\text{AGN}}(5-40\mu\text{m}) \leq 0.4$ sub-sample, there is a better agreement between the relations, which are comparable within the errors. A significant difference in slope is present for the [NeV]24.3 μ m line, for which the relations are comparable within 3σ , and for the [SIV]10.5 μ m line, comparable within 2σ .

If we compare the results obtained using the Gruppioni et al. sample with the results we obtain using the SFG sample, we find comparable results only for the PAH features for the $f_{\text{AGN}}(5-40\mu\text{m}) > 0.4$, and for all lines in the $f_{\text{AGN}}(5-40\mu\text{m}) \leq 0.4$ sub-sample, excluding the [SIV]10.5 μ m and [OIV]25.9 μ m lines.




Fig. E.4. (a: left) The PAH 6.2 μ m luminosity versus the total IR luminosity. Blue squares represent AGN, red stars SFG and black circles are the sample by Gruppioni et al. (2016). The dashed blue line represents the linear relation for AGN, the dot-dashed red line the relation for SFG, and the black solid line the relation for the Gruppioni et al. sample. **(b: centre)** The [SIV]10.5 μ m luminosity versus the total IR luminosity. **(c: right)** The PAH 11.3 μ m luminosity versus the total IR luminosity. In all three figures, the Gruppioni et al. sample was limited to AGN fractions > 0.4 in the 5-40 μ m spectral interval (see Sect.4.3).




Fig. E.5. (a: left) The [NeII] 12.8 μm line luminosity versus the total IR luminosity. (b: centre) The [NeV] 14.3 μm line versus the total IR luminosity. (c: right) The [NeIII] 15.6 μm line luminosity versus the total IR luminosity. In all three figures, the Gruppioni et al. sample was limited to AGN fractions > 0.4 in the 5-40 μm spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.




Fig. E.6. (a: left) The [SIII] 18.7 μm line luminosity versus the total IR luminosity. (b: centre) The [NeV] 24.3 μm line luminosity versus the total IR luminosity. (c: right) The [OIV] 55.9 μm line luminosity versus the total IR luminosity. In all three figures, the Gruppioni et al. sample was limited to AGN fractions > 0.4 in the 5-40 μm spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.




Fig. E.7. (a: left) The [SIII] 33.5 μm line luminosity versus the total IR luminosity. (b: right) The [SiII] 34.9 μm line luminosity versus the total IR luminosity. In both figures, the Gruppioni et al. sample was limited to AGN fractions > 0.4 in the 5-40 μm spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.




Fig. E.8. (a: left) The luminosity of the PAH feature at 6.2 μ m versus the total IR luminosity. (b: centre) The [SIV]10.5 μ m line luminosity as a function of the total IR luminosity. (c: right) The PAH 11.3 μ m luminosity versus the total IR luminosity. In all figures, the Gruppioni et al. sample was limited to AGN fractions < 0.4 in the 5-40 μ m spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.




Fig. E.9. (a: left) The [NeII]12.8 μ m line luminosity versus the total IR luminosity. (b: centre) The [NeV]14.3 μ m line luminosity versus the total IR luminosity. (c: right) The [NeIII]15.6 μ m line luminosity versus the total IR luminosity. In all three figures, the Gruppioni et al. sample was limited to AGN fractions < 0.4 in the 5-40 μ m spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.




Fig. E.10. (a: left) The [SIII]18.7 μ m line luminosity versus the total IR luminosity. (b: centre) The [NeV]24.3 μ m line luminosity versus the total IR luminosity. (c: right) The [OIV]25.9 μ m line luminosity versus the total IR luminosity. In all three figures, the Gruppioni et al. sample was limited to AGN fractions < 0.4 in the 5-40 μ m spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.




Fig. E.11. (a: left) The [SiII]33.5 μ m line luminosity versus the total IR luminosity. **(b: right)** The [SiII]34.9 μ m line luminosity versus the total IR luminosity. In both figures, the Gruppioni et al. sample was limited to AGN fractions < 0.4 in the 5-40 μ m spectral interval (see Sect.4.3). The same legend as in Fig. E.4 was used.