

A 3-approximation list scheduling algorithm for a single-machine scheduling problem with a non-renewable resource and total weighted completion time criterion *

Susumu Hashimoto [†] Shinji Mizuno [‡]

May 3, 2021

Abstract

This paper studies a single-machine scheduling problem with a non-renewable resource (NR-SSP) and total weighted completion time criterion. The non-renewable resource is consumed when the machine starts processing a job. We consider the case where each job's weight in the objective function is proportional to its resource consumption amount. The problem is known to be NP-hard in this case. We propose a 3-approximation list scheduling algorithm for this problem. Besides, we show that the approximation ratio 3 is tight for the algorithm.

1 Introduction

This paper proposes a 3-approximation list scheduling algorithm for a single-machine scheduling problem with a non-renewable resource (NR-SSP). The non-renewable resource is consumed when the machine starts processing a job. The resource is provided at fixed times and in fixed amounts. We consider the case where each job's weight in the objective function is proportional to its resource consumption amount. Such a condition appears in some production systems with a single product. This problem is more general than NR-SSPs studied by Györgyi and Kis (2019), Györgyi and Kis (2020), and Bérczi et al. (2020). Besides, no constant-factor approximation algorithm is known for the problem to our knowledge.

*This research is supported in part by Grant-in-Aid for Science Research (A) 19H00808 of Japan Society for the Promotion of Science.

[†]Department of Industrial Engineering and Economics, Tokyo Institute of Technology, Meguro-ku, Tokyo, Japan. Email:hashimoto.s.ak@m.titech.ac.jp

[‡]Department of Industrial Engineering and Economics, Tokyo Institute of Technology, Meguro-ku, Tokyo, Japan. Email:mizuno.s.ab@m.titech.ac.jp

NR-SSP is the problem of deciding start (or completion) times of n jobs on a machine under given q times resource supplies. We denote the jobs by $\mathcal{J} = (J_1, J_2, \dots, J_n)$. Each job J_j has a processing time $p_j > 0$, a weight $w_j > 0$, and a resource requirement $a_j > 0$. Each provision plan has a supply time $u_i \geq 0$ and a supply amount $b_i > 0$. The machine cannot process multiple jobs simultaneously, and it cannot preempt any process. Besides, each job J_j consumes a_j resources to start processing. The machine cannot start processing a job if the resources are not enough. This condition is expressed as follows:

$$\sum_{j: C_j - p_j \leq T} a_j \leq \sum_{i: u_i \leq T} b_i \text{ for any } T \geq 0, \quad (1)$$

where C_1, C_2, \dots, C_n indicate completion times of jobs J_1, J_2, \dots, J_n . The objective function is total weighted completion time, which is expressed as $\sum_{j=1}^n w_j C_j$. We introduce the triplet notation $\alpha|\beta|\gamma$ for scheduling problems by Graham et al. (1979), where α stands for machine environments, β represents other conditions, and γ is the objective function. The problem mentioned above is expressed as $1|nr = 1, a_j = w_j|\sum w_j C_j$ by using this notation, where $\alpha = 1$ means 1-machine problem, nr represents the number of sorts of non-renewable resources, $a_j = w_j$ indicates $a_j = w_j$ for all $j \in \{1, 2, \dots, n\}$, and $\sum w_j C_j$ stands for that the objective function is $\sum_{j=1}^n w_j C_j$.

Next, we summarize known research results for special cases of $1|nr = 1, a_j = w_j|\sum w_j C_j$. Bérczi et al. (2020) prove that $1|nr = 1, a_j = 1|\sum C_j$ is strongly NP-hard problem. Hence $1|nr = 1, a_j = w_j|\sum w_j C_j$ is also strongly NP-hard problem. Although there are no explicit theoretical results about approximation algorithms for this problem, results for more limited problems are known. Bérczi et al. (2020) propose $\frac{3}{2}$ -approximation algorithm for $1|nr = 1, a_j = 1|\sum C_j$. Györgyi and Kis (2019) prove that $1|nr = 1, p_j = 1, w_j = a_j, q = 2|\sum w_j C_j$ is weakly NP-hard problem. Moreover, they give a 2-approximation list scheduling algorithm for $1|nr = 1, w_j = p_j = a_j|\sum w_j C_j$. Györgyi and Kis (2020) propose a 2-approximation list scheduling algorithm for $1|nr = 1, p_j = 1, w_j = a_j, q = 2|\sum w_j C_j$, and they show that the algorithm is a 3-approximation algorithm for $1|nr = 1, p_j = 1, w_j = a_j|\sum w_j C_j$. They also conjecture that the approximation ratio can be reduced to 2. Hashimoto and Mizuno (2021) prove that the list scheduling algorithm by Györgyi and Kis (2020) is actually a 2-approximation algorithm for $1|nr = 1, p_j = 1, w_j = a_j|\sum w_j C_j$.

For other NR-SSP problems, we mention some complexity and (or) approximation results. Grigoriev et al. (2005) show that $1|nr = k|C_{\max}$ is strongly NP-hard for any integer k , where C_{\max} represents the maximum completion time or makespan. Besides, they prove that any list scheduling algorithm is a 2-approximation algorithm for the problem. Gafarov et al. (2011) prove that $1|nr = 1|\sum C_j, 1|nr = 1|\sum U_j$, and $1|nr = 1|L_{\max}$ are strongly NP-hard, where $\sum U_j$ stands for the number of late jobs and L_{\max} indicates the maximum lateness. Györgyi and Kis (2015) propose PTASs for $1|nr = \text{const.}, q = \text{const.}|C_{\max}$ and $1|nr = 1, a_j = p_j, r_j|C_{\max}$, where r_j means that the problem has release dates. Kis (2015) proves that $1|nr = 1, q = 2|\sum w_j C_j$ is NP-hard,

and establishes an FPTAS for the problem. Bérczi et al. (2020) establish a 6-approximation algorithm and a PTAS for $1|nr = 1, p_j = 0| \sum w_j C_j$.

In addition to NR-SSP, scheduling problems with non-renewable resource(s) have also been studied. Carlier and Rinnooy Kan (1982) develop an exact polynomial-time algorithm for a precedence constrained scheduling problem with a non-renewable resource. Slowiński (1984) gives an exact algorithm for a preemptive parallel machine scheduling problem with a non-renewable resource. Briskorn et al. (2013) propose exact algorithms for a single-machine inventory constrained scheduling problem with total weighted completion time criterion. Morsy and Pesch (2015) give 2-approximation algorithms for limited cases of the above inventory constrained scheduling problem.

2 Algorithms for NR-SSP

We denote any instance of a single-machine scheduling problem with a non-renewable resource (NR-SSP) by $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{w}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$, where n is the number of jobs, $\mathcal{J} = (J_1, J_2, \dots, J_n)$ is a sequence of jobs, $\mathbf{p} = (p_1, p_2, \dots, p_n)^\top$ is a (column) vector of processing times, $\mathbf{w} = (w_1, w_2, \dots, w_n)^\top$ is a vector of weights, $\mathbf{a} = (a_1, a_2, \dots, a_n)^\top$ is a vector of resource requirements, q is the number of supply times, $\mathbf{b} = (b_1, b_2, \dots, b_q)^\top$ is a vector of supply amounts, and $\mathbf{u} = (u_1, u_2, \dots, u_q)^\top$ is a vector of supply times, where $(\cdot)^\top$ denotes transpose. In this paper, we deal with the case where $\mathbf{w} = \mathbf{a}$. Then the instance is simply denoted by $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$. A schedule for S is denoted by $\mathbf{C} = (C_1, C_2, \dots, C_n)^\top$ which is a vector of completion times of the jobs. The objective function of the schedule is

$$f_S(\mathbf{C}) = \sum_{j=1}^n w_j C_j = \mathbf{w}^\top \mathbf{C} = \mathbf{a}^\top \mathbf{C}.$$

The problem NR-SSP is to find a feasible schedule which minimize this objective function. Throughout the paper, we assume that there are no resources other than \mathbf{b} ,

$$n \geq 1, \mathbf{p} > \mathbf{0}_n, \mathbf{w} = \mathbf{a} > \mathbf{0}_n, q \geq 1, \mathbf{b} > \mathbf{0}_q, u_1 \geq 0,$$

and

$$u_{j+1} > u_j \quad \text{for any } j \in \{1, 2, \dots, q-1\},$$

where $\mathbf{0}_n$ denotes the n -dimensional zero vector.

Let $N = \{1, 2, \dots, n\}$ and $\mathbf{r} = (r_1, r_2, \dots, r_n)^\top$ be the vector of $r_i = a_i/p_i$ for $i \in N$. We denote an order of jobs $J_i, i \in N$ by using a permutation \mathbf{o} of $(1, 2, \dots, n)$. Let \mathcal{O} be the set of all the permutations of $(1, 2, \dots, n)$. Then we define a subset of \mathcal{O} for two vectors \mathbf{a} and \mathbf{p} in any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ as follows.

Definition 1. For \mathbf{a} and \mathbf{p} in any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$, we define a set $O(\mathbf{a}, \mathbf{p}) \subset \mathcal{O}$ of permutations \mathbf{o} satisfying the following conditions:

- (i) $a_{o(n)} = \min_{k \in N} a_k$,
- (ii) for any $j \in \{1, 2, \dots, n-1\}$, if $a_{o(j)} > \sum_{k=j+1}^n a_{o(k)}$, then

$$a_{o(i)} \geq a_{o(j)} \text{ for any } i \in \{1, 2, \dots, j\},$$

- (iii) for any $j \in \{1, 2, \dots, n-1\}$, if $a_{o(j)} \leq \sum_{k=j+1}^n a_{o(k)}$, then

$$r_{o(i)} \geq r_{o(j)} \text{ for any } i \in \{1, 2, \dots, j\}$$

$$\text{such that } a_{o(i)} \leq \sum_{k=j+1}^n a_{o(k)}.$$

Now we propose two algorithms, Algorithm 1 and Algorithm 2. Algorithm 1 outputs a permutation $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$ for inputs \mathbf{a}, \mathbf{p} in $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$. Algorithm 2 is a simple list scheduling algorithm which outputs a vector \mathbf{C} of completion times for inputs $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ and $\mathbf{o} \in \mathcal{O}$.

Algorithm 1 is motivated by the 6-approximation algorithm in Bérczi et al. (2020). Their algorithm determines a job's processing order in inverse. It chooses the least economical ($\arg \min w_j/a_j$) job at each selection except for jobs whose weight is bigger than the total weight of selected jobs. In our algorithm, we employ $\arg \min w_j/p_j$ instead of $\arg \min w_j/a_j$ as the economic indicator.

Algorithm 1 A job-ordering algorithm for NR-SSP

Input: \mathbf{a}, \mathbf{p} in any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ of NR-SSP

Output: $\mathbf{o} = (o(1), o(2), \dots, o(n)) \in \mathcal{O}$

1: **Initialize:**

$$SUMW \leftarrow 0$$

$$N^{REST} \leftarrow N$$

2: **for** $i = n$ to 1 **do**

▷ Inverse order

3: $N' \leftarrow \{j \in N^{REST} \mid a_j \leq SUMW\}$

4: **if** $N' = \emptyset$ **then**

5: $o(i) \leftarrow \arg \min_{j \in N^{REST}} a_j$ (ties are broken arbitrarily)

6: **else**

7: $o(i) \leftarrow \arg \min_{j \in N'} r_j$ (ties are broken arbitrarily)

8: **end if**

9: $SUMW \leftarrow SUMW + a_{o(i)}$

10: $N^{REST} \leftarrow N^{REST} \setminus \{o(i)\}$

11: **end for**

Algorithm 2 A list-scheduling algorithm for NR-SSP

Input: $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ and $\mathbf{o} = (o(1), o(2), \dots, o(n)) \in \mathcal{O}$

Output: $\mathbf{C} = (C_1, C_2, \dots, C_n)^\top$ ▷ a schedule

1: **if** $\sum_{j=1}^n a_j > \sum_{q=1}^m b_q$ **then**
2: **return** INFEASIBLE ▷ The instance is infeasible
3: **end if**
4: $C_{o(1)} \leftarrow \min\{T \mid a_{o(1)} \leq \sum_{i:u_i \leq T} b_i\} + p_{o(1)}$
5: **for** $j = 2$ **to** n **do**
6:

$$C_{o(j)} \leftarrow \min \left\{ T \mid T \geq C_{o(j-1)}, \sum_{i=1}^j a_{o(i)} \leq \sum_{i:u_i \leq T} b_i \right\} + p_{o(j)}.$$

7: **end for**

For any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$, we denote an optimal schedule by $\mathbf{C}^*(S) = (C_1^*(S), C_2^*(S), \dots, C_n^*(S))^\top$ and the output schedule of Algorithm 2 for any $\mathbf{o} \in \mathcal{O}$ by $\mathbf{C}^{\mathbf{o}}(S) = (C_1^{\mathbf{o}}(S), C_2^{\mathbf{o}}(S), \dots, C_n^{\mathbf{o}}(S))^\top$.

3 Main results for Algorithms 1 and 2

In the rest of the paper, we impose the following assumptions for any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ without loss of generality:

Assumption 1. $\sum_{j=1}^n a_j = \sum_{i=1}^q b_i$.

Assumption 2. $\max_{j \in N} \frac{a_j}{p_j} \leq 1$.

Assumption 3. *Jobs are sorted such that $C_1^*(S) < C_2^*(S) < \dots < C_n^*(S)$.*

Since we do not know the optimal schedule $\mathbf{C}^*(S)$, we use Assumption 3 only for simplicity of discussion.

For any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ and any $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, we obtain the following results for the output schedule $\mathbf{C}^{\mathbf{o}}(S)$ of Algorithm 2. The results are proved in the succeeding sections.

Theorem 1. *For any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ of NR-SSP and any permutation $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, $f_S(\mathbf{C}^{\mathbf{o}}(S)) < 3f_S(\mathbf{C}^*(S))$ holds.*

Theorem 2. *For any $\epsilon > 0$, there exists an instance $\tilde{S} = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ of NR-SSP and an $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$ such that $f_S(\mathbf{C}^{\mathbf{o}}(\tilde{S})) \geq (3 - \epsilon)f_S(\mathbf{C}^*(\tilde{S}))$.*

Theorem 1 states that a combination of Algorithms 1 and 2 is a 3-approximation algorithm and Theorem 2 states that the approximation ratio 3 is tight.

The next three theorems are used to prove Theorem 1.

Theorem 3. Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ be any instance of NR-SSP. Define $S' = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C}^*(S) - \mathbf{p}), \mathbf{a} \rangle$. For any $\mathbf{o} \in \mathcal{O}$, $f_{S'}(\mathbf{C}^{\mathbf{o}}(S')) < 3f_{S'}(\mathbf{C}^*(S'))$ implies $f_S(\mathbf{C}^{\mathbf{o}}(S)) < 3f_S(\mathbf{C}^*(S))$.

Theorem 4. Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C}' - \mathbf{p}), \mathbf{a} \rangle$ be any instance of NR-SSP, where $\mathbf{C}' = (C'_1, C'_2, \dots, C'_n)^\top$ such that $C'_1 \geq p_1$ and $C'_{j+1} - C'_j \geq p_{j+1}$ for any $j \in \{1, 2, \dots, n-1\}$. Let $\boldsymbol{\sigma}(\mathbf{p}) = (p_1, p_1 + p_2, \dots, \sum_{j=1}^n p_j)^\top$ and $\bar{S} = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}), \mathbf{a} \rangle$. For any $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, $f_{\bar{S}}(\mathbf{C}^{\mathbf{o}}(\bar{S})) < 3f_{\bar{S}}(\mathbf{C}^*(\bar{S}))$ implies $f_S(\mathbf{C}^{\mathbf{o}}(S)) < 3f_S(\mathbf{C}^*(S))$.

Theorem 5. Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}), \mathbf{a} \rangle$ where $\boldsymbol{\sigma}(\mathbf{p}) = (p_1, p_1 + p_2, \dots, \sum_{j=1}^n p_j)^\top$. Define $S^\dagger = \langle n, \mathcal{J}, \mathbf{a}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{a}) - \mathbf{a}), \mathbf{a} \rangle$. Then $O(\mathbf{a}, \mathbf{p}) \subset O(\mathbf{a}, \mathbf{a})$ and, for any $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, $f_{S^\dagger}(\mathbf{C}^{\mathbf{o}}(S^\dagger)) < 3f_{S^\dagger}(\mathbf{C}^*(S^\dagger))$ implies $f_S(\mathbf{C}^{\mathbf{o}}(S)) < 3f_S(\mathbf{C}^*(S))$.

Theorem 3 states that it suffices to prove Theorem 1 only in the case of $\langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C}^*(S) - \mathbf{p}), \mathbf{a} \rangle$. Besides, Theorem 4 and 5 ensures that we may fix $\mathbf{u} = \boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}$ and $\mathbf{a} = \mathbf{p}$. From these theorems, it is enough to prove Theorem 1 for the special instance $S = \langle n, \mathcal{J}, \mathbf{a}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{a}) - \mathbf{a}), \mathbf{a} \rangle$.

The next two theorems show some properties of any permutation $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$ and we use them in the proof of Theorems 4 and 5. We introduce some notations to state the theorems. For any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ and any permutation $\mathbf{o} \in \mathcal{O}$, we define

$$A_j^* = \sum_{k=j}^n a_k \text{ and } A_j^{\mathbf{o}} = \sum_{k=j}^n a_{\mathbf{o}(k)} \text{ for } j \in N.$$

For simplicity of discussion, let $A_j^* = A_j^{\mathbf{o}} = 0$ for any $j \geq n+1$. Then we define

$$\lambda_j = \min\{k \in N \mid A_k^{\mathbf{o}} \geq A_j^*\} \text{ for any } j \in \{1, 2, \dots, n+1\}.$$

From the definition, we see that

$$A_{\lambda_j+1}^{\mathbf{o}} < A_j^* \leq A_{\lambda_j}^{\mathbf{o}} \text{ for any } j \in N.$$

Theorem 6. Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ be any instance of NR-SSP and $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, then $A_{\lambda_j}^{\mathbf{o}} \leq 2A_j^*$ for any $j \in \{1, 2, \dots, n+1\}$.

Theorem 7. Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ be any instance of NR-SSP and $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$. For any $k, l \in N$,

- (i) if $r_{\mathbf{o}(k)} \geq r_{\mathbf{o}(l)}$, then $a_{\mathbf{o}(l)}(p_{\mathbf{o}(k)} - a_{\mathbf{o}(k)}) \leq a_{\mathbf{o}(k)}(p_{\mathbf{o}(l)} - a_{\mathbf{o}(l)})$,
- (ii) if $k < l$ and $r_{\mathbf{o}(k)} < r_{\mathbf{o}(l)}$, then $a_{\mathbf{o}(k)} \geq a_{\mathbf{o}(l)}$ and $a_{\mathbf{o}(k)} \geq A_{l+1}^{\mathbf{o}}$.

4 Proof of Theorems 6 and 7

In this section, we prove Theorems 6 and 7.

Proof of Theorem 6. If $j = n + 1$, then $A_{\lambda_{n+1}}^o = 2A_{n+1}^* = 0$. Thus we assume that $j \in N$. From the definition of λ_j ,

$$A_{\lambda_j+1}^o < A_j^* = \sum_{k=j}^n a_k. \quad (2)$$

If $a_{o(\lambda_j)} \leq A_{\lambda_j+1}^o$, then $A_{\lambda_j}^o = a_{o(\lambda_j)} + A_{\lambda_j+1}^o < 2A_j^*$; therefore we assume $a_{o(\lambda_j)} > A_{\lambda_j+1}^o$. From Definition 1, we see $a_{o(\lambda_j)} \leq a_{o(k)}$ for any $k \in \{1, 2, \dots, \lambda_j\}$. The inequality (2) implies that $A_{\lambda_j+1}^o$ does not have at least one term $a_{o(k)}$ in A_j^* . Hence we see $a_{o(\lambda_j)} \leq a_{o(k)} \leq A_j^*$. From this inequality and (2), we obtain $A_{\lambda_j}^o < 2A_j^*$. \square

Proof of Theorem 7.

(i) For any $k, l \in N$, if $r_{o(k)} \geq r_{o(l)}$, then we have that

$$\begin{aligned} \frac{a_{o(l)}(p_{o(k)} - a_{o(k)})}{p_{o(l)}p_{o(k)}} &= r_{o(l)}(1 - r_{o(k)}) \\ &\leq r_{o(k)}(1 - r_{o(l)}) \\ &= \frac{a_{o(k)}(p_{o(l)} - a_{o(l)})}{p_{o(l)}p_{o(k)}}. \end{aligned}$$

(ii) Suppose that $k < l$ and $r_{o(k)} < r_{o(l)}$ for $k, l \in N$. If $a_{o(l)} > A_{l+1}^o$, then we have from Definition 1-(ii) that

$$a_{o(k)} \geq a_{o(l)} > A_{l+1}^o.$$

Otherwise, $a_{o(l)} \leq A_{l+1}^o$. If $a_{o(k)} \leq A_{l+1}^o$, we see that $r_{o(k)} \geq r_{o(l)}$ from Definition 1-(iii), which contradicts to the assumption $r_{o(k)} < r_{o(l)}$. Hence we have that

$$a_{o(k)} > A_{l+1}^o \geq a_{o(l)}.$$

\square

5 Proof of Theorems 3 and 4

Note that Theorems 3 and 4 are proved in Hashimoto and Mizuno (2021) for a simple list scheduling algorithm when $p_i = 1 (i \in N)$ and the approximation ratio is 2. Although the proofs of the theorems are similar to those in Hashimoto and Mizuno (2021), we prove them for confirmation. The next three lemmas are used in the proof of Theorem 3.

Lemma 1. (*Lemma 1 in Hashimoto and Mizuno (2021)*) Let $\mathbf{a} = (a_1, a_2, \dots, a_n)^\top$, $\mathbf{v} = (v_1, v_2, \dots, v_n)^\top$, $\mathbf{b} = (b_1, b_2, \dots, b_q)^\top$, and $\mathbf{u} = (u_1, u_2, \dots, u_q)^\top$ be nonnegative vectors such that $\sum_{i=1}^n a_i = \sum_{i=1}^q b_i$. If

$$\sum_{i:v_i \leq t} a_i \leq \sum_{i:u_i \leq t} b_i \quad \text{for any } t \geq 0,$$

then $\mathbf{b}^\top \mathbf{u} \leq \mathbf{a}^\top \mathbf{v}$.

The next two lemmas generalize results in Hashimoto and Mizuno (2021), where $p_i = 1 (i \in N)$.

Lemma 2. *Let $S' = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C} - \mathbf{p}), \mathbf{a} \rangle$ be any instance of NR-SSP, where $\mathbf{C} = (C_1, C_2, \dots, C_n)^\top$ such that $C_1 \geq p_1$ and $C_{j+1} - C_j \geq p_{j+1}$ for any $j \in \{1, 2, \dots, n-1\}$, then \mathbf{C} is optimal for S' .*

Proof. It is clear that \mathbf{C} is a feasible schedule for S' . Let \mathbf{C}^v be any feasible schedule for S' . Then we have that

$$\sum_{j: C_j^v - p_j \leq T} a_j \leq \sum_{i: u_i \leq T} b_i \text{ for any } T \geq 0$$

from (1), where $\mathbf{u} = \mathbf{C} - \mathbf{p}$ and $\mathbf{b} = \mathbf{a}$. We obtain

$$\mathbf{a}^\top (\mathbf{C} - \mathbf{p}) \leq \mathbf{a}^\top (\mathbf{C}^v - \mathbf{p}),$$

by applying Lemma 1 for \mathbf{a} , $\mathbf{v} = \mathbf{C}^v - \mathbf{p}$, $\mathbf{b} = \mathbf{a}$, and $\mathbf{u} = \mathbf{C} - \mathbf{p}$. Since the object function value is $\mathbf{a}^\top \mathbf{C}^v$, \mathbf{C} is optimal for S' . \square

Lemma 3. *Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ and $\tilde{S} = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q', \mathbf{u}', \mathbf{b}' \rangle$ be two instances of NR-SSP. If*

$$\sum_{i: u'_i \leq T} b'_i \leq \sum_{i: u_i \leq T} b_i \text{ for any } T \geq 0, \quad (3)$$

then $\mathbf{C}^\circ(S) \leq \mathbf{C}^\circ(\tilde{S})$ and $f_S(\mathbf{C}^\circ(S)) \leq f_{\tilde{S}}(\mathbf{C}^\circ(\tilde{S}))$ for any $\mathbf{o} \in \mathcal{O}$.

Proof. From condition (3), Algorithm 2 for any permutation \mathbf{o} outputs $\mathbf{C}^\circ(S)$ for S and $\mathbf{C}^\circ(\tilde{S})$ for \tilde{S} so that

$$\mathbf{C}^\circ(S) \leq \mathbf{C}^\circ(\tilde{S}).$$

Then we easily see that $f_S(\mathbf{C}^\circ(S)) \leq f_{\tilde{S}}(\mathbf{C}^\circ(\tilde{S}))$. \square

Proof of Theorem 3. Recall that $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, q, \mathbf{u}, \mathbf{b} \rangle$ and $S' = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C}^*(S) - \mathbf{p}), \mathbf{a} \rangle$ in Theorem 3. From Lemma 2, $\mathbf{C}^*(S)$ is an optimal schedule for S' . Since $\mathbf{C}^*(S)$ is a feasible schedule for S , (1) holds for every $T \geq 0$, namely,

$$\sum_{j: C^*(S)_j - p_j \leq T} a_j \leq \sum_{i: u_i \leq T} b_i \text{ for any } T \geq 0.$$

Hence $f_S(\mathbf{C}^\circ(S)) \leq f_{S'}(\mathbf{C}^\circ(S'))$ holds for any $\mathbf{o} \in \mathcal{O}$ from Lemma 3. If $f_{S'}(\mathbf{C}^\circ(S')) < 3f_{S'}(\mathbf{C}^*(S'))$, then we see that

$$\begin{aligned} f_S(\mathbf{C}^\circ(S)) &\leq f_{S'}(\mathbf{C}^\circ(S')) \\ &< 3f_{S'}(\mathbf{C}^*(S')) \\ &= 3\mathbf{a}^\top \mathbf{C}^*(S) \\ &= 3f_S(\mathbf{C}^*(S)). \end{aligned}$$

□

Then we prove Theorem 4 by using the following lemma.

Lemma 4. *Let $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C} - \mathbf{p}), \mathbf{a} \rangle$ be any instance of NR-SSP, where $\mathbf{C} = (C_1, C_2, \dots, C_n)^\top$ such that $C_1 \geq p_1$ and $C_{j+1} - C_j \geq p_{j+1}$ for any $j \in \{1, 2, \dots, n-1\}$. Let $\mathbf{d}_k = (\mathbf{0}_k^\top \mathbf{1}_{n-k}^\top)^\top$ be a vector whose first k elements are 0 and the latter $n-k$ elements are 1 for an integer $k \in \{0, 1, \dots, n-1\}$. Define $S^\dagger = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C} + L\mathbf{d}_k - \mathbf{p}), \mathbf{a} \rangle$ for any $L \geq 0$.*

For any $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, $f_S(\mathbf{C}^\mathbf{o}(S)) < 3f_S(\mathbf{C}^(S))$ implies $f_{S^\dagger}(\mathbf{C}^\mathbf{o}(S^\dagger)) < 3f_{S^\dagger}(\mathbf{C}^*(S^\dagger))$.*

Proof. Clearly \mathbf{C} is a feasible schedule for S . By Lemma 2, we have $\mathbf{C}^*(S) = \mathbf{C}$ and $\mathbf{C}^*(S^\dagger) = \mathbf{C} + L\mathbf{d}_k$. Then we see

$$\begin{aligned} f_{S^\dagger}(\mathbf{C}^*(S^\dagger)) - f_S(\mathbf{C}^*(S)) &= \mathbf{a}^\top(L\mathbf{d}_k) \\ &= L \sum_{j=k+1}^n a_j \\ &= LA_{k+1}^*. \end{aligned}$$

Since supply times $C_j - p_j, j \in \{1, 2, \dots, k\}$ in S^\dagger are equal to those in S , we see from Algorithm 2 that

$$C_{o(j)}^\mathbf{o}(S) = C_{o(j)}^\mathbf{o}(S^\dagger) \quad \text{for any } j \in \{1, 2, \dots, \alpha_k\},$$

where

$$\begin{aligned} \alpha_k &= \max\{j \in N \mid a_{o(1)} + \dots + a_{o(j)} \leq a_1 + \dots + a_k\} \\ &= \min\{j \in N \mid a_{o(j+1)} + \dots + a_{o(n)} \geq a_{k+1} + \dots + a_n\} \\ &= \lambda_{k+1} - 1. \end{aligned}$$

Then $\mathbf{C}' = (C'_1, C'_2, \dots, C'_n)$ define by

$$C'_{o(j)} = \begin{cases} C_{o(j)}^\mathbf{o}(S) & \text{for any } j \in \{1, 2, \dots, \lambda_{k+1} - 1\} \\ C_{o(j)}^\mathbf{o}(S) + L & \text{for any } j \in \{\lambda_{k+1}, \dots, n\} \end{cases}$$

is a feasible schedule for S^\dagger since $\mathbf{C}^\mathbf{o}(S)$ is feasible for S . We easily see that $\mathbf{C}^\mathbf{o}(S^\dagger) \leq \mathbf{C}'$ and $f_{S^\dagger}(\mathbf{C}^\mathbf{o}(S^\dagger)) \leq f_{S^\dagger}(\mathbf{C}')$ from Algorithm 2. Then we have that

$$\begin{aligned} f_{S^\dagger}(\mathbf{C}^\mathbf{o}(S^\dagger)) - f_S(\mathbf{C}^\mathbf{o}(S)) &\leq f_{S^\dagger}(\mathbf{C}') - f_S(\mathbf{C}^\mathbf{o}(S)) \\ &= \mathbf{a}^\top(\mathbf{C}' - \mathbf{C}^\mathbf{o}(S)) \\ &= L \sum_{j=\lambda_{k+1}}^n a_{o(j)} \\ &= LA_{\lambda_{k+1}}^\mathbf{o} \\ &< 2LA_{k+1}^* \\ &= 2(f_{S^\dagger}(\mathbf{C}^*(S^\dagger)) - f_S(\mathbf{C}^*(S))), \end{aligned}$$

where the strict inequality follows from Theorem 6. Therefore if $f_S(\mathbf{C}^\circ(S)) < 3f_S(\mathbf{C}^*(S))$, then we have that

$$\begin{aligned} & f_{S^\dagger}(\mathbf{C}^\circ(S^\dagger)) \\ & < f_S(\mathbf{C}^\circ(S)) + 2(f_{S^\dagger}(\mathbf{C}^*(S^\dagger)) - f_S(\mathbf{C}^*(S))) \\ & < 3f_{S^\dagger}(\mathbf{C}^*(S^\dagger)). \end{aligned}$$

□

Proof of Theorem 4. Recall that $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{C}' - \mathbf{p}), \mathbf{a} \rangle$ and $\bar{S} = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}), \mathbf{a} \rangle$ in Theorem 4. We can express that

$$\begin{aligned} \mathbf{C}' - \boldsymbol{\sigma}(\mathbf{p}) &= (C'_1 - C'_0 - p_1)\mathbf{d}_0 + (C'_2 - C'_1 - p_2)\mathbf{d}_1 \\ & \quad + \cdots + (C'_n - C'_{n-1} - p_n)\mathbf{d}_{n-1} \\ &= \sum_{k=0}^{n-1} L_k \mathbf{d}_k, \end{aligned}$$

where $C'_0 = 0$, $\mathbf{d}_k = (\mathbf{0}_k^\top \mathbf{1}_{n-k}^\top)^\top$, and $L_k = C'_{k+1} - C'_k - p_{k+1}$ for any $k \in \{0, 1, \dots, n-1\}$. From the conditions in the theorem, we have that $L_k \geq 0$ for any $k \in \{0, 1, \dots, n-1\}$. We define $\mathbf{D}_0 = \boldsymbol{\sigma}(\mathbf{p})$ and

$$\mathbf{D}_k = \mathbf{D}_{k-1} + L_{k-1} \mathbf{d}_{k-1}$$

for any $k \in N$. Then we see that

$$\mathbf{D}_n = \mathbf{C}'.$$

We also define $(n+1)$ -instances $S^k = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\mathbf{D}_k - \mathbf{p}), \mathbf{a} \rangle$ for $k \in \{0, 1, \dots, n\}$. Then we have that $S^0 = \bar{S}$ and $S^n = S$.

Now we consider two instances $S^0 = \bar{S}$ and S^1 . If $f_{S^0}(\mathbf{C}^\circ(S^0)) < 3f_{S^0}(\mathbf{C}^*(S^0))$, we have $f_{S^1}(\mathbf{C}^\circ(S^1)) < 3f_{S^1}(\mathbf{C}^*(S^1))$ from Lemma 4. For $k = 2, \dots, n$, we repeat this procedure for two instances S^{k-1} and S^k , then we finally obtain that $f_{S^n}(\mathbf{C}^\circ(S^n)) < 3f_{S^n}(\mathbf{C}^*(S^n))$, where $S^n = S$. □

6 Proof of Theorems 1, 2, and 5

In this section, we prove Theorem 5 at first. Then Theorem 1 is proved by using Theorems 3, 4, and 5. At last, we prove Theorem 2 by showing a simple instance.

Proof of Theorem 5. Recall that $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}), \mathbf{a} \rangle$ and $S^\dagger = \langle n, \mathcal{J}, \mathbf{a}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{a}) - \mathbf{a}), \mathbf{a} \rangle$. We define $\boldsymbol{\delta} = (\delta_1, \delta_2, \dots, \delta_n)^\top$ by

$$\delta_j = p_j - a_j \text{ for each } j \in N.$$

From Assumption 2, $\delta_j \geq 0$ for any $j \in N$. Then we easily see that $O(\mathbf{a}, \mathbf{p}) \subset O(\mathbf{a}, \mathbf{a})$. For any $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$, we need to prove

$$f_{S^\dagger}(\mathbf{C}^\mathbf{o}(S^\dagger)) < 3f_{S^\dagger}(\mathbf{C}^*(S^\dagger)) \Rightarrow f_S(\mathbf{C}^\mathbf{o}(S)) < 3f_S(\mathbf{C}^*(S)). \quad (4)$$

Define $D^*(S) = f_S(\mathbf{C}^*(S)) - f_{S^\dagger}(\mathbf{C}^*(S^\dagger))$ and $D^\mathbf{o}(S) = f_S(\mathbf{C}^\mathbf{o}(S)) - f_{S^\dagger}(\mathbf{C}^\mathbf{o}(S^\dagger))$. To prove (4), it suffices to show that

$$D^\mathbf{o}(S) \leq 3D^*(S). \quad (5)$$

Since the weight vector $\mathbf{w} = \mathbf{a}$ is common both in S and S^\dagger , we have that

$$\begin{aligned} D^*(S) &= \mathbf{a}^\top(\mathbf{C}^*(S) - \mathbf{C}^*(S^\dagger)), \\ D^\mathbf{o}(S) &= \mathbf{a}^\top(\mathbf{C}^\mathbf{o}(S) - \mathbf{C}^\mathbf{o}(S^\dagger)). \end{aligned}$$

From Lemma 2, $\mathbf{C}^*(S) = \boldsymbol{\sigma}(\mathbf{p})$ and $\mathbf{C}^*(S^\dagger) = \boldsymbol{\sigma}(\mathbf{a})$. Then we see that

$$\begin{aligned} D^*(S) &= \mathbf{a}^\top(\boldsymbol{\sigma}(\mathbf{p}) - \boldsymbol{\sigma}(\mathbf{a})) \\ &= \sum_{j=1}^n a_j \sum_{k=1}^j \delta_k \\ &= \sum_{k=1}^n \delta_k \sum_{j=k}^n a_j \\ &= \sum_{k=1}^n \delta_{o(k)} \sum_{j=o(k)}^n a_j \\ &= \sum_{k=1}^n A_{o(k)}^* \delta_{o(k)}. \end{aligned} \quad (6)$$

Next we evaluate $D^\mathbf{o}(S)$. Comparing S^\dagger with S , the processing time vector and the supply time vector are different. Let $k \in N$. When a_k increases by δ_k , j -th element of the supply time vector $\mathbf{u} = \boldsymbol{\sigma}(\mathbf{a}) - \mathbf{a}$ increases by δ_k for each $j \in \{k+1, \dots, n\}$. Then $C_{o(1)}^\mathbf{o}, \dots, C_{o(\alpha_k)}^\mathbf{o}$ do not change and $C_{o(\alpha_k+1)}^\mathbf{o}, \dots, C_{o(n)}^\mathbf{o}$ may increase at most δ_k , where

$$\begin{aligned} \alpha_k &= \max\{j \in N \mid a_{o(1)} + \dots + a_{o(j)} \leq a_1 + \dots + a_k\} \\ &= \lambda_{k+1} - 1. \end{aligned}$$

Figure 1 illustrates this situation.

Besides, when $o(k)$ -th element of processing time vector increases by $\delta_{o(k)}$, not only $C_{o(k)}^\mathbf{o}$ but also $C_{o(k+1)}^\mathbf{o}, \dots, C_{o(n)}^\mathbf{o}$ may increase at most $\delta_{o(k)}$. Figure 2 depicts this situation. From these facts, we have that




Figure 1: Gantt-charts of the schedules by the algorithm for S^\dagger and S_1 , where the supply times u_j ($j \geq k+1$) in S_1 increase from those in S^\dagger by δ_k .




Figure 2: Gantt-charts of the schedules by the algorithm for S^\dagger and S_2 , where the processing time $p_{o(k)}$ in S_2 increases from that in S^\dagger by $\delta_{o(k)}$.

$$\begin{aligned}
D^o(S) &\leq \sum_{k=1}^n \delta_k \sum_{l=\alpha_k+1}^n a_{o(l)} + \sum_{k=1}^n \delta_{o(k)} \sum_{l=k}^n a_{o(l)} \\
&= \sum_{k=1}^n A_{\alpha_k+1}^o \delta_k + \sum_{k=1}^n A_k^o \delta_{o(k)} \\
&= \sum_{k=1}^n A_{\lambda_{k+1}}^o \delta_k + \sum_{k=1}^n A_k^o \delta_{o(k)} \\
&= \sum_{k=1}^n A_{\lambda_{o(k)+1}}^o \delta_{o(k)} + \sum_{k=1}^n A_k^o \delta_{o(k)} \\
&\leq \sum_{k=1}^n \left(2A_{o(k)+1}^* + A_k^o \right) \delta_{o(k)}, \tag{7}
\end{aligned}$$

where the last inequality follows from Theorem 6. Comparing (7) with (6), the inequality

$$\sum_{k=1}^n A_k^o \delta_{o(k)} \leq \sum_{k=1}^n \left(2a_{o(k)} + A_{o(k)}^* \right) \delta_{o(k)},$$

implies (5). The inequality above is proved in Lemma 5 below. \square

Lemma 5. For any instance $S = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}), \mathbf{a} \rangle$ and $\mathbf{o} \in O(\mathbf{p}, \mathbf{a})$,

we define

$$M^*(S) = \sum_{k=1}^n \left(2a_{o(k)} + A_{o(k)}^* \right) \delta_{o(k)}$$

and

$$M^o(S) = \sum_{k=1}^n A_k^o \delta_{o(k)},$$

where $\delta_j = p_j - a_j$ for each $j \in N$. Then we have that

$$M^o(S) \leq M^*(S). \quad (8)$$

Proof. We show (8) by induction on n . In the case $n = 1$, the result is trivial. Then we prove (8) for S , \mathbf{o} , and $n \geq 2$ under the assumption that (8) holds for $n - 1$.

We define $S^- = \langle n-1, \mathcal{J}^-, \mathbf{p}^-, \mathbf{a}^-, n-1, (\boldsymbol{\sigma}(\mathbf{p}^-) - \mathbf{p}^-), \mathbf{a}^- \rangle$ such that $\mathcal{J}^- = (J_1, J_2, \dots, J_{o(1)-1}, J_{o(1)+1}, \dots, J_n)$, $\mathbf{p}^- = (p_1, p_2, \dots, p_{o(1)-1}, p_{o(1)+1}, \dots, p_n)^\top$, and $\mathbf{a}^- = (a_1, a_2, \dots, a_{o(1)-1}, a_{o(1)+1}, \dots, a_n)^\top$. We also define $\mathbf{o}^- = (o(2), o(3), \dots, o(n))$, $\Delta^* = M^*(S) - M^*(S^-)$, and $\Delta^o = M^o(S) - M^{o^-}(S^-)$. It is obvious that \mathbf{o}^- satisfies the conditions in Definition 1. Under the assumption $M^{o^-}(S^-) \leq M^*(S^-)$, we need to show

$$\Delta^o \leq \Delta^* \quad (9)$$

to prove (8) for n .

Next we show that (9) holds. We have that

$$M^*(S^-) = \sum_{k=2}^n \left(2a_{o(k)} + \sum_{o(k) \leq j \leq n, j \neq o(1)} a_j \right) \delta_{o(k)}$$

and

$$\begin{aligned} \Delta^* &= M^*(S) - M^*(S^-) \\ &= 2a_{o(1)}\delta_{o(1)} + A_{o(1)}^*\delta_{o(1)} + \sum_{o(k) < o(1)} a_{o(1)}\delta_{o(k)}. \end{aligned}$$

We also have that

$$M^{o^-}(S^-) = \sum_{k=2}^n \delta_{o(k)} \sum_{j=k}^n a_{o(j)}$$

and

$$\begin{aligned}
\Delta^{\circ} &= M^{\circ}(S) - M^{\circ-}(S^{-}) \\
&= \sum_{j=1}^n a_{o(j)} \delta_{o(1)} \\
&= \sum_{j=1}^{j^*-1} a_{o(j)} \delta_{o(1)} + a_{o(j^*)} \delta_{o(1)} + A_{j^*+1}^{\circ} \delta_{o(1)},
\end{aligned}$$

where

$$j^* = \min\{j \in N \mid r_{o(j)} > r_{o(1)}\}.$$

(In the case of $r_{o(j)} \leq r_{o(1)}$ for any $j \in N$, we define $j^* = n + 1$ and $a_{o(j^*)} = 0$.) Since $r_{o(j)} \leq r_{o(1)}$ for any $j < j^*$, we have from Theorem 7-(i) that

$$\begin{aligned}
\sum_{j=1}^{j^*-1} a_{o(j)} \delta_{o(1)} &= \sum_{1 \leq j \leq j^*-1, o(j) \geq o(1)} a_{o(j)} \delta_{o(1)} + \sum_{1 \leq j \leq j^*-1, o(j) < o(1)} a_{o(j)} \delta_{o(1)} \\
&\leq \sum_{1 \leq j \leq j^*-1, o(j) \geq o(1)} a_{o(j)} \delta_{o(1)} + \sum_{1 \leq j \leq j^*-1, o(j) < o(1)} a_{o(1)} \delta_{o(j)} \\
&\leq A_{o(1)}^* \delta_{o(1)} + \sum_{o(k) < o(1)} a_{o(1)} \delta_{o(k)}.
\end{aligned}$$

Since $j^* > 1$ and $r_{o(j^*)} > r_{o(1)}$, we have from Theorem 7-(ii) that

$$a_{o(j^*)} \delta_{o(1)} + A_{j^*+1}^{\circ} \delta_{o(1)} \leq 2a_{o(1)} \delta_{o(1)}.$$

Hence $\Delta^{\circ} \leq \Delta^*$ holds. \square

Proof of Theorem 1. By Theorems 3-5, it suffices to prove it for any instance $S = \langle n, \mathcal{J}, \mathbf{a}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{a}) - \mathbf{a}), \mathbf{a} \rangle$ and $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$. From Lemma 2, the optimal schedule for S is $\mathbf{C}^*(S) = \boldsymbol{\sigma}(\mathbf{a})$. Define $Q = \sum_{j=1}^n a_j$. Then we see that

$$\begin{aligned}
f_S(\mathbf{C}^*(S)) &= \mathbf{a}^T \boldsymbol{\sigma}(\mathbf{a}) \\
&= \sum_{j=1}^n a_j \sum_{i=1}^j a_i \\
&= \frac{1}{2} Q^2 + \frac{1}{2} \sum_{j=1}^n a_j^2.
\end{aligned}$$

The output schedule of Algorithm 2 is $\mathbf{C}^{\circ}(S)$ for $\mathbf{o} \in O(\mathbf{a}, \mathbf{p})$. Since all the resources are available after the time $t = Q - a_{o(n)}$, we have that

$$C_{o(j)}^{\circ}(S) \leq Q - a_{o(n)} + \sum_{i=1}^j a_{o(i)} < Q + \sum_{i=1}^j a_{o(i)}$$

for each $j \in N$. Thus we see that

$$\begin{aligned}
f_S(\mathbf{C}^\circ(S)) &= \sum_{j=1}^n a_{o(j)} C_{o(j)}^\circ(S) \\
&< \sum_{j=1}^n a_{o(j)} \left(Q + \sum_{i=1}^j a_{o(i)} \right) \\
&= Q^2 + \sum_{j=1}^n a_{o(j)} \sum_{i=1}^j a_{o(i)} \\
&= Q^2 + \frac{1}{2}Q^2 + \frac{1}{2} \sum_{j=1}^n a_j^2 \\
&< 3f_S(\mathbf{C}^*(S)).
\end{aligned}$$

□

Finally we prove Theorem 2.

Proof of Theorem 2. Let $0 < e < 0.1$ and $\tilde{S} = \langle n, \mathcal{J}, \mathbf{p}, \mathbf{a}, n, (\boldsymbol{\sigma}(\mathbf{p}) - \mathbf{p}), \mathbf{a} \rangle$ for $n = 3$, $\mathbf{p} = (e, e, 1)^\top$, and $\mathbf{a} = (1 - e, 1, 1 + e)^\top$. We easily see that

$$\begin{aligned}
\mathbf{C}^*(\tilde{S}) &= \boldsymbol{\sigma}(\mathbf{p}) = (e, 2e, 1 + 2e)^\top, \\
\mathbf{o} &= (3, 2, 1), \\
\mathbf{C}^\circ(\tilde{S}) &= (1 + 3e, 1 + 2e, 1 + e)^\top.
\end{aligned}$$

Then

$$\begin{aligned}
&\lim_{e \rightarrow 0^+} \frac{f_{\tilde{S}}(\mathbf{C}^\circ(\tilde{S}))}{f_{\tilde{S}}(\mathbf{C}^*(\tilde{S}))} \\
&= \lim_{e \rightarrow 0^+} \frac{(1 - e)(1 + 3e) + (1 + 2e) + (1 + e)(1 + e)}{(1 - e)e + 2e + (1 + e)(1 + 2e)} \\
&= 3.
\end{aligned}$$

Hence we obtain the result.

□

References

- Bérczi, K., Király, T., & Omlor, S. (2020). Scheduling with non-renewable resources: Minimizing the sum of completion times. In *International Symposium on combinatorial optimization*, pages 167–178. Springer.
- Briskorn, D., Jaehn, F., & Pesch, E. (2013). Exact algorithms for inventory constrained scheduling on a single machine. *Journal of scheduling*, 16(1), 105–115.

- Carlier, J. & Rinnooy Kan, A. H. G. (1982). Scheduling subject to nonrenewable-resource constraints. *Operations Research Letters*, 1(2), 52–55.
- Gafarov, E. R., Lazarev, A. A., & Werner, F. (2011). Single machine scheduling problems with financial resource constraints: Some complexity results and properties. *Mathematical Social Sciences*, 62(1), 7–13.
- Graham, R. L., Lawler, E. L., Lenstra, J. K., & Kan, A. R. (1979). Optimization and approximation in deterministic sequencing and scheduling: A survey. *Annals of discrete mathematics*, 5, 287–326.
- Grigoriev, A., Holthuijsen, M., & Van De Klundert, J. (2005). Basic scheduling problems with raw material constraints. *Naval Research Logistics*, 52(6), 527–535.
- Györgyi, P. & Kis, T. (2015). Approximability of scheduling problems with resource consuming jobs. *Annals of Operations Research*, 235(1), 319–336.
- Györgyi, P. & Kis, T. (2019). Minimizing total weighted completion time on a single machine subject to non-renewable resource constraints. *Journal of Scheduling*, 22(6), 623–634.
- Györgyi, P. & Kis, T. (2020). New complexity and approximability results for minimizing the total weighted completion time on a single machine subject to non-renewable resource constraints. *arXiv e-prints*, Available: <http://arxiv.org/abs/1608.02792>.
- Hashimoto, S. & Mizuno, S. (2021). A tight approximation ratio of a list scheduling algorithm for a single-machine scheduling problem with a non-renewable resource. *Journal of Scheduling*, doi: <https://doi.org/10.1007/s10951-021-00681-y>.
- Kis, T. (2015). Approximability of total weighted completion time with resource consuming jobs. *Operations Research Letters*, 43(6), 595–598.
- Morsy, E. & Pesch, E. (2015). Approximation algorithms for inventory constrained scheduling on a single machine. *Journal of Scheduling*, 18(6), 645–653.
- Slowiński, R. (1984). Preemptive scheduling of independent jobs on parallel machines subject to financial constraints. *European Journal of Operational Research*, 15(3), 366–373.