

Component Edge Connectivity of Hypercube-like Networks *

Dong Liu Pingshan Li [†] Bicheng Zhang

*Key Laboratory of Intelligent computing & Information processing of Ministry of Education
School of Mathematical and Computational Science, Xiangtan University, Xiangtan, Hunan 41105, PR China*

Abstract

As a generalization of the traditional connectivity, the g -component edge connectivity $c\lambda_g(G)$ of a non-complete graph G is the minimum number of edges to be deleted from the graph G such that the resulting graph has at least g components. Hypercube-like networks (HL-networks for short) are obtained by manipulating some pairs of edges in hypercubes, which contain several famous interconnection networks such as twisted cubes, Möbius cubes, crossed cubes, locally twisted cubes. In this paper, we determine the $(g + 1)$ -component edge connectivity of the n -dimensional HL-networks for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 8$.

Key words: HL-networks; component connectivity; component edge connectivity.

1 Introduction

As we all know, the interconnection network is an important part of the multiprocessor system. For convenience, we usually model the interconnection network by a graph, with vertices representing processors and edges representing links between processors. There are many parameters to evaluate the reliability of a network. The traditional connectivity $\kappa(G)$ of the graph G is the most crucial one among them. Generally speaking, the larger the $\kappa(G)$ of the network is, the better its reliability is. However, the traditional connectivity only indicates when the network will break but does not further depict the properties of components, which makes it impossible to accurately evaluate the reliability of the network. In order to further describe the properties of components, a lot of more precise connectivity have been proposed, such as extra connectivity [10], super connectivity [2] and restricted connectivity [8]. In 1984, Chartrand et al. [4] and Sampathkumar [14] respectively introduced g -component connectivity $c\kappa_g(G)$ and g -component edge connectivity $c\lambda_g(G)$ of the graph G .

A g -component (edge) cut of a non-complete G is a vertex (edge) set to be deleted from the graph G such that the resulting graph has at least g components. The g -component (edge) connectivity of G , written $c\kappa_g(G)$ ($c\lambda_g(G)$), is the minimum size of the g -component (edge) cut of G . Obviously, the component (edge) connectivity is a generalization of the traditional connectivity and $\kappa(G) = c\kappa_2(G)$ ($\lambda(G) = c\lambda_2(G)$). More importantly, compared with the traditional connectivity, the component (edge) connectivity can be better satisfied in practical applications. Thus, many researches have been focused on the component connectivity of some famous networks in recent years. (see, for example [5, 15, 16, 18]). But there are relatively few papers about component edge connectivity. Zhao et al. [17] determined the $(g + 1)$ -component edge connectivity of hypercubes for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 7$. Based on this nice work, we shall be centered on the $(g + 1)$ -component edge connectivity of hypercube-like networks for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 8$.

*This research is supported partially by the General Project of Hunan Provincial Education Department of China (19C1742), Hu Xiang Gao Ceng Ci Ren Cai Ju Jiao Gong Cheng-Chuang Xin Ren Cai (No. 2019RS1057), the Project of Scientific Research Fund of Hunan Provincial Science and Technology Department (No. 2018WK4006).

[†]Corresponding author.

E-mail address: lips@xtu.edu.cn (Pingshan Li).

As a popular topology for the design of the multiprocessor system, hypercubes have many excellent properties including symmetry, relatively diameter, good connectivity and recursive scalability. Improving these properties has led to the evolution of hypercube variants. A lot of hypercube variants, such as twisted cubes [11], crossed cubes [7] and Möbius cubes [6], have been introduced successively. Many of the properties of these networks are identical with that of the hypercubes. In particular, their diameter is shorter than that of the hypercubes. In order to conduct a unified study on these variants, Vaidya et al. proposed hypercube-like networks (in brief, HL-networks), which contain all of the networks mentioned above. Hence, HL-networks attracted considerable attention in the past (see, for example [3, 12]). This class of networks are sometimes called BC-networks [9]. we will use the term HL-networks in this paper.

The recursive definition of the HL-networks is as follows:

$$HL_1 = \{K_1\} \text{ and } HL_n = \{G_{n-1} \oplus G_{n-1}^* | G_{n-1}, G_{n-1}^* \in HL_{n-1}\},$$

where the symbol “ \oplus ” represents the perfect matching operation that connects G_{n-1} and G_{n-1}^* using some disjoint edges. It's easy to get that $HL_0 = \{K_1\}$, $HL_1 = \{K_2\}$, $HL_2 = \{C_4\}$, and $HL_3 = \{Q_3, G(8, 4)\}$, where C_4 is a cycle of length 4, and Q_3 and $G(8, 4)$ are shown in Figure 1. The n -dimensional HL-network G_n is n regular, and it has 2^n vertices and $n2^{n-1}$ edges.

Next, we will introduce some notations and definitions which will be used in this paper. Let $G = (V(G), E(G))$ be a graph. The size of G is the number of edges of G . The degree of v , denoted by $d_G(v)$, is the number of edges incident to v in G . For a subset X of $V(G)$, $G[X]$ is the subgraph induced by X . If D_i and D_j are two disjoint subgraphs of G , then we use $E(D_i, D_j)$ to denote the set of edges between the subgraphs D_i and D_j . Similarly, we use $E(v, D_i)$ to denote the set of edges between v and D_i , where $v \in V(G)$ but $v \notin V(D_i)$. For any $v \in V(G)$, $G - v$ denotes a subgraph obtained by deleting v and edges incident to v . Similarly, $G - D_i$ denotes a subgraph obtained by removing all vertices in $V(D_i)$ and all edges incident to vertices of $V(D_i)$, where D_i is a subgraph of G .

The rest of this paper is organized as follows. In section 2, we shall determine the maximum size of the subgraph induced by g vertices in HL-networks. In section 3, we shall determine the $(g + 1)$ -component edge connectivity of HL-networks by applying the results of section 2.




Figure 1: Two 3-dimensional HL-networks: (a) Q_3 (b) $G(8, 4)$

2 The maximum size of the subgraph induced by g vertices

It is a classical problem to determine the size (the number of edges) of the subgraph that satisfies some given property in a graph. For instance, Erdős has studied an interesting problem in [1]: What is the maximum size of the subgraph without cycles of length 4 in hypercubes. In this section, we shall determine the maximum size of the subgraph induced by g vertices in HL-networks. Furthermore, its application in the next section helps us find the $(g + 1)$ -component edge connectivity of HL-networks.

Let e_g be the maximum size of the subgraph induced by g vertices in the n -dimensional HL-network G_n , that is, $e_g = \max \{|E(G_n[X])| : X \subseteq V(G_n) \text{ and } |X| = g\}$. By the mathematical principle of




Figure 2: The HL-networks D_i and D_i^* when $s=2$

converting decimal digit to binary digit, any integer g can be written as the sum of the exponents of 2, that is, $g = \sum_{i=0}^s 2^{t_i}$, where $t_0 = \lfloor \log_2 g \rfloor$, $t_i = \lfloor \log_2(g - \sum_{r=0}^{i-1} 2^{t_r}) \rfloor$ for $i \geq 1$. Li and Yang have determined e_g of hypercubes in [13]. In order to avoid confusion, we use $e_g(Q_n)$ to denote e_g of hypercubes, where $e_g(Q_n) = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$. Furthermore, the function $e_g(Q_n)$ has the following property.

Lemma 2.1 [13][17] If $g_0 \leq g_1$, then $e_{g_0+g_1}(Q_n) \geq e_{g_0}(Q_n) + e_{g_1}(Q_n) + g_0$.

Next, we shall give an algorithm to find the subgraph H induced by g vertices in the n -dimensional HL-network G_n such that $|E(H)| = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$.

Algorithm- MS

Input: An n -dimensional network G_n , two integer g (suppose $g = \sum_{i=0}^s 2^{t_i}$) and s .

Output: A vertex set $V = V(D_0) \cup V(D_1) \dots \cup V(D_s)$ and a subgraph $H = G_n[V]$.

Initialization: $i \leftarrow 0$, $V \leftarrow \emptyset$, where \emptyset is an empty set.

Iteration: As long as $i \leq s$, we take a $t_i + 1$ -dimensional subcube T_i from D_{i-1}^* ($D_{-1}^* = G_n$). Then T_i can be written as $D_i \oplus D_i^*$, where D_i and D_i^* are two t_i -dimensional subcubes. Add $V(D_i)$ to V , $i \leftarrow i + 1$.

For ease of understanding, we list D_i as follows (see Figure 2 for $s = 2$):

- D_0 (t_0 -dimensional subcube);
- D_1 (t_1 -dimensional subcube taken from D_0^*);
- D_2 (t_2 -dimensional subcube taken from D_1^*);
-
- D_s (t_s -dimensional subcube taken from D_{s-1}^*).

We need to verify that the subgraph H found by the algorithm- MS that satisfies $|V(H)| = g$ and $|E(H)| = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$.

Note that $H = G_n[V(D_0) \cup V(D_1) \dots \cup V(D_s)]$ and $|V(D_i)| = 2^{t_i}$. Thus,

$$V(H) = \sum_{i=0}^s |V(D_i)| = \sum_{i=0}^s 2^{t_i} = g.$$

By the choice of D_i , one has $|E(D_i, D_j)| = |V(D_j)| = 2^{t_j}$ for $i < j$. Thus,

$$E(H) = \sum_{i=0}^s |E(D_i)| + \sum_{i=0}^{s-1} \sum_{j=i+1}^s |E(D_i, D_j)| = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}.$$

Theorem 2.2 For any $G_n \in HL_n$, suppose that $X \subseteq V(G_n)$ with $|X| = g = \sum_{i=0}^s 2^{t_i}$, then $e_g = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$.

Proof: For convenience, we define $f(g) = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$. Firstly, we shall prove $|E(G_n[X])| \leq f(g)$ for any $X \in V(G_n)$ with $|X| = g = \sum_{i=0}^s 2^{t_i}$ by induction on n . Clearly, the result holds for $n = 2$ since $G_2 \cong C_4$. So we assume $n \geq 3$. Note that $G_n = G_{n-1} \oplus G_{n-1}^*$. Let $X_0 = X \cap G_{n-1}$, $X_1 = X \cap G_{n-1}^*$, $|X_i| = g_i$ for $i = 0, 1$ and $g = g_0 + g_1$. Without loss of generality, we suppose $g_1 \geq g_0$. Note $f(g) = e_g(Q_n)$. By induction hypothesis, we have that

$$\begin{aligned} |E(G_n[X])| &\leq |E(G_n[X_0])| + |E(G_n[X_1])| + g_0 \\ &= |E(G_{n-1}[X_0])| + |E(G_{n-1}^*[X_1])| + g_0 \\ &\leq f(g_0) + f(g_1) + g_0 \\ &= e_{g_0}(Q_n) + e_{g_1}(Q_n) + g_0 \\ &\leq e_{g_0+g_1}(Q_n) = f(g_0 + g_1) = f(g). \end{aligned}$$

The third inequality holds because of Lemma 2.1.

Using the algorithm-*MS*, we can find a subgraph H such that $|V(H)| = g$ and $|E(H)| = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i} = f(g)$. Thus, $e_g = f(g) = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$. \square

Here are some properties of the function e_g .

Lemma 2.3 *If $g \leq 2^{n-2}$, then $(n-2)g - 2e_g \geq 0$.*

Proof: If $g \leq 2^{n-2}$, then we can take a subgraph H from an $(n-2)$ -dimensional HL-network G_{n-2} such that $|V(H)| = g$ and $|E(H)| = e_g$. Note that G_{n-2} is an $(n-2)$ -regular graph. Thus, $(n-2)g - 2e_g \geq 0$. \square

Lemma 2.4 $e_{i+1} = e_i + s + 1$, where $i = 2^{t_0} + 2^{t_1} + \dots + 2^{t_s}$.

Proof: If $t_s > 0$, then $i + 1 = 2^{t_0} + 2^{t_1} + \dots + 2^{t_s} + 2^0$. Note that $e_i = \sum_{i=0}^s t_i 2^{t_i-1} + \sum_{i=0}^s i 2^{t_i}$. We have that

$$e_{i+1} = \sum_{i=0}^s t_i 2^{t_i-1} + 0 \cdot 2^{0-1} + \sum_{i=0}^s i 2^{t_i} + (s+1) \cdot 2^0 = e_i + s + 1.$$

Otherwise, $t_s = 0$. Then there is an integer r ($0 \leq r \leq s$) such that $t_j = s - j$ for all $r \leq j \leq s$. In other words, $i = 2^{t_0} + \dots + 2^{t_{r-1}} + 2^{s-r} + 2^{s-r-1} + \dots + 2^1 + 2^0$. Then $i + 1 = 2^{t_0} + \dots + 2^{t_{r-1}} + 2^{s-r} + 2^{s-r-1} + \dots + 2^1 + 2^0 + 2^0 = 2^{t_0} + \dots + 2^{t_{r-1}} + 2^{s-r+1}$.

Using the algorithm-*MS*, we take two subgraph H_1 and H_2 from G_n , where H_1 and H_2 are induced by i and $i + 1$ vertices respectively. Clearly, $|E(H_1)| = e_i$ and $|E(H_2)| = e_{i+1}$. we can assume that

$$\begin{aligned} H_1 &= G_n [V(D_0) \cup \dots \cup V(D_{r-1}) \cup V(D_r) \cup V(D_{r+1}) \cup \dots \cup V(D_s)]; \\ H_2 &= G_n [V(D_0) \cup \dots \cup V(D_{r-1}) \cup V(D'_r)]. \end{aligned} \tag{1}$$

where D_i is a t_i -dimensional subcube, D'_r is an $(s-r+1)$ -dimensional subcube.

In fact, $|V(H_1)| = i$ and $|V(H_2)| = i + 1$. In other words, H_1 has one less vertex than H_2 . Let the vertex be v . From (1), we can see that the first r subcubes D_i in H_1 and H_2 are the same. Thus, one has $v \subseteq V(D'_r)$. Clearly, $d_{D'_r}(v) = s - r + 1$. In addition, there is only one edge between v and every D_i in H_2 for $0 \leq i \leq r - 1$. Thus, $d_{H_2}(v) = r + (s - r + 1) = s + 1$. We have that $|E(H_2)| = |E(H_1)| + d_{H_2}(v)$, that is, $e_{i+1} = e_i + s + 1$. \square

Lemma 2.5 *If $i \leq j$, then $e_{i+1} + e_j \leq e_{i+j}$.*

Proof: Suppose $i = 2^{t_0} + \dots + 2^{t_s}$, then $e_{i+1} = e_i + s + 1$ by Lemma 2.4. Note that $e_g(Q_n) = e_g$. By the Lemma 2.1, $e_{i+j} \geq e_i + e_j + i = e_{i+1} + e_j + i - (s + 1) \geq e_{i+1} + e_j$. \square

3 Component edge connectivity of HL-networks

In this section, we shall apply Theorem 2.2 to determine the $(g + 1)$ -component edge connectivity of HL-networks.

For any $X \subseteq V(G_n)$, we use E_X to denote a set of edges in which each edge has exactly one endpoint in X .

Lemma 3.1 *For any $G_n \in HL_n$, let $X \subseteq V(G_n)$ with $|X| = g = \sum_{i=0}^s 2^{t_i}$. Then $|E_X| \geq ng - 2e_g$. Moreover, $ng - e_g$ is strictly increasing (respect to g) for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 2$.*

Proof: Note that G_n is an n -regular graph. We find that $|E_X| = n|X| - 2|E(G_n[X])| \geq ng - 2e_g$. According to Lemma 2.4, if $g \leq 2^{\lceil \frac{n}{2} \rceil}$ and $n \geq 2$, then $e_{g+1} - e_g = s + 1 < n$. Hence, we have that $(n(g + 1) - e_{g+1}) - (ng - e_g) = n - (e_{g+1} - e_g) > 0$, which indicate that $ng - e_g$ is strictly increasing for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 2$. \square

Theorem 3.2 *For any $G_n \in HL_n$, $c\lambda_{g+1}(G_n) = ng - e_g$ for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 8$.*

Proof: We first prove that $c\lambda_{g+1}(G_n) \leq ng - e_g$ by constructing a $(g + 1)$ -component edge-cut F such that $|F| = ng - e_g$. Using the algorithm-*MS*, we can get a subgraph H such that $|V(H)| = g$ and $|E(H)| = e_g$. Let $F = E_{V(H)} \cup E(H)$, then we have that $|F| = ng - e_g$. Moreover, $G_n - F$ has at least $g + 1$ components. Thus, $c\lambda_{g+1}(G_n) \leq ng - e_g$.

Next, we shall prove that $c\lambda_{g+1}(G_n) \geq ng - e_g$. Assume that F is the smallest $(g + 1)$ -component edge-cut and $G_n - F$ has exactly $g + 1$ components. Denote the $g + 1$ components in $G_n - F$ by C_1, C_2, \dots, C_{g+1} . Without loss of generality, we can assume that $|V(C_1)| \leq |V(C_2)| \leq \dots \leq |V(C_{g+1})|$.

Case 1: Suppose that $|V(C_{g+1})| < 2^{n-2}$.

In this case, $|V(C_i)| < 2^{n-2}$ for all i . Note that $\sum_{i=1}^{g+1} |V(C_i)| = 2^n$. Then we can find an integer j such that $\sum_{i=1}^j |V(C_i)| < 2^{n-2}$ but $2^{n-2} \leq \sum_{i=1}^{j+1} |V(C_i)| < 2^{n-1}$. Let $X = \bigcup_{i=1}^{j+1} V(C_i)$ with $|X| = m = \sum_{i=0}^s 2^{t_i}$. Clearly, $2^{n-2} \leq m = \sum_{i=0}^s 2^{t_i} < 2^{n-1}$. It follow that $t_0 = n - 2$. We suppose that $m' = m - 2^{t_0}$, then $m' < 2^{n-2}$. By Lemma 2.3, we have that $(n - 2)m' - 2e_{m'} \geq 0$. Combining with Lemma 3.1, we have that

$$\begin{aligned} |E_X| &\geq nm - 2e_m \\ &= n2^{t_0} + n(2^{t_1} + \dots + 2^{t_s}) - t_0 2^{t_0} - (\sum_{i=1}^s t_i 2^{t_i} + \sum_{i=1}^s 2 \cdot i \cdot 2^{t_i}) \\ &= (n - t_0)2^{t_0} + (n \sum_{i=1}^s 2^{t_i} - (\sum_{i=1}^s t_i 2^{t_i} + \sum_{i=1}^s 2 \cdot i \cdot 2^{t_i})) \\ &= (n - t_0)2^{t_0} + nm' - 2e_{m'} - 2m' \\ &\geq (n - t_0)2^{t_0} \\ &= (n - t_0) \cdot 2^{(t_0 - \lceil \frac{n}{2} \rceil)} \cdot 2^{\lceil \frac{n}{2} \rceil}. \end{aligned}$$

Note that $t_0 = n - 2$. Then $(n - t_0) \cdot 2^{(t_0 - \lceil \frac{n}{2} \rceil)} > \frac{3n}{4}$ for $n \geq 8$. we have that

$$|F| \geq |E_X| > \frac{3n}{4} \cdot 2^{\lceil \frac{n}{2} \rceil} \geq n \cdot 2^{\lceil \frac{n}{2} \rceil} - e_{2^{\lceil \frac{n}{2} \rceil}} \geq ng - e_g.$$

The last inequality holds, because $ng - e_g$ is strictly increasing (respect to g) for $g \leq 2^{\lceil \frac{n}{2} \rceil}$ by Lemma 3.1.

Case 2: Suppose that $|V(C_{g+1})| \geq 2^{n-2}$.

Let $|V(C_i)| = m_i$ for $1 \leq i \leq g$ and let $m = \sum_{i=1}^g m_i$. Set $X = \bigcup_{i=1}^g V(C_i)$. If $m \geq 2^{n-2}$, we can get that $|F| \geq |E_X| > ng - e_g$ by using the proof similar to case 1. So we assume $m < 2^{n-2}$. If $m = g$, then the component C_i ($1 \leq i \leq g$) is an isolated vertex. Thus, $|F| \geq ng - e_g$. If $m > g$, then let $F' = F \cap E(G_n[X])$. We have that $|F| \geq |\bigcup_{i=1}^g E_{V(C_i)}| = |E_{V(C_1)}| + \dots + |E_{V(C_g)}| - |F'|$. Since $|E_{V(C_i)}| \geq nm_i - 2|E(C_i)|$ and $|F'| \leq e_m - |\bigcup_{i=1}^g E(C_i)|$, we have that




Figure 3: The edges between the components

$$\begin{aligned}
|F| &\geq |\bigcup_{i=1}^g E_V(C_i)| = |E_V(C_1)| + \dots + |E_V(C_g)| - |F'| \\
&\geq \sum_{i=1}^g (nm_i - 2|E(C_i)|) - e_m + \sum_{i=1}^g |E(C_i)| \\
&= nm - e_m - \sum_{i=1}^g |E(C_i)| \\
&\geq nm - e_m - \sum_{i=1}^g e_{m_i}.
\end{aligned}$$

By Lemma 2.5, $e_{m_1} + e_{m_2} + \dots + e_{m_g} \leq e_{m_1+m_2-1} + e_{m_3} + \dots + e_{m_g} \leq \dots \leq e_{m-g+1}$. Thus, we have that $|F| \geq nm - e_m - \sum_{i=1}^g e_{m_i} \geq nm - e_m - e_{m-g+1}$.

Next, we shall show that $nm - e_m - e_{m-g+1} > ng - e_g$.

Note that $G_n = G_{n-1} \oplus G_{n-1}^*$. We take a subgraph H_1 of m vertices from an $(n-2)$ -dimensional subcube in G_{n-1} by using the algorithm-*MS*, since $m < 2^{n-2}$. Clearly, $|E(H_1)| = e_m$. We take a subgraph H_2 of $m-g+1$ vertices from H_1 by using the algorithm-*MS*, since $m-g+1 \leq m$. So $|E(H_2)| = e_{m-g+1}$. Use $v_m, v_{m-1}, \dots, v_{g+1}, v_g$ to denote the vertices of H_2 . By the proof of Lemma 2.4, there exists a vertex in H_2 , say v_g , such that $d_{H_2}(v_g) = s+1$, where $m-g = 2^{t_0} + 2^{t_1} + \dots + 2^{t_s}$. Use v_1, v_2, \dots, v_{g-1} to denote vertices of $V(H_1) - V(H_2)$. We assume that (see Figure 3)

$$\begin{aligned}
f_0 &= |E(H_1 - H_2)|; & f_4 &= |E(v_g, G_n - H_1)|; \\
f_1 &= |E(H_1 - H_2, G_n - H_1)|; & f_5 &= |E(G_n - H_1, H_2 - v_g)|; \\
f_2 &= |E(H_1 - H_2, H_2 - v_g)|; & f_6 &= |E(v_g, H_2 - v_g)|; \\
f_3 &= |E(v_g, H_1 - H_2)|.
\end{aligned}$$

Then we find that

$$\begin{aligned}
f_0 + f_1 + f_2 + f_3 + f_4 + f_5 &= (f_1 + f_4 + f_5) + (f_0 + f_2 + f_3) \\
&= |E_V(H_1)| + (|E(H_1)| - |E(H_2)|) \\
&= nm - 2|E(H_1)| + (|E(H_1)| - |E(H_2)|) \\
&= nm - e_m - e_{m-g+1}.
\end{aligned} \tag{2}$$

Let $S = \{v_1, v_2, \dots, v_g\}$. Note that $|E(G_n[S])| = f_0 + f_3 \leq e_g$, and $|E_S| + 2|E(G_n[S])| = (f_1 + f_2 + f_4 + f_6) + 2(f_0 + f_3) = ng$. Thus,

$$f_0 + f_1 + f_2 + f_3 + f_4 + f_6 = (f_1 + f_2 + f_4 + f_6) + (f_0 + f_3) \geq ng - e_g. \tag{3}$$

Moreover, each vertex of H_1 has at least two neighbors out of H_1 since H_1 be taken from an $(n-2)$ -dimensional subcube. Thus, $f_5 > |V(H_2 - v_g)| = m-g$. Note that $f_6 = d_{H_2}(v_g) = s+1$ and $m-g = 2^{t_0} + 2^{t_1} + \dots + 2^{t_s}$. It is not difficult to get that $f_5 > m-g \geq s+1 = f_6$. Thus, $nm - e_m - e_{m-g+1} > ng - e_g$ by comparing (2) with (3).

To sum up, $c\lambda_{g+1}(G_n) = ng - e_g$. \square

Corollary 3.3 Let F be a $(g + 1)$ -component edge-cut of the n -dimensional HL-network G_n and $|F| = c\lambda_{g+1}(G_n)$, then $G_n - F$ contains g isolated vertices for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 8$.

Proof: By the proof of Theorem 3.2, $|F| = c\lambda_{g+1}(G_n)$ if and only if $m = g$ in case 2, that is, g components are isolated vertices. \square

4 Conclusion

Component edge connectivity is an generation of the traditional connectivity. In this paper, we studied the $(g + 1)$ -component edge connectivity of HL-networks. $c\lambda_{g+1}(G_n) = ng - e_g$ for $g \leq 2^{\lceil \frac{n}{2} \rceil}$, $n \geq 8$. But for $g > 2^{\lceil \frac{n}{2} \rceil} + 1$, The problem has not been solved.

References

- [1] A. Baker, B. Bollobas, and A. Hajnal. A tribute to paul erdős — some of my favourite unsolved problems. 10.1017/CBO9780511983917:467–478, 1990.
- [2] D. Bauer, F. Boesch, C. Suffel, and R. Tindell. Connectivity extremal problems and the design of reliable probabilistic networks. *theory & applications of graphs*, 1981.
- [3] N. W. Chang and S. Y. Hsieh. 2,3-extraconnectivities of hypercube-like networks. *Journal of Computer and System Sciences*, 79(5):669C688, 2013.
- [4] G. Chartrand, S. F. Kapoor, L. Lesniak, and D. R. Lick. Generalized connectivity in graphs. 1984.
- [5] E. Cheng, Q. Ke, and Z. Shen. *Structural Properties of Generalized Exchanged Hypercubes Generalized exchanged hypercube*. Emergent Computation, 2017.
- [6] P. Cull and S. M. Larson. The mobius cubes. *IEEE Trans Computers*, 44(5):647–659, 1995.
- [7] Efe and K. A variation on the hypercube with lower diameter. *Computers IEEE Transactions on*, 40(11):1312–1316, 1991.
- [8] A. H. Esfahanian. Generalized measures of fault tolerance with application to n -cube networks. *Computers IEEE Transactions on*, 38(11):1586–1591, 1989.
- [9] J. X. Fan and L. Q. He. Bc interconnection networks and their properties. *Chinese Journal of Computers*, 2003.
- [10] F. Harary. Conditional connectivity. *Networks*, 13, 2010.
- [11] W. T. Huang, Jjm Tan, C. N. Hung, and L. H. Hsu. Fault-tolerant hamiltonicity of twisted cubes. *Journal of Parallel & Distributed Computing*, 62(4):591–604, 2002.
- [12] Jin-Xin and Zhou. On g-extra connectivity of hypercube-like networks. *Journal of Computer & System Sciences*, 2017.
- [13] H. Li and W. Yang. Bounding the size of the subgraph induced by m vertices and extra edge-connectivity of hypercubes. *Discrete Applied Mathematics*, 161(16-17):2753–2757, 2013.
- [14] E. Sampathkumar. Connectivity of a graph - a generalization. *J.combin.inform.system Sci*, 1984.
- [15] H. Shang, E. Sabir, J. Meng, and L. Guo. Characterizations of optimal component cuts of locally twisted cubes. *The Bulletin of the Malaysian Mathematical Society Series 2*, 43(3):1–17, 2019.
- [16] Z. Shen, K. E. Qiu, and E. Cheng. Connectivity results of complete cubic networks as associated with linearly many faults. *Journal of interconnection networks*, 2015.

- [17] Zhao, Shuli, Yang, Weihua, Zhang, Shurong, Xu, and Liqiong. Component edge connectivity of hypercubes. *International Journal of Foundations of Computer Science*, 2018.
- [18] S. Zhao and W. Yang. Conditional connectivity of folded hypercubes. *Discrete Applied Mathematics*, 257:388–392, 2019.