

IMPROVING THE TRANSIENT TIMES FOR DISTRIBUTED STOCHASTIC GRADIENT METHODS*

KUN HUANG[†] AND SHI PU[†]

Abstract. We consider the distributed optimization problem where n agents each possessing a local cost function, collaboratively minimize the average of the n cost functions over a connected network. Assuming stochastic gradient information is available, we study a distributed stochastic gradient algorithm, called exact diffusion with adaptive stepsizes (EDAS) adapted from the Exact Diffusion method [37] and NIDS [11] and perform a non-asymptotic convergence analysis. We not only show that EDAS asymptotically achieves the same network independent convergence rate as centralized stochastic gradient descent (SGD) for minimizing strongly convex and smooth objective functions, but also characterize the transient time needed for the algorithm to approach the asymptotic convergence rate, which behaves as $K_T = \mathcal{O}\left(\frac{n}{1-\lambda_2}\right)$, where $1-\lambda_2$ stands for the spectral gap of the mixing matrix. To the best of our knowledge, EDAS achieves the shortest transient time when the average of the n cost functions is strongly convex and each cost function is smooth. Numerical simulations further corroborate and strengthen the obtained theoretical results.

Key words. distributed optimization, convex optimization, stochastic gradient methods

AMS subject classifications. 90C15, 90C25, 68Q25

1. Introduction. In this paper, we consider the distributed optimization problem, where a set of n agents each possessing a local cost function f_i , seek to collaboratively solve the following optimization problem:

$$(1.1) \quad \min_{x \in \mathbb{R}^p} f(x) \left(= \frac{1}{n} \sum_{i=1}^n f_i(x) \right),$$

where each $f_i : \mathbb{R}^p \rightarrow \mathbb{R}$ is assumed to have Lipschitz continuous gradients, and f is strongly convex. The agents are connected over a network, where they may only communicate and exchange information with their direct neighbors.

To solve problem (1.1), we assume that at each iteration k of the algorithm we study, each agent $i \in \mathcal{N} = \{1, 2, \dots, n\}$ is able to obtain a stochastic gradient sample of the form $g_i(x_{i,k}, \xi_{i,k})$, given input $x_{i,k} \in \mathbb{R}^p$, that satisfies the following condition:

Assumption 1.1. For all $k \geq 0$, each random vector $\xi_{i,k} \in \mathbb{R}^q$ is independent across $i \in \mathcal{N}$. Denote by \mathcal{F}_k the σ -algebra generated by $\{x_{i,0}, x_{i,1}, \dots, x_{i,k} \mid i \in \mathcal{N}\}$. Then,

$$(1.2) \quad \begin{aligned} \mathbb{E}[g_i(x_{i,k}, \xi_{i,k}) \mid \mathcal{F}_k] &= \nabla f_i(x_{i,k}), \\ \mathbb{E}[\|g_i(x_{i,k}, \xi_{i,k}) - \nabla f_i(x_{i,k})\|^2 \mid \mathcal{F}_k] &\leq \sigma_i^2, \quad \text{for some } \sigma_i > 0. \end{aligned}$$

Stochastic gradients appear in many machine learning problems. For instance, suppose $f_i(x) := \mathbb{E}_{\xi_i}[F_i(x, \xi_i)]$ represents the expected loss function for agent i , where ξ_i are independent data samples gathered continuously over time. Then for any x and ξ_i , $g_i(x, \xi_i) := \nabla F_i(x, \xi_i)$ is an unbiased estimator of $\nabla f_i(x)$. For another example, let $f_i(x) := (1/|S_i|) \sum_{\zeta_j \in S_i} F(x, \zeta_j)$ denote an empirical risk function, where S_i is the

*This work was partially supported by the Shenzhen Research Institute of Big Data (SRIBD) under Fund No. J00120190011 and by the National Natural Science Foundation of China (NSFC) under grant No. 62003287.

[†]School of Data Science, Shenzhen Research Institute of Big Data, The Chinese University of Hong Kong, Shenzhen (kunhuang@link.cuhk.edu.cn, pushi@cuhk.edu.cn).

local dataset of agent i . Under this setting, the gradient estimation of $f_i(x)$ can incur noise from various sources such as sampling and discretization [10].

Problem (1.1) has been studied extensively in the literature using various distributed algorithms [32, 18, 19, 13, 6, 7, 28, 5, 26, 17, 34, 25], among which the distributed gradient descent (DGD) method proposed in [18] has drawn the greatest attention. Recently, distributed implementation of stochastic gradient algorithms has received considerable interest. Several works have shown that distributed methods may compare with their centralized counterparts under certain conditions. For example, the work in [2, 3, 4] first showed that, with a sufficiently small constant stepsize, a distributed stochastic gradient method achieves comparable performance to a centralized method in terms of the steady-state mean-square-error.

More recent works have demonstrated that various distributed stochastic gradient methods enjoy the so-called “asymptotic network independence” property, that is, the generated iterates asymptotically converge at the same network independent rate compared to centralized stochastic gradient descent (SGD) for minimizing smooth objective functions [29, 24, 23, 12, 8, 30, 33]. Nevertheless, it always takes a certain number of updates before this network independent convergence rate can be reached, which is always network dependent and referred to as *transient time* of the algorithm. For example, when considering smooth and strongly convex objective functions, the shortest transient time that has been shown so far is achieved by the distributed stochastic gradient descent (DSGD) method [23], which behaves as $\mathcal{O}\left(\frac{n}{(1-\lambda_2)^2}\right)$, where $1 - \lambda_2$ stands for the spectral gap of the mixing matrix. For general undirected networks such as ring graphs, the transient time of DSGD is as large as $\mathcal{O}(n^5)$, which can be significant for large-scale networks. Therefore, it is critical to develop new algorithms with shorter transient times.

Recently, a class of closely related algorithms, termed EXTRA [28], D^2 [30], Exact Diffusion [37, 36] and NIDS [11] have been proposed to solve problem (1.1) under both exact and stochastic gradient settings. These methods achieve linear convergence for minimizing smooth and strongly convex objective functions under exact gradient information [11]. When only stochastic gradients are available, the work in [36] compared the performance of various distributed optimization methods under constant stepsizes and demonstrated the effectiveness of Exact Diffusion compared to other distributed stochastic gradient algorithms.

In this work, we consider a distributed stochastic gradient algorithm adapted from NIDS/Exact Diffusion/ D^2 , termed EDAS (Exact Diffusion with Adaptive Stepsizes) and perform a non-asymptotic analysis. In addition to showing the asymptotic network independence property of EDAS, we carefully identify its non-asymptotic convergence rate as a function of characteristics of the objective functions and the underlying network. Moreover, we derive the transient time needed for EDAS to achieve the network independent convergence rate, which behaves as $\mathcal{O}\left(\frac{n}{1-\lambda_2}\right)$. This is the shortest transient time for minimizing strongly convex and smooth objective functions in a distributed fashion to the best of our knowledge. Numerical experiments corroborate and strengthen the theoretical findings.

1.1. Related Work. Our work is closely related to the concept of “asymptotic network independence” in distributed optimization over networks, that is, asymptotically, the distributed algorithm converges to the optimal solution at a comparable rate to a centralized algorithm with the same computational power. In addition to [2, 3, 4] discussed before, the work in [14, 15] proved that distributed stochastic ap-

proximation methods perform asymptotically as well as centralized schemes by means of a central limit theorem. The papers [20, 21] demonstrated the advantages of distributively implementing a stochastic gradient method assuming that sampling times are random and non-negligible. The first “asymptotic network independence” result appeared in [31], where it assumed that all the local functions f_i have the same minimum. A recent paper [29] discussed an algorithm that asymptotically performs as well as the best bounds on centralized stochastic gradient descent subject to possible message losses, delays, and asynchrony. In a parallel recent work [8], a similar result was presented with a further compression technique which allowed agents to save on communication. The work in [22] considered a distributed stochastic gradient tracking method (DSGT) which asymptotically performs as well as centralized stochastic gradient descent. The result was generalized to the setting of directed communication graphs in [27]. For nonconvex objective functions, the paper [12, 30] proved that decentralized algorithms can achieve a linear speedup similar to a centralized algorithm when the number of iterates k is large enough. The work in [1] further extended the theory to directed communication networks for training deep neural networks. A recent survey [24] provided a detailed discussion on this topic.

When restricted to considering strongly convex and smooth objective functions, several recent works have not only shown asymptotic network independence for the proposed algorithms, but also provided the transient times for the algorithms to reach the network independent convergence rate. Table 1 compares the transient times for some existing algorithms, which are functions of the network characteristics. In particular, the paper [23] showed that for the distributed stochastic gradient descent (DSGD) method, it takes $\mathcal{O}\left(\frac{n}{(1-\lambda_2)^2}\right)$ ($1-\lambda_2$ stands for the spectral gap of the mixing matrix) to achieve network independence. This result was also proved to be sharp. For another distributed stochastic gradient tracking (DSGT) method, the paper [33] showed the transient time is $\mathcal{O}\left(\frac{n}{(1-\lambda_2)^3}\right)$.

	DSGD with compression	DSGD & Dif- fusion	DSGT	EDAS (this paper)
Transient Time	$\mathcal{O}\left(\frac{n}{(1-\lambda_2)^4}\right)$ [8]	$\mathcal{O}\left(\frac{n}{(1-\lambda_2)^2}\right)$ [23]	$\mathcal{O}\left(\frac{n}{(1-\lambda_2)^3}\right)$ [33]	$\mathcal{O}\left(\frac{n}{1-\lambda_2}\right)$

TABLE 1

Transient times of distributed stochastic gradient methods for minimizing strongly convex and smooth objective functions.

1.2. Main Contribution. We summarize the main contribution of the paper as follows. First, we consider a distributed stochastic gradient algorithm (EDAS) adapted from NIDS/Exact Diffusion and perform a non-asymptotic analysis. For strongly convex and smooth objective functions, we show in Theorem 4.4 that EDAS asymptotically achieves the same network independent convergence rate of centralized stochastic gradient descent (SGD). We also carefully characterize the convergence rate of EDAS as a function of the characteristics of the objective functions and the communication network.

Second, by comparing the convergence result of EDAS to that of SGD, we derive the transient time needed for EDAS to achieve the network independent rate, which is upper bounded by $K_T = \mathcal{O}\left(\frac{n}{1-\lambda_2}\right)$ under mild conditions. This is the shortest

transient time so far to the best of our knowledge, and significantly improves upon the shortest transient time previously obtained, i.e., $\mathcal{O}\left(\frac{n}{(1-\lambda_2)^2}\right)$. This is the main contribution of the paper.

Finally, we provide two simulation examples that corroborate and strengthen the obtained theoretical results. For solving a “hard” problem with EDAS, the observed transient times are consistent with the theoretical upper bounds for both the ring network topology and the square grid network topology, which verifies the sharpness of the theoretical results. For the problem of classifying handwritten digits with logistic regression, the observed transient times are even better than the upper bounds. For both problems, we compare the performance of EDAS with other existing algorithms and show the superiority of EDAS.

1.3. Notation. Throughout the paper, we use column vectors if not otherwise specified. Let each agent i hold a local copy $x_{i,k} \in \mathbb{R}^p$ of the decision variable at the k -th iteration. We denote

$$\mathbf{x}_k := (x_{1,k}, x_{2,k}, \dots, x_{n,k})^\top \in \mathbb{R}^{n \times p},$$

$$\bar{x}_k := \frac{1}{n} \sum_{i=1}^k x_{i,k}.$$

We also define the following aggregative functions,

$$F(\mathbf{x}_k) := \sum_{i=1}^n f_i(x_{i,k}),$$

$$\nabla F(\mathbf{x}_k) := (\nabla f_1(x_{1,k}), \nabla f_2(x_{2,k}), \dots, \nabla f_n(x_{n,k}))^\top \in \mathbb{R}^{n \times p},$$

$$\mathbf{g}_k := (g_1(x_{1,k}, \xi_{1,k}), g_2(x_{2,k}, \xi_{2,k}), \dots, g_n(x_{n,k}, \xi_{n,k}))^\top \in \mathbb{R}^{n \times p},$$

and write

$$\bar{\nabla} F(\mathbf{x}_k) := \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}),$$

$$\bar{\sigma}^2 := \frac{1}{n} \sum_{i=1}^n \sigma_i^2,$$

$$\bar{g}_k := \frac{1}{n} \sum_{i=1}^n g_i(x_{i,k}, \xi_{i,k}).$$

For two vectors a, b of the same dimension, $\langle a, b \rangle$ denotes the inner product. For two matrices $A, B \in \mathbb{R}^{n \times p}$, we define

$$\langle A, B \rangle := \sum_{i=1}^n \langle A_i, B_i \rangle,$$

where A_i (respectively, B_i) represents the i -th row of A (respectively, B). $\|\cdot\|$ denotes ℓ_2 -norm for vectors and Frobenius norm for matrices.

Next we specify the conditions concerning the local cost functions and the communication network.

Assumption 1.2. Each $f_i : \mathbb{R}^p \rightarrow \mathbb{R}$ has L -Lipschitz continuous gradient, and $f : \mathbb{R}^p \rightarrow \mathbb{R}$ is μ -strongly convex, i.e., $\forall x, x' \in \mathbb{R}^p$,

$$\|\nabla f_i(x) - \nabla f_i(x')\| \leq L\|x - x'\|, \quad \forall i,$$

$$\langle \nabla f(x) - \nabla f(x'), x - x' \rangle \geq \mu\|x - x'\|^2.$$

Under Assumption 1.2, problem (1.1) has a unique optimal solution $x^* \in \mathbb{R}^p$.

The agents are connected through a graph $\mathcal{G} = (\mathcal{N}, \mathcal{E})$, where \mathcal{N} is the set of nodes and $\mathcal{E} \subseteq \mathcal{N} \times \mathcal{N}$ denotes the set of edges. Regarding the graph \mathcal{G} and the corresponding mixing matrix $W = [w_{ij}] \in \mathbb{R}^{n \times n}$, we make the following standing assumption.

Assumption 1.3. The graph \mathcal{G} is undirected and strongly connected. There exists a link from i and j ($i \neq j$) in \mathcal{G} if and only if $w_{ij}, w_{ji} > 0$; otherwise, $w_{ij} = w_{ji} = 0$. The mixing matrix W is nonnegative, symmetric and stochastic, i.e., $W\mathbf{1} = \mathbf{1}$. In addition, suppose the smallest eigenvalue of W satisfies $\lambda_n \geq \underline{\lambda} > 0$ for some fixed $\underline{\lambda}$ independent of n .¹

The rest of this paper is organized as follows. In section 2 we present the EDAS algorithm and its transformed error dynamics in the form of a primal-dual-like algorithm. Next we provide some supporting lemmas and prove the sublinear convergence rate of EDAS in section 3. The main convergence results of the algorithm are presented in section 4, where we derive the convergence rate of EDAS as a function of the characteristics of the objective functions and the mixing matrix, and subsequently, obtain the transient time of EDAS. Numerical experiments are provided in section 5, and we conclude the paper in section 6.

2. Algorithm. We study the following algorithm adapted from EXTRA [28], Exact Diffusion [37], NIDS [11] and D^2 [30], where the main characteristics of EDAS lie in the diminishing stepsize policy $\alpha_k = \mathcal{O}_k(\frac{1}{k})$:

Algorithm 2.1 EDAS (Exact Diffusion with Adaptive Stepsizes)

- 1: Initialize $x_{i,0} \in \mathbb{R}^p$ for all agent $i \in \mathcal{N}$, determine $W = [w_{ij}] \in \mathbb{R}^{n \times n}$, stepsize $\alpha_k = \frac{\theta}{\mu(k+m)}$ for some $\theta > 0$ and $m > 0$.
 - 2: **for** $k = 0, 1, 2, \dots$ **do**
 - 3: **for** $i = 1, 2, \dots, n$ **do**
 - 4: Agent i acquires a stochastic gradient $g_i(x_{i,k}, \xi_{i,k}) \in \mathbb{R}^p$.
 - 5: **if** $k = 0$ **then**
 - 6: $x_{i,k+1} = \sum_{j=1}^n w_{ij} (x_{j,k} - \alpha_k g_j(x_{j,k}, \xi_{j,k}))$
 - 7: **else**
 - 8: $x_{i,k+1} = \sum_{j=1}^n w_{ij} (2x_{j,k} - x_{j,k-1} - \alpha_k g_j(x_{j,k}, \xi_{j,k}) + \alpha_{k-1} g_j(x_{j,k-1}, \xi_{j,k-1}))$
 - 9: **end if**
 - 10: **end for**
 - 11: **end for**
-

¹To satisfy such an assumption, it is convenient to let the mixing matrix W be chosen as $W = \beta I_n + (1 - \beta)\tilde{W}$ for some $\beta \in (\frac{1}{2}, 1)$ with \tilde{W} being nonnegative, symmetric and stochastic. Therefore, the condition is not restrictive.

Using the notation in [subsection 1.3](#), EDAS can be written in the following compact form:

$$(2.1a) \quad \mathbf{x}_1 = W(\mathbf{x}_0 - \alpha_0 \mathbf{g}_0),$$

$$(2.1b) \quad \mathbf{x}_{k+1} = W(2\mathbf{x}_k - \mathbf{x}_{k-1} - \alpha_k \mathbf{g}_k + \alpha_{k-1} \mathbf{g}_{k-1}), \quad \forall k \geq 1.$$

For further investigation, note that since W is nonnegative, symmetric and stochastic, it has a spectral decomposition given by

$$(2.2) \quad W = Q\Lambda Q^\top,$$

where

$$\Lambda := \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n),$$

with $1 = \lambda_1 > \lambda_2 \geq \dots \geq \lambda_n \geq -1$. Thus $I_n - W = Q(I_n - \Lambda)Q^\top$.

Let $V := (I_n - W)^{1/2}$. EDAS can be rewritten in the following equivalent form that resembles a primal-dual algorithm (see [\[37\]](#)): let $\mathbf{y}_0 = \mathbf{0}$, for $k \geq 0$,

$$(2.3a) \quad \mathbf{x}_{k+1} = W(\mathbf{x}_k - \alpha_k \mathbf{g}_k) - V\mathbf{y}_k,$$

$$(2.3b) \quad \mathbf{y}_{k+1} = \mathbf{y}_k + V\mathbf{x}_{k+1}.$$

To verify the equivalence relationship, note that from [\(2.3\)](#), when $k \geq 1$, we have

$$\begin{aligned} \mathbf{x}_{k+1} - \mathbf{x}_k &= W(\mathbf{x}_k - \mathbf{x}_{k-1} - \alpha_k \mathbf{g}_k + \alpha_{k-1} \mathbf{g}_{k-1}) - V\mathbf{y}_k + V\mathbf{y}_{k-1} \\ &= W(\mathbf{x}_k - \mathbf{x}_{k-1} - \alpha_k \mathbf{g}_k + \alpha_{k-1} \mathbf{g}_{k-1}) - V(\mathbf{y}_k - \mathbf{y}_{k-1}) \\ &= W(2\mathbf{x}_k - \mathbf{x}_{k-1} - \alpha_k \mathbf{g}_k + \alpha_{k-1} \mathbf{g}_{k-1}) - \mathbf{x}_k. \end{aligned}$$

2.1. Optimality Condition. To facilitate the analysis of EDAS, we consider the optimality condition for problem [\(1.1\)](#). From the condition we are able to construct the error dynamics for [Algorithm 2.1](#).

LEMMA 2.1 (Optimality Condition [\[35\]](#)). *Under [Assumptions 1.2 and 1.3](#), if there exists some $(\mathbf{x}^*, \mathbf{y}_k^*)$ that satisfies:²*

$$(2.4a) \quad \alpha_k W \nabla F(\mathbf{x}^*) + V\mathbf{y}_k^* = \mathbf{0}$$

$$(2.4b) \quad V\mathbf{x}^* = \mathbf{0},$$

where $\mathbf{x}^* := (x_1^*, x_2^*, \dots, x_n^*)^\top \in \mathbb{R}^{n \times p}$, then it holds that

$$x_1^* = x_2^* = \dots = x_n^* = x^*.$$

Proof. We follow the proof of Lemma 2 in [\[37\]](#), where the stepsize α_k was assumed to be a constant. First, since $\text{null}(V) = \text{span}\{\mathbf{1}_n\}$, we have

$$(2.5) \quad \mathbf{0} = V\mathbf{x}^* \iff \mathbf{x}^* \in \text{null}(V),$$

and hence $x_1^* = x_2^* = \dots = x_n^*$.

Additionally, multiplying $\mathbf{1}^\top$ on both sides of [\(2.4a\)](#) and noticing that $\mathbf{1}^\top V = \mathbf{0}$, we have

$$(2.6) \quad \alpha_k \mathbf{1}^\top W \nabla F(\mathbf{x}^*) = \alpha_k \mathbf{1}^\top \nabla F(\mathbf{x}^*) = \alpha_k \sum_{i=1}^n \nabla f_i(x_i^*) = 0,$$

²Note that \mathbf{y}_k^* depends on k .

where we use the column stochastic property of W , i.e., $\mathbf{1}^\top W = \mathbf{1}^\top$. Relation $\sum_{i=1}^n \nabla f_i(x_i^*) = 0$ then implies that $x_1^* = x_2^* = \dots = x_n^*$ must coincide with the minimizer x^* of problem (1.1). \square

We next show the existence of $(\mathbf{x}^*, \mathbf{y}_k^*)$ and how to select a unique pair of $(\mathbf{x}^*, \mathbf{y}_k^*)$ in light of the singularity of V .

LEMMA 2.2. *Under Assumptions 1.2 and 1.3, there exists $(\mathbf{x}^*, \mathbf{y}_k^*)$ that satisfies (2.4). In particular, we can choose*

$$(2.7a) \quad \mathbf{x}^* = \mathbf{1}(x^*)^\top,$$

$$(2.7b) \quad \mathbf{y}_k^* = -V^-(\alpha_k W \nabla F(\mathbf{x}^*)),$$

where V^- is the pseudoinverse of V given by

$$V^- = Q \left(\sqrt{I_n - \Lambda} \right)^- Q^\top,$$

in which Q is the same orthogonal matrix as in decomposing W , and $(\sqrt{I_n - \Lambda})^- := \text{diag}(d_1, d_2, \dots, d_n)$ is the pseudoinverse of $\sqrt{I_n - \Lambda}$ with

$$d_i = \begin{cases} \frac{1}{\sqrt{1-\lambda_i}} & \text{if } 1 - \lambda_i \neq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The proof for the existence of $(\mathbf{x}^*, \mathbf{y}_k^*)$ is inspired by Lemma 3 in [37]. Let $\mathbf{x}^* = \mathbf{1}(x^*)^\top$. Then noticing that $V\mathbf{1}_n = \mathbf{0}$, we have $V\mathbf{x}^* = \mathbf{0}$. To prove that \mathbf{y}_k^* exists for all $k \geq 0$ is equivalent to showing that the following linear system is consistent w.r.t. \mathbf{y}_k^* :

$$(2.8) \quad V\mathbf{y}_k^* = -\alpha_k W \nabla F(\mathbf{x}^*).$$

Hence we need only show that $-\alpha_k W \nabla F(\mathbf{x}^*)$ lies in $\text{range}(V)$ for all $k \geq 0$. By Assumption 1.3,

$$\mathbf{1}^\top (-\alpha_k W \nabla F(\mathbf{x}^*)) = -\alpha_k \sum_{i=1}^n f_i(x^*) = 0,$$

we have

$$\begin{aligned} & (-\alpha_k W \nabla F(\mathbf{x}^*)) \perp \text{span}\{\mathbf{1}\} = \text{null}(V), \\ \Leftrightarrow & (-\alpha_k W \nabla F(\mathbf{x}^*)) \in \text{range}(V^\top) = \text{range}(V), \end{aligned} \quad \square$$

which proves the existence of \mathbf{y}_k^* .

We next verify that $\mathbf{y}_k^* = -V^-(\alpha_k W \nabla F(\mathbf{x}^*))$ satisfies (2.8). First of all, it is easy to see that the choice of V^- is indeed a pseudoinverse of V . Then since the linear system (2.8) is consistent, we have that $V^-(\alpha_k W \nabla F(\mathbf{x}^*))$ is a solution of (2.8), or $V\mathbf{y}_k^* = -\alpha_k W \nabla F(\mathbf{x}^*)$.

Remark 2.3. Throughout the paper, we let $(\mathbf{x}^*, \mathbf{y}_k^*)$ be the unique pair given in (2.7).

2.2. Preliminary Analysis. In light of the optimality condition of problem (1.1) given in Lemma 2.1, we consider the following error dynamics as the starting point of the convergence analysis for EDAS.

LEMMA 2.4. Denote $\tilde{\mathbf{x}}_k := \mathbf{x}_k - \mathbf{x}^*$, $\tilde{\mathbf{y}}_k := \mathbf{y}_k - \mathbf{y}_k^*$ and $\mathbf{s}_k := \nabla F(\mathbf{x}_k) - \mathbf{g}_k$. We have

$$(2.9) \quad \begin{pmatrix} \tilde{\mathbf{x}}_{k+1} \\ \tilde{\mathbf{y}}_{k+1} \end{pmatrix} = B \begin{pmatrix} \tilde{\mathbf{x}}_k + \alpha_k(\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \\ \tilde{\mathbf{y}}_k \end{pmatrix} + \alpha_k B \begin{pmatrix} \mathbf{s}_k \\ \mathbf{0} \end{pmatrix} + \begin{pmatrix} \mathbf{0} \\ \mathbf{y}_k^* - \mathbf{y}_{k+1}^* \end{pmatrix},$$

where

$$(2.10) \quad B := \begin{pmatrix} W & -V \\ VW & W \end{pmatrix}.$$

Proof. From relation (2.3) and the definitions of $\tilde{\mathbf{x}}_k$, $\tilde{\mathbf{y}}_k$ and \mathbf{s}_k , we have

$$(2.11) \quad \begin{aligned} \tilde{\mathbf{x}}_{k+1} &= \mathbf{x}_{k+1} - \mathbf{x}^* \\ &= W(\mathbf{x}_k - \alpha_k \mathbf{g}_k) - V\mathbf{y}_k - \mathbf{x}^* \\ &= W[\tilde{\mathbf{x}}_k + \alpha_k(\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k))] + \alpha_k W\mathbf{s}_k - \alpha_k W\nabla F(\mathbf{x}^*) - V\mathbf{y}_k \\ &= W[\tilde{\mathbf{x}}_k + \alpha_k(\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k))] - V\tilde{\mathbf{y}}_k + \alpha_k W\mathbf{s}_k, \end{aligned}$$

and

$$(2.12) \quad \tilde{\mathbf{y}}_{k+1} = \mathbf{y}_{k+1} - \mathbf{y}_{k+1}^* = \mathbf{y}_k + V\mathbf{x}_{k+1} - \mathbf{y}_{k+1}^* = \tilde{\mathbf{y}}_k + V\tilde{\mathbf{x}}_{k+1} + (\mathbf{y}_k^* - \mathbf{y}_{k+1}^*).$$

Substituting (2.11) into relation (2.12), we obtain

$$(2.13) \quad \begin{pmatrix} \tilde{\mathbf{x}}_{k+1} \\ \tilde{\mathbf{y}}_{k+1} \end{pmatrix} = \begin{pmatrix} W & -V \\ VW & W \end{pmatrix} \begin{pmatrix} \tilde{\mathbf{x}}_k + \alpha_k(\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \\ \tilde{\mathbf{y}}_k \end{pmatrix} + \alpha_k \begin{pmatrix} W & -V \\ VW & W \end{pmatrix} \begin{pmatrix} \mathbf{s}_k \\ \mathbf{0} \end{pmatrix} + \begin{pmatrix} \mathbf{0} \\ \mathbf{y}_k^* - \mathbf{y}_{k+1}^* \end{pmatrix}. \quad \square$$

In the next lemma, we show the matrix $B \in \mathbb{R}^{2n \times 2n}$ has an eigendecomposition in the form of $B = UDU^{-1}$, inspired by the arguments in [37]. In light of the decomposition, we are able to derive the transformed error dynamics for EDAS, which is then used for performing the non-asymptotic analysis for EDAS in Section 3.

LEMMA 2.5. Under Assumption 1.3, the matrix B has an eigendecomposition given by

$$B = UDU^{-1} := \begin{pmatrix} \mathbf{1}_n & \mathbf{0} & cU_{R,u} \\ \mathbf{0} & \mathbf{1}_n & cU_{R,l} \end{pmatrix} \text{diag}(1, 1, D_1) \begin{pmatrix} \frac{1}{n}\mathbf{1}_n^\top & \mathbf{0} \\ \mathbf{0} & \frac{1}{n}\mathbf{1}_n^\top \\ \frac{1}{c}U_{L,l} & \frac{1}{c}U_{L,r} \end{pmatrix}$$

where D_1 is a diagonal matrix with complex entries, matrices $U_{R,u}, U_{R,l} \in \mathbb{R}^{n \times (n-2)}$, $U_{L,l}, U_{L,r} \in \mathbb{R}^{(n-2) \times n}$, and $c > 0$ is an arbitrary scaling parameter.

Denote

$$U_R := \begin{pmatrix} U_{R,u} \\ U_{R,l} \end{pmatrix}, \quad U_L := (U_{L,l} \quad U_{L,r}).$$

Then the product $\|U_L\|_2^2\|U_R\|_2^2$ has an upper bound that is independent of n . More specifically,

$$\|U_L\|_2^2\|U_R\|_2^2 \leq \frac{1}{\lambda_n} \leq \frac{1}{\underline{\lambda}}.$$

Proof. see [Appendix A.1](#). \square

Based on [Lemma 2.5](#), we derive the transformed error dynamics for EDAS by multiplying U^{-1} on both sides of the error dynamics (2.9). In this way, we decompose the error $(\tilde{\mathbf{x}}_k, \tilde{\mathbf{y}}_k)$ into two parts: \bar{z}_k that measures the difference between the average iterate and x^* , and \tilde{z}_k that represents the remaining error. Then we further study the relationship between \bar{z}_k and \tilde{z}_k in [section 3](#).

LEMMA 2.6. *Under [Assumptions 1.2](#) and [1.3](#), the transformed error dynamics for [Algorithm 2.1](#) is given by*

$$(2.14) \quad \begin{pmatrix} \bar{z}_{k+1} \\ \tilde{z}_{k+1} \end{pmatrix} = \begin{pmatrix} \bar{z}_k + \frac{\alpha_k}{n} \mathbf{1}_n^\top (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \\ D_1 \tilde{z}_k + \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \end{pmatrix} \\ + \alpha_k \begin{pmatrix} \frac{1}{n} \mathbf{1}_n^\top \mathbf{s}_k \\ \frac{1}{c} D_1 U_{L,l} \mathbf{s}_k \end{pmatrix} + \begin{pmatrix} 0 \\ \frac{1}{c} U_{L,r} (\mathbf{y}_k^* - \mathbf{y}_{k+1}^*) \end{pmatrix},$$

where

$$(2.15) \quad \begin{pmatrix} \bar{z}_k \\ \tilde{z}_k \end{pmatrix} := \begin{pmatrix} \frac{1}{n} \mathbf{1}_n^\top \tilde{\mathbf{x}}_k \\ \frac{1}{c} (U_{L,l} \tilde{\mathbf{x}}_k + U_{L,r} \tilde{\mathbf{y}}_k) \end{pmatrix} \quad \forall k.$$

Proof. Denote

$$(2.16) \quad \begin{pmatrix} \bar{z}_k \\ \hat{z}_k \\ \tilde{z}_k \end{pmatrix} := U^{-1} \begin{pmatrix} \tilde{\mathbf{x}}_k \\ \tilde{\mathbf{y}}_k \end{pmatrix} = \begin{pmatrix} \frac{1}{n} \mathbf{1}_n^\top \tilde{\mathbf{x}}_k \\ \frac{1}{n} \mathbf{1}_n^\top \tilde{\mathbf{y}}_k \\ \frac{1}{c} U_{L,l} \tilde{\mathbf{x}}_k + \frac{1}{c} U_{L,r} \tilde{\mathbf{y}}_k \end{pmatrix},$$

Multiplying U^{-1} on both sides of (2.9) leads to

$$\begin{pmatrix} \bar{z}_{k+1} \\ \hat{z}_{k+1} \\ \tilde{z}_{k+1} \end{pmatrix} = DU^{-1} \begin{pmatrix} \tilde{\mathbf{x}}_k + \alpha_k (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \\ \tilde{\mathbf{y}}_k \end{pmatrix} + \alpha_k DU^{-1} \begin{pmatrix} \mathbf{s}_k \\ \mathbf{0} \end{pmatrix} \\ + U^{-1} \begin{pmatrix} \mathbf{0} \\ \mathbf{y}_k^* - \mathbf{y}_{k+1}^* \end{pmatrix} \\ = \begin{pmatrix} \bar{z}_k + \frac{\alpha_k}{n} \mathbf{1}_n^\top (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \\ \hat{z}_k \\ D_1 \tilde{z}_k + \frac{1}{c} D_1 U_{L,l} \alpha_k (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \end{pmatrix} + \begin{pmatrix} \frac{\alpha_k}{n} \mathbf{1}_n^\top \mathbf{s}_k \\ \mathbf{0} \\ \frac{\alpha_k}{c} D_1 U_{L,l} \mathbf{s}_k \end{pmatrix} \\ + \begin{pmatrix} \mathbf{0} \\ \frac{1}{n} \mathbf{1}_n^\top (\mathbf{y}_k^* - \mathbf{y}_{k+1}^*) \\ \frac{1}{c} U_{L,r} (\mathbf{y}_k^* - \mathbf{y}_{k+1}^*) \end{pmatrix}. \quad \square$$

3. Analysis. In this section, we provide some useful supporting lemmas based on [Lemma 2.6](#), and then prove the sublinear convergence rate of EDAS in [Lemma 3.7](#). For the proof of sublinear convergence, we first propose [Lemmas 3.4](#) and [3.5](#) to derive two coupled recursions for $\mathbb{E}[\|\bar{z}_k\|^2]$ and $\mathbb{E}[\|\tilde{z}_k\|^2]$ respectively according to [Lemma 2.6](#).

Then we introduce a Lyapunov function $H_k := \mathbb{E}[\|\bar{z}_k\|^2] + \omega_k \mathbb{E}[\|\tilde{z}_k\|^2]$, and by constructing the recursion for H_k , we derive the sublinear convergence rate.

3.1. Supporting Lemmas. In this part, we provide two important coupled recursions regarding $\mathbb{E} [\|\bar{z}_k\|^2]$ and $\mathbb{E} [\|\check{z}_k\|^2]$, respectively. Before introducing these results, we state some preliminary lemmas.

LEMMA 3.1. *Suppose [Assumption 1.1](#) holds, then*

$$(3.1a) \quad \mathbb{E} [\bar{s}_k | \mathcal{F}_k] = \mathbf{0},$$

$$(3.1b) \quad \mathbb{E} [\|\bar{s}_k\|^2 | \mathcal{F}_k] \leq \frac{\bar{\sigma}^2}{n}.$$

The results of [Lemma 3.1](#) directly come from [Assumption 1.1](#).

LEMMA 3.2. *Under [Assumption 1.2](#), there holds*

$$\|\bar{\nabla} F(\mathbf{1}\bar{x}_k^\top) - \bar{\nabla} F(\mathbf{x}_k)\| \leq \frac{L}{\sqrt{n}} \|cU_{R,u}\check{z}_k\|.$$

Proof. By definition,

$$(3.2) \quad \|\bar{\nabla} F(\mathbf{1}\bar{x}_k^\top) - \bar{\nabla} F(\mathbf{x}_k)\| = \left\| \frac{1}{n} \sum_{i=1}^n (\nabla f_i(\bar{x}_k) - \nabla f_i(x_{i,k})) \right\|$$

$$\leq \frac{L}{n} \sum_{i=1}^n \|\bar{x}_k - x_{i,k}\|$$

$$= \frac{L}{n} \sum_{i=1}^n \|\bar{x}_k - x^* + x^* - x_{i,k}\|$$

$$(3.3) \quad \leq \frac{L}{\sqrt{n}} \|\tilde{\mathbf{x}}_k - \mathbf{1}\bar{z}_k\|$$

$$(3.4) \quad = \frac{L}{\sqrt{n}} \|cU_{R,u}\check{z}_k\|,$$

where (3.2) holds because of [Assumption 1.2](#), inequality (3.3) results from Cauchy-Schwarz's inequality, and (3.4) comes from

$$(3.5) \quad \begin{pmatrix} \tilde{\mathbf{x}}_k \\ \tilde{\mathbf{y}}_k \end{pmatrix} = U \begin{pmatrix} \bar{z}_k \\ \hat{z}_k \\ \check{z}_k \end{pmatrix},$$

which implies $\tilde{\mathbf{x}}_k = \mathbf{1}\bar{z}_k + cU_{R,u}\check{z}_k$. \square

In light of Lemma 10 in [\[26\]](#), we have the following contraction result:

LEMMA 3.3. *Under [Assumption 1.2](#), suppose $\alpha_k \leq \frac{2}{\mu+L}$. There holds*

$$(3.6) \quad \|\bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k)\| \leq (1 - \alpha_k \mu) \|\bar{z}_k\|.$$

With the above results in hand, we provide an upper bound for $\mathbb{E} [\|\bar{z}_k\|^2]$ in [Lemma 3.4](#), following [Lemma 2.6](#).

LEMMA 3.4. *Under [Algorithm 2.1](#) with [Assumptions 1.1](#) to [1.3](#), suppose $\alpha_k \leq \min\{\frac{1}{3\mu}, \frac{2}{\mu+L}\}$, we have*

$$(3.7) \quad \mathbb{E} [\|\bar{z}_{k+1}\|^2] \leq (1 - \frac{3}{2}\alpha_k \mu) \mathbb{E} [\|\bar{z}_k\|^2] + \frac{3\alpha_k L^2 c^2 \|U_R\|^2}{\mu n} \mathbb{E} [\|\check{z}_k\|] + \frac{\alpha_k^2 \bar{\sigma}^2}{n}.$$

Proof. Denote $\bar{s}_k := \frac{1}{n} \sum_{i=1}^n \mathbf{s}_{i,k}^\top$, where $s_{i,k}$ is the i -th row of \mathbf{s}_k . We have from Lemma 2.6 that

$$\begin{aligned}\bar{z}_{k+1} &= \bar{z}_k + \alpha_k (\bar{\nabla} F(\mathbf{x}^*) - \bar{\nabla} F(\mathbf{x}_k) + \bar{s}_k) \\ &= \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{x}_k) + \alpha_k \bar{s}_k,\end{aligned}$$

where the second equality follows from $\bar{\nabla} F(\mathbf{x}^*) = \sum_{i=1}^n \nabla f_i(x^*) = 0$ in light of the optimality of x^* . Hence

$$\begin{aligned}\mathbb{E} \left[\|\bar{z}_{k+1}\|^2 \mid \mathcal{F}_k \right] &= \mathbb{E} \left[\left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{x}_k) + \alpha_k \bar{s}_k \right\|^2 \mid \mathcal{F}_k \right] \\ (3.8) \quad &= \left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{x}_k) \right\|^2 + \alpha_k^2 \mathbb{E} \left[\|\bar{s}_k\|^2 \mid \mathcal{F}_k \right].\end{aligned}$$

For the first term on the right hand side of (3.8), we have

$$\begin{aligned}\left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{x}_k) \right\|^2 &= \left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k) + \alpha_k (\bar{\nabla} F(\mathbf{1}\bar{x}_k) - \bar{\nabla} F(\mathbf{x}_k)) \right\|^2 \\ &= \left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k) \right\|^2 + \alpha_k^2 \left\| \bar{\nabla} F(\mathbf{1}\bar{x}_k) - \bar{\nabla} F(\mathbf{x}_k) \right\|^2 \\ &\quad + 2\alpha_k \langle \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k), \bar{\nabla} F(\mathbf{1}\bar{x}_k) - \bar{\nabla} F(\mathbf{x}_k) \rangle \\ (3.9) \quad &\leq \left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k) \right\|^2 + \alpha_k^2 \left\| \bar{\nabla} F(\mathbf{1}\bar{x}_k) - \bar{\nabla} F(\mathbf{x}_k) \right\|^2 \\ &\quad + 2\alpha_k \left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k) \right\| \cdot \left\| \bar{\nabla} F(\mathbf{1}\bar{x}_k) - \bar{\nabla} F(\mathbf{x}_k) \right\| \\ (3.10) \quad &\leq (1 + \gamma_1) \left\| \bar{z}_k - \alpha_k \bar{\nabla} F(\mathbf{1}\bar{x}_k) \right\|^2 + \left(1 + \frac{1}{\gamma_1}\right) \alpha_k^2 \left\| \bar{\nabla} F(\mathbf{1}\bar{x}_k) - \bar{\nabla} F(\mathbf{x}_k) \right\|^2,\end{aligned}$$

where (3.9) comes from Cauchy-Schwarz's inequality and (3.10) holds because $2\|a\|\|b\| \leq \gamma_1 \|a\|^2 + \frac{1}{\gamma_1} \|b\|^2, \forall \gamma_1 > 0$. Substituting (3.10) into (3.8) and taking full expectation on both sides of (3.8), we get

$$(3.11) \quad \mathbb{E} \left[\|\bar{z}_{k+1}\|^2 \right] \leq (1 + \gamma_1)(1 - \alpha_k \mu)^2 \mathbb{E} \left[\|\bar{z}_k\|^2 \right] + (1 + \gamma_1^{-1}) \frac{\alpha_k^2 L^2 c^2 \|U_R\|^2}{n} \mathbb{E} \left[\|\bar{z}_k\|^2 \right] + \frac{\alpha_k^2 \bar{\sigma}^2}{n},$$

where we invoked relations (3.6), (3.1b), and Lemma 3.2. Let $\gamma_1 = \frac{3}{8} \alpha_k \mu$ and notice that $\alpha_k \leq \frac{1}{3\mu}$. We conclude from (3.11) that

$$E \left[\|\bar{z}_{k+1}\|^2 \right] \leq \left(1 - \frac{3}{2} \alpha_k \mu\right) \mathbb{E} \left[\|\bar{z}_k\|^2 \right] + \frac{3\alpha_k L^2 c^2 \|U_R\|^2}{\mu n} \mathbb{E} \left[\|\bar{z}_k\|^2 \right] + \frac{\alpha_k^2 \bar{\sigma}^2}{n}.$$

This finishes the proof. \square

We obtain the recursion for $\mathbb{E} [\|\bar{z}_{k+1}\|^2]$ stated in Lemma 3.5 below.

LEMMA 3.5. Under Algorithm 2.1 with Assumptions 1.1 to 1.3, suppose

$$\alpha_k \leq \frac{1 - \sqrt{\lambda_2}}{2 \|U_L\| \|U_R\| L}$$

where λ_2 is the second largest eigenvalue of W . Then for all $k \geq 1$, we have

$$\begin{aligned}\mathbb{E} [\|\bar{z}_{k+1}\|^2] &\leq \frac{3 + \sqrt{\lambda_2}}{4} \mathbb{E} [\|\bar{z}_k\|^2] + \frac{\alpha_k^2 \lambda_2 n \|U_L\|^2}{c^2} \left(\frac{4L^2}{1 - \lambda_2} \mathbb{E} [\|\bar{z}_k\|^2] + \bar{\sigma}^2 \right) \\ &\quad + \frac{4\|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{c^2 (1 - \sqrt{\lambda_2})} \|\alpha_{k+1} - \alpha_k\|^2\end{aligned}$$

Proof. By squaring and taking conditional expectation on both sides of the second recursion in [Lemma 2.6](#), we obtain

$$\begin{aligned}
\mathbb{E} [\|\tilde{z}_{k+1}\|^2 | \mathcal{F}_k] &= \|D_1 \tilde{z}_k + \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k))\|^2 \\
&\quad + \frac{\alpha_k^2 \|D_1 U_{L,l}\|^2}{c^2} \mathbb{E} [\|\mathbf{s}_k\|^2 | \mathcal{F}_k] + \frac{1}{c^2} \|U_{L,r}\|^2 \|\mathbf{y}_{k+1}^* - \mathbf{y}_k^*\|^2 \\
&\quad + 2 \left\langle D_1 \tilde{z}_k + \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)), \frac{1}{c} U_{L,r} (\mathbf{y}_k^* - \mathbf{y}_{k+1}^*) \right\rangle \\
&\leq (1 + \gamma) \|D_1 \tilde{z}_k + \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k))\|^2 \\
&\quad + \frac{\alpha_k^2 \|D_1\|^2 \|U_L\|^2}{c^2} \mathbb{E} [\|\mathbf{s}_k\|^2 | \mathcal{F}_k] + \left(1 + \frac{1}{\gamma}\right) \frac{\|U_L\|^2}{c^2} \|\mathbf{y}_{k+1}^* - \mathbf{y}_k^*\|^2,
\end{aligned} \tag{3.12}$$

where $\gamma > 0$ is arbitrary. To further bound the right hand side of (3.12) in terms of $\|\tilde{z}_k\|^2$ and $\|\tilde{z}_k\|^2$, first consider

$$\begin{aligned}
&\|D_1 \tilde{z}_k + \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k))\|^2 \\
&\leq (1 + \gamma_2) \|D_1 \tilde{z}_k\|^2 + (1 + \gamma_2^{-1}) \left\| \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)) \right\|^2
\end{aligned} \tag{3.13}$$

where $\gamma_2 > 0$ is arbitrary. Since $\lambda_2 := |\lambda_2(W)|$, it can be seen from (A.4) that

$$\|D_1\|_2 = \sqrt{|\lambda_2(W)|} = \sqrt{\lambda_2}.$$

Take $\gamma_2 = 1/\sqrt{\lambda_2} - 1$, and notice that

$$\begin{aligned}
\|\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)\|^2 &= \sum_{i=1}^n \|\nabla f_i(x_{i,k}) - \nabla f_i(x_i^*)\|^2 \\
&\leq L^2 \sum_{i=1}^n \|x_{i,k} - x_i^*\|^2 \\
&= L^2 \|\tilde{\mathbf{x}}_k\|^2 = L^2 \|\mathbf{1} \tilde{z}_k + c U_{R,u} \tilde{z}_k\|^2 \\
&\leq L^2 (2n \|\tilde{z}_k\|^2 + 2c^2 \|U_R\|^2 \|\tilde{z}_k\|^2).
\end{aligned} \tag{3.14}$$

Relation (3.13) leads to

$$\begin{aligned}
&\|D_1 \tilde{z}_k + \frac{\alpha_k}{c} D_1 U_{L,l} (\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k))\|^2 \\
&\leq \sqrt{\lambda_2} \|\tilde{z}_k\|^2 + \frac{\alpha_k^2 \lambda_2}{c^2 (1 - \sqrt{\lambda_2})} \|U_{L,l}\|^2 \|\nabla F(\mathbf{x}^*) - \nabla F(\mathbf{x}_k)\|^2 \\
&\leq \sqrt{\lambda_2} \|\tilde{z}_k\|^2 + \frac{\alpha_k^2 \lambda_2}{c^2 (1 - \sqrt{\lambda_2})} \|U_{L,l}\|^2 L^2 (2n \|\tilde{z}_k\|^2 + 2c^2 \|U_R\|^2 \|\tilde{z}_k\|^2) \\
&\leq \left(\sqrt{\lambda_2} + \frac{2\alpha_k^2 \lambda_2 \|U_L\|^2 \|U_R\|^2 L^2}{1 - \sqrt{\lambda_2}} \right) \|\tilde{z}_k\|^2 + \frac{2n\alpha_k^2 \lambda_2 \|U_L\|^2 L^2}{c^2 (1 - \sqrt{\lambda_2})} \|\tilde{z}\|^2.
\end{aligned} \tag{3.15}$$

We then bound the term $\|\mathbf{y}_k^* - \mathbf{y}_{k+1}^*\|$ on the right hand side of (3.12). As stated in [Lemma 2.2](#), $\mathbf{y}_k^* = -V^- \alpha_k W \nabla F(\mathbf{x}^*)$ where V^- is the pseudoinverse of V . Then,

$$\begin{aligned}
\|\mathbf{y}_{k+1}^* - \mathbf{y}_k^*\| &= \|V^- W \nabla F(\mathbf{x}^*) (\alpha_{k+1} - \alpha_k)\| \\
&\leq \|V^-\| \|\nabla F(\mathbf{x}^*)\| \|\alpha_{k+1} - \alpha_k\|
\end{aligned} \tag{3.16}$$

where (3.16) holds since $\|W\| = 1$.

Substituting (3.15) and (3.16) into (3.12), and taking full expectation on both sides of the inequality, we obtain

$$(3.17) \quad \begin{aligned} \mathbb{E} [\|\tilde{z}_{k+1}\|^2] &\leq (1 + \gamma) \left(\sqrt{\lambda_2} + \frac{2\alpha_k^2 \lambda_2 \|U_L\|^2 \|U_R\|^2 L^2}{1 - \sqrt{\lambda_2}} \right) \mathbb{E} [\|\tilde{z}_k\|^2] + \frac{\alpha_k^2 \lambda_2 \|U_L\|^2 n \bar{\sigma}^2}{c^2} \\ &\quad + (1 + \gamma) \frac{2n\alpha_k^2 \lambda_2 \|U_L\|^2 L^2}{c^2(1 - \sqrt{\lambda_2})} \mathbb{E} [\|\bar{z}\|^2] \\ &\quad + (1 + \frac{1}{\gamma}) \frac{\|U_L\|^2}{c^2} \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2 \|\alpha_{k+1} - \alpha_k\|^2. \end{aligned}$$

Given that $\alpha_k \leq \frac{1 - \sqrt{\lambda_2}}{2\|U_L\| \|U_R\| L}$, we have

$$\sqrt{\lambda_2} + \frac{2\alpha_k^2 \lambda_2 \|U_L\|^2 \|U_R\|^2 L^2}{1 - \sqrt{\lambda_2}} \leq \frac{\sqrt{\lambda_2} + 1}{2}.$$

Further choosing $\gamma = \frac{1 - \sqrt{\lambda_2}}{2\sqrt{\lambda_2} + 2}$ such that

$$(3.18) \quad (1 + \gamma) \frac{(\sqrt{\lambda_2} + 1)}{2} = \frac{(\sqrt{\lambda_2} + 3)}{4}.$$

We obtain the desired result. \square

3.2. Preliminary Convergence Results. We now prove the sublinear convergence rate of the EDAS algorithm under the diminishing stepsizes policy

$$\alpha_k = \frac{\theta}{\mu(k + m)},$$

where $\theta, m > 0$ are to be determined later. For simplicity, we let

$$(3.19) \quad M_k := \mathbb{E} [\|\bar{z}_k\|^2], \quad T_k := \mathbb{E} [\|\tilde{z}_k\|^2], \quad \forall k \geq 0.$$

Our analysis builds upon constructing a Lyapunov function (3.20), which is inspired by the arguments in [23]. The sublinear convergence results will then be combined with Lemma 3.5 and Lemma 3.4 respectively, so as to derive the improved convergence rates for M_k and T_k which will be stated in the next section.

The following result comes from Lemma 4.1 in [23] and will be used for bounding specific terms in the proofs. It will be used repeatedly in the analysis.

LEMMA 3.6. $\forall 1 < a < k (a \in \mathbb{N})$ and $1 < \gamma < a/2$,

$$\frac{a^{2\gamma}}{k^{2\gamma}} \leq \prod_{t=a}^{k-1} \left(1 - \frac{\gamma}{t} \right) \leq \frac{a^\gamma}{k^\gamma}$$

The Lyapunov function is defined as

$$(3.20) \quad H_k := M_k + \omega_k T_k, \quad \forall k \geq 0,$$

where

$$(3.21) \quad \omega_k := \frac{24\alpha_k c^2 \|U_R\|^2 L^2}{n\mu(1 - \sqrt{\lambda_2})}.$$

It will be made clear why ω_k takes this particular form in the proof of Lemma 3.7.

LEMMA 3.7. *Suppose*

$$(3.22) \quad m \geq \max \left\{ \frac{24\theta}{1 - \sqrt{\lambda_2}}, \frac{24\theta L^2 \|U_R\| \|U_L\|}{\mu^2(1 - \sqrt{\lambda_2})} \right\}, \quad \theta > 4.$$

Under Algorithm 2.1 with Assumptions 1.1 to 1.3, for all $k \geq 1$, we have

$$(3.23) \quad M_k \leq \frac{\hat{H}_1}{k + m} + \frac{\hat{H}_2}{(k + m)^2},$$

where

$$\hat{H}_1 := \frac{1}{\theta - 3} \left(p_2 + \frac{p_5}{m^3} \right), \quad \hat{H}_2 := m^2 H_0 + \frac{2p_3}{2\theta - 3},$$

with constants p_2, p_3, p_5 defined in (3.30).

In addition,

$$(3.24) \quad T_k \leq q_0^k T_0 + \frac{8q_2}{(1 - \sqrt{\lambda_2})(k + m)^2} + \frac{8q_3}{(1 - \sqrt{\lambda_2})(k + m)^3} + \frac{8q_4}{(1 - \sqrt{\lambda_2})(k + m)^4},$$

where constants q_0, q_2, q_3, q_4 are defined in (3.33).

Proof. Step 1: bounding H_k . From Lemmas 3.4 and 3.5 and the definition of H_{k+1} , we have for all $k \geq 0$ that

$$(3.25) \quad \begin{aligned} H_{k+1} \leq & \left[\left(1 - \frac{3}{2} \alpha_k \mu \right) + \omega_k \frac{4\alpha_k^2 \lambda_2 L^2 \|U_L\|^2 n}{c^2(1 - \lambda_2)} \right] M_k \\ & + \left[\frac{3\alpha_k c^2 \|U_R\|^2 L^2}{n\mu} + \omega_k \frac{(3 + \sqrt{\lambda_2})}{4} \right] T_k \\ & + \left(\frac{1}{n} + \frac{\omega_k \lambda_2 n \|U_L\|^2}{c^2} \right) \alpha_k^2 \bar{\sigma}^2 \\ & + \omega_k \frac{4\|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{c^2(1 - \sqrt{\lambda_2})} \|\alpha_{k+1} - \alpha_k\|^2. \end{aligned}$$

We now show the following inequalities hold for all $k \geq 0$:

$$(3.26a) \quad \left(1 - \frac{3}{2} \alpha_k \mu \right) + \omega_k \frac{4\alpha_k^2 \lambda_2 L^2 \|U_L\|^2 n}{c^2(1 - \lambda_2)} \leq 1 - \frac{4}{3} \alpha_k \mu,$$

$$(3.26b) \quad \frac{3\alpha_k c^2 \|U_R\|^2 L^2}{n\mu} + \omega_k \frac{3 + \sqrt{\lambda_2}}{4} \leq \left(1 - \frac{4}{3} \alpha_k \mu \right) \omega_k.$$

First of all, condition (3.26b) is equivalent to

$$\left(1 - \frac{4}{3} \alpha_k \mu - \frac{3 + \sqrt{\lambda_2}}{4} \right) \omega_k \geq \frac{3\alpha_k c^2 \|U_R\|^2 L^2}{n\mu}.$$

Since $m \geq \frac{24\theta}{1 - \sqrt{\lambda_2}}$, we have $\alpha_k \leq \frac{1 - \sqrt{\lambda_2}}{24\mu}$ for all $k \geq 0$, and so that

$$1 - \frac{4}{3} \alpha_k \mu - \frac{3 + \sqrt{\lambda_2}}{4} \geq \frac{1 - \sqrt{\lambda_2}}{4} - \frac{1 - \sqrt{\lambda_2}}{8} = \frac{1 - \sqrt{\lambda_2}}{8}.$$

Hence for (3.26b) to hold, it is sufficient that

$$(3.27) \quad \omega_k \geq \frac{24\alpha_k c^2 \|U_R\|^2 L^2}{n\mu(1 - \sqrt{\lambda_2})},$$

which is satisfied given the definition of ω_k in (3.21).

Second, condition (3.26a) requires

$$(3.28) \quad \omega_k \leq \frac{(1 - \lambda_2)c^2\mu}{24\lambda_2 L^2 n \|U_L\|^2} \frac{1}{\alpha_k},$$

or

$$\frac{24\alpha_k c^2 \|U_R\|^2 L^2}{n\mu(1 - \sqrt{\lambda_2})} \leq \frac{(1 - \lambda_2)c^2\mu}{24\lambda_2 L^2 n \|U_L\|^2} \frac{1}{\alpha_k}.$$

It is sufficient that

$$(3.29) \quad \alpha_k \leq \frac{\mu(1 - \sqrt{\lambda_2})}{24L^2 \|U_R\| \|U_L\|}, \forall k \geq 0 \quad \text{or} \quad m \geq \frac{24\theta L^2 \|U_R\| \|U_L\|}{\mu^2(1 - \sqrt{\lambda_2})}.$$

We have verified condition (3.26), and thus obtain the following recursion for H_{k+1} from (3.25):

$$\begin{aligned} H_{k+1} &\leq \left(1 - \frac{4}{3}\alpha_k\mu\right) H_k + \left(\frac{1}{n} + \frac{\omega_k\lambda_2 n \|U_L\|^2}{c^2}\right) \alpha_k^2 \bar{\sigma}^2 \\ &\quad + \omega_k \frac{4\|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{c^2(1 - \sqrt{\lambda_2})} \|\alpha_{k+1} - \alpha_k\|^2 \\ &\leq \left(1 - \frac{4\theta}{3(k+m)}\right) H_k + \left[\frac{24\theta L^2 \|U_L\|^2 \|U_R\|^2 \lambda_2}{\mu^2(1 - \sqrt{\lambda_2})(k+m)} + \frac{1}{n}\right] \frac{\theta^2 \bar{\sigma}^2}{\mu^2(k+m)^2} \\ &\quad + \frac{24\theta c^2 \|U_R\|^2 L^2}{n\mu^2(1 - \sqrt{\lambda_2})(k+m)} \frac{4\|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{c^2(1 - \sqrt{\lambda_2})} \frac{\theta^2}{\mu^2(k+m)^4} \\ &= \left(1 - \frac{4\theta}{3(k+m)}\right) H_k + \frac{p_2}{(k+m)^2} + \frac{p_3}{(k+m)^3} + \frac{p_5}{(k+m)^5}, \end{aligned}$$

where

$$(3.30) \quad \begin{aligned} p_2 &:= \frac{\theta^2 \bar{\sigma}^2}{n\mu^2}, \\ p_3 &:= \frac{24\lambda_2 \theta^3 \bar{\sigma}^2 \|U_L\|^2 \|U_R\|^2 L^2}{\mu^4(1 - \sqrt{\lambda_2})}, \\ p_5 &:= \frac{96\theta^3 \|U_L\|^2 \|U_R\|^2 L^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{n\mu^4(1 - \sqrt{\lambda_2})^2}. \end{aligned}$$

Then for all $k \geq 1$, we have

$$\begin{aligned}
H_k &\leq \prod_{t=0}^{k-1} \left(1 - \frac{4\theta}{3(t+m)}\right) H_0 \\
&\quad + \sum_{t=0}^{k-1} \left[\prod_{j=t+1}^{k-1} \left(1 - \frac{4\theta}{3(j+m)}\right) \right] \left[\frac{p_2}{(t+m)^2} + \frac{p_3}{(t+m)^3} + \frac{p_5}{(t+m)^5} \right] \\
&\leq \frac{m^{\frac{4\theta}{3}}}{(k+m)^{\frac{4\theta}{3}}} H_0 + \sum_{t=0}^{k-1} \frac{(m+t+1)^{\frac{4\theta}{3}}}{(k+m)^{\frac{4\theta}{3}}} \left[\frac{p_2}{(t+m)^2} + \frac{p_3}{(t+m)^3} + \frac{p_5}{(t+m)^5} \right] \\
&\leq \frac{m^{\frac{4\theta}{3}}}{(k+m)^{\frac{4\theta}{3}}} H_0 \\
&\quad + \frac{4}{3(k+m)^{\frac{4\theta}{3}}} \sum_{t=0}^{k-1} \left[p_2(m+t)^{\frac{4\theta}{3}-2} + p_3(m+t)^{\frac{4\theta}{3}-3} + p_5(m+t)^{\frac{4\theta}{3}-5} \right],
\end{aligned}$$

where we invoked [Lemma 3.6](#) for the second inequality. Additionally, since $\theta > 4$, we have

$$(3.31) \quad \sum_{t=0}^{k-1} (m+t)^{\frac{4\theta}{3}-i} \leq \int_{-1}^k (m+t)^{\frac{4\theta}{3}-i} dt \leq \frac{3}{4\theta-3i+3} (m+k)^{\frac{4\theta}{3}-i+1}, \quad i = 2, 3, 5.$$

Further noticing that $\frac{m^{\frac{4\theta}{3}}}{(k+m)^{\frac{4\theta}{3}}} \leq \frac{m^2}{(k+m)^2}$ since $\frac{m}{k+m} \leq 1$, we have

$$\begin{aligned}
H_k &\leq \frac{m^2}{(k+m)^2} H_0 + \frac{4p_2}{4\theta-3} \frac{1}{k+m} + \frac{4p_3}{4\theta-6} \frac{1}{(m+k)^2} + \frac{4p_5}{4\theta-12} \frac{1}{(m+k)^4} \\
&= \frac{1}{k+m} \left[\frac{4p_2}{4\theta-3} + \frac{p_5}{\theta-3} \frac{1}{(m+k)^3} \right] + \frac{1}{(k+m)^2} \left[m^2 H_0 + \frac{2p_3}{2\theta-3} \right] \\
&\leq \frac{1}{k+m} \left[\frac{1}{\theta-3} \left(p_2 + \frac{p_5}{m^3} \right) \right] + \frac{1}{(k+m)^2} \left[m^2 H_0 + \frac{2p_3}{2\theta-3} \right] \\
&= \frac{\hat{H}_1}{k+m} + \frac{\hat{H}_2}{(k+m)^2}.
\end{aligned}$$

Since $H_k = M_k + \omega_k T_k \geq M_k$, we get

$$(3.32) \quad M_k \leq \frac{\hat{H}_1}{k+m} + \frac{\hat{H}_2}{(k+m)^2}, \quad \forall k.$$

Step 2: bounding T_k . To bound T_k , we substitute (3.32) into [Lemma 3.5](#) and get

$$\begin{aligned}
T_{k+1} &\leq \frac{4\lambda_2 n \|U_L\|^2 \theta^2 L^2}{\mu^2 c^2 (1-\lambda_2)} \left(\frac{\hat{H}_1}{(k+m)^3} + \frac{\hat{H}_2}{(k+m)^4} \right) + \frac{\lambda_2 n \|U_L\|^2 \sigma^2 \theta^2}{c^2 \mu^2} \frac{1}{(k+m)^2} \\
&\quad + \frac{(3+\sqrt{\lambda_2})}{4} T_k + \frac{4 \|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2 \theta^2}{\mu^2 c^2 (1-\sqrt{\lambda_2})} \frac{1}{(k+m)^4} \\
&= q_0 T_k + \frac{q_2}{(k+m)^2} + \frac{q_3}{(k+m)^3} + \frac{q_4}{(k+m)^4},
\end{aligned}$$

where

$$(3.33) \quad \begin{aligned} q_0 &:= \frac{3 + \sqrt{\lambda_2}}{4}, & q_2 &:= \frac{\lambda_2 n \|U_L\|^2 \bar{\sigma}^2 \theta^2}{c^2 \mu^2}, \\ q_3 &:= \frac{4\lambda_2 n \|U_L\|^2 \theta^2 L^2 \hat{H}_1}{\mu^2 c^2 (1 - \lambda_2)}, & q_4 &:= \frac{4\lambda_2 n \|U_L\|^2 \theta^2 L^2 \hat{H}_2}{\mu^2 c^2 (1 - \lambda_2)} + \frac{4\|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2 \theta^2}{\mu^2 c^2 (1 - \sqrt{\lambda_2})}. \end{aligned}$$

Hence for all $k \geq 0$, we have

$$\begin{aligned} T_k &\leq q_0^k T_0 + \sum_{t=0}^{k-1} q_0^{k-1-t} \left[\frac{q_2}{(m+t)^2} + \frac{q_3}{(m+t)^3} + \frac{q_4}{(m+t)^4} \right] \\ &:= q_0^k T_0 + A_2(k) + A_3(k) + A_4(k), \end{aligned}$$

with

$$(3.34) \quad \begin{aligned} A_2(k) &:= \sum_{t=0}^{k-1} \frac{q_0^{k-t-1} q_2}{(m+t)^2}, & A_3(k) &:= \sum_{t=0}^{k-1} \frac{q_0^{k-t-1} q_3}{(m+t)^3}, \\ A_4(k) &:= \sum_{t=0}^{k-1} \frac{q_0^{k-t-1} q_4}{(m+t)^4}. \end{aligned}$$

Note that for $i = 2, 3, 4$,

$$A_i(k+1) = q_0 \left[\sum_{t=0}^{k-1} q_0^{k-t-1} \frac{q_i}{(m+t)^i} + \frac{q_i}{q_0(m+k)^i} \right] = q_0 A_i(k) + \frac{q_i}{(m+k)^i}, \quad \forall k \geq 0.$$

By induction we obtain

$$(3.35) \quad A_i(k) \leq \frac{1}{(k+m)^i} \frac{q_i}{(1 - \frac{1}{m+1})^i - q_0}, \quad i = 2, 3, 4.$$

Since $m \geq \frac{24\theta}{1-\sqrt{\lambda_2}}$, it can be verified that

$$(3.36) \quad \left(1 - \frac{1}{m+1} \right)^i - q_0 \geq \frac{1-q_0}{2} = \frac{1-\sqrt{\lambda_2}}{8}, \quad i = 2, 3, 4.$$

We conclude that for all $k \geq 0$, there holds

$$(3.37) \quad T_k \leq q_0^k T_0 + \frac{8q_2}{(1-\sqrt{\lambda_2})(k+m)^2} + \frac{8q_3}{(1-\sqrt{\lambda_2})(k+m)^3} + \frac{8q_4}{(1-\sqrt{\lambda_2})(k+m)^4}. \quad \square$$

4. Main Results. In this section, we present the main convergence results for EDAS. First, we show EDAS performs asymptotically as well as the centralized stochastic gradient descent (SGD) method in [Theorem 4.1](#). In other words, EDAS enjoys the so-called “asymptotic network independence” property. Then we refine the convergence results by iteratively combining the existing results with [Lemmas 3.4](#) and [3.5](#). Finally, we derive the transient time for EDAS to approach the network independent rate in [Theorem 4.6](#) by comparing with the performance of centralized SGD.

THEOREM 4.1. Under *Algorithm 2.1* with *Assumptions 1.1* to *1.3*, suppose $\theta > 4$ and m is chosen according to (3.22). We have

$$M_k \leq \frac{4\theta^2 \bar{\sigma}^2}{(3\theta - 2)n\mu^2(k + m)} + \frac{4c_0 T_0 m^{\frac{3\theta}{2}-1}}{(1 - q_0)(k + m)^{\frac{3\theta}{2}}} + \frac{m^{\frac{3\theta}{2}} M_0}{(k + m)^{\frac{3\theta}{2}}} \\ + \frac{32c_0}{1 - \sqrt{\lambda_2}} \left[\frac{q_2}{(3\theta - 4)(m + k)^2} + \frac{q_3}{(3\theta - 6)(k + m)^3} + \frac{q_4}{(3\theta - 8)(k + m)^4} \right],$$

where $c_0 := \frac{3\theta L^2 c^2 \|U_R\|^2}{n\mu^2}$

Proof. From Lemma 3.4, we have

$$M_{k+1} \leq \left(1 - \frac{3\alpha_k \mu}{2}\right) M_k + \frac{3\alpha_k L^2 c^2 \|U_R\|^2}{n\mu} T_k + \frac{\alpha_k^2 \bar{\sigma}^2}{n}.$$

Thus for all $k \geq 0$,

$$(4.1) \quad M_k \leq \prod_{t=0}^{k-1} \left(1 - \frac{3\theta}{2(t + m)}\right) M_0 \\ + \sum_{t=0}^{k-1} \left[\prod_{j=t+1}^{k-1} \left(1 - \frac{3\theta}{2(j + m)}\right) \right] \left[\frac{3\theta L^2 c^2 \|U_R\|^2}{n\mu^2(t + m)} T(t) + \frac{\theta^2 \bar{\sigma}^2}{n\mu^2(t + m)^2} \right] \\ \leq \left(\frac{m}{k + m}\right)^{\frac{3\theta}{2}} M_0 + \sum_{t=0}^{k-1} \left(\frac{t + m + 1}{k + m}\right)^{\frac{3\theta}{2}} \left[\frac{c_0 T(t)}{t + m} + \frac{\theta^2 \bar{\sigma}^2}{n\mu^2(t + m)^2} \right],$$

where Lemma 3.6 was invoked for obtaining the second inequality. In light of (3.24),

$$\sum_{t=0}^{k-1} \left(\frac{t + m + 1}{k + m}\right)^{\frac{3\theta}{2}} \left[\frac{c_0 T(t)}{t + m} + \frac{\theta^2 \bar{\sigma}^2}{n\mu^2(t + m)^2} \right] \\ \leq c_0 T_0 \sum_{t=0}^{k-1} \frac{(m + t + 1)^{\frac{3\theta}{2}-1} q_0^t}{(m + t)(k + m)^{\frac{3\theta}{2}}} + \frac{\theta^2 \bar{\sigma}^2}{n\mu^2(k + m)^{\frac{3\theta}{2}}} \sum_{t=0}^{k-1} \frac{(m + t + 1)^{\frac{3\theta}{2}}}{(m + t)^2} \\ + \frac{8c_0}{(1 - \sqrt{\lambda_2})(k + m)^{\frac{3\theta}{2}}} \sum_{t=0}^{k-1} \left[\frac{q_2(m + t + 1)^{\frac{3\theta}{2}}}{(m + t)^3} + \frac{q_3(m + t + 1)^{\frac{3\theta}{2}}}{(m + t)^4} \right. \\ \left. + \frac{q_4(m + t + 1)^{\frac{3\theta}{2}}}{(m + t)^5} \right] \\ \leq 2c_0 T_0 \sum_{t=0}^{k-1} \frac{(m + t)^{\frac{3\theta}{2}-1} q_0^t}{(k + m)^{\frac{3\theta}{2}}} + \frac{2\theta^2 \bar{\sigma}^2}{n\mu^2(k + m)^{\frac{3\theta}{2}}} \sum_{t=0}^{k-1} (m + t)^{\frac{3\theta}{2}-2} \\ + \frac{16c_0}{(1 - \sqrt{\lambda_2})(k + m)^{\frac{3\theta}{2}}} \sum_{t=0}^{k-1} \left[q_2(m + t)^{\frac{3\theta}{2}-3} + q_3(m + t)^{\frac{3\theta}{2}-4} + q_4(m + t)^{\frac{3\theta}{2}-5} \right].$$

Since $(m + t)^{\frac{3\theta}{2}-1} q_0^t$ is decreasing in t and

$$\frac{1}{\ln q_0} d((m + t)^{\frac{3\theta}{2}-1} q_0^t) = \frac{1}{\ln q_0} \left(\frac{3\theta}{2} - 1 \right) (m + t)^{\frac{3\theta}{2}-2} q_0^t dt + (m + t)^{\frac{3\theta}{2}-1} q_0^t dt \\ \geq \frac{1}{2} (m + t)^{\frac{3\theta}{2}-1} q_0^t dt,$$

we have

$$\begin{aligned} \sum_{t=0}^{k-1} (m+t)^{\frac{3\theta}{2}-1} q_0^t &\leq \int_0^{+\infty} (m+t)^{\frac{3\theta}{2}-1} q_0^t dt \\ &\leq \frac{2}{\ln q_0} \int_m^{+\infty} \frac{d(t^{\frac{3\theta}{2}-1} q_0^{t-m})}{dt} \\ &= -\frac{2m^{\frac{3\theta}{2}-1}}{\ln q_0} \leq \frac{2m^{\frac{3\theta}{2}-1}}{1-q_0}. \end{aligned}$$

In addition, from an argument similar to (3.31), we obtain

$$\sum_{t=0}^{k-1} (m+t)^{\frac{3\theta}{2}-i} \leq \frac{1}{\frac{3\theta}{2}-i+1} (m+k)^{\frac{3\theta}{2}-i+1}, \quad i = 2, 3, 4, 5.$$

Hence for all $k \geq 0$,

$$\begin{aligned} M_k &\leq \frac{4\theta^2 \bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \frac{4c_0 T_0 m^{\frac{3\theta}{2}-1}}{(1-q_0)(k+m)^{\frac{3\theta}{2}}} + \frac{m^{\frac{3\theta}{2}} M_0}{(k+m)^{\frac{3\theta}{2}}} \\ &\quad + \frac{32c_0}{1-\sqrt{\lambda_2}} \left[\frac{q_2}{(3\theta-4)(m+k)^2} + \frac{q_3}{(3\theta-6)(k+m)^3} + \frac{q_4}{(3\theta-8)(k+m)^4} \right]. \end{aligned}$$

From [Theorem 4.1](#), we can see that the asymptotic convergence rate of EDAS, which behaves as $\frac{4\theta^2 \bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)}$, is of the same order as that of the centralized SGD method [23]. Our next goal is to derive the transient time needed for EDAS to reach this asymptotic convergence rate. In particular, we would like to characterize the dependence relationship between the transient time and the network characteristics as well as function properties. For this purpose, we first simplify the presentation of [Theorem 4.1](#) with $\mathcal{O}(\cdot)$ notation in [Corollary 4.3](#) and then improve the convergence results by iteratively substituting the existing upper bounds into [Lemmas 3.4](#) and [3.5](#).

In the following lemma, we estimate the constants $q_i, i = 2, 3, 4, M_0, T_0, H_0$, and c_0 appearing in the statement of [Theorem 4.1](#) with $\mathcal{O}(\cdot)$ notation.

LEMMA 4.2. *Let the free scaling parameter c in [Lemma 2.5](#) be chosen as*

$$c^2 = n \|U_L\|^2.$$

Then we have

$$\begin{aligned} M_0 &= \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n} \right), \quad T_0 = \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + (1-\lambda_2) \|\nabla F(\mathbf{x}^*)\|^2}{n} \right) \\ H_0 &= \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + (1-\lambda_2) \|\nabla F(\mathbf{x}^*)\|^2}{n} \right), \quad c_0 = \mathcal{O}(1) \\ \frac{1}{1-q_0} &= \mathcal{O} \left(\frac{1}{1-\lambda_2} \right), \quad q_2 = \mathcal{O}(1) \\ q_3 &= \mathcal{O} \left(\frac{1 + \|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)} \right), \quad q_4 = \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^3} + \frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^2} + \frac{1}{(1-\lambda_2)^2} \right). \end{aligned}$$

Proof. See [Appendix A.2](#). □

In light of Lemma 4.2, we again iteratively apply Lemmas 3.4 and 3.5 to derive Corollary 4.3 based on Theorem 4.1.

COROLLARY 4.3. *Under Algorithm 2.1 with Assumptions 1.1 to 1.3, suppose $\theta > 4$ and m be chosen according to (3.22). We have for all $k \geq 0$,*

$$\begin{aligned} M_k &\leq \frac{4\theta^2\bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} \\ &\quad + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^3}\right) \frac{1}{(k+m)^4} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^6}\right) \frac{1}{(k+m)^6}. \end{aligned}$$

and

$$\begin{aligned} T_k &\leq \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\nabla F(\mathbf{x}^*)\|^2}{n}\right) \left(\frac{3+\sqrt{\lambda_2}}{4}\right)^k + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} \\ &\quad + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^3}\right) \frac{1}{(k+m)^4} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^6}\right) \frac{1}{(k+m)^6}. \end{aligned}$$

Proof. From Theorem 4.1, noticing that $m = \Omega(\frac{1}{1-\lambda_2})$, we have when $k \geq 0$,

$$\begin{aligned} (4.2) \quad M_k &\leq \frac{4\theta^2\bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \frac{4c_0T_0m^{\frac{3\theta}{2}-1}}{(1-q_0)(k+m)^{\frac{3\theta}{2}}} + \frac{m^{\frac{3\theta}{2}}M_0}{(k+m)^{\frac{3\theta}{2}}} \\ &\quad + \frac{32c_0}{1-\sqrt{\lambda_2}} \left[\frac{q_2}{(3\theta-4)(k+m)^2} + \frac{q_3}{(3\theta-6)(k+m)^3} + \frac{q_4}{(3\theta-8)(k+m)^4} \right] \\ &= \frac{4\theta^2\bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + (1-\lambda_2)\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^{\frac{3\theta}{2}}}\right) \frac{1}{(k+m)^{\frac{3\theta}{2}}} \\ &\quad + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} + \mathcal{O}\left(\frac{1+\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^2}\right) \frac{1}{(k+m)^3} \\ &\quad + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^4} + \frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^3} + \frac{1}{(1-\lambda_2)^3}\right) \frac{1}{(k+m)^4} \\ &= \frac{4\theta^2\bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} \\ &\quad + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^2}\right) \frac{1}{(k+m)^3} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^4}\right) \frac{1}{(k+m)^4} \\ &= \frac{4\theta^2\bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \frac{c_2}{(k+m)^2} + \frac{c_3}{(k+m)^3} + \frac{c_4}{(k+m)^4}, \end{aligned}$$

where

$$c_2 = \mathcal{O}\left(\frac{1}{1-\lambda_2}\right), \quad c_3 = \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^2}\right), \quad c_4 = \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^4}\right).$$

Substituting (4.2) into Lemma 3.5, we have

$$\begin{aligned}
T_{k+1} &\leq q_0 T_k + \frac{\theta^2 \lambda_2}{\mu^2 (k+m)^2} \left[\frac{4L^2}{1-\lambda_2} \left(\frac{4\theta^2 \bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \frac{c_2}{(k+m)^2} \right. \right. \\
&\quad \left. \left. + \frac{c_3}{(k+m)^3} + \frac{c_4}{(k+m)^4} \right) + \bar{\sigma}^2 \right] + \frac{4\theta^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{n\mu^2(1-\sqrt{\lambda_2})} \frac{1}{(k+m)^4} \\
&= q_0 T_k + \frac{\theta^2 \lambda_2 \bar{\sigma}^2}{\mu^2} \frac{1}{(k+m)^2} + \frac{16\theta^4 \bar{\sigma}^2 \lambda_2 L^2}{(3\theta-2)n\mu^4(1-\lambda_2)} \frac{1}{(k+m)^3} \\
&\quad + \left[\frac{4\theta^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{n\mu^2(1-\sqrt{\lambda_2})} + \frac{4\theta^2 \lambda_2 c_2 L^2}{\mu^2(1-\lambda_2)} \right] \frac{1}{(k+m)^4} \\
&\quad + \frac{4\theta^2 \lambda_2 L^2 c_3}{\mu^2(1-\lambda_2)} \frac{1}{(k+m)^5} + \frac{4\theta^2 \lambda_2 L^2 c_4}{\mu^2(1-\lambda_2)} \frac{1}{(k+m)^6} \\
&= q_0 T_k + \frac{b_2}{(k+m)^2} + \frac{b_4}{(k+m)^4} + \frac{b_6}{(k+m)^6},
\end{aligned}$$

where

$$b_2 = \mathcal{O}(1), \quad b_4 = \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^2}\right), \quad b_6 = \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^5}\right).$$

Then,

$$T_k \leq q_0^k T_0 + \sum_{t=0}^{k-1} q_0^{k-t-1} \left[\sum_{i=2,4,6} \frac{b_i}{(k+m)^i} \right].$$

Similar to the derivation of (3.35), we obtain that for all $k \geq 0$,

$$\begin{aligned}
(4.3) \quad T_k &\leq q_0^k T_0 + \frac{8}{1-\sqrt{\lambda_2}} \left[\frac{b_2}{(k+m)^2} + \frac{b_4}{(k+m)^4} + \frac{b_6}{(k+m)^6} \right] \\
&\leq q_0^k T_0 + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^3}\right) \frac{1}{(k+m)^4} \\
&\quad + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^6}\right) \frac{1}{(k+m)^6} \\
&= q_0^k T_0 + \sum_{i=2,4,6} \frac{e_i}{(k+m)^i},
\end{aligned}$$

where each e_i is of the same order of $\frac{b_i}{(1-\lambda_2)}$ for $i = 2, 4, 6$.

Substituting (4.3) into Lemma 3.4, we obtain

$$(4.4) \quad M_{k+1} \leq \left(1 - \frac{3\theta}{2(k+m)}\right) M_k + \frac{\theta^2 \bar{\sigma}^2}{n\mu^2(k+m)^2} + \frac{c_0}{k+m} \left[q_0^k T_0 + \sum_{i=2,4,6} \frac{e_i}{(k+m)^i} \right].$$

Similar to relation (4.1), we derive

$$\begin{aligned}
M_k &\leq \frac{m^{\frac{3\theta}{2}}}{(k+m)^{\frac{3\theta}{2}}} M_0 + \sum_{t=0}^{k-1} \frac{(m+t+1)^{\frac{3\theta}{2}}}{(k+m)^{\frac{3\theta}{2}}} \left[\frac{c_0 q_0^t T_0}{t+m} + \sum_{i=2,4,6} \frac{c_0 e_i}{(t+m)^{(i+1)}} \right] \\
&\quad + \frac{\theta^2 \bar{\sigma}^2}{n\mu^2(k+m)^{\frac{3\theta}{2}}} \sum_{t=0}^{k-1} \frac{(m+t+1)^{\frac{3\theta}{2}}}{(t+m)^2} \\
&\leq \frac{4\theta^2 \bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \frac{m^{\frac{3\theta}{2}}}{(k+m)^{\frac{3\theta}{2}}} M_0 + \frac{2c_0 T_0 m^{\frac{3\theta}{2}-1}}{(1-q_0)(k+m)^{\frac{3\theta}{2}}} \\
&\quad + \frac{4c_0 e_2}{(3\theta-4)(k+m)^2} + \frac{4c_0 e_4}{(3\theta-8)(k+m)^4} + \frac{4c_0 e_6}{(3\theta-12)(k+m)^6} \\
&= \frac{4\theta^2 \bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} \\
&\quad + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^3}\right) \frac{1}{(k+m)^4} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^6}\right) \frac{1}{(k+m)^6}.
\end{aligned}$$

Again we considered the fact that $(k+m) \geq \Omega(\frac{1}{1-\lambda_2})$ with m chosen according to (3.22) for all $k \geq 1$. \square

We now state the convergence rate for the EDAS algorithm under the measure $\frac{1}{n} \sum_{i=1}^n \mathbb{E} [\|x_{i,k} - x^*\|^2] = \mathbb{E} [\|\tilde{\mathbf{x}}_k\|^2]$, i.e., the average expected optimization error over all the agents in the network. Note that from Corollary 4.3, the error $\tilde{\mathbf{x}}_k$ can be decomposed into $\tilde{\mathbf{x}}_k = \mathbf{1}\bar{z}_k + cU_{R,u}\tilde{z}_k$. Therefore,

$$(4.5) \quad \|\tilde{\mathbf{x}}_k\|^2 \leq 2n\|\bar{z}_k\|^2 + 2c^2\|U_R\|^2\|\tilde{z}_k\|^2.$$

Taking full expectation on both sides of (4.5) and noting the choice $c^2 = n\|U_L\|^2$, we obtain

$$(4.6) \quad \mathbb{E} [\|\tilde{\mathbf{x}}_k\|^2] \leq 2nM_k + 2n\|U_L\|^2\|U_R\|^2T_k.$$

Combining Corollary 4.3 and (4.6) leads to the improved convergence rate of EDAS stated in Theorem 4.4.

THEOREM 4.4. *Under Algorithm 2.1 with Assumptions 1.1 to 1.3, suppose $\theta > 5$, and m is chosen as in (3.22). We have for all $k \geq 1$,*

$$\begin{aligned}
\frac{1}{n} \sum_{i=1}^n \mathbb{E} [\|x_{i,k} - x^*\|^2] &\leq \frac{4\theta^2 \bar{\sigma}^2}{(3\theta-2)n\mu^2(k+m)} + \mathcal{O}\left(\frac{1}{1-\lambda_2}\right) \frac{1}{(k+m)^2} \\
&\quad + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\nabla F(\mathbf{x}^*)\|^2}{n}\right) \left(\frac{3+\lambda_2}{4}\right)^k \\
&\quad + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1-\lambda_2)^3}\right) \frac{1}{(k+m)^4} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1-\lambda_2)^6}\right) \frac{1}{(k+m)^6}.
\end{aligned}$$

Proof. Substitute the bounds on M_k and T_k in Corollary 4.3 into (4.6), and notice that the product $\|U_R\|^2\|U_L\|^2$ is independent of n according to Lemma 2.5. We obtain the result. \square

4.1. Transient Time. In this part, we estimate how long it takes for EDAS to achieve the convergence rate of centralized stochastic gradient descent (SGD) method, i.e., *transient time* of the algorithm. First, we state the convergence rate of SGD [23].

THEOREM 4.5. *Under the centralized stochastic gradient descent method (SGD) with stepsize policy $\alpha_k = \frac{\theta}{\mu(k+m)}$, suppose $m \geq \left\lceil \frac{\theta L}{\mu} \right\rceil$. We have*

$$\mathbb{E} [\|x_k - x^*\|^2] \leq \frac{\theta^2 \bar{\sigma}^2}{(2\theta - 1)n\mu^2(k+m)} + \mathcal{O}\left(\frac{1}{n}\right) \frac{1}{(k+m)^2}.$$

In the next theorem, we derive the transient time for EDAS.

THEOREM 4.6. *Under Algorithm 2.1 with Assumptions 1.1 to 1.3, suppose $\theta > 5$ and m is chosen according to (3.22). Then it takes*

$$K_T = \max \left\{ \mathcal{O}\left(\frac{n}{1-\lambda_2}\right), \mathcal{O}\left(\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{(1-\lambda_2)^3}\right)^{\frac{1}{3}}\right), \mathcal{O}\left(\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{(1-\lambda_2)^6}\right)^{\frac{1}{6}}\right), \right. \\ \left. \mathcal{O}\left(\frac{\max\left\{\log\left(\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\nabla F(\mathbf{x}^*)\|^2\right), -\log(1-\lambda_2)\right\}}{1-\lambda_2}\right) \right\}$$

time for Algorithm 2.1 to reach the asymptotic, network independent convergence rate, that is, when $k \geq K_T$, we have $\frac{1}{n} \sum_{i=1}^n \mathbb{E} [\|x_{i,k} - x^*\|^2] \leq \frac{\theta^2 \bar{\sigma}^2}{(2\theta-1)n\mu^2 k} \mathcal{O}(1)$.

Proof. From Theorem 4.4, we have

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [\|x_{i,k} - x^*\|^2] &\leq \frac{\theta^2 \bar{\sigma}^2}{(2\theta - 1)n\mu^2 k} \left[\frac{4(2\theta - 1)}{3\theta - 2} + \mathcal{O}\left(\frac{n}{1-\lambda_2}\right) \frac{1}{k} \right. \\ &\quad + \mathcal{O}\left(\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\nabla F(\mathbf{x}^*)\|^2\right) \left(\frac{3+\lambda_2}{4}\right)^k k \\ &\quad \left. + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{(1-\lambda_2)^3}\right) \frac{1}{k^3} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{(1-\lambda_2)^6}\right) \frac{1}{k^5} \right]. \end{aligned}$$

Let K_T be such that

$$\begin{aligned} \mathcal{O}(1) &= \frac{4(2\theta - 1)}{3\theta - 2} + \mathcal{O}\left(\frac{n}{1-\lambda_2}\right) \frac{1}{K_T} \\ &\quad + \mathcal{O}\left(\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\nabla F(\mathbf{x}^*)\|^2\right) \left(\frac{3+\lambda_2}{4}\right)^{K_T} K_T \\ &\quad + \mathcal{O}\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{(1-\lambda_2)^3}\right) \frac{1}{K_T^3} + \mathcal{O}\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{(1-\lambda_2)^6}\right) \frac{1}{K_T^5}. \end{aligned}$$

We obtain

$$K_T = \max \left\{ \mathcal{O} \left(\frac{n}{1-\lambda_2} \right), \mathcal{O} \left(\left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{(1-\lambda_2)^3} \right)^{\frac{1}{3}} \right), \mathcal{O} \left(\left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{(1-\lambda_2)^6} \right)^{\frac{1}{5}} \right), \right. \\ \left. \mathcal{O} \left(\frac{\max \left\{ \log \left(\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\nabla F(\mathbf{x}^*)\|^2 \right), -\log(1-\lambda_2) \right\}}{1-\lambda_2} \right) \right\},$$

which finishes the proof. \square

Under mild additional conditions, we can obtain a cleaner expression for the transient time of EDAS in the following corollary.

COROLLARY 4.7. *Under [Algorithm 2.1](#) with [Assumptions 1.1 to 1.3](#), suppose $\theta > 5$ and m is chosen according to [\(3.22\)](#). Assume in addition that $\|\mathbf{x}_0 - \mathbf{x}^*\|^2 = \mathcal{O}(n^3)$, $\|\nabla F(\mathbf{x}^*)\|^2 = \mathcal{O}(n^3)$ and $\frac{1}{1-\lambda_2} = \mathcal{O}(n^q)$ for some $q > 0$. Then it takes*

$$K_T = \mathcal{O} \left(\frac{n}{1-\lambda_2} \right)$$

steps for [Algorithm 2.1](#) to reach the asymptotic, network independent convergence rate.

Remark 4.8. Assuming that $\|\mathbf{x}_0 - \mathbf{x}^*\|^2 = \mathcal{O}(n^3)$ and $\|\nabla F(\mathbf{x}^*)\|^2 = \mathcal{O}(n^3)$ is mild. This condition can be satisfied for many problem settings including the ones we consider in [Section 5](#). In addition, $\frac{1}{1-\lambda_2} = \mathcal{O}(n^q)$ is generally satisfied since $\frac{1}{1-\lambda_2} = \mathcal{O}(n^2)$ holds under common choices of the mixing weights for undirected graphs [\[16\]](#).


5. Numerical Examples. In this section, we present two numerical examples to verify and complement our theoretical results. To begin with, we define the transient time for the algorithms of interest as follows for practical consideration:

$$(5.1) \quad \inf_k \left\{ k > 100 : \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\|x_{i,k} - x^*\|^2 \right] \leq 2\mathbb{E} \left[\|x_k - x^*\|^2 \right] \right\},$$

where $x_k \in \mathbb{R}^p$ stands for the k -th iterate for the centralized SGD algorithm.

In the first experiment, we construct a “hard” optimization problem for which the experimental transient time of EDAS is of the order of $\mathcal{O} \left(\frac{n}{1-\lambda_2} \right)$, which agrees with the upper bound given in [Theorem 4.6](#). This verifies the sharpness of the obtained theoretical results. Then we consider the problem of logistic regression for classifying handwritten digits from the MNIST dataset [\[9\]](#). In this case, EDAS achieves a transient time close to $\mathcal{O} \left(\frac{n}{(1-\lambda_2)^{0.5}} \right)$, which is better than the upper bound in [Corollary 4.7](#). Hence in practice, the performance of EDAS depends on the specific problem instances and can be better than the worse-case scenario.

For both problems, we consider two different types of network topologies, i.e., ring network shown in [Figure 1a](#) and grid network shown in [Figure 1b](#). The mixing matrices compliant with the networks are constructed under the Lazy Metropolis rule [\[16\]](#). In addition to presenting the transient times for EDAS, we also compare its performance with other algorithms enjoying the asymptotic network independent property, i.e., distributed stochastic gradient descent (DSGD) [\[23, 37\]](#) and distributed stochastic gradient tracking method (DSGT) [\[22\]](#).



(a) Ring network topology with 8 nodes

(b) Grid network topology with 16 nodes


FIG. 1. *Two network topologies.*

5.1. A Near-Sharp Quadratic Problem. Consider the following problem:

$$(5.2) \quad \min_{x \in \mathbb{R}^p} f(x) = \frac{1}{n} \sum_{i=1}^n \frac{\sqrt{n}}{2} \|x - d_i\|^2,$$

where $d_i = a_i \sqrt{i} \mathbf{1}$ and a_i is the i -th smallest eigenvalue of W . The unique solution to problem (5.2) is given by $x^* = \frac{1}{n} \sum_{i=1}^n d_i$. In the experiments, we set the initial solution to $\mathbf{0}$ for each node in the network; for centralized SGD we use the same initialization. Let $p = 1$, and each node i is able to obtain noisy gradients in the form of $g_i(x_i) = \sqrt{n}(x_i - d_i) + \omega$, where $\omega \sim \mathcal{N}(0, 0.1^2)$ denotes the noise. Stepsizes are chosen as $\alpha_k = 20/(k + 200), \forall k$.

Fist, we illustrate the property of “asymptotic network independence” by comparing the performance of EDAS and SGD in Figure 2a. We can see that the error of EDAS gets close to that of centralized SGD as the number of iteration grows. Eventually, it is as if the network were not there and the convergence rate for EDAS is exactly the same as SGD. Hence the phenomenon is called “asymptotic network independence”.




(a) Error terms for centralized and decentralized methods.

FIG. 2. *An illustration of asymptotic network independence.*

In Figure 3a, we plot the observed transient times of EDAS and $\frac{6n}{1-\lambda_2}$ against

the network size n for the ring network topology.³ We then consider the grid network topology in Figure 3b. It can be seen that the two curves are close in both cases, which suggests that the theoretical upper bound $K_T = \mathcal{O}\left(\frac{n}{1-\lambda_2}\right)$ given in Corollary 4.7 is sharp.



(a) Transient time for the ring network topology. (b) Transient time for the grid network topology.

FIG. 3. Comparison of the transient time for Algorithm 2.1 and multiples of $\frac{n}{1-\lambda_2}$. The expected errors are approximated by averaging 500 simulation results.

In Figure 4, we further compare the early stage performance of DSGT, DSGD, and EDAS with the same initialization. It can be seen that the error term of EDAS decreases faster than that of the other two decentralized methods and quickly gets close to the error of SGD. This is particularly true when the network is not well connected, e.g., when the network has a ring topology with a large number of nodes (see Figure 4c).

5.2. Logistic Regression. We consider classifying handwritten digits 1 and 2 in the MNIST dataset using logistic regression. There are 12700 data points $\mathcal{S} = \{(u, v)\}$ with $u \in \mathbb{R}^{785}$ denoting the image input and $v \in \{-1, 1\}$ being the label.⁴ Each agent possesses a distinct local dataset \mathcal{S}_i randomly selected from \mathcal{S} . The classifier can then be obtained by solving the following optimization problem using all the agents' local datasets $\mathcal{S}_i, i = 1, 2, \dots, n$:


$$(5.3a) \quad \min_{x \in \mathbb{R}^{785}} f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x),$$

$$(5.3b) \quad f_i(x) := \frac{1}{|\mathcal{S}_i|} \sum_{j \in \mathcal{S}_i} \log [1 + \exp(-x^\top u_j v_j)] + \frac{\rho}{2} \|x\|^2,$$


where $\rho > 0$ is a regularization parameter (hyperparameter). To solve the problem, each agent can obtain an unbiased estimate of $\nabla f_i(x)$ using a minibatch of randomly selected data points from its local dataset \mathcal{S}_i . In the experiment, we let $\rho = 1$, $\alpha_k = 6/(k+20), \forall k$ for all the considered algorithms with the same initial solutions:

³For problem (5.2), the upper bound on the transient time is $K_T = \mathcal{O}\left(\frac{n}{1-\lambda_2}\right)$ from Corollary 4.7 since we have $\|x_0 - x^*\|^2 = \|\nabla F(x^*)\|^2 = \mathcal{O}(n^3)$.

⁴Label -1 represents handwritten digit 2 and label 1 represents digit 1.



(a) Error terms over a ring net- (b) Error terms over a ring net- (c) Error terms over a ring net-
work with 20 nodes. work with 40 nodes. work with 60 nodes.



(d) Error terms over a grid net- (e) Error terms over a grid net- (f) Error terms over a grid net-
work with 49 nodes. work with 81 nodes. work with 121 nodes.

FIG. 4. Comparison among DSGT, DSGD, and EDAS over ring and grid network topologies for problem (5.2). The curves stand for $\mathbb{E}(\|x(k) - x^*\|^2)$ for SGD and $\frac{1}{n} \sum_{i=1}^n \mathbb{E}(\|x_{i,k} - x^*\|^2)$ for decentralized methods. The expected errors are approximated by averaging over 100 simulation results.

$x_{i,0} = \mathbf{0}, \forall i$ (distributed methods) and $x_0 = \mathbf{0}$ (centralized SGD). Each local dataset has $|\mathcal{S}_i| = 100$ data points and the mini-batch size is set to be 1.

In Figure 5, we can see that the observed transient times of EDAS are close to the multiple of $\frac{n}{(1-\lambda_2)^{0.5}}$. Hence in practice EDAS may achieve a better transient time than the derived worst-case upper bound.

We also compare the performance of EDAS with DSGD and DSGT for the logistic regression problem in Figure 6. On one hand, similar to problem (5.2), EDAS achieves significantly shorter transient times for ring networks when the network size is moderately large; see Figures 6a to 6c. On the other hand, for grid networks, EDAS and DSGT both achieve better transient times than DSGD; see Figures 6d to 6f.


6. Conclusions. In this paper, we consider a distributed stochastic gradient algorithm (EDAS) and perform a non-asymptotic analysis. In addition to showing that EDAS achieves asymptotic network independence, we identify its non-asymptotic convergence rate as a function of characteristic of the objective functions and the network. We show that EDAS achieves the fastest transient time so far to the best of our knowledge. Future work will explore the lower bound of transient times for a general class of distributed stochastic gradient methods.

Appendix A. Proofs.

A.1. Proof of Lemma 2.5.


Proof. Since W is a stochastic symmetric matrix, the matrices W and V have the following spectral decomposition:

$$W = Q\Lambda Q^\top, \quad V = Q\sqrt{I_n - \Lambda}Q^\top$$




(a) Transient time for the ring network topology. (b) Transient time for the grid network topology.

FIG. 5. Comparison between the transient times for EDAS and multiples of $\frac{n}{(1-\lambda_2)^{0.5}}$. The expected errors are approximated by averaging over 200 simulation results.



(a) Error terms over a ring net- (b) Error terms over a ring net- (c) Error terms over a ring net-
work with 20 nodes. work with 40 nodes. work with 60 nodes.



(d) Error terms over a grid net- (e) Error terms over a grid net- (f) Error terms over a grid net-
work with 49 nodes. work with 81 nodes. work with 121 nodes.

FIG. 6. Comparison among DSGT, DSGD, and EDAS over ring and grid network topologies for problem (5.3). The y-axis stands for $\mathbb{E}(\|x(k) - x^*\|^2)$ for SGD and $\frac{1}{n} \sum_{i=1}^n \mathbb{E}(\|x_{i,k} - x^*\|^2)$ for decentralized methods. The expected errors are approximated by averaging 100 simulation results.

where Q is an orthogonal matrix, and

$$\Lambda := \text{diag}(1, \lambda_2, \dots, \lambda_n).$$

Therefore,

$$(A.1) \quad B = \begin{pmatrix} W & -V \\ VW & W \end{pmatrix} = \begin{pmatrix} Q & \\ & Q \end{pmatrix} \begin{pmatrix} \Lambda & -\sqrt{I_n - \Lambda} \\ \sqrt{I_n - \Lambda} & \Lambda \end{pmatrix} \begin{pmatrix} Q^\top & \\ & Q^\top \end{pmatrix}.$$

Noticing the particular form of the middle term on the right hand side of (A.1),

there exists a permutation operator $\Pi_1 \in \mathbb{R}^{2n \times 2n}$ such that

$$\begin{pmatrix} \Lambda & -\sqrt{I_n - \Lambda} \\ \sqrt{I_n - \Lambda} & \Lambda \end{pmatrix} = \Pi_1 \text{diag}(E_1, E_2, \dots, E_n) \Pi_1^\top,$$

where

$$(A.2) \quad E_1 := I_2, \quad E_k := \begin{pmatrix} \lambda_k & -\sqrt{1 - \lambda_k} \\ \lambda_k \sqrt{1 - \lambda_k} & \lambda_k \end{pmatrix}, \quad k = 2, 3, \dots, n.$$

Hence

$$(A.3) \quad B = \begin{pmatrix} Q & \\ & Q \end{pmatrix} \Pi_1 \text{diag}(I_2, E_2, \dots, E_n) \Pi_1^\top \begin{pmatrix} Q^\top & \\ & Q^\top \end{pmatrix}.$$

We then consider the eigendecomposition of E_k for $k = 2, 3, \dots, n$. It can be verified that

$$(A.4) \quad E_k = Z_k \begin{pmatrix} \lambda_k + i\sqrt{\lambda_k - \lambda_k^2} & 0 \\ 0 & \lambda_k - i\sqrt{\lambda_k - \lambda_k^2} \end{pmatrix} Z_k^{-1},$$

where $i^2 = -1$, and

$$(A.5) \quad Z_k = \begin{pmatrix} 1 & 1 \\ -i\sqrt{\lambda_k} & i\sqrt{\lambda_k} \end{pmatrix}, \quad Z_k^{-1} = \begin{pmatrix} \frac{1}{2} & \frac{i}{2\sqrt{\lambda_k}} \\ \frac{1}{2} & -\frac{i}{2\sqrt{\lambda_k}} \end{pmatrix}.$$

Therefore, from (A.3), we have

$$(A.6) \quad B = \begin{pmatrix} Q & \\ & Q \end{pmatrix} \Pi_1 Z (\text{diag}(I_2, D_1)) Z^{-1} \Pi_1^\top \begin{pmatrix} Q^\top & \\ & Q^\top \end{pmatrix},$$

where $Z = \text{diag}(I_2, Z_1, Z_2, \dots, Z_n)$ and D_1 is a complex diagonal matrix.

Additionally, note that $(\mathbf{1}_n^\top, \mathbf{0}^\top)^\top$ and $(\mathbf{0}^\top, \mathbf{1}_n^\top)^\top$ are the two eigenvectors of B w.r.t. eigenvalue 1. Thus after a permutation Π_2 , we reach a decomposition for B given by $B = \tilde{U} D \tilde{U}^{-1}$, where

$$D := \text{diag}(I_2, D_1),$$

$$\tilde{U} := \begin{pmatrix} Q & \\ & Q \end{pmatrix} \Pi_1 Z \Pi_2 = \begin{pmatrix} \mathbf{1}_n & \mathbf{0} & U_{R,u} \\ \mathbf{0} & \mathbf{1}_n & U_{R,l} \end{pmatrix},$$

for some $U_{R,u}, U_{R,l} \in \mathbb{R}^{n \times (n-2)}$. We then further choose $c > 0$ as a free parameter, to obtain $B = U D U^{-1}$, where

$$U := \begin{pmatrix} \mathbf{1}_n & \mathbf{0} & cU_{R,u} \\ \mathbf{0} & \mathbf{1}_n & cU_{R,l} \end{pmatrix}, \quad U^{-1} := \begin{pmatrix} \frac{1}{n}\mathbf{1}_n^\top & \mathbf{0} \\ \mathbf{0} & \frac{1}{n}\mathbf{1}_n^\top \\ \frac{1}{c}U_{L,l} & \frac{1}{c}U_{L,r} \end{pmatrix}.$$

We next bound $\|U_R\|_2$ and $\|U_L\|_2$. By definition, we have

$$\|U_R\|_2 \leq \|\tilde{U}\|_2 \leq \left\| \begin{pmatrix} Q & \\ & Q \end{pmatrix} \right\|_2 \|\Pi_1\|_2 \|Z\|_2 \|\Pi_2\|_2 = \|Z\|_2.$$

Since

$$\|Z_k\|_2 = \sqrt{\lambda_{\max}(Z_k Z_k^*)} = \sqrt{\lambda_{\max} \begin{pmatrix} 2 & 0 \\ 0 & 2|\lambda_k| \end{pmatrix}} = \sqrt{2}, \quad \forall k = 2, 3, \dots, n,$$

we obtain $\|U_R\|_2 \leq \|Z\|_2 = \sqrt{2}$.

Similarly,

$$\|U_L\|_2 \leq \|\tilde{U}^{-1}\|_2 \leq \|Z^{-1}\|_2.$$

Given that

$$Z_k^{-1}(Z_k^{-1})^* = \frac{1}{4} \begin{pmatrix} 1 + \frac{1}{|\lambda_k|} & 1 - \frac{1}{|\lambda_k|} \\ 1 - \frac{1}{|\lambda_k|} & 1 + \frac{1}{|\lambda_k|} \end{pmatrix},$$

we have

$$\|Z_k^{-1}\|_2 = \sqrt{\lambda_{\max}(Z_k^{-1}(Z_k^{-1})^*)} = \sqrt{\frac{1}{2|\lambda_k|}}.$$

Hence $\|U_L\|_2 \leq \|Z^{-1}\|_2 = \sqrt{\frac{1}{2|\lambda_n|}} \leq \sqrt{\frac{1}{2\lambda}} by assuming $\lambda_n \geq \lambda > 0$. $\square$$

A.2. Proof of Lemma 4.2.

Proof. From the definitions of M_k, T_k in (3.19), we have

$$\begin{aligned} M_0 &= \|\bar{z}_0\|^2 = \left\| \frac{1}{n} \mathbf{1}_n^\top (\mathbf{x}_0 - \mathbf{x}^*) \right\|^2 = \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n} \right), \\ T_0 &= \left\| \frac{1}{c} (U_{L,l} \tilde{\mathbf{x}}_0 + U_{L,r} \tilde{\mathbf{y}}_0) \right\|^2 \leq \frac{2\|U_L\|^2}{c^2} (\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + \|\mathbf{y}_0 - \mathbf{y}_0^*\|^2). \end{aligned}$$

For $\|\mathbf{y}_0 - \mathbf{y}_0^*\|^2$, notice that $\mathbf{y}_0 = \mathbf{0}$ in Algorithm 2.1, and $\mathbf{y}_0^* = -V^{-1}\alpha_0 W \nabla F(\mathbf{x}^*)$ by Lemma 2.2, we have

$$\|\mathbf{y}_0 - \mathbf{y}_0^*\| = \|V^{-1}\alpha_0 W \nabla F(\mathbf{x}^*)\|.$$

Given that $\|V^{-1}\| = \left\| Q \left(\sqrt{I_n - \bar{\Lambda}} \right)^{-1} Q^\top \right\| = \sqrt{\frac{1}{1-\lambda_2}}$ and $\alpha_0 \leq \frac{1-\lambda_2}{\mu}$ from (3.22), we derive

$$T_0 = \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + (1 - \lambda_2) \|\nabla F(\mathbf{x}^*)\|^2}{n} \right).$$

Noting that $\omega_0 = \frac{24\theta c^2 \|U_R\|^2 L^2}{n\mu^2 m(1-\sqrt{\lambda_2})} = \mathcal{O}(1)$, it follows that

$$(A.7) \quad H_0 = M_0 + \omega_0 T_0 = \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2 + (1 - \lambda_2) \|\nabla F(\mathbf{x}^*)\|^2}{n} \right).$$

To estimate q_0, q_2, q_3, q_4 , consider

$$(A.8a) \quad p_2 = \frac{\theta^2 \bar{\sigma}^2}{n\mu^2} = \mathcal{O} \left(\frac{1}{n} \right),$$

$$(A.8b) \quad p_3 = \frac{24\lambda_2 \theta^3 \bar{\sigma}^2 \|U_L\|^2 \|U_R\|^2 L^2}{\mu^4 (1 - \sqrt{\lambda_2})} = \mathcal{O} \left(\frac{1}{1 - \lambda_2} \right),$$

$$(A.8c) \quad p_5 = \frac{96\theta^3 \|U_L\|^2 \|U_R\|^2 L^2 \|V^{-1}\|^2 \|\nabla F(\mathbf{x}^*)\|^2}{n\mu^4 (1 - \sqrt{\lambda_2})^2} = \mathcal{O} \left(\frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1 - \lambda_2)^3} \right)$$

Combining (A.8) and (A.7) leads to

$$\begin{aligned}\hat{H}_1 &= \frac{1}{\theta - 3} \left(p_2 + \frac{p_5}{m^3} \right) = \mathcal{O} \left(\frac{1 + \|\nabla F(\mathbf{x}^*)\|^2}{n} \right), \\ \hat{H}_2 &= m^2 H_0 + \frac{2p_3}{2\theta - 3} = \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1 - \lambda_2)^2} + \frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1 - \lambda_2)} + \frac{1}{1 - \lambda_2} \right).\end{aligned}$$

Therefore,

$$\begin{aligned}c_0 &= \frac{3\theta L^2 c^2 \|U_R\|^2}{n\mu^2} = \mathcal{O}(1), \quad \frac{1}{1 - q_0} = \frac{4}{1 - \sqrt{\lambda_2}} = \mathcal{O} \left(\frac{1}{1 - \lambda_2} \right), \\ q_2 &= \frac{\lambda_2 n \|U_L\|^2 \bar{\sigma}^2 \theta^2}{c^2 \mu^2} = \mathcal{O}(1), \quad q_3 = \frac{4\lambda_2 n \|U_L\|^2 \theta^2 L^2 \hat{H}_1}{\mu^2 c^2 (1 - \lambda_2)} = \mathcal{O} \left(\frac{1 + \|\nabla F(\mathbf{x}^*)\|^2}{n(1 - \lambda_2)} \right), \\ q_4 &= \frac{4\lambda_2 n \|U_L\|^2 \theta^2 L^2 \hat{H}_2}{\mu^2 c^2 (1 - \lambda_2)} + \frac{4\|U_L\|^2 \|V^-\|^2 \|\nabla F(\mathbf{x}^*)\|^2 \theta^2}{\mu^2 c^2 (1 - \sqrt{\lambda_2})} \\ &= \mathcal{O} \left(\frac{\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{n(1 - \lambda_2)^3} + \frac{\|\nabla F(\mathbf{x}^*)\|^2}{n(1 - \lambda_2)^2} + \frac{1}{(1 - \lambda_2)^2} \right).\end{aligned}$$

Acknowledgment. We would like to thank Bingyu Wang from The Chinese University of Hong Kong, Shenzhen for providing helpful feedback.

REFERENCES

- [1] M. ASSRAN, N. LOIZOU, N. BALLAS, AND M. RABBAT, *Stochastic gradient push for distributed deep learning*, arXiv preprint arXiv:1811.10792, (2018).
- [2] J. CHEN AND A. H. SAYED, *On the limiting behavior of distributed optimization strategies*, in 2012 50th Annual Allerton Conference on Communication, Control, and Computing (Allerton), IEEE, 2012, pp. 1535–1542.
- [3] J. CHEN AND A. H. SAYED, *On the learning behavior of adaptive networks—part i: Transient analysis*, IEEE Transactions on Information Theory, 61 (2015), pp. 3487–3517.
- [4] J. CHEN AND A. H. SAYED, *On the learning behavior of adaptive networks—part ii: Performance analysis*, IEEE Transactions on Information Theory, 61 (2015), pp. 3518–3548.
- [5] P. DI LORENZO AND G. SCUTARI, *Next: In-network nonconvex optimization*, IEEE Transactions on Signal and Information Processing over Networks, 2 (2016), pp. 120–136.
- [6] D. JAKOVETIĆ, J. XAVIER, AND J. M. MOURA, *Fast distributed gradient methods*, IEEE Transactions on Automatic Control, 59 (2014), pp. 1131–1146.
- [7] S. S. KIA, J. CORTÉS, AND S. MARTÍNEZ, *Distributed convex optimization via continuous-time coordination algorithms with discrete-time communication*, Automatica, 55 (2015), pp. 254–264.
- [8] A. KOLOSKOVA, S. U. STICH, AND M. JAGGI, *Decentralized stochastic optimization and gossip algorithms with compressed communication*, arXiv preprint arXiv:1902.00340, (2019).
- [9] Y. LECUN, L. BOTTOU, Y. BENGIO, AND P. HAFFNER, *Gradient-based learning applied to document recognition*, Proceedings of the IEEE, 86 (1998), pp. 2278–2324, <https://doi.org/10.1109/5.726791>.
- [10] Z. LI, Y. LIAO, K. HUANG, AND S. PU, *Compressed gradient tracking for decentralized optimization with linear convergence*, arXiv preprint arXiv:2103.13748, (2021).
- [11] Z. LI, W. SHI, AND M. YAN, *A decentralized proximal-gradient method with network independent step-sizes and separated convergence rates*, IEEE Transactions on Signal Processing, 67 (2019), pp. 4494–4506.
- [12] X. LIAN, C. ZHANG, H. ZHANG, C.-J. HSIEH, W. ZHANG, AND J. LIU, *Can decentralized algorithms outperform centralized algorithms? a case study for decentralized parallel stochastic gradient descent*, in Advances in Neural Information Processing Systems, 2017, pp. 5336–5346.

- [13] I. LOBEL, A. OZDAGLAR, AND D. FEIJER, *Distributed multi-agent optimization with state-dependent communication*, Mathematical programming, 129 (2011), pp. 255–284.
- [14] G. MORRAL, P. BIANCHI, AND G. FORT, *Success and failure of adaptation-diffusion algorithms for consensus in multi-agent networks*, in 53rd IEEE Conference on Decision and Control, 2014, pp. 1476–1481.
- [15] G. MORRAL, P. BIANCHI, AND G. FORT, *Success and failure of adaptation-diffusion algorithms with decaying step size in multiagent networks*, IEEE Transactions on Signal Processing, 65 (2017), pp. 2798–2813.
- [16] A. NEDIĆ, A. OLSHEVSKY, AND M. G. RABBAT, *Network topology and communication-computation tradeoffs in decentralized optimization*, Proceedings of the IEEE, 106 (2018), pp. 953–976.
- [17] A. NEDIĆ, A. OLSHEVSKY, AND W. SHI, *Achieving geometric convergence for distributed optimization over time-varying graphs*, SIAM Journal on Optimization, 27 (2017), pp. 2597–2633.
- [18] A. NEDIC AND A. OZDAGLAR, *Distributed subgradient methods for multi-agent optimization*, IEEE Transactions on Automatic Control, 54 (2009), pp. 48–61.
- [19] A. NEDIĆ, A. OZDAGLAR, AND P. A. PARRILO, *Constrained consensus and optimization in multi-agent networks*, IEEE Transactions on Automatic Control, 55 (2010), pp. 922–938.
- [20] S. PU AND A. GARCIA, *A flocking-based approach for distributed stochastic optimization*, Operations Research, 1 (2018), pp. 267–281.
- [21] S. PU AND A. GARCIA, *Swarming for faster convergence in stochastic optimization*, SIAM Journal on Control and Optimization, 56 (2018), pp. 2997–3020.
- [22] S. PU AND A. NEDIĆ, *Distributed stochastic gradient tracking methods*, Mathematical Programming, (2020), pp. 1–49.
- [23] S. PU, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *A sharp estimate on the transient time of distributed stochastic gradient descent*, 2019, <https://arxiv.org/abs/1906.02702>.
- [24] S. PU, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *Asymptotic network independence in distributed stochastic optimization for machine learning: Examining distributed and centralized stochastic gradient descent*, IEEE Signal Processing Magazine, 37 (2020), pp. 114–122.
- [25] S. PU, W. SHI, J. XU, AND A. NEDIĆ, *Push-pull gradient methods for distributed optimization in networks*, IEEE Transactions on Automatic Control, (2020).
- [26] G. QU AND N. LI, *Harnessing smoothness to accelerate distributed optimization*, IEEE Transactions on Control of Network Systems, (2017).
- [27] M. I. QURESHI, R. XIN, S. KAR, AND U. A. KHAN, *S-adapt: Decentralized stochastic first-order optimization over directed graphs*, IEEE Control Systems Letters, 5 (2020), pp. 953–958.
- [28] W. SHI, Q. LING, G. WU, AND W. YIN, *Extra: An exact first-order algorithm for decentralized consensus optimization*, SIAM Journal on Optimization, 25 (2015), pp. 944–966.
- [29] A. SPIRIDONOFF, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *Robust asynchronous stochastic gradient-push: Asymptotically optimal and network-independent performance for strongly convex functions.*, Journal of Machine Learning Research, 21 (2020), pp. 1–47.
- [30] H. TANG, X. LIAN, M. YAN, C. ZHANG, AND J. LIU, *D2: Decentralized training over decentralized data*, in International Conference on Machine Learning, 2018, pp. 4848–4856.
- [31] Z. J. TOWFIC, J. CHEN, AND A. H. SAYED, *Excess-risk of distributed stochastic learners*, IEEE Transactions on Information Theory, 62 (2016), pp. 5753–5785.
- [32] J. TSITSIKLIS, D. BERTSEKAS, AND M. ATHANS, *Distributed asynchronous deterministic and stochastic gradient optimization algorithms*, IEEE transactions on automatic control, 31 (1986), pp. 803–812.
- [33] R. XIN, U. A. KHAN, AND S. KAR, *An improved convergence analysis for decentralized online stochastic non-convex optimization*, arXiv preprint arXiv:2008.04195, (2020).
- [34] J. XU, S. ZHU, Y. C. SOH, AND L. XIE, *Convergence of asynchronous distributed gradient methods over stochastic networks*, IEEE Transactions on Automatic Control, 63 (2017), pp. 434–448.
- [35] K. YUAN, S. A. ALGHUNAIM, B. YING, AND A. H. SAYED, *On the performance of exact diffusion over adaptive networks*, in 2019 IEEE 58th Conference on Decision and Control (CDC), IEEE, 2019, pp. 4898–4903.
- [36] K. YUAN, S. A. ALGHUNAIM, B. YING, AND A. H. SAYED, *On the influence of bias-correction on distributed stochastic optimization*, IEEE Transactions on Signal Processing, 68 (2020), pp. 4352–4367.
- [37] K. YUAN, B. YING, X. ZHAO, AND A. H. SAYED, *Exact diffusion for distributed optimization and learning—part ii: Convergence analysis*, IEEE Transactions on Signal Processing, 67 (2018), pp. 724–739.