Fusion of ¹⁶O+¹⁶⁵Ho at deep sub-barrier energies

Saikat Bhattacharjee,^{1,3} A. Mukherjee,^{1,3,*} Ashish Gupta,^{1,3} Rajkumar Santra,^{1,3} D.

Chattopadhyay,^{1,†} N. Deshmukh,^{1,‡} Sangeeta Dhuri,^{2,3} Shilpi Gupta,^{2,3} V.V. Parkar,^{2,3} S.K.

Pandit,² K. Ramachandran,² K. Mahata,^{2,3} A.Shrivastava,^{2,3} Rebecca Pachuau,^{4,§} and S.Rathi⁵

¹Saha Institute of Nuclear Physics, 1/AF, Bidhan Nagar, Kolkata - 700064, India

²Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai - 400085, India

³Homi Bhabha National Institute, Anushaktinagar, Mumbai - 400094, India

⁵ Vivekanand Education Society's College of Arts, Science & Commerce, Mumbai - 400071, India

Fusion cross-sections have been measured for the asymmetric system ${}^{16}O+{}^{165}Ho$ at energies near and deep below the Coulomb barrier with an aim to investigate the occurrence of fusion hindrance for the system. Fusion cross sections down to ~ 700 nb have been measured using the off-beam γ ray technique. The fusion cross sections have been compared with the coupled channel calculations. Although the onset of fusion hindrance could not be observed experimentally, an indication of a small deviation of the experimental fusion cross-sections with respect to the calculated cross-sections could be observed at the lowest energy measured. However, the energy onset of fusion hindrance has been obtained from the extrapolation technique and is found to be about 2 MeV below the lowest energy of the present measurement.

I. INTRODUCTION

Extensive studies on fusion reactions at sub-barrier energies in different mass regions have unraveled the fundamentals of quantum mechanical tunnelling and distribution of potential barriers between two interacting nuclei. Fusion in the vicinity of Coulomb barrier is a probe to discern the indispensable role of different intrinsic features of the interacting nuclei on the reaction process. Enhancement of fusion cross sections observed in heavy-ion collisions at sub-barrier energies has been well explained by the coupled channels model [1, 2].

On extending the measurements from sub-barrier down to deep sub-barrier energies, for a wide range of reactions [3–19], a steep fall-off of fusion excitation function has been observed compared to the standard coupled channels calculations, although fusion cross sections are still enhanced with respect to single-barrier penetration model calculations. This phenomenon of change of slope in the fusion excitation function at deep sub-barrier energies is termed as "fusion hindrance". Observation of fusion hindrance at deep sub-barrier energies, especially in light systems like ${}^{12}C+{}^{12}C$ and ${}^{16}O+{}^{16}O$ [4] have astrophysical implications, as some of the light systems transpire in the late evolutionary stages of massive stars.

Fusion hindrance was initially observed in the symmetric medium-heavy system ${}^{60}\text{Ni}+{}^{89}\text{Y}$ [3], having negative Q-value. Subsequently, studies of several symmetric and nearly-symmetric [5–13] and asymmetric [14, 15]

systems over a wide range of mass and fusion Q-values have also exhibited similar observation. By contrast, fusion of weakly bound light projectiles ^{6,7}Li with ¹⁹⁸Pt does not manifest hindrance at deep sub-barrier energies [16, 17]. However, in fusion with relatively heavier projectiles ¹¹B and ¹²C on the targets ¹⁹⁷Au [18] and ¹⁹⁸Pt [16], respectively, hindrance has been observed. From these studies, it appears that for asymmetric heavy systems, fusion hindrance becomes increasingly significant with increasing mass and charge of the projectiles.

Several models with different physical foundations have been proposed to describe this phenomenon. Among them, the model developed by Misicu and Esbensen [20, 21] is based on sudden approximation. A soft repulsive core was incorporated with the density folded M3Y potential in this model, to consider the nuclear incompressibility in the overlapping region of the two interacting nuclei. The adiabatic model proposed by Ichikawa et al. [22, 23] introduces an additional damping factor on the nuclear coupling potential. The damping factor is a function of the internuclear distance, which takes into consideration of the smooth change from sudden to adiabatic transition while the two nuclei are going through fusion deep below the barrier. More recently, Simenel et al. have studied the effect of Pauli repulsion on heavy ion fusion by implementing the density-constrained frozen Hartree-Fock method [24]. Despite having different physical origins, these models have been quite successful in reproducing the experimental results for different systems at deep sub-barrier energies [4].

In the light of the problem of fusion hindrance, we have recently measured fusion cross-sections for ${}^{16}\text{O}+{}^{165}\text{Ho}$ system at energies near and deep below the Coulomb barrier. Owing to the large deformation of nuclei in the rare-earth region, systems involving such nuclei usually exhibit strong coupling effects between relative motion and internal degrees of freedom in the sub barrier fusion mechanism. The target nucleus ${}^{165}\text{Ho}$ has a large

⁴ Tata Institute of Fundamental Research, Mumbai - 400005, India

^{*} anjali.mukherjee@saha.ac.in

[†] Present adddress: Tata Institute of Fundamental Research, Mumbai - 400005, India

[‡] Present address: School of Sciences, Auro University, Surat, Gujarat - 394510, India

[§] Present address: Department of Physics, Banaras Hindu University, Varanasi 221005, India.

deformation parameter [25], whereas the projectile ¹⁶O is a tightly bound spherical nucleus and therefore subdues the projectile effect of different coupling schemes on the fusion mechanism. It has been perceived that stiff systems, where coupling effects are small, typically show fusion hindrance more readily than soft systems [4]. The dominance of the influence of different direct reaction channels, like inelastic scattering, transfer and breakup, in reactions involving soft nuclei are believed to be responsible for the occurrence of fusion hindrance at much lower energies than stiff systems [4]. The system $^{16}\mathrm{O}+^{165}\mathrm{Ho}$ lies between stiff and soft systems having a negative Q-value (Q=-23.1 MeV) for fusion. It would be interesting to study fusion mechanism in the system of tightly bound projectile ¹⁶O on the deformed nucleus ¹⁶⁵Ho, especially at deep sub-barrier energies.

The complete fusion(CF) cross sections of ${}^{16}\text{O}+{}^{165}\text{Ho}$ at above-barrier energies have been reported in the literature [26]. Present study overlaps some of the energies of the reported measurement. The measurement has been extended towards below barrier to deep sub-barrier energy region. An off-beam γ -ray detection technique has been implemented to measure the cross sections of the β -active evaporation residues. Section II recounts the detailed experimental method that have been implemented to perform the experiment. The procedure of data analysis have been explained in Section III. The details of theoretical coupled channel calculations and comparison with experimental results have been described in Section IV. Section V consists of a discussion followed by a summary of the present work in Section VI.

II. EXPERIMENTAL DETAILS

The experiment was performed at the 14UD BARC-TIFR Pelletron-LINAC facility, Mumbai. Selfsupporting, rolled, natural foils of ¹⁶⁵Ho, having thick-



FIG. 1. γ -ray spectrum of the evaporation residues arising from the complete fusion(CF) of $^{16}O+^{165}Ho$ system at 84 MeV beam energy. The red line is the background estimated by ROOT data analysis package [28].



FIG. 2. Activity of the ¹⁷⁸Re evaporation residue as a function of progressing time at $E_{lab}=70$ MeV. The dotted line corresponds to the activity obtained by fitting the data. The half life from the fit is mentioned in the graph, which is consistent with the known value.

ness in the range ~ 1.02 -1.9 mg/cm² were irradiated by beams of ¹⁶O, in the energy range $E_{lab}=62-85$ MeV. Each target foil of ¹⁶⁵Ho was followed by an Al catcher foil, sufficiently thick to stop the heavy evaporation residues (ERs) produced in the reaction. The thickness of each target and catcher foil was measured by the α -transmission method. For each irradiation, a fresh target-catcher foil assembly was used. Typical beam current during the irradiations was $\sim 2-10$ pnA. To correct for beam fluctuations during the irradiation, the beam current was recorded at regular intervals of 1 min using a CAMAC scaler. The energies of the incident beam were corrected for the loss of energy in the target material by employing SRIM [27], at half-thickness of the target. As all the ERs were β -active and yielded delayed γ -rays, the activation technique was employed to determine the fusion cross sections for the system. After each irradiation, the target-catcher foil assembly was removed from the chamber and placed in front of an efficiency calibrated HPGe detector, which detected the delayed γ -rays emitted by the ERs.

The target-catcher foil assembly was placed either at a distance of 10 cm from the face of the detector or on the face of the detector, depending on the activity of the irradiated sample. The energy calibration and absolute efficiency measurement of the detector were carried out using the standard radioactive sources, ¹⁵²Eu, ¹³³Ba and ⁶⁰Co, mounted in the same geometry as the target. The measurement was done in a low background setup with Pb-Cu graded shielding to reduce the background γ -rays. Data were recorded using a digital data acquisition system employing a CAEN N6724 digitizer and the data were analyzed using the ROOT data analysis framework [28]. A typical off-beam γ -ray spectrum, after the



FIG. 3. The cross-sections of different evaporation residues of compound nucleus for ${}^{16}\text{O}+{}^{165}\text{Ho}$ system. The dotted lines are the statistical model calculation (PACE4) results.

irradiation at $E_{lab} = 84$ MeV, is shown in Fig. 1. The complete fusion ERs ¹⁷⁶⁻¹⁷⁸Re occurring from the decay of the compound nucleus ¹⁸¹Re were uniquely identified from the characteristic γ -rays emitted by their daughter nuclei and by following the half-lives. Half lives of the ERs were measured from the time sliced yields of the characteristic γ -rays and compared with the previously measured values [29], to ensure the absence of any contribution from sources other than the complete fusion residues. The half-life plot for ¹⁷⁸Re at $E_{lab} = 70$ MeV has been shown in Fig. 2.

III. ANALYSIS AND RESULTS

If N_{γ} represents the number of counts under a particular γ -ray peak, corresponding to a given ER in the spectrum, then from the principle of radioactive decay the corresponding ER cross-section (σ_{ER}) is given as [30]:

$$\sigma_{ER} = \frac{N_{\gamma} \lambda e^{\lambda t_w}}{N_B N_T \epsilon_{\gamma} F_{\gamma} (1 - e^{-\lambda t_c}) (1 - e^{-\lambda t_{irr}})} \qquad (1)$$

where N_B is the number of incident nuclei, N_T is the number of target nuclei per unit area, λ is the decay constant of the ER, t_{irr} is the irradiation time, t_w is the time elapsed between the end of irradiation and the beginning of counting, t_c is the counting time, ϵ_{γ} is the efficiency of the detector for a given γ -ray energy, and F_{γ} is the absolute intensity of a γ -ray decay.

The γ -ray peak corresponding to ³⁴Cl, seen in the spectrum (Fig. 1), arises from the reaction of ¹⁶O with the Al catcher foil. Different pxn and αxn channels, corresponding to isotopes of W and Ta respectively, populated in the reaction of ¹⁶O+¹⁶⁵Ho are marked in the spectrum. The pxn channel can be populated as an ER of compound nucleus and also from the decay product of Re (e.g. ¹⁷⁷Re $\stackrel{\epsilon}{\rightarrow}$ ¹⁷⁷W). The procedure to calculate



FIG. 4. Complete fusion excitation function measured in this experiment and the previous measurement has been compared with CC calculations using Woods-Saxon potential form. The solid lines are the results obtained from the CC calculations. The arrow corresponds to the Coulomb barrier for this system. (Inset) Fusion cross-section at above barrier energies are fitted with Wong formula to obtain the optical model potential parameters of the system. The solid line is the fitted result.

TABLE I. Spectroscopic properties of the evaporation residues, resulting via CF process that have been used to calculate the CF cross-sections for the $\rm ^{16}O+\rm ^{165}Ho$ system.

Residue	$T_{1/2}(\min)$	J^{π}	$E_{\gamma}(\text{keV})$	$I^{\gamma}(\%)$
$^{178}\mathrm{Re}(3n)$	13.2	3^{+}	237.0	44.5
			106.0	23.4
$^{177}\mathrm{Re}(4n)$	14.0	$5/2^-$	196.9	8.4
			209.8	2.8
$^{176}\mathrm{Re}(5n)$	5.3	3^{+}	240.3	54.0
			109.1	25.0
174 Ta $(\alpha 3n)$	68.4	3^{+}	206.5	60.0

the true weight of pxn ERs from the cumulative crosssections have been discussed by Cavinato *et al.* [31]. In the energy region of the present measurement, it has been found that majority of the γ -ray peaks corresponding to pxn channels are due to the decay of Re nuclei. The contribution of pxn ERs, i.e. the direct decay product of compound nucleus falls within the error limit of respective cross-sections at different energies and thus have not been estimated rigorously. The αxn channels are populated either by incomplete fusion (ICF) or via the decay of compound nucleus(CF). To estimate the contribution of αxn channels arising from the CF residues, statistical model calculation has been performed using the PACE4 code [32].

To obtain a relatively better agreement between theo-



FIG. 5. Astrophysical S-factor for ${}^{16}O+{}^{165}Ho$ system compared with S-factor obtained from CC calculation.

retical and experimental cross-sections, the neutron and alpha potentials in the code have been modified according to Ref. [33] and Ref. [34], respectively. Fission barrier was fixed at 21.89 MeV following Ref. [26] and the level density parameter "k" was taken to be 9. The other parameters in the code were set to the default values. Calculated cross section for each partial wave by coupled channel calculation (detailed discussion in next section) has been fed as input to the PACE4 calculations. The production of ¹⁷⁴Ta via ICF is energetically possible at higher energies (80-85 MeV) but at lower energies, it becomes unlikely that ¹⁷⁴Ta will be populated as an ICF product. It has also been observed that within the energy range of the measurement, the experimental cross-sections of $\alpha 3n$ channel (¹⁷⁴Ta) reasonably agree with the statistical model, which exclusively deals with CF process. The experimentally measured cross-sections of other αxn channels are distinctly underestimated by the statistical model calculations. Similar trend has also been observed in Ref. [26]. As the PACE4 code has been able to reproduce the cross-sections of xn and $\alpha 3n$ channels relatively well, it can be inferred that all the other αxn channels, or at least an exceedingly significant part of them have been populated via incomplete fusion(ICF). Fig.3 shows a comparison of the statistical model calculations with the measured ER channel cross-sections.

The total fusion cross-sections have been determined by adding the measured cross sections of xn and $\alpha 3n$ channels and are shown in Fig. 4. Fusion cross section down to 678 nb has been measured in this work. Statistical errors, as well as errors ensuing from the measurement of beam current, target thickness and detector efficiency have been taken into account. The astrophysical S-factor, often used in nuclear astrophysics to study the low-energy behavior of nuclear reactions, is defined as,

$$S(E) = E\sigma(E)exp(2\pi\eta), \qquad (2)$$

where E is the center-of-mass energy, σ is the fusion cross-section and $\eta = Z_1 Z_2 e^2/\hbar v$ is Sommerfield parameter with v being the beam velocity. The experimental S-factor curve corresponding to the measured fusion cross-sections for the present system has been plotted in Fig. 5. Appearance of a maximum in the S-factor curve has been presented in previous studies at an energy where the hindrance in fusion cross-section sets in. In the present study, although no clear maximum could be seen in the S-factor curve within the measured energy range, an indication of a change of slope at the lowest energy could be observed. An alternative representation, the logarithmic slope of the fusion excitation function is defined as [3],

$$L(E) = d[ln(E\sigma)]/dE = \frac{1}{E\sigma} \frac{d(E\sigma)}{dE}$$
(3)

The values of L(E), extracted from the measured fusion cross sections for ${}^{16}O+{}^{165}Ho$ are plotted in Fig. 6. These representations are independent of any theoretical model and are alternative approaches to manifest any deviation in the slope of excitation function. From eqns. (2) and (3), one gets the relation,

$$\frac{dS}{dE} = S(E)[L(E) - \frac{\pi\eta}{E}]$$
(4)

The derivative $\frac{dS}{dE}$ becomes zero when S-factor becomes maximum, and from eqn. (3) one finds that this corresponds to the logarithmic derivative for constant Sfactor, $L_{cs}(E)$, given by:

$$L_{cs}(E) = \frac{\pi\eta}{E} \tag{5}$$

The dashed curve $L_{cs}(E)$ for the present system is shown in Fig. 6 by the dashed line. The experimental values of L(E) have been fitted with a function $A + B/E^{\frac{3}{2}}$ [36, 37], and is shown in Fig. 6 by the dotted line. The crossover point of the curves $L_{cs}(E)$ and fitted L(E) corresponds to the maximum of the S-factor curve and is related to the threshold energy (E_s) for the occurance of fusion hindrance [4]. In the measured energy regime of the present work, the fitted L(E) curve does not intersect the $L_{cs}(E)$ curve. However, it can be seen from Fig. 6 that an extrapolation of the fitted L(E) curve intersects the curve $L_{cs}(E)$ at the energy $E_s = 52.78 \pm 0.78$ MeV. The value of E_s calculated from the empirical equation [38] is 53.17 MeV. The estimation of E_s obtained from the touching point configuration in adiabatic picture [22, 23] is 56.1 MeV. The present measurement has been performed down to E = 54.78 MeV, which is ≈ 2 MeV above the extrapolated $\mathrm{E_s}$ value and 1.3 MeV below the prediction from adiabatic model. This shows that the E_s value for $^{16}\text{O}+^{165}\text{Ho}$ system will be lower than that predicted by the adiabatic model.

IV. COMPARISON WITH COUPLED CHANNELS CALCULATIONS

The data obtained in the present work have been analyzed in the framework of coupled channels (CC) calcu-



FIG. 6. Logarthmic derivative L(E) of the experimental fusion excitation function compared with CC calculations(dashed dot lines). The extrapolation of L(E) (dotted lines) intersect the L_{cs} curve at 52.78 MeV.

lations, using the code CCFULL [39]. These calculations require an initial set of potential parameters. They were obtained by fitting the fusion cross sections well above the barrier using Wong's formula[41], as the fusion cross sections in this energy regime are expected to be fairly insensitive to the form or magnitude of the couplings. The nuclear potential was taken to be of Woods-Saxon form,

$$V_n(r) = \frac{-V_0}{1 + exp[(r - r_0 A_P^{1/3} - r_0 A_T^{1/3})/a]}$$
(6)

where V_0 is the depth, r_0 is the radius parameter, and a is the diffuseness of the nuclear potential. The potential parameters were obtained by fixing a to be 0.63 fm, and varying r_0 and V_0 to obtain a good fit to the high energy part of the cross sections. The parameters thus obtained are: $V_0 = 102$ MeV, $r_0 = 1.15$ fm and $a_0 = 0.63$ fm. The inset in Fig. 4 shows the resulting fit. The potential parameters for the same system has also been compared with Ref. [40]; both results are found to be identical.

The CCFULL calculations in the no-coupling limit are shown by the dotted curve in Fig. 4, and are seen to underpredict the data, suggesting strong effects of deformation in the target nucleus. The CCFULL code estimates the effects of deformation by linear as well as non linear coupling to the pure rotational bands of the deformed nucleus. The target nucleus ¹⁶⁵Ho is deformed with a valence proton. The rotational states of ¹⁶⁵Ho may be considered to be built up by the coupling of the unpaired valence proton particle (or proton hole) with the $0^+, 2^+, 4^+, \dots$ rotational states of the neighboring even-even nucleus ¹⁶⁴Dy (or ¹⁶⁶Er). To remain within the model space of CCFULL, the excitation energies and deformation parameters for the target nuclues ¹⁶⁵Ho were taken to be the averages of those of the neighboring even-even



FIG. 7. Systematic representation of E_s as a function of system parameter ζ . The solid circles are measured and hollow circles are extrapolated values of E_s , taken from the tabulated values of Ref.[4, 16, 36]. The solid square represents the extrapolated value in this work. The hollow squares correspond to Ref.[10, 45] where the value of E_s can not be obtained either by extrapolation or direct measurements and an upper bound has been mentioned in Ref.[4].

nuclei ^{164}Dy and ^{166}Er [42, 43]. The resulting ground state rotational band upto 12^+ state $(\beta_2{=}0.32$, $\beta_4{=}0.02$) was included in the calculations. The results of these calculations are shown by the solid curve in the figure. They are seen to be in fairly good agreement with the data, except the lowest energy data which appears to be slightly below the calculated value.

The solid curve in Fig. 5 shows the calculated astrophysical S-factor values extracted from the CC cross sections and are found to agree well with the experimental value, except the lowest energy point. Although no clear maximum in the experimental S-factor plot or deviation from the the CC calculations could be observed in the measured energy range, an indication of a small deviation could be seen at the lowest energy measured. The dot-dashed curve in Fig. 6 shows the calculated L(E) values extracted from the CC cross sections and and are also seen to agree well with the experimental values, although the trend of the two extreme low energy points show an indication of possible deviation at still lower energies.

V. DISCUSSION

For a systematic study of the energy onset of fusion hindrance, the value of E_s predicted for the system $^{16}O+^{165}Ho$ in the present work has been compared in Fig. 7 with the E_s values of other heavy systems as a function of the system parameter $\zeta = Z_p Z_t \sqrt{\frac{A_p A_t}{A_p + A_t}}$. The parameters A_p, Z_p, A_t, Z_t are the mass number and atomic number of projectile and target. In this comparison, we have considered only the systems in the ζ range \sim 1500-3000, for which E_s are available

in the literature. The parameter ζ contains information of the mass and Coulomb barrier of the targetprojectile system. The dashed line in the figure corresponds to the empirical form of E_s [38]. The extrapolated value of E_s obtained in the present work lies very close to the result obtained from the empirical form $E_s^{emp} = [0.495\zeta/(2.33 + 580/\zeta)]^{2/3}$; although it should be pointed out that an actual measurement down to the predicted threshold energy will confirm the result. On the other hand, the E_s limit, obtained from the adiabatic model, has been crossed but hindrance for this system was not experimentally observed within the measured energy range.



FIG. 8. Comparison of normalized fusion excitation function between $^{16}\rm O+^{165}Ho$, $^{19}\rm F+^{165}Ho$ [44], and $^{7}\rm Li+^{165}Ho$ [46] system.

A systematic comparison of fusion excitation functions have been carried out for systems involving the target nucleus ¹⁶⁵Ho and different projectiles and is shown in Fig. 8. The systems involving ¹⁶⁵Ho target nucleus, for which sub-barrier fusion cross sections have been reported in literature are ¹⁹F+¹⁶⁵Ho [44] and ⁷Li+¹⁶⁵Ho [46]. The projectile ¹⁶O in the present system is a stiff nucleus, while ¹⁹F is relatively heavier and less bound than ¹⁶O, and ⁷Li is a well known weakly bound stable nucleus. For comparison of different projectile-target systems, the fusion excitation functions have been plotted in a reduced scale. Barrier radii $R_b=r_0(A_p^{\frac{1}{3}}+A_t^{\frac{1}{3}})$ and Coulomb barrier V_b for each system was obtained by performing 1DBPM model calculation, using the Akyüz-Winther parametrization of Woods-Saxon potential [47].

Fig. 8 shows that the reduced fusion excitation func-

tions for the three systems overlap reasonably well with each other at above barrier energies. But there are no reported fusion cross sections at deep-sub-barrier energies for the systems $^{19}\mathrm{F}+^{165}\mathrm{Ho}$ and $^{7}\mathrm{Li}+^{165}\mathrm{Ho}$. At sub-barrier energies, the fusion cross-sections for $^{19}\mathrm{F}+^{165}\mathrm{Ho}$ and $^{16}\mathrm{O}+^{165}\mathrm{Ho}$ agree fairly well with each other; while

TABLE II. V_b and R_b of the systems that have been used in the reduced plot (Fig. 8).

System	V_b (MeV)	R_b (fm)
$^{16}\mathrm{O}{+}^{165}\mathrm{Ho}$	66.34	10.93
$^{19}{ m F}{+}^{165}{ m Ho}$	72.24	11.26
$^{7}\mathrm{Li}\mathrm{+}^{165}\mathrm{Ho}$	25.0	10.76

the fusion cross-sections for ${}^{7}\text{Li}+{}^{165}\text{Ho}$ are enhanced in the reduced scale. It would be interesting to see how the deep sub-barrier fusion cross sections for the two systems compare with the present ${}^{16}\text{O}+{}^{165}\text{Ho}$ system.

VI. SUMMARY

The fusion cross-sections for ${}^{16}O+{}^{165}Ho$ have been measured down to 678 nb, from above-barrier to deep sub-barrier energies. The present measurement agrees with the earlier reported data [26], in the overlapping above-barrier energy region. Statistical model calculation of different evaporation residues occurring from CF vielded similar results with experimental data. The experimental CF cross-sections have been well reproduced by the coupled channels calculations. No clear evidence of fusion hindrance has been observed in the fusion excitation function in the measured energy range of the present work. However, the trend of the extreme low energy points in the S-factor and L(E) plots show an indication of possible deviation at still lower energies. The extrapolation of L(E) suggests the onset of fusion hindrance at 52.78 ± 0.78 MeV, approximately 2 MeV lower than the lowest energy measured in the present work, in accordance with the prediction from systematics. Further measurement at lower energies may confirm the result.

Acknowledgments

We sincerely thank the staff of BARC-TIFR Pelletron facility for an uninterrupted supply of beam. We would like to extend our gratitude towards Dr. Sanjoy Pal of TIFR, Mumbai; for helping with the digital data acquisition.

- A.B. Balantekin and N. Takigawa, Rev. Mod. Phys. 70, 77 (1998), and references therein.
- [2] M. Dasgupta, D.J. Hinde, N. Rowley, and A.M. Stefanini, Annu. Rev. Nucl. Part. Sci. 48, 401 (1998), and refer-

ences therein.

- [3] C. L. Jiang, H. Esbensen, K. E. Rehm, B. B. Back, R. V. F. Janssens, J. A. Caggiano, P. Collon, J. Greene, A. M. Heinz, D. J. Henderson, I. Nishinaka, T. O. Pennington, and D. Seweryniak, Phys. Rev. Lett. 89, 052701 (2002).
- [4] B. B. Back, H. Esbensen, C. L. Jiang, and K. E. Rehm, Rev. Mod. Phys. 86, 317 (2014), and references therein.
- [5] C. L. Jiang, K. E. Rehm, R. V. F. Janssens, H. Esbensen, I. Ahmad, B. B. Back, P. Collon, C. N. Davids, J. P. Greene, D. J. Henderson, G. Mukherjee, R. C. Pardo, M. Paul, T. O. Pennington, D. Seweryniak, S. Sinha, and Z. Zhou, Phys. Rev. Lett. **93**, 012701 (2004).
- [6] C. L. Jiang, K. E. Rehm, H. Esbensen, R. V. F. Janssens, B. B. Back, C. N. Davids, J. P. Greene, D. J. Henderson, C. J. Lister, R. C. Pardo, T. Pennington, D. Peterson, D. Seweryniak, B. Shumard, S. Sinha, X. D. Tang, I. Tanihata, S. Zhu, P. Collon, S. Kurtz, and M. Paul, Phys. Rev. C 71, 044613 (2005).
- [7] A. M. Stefanini, G. Montagnoli, L. Corradi, S. Courtin, E. Fioretto, A. Goasduff, F. Haas, P. Mason, R. Silvestri, P. P. Singh, F. Scarlassara, and S. Szilner, Phys. Rev. C 82, 014614 (2010).
- [8] C. L. Jiang, A. M. Stefanini, H. Esbensen, K. E. Rehm, L. Corradi, E. Fioretto, P. Mason, G. Montagnoli, F. Scarlassara, R. Silvestri, P. P. Singh, S. Szilner, X. D. Tang, and C. A. Ur, Phys. Rev. C 82, 041601 (2010).
- [9] G. Montagnoli, A. M. Stefanini, C. L. Jiang, H. Esbensen et al., L. Corradi, S. Courtin, E. Fioretto, A. Goasduff, F. Haas, A. F. Kifle, C. Michelagnoli, D. Montanari, T. Mijatović, K. E. Rehm, R. Silvestri, Pushpendra P. Singh, F. Scarlassara, S. Szilner, X. D. Tang, and C. A. Ur, Phys. Rev. C 85, 024607 (2012).
- [10] G. Montagnoli, A. M. Stefanini, L. Corradi, S. Courtin, E. Fioretto, F. Haas, D. Lebhertz, F. Scarlassara, R. Silvestri, and S. Szilner, Phys. Rev. C 82, 064609 (2010).
- [11] G. Montagnoli, A. M. Stefanini, H. Esbensen, C. L. Jiang, L. Corradi, S. Courtin, E. Fioretto, A. Goasduff, J. Grebosz, F. Haas, M. Mazzocco, C. Michelagnoli, T. Mijatovic, D. Montanari, C. Parascandolo, K. E. Rehm, F. Scarlassara, S. Szilner, X. D. Tang, and C. A. Ur, Phys. Rev. C 87, 014611 (2013).
- [12] C. L. Jiang, A. M. Stefanini, H. Esbensen, K. E. Rehm, S. Almaraz-Calderon, B. B. Back, L. Corradi, E. Fioretto, G. Montagnoli, F. Scarlassara, D. Montanari, S. Courtin, D. Bourgin, F. Haas, A. Goasduff, S. Szilner, and T. Mijatovic, Phys. Rev. Lett. **113**, 022701 (2014).
- [13] C. L. Jiang, B. B. Back, H. Esbensen *et al.*, J. P. Greene, R. V. F. Janssens, D. J. Henderson, H. Y. Lee, C. J. Lister, M. Notani, R. C. Pardo, N. Patel, K. E. Rehm, D. Seweryniak, B. Shumard, X. Wang, S. Zhu, S. Mişicu, P. Collon, and X. D. Tang, Phys. Rev. C **78**, 017601 (2008).
- [14] M. Dasgupta, D. J. Hinde, A. Diaz-Torres, B. Bouriquet, Catherine I. Low, G. J. Milburn, and J. O. Newton, Phys. Rev. Lett. 99, 192701 (2007).
- [15] A. M. Stefanini, G. Montagnoli, H. Esbensen, L. Corradi, S. Courtin, E. Fioretto, A. Goasduff, J. Grebosz, F. Haas, M. Mazzocco, C. Michelagnoli, T. Mijatovic, D. Montanari, G. Pasqualato, C. Parascandolo, F. Scarlassara, E. Strano, S. Szilner, and D. Torresi, Phys. Lett. B **728**, 639 (2014).
- [16] A. Shrivastava, K. Mahata, S. K. Pandit, V. Nanal, T. Ichikawa, K. Hagino, A. Navin, C. S. Palshetkar, V. V. Parkar, K. Ramachandran, P. C. Rout, A. Kumar, A.

Chatterjee, and S. Kailas, Phys. Lett. B **755**, 332 (2016).

- [17] A. Shrivastava, A. Navin, A. Lemasson, K. Ramachandran, V. Nanal, M. Rejmund, K. Hagino, T. Ichikawa, S. Bhattacharyya, A. Chatterjee, S. Kailas, K. Mahata, V. V. Parkar, R. G. Pillay, and P. C. Rout, Phys. Rev. Lett. **103**, 232702 (2009).
- [18] A. Shrivastava, K. Mahata, V. Nanal, S. K. Pandit, V. V. Parkar, P. C. Rout, N. Dokania, K. Ramachandran, A. Kumar, A. Chatterjee and S. Kailas, Phys. Rev. C 96, 034620 (2017).
- [19] G. Montagnoli, A. M. Stefanini, C. L. Jiang, K. Hagino, F. Galtarossa, G. Colucci, S. Bottoni, C. Broggini, A. Caciolli et al., Phys. Rev. C 97, 024610 (2018).
- [20] S. Mişicu and H. Esbensen, Phys. Rev. Lett. 96, 112701 (2006).
- [21] S. Mişicu and H. Esbensen, Phys. Rev. C 75, 034606 (2007).
- [22] T. Ichikawa, K. Hagino, and A. Iwamoto, Phys. Rev. Lett. 103, 202701 (2009).
- [23] T. Ichikawa, Phys. Rev. C 92, 064604 (2015).
- [24] C. Simenel, A. S. Umar, K. Godbey, M. Dasgupta, and D. J. Hinde, Phys. Rev. C 95, 031601(R) (2017).
- [25] P. Möller, A. J. Sierk, T. Ichikawa, and H. Sagawa, At. Data Nucl. Data Tables, 109-110, 1 (2016).
- [26] Kamal Kumar, Tauseef Ahmad, Sabir Ali, I. A. Rizvi, Avinash Agarwal, R. Kumar, K. S. Golda and A. K. Chaubey, Phys. Rev. C 87, 044608 (2013).
- [27] J. F.Ziegler, M. D. Ziegler, and J. P. Biersack, Nucl. Instrum. Methods B 268, 1818 (2010).
- [28] Rene Brun, and Fons Rademakers, Nucl. Instrum. Methods Phys. Res. A 389, 81 (1997).
- [29] https://www.nndc.bnl.gov.
- [30] P.R.S. Gomes, and T.J.P. Penna, Nucl. Instrum. Methods Phys. Res. A 280, 395 (1989).
- [31] M. Cavinato, E. Fabrici, E. Gadioli, E. Gadioli Erba, P. Vergani, M. Crippa, G. Colombo, I. Redaelli, and M. Ripamonti, Phys. Rev C 52, 2577 (1995).
- [32] A. Gavron, Phys. Rev. C 21, 230 (1980).
- [33] R. L. Walter and P. P. Guss, Radiafron Effects, 95/1-4, 73 (1986).
- [34] L. McFadden, and G. R. Satchler, Nucl. Phys. 84, 177 (1966).
- [35] N. Rowley, G. R. Satchler and P. H. Stelson, Phys. A Lett. B 254, 25 (1991).
- [36] C. L. Jiang, B. B. Back, H. Esbensen, R. V. F. Janssens, and K. E. Rehm, Phys. Rev. C 73, 014613 (2006).
- [37] C. L. Jiang, B. B. Back, R. V. F. Janssens, and K. E. Rehm, Phys. Rev. C 75, 057604 (2007).
- [38] C. L. Jiang, K. E. Rehm, B. B. Back, and R. V. F. Janssens, Phys. Rev. C 79, 044601 (2009).
- [39] K. Hagino, N. Rowley, and A. T. Kruppa, Comput. Phys. Commun. **123**, 143 (1999).
- [40] K. Hagino and S. Sakaguchi, Phys. Rev. C, 100, 064614 (2019).
- [41] C. Y. Wong, Phys. Rev. Lett. **31**, 12 (1973).
- [42] A. Mukherjee, Subinit Roy, M.K. Pradhan, M. Saha Sarkar, P. Basu, B. Dasmahapatra, T. Bhattacharya, S. Bhattacharya, S.K. Basu, A. Chatterjee, V. Tripathi, S. Kailas, Phys. Lett. B 636, 91 (2006).
- [43] S. Raman, C. H. Malarkey, W. T. Milner, C. W. Nestor, Jr, and, P. H. Stelson, At. Data Nucl. Data Tables 36, 1 (1987).
- [44] A. Navin, A. Chatterjee, S. Kailas, A. Shrivastava, P. Singh, and S. S. Kapoor, Phys. Rev. C54, 767(1996).

- [45] A.M. Stefanini, G. Montagnoli, R. Silvestri, L. Corradi, S. Courtin, E. Fioretto, B. Guiot, F. Haas, D. Lebhertz, P. Mason, F. Scarlassara, S. Szilner, Phys. Lett. B 679, 95 (2009).
- [46] V. Tripathi, A. Navin, K. Mahata, K. Ramachandran, A. Chatterjee, and S. Kailas, Phys. Rev. Lett. 88, 17 (2002).
- [47] O. Akyüz and A. Winther, Proceedings of the International School of Physics "Enrico Fermi", Course LXXVII, edited by R. A. Broglia, R. A. Ricci, and C. H. Dasso (North-Holland, Amsterdam), p. 492 (1981).