WELL-POSEDNESS FOR HYPERBOLIC EQUATIONS WHOSE COEFFICIENTS LOSE REGULARITY AT ONE POINT

DANIELE DEL SANTO AND MARTINO PRIZZI

ABSTRACT. We prove some C^∞ and Gevrey well-posedness results for hyperbolic equations whose coefficients lose regularity at one point.

1. INTRODUCTION

In this paper we deal with the well-posedness of the Cauchy problem for a linear hyperbolic operator whose coefficients depend only on time. Namely, we consider the equation

(1.1)
$$u_{tt} - \sum_{i,j=1}^{n} a_{ij}(t) u_{x_i x_j} = 0$$

in $[0,T] \times \mathbb{R}^n$, with initial data

(1.2)
$$u(0,x) = u_0(x), \quad u_t(0,x) = u_1(x)$$

in \mathbb{R}^n . The matrix $(a_{ij})_{i,j}$ is supposed to be real and symmetric. Setting

(1.3)
$$a(t,\xi) := \sum_{i,j=1}^{n} a_{ij}(t)\xi_i\xi_j/|\xi|^2, \quad (t,\xi) \in [0,T] \times (\mathbb{R}^n \setminus \{0\}),$$

we assume throughout that $a(\cdot,\xi) \in L^{\infty}(0,T)$ for all $\xi \in \mathbb{R}^n \setminus \{0\}$. Moreover, we suppose that the equation (1.1) is strictly hyperbolic, i.e.

(1.4)
$$\Lambda_0 \ge a(t,\xi) \ge \lambda_0 > 0$$

for all $(t,\xi) \in [0,T] \times (\mathbb{R}^n \setminus \{0\}).$

It is a classical result that if the coefficients $a_{ij}(t)$'s are real integrable functions, then the Cauchy problem (1.1), (1.2) is well posed in $\mathcal{A}'(\mathbb{R}^n)$, the space of real analytic functionals; moreover, if the initial data vanish in a ball, then the solution vanishes in a cone, whose slope depends on the coefficients $a_{ij}(t)$'s (see [1, Theorems 1 and 3.a]). On this basis,

²⁰²⁰ Mathematics Subject Classification. 35L10, 35A22, 46F12.

Key words and phrases. Gevrey space, well-posedness, strictly hyperbolic, modulus of continuity.

various well-posedness results can be proved by mean of the Paley-Wiener theorem (in the version of [1, p. 517], to which we refer here and throughout) and some energy estimates. If the coefficients $a_{ij}(t)$'s are Lipschitz-continuous then the Cauchy problem (1.1), (1.2) is well posed in Sobolev spaces. Relaxing this regulatity assumption, one has that if the $a_{ij}(t)$'s are Log-Lipschitz-continuous or Hölder-continuous of index α , then (1.1), (1.2) is well posed in C^{∞} or in the Gevrey space $\gamma^{(s)}$ for $s < \frac{1}{1-\alpha}$ respectively (see [1, Theorem 3.b,c]). Suitable counterexamples show that in each case the regularity assumption on the $a_{ij}(t)$'s is sharp for the well posedness of (1.1), (1.2) in the corresponding function space.

It is a remarkable fact that in the above mentioned counterexamples the coefficients $a_{ij}(t)$'s are in fact C^{∞} for $t \neq 0$, and each time the specific regularity fails only at t = 0. In [2] the authors showed that a control on the rate of the loss of Lipschitz regularity of the $a_{ij}(t)$'s as $t \to 0$ allows to recover well-posedness of (1.1), (1.2) in suitable function spaces. To be more specific, if the $a_{ij}(t)$'s are of class C^1 in [0,T] and $|a'_{ij}(t)| \leq Ct^{-p}$, then (1.1), (1.2) is well posed in C^{∞} when p = 1, and in the Gevrey space $\gamma^{(s)}$ for $s < \frac{p}{p-1}$ when p > 1. Concerning C^{∞} well-posedness, it was proved in [3] that a control on the second derivative of the a_{ij} 's as $t \to 0$ allows to relax slightly the growth assumption on the first derivative up to $|a'_{ij}(t)| \leq Ct^{-1}|\log t|$. In [5] some of the above results were extended to the case in which the coefficients a_{ij} 's depend also on the x variable in C^{∞} fashion.

In this paper we consider non Lipschitz coefficients whose regularity is ruled by a modulus of continuity μ , with a constant which blows up as $t \to 0$. More precisely, we assume that

(1.5)
$$|a_{ij}(t+\tau) - a_{ij}(t)| \le \frac{C}{\nu(t)}\mu(\tau), \quad 0 \le \tau \le \tau_0, \quad t, t+\tau \in [0,T],$$

where $\nu(t)^{-1}$ is possibly non integrable at t = 0 and where μ -continuity is possibly strictly weaker than Lipschitz continuity. We investigate how the interaction between ν and μ affects the well-posedness of (1.1), (1.2).

In Section 2 we prove a technical regularization result for the coefficients a_{ii} 's.

In Section 3 we consider locally Hölder continuous coefficients satisfying

(1.6)
$$|a_{ij}(t+\tau) - a_{ij}(t)| \le \frac{C}{t^p} \tau^{\alpha}, \quad 0 \le \tau, \quad t, t+\tau \in]0, T]$$

with $0 < \alpha < 1$ and p > 1, and we obtain well-posedness in the Gevrey space $\gamma^{(\sigma)}$ for $\sigma < \frac{p}{p-\alpha}$, a condition which fits perfectly with the ones of [1] and [2].

In Section 4 we consider the problem of C^{∞} well-posedness and we identify a precise relation between μ and ν which guarantees the latter. In particular we obtain well-posedness for coefficients satisfying (1.7)

$$|a_{ij}(t+\tau) - a_{ij}(t)| \le \frac{C}{t|\log t|} \frac{\tau|\log \tau|}{\log|\log \tau|}, \quad 0 \le \tau, \quad t, t+\tau \in]0, T],$$

where one can easily see that $\nu(t)^{-1}$ is non integrable and μ -continuity is strictly weaker than Lipschitz continuity. Also in this situation the results fits with the ones contained in [1] and [2] and contain them as particular cases.

2. Approximation

We begin by recalling the notion of *modulus of continuity*.

Definition 1. Let $\tau_0 > 0$. A function $\mu : [0, \tau_0] \rightarrow [0, +\infty[$ is a modulus of continuity if it is continuous, concave, strictly increasing and $\mu(0) = 0$.

Let μ be a modulus of continuity and let $a: [0,T] \to \mathbb{R}$ be a bounded function. Without loss of generality we can assume that $\tau_0 \leq T$. We assume that

(2.1)
$$|a(t+\tau) - a(t)| \le \frac{C}{\nu(t)}\mu(\tau), \quad 0 \le \tau \le \tau_0, \quad t, t+\tau \in [0,T],$$

where $\nu: [0,T] \to [0,+\infty[$ is a non-decreasing continuous function such that, for some $\kappa > 0$,

(2.2)
$$\nu(t/2) \ge \kappa \nu(t), \quad t \in]0, T].$$

Remark 1. Condition (2.2) is satisfied whenever ν is concave. Moreover, it is satisfied by $\nu(t) = t^p$ for every real exponent p > 0. On the other hand, it is not satisfied if $\nu(t)$ tends to 0 too fast as $t \to 0$, e.g. by $\nu(t) = e^{-1/t}$.

Now let $0 < \epsilon \leq \tau_0 \leq T$ and define

(2.3)
$$\tilde{a}_{\epsilon}(t) := \begin{cases} a(\epsilon) & \text{for } t \leq \epsilon, \\ a(t) & \text{for } \epsilon \leq t \leq T, \\ a(T) & \text{for } T \leq t. \end{cases}$$

Let $\rho \in C^{\infty}(\mathbb{R})$ with $\operatorname{supp} \rho \subset [-1,1]$, $\rho(s) \ge 0$, $\int_{\mathbb{R}} \rho(s) ds = 1$, set $\rho_{\epsilon}(s) := \frac{1}{\epsilon} \rho(\frac{s}{\epsilon})$, and define

(2.4)
$$a_{\epsilon}(t) := \int_{-\epsilon}^{\epsilon} \rho_{\epsilon}(s) \tilde{a}_{\epsilon}(t-s) \, ds, \quad t \in \mathbb{R}.$$

We have the following

Proposition 2. Under the above hypotheses, there exist constants C' and C'' > 0 such that, for $0 < \epsilon \leq \tau_0$,

(2.5)
$$|a_{\epsilon}(t) - \tilde{a}_{\epsilon}(t)| \le C' \min\left\{1, \frac{1}{\nu(t)}\mu(\epsilon)\right\}, \quad t \in]0, T]$$

and

(2.6)
$$|a'_{\epsilon}(t)| \leq \frac{C''}{\epsilon} \min\left\{1, \frac{1}{\nu(t)}\mu(\epsilon)\right\}, \quad t \in [0, T].$$

The constants C' and C'' depend only on C, ρ , κ and $||a||_{\infty}$.

Proof. We have

$$|a_{\epsilon}(t) - \tilde{a}_{\epsilon}(t)| = \left| \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s)(\tilde{a}_{\epsilon}(s) - \tilde{a}_{\epsilon}(t)) \, ds \right|$$
$$\leq \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s)|\tilde{a}_{\epsilon}(s) - \tilde{a}_{\epsilon}(t)| \, ds.$$

If $t \ge 2\epsilon$, then $t - \epsilon \ge t/2 \ge \epsilon$, so $\nu(t - \epsilon) \ge \nu(t/2) \ge \kappa \nu(t)$. Therefore, we have

$$\begin{aligned} |a_{\epsilon}(t) - \tilde{a}_{\epsilon}(t)| &\leq \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) |a(s) - a(t)| \, ds \\ &\leq \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) \frac{C}{\nu(t-\epsilon)} \mu(|s-t|) \, ds \\ &\leq \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) \frac{C}{\nu(t/2)} \mu(|s-t|) \, ds \\ &\leq \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) \frac{C/\kappa}{\nu(t)} \mu(|s-t|) \, ds \leq \frac{C/\kappa}{\nu(t)} \mu(\epsilon). \end{aligned}$$

If $0 < t \le \epsilon$, then $\tilde{a}_{\epsilon}(s) = \tilde{a}_{\epsilon}(t) = a(\epsilon)$ for $s \le \epsilon$, and therefore,

$$\begin{aligned} |a_{\epsilon}(t) - \tilde{a}_{\epsilon}(t)| &\leq \int_{\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) |a(s) - a(\epsilon)| \, ds \\ &\leq \int_{\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) \frac{C}{\nu(\epsilon)} \mu(|s-\epsilon|) \, ds \\ &\leq \int_{\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) \frac{C}{\nu(t)} \mu(\epsilon) \, ds \leq \frac{C}{\nu(t)} \mu(\epsilon). \end{aligned}$$

If $\epsilon \leq t \leq 2\epsilon$, then $0 \leq t - \epsilon \leq \epsilon \leq t$. Therefore, we have

$$\begin{aligned} |a_{\epsilon}(t) - \tilde{a}_{\epsilon}(t)| &\leq \int_{t-\epsilon}^{\epsilon} \rho_{\epsilon}(t-s) |a(\epsilon) - a(t)| \, ds + \int_{\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) |a(s) - a(t)| \, ds \\ &\leq \int_{t-\epsilon}^{\epsilon} \rho_{\epsilon}(t-s) \frac{C}{\nu(\epsilon)} \mu(|t-\epsilon|) \, ds + \int_{\epsilon}^{t+\epsilon} \rho_{\epsilon}(t-s) \frac{C}{\nu(\epsilon)} \mu(|t-s|) \, ds \\ &\leq \frac{C}{\nu(\epsilon)} \mu(\epsilon) \leq \frac{C}{\nu(t/2)} \mu(\epsilon) \leq \frac{C/\kappa}{\nu(t)} \mu(\epsilon). \end{aligned}$$

The thesis follows setting $C' := \max\{C, C/\kappa, 2 \|a\|_{\infty}\}.$

In order to estimate a'_{ϵ} , we observe that

$$\begin{aligned} |a_{\epsilon}'(t)| &= \left| \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}'(t-s)\tilde{a}_{\epsilon}(s) \, ds \right| = \left| \int_{t-\epsilon}^{t+\epsilon} \rho_{\epsilon}'(t-s)(\tilde{a}_{\epsilon}(s) - \tilde{a}_{\epsilon}(t)) \, ds \right| \\ &\leq \int_{t-\epsilon}^{t+\epsilon} |\rho_{\epsilon}'(t-s)| |\tilde{a}_{\epsilon}(s) - \tilde{a}_{\epsilon}(t)| \, ds. \end{aligned}$$

Then we proceed as above, noticing that $\rho'_{\epsilon}(t) = \frac{1}{\epsilon^2} \rho'(\frac{t}{\epsilon})$, and hence

$$\int_{t-\epsilon}^{t+\epsilon} |\rho_{\epsilon}'(t-s)| \, ds = \frac{\|\rho'\|_{L^1}}{\epsilon}.$$

The thesis follows setting $C'' := \|\rho'\|_{L^1} \max\{C, C/\kappa, \|a\|_{\infty}\}.$

3. Well posedness in Gevrey spaces

In this section we shall prove that if the coefficients a_{ij} 's are locally Hölder continuous of exponent α , with a Hölder constant which grows like t^{-p} as $t \to 0$, then the Cauchy problem (1.1), (1.2) is well posed in a suitable Gevrey space $\gamma^{(\sigma)}$, where σ depends on α and p.

As we pointed out in the Introduction, since the coefficients a_{ij} 's are real integrable functions, the Cauchy problem (1.1), (1.2) is well posed in $\mathcal{A}'(\mathbb{R}^n)$, the space of real analytic functionals (which have by definition compact support). Moreover, if the initial data vanish in a ball, then the solution vanishes in a cone, whose basis is the same ball

and whose slope depends on the coefficients a_{ij} 's. Therefore, it will be sufficient to show that if u_0 and u_1 belong to a suitable Gevrey space $\gamma^{(\sigma)}$ and have compact support, then the corresponding solution u is not only in $W^{2,1}([0,T], \mathcal{A}'(\mathbb{R}^n))$, but it belongs to the same Gevrey space in the x variable for all $t \in [0,T]$. The result for initial data which do not have compact support follows by an exhaustion argument. Our main tools in the proof will be the Paley-Wiener theorem and energy estimates.

Theorem 3. Let p > 1 and $0 < \alpha < 1$, and assume that there exists a constant C > 0 such that the function $a = a(t,\xi)$ defined by (1.3) satisfies

(3.1)
$$|a(t+\tau,\xi) - a(t,\xi)| \le \frac{C}{t^p} \tau^{\alpha}, \quad 0 \le \tau, \quad t, t+\tau \in]0,T]$$

for all $\xi \in \mathbb{R}^n \setminus \{0\}$. Then the Cauchy problem (1.1), (1.2) is $\gamma^{(\sigma)}$ -well-posed for $1 \leq \sigma < \frac{p}{p-\alpha}$.

Remark 2. For a fixed p > 1, passing to the limit as $\alpha \to 1$ we regain the result of [2]. In the same way, for a fixed $\alpha < 1$, passing to the limit as $p \to 1$ we extend to p = 1 the result of [1] which was valid only for p < 1. The case $\alpha = 1$, p = 1 was considered in [2] and will be reconsidered here in a more general context: in this case one has well posedness in C^{∞} .

Remark 3. The result in Theorem 3 can be considered sharp in the following sense. Let $p_0 > 1$ and $0 < \alpha_0 < 1$. It is possible to construct a positive function $a \in C^{\infty}([0,T]) \cap C([0,T])$ such that

$$|a(t+\tau) - a(t)| \le \frac{C}{t^{p_0}} \tau^{\alpha_0}, \quad 0 \le \tau, \quad t, t+\tau \in]0, T],$$

and it is possible to construct two functions u_0 , $u_1 \in \gamma^{(s)}(\mathbb{R})$, for all $s > \frac{p_0}{p_0 - \alpha_0}$ such that the Cauchy problem

$$\begin{cases} u_{tt} - a(t)u_{xx} = 0\\ u(0, x) = u_0(x), \ u_t(0, x) = u_1(x) \end{cases}$$

has no solution in $C^1([0,r[;D'^{(s)}))$, for all $s > \frac{p_0}{p_0-\alpha_0}$ and for all r > 0 (here $D'^{(s)}$ denotes the set of Gevrey-ultradistributions of index s). The construction of such a counterexample is exactly the same as that contained in Theorem 5 in [2].

Remark 4. A result analogous to that of Theorem 3 can be proved if the singularity of the a_{ij} 's is located at t = T, with only minor obvious changes in the proof. As a consequence, the result is still valid if the coefficients have a finite number of singularities, where the loss of regularity is controlled as in (3.1).

Proof of Theorem 3. We take the Fourier transform of u with respect to x, and we denote it by \hat{u} . Equation (1.1) then transforms to

(3.2)
$$\hat{u}_{tt}(t,\xi) + a(t,\xi)|\xi|^2 \hat{u}(t,\xi) = 0.$$

Let ϵ be a positive parameter and for each ϵ let $a_{\varepsilon} \colon [0,T] \times (\mathbb{R}^n \setminus \{0\}) \to \mathbb{R}$ be defined according to (2.3)-(2.4).

We define the *approximate energy* of \hat{u} by (3.3)

$$E_{\varepsilon}(t,\xi) := a_{\varepsilon}(t,\xi)|\xi|^{2}|\hat{u}(t,\xi)|^{2} + |\hat{u}_{t}(t,\xi)|^{2}, \quad (t,\xi) \in [0,T] \times (\mathbb{R}^{n} \setminus \{0\}).$$

Differentiating E_{ε} with respect to t and using (3.2) we get

$$E_{\varepsilon}'(t,\xi) = a_{\varepsilon}'(t,\xi)|\xi|^{2}|\hat{u}(t,\xi)|^{2} + 2a_{\varepsilon}(t,\xi)|\xi|^{2}\operatorname{Re}(\hat{u}_{t}(t,\xi)\bar{\hat{u}}(t,\xi)) + 2\operatorname{Re}(\hat{u}_{tt}(t,\xi)\bar{\hat{u}}_{t}(t,\xi)) \leq \left(\frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} + \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}}|\xi|\right)E_{\varepsilon}(t,\xi).$$

By Gronwall's lemma we obtain (3.4)

$$E_{\varepsilon}(t,\xi) \le E_{\varepsilon}(0,\xi) \exp\left(\int_0^T \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + |\xi| \int_0^T \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} dt\right)$$

for all $t \in [0, T]$ and for all $\xi \in \mathbb{R}^n$, $|\xi| \ge 1$. By Proposition 2 with $\mu(\tau) = \tau^{\alpha}$ and $\nu(t) = t^p$ we have

$$\begin{split} \int_{0}^{T} \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + |\xi| \int_{0}^{T} \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} dt \\ \leq \int_{0}^{T} \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + |\xi| \int_{0}^{T} \left(\frac{|a(t,\xi) - \tilde{a}_{\epsilon}(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} + \frac{|a_{\varepsilon}(t,\xi) - \tilde{a}_{\epsilon}(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} \right) dt \\ \leq \int_{0}^{\epsilon^{\alpha/p}} \frac{C''}{\lambda_{0}\epsilon} dt + \int_{\epsilon^{\alpha/p}}^{T} \frac{C''}{\lambda_{0}\epsilon} t^{-p} \epsilon^{\alpha} dt + \frac{2\Lambda_{0}}{\lambda_{0}^{1/2}} |\xi| \epsilon \\ &+ |\xi| \left(\int_{0}^{\epsilon^{\alpha/p}} \frac{C'}{\lambda_{0}^{1/2}} dt + \int_{\epsilon^{\alpha/p}}^{T} \frac{C'}{\lambda_{0}^{1/2}} t^{-p} \epsilon^{\alpha} dt \right) \\ \leq M |\xi| \epsilon + M \left(|\xi| + \frac{1}{\epsilon} \right) \left(\epsilon^{\alpha/p} + (\epsilon^{\alpha/p})^{1-p} \epsilon^{\alpha} \right) \\ &= M |\xi| \epsilon + 2M \left(|\xi| + \frac{1}{\epsilon} \right) \epsilon^{\alpha/p}, \end{split}$$

where M depends on C', C'', λ_0 , Λ_0 , α and p. Choosing $\epsilon = |\xi|^{-1}$ we get (3.5)

$$\left[\int_0^T \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + |\xi| \int_0^T \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} dt\right]_{\epsilon = |\xi|^{-1}} \le M + 4M|\xi|^{\frac{p-\alpha}{p}}.$$

Putting together (3.4) and (3.5) we get

(3.6)
$$E_{1/|\xi|}(t,\xi) \le e^M e^{4M|\xi|^{\frac{p-\alpha}{p}}} E_{1/|\xi|}(0,\xi)$$

and, finally, (3.7)

$$|\hat{u}_t(t,\xi)|^2 + |\xi|^2 |\hat{u}(t,\xi)|^2 \le \frac{e^M \Lambda_o}{\lambda_0} e^{4M|\xi|^{\frac{p-\alpha}{p}}} \left(|\hat{u}_t(0,\xi)|^2 + |\xi|^2 |\hat{u}(0,\xi)|^2 \right)$$

Now if $u_0, u_1 \in \gamma^{(\sigma)} \cap C_0^{\infty}$, the Paley-Wiener theorem ensures that there exist $K, \delta > 0$ such that

(3.8)
$$|\hat{u}(0,\xi)|^2 + |\hat{u}_t(0,\xi)|^2 \le K \exp(-\delta|\xi|^{1/\sigma})$$

for all $\xi \in \mathbb{R}^n$, $|\xi| \ge 1$. It follows from (3.7) that if $\sigma < p/(p-\alpha)$, then there exist $K', \delta' > 0$ such that

(3.9)
$$|\hat{u}(t,\xi)|^2 + |\hat{u}_t(t,\xi)|^2 \le K' \exp(-\delta' |\xi|^{1/\sigma})$$

for all $t \in [0,T]$ and for all $\xi \in \mathbb{R}^n$, $|\xi| \ge 1$ and, therefore, $u \in W^{2,1}([0,T], \gamma^{(\sigma)})$. The proof is complete.

4. Well posedness in C^{∞}

Let $\psi: [1, +\infty[\rightarrow]0, +\infty[$ be a strictly increasing continuous function, such that ψ' is non-increasing and $e^r \psi'(r)$ is non-decreasing. Moreover, we assume that

(1)
$$\lim_{r\to\infty} \psi(r) = \chi, \ 0 < \chi \le +\infty;$$

(2) $\lim_{r\to\infty} \psi'(r) = \eta, \ 0 \le \eta < +\infty;$

We set

(4.1)
$$\nu(t) := \begin{cases} \frac{t}{\psi'(|\log t|)} & \text{for } 0 < t \le e^{-1}, \\ \\ \frac{e^{-1}}{\psi'(1)} & \text{for } e^{-1} \le t. \end{cases}$$

A direct computation shows that ν is a non-decreasing continuous function and that $\nu(t/2) \ge (1/2)\nu(t)$ for $t \in [0, T]$. We define

(4.2)
$$\mu(\tau) := \frac{\tau |\log \tau|}{\psi(|\log \tau|)}$$

and we assume that μ is strictly increasing and concave in $[0, \tau_0]$ for a suitable $\tau_0 > 0$, so it is a modulus of continuity.

Theorem 4. Let $\nu = \nu(t)$ and $\mu = \mu(\tau)$ be as above, and assume that there exists a constant C > 0 such that the function $a = a(t, \xi)$ defined by (1.3) satisfies

(4.3)
$$|a(t+\tau,\xi) - a(t,\xi)| \le \frac{C}{\nu(t)}\mu(\tau), \quad 0 \le \tau \le \tau_0, \quad t, t+\tau \in [0,T],$$

for all $\xi \in \mathbb{R}^n \setminus \{0\}$. Then the Cauchy problem (1.1), (1.2) is well-posed in C^{∞} .

Remark 5. Examples of functions satisfying all the above properties are $\psi(r) = 1 - e^{-\alpha r}$ with $0 < \alpha \leq 1$, $\psi(r) = 1 + \log r$ and $\psi(r) = r^{\beta}$ with $0 < \beta \leq 1$. In particular, we have:

- if $\psi(r) = r$ we have $\eta = 1$ and $\chi = +\infty$ and we get $\mu(\tau) = \tau$ and $\nu(t) = t$, that is the situation considered in [2];
- if $\psi(r) = 1 e^{-\alpha r}$ we have $\eta = 0$ and $\chi = 1$ and we get $\mu(\tau) = \tau |\log \tau|/(1-\tau^{\alpha})$, which is equivalent to $\mu(\tau) = \tau |\log \tau|$, and $\nu(t) = \alpha t^{1-\alpha}$, that is a situation covered by the result of [1], since $\nu(t)^{-1}$ is integrable;
- if $\psi(r) = 1 + \log r$ or $\psi(r) = r^{\beta}$ with $0 < \beta < 1$, we have $\eta = 0$ and $\chi = +\infty$, and we get $\mu(\tau) = \tau |\log \tau|/(1 + \log |\log \tau|)$ or $\mu(\tau) = \tau |\log \tau|^{1-\beta}$. In both cases μ -continuity is weaker than Lipschitz continuity. Moreover we have $\nu(t) = t |\log t|$ or $\nu(t) = t |\log t|^{1-\beta}$, so in both cases $\nu(t)^{-1}$ is not integrable.

The case in which $\mu(\tau) = \tau |\log \tau|$ and $\nu(t)^{-1}$ is not integrable is not covered by Theorem 4, and we were not able to find a counterexample to C^{∞} well posedness either, so the question remains open. On the other hand, when $\mu(\tau) = \tau |\log \tau|$ and $\nu(t) = t$ by Theorem 3 we get authomatically $\gamma^{(\infty)}$ well posedness.

Remark 6. A result analogous to that of Theorem 4 can be proved if the singularity of the a_{ij} 's is located at t = T, with only minor obvious changes in the proof. As a consequence, the result is still valid if the coefficients have a finite number of singularities, where the loss of regularity is controlled as in (4.3).

Proof of Theorem 4. Like in the proof of Theorem 3, we take the Fourier transform \hat{u} of u. Equation (1.1) then transforms to

(4.4)
$$\hat{u}_{tt}(t,\xi) + a(t,\xi)|\xi|^2 \hat{u}(t,\xi) = 0.$$

For $0 < \epsilon \leq \tau_1 := \min\{\tau_0, T, e^{-1}\}$ we define $a_{\varepsilon} : [0, T] \times (\mathbb{R}^n \setminus \{0\}) \to \mathbb{R}$ according to (2.3)-(2.4). Again, we define an *approximate energy* of \hat{u} by

(4.5) $E_{\varepsilon}(t,\xi) := a_{\varepsilon}(t,\xi)|\xi|^{2}|\hat{u}(t,\xi)|^{2} + |\hat{u}_{t}(t,\xi)|^{2}, \quad (t,\xi) \in [0,T] \times (\mathbb{R}^{n} \setminus \{0\}).$ Differentiating E_{ε} with respect to t and using (4.4) we get

$$\begin{split} E_{\varepsilon}'(t,\xi) &= a_{\varepsilon}'(t,\xi) |\xi|^2 |\hat{u}(t,\xi)|^2 + 2a_{\varepsilon}(t,\xi) |\xi|^2 \operatorname{Re}(\hat{u}_t(t,\xi)\bar{\hat{u}}(t,\xi)) \\ &+ 2\operatorname{Re}(\hat{u}_{tt}(t,\xi)\bar{\hat{u}}_t(t,\xi)) \\ &\leq \left(\frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} + \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} |\xi|\right) E_{\varepsilon}(t,\xi). \end{split}$$

By Gronwall's lemma we obtain (4.6)

$$E_{\varepsilon}(t,\xi) \le E_{\varepsilon}(0,\xi) \exp\left(\int_0^T \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + |\xi| \int_0^T \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} dt\right)$$

for all $t \in [0, T]$ and for all $\xi \in \mathbb{R}^n$, $|\xi| \ge 1$. By Proposition 2 with $\mu(\tau)$ and $\nu(t)$ given by (4.2) and (4.1), we have

$$\int_{0}^{T} \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt = \int_{0}^{\epsilon} \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + \int_{\epsilon}^{e^{-1}} \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + \int_{e^{-1}}^{T} \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt$$
$$\leq \frac{C''}{\lambda_{0}\epsilon} \left(\epsilon + \int_{\epsilon}^{e^{-1}} \frac{\psi'(|\log t|)}{t} \frac{\epsilon|\log \epsilon|}{\psi(|\log \epsilon|)} dt + \int_{e^{-1}}^{T} \frac{\psi'(1)}{e^{-1}} \frac{\epsilon|\log \epsilon|}{\psi(|\log \epsilon|)} dt\right).$$

Since

$$\frac{\psi'(|\log t|)}{t} = -\frac{d}{dt}\psi(|\log t|)$$

and $\psi(|\log \epsilon|) \ge \psi(|\log \tau_1|)$, we obtain

(4.7)
$$\int_0^T \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt \le M'' \left(1 + |\log \epsilon|\right).$$

On the other hand

$$\begin{split} \int_0^T \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} \, dt &= \int_0^T \left(\frac{|a(t,\xi) - \tilde{a}_{\epsilon}(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} + \frac{|a_{\varepsilon}(t,\xi) - \tilde{a}_{\epsilon}(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} \right) \, dt \\ &\leq \frac{2\Lambda_0}{\lambda_0^{1/2}} \epsilon + \frac{C'}{\lambda_0^{1/2}} \left(\epsilon + \int_{\epsilon}^{e^{-1}} \frac{\psi'(|\log t|)}{t} \frac{\epsilon |\log \epsilon|}{\psi(|\log \epsilon|)} \, dt + \int_{e^{-1}}^T \frac{\psi'(1)}{e^{-1}} \frac{\epsilon |\log \epsilon|}{\psi(|\log \epsilon|)} \, dt \right). \end{split}$$

Arguing as above we, get

(4.8)
$$\int_0^T \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} dt \le M' \epsilon \left(1 + |\log \epsilon|\right).$$

10

Choosing
$$\epsilon = |\xi|^{-1}$$
 we get
(4.9)
$$\left[\int_0^T \frac{|a_{\varepsilon}'(t,\xi)|}{a_{\varepsilon}(t,\xi)} dt + |\xi| \int_0^T \frac{|a_{\varepsilon}(t,\xi) - a(t,\xi)|}{a_{\varepsilon}(t,\xi)^{1/2}} dt\right]_{\epsilon = |\xi|^{-1}} \le M((1 + \log|\xi|))$$

for $|\xi| \ge \tau_1^{-1}$.

Putting together (4.6) and (4.9) we get

(4.10)
$$E_{1/|\xi|}(t,\xi) \le e^M |\xi|^M E_{1/|\xi|}(0,\xi)$$

and, finally,

(4.11)

$$|\hat{u}_t(t,\xi)|^2 + |\xi|^2 |\hat{u}(t,\xi)|^2 \le \frac{e^M \Lambda_o}{\lambda_0} |\xi|^M \left(|\hat{u}_t(0,\xi)|^2 + |\xi|^2 |\hat{u}(0,\xi)|^2 \right).$$

Now if $u_0, u_1 \in C_0^{\infty}$, the Paley-Wiener theorem ensures that for all $\zeta > 0$ there exists $K_{\zeta} > 0$ such that

(4.12)
$$|\hat{u}(0,\xi)|^2 + |\hat{u}_t(0,\xi)|^2 \le K_{\zeta}|\xi|^{-\zeta}$$

for all $\xi \in \mathbb{R}^n$, $|\xi| \ge \tau_1^{-1}$. It follows from (4.11) that for all $\theta > 0$ there exist $K'_{\theta} > 0$ such that

(4.13)
$$|\hat{u}(t,\xi)|^2 + |\hat{u}_t(t,\xi)|^2 \le K'_{\theta}|\xi|^{-\theta}$$

for all $t \in [0,T]$ and for all $\xi \in \mathbb{R}^n$, $|\xi| \ge \tau_1^{-1}$, and therefore, $u \in W^{2,1}([0,T], C_0^{\infty})$. The proof is complete.

References

- F. Colombini, E. De Giorgi and S. Spagnolo, Sur les équations hyperboliques avec des coefficients qui ne dépendent que du temp, Ann. Sc. Norm. Sup. Pisa (4) 6 (1979), no. 3, 511-559.
- [2] F. Colombini, D. Del Santo and T. Kinoshita, Well-posedness of the Cauchy problem for a hyperbolic equation with non-Lipschitz coefficients, Ann. Sc. Norm. Sup. Pisa (5) 1 (2002), no. 2, 327-358.
- [3] F. Colombini, D. Del Santo and M. Reissig, On the optimal regularity of coefficients in hyperbolic Cauchy problems, Bull. Sci. Math. 127 (2003), no. 4, 328-347.
- [4] F. Colombini and N. Lerner, Hyperbolic operators with non-Lipschitz coefficients, Duke Math. J., 77 (1995), no. 3, 657-698.
- [5] T. Kinoshita and M. Reissig, About the loss of derivatives for strictly hyperbolic equations with non-Lipschitz coefficients, Adv. Differential Equations 10 (2005), no. 2, 191-222.

DANIELE DEL SANTO, UNIVERSITÀ DI TRIESTE, DIPARTIMENTO DI MATEM-ATICA E GEOSCIENZE, VIA VALERIO 12/1, 34127 TRIESTE, ITALY

Email address: delsanto@units.it

MARTINO PRIZZI, UNIVERSITÀ DI TRIESTE, DIPARTIMENTO DI MATEMATICA E GEOSCIENZE, VIA VALERIO 12/1, 34127 TRIESTE, ITALY *Email address*: mprizzi@units.it