Global Solutions of Three-dimensional Inviscid MHD Fluids with Velocity Damping in Horizontally Periodic Domains

Fei Jiang^a, Song Jiang^b, Youyi Zhao^{a,*}

^aCollege of Mathematics and Computer Science, Fuzhou University, Fuzhou, 350108, China. ^bInstitute of Applied Physics and Computational Mathematics, Beijing, 100088, China.

Abstract

The two-dimensional (2D) existence result of global(-in-time) solutions for the motion equations of incompressible, inviscid, non-resistive magnetohydrodynamic (MHD) fluids with velocity damping had been established in [Wu–Wu–Xu, SIAM J. Math. Anal. 47 (2013), 2630–2656]. This paper further studies the existence of global solutions for the three-dimensional (a dimension of real world) initial-boundary value problem in a horizontally periodic domain with finite height. Motivated by the multi-layers energy method introduced in [Guo–Tice, Arch. Ration. Mech. Anal. 207 (2013), 459–531], we develop a new type of two-layer energy structure to overcome the difficulty arising from three-dimensional nonlinear terms in the MHD equations, and thus prove the initial-boundary value problem admits a unique global solution. Moreover the solution has the exponential decay-in-time around some rest state. Our two-layer energy structure enjoys two features: (1) the lower-order energy (functional) can not be controlled by the higher-order energy. (2) under the *a priori* smallness assumption of lower-order energy, we first close the higher-order energy estimates, and then further close the lower-energy estimates in turn.

Keywords: non-resistive MHD fluids; incompressible; inviscid; damping; global well-posedness; exponential decay-in-time.

1. Introduction

In this paper, we investigate the global(-in-time) well-posdedness of the following *threedimensional* initial-boundary value problem for the incompressible, inviscid, non-resistive magnetohydrodynamic (MHD) fluid with velocity damping in a horizontally periodic domain Ω with finite height:

$$\begin{cases} \rho v_t + \rho v \cdot \nabla v + \nabla (P + \lambda |M|^2/2) + a\rho v = \lambda M \cdot \nabla M, \\ M_t + v \cdot \nabla M = M \cdot \nabla v, \\ \operatorname{div} v = \operatorname{div} M = 0, \\ (v, M)|_{t=0} = (v^0, M^0), \\ v_3|_{\partial\Omega} = 0. \end{cases}$$
(1.1)

Next we shall explain the mathematical notations appearing in the problem above.

^{*}Corresponding author.

Email addresses: jiangfei0591@163.com (Fei Jiang), jiang@iapcm.ac.cn (Song Jiang), zhaoyouyi957@163.com (Youyi Zhao)

The unknowns v := v(x, t), M := M(x, t) and P := P(x, t) denotes the velocity and magnetic field and kinetic pressure of incompressible MHD fluids, resp., where $(x, t) \in \mathbb{R}^3 \times \mathbb{R}^+_0$ and $\mathbb{R}^+_0 := [0, \infty)$. The positive constants ρ , a and λ stands for the density of fluids, damping coefficient and permeability of the vacuum, resp. The horizontally periodic domain Ω with finite height is defined as follows:

$$\Omega := \{ (x_{\rm h}, x_3)^{\rm T} \mid x_{\rm h} := (x_1, x_2)^{\rm T} \in \mathbb{T}, \ x_3 \in (0, h) \} \subset \mathbb{R}^3,$$
(1.2)

where $\mathbb{T} = (l_1 \mathbb{R}/\mathbb{Z}) \times (l_2 \mathbb{R}/\mathbb{Z})$, \mathbb{R}/\mathbb{Z} represents the usual 1-torus, and the superscript T denotes the transposition. Since Ω is horizontally periodic, the set $\mathbb{T} \times \{0, h\}$, denoted by $\partial \Omega$, customarily represents the boundary of Ω . Without loss of generality, we assume that $l_1 = l_2 = h = 1$ for the sake of simplicity.

For the the viscous case, i.e. the equations $(1.1)_1-(1.1)_3$ with the viscous term $-\nu\Delta v$ in place of the velocity damping term $a\rho v$, the global well-posedness, in which the initial data is a small perturbation around a non-zero trivial stationary state (i.e., $(v, M) = (0, \overline{M})$, where \overline{M} is a non-zero constant vector and \overline{M} often called the impressive magnetic field), has been widely investigated, see [23, 35, 36] and [1, 34] for the 2D and 3D Cauchy problems resp., and see [26] and [29] for the 2D and 3D initial-boundary value problems resp.. The existence of global solutions to the 2D Cauchy problem with large initial perturbations was obtained by Zhang under a strong impressive magnetic field [36]. As for the well-poseness of the 3D Cauchy and initial-boundary value problems with large initial perturbations, to our best knowledge, all available results are about the local(-in-time) existence, see [5, 9, 10] for examples. We mention here that the corresponding compressible case has been also widely studied, see [21, 22, 32] and the references cited therein.

For the inviscid case, Bardos–Sulem–Sulem used hyperbolicity of $(1.1)_1-(1.1)_3$ with $\mu = 0$ to establish an interesting global existence result of classical solutions of the Cauchy problem with small initial data in the Hölder space $H^s(\mathbb{R}^3)$ [3]; also see [4, 13] for the case of Sobolev spaces). We remark that such a result is not known in the 3D incompressible Euler equations (i.e. the case of the absence of magnetic field).

For the inviscid case with a velocity damping term, Wu–Wu–Xu first given the existence of a unique global solution with algebraic decay-in-time for the 2D Cauchy problem of $(1.1)_1-(1.1)_4$ with small initial perturbation [33]. Recently, Du–Yang–Zhou also obtained the existence of a unique global solution with exponential decay-in-time for the initial-boundary value problem in a 2D slab domain with small initial perturbation around some non-trivial equilibrium [8]. To our best knowledge, there are not any available result for the 3D case, thought the 3D wellposedness problem of inviscid fluids with velocity damping had been widely investigated, see [14, 25, 27, 28, 30] for examples. The goal of this paper is devoted to further providing the first 3D result by developing a new type of two-layer energy structure to overcome the difficulty arising from three-dimensional nonlinear terms in the MHD equations. It should be noted that the mathematical methods adopted in [8, 33] for the well-posedness problem in Eulerian coordinates cannot be applied to our 3D problem, therefore next we shall reformulate the initial-boundary value problem (1.1) in Lagrangian coordinates as in [16, 19, 31].

1.1. Reformulation in Lagrangian coordinates.

Let the flow map ζ to be the solution to the initial-value problem

$$\begin{cases} \zeta_t(y,t) = v(\zeta(y,t),t) & \text{in } \Omega, \\ \zeta(y,0) = \zeta^0(y) & \text{in } \Omega, \end{cases}$$

where the invertible mapping $\zeta^0 := \zeta^0(y) : \Omega \to \Omega$ satisfies

$$J^0 := \det(\nabla \zeta^0) = 1 \text{ in } \Omega, \tag{1.3}$$

$$\zeta_3^0 = y_3 \text{ on } \partial\Omega. \tag{1.4}$$

Here and in what follows det denotes a determinant of matrix, and f^0 with the superscript 0 represents the initial data of f. Sometimes we also use f_0 with the subscript 0 to represent the initial data of f.

We denote the Eulerian coordinates by (x, t) with $x = \zeta(y, t)$ and the Lagrangian coordinates by $(y, t) \in \Omega \times \mathbb{R}_0^+$. We further assume that, for each fixed t > 0,

$$\zeta|_{y_3=i} : \mathbb{R} \to \mathbb{R} \text{ is a } C^1(\mathbb{R}) \text{-diffeomorphism mapping for } i = 0, 1,$$
 (1.5)

$$\zeta: \overline{\Omega} \to \overline{\Omega} \text{ is a } C^1(\overline{\Omega}) \text{-diffeomorphism mapping}, \tag{1.6}$$

where $\overline{\Omega} := \mathbb{R}^2 \times [0, 1].$

Since v satisfies the divergence-free condition, and slip boundary condition $(1.1)_5$, we can deduce from (1.3)–(1.4) that

$$J := \det(\nabla \zeta) = 1 \text{ in } \Omega,$$

$$\zeta_3 = y_3 \text{ on } \partial \Omega.$$

We introduce the matrix $\mathcal{A} = (\mathcal{A}_{ij})_{3\times 3}$, which is defined via

$$\mathcal{A}^{\mathrm{T}} = (\nabla \zeta)^{-1} := (\partial_j \zeta_i)_{3 \times 3}^{-1}.$$

Then we further define the differential operators $\nabla_{\mathcal{A}}$, div_{\mathcal{A}} and curl_{\mathcal{A}} as follows: for a given scalar function f and a given vector function $X := (X_1, X_2, X_3)^{\mathrm{T}}$,

$$\nabla_{\mathcal{A}}f := (\mathcal{A}_{1k}\partial_{k}f, \mathcal{A}_{2k}\partial_{k}f, \mathcal{A}_{3k}\partial_{k}f)^{\mathrm{T}}, \ \nabla_{\mathcal{A}}X := (\nabla_{\mathcal{A}}X_{1}, \nabla_{\mathcal{A}}X_{2}, \nabla_{\mathcal{A}}X_{3})^{\mathrm{T}},$$
$$\operatorname{curl}_{\mathcal{A}}X := (\mathcal{A}_{2k}\partial_{k}X_{3} - \mathcal{A}_{3k}\partial_{k}X_{2}, \mathcal{A}_{3k}\partial_{k}X_{1} - \mathcal{A}_{1k}\partial_{k}X_{3}, \mathcal{A}_{1k}\partial_{k}X_{2} - \mathcal{A}_{2k}\partial_{k}X_{1})^{\mathrm{T}},$$
$$\operatorname{div}_{\mathcal{A}}(X_{1}, X_{2}, X_{3})^{\mathrm{T}} := \mathcal{A}_{lk}\partial_{k}X_{l},$$

where we have used the Einstein convention of summation over repeated indices, and $\partial_k = \partial_{y_k}$. In particular, $\operatorname{curl} X := \operatorname{curl}_I X$, where I is a 3×3 identity matrix. In addition, we will denote $(\operatorname{curl}_{\mathcal{A}} X^1, \ldots, \operatorname{curl}_{\mathcal{A}} X^n)$ by $\operatorname{curl}_{\mathcal{A}} (X^1, \ldots, X^n)$ for simplicity, where $X^i = (X_1^i, X_2^i, X_3^i)^{\mathrm{T}}$ is a vector function for $1 \leq i \leq n$.

Defining the Lagrangian unknowns

$$(u, Q, B)(y, t) = (v, P + \lambda |M|^2/2, M)(\zeta(y, t), t) \text{ for } (y, t) \in \Omega \times \mathbb{R}_0^+,$$

then in Lagrangian coordinates, the initial-boundary value problem (1.1) reads as follows:

$$\begin{cases} \zeta_t = u, \ \operatorname{div}_{\mathcal{A}} u = 0 & \operatorname{in} \Omega, \\ \rho u_t + \nabla_{\mathcal{A}} Q + a \rho u = \lambda B \cdot \nabla_{\mathcal{A}} B & \operatorname{in} \Omega, \\ B_t = B \cdot \nabla_{\mathcal{A}} u & \operatorname{in} \Omega, \\ \operatorname{div}_{\mathcal{A}} B = 0 & \operatorname{in} \Omega, \\ (u, \zeta, B)|_{t=0} = (u^0, \zeta^0, B^0) & \operatorname{in} \Omega, \\ (\zeta_3 - y_3) = u_3 = 0 & \operatorname{on} \partial\Omega. \end{cases}$$
(1.7)

We can derive from $(1.7)_3$ the differential version of magnetic flux conservation [18]:

$$\mathcal{A}^{\mathrm{T}}B = \mathcal{A}_{0}^{\mathrm{T}}B^{0}$$

which yields

$$B = \nabla \zeta \mathcal{A}_0^{\mathrm{T}} B^0. \tag{1.8}$$

Let $\overline{M} = (\overline{M}_1, 0, 0)^{\mathrm{T}}$ and \overline{M}_1 is a nonzero constant. If we assume

$$\mathcal{A}_0^{\mathrm{T}} B^0 = \bar{M} \text{ (i.e. } B^0 = \nabla \zeta^0 \bar{M} = \partial_{\bar{M}} \zeta^0 \text{)}, \tag{1.9}$$

then (1.8) reduces to

$$B = \partial_{\bar{M}} \zeta. \tag{1.10}$$

It should be noted that B given by (1.10) automatically satisfies $(1.7)_3$ and $(1.7)_4$. Moreover, we see from (1.10) that the magnetic tension in Lagrangian coordinates has the relation

$$\lambda B \cdot \nabla_{\mathcal{A}} B = \lambda \bar{M}_1^2 \partial_1^2 \zeta.$$

Let

$$e_1 = (1, 0, 0)^{\mathrm{T}}, \ m = \lambda \bar{M}_1^2 / \rho, \ q = Q / \rho, \ \eta = \zeta - y, \ \mathcal{A} = (\nabla \eta + I)^{\mathrm{T}} \text{ and } \eta^0 = \zeta^0 - y.$$

Consequently, under the assumption (1.9), the problem (1.7) is equivalent to the following initialboundary value problem:

$$\begin{cases} \eta_t = u & \text{in } \Omega, \\ u_t + \nabla_{\mathcal{A}} q + au = m\partial_1^2 \eta & \text{in } \Omega, \\ \operatorname{div}_{\mathcal{A}} u = 0 & \text{in } \Omega, \\ (u, \eta)|_{t=0} = (u^0, \eta^0) & \text{in } \Omega, \\ (u_3, \eta_3) = 0 & \text{on } \partial\Omega, \end{cases}$$
(1.11)

with the expression of magnetic field

$$B = \overline{M}_1(\partial_1 \eta + e_1). \tag{1.12}$$

1.2. Main results.

Before stating our main result, we introduce some notations which will be frequently used throughout this paper.

(1) Simplified basic notations: $I_a := (0, a)$, in particular, $I_{\infty} = \mathbb{R}^+$. \overline{S} denotes the closure of the set $S \subset \mathbb{R}^n$ with $n \ge 1$, in particular $\overline{I_T} := [0, T]$. $\Omega_a := \Omega \times I_a$. $\vec{\mathbf{n}} := (\mathbf{n}_1, \mathbf{n}_2, \mathbf{n}_3)^{\mathrm{T}}$ denotes the outward unit normal vector on $\partial \Omega$. $\int := \int_{\Omega} = \int_{(0,1)^3} (u)_{\Omega} = \int u \mathrm{d}y$.

 $a \leq b$ means that $a \leq cb$. If not stated explicitly, the positive constant c may depend on a, m and Ω , and may vary from one place to other place. Sometimes we use c_i for $i \geq 1$ to replace c in order to emphasize that c_i is fixed value.

 α always denotes the multiindex with respect to the variable y, $|\alpha| = \alpha_1 + \alpha_2 + \alpha_3$ is called the order of multiindex and $\partial^{\alpha} := \partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}$. " $\nabla^i f \in \mathbb{X}$ " represents that $\partial^{\alpha} f \in \mathbb{X}$ for any multiindex α satisfying $|\alpha| = i$, where \mathbb{X} denotes some set of functions.

(2) Simplified Banach spaces:

$$\begin{split} L^{p} &:= L^{p}(\Omega) = W^{0,p}(\Omega), \quad H^{i} := W^{i,2}(\Omega), \quad L^{p}_{T}H^{i} := L^{p}(I_{T}, H^{i}), \\ H^{l,i} &:= \{ w \in H^{i} \mid \partial_{1}^{k}w \in H^{i}, \; 0 \leqslant k \leqslant l \}, \; H^{l,j}_{s} := \{ w \in H^{l,j} \mid w_{3}|_{\partial\Omega} = 0 \}, \\ H^{j}_{1} &:= \{ w \in H^{j} \mid \det(\nabla w + I) = 1 \}, \; H^{l,j}_{1,s} := H^{l,j}_{s} \cap H^{j}_{1}, \\ C^{0}_{B,\text{weak}}(\overline{I_{T}}, L^{2}) &:= L^{\infty}_{T}L^{2} \cap C^{0}(\overline{I_{T}}, L^{2}_{\text{weak}}), \; \underline{H}^{i} := \{ w \in H^{i} \mid (w)_{\Omega} = 0 \}, \end{split}$$

where $1 \leq p \leq \infty$, and $i, l \geq 0$ and $j \geq 1$ are integers. In addition, $||(f^1, \cdot, f^n)||_B := \sqrt{\sum_{1 \leq i \leq n} ||f^i||_B^2}$, where f^i may be a scalar function, a vector or a matrix function for $1 \leq i \leq n$. (3) Simplified function classes: for any given integer $j \geq 1$,

$$\begin{split} H^j_* &:= \{\xi \in H^j \mid \xi(y) + y : \overline{\Omega} \to \overline{\Omega} \text{ satisfies the properties} \\ &\text{ of diffeomorphism (1.5) and (1.6) as } \zeta \}, \\ \mathfrak{C}^0(\overline{I_T}, H^{2,3}_{\mathrm{s}}) &:= \{\eta \in C^0(\overline{I_T}, H^{1,3}_{\mathrm{s}}) \mid \nabla^3 \partial_1^2 \eta \in C^0_{B,\mathrm{weak}}(\overline{I_T}, L^2) \}, \\ \underline{\mathfrak{H}}^{2,3}_{1,*,T} &:= \{\eta \in \mathfrak{C}^0(\overline{I_T}, H^{2,3}_{\mathrm{s}}) \mid \eta(t) \in H^3_1 \cap H^3_* \text{ for each } t \in \overline{I_T} \}, \\ \mathfrak{U}^{1,3}_T &:= \{u \in C^0(\overline{I_T}, H^3_{\mathrm{s}}) \mid \nabla^3 \partial_1 u \in C^0_{B,\mathrm{weak}}(\overline{I_T}, L^2), \\ u_t \in C^0(\overline{I_T}, H^2_{\mathrm{s}}), u_t \in L^\infty_T H^3 \}. \end{split}$$

(4) Simplified norms and semi-norms: for integers $i \ge 0$ and $j \ge 0$,

$$\|\cdot\|_{i} := \|\cdot\|_{H^{i}}, \quad \|\cdot\|_{j,i} := \|\partial_{1}^{j}\cdot\|_{i}, \quad \|\cdot\|_{\underline{j},i} := \sqrt{\sum_{0 \le k \le j} \|\cdot\|_{k,i}^{2}}.$$

(5) Simplified functionals:

$$\begin{aligned} \mathcal{E}^{L} &:= \|(u,\eta,\partial_{1}\eta)\|_{3}^{2} + \|(u_{t},\nabla q)\|_{2}^{2}, \quad \mathcal{D}^{L} := \|(u,\partial_{1}\eta)\|_{3}^{2} + \|(u_{t},\nabla q)\|_{2}^{2}, \\ \mathcal{E}^{H} &:= \|(u,\partial_{1}\eta)\|_{\underline{1},3}^{2} + \|(u_{t},\nabla q)\|_{2}^{2}, \quad \mathcal{D}^{H} := \|(u,\partial_{1}u,\partial_{1}^{2}\eta)\|_{3}^{2} + \|(u_{t},\nabla q)\|_{2}^{2}, \\ I_{L}^{0} &:= \|(u^{0},\eta^{0},\partial_{1}\eta^{0})\|_{3}^{2}, \quad I_{H}^{0} := \|(u^{0},\partial_{1}\eta^{0})\|_{\underline{1},3}. \end{aligned}$$

We call \mathcal{E}^L , resp. \mathcal{D}^L the lower energy, resp. dissipation functionals, and \mathcal{E}^H , resp. \mathcal{D}^H the higher energy, resp. dissipation functionals.

Now we state the existence result of unique global solutions for the initial-boundary value problem (1.11) with small initial data.

Theorem 1.1. Let $(u^0, \eta^0) \in H^{1,3}_s \times (H^{2,3}_{1,s} \cap H^3_*)$ satisfy the incompressible condition $\operatorname{div}_{\mathcal{A}^0} u^0 = 0$ in Ω ,

where $\mathcal{A}^0 := (\nabla \eta^0 + I)^{-T}$. Then there exist a sufficiently small constant $\delta > 0$ and a positive constant c, such that, for any (u^0, η^0) satisfying

$$e^{c\sqrt{I_H^0}} \|(u^0,\eta^0,\partial_1\eta^0)\|_3^2 \leqslant \delta^2,$$

the initial-boundary problem (1.11) admits a unique global classical solution $(u, \eta, q) \in \mathfrak{U}^{1,3}_{s,\infty} \times \underline{\mathfrak{H}}^{2,3,*}_{1,s,\infty} \times C(\mathbb{R}^+_0, \underline{H}^3)$. Moreover the solution enjoys the following stability estimates:

$$\mathcal{E}^{L}(t) + \int_{0}^{t} \mathcal{D}^{L}(s) \mathrm{d}s \lesssim e^{c\sqrt{I_{H}^{0}}} I_{L}^{0}, \qquad (1.13)$$

$$e^{c_1 t} \mathcal{E}^H(t) + \int_0^t e^{c_1 \tau} \mathcal{D}^H(s) \mathrm{d}s \lesssim I_H^0, \qquad (1.14)$$

$$e^{c_1 t} \|\eta(t) - \eta^{\infty}\|_3^2 \lesssim I_H^0,$$
 (1.15)

where η^{∞} only depends on y_2 and y_3 . In addition, $\nabla_{\mathcal{A}} q \in L^{\infty}_{\infty} H^3$.

Remark 1.1. In this paper, we only consider the impressive horizontal magnetic field \overline{M} . However it is not clear to the authors that whether there exists a global well-posdness result for non-horizontal magnetic field.

Remark 1.2. For the case $\Omega = \mathbb{R}^2 \times (0, 1)$, we also obtain similar result, where the exponential stability estimates (1.14) and (1.15) should be replaced by algebraic stability estimates. We will verify this assertion in a forthcoming paper.

Remark 1.3. For each fixed $t \in \mathbb{R}_0^+$, the solution $\eta(y, t)$ in Theorem 1.1 belongs to H^3_* . Let $\zeta = \eta + y$, then ζ satisfies (1.5) and (1.6) for each fixed $t \in \mathbb{R}_0^+$. We denote the inverse transformation of ζ by ζ^{-1} , and then define that

$$(v, N, Q)(x, t) := (u(y, t), M_1 \partial_1 \eta(y, t), \rho q(y, t))|_{y = \zeta^{-1}(x, t)},$$

$$M := N + \bar{M} \text{ and } P := Q - \lambda |M|^2 / 2.$$

Consequently (v, M, P) is a global solution for (1.1); moreover, making use of (1.13), (1.14) and the fifth conclusion in Lemma A.5, (v, N, Q) enjoys stability estimates, which are similar to (1.13) and (1.14).

Next we briefly sketch the proof of Theorem 1.1, and the details will be presented in Section 2.

The key proof for the existence of global small solutions is to derive an *a priori* energy inequalities (1.13) and (1.14). To this purpose, let (η, u) be a solution to (1.11) and satisfy

$$\sup_{t\in\overline{I_T}} \|(u,\eta,\partial_1\eta)(t)\|_3 \leqslant \delta \in (0,1] \text{ for some } T,$$
(1.16)

$$\det(\nabla \eta + I) = 1, \tag{1.17}$$

where δ is sufficiently small.

Under the *a priori* assumption (1.16) of lower-order energy, we first derive the both energy inequalities of horizontal-type and vorticity-type, see (2.27) and (2.41). Summing up the two energy inequalities, we can arrive at the lower-order energy inequality

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{\mathcal{E}}^L + \mathcal{D}^L \lesssim (\|u\|_3 + \|\nabla q\|_0) \|\eta\|_3^2 \tag{1.18}$$

for some energy functional $\tilde{\mathcal{E}}^L$, which is equivalent to \mathcal{E}^L .

Obviously we can not directly close the lower-order energy by using (1.18), unless the term " $||u||_3 + ||\nabla q||_0$ " decays in time. It should be noted that the term " $(||u||_3 + ||\nabla q||_0)||\eta||_3^2$ " (arising from nonlinear terms, see (2.25) and (2.39)) does not appear in (1.18) for the two dimensional cases. Motivated by the multi-layers energy method introduced in [12], in which Guo–Tice investigated the global well-posdedness of surface wave problem, we naturally look for another energy inequality, which includes the decay-in-time of $||u||_3$.

By careful analysis of energy estimates, we find that, if (u^0, η^0) additionally satisfies the regularity

$$\partial_1(u^0, \partial_1\eta^0) \in H^3,$$

the following higher-order energy inequality can be established:

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{\mathcal{E}}^H + \mathcal{D}^H \leqslant 0, \tag{1.19}$$

where $\tilde{\mathcal{E}}^H$ is equivalent to \mathcal{E}^H . Moreover, thanks to the estimate

$$\|\partial_1\eta\|_3 \lesssim \|\partial_1^2\eta\|_3,$$

we have

 \mathcal{E}^H is equivalent to \mathcal{D}^H .

Thus the higher-order energy inequality (1.19), together with the above equivalence, immediately implies the exponential stability of higher-order energy (1.14), which particularly presents the exponential decay-in-timely of $||u||_3$ and $||\nabla q||_0$. Hence the *a priori* stability estimate (1.13) further follows from (1.18).

Thanks to the *a priori* stability estimates (1.13)-(1.14) and the unique local solvability of the initial-boundary problem (1.11), we immediately get the unique global solvability of the initial-boundary problem (1.11). Finally, (1.15) can be deduced from (1.14) by an asymptotic analysis method.

The rest of this paper is organized as follows. In Sections 2, we will provide the proof of Theorem 1.1. In Section 3, we will establish the local well-posdedness result for the initialboundary problem (1.11). Finally, in Appendix A, we will list some well-known mathematical results, which will be used in Sections 2–3.

2. Proof of Theorem 1.1

This section is devoted to the proof of Theorem 1.1. The key step is to derive the a *priori* stability estimates (1.13)-(1.14) for the initial-boundary value problem (1.11). To this end, let (u, η, q) be a solution to the problem (1.11), and satisfy (1.16) and (1.17) with sufficiently small δ . Next we proceed the derivation of *a priori* estimates.

2.1. Preliminary estimates

To begin with, we shall establish some preliminary estimates.

Lemma 2.1. Let $0 \leq i \leq 2$ and $1 \leq j \leq 2$. Under the assumptions (1.16) and (1.17), we have

(1) the estimates for \mathcal{A} and $\tilde{\mathcal{A}}$:

$$\|\mathcal{A}\|_{C^0(\overline{\Omega})} + \|\mathcal{A}\|_2 \lesssim 1, \tag{2.1}$$

$$\|\tilde{\mathcal{A}}\|_i \lesssim \|\eta\|_{i+1} \tag{2.2}$$

$$\|\tilde{\mathcal{A}}\|_{j,i} \lesssim \|\eta\|_{j,i+1},\tag{2.3}$$

 $\|\mathcal{A}_t\|_i \lesssim \|u\|_{i+1},\tag{2.4}$

$$\|\tilde{\mathcal{A}}_t\|_{1,i} \lesssim \|(u,\eta)\|_{1,i+1}.$$
(2.5)

Here and in what follows $\tilde{\mathcal{A}} := \mathcal{A} - I$.

(2) the estimates of divu:

$$\|\operatorname{div} u\|_i \lesssim \|\eta\|_3 \|u\|_{i+1},$$
 (2.6)

 $\|\operatorname{div} u\|_{1,i} \lesssim \|(u,\eta)\|_3\|(u,\eta)\|_{1,i+1}.$ (2.7)

(3) the estimates of div η :

$$|\operatorname{div}\eta||_i \lesssim ||\eta||_3 ||\eta||_{i+1},$$
(2.8)

$$\|\operatorname{div}\eta\|_{1,i} \lesssim \|\eta\|_3 \|\eta\|_{1,i+1},\tag{2.9}$$

$$\|\operatorname{div}\eta\|_{2,2} \lesssim \|\eta\|_{3} \|\eta\|_{2,3} + \|\eta\|_{1,3}^{2}.$$
(2.10)

PROOF. Recalling the condition (1.17) and the definition of \mathcal{A} , we have

$$\mathcal{A} = (A_{ij}^*)_{3\times 3},\tag{2.11}$$

where A_{ij}^* is the algebraic complement minor of (i, j)-th entry of matrix $(\partial_j \zeta_i)_{3\times 3}$. Exploiting $(1.11)_1$, (1.16), the produce estimate (A.2) and Poincáre's inequality (A.3), we easily derive (2.1)-(2.5) from the expression (2.11). In addition, we easily derive (2.6) and (2.7) from the incompressible condition $\operatorname{div}_{\mathcal{A}} u = 0$ by (A.2). Similarly, we easily get (2.8)–(2.10) from the relation (1.17). Interesting readers can refer to [15] for the detailed derivation.

Lemma 2.2. Let i = 0, 1, and the multindex α satisfy $|\alpha| \leq 2, w(t) \in H^3$ for each $t \in \overline{I_T}$ and

$$W^{i,\alpha} := \partial_1^i \partial^\alpha \left(\operatorname{curl}_{\mathcal{A}_t} u - m \operatorname{curl}_{\partial_1 \mathcal{A}} \partial_1 \eta \right).$$
(2.12)

Under the assumptions (1.16) and (1.17), we have

$$\left\|\partial^{\alpha}\left(\operatorname{curl}_{\mathcal{A}} w - \operatorname{curl} w\right)\right\|_{0} \lesssim \|\eta\|_{3} \|w\|_{3},\tag{2.13}$$

$$\begin{aligned} \|\partial^{\alpha} \left(\operatorname{curl}_{\mathcal{A}} w - \operatorname{curl} w\right)\|_{1,0} &\lesssim \|(\eta, w)\|_{3} \|(\eta, w)\|_{1,3}, \end{aligned} \tag{2.14} \\ \|W^{0,\alpha}\|_{0} &\lesssim \|(u, \partial_{1}\eta)\|_{3}^{2}, \end{aligned} \tag{2.15}$$

$$W^{0,\alpha} \Big\|_{0} \lesssim \|(u,\partial_{1}\eta)\|_{3}^{2},$$
 (2.15)

$$\left\| W^{1,\alpha} \right\|_{0} \lesssim \|u\|_{3} \|(u,\eta)\|_{1,3} + \|\eta\|_{1,3} \|\eta\|_{2,3}.$$
(2.16)

PROOF. By (2.2), (2.3) and the product estimate (A.2), we easily get (2.13) and (2.14). Similarly, making use of (2.3)–(2.5) and (A.2), we easily derive (2.15)–(2.16) from the definition of $W^{i,\alpha}$. The proof is complete.

2.2. Horizontal spatial estimates

Now we derive the estimates of horizontally spatial derivatives of (u, η) .

Lemma 2.3. Let $0 \leq k \leq 1$. Under the assumptions (1.16) and (1.17), we have

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\|\partial_1^k u\|_0^2 + m \|\partial_1^{k+1} \eta\|_0^2 \right) + a \|\partial_1^k u\|_0^2 \lesssim \begin{cases} 0 & \text{for } k = 0; \\ \sqrt{\mathcal{E}^L} \mathcal{D}^H & \text{for } k = 1, \end{cases}$$
(2.17)

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\int \partial_1^k \eta \cdot \partial_1^k u \mathrm{d}y + \frac{a}{2} \|\partial_1^k \eta\|_0^2 \right) + m \|\partial_1^{k+1} \eta\|_0^2$$

$$\lesssim \|\partial_1^k u\|_0^2 + \begin{cases} \|\eta\|_2^2 \|\nabla q\|_0 & \text{for } k = 0; \\ \sqrt{\mathcal{E}^L} \mathcal{D}^H & \text{for } k = 1. \end{cases}$$
(2.18)

PROOF. Applying ∂_1^k to (1.11) yields

$$\begin{cases} \partial_1^k \eta_t = \partial_1^k u & \text{in } \Omega, \\ \partial_1^k u_t + \partial_1^k \nabla_{\mathcal{A}} q + a \partial_1^k u = m \partial_1^2 \partial_1^k \eta & \text{in } \Omega, \\ \partial_1^k \operatorname{div}_{\mathcal{A}} u = 0 & \text{in } \Omega, \\ \partial_1^k (u_3, \eta_3) = 0 & \text{on } \partial\Omega, \end{cases}$$
(2.19)

Multiplying $(2.19)_2$ by $\partial_1^k u$ and $\partial_1^k \eta$ in L^2 , resp., and then using the integration by parts and $(1.11)_1$, we have

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\left(\|\partial_1^k u\|_0^2 + m\|\partial_1\partial_1^k \eta\|_0^2\right) + a\|\partial_1^k u\|_0^2 = -\int \partial_1^k \nabla_{\mathcal{A}} q \cdot \partial_1^k u \mathrm{d}y$$
(2.20)

and

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\int \partial_1^k \eta \cdot \partial_1^k u \mathrm{d}y + \frac{a}{2} \|\partial_1^k \eta\|_0^2 \right) + m \|\partial_1 \partial_1^k \eta\|_0^2 = \|\partial_1^k u\|_0^2 - \int \partial_1^k \nabla_\mathcal{A} q \cdot \partial_1^k \eta \mathrm{d}y.$$
(2.21)

Next we estimate for the integrals involving the pressure in (2.20) and (2.21) by two cases.

(1) Case k = 0.

Noting that $\eta_3|_{\partial\Omega} = 0$, thus

$$\mathcal{A}_{13}|_{\partial\Omega} = \mathcal{A}_{23}|_{\partial\Omega} = 0. \tag{2.22}$$

Exploiting the integration by parts, $(1.11)_3$ and the boundary-value conditions of $u_3|_{\partial\Omega} = 0$ and (2.22), we have

$$-\int \nabla_{\mathcal{A}} q \cdot u \mathrm{d}y = \int q \mathrm{div}_{\mathcal{A}} u \mathrm{d}y = 0.$$

Putting the above identity into (2.20) with k = 0 yields (2.17) with k = 0.

Now we turn to estimating for the last integral with k = 0 in (2.21). Since $det(\nabla \eta + I) = 1$, we have

$$\operatorname{div}\eta = \operatorname{div}\Psi(\eta),\tag{2.23}$$

where

$$\Psi(\eta) := - \begin{pmatrix} \eta_1(\partial_2\eta_2 + \partial_3\eta_3) - \eta_1(\partial_2\eta_3\partial_3\eta_2 - \partial_2\eta_2\partial_3\eta_3) \\ \eta_2\partial_3\eta_3 - \eta_1\partial_1\eta_2 - \eta_1(\partial_1\eta_2\partial_3\eta_3 - \partial_1\eta_3\partial_3\eta_2) \\ -\eta_1\partial_1\eta_3 - \eta_2\partial_2\eta_3 - \eta_1(\partial_1\eta_3\partial_2\eta_2 - \partial_1\eta_2\partial_2\eta_3) \end{pmatrix}.$$

It is easy to estimate that

$$\|\Psi(\eta)\|_0 \lesssim \|\eta\|_2^2. \tag{2.24}$$

Thanks to (2.23), (2.24) and the boundary-value conditions

$$\eta_3|_{\partial\Omega} = 0, \ \Psi_3(\eta)|_{\partial\Omega} = 0,$$

we have

$$\int q \operatorname{div} \eta \mathrm{d}y \leqslant \|\Psi(\eta)\|_0 \|\nabla q\|_0 \lesssim \|\eta\|_2^2 \|\nabla q\|_0.$$

In addition, we can estimate that

$$\left| \int \nabla_{\tilde{\mathcal{A}}} q \cdot \eta \mathrm{d}y \right| \lesssim \|\eta\|_2^2 \|\nabla q\|_0$$

Exploiting the above two estimates and integration by parts, we get

$$-\int \nabla_{\mathcal{A}} q \cdot \eta \mathrm{d}y = \int (q \mathrm{div}\eta - \nabla_{\tilde{\mathcal{A}}} q \cdot \eta) \mathrm{d}y \lesssim \|\eta\|_2^2 \|\nabla q\|_0.$$
(2.25)

Putting the above estimate into (2.21) yields (2.18) with k = 0.

(2) Case k = 1.

Thanks to (2.2), (2.7) and (A.3), we can estimate that

$$-\int \partial_1 \nabla_{\mathcal{A}} q \cdot \partial_1 u dy = \int (\partial_1 q \partial_1 \operatorname{div} u + \nabla_{\tilde{\mathcal{A}}} q \cdot \partial_1^2 u) dy$$
$$\lesssim \|(u,\eta)\|_3 \|(u,\eta)\|_{1,1} \|\partial_1 q\|_0 + \|\partial_1^2 u\|_0 \|\eta\|_3 \|\nabla q\|_0 \lesssim \sqrt{\mathcal{E}^L} \mathcal{D}^H$$

Putting the above estimate into (2.20) yields (2.17) with k = 1.

Thanks to (2.2), (2.9) and (A.3), we have

$$-\int \partial_1 \nabla_{\mathcal{A}} q \cdot \partial_1 \eta \, \mathrm{d}y = \int (\nabla_{\tilde{\mathcal{A}}} q \cdot \partial_1^2 \eta + \partial_1 q \, \mathrm{div} \partial_1 \eta) \, \mathrm{d}y$$

$$\lesssim \|\eta\|_{2,0} \|\eta\|_3 \|\nabla q\|_0 + \|\eta\|_{1,1} \|\eta\|_3 \|q\|_1 \lesssim \sqrt{\mathcal{E}^L} \mathcal{D}^H.$$

Plugging the above estimate into (2.21) yields (2.18) with k = 1. The proof is complete.

Lemma 2.3 immediately implies the following two horizontal-type energy inequalities.

Proposition 2.1. Under the assumptions (1.16) and (1.17), there exist two energy functionals \mathcal{E}_{h}^{H} and \mathcal{E}_{h}^{L} of (η, u) such that

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathcal{E}_{h}^{L} + \|(u,\partial_{1}\eta)\|_{0}^{2} \lesssim \|\eta\|_{2}^{2} \|\nabla q\|_{0}, \qquad (2.26)$$

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{E}_{h}^{H} + \|(u,\partial_{1}u,\partial_{1}^{2}\eta)\|_{0}^{2} \lesssim \sqrt{\mathcal{E}^{L}}\mathcal{D}^{H}, \qquad (2.27)$$

where, for each t > 0,

 \mathcal{E}_{h}^{L} and \mathcal{E}_{h}^{H} are equivalent to $\|(u,\eta,\partial_{1}\eta)\|_{0}^{2}$ and $\|(u,\partial_{1}\eta)\|_{\underline{1},0}^{2}$, resp..

2.3. Vorticity estimates

Let the multindex α satisfy $|\alpha| \leq 2$. Applying $\partial^{\alpha} \partial_1^i \operatorname{curl}_{\mathcal{A}}$ to $(1.11)_2$ yields

$$\partial_t \partial^\alpha \partial_1^i \operatorname{curl}_{\mathcal{A}} u + a \partial^\alpha \partial_1^i \operatorname{curl}_{\mathcal{A}} u = m \partial_1 (\partial^\alpha \partial_1^i \operatorname{curl}_{\mathcal{A}} \partial_1 \eta) + W^{i,\alpha}, \qquad (2.28)$$

where $W^{i,\alpha}$ is defined by (2.12).

Now we derive the following vorticity estimates for u and η .

Lemma 2.4. Let $|\alpha| \leq 2$. Under the assumptions (1.16) and (1.17), we have

$$\frac{1}{2} \frac{d}{dt} \int \left(|\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} u|^{2} + m |\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} \partial_{1} \eta|^{2} \right) dy + a \|\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} u\|_{0}^{2}
\lesssim \begin{cases} \|u\|_{3}\|(u,\partial_{1}\eta)\|_{3}^{2} & \text{for } i = 0; \\ \|\eta\|_{2,3}(\|u\|_{3}\|\eta\|_{2,3} + \|(u,\eta)\|_{1,3}\|\eta\|_{1,3}) + \|u\|_{3}\|(u,\eta)\|_{1,3}^{2} & \text{for } i = 1, \end{cases}$$

$$\frac{d}{dt} \int \left(\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} \eta \cdot \partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} u + \frac{a}{2} |\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} \eta|^{2} \right) dy + m \|\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} \partial_{1} \eta\|_{0}^{2}
\lesssim \|\partial^{\alpha} \partial_{1}^{i} \operatorname{curl}_{\mathcal{A}} u\|_{0}^{2} & \text{for } i = 0; \\ \|\partial^{\alpha} \partial_{1}^{i} \|(u,\eta)\|_{3}^{2} + \|u\|_{3} \|\eta\|_{3}^{2} & \text{for } i = 0; \\ \|(u,\eta)\|_{3} \|(u,\eta)\|_{1,3}^{2} + \|\eta\|_{2,3}(\|\eta\|_{3} \|\eta\|_{2,3}^{2} + \|\eta\|_{1,3}^{2}) & \text{for } i = 1. \end{cases}$$

$$(2.30)$$

PROOF. (1) Multiplying (2.28) by $\partial^{\alpha}\partial_1^i \operatorname{curl}_{\mathcal{A}} u$ in L^2 and then using the integration by parts, we get

$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \int \left(|\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u|^{2} + m |\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \partial_{1} \eta|^{2} \right) \mathrm{d}y + a \|\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u\|_{0}^{2}$$

$$= m \int \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \partial_{1} \eta \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}_{t}} \partial_{1} \eta \mathrm{d}y - m \int \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \partial_{1} \eta \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\partial_{1}\mathcal{A}} u \mathrm{d}y$$

$$+ \int W^{i,\alpha} \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u \mathrm{d}y =: \sum_{j=1}^{3} I_{j}^{i,\alpha}.$$
(2.31)

Making use of (2.1), (2.3), (2.4), and (2.15), the three integrals $I_1^{0,\alpha} - I_3^{0,\alpha}$ can be estimated as follows:

$$I_1^{0,\alpha} \lesssim \|\mathrm{curl}_{\mathcal{A}} \partial_1 \eta\|_2 \|\mathrm{curl}_{\mathcal{A}_t} \partial_1 \eta\|_2 \lesssim \|u\|_3 \|\eta\|_{1,3}^2, \tag{2.32}$$

$$I_{2}^{0,\alpha} \lesssim \|\mathrm{curl}_{\mathcal{A}}\partial_{1}\eta\|_{2}\|\mathrm{curl}_{\partial_{1}\mathcal{A}}u\|_{2} \lesssim \|u\|_{3}\|\eta\|_{1,3}^{2}, \tag{2.33}$$

$$I_3^{0,\alpha} \lesssim \|W^{0,\alpha}\|_0 \|\operatorname{curl}_{\mathcal{A}} u\|_2 \lesssim \|u\|_3 \|(u,\partial_1\eta)\|_3^2.$$
(2.34)

Exploiting (1.16), (2.1), (2.3)–(2.5), (2.16) and (A.3), the three integrals $I_1^{1,\alpha}-I_3^{1,\alpha}$ can be estimated as follows:

$$I_{1}^{1,\alpha} \lesssim \|\eta\|_{2,3}(\|u\|_{3}\|\eta\|_{2,3} + \|(u,\eta)\|_{1,3}\|\eta\|_{1,3}),$$
(2.35)

$$I_2^{1,\alpha} \lesssim \|\eta\|_{2,3} (\|u\|_3 \|\eta\|_{2,3} + \|u\|_{1,3} \|\eta\|_{1,3}),$$
(2.36)

$$I_{3}^{1,\alpha} \lesssim \|(u,\eta)\|_{1,3}(\|u\|_{3}\|(u,\eta)\|_{1,3} + \|\eta\|_{1,3}\|\eta\|_{2,3}).$$

$$(2.37)$$

Thanks to the above six estimates and (A.3), we immediately get (2.29) from (2.31).

(2) Multiplying (2.28) by $\partial^{\alpha}\partial_1^i \operatorname{curl}_{\mathcal{A}}\eta$ in L^2 , and then using the integration by parts, we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t} \int \left(\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \eta \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u + \frac{a}{2} |\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \eta|^{2}\right) \mathrm{d}y + m \|\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \partial_{1} \eta\|_{0}^{2}$$

$$= \|\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u\|_{0}^{2} + \int \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}_{t}} \eta \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u \mathrm{d}y + a \int \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}_{t}} \eta \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \eta \mathrm{d}y$$

$$- m \int (\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \partial_{1} \eta) \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\partial_{1}\mathcal{A}} \eta \mathrm{d}y + \int W^{i,\alpha} \cdot \partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} \eta \mathrm{d}y$$

$$=: \|\partial^{\alpha} \partial_{1}^{i} \mathrm{curl}_{\mathcal{A}} u\|_{0}^{2} + \sum_{j=4}^{7} I_{j}^{i,\alpha}.$$
(2.38)

Similarly to (2.32)–(2.37), the integrals $I_4^{i,\alpha} - I_7^{i,\alpha}$ can be estimated as follows:

$$I_{4}^{0,\alpha} \lesssim \|u\|_{3}^{2} \|\eta\|_{3}, \ I_{5}^{0,\alpha} \lesssim \|u\|_{3} \|\eta\|_{3}^{2}, \tag{2.39}$$

$$I_{6}^{0,\alpha} \lesssim \|\eta\|_{3} \|\eta\|_{1,3}^{2}, \ I_{7}^{0,\alpha} \lesssim \|\eta\|_{3} \|(u,\partial_{1}\eta)\|_{3}^{2}, \qquad (2.39)$$

$$I_{4}^{1,\alpha}, \ I_{5}^{1,\alpha} \lesssim \|(u,\eta)\|_{3} \|(u,\eta)\|_{1,3}^{2}, \qquad I_{6}^{1,\alpha} \lesssim \|\eta\|_{2,3} (\|\eta\|_{3} \|\eta\|_{2,3} + \|\eta\|_{1,3}^{2}), \qquad I_{7}^{1,\alpha} \lesssim \|\eta\|_{1,3} (\|u\|_{3} \|\eta\|_{2,3} + \|\eta\|_{1,3}^{2}), \qquad I_{7}^{1,\alpha} \lesssim \|\eta\|_{1,3} (\|u\|_{3} \|(u,\eta)\|_{1,3} + \|\eta\|_{1,3} \|\eta\|_{2,3}).$$

Thanks to the above eight estimates, we immediately get (2.30) from (2.38). This completes the proof of Lemma 2.4.

Next we further establish the vorticity-type energy inequalities.

Proposition 2.2. Under the assumptions (1.16) and (1.17), there exist two energy functionals \mathcal{E}_{cul}^{H} and \mathcal{E}_{cul}^{L} of (η, u) such that

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathcal{E}_{\mathrm{cul}}^{L} + \|\mathrm{curl}(u,\partial_{1}\eta)\|_{2}^{2} \lesssim \|(u,\eta)\|_{3}\|(u,\partial_{1}\eta)\|_{3}^{2} + \|u\|_{3}\|\eta\|_{3}^{2}$$
(2.40)

and

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{E}_{\mathrm{cul}}^{H} + \|\mathrm{curl}(u,\partial_{1}u,\partial_{1}^{2}\eta)\|_{2}^{2} \lesssim \|(u,\eta,\partial_{1}\eta)\|_{3}\|(u,\partial_{1}\eta)\|_{\underline{1},3}^{2}, \qquad (2.41)$$

where, for each $t \in \overline{I_T}$,

$$\mathcal{E}_{\text{cul}}^L$$
 is equivalent to $\|\text{curl}_{\mathcal{A}}(u,\eta,\partial_1\eta)\|_2^2$, (2.42)

$$\mathcal{E}_{\text{cul}}^{H} \text{ is equivalent to } \|\text{curl}_{\mathcal{A}}(u,\partial_{1}\eta)\|_{\underline{1},2}^{2} + \|\text{curl}_{\mathcal{A}}\eta\|_{\underline{1},2}^{2}.$$
(2.43)

PROOF. We easily derive from (2.29) and (2.30) with i = 0 that

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathcal{E}_{\mathrm{cul}}^{L} + \|\mathrm{curl}_{\mathcal{A}}(u,\partial_{1}\eta)\|_{2}^{2} \lesssim \|(u,\eta)\|_{3}\|(u,\partial_{1}\eta)\|_{3}^{2} + \|u\|_{3}\|\eta\|_{3}^{2},$$
(2.44)

where \mathcal{E}_{cul}^{L} satisfies (2.42). Employing (2.13) with w = u and $\partial_1 \eta$, we further get (2.40) from (2.44).

We derive from (2.29) and (2.30) with i = 1 that

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathcal{E}_{\mathrm{cul}}^{H} + \|\mathrm{curl}_{\mathcal{A}} u\|_{2}^{2} + \|\mathrm{curl}_{\mathcal{A}} (u, \partial_{1}\eta)\|_{1,2}^{2} \lesssim \|(u, \eta, \partial_{1}\eta)\|_{3}(\|u\|_{3}^{2} + \|(u, \eta, \partial_{1}\eta)\|_{1,3}^{2}), \qquad (2.45)$$

where \mathcal{E}_{cul}^{H} satisfies (2.43). Consequently, making use of (2.13) and (2.14), we derive (2.40) and (2.41) from (2.44) and (2.45). This completes the proof of Proposition 2.2.

2.4. Equivalent estimates

This section is devoted to establishing the equivalent estimates for \mathcal{E}^L , \mathcal{D}^L , \mathcal{E}^H and \mathcal{D}^H .

Lemma 2.5. Under assumptions (1.16) and (1.17), we have

$$\mathcal{E}^L$$
 is equivalent to $\|(u,\eta,\partial_1\eta)\|_3^2$, (2.46)

$$\mathcal{D}^L$$
 is equivalent to $\|(u,\partial_1\eta)\|_3^2$, (2.47)

$$\mathcal{E}^H, \ \mathcal{D}^H \ and \ \|(u, \partial_1 u, \partial_1^2 \eta)\|_3^2 \ are \ equivalent.$$
 (2.48)

PROOF. To obtain (2.46)–(2.48), the key step is to establish the estimates of ∇q and u_t . By (1.11)₂, we have

$$\begin{cases} -\Delta q = f^1 & \text{in } \Omega, \\ \partial_3 q = f^2 \vec{\mathbf{n}}_3 & \text{on } \partial\Omega, \end{cases}$$
(2.49)

where $f^1 := \operatorname{div} \left(u_t + au - m\partial_1^2 \eta + \nabla_{\tilde{\mathcal{A}}} q \right)$ and $f^2 := -\tilde{\mathcal{A}}_{3j} \partial_j q$. Note that

$$\int f^1 \mathrm{d}y + \int_{\partial\Omega} f^2 \vec{\mathbf{n}}_3 \mathrm{d}y_\mathrm{h} = 0,$$

where $\vec{\mathbf{n}}_3$ is the third component of the outward unit normal vector $\vec{\mathbf{n}}$ on $\partial\Omega$, thus applying the elliptic estimate (A.16) to (2.49), and then using (1.11)₃, we have

$$\begin{aligned} \|\nabla q\|_{2} &\lesssim \|f^{1}\|_{1} + \|f^{2}\|_{2} \\ &\lesssim \|(\operatorname{div}_{\mathcal{A}_{t}}u, \operatorname{div}_{\tilde{\mathcal{A}}}u_{t}, \operatorname{div}_{\tilde{\mathcal{A}}}u)\|_{1} + \|(\partial_{1}^{2}\eta, \nabla_{\tilde{\mathcal{A}}}q)\|_{2} \\ &\lesssim \|\eta\|_{2,2} + \|(u,\eta)\|_{3}\|(u, \nabla q, u_{t})\|_{2}. \end{aligned}$$

$$(2.50)$$

In addition, by $(1.11)_2$, we have

$$\|u_t\|_2 \lesssim \|(u,\partial_1^2\eta,\nabla q)\|_2,$$

which, together with (2.50), yields, for sufficiently small δ ,

$$\|(u_t, \nabla q)\|_2 \lesssim \|(u, \partial_1^2 \eta)\|_2.$$
(2.51)

Consequently, we conclude the desired results by (2.51) and (A.3). The proof is complete.

2.5. Stability estimates

Now we are in the position to derive a priori stability estimates (1.13) and (1.14).

Making use of (2.6)–(2.10), and Hodge-type elliptic estimate (A.4), we can deduce that, for $0 \leq i \leq 1$ and $0 \leq j \leq 2$,

$$\|u\|_{i,3} \lesssim \|u\|_{i,0} + \|\operatorname{curl} u\|_{i,2} + \begin{cases} \|\eta\|_3 \|u\|_3 & \text{for } i = 0, \\ \|(u,\eta)\|_3 \|(u,\eta)\|_{1,3} & \text{for } i = 1 \end{cases}$$

$$(2.52)$$

and

$$\|\eta\|_{j,3} \lesssim \|\eta\|_{j,0} + \|\operatorname{curl}\eta\|_{j,2} + \begin{cases} \|\eta\|_3^2 & \text{for } j = 0; \\ \|\eta\|_3 \|\eta\|_{1,3} & \text{for } j = 1; \\ \|\eta\|_3 \|\eta\|_{2,3} + \|\eta\|_{1,3}^2 & \text{for } j = 2. \end{cases}$$
(2.53)

By (2.27), (2.41) and (A.3), we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{\mathcal{E}}^{H} + c\tilde{\mathcal{D}}^{H} \lesssim \sqrt{\mathcal{E}^{L}}\mathcal{D}^{H}, \qquad (2.54)$$

where

 $\tilde{\mathcal{E}}^H := \mathcal{E}_h^H + \mathcal{E}_{\text{cul}}^H \text{ and } \tilde{\mathcal{D}}^H := \|(u, \partial_1 u, \partial_1^2 \eta)\|_0^2 + \|\text{curl}(u, \partial_1 u, \partial_1^2 \eta)\|_2^2.$

In addition, it follows from (2.13), (2.14), (2.48), (2.52) and (2.53) that for sufficiently small δ ,

$$\tilde{\mathcal{E}}^{H}, \ \mathcal{E}^{H}, \ \tilde{\mathcal{D}}^{H}, \ \mathcal{D}^{H} \text{ and } \|(u, \partial_{1}u, \partial_{1}^{2}\eta)\|_{3}^{2} \text{ are equivalent.}$$
(2.55)

Thus we deduce from (2.54)–(2.55) that, for any sufficiently small δ ,

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{\mathcal{E}}^H + \mathcal{D}^H \leqslant 0$$

Exploiting (2.55), we further deduce from the above inequality that there exists a constant $c_1 > 0$ such that

$$e^{c_1 t} \mathcal{E}^H(t) + \int_0^t e^{c_1 s} \mathcal{D}^H(s) \mathrm{d}s \lesssim I_H^0.$$
(2.56)

In addition, by (2.26) and (2.40), we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{\mathcal{E}}^{L} + c\tilde{\mathcal{D}}^{L} \lesssim \|(u,\eta)\|_{3}\|(u,\partial_{1}\eta)\|_{3}^{2} + (\|u\|_{3} + \|\nabla q\|_{0})\|\eta\|_{3}^{2},$$
(2.57)

where

$$\tilde{\mathcal{E}}^L := \mathcal{E}_h^L + \mathcal{E}_{cul}^L$$
 and $\tilde{\mathcal{D}}^L := \|(u, \partial_1 \eta)\|_0^2 + \|curl(u, \partial_1 \eta)\|_2^2$.

In addition, thanks to (2.13), (2.46), (2.47), (2.52) and (2.53), we have, for sufficiently small δ ,

$$\tilde{\mathcal{E}}^L, \ \mathcal{E}^L \text{ and } \|(u,\eta,\partial_1\eta)\|_3^2 \text{ are equivalent}$$
 (2.58)

and

$$\tilde{\mathcal{D}}^L$$
 is equivalent to \mathcal{D}^L . (2.59)

Thanks to (2.59), we get from (2.57) that, for sufficiently small δ ,

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{\mathcal{E}}^L + c\tilde{\mathcal{D}}^L \lesssim (\|u\|_3 + \|\nabla q\|_0) \|\eta\|_3^2.$$

Exploiting (2.56), (2.58) and Gronwall's lemma, we deduce from the above estimate that

$$\mathcal{E}^{L}(t) + \int_{0}^{t} \mathcal{D}^{L}(s) \mathrm{d}s \lesssim e^{c\sqrt{I_{H}^{0}}} I_{L}^{0}.$$
(2.60)

In particular, by (2.56) and (2.60), there exist constants $c_2 \ge 1$ and $\delta_1 \in (0, 1]$ such that, for any $\delta \le \delta_1$,

$$\mathcal{E}^{L}(t) + \int_{0}^{t} \mathcal{D}^{L}(s) \mathrm{d}s \leqslant c_{2}^{2} e^{c_{2}\sqrt{I_{H}^{0}}} I_{L}^{0}, \qquad (2.61)$$

$$e^{c_1 t} \mathcal{E}^H(t) + \int_0^t e^{c_1 \tau} \mathcal{D}^H(s) \mathrm{d}s \leqslant c_2^2 I_H^0.$$
(2.62)

2.6. Proof of Theorem 1.1

Now we state the local well-posedness result for the initial-boundary value problem (1.11).

Proposition 2.3. Let b > 0 be a constant and $\iota > 0$ be the constant in Lemma A.4. We assume that $(u^0, \eta^0) \in H^{1,3}_{s} \times (H^{2,3}_{s} \cap H^3_{*})$, $\|(u^0, \partial_1 u^0, \partial_1^2 \eta^0)\|_3 \leq b$ and $\operatorname{div}_{\mathcal{A}^0} u^0 = 0$, where \mathcal{A}^0 is defined by ζ^0 and $\zeta^0 = \eta^0 + y$. Then there exist a constant $\delta_2 \in (0, \iota/2]$ such that if η^0 satisfies

$$\|\eta^0\|_{\underline{1},3} \leqslant \delta_2,$$

the initial-boundary value problem (1.11) admits a unique local classical solution $(u, \eta, q) \in \mathfrak{U}_T^{1,3} \times \mathfrak{C}^0(\overline{I_T}, H_s^{2,3}) \times C(\overline{I_T}, \underline{H}^3)$ for some local existence time T > 0. Moreover, (u, η, q) satisfies

$$\sup_{t\in\overline{I_T}} \|\eta\|_3 \leqslant 2\delta_2,$$

$$\sup_{t\in\overline{I_T}} \|(u,\eta,\partial_1\eta)\|_3 \leqslant c_3\sqrt{I_L^0} \text{ and } \sup_{t\in\overline{I_T}} \|(u,\partial_1\eta)\|_{\underline{1},3} \leqslant c_3\sqrt{I_H^0}$$
(2.63)

for some positive constant $c_3 \ge 1$. It should be noted that δ_2 and T depend on a, m and Ω ; moreover T further depends on b.

Remark 2.1. Here the uniqueness means that if there is another solution $(\tilde{u}, \tilde{\eta}, \tilde{q}) \in \mathfrak{U}_T^{1,3} \times \mathfrak{C}^0(\overline{I_T}, H_s^{2,3}) \times C(\overline{I_T}, \underline{H}^3)$ satisfying $0 < \inf_{(y,t) \in \mathbb{R}^2 \times \overline{I_T}} \det(\nabla \tilde{\eta} + I)$ and $\nabla_{(\nabla \tilde{\eta} + I)^{-T}} \tilde{q} \in L_T^{\infty} H^3$, then $(\tilde{\eta}, \tilde{u}, \tilde{q}) = (\eta, u, q)$ by virtue of the smallness condition " $\sup_{t \in \overline{I_T}} \|\eta\|_3 \leq 2\delta_2$ ". In addition, we have, by the fact " $\sup_{t \in \overline{I_T}} \|\eta\|_3 \leq \iota$ " and Lemma A.4,

$$\inf_{(y,t)\in\mathbb{R}^2\times I_T} \det(\nabla\eta + I) \ge 1/4.$$

PROOF. The proof of Proposition 2.3 will be provided in Section 3.

Thanks to the *priori* estimates (2.61), (2.62) and Proposition 2.3, we can easily establish Theorem 1.1. Next we briefly give the proof.

Let (u^0, η^0) satisfies the assumptions in Theorem 1.1,

$$e^{c_2 c_3} \sqrt{I_H^0} I_L^0 \leqslant \delta^2$$
 and $\delta = \min \left\{ \delta_1, \delta_2 \right\} / c_2 c_3^2$.

By Proposition 2.3, there exists a unique local solution (u, η, q) to the initial-boundary value problem (1.11) with a maximal existence time T^{\max} , which satisfies that

• for any $\tau \in I_{T^{\max}}$, the solution (u, η, q) belongs to $\mathfrak{U}^{1,3}_{\tau} \times \mathfrak{H}^{2,3}_{1,*,\tau} \times C^0(\overline{I_{\tau}}, \underline{H}^3)$,

$$\sup_{t\in I_{\tau}} \|\eta\|_3 \leqslant 2\delta_2;$$

• $\lim \sup_{t \to T^{\max}} \|\eta(t)\|_{\underline{1},3} > \delta_2 \text{ or } \limsup_{t \to T^{\max}} \|(u,\partial_1 u,\partial_1 \eta)(t)\|_3 = \infty, \text{ if } T^{\max} < \infty.$

Let

$$T^* := \sup \left\{ \tau \in I_{T^{\max}} \mid \| (u, \eta, \partial_1 \eta)(t) \|_3 \leqslant c_2 c_3^2 \delta \text{ for any } t \leqslant \tau \right\}.$$

We easily see that the definition of T^* makes sense and $T^* > 0$. Thus, to obtain the existence of global solutions, it suffices to verify $T^* = \infty$. Next we shall prove this fact by contradiction.

We assume that $T^* < \infty$. Then, for any given $T^{**} \in I_{T^*}$,

$$\sup_{t\in\overline{I_T^{**}}} \|(u,\eta,\partial_1\eta)(t)\|_3 \leqslant c_2 c_3^2 \delta \leqslant \delta_1.$$
(2.64)

Thanks to the estimate (2.64), we can follow the arguments of (2.61)–(2.62) and use a regularity method and (2.63) to derive that

$$\|e^{c_1 t} \mathcal{E}^H(t)\|_{L^{\infty}(I_{T^{**}})} + \int_0^{T^{**}} e^{c_1 \tau} \mathcal{D}^H(s) \mathrm{d}s \leqslant c_2^2 \lim_{t \to 0} \sup_{\tau \in (0,t)} \|(u,\partial_1 \eta)(\tau)\|_{\underline{1},3}^2 \leqslant c_2^2 c_3^2 I_H^0, \quad (2.65)$$

$$\|\mathcal{E}^{L}(t)\|_{L^{\infty}(I_{T^{**}})} + \int_{0}^{T^{**}} \mathcal{D}^{L}(s) \mathrm{d}s \leqslant c_{2}^{2} e^{c_{2}c_{3}} \sqrt{I_{H}^{0}} \lim_{t \to 0} \sup_{\tau \in (0,t)} \|(u,\eta,\partial_{1}\eta)\|_{3}^{2} \leqslant c_{2}^{2} c_{3}^{2} \delta^{2}.$$
(2.66)

By the weak continuity of (η, u) and the fact

$$\sup_{t\in\overline{I_{\tau}}} \|f\|_{0} = \|f\|_{L^{\infty}(I_{\tau},L^{2})} \text{ for any } f \in C(\overline{I_{\tau}},L^{2}_{\text{weak}}) \text{ with } \tau > 0,$$

we further derive from (2.65) and (2.66) that

$$\sup_{t\in\overline{I_T^{**}}} \|(u,\partial_1\eta)(t)\|_{\underline{1},3} \leqslant c_2 c_3 \sqrt{I_H^0},$$

$$\sup_{t\in\overline{I_T^{**}}} \|(u,\eta,\partial_1\eta)(t)\|_3 \leqslant c_2 c_3 \delta.$$
(2.67)

We take $(u(T^{**}, \eta(T^{**})))$ as a initial data. Noting that

$$||(u, \partial_1 u, \partial_1^2 \eta)(T^{**})||_3 \leq b := c_2 c_3 \sqrt{I_H^0} \text{ and } ||\eta(T^{**})||_{\underline{1},3} \leq c_2 c_3 \delta \leq \delta_2,$$

then, by Proposition 2.3 and (2.67), there exists a unique local classical solution, denoted by (u^*, η^*, q^*) , to the initial-boundary value problem (1.11) with $(u(T^{**}), \eta(T^{**}))$ in place of (u^0, η^0) ; moreover,

$$\sup_{t \in [T^{**},T]} \|(u^*,\eta^*,\partial_1\eta^*)\|_3 \leqslant c_2 c_3^2 \delta \text{ and } \sup_{t \in [T^{**},T]} \|\eta^*\|_3 \leqslant 2\delta_2,$$

where the local existence time T > 0 depends on b, a, m and Ω .

In view of the existence result of (u^*, η^*, q^*) , the uniqueness conclusion in Proposition 2.3 and the fact that T^{\max} denotes the maximal existence time, we immediately see that $T^{\max} > T^* + T/2$ and $\sup_{t \in [0, T^* + T/2]} ||(u, \eta, \partial_1 \eta)||_3 \leq c_2 c_3^2 \delta$. This contradicts with the definition of T^* . Hence $T^* = \infty$ and thus $T^{\max} = \infty$. This completes the proof of the existence of global solutions. The uniqueness of global solution is obvious due to the uniqueness of local solutions in Proposition 2.3 and the fact $\sup_{t \geq 0} ||\eta||_{1,3} \leq 2\delta_2$. In addition, it is obvious that (η, u) indeed enjoys (1.13) and (1.14) by the derivation of (2.65) and (2.66).

Finally, we derive (1.15). By (1.14), we easily see that

$$\partial_1 \eta(t) \to 0 \text{ in } H^3 \text{ as } t \to \infty$$
 (2.68)

and

$$\left\| \int_{0}^{t} u \mathrm{d}\tau \right\|_{3} \lesssim \int_{0}^{t} \|u\|_{3} \mathrm{d}\tau \lesssim \|(u^{0}, \partial_{1}\eta^{0})\|_{\underline{1},3} \text{ for any } t > 0$$
(2.69)

Thanks to (2.69), there exist a subsequence $\{t_n\}_{n=1}^{\infty}$ and some function $\eta^{\infty} \in H^3$ such that

$$\eta(t_n) - \eta^0 = \int_0^{t_n} u d\tau \to \eta^\infty - \eta^0$$
 weakly in H^3 .

Exploiting $(1.11)_1$ and (1.14), we have

$$\|\eta(t) - \eta^{\infty}\|_{3} \leq \liminf_{t_{n} \to \infty} \left\| \int_{t}^{t_{n}} u \mathrm{d}\tau \right\|_{3} \lesssim \sqrt{I_{H}^{0}} \liminf_{t_{n} \to \infty} \int_{t}^{t_{n}} e^{-c_{1}t/2} \mathrm{d}\tau \lesssim \sqrt{I_{H}^{0}} e^{-c_{1}t/2},$$

which, together with (2.68), yields that (1.15) holds and η^{∞} only depends on y_2 and y_3 . This completes the proof of Theorem 1.1.

3. Local well-posedness

This section is devoted to the proof of local well-posedness of the initial-boundary value problem (1.11). Next we roughly sketch the proof frame to establish the local well-posedness result.

Similarly to [11], in which the authors investigated the well-posdeness problem of incompressible inviscid MHD fluid equations with free boundary, we first alternatively use an iteration method to establish the existence result of the unique local solution (u, η, q) of the linear κ approximate problem

$$\begin{cases}
\eta_t - \kappa \partial_1^2 \eta = u, \\
u_t + \nabla_{\mathcal{B}} q + au = m \partial_1^2 \eta, \\
\operatorname{div}_{\mathcal{B}} u = 0, \\
(u, \eta)|_{t=0} = (u^0, \eta^0), \\
(u_3, \eta_3)|_{\partial\Omega} = 0, \\
\mathcal{B}_{3j} \partial_j q|_{\partial\Omega} = 0,
\end{cases}$$
(3.1)

where $\kappa > 0$, $\mathcal{B} := (\nabla \varsigma + I)^{-\mathrm{T}}$, and the given function ς belongs to some proper function space, see Proposition 3.4 in Section 3.1 for details. It should be noted that the Neumann boundary-value condition (3.1)₆ makes sure the solvability of q.

Then we derive the κ -independent estimates of the solutions, denoted by $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$ for $\kappa \in \mathbb{R}^+$, of the κ -approximate problem above, and then take the limit of $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$ with respect to $\kappa \to 0$ in a common definition domain. In particular, the limit function, denoted by (u, η, q) , is the unique local solution to the linearized problem

$$\begin{cases} \eta_t = u, \\ u_t + \nabla_{\mathcal{B}} q + au = m \partial_1^2 \eta, \\ \operatorname{div}_{\mathcal{B}} u = 0, \\ (u, \eta)|_{t=0} = (u^0, \eta^0), \\ (u_3, \eta_3)|_{\partial\Omega} = 0, \end{cases}$$
(3.2)

see Proposition 3.5 in Section 3.2 for details. It should be noted that the solution q of (3.2) automatically satisfies $(3.1)_6$ due to u_3 satisfying $\partial_t u_3|_{\partial\Omega} = 0$.

Finally, since the linearized problem (3.2) admits a unique local solution for any given ς , thus we easily arrive at Proposition 2.3 by a standard iteration method as in [6, 7], in which the authors investigated the well-posdeness problem of incompressible Euler equations with free boundary.

Now we turn to introducing some new notations appearing in this section. $P(x_1, \ldots, x_n)$ and $\dot{P}(x_1, \ldots, x_n)$ represent the generic polynomials with respect to the parameters x_1, \ldots, x_n , where all the coefficients in P and \dot{P} are equal one, and \dot{P} satisfies $\dot{P}(0, \ldots, 0) = 0$; $a \leq_{\kappa} b$ means that $a \leq c^{\kappa}b$, where c^{κ} denotes a generic positive constant, which may depend on κ , a, m and Ω . It should be noted that $P(x_1, \ldots, x_n)$, $\dot{P}(x_1, \ldots, x_n)$ and c^{κ} may vary from line to line.

We define the difference quotient with respect variable y_1 as follows:

$$D_1^{\tau} f(y_1, y_2, y_3) = (f(y_1 + \tau, y_2, y_3) - f(y_1, y_2, y_3)) / \tau \text{ for } |\tau| \in (0, 1).$$
(3.3)

The notation \mathcal{X} represents Id or D_1^{τ} , where Id denotes and identity mapping, and D_1^{τ} the difference quotient with respect to the variable y_1 .

We always use the notations \mathcal{B} , resp. \mathcal{J} to represent $(\nabla \varsigma + I)^{-T}$, resp. det $(\nabla \varsigma + I)$, where ς at least satisfies

$$\varsigma \in C^0(\overline{I_T}, H^3)$$
 and $\inf_{(y,t)\in\Omega_T} \det(\nabla\varsigma + I) \ge 1/4$ for some $T > 0.$ (3.4)

In addition, we define that

$$\mathbb{A}_{T,\iota}^{3,1/4} := \{ \psi \in C^0(\overline{I_T}, H_s^3) \mid \|\psi\|_3 \leqslant \iota \}$$
(3.5)

and

$$\mathbb{S}_T := \{ (w,\xi,\beta) \in C^0(\overline{I_T}, H^3_{\mathrm{s}}) \times C^0(\overline{I_T}, H^3_{\mathrm{s}}) \times (C^0(\overline{I_T}, \underline{H}^3)) \mid \partial_1^2 \xi \in C^0(\overline{I_T}, H^2_{\mathrm{s}}), \\ \partial_1^2 \xi \in L^2_T H^3, \ \nabla^3 \partial_1 \xi \in C^0_{B,\mathrm{weak}}(\overline{I_T}, L^2), \ \nabla_{\mathcal{B}}\beta \in L^2_T H^3, \ \mathcal{B}_{3j}\partial_j\beta|_{\partial\Omega} = 0 \},$$
(3.6)

where T > 0 and ι is the positive constant provided in Lemma A.4. It should be noted that the function ς , which belongs to $\mathbb{A}_{T,\iota}^{3,1/4}$, automatically satisfies (3.4) by Lemma A.4.

Finally, some preliminary estimates for \mathcal{B} and \mathcal{J} are collected as follows.

Lemma 3.1. Let ς satisfy (3.4) and i = 0, 1. Then

$$\|\mathcal{B}\|_{C^0(\bar{\Omega})} + \|(\mathcal{B}, \mathcal{J}^{-1})\|_2 \lesssim P(\|\varsigma\|_3), \tag{3.7}$$

$$\|\mathcal{B} - I\|_{i,2} \lesssim \|\varsigma\|_{i,3} P((\|\varsigma\|_3) \text{ for any } t \in \overline{I_T}.$$
(3.8)

If additionally ς further satisfies $(\partial_1 \varsigma, \varsigma_t) \in L^{\infty}_T H^{i,3}$, then, for a.e. $t \in I_T$,

$$\|\partial_t(\mathcal{B},\mathcal{J})\|_{i,2} \lesssim \dot{P}(\|(\varsigma,\varsigma_t)\|_{\underline{i},3}),\tag{3.9}$$

$$\|\mathcal{B}\|_{1+i,3} \lesssim P(\|\varsigma\|_{\underline{1+i},3}).$$
 (3.10)

PROOF. Since the derivation of the estimates (3.7)–(3.10) are very elementary, we omit it. \Box

3.1. Solvability of the linear κ -approximate problem (3.1)

In this section, we construct the unique local solution of the linear κ -approximate problem (3.1). To this purpose, we shall rewrite the linear κ -approximate problem (3.1) as the following equivalent problem (in the sense of classical solutions):

$$\begin{cases} \eta_t - \kappa \partial_1^2 \eta = u, \\ u_t + \nabla_{\mathcal{B}} q + au = m \partial_1^2 \eta, \\ -\Delta_{\mathcal{B}} q = K^1, \\ (u, \eta)|_{t=0} = (u^0, \eta^0), \\ (u_3, \eta_3)|_{\partial\Omega} = 0, \ \mathcal{B}_{3j} \partial_j q|_{\partial\Omega} = 0, \end{cases}$$
(3.11)

where $\Delta_{\mathcal{B}} := \operatorname{div}_{\mathcal{B}} \nabla_{\mathcal{B}}, \varsigma|_{t=0} = \eta^0, (u^0, \eta^0)$ satisfies $\operatorname{div}_{(\nabla \eta^0 + I)^{-T}} u^0 = 0, \operatorname{inf}_{y \in \overline{\Omega}} \det (\nabla \eta^0 + I)^{-T} > 0$, and

$$K^{1} := a \operatorname{div}_{\mathcal{B}} u - \operatorname{div}_{\mathcal{B}_{t}} u - \mathcal{J}_{t} \operatorname{div}_{\mathcal{B}} u / \mathcal{J} - m \operatorname{div}_{\mathcal{B}} \partial_{1}^{2} \eta.$$
(3.12)

Then the solvability of the linear κ -approximate problem (3.1) reduces to the solvability of the initial-boundary value problem (3.11).

We want to establish the local well-posteness result for (3.11) by an iteration method. To begin with, we shall investigate the solvability of the linear problem

$$\begin{cases} \eta_t - \kappa \partial_1^2 \eta = w, \\ u_t + au = K^2, \\ -\Delta_{\mathcal{B}} q = K^1, \\ (u, \eta)|_{t=0} = (u^0, \eta^0), \\ (u_3, \eta_3)|_{\partial\Omega} = 0, \ \mathcal{B}_{3j} \partial_j q|_{\partial\Omega} = 0, \end{cases}$$
(3.13)

where $(\xi, w, \theta) \in \mathbb{S}_T$ is given, and $K^2 := m\partial_1^2 \xi - \nabla_{\mathcal{B}} \theta$.

It is easy to see that the above linear problem is equivalent to the following three sub-problems: the initial-boundary value problem of partly dissipative equation for η

$$\begin{cases} \eta_t - \kappa \partial_1^2 \eta = w, \\ \eta|_{t=0} = \eta^0, \\ \eta_3|_{\partial\Omega} = 0, \end{cases}$$
(3.14)

the initial-value problem of ODE for u

$$\begin{cases} u_t + au = K^2, \\ u_{|t=0} = u^0, \\ u_{3|\partial\Omega} = 0 \end{cases}$$
(3.15)

and the Neumann boundary-value problem of elliptic equation for q

$$\begin{cases} -\Delta_{\mathcal{B}}q = K^1 & \text{in } \Omega, \\ \mathcal{B}_{3j}\partial_j q = 0 & \text{on } \partial\Omega. \end{cases}$$
(3.16)

Thus the solvability of the linear problem (3.13) reduces to the solvability of the three subproblems above. Next we establish the global well-posedness results for the above three subproblems in sequence.

Proposition 3.1. Let $\kappa > 0$, T > 0, $w \in L_T^2 H_s^3$ and $\eta^0 \in H_s^{1,3}$. Then the initial-boundary value problem (3.14) admits a unique solution $\eta \in C^0(\overline{I_T}, H_s^3)$, which satisfies $\partial_1^2 \eta \in C^0(\overline{I_T}, H_s^2)$, $\nabla^3 \partial_1 \eta \in C_{B,\text{weak}}^0(\overline{I_T}, L^2)$, and enjoys the estimates

$$\sup_{t \in \overline{I_T}} (\|\eta\|_3 + \sqrt{\kappa} \|\eta\|_{1,3}) + \|(\kappa \partial_1^2 \eta, \eta_t)\|_{L^2_T H^3} \lesssim \|\eta^0\|_3 + \sqrt{\kappa} \|\eta^0\|_{1,3} + (1 + \sqrt{T}) \|w\|_{L^2_T H^3},$$
(3.17)

$$\kappa \sup_{t \in \overline{I_s}} \|\eta^{\varepsilon}\|_{1,3}^2 \lesssim \kappa \|\eta^0\|_{1,3}^2 + \|w\|_{L_s^2 H^3}^2 \text{ for any } s \in (0,T].$$
(3.18)

PROOF. Let $\varepsilon \in (0,1)$, χ be a one-dimensional standard mollifier (see [24, pp. 38] for the definition), and $\chi^{\varepsilon}(s) := \chi(s/\varepsilon)/\varepsilon$. Let $\tilde{w} = w$ in Ω_T and $\tilde{w} = 0$ outside $\Omega \times (\mathbb{R} \setminus I_T)$. We define the mollification of \tilde{w} with respect to t:

$$S^t_{\varepsilon}(\tilde{w}) := \chi^{\varepsilon} * \tilde{w}$$

Then $S^t_{\varepsilon}(\tilde{w}) \in C^{\infty}(\mathbb{R}, H^{1,3}_s)$. We can check that

$$\begin{split} \|S^t_{\varepsilon}(\tilde{w})\|_{L^2_T H^3} &\lesssim \|w\|_{L^2_T H^3}, \\ S^t_{\varepsilon}(\tilde{w}) \to w \text{ strongly in } L^2_T H^3, \end{split}$$
(3.19)

Now we consider the τ -approximate problem for (3.14):

$$\begin{cases} \eta_t^{\tau} = L(\eta^{\tau}) + S_{\varepsilon}^t(\tilde{w}), \\ \eta^{\tau}|_{t=0} = \eta^0, \\ \eta_3^{\tau}|_{\partial\Omega} = 0, \end{cases}$$
(3.20)

where $L: H_{s}^{1,3} \to H_{s}^{1,3}$ by the rule $L(f) = \kappa D_{1}^{-\tau}(D_{1}^{\tau}f)$ for $f \in H_{s}^{1,3}$, see (3.3) for the definition of difference quotient D_{1}^{τ} .

It is easy to see that $L(\varpi) \in C^0(\overline{I_T}, H_s^{1,3})$ for $\varpi \in C^0(\overline{I_T}, H_s^{1,3})$ and $L \in \mathcal{L}(H_s^{1,3})$, where $\mathcal{L}(H_s^{1,3})$ is a set of all linear bounded operators of $H_s^{1,3}$. In particular, $L \in C^0(\overline{I_T}, \mathcal{L}(H_s^{1,3}))$. By existence theory of the initial-value problem of a abstract ODE equation (see [24, Proposition 2.17]), there exists a unique solution $\eta^{\tau} \in C^0(\overline{I_T}, H_s^{1,3}) \cap C^1(\overline{I_T}, H_s^{1,3})$ to (3.20). Obviously $\eta^{\tau}, \eta_t^{\tau}$ automatically belong to $L_T^2 H_s^{1,3}$ by the second conclusion in Lemma A.5 and the separability of $H_s^{1,3}$.

Let α satisfy $0 \leq |\alpha| \leq 3$. Applying ∂^{α} to $(3.20)_1$, and then multiplying the resulting identity by $\partial^{\alpha} \eta^{\tau}$ in L^2 , we have

$$\frac{\mathrm{d}}{\mathrm{d}t} \int |\partial^{\alpha} \eta^{\tau}|^{2} \mathrm{d}y + \kappa \int |D_{1}^{\tau} \partial^{\alpha} \eta^{\tau}|^{2} \mathrm{d}y = \int S_{\varepsilon}^{t} (\partial^{\alpha} \tilde{w}) \cdot \partial^{\alpha} \eta^{\tau} \mathrm{d}y.$$
(3.21)

Making use of (3.19), we easily deduce from (3.21) that,

$$\|\eta^{\tau}\|_{C^{0}(\overline{I_{T}},H^{3})} \lesssim \|\eta^{0}\|_{3} + \sqrt{T} \|w\|_{L^{2}_{T}H^{3}}.$$
(3.22)

In addition, we easily deduce from $(3.20)_1$ that

$$\frac{\kappa}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|D_1^{\tau} \partial^{\alpha} \eta^{\tau}\|_0^2 + \int |\partial^{\alpha} \eta_t^{\tau}|^2 \mathrm{d}y = \int (\partial^{\alpha} S_{\varepsilon}^t(\tilde{w})) \cdot \partial^{\alpha} \eta_t^{\tau} \mathrm{d}y.$$
(3.23)

Noting that

$$\|D_{1}^{\tau}\partial^{\alpha}\eta^{\tau}|_{t=0}\|_{0} \lesssim \|\partial^{\alpha}\eta^{0}\|_{1,0}, \qquad (3.24)$$

thus, making use of Young's inequality, (3.19), $(3.20)_1$ and (3.24), we deduce from (3.23) that, for any $s \in (0, T]$

$$\kappa \sup_{t \in \overline{I_s}} \|D_1^{\tau} \eta^{\tau}\|_3^2 + \|(L(\eta^{\tau}), \eta_t^{\tau})\|_{L_s^2 H^3}^2 \lesssim \kappa \|\eta^0\|_{1,3}^2 + \|w\|_{L_s^2 H^3}^2.$$
(3.25)

Thanks to the regularity of η^{τ} , we can we easily derive $(3.20)_1$ that, for any $\varphi \in H^1$ and for any $r, s \in I_T$,

$$\int D_1^{\tau} \partial^{\alpha} (\eta^{\tau}(y,t) - \eta^{\tau}(y,r)) \cdot \varphi dy$$

= $-\int_r^t \int (S_{\varepsilon}^t (\partial^{\alpha} \tilde{w}) + L(\partial^{\alpha} \eta^{\tau})) \cdot D_1^{-\tau} \varphi dy ds.$ (3.26)

Exploiting (3.19), the uniform estimate of $L(\eta^{\tau})$ in (3.25) and the fact

$$\|D_1^{-\tau}\varphi\|_0 \lesssim \|\varphi\|_{1,0},$$

we easily derive from the identity (3.26) that

$$D_1^{\tau} \nabla^3 \eta^{\tau}$$
 is uniformly continuous in H^{-1} . (3.27)

Here and in what follows, H^{-1} denotes the dual space of $H_0^1 := \{ w \in H^1 \mid w |_{\partial\Omega} = 0 \}.$

Making use of (3.22), (3.25), (3.27) and (A.7)–(A.9) there exists a sequence of $\{\eta^{\tau}\}_{|\tau|\in(0,1)}$ (still denoted by η^{τ} for simplicity) such that, as $\tau \to 0$,

$$\begin{aligned} &(\eta_t^{\tau}, D_1^{-\tau}(D_1^{\tau}\eta^{\tau})) \rightharpoonup (\eta_t^{\varepsilon}, \partial_1^2 \eta^{\varepsilon}) \text{ weakly in } L_T^2 H_{\mathrm{s}}^3, \ \eta^{\tau} \to \eta^{\varepsilon} \text{ strongly in } C^0(\overline{I_T}, H^2), \\ &(\eta^{\tau}, D_1^{\tau}\eta^{\tau}) \rightharpoonup (\eta^{\varepsilon}, \partial_1 \eta^{\varepsilon}) \text{ weakly-* in } L_T^{\infty} H_{\mathrm{s}}^3, \ D_1^{\tau} \nabla^3 \eta^{\tau} \to \nabla^3 \partial_1 \eta^{\varepsilon} \text{ in } C^0(\overline{I_T}, L_{\mathrm{weak}}^2). \end{aligned}$$

Moreover, the limit function η^{ε} is just the unique solution to the problem

$$\begin{cases} \eta^{\varepsilon}_t = \kappa \partial_1^2 \eta^{\varepsilon} + S^t_{\varepsilon}(\tilde{w}), \\ \eta^{\varepsilon}|_{t=0} = \eta^0, \\ \eta^{\varepsilon}_3|_{\partial\Omega} = 0 \end{cases}$$

and satisfies the following estimates:

$$\sup_{t \in \overline{I_T}} \|\eta^{\varepsilon}\|_3 \lesssim \|\eta^0\|_3 + \sqrt{T} \|w\|_{L^2_T H^3},$$

$$\kappa \sup_{t \in \overline{I_s}} \|\eta^{\varepsilon}\|_{1,3}^2 + \|(\kappa \partial_1^2 \eta^{\varepsilon}, \eta^{\varepsilon}_t)\|_{L^2_x H^3}^2 \lesssim \kappa \|\eta^0\|_{1,3}^2 + \|w\|_{L^2_x H^3}^2 \text{ for any } s \in (0,T].$$

$$(3.29)$$

Thus we further have $\partial_1^2 \eta^{\varepsilon} \in C^0(\overline{I_T}, H_s^2)$ and $\eta^{\varepsilon} \in C^0(\overline{I_T}, H_s^3)$.

Noting that η^{ε} enjoys the estimates (3.28) and (3.29), thus we get a limit function η , which satisfies the desired conclusion in Proposition 3.1, by taking limit of some sequence of $\{\eta^{\varepsilon}\}_{\varepsilon>0}$. We omit such limit process, since it is very similar to the argument of obtaining η^{ε} .

Proposition 3.2. Let $a \in \mathbb{R}$, $K^2 \in L_T^1 H_s^3$ and $u^0 \in H_s^3$, then the initial-boundary value problem (3.15) admits a unique solution $u \in C^0(\overline{I_T}, H_s^3)$, which satisfies $u_t \in L_T^1 H_s^3$ and

$$||u(t)||_3 \lesssim ||u^0||_3 + \int_0^t ||K^2(s)||_3 \mathrm{d}s.$$
 (3.30)

PROOF. Let

$$u = u^{0}e^{-at} + \int_{0}^{t} K^{2}(s)e^{-a(t-\tau)} \mathrm{d}s.$$
(3.31)

Since $K^2 \in L_T^1 H_s^3$ and $u^0 \in H_s^3$, it is easy to check that u given by (3.31) is the unique solution of (3.15); moreover, u belongs to $C^0(\overline{I_T}, H_s^3)$ and satisfies (3.30).

Proposition 3.3. Let T > 0, ι be the positive constant provided in Lemma A.4 and $\varsigma \in \mathbb{A}_{T,\iota}^{3,1/4}$ defined by (3.5). If $K^1 \in C^0(\overline{I_T}, H^1)$ and satisfies $\int K^1 \mathcal{J} dy = 0$ for $t \in \overline{I_T}$, then there exist a unique solution $q \in C^0(\overline{I_T}, \underline{H}^3)$, which satisfies the boundary-value problem (3.16) for each $t \in I_T$ and enjoys the estimate

$$\|q\|_{C^{0}(\overline{I_{T}},H^{3})} \lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}}) \|K^{1}\|_{L^{\infty}_{T}H^{1}}.$$
(3.32)

If we additionally assume that $K^1 \in L^r_T H^2$ for r = 2 or ∞ , then

$$\|\nabla_{\mathcal{B}}q\|_{L^{r}_{T}H^{3}} \lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}})\|K^{1}\|_{L^{r}_{T}H^{2}}.$$
(3.33)

PROOF. Let $\varphi = \varsigma + y$, then φ is a C^1 -diffeomorphism mapping by Lemma A.4. We denote φ^{-1} the inverse mapping of φ with respect to variable y, and then define $\tilde{K}^1 := K^1(\varphi^{-1}, t)$. Obviously $\tilde{K}^1 \in C^0(\overline{I_T}, H^1)$ by (A.11).

Now we consider the following Neumann boundary-value problem of elliptic equation

$$\begin{cases} -\Delta p = \tilde{K}^1 & \text{in } \Omega, \\ \partial_3 p = 0 & \text{on } \partial\Omega. \end{cases}$$
(3.34)

It is easy to see that

$$\int \tilde{K}^1 \mathrm{d}x = \int K^1 \mathcal{J} \mathrm{d}y = 0.$$

By Lemma A.6, there exists a unique solution $p \in C^0(\overline{I_T}, \underline{H}^3)$ of the boundary-value problem (3.34) such that

$$\|p\|_{C^0(\overline{I_T}, H^3)} \lesssim \|\tilde{K}^1\|_{L^{\infty}_T H^1}.$$
(3.35)

Let $\tilde{q} = p(\varphi)$, then $\tilde{q} \in C^0(\overline{I_T}, H^3)$ by (A.10). We further define that $q := \tilde{q} - (\tilde{q})_{\Omega}$, then $q \in C^0(\overline{I_T}, \underline{H}^3)$ and satisfies (3.32) by (3.35) and (A.13). Since p satisfies (3.34), we see that q is the solution of the boundary-value problem (3.16). In addition, the uniqueness of the solution of the problem (3.16) is obvious in the class $C^0(\overline{I_T}, \underline{H}^3)$.

If we additionally assume that $K^1 \in L^r_T H^2$, then $\tilde{K}^1 \in L^r_T H^2$, where r = 2 and ∞ . By the second assertion in Lemmas A.5–A.6, (A.6) and the separability of \underline{H}^3 , we have $p \in L^r_T H^3$ and

$$\|p\|_{L^{r}_{T}H^{4}} \lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}}) \|\tilde{K}^{1}\|_{L^{r}_{T}H^{2}}.$$
(3.36)

By (3.36), (A.12) and (A.13),

$$\begin{aligned} |\nabla_{\mathcal{B}}q\|_{L^{r}_{T}H^{3}} &= \|\nabla p(x,t)|_{x=\varphi(y,t)}\|_{L^{r}_{T}H^{3}} \\ &\lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}})\|\nabla p\|_{L^{r}_{T}H^{3}} \lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}})\|K^{1}\|_{L^{r}_{T}H^{2}} \end{aligned}$$

which yields (3.33).

With Propositions 3.1–3.3 in hand, next we will use an iterative method to establish the existence result of a unique local solution to the linear κ -approximate problem (3.1).

Proposition 3.4. Let $\mathbb{A}_{\alpha,\iota}^{3,1/4}$, resp. \mathbb{S}_{α} be defined by (3.5) resp. (3.6) with some positive constant α in place of T. We assume that $\kappa > 0$, $a \ge 0$, $(u^0, \eta^0) \in H_s^3 \times H_s^{1,3}$ and $\varsigma \in \mathbb{A}_{\alpha,\iota}^{3,1/4}$ satisfies $\varsigma_t \in C^0(\overline{I_{\alpha}}, H^2) \cap L_{\alpha}^{\infty}H^3$, then the κ -approximate problem (3.1) defined on Ω_{α} admits a unique solution, denoted by $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$, which belongs to \mathbb{S}_{α} .

PROOF. Let $T \leq \alpha$. In view of Propositions 3.1–3.3, we easily see that there exists a solution sequence $\{(u^n, \eta^n, q^n)\}_{n=1}^{\infty}$ defined on $\overline{I_T}$. Moreover,

- (1) $(u^1, \eta^1, q^1) = 0;$
- (2) $(u^n, \eta^n, Q^n) \in \mathbb{S}_T$ for $n \ge 1$;
- (3) (u^n, η^n, q^n) for $n \ge 2$ enjoys the following relations:

$$\begin{cases} \eta_t^n - \kappa \partial_1^2 \eta^n = u^{n-1}, \\ u_t^n + a u^n = m \partial_1^2 \eta^{n-1} - \nabla_{\mathcal{B}} q^{n-1} =: K^{2,n}, \\ -\Delta_{\mathcal{B}} q^n = a \operatorname{div}_{\mathcal{B}} u^n - \operatorname{div}_{\mathcal{B}_t} u^n - \mathcal{J}_t \operatorname{div}_{\mathcal{B}} u^n / \mathcal{J} - m \operatorname{div}_{\mathcal{B}} \partial_1^2 \eta^n =: K^{1,n}, \\ (u^n, \eta^n)|_{t=0} = (u^0, \eta^0), \\ (u_3^n, \eta_3^n)|_{\partial\Omega} = 0, \ \mathcal{B}_{3j} \partial_j q^n|_{\partial\Omega} = 0, \end{cases}$$

where $K^{1,n} \in C^0(\overline{I_T}, H^1) \cap L^2_T H^2$, $K^{2,n} \in L^2_T H^3_s$ and $\int K^{1,n} \mathcal{J} dy = 0 \text{ for each } t \in \overline{I_T}.$

It should be noted that we have used the relation $\partial_j(\mathcal{JB}_{ij}) = 0$ for i = 1, 2, and the regularity $\varsigma_t \in C^0(\overline{I_\alpha}, H^1)$ in the derivation of the above identity.

We further define $(\bar{u}^{n+1}, \bar{\eta}^{n+1}, \bar{q}^{n+1}) := (u^{n+1} - u^n, \eta^{n+1} - \eta^n, q^{n+1} - q^n)$ and $(\bar{u}^n, \bar{\eta}^n, \bar{q}^n) := (u^n - u^{n-1}, \eta^n - \eta^{n-1}, q^n - q^{n-1})$ for $n \ge 2$. Then we have

$$\begin{cases} \bar{\eta}_{t}^{n+1} - \kappa \partial_{1}^{2} \bar{\eta}^{n+1} = \bar{u}^{n}, \\ \bar{u}_{t}^{n+1} + a \bar{u}^{n+1} = m \partial_{1}^{2} \bar{\eta}^{n} - \nabla_{\mathcal{B}} \bar{q}^{n}, \\ -\Delta_{\mathcal{B}} \bar{q}^{n+1} = a \mathrm{div}_{\mathcal{B}} \bar{u}^{n+1} - \mathrm{div}_{\mathcal{B}_{t}} \bar{u}^{n+1} - \mathcal{J}_{t} \mathrm{div}_{\mathcal{B}} \bar{u}^{n+1} / \mathcal{J} - m \mathrm{div}_{\mathcal{B}} (\partial_{1}^{2} \bar{\eta}^{n+1}), \\ (\bar{u}^{n+1}, \bar{\eta}^{n+1})|_{t=0} = (0, 0), \\ (\bar{u}_{3}^{n+1}, \bar{\eta}_{3}^{n+1})|_{\partial\Omega} = 0, \\ \mathcal{B}_{3j} \partial_{i} \bar{q}^{n+1}|_{\partial\Omega} = 0. \end{cases}$$

$$(3.37)$$

Applying Propositions 3.1-3.3 to (3.37), and then using (3.7) and (3.9), we get

$$\begin{split} \|\bar{\eta}^{n+1}\|_{C^{0}(\overline{L_{T}},H^{3})} &+ \|\partial_{1}\bar{\eta}^{n+1}\|_{L_{T}^{\infty}H^{3}} + \|\partial_{1}^{2}\bar{\eta}^{n+1}\|_{L_{T}^{2}H^{3}} \lesssim_{\kappa} T(1+\sqrt{T})\|\bar{u}^{n}\|_{L_{T}^{\infty}H^{3}}, \\ \|\bar{u}^{n+1}\|_{C^{0}(\overline{L_{T}},H^{3})} &\lesssim \sqrt{T}\|(\partial_{1}^{2}\bar{\eta}^{n},\nabla_{\mathcal{B}}\bar{q}^{n})\|_{L_{T}^{2}H^{3}}, \\ \|\bar{q}^{n+1}\|_{C^{0}(\overline{L_{T}},H^{3})} &\lesssim_{\kappa} P(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\|(\bar{u}^{n+1},\partial_{1}^{2}\bar{\eta}^{n+1})\|_{L_{T}^{\infty}H^{2}}, \\ \|\nabla_{\mathcal{B}}\bar{q}^{n+1}\|_{L_{T}^{2}H^{3}} \lesssim_{\kappa} P(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\left(\sqrt{T}\|\bar{u}^{n+1}\|_{L_{T}^{\infty}H^{3}} + \|\partial_{1}^{2}\bar{\eta}^{n+1}\|_{L_{T}^{2}H^{3}}\right). \end{split}$$

In addition, by $(3.37)_1$ and $(3.37)_2$,

$$\|(\bar{\eta}_t^{n+1}, \bar{u}_t^{n+1})\|_{C^0(\overline{I_T}, H^2)} \lesssim \|(\bar{u}^{n+1}, \bar{u}^n, \partial_1^2 \bar{\eta}^{n+1}, \partial_1^2 \bar{\eta}^n, \nabla \bar{q}^n)\|_{L^{\infty}_T H^2}.$$

Thus we immediately see from the above five estimates that

$$\{(u^n,\eta^n,q^n,\eta^n_t,u^n_t)\}_{n=1}^{\infty}$$

is a Cauchy sequence in $C^0(\overline{I_T}, H_s^3) \times C^0(\overline{I_T}, H_s^3) \times C^0(\overline{I_T}, \underline{H}^3) \times C^0(\overline{I_T}, H_s^2) \times C^0(\overline{I_T}, H_s^2)$. Hence there exists one limit function $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$, which is the unique local solution of the linear κ approximate problem (3.11) with $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$ in place of (η, u, q) , and also the unique local solution of the linear κ -approximate problem (3.1). In addition, it is easy to see that the solution $(u^{\kappa}, \eta^{\kappa}, q^{\kappa}) \in \mathbb{S}_T$ by further using Propositions 3.1 and 3.3.

Noting that the local time T_1 is independent of the initial data and the local solution constructed above satisfies $(u, \eta)|_{t=T} \in H^3_s \times H^{1,3}_s$, thus, if $T < \alpha$, we can further extend the local solution to be a global solution defined on $\overline{I_\alpha}$ by finite steps; moreover the obtained global solution is the unique solution of the linear κ -approximate problem (3.1) and belongs to \mathbb{S}_α . This completes the proof of Proposition 3.4.

3.2. Solvability of linearized problem (3.1)

To investigate the solvability of linearized problem (3.2), we shall first κ -independent estimates of the solutions of the κ -approximate problem.

Lemma 3.2. Under the assumptions of Proposition 3.4, we further assume that $\kappa \in (0,1)$, $\|\eta^0\|_{1,3} \leq \delta \in \mathbb{R}^+$, $(\partial_1 u^0, \partial_1^2 \eta^0) \in H^3$ and $\varsigma \in \mathbb{A}^{3,1/4}_{\alpha,\varsigma}$ satisfies

$$\partial_1 \varsigma \in L^\infty_\alpha H^{1,3}, \ \varsigma_t \in L^\infty_\alpha H^{1,3} \cap C^0(\overline{I_T}, H^2), \ \varsigma|_{t=0} = \eta^0.$$

Then there exist polynomials $P(I_H^0, \|(\varsigma, \partial_1\varsigma, \varsigma_t)\|_{L^{\infty}_{\alpha}H^3})$, $P(\|(\varsigma, \partial_1\varsigma, \varsigma_t)\|_{L^{\infty}_{\alpha}H^{1,3}})$, positive constants $c, c_3 \ge 1$, and $\delta_0 > 0$ (may depending on a, m and Ω), such that, for any $\delta \le \delta_0$, the local

solution $(\eta^{\kappa}, u^{\kappa}, q^{\kappa})$ provided by Proposition 3.4 belongs to $\mathfrak{U}_{T_1}^{1,3} \times \mathfrak{C}^0(\overline{I_{T_1}}, H_s^{2,3}) \times C^0(\overline{I_{T_1}}, \underline{H}^3)$ and enjoys the following estimates:

$$\sup_{t\in\overline{I_{T_1}}}\mathcal{E}^L_\kappa(t) + \kappa \|\partial_1^2\eta\|_{L^2_TH^3}^2 \leqslant 4c_3^2 I_L^0$$

$$(3.38)$$

$$\sup_{t\in\overline{I_{T_1}}}\mathcal{E}^H_\kappa(t)\leqslant 4c_3^2I_H^0,\tag{3.39}$$

$$\|q^{\kappa}\|_{C^{0}(\overline{I_{T_{1}}},H^{3})} + \|(u_{t}^{\kappa},\eta_{t}^{\kappa},\nabla_{\mathcal{B}}q^{\kappa})\|_{L^{\infty}_{T_{1}}H^{3}} \lesssim P\left(I^{0}_{H},\|(\varsigma,\varsigma_{t})\|^{2}_{L^{\infty}_{T_{1}}H^{3}}\right),$$
(3.40)

where $\mathcal{E}_{\kappa}^{L}(t) := \|(\eta^{\kappa}, \partial_{1}\eta^{\kappa}, u^{\kappa})(t)\|_{3}^{2}, \ \mathcal{E}_{\kappa}^{H}(t) := \|(u^{\kappa}, \partial_{1}\eta^{\kappa})(t)\|_{\underline{1},3}^{2}$ and

$$T_1 := \min\{1/3c_5 P(\|(\varsigma, \partial_1\varsigma, \varsigma_t)\|_{L^{\infty}_{\alpha}H^{1,3}}), \alpha, 1\}.$$
(3.41)

Moreover, for $(\eta^{\kappa}, u^{\kappa})$ restricted on $[0, T_1]$,

$$\partial_1 \nabla^3(u^{\kappa}, \partial_1 \eta^{\kappa})$$
 is uniformly continuous in H^{-1} . (3.42)

PROOF. Let $T \leq \min\{\alpha, 1\}$,

$$\|\eta^0\|_{\underline{1},3} \leqslant \delta \in (0,1]. \tag{3.43}$$

We define that

$$\|\cdot\|_{*,i}^2 := \|(\cdot, D_1^{\tau} \cdot)\|_i^2$$
 and $\mathcal{E}_{\kappa}^{\tau}(t) := \|(u^{\kappa}, \partial_1 \eta^{\kappa})(t)\|_{*,3}^2$.

In addition, we denote $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$ by (u, η, q) for simplicity. Next we establish the desired uniform estimates for (u, η, q) by seven steps.

(1) Estimates of η

Recalling (3.17), it is easy to see that

$$\sup_{t \in \overline{I_T}} \|\eta(t)\|_3^2 + \kappa^2 \|\partial_1^2 \eta\|_{L^2_T H^3}^2 \lesssim I_L^0 + T \sup_{t \in \overline{I_T}} \|u(t)\|_3^2.$$
(3.44)

By (3.18), we have

$$\limsup_{s \to 0^+} \sup_{t \in \overline{I_s}} \|\eta(t)\|_{1,3}^2 \lesssim \|\eta^0\|_{1,3}^2.$$
(3.45)

Since

$$\begin{cases} D_{1}^{\tau}\eta_{t} - \kappa D_{1}^{\tau}\partial_{1}^{2}\eta = D_{1}^{\tau}w, \\ D_{1}^{\tau}\eta|_{t=0} = \eta^{0}, \\ D_{1}^{\tau}\eta_{3}|_{\partial\Omega} = 0, \end{cases}$$
(3.46)

applying Proposition 3.1 to (3.46) yields that

$$\limsup_{s \to 0^+} \sup_{t \in \overline{I_s}} \|\partial_1 D_1^{\tau} \eta(t)\|_3^2 \lesssim \|\eta^0\|_{2,3}^2.$$
(3.47)

(2) Estimates of q

Noting that q satisfies

$$\begin{cases} -\Delta_{\mathcal{B}}q = K^1 & \text{in } \Omega, \\ \mathcal{B}_{3j}\partial_i q = 0 & \text{on } \partial\Omega, \end{cases}$$
(3.48)

where K^1 is defined in (3.12). Applying the estimate (3.32) to the above boundary-value problem, and then using (3.7) and (3.9), we have

$$\|q\|_{C^{0}(\overline{I_{T}}H^{3})} \lesssim P\left(\|\varsigma\|_{L_{T}^{\infty}H^{3}}\right) \|K^{1}\|_{L_{T}^{\infty}H^{1}} \lesssim P\left(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}\right) \sup_{t\in\overline{I_{T}}} \|(u,\partial_{1}^{2}\eta)(t)\|_{2}.$$
 (3.49)

(3) L^2 -norm estimates of $(u, \partial_1 \eta)$.

Applying \mathcal{X} to $(3.1)_2$ yields that

$$\partial_t \mathcal{X} u + a \mathcal{X} u = m \mathcal{X} \partial_1^2 \eta - \mathcal{X} \nabla_{\mathcal{B}} q \tag{3.50}$$

Multiplying (3.50) by $\mathcal{X}u$ in L^2 yields that (for a.e. $t \in I_T$)

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int |\mathcal{X}u|^2 \mathrm{d}y + \int (a|\mathcal{X}u|^2 \mathrm{d}y - m(\mathcal{X}\partial_1^2\eta) \cdot \mathcal{X}u) \mathrm{d}y = -\int (\mathcal{X}\nabla_{\mathcal{B}}q) \cdot \mathcal{X}u \mathrm{d}y.$$
(3.51)

By (3.7), (3.10) and (3.49), we can estimate that

$$-\int \mathcal{X}\nabla_{\mathcal{B}}q \cdot \mathcal{X}u \mathrm{d}y \lesssim \|\mathcal{B}\|_{\underline{1},2} \|\nabla q\|_{\underline{1},0} \|u\|_{1,0} \lesssim P\left(\|(\varsigma,\partial_1\varsigma,\varsigma_t)\|_{L^{\infty}_T H^3}\right) \sup_{t\in\overline{I_T}} \|(u,\partial_1^2\eta)\|_2^2.$$
(3.52)

In addition, by $(3.1)_1$ and the integration by parts, we get

$$-m\int (\mathcal{X}\partial_1^2\eta) \cdot \mathcal{X}u \mathrm{d}y = \frac{m}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int |\mathcal{X}\partial_1\eta|^2 \mathrm{d}y + m\kappa\int |\mathcal{X}\partial_1^2\eta|^2 \mathrm{d}y.$$
(3.53)

Plugging (3.52)–(3.53) into (3.51) and then integrating the resulting inequality over (0, t), we arrive at

$$\begin{aligned} &\|\mathcal{X}\left(u,\partial_{1}\eta\right)\left(t\right)\|_{0}^{2}+\kappa\int_{0}^{t}\|\mathcal{X}\partial_{1}^{2}\eta(s)\|_{0}^{2}\mathrm{d}s\\ &\lesssim\|\left(u^{0},\partial_{1}\eta^{0}\right)\|_{\underline{1},0}^{2}+TP\left(\|(\varsigma,\partial_{1}\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}\right)\sup_{t\in\overline{I_{T}}}\|(u,\partial_{1}^{2}\eta)(t)\|_{2}^{2}. \end{aligned}$$
(3.54)

(4) Curl estimates of $(u, \partial_1 \eta)$.

Applying $\operatorname{curl}_{\mathcal{B}}$ to $(3.1)_2$ yields that

 $\partial_t \operatorname{curl}_{\mathcal{B}} u + a \operatorname{curl}_{\mathcal{B}} u = m \operatorname{curl}_{\mathcal{B}} \partial_1^2 \eta + \operatorname{curl}_{\mathcal{B}_t} u.$

Let the multiindex α satisfy $|\alpha| \leq 2$. Applying $\partial^{\alpha} \mathcal{X}$ to the above identity yields

$$\partial_t \partial^\alpha \mathcal{X} \mathrm{curl}_{\mathcal{B}} u + a \partial^\alpha \mathcal{X} \mathrm{curl}_{\mathcal{B}} u = m \partial_1 \partial^\alpha \mathcal{X} \mathrm{curl}_{\mathcal{B}} \partial_1 \eta + K_3^{\mathcal{X},\alpha}, \tag{3.55}$$

where $K_3^{\mathcal{X},\alpha} := \partial^{\alpha} \mathcal{X} \operatorname{curl}_{\mathcal{B}_t} u - m \partial^{\alpha} \mathcal{X} \operatorname{curl}_{\partial_1 \mathcal{B}} \partial_1 \eta$. Multiplying (3.55) by $\partial^{\alpha} \mathcal{X} \operatorname{curl}_{\mathcal{B}} u$ in L^2 , we obtain

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int |\partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}u|^{2}\mathrm{d}y + a\int |\partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}u|^{2}\mathrm{d}y - m\int (\partial_{1}\partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}\partial_{1}\eta) \cdot \partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}u\mathrm{d}y = \int K_{3}^{\mathcal{X},\alpha} \cdot \partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}u\mathrm{d}y =: I_{8}.$$
(3.56)

Using the integration by parts, $(3.1)_1$ and a regularity method, we obtain

$$-m\int \left(\partial_{1}\partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}\partial_{1}\eta\right)\cdot\partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}u\mathrm{d}y$$
$$=\frac{m}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int |\partial^{\alpha}\mathcal{X}\mathrm{curl}_{\mathcal{B}}\partial_{1}\eta|^{2}\mathrm{d}y+m\kappa\int |\partial^{\alpha}\mathcal{X}\mathrm{curl}\partial_{1}^{2}\eta|^{2}\mathrm{d}y+I_{9}+I_{10},\qquad(3.57)$$

where we have defined that

$$I_{9} := m \int (\partial^{\alpha} \mathcal{X} \operatorname{curl}_{\mathcal{B}} \partial_{1} \eta) \cdot \partial^{\alpha} \mathcal{X} \left(\operatorname{curl}_{\partial_{1} \mathcal{B}} u - \operatorname{curl}_{\partial_{t} \mathcal{B}} \partial_{1} \eta \right) \mathrm{d}y,$$

$$I_{10} := m \kappa \int \left(\left(\partial^{\alpha} \mathcal{X} \operatorname{curl}_{\partial_{1} \mathcal{B}} \partial_{1} \eta \right) \cdot \partial^{\alpha} \mathcal{X} \operatorname{curl}_{\mathcal{B}} \partial_{1}^{2} \eta + |\partial^{\alpha} \mathcal{X} \operatorname{curl}_{\mathcal{B}-I} \partial_{1}^{2} \eta|^{2} + 2 \left(\partial^{\alpha} \mathcal{X} \operatorname{curl}_{\mathcal{B}-I} \partial_{1}^{2} \eta \right) \cdot \partial^{\alpha} \mathcal{X} \operatorname{curl}_{\partial_{1}^{2}} \eta \right) \mathrm{d}y.$$

By the estimates (3.7), (3.9) and (3.10), we can estimate that,

• for the case $\mathcal{X} = \mathrm{Id}$,

$$I_{8} + I_{9} \lesssim P(\|(\varsigma, \partial_{1}\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\|(u, \partial_{1}\eta)\|_{3}^{2},$$

$$I_{10} \lesssim \kappa \left(P(\|\varsigma\|_{L_{T}^{\infty}H^{1,3}})\|\eta\|_{1,3}(\|\operatorname{curl}\eta\|_{2,2} + \|\mathcal{B} - I\|_{2}\|\eta\|_{2,3}) + \|\mathcal{B} - I\|_{2}\left(\|\mathcal{B} - I\|_{2}\|\eta\|_{2,3}^{2} + \|\eta\|_{2,3}\|\operatorname{curl}\eta\|_{2,2}\right)\right).$$

$$(3.59)$$

• for the case $\mathcal{X} = D_1^{\tau}$,

$$I_8 + I_9 \lesssim P(\|(\varsigma, \partial_1 \varsigma, \varsigma_t)\|_{L^{\infty}_T H^{1,3}}) \|(u, \partial_1 \eta)\|^2_{*,3},$$
(3.60)

$$I_{10} \lesssim \kappa \left(P(\|\varsigma\|_{L_T^{\infty} H^{2,3}}) \|\partial_1 \eta\|_{*,3} (\|\operatorname{curl} \partial_1^2 \eta\|_{*,2} + \|\mathcal{B} - I\|_{\underline{1},2} \|\partial_1^2 \eta\|_{*,3}) + \|\mathcal{B} - I\|_{\underline{1},2} (\|\mathcal{B} - I\|_{\underline{1},2} \|\partial_1^2 \eta\|_{*,3}^2 + \|\partial_1^2 \eta\|_{*,3} \|\operatorname{curl} \partial_1^2 \eta\|_{*,2}) \right).$$
(3.61)

Noting that $\varsigma|_{t=0} = \eta^0$, by Newton–Leibniz formula, (3.8) with t = 0, (3.9) and (3.43), we have

$$\|(\mathcal{B}(t) - I)\|_{\underline{i},2} \lesssim \|(\mathcal{B}^{0} - I)\|_{\underline{i},2} + \int_{0}^{t} \|\mathcal{B}_{\tau}(s)\|_{\underline{i},2} \,\mathrm{d}s$$

$$\lesssim \begin{cases} \|\eta^{0}\|_{3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}) & \text{for } i = 0; \\ \|\eta^{0}\|_{1,3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}}) & \text{for } i = 1. \end{cases}$$
(3.62)

Making use of the above estimates (3.58)–(3.62), Young's inequality and the fact " $\kappa \in (0, 1)$ ", we deduce from (3.56)–(3.57) that

$$\frac{\mathrm{d}}{\mathrm{d}t} \|\mathrm{curl}_{\mathcal{B}}(u,\partial_{1}\eta)\|_{2}^{2} + c\kappa \|\mathrm{curl}\eta\|_{2,2}^{2}
\lesssim P(\|(\varsigma,\partial_{1}\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\|(u,\partial_{1}\eta)\|_{3}^{2}
+ \kappa \left(\|\eta^{0}\|_{3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\right)\|\eta\|_{2,3}^{2}$$
(3.63)

and

$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \| \mathrm{curl}_{\mathcal{B}}(u, \partial_{1}\eta) \|_{*,2}^{2} + c\kappa \| \mathrm{curl} \partial_{1}^{2}\eta \|_{*,2}^{2}
\lesssim P(\|(\varsigma, \partial_{1}\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}}) \| (u, \partial_{1}\eta) \|_{*,3}^{2}
+ \kappa \left(\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}}) \right) \|\partial_{1}^{2}\eta \|_{*,3}^{2}.$$
(3.64)

In addition, similarly to (3.62), we have

$$\begin{split} \|\mathcal{X}\mathrm{curl}f(t)\|_{2} &\lesssim \left\|\mathcal{X}\left(\mathrm{curl}_{\mathcal{B}^{0}-I}f(t),\mathrm{curl}_{\mathcal{B}}f(t),\mathrm{curl}_{\int_{0}^{t}\mathcal{B}_{s}\mathrm{d}s}f(t)\right)\right\|_{2} \\ &\lesssim \|\mathcal{X}\mathrm{curl}_{\mathcal{B}}f(t)\|_{2} \\ &+ \begin{cases} (\|\eta^{0}\|_{3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}))\|f(t)\|_{3} & \text{for } \mathcal{X} = \mathrm{Id}; \\ (\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}})\|f(t)\|_{*,3} & \text{for } \mathcal{X} = D_{1}^{\tau}. \end{cases}$$

Integrating (3.63) and (3.64) over (0, t), resp., and then using (3.43), (3.45) and the above estimate, we conclude that

$$\begin{aligned} \|(\operatorname{curl} u, \operatorname{curl} \partial_{1} \eta)(t)\|_{2}^{2} + \kappa \int_{0}^{t} \|\operatorname{curl} \eta(s)\|_{2,2}^{2} \mathrm{d}s \\ &\lesssim I_{L}^{0} + (\|\eta^{0}\|_{3}^{2} + TP(\|(\varsigma, \partial_{1}\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{3}})) \sup_{t \in \overline{I_{T}}} \|(u, \partial_{1}\eta)(t)\|_{3}^{2} \\ &+ \kappa \left(\|\eta^{0}\|_{3} + TP(\|(\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\right) \int_{0}^{t} \|\eta(s)\|_{2,3}^{2} \mathrm{d}s \end{aligned}$$
(3.65)

and

$$\begin{aligned} \|(\operatorname{curl} u, \operatorname{curl} \partial_{1} \eta)(t)\|_{*,2}^{2} + \kappa \int_{0}^{t} \|\operatorname{curl} \partial_{1}^{2} \eta(s)\|_{*,2}^{2} \mathrm{d}s \\ &\lesssim I_{H}^{0} + (\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma, \partial_{1}\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}})) \sup_{t \in \overline{I_{T}}} \|(u, \partial_{1}\eta)(t)\|_{*,3}^{2} \\ &+ \kappa P\left(\|\eta^{0}\|_{\underline{1},3} + T(\|(\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}})\right) \int_{0}^{t} \|\partial_{1}^{2} \eta(s)\|_{*,3}^{2} \mathrm{d}s. \end{aligned}$$
(3.66)

(5) Divergence estimates of $(u, \partial_1 \eta)$. Similarly to (3.62), we have

$$\|\mathcal{X}\operatorname{div} f(t)\|_{2} \lesssim \left\|\mathcal{X}\left(\operatorname{div}_{\mathcal{B}^{0}-I}f(t),\operatorname{div}_{\mathcal{B}}f(t),\operatorname{div}_{\int_{0}^{t}\mathcal{B}_{\tau}\mathrm{d}\tau}f(t)\right)\right\|_{2}.$$

Noting that $\operatorname{div}_{\mathcal{B}} u = 0$, thus taking f = u in the above estimate yields

$$\|\mathcal{X} \text{div} u\|_{2} \lesssim \begin{cases} (\|\eta^{0}\|_{3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}))\|u\|_{3} & \text{for } \mathcal{X} = \text{Id}; \\ (\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}}))\|u\|_{*,3} & \text{for } \mathcal{X} = D_{1}^{\tau}. \end{cases}$$
(3.67)

Applying $\operatorname{div}_{\mathcal{B}}\partial_1$ to $(3.1)_1$ and then use $(3.1)_3$ yields

$$\partial_t \mathrm{div}_{\mathcal{B}} \partial_1 \eta - \kappa \partial_1 \mathrm{div}_{\mathcal{B}} \partial_1^2 \eta = \mathrm{div}_{\mathcal{B}_t} \partial_1 \eta - \mathrm{div}_{\partial_1 \mathcal{B}} u - \kappa \mathrm{div}_{\partial_1 \mathcal{B}} \partial_1^2 \eta =: K^4.$$

We further derive from the above identity that

$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|\partial^{\alpha} \mathcal{X} \mathrm{div}_{\mathcal{B}} \partial_{1} \eta\|_{0}^{2} + \kappa \|\partial^{\alpha} \mathcal{X} \mathrm{div}_{\mathcal{B}} \partial_{1}^{2} \eta\|_{2}^{2}
= \int \left(\partial^{\alpha} \mathcal{X} K^{4} \partial^{\alpha} \mathcal{X} \mathrm{div}_{\mathcal{B}} \partial_{1} \eta - \kappa \partial^{\alpha} \mathcal{X} \mathrm{div}_{\mathcal{B}} (\partial_{1}^{2} \eta \partial^{\alpha} \mathcal{X} \mathrm{div}_{\partial_{1} \mathcal{B}} \partial_{1} \eta) \right) \mathrm{d}y.$$
(3.68)

Following the arguments of (3.65) and (3.66) by further using (3.45) and (3.47), we can derive from (3.68) that

$$\begin{aligned} \|\operatorname{div}\eta(t)\|_{1,2}^{2} + \kappa \int_{0}^{t} \|\operatorname{div}\eta(s)\|_{2,2}^{2} \mathrm{d}s \\ \lesssim \|\eta^{0}\|_{1,3} + (\|\eta^{0}\|_{3} + TP(\|(\varsigma,\partial_{1}\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}) \sup_{t\in\overline{I_{T}}} \|(u,\partial_{1}\eta)(t)\|_{3}^{2} \\ + \kappa \left(\|\eta^{0}\|_{3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\right) \int_{0}^{t} \|\eta(s)\|_{2,3}^{2} \mathrm{d}s \\ + \kappa P(\|\varsigma\|_{L_{T}^{\infty}H^{1,3}}) \int_{0}^{t} \|\eta(s)\|_{1,3} \|\eta(s)\|_{2,3} \mathrm{d}s \end{aligned}$$
(3.69)

and

$$\begin{aligned} \|\operatorname{div}\partial_{1}\eta(t)\|_{*,2}^{2} + \kappa \int_{0}^{t} \|\operatorname{div}\partial_{1}^{2}\eta(s)\|_{*,2}^{2} \mathrm{d}s \\ \lesssim \|\partial_{1}\eta^{0}\|_{\underline{1},3} + (\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma,\partial_{1}\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}}) \sup_{t\in\overline{I_{T}}} \|(u,\partial_{1}\eta)(t)\|_{*,3}^{2} \\ + \kappa \left(\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}})\right) \int_{0}^{t} \|\partial_{1}^{2}\eta(s)\|_{*,3}^{2} \mathrm{d}s \\ + \kappa P(\|\varsigma\|_{L_{T}^{\infty}H^{2,3}}) \int_{0}^{t} \|\partial_{1}\eta(s)\|_{*,3} \|\partial_{1}^{2}\eta(s)\|_{*,3} \mathrm{d}s. \end{aligned}$$
(3.70)

Consequently, we immediately deduce from (3.65)-(3.67), (3.69) and (3.70) that

$$\begin{aligned} \|(\operatorname{div} u, \operatorname{curl} u, \operatorname{div} \partial_{1} \eta, \operatorname{curl} \partial_{1} \eta)(t)\|_{2}^{2} + \kappa \int_{0}^{t} \|(\operatorname{div} \eta, \operatorname{curl} \eta)(s)\|_{2,2}^{2} \mathrm{d}s \\ &\lesssim I_{L}^{0} + (\|\eta^{0}\|_{3} + TP(\|(\varsigma, \partial_{1}\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{3}}) \sup_{t \in \overline{I_{T}}} \|(u, \partial_{1}\eta)(t)\|_{3}^{2} \\ &+ \kappa \left(\|\eta^{0}\|_{3} + TP(\|(\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{3}})\right) \int_{0}^{t} \|\eta(s)\|_{2,3}^{2} \mathrm{d}s \\ &+ \kappa P(\|\varsigma\|_{L_{T}^{\infty}H^{1,3}}) \int_{0}^{t} \|\eta(s)\|_{1,3} \|\eta(s)\|_{2,3} \mathrm{d}s \end{aligned}$$
(3.71)

and

$$\begin{aligned} \|(\operatorname{div} u, \operatorname{curl} u, \operatorname{div} \partial_{1} \eta, \operatorname{curl} \partial_{1} \eta)(t)\|_{*,2}^{2} + \kappa \int_{0}^{t} \|(\operatorname{div} \partial_{1}^{2} \eta, \operatorname{curl} \partial_{1}^{2} \eta)(s)\|_{*,2}^{2} \mathrm{d}s \\ &\lesssim I_{H}^{0} + (\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma, \partial_{1}\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}}) \sup_{t \in \overline{I_{T}}} \|(u, \partial_{1}\eta)(t)\|_{*,3}^{2} \\ &+ \kappa \left(\|\eta^{0}\|_{\underline{1},3} + TP(\|(\varsigma, \varsigma_{t})\|_{L_{T}^{\infty}H^{1,3}})\right) \int_{0}^{t} \|\partial_{1}^{2}\eta(s)\|_{*,3}^{2} \mathrm{d}s \\ &+ \kappa P(\|\varsigma\|_{L_{T}^{\infty}H^{2,3}}) \int_{0}^{t} \|\partial_{1}\eta(s)\|_{*,3} \|\partial_{1}^{2}\eta(s)\|_{*,3} \mathrm{d}s. \end{aligned}$$
(3.72)

(6) Summing up the estimates of (u, η, q) .

Thanks to the estimates (3.44), (3.54), (3.71), (3.72), the Hodge-type elliptic estimate (A.4) and Young's inequality, we have, for sufficiently small δ ,

$$\sup_{t\in\overline{I_T}} \mathcal{E}^L_{\kappa}(t) + \kappa \|\partial_1^2 \eta^{\kappa}\|_{L^2_T H^3}^2$$

$$\leq c_5 I^0_L + c_5 TP(\|(\varsigma, \partial_1\varsigma, \varsigma_t)\|_{L^\infty_T H^3}) \left(\sup_{t\in\overline{I_T}} \mathcal{E}^L_{\kappa}(t) + \kappa \|\partial_1^2 \eta^{\kappa}\|_{L^2_T H^3}^2\right)$$
(3.73)

and

$$\sup_{t\in\overline{I_T}} \mathcal{E}^{\tau}_{\kappa}(t) + \kappa \int_0^t \|\partial_1^2 \eta^{\kappa}(s)\|_{*,3}^2 \mathrm{d}s$$

$$\leqslant c_5 I_H^0 + c_5 TP(\|(\varsigma,\partial_1\varsigma,\varsigma_t)\|_{L_T^\infty H^{1,3}}) \bigg(\sup_{t\in\overline{I_T}} \mathcal{E}^{\tau}_{\kappa}(t) + \kappa \int_0^t \|\partial_1^2 \eta^{\kappa}(s)\|_{*,3}^2 \mathrm{d}s \bigg).$$
(3.74)

Obviously, we can choose a new polynomial, still denoted by $P(\|(\varsigma, \partial_1\varsigma, \varsigma_t)\|_{L^{\infty}_{\alpha}H^{1,3}})$, which is not less than the two polynomials in (3.73) and (3.74). Now we use c_5 and the new polynomial P to define T_1 by (3.41). Taking $T = T_1$ in (3.73) and (3.74), thus we get (3.38) and

$$\sup_{t\in\overline{I_T}}\mathcal{E}^{\tau}_{\kappa}(t) + \kappa \int_0^t \|\partial_1^2\eta^{\kappa}(s)\|_{*,3}^2 \mathrm{d}s \leqslant 4c_3^2 I_H^0$$

which, together with (A.7), yields that

$$\operatorname{ess\,sup}_{t \in I_T} \mathcal{E}^H_{\kappa}(t) + \kappa \|\partial_1^2 \eta^{\kappa}\|_{L^2_T H^{1,3}}^2 \leqslant 4c_3^2 I_H^0.$$
(3.75)

We further derive from $(3.11)_1$, $(3.11)_2$, (3.75) and the fact $(u_3, \partial_1^2 \eta_3, \mathcal{B}_{3j} \partial_j q)|_{\partial\Omega} = 0$ that

$$\eta \in C^0(\overline{I_T}, H^{1,3}) \text{ and } u_t \in C^0(\overline{I_T}, H_s^2).$$

$$(3.76)$$

By (3.38) and (3.49), we get

$$\|q\|_{C^{0}(\overline{I_{T}},H^{3})}^{2} \lesssim P\left(I_{H}^{0},\|(\varsigma,\varsigma_{t})\|_{L_{T}^{\infty}H^{3}}^{2}\right).$$
(3.77)

Applying (3.33) to (3.48), and then using (3.75), we have

$$\|\nabla_{\mathcal{B}}q\|_{L^{\infty}_{T}H^{3}} \lesssim P\left(\|\varsigma\|_{L^{\infty}_{T}H^{3}}\right) \|K^{1}\|_{L^{\infty}_{T}H^{2}} \lesssim P\left(I^{0}_{H}, \|(\varsigma,\varsigma_{t})\|^{2}_{L^{\infty}_{T}H^{3}}\right).$$
(3.78)

In addition, exploiting (3.75) and (3.78), we deduce from $(3.1)_1$ and $(3.1)_2$ that

$$\|(u_t, \eta_t)\|_{L^{\infty}_T H^3} \lesssim P\left(I^0_H, \|(\varsigma, \varsigma_t)\|^2_{L^{\infty}_T H^3}\right).$$
(3.79)

Putting the estimates (3.77)-(3.79) together yields (3.40) immediately.

(7) Estimates of uniform continuity of $\partial_1 \nabla^3(u^{\kappa}, \partial_1 \eta^{\kappa})$ in H^{-1} .

Let the multiindex β satisfy $|\beta| = 3$. Similarly to (3.26), we easily deduce from $(3.1)_1$ and $(3.1)_2$ that, for any $\varphi \in H^1$ and for any $t, r \in I_T$,

$$\int \partial^{\beta} D_{1}^{\tau} \partial_{1} (\eta(y,t) - \eta(y,r)) \cdot \varphi dy = -\int_{r}^{t} \int \partial^{\beta} D_{1}^{\tau} (u + \kappa \partial_{1}^{2} \eta) \cdot \partial_{1} \varphi dy ds, \qquad (3.80)$$
$$\int \partial^{\beta} D_{1}^{\tau} (u(y,t) - u(y,r)) \cdot \varphi dy$$
$$= \int_{r}^{t} \int (\partial^{\beta} \nabla_{\mathcal{B}} q \cdot D_{1}^{-\tau} \varphi - m \partial^{\beta} D_{1}^{\tau} \partial_{1} \eta \cdot \partial_{1} \varphi - a \partial^{\beta} D_{1}^{\tau} u \cdot \varphi) dy ds. \qquad (3.81)$$

Exploiting the uniform estimates (3.75) and (3.78), we easily deduce that

$$D_1^{\tau} \nabla^3(u, \partial_1 \eta)$$
 is uniformly continuous in H^{-1} . (3.82)

In addition, it is easy to check that

$$\|D_1^{\tau}\nabla^3(u,\partial_1\eta)\|_{L^{\infty}_T L^2} \lesssim \|\partial_1\nabla^3(u,\partial_1\eta)\|_{L^{\infty}_T L^2}.$$
(3.83)

Making use of (3.82), (3.83), (A.8) and the regularity $(u, \partial_1 \eta) \in C^0(\overline{I_T}, H^3)$, we arrive at

$$\partial_1 \nabla^3(u, \partial_1 \eta) \in C^0(\overline{I_T}, L^2_{\text{weak}}), \tag{3.84}$$

which, together with (3.75), yields (3.39). Noting that (3.80) and (3.81) also hold with ∂_1 in place of D_1^{τ} , thus we further derive the assertion in (3.42).

Thanks to (3.76), (3.79), (3.84) and the fact $(u, \eta, q) \in \mathbb{S}_{T_1}$, we easily see that

$$(u,\eta,q) \in \mathfrak{U}_{T_1}^{1,3} \times \mathfrak{C}^0(\overline{I_{T_1}}, H_{\mathrm{s}}^{2,3}) \times C^0(\overline{I_{T_1}}, \underline{H}^3).$$

This completes the proof.

Thanks to Lemma 3.2, we easily get a unique local solution to the linearized problem (3.2) by a compactness argument and a regularity argument.

Proposition 3.5. Let the assumptions of Lemma 3.2 be satisfied, and δ_0 , T_1 be provided by Lemma 3.2. Then, for any $\delta \leq \delta_0$, the linearized problem (3.2) defined on Ω_{T_1} admits a unique solution $(u^L, \eta^L, q^L) \in \mathfrak{U}_{T_1}^{1,3} \times \mathfrak{C}^0(\overline{I_{T_1}}, H_s^{2,3}) \times C^0(\overline{I_{T_1}}, \underline{H}^3)$; moreover the solution satisfies

$$\|\eta^L\|_{C^0(\overline{I_{T_1}},H^3)} \leqslant \|\eta^0\|_3 + 2c_3\sqrt{I_L^0}T, \tag{3.85}$$

$$\|(u^{L}, \eta^{L}, \partial_{1}\eta^{L})\|_{C^{0}(\overline{I_{T_{1}}}, H^{3})} \leq 2c_{3}\sqrt{I_{L}^{0}},$$
(3.86)

$$\sup_{t\in\overline{I_{T_1}}} \|(u^L,\partial_1\eta^L)\|_{\underline{1},3} \leqslant 2c_3\sqrt{I_H^0},\tag{3.87}$$

$$\|q^{L}\|_{C^{0}(\overline{I_{T_{1}}},H^{3})} + \|(u_{t}^{L},\nabla_{\mathcal{B}}q^{L})\|_{L^{\infty}_{T_{1}}H^{3}} \leqslant P\left(I^{0}_{H},\|(\varsigma,\varsigma_{t})\|_{L^{\infty}_{T_{1}}H^{3}}\right).$$
(3.88)

PROOF. Let $\kappa \in (0,1)$ and $(u^{\kappa}, \eta^{\kappa}, q^{\kappa}) \in \mathbb{S}_{T_1}$ be the local solution stated as in Lemma 3.2. Thanks to the κ -independent estimates (3.38)–(3.40) and (3.42) satisfied by $(u^{\kappa}, \eta^{\kappa}, q^{\kappa})$, we can easily follow the compactness argument as in the proof of Proposition 3.1 to obtain a limit function (u^L, η^L, q^L) , which is the solution of the linearized problem (3.2) and satisfies

• the estimates

$$\|(u^L, \eta^L, \partial_1 \eta^L)\|_{L^{\infty}_{T_1}H^3} \leqslant 2c_3 \sqrt{I^0_L},$$
(3.89)

$$\|(u^{L},\partial_{1}\eta^{L})\|_{L^{\infty}_{T_{1}}H^{1,3}} \leq 2c_{3}\sqrt{I^{0}_{H}},\tag{3.90}$$

$$\|(u_t^L, q, \nabla_{\mathcal{B}} q^L)\|_{L^{\infty}_{T_1} H^3} \leqslant P\left(I^0_H, \|(\varsigma, \varsigma_t)\|_{L^{\infty}_{T_1} H^3}\right).$$
(3.91)

• the additional regularity

$$u^{L} \in C^{0}(I_{T_{1}}H_{s}^{2}), \ \nabla^{3}\partial_{1}u \in C^{0}(\overline{I_{T}}, L_{weak}^{2}), \ \eta^{L} \in \mathfrak{C}^{0}(\overline{I_{T_{1}}}, H_{s}^{2,3}).$$
 (3.92)

In addition,

$$(q)_{\Omega} = 0$$
 and $u_t|_{\partial\Omega} = 0$ for a.e. $t \in I_T$.

Thanks to Proportion 3.2 and the regularity of (u, η, q) , we further have

$$u^{L} \in C^{0}(I_{T_{1}}H_{s}^{3}). \tag{3.93}$$

By $(3.2)_2$ and $(3.2)_3$, q satisfies

$$\begin{cases} \Delta_{\mathcal{B}}q = K^4 := m \mathrm{div}_{\mathcal{B}} \partial_1^2 \eta + \mathrm{div}_{\mathcal{B}_t} u & \text{in } \Omega, \\ \mathcal{B}_{3j} \partial_i q = 0 & \text{on } \partial\Omega, \end{cases}$$

where $K^4 \in C^0(I_{T_1}H^1)$. Applying Proposition 3.3 to the above boundary-value problem, we easily obtain

$$q \in C^0(\overline{I_{T_1}}, \underline{H}^3). \tag{3.94}$$

Thus we further derive from $(3.2)_2$ that

$$u_t \in C^0(\overline{I_{T_1}}, H_s^2).$$
 (3.95)

Collecting the regularity of (u, η, q) in (3.90)–(3.95), we easily see that $(u^L, \eta^L, q^L) \in \mathfrak{U}_{T_1}^{1,3} \times \mathfrak{C}^0(\overline{I_{T_1}}, H_s^{2,3}) \times C(\overline{I_{T_1}}, \underline{H}^3)$. Thus we further get (3.86)–(3.88) from (3.89)–(3.91).

The estimate (3.85) is obvious by $(3.2)_1$ satisfied by (η^L, u^L) and the estimate (3.86). In addition, the uniqueness can be easily verified by a standard energy method. The proof of Proposition 3.5 is complete.

3.3. Proof of Proposition 2.3

Now we are in the position to the proof of Proposition 2.3. Let (u^0, η^0) satisfy the assumptions in Proposition 2.3, $\|\eta^0\|_{\underline{1},3} \leq \delta \leq \{\delta_0, \iota/2\}$ and

$$T_2 = \min\{1/3c_3P(4c_3(b+1), \delta/2c_3(b+1), 1)\},\$$

where the positive constants $c_3 \ge 1$ and δ_0 are provided by Proposition 3.5 with $\alpha = T_2$. By Proposition 3.5 and Lemma A.4, for any $T \le T_2$, we can construct a solution sequence

$$\{(u^n,\eta^n,Q^n)\}_{n=1}^{\infty} \subset \mathfrak{U}_T^{1,3} \times \mathfrak{C}^0(\overline{I_T},H_{\mathrm{s}}^{2,3}) \times C^0(\overline{I_{T_1}},\underline{H}^3)$$

such that

(1)
$$(u^1, \eta^1, q^1) = (\eta^0, u^0, 0);$$

(2) $(u^{n+1}, \eta^{n+1}, q^{n+1})$ satisfies

$$\begin{cases} \eta_t^{n+1} = u^{n+1}, \\ u_t^{n+1} + \nabla_{\mathcal{A}^n} q^{n+1} + a u^{n+1} = m \partial_1^2 \eta^{n+1}, \\ \operatorname{div}_{\mathcal{A}^n} u^{n+1} = 0, \\ (u^{n+1}, \eta^{n+1})|_{t=0} = (u^0, \eta^0), \\ (u_3^{n+1}, \eta_3^{n+1})|_{\partial\Omega} = \mathcal{A}_{3j}^n \partial_j q^{n+1}|_{\partial\Omega} = 0 \end{cases}$$

$$(3.96)$$

for $n \ge 1$, where $\mathcal{A}^n = (\nabla(\eta^n + I))^{-\mathrm{T}}$.

(3) (u^n, η^n, q^n) enjoys the estimates, for $n \ge 1$,

$$\|\eta^n\|_{C^0(\overline{I_{T_1}},H^3)}^2 \leqslant 2\delta \leqslant \iota \leqslant 1,\tag{3.97}$$

$$\|(u^{n},\eta^{n},\partial_{1}\eta^{n})\|_{C^{0}(\overline{I_{T_{1}}},H^{3})} \leqslant 2c_{3}\sqrt{I_{L}^{0}} \lesssim b+1,$$
(3.98)

$$\sup_{t\in\overline{I_T}} \|(u^n,\partial_1\eta^n)(t)\|_{\underline{1},3} \leqslant 2c_3\sqrt{I_H^0} \lesssim b+1,$$
(3.99)

$$\|q^n\|_{C^0(\overline{I_T},H^3)} + \|(u^n_t,\nabla_{\mathcal{A}^{n-1}}q^n)\|_{L^{\infty}_TH^3} \lesssim P(b).$$
(3.100)

Similarly to (3.7) and (3.9), by using (3.97) and (3.98), we have

$$\sup_{t \in I_T} \|(\mathcal{A}^n_t, \mathcal{A}^n, \partial_1 \mathcal{A}^n, 1/J^n)\|_2 \lesssim P(b),$$
(3.101)

$$\sup_{t \in I_T} \|\mathcal{A}^n - I\|_2 \lesssim \|\eta^n\|_3. \tag{3.102}$$

Similarly to (3.80) and (3.81), by a regularity method, we can easily verify that, for any $\varphi \in H^1$ and for any $t, r \in \overline{I_T}, (\eta^{n+1}, u^{n+1})$ satisfies

$$\int \partial^{\beta} \partial_{1}^{2} (\eta^{n+1}(t) - \eta^{n+1}(r)) \cdot \varphi dy = -\int_{r}^{t} \int \partial^{\beta} \partial_{1} u^{n+1} \cdot \partial_{1} \varphi dy ds,$$
$$\int \partial^{\beta} \partial_{1} (u^{n+1}(t) - u^{n+1}(r)) \cdot \varphi dy$$
$$= \int_{r}^{t} \int (\partial^{\beta} \nabla_{\mathcal{A}^{n}} q^{n+1} - m \partial^{\beta} \partial_{1}^{2} \eta^{n+1} + a \partial^{\beta} u^{n+1}) \cdot \partial_{1} \varphi dy ds.$$

Thus we further derive from the above two identities that, for $n \ge 1$,

$$\partial_1 \nabla^3 (u^n, \partial_1 \eta^n)$$
 are uniformly continuous in H^{-1} . (3.103)

Next we shall prove that $\{(\eta^n, u^n, q^n)\}_{m=1}^{\infty}$ is a Cauchy sequence. From now on, we always assume $n \ge 2$. We define that

$$\bar{\eta}^{n+1} := \eta^{n+1} - \eta^n, \quad \bar{u}^{n+1} := u^{n+1} - u^n, \quad \bar{q}^{n+1} := q^{n+1} - q^n \text{ and } \bar{\mathcal{A}}^n := \mathcal{A}^n - \mathcal{A}^{n-1}.$$

Then it follows from (3.96) that

$$\begin{cases} \bar{\eta}_{t}^{n+1} = \bar{u}^{n+1}, \\ \bar{u}_{t}^{n+1} + \nabla_{\mathcal{A}^{n}} \bar{q}^{n+1} + a \bar{u}^{n+1} = m \partial_{1}^{2} \bar{\eta}^{n+1} - \nabla_{\bar{\mathcal{A}}^{n}} q^{n}, \\ \operatorname{div}_{\mathcal{A}^{n}} \bar{u}^{n+1} = -\operatorname{div}_{\bar{\mathcal{A}}^{n}} u^{n}, \\ (\bar{u}^{n+1}, \bar{\eta}^{n+1})|_{t=0} = (0, 0), \\ (\bar{u}_{3}^{n+1}, \bar{\eta}_{3}^{n+1})|_{\partial\Omega} = 0. \end{cases}$$

$$(3.104)$$

By $(3.104)_1$ and $(3.104)_4$, we have

$$\|\bar{\eta}^{n+1}(t)\|_{C^0(\overline{I_T},H^3)} \leqslant \int_0^t \|\bar{u}^{n+1}(s)\|_3 \mathrm{d}s \leqslant T \|\bar{u}^{n+1}\|_{L^\infty_T H^3}.$$
(3.105)

Similarly to (3.101), we can estimate that

 $\sup_{t\in I_T} \|\bar{\mathcal{A}}^n\|_2 \lesssim \|\bar{\eta}^n\|_{L^{\infty}_T H^3},$ (3.106)

$$\sup_{t \in I_T} \|\bar{\mathcal{A}}^n\|_{1,2} \lesssim \|\bar{\eta}^n\|_{L^{\infty}_T H^{1,3}}, \tag{3.107}$$

 $\sup_{t\in I_{\mathcal{T}}} \|\bar{\mathcal{A}}^n_t\|_2 \lesssim P(b) \|(\bar{u}^n, \bar{\eta}^n)\|_{L^{\infty}_{\mathcal{T}}H^3}.$ (3.108) Thanks to $(3.104)_2$, \bar{q}^{n+1} satisfies

$$\begin{cases} \Delta_{\mathcal{A}^n} \bar{q}^{n+1} = \operatorname{div}_{\mathcal{A}^n} \left(m \partial_1^2 \bar{\eta}^{n+1} - \nabla_{\bar{\mathcal{A}}^n} q^n - \bar{u}_t^{n+1} - a \bar{u}^{n+1} \right) & \text{in } \Omega, \\ \nabla_{\mathcal{A}^n} \bar{q}^{n+1} \cdot \vec{\mathbf{n}} = -\nabla_{\bar{\mathcal{A}}^n} q^n \cdot \vec{\mathbf{n}} & \text{on } \partial\Omega, \end{cases}$$
(3.109)

where $\Delta_{\mathcal{A}^n} := \operatorname{div}_{\mathcal{A}^n} \nabla_{\mathcal{A}^n}$. Let $\zeta^n = \eta^n + y$. $(\zeta^n)^{-1}$ denotes the inverse function of ζ^n with respect to the variable y. We define that

$$\begin{split} K^5 &:= -\nabla_{\bar{\mathcal{A}}^n} q^n, \ K^6 &:= m\partial_1^2 \bar{\eta}^{n+1} - \bar{u}_t^{n+1} - a\bar{u}^{n+1} - \nabla_{\bar{\mathcal{A}}^n} q^n, \\ (\beta, \tilde{K}^5, \tilde{K}^6) &:= (\bar{q}^{n+1}, K^5, K^6)|_{y = (\zeta^n)^{-1}(x,t)}, \end{split}$$

then, by (3.109),

$$\begin{cases} \Delta\beta = \operatorname{div} \tilde{K}^6 & \text{in } \Omega, \\ \nabla\beta \cdot \vec{\mathbf{n}} = \tilde{K}^5 \cdot \vec{\mathbf{n}} & \text{on } \partial\Omega. \end{cases}$$

Applying the elliptic estimate (A.16) to the above boundary-value problem and then making use of (A.12) and (A.13), we have, for a.e. $t \in I_T$,

$$\|\bar{q}^{n+1}\|_{3} \lesssim P(b) \|\beta\|_{3} \lesssim P(b) \left(\|\operatorname{div} \tilde{K}^{6}\|_{1} + \|\tilde{K}^{5}\|_{2} \right) \\ \lesssim P(b) \left(\|\operatorname{div}_{\mathcal{A}^{n}} K^{6}\|_{1} + \|K^{5}\|_{2} \right).$$
(3.110)

Noting that

$$\operatorname{div}_{\mathcal{A}^n} K^6 = \operatorname{div}_{\mathcal{A}^n} \left(m \partial_1^2 \bar{\eta}^{n+1} - \nabla_{\bar{\mathcal{A}}^n} q^n - a \bar{u}^{n+1} \right) + \operatorname{div}_{\mathcal{A}^n_t} \bar{u}^{n+1} + \partial_t \operatorname{div}_{\bar{\mathcal{A}}^n} u^n,$$

thus, making use of (3.100), (3.101), (3.106) and (3.108), we easily estimate that

$$\|\operatorname{div}_{\mathcal{A}^n} K^6\|_1 + \|K^5\|_2 \lesssim P(b)\| \left(\bar{u}^n, \bar{u}^{n+1}, \bar{\eta}^n, \partial_1 \bar{\eta}^{n+1}\right)\|_{L^{\infty}_T H^3}.$$

Putting the above estimate into (3.110) yields

$$\|\bar{q}^{n+1}\|_{C^0(\overline{I_T},H^3)} \lesssim P(b) \| \left(\bar{u}^n, \bar{u}^{n+1}, \bar{\eta}^n, \partial_1 \bar{\eta}^{n+1}\right) \|_{L^{\infty}_T H^3}.$$
(3.111)

Similarly to the derivation of (3.54), we have that

$$\|(\bar{u}^{n+1},\partial_1\bar{\eta}^{n+1})(t)\|_{C^0(\overline{I_T},L^2)} \lesssim P(b)T\| \left(\bar{u}^n,\bar{u}^{n+1},\bar{\eta}^n,\partial_1\bar{\eta}^{n+1}\right)\|_{L^{\infty}_T H^3}.$$
(3.112)

By $(3.104)_1 - (3.104)_2$, we have

$$\begin{aligned} \partial_t \operatorname{div}_{\mathcal{A}^n} \partial_1 \bar{\eta}^{n+1} &= \operatorname{div}_{\mathcal{A}^n_t} \partial_1 \bar{\eta}^{n+1} - \partial_1 \operatorname{div}_{\bar{\mathcal{A}}^n} u^n - \operatorname{div}_{\partial_1 \mathcal{A}^n} \bar{u}^{n+1}, \\ \partial_t \operatorname{curl}_{\mathcal{A}^n} \bar{u}^{n+1} + a \operatorname{curl}_{\mathcal{A}^n} \bar{u}^{n+1} - m \partial_1 \operatorname{curl}_{\mathcal{A}^n} \partial_1 \bar{\eta}^{n+1} \\ &= \operatorname{curl}_{\mathcal{A}^n_t} \bar{u}^{n+1} - m \operatorname{curl}_{\partial_1 \mathcal{A}^n} \partial_1 \bar{\eta}^{n+1} + \operatorname{curl}_{\bar{\mathcal{A}}^n} (\nabla_{\mathcal{A}^{n-1}} q^n). \end{aligned}$$

Following the argument of (3.71) by further making use of (3.99), (3.101), (3.102), (3.106) and (3.107), we derive can from $(3.104)_3$ and the above identities that

$$\| (\operatorname{div}\bar{u}^{n+1}, \operatorname{curl}\bar{u}^{n+1}, \operatorname{div}\partial_{1}\bar{\eta}^{n+1}, \operatorname{curl}\partial_{1}\bar{\eta}^{n+1}) \|_{C^{0}(\overline{I_{T}}, H^{3})} \lesssim P(b)T \| \left(\bar{u}^{n+1}, \bar{\eta}^{n}, \partial_{1}\bar{\eta}^{n}, \partial_{1}\bar{\eta}^{n+1} \right) \|_{L_{T}^{\infty}H^{3}} + \| \eta^{0} \|_{3} \| \left(\bar{u}^{n+1}, \partial_{1}\bar{\eta}^{n+1} \right) \|_{L_{T}^{\infty}H^{3}},$$
(3.113)

where we have used (3.106) in the last inequality of (3.113).

Summing up the estimates (3.105), (3.112) and (3.113) and then using Hodge-type elliptic estimate (A.4), we obtain, for sufficiently small δ ,

$$\|\left(\bar{u}^{n+1},\bar{\eta}^{n+1},\partial_{1}\bar{\eta}^{n+1}\right)\|_{C^{0}(\overline{I_{T}},H^{3})} \lesssim P(b)T\|\left(\bar{u}^{n},\bar{u}^{n+1},\bar{\eta}^{n},\partial_{1}\bar{\eta}^{n},\partial_{1}\bar{\eta}^{n+1}\right)\|_{L^{\infty}_{T}H^{3}}.$$

In particular

$$\|\left(\bar{u}^{n+1}, \bar{\eta}^{n+1}, \partial_1 \bar{\eta}^{n+1}\right)\|_{C^0(\overline{I_T}, H^3)} \leqslant \|\left(\bar{u}^n, \bar{\eta}^n, \partial_1 \bar{\eta}^n\right)\|_{L^{\infty}_T H^3}/2$$
(3.114)

for sufficiently small $T \in (0, T_2]$, where the smallness depends on b, a, m and Ω .

In addition, by $(3.104)_2$ and (3.101), we get that

$$\|\bar{u}_t^{n+1}\|_{C^0(\overline{I_T},H^2)} \lesssim P(b) \| \left(\bar{u}^{n+1}, \bar{\eta}^n, \partial_1 \bar{\eta}^{n+1}, \bar{q}^{n+1} \right) \|_{L^{\infty}_T H^3}.$$
(3.115)

We immediately see from (3.111), (3.114) and (3.115) that the sequence $\{(u^n, \eta^n, q^n, u_t^n)\}_{n=1}^{\infty}$ is a Cauchy sequence in $C^0(\overline{I_T}, H_s^3 \times H_s^{1,3} \times \underline{H}^3 \times H_s^2)$. Thus

$$(u^n, \eta^n, q^n, u^n_t) \to (u, \eta, q, u_t)$$
 strongly in $C^0(\overline{I_T}, H^3_s \times H^{1,3}_s \times \underline{H}^3 \times H^2_s)$.

Thanks to the strong convergence of $(u^n, \eta^n, q^n, u_t^n)$, we can pass to limits in (3.96), and then get a solution (u, η, q, u_t) to the problem (1.11).

Finally, thanks to (3.98)–(3.100) and (3.103), we easily follow the compactness argument and the regularity argument in Proposition 3.5 to further derive that

$$(u,\eta,q) \in \mathfrak{U}_{T}^{1,3} \times \mathfrak{C}^{0}(\overline{I_{T}}, H^{2,3}) \times C^{0}(\overline{I_{T}}, \underline{H}^{3}), \ \nabla_{\mathcal{A}}q \in L_{T}^{\infty}H^{3},$$
$$\|(u,\eta,\partial_{1}\eta)\|_{C^{0}(\overline{I_{T_{1}}}, H^{3})} \leqslant 2c_{3}\sqrt{I_{L}^{0}},$$
(3.116)

$$\sup_{t\in\overline{I_T}} \|(u,\partial_1\eta)\|_{\underline{1},3} \leqslant 2c_3\sqrt{I_H^0}.$$
(3.117)

Finally it is easy to check that (η, u, q) is the unique solution to the problem (1.11), provided that δ is sufficiently small. This completes the proof of Proposition 2.3.

Appendix A. Analytic tools

This appendix is devoted to providing some mathematical results, which have been used in previous sections. It should be noted that Ω appearing in what follows is still defined by (1.2) with h > 0 and we will also use the simplified notations appearing in Section 1.2. In addition, $a \leq b$ still denotes $a \leq cb$, however the positive constant c depends on the parameters and domain in the lemma, in which c appears.

Lemma A.1. (1) Embedding inequality (see [2, 4.12 Theorem]): Let $D \subset \mathbb{R}^2$ be a domain satisfying the cone condition, then

$$\|f\|_{C^0(\overline{D})} = \|f\|_{L^\infty(D)} \lesssim \|f\|_{H^2(D)}.$$
(A.1)

(2) Product estimates (see Section 4.1 in [17]): Let $D \in \mathbb{R}^2$ be a domain satisfying the cone condition, and the functions f, g are defined in D. Then

$$\|fg\|_{H^{i}(D)} \lesssim \begin{cases} \|f\|_{H^{1}(D)} \|g\|_{H^{1}(D)} & \text{for } i = 0; \\ \|f\|_{H^{i}(D)} \|g\|_{H^{2}(D)} & \text{for } 0 \leq i \leq 2. \end{cases}$$
(A.2)

Lemma A.2. Poincaré inequality (see [24, Lemma 1.43]): Let $1 \leq p < \infty$, and D be a bounded Lipchitz domain in \mathbb{R}^n for $n \geq 2$ or a finite interval in \mathbb{R} . Then for any $w \in W^{1,p}(D)$,

$$\|w\|_{L^p(D)} \lesssim \|\nabla w\|_{L^p(D)}^p + \left|\int_D w \mathrm{d}y\right|^p$$

Remark A.1. By Poincaré inequality, we have, for any given $i \ge 0$,

$$\|w\|_{1,i} \lesssim \|w\|_{2,i} \text{ for any } w \text{ satisfying } \partial_1 w, \ \partial_1^2 w \in H^i.$$
(A.3)

Lemma A.3. Hodge-type elliptic estimate: Let $r \ge 1$, then it holds that

$$||u||_{r} \lesssim ||u||_{0} + ||(\operatorname{curl} u, \operatorname{div} u)||_{r-1} + ||u_{3}|_{\partial\Omega}||_{H^{r-1/2}(\partial\Omega)}$$
(A.4)

PROOF. Please refer to [6, Proposition 5.1] for the proof.

Lemma A.4. Diffeomorphism mapping theorem: There exists a sufficiently small constant $\iota \in (0,1]$, depending on Ω , such that, for any $\varsigma \in H^3_s$ satisfying $\|\varsigma\|_3 \leq \iota, \psi := \varsigma + y$ satisfies $\inf_{y \in \Omega} \det(\nabla \varsigma + I) \ge 1/4$ and the diffeomorphism properties

 $\psi|_{y_3=i}: \mathbb{R} \to \mathbb{R} \text{ is a } C^1(\mathbb{R})\text{-diffeomorphism mapping for } i=0, 1,$ $\psi:=\varsigma+y:\overline{\Omega}\to\overline{\Omega} \text{ is a } C^1\text{-diffeomorphism mapping.}$

PROOF. Please refer to [20, Lemma A.10] for the proof.

Lemma A.5. Some results for functions with values in Banach spaces: Let T > 0, integers i, $j \ge 1$ be given and $1 \le p \le \infty$.

- (1) Assume $f \in L^p_T H^i$, $\partial^k_1 \nabla^i f \in L^p_T L^2$ for any $1 \leq k \leq j$, then $f \in L^p_T H^{j,i}$.
- (2) Let X be a separable Banach space and T > 0. If $w \in C^0(\overline{I_T}, X)$, then $w: I_T \to X$ is a strongly measurable function and

$$\|w\|_X \in C^0(\overline{I_T}). \tag{A.5}$$

(3) Let $j \ge i+1$ and c be a constant. If $f \in L^p(I_T, H^i)$ with $1 , <math>||f||_j \le c ||g||_j$ holds for a.e. $t \in I_T$ and $g \in L^p(I_T, H^j)$, then

$$f \in L^p_T H^i. \tag{A.6}$$

- (4) Assume $f \in L^p_T H^i$ and $\{D^{\tau}_1 f\}_{|\tau| \in (0,1)}$ is uniformly bounded in $L^p_T H^i$.
 - (a) There exists a sequence (still denoted by $D_1^{\tau}f$) of $\{D_1^{\tau}f\}_{|\tau|\in(0,1)}$ such that, if $\tau \to 0$, then

$$D_1^{\tau} f \rightharpoonup \partial_1 f \begin{cases} weakly \text{-}^* \text{ in } L_T^{\infty} H^i & \text{for } p = \infty, \\ weakly \text{ in } L_T^p H^i & \text{for } 1 (A.7)$$

Moreover, $f \in L^{\infty}_{T} H^{1,i}$.

(b) Let $p = \infty$. If additionally $\partial^{\alpha} f \in C^{0}(\overline{I_{T}}, L^{2}_{weak})$ and $D^{\tau}_{1} \partial^{\alpha} f$ is uniformly continuous in H^{-1} , where the multiindex α satisfying $|\alpha| = i$, then

$$D_1^{\tau} \partial^{\alpha} f \to \partial^{\alpha} \partial_1 f \text{ in } C^0(\overline{I_T}, L_{\text{weak}}^2) \text{ (some sequence).}$$
 (A.8)

(c) We additionally assume $f \in L^p_T H^i$ and $\{D_1^{-\tau} D_1^{\tau} f\}_{|\tau| \in (0,1)}$ is uniformly bounded in $L^p_T H^i$ with 1 , then (some sequence)

$$D_1^{-\tau}(D_1^{\tau}f) \rightharpoonup \partial_1^2 f \text{ weakly in } L_T^p H^i \text{ (some sequence)}.$$
 (A.9)

(5) Let $\varphi = \varsigma + y$ and $\varsigma \in \mathbb{A}_T^{3,1/4}$ defined by (3.5). (a) If $C^0(\overline{I_T}, H^i)$ or $f \in L^2_T H^i$ with $0 \leq i \leq 3$, then

$$F := f(\varphi, t) \in C^0(\overline{I_T}, H^i) \text{ or } L^p_T H^i$$
(A.10)

and

$$\mathcal{F} := f(\varphi^{-1}, t) \in C^0(\overline{I_T}, H^i) \text{ or } L^p_T H^i.$$
(A.11)

Moreover,

$$\|F\|_{L^{p}_{T}H^{i}} \lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}}) \|f\|_{L^{p}_{T}H^{i}}, \tag{A.12}$$

$$\|\mathcal{F}\|_{L^{p}_{T}H^{i}} \lesssim P(\|\varsigma\|_{L^{\infty}_{T}H^{3}}) \|f\|_{L^{p}_{T}H^{i}}.$$
(A.13)

(b) If ς additionally satisfies $\varsigma_t \in L_T^{\infty} H^2$, then for any f satisfying $f \in L^p(I_T, H^i)$ and $f_t \in L^p(I_T, H^{i-1})$, where $1 \leq i \leq 3$, then

$$F_t = (f_t(x,t) + \varsigma_t \cdot \nabla f(y,t))|_{x=\varphi} \in L^p_T H^{i-1}$$

and

$$\mathcal{F}_t = (f_t(y,t) - (\nabla \varphi)^{-1}\varsigma_t \cdot \nabla f(y,t))|_{y=\varphi^{-1}} \in L^p_T H^{i-1}$$

PROOF. Please refer to [20, Lemma A.10] for the proof.

Lemma A.6. Elliptic estimates for Neumann boundary-value problem: Let $i \ge 2$ be an integer. We further assume A is a 3×3 matrix function, each element of which belongs to $W^{i-1,\infty}(\Omega)$, and satisfies positivity condition, i.e., there exists a positive constant θ such that $(A\xi) \cdot \xi \ge \theta |\xi|$ for a.e. $y \in \Omega$ and all $\xi \in \mathbb{R}^3$. If $f_1 \in H^{i-2}$ and $f_2 \in H^{i-1}$ satisfy the following compatibility condition

$$\int f_1 dy + \int_{\partial \Omega} f_2 \vec{\mathbf{n}}_3 dy_h = 0, \qquad (A.14)$$

where $\vec{\mathbf{n}}_3$ denotes the outward unit normal vector $\vec{\mathbf{n}}$ on the boundary $\partial \Omega$, there exists a unique strong solution $p \in \underline{H}^j$ to the Neumann boundary-value problem of elliptic equations:

$$\begin{cases} -\operatorname{div} \left(A\nabla p\right) = f_1 & \text{in } \Omega, \\ A_{3k}\partial_k p = f_2 & \text{on } \partial\Omega. \end{cases}$$
(A.15)

Moreover *p* satisfies

$$\|p\|_{i} \lesssim (1 + \|A\|_{W^{i-1,\infty}(\Omega)})(\|f_{1}\|_{i-2} + \|f_{2}\|_{i-1}).$$
(A.16)

PROOF. Please refer to [20, Lemma A.7] for the proof.

Acknowledgements. The research of Fei Jiang was supported by NSFC (Grant Nos. 12022102) and the Natural Science Foundation of Fujian Province of China (2020J02013), and the research of Song Jiang by National Key R&D Program (2020YFA0712200), National Key Project (GJXM92579), and NSFC (Grant No. 11631008), the Sino-German Science Center (Grant No. GZ 1465) and the ISF-NSFC joint research program (Grant No. 11761141008).

References

- H. Abidi, P. Zhang, On the global solution of a 3-D MHD system with initial data near equilibrium, Comm. Pure Appl. Math. 70 (2017) 1509–1561.
- [2] R.A. Adams, J.J.F. Fournier, Sobolev Space, Academic Press: New York, 2005.
- [3] C. Bardos, C. Sulem, P. Sulem, Longtime dynamics of a conductive fluid in the presence of a strong magnetic field, Trans. Am. Math. Soc. 305 (1988) 175–191.
- [4] Y. Cai, Z. Lei, Global well-posedness of the incompressible magnetohydrodynamics, Arch. Ration. Mech. Anal. 228 (2018) 969–993.
- [5] J.Y. Chemin, D.S. McCormick, J.C. Robinson, J.L. Rodrigo, Local existence for the non-resistive mhd equations in besov spaces, Adv. Math. 286 (2016) 1–31.
- [6] D. Coutand, S. Shkoller, Well-posedness of the free-surface incompressible Euler equations with or without surface tension, J. Amer. Math. Soc. 20 (2007) 829–930.
- [7] D. Coutand, S. Shkoller, A simple proof of well-posedness for the free-surface incompressible Euler equations, Discrete Contin. Dyn. Syst., Ser. S 3 (2010) 429–449.
- [8] Y. Du, W. Yang, Y. Zhou, On the exponential stability of a stratified flow to the 2D ideal MHD equations with damping, SIAM J. Math. Anal. 51 (2019) 5077–5102.
- [9] C.L. Fefferman, D.S. McCormick, J.C. Robinson, J.L. Rodrigo, Higher order commutator estimates and local existence for the non-resistive MHD equations and related models, J. Funct. Anal. 267 (2014) 1035–1056.
- [10] C.L. Fefferman, D.S. McCormick, J.C. Robinson, J.L. Rodrigo, Local existence for the non-resistive mhd equations in nearly optimal sobolev spaces, Arch. Ration. Mech. Anal. 223 (2017) 677–691.
- [11] X.M. Gu, Y.J. Wang, On the construction of solutions to the free-surface incompressible ideal magnetohydrody-namic equations, J. Math. Pures Appl. 128 (2019) 1–41.
- [12] Y. Guo, I. Tice, Almost exponential decay of periodic viscous surface waves without surface tension, Arch. Ration. Mech. Anal. 207 (2013) 459–531.
- [13] L.B. He, L. Xu, P. Yu, On global dynamics of three dimensional magnetohydrodynamics: nonlinear stability of Alfvén waves, Anal. PDE 4 (2018) Paper No. 5, 105 pp.
- [14] F.M. Huang, R.H. Pan, Convergence rate for compressible Euler equations with damping and vacuum, Arch. Ration. Mech. Anal. 166 (2003) 359–376.
- [15] F. Jiang, S. Jiang, On the stabilizing effect of the magnetic fields in the magnetic Rayleigh-Taylor problem, SIAM J. Math. Anal. 50 (2018) 491–540.
- [16] F. Jiang, S. Jiang, On the dynamical stability and instability of Parker problem, Physica D 391 (2019) 17–51.
- [17] F. Jiang, S. Jiang, Nonlinear stability and instability in Rayleight–Taylor problem of stratisfied compressible MHD fluids, Calc. Var. Partial Differential Equations 58 (1) (2019) Art. 29, 61 pp.
- [18] F. Jiang, S. Jiang, On magnetic inhibition theory in non-resistive magnetohydrodynamic fluids, Arch. Ration. Mech. Anal. 233 (2019) 749–798.
- [19] F. Jiang, S. Jiang, On the inhibition of thermal convection by a magnetic field under zero resistivity, J. Math. Pures Appl. 141 (2020) 220–265.
- [20] F. Jiang, S. Jiang, Y.Y. Zhao, On inhibition of Rayleigh–Taylor instability by horizontal magnetic field in an inviscid MHD fluid with velocity damping, Submmited (2021).
- [21] X.L. Li, N. Su, D.H. Wang, Local strong solution to the compressible magnetohydrodynamic flow with large data, J. Hyperbolic Differ. Equ. 8 (2011) 415–436.
- [22] Y. Li, Y.Z. Sun, Global weak solutions to a two-dimensional compressible mhd equations of viscous non-resistive fluids, J. Differential Equations. 267 (2019) 3827–3851.
- [23] F.H. Lin, L. Xu, P. Zhang, Global small solutions of 2-D incompressible MHD system, J Differ. Equations 259 (2015) 5440–5485.
- [24] A. Novotnỳ, I. Straškraba, Introduction to the Mathematical Theory of Compressible Flow, Oxford University Press, USA, 2004.

- [25] R.H. Pan, K. Zhao, The 3D compressible Euler equations with damping in a bounded domain, J. Differential Equations 246 (2009) 581–596.
- [26] X.X. Ren, Z.Y. Xiang, Z.F. Zhang, Global well-posedness for the 2d mhd equations without magnetic diffusion in a strip domain, Nonlinearity 29 (2016) 1257.
- [27] T.C. Sideris, B. Thomases, D.H. Wang, Long Time Behavior of Solutions to the 3D Compressible Euler Equations with Damping, Comm. Partial Differential Equations 28 (2003) 795–816.
- [28] Z. Tan, Y. Wang, Global solution and large-time behavior of the 3D compressible Euler equations with damping, J. Differential Equations 254 (2013) 1686–1704.
- [29] Z. Tan, Y.J. Wang, Global well-posedness of an initial-boundary value problem for viscous nonresistive MHD systems, SIAM J. Math. Anal. 50 (2018) 1432–1470.
- [30] W.K. Wang, T. Yang, The pointwise estimates of solutions for Euler equations with damping in multi-dimensions, J. Differential Equations 173 (2001) 410–450.
- [31] Y.J. Wang, Sharp nonlinear stability criterion of viscous non-resistive MHD internal waves in 3D, Arch. Ration. Mech. Anal. 231 (2019) 1675–1743.
- [32] J.H. Wu, Y.F. Wu, Global small solutions to the compressible 2D magnetohydrodynamic system without magnetic diffusion, Adv. Math. 310 (2017) 759–888.
- [33] J.H. Wu, Y.F. Wu, X.J. Xu, Global small solution to the 2D MHD system with a velocity damping term, SIAM J. Math. Anal. 47 (2013) 2630–2656.
- [34] L. Xu, P. Zhang, Global small solutions to three-dimensional incompressible magnetohydrodynamical system, SIAM J. Math. Anal. 47 (2015) 26–65.
- [35] T. Zhang, An elementary proof of the global existence and uniqueness theorem to 2-d incompressible non-resistive mhd system, arXiv preprint arXiv:1404.5681 (2014).
- [36] T. Zhang, Global solutions to the 2D viscous, non-resistive MHD system with large background magnetic field, J. Differential Equations 260 (2016) 5450–5480.