Asymptotic completeness of wave operators for Schrödinger operators with time-periodic magnetic fields

by Masaki Kawamoto.

Graduate School of Science and Engineering, Ehime University, 3 Bunkyo-cho Matsuyama, Ehime 790-8577. Japan. Email: kawamoto.masaki.zs@ehime-u.ac.jp,

Abstract Under the effect of suitable time-periodic magnetic fields, the velocity of a charged particle grows exponentially in t; this phenomenon provides the asymptotic completeness for wave operators with slowly decaying potentials. These facts were shown under some restrictions for time-periodic magnetic fields and the range of wave operators. In this study, we relax these restrictions and finally obtain the asymptotic completeness without any restrictions on the range of wave operators. Additionally, we show them under generalized conditions, which are truly optimal for time-periodic magnetic fields. Moreover, we provide a uniform resolvent estimate for the perturbed Floquet Hamiltonian.

Keywords: quantum scattering theory; time-periodic magnetic fields; Floquet Hamiltonian; Time-periodic systems

MSC classification 2020: Primary 81U05; Second 35P25, 47A55.

1 Introduction

We study a scattering problem for a charged particle under the effect of a time-periodic magnetic field. In this study, we assume that the charged particle moves on a plane \mathbf{R}^2 in the presence of a time-periodic magnetic field $\mathbf{B}(t) = (0, 0, B(t))$ with B(t + T) = B(t), which is always perpendicular to the plane. Subsequently, the free Hamiltonian for this system is given by

$$H_0(t) = (p - qA(t, x))^2 / (2m), \quad A(t, x) = (-B(t)x_2, B(t)x_1)/2,$$

where $x = (x_1, x_2) \in \mathbf{R}^2$, $p = (p_1, p_2) = -i(\partial_1, \partial_2)$, m > 0 and $q \in \mathbf{R} \setminus \{0\}$ are position, momentum, mass, and charge of a particle, respectively. $B(t) \in L^{\infty}(\mathbf{R})$ is the intensity of the magnetic field at time t. The wave function $\psi(t, x)$ described by this system satisfies the following time-dependent Schrödinger equation;

$$\begin{cases} i\partial_t \psi(t,x) &= H_0(t)\psi(t,x), \\ \psi(0,x) &= \psi_0 \end{cases}$$

By defining a propagator for $H_0(t)$ as $U_0(t, s)$, the wave function $\psi(t, x)$ is denoted by $\psi(t, x) = U_0(t, 0)\psi_0$, which we refer to as family of unitary operators $\{U_0(t, s)\}_{(t,s)\in\mathbf{R}^2}$ a propagator for $H_0(t)$ if each component satisfies

$$i\partial_t U_0(t,s) = H_0(t)U_0(t,s), \quad i\partial_s U_0(t,s) = -U_0(t,s)H_0(s)$$
$$U_0(t,\theta)U_0(\theta,s) = U_0(t,s), \quad U_0(s,s) = \mathrm{Id}_{L^2(\mathbf{R}^2)}.$$

Under these settings, the classical trajectory x(t) and p(t) of this system can be expressed in the form

$$x(t) := U_0(0,t)xU_0(t,0), \quad p(t) := U_0(0,t)pU_0(t,0).$$

Let us define $L := x_1 p_2 - x_2 p_1$, $\omega(t) = qB(t)/m$, and $\Omega(t) = \int_0^t \omega(s) ds$. It can be noted that $\tilde{U}_0(t,0) := e^{-i\Omega(t)L}U_0(t,0)$ is a propagator for $\tilde{H}_0(t) := p^2/(2m) + q^2B^2(t)x^2/(8m)$ because L commutes with p^2 and x^2 . By defining $\tilde{x}(t)$ and $\tilde{p}(t)$ as $\tilde{U}_0(0,t)x\tilde{U}_0(t,0)$ and $\tilde{U}_0(0,t)p\tilde{U}_0(t,0)$, respectively, the straightforward calculation shows that

$$\tilde{x}'(t) = \tilde{U}_0(0,t)i[\tilde{H}_0(t),x]\tilde{U}_0(t,0) = \tilde{p}(t)/m,$$

$$\tilde{p}'(t)/m = \tilde{U}_0(0,t)i[\tilde{H}_0(t),p/m]\tilde{U}_0(t,0) = -(qB(t)/(2m))^2\tilde{x}(t).$$

hold on $\mathscr{S}(\mathbf{R}^2)$, where $[\cdot, \cdot]$ denotes the commutator of operators, and these equations yield *Hill's equation*

$$\tilde{x}''(t) + \left(\frac{qB(t)}{2m}\right)^2 \tilde{x}(t) = 0, \quad \begin{cases} \tilde{x}(0) = 0, \\ \tilde{x}'(0) = \tilde{p}(0)/m. \end{cases}$$

and a differential equation

$$\tilde{p}(t) = m\tilde{x}'(t),$$

refer to Kawamoto [7] §3. Hence, by introducing the fundamental solutions of the Hill's equation, $\zeta_1(t)$ and $\zeta_2(t)$ as

$$\zeta_j''(t) + \left(\frac{qB(t)}{2m}\right)^2 \zeta_j(t) = 0, \quad \begin{cases} \zeta_1(0) = 1, \\ \zeta_1'(0) = 0, \end{cases} \quad \begin{cases} \zeta_2(0) = 0, \\ \zeta_2'(0) = 1, \end{cases}$$
(1)

we obtain

$$\begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix} = \begin{pmatrix} \zeta_1(t) & \zeta_2(t)/m \\ m\zeta_1'(t) & \zeta_2'(t) \end{pmatrix} \begin{pmatrix} x \\ p \end{pmatrix}.$$

Noting

$$e^{-i\Omega(t)L} \begin{pmatrix} x \\ p \end{pmatrix} e^{i\Omega(t)L} = \begin{pmatrix} \hat{R}(\Omega(t)/2)x \\ \hat{R}(\Omega(t)/2)p \end{pmatrix}, \quad \hat{R}(t) = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix},$$

(refer to Adachi-Kawamoto [1] or [7]), it can be deduced that

$$\begin{pmatrix} x(t) \\ p(t) \end{pmatrix} = \begin{pmatrix} \zeta_1(t) & \zeta_2(t)/m \\ m\zeta_1'(t) & \zeta_2'(t) \end{pmatrix} \begin{pmatrix} \hat{R}(\Omega(t)/2)x \\ \hat{R}(\Omega(t)/2)p \end{pmatrix}$$

holds. Here we let

$$\mathcal{L} := \begin{pmatrix} \zeta_1(T) & \zeta_2(T)/m \\ m\zeta_1'(T) & \zeta_2'(T) \end{pmatrix}.$$

Then, for t = NT, $N \in \mathbb{Z}$, we have

$$\begin{pmatrix} x(NT) \\ p(NT) \end{pmatrix} = \mathcal{L}^N \begin{pmatrix} \hat{R}(\Omega(NT)/2)x \\ \hat{R}(\Omega(NT)/2)p \end{pmatrix}.$$
 (2)

Using $\zeta_1(t)\zeta'_2(t) - \zeta'_1(t)\zeta_2(t) = 1$ for all $t \in \mathbf{R}$, we have

$$\det(\mathcal{L} - \lambda) = \lambda^2 - \mathcal{D}\lambda + 1,$$

where $\mathcal{D} = \zeta_1(T) + \zeta'_2(T)$ is referred to as the *discriminant of the Hill's equation*, and together with (2), we find the following lemma:

Lemma 1.1. Let $N \in \mathbb{Z}$, $|N| \gg 1$. Then, for all $\phi \in C_0^{\infty}(\mathbb{R}^2)$, there exist constants λ_e , $C_e, C_p, C_m > 0$ such that

$$(C_e)^{-1} e^{\lambda_e N} \leq \|x U_0(NT, 0)\phi\|_{(L^2(\mathbf{R}^2))^2} \leq C_e e^{\lambda_e N}, \quad \text{if } \mathcal{D}^2 > 4, (C_p)^{-1} N \leq \|x U_0(NT, 0)\phi\|_{(L^2(\mathbf{R}^2))^2} \leq C_p N, \quad \text{if } \mathcal{D}^2 = 4, \|x U_0(NT, 0)\phi\|_{(L^2(\mathbf{R}^2))^2} \leq C_m, \quad \text{if } \mathcal{D}^2 < 4,$$

hold.

This lemma implies that for $\mathcal{D}^2 = 4$, the particle assumes a uniform linear motion; however, for $\mathcal{D}^2 > 4$, the asymptotic velocity of the particle grows exponentially in t. Such physical phenomena were reported by Korotyaev [10] and the scattering theory for the case where $\mathcal{D}^2 = 4$ with dimensions n = 3 and $\mathcal{D}^2 > 4$ with dimension n = 2, 3 have been considered. Conversely, [7] found a relationship between the repulsive Hamiltonian $p^2 - x^2$ and $\mathcal{D}^2 > 4$; Schrödinger operator p^2 and $\mathcal{D}^2 = 4$, as defined by the Floquet operator.

In this study, we focus on the case $\mathcal{D}^2 > 4$ and prove the asymptotic completeness of the wave operators. The scattering theory for a time-periodic magnetic field was first considered by [10], who proved the asymptotic completeness under some technical conditions of the magnetic field; thereafter, Adachi-Kawamoto [1] proved the asymptotic completeness of wave operators for the case where the magnetic field is pulsed. In the study of [1], they discovered an explicit formula for the integral kernel of free propagators, which indicated that $\mathcal{D}^2 > 4$ and $\zeta_2(T) \neq 0$ is the best possible condition for showing asymptotic completeness in the pulsed case; if $\zeta_2(T) = 0$, the absolute value of the integral kernel of $U_0(nT, 0)\phi$, $n \in \mathbb{Z}$, diverges infinity for any $\phi \in L^2(\mathbb{R}^2)$. Hence, it still remains important to determine the asymptotic completeness for general time-periodic magnetic fields with only two conditions $\mathcal{D}^2 > 4$ and $\zeta_2(T) \neq 0$; this factor is considered herein.

Assumption 1.2. Suppose that $\zeta_1(t), \zeta'_1(t), \zeta_2(t)$, and $\zeta'_2(t)$ are continuous functions on $t \in [0, T)$, and that for all $t \in [0, T)$ and $N \in \mathbb{Z}$ there exist $A_{1,N}, A_{2,N}, A_{3,N}, A_{4,N}$ such that the solutions to (1) satisfy

$$\begin{pmatrix} \zeta_1(t+NT) \\ \zeta_2(t+NT) \end{pmatrix} = \begin{pmatrix} A_{1,N} & A_{2,N} \\ A_{3,N} & A_{4,N} \end{pmatrix} \begin{pmatrix} \zeta_1(t) \\ \zeta_2(t) \end{pmatrix}$$

Moreover, for some $\lambda > 0$, $\tilde{\lambda} \leq \lambda$, and for $|N| \gg 1$, there exist $0 < c_3 < C_3$ and $0 < c_4 < C_4$ such that

$$c_3 e^{\lambda N} \le |A_{3,N}| \le C_3 e^{\lambda N}, \quad c_4 e^{\lambda N} \le |A_{4,N}| \le C_4 e^{\lambda N}.$$

holds.

Remark 1.3. Owing to Lemma 8 in [7], this assumption will be true for $\mathcal{D}^2 = (\zeta_1(T) + \zeta'_2(T))^2 > 4$ and $\zeta_2(T) \neq 0$, and according to Kargel-Korotyaev [5], the model of B(t) such that $\zeta_2(T) \neq 0$ is known if B(t) is even. To simplify the proofs, there is need to handle this assumption. In this case, where $\mathcal{D}^2 > 4$ and $\zeta_2(T) \neq 0$, it is possible that $A_{4,N} \sim e^{-\lambda N}$ as $|N| \gg 1$ (refer also to [10]). The assumption on $A_{4,N}$ is stated to admit such cases.

We assume the following on the potential V;

Assumption 1.4. V is a multiplication operator of V(x), and $V(x) = \rho_1(x)\rho_2(x)$, where $\rho_1(x) = |V(x)|^{1/2}$ and $\rho_2(x) = \operatorname{sign}(V)|V|^{1/2}$ satisfies the following: V is in $C(\mathbf{R}^2)$ and is bounded. Moreover, there exists p > 4 such that $\rho_1, \rho_2 \in L^p(\mathbf{R}^2)$.

Remark 1.5. To simplify the proof, we used this assumption in this study. In the case where $|V(x)| \leq C \langle x \rangle^{-\rho}$, $\rho > 0$, Assumption 1.4 demands $\rho > 2/p$ and noting that we can consider p > 4 to be sufficiently large, this assumption allows any small $\rho > 0$. In this sense, we can consider the scattering theory even if V decays slowly in x.

Here, let us define U(t,0) as a propagator for $H(t) = H_0(t) + V$, and the unique existence of propagator U(t,0) under this assumption is guaranteed by Yajima [13].

Under these settings, we can define the Floquet Hamiltonian associated with H_0 and H. Let $\mathscr{K} := L^2([0,T]; L^2(\mathbf{R}^2))$ and define the self-adjoint operators acting on \mathscr{K} as

$$\hat{H} = \hat{H}_0 + V, \quad \hat{H}_0 = -i\partial_t + H_0$$

and term \hat{H} and \hat{H}_0 as the Floquet Hamiltonian associated with $H_0(t)$ and $H(t) = H_0(t) + V$, respectively. Under these assumptions, we can prove the compactness of $V(\hat{H}-i)^{-1}$ on \mathscr{K} (refer to, e.g., [7] (refer also to [10] and [1])). However, our scheme does not demand this property because we do not employ energy cut-off $\varphi(\hat{H}), \varphi \in C_0^{\infty}(\mathbf{R})$ at all in this study.

As a theorem of this study, we first obtain the following Theorem:

Theorem 1.6. Under assumptions 1.2 and 1.4, $\sigma_{\text{sing}}(\hat{H}) = \sigma_{\text{pp}}(\hat{H}) = \emptyset$, where $\sigma_{\text{sing}}(A)$ denotes the set of the singular continuous spectrum of A, and $\sigma_{\text{pp}}(A)$ denotes the set of the pure point spectrum of A.

Owing to the correspondence of the spectrum sets between \hat{H} and the monodromy operator U(T,0), refer to, for example, Proposition 3.3 of Møller [11], we also have the following corollary:

Corollary 1.7. Under Assumptions 1.2 and 1.4, $\sigma_{\text{sing}}(U(T,0)) = \sigma_{\text{pp}}(U(T,0)) = \emptyset$.

Remark 1.8. Recently, [7] showed the absence of singular continuous spectra of U(T, 0) using the Mourre theory. However, in this study, $V \in C^2(\mathbf{R}^2)$ is required, and hence, our result is successful in relaxing this condition. Moreover, as the author's knowledge, the absence of embedded eigenvalues for U(T, 0) is stated as an open problem.

Under these assumptions, we can obtain the existence and completeness of wave operators: **Theorem 1.9.** Under the assumption 1.2 and the assumption 1.4, the wave operators

$$W^{\pm} = s - \lim_{t \to \pm \infty} U(t, 0)^* U_0(t, 0)$$
(3)

exist and complete, that is,

$$\operatorname{Ran}\left(W^{\pm}\right) = L^2(\mathbf{R}^2)$$

holds.

Moreover, because \hat{H} has only an absolutely continuous spectrum, the uniform resolvent estimate for \hat{H} can be considered. As a result, the following uniform resolvent estimate holds:

Theorem 1.10. Under assumptions 1.2 and 1.4, for all $\phi \in \mathscr{K}$, there exists C > 0 such that

$$\sup_{\lambda \in \mathbf{R}, \mu > 0} \left\| |V|^{1/2} (\hat{H} - \lambda \mp i\mu)^{-1} |V|^{1/2} \phi \right\|_{\mathscr{H}} \le C \|\phi\|_{\mathscr{H}}$$

holds.

Remark 1.11. For the case where V has singularities, the approach proposed by [10] works well if $V \in L^{\tilde{p}}(\mathbb{R}^2)$ with $\tilde{p} > 2$. However, as in our approach in such a case, the additional term $[\rho_1, (\hat{H} + i)^{-1}]$ should be calculated. However, showing the boundedness of $[\rho_1, (\hat{H} + i)^{-1}]$ is difficult because of the operator $-i\partial_t$. Moreover, even if this difficulty can be handled, only the potential satisfying $V \in L^{\tilde{p}}(\mathbb{R}^2)$, that is, the potential decays slowly, and the singularity is also weak. In this sense, we do not discuss this issue.

The first approach of the proof is to show the uniform resolvent estimate (URE) for \hat{H}_0 , in which we imitate the approach in [10]. Owing to the URE and Kato's smooth perturbation method, Kato [6], we have

$$\int_{\mathbf{R}} \left\| |V|^{1/2} e^{-i\sigma \hat{H}_0} \phi \right\|_{\mathscr{H}}^2 d\sigma \le C \|\phi\|_{\mathscr{H}}^2.$$

$$\tag{4}$$

Here, Korotaev's strong propagation estimate (Proposition 2.1) enables us to extend this estimate to the estimate of \hat{H} , which yields the nonexistence of eigenvalues of \hat{H} . By employing Kato's smooth perturbation method in (4) with replacement $\hat{H}_0 \rightarrow \hat{H}$, we obtain a resolvent estimate for \hat{H} , and using this, we can prove the nonexistence of singular continuous spectra of \hat{H} . Moreover, two estimates (4) and (4) with replacement $\hat{H}_0 \rightarrow \hat{H}$ immediately prove the asymptotic completeness of the wave operators. For \hat{H} , we employ the approach of Herbst [3]. Here, we employ the strong propagation estimate (Proposition 2.1).

The key approach for characterizing the range of wave operators for time periodic systems is the *Howland-Yajima method*; if the wave operators W^{\pm} exist and that wave operators in the sense of the Floquet Hamiltonian

$$\hat{W}^{\pm} := \mathbf{s} - \lim_{\sigma \to \pm \infty} e^{i\sigma \hat{H}} e^{-i\sigma \hat{H}_0}$$

exist and satisfy $\operatorname{Ran}\left(\hat{W}^{\pm}\right) = \mathscr{K}_{\operatorname{ac}}(\hat{H})$, then the asymptotic completeness

$$\operatorname{Ran}\left(W^{\pm}\right) = L_{\operatorname{ac}}(U(T,0))$$

holds, where $\mathscr{K}_{ac}(\hat{H}) \subset \mathscr{K}$ is the space of the absolutely continuous spectrum of \hat{H} . Hence, to prove Theorem 1.9, we should show the existence of W^{\pm} , \hat{W}^{\pm} , and $\operatorname{Ran}\left(\hat{W}^{\pm}\right) = \mathscr{K}_{ac}(\hat{H})$. Here, we remark that Enss-Veserić [2] and Kitada-Yajima [9] show the equivalence

$$L^{2}(\mathbf{R}^{2}) = L_{\rm sc}(U(T,0)) \oplus L_{\rm ac}(U(T,0)) \oplus U_{\rm pp}(U(T,0)).$$

where $L_{\rm sc}(U(T,0))$, $L_{\rm ac}(U(T,0))$, and $U_{\rm pp}(U(T,0))$ denote the space of the singular continuous spectral, absolutely continuous spectral, and pure point spectra, respectively. Hence, to obtain Theorem 1.9, we additionally should show that $L_{\rm sc}(U(T,0)) \oplus U_{\rm pp}(U(T,0)) = \emptyset$, which can be fulfilled by Corollary 1.7.

2 Uniform resolvent estimates for H_0

In this section, we consider the URE for \hat{H}_0 under assumption 1.2. In the following, $\|\cdot\|_p$, $1 \leq p \leq \infty$ denotes $\|\cdot\|_{L^p(\mathbf{R}^2)}$. The estimation key for showing the URE is the following dispersive estimates for the free propagator $U_0(t, s)$:

Proposition 2.1. For all $\phi \in L^1(\mathbf{R}^2)$, the dispersive estimates

$$\left\|U_0(\tau,s)\phi\right\|_{\infty} \le \frac{C}{\left|\Gamma(\tau,s)\right|} \left\|\phi\right\|_1 \tag{5}$$

holds, where

$$\Gamma(\tau, s) = |\zeta_1(s)\zeta_2(\tau) - \zeta_1(\tau)\zeta_2(s)| \tag{6}$$

Moreover, for $\eta_1, \eta_2 \in L^{\tilde{p}}(\mathbf{R})$ with $2 \leq \tilde{p} < \infty$ and $\psi \in L^2(\mathbf{R}^2)$,

$$\|\eta_1 U_0(\tau, s)\eta_2 \psi\|_2 \le C |\Gamma(\tau, s)|^{-2/\tilde{p}} \|\eta_1\|_{\tilde{p}} \|\eta_2\|_{\tilde{p}} \|\psi\|_2.$$
(7)

Proof. This proposition was proven by [10] for a special case of magnetic fields. Inequality (5) can be shown using (35) in [7] (or refer also to (7.7) of [1] and Lemma 2.3 in Kawamoto [8]). Hence, we only show (7). Owing to the Riesz-Thorin interpolation theorem, we have for $2 \leq Q \leq \infty$ and $\phi \in L^{Q/(Q-1)}(\mathbf{R}^2)$

$$\|U_0(\tau, s)\phi\|_Q \le C |\Gamma(\tau, s)|^{-2(1/2 - 1/Q)} \|\phi\|_{(Q/(Q-1))}$$

holds. Hence, for $\psi \in L^2(\mathbf{R}^2)$ and $Q = 2\tilde{p}/(\tilde{p}-2)$,

$$\begin{aligned} \|\eta_{1}U_{0}(\tau,s)\eta_{2}\psi\|_{2} &\leq C \|\eta_{1}\|_{\tilde{p}} \|U_{0}(\tau,s)\eta_{2}\psi\|_{Q} \\ &\leq C|\Gamma(\tau,s)|^{-2/\tilde{p}} \|\eta_{1}\|_{\tilde{p}} \|\eta_{2}\psi\|_{2\tilde{p}/(\tilde{p}+2)} \\ &\leq C|\Gamma(\tau,s)|^{-2/\tilde{p}} \|\eta_{1}\|_{\tilde{p}} \|\eta_{2}\|_{\tilde{p}} \|\psi\|_{2}. \end{aligned}$$

$$(8)$$

Before we show the URE for \hat{H}_0 , we remark some properties for $\zeta_j(t)$, j = 1, 2. By the definition of $\zeta_j(t)$, one discovery

$$\zeta_1(t)\zeta_2'(t) - \zeta_1'(t)\zeta_2(t) = 1.$$
(9)

Owing to this equation, one sees that zero points for $\zeta_j(t)$ and $t \in [0, T)$ are unique. Indeed, if $t_0 \in [0, T)$ exists such that $\zeta_1(t_0) = 0$, then (9) yields $\zeta'_1(t_0)\zeta_2(t_0) = -1 \neq 0$, that is, $\zeta'_1(t_0) \neq 0$. The same is true for $\zeta_2(t)$. Now, we divide [0, T) into

$$[0,T) = \Omega_1^l \cup \Omega_2^l \cup \Omega_{J_l}^l \cup \{0, t_1^{(l)}, t_2^{(l)}, \dots, t_{J_l-1}^{(l)}\}, \quad J_l \in \mathbf{N}, \quad l = 1,2$$
(10)

where $\zeta_l(t_k^{(l)}) = 0$, $k \in \{1, 2, ..., J_l - 1\}$, and $\zeta_l(t) \neq 0$ for all $t \in \Omega_j^l$, $j \in \{1, ..., J_l\}$. Noting (9), we also notice that $\zeta_1(t)/\zeta_2(t)$ (resp. $\zeta_2(t)/\zeta_1(t)$) is a monotonically increasing function (resp. the monotone decreasing function) on Ω_j^2 (resp. Ω_j^1) and satisfies

$$\frac{d}{dt}\frac{\zeta_1(t)}{\zeta_2(t)} = \frac{1}{(\zeta_2(t))^2} > 0, \quad \left(\text{resp.} \quad \frac{d}{dt}\frac{\zeta_2(t)}{\zeta_1(t)} = -\frac{1}{(\zeta_1(t))^2} < 0\right)$$

Proposition 2.2. For all $\phi \in \mathscr{K}$, there exists a constant C > 0 such that

$$\sup_{\lambda \in \mathbf{R}, \, \mu > 0} \left\| \rho_1 (\hat{H}_0 - \lambda \mp i\mu)^{-1} \rho_2 \phi \right\|_{\mathscr{H}} \le C \|\phi\|_{\mathscr{H}}$$

Proof. The fundamental proof is based on [10]. Owing to the Laplace transform, we have:

$$(\rho_1(\hat{H}_0 - \lambda - i\mu)^{-1}\rho_2\phi)(t, x) = -i\rho_1(x)\int_0^\infty \left(e^{-i\sigma(\hat{H}_0 - \lambda - i\mu)}\rho_2\phi\right)(t, x)d\sigma.$$
(11)

Then, using the formula of $e^{-i\sigma \hat{H}_0}$, consider, for example, Yajima [12], we have

$$(\rho_1(\hat{H}_0 - \lambda - i\mu)^{-1}\rho_2\phi)(t, x) = i\rho_1(x)\sum_{N=1}^{\infty}\int_0^T e^{i(t+NT-s)(\lambda+i\mu)}U_0(t+NT, s)(\rho_2\phi)(s, x)ds + i\rho_1(x)\int_0^t e^{i(t-s)(\lambda+i\mu)}U_0(t, s)(\rho_2\phi)(s, x)ds.$$

Then, for p > 4, $\lambda \in \mathbf{R}$ and $\mu > 0$, using Proposition 2.1, we have

$$\begin{aligned} \|(\rho_{1}(\hat{H}_{0} - \lambda \mp i\mu)^{-1}\rho_{2}\phi)\|_{\mathscr{K}} \\ &\leq C \sum_{N=0}^{\infty} \left\| \int_{0}^{T} \|(\rho_{1}U_{0}(t + NT, s)\rho_{2}\phi(s))\|_{2}ds \right\|_{L^{2}([0,T])} \\ &\leq C \sum_{N=0}^{\infty} \|\rho_{1}\|_{p} \|\rho_{2}\|_{p} \left\| \int_{0}^{T} |\Gamma(t + NT, s)|^{-2/p} \|\phi(s)\|_{2}ds \right\|_{L^{2}([0,T])} \\ &\leq C \sum_{N=0}^{\infty} \|\rho_{1}\|_{p} \|\rho_{2}\|_{p} \|\phi\|_{\mathscr{K}} \left(\int_{[0,T]^{2}} |\Gamma(t + NT, s)|^{-4/p} ds dt \right)^{1/2}. \end{aligned}$$
(12)

By (6),

$$\int_0^T \int_0^T |\Gamma(t + NT, s)|^{-4/p} ds dt$$

= $\int_0^T |\zeta_2(t + NT)|^{-4/p} \int_0^T |\zeta_2(s)|^{-4/p} |(\zeta_1(s)/\zeta_2(s) - \alpha)|^{-4/p} ds dt$

holds, where $\alpha = \alpha(t) = \zeta_1(t + NT)/\zeta_2(t + NT)$. We decompose

$$\int_0^T |\zeta_2(s)|^{-4/p} |(\zeta_1(s)/\zeta_2(s) - \alpha)|^{-4/p} ds = \sum_{J=1}^{J_2} \int_{\Omega_J^2} |\zeta_2(s)|^{-4/p} |(\zeta_1(s)/\zeta_2(s) - \alpha)|^{-4/p} ds,$$

where J_2 and Ω_J^2 are defined by the same rules as in (10). Here, we remark that J_2 can be considered a finite integer. Then, by denoting $\tau = \zeta_1(s)/\zeta_2(s)$ and employing $d\tau/ds = (\zeta_2(s))^{-2}$, we have

$$\int_{\Omega_J^2} |\zeta_2(s)|^{-4/p} |(\zeta_1(s)/\zeta_2(s) - \alpha)|^{-4/p} ds \le I_1 + I_2 + I_3$$

with

$$I_{1} := \int_{|\tau-\alpha| \le 1} |\zeta_{2}(s)|^{2-4/p} |\tau-\alpha|^{-4/p} d\tau \le C \int_{|\tau-\alpha| \le 1} |\tau-\alpha|^{-4/p} d\tau \le C,$$

$$I_{2} := \int_{|\tau-\alpha| \ge 1, \ |\tau| \le 1} |\zeta_{2}(s)|^{2-4/p} |\tau-\alpha|^{-4/p} d\tau \le C \int_{|\tau| \le 1} d\tau \le C$$

and

$$I_3 := \int_{|\tau - \alpha| \ge 1, \ |\tau| \ge 1} |\zeta_2(s)|^{2-4/p} |\tau - \alpha|^{-4/p} d\tau$$

where we use p > 4. Furthermore, $|\zeta_2(s)|^{-1} \leq |\tau|^{-1} |\zeta_1(s)| \leq C |\tau|^{-1}$, we also have

$$I_3 \le C \int_{|\tau-\alpha|\ge 1, |\tau|\ge 1} |\tau|^{-2+4/p} |\tau-\alpha|^{-4/p} d\tau \le C$$

by p > 4. Combining all, one can obtain a constant C > 0, independent of t, such that

$$\sum_{J=1}^{J_2} \int_{\Omega_J^2} |\zeta_2(s)|^{-4/p} |(\zeta_1(s)/\zeta_2(s) - \alpha)|^{-4/p} ds \le C$$

Hence, one has

$$\|(\rho_1(\hat{H}_0 - \lambda \mp i\mu)^{-1}\rho_2\phi)\|_{\mathscr{H}} \le C \, \||V|^{1/2}\|_p^2 \|\phi\|_{\mathscr{H}} \sum_{N=0}^\infty \left(\int_0^T |\zeta_2(t+NT)|^{-4/p} dt\right)^{1/2}.$$
(13)

We now calculate the integral in t. By $\zeta_2(t + NT) = A_{3,N}\zeta_1(t) + A_{4,N}\zeta_2(t)$, we have

$$\int_{0}^{T} |\zeta_{2}(t+NT)|^{-4/p} dt \le |A_{3,N}|^{-4/p} \int_{0}^{T} |\zeta_{1}(t)|^{-4/p} |1+r_{0}(\zeta_{2}(t)/\zeta_{1}(t))|^{-4/p} dt$$
(14)

where $r_0 = A_{4,N}/A_{3,N}$. We again decompose

$$\int_0^T |\zeta_1(t)|^{-4/p} \left|1 + r_0(\zeta_2(t)/\zeta_1(t))\right|^{-4/p} dt = \sum_{J=1}^{J_1} \int_{\Omega_J^1} |\zeta_1(t)|^{-4/p} \left|1 + r_0(\zeta_2(t)/\zeta_1(t))\right|^{-4/p} dt,$$

where J_1 and Ω_J^1 are defined by the same rules as in (10). By denoting $\sigma = \zeta_2(t)/\zeta_1(t)$, we have $d\sigma/(dt) = -(\zeta_1(t))^{-2}$ and

$$\int_{\Omega_J^1} |\zeta_1(t)|^{-4/p} |1 + r_0(\zeta_2(t)/\zeta_1(t))|^{-4/p} dt \le I_4 + I_5 + I_6$$

with

$$I_4 := \int_{|1+r_0\sigma| \ge 1/2, |\sigma| \le 1} |\zeta_1(t)|^{2-4/p} |1+r_0\sigma|^{-4/p} d\sigma \le C,$$

$$I_5 := \int_{|1+r_0\sigma| \ge 1/2, |\sigma| \ge 1} |\sigma|^{-2+4/p} |1+r_0\sigma|^{-4/p} d\sigma \le C.$$

and

$$I_6 := \int_{|1+r_0\sigma| \le 1/2} |\zeta_1(t)|^{2-4/p} |1+r_0\sigma|^{-4/p} d\sigma.$$

Now, we estimate I_6 . Based on assumption 1.2, there are $\lambda > 0$ and $\tilde{\lambda} \leq \lambda$ such that

$$(c_4/C_3)e^{-(\lambda-\tilde{\lambda})N} \le |r_0| \le (C_4/c_3)e^{-(\lambda-\tilde{\lambda})N}.$$

holds. Then, it can be calculated that $|\sigma| \geq Ce^{(\lambda - \tilde{\lambda})N}$, that is, $|\sigma|^{-1} \leq Ce^{-(\lambda - \tilde{\lambda})N}$ on the support of $|1 + r_0\sigma| \leq 1/2$, which yields $|\zeta_1(t)| \leq |\sigma|^{-1}|\zeta_2(t)| \leq Ce^{-(\lambda - \tilde{\lambda})N}$. Thus, we also have

$$I_{6} \leq C e^{-N(\lambda - \tilde{\lambda})(2 - 4/p)} \int_{|1 + r_{0}\sigma| \leq 1/2} |1 + r_{0}\sigma|^{-4/p} d\sigma$$

$$\leq C e^{-N(\lambda - \tilde{\lambda})(2 - 4/p)} |r_{0}|^{-1}$$

$$\leq C e^{-N(\lambda - \tilde{\lambda})(1 - 4/p)}$$

$$\leq C.$$

Thus, we finally obtain that for some C > 0,

$$\int_0^T |\zeta_2(t+NT)|^{-4/p} dt \le C e^{-4\lambda N/p} \tag{15}$$

holds for p > 4. Finally, from (12), (13), and (15), we obtain

$$\left\| \rho_1 (\hat{H}_0 - \lambda \mp i\mu)^{-1} \rho_2 \phi \right\|_{\mathscr{K}} \le C \left\| |V|^{1/2} \right\|_p^2 \|\phi\|_{\mathscr{K}} \sum_{N=0}^{\infty} e^{-2\lambda N/p} \le C \|\phi\|_{\mathscr{K}}$$

It proves Proposition 2.2.

3 Non existence of embedded eigenvalues and singular continuous spectra

In this section, we show that \hat{H} has no embedded eigenvalues or singular spectra. From Proposition 2.2 and the smooth perturbation method according to [6], we have that for all $\phi \in \mathcal{K}$,

$$\int_0^\infty \left\| |V|^{1/2} e^{-i\sigma \hat{H}_0} \phi \right\|_{\mathscr{H}}^2 d\sigma \le C \left\| \phi \right\|_{\mathscr{H}}^2 \tag{16}$$

holds. The idea for proof is quite simple; we replace H_0 in the estimate (16) with H. Then, if H has eigenvalues, this estimate fails. Hence, we present the following theorem:

Theorem 3.1. Suppose Assumptions 1.2 and 1.4. Then, for all $\phi \in \mathscr{K}$,

$$\int_{0}^{\infty} \left\| |V|^{1/2} e^{-i\sigma \hat{H}} \phi \right\|_{\mathscr{H}}^{2} d\sigma \leq C \left\| \phi \right\|_{\mathscr{H}}^{2}$$

$$\tag{17}$$

holds.

Proof. By the Duhamel formula we have

$$|V|^{1/2}e^{-i\sigma\hat{H}}\phi = |V|^{1/2}e^{-i\sigma\hat{H}_0}\phi - i\int_0^\sigma |V|^{1/2}e^{-i(\sigma-s)\hat{H}_0}Ve^{-is\hat{H}}\phi ds$$

and which leads

$$\begin{split} \left\| |V|^{1/2} e^{-i\sigma \hat{H}} \phi \right\|_{\mathscr{H}} &\leq \left\| |V|^{1/2} e^{-i\sigma \hat{H}_0} \phi \right\|_{\mathscr{H}} \\ &+ C \int_0^{\sigma} \left\| |V|^{1/2} e^{-i(\sigma-s)\hat{H}_0} |V|^{1/2} \right\|_{\mathscr{B}(\mathscr{H})} \left\| |V|^{1/2} e^{-is\hat{H}} \phi \right\|_{\mathscr{H}} ds, \end{split}$$

where $\|\cdot\|_{\mathscr{B}(\mathscr{K})}$ denotes operator norm. Now, we have $F(\sigma, s) := \left\| |V|^{1/2} e^{-i(\sigma-s)\hat{H}_0} |V|^{1/2} \right\|_{\mathscr{B}(\mathscr{K})}$. Then, using the Gronwall inequality

$$\begin{split} \left\| |V|^{1/2} e^{-i\sigma \hat{H}} \phi \right\|_{\mathscr{K}} &\leq C \left\| |V|^{1/2} e^{-i\sigma \hat{H}_0} \phi \right\|_{\mathscr{K}} \\ &+ \int_0^\sigma \left\| |V|^{1/2} e^{-is \hat{H}_0} \phi \right\|_{\mathscr{K}} F(\sigma, s) \exp\left(\int_s^\sigma F(\sigma, \tau) d\tau\right) ds \end{split}$$

Here, we remark that

$$\int_{s}^{\sigma} F(\sigma,\tau) d\tau = \sup_{\|\psi\|_{\mathscr{K}}=1} \int_{0}^{\sigma-s} \left\| |V|^{1/2} e^{-i\gamma \hat{H}_{0}} |V|^{1/2} \psi \right\|_{\mathscr{K}} d\gamma$$

$$\leq \sup_{\|\psi\|_{\mathscr{K}}=1} \int_{0}^{\infty} \left\| |V|^{1/2} e^{-i\gamma \hat{H}_{0}} |V|^{1/2} \psi \right\|_{\mathscr{K}} d\gamma$$

$$\leq C, \qquad (18)$$

where we use (11) and (13) to obtain

$$\left\| |V|^{1/2} e^{-i\sigma \hat{H}} \phi \right\|_{\mathscr{X}}^{2} \leq C \left\| |V|^{1/2} e^{-i\sigma \hat{H}_{0}} \phi \right\|_{\mathscr{X}}^{2} + C \left(\int_{0}^{\sigma} \left\| |V|^{1/2} e^{-is \hat{H}_{0}} \phi \right\|_{\mathscr{X}} F(\sigma, s) ds \right)^{2}$$

and by the Cauchy-Swartz inequality we also have

$$\begin{split} & \left(\int_0^{\sigma} \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{H}} F(\sigma, s) ds \right)^2 \\ & \leq C \left(\int_0^{\sigma} \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{H}}^2 F(\sigma, s) ds \right) \left(\int_0^{\sigma} F(\sigma, s) ds \right) \\ & \leq C \int_0^{\sigma} \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{H}}^2 F(\sigma, s) ds. \end{split}$$

Moreover, by the Fubini theorem and (18) one obtains

$$\begin{split} &\int_0^\infty \int_0^\sigma \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{X}}^2 F(\sigma, s) ds d\sigma \\ &= \int_0^\infty \int_s^\infty \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{X}}^2 F(\sigma, s) d\sigma ds \\ &= \int_0^\infty \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{X}}^2 \left(\int_s^\infty F(\sigma, s) d\sigma \right) ds \\ &\leq C \int_0^\infty \left\| |V|^{1/2} e^{-is\hat{H}_0} \phi \right\|_{\mathscr{X}}^2 ds \\ &\leq C \|\phi\|_{\mathscr{X}}^2. \end{split}$$

Hence, together with (16), we obtain (17).

By the same approaches, we also have

$$\int_{-\infty}^{\infty} \left\| |V|^{1/2} e^{-i\sigma \hat{H}_0} \phi \right\|_{\mathscr{H}}^2 d\sigma \le C \left\| \phi \right\|_{\mathscr{H}}^2$$
(19)

and

$$\int_{-\infty}^{\infty} \left\| |V|^{1/2} e^{-i\sigma \hat{H}} \phi \right\|_{\mathscr{H}}^{2} d\sigma \leq C \left\| \phi \right\|_{\mathscr{H}}^{2}$$

$$\tag{20}$$

3.1 Proof of Theorem 1.6

Now, we prove Theorem 1.6. First, we show the nonexistence of eigenvalues. If there exists $\psi \in \mathscr{D}(\hat{H})$ such that $\hat{H}\psi = \lambda\psi$ with $\lambda \in \mathbf{R}$ and $\|\psi\|_{\mathscr{K}} \neq 0$. Then, using $e^{-i\sigma\hat{H}}\psi = e^{-i\sigma\lambda}\psi$ and (17), we obtain

$$\int_0^\infty \left\| |V|^{1/2} \psi \right\|_{\mathscr{H}}^2 d\sigma = \int_0^\infty \left\| |V|^{1/2} e^{-i\sigma \hat{H}} \psi \right\|_{\mathscr{H}}^2 d\sigma \le C \|\psi\|_{\mathscr{H}}^2.$$

However, this inequality is true only for the case $\|\psi\| \equiv 0$, which contradicts $\|\psi\| \neq 0$.

Next, we show the absence of a singular continuous spectrum of \hat{H} . In the case where $V \in C^2(\mathbb{R}^2)$, [7] showed this issue using the Mourre inequality. Hence, we relaxed this condition. From (20) and Kato's smooth perturbation theory, we have that

$$\sup_{\lambda \in \mathbf{R}, \varepsilon > 0} \left\| |V|^{1/2} \left((\hat{H} - \lambda - i\varepsilon)^{-1} - (\hat{H} - \lambda + i\varepsilon)^{-1} \right) |V|^{1/2} \phi, \right\|_{\mathscr{H}} \le C \left\| \phi \right\|_{\mathscr{H}}.$$

Here, we let (\cdot, \cdot) be an inner product of $L^2(\mathbf{R}^2)$. Then, for all $\lambda_1, \lambda_2 \in \mathbf{R}$,

$$\begin{split} &\left|\left((E_{\hat{H}}(\lambda_1) - E_{\hat{H}}(\lambda_2))|V|^{1/2}\phi, |V|^{1/2}\phi\right)\right|.\\ &= \left|\lim_{\varepsilon \to 0} \frac{1}{2\pi i} \int_{\lambda_2}^{\lambda_1} \left(|V|^{1/2} \left((\tau - \lambda - i\varepsilon)^{-1} - (\tau - \lambda + i\varepsilon)^{-1}\right)|V|^{1/2}\phi, \phi\right) d\tau \\ &\leq C|\lambda_2 - \lambda_1| \left\|\phi\right\|_{\mathscr{H}}^2 \end{split}$$

holds, where $E_{\hat{H}}(\cdot)$ is the spectral decomposition of \hat{H} , implying that for the Borel set $I := (\lambda_1, \lambda_2) \subset \mathbf{R}, \ E(I)|V|^{1/2}\phi \in \mathscr{K}_{\mathrm{ac}}(\hat{H})$. Here, $|V|^{1/2}$ is the closed operator, which yields for all $u \in \mathscr{K}, \ E(I)u \in \mathscr{K}_{\mathrm{ac}}(\hat{H})$. Because we can arbitrarily take λ_1, λ_2 , we find $\mathscr{K} = \mathscr{K}_{\mathrm{ac}}(\hat{H})$, that is, $\mathscr{K}_{\mathrm{sing}}(\hat{H}) = \emptyset$.

4 Asymptotic completeness

Finally, the main theorem is presented. For all $\phi \in \mathscr{K}$, $\sigma_1, \sigma_2 \in \mathbf{R}_{\pm} := \{a \in \mathbf{R} \mid \pm a \ge 0\}$,

$$\begin{split} \left\| e^{i\sigma_{1}\hat{H}_{0}}e^{-i\sigma_{1}\hat{H}}\phi - e^{i\sigma_{2}\hat{H}_{0}}e^{-i\sigma_{2}\hat{H}}\phi \right\|_{\mathscr{H}} \\ &\leq \sup_{\|\psi\|_{\mathscr{H}=1}} \int_{\sigma_{2}}^{\sigma_{1}} \left| \left(\rho_{1}e^{-i\sigma\hat{H}}\phi, \rho_{2}e^{-i\sigma\hat{H}_{0}}\psi \right) \right| d\sigma \\ &\leq \sup_{\|\psi\|_{\mathscr{H}=1}} \left(\int_{\sigma_{2}}^{\sigma_{1}} \left\| |V|^{1/2}e^{-i\sigma\hat{H}}\phi \right\|_{\mathscr{H}}^{2} d\sigma \right)^{1/2} \left(\int_{\sigma_{2}}^{\sigma_{1}} \left\| |V|^{1/2}e^{-i\sigma\hat{H}_{0}}\psi \right\|_{\mathscr{H}}^{2} d\sigma \right)^{1/2} \\ &\to 0, \quad \text{as } \sigma_{1}, \sigma_{2} \to \infty \end{split}$$

holds, where we use (16) and (17). Hence, there is a strong limit: $s - \lim_{\sigma \to \pm \infty} e^{i\sigma \hat{H}_0} e^{-i\sigma \hat{H}}$ exists without any restrictions on the domain. Similarly, we have a strong limit: $s - \lim_{\sigma \to \pm \infty} e^{i\sigma \hat{H}} e^{-i\sigma \hat{H}_0}$. Consequently, we obtain the unitary equivalence of the wave operators on \mathscr{K} :

$$\operatorname{Ran}\left(\mathrm{s-}\lim_{\sigma\to\pm\infty}e^{i\sigma\hat{H}}e^{-i\sigma\hat{H}_{0}}\right)=\mathscr{K}.$$

Using the Howland-Yajima method, the proof for Theorem 1.10 is completed by showing the existence of W^{\pm} ; more precisely, refer to §4 of [12] or the end §6 of [1]. We now show the existence of W^{\pm} . Owing to the Cook-Kuroda method, it suffices to show that for all $\phi \in \mathscr{S}(\mathbf{R}^2)$,

$$\int_{1}^{\infty} \left\| |V|^{1/2} U_0(t,0)\phi \right\|_{L^2(\mathbf{R}^2)} dt \le C.$$
(21)

Hence, we show that

$$\sum_{N=1}^{\infty} \int_0^T \left\| |V|^{1/2} U_0(t+NT,0)\phi \right\|_2 dt \le C.$$
(22)

holds. Then, (21) holds. By noting (8), for p > 4, the left hand side of (22) is smaller than

$$C \left\| |V|^{1/2} \right\|_p \|\phi\|_{2p/(p+2)} \sum_{N=0}^{\infty} \int_0^T |\zeta_2(t+NT)|^{-2/p} dt$$

where we also use (5) and (7) with $\tau = t + NT$ and s = 0. From (15), we obtain (22).

5 Uniform resolvent estimate for H

In this section, we present the URE for \hat{H} . To demonstrate this, we employ the approach according to Herbst [3]. However, in [3], among the specific conditions, only the Stark Hamiltonian has been fully used, and imitating this approach may be difficult. To overcome this difficulty, we should find the alternative condition of \hat{H} , as well as the following *Lipschitz continuity for resolvent of* \hat{H}_0 , which plays a crucial role in mimicking the approach of [3]:

Theorem 5.1. Let $z_{\pm}, w_{\pm} \in \mathbf{C}_{\pm}$. Then, for all $\phi \in \mathscr{K}$,

$$\left\| \left(\rho_1 (\hat{H}_0 - z_{\pm})^{-1} \rho_2 - \rho_1 (\hat{H}_0 - w_{\pm})^{-1} \rho_2 \right) \phi \right\|_{\mathscr{H}} \le C |z_{\pm} - w_{\pm}| \|\phi\|_{\mathscr{H}}.$$

holds.

Remark 5.2. In the case where $k_0 = -\Delta$, the Hölder continuity

$$\left\| \langle x \rangle^{-s} \left((k_0 - z_{\pm})^{-1} - (k_0 - w_{\pm})^{-1} \right) \langle x \rangle^{-s} \phi \right\|_2 \le C |z_{\pm} - w_{\pm}|^{(2s-1)/(2s+1)} \|\phi\|_2$$
(23)

holds. Here, formally, we consider $s \to \infty$; then, the power on $|z_{\pm} - w_{\pm}|$ in (23) tends to 1 (that is, the Lipschitz continuity). In our model, compared with the exponential growth of |x(t)| in t, the decay for the potential $\rho_j \in L^p(\mathbf{R}^2)$ is fast. In the words of k_0 , the potential decays exponentially in x. Thus, it is not surprising that Lipschitz continuation (23) holds.

This theorem can be obtained as the sub-consequence of the following lemma;

Lemma 5.3. Suppose Assumption 1.2. For a sufficiently small $\varepsilon > 0$, let $\tau \in [-\varepsilon, \varepsilon]$. Define

$$\Sigma_R = \int_0^R \left\| \sigma \rho_1 e^{-i(H_0 + \lambda \pm \tau i)\sigma}(\rho_2 \phi) \right\|_{\mathscr{H}} d\sigma$$

for R > 0. Then, there exists $\varepsilon > 0$ such that for all $\tau \in [-\varepsilon, \varepsilon]$ and $\lambda \in \mathbf{R}$, the limit $\lim_{R\to\infty} \Sigma_R$ exists, satisfying

$$\lim_{R \to \infty} \Sigma_R < C \|\phi\|_{\mathscr{K}}$$

Furthermore,

$$s - \lim_{\sigma \to \infty} \sigma \rho_1 e^{-i(H_0 + \lambda \pm \tau i)\sigma} \rho_2 = 0$$

holds.

Proof. Let $R = N_0T + s$ with $N_0 \in \mathbb{N}$ and $s \in [0, T]$. Then, the same calculations in the proof of Proposition 2.2 yield

$$\Sigma_R \le C \sum_{N=0}^{N_0+1} \int_0^T NT e^{\varepsilon NT} \|\rho_1 U_0(t+NT,s)(\rho_2 \phi)\|_{\mathscr{H}} ds$$
$$\le C \sum_{N=0}^{N_0+1} |NT| e^{\varepsilon NT} e^{-2\lambda N/p} \|\phi\|_{\mathscr{H}}$$

with p > 4. Then, if $|\varepsilon|$ is sufficiently small, there exists $\varepsilon_1 > 0$ such that $(|NT|e^{\varepsilon NT}e^{-2\lambda/p}) \le Ce^{-\varepsilon_1 N}$. This implies that $\lim \Sigma_R$ exists and $\lim \Sigma_R < \infty$.

Because

$$\rho_{1}(\hat{H}_{0}-z_{\pm})^{-1}\rho_{2}\phi - \rho_{1}(\hat{H}_{0}-w_{\pm})^{-1}\rho_{2}\phi = \mp i\rho_{1}\int_{0}^{\infty} \left(e^{\mp i\sigma(\hat{H}_{0}-z_{\pm})} - e^{\mp i\sigma(\hat{H}_{0}-w_{\pm})}\right)\rho_{2}\phi d\sigma$$
$$= \mp i\int_{0}^{\infty}\sigma\rho_{1}e^{\mp i\sigma\hat{H}_{0}}\rho_{2}\left(\frac{e^{\mp i\sigma z_{\pm}} - e^{\mp i\sigma w_{\pm}}}{\sigma}\right)\phi d\sigma,$$

and Lemma 5.3 with $\lambda = \tau = 0$, Theorem 5.1 can be proven.

5.1 Proof of Theorem 1.10

Now, we prove Theorem 1.10. The fundamental approach is based on the approach described by Herbst [3]. By the resolvent formula, we have that

$$\rho_1(\hat{H} - \lambda - i\varepsilon)^{-1}\rho_2 = (1 + \rho_1(\hat{H} - \lambda - i\varepsilon)^{-1}\rho_2)^{-1} \cdot \rho_1(\hat{H}_0 - \lambda - i\varepsilon)^{-1}\rho_2$$

holds. Hence, we prove that for any $\varepsilon \geq 0$, if $\psi = \psi_{\varepsilon} \in \mathscr{K}$ such that $(1 + \rho_1(\hat{H} - \lambda - i\varepsilon)^{-1}\rho_2)\psi = 0$, then $\psi \equiv 0$. Therefore, we have that $(1 + \rho_1(\hat{H} - \lambda - i\varepsilon)^{-1}\rho_2)$ is invertible for any $\varepsilon \geq 0$, which proves that Theorem 1.10 holds.

Let $\varphi = (\hat{H}_0 - \lambda - i\varepsilon)^{-1} \rho_2 \psi$, then

$$(\hat{H} - \lambda - i\varepsilon)\varphi = 0 \tag{24}$$

holds. Here, remarking \hat{H} is a selfadjoint operator and $\lambda \notin \sigma_{pp}(\hat{H})$, (24) implies $\varphi \equiv 0$, (i.e., $\psi \equiv 0$) if $\varphi \in \mathscr{D}(\hat{H})$. Thus, we prove $\varphi \in \mathscr{D}(\hat{H})$, as follows: Using the resolvent formula

$$\varphi = (\hat{H}_0 - i)^{-1} \rho_2 \psi + (\lambda + i\varepsilon - i)(\hat{H}_0 - i)^{-1} \varphi,$$

we notice that it suffices to show that $\varphi \in \mathscr{K}$ shows $\varphi \in \mathscr{D}(\hat{H})$. Hence, we show $\varphi \in \mathscr{K}$. In the case where $\varepsilon > 0$, it is clear that $\varphi \in \mathscr{K}$; hence, we only consider the case where

$$\begin{split} \varepsilon &\to 0, \text{ that is, } \psi = -\rho_1 (\hat{H}_0 - \lambda - i0)^{-1} \rho_2 \psi \text{ and } \varphi = (\hat{H}_0 - \lambda - i0)^{-1} \rho_2 \psi: \\ \|\varphi\|_{\mathscr{K}}^2 &= \lim_{\varepsilon \to 0} \left\| (\hat{H}_0 - \lambda - i\varepsilon)^{-1} \rho_2 \psi \right\|_{\mathscr{K}}^2 \\ &= \lim_{\varepsilon \to 0} \left\{ \left(\rho_2 (\hat{H}_0 - \lambda + i\varepsilon)^{-1} (\hat{H}_0 - \lambda - i\varepsilon)^{-1} \rho_2 \psi, \psi \right) \right\} \\ &= \lim_{\varepsilon \to 0} \left\{ \frac{-i}{2\varepsilon} \left(\rho_2 \left((\hat{H}_0 - \lambda - i\varepsilon)^{-1} - (\hat{H}_0 - \lambda + i\varepsilon)^{-1} \right) \rho_2 \psi, \psi \right) - \frac{i}{2\varepsilon} ((\psi, \psi) - (\psi, \psi)) \right\} \\ &= \lim_{\varepsilon \to 0} \lim_{\delta_1 \to 0} \left\{ \frac{-i}{2\varepsilon} \left(\left(\rho_2 (\hat{H}_0 - \lambda - i\varepsilon)^{-1} - (\hat{H}_0 - \lambda + i\varepsilon)^{-1} \right) \rho_2 \psi, \psi \right) - \frac{i}{2\varepsilon} ((\psi, \psi) - (\psi, \psi)) \right\} \\ &- \lim_{\varepsilon \to 0} \lim_{\delta_2 \to 0} \left\{ \frac{-i}{2\varepsilon} \left(\left(\rho_2 (\hat{H}_0 - \lambda - i\varepsilon)^{-1} \rho_2 - \rho_1 (\hat{H}_0 - \lambda - i\delta_1)^{-1} \rho_2 \right) \psi, \psi \right) \right\} \\ &= :\lim_{\varepsilon \to 0} \frac{-i}{2\varepsilon} (I_1(\varepsilon) - I_2(\varepsilon)) . \end{split}$$

We have

_

$$I_1(\varepsilon) = \lim_{\delta_1 \to 0} \left(\left(\rho_2 \left((\hat{H}_0 - \lambda - i\varepsilon)^{-1} - (\hat{H}_0 - \lambda - i\delta_1)^{-1} \right) \rho_2 \right) \psi, \psi \right) + \left(\frac{\rho_1 - \rho_2}{\rho_1} \psi, \psi \right).$$

and

$$I_2(\varepsilon) = \lim_{\delta_2 \to 0} \left(\psi, \left(\rho_2 \left((\hat{H}_0 - \lambda - i\varepsilon)^{-1} - (\hat{H}_0 - \lambda - i\delta_2)^{-1} \right) \rho_2 \right) \psi \right) + \left(\psi, \frac{\rho_1 - \rho_2}{\rho_1} \psi \right).$$

These and (23) yield

$$\left\|\varphi\right\|_{\mathscr{K}}^{2} \leq C\left(\lim_{\varepsilon \to 0} \lim_{\delta_{1} \to 0} \frac{|\varepsilon - \delta_{1}|}{\varepsilon} + \lim_{\varepsilon \to 0} \lim_{\delta_{2} \to 0} \frac{|\varepsilon - \delta_{2}|}{\varepsilon}\right) \left\|\psi\right\|_{\mathscr{K}}^{2} \leq C \left\|\psi\right\|_{\mathscr{K}}^{2},$$

where we use

$$\left(\frac{\rho_1-\rho_2}{\rho_1}\psi,\psi\right)-\left(\psi,\frac{\rho_1-\rho_2}{\rho_1}\psi\right)=0.$$

Therefore, we obtain $\varphi \equiv 0$ and which implies $\psi \equiv 0$.

References

- [1] Adachi, T., Kawamoto, M.: Quantum scattering theory in a periodically pulsed magnetic field, Annales Henri Poincaré, 17, 2409–2438, (2016).
- [2] Enss, V., Veselić, K.: Bound states and propagating states for time-dependent hamiltonians, Ann., de l'I.H.P, section A, **39**, 159–191, (1983).
- [3] Herbst, I.W.: Unitary equivalence of Stark Hamiltonians, Math. Z., 155, 55–70, (1977).

- [4] Howland, J.S.: Scattering theory for hamiltonians periodic in time, Indiana Univ. Math. J. 28, 471–494, (1979).
- [5] Kargaev, P., Korotyaev, E.L.: The inverse problem for the Hill operator, a direct approach. Inv. Math., 129, 567–593, (1997).
- [6] Kato, T.: Wave operators and similarity for some non-self-adjoint operators, math. Ann. 162, 258–279, (1966).
- [7] Kawamoto, M.: Mourre theory for time-periodic magnetic fields, J. Func., Anal., 277, 1–30, (2019).
- [8] Kawamoto, M.: Strichartz estimates for Schrödinger operators with square potential with time-dependent coefficients, Differ. Equ. Dyn. Syst., (2020).
- [9] Kitada, H., Yajima, K.: Bound states and scattering states for time periodic hamiltonians, Ann., de l'I.H.P, section A, 39, 145–157, (1983).
- [10] Korotyaev, E. L.: On scattering in an external, homogeneous, time-periodic magnetic field, Math. USSR-Sb. 66, 499–522 (1990)
- [11] Møller, J. S.: Two-body short-range systems in a time-periodic electric field, Duke Math. J. 105 135-166 (2000).
- [12] Yajima, K.: Schrödinger equations with potentials periodic in time, J. Math. Soc. Japan. 29, 729–743 (1977).
- [13] Yajima, K.: Schrödinger evolution equations with magnetic fields, J. Anal. Math. 56, 29–76 (1991).