

Observing intermediate-mass black holes and the upper–stellar-mass gap with LIGO and Virgo

AJIT KUMAR MEHTA,¹ ALESSANDRA BUONANNO,^{1,2} JONATHAN GAIR,¹ M. COLEMAN MILLER,³ EBRAHEEM FARAG,^{4,5}
R. J. DEBOER,^{6,5} M. WIESCHER,^{6,5} AND F.X. TIMMES^{4,5}

¹*Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, Potsdam 14476, Germany*

²*Department of Physics, University of Maryland, College Park, MD 20742-2421, USA*

³*Department of Astronomy and Joint Space-Science Institute, University of Maryland, College Park, MD 20742-2421, USA*

⁴*School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85287, USA*

⁵*Joint Institute for Nuclear Astrophysics - Center for the Evolution of the Elements, USA*

⁶*Department of Physics, University of Notre Dame, Notre Dame, Indiana 46556, USA*

ABSTRACT

We investigate the potential of ground-based gravitational-wave detectors to probe the mass function of intermediate-mass black holes (IMBHs) wherein we also include BHs in the upper mass gap $\sim 60 - 130 M_{\odot}$. Using the noise spectral density of the upcoming LIGO and Virgo fourth observing (O4) run, we perform Bayesian analysis on quasi-circular non-precessing, spinning IMBH binaries (IMBHBs) with total masses $50-500 M_{\odot}$, mass ratios 1.25, 4, and 10, and (dimensionless) spins up to 0.95, and estimate the precision with which the source-frame parameters can be measured. We find that, at 2σ , the source-frame mass of the heavier component of the IMBHBs can be constrained with an uncertainty of $\sim 10 - 40\%$ at a signal to noise ratio of 20. Focusing on the stellar-mass gap, we first evolve stars with massive helium cores using the open-source MESA software instrument to establish the upper and lower edges of the mass gap. We determine that the lower edge of the mass gap is $\simeq 59^{+34}_{-13} M_{\odot}$, while the upper edge is $\simeq 139^{+30}_{-14} M_{\odot}$, where the error bars indicate the mass range that follows from the $\pm 3\sigma$ uncertainty in the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ nuclear rate. We then study IMBHBs with components lying in the mass gap and show that the O4 run will be able to robustly identify most such systems. In this context, we also re-analyze the GW190521 event and show that the 90% confidence interval of the primary-mass measurement lies inside the mass gap. Finally, we show that the precision achieved with the O4 run (and future O5 run) could be crucial for understanding the mass function, the formation mechanism, and evolution history of IMBHs.

Keywords: Gravitational-wave astronomy (675); Stellar-mass black holes (1611); Nuclear astrophysics (1129); Stellar physics (1621);

1. INTRODUCTION

The LIGO and Virgo detectors (Aasi et al. 2015; Acernese et al. 2015) have opened the gravitational-wave (GW) window onto the universe, reporting, as of today, 48 GW signals from binary black-hole (BH) mergers (Abbott et al. 2019a, 2020b). They have also opened the era of multi-messenger astronomy with GWs, shedding light on the origin of short-hard gamma-ray bursts with the observation of a coalescing binary neutron star (Abbott et al. 2017a,b). In addition, independent claims of GW observations have also been made (Nitz

et al. 2019, 2020; Venumadhav et al. 2020; Zackay et al. 2019). Detections of these compact-object binaries have allowed us to probe various problems pertaining to astrophysics, astronomy and cosmology, such as measuring the Hubble parameter (Abbott et al. 2017c), establishing the rates and population of compact binaries (Abbott et al. 2020c), and constraining the neutron-star radius and equation of states (Abbott et al. 2018, 2020b). The detections of binary black-hole (BBH) mergers with masses $\gtrsim 40 M_{\odot}$, in particular, can allow us to probe the physical processes that are involved in the evolution of massive stars — for example the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ nuclear rate that significantly affects the mass of a BH formed through the collapse of a massive star (Brown

et al. 2001; Woosley et al. 2002; Farmer et al. 2020; Woosley & Heger 2021).

The GW detections during the first and second observing (O1 and O2) runs (Abbott et al. 2019a) revealed a population of BBHs with component source masses $\lesssim 50M_\odot$ and total source mass $\lesssim 84M_\odot$. These component masses are mostly consistent with the definition of stellar-mass BHs. However, in the first half of the third (O3a) observing run, an event (GW190521) (Abbott et al. 2020d,e) was detected with a pre-merger binary total source mass of $\sim 150M_\odot$ and a remnant source mass of $\sim 140M_\odot$. The best estimates of the component BH source masses, obtained with quasi-circular spinning, precessing waveforms, are $\sim 85M_\odot$ and $\sim 65M_\odot$ (Abbott et al. 2020d). The remnant of this GW event falls in the category of intermediate-mass black holes (IMBHs), which are usually defined as BHs with mass between $\sim 10^2$ – 10^5M_\odot . The latter is not a strict definition, and since here we are also interested in studying BHs in the upper-stellar-mass gap (i.e., the pair-instability-supernova mass gap), we adopt a threshold of $M \gtrsim 60M_\odot$ for IMBHs. Minimal-assumption LIGO-Virgo pipelines dedicated to searches for IMBHs did not report any detection except for GW190521, so far (Abbott et al. 2017d, 2019b).

We note that Nitz & Capano (2021) have recently found evidence, using a waveform model that was not employed in Abbott et al. (2020d,e), that the source-mass posterior distributions of GW190521 are multimodal, opening the possibility that GW190521 was an intermediate mass-ratio binary (i.e., a binary with mass ratio around 10). We will comment on their analysis, and we re-analyse this GW event as well in Sec. 4.5 below. A similar conclusion was drawn by Fishbach & Holz (2020), in which the authors employed a different, namely, population-informed prior on the secondary mass rather than a uniform prior on the secondary mass. Given the very short signal, other analyses in the literature (Romero-Shaw et al. 2020; Calderón Bustillo et al. 2020; Gayathri et al. 2020b; Bustillo et al. 2021) pointed out the importance of re-analysing GW190521 with waveform models of eccentric compact binaries, which, however, are not yet available.

IMBHs are difficult to observe. Indeed, there is no definitive electromagnetic evidence for their existence. Thus, their formation channels and mass function are highly uncertain (Miller & Colbert 2004; Amaro-Seoane et al. 2007; Gair et al. 2011; Belczynski et al. 2014). By contrast, GW observations have the potential to solve these mysteries, providing accurate measurements of their properties, such as their masses, spins, and location. This, of course, depends on the rate at which

upcoming LIGO and Virgo runs would be observing IMBHB mergers (See Ezquiaga & Holz (2021) for a study when BH’s masses are above $120M_\odot$). Based on the observation of one event, GW190521, Abbott et al. (2020e) estimated a merger rate of $0.13^{+0.30}_{-0.11} \text{ Gpc}^{-3} \text{ yr}^{-1}$.

Graff et al. (2015); Veitch et al. (2015a); Haster et al. (2016) studied the precision with which IMBH masses and spins could be measured with LIGO and Virgo detectors. Graff et al. (2015) used multipolar waveform models that describe the entire coalescence process of non-spinning IMBHBs, and performed a Bayesian analysis to estimate the precision with which the parameters of IMBHBs can be estimated with LIGO detectors. Veitch et al. (2015a) employed spinning, non-precessing waveforms, but did not include subdominant modes; these are relevant for high total-mass binaries, because they break degeneracies between parameters and reduce the measurement uncertainties. Here we extend these analyses in several directions. We consider state-of-the-art multipolar spinning, non-precessing models for gravitational waves from IMBHBs, including the five strongest gravitational modes, and explore a larger region of the parameter space (e.g., mass ratios 1 – 10 and dimensionless spin values up to 0.95). We focus on masses in the source frame rather than the detector frame, since the former provides us with information about the upper-stellar-mass gap and, more generally, about the IMBH mass function. We employ for our study the projected noise-spectral densities (Abbott et al. 2020a) of the upcoming fourth observing (O4) run (scheduled to start not earlier than the second half of 2022). We also comment on results that could be obtained during the fifth observing (O5) run (Abbott et al. 2020a) (expected to start in 2025).

To set the stage, we show in Figure 1 the distance reach of the LIGO-Virgo-detector network during the O4 run to binaries with signal-to-noise (SNR) of 20. We display results using multipolar spinning, non-precessing waveforms, for a variety of spin values and for mass ratios of 1.25 and 4, and binary inclinations $\iota = 0$ (face-on) and $\iota = \pi/3$ (close to edge-on). As we can see, the distance reach for face-on binaries even at SNR=20 could go up to a redshift $z \sim 1.4$ (11 Gpc), while for near edge-on binaries this reduces a bit but sources at a redshift of ~ 0.9 (6 Gpc) can still be probed. As stated previously, though highly uncertain, as it is based on just a single observation, Abbott et al. (2020e) estimated a merger rate of $0.13^{+0.30}_{-0.11} \text{ Gpc}^{-3} \text{ yr}^{-1}$ for IMBH mergers. This implies that, at best, we might expect a detection rate of $\sim 42.86^{+184.63}_{-36.26} \text{ yr}^{-1}$ for face-on binaries and $\sim 21.32^{+91.86}_{-18.04} \text{ yr}^{-1}$ for near edge-on IMBH binaries with SNR ~ 20 at O4 sensitivity. At O5 sensitivity at the




Figure 1. The distance reach as a function of the total mass M_{tot} for spinning, non-precessing IMBHBs with mass ratio $q = 1.25$ (solid lines) and $q = 4$ (dashed lines) at inclination angle $\iota = \pi/3$ (left panel) and $\iota = 0$ (right panel), and SNR = 20. The different curves correspond to different values of the component spins (χ_1, χ_2) , assumed to be equal for the two BHs. We use the noise spectral densities expected for the upcoming O4 run (Abbott et al. 2020a) and the spinning, non-precessing SEOBNRHM waveform model. The distance reach shown here is computed by averaging over the antenna pattern functions (see Equation 4), that is the angles that specify the location of the source in the sky and the polarization angle. The maximum redshift (distance reach) for IMBHBs with inclination $\iota = 0$ (face-on) and $\pi/3$ (near edge-on) is $z \sim 1.4$ (11 Gpc) and ~ 0.9 (6 Gpc), respectively.

same SNR, the maximum redshift reach for face-on binaries can go up to ~ 2.5 , while for near edge-on binaries it is ~ 1.6 . The previous numbers then increase to $\sim 115.96^{+499.54}_{-98.12} \text{ yr}^{-1}$ and $\sim 54.56^{+235.03}_{-46.16} \text{ yr}^{-1}$. Note that these numbers are just for illustrative purposes. A real computation of these numbers would require a knowledge of the mass distribution of the IMBHBs in binaries as a function of redshift, in addition to the merger rate density of IMBHBs. These are not yet very accurately constrained.

As said earlier, inferring IMBH parameters and estimating the corresponding measurement uncertainties will have important implications in understanding the formation of high-mass BHs and the evolution of massive stars. The theory of stellar evolution predicts that stars with zero-age main-sequence (ZAMS) masses $100 M_{\odot} \lesssim M_{\text{ZAMS}} \lesssim 130 M_{\odot}$ are subject to pulsational pair-instability (PI, Fowler & Hoyle 1964; Barkat et al. 1967; Rakavy & Shaviv 1967), which causes the stars to lose mass and leave behind a remnant with a mass smaller than $\sim 65 M_{\odot}$ (Heger et al. 2003; Blinnikov 2010; Chatzopoulos & Wheeler 2012; Yoshida et al. 2016; Woosley 2017; Umeda et al. 2020). The stars with masses $130 M_{\odot} \lesssim M_{\text{ZAMS}} \lesssim 250 M_{\odot}$ are subject to a PI, which disrupts them completely, and hence no BH forms. Stars with $M_{\text{ZAMS}} \gtrsim 250 M_{\odot}$ can collapse directly to IMBHBs with a mass $\gtrsim 135 M_{\odot}$. Thus, in the standard picture, there should be an upper-stellar-mass BH gap in the range $[65, 135] M_{\odot}$, and any BH observed

in this range (e.g., the primary BH of GW190521) have to form via other formation channels — for example through hierarchical coalescence of smaller BHs or direct collapse of a stellar merger between an evolved star and a main-sequence companion (e.g., O’Leary et al. 2006; Gerosa & Berti 2017; Di Carlo et al. 2019; Antonini et al. 2019; Rodriguez et al. 2019; Gayathri et al. 2020a; Kimball et al. 2020; Di Carlo et al. 2020; Mapelli et al. 2021; Quinlan & Shapiro 1989; Portegies Zwart & McMillan 2000; Ebisuzaki et al. 2001; Miller & Hamilton 2002). However, the exact mass boundaries of the gap depend on parameters that are uncertain. For example, the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ nuclear reaction rate, which converts carbon to oxygen in the core, can affect the boundary significantly (Takahashi 2018; Farmer et al. 2020; Woosley & Heger 2021; Costa et al. 2021). Here, to better determine whether future GW observations will be able to observe BHs in the mass gap, we re-compute the mass-gap boundaries with updated $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate, and much finer mass and time grids. The complexity of this system has made a reliable analysis

of the reaction a decades old challenge¹. The rapidly declining cross section at low energies has prohibited a direct measurement of the reaction at stellar temperatures and the reaction rate is entirely based on the theoretical analysis and extrapolation of the experimental data towards lower energies. The newly derived rate by deBoer et al. (2017) using a multi-channel analysis approach, derives for the first time a reliable prediction for the interference patterns within the reaction components by taking into account all available experimental data sets that cover the near threshold energy range of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ process.

The paper is organized as follows. In Section 2 we introduce the gravitational waveform models that we employ for our parameter-estimation studies, and briefly review the Bayesian-analysis method that we use to infer the source properties from the GW signals. In Section 3 we first describe the parameter space of the binary simulations that we investigate and the choice of priors. Then, we present the results for the expected measurement uncertainties that could be obtained with observations made during the LIGO-Virgo O4 run and also comment on results that could be obtained during the O5 run. We also discuss the bi-modality that appears in the posterior distributions for some parameters in some regions of the parameter space. In Section 4, after a brief review of the BH mass gap and a discussion of the current estimate of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rates, we evolve massive He stars by incorporating new uncertainties in the nuclear-reaction rates and we establish new bounds on the lower and upper edges of the mass gap. Then, using these results, we estimate the probability with which LIGO-Virgo O4 and O5 runs can identify IMBHB systems whose primary and secondary masses lie in the BH mass gap. We also re-analyse GW190521 with the spinning, non-precessing waveform models employed in this work, and find that, although a bi-modality in the posterior distributions of the detector-frame masses is present, it is absent in the posteriors of the source-mass parameters. To contrast these findings with precessing waveforms, we also analyze GW190521 with one precessing waveform model, whose former public version was employed in Nitz & Ca-

pano (2021). Finally, in Section 5, we present our main conclusions and discuss possible future research directions.

2. SETUP

2.1. Waveform models

We focus our study on GW signals generated by BBHs with non-precessing spins, moving on quasi-circular orbits. Such signals are described by 11 parameters: $\theta \equiv \{m_1, m_2, \chi_1, \chi_2, d_L, t_c, \delta, \alpha, \iota, \psi, \phi_c\}$. The parameters $m_{1,2}$ are the redshifted (i.e., detector-frame) component masses $m_i = (1+z)m_i^s$ for $i = 1, 2$, where m_i^s is the source-frame mass and z the redshift. The quantities $\chi_{1,2}$ are the dimensionless component spins along the orbital angular momentum \mathbf{L} of the binary (i.e., $\chi_i = \mathbf{S}_i \cdot \mathbf{L}/m_i^2$ for $i = 1, 2$). The parameter d_L is the luminosity distance to the binary, which along with the declination δ and the right ascension α define the location of the binary in the sky. The parameter t_c is the merger time or more specifically, it is the peak time of the $\ell = 2, m = 2$ gravitational mode at the geocenter. The angle ι measures the inclination of the binary's total angular momentum \mathbf{J} (which, for non-precessing binary systems, has the same direction as the orbital angular momentum \mathbf{L}) with respect to the line of sight from the detector at the geocenter. The remaining parameters ϕ_c and ψ are the merger phase and the gravitational wave polarization, respectively.

It is also useful to define the following binary parameters: the total mass $M_{\text{tot}} = m_1 + m_2$, the mass ratio $q = m_1/m_2 \geq 1$, the symmetric mass ratio $\nu = q/(1+q)^2$, the chirp mass $M_c = M_{\text{tot}}\nu^{3/5}$ and the effective spin of the binary $\chi_{\text{eff}} = (m_1\chi_1 + m_2\chi_2)/M_{\text{tot}}$.

In general relativity, GWs are described by the two polarizations $h_+(t)$ and $h_\times(t)$. The complex waveform defined by $h(t) \equiv h_+(t) - ih_\times(t)$ can be conveniently decomposed in a basis of -2 spin-weighted spherical harmonics (Pan et al. 2011):

$$h(t; \lambda, \iota, \varphi_c) = \frac{1}{d_L} \sum_{\ell \geq 2} \sum_{|m| \leq \ell} {}_{-2}Y_{\ell m}(\iota, \varphi_c) h_{\ell m}(t, \lambda), \quad (1)$$

where λ denotes a subset of the θ parameters, namely, the intrinsic parameters of the binary systems such as masses (m_1, m_2) and spins (χ_1, χ_2).

The GW signal emitted throughout the coalescence of a BBH can be divided into three phases: inspiral, merger and ringdown (IMR). The inspiral phase describes the steady, adiabatic evolution of the system where the component BHs come closer and closer to each other, losing orbital energy because of GW emission. At the end of the inspiral, the BHs plunge into each other, form a common apparent horizon and merge. The ringdown phase

¹ The reaction rate for $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ is determined by the quantum structure of the compound nucleus ^{16}O as an α cluster system. It is characterized by the interfering $\ell=1$ waves of the $J^\pi=1^-$ resonances and sub-threshold levels defining an E1 component for the reaction cross section as well as by the $\ell=2$ components and interference from broad $J^\pi=2^+$ resonances and the non-resonant E2 external capture to the ground state of ^{16}O . In addition to these two main E1 and E2 ground state components, transitions to higher lying excited states occur that also add to the total cross section.

describes the evolution of the system as the remnant object settles down to a stationary (Kerr) BH.

Here, we employ, as the main IMR waveform model, the one developed within the effective one-body formalism (EOB), which is a semi-analytical method that combines results from post-Newtonian (PN) theory for the inspiral, BH perturbation theory for the ringdown, and numerical relativity (NR) for the merger stage. More specifically, since we are interested in studying high-mass BBHs with mass ratio as large as 10, we employ the quasi-circular, non-precessing spinning waveform models with gravitational modes beyond the dominant, quadrupolar one (Cotesta et al. 2018) (henceforth, **SEOBNRHM**)². The **SEOBNRHM** model contains the five strongest modes $(\ell, m) = (2, \pm 1), (2, \pm 2), (3, \pm 3), (4, \pm 4), (5, \pm 5)$ (see Equation (1)). We shall also use for synthetic (injection) signals a spinning, precessing IMR model built directly interpolating NR waveforms **NRSurPHM** (Varma et al. 2019a)³. Such a model contains all modes with $\ell \leq 5$. However, the extra modes present in **NRSurPHM** and absent in **SEOBNRHM** are not expected to contribute significantly for our choices of parameters. Finally, we shall also use a phenomenological IMR waveform model built in the frequency domain by combining EOB and NR waveforms, **PhenomHM** (García-Quirós et al. 2020)⁴. We stress that, in general, the higher-order (or subdominant) modes become important in the parameter estimation of the binaries when the inclination angle is large and the mass ratio is large ($q \equiv m_1/m_2 \geq 1$) (see, e.g., Cotesta et al. (2018)). Moreover, higher total mass and higher spins also increase the amplitude of subdominant modes, especially close to the merger of the binary. We will see in Sections 3.2 and 3.3 that higher modes can be important to precisely infer the parameters even when the mass ratio is as low as $q \sim 1.25$, but the total mass and/or the spins are high.

2.2. Bayesian statistics

The Bayes' theorem allows us to construct the probability distribution of parameters θ given a hypothesis (or a model) \mathcal{H} and a data set d . It states:

$$P(\theta|d, \mathcal{H}) = \frac{P(d|\theta, \mathcal{H})P(\theta|\mathcal{H})}{P(d|\mathcal{H})}, \quad (2)$$

² In the LIGO Algorithm Library (LAL) the technical name of this waveform model is **SEOBNRv4HM.ROM**. It is the reduced-order model (ROM) of the time-domain waveform model **SEOBNRv4HM** (Cotesta et al. 2018, 2020).

³ In LAL this waveform model is denoted **NRSur7dq4** (Varma et al. 2019a).

⁴ In LAL this waveform model is denoted **IMRPhenomXHM** (García-Quirós et al. 2020).

where $P(\theta|d, \mathcal{H})$ is the posterior probability distribution of parameters θ , given a data set d , under the hypothesis \mathcal{H} . The quantity $P(\theta|\mathcal{H})$ in Equation (2) is the prior probability distribution of the parameters θ under the hypothesis \mathcal{H} . The function $P(d|\theta, \mathcal{H})$ is the likelihood (very often denoted as $\mathcal{L}(\theta)$) for obtaining the data set d with a specific parameter set θ , under the hypothesis \mathcal{H} . Bayes' theorem updates our prior knowledge of the parameters using the likelihood of the data to finally provide us with the posterior probability distribution. Lastly, the quantity $P(d|\mathcal{H})$ in Equation (2) is known as the evidence for the data set d under the hypothesis \mathcal{H} . It is the normalization factor of the posterior probability distribution in Equation (2) and, thus, it does not matter in our parameter-estimation study. It is, however, widely used for comparing different hypotheses.

In this work, we assume that we have already detected a GW signal and our job is to extract the parameters that describe this signal most closely. The data output from the GW detectors can be written as follows (under the assumption of additive noise):

$$d = n + h, \quad (3)$$

where n represents the noise realization from the GW detectors and h is the GW strains measured at the different detectors. Here

$$h = F_+(\alpha, \delta, \psi) h_+ + F_\times(\alpha, \delta, \psi) h_\times, \quad (4)$$

where $F_{+, \times}(\alpha, \delta, \psi)$ denote the antenna-pattern functions (Finn & Chernoff 1993) that account for the angular sensitivity of the GW detectors and thus depend on the source location (α, δ) , and on the polarization angle, ψ , that defines the relative orientation of the polarization axes with respect to which the polarization states $h_{+, \times}$ are defined. To construct the posterior distribution of parameters θ , which describe the GW signal h , we must first write down the likelihood function.

2.2.1. Likelihood function

Let d_a denote a data stream in a particular detector a . Then the likelihood $\mathcal{L}_a(\theta)$ or $P(d_a|\theta, \mathcal{H}_S)$ is, by definition, the probability of obtaining the data d_a with a specified set of parameters θ . Thus, for a GW signal $h_a(\theta)$, the likelihood should be given by the probability of observing the noise realization, $n_a = d_a - h_a(\theta)$. The noise realizations in GW detectors are modeled as independent Gaussian distributions in each frequency bin with a zero mean and a variance given by the detector's power spectral density (PSD). Thus, up to an additive constant,

$$\log \mathcal{L}_a(\theta) \propto -\frac{1}{2} \sum_i \left[\frac{4}{T_a} \frac{|\tilde{d}_a(f_i) - \tilde{h}_a(f_i, \theta)|^2}{S_n^{(a)}(f_i)} \right], \quad (5)$$

where i runs over each frequency bin, T_a is the duration of the GW signal in detector a , and a tilde represents the Fourier transform of the data series. Here $S_n^{(a)}(f)$ is the (one-sided) noise PSD associated with the detector a . The first term in Equation (5) is the discrete approximation to the inner product between the data d and the waveform model (or template) h , denoted by (d, h) , where

$$(d, h) \equiv 4\text{Re} \int_{f_{\min}}^{f_{\max}} df \frac{\tilde{d}(f)\tilde{h}^*(f)}{S_n(f)}, \quad (6)$$

the asterisk denotes the complex conjugate, and f_{\min} and f_{\max} are the minimum and maximum frequency over which the integration is performed. Generally, the frequency limits are different for different detectors. The signal to noise ratio (SNR) of a signal h is defined as $\sqrt{(h, h)}$.

The overlap between two signals h_1 and h_2 is

$$\mathcal{O}_{12} = \max_{t_c, \phi_c} \left[\frac{(h_1, h_2)}{\sqrt{(h_1, h_1)(h_2, h_2)}} \right]. \quad (7)$$

By definition, the overlap varies between 0 and 1, with the latter representing the case when the two signals are scaled versions of each other.

Assuming that the data streams in the different detectors are independent, we can construct the total log likelihood function by summing the individual log likelihoods of the detectors, that is

$$\log \mathcal{L}(\theta) = \sum_a \log \mathcal{L}_a(\theta). \quad (8)$$

From Equation (8), one can show that the total SNR from all the detectors is the sum in quadrature of the individual SNRs,

$$\rho = \sqrt{\rho_1^2 + \rho_2^2 + \cdots + \rho_N^2}, \quad (9)$$

where ρ_a represents the SNR of a signal in detector a , and N is the number of detectors.

In this work, when we evaluate Equations (5)–(9), we use $f_{\min} = 11$ Hz, and we set f_{\max} to half the sampling frequency, which is 4096 Hz. For the results displayed in figures and tables, we use the projected PSDs of the LIGO-Virgo detector network for the upcoming O4 run (Abbott et al. 2020a). We also regenerated most of the results with O5 PSDs (Abbott et al. 2020a) at the same SNR. The duration of the signal T in Equation (5) is chosen to vary between 4 s to 128 s depending on the total mass of the injected signal.

2.2.2. Priors

In order to construct the posterior distribution of the parameters θ using Equation (2), we also need to specify our prior probability distribution $P(\theta|\mathcal{H})$.

For our analysis, we assume a flat prior in the component (detector-frame) masses $m_{1,2} \in [1, 600]M_\odot$ with $m_1 \geq m_2$. For the location of the binary in the sky, we use flat priors in $\cos \delta$ and α with $\delta \in [0, \pi]$ rad and $\alpha \in [0, 2\pi]$ rad. We assume an isotropic distribution for the orientation of the binary with respect to the observer. This implies that the orientation-angle priors are flat in $\cos(\iota)$ for $\iota \in [0, \pi]$ rad, flat in ψ and ϕ_c for $\psi \in [0, \pi]$ rad and $\phi_c \in [0, 2\pi]$ rad. The dimensionless spins, $\chi_{1,2}$, are chosen to follow a uniform distribution between -0.99 and 0.99 .

We make use of two distinct distance priors: i) uniform in Euclidean volume (i.e., flat in d_L^2); and ii) uniform in comoving volume (V_c), i.e.,

$$P(d_L) \propto \frac{1}{1+z} \frac{dV_c}{dz} \left(\frac{dd_L}{dz} \right)^{-1}, \quad (10)$$

where the conversion from redshift, z , to luminosity distance, d_L , depends on the cosmology under consideration. In this work, we use the standard Λ CDM model of the universe (Planck Collaboration et al. 2016). For a spatially flat universe, we have

$$\frac{dd_L}{dz} = d_C + (1+z) \frac{d_H}{E(z)}, \quad (11)$$

where d_C is the comoving distance, $d_H = c/H_0$ is the Hubble distance and $E(z)$ is the normalized Hubble parameter at the redshift z . For both distance priors, we have $d_L \in [100 \text{ Mpc}, 12 \text{ Gpc}]$. By inverting Equation (11), we can obtain the redshift of the source for a given luminosity distance d_L . The source-frame masses are obtained from the detector-frame masses via $m^s = m/(1+z)$.

To sample the posterior distribution we use the LAL-INFERENCENEST code (Veitch et al. 2015b). This is a software package for sampling posterior distributions of the parameters of compact-binary GW sources that is part of LAL. It uses nested sampling to explore the posterior distribution. Nested sampling (Skilling 2006) was originally introduced as an efficient way to compute the Bayesian evidence, as a tool for model selection, but also returns independent samples from the posterior distribution on the parameters. The algorithm evolves a set of live points, replacing the lowest likelihood point at each step by another point of higher likelihood chosen uniformly from the prior distribution. The LALINFERENCENEST code achieves these updates using short Markov-chain Monte-Carlo evolutions. We refer the reader to Veitch et al. (2015b) for further details of the implementation.

3. MEASUREMENT OF IMBH PROPERTIES




Figure 2. The 95% relative width in the measurement of the parameters $\Theta = \{M_c, M_{\text{tot}}, d_L\}$ defined by $\Delta_\Theta = \Delta\Theta^{95\%}/\Theta^{\text{inj}} \times 100$, where $\Delta\Theta^{95\%}$ is the 95% absolute width and Θ^{inj} is the true value of the parameter Θ . These results correspond to spinning, non-precessing BBH systems with $q = 4$, $\iota = \pi/3$ at $\text{SNR} = 20$. The different color lines represent the different values of component spins (with $\chi_1 = \chi_2$), as illustrated in the legend in the rightmost plot. We use **SEOBNRHM** waveforms for the injection and recovery. For $\chi_1 = \chi_2 = 0.95$ systems, the worsening in the accuracy of the chirp mass (left panel) for $M_{\text{tot}} \geq 450 M_\odot$ is due to the occurrence of bi-modality in the posterior of M_c .

3.1. Parameter space of simulations

To understand the uncertainty with which the parameters of GW signals from IMBHBs could be constrained with upcoming observations (Abbott et al. 2020a), we simulate a set of (synthetic) GW events and analyze them using Equation (2). For simplicity, the signals are simulated (injected) in a zero-noise background. The addition of noise is expected mainly to change the peak of the posteriors not the widths (or the uncertainties), which are our primary interest. Unless otherwise stated, we choose simulated signals that have LIGO-Virgo network SNR of 20. We restrict our analysis to high SNRs, mainly because we are looking for the best-constrained events given that these are the ones that will be most confidently identified as IMBHs.

We work with IMBHBs with mass ratios $q = 1.25$, $q = 4$ and $q = 10$. Based on the trend that we observe in the results, we expect that for any other mass ratio between them, the associated uncertainty is contained within the uncertainties of these three mass ratios. We fix the inclination angle to 60° (i.e., $\iota = \pi/3$) and 0° (face-on). We vary the (detector-frame) total mass in the range, $M_{\text{tot}} \in [50, 500] M_\odot$ with steps of $50 M_\odot$. Given that we still do not know very accurately the spin distribution for IMBHs, we choose a very wide range of values for the component spins, namely, $\chi_1 = \chi_2 = \{-0.8, -0.5, 0., 0.5, 0.8, 0.95\}$. For the same reason, we also explore binaries with opposite spins — for example, $\chi_1 = 0.5, \chi_2 = -0.5$. We find that the results are contained within the range of results set by the equally spinning binaries.

3.2. Results using Bayesian analysis

As a cross-check of our analysis, we start our study by reproducing the results of Graff et al. (2015). They fo-

cused on IMBHBs with (detector-frame) total masses in the range $[50, 500] M_\odot$, mass ratios $q = 1.25$ and $q = 4$, and $\text{SNR} = 12$ and used a version of the LIGO noise-spectral density at design sensitivity available at that time. They employed for the Bayesian analysis the multipolar non-spinning waveforms, **EOBNRHM** (Pan et al. 2011)⁵. Using our waveform model **SEOBNRHM** in the non-spinning limit, we could recover the results of Graff et al. 2015 with some small differences — for example, we find that the maximum discrepancy (i.e., the absolute difference between the estimated precisions) is 8% in M_c and ν for high total masses ($M_{\text{tot}} \gtrsim 300 M_\odot$), where the merger and ringdown phase of the signal dominates in the most sensitive frequency band of the detectors. These discrepancies are mainly due to differences between the waveform models. The waveform model (Cotesta et al. 2018, 2020) used in this work is more accurate than the one employed in Graff et al. (2015), since it was calibrated to a much larger set of NR simulations and contains more information from PN theory.

As described in Section 3.1, here we extend the study of Graff et al. (2015) in several directions. We consider multipolar spinning, non-precessing IMBHB systems with (detector-frame) total masses in the range $[50, 500] M_\odot$, but mass ratios up to 10 (i.e., $q \in [1.25, 10]$) and $\text{SNR} = 20$. We also use updated LIGO and Virgo PSDs, notably the ones for the upcoming O4 and O5 runs (Abbott et al. 2020a). In Figures 2, 3, 4 and 5 we summarize our results for mass ratios $q = 1.25$ and $q = 4$, inclination $\pi/3$, and a variety of spin values,

⁵ In LAL this waveform model is denoted **EOBNRv2HM** (Pan et al. 2011).




Figure 3. The 95% relative widths in the measurement of the parameters for the binaries with mass ratio, $q = 1.25$. The definitions are same as in Figure 2. For $\chi_1 = \chi_2 = 0.95$ systems, the worsening in the accuracy of the chirp mass, due to bi-modality, starts at $M_{\text{tot}} \sim 250M_\odot$.

$\chi_1 = \chi_2 = -0.8, -0.5, 0, 0.5, 0.8, 0.95$, while in Tables 1, 2 and 3, we provide results for zero inclination (i.e., face-on configuration) and for larger mass ratio (i.e., $q = 10$). Furthermore, we do not show the $\chi_1 = \chi_2 = -0.8$ results in our figures because, sometimes, they have much higher posterior widths compared to the other cases owing to their very small number of GW cycles in the detectors’ bandwidth. The results also depend on the specific prior choices on the luminosity distance — for example priors flat in d_L^2 or flat in the comoving volume produce noticeably different results for higher total masses. For all other spin configurations, we do not see any significant differences between the results of the two distance priors. In all figures and Tables we display results with a prior flat in comoving volume.

Figure 2 shows the 95% relative widths of the posteriors of different parameters of the binary systems with mass ratio $q = 4$ and several choices for the BH spins. It can be seen that all the systems at the lowest total mass (i.e., $M_{\text{tot}} = 50M_\odot$) measure chirp mass (M_c) better than the total mass. This is because the waveform from the systems at such a low total mass is dominated by its inspiral phase in the most sensitive frequency band of the detectors. At leading PN order, the phase of inspiral gravitational waveforms depends only on the chirp mass M_c (Sathyaprakash & Dhurandhar 1991). At higher total mass, the waveform is dominated by its post-inspiral phase (i.e., by the merger), which is better described by M_{tot} , and thus measures the total mass better (Graff et al. 2015). We find that the total mass at which M_{tot} starts to be measured better than M_c depends also on the spins of the components. We note that mergers of binaries with component spins aligned (anti-aligned) in the direction of the orbital angular momentum \mathbf{L} are delayed (accelerated) compared to their non-spinning counterparts (Campanelli et al. 2006), and thus, the binary remains in the inspiral phase

longer (shorter) than the non-spinning systems. For example, at $M_{\text{tot}} \sim 100M_\odot$, all aligned systems measure the chirp mass better than the total mass, whereas the anti-aligned systems (see, e.g., the magenta curve) measure M_{tot} better. For $M_{\text{tot}} \gtrsim 150M_\odot$ onward, independently of spins, all systems measure the total mass better than the chirp mass.

As can be seen from Figure 2, the precision of the chirp mass (M_c) measurement initially degrades as the total mass is increased, before starting to improve for sufficiently high masses. The initial increase in the uncertainties is because of the decrease in the number of GW cycles, as we increase the total mass of the systems. However, after a certain total mass (depending on the spins), the uncertainty starts to decrease as the merger-ringdown phase of the waveform starts to match well with the minimum of the PSDs, and also the subleading modes reach the most sensitive frequency range of the detector. The $\chi_1 = \chi_2 = 0.95$ systems, in particular, behave somewhat unexpectedly above $M_{\text{tot}} \gtrsim 450M_\odot$. We find that these systems have a bi-modal distribution in the parameters M_c , ν , $m_{1,2}$, and χ_2 , which cause the width of the posteriors to increase significantly. We shall discuss these features in more detail in Section 3.3 below.

Figure 3 shows the posterior relative widths for the parameters when the mass ratio of the binaries is $q = 1.25$. As with the $q = 4$ binaries, there is a trend for each spin, with a few exceptions. First, we find that systems with $\chi_1 = \chi_2 = 0.95$ show bi-modality even when the total mass is as low as $200M_\odot$, and continue doing so for higher total masses. The bi-modality in χ_2 appears throughout (i.e., even when the total mass is $\sim 100M_\odot$). Second, some systems show an unusual trend with total mass (e.g., see the total mass uncertainty plot in Figure 3). The uncertainty increases again after $350M_\odot$, where we would expect it to decrease. We find that




Figure 4. The 95% relative width in the measurement of the source-frame masses $\{m_1^s, m_2^s, M_{\text{tot}}^s\}$ of the injected signals from Figure 2 (i.e., for $q = 4$). The upper limits of the x -axes are restricted to the final source-frame masses associated with $\chi_1 = \chi_2 = 0.95$ IMBHB systems which, because of higher amplitude and hence higher redshift reach, provide the smallest source-frame mass for a given detector-frame mass (e.g., at $M_{\text{tot}} = 500M_\odot$). The shaded region represents the BH’s mass gap $[60, 130]M_\odot$ derived in Section 4.2 and computed at the median ($\sigma = 0$) of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate (see Figure 9). The dots represent the uncertainties when we inject spinning, precessing signals **NRSurPHM**, and recover them with spinning, non-precessing **SEOBNRHM** waveforms. The 95% uncertainty of each dot should be compared to the one of the curve with the same color at the same value of the source-frame parameter. The primary mass can be estimated with the precision ~ 15 –25% while the total mass can be estimated with a slightly better precision ~ 12 –22%.

for such parameter choices, there is a strong correlation between the total mass M_{tot} and the primary spin χ_1 . Additionally, χ_1 also develops a mild bi-modality.

As stressed before, we are mostly interested in characterizing the uncertainty in the measurements of source-frame masses, as they could help us understand the precision with which, e.g., the mass function of IMBHs can be constructed from GW measurements in upcoming observations, and also determine the probability that the observed BH’s mass is in the mass gap. This information can be extracted from Figures 4 and 5. The primary mass for all the systems with $q = 4$ can be measured with an uncertainty of ~ 17 –25%⁶ while for binaries with $q = 1.25$ the uncertainties are ~ 30 –40% except when $\chi_1 = \chi_2 > 0.8$ where bi-modality further worsens the precision, pushing the uncertainties to $\sim 60\%$. The improvement in the precision for $q = 4$ binaries is due to the presence of higher modes in the gravitational signal, which break the degeneracy among parameters and lead to a better measurement of m_1 and d_L measurement. For high mass-ratio ($q = 10$), the precision improves further, bringing the uncertainties down to ~ 11 –20% (see Table 3) except when $M_{\text{tot}} < 100M_\odot$ where we see a very strong correlation among the parameters M_{tot} , m_1 , m_2 and q . This is also true for even smaller mass ratios. We can see this from Figure 2 for $q = 4$, where, contrary to our expectation, the precision of the total-mass measurement at $50M_\odot$ is poorer than the one at $100M_\odot$. We observe a similar trend in the

measurement of the primary mass m_1 . The secondary mass, on the other hand, is measured with relatively poor precision: for symmetric systems ($q \sim 1$), the uncertainties lie between 30–60% except for binaries with $\chi_1 = \chi_2 \geq 0.8$, for which the uncertainty can reach 80%. For higher mass-ratio signals, the uncertainty can exceed 100%, except for a few highly spinning aligned systems (e.g., $\chi_1 = \chi_2 \geq 0.5$) where the uncertainty can reduce to ~ 10 –30% (see Table 3). The total source mass (M_{tot}^s), however, is measured with a much better precision, with uncertainties in the range ~ 10 –30%, regardless of the details of the signals.

From Figures 2 and 3, we also see that component spins can only be measured poorly. The primary spin (χ_1) can be measured better than $\sim 50\%$ only for asymmetric binaries which have high component spins, $\chi_1 = \chi_2 > 0.5$. Figure 3 shows that for nearly symmetric systems, unless the spins are $\chi_1 = \chi_2 > 0.8$, we might not be able to measure the primary component spin better than 50%. But, as expected, for very high mass-ratio signals, we can measure the primary spin with an uncertainty lower than $\sim 30\%$ if the systems are aligned (see Table 3). Measuring the secondary spin, however, seems to be difficult for almost all of the IMBH binaries.

To understand how much the neglect of spin-precession in our waveform model affects these precisions, we simulate a few moderately spinning, precessing GW signals with the **NRSurPHM** waveform model (Varma et al. 2019b), and analyze them with the spinning, non-precessing **SEOBNRPHM** model. Mild spin-precession is motivated from the LIGO-Virgo observations (Abbott et al. 2019a, 2020b), so far. More specifically, we fix the

⁶ Including the $\chi_1 = \chi_2 = -0.8$ results from the Table 2.




Figure 5. The 95% relative width in the measurement of the source-frame masses $\{m_1^s, m_2^s, M_{\text{tot}}^s\}$ of the injected signals from Figure 3 (i.e., for $q = 1.25$). The other definitions are same as Figure 4. Even for symmetric IMBHBs, except when spins are high ($\chi_1 = \chi_2 > 0.8$), for which bi-modality occurs, the primary mass (total mass) can still be estimated with a precision below 40% (30%).

angle between the primary spin χ_1 and the total angular momentum \mathbf{J} (i.e., the tilt angle) to be 30° except for anti-aligned binaries for which the tilt angle is taken to be 210° , while the magnitude of the component spin vectors are taken to be the same as their non-precessing counterparts. The other angles required to define the components of the spins on the orbital plane are taken to be zero. All of these quantities are defined at a reference frequency, that we choose to be $f_{\text{ref}} = 11$ Hz. These results are indicated by the small dots in Figures 4 and 5. We can see that the uncertainties in the component-mass measurement change only by $\lesssim 5\%$. The additional systematic bias introduced by the neglect of precession in the recovery model is also $\lesssim 5\%$. Thus, a mild precession in the signals is not expected to change the results established here significantly, as long as they are recovered with spinning, non-precessing waveforms. We plan in the future to carry out a comprehensive study that will analyze spinning, precessing GW signals with precessing waveforms.

We also produced results with the noise-spectral density projected for the O5 run (Abbott et al. 2020a), but at the same SNR=20, since we a priori do not know what the distribution of the observed SNRs is going to be. However, we do expect that during O5 a larger number of IMBHBs should be observed at a given SNR compared to O4, as also shown in Section 1. We find that, at O5 sensitivity and SNR = 20, the precision of detector-frame masses improves only by a few percent (e.g., for the chirp mass M_c it improves by $\sim 6\%$), while for the source-frame component masses, which we are mainly interested in, it remains mostly unchanged.

When considering also the results in Tables 1, 2 and 3, obtained at inclination angle of $\pi/3$ and mass ratio $q = 10$, and at zero inclination angle for mass ratios $q = 1, 4, 10$, we can summarize the main findings as fol-

lows. At inclination $\pi/3$, for quite asymmetric IMBHB systems (i.e., $q \sim 10$), we could measure the primary mass with uncertainties 11 – 25%, whereas for symmetric binaries ($q \sim 1$), the uncertainties are expected to be $\sim 30 - 40\%$ except for systems with component spins $\chi_1 = \chi_2 \gtrsim 0.8$, where the uncertainties can reach $\sim 60\%$ due to the presence of bi-modality. At zero inclination, however, the uncertainty in the primary mass can also go up to $\sim 60\%$ independently of the mass ratio (except for $\chi_{1,2} \gtrsim 0.80$ systems where the precision becomes even worse due to bi-modality). However, for the bulk of the parameter space, even at zero inclination, the uncertainty is $\lesssim 40\%$ for the primary mass, where this upper limit is set by the high-mass IMBHBs (e.g., $M_{\text{tot}} \sim 500M_\odot$). An important question in astrophysics concerns the nature of the IMBH mass function. Although upcoming LIGO and Virgo observations may not be capable of observing enough IMBHs to reconstruct their mass function, they hold the potential to provide us with much better mass measurements than what might be possible with electromagnetic observations. In Section 4.4, we shall study in more detail the implications of the component-mass measurements in assessing the BH upper-stellar-mass gap.

From Figures 2 and 3, we can see that measuring the luminosity distance (d_L) with precision better than 50% may not be possible at high inclination ($i = \pi/3$). On the other hand, Table 1 shows that symmetric face-on IMBH binaries, which emit GW signals with the highest amplitudes, can allow us to constrain d_L with uncertainty $< 50\%$. When the total mass and spins are high (e.g., $M_{\text{tot}} \gtrsim 400M_\odot$, $\chi_1 = \chi_2 \gtrsim 0.5$), we can even constrain the luminosity distance with uncertainties less than 40% (20%) at O4 (O5) sensitivity. Given the large detection horizon distance for IMBH binaries, these luminosity distance measurements could be valu-




Figure 6. Posterior distributions of the parameters for an injected signal with $M_{\text{tot}} = 500M_{\odot}$, $q = 1.25$, $\chi_1 = \chi_2 = 0.95$, $\iota = \pi/3$ at SNR = 20. The dashed lines show the posteriors when **SEOBNRHM** injected signal is recovered with 22 mode waveform model **SEOBNR**. In each panel the vertical dashed lines indicate the true (injected) value of the parameter. Both **SEOBNRHM** and **PhenomHM** models show bi-modality in various parameters (e.g., the component masses $m_{1,2}$). The **SEOBNR** model, which only contain the dominant (2,2) mode, hardly shows bi-modality in most of the parameters, but the posteriors peak away from the true (injected) values. Higher modes can, thus, be important even when the mass ratio is close to 1, but the spins and total mass are high.




Figure 7. Whitened (left panel) and unwhitened (right panel) strains, at the LIGO-Livingston detector, for the parameters of the two maximum-likelihood points of the injection study shown in Figure 6 (for **SEOBNRHM**-**SEOBNRHM** model), that is from the region $q \leq 2.5$ and $q > 2.5$. The scale of the y -axis is irrelevant here. Note that we shift the waveforms in time so that $t = 0$ represents the coalescence time for both waveforms. The two whitened waveforms match very well (left panel) even when the unwhitened waveforms differ before the coalescence (right panel), notably at earlier and earlier times.

able for statistical constraints on cosmological parameters (Schutz 1986; Abbott et al. 2021a), but it will depend on the rate of observed events.

3.3. Bi-modality in component masses and spins

As we discussed in the previous section, for sufficiently high total masses, highly spinning IMBHB systems (i.e., $\chi_1 = \chi_2 \gtrsim 0.8$) can exhibit bi-modality in the posterior distributions of some parameters. As an example, we show in Figure 6 the results obtained for an IMBHB system with $M_{\text{tot}} = 500M_{\odot}$, $q = 1.25$, $\chi_1 = \chi_2 = 0.95$, $\iota = \pi/3$ at SNR = 20. To better under-

stand the bimodality we inject and run the Bayesian analysis with two spinning, non-precessing waveform models: **SEOBNRHM** and **PhenomHM** (García-Quirós et al. 2020). For both models we observe bi-modal posterior distributions in the component masses $m_{1,2}^s$, the secondary component spin χ_2 , and the inclination angle ι .

To understand the results, we compare in the (left) right panel of Figure 7 the (whitened) waveforms for the **SEOBNRHM** model corresponding to the two maximum-likelihood points defined in the region $q \leq 2.5$ (low q) and $q > 2.5$ (high q). To obtain whitened waveforms in the time domain we first divide the waveform in the fre-

quency domain by $\sqrt{S_n(f)}$, and then we inverse Fourier transform them into the time domain. Whitening of the waveforms helps to better understand the matching of the signal with the waveform, because we can see from Equation (5) that the power spectral density $S_n(f)$ appears inverse weighted in the likelihood function.

From the left panel of Figure 7, we can see that there is a very good agreement between the two whitened waveforms even though the unwhitened waveforms, shown in the right panel of Figure 7, have differences at earlier and earlier times before coalescence. Thus the bi-modality appears to stem from a conspiracy: the total mass is very high and hence the number of GW cycles is already just a few within the detectors' bandwidth, and the early cycles of the signals, where they differ significantly, are being suppressed by the worsening of the PSDs⁷. We also compute overlaps using Equation (7) between the maximum likelihood signal and other points from the posterior samples. We find similar bi-modal behaviour in the distribution of the overlaps. This indicates that there are two points in the parameter space that have larger match, as we also see visually in the left panel of Figure 7. Moreover, when we recover the signal with just the SEOBNR model, which only contains the (2,2) mode, we find that it is the high q region which has the point with the highest posterior probability rather than the low q region, where actually the injection lies. The SNRs recovered with the SEOBNR model at the maximum likelihood points in these two regions are pretty close, 19.49 and 19.63, respectively. Note that the injected SNR is 20. Thus, almost all of the SNR is being recovered by the 22 mode waveform, yet the recovered posteriors are significantly different. This suggests that the inclusion of higher modes can matter for inferring the properties of the source (IMBHBs) even when the mass ratio is close to 1, but the spins and the total mass are high. In particular, the global maximum likelihood, is still recovered in the region around the injection (low q).

4. UPPER-STELLAR-MASS GAP

Before discussing the implications of our parameter-estimation study of IMBHB systems on the measurement of the BH's upper-stellar-mass gap, we first review the main results in the literature on this topic, and then perform, using updated $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rates, a new study aimed at establishing more robustly the uncertainties in the upper and lower edges of the mass gap.

⁷ However, an unresolved question is why bi-modality does not seem to occur when spins $\chi_{1,2} < 0.8$ at high total masses.

4.1. What is the mass gap?

As mentioned in Section 1, single stars with masses of $20 M_\odot \lesssim M_{\text{ZAMS}} \lesssim 100 M_\odot$ end their lives in core-collapse supernovae and are thought to exclusively form BHs (Timmes et al. 1996; Fryer & Kalogera 2001; Zhang et al. 2008; Sukhbold & Woosley 2014; Sukhbold et al. 2018). Stars with $M_{\text{ZAMS}} \gtrsim 100 M_\odot$ reach core temperatures of $\gtrsim 7 \times 10^8$ K that allows for the production of electron-positron pairs from photons, $\gamma + \gamma \rightarrow e^- + e^+$ (Fowler & Hoyle 1964; Barkat et al. 1967; Rakavy & Shaviv 1967). The production of $e^- e^+$ pairs removes photons, softening the equation of state. These stars are expected to become dynamically unstable before core-oxygen depletion, as the pair production leads to regions where the adiabatic index $\Gamma_1 \leq 4/3$ (Fraley 1968; Ober et al. 1983; Bond et al. 1984; Woosley et al. 2002; Heger et al. 2003; Takahashi 2018; Farmer et al. 2019; Marchant & Moriya 2020). The ensuing dynamical collapse results in vigorous oxygen burning whose outcome depends, in part, on the mass of the star and the adopted $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rate.

Stars with $100 M_\odot \lesssim M_{\text{ZAMS}} \lesssim 130 M_\odot$ can undergo a cyclic pattern of entering the pair instability region; contracting, undergoing oxygen burning, and expanding (Heger et al. 2003; Blinnikov 2010; Chatzopoulos & Wheeler 2012; Yoshida et al. 2016; Woosley 2017; Umeda et al. 2020). This process yields a series of pulsations that removes large amounts of mass from the star, leading to a pulsation-pair-instability supernova (PPISN) whose core collapse leaves significantly lower mass BHs. PPISN set the lower edge of the BH mass gap. Stars with $130 M_\odot \lesssim M_{\text{ZAMS}} \lesssim 250 M_\odot$ can produce a pair-instability supernova (PISN) where the energy injected from the first explosive oxygen-burning event completely unbinds the star (Joggerst & Whalen 2011; Chatzopoulos et al. 2013; Kozyreva et al. 2014; Gilmer et al. 2017; Marchant & Moriya 2020; Renzo et al. 2020). PISN leave no compact object, making them responsible for the existence of the BH mass gap. Stars with $M_{\text{ZAMS}} \gtrsim 250 M_\odot$ reach core temperatures of $\log(T_c/\text{K}) \approx 9.8$, where the rate of endothermic photodisintegration reactions absorbs enough energy to prevent the star from unbinding (Heger et al. 2003). The star, once again, can reach core collapse. This sets the upper boundary of the mass gap.

In addition to the ZAMS mass, the $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rate plays a central role in determining the final outcome by setting the C/O ratio in the core after helium burning. Takahashi (2018) found that cores with reduced $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ rates have larger C/O ratios, develop shell convection during central carbon burning, and sufficiently avoid the PI regime to collapse as BHs. Farmer

et al. (2020) found that cores with reduced $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ rates can have C/O ratios $\simeq 0.4$. These cores undergo a sequence of central carbon burning, off-center carbon burning, central oxygen burning, and core collapse to produce a BH. Cores with median $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ rates can have C/O ratios $\simeq 0.1$. These cores effectively skip central carbon burning to yield PPISN with smaller BH masses. Cores with large $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ rates can have C/O ratios $\ll 0.1$. These cores effectively skip central and shell carbon burning. They proceed directly to explosive oxygen burning and result in a PISN. Given this sensitivity, we thus undertake a new exploration of the $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rate and its uncertainties.

4.2. Updated $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rates

The C/O content of stellar cores is determined by the competition between triple- α to ^{12}C and $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ nuclear- reaction rates during helium burning (deBoer et al. 2017). Because of the complexity of the calculations, most analyses have only considered a subset of the reaction channels and a few representative data sets. In order to obtain a comprehensive evaluation, deBoer et al. (2017) considered the entirety of existing experimental data related to the determination of the low energy $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ cross sections, aggregating 60 year of experimental data consisting of more than 50 independent experimental studies. The more than 10,000 data points were then incorporated into a complete multi-channel phenomenological R -matrix analysis (Lane & Thomas 1958; Descouvemont & Baye 2010; Azuma et al. 2010) using the code AZURE2 (Azuma et al. 2010; Uberseder & deBoer 2015). A main result was the characterization of the uncertainty in the reaction rate, which was accomplished through a Monte-Carlo-uncertainty analysis of the data and the extrapolation to low energy using the R -matrix model. This resulted in a rate uncertainty that had statistical significance, which had only been accomplished in a few other previous works (e.g., Gialanella et al. 2001; Schürmann et al. 2012), and there with much more limited data sets. In deBoer et al. (2017), the 1σ uncertainty of the reaction rate was given after finding an approximately Gaussian underlying probability distribution for the rate.

In the recent work by Farmer et al. (2020), the main rate that was used was that of Kunz et al. (2002), but with the central value adjusted to be the geometric mean of the upper and lower 1σ uncertainty estimates, with the assumption that these 1σ values reflect an underlying probability distribution that is approximately Gaussian. This modified rate was implemented in the STARLIB rate library (Sallaska et al. 2013a). The work of deBoer et al. (2017) was very much in the spirit of that

of Kunz et al. (2002), but used a Monte-Carlo method to estimate the uncertainties (instead of χ^2) and had at its disposal the significantly increased amount of experimental measurements that had accrued in the intervening time. In particular, the significantly more stringent constraints for the values of the sub-threshold asymptotic normalization coefficients determined through sub-Coulomb transfer measurements (Brune et al. 1999; Avila et al. 2015) and the facilitation of them using the alternative R -matrix parameterization of Brune (2002).

To facilitate the present calculations, and future ones, we expand the tabulated reaction rate to a much finer temperature grid. This is done to ensure that no temperature step results in variations in the rate of more than an order of magnitude. The expanded reaction rates of deBoer et al. (2017) are shown in Figure 8, over a region of $\pm 3\sigma$. As shown previously by Farmer et al. (2020), the uncertainty present in this nuclear-reaction rate translate into one of the primary sources of uncertainty in the location of the BH mass gap boundaries.




Figure 8. Relative uncertainties in the $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rate of this work, expanded from those presented in (deBoer et al. 2017). The uncertainties are normalized to the central value for clearer presentation. The regions of fading blue color represent 0.5σ steps in the Gaussian uncertainty distribution.

4.3. The black-hole mass gap with updated $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rates

We use MESA version r11701 (Paxton et al. 2011, 2013, 2015, 2018, 2019) to evolve massive helium cores until they either collapse to form a BH or explode as a PISN without leaving a compact remnant. We use the same inlists and `run_star_extras.f` used in Farmer et al.

(2020) to calculate the boundaries of the BH mass gap with respect to the updated $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction-rate uncertainties.

Figure 9 shows the location of the PISN BH mass gap as a function of the uncertainty in the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate. As the reaction rate increases, through increasing $\sigma[^{12}\text{C}(\alpha, \gamma)^{16}\text{O}]$, both the lower and upper edges of the BH mass gap shift to lower masses while maintaining a roughly constant width, of $\simeq 80^{+9}_{-5} M_{\odot}$. For the updated $\sigma[^{12}\text{C}(\alpha, \gamma)^{16}\text{O}]$ rates adopted in this work, the location of the lower and upper edge over the $\pm 3\sigma$ range is $\simeq 59^{+34}_{-13} M_{\odot}$ and $\simeq 139^{+30}_{-14} M_{\odot}$ respectively. These results are commensurate with Farmer et al. (2020) at the $\simeq 20\%$ level for the lower edge of the BH mass gap and at the $\simeq 5\%$ for the upper edge of the BH mass gap. We next discuss the main reasons why our results slightly differ and put them in context with previous studies.

Marchant & Moriya (2020) found that the efficiency of angular-momentum transport changes the lower edge of the BH’s mass gap at the $\simeq 10\%$ level. Farmer et al. (2019) found that the lower edge of mass gap was robust at the $\simeq 10\%$ level to changes in the metallicity, wind mass loss prescription, and treatment of chemical mixing. We find that the primary uncertainty in our models is the mass and time resolution. To ensure the results shown in Figure 9 are robust at the $\simeq 10\%$ level, our models use 2 times the mass resolution and about 2.5 times the temporal resolution as those used in Farmer et al. (2020). For each $\sigma[^{12}\text{C}(\alpha, \gamma)^{16}\text{O}]$, our $\Delta M = 1 M_{\odot}$ mass grid of MESA models consumed $\simeq 60,000$ core-hours, with Figure 9 thus costing $\simeq 780,000$ core-hours.

We also calculate new $\pm 3\sigma$ rates for the $^{12}\text{C} + ^{12}\text{C}$, $^{12}\text{C} + ^{16}\text{O}$, $^{16}\text{O} + ^{16}\text{O}$ reactions. Consistent with Farmer et al. (2020), we find that these reaction rates move the the BH’s mass-gap boundary by $\lesssim 1 M_{\odot}$. Evidently, in this case, the total energy liberated by C-burning is more important than how quickly or slowly the C-burning energy is liberated. A large carbon fuel reservoir from a small $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate leads to a more massive BH, an intermediate carbon fuel reservoir from the recommended (i.e., median) $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate leads to less massive BHs, and a small carbon-mass fraction from a large $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate leads to no compact object being formed (i.e., a PISN).

4.4. Measurement of mass gap with upcoming LIGO-Virgo observations

Figure 9 shows that at the median ($\sigma = 0$) of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate, the mass gap would typically fall in the range $60\text{--}130 M_{\odot}$ considering the overlapping parts of the range from the two $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rates. Including such a range in Figures 4 and 5 (see shaded gray




Figure 9. The location of the BH’s mass gap as a function of the temperature-dependent uncertainty in the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate. The blue lines mark the mass-gap boundaries predicted by our updated $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate uncertainties. The orange lines mark the mass-gap boundaries, as found in Figure 5 of Farmer et al. (2020), predicted by the Kunz et al. (2002) rate as expressed in the STARLIB reaction-rate library (Sallaska et al. 2013b). The white region denotes the mass gap, the purple regions highlight differences of the adopted $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rates, and the labeled gray horizontal bars denote the mass range where a BH does not form for any value of the adopted $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate.

region), we find that the uncertainties in the primary-mass measurements for asymmetric IMBHB systems are $\sim 17\text{--}25\%$, while for nearly symmetric systems they are $\sim 30\text{--}50\%$. For the latter systems, the secondary mass also quite often falls in the mass gap, and can be constrained with an uncertainty of $\sim 40\text{--}70\%$.

How confidently the upcoming O4 run will be able to identify that the component masses of IMBHB systems lie in the mass gap? To address this question we need to account for the uncertainties in the boundaries of the mass gap shown in Figure 9, which are caused by uncertainties in the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate. The probability that a component mass of the IMBHB system lies in the mass gap can be computed as:

$$P(\text{MG}) = P(m_{\text{MG}}^{\text{L}} < m_i^s < m_{\text{MG}}^{\text{U}}), \\ = \int P(x) dx \int_{m_{\text{MG}}^{\text{L}}(x)}^{m_{\text{MG}}^{\text{U}}(x)} P(m_i^s | d) dm_i^s, \quad (12)$$

where $i = 1, 2$, x denotes the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate, m_{MG}^{L} and m_{MG}^{U} denote the lower and upper edges of the mass gap and d represents the data (i.e., the simulated GW signal). We can approximate the above equation as a




Figure 10. The source-frame component masses and their associated 90% confidence intervals of all events from GWTC-1 and GWTC-2 with the restriction that the mean estimated mass of the primary is $\geq 10M_\odot$. The red region shows the mass gap at the $\sigma = 0$ for the updated $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate (see Figure 8). The primary mass of the GW190521 event along with its associated 90% confidence interval lies well inside the red region indicating that this could be a BH in the mass gap. Our re-analysis of GW190521 in Section 4.5, confirms this result with the waveform models used in this work. We note that there are also 5 more component masses, including the secondary mass of GW190521, whose mean lie in the mass gap.

discrete sum in x :

$$P(\text{MG}) = P(m_{\text{MG}}^{\text{L}} < m_i^s < m_{\text{MG}}^{\text{U}}), \\ = \sum_{j=1}^{N=13} P(x_j) \int_{m_{\text{MG}}^{\text{L}}(x_j)}^{m_{\text{MG}}^{\text{U}}(x_j)} P(m_i^s | d) dm_i^s, \quad (13)$$

where j runs from 1 to the 13 grid points of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate shown in Figure 9 and $P(x_j)$ is the corresponding probability⁸.

If $P(\text{MG})$ is large, then the source could be a mass-gap event. However, events that are not in the mass gap will occasionally have large $P(\text{MG})$ simply due to noise fluctuations. Thus, the probability necessary to claim a confident detection of a mass-gap event depends on the relative rate of mass-gap and no-mass-gap events. To test the hypothesis that there are events with components in the mass gap, we should compute the Bayesian evidence for the no-mass-gap versus mass-gap hypotheses, marginalized over the uncertain proportion of events that are in the mass gap. This approach leverages information from all events, not just those with high $P(\text{MG})$, and therefore has greater statistical power. However, such an analysis is beyond the scope of the current paper. Here we will only compute $P(\text{MG})$ for noise-free

data with a variety of injected IMBHB systems. Signals with high $P(\text{MG})$ are more likely to be robustly identified as mass-gap events, but we warn the reader against over-interpreting our numbers.

We show in Figure 11 $P(\text{MG})$ for IMBHB systems with primary mass $m_1^s = [70, 80, 90, 110, 120]M_\odot$ and several spin values, observed at inclination angle of $\pi/3$, computed using the updated $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate. As we can see, for asymmetric IMBHB systems (i.e., $q > 1.25$), if the primary mass is well within the median mass gap (e.g., $m_1^s \in [80, 120]M_\odot$), then the primary mass has probability of being in the mass gap $> 95\%$ (i.e., a single observation could be sufficient to robustly identify the existence of sources in the mass gap). For systems with primary mass close to the lower edge of the median mass gap (e.g., $m_1^s \sim 70M_\odot$), $P(\text{MG})$ reduces to ~ 0.85 for anti-aligned systems. For nearly symmetric IMBHB systems (i.e., $q = 1.25$), $P(\text{MG}) > 95\%$ for systems with somewhat higher primary mass, namely, $m_1^s \gtrsim 85M_\odot$. Symmetric systems can also have high probability that the secondary mass lies in the mass gap. The right panel in Figure 11 indicates that for IMBHB systems with $m_2^s \in [90, 120]M_\odot$, the posterior probability that the secondary mass lies within the mass gap exceeds 90% for systems with $\chi_{1,2} < 0.80$. By contrast, high spin systems with $\chi_{1,2} > 0.80$ exhibit bi-modality which worsens the precision of the mass measurement, as discussed in Section 3.3.

For zero inclination-angle (face-on) IMBHBs, our study shows that $P(\text{MG})$ is generally lower for measure-

⁸ We compute these probabilities from the standard normal distribution, evaluated at the given σ uncertainty and then normalize them so that they add up to 1. Alternatively, one could interpolate between these grid points and make the approximation in Equation 13 more and more precise by increasing N , but we do not expect that the results will change very significantly.




Figure 11. Probability of being in the BH’s mass gap (see Equation (13)) as function of source-frame component masses, for several IMBHB systems with primary mass $m_1^s = [70, 80, 90, 110, 120]M_\odot$, different spin values (as illustrated in the legend in the left panel), at inclination angle of $\pi/3$ and SNR of 20. Results are obtained using the spinning, nonprecessing **SEOBNRHM** model. Independently of the mass ratio, when IMBHBs have primary mass in the range $[80, 120]M_\odot$, the signal can be identified as a BH in the mass gap with probability $> 90\%$.

ments of the primary mass, reaching values as low as ~ 0.6 in some cases (e.g., at the upper edge of the mass gap $\sim 120M_\odot$, where bi-modality occurs). Nonetheless, systems with $m_1^s \in [80, 110]M_\odot$, would still yield $P(\text{MG})$ exceeding ~ 0.8 . Measurements of the secondary mass only provide $P(\text{MG}) > 0.8$ when the binary system has $m_2^s > 90M_\odot$.

4.5. Re-analysis of GW190521

As discussed in Section 1, [Abbott et al. \(2020d\)](#) reported the detection of the first IMBHB system, namely, GW190521. Using spinning, precessing waveform models, the primary mass of GW190521 was estimated to lie in the mass gap $[65, 120]M_\odot$ with probability above 99%, i.e., $P(\text{MG}) > 0.99$ (see this event in the Figure 10), when using the **NRSurPHM** model. When taking into account the uncertainty in the mass gap boundaries itself (see Equation 13), we still find that the probability is above 98%. Recently, [Nitz & Capano \(2021\)](#), using the spinning, precessing waveform model **PhenomPHM** ([Pratten et al. 2020](#))⁹, which was not employed in [Abbott et al. \(2020d\)](#), re-analysed GW190521. They extended the priors on the mass ratio, observed multi-modality in the component masses, and also found that the maximum likelihood parameter lies at high mass ratio, $q \sim 10$ indicating that the event might be an intermediate mass-ratio inspiral. Using their public data, namely, the results with the prior flat in the source-frame component masses, we find that their results would lead to a probability that the primary mass of GW190521 is in the mass gap of 55%. We shall comment again on the results of [Nitz & Capano \(2021\)](#) at the end of this section.

Here, we re-analyze GW190521, but mainly with the spinning, non-precessing **SEOBNRHM** and **PhenomHM** waveform models used in this work. We employ the same settings as used for the results publicly released by LIGO and Virgo ([GWOSC 2020](#); [Abbott et al. 2021b](#); [Vallisneri et al. 2015](#)), except for two modifications: i) we extend the prior ranges in the component masses and the mass ratio; and ii) we use a luminosity-distance prior uniform in comoving volume (see Equation 10). We show the results in Figure 12. With non-precessing waveforms, we find bi-modality in the posterior distributions of some parameters — for example, in the secondary-component mass (m_2), the luminosity distance (d_L), and the total mass (M_{tot}). However, the posteriors of the source-frame component masses ($m_{1,2}^s$) do not show bi-modality. Our non-precessing analysis also shows that irrespective of the waveform used, the posterior probability for the primary mass to lie in the mass gap is $\sim 99\%$, while the posterior probability that the secondary mass lies in the mass gap is $\sim 52\%$.

Furthermore, to contrast the results from the non-precessing waveforms to the precessing case, we analyze GW190521 with the **PhenomPHM** model. We note that after the paper [Nitz & Capano \(2021\)](#) came out, it was realized that the **PhenomPHM** model had an issue in modeling properly the merger-ringdown waveforms of spinning BBHs when the primary-spin is (close to) anti-aligned with the orbital angular momentum at merger. Here, we first perform our analysis with the same version of the **PhenomPHM** model used by [Nitz & Capano \(2021\)](#), and find full agreement with their results. Then, we re-analyze GW190521 with the new publicly released version of **PhenomPHM**. We display the results in Figure 12. As it can be seen, the posteriors with the precessing **PhenomPHM** waveforms show an additional peak, though small, at, e.g., $q \sim 5$ in the mass

⁹ In LAL this waveform model is denoted **IMRPhenomXPHM** ([Pratten et al. 2020](#)).




Figure 12. Posterior distributions of the parameters for the GW190521 event observed by LIGO and Virgo detectors (Abbott et al. 2020d,e). The parameter M_f^s and a_f denote the mass and spin of the remnant BH, while the other parameters have been introduced in Section 2.1. The vertical dashed lines in each plot indicate the 90% confidence interval for each posterior shown by the same color. The shaded region represents the BH’s mass gap $[60, 130]M_\odot$ derived in Section 4.2 and computed at the median of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate (see Figure 9). The non-precessing SEOBNRHM and PhenomHM models do not show bi-modality in the source-frame masses $m_{1,2}^s$, while the precessing PhenomPHM model shows additional small bumps in the secondary-mass posterior at $m_2^s \sim 30M_\odot$ and $m_2^s \sim 110M_\odot$. Except for the latter, there is good agreement between the results of non-precessing and precessing waveforms, and, in all cases, the 90% confidence interval of the primary-mass posterior lies inside the mass gap.

ratio. The secondary source-frame mass posterior, however, has two additional small bumps at $m_2^s \sim 30M_\odot$ and $m_2^s \sim 110M_\odot$. Besides those differences, the agreement between the posteriors from the precessing and non-precessing waveforms is quite good. With the precessing PhenomPHM model in Figure 12, we find that $P(\text{MG})$ for the primary mass is $\sim 98\%$, while for the secondary mass it is $\sim 54\%$. To further understand the robustness of these findings, we are currently finalizing a comprehensive analysis with other waveform models with spin-precession, namely, the time-domain spinning, precessing model from the EOB family (Ossokine et al. 2020). Moreover, such an analysis has also been carried out with a new time-domain phenomenological IMR model in Estelles (2021).

5. CONCLUSIONS

In this paper, we used the spinning, non-precessing SEOBNRHM waveform model to estimate the precision with which the parameters of non-precessing IMBHBs could be estimated in the upcoming LIGO-Virgo O4 and O5 runs. We simulated IMBHBs for total detector-frame masses in the range $M_{\text{tot}} = [50, 500]M_\odot$, mass ratios $q = \{1.25, 4, 10\}$ and component spins $\chi_1 = \chi_2 = [-0.8, 0.95]$ at a SNR of 20. We showed that for high mass-ratio binaries with relatively high inclination ($\iota = \pi/3$), the mass of the heavier component (i.e., the primary BH) can be constrained with an uncertainty $\sim 11 - 25\%$. These precisions are much better than

what is expected from electromagnetic observations. We also showed that the total source mass of IMBHBs can be constrained with uncertainties $\sim 10 - 30\%$, independently of the parameters of the IMBHB systems. These results suggest that future LIGO and Virgo observations hold the potential to measure the mass function of IMBHs and IMBHBs, which remains an important open question in astrophysics.

We also focused on IMBHB systems whose component BHs fall in the upper-stellar-mass gap, which is predicted from stellar-evolution theory for massive stars. We first studied the sensitivity of the mass-gap edges to the uncertainties in the most relevant nuclear rates, such as $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$, $^{12}\text{C} + ^{12}\text{C}$, and $^{16}\text{O} + ^{16}\text{O}$, using MESA. We confirmed the results of Farmer et al. (2020) that the boundaries of the mass gap are dependent on the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate, e.g., the lower edge of the mass gap can vary between $\sim 40 - 90M_\odot$ while the upper edge can lie anywhere between $\sim 125 - 170M_\odot$. The other nuclear rates, $^{12}\text{C} + ^{12}\text{C}$, $^{12}\text{C} + ^{16}\text{O}$, and $^{16}\text{O} + ^{16}\text{O}$ move the BH mass gap boundary only by $\lesssim 1M_\odot$. The main difference between our results and Farmer et al. (2020) is at the $\simeq 20\%$ level for the lower edge of the BH mass gap and at the $\simeq 5\%$ level for the upper edge of the BH mass gap. This is primarily due to an increased time resolution in our calculations.

Having updated the boundaries of the mass gap, we analyzed a few IMBHB systems with component masses

in the most probable range of the mass gap, and computed the posterior probability of them being in the mass gap based on their single-event observations. We found that for asymmetric inclined IMBHs whose primary component mass m_1^s lies in the range $[80, 120]M_\odot$, the posterior probability, $P(\text{MG})$, that the primary mass lies in the mass gap is $\gtrsim 95\%$. For symmetric inclined IMBHs, the same holds when $m_1^s \in [85, 120]M_\odot$. Lowering the inclination reduces the precision of the mass measurement and hence the posterior probability of being in the mass gap reduces. However, face-on IMBHs with $m_1^s \in [80, 110]M_\odot$, would still yield $P(\text{MG})$ exceeding ~ 0.8 ; systems with $m_1^s \sim 120M_\odot$ and high spins fall within the parameter space where bimodality occurs (see section 3.3) and hence the precision (width) of the mass posteriors decreases (increases). The secondary mass can also fall in the mass gap especially when the mass ratio of the IMBHs is near one ($q \sim 1$). However, since the measurement of the secondary mass is relatively poor, $P(\text{MG}) > 0.9$ can be achieved only when the secondary mass lies far above the lower edge of the mass gap, $m_2^s \in [90, 120]M_\odot$, given also that the inclination is higher. Within this context, we re-analysed the GW190521 event with spinning, non-precessing waveforms **SEOBNRHM** and **PhenomHM** and also with the latest version of (frequency-domain) precessing **PhenomPHM** waveform. We found that there is a very good agreement between the results of non-precessing and precessing waveforms, at least for the source frame component masses, which we are interested in. However, the precessing **PhenomPHM** waveform shows two additional, though really small, bumps in the secondary source-frame component mass (m_2^s). We showed that the $P(\text{MG})$ for the primary mass (m_1^s) with the non-precessing waveforms is $\sim 99\%$ while with the precessing waveform it still is $\sim 98\%$. The $P(\text{MG})$ for the secondary mass is $\sim 52\%$ with the non-precessing waveforms while with the precessing waveform it increases a bit, $\sim 54\%$ because one of the additional bumps occurs in the mass gap. To complete the re-analysis of GW190521 with state-of-the-art precessing waveforms, we are further investigating this event with more accurate (when compared to numerical-relativity simulations) waveforms, notably the time-domain spinning, precessing model from the EOB family (Ossokine et al. 2020) (see also Estelles (2021), which uses a new time-domain phenomenological model). We further note that the posterior probabilities were obtained using our default priors (see Section 2.2.2), notably priors flat in detector-frame component masses and uniform in co-moving volume for the luminosity distance. Employing different priors may change the results.

Furthermore, the probability $P(\text{MG})$ is a guide to how confidently an event can be identified as being a mass-gap event, but the number should not be over-interpreted, as noise fluctuations can lead to large $P(\text{MG})$'s for events that are not in the mass gap. The value required for an event to be confidently identified as being a mass gap event depends on the relative, and unknown, rates of events inside and outside the mass gap. To properly address the question of how many events would be required for a robust identification of a mass-gap population, would require a Bayesian model comparison between a “mass-gap” model and a “no mass-gap model”, using all observed events. We leave this work for future studies. However, systems with high $P(\text{MG})$ are more likely to be robustly identified as mass-gap events.

The inference studies in this work were limited to multipolar spinning, non-precessing quasi-circular waveform models, although we performed an analysis where we injected mildly precessing quasi-circular signals and found that the measurement uncertainties on the component masses only changed by $\sim 5\%$ when recovering those signals with non-precessing waveforms. In view also of possible multi-modal posterior distributions for IMBH systems, we plan in the future to carry out a comprehensive investigation using a larger set of injections of spinning, precessing signals and recover them with precessing (instead of non-precessing) waveforms. Furthermore, it will be crucial to extend the parameter-estimation study to IMBHs moving on eccentric orbits. Indeed, given the shortness of IMBH signals, all physical effects need to be included to avoid misinterpreting the properties of the source (Calderón Bustillo et al. 2020; Romero-Shaw et al. 2020). Recently, a few examples of multipolar spinning, non-precessing waveform models with mild eccentricity have been developed (Nagar et al. 2021; Liu et al. 2021; Yun et al. 2021; Khalil et al. 2021) that could be used for such studies.

ACKNOWLEDGMENTS

We are grateful to Andrew Matas for providing us with comments on this manuscript. We thank Rob Farmer for sharing his MESA version r11701 materials and his insights on our mass gap models. We also thank Pablo Marchant for useful discussions of PISN. The MESA project is supported by the National Science Foundation (NSF) under the Software Infrastructure for Sustained Innovation program grants (ACI-1663684, ACI-1663688, ACI-1663696). This research was also supported by the NSF under grant PHY-1430152 for the Physics Frontier Center “Joint Institute for Nuclear Astrophysics - Center for the Evolution of the Elements” (JINA-CEE). A.T. is a Research Associate at the Belgian Scientific Research Fund (F.R.S-FNRS).

This research has made use of data, software and/or web tools obtained from the Gravitational Wave Open Science Center (<https://www.gw-openscience.org>), a service of LIGO Laboratory, the LIGO Scientific Collaboration and the Virgo Collaboration. LIGO is funded by the U.S. National Science Foundation. Virgo is funded by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale della Fisica Nucleare (INFN) and the Dutch Nikhef, with contributions by Polish and Hungarian institutes. The authors are grateful for computational resources at the AEI, specifically the Hypatia cluster where all the the computations were carried out. The research of R.J.D utilized resources from the Notre Dame Center for Research Computing. This research made extensive use of the SAO/NASA Astrophysics Data System (ADS). This material is based upon work supported by NSF’s LIGO Laboratory which is a major facility fully funded by the National Science Foundation.

Software: AZURE2 (Azuma et al. 2010; Uberseder & deBoer 2015, <http://azure.nd.edu>), MESA (Paxton et al. 2011, 2013, 2015, 2018, 2019, <http://mesa.sourceforge.net>), MESASDK 20190830 (Townsend 2019a,b), matplotlib (Hunter 2007), and NumPy (van der Walt et al. 2011).

Table 1. The 95% width of one-dimensional marginalized posteriors scaled by the true value for injections and recoveries with SEOBNRHM waveforms. For all runs, $q = 1.25$, $\iota = \pi/3$ and $\text{SNR} = 20$. The spin column represents the equal component spins (i.e., $\chi_1 = \chi_2$) along the orbital angular momentum \mathbf{L} . The results in the parenthesis denote the corresponding face-on case (i.e., $\iota = 0$).

Spin	$M_{\text{tot}} = 50M_{\odot}$	$100M_{\odot}$	$150M_{\odot}$	$200M_{\odot}$	$250M_{\odot}$	$300M_{\odot}$	$400M_{\odot}$	$500M_{\odot}$
$\Delta M_c/M_c$								
-0.80	6.53 (6.69)	13.76 (14.85)	23.65 (25.91)	38.73 (38.79)	44.03 (40.90)	47.66 (44.36)	43.84 (39.59)	35.97 (36.33)
-0.50	6.08 (6.11)	21.45 (20.69)	22.99 (23.35)	24.03 (26.60)	27.97 (30.15)	28.52 (32.96)	22.75 (34.83)	18.11 (31.63)
0.00	4.20 (4.18)	16.02 (16.33)	20.32 (20.85)	20.22 (22.43)	21.07 (22.91)	23.56 (25.01)	23.64 (25.95)	21.00 (26.02)
0.50	3.33 (3.30)	13.43 (13.44)	20.55 (20.62)	23.49 (23.58)	23.93 (24.49)	22.59 (23.68)	22.52 (23.53)	23.66 (24.29)
0.80	3.08 (3.02)	11.25 (10.95)	19.18 (18.55)	22.26 (22.20)	23.64 (24.55)	24.26 (25.92)	21.96 (25.76)	19.83 (25.99)
0.95	2.34 (2.33)	8.49 (8.56)	16.37 (16.82)	20.90 (22.05)	24.29 (25.20)	26.79 (26.26)	30.81 (30.68)	34.79 (40.10)
$\Delta \nu/\nu$								
-0.80	13.99 (14.95)	8.43 (7.59)	17.36 (17.43)	30.87 (26.84)	34.16 (29.24)	39.47 (32.44)	36.78 (28.54)	27.04 (25.92)
-0.50	12.33 (12.60)	8.02 (7.42)	9.50 (10.35)	11.41 (15.06)	15.56 (18.93)	17.12 (22.32)	9.05 (24.82)	4.48 (24.23)
0.00	8.87 (8.50)	9.14 (8.04)	11.67 (11.21)	12.77 (13.82)	13.59 (14.68)	14.86 (17.10)	12.12 (17.61)	7.49 (17.78)
0.50	10.59 (10.10)	10.28 (9.98)	14.09 (12.99)	16.75 (16.13)	18.04 (17.57)	17.50 (17.58)	15.28 (17.45)	13.99 (17.17)
0.80	12.37 (11.36)	12.23 (11.90)	15.76 (15.13)	18.34 (17.92)	20.63 (21.55)	22.24 (23.61)	22.50 (26.68)	19.50 (27.49)
0.95	13.13 (12.36)	12.22 (11.84)	17.53 (18.10)	21.45 (21.85)	24.48 (25.48)	28.69 (28.25)	35.40 (35.79)	41.52 (49.53)
$\Delta d_L/d_L$								
-0.80	109.24 (48.12)	124.30 (51.67)	114.53 (51.11)	114.46 (63.14)	125.25 (66.61)	127.59 (71.34)	121.86 (66.87)	117.86 (68.45)
-0.50	103.65 (48.38)	129.83 (54.84)	131.48 (51.71)	130.28 (52.81)	135.20 (54.66)	136.64 (55.23)	124.22 (56.76)	109.45 (57.42)
0.00	99.20 (49.37)	113.16 (50.23)	114.13 (48.00)	112.25 (47.21)	108.44 (45.96)	110.75 (47.10)	107.76 (46.74)	97.85 (47.04)
0.50	103.31 (48.77)	110.10 (49.15)	110.33 (47.05)	106.26 (45.79)	104.54 (46.31)	104.99 (44.27)	102.17 (45.73)	97.40 (44.49)
0.80	103.08 (49.66)	105.15 (47.99)	103.31 (44.34)	100.17 (42.86)	104.40 (42.41)	101.70 (42.34)	103.96 (42.39)	100.91 (42.72)
0.95	102.05 (47.09)	99.08 (44.33)	88.38 (36.40)	85.40 (36.85)	88.10 (39.02)	88.58 (39.61)	88.39 (42.79)	91.10 (51.23)
$\Delta m_1^s/m_1^s$								
-0.80	40.98 (45.62)	31.75 (30.72)	35.04 (39.82)	39.54 (50.16)	42.04 (54.91)	49.19 (60.53)	48.16 (59.19)	38.07 (58.29)
-0.50	37.89 (41.78)	29.41 (30.81)	30.78 (33.11)	31.96 (38.60)	35.49 (45.52)	36.42 (49.73)	33.02 (53.81)	28.95 (55.42)
0.00	30.70 (32.47)	29.80 (30.15)	30.29 (32.31)	31.57 (35.59)	31.46 (37.38)	31.79 (40.52)	28.29 (43.06)	23.98 (44.56)
0.50	34.68 (36.27)	32.27 (34.38)	33.18 (36.65)	34.18 (38.88)	34.82 (41.31)	35.44 (42.00)	33.88 (43.33)	28.58 (43.98)
0.80	37.22 (38.46)	34.15 (37.63)	34.39 (39.69)	34.34 (42.72)	36.02 (47.98)	37.41 (51.04)	38.86 (57.85)	37.79 (61.08)
0.95	37.64 (39.72)	33.05 (36.85)	32.69 (43.14)	35.02 (46.72)	37.59 (52.18)	40.99 (58.19)	48.21 (71.12)	54.54 (92.24)
$\Delta m_2^s/m_2^s$								
-0.80	37.94 (41.76)	37.66 (39.26)	52.09 (56.24)	63.16 (66.60)	66.53 (68.92)	72.07 (71.17)	74.50 (66.15)	69.50 (62.66)
-0.50	33.88 (37.77)	40.83 (42.93)	45.74 (48.60)	49.70 (55.92)	56.81 (60.14)	59.23 (62.33)	49.08 (62.41)	36.17 (59.93)
0.00	29.69 (34.13)	38.23 (40.26)	45.75 (48.84)	47.74 (51.72)	50.02 (52.34)	53.09 (54.47)	50.45 (53.06)	43.21 (51.57)
0.50	32.82 (36.51)	37.25 (41.91)	45.75 (49.90)	50.89 (55.81)	52.98 (57.25)	53.51 (55.19)	51.94 (53.01)	50.32 (51.12)
0.80	35.37 (38.93)	38.37 (44.42)	45.55 (51.92)	50.55 (56.62)	55.09 (60.72)	57.34 (61.86)	59.69 (63.35)	58.78 (62.21)
0.95	34.17 (37.98)	35.93 (42.42)	41.92 (48.82)	48.28 (54.91)	54.21 (61.09)	58.86 (64.34)	65.04 (69.39)	70.40 (80.83)
$\Delta M_{\text{tot}}^s/M_{\text{tot}}^s$								
-0.80	12.96 (12.72)	19.25 (16.84)	21.39 (17.91)	22.59 (19.26)	24.02 (19.86)	24.56 (19.20)	26.05 (18.56)	25.47 (18.92)
-0.50	12.53 (12.35)	20.71 (19.89)	23.45 (20.43)	24.91 (20.24)	26.94 (20.46)	27.89 (19.28)	28.43 (18.05)	25.42 (18.20)
0.00	11.54 (11.66)	19.06 (17.57)	22.15 (19.70)	22.99 (19.14)	24.28 (18.83)	24.71 (17.86)	24.89 (17.10)	23.40 (16.57)
0.50	13.03 (12.68)	18.43 (17.24)	21.77 (19.41)	23.44 (19.16)	24.66 (19.09)	26.28 (17.62)	26.56 (17.07)	25.75 (16.66)
0.80	13.57 (13.25)	18.27 (17.51)	21.49 (18.76)	23.30 (19.23)	25.58 (19.08)	26.69 (17.83)	29.45 (17.68)	30.97 (17.81)
0.95	13.72 (13.18)	17.70 (16.37)	18.37 (15.27)	19.96 (15.18)	22.76 (16.32)	23.48 (16.41)	25.38 (18.74)	27.09 (24.16)
$\Delta \chi_1/\chi_1$								
-0.80	57.79 (59.56)	62.54 (68.11)	69.60 (84.77)	113.42 (119.57)	125.66 (125.30)	134.07 (124.58)	142.31 (131.19)	145.65 (134.80)
-0.50	167.26 (167.97)	190.27 (193.72)	191.59 (192.84)	200.30 (202.10)	203.99 (203.28)	212.54 (207.80)	206.81 (211.35)	199.01 (222.16)
0.50	142.76 (146.91)	161.14 (165.53)	155.77 (170.61)	164.73 (171.53)	171.11 (176.87)	176.26 (180.27)	188.26 (198.70)	187.56 (201.88)
0.80	38.25 (38.19)	48.82 (50.35)	53.32 (54.15)	54.50 (54.87)	57.82 (55.19)	59.03 (57.63)	65.26 (65.47)	71.83 (70.79)
0.95	10.42 (10.35)	17.02 (17.86)	20.70 (21.17)	21.32 (22.72)	21.71 (23.13)	21.20 (22.56)	19.83 (23.97)	20.31 (27.11)
$\Delta \chi_2/\chi_2$								
-0.80	100.77 (114.01)	79.72 (83.05)	117.00 (118.74)	143.12 (145.64)	150.22 (156.60)	163.02 (158.71)	167.28 (161.52)	156.74 (166.48)
-0.50	207.89 (208.63)	203.40 (200.99)	212.48 (213.34)	220.37 (227.68)	242.12 (244.35)	248.53 (252.90)	255.91 (260.95)	249.87 (258.30)
0.50	207.34 (208.59)	222.72 (218.71)	225.93 (225.32)	229.50 (225.50)	234.26 (224.77)	232.81 (231.76)	230.68 (237.31)	234.28 (237.26)
0.80	98.71 (90.01)	118.82 (117.89)	135.50 (131.24)	138.07 (136.65)	138.26 (149.40)	140.59 (137.77)	129.21 (146.30)	131.95 (143.18)
0.95	42.45 (40.38)	68.89 (67.14)	108.41 (112.57)	123.05 (124.62)	120.47 (132.93)	126.90 (127.62)	131.70 (136.27)	134.21 (136.63)

Table 2. The 95% width of one-dimensional marginalized posteriors scaled by the true value for injections and recoveries with SEOBNRHM waveforms. For all runs, $q = 4$, $\iota = \pi/3$ and $\text{SNR} = 20$. The spin column represents the equal component spins (i.e., $\chi_1 = \chi_2$) along the orbital angular momentum \mathbf{L} . The results in the parenthesis denote the corresponding face-on case (i.e., $\iota = 0$).

Spin	$M_{\text{tot}} = 50M_\odot$	$100M_\odot$	$150M_\odot$	$200M_\odot$	$250M_\odot$	$300M_\odot$	$400M_\odot$	$500M_\odot$
$\Delta M_c/M_c$								
-0.80	3.06 (5.10)	24.23 (64.35)	38.81 (62.92)	47.22 (68.48)	60.64 (84.09)	63.12 (74.90)	57.12 (73.01)	51.07 (72.81)
-0.50	2.44 (4.06)	35.08 (50.82)	55.56 (62.38)	61.89 (58.60)	62.30 (74.33)	59.85 (64.66)	56.14 (74.01)	48.41 (60.82)
0.00	1.81 (2.93)	11.28 (18.71)	33.00 (73.94)	48.90 (59.53)	51.53 (42.28)	55.06 (47.46)	49.16 (45.79)	40.34 (45.50)
0.50	1.44 (2.22)	7.91 (11.24)	18.25 (27.65)	27.16 (46.72)	31.72 (80.82)	33.31 (73.86)	31.89 (41.36)	29.96 (36.71)
0.80	1.25 (1.82)	5.32 (7.62)	11.20 (17.35)	17.42 (27.00)	21.86 (35.60)	25.20 (45.22)	27.15 (75.79)	25.94 (54.37)
0.95	1.17 (1.36)	4.06 (5.82)	8.83 (12.06)	12.99 (16.91)	15.84 (20.42)	17.84 (23.91)	19.32 (34.02)	23.16 (42.01)
$\Delta \nu/\nu$								
-0.80	25.59 (28.17)	29.72 (57.01)	41.66 (51.78)	51.33 (55.62)	57.07 (73.90)	58.96 (65.28)	53.84 (65.64)	49.11 (67.59)
-0.50	25.95 (34.98)	40.70 (56.41)	47.91 (48.07)	52.23 (45.70)	57.14 (66.92)	56.23 (55.03)	53.97 (68.97)	47.95 (57.68)
0.00	23.03 (36.59)	25.14 (33.43)	34.70 (64.77)	45.75 (49.93)	48.53 (32.46)	49.03 (39.06)	48.30 (39.64)	42.00 (38.73)
0.50	25.20 (35.75)	21.09 (24.83)	21.08 (27.57)	24.84 (41.79)	29.09 (80.86)	33.28 (72.85)	35.44 (35.61)	35.33 (31.75)
0.80	18.84 (24.80)	16.15 (18.05)	14.86 (18.50)	16.11 (22.68)	18.42 (31.15)	21.69 (43.34)	25.10 (90.22)	25.53 (67.91)
0.95	15.18 (16.45)	11.74 (13.57)	13.71 (15.90)	16.10 (18.48)	18.62 (21.99)	19.80 (25.81)	20.08 (40.85)	30.16 (54.17)
$\Delta d_L/d_L$								
-0.80	80.23 (26.19)	79.94 (113.39)	85.19 (118.92)	88.45 (132.54)	151.00 (163.15)	144.86 (151.74)	118.05 (158.19)	103.42 (180.17)
-0.50	87.40 (25.58)	123.58 (90.76)	150.47 (119.08)	152.48 (114.32)	144.31 (142.43)	132.20 (126.29)	117.51 (146.26)	102.48 (143.31)
0.00	85.53 (25.60)	81.41 (42.10)	96.18 (128.58)	107.68 (107.21)	103.42 (91.97)	106.08 (92.65)	93.52 (92.39)	80.20 (98.80)
0.50	88.50 (25.37)	80.90 (31.63)	77.94 (50.67)	79.40 (74.99)	78.03 (121.11)	75.25 (115.71)	71.23 (81.83)	64.99 (76.73)
0.80	86.90 (24.40)	80.97 (26.64)	75.31 (35.61)	73.42 (46.28)	71.99 (56.41)	70.28 (67.01)	68.52 (108.58)	66.31 (86.28)
0.95	91.11 (24.53)	81.52 (24.88)	73.04 (28.32)	66.87 (32.59)	62.08 (35.68)	59.68 (38.67)	55.85 (49.72)	53.80 (58.03)
$\Delta m_1^s/m_1^s$								
-0.80	28.61 (30.60)	18.60 (35.26)	17.46 (37.51)	18.79 (39.56)	23.28 (48.99)	27.40 (47.92)	26.40 (48.43)	25.64 (47.64)
-0.50	24.55 (34.49)	24.39 (32.05)	25.06 (35.54)	24.97 (36.08)	25.56 (45.59)	25.68 (41.53)	25.83 (48.42)	25.85 (46.06)
0.00	20.64 (33.46)	19.94 (25.12)	19.14 (41.07)	20.05 (37.02)	19.59 (30.61)	18.69 (34.05)	18.43 (35.20)	16.94 (34.39)
0.50	23.68 (33.91)	21.06 (23.13)	19.97 (20.39)	20.07 (21.57)	20.00 (48.40)	20.48 (48.48)	20.80 (32.73)	18.87 (30.40)
0.80	18.45 (24.90)	18.34 (18.24)	18.01 (17.19)	17.80 (16.71)	17.97 (17.91)	18.26 (22.75)	18.79 (60.33)	18.89 (50.79)
0.95	16.46 (17.03)	15.82 (14.01)	16.84 (14.48)	16.90 (15.31)	17.05 (16.40)	17.10 (18.18)	16.96 (27.45)	17.88 (37.37)
$\Delta m_2^s/m_2^s$								
-0.80	18.69 (20.44)	45.03 (146.80)	69.64 (153.41)	81.87 (150.82)	83.61 (169.60)	79.82 (154.00)	70.84 (155.98)	65.17 (158.58)
-0.50	20.53 (25.88)	59.73 (99.74)	91.40 (153.57)	96.89 (143.49)	97.26 (165.00)	91.11 (145.36)	84.40 (159.96)	72.80 (145.10)
0.00	19.50 (28.76)	28.04 (35.48)	52.90 (169.22)	76.55 (155.11)	80.97 (124.01)	81.61 (129.71)	76.62 (124.83)	64.40 (122.44)
0.50	20.67 (27.67)	22.16 (21.72)	28.85 (33.26)	38.10 (55.15)	45.59 (165.68)	51.95 (168.77)	55.17 (116.65)	55.18 (109.47)
0.80	16.46 (18.60)	19.14 (15.52)	21.64 (20.14)	25.62 (28.14)	29.83 (38.69)	34.78 (53.36)	40.35 (167.81)	39.92 (146.18)
0.95	16.01 (12.87)	18.15 (12.95)	23.36 (17.06)	27.19 (20.32)	30.24 (24.29)	31.58 (27.97)	32.24 (42.50)	39.73 (55.75)
$\Delta M_{\text{tot}}^s/M_{\text{tot}}^s$								
-0.80	19.86 (20.83)	15.32 (18.74)	17.34 (18.87)	18.85 (19.11)	19.63 (19.91)	19.72 (19.39)	18.53 (19.57)	18.23 (19.92)
-0.50	16.36 (22.69)	16.73 (16.98)	21.36 (20.11)	21.45 (19.54)	21.09 (19.25)	20.73 (18.17)	20.92 (18.55)	21.39 (18.63)
0.00	13.90 (21.34)	14.07 (15.35)	15.61 (17.69)	17.98 (18.57)	18.08 (18.44)	17.90 (16.97)	16.82 (16.23)	14.80 (16.03)
0.50	16.09 (21.84)	16.26 (15.65)	17.38 (14.79)	19.10 (14.29)	19.81 (15.09)	20.16 (16.45)	19.97 (15.94)	18.11 (15.28)
0.80	13.25 (16.61)	15.32 (12.96)	16.36 (12.98)	17.08 (13.02)	17.66 (12.65)	18.30 (12.67)	18.84 (21.24)	18.74 (18.72)
0.95	13.21 (11.76)	14.63 (10.25)	16.24 (10.73)	16.89 (11.19)	17.27 (11.41)	17.40 (11.86)	16.95 (15.60)	16.54 (20.55)
$\Delta \chi_1/\chi_1$								
-0.80	68.73 (57.12)	50.48 (89.33)	49.23 (101.19)	54.29 (114.19)	115.62 (119.20)	119.17 (120.38)	112.42 (125.58)	109.84 (127.39)
-0.50	100.66 (115.24)	122.17 (140.52)	151.05 (184.85)	164.38 (190.73)	158.42 (196.81)	150.45 (196.82)	148.16 (206.80)	151.68 (208.22)
0.50	46.73 (48.52)	46.42 (51.89)	58.35 (68.90)	67.49 (81.77)	69.50 (102.69)	69.81 (168.79)	71.77 (197.82)	69.04 (211.44)
0.80	14.41 (15.49)	14.98 (17.20)	17.47 (23.66)	22.27 (29.43)	25.49 (31.29)	26.77 (30.52)	26.04 (40.26)	23.95 (62.71)
0.95	4.36 (4.54)	4.40 (5.49)	4.95 (7.04)	5.51 (8.29)	6.14 (8.50)	6.54 (8.38)	6.91 (8.43)	7.15 (8.11)
$\Delta \chi_2/\chi_2$								
-0.80	159.54 (162.59)	152.68 (146.97)	148.79 (149.74)	156.49 (161.73)	165.62 (168.52)	171.08 (169.93)	171.03 (169.06)	170.25 (174.42)
-0.50	262.59 (270.20)	256.04 (258.78)	245.29 (243.53)	257.00 (252.92)	264.46 (260.38)	273.12 (271.42)	271.56 (266.43)	271.45 (268.27)
0.50	266.38 (257.87)	259.36 (265.83)	256.61 (268.90)	255.57 (273.41)	245.65 (277.63)	250.18 (239.98)	244.83 (248.33)	238.23 (252.89)
0.80	143.92 (149.29)	148.14 (157.34)	152.42 (165.12)	153.04 (170.84)	151.81 (170.30)	149.25 (172.16)	145.39 (173.50)	142.27 (152.74)
0.95	67.77 (73.63)	83.15 (97.37)	103.08 (124.20)	114.54 (139.45)	121.39 (144.29)	117.70 (143.50)	109.63 (147.88)	115.18 (145.90)

Table 3. The 95% width of one-dimensional marginalized posteriors scaled by the true value for injections and recoveries with SEOBNRHM waveforms. For all runs, $q = 10$, $\iota = \pi/3$ and $\text{SNR} = 20$. The spin column represents the equal component spins (i.e., $\chi_1 = \chi_2$) along the orbital angular momentum \mathbf{L} . The results in the parenthesis denote the corresponding face-on case (i.e., $\iota = 0$).

Spin	$M_{\text{tot}} = 50M_{\odot}$	$100M_{\odot}$	$150M_{\odot}$	$200M_{\odot}$	$250M_{\odot}$	$300M_{\odot}$	$400M_{\odot}$	$500M_{\odot}$
$\Delta M_c/M_c$								
-0.80	1.83 (3.14)	12.19 (72.67)	83.57 (130.25)	219.60 (147.28)	123.69 (136.57)	50.09 (137.15)	30.90 (152.80)	26.94 (117.98)
-0.50	2.48 (2.79)	10.31 (30.93)	49.03 (163.31)	65.12 (122.41)	68.00 (127.08)	66.58 (126.37)	55.53 (134.03)	48.18 (105.43)
0.00	1.49 (1.95)	4.72 (12.00)	13.82 (54.59)	25.68 (129.93)	36.25 (92.34)	38.47 (87.81)	35.76 (92.42)	31.84 (72.31)
0.50	1.22 (1.64)	3.08 (5.80)	7.63 (18.03)	14.83 (42.78)	21.30 (82.63)	26.75 (148.46)	25.19 (64.02)	22.69 (62.52)
0.80	0.99 (1.36)	2.36 (4.26)	4.92 (10.41)	8.16 (17.92)	11.66 (26.27)	16.05 (40.96)	22.34 (116.26)	27.49 (136.58)
0.95	0.64 (0.81)	1.68 (3.20)	3.05 (6.38)	4.31 (9.82)	5.28 (12.81)	6.26 (17.34)	7.83 (36.98)	9.82 (62.95)
$\Delta\nu/\nu$								
-0.80	28.31 (36.62)	25.15 (103.99)	95.19 (136.93)	198.58 (160.93)	120.07 (148.38)	61.28 (147.45)	37.00 (174.71)	32.79 (143.79)
-0.50	53.84 (50.44)	36.63 (55.97)	59.75 (177.13)	73.93 (131.25)	78.65 (136.07)	77.11 (135.84)	67.40 (148.57)	60.67 (126.30)
0.00	41.79 (49.66)	24.27 (36.95)	26.34 (72.55)	35.73 (136.06)	47.88 (94.84)	51.30 (91.34)	50.82 (100.71)	51.37 (73.80)
0.50	41.66 (72.24)	24.14 (35.21)	18.86 (31.66)	20.92 (49.91)	24.17 (94.04)	28.02 (188.81)	26.42 (59.79)	25.21 (58.16)
0.80	32.96 (47.31)	18.67 (26.10)	14.35 (20.61)	13.06 (22.18)	13.93 (27.57)	16.08 (43.81)	20.59 (160.46)	25.10 (205.87)
0.95	14.88 (16.97)	10.56 (13.90)	9.65 (12.63)	9.70 (14.14)	9.57 (16.04)	9.91 (20.75)	10.24 (47.39)	11.32 (87.60)
$\Delta d_L/d_L$								
-0.80	73.11 (25.06)	65.60 (154.89)	231.74 (311.80)	779.20 (361.17)	396.09 (359.12)	107.18 (374.55)	54.23 (448.51)	36.62 (443.06)
-0.50	72.74 (25.24)	66.74 (65.21)	104.59 (373.35)	123.77 (306.22)	125.95 (326.62)	124.00 (329.27)	103.44 (383.32)	93.69 (363.36)
0.00	74.63 (24.48)	63.99 (34.19)	59.96 (103.53)	66.07 (286.97)	75.47 (224.17)	74.46 (215.62)	68.73 (240.12)	60.79 (224.08)
0.50	78.26 (24.18)	68.11 (26.58)	62.94 (40.44)	62.63 (76.50)	62.80 (139.50)	64.71 (270.90)	57.53 (163.59)	54.76 (160.13)
0.80	76.81 (23.87)	69.44 (24.91)	64.64 (29.68)	62.21 (38.27)	58.95 (47.82)	59.18 (66.28)	58.95 (188.20)	63.14 (243.47)
0.95	76.34 (21.90)	69.09 (24.14)	64.40 (24.37)	59.30 (27.87)	56.62 (30.44)	52.64 (33.92)	48.95 (58.46)	49.68 (96.24)
$\Delta m_1^s/m_1^s$								
-0.80	24.98 (34.85)	13.70 (26.92)	13.84 (34.98)	34.42 (39.55)	14.71 (40.06)	9.44 (40.44)	5.93 (47.07)	4.87 (43.39)
-0.50	47.98 (41.47)	25.17 (24.95)	20.87 (43.39)	19.63 (37.73)	18.56 (38.88)	17.75 (39.69)	17.34 (42.20)	17.63 (39.50)
0.00	29.90 (35.83)	17.34 (22.64)	13.98 (21.61)	12.42 (38.02)	12.10 (32.02)	11.74 (31.30)	11.83 (33.34)	12.78 (28.82)
0.50	28.84 (44.84)	18.27 (26.33)	14.86 (21.08)	13.98 (18.85)	13.02 (21.13)	12.80 (49.78)	11.73 (23.69)	11.33 (23.00)
0.80	24.22 (31.94)	15.52 (21.40)	13.52 (17.59)	12.97 (16.11)	12.57 (14.71)	12.73 (14.82)	11.96 (39.63)	11.48 (58.17)
0.95	12.57 (13.74)	11.36 (12.32)	11.96 (11.08)	12.39 (11.18)	12.77 (10.91)	12.55 (11.05)	12.80 (18.07)	13.71 (31.41)
$\Delta m_2^s/m_2^s$								
-0.80	14.96 (18.74)	22.50 (123.95)	130.83 (346.02)	403.11 (353.57)	166.47 (323.09)	75.83 (311.73)	45.95 (359.18)	41.34 (330.96)
-0.50	27.13 (24.63)	23.81 (51.06)	73.94 (421.33)	98.56 (336.48)	101.92 (328.70)	98.20 (318.44)	80.91 (338.26)	69.20 (322.25)
0.00	20.90 (23.89)	15.61 (24.89)	24.92 (78.17)	40.38 (367.98)	55.33 (293.82)	58.81 (277.45)	56.95 (283.17)	54.70 (249.85)
0.50	21.44 (34.68)	14.79 (17.94)	15.59 (24.15)	22.33 (51.83)	29.00 (105.19)	35.61 (363.76)	33.97 (217.18)	31.52 (210.56)
0.80	18.00 (23.10)	13.30 (12.75)	13.55 (13.28)	15.24 (19.26)	17.35 (27.98)	21.03 (46.00)	26.23 (173.56)	30.84 (347.95)
0.95	10.50 (9.01)	11.94 (8.56)	13.74 (9.74)	15.11 (12.40)	16.27 (15.10)	16.88 (19.80)	17.79 (43.84)	19.40 (77.47)
$\Delta M_{\text{tot}}^s/M_{\text{tot}}^s$								
-0.80	21.50 (30.09)	11.63 (19.21)	12.20 (19.37)	15.49 (18.28)	11.32 (18.91)	10.10 (18.95)	7.65 (20.27)	7.20 (20.36)
-0.50	41.23 (35.53)	21.40 (20.46)	19.88 (19.78)	20.04 (19.07)	18.74 (18.07)	17.64 (17.82)	15.60 (18.67)	15.06 (18.98)
0.00	25.48 (30.52)	14.95 (18.87)	12.22 (16.48)	11.54 (18.14)	11.09 (17.21)	10.44 (16.04)	9.66 (16.23)	9.58 (15.87)
0.50	24.46 (37.56)	15.99 (22.54)	13.41 (18.22)	13.03 (16.10)	12.67 (14.84)	12.97 (18.99)	11.90 (14.01)	11.40 (14.05)
0.80	20.81 (27.06)	13.89 (18.54)	12.56 (15.57)	12.42 (14.39)	12.23 (13.27)	12.53 (12.34)	11.98 (21.73)	11.60 (27.24)
0.95	11.20 (11.86)	10.79 (10.84)	11.68 (9.89)	12.21 (10.13)	12.73 (9.80)	12.58 (9.62)	12.91 (13.58)	13.74 (22.32)
$\Delta\chi_1/\chi_1$								
-0.80	53.72 (68.55)	33.18 (64.17)	51.25 (97.30)	125.65 (110.64)	96.31 (114.81)	54.26 (116.72)	33.30 (118.64)	28.77 (116.68)
-0.50	137.67 (126.95)	81.82 (87.06)	105.28 (167.70)	112.60 (186.32)	117.31 (187.19)	123.53 (192.92)	137.55 (194.92)	140.38 (197.06)
0.50	36.01 (54.61)	25.42 (33.53)	25.75 (39.63)	32.19 (54.47)	37.47 (64.16)	41.37 (124.42)	36.68 (206.31)	38.38 (216.32)
0.80	7.57 (7.38)	7.90 (11.10)	8.45 (14.28)	9.75 (17.74)	11.05 (19.76)	13.72 (20.67)	16.18 (22.60)	18.33 (49.87)
0.95	2.39 (3.19)	2.09 (3.95)	1.95 (4.50)	1.82 (5.46)	1.74 (5.25)	1.72 (5.00)	1.71 (5.34)	1.88 (5.57)
$\Delta\chi_2/\chi_2$								
-0.80	170.25 (172.21)	170.05 (170.02)	171.44 (161.66)	170.63 (167.33)	170.89 (173.95)	174.03 (173.59)	169.61 (173.30)	169.13 (173.63)
-0.50	271.18 (276.01)	272.48 (277.39)	275.31 (260.76)	270.29 (269.17)	273.27 (271.54)	276.18 (272.73)	277.80 (275.65)	278.82 (271.75)
0.50	281.33 (264.94)	268.18 (274.95)	261.38 (270.31)	252.48 (273.60)	245.65 (274.66)	246.45 (266.32)	250.92 (253.02)	269.66 (256.35)
0.80	151.86 (124.07)	143.18 (164.41)	140.32 (163.40)	142.57 (166.37)	141.21 (168.46)	145.89 (170.90)	144.07 (177.06)	148.05 (161.97)
0.95	86.72 (104.97)	82.09 (103.09)	96.21 (117.98)	100.94 (136.49)	104.87 (144.37)	106.65 (143.13)	105.35 (146.09)	108.04 (146.61)

REFERENCES

- Aasi, J., Abbott, B. P., Abbott, R., et al. 2015, *Classical and Quantum Gravity*, 32, 074001, doi: [10.1088/0264-9381/32/7/074001](https://doi.org/10.1088/0264-9381/32/7/074001)
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017a, *PhRvL*, 119, 161101, doi: [10.1103/PhysRevLett.119.161101](https://doi.org/10.1103/PhysRevLett.119.161101)
- . 2017b, *ApJL*, 848, L12, doi: [10.3847/2041-8213/aa91c9](https://doi.org/10.3847/2041-8213/aa91c9)
- . 2017c, *Nature*, 551, 85, doi: [10.1038/nature24471](https://doi.org/10.1038/nature24471)
- . 2017d, *PhRvD*, 96, 022001, doi: [10.1103/PhysRevD.96.022001](https://doi.org/10.1103/PhysRevD.96.022001)
- . 2018, *PhRvL*, 121, 161101, doi: [10.1103/PhysRevLett.121.161101](https://doi.org/10.1103/PhysRevLett.121.161101)
- . 2019a, *Physical Review X*, 9, 031040, doi: [10.1103/PhysRevX.9.031040](https://doi.org/10.1103/PhysRevX.9.031040)
- . 2019b, arXiv e-prints, arXiv:1906.08000, <https://arxiv.org/abs/1906.08000>
- . 2020a, *Living Reviews in Relativity*, 23, 3, doi: [10.1007/s41114-020-00026-9](https://doi.org/10.1007/s41114-020-00026-9)
- . 2021a, *ApJ*, 909, 218, doi: [10.3847/1538-4357/abdc67](https://doi.org/10.3847/1538-4357/abdc67)
- Abbott, R., Abbott, T. D., Abraham, S., & Acernese, F. 2021b, *SoftwareX*, 13, 100658, doi: [10.1016/j.softx.2021.100658](https://doi.org/10.1016/j.softx.2021.100658)
- Abbott, R., Abbott, T. D., Abraham, S., et al. 2020b, arXiv e-prints, arXiv:2010.14527, <https://arxiv.org/abs/2010.14527>
- . 2020c, arXiv e-prints, arXiv:2010.14533, <https://arxiv.org/abs/2010.14533>
- . 2020d, *PhRvL*, 125, 101102, doi: [10.1103/PhysRevLett.125.101102](https://doi.org/10.1103/PhysRevLett.125.101102)
- . 2020e, *ApJL*, 900, L13, doi: [10.3847/2041-8213/aba493](https://doi.org/10.3847/2041-8213/aba493)
- Acernese, F., Agathos, M., Agatsuma, K., et al. 2015, *Classical and Quantum Gravity*, 32, 024001, doi: [10.1088/0264-9381/32/2/024001](https://doi.org/10.1088/0264-9381/32/2/024001)
- Amaro-Seoane, P., Gair, J. R., Freitag, M., et al. 2007, *Classical and Quantum Gravity*, 24, R113, doi: [10.1088/0264-9381/24/17/R01](https://doi.org/10.1088/0264-9381/24/17/R01)
- Antonini, F., Gieles, M., & Gualandris, A. 2019, *MNRAS*, 486, 5008, doi: [10.1093/mnras/stz1149](https://doi.org/10.1093/mnras/stz1149)
- Avila, M. L., Rogachev, G. V., Koshchiy, E., et al. 2015, *PhRvL*, 114, 071101, doi: [10.1103/PhysRevLett.114.071101](https://doi.org/10.1103/PhysRevLett.114.071101)
- Azuma, R. E., Uberseder, E., Simpson, E. C., et al. 2010, *PhRvC*, 81, 045805, doi: [10.1103/PhysRevC.81.045805](https://doi.org/10.1103/PhysRevC.81.045805)
- Barkat, Z., Rakavy, G., & Sack, N. 1967, *PhRvL*, 18, 379, doi: [10.1103/PhysRevLett.18.379](https://doi.org/10.1103/PhysRevLett.18.379)
- Belczynski, K., Buonanno, A., Cantiello, M., et al. 2014, *ApJ*, 789, 120, doi: [10.1088/0004-637X/789/2/120](https://doi.org/10.1088/0004-637X/789/2/120)
- Blinnikov, S. I. 2010, *Physics of Atomic Nuclei*, 73, 604, doi: [10.1134/S1063778810040046](https://doi.org/10.1134/S1063778810040046)
- Bond, J. R., Arnett, W. D., & Carr, B. J. 1984, *ApJ*, 280, 825, doi: [10.1086/162057](https://doi.org/10.1086/162057)
- Brown, G. E., Heger, A., Langer, N., et al. 2001, *NewA*, 6, 457, doi: [10.1016/S1384-1076\(01\)00077-X](https://doi.org/10.1016/S1384-1076(01)00077-X)
- Brune, C. R. 2002, *PhRvC*, 66, 044611, doi: [10.1103/PhysRevC.66.044611](https://doi.org/10.1103/PhysRevC.66.044611)
- Brune, C. R., Geist, W. H., Kavanagh, R. W., & Veal, K. D. 1999, *Phys. Rev. Lett.*, 83, 4025, doi: [10.1103/PhysRevLett.83.4025](https://doi.org/10.1103/PhysRevLett.83.4025)
- Bustillo, J. C., Sanchis-Gual, N., Torres-Forné, A., et al. 2021, *PhRvL*, 126, 081101, doi: [10.1103/PhysRevLett.126.081101](https://doi.org/10.1103/PhysRevLett.126.081101)
- Calderón Bustillo, J., Sanchis-Gual, N., Torres-Forné, A., & Font, J. A. 2020, arXiv e-prints, arXiv:2009.01066, <https://arxiv.org/abs/2009.01066>
- Campanelli, M., Lousto, C. O., & Zlochower, Y. 2006, *PhRvD*, 74, 041501, doi: [10.1103/PhysRevD.74.041501](https://doi.org/10.1103/PhysRevD.74.041501)
- Chatzopoulos, E., & Wheeler, J. C. 2012, *ApJ*, 760, 154, doi: [10.1088/0004-637X/760/2/154](https://doi.org/10.1088/0004-637X/760/2/154)
- Chatzopoulos, E., Wheeler, J. C., & Couch, S. M. 2013, *ApJ*, 776, 129, doi: [10.1088/0004-637X/776/2/129](https://doi.org/10.1088/0004-637X/776/2/129)
- Costa, G., Bressan, A., Mapelli, M., et al. 2021, *MNRAS*, 501, 4514, doi: [10.1093/mnras/staa3916](https://doi.org/10.1093/mnras/staa3916)
- Cotesta, R., Buonanno, A., Bohé, A., et al. 2018, *PhRvD*, 98, 084028, doi: [10.1103/PhysRevD.98.084028](https://doi.org/10.1103/PhysRevD.98.084028)
- Cotesta, R., Marsat, S., & Pürrer, M. 2020, *PhRvD*, 101, 124040, doi: [10.1103/PhysRevD.101.124040](https://doi.org/10.1103/PhysRevD.101.124040)
- deBoer, R. J., Görres, J., Wiescher, M., et al. 2017, *Reviews of Modern Physics*, 89, 035007, doi: [10.1103/RevModPhys.89.035007](https://doi.org/10.1103/RevModPhys.89.035007)
- Descouvemont, P., & Baye, D. 2010, *Reports on Progress in Physics*, 73, 036301, doi: [10.1088/0034-4885/73/3/036301](https://doi.org/10.1088/0034-4885/73/3/036301)
- Di Carlo, U. N., Giacobbo, N., Mapelli, M., et al. 2019, *MNRAS*, 487, 2947, doi: [10.1093/mnras/stz1453](https://doi.org/10.1093/mnras/stz1453)
- Di Carlo, U. N., Mapelli, M., Bouffanais, Y., et al. 2020, *MNRAS*, 497, 1043, doi: [10.1093/mnras/staa1997](https://doi.org/10.1093/mnras/staa1997)
- Ebisuzaki, T., Makino, J., Tsuru, T. G., et al. 2001, *ApJL*, 562, L19, doi: [10.1086/338118](https://doi.org/10.1086/338118)
- Estelles, H. 2021, <https://pnp.ligo.org/P2100149/>
- Ezquiaga, J. M., & Holz, D. E. 2021, *Astrophys. J. Lett.*, 909, L23, doi: [10.3847/2041-8213/abe638](https://doi.org/10.3847/2041-8213/abe638)
- Farmer, R., Renzo, M., de Mink, S. E., Fishbach, M., & Justham, S. 2020, *ApJL*, 902, L36, doi: [10.3847/2041-8213/abbadd](https://doi.org/10.3847/2041-8213/abbadd)
- Farmer, R., Renzo, M., de Mink, S. E., Marchant, P., & Justham, S. 2019, *ApJ*, 887, 53, doi: [10.3847/1538-4357/ab518b](https://doi.org/10.3847/1538-4357/ab518b)
- Finn, L. S., & Chernoff, D. F. 1993, *PhRvD*, 47, 2198, doi: [10.1103/PhysRevD.47.2198](https://doi.org/10.1103/PhysRevD.47.2198)

- Fishbach, M., & Holz, D. E. 2020, *ApJL*, 904, L26, doi: [10.3847/2041-8213/abc827](https://doi.org/10.3847/2041-8213/abc827)
- Fowler, W. A., & Hoyle, F. 1964, *ApJS*, 9, 201, doi: [10.1086/190103](https://doi.org/10.1086/190103)
- Fraley, G. S. 1968, *Ap&SS*, 2, 96, doi: [10.1007/BF00651498](https://doi.org/10.1007/BF00651498)
- Fryer, C. L., & Kalogera, V. 2001, *ApJ*, 554, 548, doi: [10.1086/321359](https://doi.org/10.1086/321359)
- Gair, J. R., Mandel, I., Miller, M. C., & Volonteri, M. 2011, *General Relativity and Gravitation*, 43, 485, doi: [10.1007/s10714-010-1104-3](https://doi.org/10.1007/s10714-010-1104-3)
- García-Quirós, C., Colleoni, M., Husa, S., et al. 2020, *PhRvD*, 102, 064002, doi: [10.1103/PhysRevD.102.064002](https://doi.org/10.1103/PhysRevD.102.064002)
- Gayathri, V., Bartos, I., Haiman, Z., et al. 2020a, *ApJL*, 890, L20, doi: [10.3847/2041-8213/ab745d](https://doi.org/10.3847/2041-8213/ab745d)
- Gayathri, V., Healy, J., Lange, J., et al. 2020b, *arXiv e-prints*, arXiv:2009.05461, <https://arxiv.org/abs/2009.05461>
- Gerosa, D., & Berti, E. 2017, *PhRvD*, 95, 124046, doi: [10.1103/PhysRevD.95.124046](https://doi.org/10.1103/PhysRevD.95.124046)
- Gialanella, L., Rogalla, D., Strieder, F., et al. 2001, *Eur. Phys. J. A*, 11, 357, doi: [10.1007/s100500170075](https://doi.org/10.1007/s100500170075)
- Gilmer, M. S., Kozyreva, A., Hirschi, R., Fröhlich, C., & Yusof, N. 2017, *ApJ*, 846, 100, doi: [10.3847/1538-4357/aa8461](https://doi.org/10.3847/1538-4357/aa8461)
- Graff, P. B., Buonanno, A., & Sathyaprakash, B. S. 2015, *PhRvD*, 92, 022002, doi: [10.1103/PhysRevD.92.022002](https://doi.org/10.1103/PhysRevD.92.022002)
- GWOSC. 2020, https://www.gw-openscience.org/o3_eventdata_docs/
- Haster, C.-J., Wang, Z., Berry, C. P. L., et al. 2016, *MNRAS*, 457, 4499, doi: [10.1093/mnras/stw233](https://doi.org/10.1093/mnras/stw233)
- Heger, A., Fryer, C. L., Woosley, S. E., Langer, N., & Hartmann, D. H. 2003, *ApJ*, 591, 288, doi: [10.1086/375341](https://doi.org/10.1086/375341)
- Hunter, J. D. 2007, *Computing In Science & Engineering*, 9, 90
- Joggerst, C. C., & Whalen, D. J. 2011, *ApJ*, 728, 129, doi: [10.1088/0004-637X/728/2/129](https://doi.org/10.1088/0004-637X/728/2/129)
- Khalil, M., Buonanno, A., Steinhoff, J., & Vines, J. 2021, *arXiv e-prints*, arXiv:2104.11705, <https://arxiv.org/abs/2104.11705>
- Kimball, C., Talbot, C., Berry, C. P. L., et al. 2020, *arXiv e-prints*, arXiv:2011.05332, <https://arxiv.org/abs/2011.05332>
- Kozyreva, A., Yoon, S.-C., & Langer, N. 2014, *A&A*, 566, A146, doi: [10.1051/0004-6361/201423641](https://doi.org/10.1051/0004-6361/201423641)
- Kunz, R., Fey, M., Jaeger, M., et al. 2002, *Astrophys. J.*, 567, 643, doi: [10.1086/338384](https://doi.org/10.1086/338384)
- Kunz, R., Fey, M., Jaeger, M., et al. 2002, *ApJ*, 567, 643, doi: [10.1086/338384](https://doi.org/10.1086/338384)
- Lane, A. M., & Thomas, R. G. 1958, *Rev. Mod. Phys.*, 30, 257, doi: [10.1103/RevModPhys.30.257](https://doi.org/10.1103/RevModPhys.30.257)
- Liu, X., Cao, Z., & Zhu, Z.-H. 2021, *arXiv e-prints*, arXiv:2102.08614, <https://arxiv.org/abs/2102.08614>
- Mapelli, M., Dall’Amico, M., Bouffanais, Y., et al. 2021, *arXiv e-prints*, arXiv:2103.05016, <https://arxiv.org/abs/2103.05016>
- Marchant, P., & Moriya, T. J. 2020, *A&A*, 640, L18, doi: [10.1051/0004-6361/202038902](https://doi.org/10.1051/0004-6361/202038902)
- Miller, M. C., & Colbert, E. J. M. 2004, *International Journal of Modern Physics D*, 13, 1, doi: [10.1142/S0218271804004426](https://doi.org/10.1142/S0218271804004426)
- Miller, M. C., & Hamilton, D. P. 2002, *MNRAS*, 330, 232, doi: [10.1046/j.1365-8711.2002.05112.x](https://doi.org/10.1046/j.1365-8711.2002.05112.x)
- Nagar, A., Bonino, A., & Rettegno, P. 2021, *arXiv e-prints*, arXiv:2101.08624, <https://arxiv.org/abs/2101.08624>
- Nitz, A. H., Capano, C., Nielsen, A. B., et al. 2019, *ApJ*, 872, 195, doi: [10.3847/1538-4357/ab0108](https://doi.org/10.3847/1538-4357/ab0108)
- Nitz, A. H., & Capano, C. D. 2021, *ApJL*, 907, L9, doi: [10.3847/2041-8213/abccc5](https://doi.org/10.3847/2041-8213/abccc5)
- Nitz, A. H., Dent, T., Davies, G. S., et al. 2020, *ApJ*, 891, 123, doi: [10.3847/1538-4357/ab733f](https://doi.org/10.3847/1538-4357/ab733f)
- Ober, W. W., El Eid, M. F., & Fricke, K. J. 1983, *A&A*, 119, 61
- O’Leary, R. M., Rasio, F. A., Fregeau, J. M., Ivanova, N., & O’Shaughnessy, R. 2006, *ApJ*, 637, 937, doi: [10.1086/498446](https://doi.org/10.1086/498446)
- Ossokine, S., Buonanno, A., Marsat, S., et al. 2020, *PhRvD*, 102, 044055, doi: [10.1103/PhysRevD.102.044055](https://doi.org/10.1103/PhysRevD.102.044055)
- Pan, Y., Buonanno, A., Boyle, M., et al. 2011, *PhRvD*, 84, 124052, doi: [10.1103/PhysRevD.84.124052](https://doi.org/10.1103/PhysRevD.84.124052)
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, 192, 3, doi: [10.1088/0067-0049/192/1/3](https://doi.org/10.1088/0067-0049/192/1/3)
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, *ApJS*, 208
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, *ApJS*, 220, 15, doi: [10.1088/0067-0049/220/1/15](https://doi.org/10.1088/0067-0049/220/1/15)
- Paxton, B., Schwab, J., Bauer, E. B., et al. 2018, *ApJS*, 234, 34, doi: [10.3847/1538-4365/aaa5a8](https://doi.org/10.3847/1538-4365/aaa5a8)
- Paxton, B., Smolec, R., Schwab, J., et al. 2019, *ApJS*, 243, 10, doi: [10.3847/1538-4365/ab2241](https://doi.org/10.3847/1538-4365/ab2241)
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13, doi: [10.1051/0004-6361/201525830](https://doi.org/10.1051/0004-6361/201525830)
- Portegies Zwart, S. F., & McMillan, S. L. W. 2000, *ApJL*, 528, L17, doi: [10.1086/312422](https://doi.org/10.1086/312422)
- Pratten, G., García-Quirós, C., Colleoni, M., et al. 2020, *arXiv e-prints*, arXiv:2004.06503, <https://arxiv.org/abs/2004.06503>
- Quinlan, G. D., & Shapiro, S. L. 1989, *ApJ*, 343, 725, doi: [10.1086/167745](https://doi.org/10.1086/167745)

- Rakavy, G., & Shaviv, G. 1967, *ApJ*, 148, 803, doi: [10.1086/149204](https://doi.org/10.1086/149204)
- Renzo, M., Farmer, R., Justham, S., et al. 2020, *A&A*, 640, A56, doi: [10.1051/0004-6361/202037710](https://doi.org/10.1051/0004-6361/202037710)
- Rodriguez, C. L., Zevin, M., Amaro-Seoane, P., et al. 2019, *PhRvD*, 100, 043027, doi: [10.1103/PhysRevD.100.043027](https://doi.org/10.1103/PhysRevD.100.043027)
- Romero-Shaw, I., Lasky, P. D., Thrane, E., & Calderón Bustillo, J. 2020, *ApJL*, 903, L5, doi: [10.3847/2041-8213/abbe26](https://doi.org/10.3847/2041-8213/abbe26)
- Sallaska, A. L., Iliadis, C., Champange, A. E., et al. 2013a, *ApJS*, 207, 18, doi: [10.1088/0067-0049/207/1/18](https://doi.org/10.1088/0067-0049/207/1/18)
- . 2013b, *ApJS*, 207, 18, doi: [10.1088/0067-0049/207/1/18](https://doi.org/10.1088/0067-0049/207/1/18)
- Sathyaprakash, B. S., & Dhurandhar, S. V. 1991, *PhRvD*, 44, 3819, doi: [10.1103/PhysRevD.44.3819](https://doi.org/10.1103/PhysRevD.44.3819)
- Schürmann, D., Gialanella, L., Kunz, R., & Strieder, F. 2012, *Phys. Lett. B*, 711, 35, doi: [10.1016/j.physletb.2012.03.064](https://doi.org/10.1016/j.physletb.2012.03.064)
- Schutz, B. F. 1986, *Nature*, 323, 310, doi: [10.1038/323310a0](https://doi.org/10.1038/323310a0)
- Skilling, J. 2006, *Bayesian Analysis*, 1, 833, doi: [10.1214/06-BA127](https://doi.org/10.1214/06-BA127)
- Sukhbold, T., & Woosley, S. E. 2014, *ApJ*, 783, 10, doi: [10.1088/0004-637X/783/1/10](https://doi.org/10.1088/0004-637X/783/1/10)
- Sukhbold, T., Woosley, S. E., & Heger, A. 2018, *ApJ*, 860, 93, doi: [10.3847/1538-4357/aac2da](https://doi.org/10.3847/1538-4357/aac2da)
- Takahashi, K. 2018, *ApJ*, 863, 153, doi: [10.3847/1538-4357/aad2d2](https://doi.org/10.3847/1538-4357/aad2d2)
- Timmes, F. X., Woosley, S. E., & Weaver, T. A. 1996, *ApJ*, 457, 834
- Townsend, R. H. D. 2019a, MESA SDK for Linux, 20190503, Zenodo, doi: [10.5281/zenodo.2669541](https://doi.org/10.5281/zenodo.2669541), <https://doi.org/10.5281/zenodo.2669541>
- . 2019b, MESA SDK for Mac OS, 20190503, Zenodo, doi: [10.5281/zenodo.2669543](https://doi.org/10.5281/zenodo.2669543), [http://doi.org/10.5281/zenodo.2669543](https://doi.org/10.5281/zenodo.2669543)
- Uberseder, E., & deBoer, R. J. 2015, *AZURE2 User Manual*
- Umeda, H., Yoshida, T., Nagele, C., & Takahashi, K. 2020, *ApJL*, 905, L21, doi: [10.3847/2041-8213/abcb96](https://doi.org/10.3847/2041-8213/abcb96)
- Vallisneri, M., Kanner, J., Williams, R., Weinstein, A., & Stephens, B. 2015, in *Journal of Physics Conference Series*, Vol. 610, *Journal of Physics Conference Series*, 012021
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, *Computing in Science Engineering*, 13, 22, doi: [10.1109/MCSE.2011.37](https://doi.org/10.1109/MCSE.2011.37)
- Varma, V., Field, S. E., Scheel, M. A., et al. 2019a, *Physical Review Research*, 1, 033015, doi: [10.1103/PhysRevResearch.1.033015](https://doi.org/10.1103/PhysRevResearch.1.033015)
- . 2019b, *PhRvD*, 99, 064045, doi: [10.1103/PhysRevD.99.064045](https://doi.org/10.1103/PhysRevD.99.064045)
- Veitch, J., Pürrer, M., & Mandel, I. 2015a, *PhRvL*, 115, 141101, doi: [10.1103/PhysRevLett.115.141101](https://doi.org/10.1103/PhysRevLett.115.141101)
- Veitch, J., Raymond, V., Farr, B., et al. 2015b, *PhRvD*, 91, 042003, doi: [10.1103/PhysRevD.91.042003](https://doi.org/10.1103/PhysRevD.91.042003)
- Venumadhav, T., Zackay, B., Roulet, J., Dai, L., & Zaldarriaga, M. 2020, *PhRvD*, 101, 083030, doi: [10.1103/PhysRevD.101.083030](https://doi.org/10.1103/PhysRevD.101.083030)
- Woosley, S. E. 2017, *ApJ*, 836, 244, doi: [10.3847/1538-4357/836/2/244](https://doi.org/10.3847/1538-4357/836/2/244)
- Woosley, S. E., & Heger, A. 2021, arXiv e-prints, arXiv:2103.07933. <https://arxiv.org/abs/2103.07933>
- Woosley, S. E., Heger, A., & Weaver, T. A. 2002, *Rev. Mod. Phys.*, 74, 1015, doi: [10.1103/RevModPhys.74.1015](https://doi.org/10.1103/RevModPhys.74.1015)
- Yoshida, T., Umeda, H., Maeda, K., & Ishii, T. 2016, *MNRAS*, 457, 351, doi: [10.1093/mnras/stv3002](https://doi.org/10.1093/mnras/stv3002)
- Yun, Q., Han, W.-B., Zhong, X., & Benavides-Gallego, C. A. 2021, arXiv e-prints, arXiv:2104.03789. <https://arxiv.org/abs/2104.03789>
- Zackay, B., Dai, L., Venumadhav, T., Roulet, J., & Zaldarriaga, M. 2019, arXiv e-prints, arXiv:1910.09528. <https://arxiv.org/abs/1910.09528>
- Zhang, W., Woosley, S. E., & Heger, A. 2008, *ApJ*, 679, 639, doi: [10.1086/526404](https://doi.org/10.1086/526404)