

STRUCTURAL FORMULAS FOR MATRIX-VALUED ORTHOGONAL POLYNOMIALS RELATED TO 2×2 HYPERGEOMETRIC OPERATORS

C. CALDERÓN AND M. M. CASTRO

ABSTRACT. We give some structural formulas for the family of matrix-valued orthogonal polynomials of size 2×2 introduced by C. Calderón et al. in an earlier work, which are common eigenfunctions of a differential operator of hypergeometric type. Specifically, we give a Rodrigues formula that allows us to write this family of polynomials explicitly in terms of the classical Jacobi polynomials, and write, for the sequence of orthonormal polynomials, the three-term recurrence relation and the Christoffel-Darboux identity. We obtain a Pearson equation, which enables us to prove that the sequence of derivatives of the orthogonal polynomials is also orthogonal, and to compute a Rodrigues formula for these polynomials as well as a matrix-valued differential operator having these polynomials as eigenfunctions. We also describe the second-order differential operators of the algebra associated with the weight matrix.

1. INTRODUCTION

In the last few years, the search for examples of matrix-valued orthogonal polynomials that are common eigenfunctions of a second order differential operator, that is to say, satisfying a *bispectral property* in the sense of [13], has received a lot of attention after the seminal work of A. Durán in [15].

The theory of matrix-valued orthogonal polynomials was started by Krein in 1949 [37], [38] (see also [1] and [2]), in connection with spectral analysis and moment problems. Nevertheless, the first examples of orthogonal matrix polynomials satisfying this extra property and non reducible to scalar case, appeared more recently in [27, 28, 29, 25] and [19]. The collection of examples has been growing lately (see for instance [16, 17, 21, 22, 26, 41, 40, 3, 35, 36, 42, 34, 4]). Moreover, the problem of giving a general classification of these families of matrix-valued orthogonal polynomials as solutions of the so called *Matrix Bochner Problem* has been also recently addressed in [8] and in [7] for the special case of 2×2 hypergeometric matrix differential operators.

As the case of classical orthogonal polynomials, the families of matrix-valued orthogonal polynomials satisfy many formal properties such as structural formulas (see for instance [20, 24, 18, 3, 34]), which have been very useful to compute explicitly the orthogonal polynomials related with several of these families. Having these explicit formulas is essential when one is looking for applications of

2010 *Mathematics Subject Classification.* 42C05; 47S10; 33C45.

Key words and phrases. Matrix-valued orthogonal polynomials; Matrix-valued differential operators; Rodrigues formula.

The research of the first author was partially supported by CONICET and SECTYP-UNCUYO Argentina grant 06\M125. The research of the second author was partially supported by PGC2018-096504-B-C31 (FEDER(EU)/Ministerio de Ciencia e Innovación-Agencia Estatal de Investigación), FQM-262 and Feder-US-1254600 (FEDET(EU)/Junta de Andalucía).

these matrix-valued bispectral polynomials, such as in the problem of *time and band limiting* over a non-commutative ring and matrix-valued commuting operators, see [30, 10, 31, 11, 12, 32].

Recently, in [4], a new family of matrix-valued orthogonal polynomials of size 2×2 was introduced, which are common eigenfunctions of a differential operator of *hypergeometric type* (in the sense defined by Juan A. Tirao in [44]):

$$D = \frac{d^2}{dt^2} t(1-t) + \frac{d}{dt} (C - tU) - V, \quad \text{with } U, V, C \in \mathbb{C}^{2 \times 2}.$$

In particular, the polynomials $(P_n^{(\alpha, \beta, v)})_{n \geq 0}$ introduced in [4], orthogonal with respect to the weight matrix $W^{(\alpha, \beta, v)}$ given in (2.4) and (2.5), are common eigenfunctions of an hypergeometric operator with matrix eigenvalues Λ_n , which are diagonal matrices with no repetition in their entries. This fact could be especially useful if one intends to use this family of polynomials in the context of *time and band limiting*, where the commutativity of the matrix-valued eigenvalues $(\Lambda_n)_n$ could play an important role.

In this paper we give some structural formulas for the family of matrix-valued orthogonal polynomials introduced in [4]. In particular, in Section 3 we give a Rodrigues formula (see Theorem 3.1), which allows us to write this family of polynomials explicitly in terms of the classical Jacobi polynomials (see Corollary 3.3).

In Section 4, this Rodrigues formula allows us to compute the norms of the sequence of monic orthogonal polynomials and therefore, we can find the coefficients of the three-term recurrence relation and the Christoffel-Darboux identity for the sequence of orthonormal polynomials.

In Section 5, we obtain a Pearson equation (see Proposition 5.4), which allows us to prove that the sequence of derivatives of k -th order, $k \geq 1$, of the orthogonal polynomials is also orthogonal with respect to the weight matrix given explicitly in Proposition 5.3.

In Section 6, following the ideas in [34, Section 5.1], we use the Pearson equation to give explicit lowering and rising operators for the sequence of derivatives. Thus, we deduce a Rodrigues formula for these polynomials and find a matrix-valued differential operator that has these matrix-valued polynomials as common eigenfunctions.

Finally, in Section 7, we describe the algebra of second order differential operators associated with the weight matrix $W^{(\alpha, \beta, v)}$ given in (2.4) and (2.5). Indeed, for a given weight matrix W , the analysis of the algebra $D(W)$ of all differential operators that have a sequence of matrix-valued orthogonal polynomials with respect to W as eigenfunctions has received much attention in the literature in the last fifteen years [9, 33, 45, 42, 47, 6, 8]. While for classical orthogonal polynomials the structure of this algebra is very well known (see [39]), in the matrix setting, where this algebra is non commutative, the situation is highly non trivial.

2. PRELIMINARIES

In this section we give some background on matrix-valued orthogonal polynomials (see [23] for further details). A weight matrix W is a complex $N \times N$ matrix-valued integrable function on the interval (a, b) , such that W is positive definite almost everywhere and with finite moments of all orders, i.e., $\int_a^b t^n dW(t) \in \mathbb{C}^{N \times N}$, $n \in \mathbb{N}$. The weight matrix W induces a Hermitian sesquilinear form,

$$\langle P, Q \rangle_W = \int_a^b P(t) W(t) Q^*(t) dt,$$

for any pair of $N \times N$ matrix-valued functions $P(t)$ and $Q(t)$, where $Q^*(t)$ denotes the conjugate transpose of $Q(t)$.

A sequence $(P_n)_{n \geq 0}$ of orthogonal polynomials with respect to a weight matrix W is a sequence of matrix-valued polynomials satisfying that $P_n(t)$, $n \geq 0$, is a matrix polynomial of degree n with non-singular leading coefficient, and $\langle P_n, P_m \rangle_W = \Delta_n \delta_{n,m}$, where Δ_n , $n \geq 0$, is a positive definite matrix. When $\Delta_n = I$, here I denotes the identity matrix, we say that the polynomials $(P_n)_{n \geq 0}$ are orthonormal. In particular, when the leading coefficient of $P_n(t)$, $n \geq 0$, is the identity matrix, we say that the polynomials $(P_n)_{n \geq 0}$ are monic.

Given a weight matrix W , there exists a unique sequence of monic orthogonal polynomials $(P_n)_{n \geq 0}$ in $\mathbb{C}^{N \times N}[t]$, any other sequence of orthogonal polynomials $(Q_n)_{n \geq 0}$ can be written as $Q_n(t) = K_n P_n(t)$ for some non-singular matrix K_n .

Any sequence of monic orthogonal matrix-valued polynomials $(P_n)_{n \geq 0}$ satisfies a three-term recurrence relation

$$tP_n(t) = P_{n+1}(t) + B_n P_n(t) + A_n P_{n-1}(t), \quad \text{for } n \in \mathbb{N}_0,$$

where $P_{-1}(t) = 0$, $P_0(t) = I$. The $N \times N$ matrix coefficients A_n and B_n enjoy certain properties: in particular, A_n is non-singular for any n .

Two weights W and \widetilde{W} are said to be *equivalent* if there exists a non-singular matrix M , which does not depend on t , such that

$$\widetilde{W}(t) = MW(t)M^*, \quad \text{for all } t \in (a, b).$$

A weight matrix W *reduces* to a smaller size if there exists a non-singular matrix M such that

$$MW(t)M^* = \begin{pmatrix} W_1(t) & 0 \\ 0 & W_2(t) \end{pmatrix}, \quad \text{for all } t \in (a, b),$$

where W_1 and W_2 are weights of smaller size. A weight matrix W is said to be *irreducible* if it does not reduce to a smaller size (see [19], [46]).

Let D be a right-hand side ordinary differential operator with matrix valued polynomial coefficients,

$$D = \sum_{i=0}^s \partial^i F_i(t), \quad \partial^i = \frac{d^i}{dt^i}.$$

The operator D acts on a polynomial function $P(t)$ as $PD = \sum_{i=0}^s \partial^i P F_i(t)$.

We say that the differential operator D is symmetric with respect to W if

$$(2.1) \quad \langle PD, Q \rangle_W = \langle P, QD \rangle_W, \quad \text{for all } P, Q \in \mathbb{C}^{N \times N}[t].$$

The differential operator $D = \frac{d^2}{dt^2} F_2(t) + \frac{d}{dt} F_1(t) + F_0$ is symmetric with respect to W if and only if ([19, Theorem 3.1])

$$(2.2) \quad \begin{aligned} F_2 W &= W F_2^*, \\ 2(F_2 W)' &= F_1 W + W F_1^*, \\ (F_2 W)'' - (F_1 W)' &= W F_0^* - F_0 W, \end{aligned}$$

and

$$(2.3) \quad \lim_{t \rightarrow a, b} F_2(t) W(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow a, b} (F_1(t) W(t) - W(t) F_1^*(t)) = 0.$$

We will need the following result to find the Rodrigues' formula for the sequence of orthogonal polynomials with respect to a weight matrix W .

Theorem 2.1. ([18, Lemma 1.1]) *Let F_2 , F_1 and F_0 be matrix polynomials of degrees not larger than 2, 1, and 0, respectively. Let W , R_n be $N \times N$ matrix functions twice and n times differentiable, respectively, in an open set of the real line Ω . Assume that $W(t)$ is non-singular for $t \in \Omega$ and that satisfies the identity and the differential equations in (2.2). Define the functions P_n , $n \geq 1$, by*

$$P_n = R_n^{(n)} W^{-1}.$$

If for a matrix Λ_n , the function R_n satisfies

$$(R_n F_2^*)'' - (R_n [F_1^* + n (F_2^*)'])' + R_n [F_0^* + n (F_1^*)' + \binom{n}{2} (F_2^*)''] = \Delta_n R_n,$$

then the function P_n satisfies

$$P_n''(t) F_2(t) + P_n'(t) F_1(t) + P_n(t) F_0(t) = \Delta_n P_n(t).$$

2.1. The family of matrix-valued orthogonal polynomials. In [4] the authors introduce a Jacobi type weight matrix $W^{(\alpha, \beta, v)}(t)$ and a differential operator $D^{(\alpha, \beta, v)}$ such that $D^{(\alpha, \beta, v)}$ is symmetric with respect to the weight matrix $W^{(\alpha, \beta, v)}(t)$.

Let $\alpha, \beta, v \in \mathbb{R}$, $\alpha, \beta > -1$ and $|\alpha - \beta| < |v| < \alpha + \beta + 2$. We consider the weight matrix function

$$(2.4) \quad W^{(\alpha, \beta, v)}(t) = t^\alpha (1 - t)^\beta \widetilde{W}^{(\alpha, \beta, v)}(t), \quad \text{for } t \in (0, 1),$$

with

$$(2.5) \quad \widetilde{W}^{(\alpha, \beta, v)}(t) = \begin{pmatrix} \frac{v(\kappa_{v, \beta} + 2)}{\kappa_{v, -\beta}} t^2 - (\kappa_{v, \beta} + 2)t + (\alpha + 1) & (\alpha + \beta + 2)t - (\alpha + 1) \\ (\alpha + \beta + 2)t - (\alpha + 1) & -\frac{v(\kappa_{-v, -\beta} + 2)}{\kappa_{-v, -\beta}} t^2 - (\kappa_{-v, \beta} + 2)t + (\alpha + 1) \end{pmatrix},$$

where, for the sake of clearness in the rest of the paper, we use the notation:

$$(2.6) \quad \kappa_{\pm v, \pm \beta} = \alpha \pm v \pm \beta.$$

$W^{(\alpha, \beta, v)}$ is an irreducible weight matrix and the hypergeometric type differential operator given by

$$(2.7) \quad D^{(\alpha, \beta, v)} = \frac{d^2}{dt^2} F_2(t) + \frac{d}{dt} F_1(t) + F_0(t),$$

where

$$(2.8) \quad F_2(t) = t(1 - t), \quad F_1(t) = C^* - tU \quad \text{and} \quad F_0(t) = -V,$$

and

$$(2.9) \quad C = \begin{pmatrix} \alpha + 1 - \frac{\kappa_{-v, -\beta}}{v} & \frac{\kappa_{v, -\beta}}{v} \\ -\frac{\kappa_{-v, -\beta}}{v} & \alpha + 1 + \frac{\kappa_{v, -\beta}}{v} \end{pmatrix}, \quad U = (\alpha + \beta + 4)I \quad \text{and} \quad V = \begin{pmatrix} v & 0 \\ 0 & 0 \end{pmatrix},$$

is symmetric with respect to the weight matrix $W^{(\alpha, \beta, v)}$.

In the same paper, the authors also give the corresponding monic orthogonal polynomials in terms of the hypergeometric function ${}_2H_1(C, U, V; t)$ defined by J. A. Tirao in [44] and their three-term recurrence relation.

Proposition 2.2. [4, Theorem 4.3] *Let $(P_n^{(\alpha, \beta, v)})_{n \geq 0}$ be the sequence of matrix-valued monic orthogonal polynomials associated with the weight function $W^{(\alpha, \beta, v)}(t)$. Then, $P_n^{(\alpha, \beta, v)}$ is an eigenfunction of the differential operator $D^{(\alpha, \beta, v)}$ with diagonal eigenvalue*

$$(2.10) \quad \Lambda_n = \begin{pmatrix} \lambda_n & 0 \\ 0 & \mu_n \end{pmatrix}, \quad \begin{aligned} \lambda_n &= -n(n-1) - n(\alpha + \beta + 4) - v, \\ \mu_n &= -n(n-1) - n(\alpha + \beta + 4). \end{aligned}$$

Moreover ([4, Section 4.2]), the matrix-valued monic orthogonal polynomials $(P_n^{(\alpha, \beta, v)})_{n \geq 0}$ are given by

$$(2.11) \quad \begin{aligned} (P_n^{(\alpha, \beta, v)}(t))^* &= {}_2H_1(C, U, V + \lambda_n I; t) n! [C, U, V + \lambda_n I]_n^{-1} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \\ &\quad + {}_2H_1(C, U, V + \mu_n I; t) n! [C, U, V + \mu_n I]_n^{-1} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \end{aligned}$$

where

$${}_2H_1(C, U, V; t) = \sum_{k \geq 0} [C, U, V]_k \frac{t^k}{k!},$$

and $[C, U, V]_k$ is defined inductively as $[C, U, V]_0 = I$ and

$$(2.12) \quad [C, U, V]_{k+1} = (C + kI)^{-1} (k(k-1)I + kU + V) [C, U, V]_k.$$

Proposition 2.3. ([4, Theorem 3.12]) *The monic orthogonal polynomials $(P_n^{(\alpha, \beta, v)})_{n \geq 0}$ satisfy the three-term recurrence relation*

$$(2.13) \quad tP_n^{(\alpha, \beta, v)}(t) = P_{n+1}^{(\alpha, \beta, v)}(t) + B_n^{(\alpha, \beta, v)}P_n^{(\alpha, \beta, v)}(t) + A_n^{(\alpha, \beta, v)}P_{n-1}^{(\alpha, \beta, v)}(t)$$

where

$$(2.14) \quad A_n^{(\alpha, \beta, v)} = a_n^{(\alpha, \beta, v)} \begin{pmatrix} (4 + 2n + \kappa_{-v, \beta})(2n + \kappa_{v, \beta}) & 0 \\ 0 & (4 + 2n + \kappa_{v, \beta})(2n + \kappa_{-v, \beta}) \end{pmatrix},$$

with

$$(2.15) \quad a_n^{(\alpha, \beta, v)} = \frac{n(1+n+\alpha)(1+n+\beta)(2+n+\alpha+\beta)}{(1+2n+\alpha+\beta)(2+2n+\alpha+\beta)^2(3+2n+\alpha+\beta)(2+2n+\kappa_{-v, \beta})(2+2n+\kappa_{v, \beta})}, \quad n \geq 0,$$

and the entries of $B_n = B_n^{(\alpha, \beta, v)}$, $n \geq 0$, are

$$(2.16) \quad \begin{aligned} (B_n)_{11} &= -n \frac{(\alpha + n)v - \kappa_{-v, -\beta}}{(\alpha + \beta + 2n + 2)v} + (n + 1) \frac{(\alpha + n + 1)v - \kappa_{-v, -\beta}}{(\alpha + \beta + 2n + 4)v}, \\ (B_n)_{21} &= \frac{\kappa_{v, -\beta} (\kappa_{-v, \beta} + 2)}{v (\kappa_{-v, \beta} + 2n + 2) (\kappa_{-v, \beta} + 2n + 4)}, \\ (B_n)_{12} &= \frac{-\kappa_{-v, -\beta} (\kappa_{v, \beta} + 2)}{v (\kappa_{v, \beta} + 2n + 2) (\kappa_{v, \beta} + 2n + 4)}, \\ (B_n)_{22} &= -n \frac{(\alpha + n)v + \kappa_{v, -\beta}}{(\alpha + \beta + 2n + 2)v} + (n + 1) \frac{(\alpha + n + 1)v + \kappa_{v, -\beta}}{(\alpha + \beta + 2n + 4)v}. \end{aligned}$$

Using the symmetry condition (2.1) and the three-term recurrence relation (2.13) one can easily see that the coefficients $A_n^{(\alpha, \beta, v)}$ and $B_n^{(\alpha, \beta, v)}$ satisfy the identities:

$$(2.17) \quad A_n^{(\alpha, \beta, v)} \left\| P_{n-1}^{(\alpha, \beta, v)}(t) \right\|^2 = \left\| P_n^{(\alpha, \beta, v)}(t) \right\|^2,$$

$$(2.18) \quad \left(B_n^{(\alpha, \beta, v)} \left\| P_n^{(\alpha, \beta, v)}(t) \right\|^2 \right)^* = B_n^{(\alpha, \beta, v)} \left\| P_n^{(\alpha, \beta, v)}(t) \right\|^2.$$

3. RODRIGUES FORMULA

In this section we will provide a Rodrigues formula for the sequence of monic orthogonal polynomials $\left(P_n^{(\alpha, \beta, v)} \right)_{n \geq 0}$ with respect to the weight matrix $W = W^{(\alpha, \beta, v)}$ in (2.4). Moreover, the Rodrigues formula will allow us to find an explicit expression for the polynomials in terms of Jacobi polynomials.

Theorem 3.1. *Consider the weight matrix $W(t) = W^{(\alpha, \beta, v)}(t)$ given by the expression in (2.4) and (2.5). Consider the matrix-valued functions $(P_n)_{n \geq 0}$ and $(R_n)_{n \geq 0}$ defined by*

$$(3.1) \quad P_n(t) = (R_n(t))^{(n)} (W(t))^{-1},$$

$$(3.2) \quad R_n(t) = R_n^{(\alpha, \beta, v)}(t) = t^{n+\alpha} (1-t)^{n+\beta} \left(R_{n,2}^{(\alpha, \beta, v)} t^2 + R_{n,1}^{(\alpha, \beta, v)} t + R_{n,0}^{(\alpha, \beta, v)} \right),$$

with

$$\begin{aligned} R_{n,2}^{(\alpha, \beta, v)} = R_{n,2} &= \begin{pmatrix} c_n & 0 \\ 0 & d_n \end{pmatrix}, \\ R_{n,1}^{(\alpha, \beta, v)} = R_{n,1} &= \frac{1}{v} \begin{pmatrix} -c_n \kappa_{v, -\beta} & \frac{c_n (\alpha + 2n + 2 + \beta) \kappa_{v, -\beta}}{(\kappa_{v, \beta} + 2n + 2)} \\ -\frac{d_n (\alpha + 2n + 2 + \beta) \kappa_{-v, -\beta}}{(\kappa_{-v, \beta} + 2n + 2)} & d_n \kappa_{-v, -\beta} \end{pmatrix}, \\ R_{n,0}^{(\alpha, \beta, v)} = R_{n,0} &= \frac{1 + n + \alpha}{v} \begin{pmatrix} c_n \frac{\kappa_{v, -\beta}}{(\kappa_{v, \beta} + 2n + 2)} & -c_n \frac{\kappa_{v, -\beta}}{(\kappa_{v, \beta} + 2n + 2)} \\ d_n \frac{\kappa_{-v, -\beta}}{(\kappa_{-v, \beta} + 2n + 2)} & -d_n \frac{\kappa_{-v, -\beta}}{(\kappa_{-v, \beta} + 2n + 2)} \end{pmatrix}, \end{aligned}$$

where $(c_n)_n$ and $(d_n)_n$ are arbitrary sequences of complex numbers. Then $P_n(t)$ is a polynomial of degree n with non-singular leading coefficient equal to

$$\begin{pmatrix} \frac{\kappa_{v,-\beta}(\alpha + \beta + n + 3)_n c_n}{(-1)^n v(\kappa_{v,\beta} + 2)} & 0 \\ 0 & \frac{\kappa_{-v,-\beta}(\alpha + \beta + n + 3)_n d_n}{(-1)^{n+1} v(\kappa_{-v,\beta} + 2)} \end{pmatrix},$$

where $(a)_n = a(a+1)\dots(a+n-1)$ denotes the usual Pochhammer symbol. Moreover, if we put

$$(3.3) \quad c_n = \frac{(-1)^n v(\kappa_{v,\beta} + 2)}{\kappa_{v,-\beta}(\alpha + \beta + n + 3)_n}, \quad d_n = \frac{(-1)^{n+1} v(\kappa_{-v,\beta} + 2)}{\kappa_{-v,-\beta}(\alpha + \beta + n + 3)_n},$$

then $(P_n)_{n \geq 0}$ is a sequence of monic orthogonal polynomials with respect to W and $P_n = P_n^{(\alpha, \beta, v)}$.

Proof. Let W be the weight matrix given in (2.4) and F_2, F_1, F_0 and Λ_n are the polynomials coefficients defined in (2.8)-(2.10).

Following straightforward computations, we can prove that the matrix-valued function $R_n(t)$ satisfies the equation

$$(R_n F_2^*)'' - (R_n [F_1^* + n(F_2^*)'])' + R_n [F_0^* + n(F_1^*)'] + \binom{n}{2} (F_2^*)'' = \Lambda_n R_n.$$

Theorem 2.1 guarantees that the function $P_n(t) = (R_n(t))^n (W(t))^{-1}$ is an eigenfunction of $D^{(\alpha, \beta, v)}$ with eigenvalue Λ_n given in (2.10). Then $P_n^{(\alpha, \beta, v)}(t)$ and $P_n(t)$ satisfy the same differential equation.

We will prove that P_n is a polynomial of degree n with non-singular leading coefficient. We will use the following Rodrigues formula for the classical Jacobi polynomial $p_n^{(\alpha, \beta)}(t)$ ([43, Chapter IV])

$$\frac{d^n}{dt^n} [t^{n+\alpha} (1-t)^{n+\beta}] = n! t^\alpha (1-t)^\beta p_n^{(\alpha, \beta)}(1-2t),$$

where

$$(3.4) \quad p_n^{(\alpha, \beta)}(1-2t) = \frac{\Gamma(n+\alpha+1)}{n! \Gamma(n+\alpha+\beta+1)} \sum_{j=0}^n \binom{n}{j} \frac{\Gamma(n+\alpha+\beta+1+j)}{\Gamma(j+\alpha+1)} (-1)^j t^j.$$

Thus, we obtain

$$R_n^{(n)}(t) = n! t^\alpha (1-t)^\beta \left(p_n^{(\alpha+2, \beta)}(1-2t) R_{n,2} t^2 + p_n^{(\alpha+1, \beta)}(1-2t) R_{n,1} t + p_n^{(\alpha, \beta)}(1-2t) R_{n,0} \right).$$

We can rewrite $(W(t))^{-1}$ as

$$(W(t))^{-1} = t^{-\alpha-2} (1-t)^{-\beta-2} (J_2 t^2 + J_1 t + J_0),$$

with

$$J_2 = \begin{pmatrix} \frac{\kappa_{v,-\beta}}{v(\kappa_{v,\beta} + 2)} & 0 \\ 0 & -\frac{\kappa_{-v,-\beta}}{v(\kappa_{-v,\beta} + 2)} \end{pmatrix}, \quad J_0 = \frac{-\kappa_{v,-\beta}\kappa_{-v,-\beta}(\alpha + 1)}{v^2(\kappa_{v,\beta} + 2)(\kappa_{-v,\beta} + 2)} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix},$$

$$J_1 = \frac{\kappa_{v,-\beta}\kappa_{-v,-\beta}}{v^2} \begin{pmatrix} \frac{1}{(\kappa_{v,\beta} + 2)(\kappa_{-v,\beta} + 2)} & \frac{(\alpha + \beta + 2)}{(\kappa_{v,\beta} + 2)(\kappa_{-v,\beta} + 2)} \\ \frac{(\kappa_{v,\beta} + 2)}{(\alpha + \beta + 2)} & \frac{1}{(\kappa_{-v,\beta} + 2)} \end{pmatrix}.$$

Observe that $R_{n,0}J_0 = 0$. Thus, $P_n(t)$ becomes

$$P_n(t) = n!t^{-1}(1-t)^{-2} \left[p_n^{(\alpha+2,\beta)}(1-2t)R_{n,2}t(J_2t^2 + J_1t + J_0) \right. \\ \left. + p_n^{(\alpha+1,\beta)}(1-2t)R_{n,1}(J_2t^2 + J_1t + J_0) + p_n^{(\alpha,\beta)}(1-2t)R_{n,0}(J_2t + J_1) \right].$$

Hence, $P_n(t)$ is a polynomial of degree n if and only if $t = 0$ and $t = 1$ are zeros of the following polynomial

$$Q(t) = p_n^{(\alpha+2,\beta)}(1-2t)R_{n,2}t(J_2t^2 + J_1t + J_0) + p_n^{(\alpha+1,\beta)}(1-2t)R_{n,1}(J_2t^2 + J_1t + J_0) \\ + p_n^{(\alpha,\beta)}(1-2t)R_{n,0}(J_2t + J_1)$$

and $t = 1$ has multiplicity two, i.e., $Q(0) = Q(1) = Q'(1) = 0$.

Taking into account that $p_n^{(\alpha,\beta)}(1) = \frac{\Gamma(n+\alpha+1)}{n!\Gamma(\alpha+1)}$ and $p_n^{(\alpha,\beta)}(-1) = (-1)^n \frac{\Gamma(n+\beta+1)}{n!\Gamma(\beta+1)}$ we have

$$Q(0) = p_n^{(\alpha+1,\beta)}(1)R_{n,1}J_0 + p_n^{(\alpha,\beta)}(1)R_{n,0}J_1 = \frac{\Gamma(n+\alpha+1)}{n!\Gamma(\alpha+1)} \left(\frac{n+\alpha+1}{\alpha+1}R_{n,1}J_0 + R_{n,0}J_1 \right) = \mathbf{0},$$

$$Q(1) = (-1)^n \frac{\Gamma(n+\beta+1)}{n!\Gamma(\beta+1)} (R_{n,2} + R_{n,1} + R_{n,0})(J_2 + J_1 + J_0) = \mathbf{0}.$$

Now, by taking derivative of $Q(t)$ with respect to t and considering the identity

$$\left(p_n^{(\alpha,\beta)} \right)'(-1) = \frac{(\beta + \alpha + n + 1)(-1)^{n-1}\Gamma(n+\beta+1)}{2(n-1)!\Gamma(\beta+2)},$$

we obtain

$$Q'(1) = -2 \left(\left(p_n^{(\alpha+2,\beta)} \right)'(-1)R_{n,2} + \left(p_n^{(\alpha+1,\beta)} \right)'(-1)R_{n,1} + \left(p_n^{(\alpha,\beta)} \right)'(-1)R_{n,0} \right) (J_2 + J_1 + J_0) \\ + p_n^{(\alpha+2,\beta)}(-1)R_{n,2}(3J_2 + 2J_1 + J_0) + p_n^{(\alpha+1,\beta)}(-1)R_{n,1}(2J_2 + J_1) + p_n^{(\alpha,\beta)}(-1)R_{n,0}J_2 = 0.$$

This shows that $Q(t)$ is divisible by $t(t-1)^2$ therefore, $P_n(t)$ is a polynomial of degree n since $\deg(Q(t)) = n + 3$.

Observe that the leading coefficient of $P_n(t)$ is determined by the leading coefficient of $n!p_n^{(\alpha+2,\beta)}(1-2t)R_{n,2}J_2t^4$. Considering (3.4) we have

$$\frac{(-1)^n \Gamma(2n + \alpha + \beta + 3)}{\Gamma(n + \alpha + \beta + 3)} R_{n,2} J_2 = \frac{(-1)^n (\alpha + \beta + n + 3)_n}{v} \begin{pmatrix} \frac{\kappa_{v,-\beta}}{\kappa_{v,\beta} + 2} c_n & 0 \\ 0 & -\frac{\kappa_{-v,-\beta}}{\kappa_{-v,\beta} + 2} d_n \end{pmatrix}.$$

The previous matrix coefficient is non-singular since $|\alpha - \beta| < |v| < \alpha + \beta + 2$.

Moreover, if (3.3) holds true then $P_n(t)$ is a monic polynomial and equal to $P_n^{(\alpha,\beta,v)}(t)$. \square

Corollary 3.2. *Consider the weight matrix $W^{(\alpha,\beta,v)}(t)$ given in (2.4) and (2.5). Then, the monic orthogonal polynomials $P_n^{(\alpha,\beta,v)}(t)$ satisfy the Rodrigues formula*

$$P_n^{(\alpha,\beta,v)}(t) = (R_n^{(\alpha,\beta,v)}(t))^{(n)} \left(W^{(\alpha,\beta,v)}(t) \right)^{-1}.$$

We can see in the proof of Theorem 3.1 that Rodrigues' formula allows us to find an explicit expression for the polynomials in terms of the classical Jacobi polynomials.

Corollary 3.3. *Consider the matrix-valued function $\widetilde{W}^{(\alpha,\beta,v)}(t)$ given in (2.5) and let $R_{n,i}^{(\alpha,\beta,v)}$, $i = 0, 1, 2$, be as in Theorem 3.1. Define the coefficients c_n and d_n as in (3.3). Then, the sequence of monic orthogonal polynomials $(P_n)_{n \geq 0}$, defined by (3.1) can be written as*

$$(3.5) \quad P_n(t) = n! \left(p_n^{(\alpha+2,\beta)}(1-2t)R_{n,2}^{(\alpha,\beta,v)}t^2 + p_n^{(\alpha+1,\beta)}(1-2t)R_{n,1}^{(\alpha,\beta,v)}t + p_n^{(\alpha,\beta)}(1-2t)R_{n,0}^{(\alpha,\beta,v)} \right) (\widetilde{W}^{(\alpha,\beta,v)}(t))^{-1}.$$

Moreover,

$$(3.6) \quad P_n(t) = n! \left(p_n^{(\alpha,\beta)}(1-2t)\mathcal{C}_{n,2}^{(\alpha,\beta,v)} + p_{n+1}^{(\alpha,\beta)}(1-2t)\mathcal{C}_{n,1}^{(\alpha,\beta,v)} + p_{n+2}^{(\alpha,\beta)}(1-2t)\mathcal{C}_{n,0}^{(\alpha,\beta,v)} \right) (\widetilde{W}^{(\alpha,\beta,v)}(t))^{-1},$$

with

$$\begin{aligned} \mathcal{C}_{n,2}^{(\alpha,\beta,v)} &= \frac{(\beta + n + 1)(\alpha + n + 1)}{(\alpha + \beta + 2n + 2)(\alpha + \beta + 2n + 3)} \begin{pmatrix} \frac{c_n(\kappa_{-v,\beta} + 2n + 4)}{\kappa_{v,\beta} + 2n + 2} & 0 \\ 0 & \frac{d_n(\kappa_{v,\beta} + 2n + 4)}{\kappa_{-v,\beta} + 2n + 2} \end{pmatrix}, \\ \mathcal{C}_{n,1}^{(\alpha,\beta,v)} &= \frac{n+1}{v} \begin{pmatrix} \frac{(\alpha - \beta)(\kappa_{-v,\beta} + 2n + 4)c_n}{(\alpha + \beta + 2n + 2)(\alpha + \beta + 2n + 4)} & -\frac{c_n\kappa_{v,-\beta}}{\kappa_{v,\beta} + 2n + 2} \\ \frac{d_n\kappa_{-v,-\beta}}{\kappa_{-v,\beta} + 2n + 2} & -\frac{(\alpha - \beta)(\kappa_{v,\beta} + 2n + 4)d_n}{(\alpha + \beta + 2n + 2)(\alpha + \beta + 2n + 4)} \end{pmatrix}, \\ \mathcal{C}_{n,0}^{(\alpha,\beta,v)} &= \frac{(n+1)(n+2)}{(\alpha + \beta + 2n + 4)(\alpha + \beta + 2n + 3)} \begin{pmatrix} c_n & 0 \\ 0 & d_n \end{pmatrix}. \end{aligned}$$

Proof. The expression in (3.5) follows from the proof above and to obtain (3.6) we use the following property for the classical Jacobi polynomials $p_n^{(\alpha,\beta)}(t)$ ([43, Section 4.5]):

$$p_n^{(\alpha+1,\beta)}(1-2t) = \frac{(n + \alpha + 1)p_n^{(\alpha,\beta)}(1-2t) - (n + 1)p_{n+1}^{(\alpha,\beta)}(1-2t)}{(2n + \alpha + \beta + 2)t}$$

in (3.5). □

4. ORTHONORMAL POLYNOMIALS

In this section we give an explicit expression for the norm of the matrix-valued polynomials $\left(P_n^{(\alpha, \beta, v)}\right)_{n \geq 0}$. In addition, for the sequence of orthonormal polynomials we show the three-term recurrence relation and the Christoffel-Darboux formula, introduced for a general sequence of matrix-valued orthogonal polynomials in [14] (see also [23]).

Proposition 4.1. *The norm of the monic orthogonal polynomials $P_n^{(\alpha, \beta, v)}(t)$, $n \geq 0$, is determined by*

$$(4.1) \quad \left\|P_n^{(\alpha, \beta, v)}\right\|^2 = \frac{n!vB(\alpha+n+2, \beta+n+2)}{(\alpha+n+3+\beta)_n} \begin{pmatrix} \frac{(\kappa_{v, \beta}+2)(\kappa_{-v, \beta}+2n+4)}{\kappa_{v, -\beta}(\kappa_{v, \beta}+2n+2)} & 0 \\ 0 & -\frac{(\kappa_{-v, \beta}+2)(\kappa_{v, \beta}+2n+4)}{\kappa_{-v, -\beta}(\kappa_{-v, \beta}+2n+2)} \end{pmatrix}.$$

Therefore the sequence of polynomials

$$\tilde{P}_n^{(\alpha, \beta, v)}(t) = \left\|P_n^{(\alpha, \beta, v)}\right\|^{-1} P_n^{(\alpha, \beta, v)}(t)$$

is orthonormal with respect to W .

Proof. Let $P_n^{(\alpha, \beta, v)}(t) = \sum_{k=0}^n \mathcal{P}_n^k t^k$, using Rodrigues' formula we have

$$\left\|P_n^{(\alpha, \beta, v)}\right\|^2 = \int_0^1 P_n^{(\alpha, \beta, v)}(t) W(t) \left(P_n^{(\alpha, \beta, v)}(t)\right)^* dt = \sum_{k=0}^n \int_0^1 (R_n(t))^{(n)} \left(\mathcal{P}_n^k\right)^* t^k dt.$$

Integrating by parts n times we have,

$$\begin{aligned} \left\|P_n^{(\alpha, \beta, v)}\right\|^2 &= (-1)^n \sum_{k=0}^n \int_0^1 R_n(t) \frac{d^n}{dt^n} \left(\mathcal{P}_n^k\right)^* t^k dt = (-1)^n \int_0^1 R_n(t) \frac{d^n}{dt^n} t^n dt \\ &= (-1)^n n! \int_0^1 R_n(t) dt = (-1)^n n! \int_0^1 t^{n+\alpha} (1-t)^{n+\beta} (R_{n,2} t^2 + R_{n,1} t + R_{n,0}) dt \\ &= (-1)^n n! [B(\alpha+n+3, \beta+n+1) R_{n,2} + B(\alpha+n+2, \beta+n+1) R_{n,1} \\ &\quad + B(\alpha+n+1, \beta+n+1) R_{n,0}], \end{aligned}$$

where $B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt$ is the Beta function. Using the following property,

$$B(x+1, y) = \frac{x}{x+y} B(x, y)$$

we obtain,

$$\begin{aligned} \left\|P_n^{(\alpha, \beta, v)}\right\|^2 &= (-1)^n n! B(\alpha+n+1, \beta+n+1) \\ &\quad \left(\frac{(\alpha+n+1)(\alpha+n+2)}{(\alpha+\beta+2n+2)(\alpha+\beta+2n+3)} R_{n,2} + \frac{\alpha+n+1}{\alpha+\beta+2n+2} R_{n,1} + R_{n,0} \right). \end{aligned}$$

Using the expressions in (3.2), after some straightforward computations we complete the proof. □

The sequence of orthonormal polynomials satisfies the following properties.

Proposition 4.2. *The orthonormal polynomials $\left(\tilde{P}_n^{(\alpha,\beta,v)}\right)_{n \geq 0}$ satisfy the three-term recurrence relation*

$$(4.2) \quad t\tilde{P}_n^{(\alpha,\beta,v)}(t) = \tilde{A}_{n+1}^{(\alpha,\beta,v)} \tilde{P}_{n+1}^{(\alpha,\beta,v)}(t) + \tilde{B}_n^{(\alpha,\beta,v)} \tilde{P}_n^{(\alpha,\beta,v)}(t) + \left(\tilde{A}_n^{(\alpha,\beta,v)}\right)^* \tilde{P}_{n-1}^{(\alpha,\beta,v)}(t),$$

with

$$\begin{aligned} \tilde{A}_{n+1}^{(\alpha,\beta,v)} &= \left\|P_n^{(\alpha,\beta,v)}\right\|^{-1} \left\|P_{n+1}^{(\alpha,\beta,v)}\right\|, \\ \tilde{B}_n^{(\alpha,\beta,v)} &= \left\|P_n^{(\alpha,\beta,v)}\right\|^{-1} B_n^{(\alpha,\beta,v)} \left\|P_n^{(\alpha,\beta,v)}\right\|, \end{aligned}$$

where $B_n^{(\alpha,\beta,v)}$ is the coefficient of the three-term recurrence relation for the monic orthogonal polynomials $\left(P_n^{(\alpha,\beta,v)}\right)_{n \geq 0}$ (2.13). Clearly, $\tilde{B}_n^{(\alpha,\beta,v)}$ is a symmetric matrix.

Proof. By replacing the identity $\tilde{P}_n^{(\alpha,\beta,v)}(t) = \left\|P_n^{(\alpha,\beta,v)}\right\|^{-1} P_n^{(\alpha,\beta,v)}(t)$ in the three-term recurrence relation (2.13) and using identity (2.17) we obtain (4.2), and by (2.18) one verifies that $\left(\tilde{B}_n^{(\alpha,\beta,v)}\right)^* = \tilde{B}_n^{(\alpha,\beta,v)}$. \square

We also have the following Christoffel-Darboux formula for the sequence of orthonormal polynomials $\left(\tilde{P}_n^{(\alpha,\beta,v)}\right)_{n \geq 0}$:

$$\begin{aligned} \sum_{k=0}^n \left(\tilde{P}_k^{(\alpha,\beta,v)}\right)^*(y) \tilde{P}_k^{(\alpha,\beta,v)}(x) &= \frac{\left(\tilde{P}_n^{(\alpha,\beta,v)}\right)^*(y) \left(\tilde{A}_{n+1}^{(\alpha,\beta,v)}\right)^* \tilde{P}_{n+1}^{(\alpha,\beta,v)}(x)}{x - y} \\ &\quad - \frac{\left(\tilde{P}_{n+1}^{(\alpha,\beta,v)}\right)^*(y) \tilde{A}_{n+1}^{(\alpha,\beta,v)} \tilde{P}_n^{(\alpha,\beta,v)}(x)}{x - y}. \end{aligned}$$

Hence, the sequence of monic polynomials $\left(P_n^{(\alpha,\beta,v)}\right)_{n \geq 0}$ satisfies

$$\begin{aligned} \sum_{k=0}^n \left(P_k^{(\alpha,\beta,v)}\right)^*(y) \left\|P_k^{(\alpha,\beta,v)}\right\|^{-2} P_k^{(\alpha,\beta,v)}(x) &= \frac{\left(P_n^{(\alpha,\beta,v)}\right)^*(y) \left\|P_n^{(\alpha,\beta,v)}\right\|^{-2} P_{n+1}^{(\alpha,\beta,v)}(x)}{x - y} \\ &\quad - \frac{\left(P_{n+1}^{(\alpha,\beta,v)}\right)^*(y) \left\|P_n^{(\alpha,\beta,v)}\right\|^{-2} P_n^{(\alpha,\beta,v)}(x)}{x - y}, \end{aligned}$$

where the explicit expression of $\left\|P_n^{(\alpha,\beta,v)}\right\|^{-2}$ follows from (4.1).

5. THE DERIVATIVES OF THE ORTHOGONAL MATRIX-VALUED POLYNOMIALS

In this section we prove that polynomials in the sequence of derivatives of the orthogonal matrix polynomials $\left(P_n^{(\alpha,\beta,v)}\right)_{n \geq 0}$ are also orthogonal by obtaining a Pearson equation for the weight matrix $W^{(\alpha,\beta,v)}(t)$.

Let $\frac{d^k}{dt^k} P_n^{(\alpha,\beta,v)}(t)$ be the derivative of order k of the monic polynomial $P_n^{(\alpha,\beta,v)}(t)$, for $n \geq k$. Then

$$(5.1) \quad P_n^{(\alpha,\beta,v,k)}(t) = \frac{(n-k)!}{n!} \frac{d^k}{dt^k} P_n^{(\alpha,\beta,v)}(t)$$

are monic polynomials of degree $n-k$ for all $n \geq k$.

The polynomial $P_n^{(\alpha,\beta,v)}(t)$ is an eigenfunction of the operator $D^{(\alpha,\beta,v)}$ given above in (2.7)-(2.9).

Taking derivative k times we have that $P_n^{(\alpha,\beta,v,k)}(t)$ is an eigenfunction of the differential hypergeometric operator

$$(5.2) \quad D^{(k)} = D^{(\alpha,\beta,v,k)} = \frac{d^2}{dt^2} t(1-t) + \frac{d}{dt} ((C^{(k)})^* - tU^{(k)}) - V,$$

with

$$C^{(k)} = C + kI, \quad U^{(k)} = U + 2kI = (\alpha + \beta + 4 + 2k) I,$$

where C , U and V are the matrix entries of the operator $D^{(\alpha,\beta,v)}$ given in (2.9). One has that

$$P_n^{(\alpha,\beta,v,k)} D^{(\alpha,\beta,v,k)} = \Lambda_n^{(k)} P_n^{(\alpha,\beta,v,k)}, \quad n \geq k,$$

where $\Lambda_n^{(k)} = \Lambda_n + kU + k(k-1)I$, with Λ_n given in (2.10). One has in particular, the standard expression for the eigenvalue shown in [4, Proposition 3.3], $\Lambda_n^{(k)} = -(n-k)(n-k-1)I - (n-k)U^{(k)} - V$. More explicitly,

$$(5.3) \quad \Lambda_n^{(k)} = \begin{pmatrix} \lambda_n^{(k)} & 0 \\ 0 & \mu_n^{(k)} \end{pmatrix}, \quad \begin{aligned} \lambda_n^{(k)} &= -(n-k)(\alpha + \beta + 3 + n + k) - v, \\ \mu_n^{(k)} &= -(n-k)(\alpha + \beta + 3 + n + k). \end{aligned}$$

Remark 5.1. One notices that $D^{(\alpha,\beta,v,k)} = D^{(\alpha+k,\beta+k,v)}$. Thus, the sequence of derivatives are still common eigenfunctions of an hypergeometric operator with diagonal matrix eigenvalues $\Lambda_n^{(k)}$, with no repetition in their entries.

Proposition 5.2. As in (2.11), we have the following explicit expression for the sequence of polynomials $\left(P_n^{(\alpha,\beta,v,k)}\right)_{n \geq k}$ in terms of hypergeometric functions

$$(5.4) \quad \begin{aligned} \left(P_n^{(\alpha,\beta,v,k)}(t)\right)^* &= {}_2H_1\left(C^{(k)}, U^{(k)}, V + \lambda_n^{(k)} I; t\right) (n-k)! \left[C^{(k)}, U^{(k)}, V + \lambda_n^{(k)} I\right]_{n-k}^{-1} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \\ & {}_2H_1\left(C^{(k)}, U^{(k)}, V + \mu_n^{(k)} I; t\right) (n-k)! \left[C^{(k)}, U^{(k)}, V + \mu_n^{(k)} I\right]_{n-k}^{-1} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \end{aligned}$$

where $C^{(k)}$, $U^{(k)}$ and V are the entries of the differential operator in (5.2) and $\lambda_n^{(k)}$ and $\mu_n^{(k)}$ are the diagonal entries of the matrix eigenvalue $\Lambda_n^{(k)}$ given in (5.3).

We include the proof for completeness.

Proof. Indeed, the polynomials $\left(P_n^{(\alpha, \beta, v, k)}\right)_{n \geq k}$ are common eigenfunctions of the matrix hypergeometric type operator (5.2) with diagonal eigenvalue $\Lambda_n^{(k)}$.

The fact that the eigenvalue is diagonal implies that the matrix equation can be written as two vectorial hypergeometric equations as in [44, Theorem 5] and the solutions of these equations are the columns of $\left(P_n^{(\alpha, \beta, v, k)}\right)_{n \geq k}$. Since the eigenvalues of the matrices $C^{(k)}$, $3 + \alpha + k$ and $1 + \alpha + k$, are non negative integers for all $k \geq 1$, then these solutions are hypergeometric vector functions.

Moreover, the vectorial functions are polynomials of degree $n - k$ since the form of the factor $\left((n - k)(n - k - 1)I + (n - k)U^{(k)} + V + \lambda_n^{(k)}I\right) = -\Lambda_n^{(k)} + \lambda_n^{(k)}I$ appearing in the expression of $\left[C^{(k)}, U^{(k)}, V + \lambda_n^{(k)}I\right]_{n-k+1}$ (see 2.12), makes its first column equal to zero. Analogously for the second column of $\left[C^{(k)}, U^{(k)}, V + \mu_n^{(k)}I\right]_{n-k+1}$.

The matrices $\left[C^{(k)}, U^{(k)}, V + \mu_n^{(k)}I\right]_{n-k}$ and $\left[C^{(k)}, U^{(k)}, V + \lambda_n^{(k)}I\right]_{n-k}$ are non-singular, since $\lambda_q^{(k)} \neq \mu_\ell^{(k)}$, $\lambda_q^{(k)} \neq \lambda_\ell^{(k)}$ and $\mu_q^{(k)} \neq \mu_\ell^{(k)}$ for all $q \neq \ell$. □

Proposition 5.3. *Let $\alpha, \beta > -(k + 1)$ and $|\alpha - \beta| < |v| < \alpha + \beta + 2(k + 1)$. We write*

$$(5.5) \quad W^{(k)}(t) = W^{(\alpha, \beta, v, k)}(t) = t^{\alpha+k} (1-t)^{\beta+k} \widetilde{W}^{(\alpha, \beta, v, k)}(t), \text{ where } \widetilde{W}^{(\alpha, \beta, v, k)}(t) = W_2^{(k)} t^2 + W_1^{(k)} t + W_0^{(k)},$$

with

$$\begin{aligned} W_2^{(k)} &= v \begin{pmatrix} \frac{\kappa_{v, \beta} + 2(k+1)}{\kappa_{v, -\beta}} & 0 \\ 0 & -\frac{\kappa_{-v, \beta} + 2(k+1)}{\kappa_{-v, -\beta}} \end{pmatrix}, \quad W_0^{(k)} = (\alpha + k + 1) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \\ W_1^{(k)} &= \begin{pmatrix} -\kappa_{v, \beta} & \alpha + \beta \\ \alpha + \beta & -\kappa_{-v, \beta} \end{pmatrix} + 2(k+1) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}. \end{aligned}$$

Then $W^{(k)}$ is an irreducible weight matrix and the differential hypergeometric operator $D^{(k)}$ in (5.2) is symmetric with respect to the weight matrix $W^{(k)}$. Moreover, it holds that $W^{(k)}(t) = W^{(\alpha+k, \beta+k, v)}(t)$.

Proof. Taking into account Remark 5.1 and the fact that $W^{(k)}(t) = W^{(\alpha+k, \beta+k, v)}(t)$, from Proposition 4.1 in [4] one has that $D^{(\alpha+k, \beta+k, v)}$ is symmetric with respect to $W^{(\alpha+k, \beta+k, v)}$ and $W^{(\alpha+k, \beta+k, v)}$ is an irreducible weight matrix if and only if $\alpha + k > -1$, $\beta + k > -1$ and $|(\alpha + k) - (\beta + k)| < |v| < (\alpha + k) + (\beta + k) + 2$. □

We will use the following Pearson equation to prove that the sequence of polynomials $\left(P_n^{(\alpha, \beta, v, k)}\right)_{n \geq k}$ is orthogonal with respect to $W^{(k)}$.

Theorem 5.4. *The weight matrix $W^{(k)}$ satisfies the following Pearson equation,*

$$(5.6) \quad \left(W^{(k)}(t) \Phi^{(k)}(t) \right)' = W^{(k)}(t) \Psi^{(k)}(t), \quad k \in \mathbb{N}, \quad \text{with}$$

$$\Phi^{(k)}(t) = \mathcal{A}_2^k t^2 + \mathcal{A}_1^k t + \mathcal{A}_0^k \quad \text{and} \quad \Psi^{(k)}(t) = \mathcal{B}_1^k t + \mathcal{B}_0^k,$$

where

$$(5.7) \quad \mathcal{A}_2^k = \begin{pmatrix} -\frac{\kappa_{v,\beta} + 2(k+2)}{\kappa_{v,\beta} + 2(k+1)} & 0 \\ 0 & -\frac{\kappa_{-v,\beta} + 2(k+2)}{\kappa_{-v,\beta} + 2(k+1)} \end{pmatrix},$$

$$(5.8) \quad \mathcal{A}_1^k = \frac{2}{(\kappa_{-v,\beta} + 2(k+1))(\kappa_{v,\beta} + 2(k+1))} \begin{pmatrix} 0 & \kappa_{v,-\beta} \\ \kappa_{-v,-\beta} & 0 \end{pmatrix} - \mathcal{A}_2^k,$$

$$(5.9) \quad \mathcal{A}_0^k = \frac{\kappa_{v,-\beta} \kappa_{-v,-\beta}}{v(\kappa_{-v,\beta} + 2(k+1))(\kappa_{v,\beta} + 2(k+1))} \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix},$$

$$(5.10) \quad \mathcal{B}_1^k = (\alpha + \beta + 4 + 2k) \mathcal{A}_2^k,$$

$$(5.11) \quad \mathcal{B}_0^k = \left(-(\alpha + k + 1)I - \frac{1}{v} \begin{pmatrix} -\kappa_{-v,-\beta} & 0 \\ 0 & \kappa_{v,-\beta} \end{pmatrix} \right) \mathcal{A}_2^k$$

$$+ \frac{1}{2v} \left(\frac{\alpha + \beta + 2k + 4}{v} \mathcal{A}_1^k + \mathcal{B}_1^k \right) \begin{pmatrix} -\kappa_{-v,\beta} - 2(k+1) & 0 \\ 0 & \kappa_{v,\beta} + 2(k+1) \end{pmatrix}.$$

Proof. By replacing the expression of $\Phi^{(k)}(t)$ and $\Psi^{(k)}(t)$ in (5.6) and taking derivative we obtain

$$(5.12) \quad \left(W^{(k)}(t) \right)' \left(\mathcal{A}_2^k t^2 + \mathcal{A}_1^k t + \mathcal{A}_0^k \right) - W^{(k)}(t) \left(\left(\mathcal{B}_1^k - 2\mathcal{A}_2^k \right) t + \mathcal{B}_0^k - \mathcal{A}_1^k \right) = 0.$$

The derivative of $W^{(k)}(t)$ is

$$\left(W^{(k)}(t) \right)' = -t^{\alpha+k-1} (1-t)^{\beta+k-1} (\alpha + \beta + 2k + 2) W_2^{(k)} t^3 + (\alpha + k) W_0^{(k)}$$

$$+ \left[(\alpha + k + 2) W_2^{(k)} - (\alpha + \beta + 2k + 1) W_1^{(k)} \right] t^2 + \left[(\alpha + k + 1) W_1^{(k)} - (\alpha + \beta + 2k) W_0^{(k)} \right] t.$$

Hence, the left hand side of (5.12) is a product between a polynomial of degree five and $t^{(\alpha+k-1)} (1-t)^{(\beta+k-1)}$. Therefore, equating to zero the entries of this polynomial, taking into account (5.10) and the equality $W_0^{(k)} \mathcal{A}_0^k = 0$, it only remains to verify the identities below, which

follow immediately by straightforward computations.

$$\begin{aligned}
& (\alpha + k + 4) W_2^{(k)} \mathcal{A}_2^k - (\alpha + \beta + 2k + 3) \left(W_2^{(k)} \mathcal{A}_1^k + W_1^{(k)} \mathcal{A}_2^k \right) \\
& \quad + W_2^{(k)} \left(\mathcal{B}_0^k - \mathcal{B}_1^k \right) + W_1^{(k)} \mathcal{B}_1^k = 0, \\
& (\alpha + k + 3) \left(W_2^{(k)} \mathcal{A}_1^k + W_1^{(k)} \mathcal{A}_2^k \right) - (\alpha + \beta + 2k + 2) \left(W_2^{(k)} \mathcal{A}_0^k + W_1^{(k)} \mathcal{A}_1^k \right) - W_2^{(k)} \mathcal{B}_0^k \\
& \quad + W_1^{(k)} \left(\mathcal{B}_0^k - \mathcal{B}_1^k \right) + 2W_0^{(k)} \mathcal{A}_2^k = 0, \\
& (\alpha + k + 2) \left(W_2^{(k)} \mathcal{A}_0^k + W_1^{(k)} \mathcal{A}_1^k + W_0^{(k)} \mathcal{A}_2^k \right) - (\alpha + \beta + 2k + 1) \left(W_1^{(k)} \mathcal{A}_0^k + W_0^{(k)} \mathcal{A}_1^k \right) \\
& \quad - W_1^{(k)} \mathcal{B}_0^k + W_0^{(k)} \left(\mathcal{B}_0^k - \mathcal{B}_1^k \right) = 0, \\
& (\alpha + k + 1) \left(W_1^{(k)} \mathcal{A}_0^k + W_0^{(k)} \mathcal{A}_1^k \right) - W_0^{(k)} \mathcal{B}_0^k = 0.
\end{aligned}$$

□

Remark 5.5. Let us consider the matrix-valued functions $W^{(\alpha, \beta, v, k)}(t) = W^{(k)}(t)$, $\Phi^{(k)}(t)$ and $\Psi^{(k)}(t)$, $k \in \mathbb{N}$, defined in (5.5) and Theorem 5.4 respectively. Then, by straightforward computations, one can verify the following identities:

$$(5.13) \quad W^{(\alpha, \beta, v, k+1)}(t) = W^{(\alpha, \beta, v, k)}(t) \Phi^{(k)}(t),$$

$$(5.14) \quad \left(W^{(\alpha, \beta, v, k+1)}(t) \right)' = W^{(\alpha, \beta, v, k)}(t) \Psi^{(k)}(t),$$

Taking into account that $\deg(\Phi^{(k)}(t)) = 2$ and $\deg(\Psi^{(k)}(t)) = 1$, we obtain from [5, Corollary 3.10] the following

Corollary 5.6. The sequence of polynomials $\left(P_n^{(\alpha, \beta, v, k)} \right)_{n \geq k}$ is orthogonal with respect to the weight matrix $W^{(k)} = W^{(\alpha, \beta, v, k-1)}(t) \Phi^{(k-1)}(t)$.

The following results are obtained in a similar way than in Theorem 3.1 and Corollary 3.3.

Proposition 5.7. Let $W^{(k)}(t)$ be defined as in (5.5). A Rodrigues formula for the sequence of polynomials $\left(P_n^{(\alpha, \beta, v, k)} \right)_{n \geq k}$ is

$$P_n^{(\alpha, \beta, v, k)}(t) = \left(R_n^{(\alpha, \beta, v, k)}(t) \right)^{(n-k)} \left(W^{(k)}(t) \right)^{-1}, \text{ where } R_n^{(\alpha, \beta, v, k)}(t) = R_{n-k}^{(\alpha+k, \beta+k, v)}(t).$$

Corollary 5.8. Let the matrix-valued function $W^{(k)}(t)$, and the matrices $R_{n-k,2}^{(\alpha, \beta, v)}$, $R_{n-k,1}^{(\alpha, \beta, v)}$ and $R_{n-k,0}^{(\alpha, \beta, v)}$ be defined as in (5.5) and (3.2). From the Rodrigues formula we get the explicit expressions

for the sequence of polynomials $\left(P_n^{(\alpha,\beta,v,k)}\right)_{n \geq k}$ in terms of the classical Jacobi polynomials $p_n^{(\alpha,\beta)}(t)$,

$$P_n^{(\alpha,\beta,v,k)}(t) = (n-k)! \left(p_{n-k}^{(\alpha+2+k,\beta+k)}(1-2t) R_{n-k,2}^{(\alpha+k,\beta+k,v)} t^2 + p_{n-k}^{(\alpha+1+k,\beta+k)}(1-2t) R_{n-k,1}^{(\alpha+k,\beta+k,v)} t \right. \\ \left. + p_{n-k}^{(\alpha+k,\beta+k)}(1-2t) R_{n-k,0}^{(\alpha+k,\beta+k,v)} \right) \left(\widetilde{W}^{(k)} \right)^{-1}$$

and

$$P_n^{(\alpha,\beta,v,k)}(t) = (n-k)! \left(p_{n-k}^{(\alpha+k,\beta+k)}(1-2t) \mathcal{C}_{n-k,2}^{(\alpha+k,\beta+k,v)} + p_{n-k+1}^{(\alpha+k,\beta+k)}(1-2t) \mathcal{C}_{n-k,1}^{(\alpha+k,\beta+k,v)} \right. \\ \left. + p_{n-k+2}^{(\alpha+k,\beta+k)}(1-2t) \mathcal{C}_{n-k,0}^{(\alpha+k,\beta+k,v)} \right) \left(\widetilde{W}^{(k)} \right)^{-1},$$

with $\mathcal{C}_{n-k,i}^{(\alpha+k,\beta+k,v)}$, $i = 0, 1, 2$, given by (3.6).

Proposition 5.9. *The orthogonal monic polynomials $\left(P_n^{(\alpha,\beta,v,k)}\right)_{n \geq k}$ satisfy the three-term recurrence relation*

$$tP_n^{(\alpha,\beta,v,k)}(t) = P_{n+1}^{(\alpha,\beta,v,k)}(t) + B_n^{(\alpha,\beta,v,k)} P_n^{(\alpha,\beta,v,k)}(t) + A_n^{(\alpha,\beta,v,k)} P_{n-1}^{(\alpha,\beta,v,k)}(t)$$

with

$$B_n^{(\alpha,\beta,v,k)} = B_{n-k}^{(\alpha+k,\beta+k,v)}, \quad A_n^{(\alpha,\beta,v,k)} = A_{n-k}^{(\alpha+k,\beta+k,v)}, \quad n \geq k.$$

The explicit expressions of $B_n^{(\alpha,\beta,v)}$ and $A_n^{(\alpha,\beta,v)}$ are given in (2.14)-(2.16).

Considering that $W^{(k)}(t) = W^{(\alpha+k,\beta+k,v)}(t)$ (see Proposition 5.3), the previous recurrence follows directly from (2.3). Notwithstanding, we include the following proof for completeness.

Proof. If we write $P_n^{(\alpha,\beta,v,k)}(t) = \sum_{s=0}^{n-k} \mathcal{P}_{n-k}^s t^s$, from (5.4) we have the following explicit expressions,

$$(5.15) \quad \mathcal{P}_{n-k}^{n-k-1} = \frac{(n-k)}{v} \begin{pmatrix} -\frac{(\alpha+n)v - \kappa_{-v,-\beta}}{(\alpha+\beta+2n+2)} & \frac{\kappa_{-v,-\beta}}{(\kappa_{v,\beta}+2n+2)} \\ -\frac{\kappa_{v,-\beta}}{(\kappa_{-v,\beta}+2n+2)} & -\frac{(\alpha+n)v + \kappa_{-v,-\beta}}{(\alpha+\beta+2n+2)} \end{pmatrix},$$

(5.16)

$$\mathcal{P}_{n-k}^{n-k-2} = \frac{(n-k)(n-k-1)(\alpha+n+1)}{(\alpha+\beta+2n+2)} \left[\frac{\alpha+n}{2(\alpha+\beta+2n+1)} \begin{pmatrix} \frac{\kappa_{v,\beta}+2n}{\kappa_{v,\beta}+2n+2} & 0 \\ 0 & \frac{\kappa_{-v,\beta}+2n}{\kappa_{-v,\beta}+2n+2} \end{pmatrix} \right. \\ \left. + \frac{n+\beta+1}{\alpha+\beta+2n+1} \begin{pmatrix} 1 & 0 \\ \frac{1}{\kappa_{v,\beta}+2n+2} & 1 \end{pmatrix} \right. \\ \left. + \frac{1}{v} \begin{pmatrix} -\frac{\alpha-\beta}{(\kappa_{v,\beta}+2n+2)} & -\frac{\kappa_{-v,-\beta}}{\kappa_{v,\beta}+2n+2} \\ \frac{\kappa_{v,-\beta}}{\kappa_{-v,\beta}+2n+2} & \frac{\alpha-\beta}{(\kappa_{-v,\beta}+2n+2)} \end{pmatrix} \right]. \quad (5.17)$$

If we consider the coefficient of order $n - k$ and $n - k - 1$ in the three-term recurrence relation we have,

$$\begin{aligned} B_n^{(\alpha, \beta, v, k)} &= \mathcal{P}_{n-k}^{n-k-1} - \mathcal{P}_{n+1-k}^{n-k-1}, \\ A_n^{(\alpha, \beta, v, k)} &= \mathcal{P}_{n-k}^{n-k-2} - \mathcal{P}_{n+1-k}^{n-k-1} - B_n^{(\alpha, \beta, v, k)} \mathcal{P}_{n-k}^{n-k-1}, \quad n \in \mathbb{N}, \end{aligned}$$

respectively. Comparing with the expressions of $B_{n-k}^{(\alpha+k, \beta+k, v)}$ and $A_{n-k}^{(\alpha+k, \beta+k, v)}$ given by substituting properly in (2.14)-(2.16), the proposition follows. \square

6. SHIFT OPERATORS

In this section we use Pearson equation (5.6) to give explicit lowering and rising operators for the monic n -degree polynomials $P_{n+k}^{(\alpha, \beta, v, k)}(t)$, $n \geq 0$, defined in (5.1). Moreover, from the existence of the shift operators we deduce a Rodrigues formula for the sequence of derivatives $\left(P_{n+k}^{(\alpha, \beta, v, k)}\right)_{n \geq 0}$, and we find a matrix-valued differential operator for which these matrix-valued polynomials are eigenfunctions. In what follows, we will consider the matrix-valued functions $W^{(k)}(t)$, $\Phi^{(k)}(t)$ and $\Psi^{(k)}(t)$, $k \in \mathbb{N}$ as defined above in Theorem 5.4.

For any pair of matrix-valued functions P and Q , we denote

$$\langle P, Q \rangle_k = \int_0^1 P(t) W^{(k)}(t) Q^*(t) dt.$$

Proposition 6.1. *Let $\eta^{(k)}$ be the first order matrix-valued right differential operator*

$$(6.1) \quad \eta^{(k)} = \frac{d}{dt}(\Phi^{(k)}(t))^* + (\Psi^{(k)}(t))^*.$$

Then $\frac{d}{dt} : L^2(W^{(k)}) \rightarrow L^2(W^{(k+1)})$ and $\eta^k : L^2(W^{(k+1)}) \rightarrow L^2(W^{(k)})$ satisfy

$$\left\langle \frac{dP}{dt}, Q \right\rangle_{k+1} = - \left\langle P, Q \eta^{(k)} \right\rangle_k.$$

Proof. From $\left\langle \frac{dP}{dt}, Q \right\rangle_{k+1} = \int_0^1 \frac{dP(t)}{dt} W^{(k+1)}(t) Q^*(t) dt$, integrating by parts and taking into account equalities (5.13) and (5.14) in Remark 5.5 we get,

$$\begin{aligned} \left\langle \frac{dP}{dt}, Q \right\rangle_{k+1} &= - \int_0^1 P(t) \frac{d}{dt} \left(W(t)^{(k+1)} \right) Q^*(t) dt - \int_0^1 P(t) W^{(k+1)}(t) \left(\frac{dQ(t)}{dt} \right)^* dt \\ &= - \int_0^1 P(t) W^{(k)}(t) \Psi^{(k)}(t) Q^*(t) dt - \int_0^1 P(t) W^{(k)}(t) \Phi^{(k)}(t) \left(\frac{dQ(t)}{dt} \right)^* dt \\ &= - \int_0^1 P(t) W^{(k)}(t) \left(\Psi^{(k)}(t) Q^*(t) + \Phi^{(k)}(t) \left(\frac{dQ(t)}{dt} \right)^* \right) Q^*(t) dt \\ &= - \left\langle P, Q \eta^{(k)} \right\rangle_k. \end{aligned}$$

\square

Lemma 6.2. *The following identity holds true*

$$I\eta^{(k+n-1)} \dots \eta^{(k+1)} \eta^{(k)} = \mathcal{C}_n^k P_{n+k}^{(\alpha, \beta, v, k)}, \quad n \geq 1,$$

for a given $k \geq 0$, where

$$(6.2) \quad \mathcal{C}_n^k = (-1)^n (\alpha + \beta + 3 + 2k + n)_n \begin{pmatrix} \frac{(\kappa_{v, \beta} + 2(k+1+n))}{(\kappa_{v, \beta} + 2(k+1))} & 0 \\ 0 & \frac{(\kappa_{-v, \beta} + 2(k+1+n))}{(\kappa_{-v, \beta} + 2(k+1))} \end{pmatrix}, \quad n \geq 1.$$

Proof. It holds that $I\eta^{(k+n-1)} \dots \eta^{(k+1)} \eta^{(k)}$ is a polynomial of degree n . From the definition of the monic sequence of derivatives in (5.1) one has

$$\frac{d}{dt} P_{n+k}^{(\alpha, \beta, v, k)}(t) = n P_{n+k}^{(\alpha, \beta, v, k+1)}(t).$$

Thus, Proposition 6.1 implies that $P_{n+k}^{(\alpha, \beta, v, k+1)} \eta^{(k)}$ is a multiple of $P_{n+k}^{(\alpha, \beta, v, k)}$.

Therefore, applying the raising operators $\eta^{(k+n-1)} \dots \eta^{(k+1)} \eta^{(k)}$ to $P_{n+k}^{(\alpha, \beta, v, k+n)} = I$ we get a multiple of $P_{n+k}^{(\alpha, \beta, v, k)}$. For the leading coefficient \mathcal{C}_n^k of the polynomial $I\eta^{(k+n-1)} \dots \eta^{(k+1)} \eta^{(k)}$ one obtains the expression

$$\mathcal{C}_n^k = \prod_{i=1}^n \left((i-1) \mathcal{A}_2^{k+n-i} + \mathcal{B}_1^{k+n-i} \right).$$

The diagonal matrix entries \mathcal{A}_2^k and \mathcal{B}_1^k are defined in (5.7) and (5.10). Then, by replacing $\mathcal{B}_1^k = (\alpha + \beta + 4 + 2k) \mathcal{A}_2^k$ in the identity above we have

$$\begin{aligned} \mathcal{C}_n^k &= \prod_{i=1}^n \left((2n + \alpha + \beta + 3 + 2k - i) \mathcal{A}_2^{k+n-i} \right) \\ &= (-1)^n \prod_{i=1}^n (2n + \alpha + \beta + 3 + 2k - i) \begin{pmatrix} \prod_{i=1}^n \frac{(\kappa_{v, \beta} + 2(k+n-i+2))}{\kappa_{v, \beta} + 2(k+n-i+1)} & 0 \\ 0 & \prod_{i=1}^n \frac{(\kappa_{-v, \beta} + 2(k+n-i+2))}{\kappa_{-v, \beta} + 2(k+n-i+1)} \end{pmatrix}. \end{aligned}$$

Hence, the proof follows.

Note that \mathcal{C}_n^k is non-singular since $|\alpha - \beta| < |v| < \alpha + \beta + 2(k+1)$. \square

From the proposition and the lemma above we obtain another expression for the Rodrigues formula.

Proposition 6.3. *The polynomials $\left(P_{n+k}^{(\alpha, \beta, v, k)} \right)_{n \geq 0}$ satisfy the following Rodrigues formula,*

$$P_{n+k}^{(\alpha, \beta, v, k)}(t) = \left(\mathcal{C}_n^k \right)^{-1} \left(\frac{d^n}{dt^n} W^{(k+n)}(t) \right) \left(W^{(k)}(t) \right)^{-1}, \quad n \geq 1,$$

where the matrices \mathcal{C}_n^k are given by the expression in (6.2).

Proof. Let Q be a matrix-valued function and $\eta^{(k)}$ the raising operator in (6.1), then

$$Q\eta^{(k)} = \frac{dQ}{dt}(\Phi^{(k)})^* + Q(\Psi^{(k)})^*.$$

Using the identities (5.13) and (5.14) we obtain

$$Q\eta^{(k)} = \frac{d}{dt} \left(QW^{(k+1)} \right) \left(W^{(k)} \right)^{-1}.$$

Iterating, it gives

$$Q\eta^{(k+n-1)} \dots \eta^{(k+1)} \eta^{(k)} = \frac{d^n}{dt^n} \left(QW^{(k+n)} \right) \left(W^{(k)} \right)^{-1}.$$

Now, taking $Q(t) = I$ and using Lemma 6.2 we have

$$P_{n+k}^{(\alpha, \beta, v, k)}(t) = \left(C_n^k \right)^{-1} \frac{d^n}{dt^n} \left(W^{(k+n)}(t) \right) \left(W^{(k)}(t) \right)^{-1}.$$

□

Corollary 6.4. *Let $W^{(k)}(t)$ be the weight matrix (5.5). Then, the differential operator*

$$(6.3) \quad E^{(k)} = \frac{d}{dt} \circ \eta^{(k)} = \frac{d^2}{dt^2} (\Phi^{(k)}(t))^* + \frac{d}{dt} (\Psi^{(k)}(t))^*$$

is symmetric with respect to $W^{(k)}(t)$ for all $k \in \mathbb{N}_0$. Moreover, the polynomials $\left(P_{n+k}^{(\alpha, \beta, v, k)} \right)_{n \geq 0}$ are eigenfunctions of the operator $E^{(k)}$ with eigenvalue

$$\Lambda_n \left(E^{(k)} \right) = n(n + \alpha + \beta + 3 + 2k) \mathcal{A}_2^k,$$

where \mathcal{A}_2^k is given by (5.7).

Proof. From Proposition 6.1 and the factorization $E^{(k)} = \frac{d}{dt} \circ \eta^{(k)}$ follows directly that $E^{(k)}$ is symmetric with respect to $W^{(k)}$.

The eigenvalue is obtained by looking at the leading coefficients of $\Phi^{(k)}(t)$ and $\Psi^{(k)}(t)$ in (5.6). Thus, we obtain $\Lambda_n \left(E^{(k)} \right) = n(n-1) \mathcal{A}_2^k + n \mathcal{B}_1^k = n(n + \alpha + \beta + 3 + 2k) \mathcal{A}_2^k$. □

Remark 6.5. *The operators $E^{(k)}$ and $D^{(k)}$ in (5.2) commute. This result follows from the fact that the corresponding eigenvalues $\Lambda_n \left(E^{(k)} \right)$ and $\Lambda_{n+k}^{(k)}$ in (5.3) commute, and the linear map that assigns to each differential operator in the algebra of differential operators $D(W^{(k)})$ its corresponding sequence of eigenvalues, is an isomorphism (see [33, Propositions 2.6 and 2.8]).*

Remark 6.6. *The Darboux transform $\tilde{E}^{(k)} = \eta^{(k)} \circ \frac{d}{dt}$ of the operator $E^{(k)}$ is not symmetric with respect to $W^{(k)}$. Moreover, it is symmetric with respect to $W^{(k+1)}$. Indeed,*

$$\begin{aligned} \eta^{(k)} \circ \frac{d}{dt} &= \frac{d^2}{dt^2} \left(\Phi^{(k)}(t) \right)^* + \frac{d}{dt} \left(\frac{d}{dt} \left(\Phi^{(k)}(t) \right)^* + \frac{d}{dt} \left(\Psi^{(k)}(t) \right)^* \right) + \frac{d}{dt} \left(\Psi^{(k)}(t) \right)^* \\ &= \frac{d^2}{dt^2} \left(\mathcal{A}_2^k t^2 + \mathcal{A}_1^k t + \mathcal{A}_0^k \right)^* + \frac{d}{dt} \left(\left(2\mathcal{A}_2^k + \mathcal{B}_1^k \right) t + \mathcal{A}_1^k + \mathcal{B}_0^k \right)^* + (\mathcal{B}_1^k)^*. \end{aligned}$$

In fact, if we substitute the coefficient of the second derivative in the first symmetry condition in (2.2) we obtain

$$W^{(k)}(t) \left(\mathcal{A}_2^k t^2 + \mathcal{A}_1^k t + \mathcal{A}_0^k \right) = \left(\mathcal{A}_2^k t^2 + \mathcal{A}_1^k t + \mathcal{A}_0^k \right)^* W^{(k)}(t),$$

which does not hold. Taking the main coefficient $W_2^{(k)}$ of $\widetilde{W}^{(\alpha, \beta, v, k)}$ in (5.5), one has in particular

$$W_2^k \mathcal{A}_1^k - \left(\mathcal{A}_1^k \right)^* W_2^k = \frac{4v(\alpha + \beta + 2(k+1))}{(\kappa_{-v, \beta} + 2(k+1))(\kappa_{v, \beta} + 2(k+1))} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \neq \mathbf{0}.$$

The second statement follows from Proposition 6.1.

7. THE ALGEBRA $D(W)$

In this section we will discuss some properties of the structure of the algebra of matrix differential operators having as eigenfunctions a sequence of polynomials $(P_n)_{n \geq 0}$, orthogonal with respect to the weight matrix $W = W^{(\alpha, \beta, v)}$, i.e.,

$$D(W) = \{D : P_n D = \Lambda_n(D) P_n, \Lambda_n(D) \in \mathbb{C}^{N \times N} \text{ for all } n \geq 0\}.$$

The definition of $D(W)$ does not depend on the particular sequence of orthogonal polynomials (see [33, Corollary 2.5]).

Theorem 7.1. *Consider the weight matrix function $W = W^{(\alpha, \beta, v)}(t)$. Then, the differential operators of order at most two in $D(W)$ are of the form*

$$(7.1) \quad D = \frac{d^2}{dt^2} (\mathcal{A}_2 t^2 + \mathcal{A}_1 t + \mathcal{A}_0) + \frac{d}{dt} (\mathcal{B}_1 t + \mathcal{B}_0) + \mathcal{C}_0,$$

where

$$\mathcal{A}_2 = \begin{pmatrix} a & c \\ b & d \end{pmatrix}, \quad a, b, c, d \in \mathbb{C},$$

$$\mathcal{A}_1 = \frac{1}{2v} \left[\begin{pmatrix} -2va & (a-d)\kappa_{-v, -\beta} \\ (a-d)\kappa_{v, -\beta} & -2vd \end{pmatrix} + b\kappa_{-v, -\beta} \begin{pmatrix} -1 & 0 \\ 2 & 1 \end{pmatrix} + c\kappa_{v, -\beta} \begin{pmatrix} -1 & -2 \\ 0 & 1 \end{pmatrix} \right],$$

$$\mathcal{A}_0 = \frac{(a-d)\kappa_{v, -\beta}\kappa_{-v, -\beta} + b\kappa_{-v, -\beta}^2 - c\kappa_{v, -\beta}^2}{4v^2} \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix},$$

$$\mathcal{B}_1 = \begin{pmatrix} a(\alpha + \beta + 4) & (\kappa_{-v, \beta} + 4)c \\ (\kappa_{v, \beta} + 4)b & (\alpha + \beta + 4)d \end{pmatrix},$$

$$\mathcal{B}_0 = \frac{1}{4v} \left[a \begin{pmatrix} -4((\alpha + 1)v - \kappa_{-v, -\beta}) & \kappa_{-v, -\beta}(\kappa_{-v, \beta} + 6) \\ \kappa_{v, -\beta}(\kappa_{v, \beta} + 2) & 0 \end{pmatrix} + b\kappa_{-v, -\beta} \begin{pmatrix} -(\kappa_{v, \beta} + 2) & 0 \\ 2(\kappa_{v, \beta} + 4) & \kappa_{v, \beta} + 6 \end{pmatrix} \right. \\ \left. + c\kappa_{v, -\beta} \begin{pmatrix} -(\kappa_{-v, \beta} + 6) & -2(\kappa_{-v, \beta} + 4) \\ 0 & \kappa_{-v, \beta} + 2 \end{pmatrix} + d \begin{pmatrix} 0 & -\kappa_{-v, -\beta}(\kappa_{-v, \beta} + 2) \\ -\kappa_{v, -\beta}(\kappa_{v, \beta} + 6) & -4((\alpha + 1)v + \kappa_{v, -\beta}) \end{pmatrix} \right],$$

$$\mathcal{C}_0 = \frac{1}{4}(\kappa_{v, \beta} + 4)(\kappa_{v, \beta} + 2) \begin{pmatrix} a \frac{(\kappa_{-v, \beta} + 4)}{\kappa_{v, \beta} + 4} - d \frac{\kappa_{-v, \beta} + 2}{\kappa_{v, \beta} + 2} & c \frac{(\kappa_{-v, \beta} + 4)(\kappa_{-v, \beta} + 2)}{(\kappa_{v, \beta} + 4)(\kappa_{v, \beta} + 2)} \\ b & 0 \end{pmatrix} + eI, \quad e \in \mathbb{C}.$$

Proof. Let $(P_n^{(\alpha, \beta, v)})_{n \geq 0}$ be the monic sequence of orthogonal polynomials with respect to $W^{(\alpha, \beta, v)}$. The polynomial $P_n^{(\alpha, \beta, v)}$ is an eigenfunction of the operator D (7.1) if

$$P_n^{(\alpha, \beta, v)} D = \Lambda_n P_n^{(\alpha, \beta, v)},$$

with $\Lambda_n = n(n-1)\mathcal{A}_2 + n\mathcal{B}_1 + \mathcal{C}_0$. This equation holds if and only if

$$(7.2) \quad \begin{aligned} & k(k-1)\mathcal{P}_n^k \mathcal{A}_2 + (k+1)k\mathcal{P}_n^{k+1} \mathcal{A}_1 + (k+2)(k+1)\mathcal{P}_n^{k+2} \mathcal{A}_0 + k\mathcal{P}_n^k \mathcal{B}_1 \\ & + (k+1)\mathcal{P}_n^{k+1} \mathcal{B}_0 + \mathcal{P}_n^k \mathcal{C}_0 - (n(n-1)\mathcal{A}_2 + n\mathcal{B}_1 + \mathcal{C}_0)\mathcal{P}_n^k = 0, \end{aligned}$$

where \mathcal{P}_n^k denotes the k -th coefficient of P_n , $k = 0, 1, 2, \dots, n$.

To prove the theorem we need to solve equation (7.2) for $k = n-1$ and $k = n-2$ to find relations between the parameters of the matrix-valued coefficients $\mathcal{A}_2, \mathcal{A}_1, \mathcal{A}_0, \mathcal{B}_1, \mathcal{B}_0$ and \mathcal{C}_0 .

We obtain the explicit expressions of \mathcal{P}_n^{n-1} and \mathcal{P}_n^{n-2} by substituting $k = 0$ in the equalities (5.15) and (5.16) respectively.

From equation (7.2) for $k = n-1$ we get

$$(7.3) \quad (P_n^{n-1} \Lambda_n - \Lambda_n P_n^{n-1}) - P_n^{n-1} (2(n-1)\mathcal{A}_2 + \mathcal{B}_1) + [n(n-1)\mathcal{A}_1 + n\mathcal{B}_0] = 0.$$

Multiplying equation (7.3) by

$$\frac{v(\alpha + \beta + 2(n+1))(\kappa_{v, \beta} + 2(n+1))(\kappa_{-v, \beta} + 2(n+1))}{n}$$

one obtains a matrix polynomial on n of degree four, where each coefficient must be equal to zero. From the expression of the coefficient of n^4 we get the expression for \mathcal{A}_1 given above and from the expression of the coefficient of n^3 we get \mathcal{B}_0 in terms of \mathcal{A}_2 and \mathcal{B}_1 . Looking at the entries $(1, 1)$, $(1, 2)$ and $(2, 2)$ of the coefficient of n^2 and the fact that $\kappa_{v, -\beta}$ and $\kappa_{-v, -\beta}$ are not zero we get $(\mathcal{C}_0)_{12}$, $(\mathcal{C}_0)_{11}$ and $(\mathcal{B}_1)_{12}$ respectively in terms of \mathcal{A}_2 and the other entries of \mathcal{C}_0 and \mathcal{B}_1 . Finally, looking at the coefficient of n we get the values of $(\mathcal{B}_1)_{11}$, $(\mathcal{B}_1)_{21}$, $(\mathcal{B}_1)_{22}$ and $(\mathcal{C}_0)_{21}$, consequently, we obtain the values of \mathcal{B}_1 , \mathcal{B}_0 and \mathcal{C}_0 written above.

Analogously, from equation (7.2) for $k = n-2$ we obtain

$$(7.4) \quad (P_n^{n-2} \Lambda_{n-2} - \Lambda_{n-2} P_n^{n-2}) + (n-1)P_n^{n-1}((n-2)\mathcal{A}_1 + \mathcal{B}_0) + n(n-1)\mathcal{A}_0 = 0.$$

Multiplying equation (7.4) by

$$\begin{aligned} & v^2(\alpha + \beta + 2n + 1)(\alpha + \beta + 2(n+1))(\kappa_{v, \beta} + 2(n+1)) \\ & (\kappa_{v, \beta} + \alpha + \beta + 2(2n+1))(\kappa_{-v, \beta} + \alpha + \beta + 2(2n+1))(\kappa_{-v, \beta} + 2(n+1)) \end{aligned}$$

one obtains a matrix polynomial on n of degree eight, where each coefficient must be equal to zero. We get the expression of \mathcal{A}_0 from the coefficient of n^8 .

Thus, if we replace the expressions of \mathcal{A}_0 , \mathcal{A}_1 , \mathcal{B}_1 , \mathcal{B}_0 and the entries $(1, 1)$, $(1, 2)$ and $(2, 1)$ of \mathcal{C}_0 in (7.3) and (7.4) both equations hold true.

Let \mathcal{D}_2 be the complex vector space of differential operators in $D(W)$ of order at most two. We have already proved that $\dim \mathcal{D}_2 \leq 5$.

If D is symmetric then $D \in D(W)$. Using symmetry equations in (2.2) one verifies that the operator D in (7.1) is symmetric with respect to W if and only if $a, d, e \in \mathbb{R}$ and condition

$$(7.5) \quad b \frac{\kappa_{v,\beta} + 2}{\kappa_{v,-\beta}} = -\bar{c} \frac{\kappa_{-v,\beta} + 2}{\kappa_{-v,-\beta}}$$

holds true. Indeed, writing $W(t) = W^{(\alpha,\beta,v)} = W_2 t^2 + W_1 t + W_0$, from the first equation of symmetry in (2.2), we have that $W_2 \mathcal{A}_2^* - \mathcal{A}_2 W_2 = 0$, i.e.,

$$(7.6) \quad \begin{pmatrix} 2\operatorname{Im}(a) \frac{(\kappa_{v,\beta} + 2)}{\kappa_{v,-\beta}} & -\bar{b} \frac{(\kappa_{v,\beta} + 2)}{\kappa_{v,-\beta}} - c \frac{(\kappa_{-v,\beta} + 2)}{\kappa_{-v,-\beta}} \\ b \frac{(\kappa_{v,\beta} + 2)}{\kappa_{v,-\beta}} + \bar{c} \frac{(\kappa_{-v,\beta} + 2)}{\kappa_{-v,-\beta}} & -2\operatorname{Im}(d) \frac{(\kappa_{v,\beta} + 2)}{\kappa_{-v,-\beta}} \end{pmatrix} = 0,$$

where $\operatorname{Im}(z)$ denotes the imaginary part of a complex number z . Then, since $\kappa_{v,\beta} + 2 > 0$ because of the restrictions of the parameters α, β and v in the definition of $W^{(\alpha,\beta,v)}$ in (2.4), to verify (7.6) one needs to have $a, d \in \mathbb{R}$ and condition (7.5).

In addition, from the third symmetry equation (2.2) we have that $e \in \mathbb{R}$.

Thus, there exists at least five linearly independent symmetric operators of order at most two in $D(W)$. Therefore $\dim D_2 = 5$. \square

By taking as the only non zero parameters $a = 1$ and $d = 1$ respectively in the expression of the operator in (7.1), we write the operators:

$$\begin{aligned} D_1 &= \frac{d^2}{dt^2} \left[\begin{pmatrix} t^2 - t & \frac{\kappa_{-v,-\beta}}{2v} t \\ \frac{\kappa_{v,-\beta}}{2v} t & 0 \end{pmatrix} + \frac{\kappa_{v,-\beta} \kappa_{-v,-\beta}}{4v^2} \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix} \right] \\ &+ \frac{d}{dt} \left(\begin{pmatrix} (\alpha + \beta + 4)t + \frac{\kappa_{-v,-\beta}}{v} & -(\alpha + 1) \frac{\kappa_{-v,-\beta}(\kappa_{-v,\beta} + 6)}{4v} \\ \frac{\kappa_{v,-\beta}(\kappa_{v,\beta} + 2)}{4v} & 0 \end{pmatrix} + \begin{pmatrix} \frac{(\kappa_{-v,\beta} + 4)(\kappa_{v,\beta} + 2)}{4} & 0 \\ 0 & 0 \end{pmatrix} \right), \\ D_2 &= \frac{d^2}{dt^2} \left[\begin{pmatrix} 0 & -\frac{\kappa_{-v,-\beta}}{2v} t \\ -\frac{\kappa_{v,-\beta}}{2v} t & t^2 - t \end{pmatrix} - \frac{\kappa_{-v,-\beta} \kappa_{v,-\beta}}{4v^2} \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix} \right] \\ &+ \frac{d}{dt} \left(\begin{pmatrix} 0 & -\frac{\kappa_{-v,-\beta}(\kappa_{-v,\beta} + 2)}{4v} \\ -\frac{\kappa_{v,-\beta}(\kappa_{v,\beta} + 6)}{4v} & (\alpha + \beta + 4)t - \frac{\kappa_{v,-\beta}}{v} - (\alpha + 1) \end{pmatrix} + \begin{pmatrix} -\frac{1}{4}(\kappa_{v,\beta} + 4)(\kappa_{-v,\beta} + 2) & 0 \\ 0 & 0 \end{pmatrix} \right). \end{aligned}$$

Analogously, by choosing as nonzero parameters $c = 1$, $b = -\frac{\kappa_{v,-\beta}(\kappa_{-v,\beta} + 2)}{\kappa_{-v,-\beta}(\kappa_{v,\beta} + 2)}$ and $c = i$,

$b = i \frac{\kappa_{v,-\beta}(\kappa_{-v,\beta} + 2)}{\kappa_{-v,-\beta}(\kappa_{v,\beta} + 2)}$ respectively, we define the operators:

$$\begin{aligned}
D_3 = & \frac{d^2}{dt^2} \left\{ \begin{pmatrix} 0 & 1 \\ -\frac{\kappa_{v,-\beta}(\kappa_{-v,\beta}+2)}{\kappa_{-v,-\beta}(\kappa_{v,\beta}+2)} & 0 \end{pmatrix} t^2 + \frac{\kappa_{v,-\beta}}{\kappa_{v,\beta}+2} \left[\begin{pmatrix} -1 & -\frac{\kappa_{v,\beta}+2}{v} \\ -\frac{\kappa_{-v,\beta}+2}{v} & 1 \end{pmatrix} t \right. \right. \\
& + \frac{1}{2} \left(\frac{(\alpha+\beta+2)(\alpha-\beta)}{v^2} + 1 \right) \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix} \left. \right] \Big\} + \frac{d}{dt} \left[\begin{pmatrix} 0 & \kappa_{-v,\beta}+4 \\ -\frac{\kappa_{v,-\beta}(\kappa_{-v,\beta}+2)(\kappa_{v,\beta}+4)}{\kappa_{-v,-\beta}(\kappa_{v,\beta}+2)} & 0 \end{pmatrix} t \right. \\
& + \frac{\kappa_{v,-\beta}}{v} \begin{pmatrix} -1 & -\frac{(\kappa_{-v,\beta}+4)}{2} \\ -\frac{(\kappa_{v,\beta}+4)(\kappa_{-v,\beta}+2)}{2(\kappa_{v,\beta}+2)} & -\frac{\kappa_{-v,\beta}+2}{\kappa_{v,\beta}+2} \end{pmatrix} \left. \right] + \frac{1}{4}(\kappa_{-v,\beta}+2) \begin{pmatrix} 0 & \kappa_{-v,\beta}+4 \\ -\frac{(\kappa_{v,\beta}+4)\kappa_{v,-\beta}}{\kappa_{-v,-\beta}} & 0 \end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
iD_4 = & \frac{d^2}{dt^2} \left\{ \begin{pmatrix} 0 & -1 \\ -\frac{\kappa_{v,-\beta}(\kappa_{-v,\beta}+2)}{\kappa_{-v,-\beta}(\kappa_{v,\beta}+2)} & 0 \end{pmatrix} t^2 + \frac{\kappa_{v,-\beta}}{v(\kappa_{v,\beta}+2)} \left[\begin{pmatrix} \alpha+\beta+2 & \kappa_{v,\beta}+2 \\ -(\kappa_{-v,\beta}+2) & -(\alpha+\beta+2) \end{pmatrix} t \right. \right. \\
& - (\alpha+1) \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix} \left. \right] \Big\} + \frac{d}{dt} \left[\begin{pmatrix} 0 & -(\kappa_{-v,\beta}+4) \\ -\frac{\kappa_{v,-\beta}(\kappa_{-v,\beta}+2)(\kappa_{v,\beta}+4)}{\kappa_{-v,-\beta}(\kappa_{v,\beta}+2)} & 0 \end{pmatrix} t \right. \\
& + \frac{\kappa_{v,-\beta}}{2v} \begin{pmatrix} \kappa_{-v,\beta}+4 & \kappa_{-v,\beta}+4 \\ -\frac{(\kappa_{v,\beta}+4)(\kappa_{-v,\beta}+2)}{(\kappa_{v,\beta}+2)} & -\frac{(\kappa_{v,\beta}+4)(\kappa_{-v,\beta}+2)}{(\kappa_{v,\beta}+2)} \end{pmatrix} \left. \right] \\
& - \frac{\kappa_{-v,\beta}+2}{4} \begin{pmatrix} 0 & \kappa_{-v,\beta}+4 \\ \frac{(\kappa_{v,\beta}+4)\kappa_{v,-\beta}}{\kappa_{-v,-\beta}} & 0 \end{pmatrix}.
\end{aligned}$$

One has the following

Corollary 7.2. *The set of symmetric operators $\{D_1, D_2, D_3, D_4, I\}$ is a basis of the space of differential operators of order at most two in $D(W)$. Moreover, the corresponding eigenvalues for the differential operators D_1, D_2, D_3 and D_4 are*

$$\begin{aligned}
\Lambda_n(D_1) &= \frac{1}{4} \begin{pmatrix} (\kappa_{v,\beta} + 2(n+1))(\kappa_{-v,\beta} + 2(n+2)) & 0 \\ 0 & 0 \end{pmatrix}, \\
\Lambda_n(D_2) &= \begin{pmatrix} -\frac{1}{4}(\kappa_{-v,\beta} + 2)(\kappa_{v,\beta} + 4) & 0 \\ 0 & (n + \alpha + \beta + 3)n \end{pmatrix}, \\
\Lambda_n(D_3) &= \frac{1}{4}(\kappa_{-v,\beta} + 2(1+n))(\kappa_{-v,\beta} + 2(2+n)) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\
&\quad - \frac{(\kappa_{v,\beta} + 2(1+n))(\kappa_{v,\beta} + 2(2+n))(\kappa_{-v,\beta} + 2)\kappa_{v,-\beta}}{4\kappa_{-v,-\beta}(\kappa_{v,\beta} + 2)} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \\
\Lambda_n(iD_4) &= -\frac{1}{4}(\kappa_{-v,\beta} + 2(1+n))(\kappa_{-v,\beta} + 2(2+n)) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\
&\quad - \frac{(\kappa_{v,\beta} + 2(1+n))(\kappa_{v,\beta} + 2(2+n))(\kappa_{-v,\beta} + 2)\kappa_{v,-\beta}}{4\kappa_{-v,-\beta}(\kappa_{v,\beta} + 2)} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.
\end{aligned}$$

Corollary 7.3. *The differential operators appearing in (2.7) and (6.3) are $D^{(\alpha,\beta,v)} = -D_1 - D_2$ and $E^{(0)} = -\frac{\kappa_{v,\beta} + 4}{\kappa_{v,\beta} + 2}D_1 - \frac{\kappa_{-v,\beta} + 4}{\kappa_{-v,\beta} + 2}D_2$ respectively.*

Corollary 7.4. *There are no operators of order one in the algebra $D(W)$.*

Proof. Suppose that there exists a right differential operator of order one, such that $D = aD_1 + bD_2 + cD_3 + d(iD_4) + eI$, with $a, b, c, d, e \in \mathbb{R}$. Equating to zero the matrix-valued coefficient of $\frac{d^2}{dt^2}$ one obtains:

$$\begin{pmatrix} a & c - di \\ -\frac{\kappa_{v,-\beta}(\kappa_{-v,\beta} + 2)}{\kappa_{-v,-\beta}(\kappa_{v,\beta} + 2)}(c + di) & b \end{pmatrix} = \mathbf{0}.$$

Therefore $a = b = c = d = 0$. □

Corollary 7.5. *The algebra $D(W)$ is not commutative.*

Proof. Using the isomorphism between the algebra of differential operators and the algebra of matrix-valued functions of n generated by the eigenvalues going with this operators we have that $D_1D_3 \neq D_3D_1$ since $\Lambda_n(D_1)\Lambda_n(D_3) \neq \Lambda_n(D_3)\Lambda_n(D_1)$. □

Remark 7.6. *In [42] the authors study the algebra $D(W^{(p,q)})$, where $W^{(p,q)}$ is, for $p \neq \frac{q}{2}$, the irreducible weight matrix*

$$W^{(p,q)}(t) = (t(1-t)) \frac{q-2}{2} \begin{pmatrix} 2pt^2 - 2pt + \frac{q}{2} & qt - \frac{q}{2} \\ qt - \frac{q}{2} & -2(p-q)t^2 + 2(p-q)t + \frac{q}{2} \end{pmatrix}, \quad t \in [0, 1].$$

Let us denote by $D_1^{(p,q)}, D_2^{(p,q)}, D_3^{(p,q)}$ and $D_4^{(p,q)}$ the differential operators appearing in [42]. Then, taking $\alpha = \beta = \frac{q}{2} - 1$ in (2.4) and writing $v = 2p - q$, we have the following relations with the

operators D_i , $i = 1, 2, 3, 4$, defined above:

$$\begin{aligned} D_1^{(p,q)} &= D_1, & D_2^{(p,q)} &= D_2 + (q-p)(p+1)I, \\ D_3^{(p,q)} &= \frac{p}{2(q-p)}(D_3 + iD_4), & D_4^{(p,q)} &= \frac{1}{2}(D_3 - iD_4). \end{aligned}$$

8. ACKNOWLEDGEMENTS

The authors would like to thank Pablo Román for useful suggestions on earlier versions of the paper. The authors also thank the anonymous referees for their careful reading and remarks.

REFERENCES

- [1] F. V. Atkinson, Discrete and continuous boundary problems, Academic Press, New York, 1964.
- [2] Ju. M. Berezanskii, Expansions in eigenfunctions of selfadjoint operators, *Transl. Math. Monographs*, AMS **17** 1968.
- [3] J. Borrego, M. Castro, and A. J. Durán, Orthogonal matrix polynomials satisfying differential equations with recurrence coefficients having non-scalar limits, *Integral Transforms Spec. Funct.*, **23**, 685–700 (2012).
- [4] C. Calderón, Y. González, I. Pacharoni, S. Simondi, and I. Zurrián, 2×2 hypergeometric operators with diagonal eigenvalues *J. Approx. Theory*, 248:105299, 17pp (2019).
- [5] M. J. Cantero, L. Moral and L. Velázquez, Matrix orthogonal polynomials whose derivatives are also orthogonal, *J. Approx. Theory*, 146(2):174–211 (2007).
- [6] W. R. Casper, Elementary examples of solutions to Bochner’s problem for matrix differential operators, *J. Approx. Theory*, **229**, 36–71 (2018).
- [7] W. R. Casper, The symmetric 2×2 hypergeometric matrix differential operators, *arXiv:1907.12703*, (2019).
- [8] W. R. Casper and M. Yakimov, The matrix Bochner problem, *Amer. J. Math.* (2021), to appear, *arXiv:1803.04405*.
- [9] M.M. Castro and F. A. Grünbaum, The algebra of differential operators associated to a family of matrix-valued orthogonal polynomials: five instructive examples, *Int. Math. Res. Not.*, **7**, 1–33 (2006).
- [10] M. Castro and F. A. Grünbaum, The Darboux process and time-and-band limiting for matrix orthogonal polynomials, *Linear Algebra Appl.*, **487**, 328–341 (2015).
- [11] M. Castro and F. A. Grünbaum, Time-and-band limiting for matrix orthogonal polynomials of Jacobi type, *Random Matrices: Theory and Applications*, 06(04):1740001, 12pp (2017).
- [12] M. Castro, F. A. Grünbaum, I. Pacharoni, and I. Zurrián, A further look at time-and-band limiting for matrix orthogonal polynomials, in M. Nashed and X. Li, editors, *Frontiers in Orthogonal Polynomials and q-Series*. World Scientific, 2018.
- [13] J. J. Duistermaat and F. A. Grünbaum, Differential equations in the spectral parameter, *Comm. Math. Phys.*, **103**(2):177–240 (1986).
- [14] A. J. Durán, Markov’s theorem for orthogonal matrix polynomials, *Canad. J. Math.*, **48**:1180–1195 (1996).
- [15] A. J. Durán, Matrix inner product having a matrix symmetric second-order differential operator, *Rocky Mt. J. Math.*, **27**(2):585–600 (1997).
- [16] A. J. Durán, Generating orthogonal matrix polynomials satisfying second order differential equations from a trio of triangular matrices, *J. Approx. Theory*, **161**, 88–113 (2009).
- [17] A. J. Durán, A method to find weight matrices having symmetric second-order differential operators with matrix leading coefficients, *Const. Approx.*, **29**, 181–205 (2009).
- [18] A. J. Durán, Rodrigues’ formulas for orthogonal matrix polynomials satisfying second-order differential equations, *Int. Math. Res. Not.*, **5**, 824–855 (2010).
- [19] A. J. Durán and F. A. Grünbaum, Orthogonal matrix polynomials satisfying second-order differential equations, *Int. Math. Res. Not.*, **10**:461–484 (2004).
- [20] A. J. Durán and F. A. Grünbaum, Structural formulas for orthogonal matrix polynomials satisfying second order differential equations, I, *Constr. Approx.*, **22**, 255–271 (2005).

- [21] A. J. Durán and F. A. Grünbaum, Matrix orthogonal polynomials satisfying second order differential equations: coping without help from group representation theory, *J. Approx. Theory*, 148, 35–48 (2007).
- [22] A. J. Durán, and M. D. de la Iglesia, Some examples of orthogonal matrix polynomials satisfying odd order differential equations, *J. Approx. Theory*, 150, 153–174 (2008).
- [23] A. J. Durán and P. López, Orthogonal matrix polynomials, *Laredo Lectures on Orthogonal Polynomials and Special Functions*, 13–44, Adv. Theory Spec. Funct. Orthogonal Polynomials, Nova Sci. Publ., Hauppauge, NY, 2004.
- [24] A. J. Durán and P. López, Structural formulas for orthogonal matrix polynomials satisfying second order differential equations, II, *Constr. Approx.*, 26, No. 1, 29–47 (2007).
- [25] F. A. Grünbaum, Matrix-valued Jacobi polynomials, *Bull. Sci. Math.*, 127(3):207–214 (2003).
- [26] F.A. Grünbaum and M. D. de la Iglesia, Matrix-valued orthogonal polynomials arising from group representation theory and a family of quasi-birth-and death processes, *SIAM J. Matrix Anal. Appl.*, 30, 741–761 (2008).
- [27] F. A. Grünbaum, I. Pacharoni, and J. Tirao, A matrix-valued solution to Bochner’s problem, *J. Phys. A*, 34(48):10647–10656 (2001).
- [28] F. A. Grünbaum, I. Pacharoni, and J. Tirao, Matrix-valued spherical functions associated to the complex projective plane, *J. Funct. Anal.*, 188(2):350–441 (2002).
- [29] F. A. Grünbaum, I. Pacharoni, and J. Tirao, Matrix-valued orthogonal polynomials of the Jacobi type, *Indag. Math. (N.S.)*, 14(3-4):353–366 (2003).
- [30] F. A. Grünbaum, I. Pacharoni, and I. Zurrián, Time and band limiting for matrix-valued functions, an example, *SIGMA, Symmetry Integrability Geom. Methods Appl.*, 11(044):1–14 (2015).
- [31] F. A. Grünbaum, I. Pacharoni, and I. Zurrián, Time and band limiting for matrix-valued functions, *Inverse Problems*, 33(2):1–26 (2017).
- [32] F. A. Grünbaum, I. Pacharoni, and I. Zurrián, Bispectrality and time-band-limiting: Matrix-valued polynomials, *International Math. Research Notices* 13, 4016–4036 (2020).
- [33] F. A. Grünbaum and J. Tirao, The algebra of differential operators associated to a weight matrix, *Integral Equations Operator Theory*, 58(4):449–475 (2007).
- [34] E. Koelink, A. de los Ríos, and P. Román, Matrix-valued Gegenbauer-type polynomials, *Constr. Approx.*, 46(3):459–487 (2017).
- [35] E. Koelink, M. van Pruijssen, and P. Román, Matrix-valued orthogonal polynomials related to $(\mathrm{SU}(2) \times \mathrm{SU}(2), \mathrm{diag})$, *Int. Math. Res. Not.*, 2012(24):5673–5730 (2012).
- [36] E. Koelink, M. van Pruijssen, and P. Román, Matrix-valued orthogonal polynomials related to $(\mathrm{SU}(2) \times \mathrm{SU}(2), \mathrm{SU}(2))$, II, *PRIMS*, 49(2):271–312 (2013).
- [37] M. G. Krein, Infinite j-matrices and a matrix moment problem, *Dokl. Akad. Nauk SSSR*, 69(2):125–128 (1949).
- [38] M. G. Krein, Fundamental aspects of the representation theory of hermitian operators with deficiency index (m, m) , *AMS Translations, series 2*, 97:75–143 (1971).
- [39] L. Miranian, On classical orthogonal polynomials and differential operators, *J. Phys. A: Math Gen*, 38, 6379–6383 (2005).
- [40] I. Pacharoni and P. Román, A sequence of matrix-valued orthogonal polynomials associated to spherical functions, *Constr. Approx.*, 28(2):127–147 (2008).
- [41] I. Pacharoni and J. Tirao, Matrix-valued orthogonal polynomials arising from the complex projective space, *Constr. Approx.*, 25(2):177–192 (2006).
- [42] I. Pacharoni and I. Zurrián, Matrix Gegenbauer Polynomials: The 2×2 Fundamental Cases, *Constr. Approx.*, 43(2):253–271 (2016).
- [43] G. Szegő, Orthogonal Polynomials, *Coll. Publ., XXIII, American Mathematical Society*, Providence, RI, 1975.
- [44] J. Tirao, The matrix-valued hypergeometric equation, *Proc. Natl. Acad. Sci. U.S.A.*, 100(14):8138–8141 (2003).
- [45] J. Tirao, The algebra of differential operators associated to a weight matrix: a first example, Polcino Milies, César (ed.), Groups, algebras and applications. XVIII Latin American algebra colloquium, São

Pedro, Brazil, August 3–8, 2009. Proceedings. Providence, RI: American Mathematical Society (AMS). Contemporary Mathematics 537, 291-324 2011.

- [46] J. Tirao and I. Zurrián, Reducibility of Matrix Weights, *Ramanujan J.*, 45, no. 2, 349-374 (2018).
- [47] I. Zurrián, The Algebra of Differential Operators for a Gegenbauer Weight Matrix, *Int. Math. Res. Not.*, 8, 2402-2430 (2016).

Email address, C. Calderón: `celeste.calderon@fce.uncu.edu.ar`

Email address, M.M. Castro: `mirta@us.es`

(C. Calderón) FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD NACIONAL DE CUYO, 5500 MEN-DOZA, ARGENTINA

(M. M. Castro) DEPARTAMENTO DE MATEMÁTICA APLICADA II AND IMUS, UNIVERSIDAD DE SEVILLA, ESCUELA POLITÉCNICA SUPERIOR, CALLE VIRGEN DE ÁFRICA 7, 41011, SEVILLA, SPAIN