

The impact of Friedmann’s work on cosmology

Claus Kiefer¹ and Hermann Nicolai²

Historical introduction. The impact of Friedmann’s work on cosmology can hardly be overestimated. By training, Friedmann was a mathematician, but one of exceptional versatility, who made important contributions also in other fields, such as meteorology. In the summer of 1917 and in the middle of tumultuous events in Russia, he founded and was the first director of the “Aeroprabor” factory in Moscow which produced tools for airplanes, and which still exists to this day. Nevertheless, his greatest contribution to science is undoubtedly contained in the two pioneering paper in 1922 and 1924 which appeared in the German journal *Zeitschrift für Physik* [Friedmann 1922, 1924].³ In these papers, he demonstrated that Einstein’s field equations with a cosmological constant (called by him *Weltgleichungen*, i.e. world equations) do not only allow Einstein’s 1917 static solution with matter and de Sitter’s 1917 apparent static vacuum solution, but also dynamical solutions describing an expanding or collapsing Universe. The corresponding equations, today called Friedmann or Friedmann–Lemaître equations, form the basis of modern cosmology. In 1923, Friedmann published a book on cosmology in which he also presents insights into his general philosophical ideas [Friedmann 2000].⁴

In the 1920s, Friedmann’s work had little impact [Ellis 1989]. The main question in those years was trying to find out whether there is an observational difference between Einstein’s and de Sitter’s solution. Friedmann’s papers were apparently also unknown to Georges Lemaître, who in 1927 wrote another groundbreaking paper that was little appreciated at the time: he related the formal solutions for an expanding or contracting Universe to redshifts and thus to observations. Einstein, after

having read Friedmann’s first paper, first thought that the solutions are wrong. Later he admitted that the solutions are mathematically correct, but (in his opinion) physically irrelevant. This demonstrates how deeply the idea of a static Universe was rooted in people’s imagination at the time.

It is often stated that Friedmann was only interested in the mathematics of the equations, not in their physical content. In our opinion, this is only partially true. He was certainly strongly influenced by the mathematicians Weyl and Hilbert, especially the latter’s idea of axiomatization.⁵ But in his work he strongly emphasized that the geometry of the world should be determined by theoretical physics *and* observational astronomy.⁶ At the end of his 1922 paper, he gives an estimate of 10^{10} years for the duration of a recollapsing Universe, which is close to the current estimate for the age of our Universe.

Friedmann was, in particular, interested in the question whether the world (three-dimensional space) is finite or infinite. This motivated him to study the case of negative curvature in 1924 [Friedmann 1924]. He found that, in contrast to the spatially closed case discussed in 1922, the case of negative curvature leaves this question open. He concludes the 1924 paper with the words: “This is the reason why, according to our opinion, Einstein’s world equations without additional assumptions are not yet sufficient to draw a conclusion about the finiteness of our world.”⁷ The question whether it makes sense to talk about actual infinities in physics (in contrast to mathematics) is an intriguing one and continues to be discussed up to the present day [Ellis 2018], as Friedmann’s insights continue to inspire modern research.

¹University of Cologne, Faculty of Mathematics and Natural Sciences, Institute for Theoretical Physics, Cologne, Germany.

²Max Planck Institute for Gravitational Physics, Albert Einstein Institute, Potsdam, Germany.

³In [Friedmann 1922], the German transcription of the Russian name was chosen “Friedman”, but we stick to the common practice of writing “Friedmann”. For an editorial note to the English translation of these papers, see [Ellis 1999].

⁴The editor of the German translation [Friedmann 2000] speculates that the title *Мир как пространство и время* (Die Welt als Raum und Zeit) alludes to Schopenhauer’s opus magnum *Мир как воля и представление* (Die Welt als Wille und Vorstellung).

⁵Friedmann had paid a visit to Göttingen in 1923.

⁶In 1924, he even gave a thesis topic to his student A.B. Schechter dealing with the question whether trigonometric measurements at astronomical dimensions can lead to a decision between different world geometries. A paper on this was published three years after Friedmann’s death by Frédericksz and Schechter [Ellis 1989].

⁷The German original reads: “Dies ist der Grund dafür, daß, unserer Meinung nach, Einsteins Weltgleichungen ohne ergänzende Annahmen noch nicht hinreichen, um einen Schluß über die Endlichkeit unserer Welt zu ziehen.”

⁸A comprehensive discussion of the material in this and the following section can be found in many reviews and textbooks, see e.g. [Weinberg 1972], [Mukhanov 2005], [Ellis 2012] and [Calcagni 2017].

Friedmann’s equations.⁸ Starting point is Einstein’s field equation

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^2}T_{\mu\nu}. \quad (1)$$

Observations indicate that the Universe is approximately isotropic around our position. These come mainly from the anisotropy spectrum of the Cosmic Microwave Background (CMB). Adopting the Cosmological Principle (“all places in the Universe are alike”), one is led to assume (approximate) isotropy around *every* position. One can then mathematically prove that our Universe must also be (approximately) homogeneous. The geometry of a homogeneous and isotropic spacetime is characterised by the line element

$$ds^2 = -c^2 dt^2 + a^2(t) \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right), \quad (2)$$

where $a(t)$ is the scale factor. For the parameter k , we have the possible choices $k = 0$ (flat spatial geometry), $k = 1$ (positive curvature), $k = -1$ (negative curvature); only the latter two cases were treated by Friedmann. Current observations favour a spatially flat Universe, although there is still a controversy [Di Valentino 2020]. A given value for k does not fix the topology of our (spatial) Universe, and it is a most intriguing question to determine the cosmic topology from observations [Luminet 2015].

Inserting the ansatz (2) into (1), one is led to Friedmann’s equations.⁹ The first equation is the restriction of the general Raychaudhuri equation to a homogeneous and isotropic Universe,

$$\ddot{a} = -\frac{4\pi G}{3}(\rho + 3p)a, \quad (3)$$

where ρ and p denote energy density and pressure of matter, respectively. If matter obeys the strong energy condition $\rho + 3p \geq 0$, (3) leads to concave solutions for $a(t)$, that is, to a world model with a singular origin. The second Friedmann equation reads

$$\dot{a}^2 = \frac{8\pi G}{3}\rho a^2 - k. \quad (4)$$

In contrast to (3), this equation only contains temporal derivatives up to *first* order, so it has the interpretation of a *constraint*. In fact, it is the

Friedmann version of the Hamiltonian constraint in general relativity [Kiefer 2012].

From (3) and (4), one can derive a third equation,

$$\dot{\rho} + 3H(\rho + p) = 0, \quad (5)$$

where $H := \dot{a}/a$ is the Hubble parameter (its evaluation at the present day is called Hubble constant, denoted by H_0). The combination $\rho + p$ occurring in (5) is called inertial mass density. In these Friedmann equations, we have followed the modern practice of including the cosmological constant Λ into the density ρ (although this was already suggested by Schrödinger in 1919), because it contributes an ‘energy density of the vacuum’ $\rho_\Lambda := \Lambda/8\pi G$. Its equation of state reads $p_\Lambda = -\rho_\Lambda$, so from (5) we see that ρ_Λ is constant. For barotropic equations of state $p = w\rho$, $w \neq -1$, we find from (5) the solution

$$\rho a^{3(1+w)} = \text{constant}, \quad (6)$$

which includes as particular cases:

- dust ($p = 0$) $\longrightarrow \rho \propto a^{-3}$,
- radiation ($p = \rho/3$) $\longrightarrow \rho \propto a^{-4}$,
- stiff matter ($p = \rho$) $\longrightarrow \rho \propto a^{-6}$.

By the kinematic relation $a_0/a = 1 + z$, with a_0 as the present scale factor, we can relate ρ to the observable redshift z of objects. The case of radiation is relevant for the early Universe, while stiff matter so far seems unrealistic. Today, the Universe is dominated by dust (about one third) and vacuum energy (about two thirds), leading to the temporal evolution

$$a(t) = a_0 \left(\frac{3\Omega_0 H_0}{\Lambda} \right)^{1/3} \sinh^{2/3} \left(\frac{3}{2} \sqrt{\frac{\Lambda}{3}} t \right), \quad (7)$$

where Ω_0 is today’s matter density in terms of the critical density, observationally determined to be about 1/3. Observations also indicate that the age of our Universe is about 13.8 billion years. For large times, the evolution law (7) asymptotes to de Sitter space.¹⁰ From observations of the CMB, there are strong indications that our Universe underwent a quasi-exponential expansion (with very large Λ) already very early in its history, a phase called inflation. Inflation offers the means to explain the origin of structure in the Universe.

Instead of barotropic equations of state, one often employs dynamical matter models, typically with a scalar field ϕ . In the Friedmann limit, this field

⁸From here on, we set $c = 1$.

¹⁰For late time expansion with constant positive ρ_Λ one speaks of *dark energy*, but there is also the possibility that the effective vacuum energy density varies with time, in which case one speaks of *quintessence*. The latter is thought to originate from matter sources and is often modelled by means of a time-dependent scalar field ϕ .

depends, of course, only on time. In the case of a massless field, it corresponds in (4) to the choice of a density $\rho_\phi = \dot{\phi}^2/2$.

Beyond the Friedmann approximation. Beyond the immediate and obvious utility of the Friedmann equations for cosmological applications there are several important and promising directions for future development that build on Friedmann’s achievements. For lack of space we here mention only two of these, namely (i) their use for taking first steps towards a theory of quantum gravity, and (ii) the generalization of the isotropic ansatz (2) in order to search for a fundamental symmetry of Nature.

When adapting the ansatz (2) to a quantum mechanical context one speaks of the so-called *minisuperspace approximation*, in which the full superspace of geometrodynamics, being the moduli space of all three-metrics modulo spatial diffeomorphisms, is restricted to few homogeneous degrees of freedom such as the scale factor a . This limit was first discussed by DeWitt in his pioneering paper on canonical quantum gravity [DeWitt 1967]. This is a huge simplification because key technical issues such as the non-renormalizability of perturbative quantum gravity can be ignored in this approximation. Furthermore, various conceptual issues of quantum gravity and quantum cosmology can be studied. Namely, the direct canonical quantization of the second Friedmann equation (4) leads to a special case of the Wheeler-DeWitt equation [Kiefer 2012, Calcagni 2017], here given for the case of a massless homogeneous scalar field,

$$\left[\frac{4G\hbar^2}{3\pi a^2} \frac{\partial}{\partial a} \left(a \frac{\partial}{\partial a} \right) - \frac{\hbar^2}{a^3} \frac{\partial^2}{\partial \phi^2} - \frac{3\pi}{4G} ka \right] \Psi(a, \phi) = 0,$$

where a particular factor ordering has been adopted. The wave function $\Psi(a, \phi)$ is a simple example of the ‘wave function of the universe’. It is, in particular, possible to analyse the behaviour of Ψ near the singularity where $a \rightarrow 0$. The Wheeler-DeWitt equation has no external time parameter, but one can employ the scale factor a so as to track the evolution of the matter degrees of freedom with respect to this “intrinsic time”. (Note that the minisuperspace Wheeler-DeWitt equation is hyperbolic with respect to a .) Key open issues concern the physical interpretation of Ψ , the construction of a suitable Hilbert space, and the meaning of observables; for a survey and further discussion, see [Kiefer 2012].

The other extension concerns the inclusion of *non-homogeneous* degrees of freedom. On the phenomenological side, the evolution of our Universe, if

approximated by a homogeneous and isotropic spatial part, is successfully described by Friedmann’s equations, but small inhomogeneities *must* be taken into account in order to understand properties of the CMB in the framework of cosmological perturbation theory. Furthermore, a precise understanding of the formation of galaxies and clusters of galaxies requires the numerical treatment of the Einstein equations (1) and their Newtonian limit. Incorporating inhomogeneities is likewise crucial for a better understanding of the origin of the universe, because inhomogeneities scale like a^{-6} , like stiff matter, and thus dominate very close to the Big Bang singularity. This is a crucial issue for inflationary cosmology, which hinges on the ansatz (2). Finally, there remain difficult issues related to defining a generally covariant averaging procedure in Einstein’s theory that would provide a rigorous basis for the Friedmann approximation [Buchert 2015].

On the more mathematical side, a key insight came from the Belinski-Khalatnikov-Lifshitz (BKL) analysis [Belinski 1970, 1982] of the generic behaviour of solutions of Einstein’s equations near a spacelike singularity. There, one generalizes the ansatz (2) to

$$ds^2 = -dt^2 + a^2(t)dx^2 + b^2(t)dy^2 + c^2(t)dz^2,$$

thus giving up isotropy, but retaining spatial homogeneity. A surprising result to come out of this analysis is the appearance of chaotic oscillations of the metric coefficients a, b, c as one approaches the singularity. This result indicates that the ‘near singularity limit’ of the metric exhibits a far more complicated behaviour than inspection of, say, the Schwarzschild metric would suggest, thus also showing the limitations of the assumption of isotropy.

The BKL analysis has been generalised in many directions, in particular also to accommodate inhomogeneities [Krasinski 1997]. Furthermore, closer study of the BKL limit has revealed evidence for a huge infinite-dimensional symmetry of indefinite Kac-Moody type, vastly generalizing the known duality symmetries of supergravity and string theory. This novel symmetry can possibly serve as a guiding principle towards unifying the fundamental interactions [Damour 2002].

We thank Alexander Kamenshchik for his comments on our manuscript.

References

- [Belinski 1970] V. A. Belinski, I.M. Khalatnikov, I.M. Lifshitz,
Oscillatory approach to a singular point in the relativistic cosmology, *Advances in Physics*. 19 (1970) 525–573.
- [Belinski 1982] V. A. Belinski, I.M. Khalatnikov, I.M. Lifshitz,
A general solution of the Einstein equations with a time singularity, *Advances in Physics* 31 (1982) 639–667.
- [Buchert 2015] T. Buchert, M. Carfora, G.F.R. Ellis, E.W. Kolb, M.A.H. MacCallum, J.J. Ostrowski, S. Räsänen, B.F. Roukema, L. Andersson, A.A. Coley, D.L. Wiltshire, Is there proof that backreaction of inhomogeneities is irrelevant in cosmology?, *Classical and Quantum Gravity*, 32 (2015) 215021 (38pp).
- [Calcagni 2017] G. Calcagni, *Classical and Quantum Cosmology*, Graduate Texts in Physics, Springer International Publishing Switzerland, 2017.
- [Damour 2002] T. Damour, M. Henneaux, H. Nicolai,
 E_{10} and a Small Tension Expansion of M Theory, *Physical Review Letters* 89 (2002) 221601 (4pp).
- [DeWitt 1967] B.S. DeWitt, Quantum theory of gravity. I. The canonical theory, *Physical Review*, 160 (1967) 1113–1148.
- [Di Valentino 2020] E. Di Valentino et al., Cosmology Intertwined IV: The Age of the Universe and its Curvature, arXiv:2008.11286v4 [astro-ph.CO].
- [Ellis 1989] G.F.R. Ellis, The Expanding Universe: A History of Cosmology from 1917 to 1960, in: *Einstein and the History of General Relativity*, ed. by D. Howard and J. Stachel, Birkhäuser, Boston, 1989, pp. 367–431.
- [Ellis 1999] G.F.R. Ellis and A. Krasinski, *General Relativity and Gravitation* 31 (1999) 1985–1990; Addendum: *General Relativity and Gravitation* 32 (2000) 1937–1938.
- [Ellis 2012] G.F.R. Ellis, R. Maartens, M.A.H. MacCallum, *Relativistic Cosmology*, Cambridge University Press, Cambridge, 2012.
- [Ellis 2018] G.F.R. Ellis, K. Meissner, H. Nicolai, The physics of infinity, *Nature Physics*, 14 (2018) 770–772 .
- [Friedmann 1922] A. Friedman, Über die Krümmung des Raumes, *Zeitschrift für Physik*, 10 (1922) 377–386.
- [Friedmann 1924] A. Friedmann, Über die Möglichkeit einer Welt mit konstanter negativer Krümmung des Raumes, *Zeitschrift für Physik*, 21 (1924) 326–332.
- [Friedmann 2000] A. Friedmann, *Die Welt als Raum und Zeit*, translated from the Russian by G. Singer, Verlag Harri Deutsch, Thun und Frankfurt am Main, 2000. The Russian original *Мир как пространство и время* appeared 1923 with Academia, Petrograd. In English, the title is “The world as space and time”, but an English translation of the book is not known to us.
- [Kiefer 2012] C. Kiefer, *Quantum Gravity*, 3rd ed., Oxford University Press, Oxford, 2012.
- [Krasinski 1997] A. Krasinski, *Inhomogeneous cosmological models*, Cambridge University Press, Cambridge, 1997.
- [Luminet 2015] J.-P. Luminet, *Cosmic Topology*, *Scholarpedia*, 10(8):31544.
- [Mukhanov 2005] V. Mukhanov, *Physical Foundations of Cosmology*, Cambridge University Press (2005).
- [Weinberg 1972] S. Weinberg, *Gravitation and Cosmology*, John Wiley and Sons (1972).