ONE WEIGHT INEQUALITY FOR BERGMAN PROJECTION AND CALDERÓN OPERATOR INDUCED BY RADIAL WEIGHT

FRANCISCO J. MARTÍN REYES, PEDRO ORTEGA, JOSÉ ÁNGEL PELÁEZ, AND JOUNI RÄTTYÄ

ABSTRACT. Let ω and ν be radial weights on the unit disc of the complex plane such that ω admits the doubling property $\sup_{0\leqslant r<1} \frac{\int_{1}^{1}\omega(s)\,ds}{\int_{1+r}^{1}\omega(s)\,ds}<\infty$. Consider the one weight inequality

$$||P_{\omega}(f)||_{L^{p}_{\nu}} \le C||f||_{L^{p}_{\nu}}, \quad 1 (†$$

for the Bergman projection P_{ω} induced by ω . It is shown that the Muckenhoupt-type condition

$$A_p(\omega, \nu) = \sup_{0 \leqslant r < 1} \frac{\left(\int_r^1 s\nu(s) \, ds\right)^{\frac{1}{p}} \left(\int_r^1 s \left(\frac{\omega(s)}{\nu(s)^{\frac{1}{p}}}\right)^{p'} \, ds\right)^{\frac{1}{p'}}}{\int_r^1 s\omega(s) \, ds} < \infty,$$

is necessary for (†) to hold, and sufficient if ν is of the form $\nu(s) = \omega(s) \left(\int_r^1 s \omega(s) \, ds \right)^{\alpha}$ for some $-1 < \alpha < \infty$. This result extends the classical theorem due to Forelli and Rudin for a much larger class of weights. In addition, it is shown that for any pair (ω, ν) of radial weights the Calderón operator

$$H_{\omega}^{\star}(f)(z) + H_{\omega}(f)(z) = \int_{0}^{|z|} f\left(s\frac{z}{|z|}\right) \frac{s\omega(s) ds}{\int_{s}^{1} t\omega(t) dt} + \frac{\int_{|z|}^{1} f\left(s\frac{z}{|z|}\right) s\omega(s) ds}{\int_{|z|}^{1} s\omega(s) ds} ds$$

is bounded on L^p_{ν} if and only if $A_p(\omega, \nu) < \infty$.

1. Introduction and main results

Let $\mathcal{H}(\mathbb{D})$ denote the space of analytic functions in the unit disc $\mathbb{D}=\{z\in\mathbb{C}:|z|<1\}$. For a nonnegative function $\omega\in L^1([0,1))$, its extension to \mathbb{D} , defined by $\omega(z)=\omega(|z|)$ for all $z\in\mathbb{D}$, is called a radial weight. For $0< p<\infty$ and such an ω , the Lebesgue space L^p_ω consists of complex-valued measurable functions f on \mathbb{D} such that

$$||f||_{L^p_\omega}^p = \int_{\mathbb{D}} |f(z)|^p \omega(z) \, dA(z) < \infty,$$

where $dA(z)=\frac{dx\,dy}{\pi}$ is the normalized Lebesgue area measure on \mathbb{D} . The corresponding weighted Bergman space is $A^p_\omega=L^p_\omega\cap\mathcal{H}(\mathbb{D})$. Throughout this paper we assume $\hat{\omega}(z)=\int_{|z|}^1\omega(s)\,ds>0$ for all $z\in\mathbb{D}$, for otherwise $A^p_\omega=\mathcal{H}(\mathbb{D})$. As usual, we write A^p_α for the classical weighted Bergman space induced by the standard radial weight $(\alpha+1)(1-|z|^2)^\alpha$, where $-1<\alpha<\infty$.

The Bergman Hilbert space A^2_{ω} is a closed subspace of L^2_{ω} , and the orthogonal Bergman projection from L^2_{ω} to A^2_{ω} is the integral operator

$$P_{\omega}(f)(z) = \int_{\mathbb{D}} f(\zeta) \overline{B_z^{\omega}(\zeta)} \,\omega(\zeta) dA(\zeta), \quad z \in \mathbb{D},$$

Date: December 5, 2021.

²⁰¹⁰ Mathematics Subject Classification. 30H20, 47B34, 42A38.

 $[\]it Key\ words\ and\ phrases.$ Bergman space, Bergman projection, Calderón operator, Stieltjes transform, radial doubling weight.

The first three authors are supported in part by Ministerio de Economía y Competitividad, Spain, projects PGC2018-096166-B-100; Junta de Andalucía, projects FQM210, FQM354 and UMA18-FEDERJA-002.

where $\{B_z^\omega\}_{z\in\mathbb{D}}$ are the reproducing kernels of A_ω^2 . The kernel of the classical weighted Bergman space A_α^2 is denoted by B_z^α , and P_α stands for the corresponding Bergman projection.

This paper concerns the question of when, for a given 1 , we have

$$||P_{\omega}(f)||_{L^{p}_{\nu}} \le C||f||_{L^{p}_{\nu}}, \quad f \in L^{p}_{\nu}.$$
 (1.1)

We also consider the one-weight inequality analogous to (1.1) for the maximal Bergman projection

$$P_{\omega}^{+}(f)(z) = \int_{\mathbb{D}} f(\zeta) |B_{z}^{\omega}(\zeta)| \omega(\zeta) dA(\zeta), \quad z \in \mathbb{D}.$$

The boundedness of projections on L^p -spaces is a compelling topic which has attracted a lot of attention during the last decades. Regarding the inequality (1.1), we refer to [2, 3, 4, 16, 19, 18]. The most known result concerning (1.1) is due to Bekollé and Bonami [3, 4], and it concerns the case when ν is an arbitrary weight and the inducing weight ω is standard. See [1, 19, 21] for recent extensions of this result. Some of the key tools in the proof of the aforementioned result of Bekollé and Bonami are based on the classical harmonic analysis, and rely strongly on the nice properties that the standard kernel has. In particular, the neat explicit formula $B_z^{\alpha}(\zeta) = (1-\overline{z}\zeta)^{-(2+\alpha)}$ shows that each B_z^{α} is zero-free and its modulus is essentially constant in hyperbolically bounded regions. However, these properties are no longer true for the Bergman reproducing kernel induced by an arbitrary radial weight. Namely, for a given radial weight ω there exists another radial weight ν such that $A_{\omega}^2 = A_{\nu}^2$, but B_z^{ν} have zeros, see the proof of [5, Theorem 2]. Going further, one of the main obstacles to tackle the inequality (1.1) is the lack of explicit expressions for B_z^{ω} . By and large, this kernel has the representation $B_z^{\omega}(\zeta) = \sum \overline{e_n(z)} e_n(\zeta)$ for each orthonormal basis $\{e_n\}$ of A_{ω}^2 . Therefore the normalized monomials show that the properties of the weight ω are transmitted to the projection P_{ω} through the representation $B_z^{\omega}(\zeta) = \sum_{n=0}^{\infty} \frac{(\overline{z}\zeta)^n}{2\omega_{2n+1}}$ of the kernel via its odd moments ω_{2n+1} . In general, from now on we write $\omega_x = \int_0^1 r^x \omega(r) dr$ for all $x \geqslant 0$.

To the best of our knowledge, describing the radial weights ω such that P_{ω} is bounded on L^p_{ω} is an open problem that was formally posed by Dostanić in [6] almost two decades ago. Even though this particular case $\omega = \nu$ of (1.1) remains unsettled, it was recently proved in [18, Theorem 7] that $P_{\omega}: L_{\omega}^{p} \to L_{\omega}^{p}$, $1 , is bounded if <math>\omega$ belongs to $\widehat{\mathcal{D}}$. The class $\widehat{\mathcal{D}}$ consists of the radial weights ω having the doubling property $\sup_{0 \leqslant r < 1} \frac{\widehat{\omega}(r)}{\widehat{\omega}(\frac{1+r}{2})} < \infty$. Further, mild extra regularity hypothesis on ω guarantee the necessity of the doubling condition $\omega \in \widehat{\mathcal{D}}$ for $P_{\omega}: L_{\omega}^{p} \to L_{\omega}^{p}$ to be bounded [18, Corollary 8]. Due to these results we will study the inequality (1.1) under the initial hypothesis $\omega \in \widehat{\mathcal{D}}$. Another class of doubling weights that naturally appears in our study is denoted by $\check{\mathcal{D}}$, and it consists of radial weights ω for which there exist constants $K = K(\omega) > 1$ and $C = C(\omega) > 1$ such that $\widehat{\omega}(r) \ge C\widehat{\omega}\left(1 - \frac{1-r}{K}\right)$ for all $0 \le r < 1$. The intersection $\hat{\mathcal{D}} \cap \check{\mathcal{D}}$ of these classes of weights is denoted by \mathcal{D} . The next few lines are dedicated to offer a brief insight to these classes. Each standard radial weight $(\alpha+1)(1-|z|^2)^{\alpha}$ belongs \mathcal{D} , while $\mathcal{D}\setminus\mathcal{D}$ contains weights that tend to zero exponentially. The class of rapidly increasing weights, introduced in [15], lies entirely within $\widehat{\mathcal{D}} \setminus \mathcal{D}$, and a typical example of such a weight is $\nu(z) = (1 - |z|^2)^{-1} \left(\log \frac{e}{1 - |z|^2}\right)^{-\alpha}$, where $\alpha > 1$. To this end we emphasize that the containment in $\hat{\mathcal{D}}$ or $\check{\mathcal{D}}$ does not require continuity neither positivity. In fact, weights in these classes may vanish on a relatively large part of each outer annulus $\{z:r\leqslant |z|<1\}$ of $\mathbb D$. For basic properties of aforementioned classes, concrete nontrivial examples and more, see [13, 15, 18] and the relevant references therein.

The recent papers [10, 20] show that the (radial) averaging operator

$$H_{\omega}(f)(z) = \frac{\int_{|z|}^{1} f\left(s\frac{z}{|z|}\right) \omega_{1}(s) ds}{\widehat{\omega}_{1}(|z|)}, \quad z \in \mathbb{D} \setminus \{0\}, \quad \omega_{1}(s) = s\omega(s),$$

can be considered as a toy model to study the pairs of radial weights (ω, ν) such that (1.1) holds. In order to make this statement precise, for $1 and radial weights <math>\omega$ and ν ,

write
$$\sigma(r) = \sigma_{p,\omega,\nu}(r) = \left(\frac{\omega(r)}{\nu(r)^{\frac{1}{p}}}\right)^{p'}$$
, and define

$$M_{p}(\omega,\nu) = \sup_{0 < r < 1} \left(\int_{0}^{r} \frac{\nu_{1}(s)}{\widehat{\omega}_{1}(s)^{p}} ds + 1 \right)^{\frac{1}{p}} \widehat{\sigma}_{1}(r)^{\frac{1}{p'}}.$$
 (1.2)

With this notation the classical result of Muckenhoupt [11, Theorem 2] states that $H_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded if and only if $M_p(\omega, \nu) < \infty$. This together with minor modifications in the proof of [18, Theorem 13] yields the following result, which is a substantial improvement of [10, Corollary 1.2].

Theorem A. Let $1 and <math>\omega \in \widehat{\mathcal{D}}$, and let ν be a radial weight. Then the following statements are equivalent:

- (i) $P_{\omega}^+: L_{\nu}^p \to L_{\nu}^p$ is bounded;
- (ii) $P_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded and $\nu \in \check{\mathcal{D}}$;
- (iii) $H_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded and $\omega, \nu \in \mathcal{D}$;
- (iv) $M_p(\omega, \nu) < \infty \text{ and } \omega, \nu \in \mathcal{D}.$

This result characterizes the one-weight inequality (1.1) whenever $\omega \in \hat{\mathcal{D}}$ and $\nu \in \check{\mathcal{D}}$, and establishes clear connections between the projections P_{ω} and P_{ω}^{+} , and the averaging operator H_{ω} . What remains unsettled is (1.1) when $\omega \in \hat{\mathcal{D}}$ but the hypothesis $\nu \in \check{\mathcal{D}}$ is removed. In order to approach this question, we write

$$A_p(\omega, \nu) = \sup_{0 \le r < 1} \frac{\widehat{\nu_1}(r)^{\frac{1}{p}} \widehat{\sigma_1}(r)^{\frac{1}{p'}}}{\widehat{\omega_1}(r)}.$$
 (1.3)

It is known by [18, Theorem 13] that also the condition $A_p(\omega, \nu) < \infty$ characterizes the bounded Bergman projections $P_{\omega}: L^p_{\nu} \to L^p_{\nu}$, provided $\omega \in \widehat{\mathcal{D}}$ and $\nu \in \widecheck{\mathcal{D}}$. Further, this condition implies $M_p(\omega, \nu) < \infty$ for any pair of radial weights (ω, ν) , see the proof of [18, Theorem 13] or Theorem 4 below. Going further, we will prove the following result.

Lemma B. Let $1 and <math>\omega \in \widehat{\mathcal{D}}$, and let ν be a radial weight. If $P_{\omega} : L^p_{\nu} \to L^p_{\nu}$ is bounded, then $A_p(\omega, \nu) < \infty$.

This lemma is essentially contained in the proof of [18, Theorem 13], but we provide the details for the convenience of the reader and the sake of completeness.

To get the big picture right, let us now consider the radial weights $\omega \equiv 1$ and $\nu(s) = (1-s^2)^{-1} \left(\log \frac{e}{1-s^2}\right)^{-\alpha}$, where $\alpha > 1$. Obviously, $\omega \in \hat{\mathcal{D}}$, and straightforward calculations show that $\nu \notin \check{\mathcal{D}}$, $M_p(\omega,\nu) < \infty$ but $A_p(\omega,\nu) = \infty$. Therefore P_ω is not bounded on L^p_ν for any $1 by Lemma B. This forces us to focus on the condition <math>A_p(\omega,\nu) < \infty$ and rules out $M_p(\omega,\nu) < \infty$ as a candidate to describe the couples of radial weights (ω,ν) such that (1.1) holds under the initial hypothesis $\omega \in \hat{\mathcal{D}}$. Unfortunately, we can not judge whether or not the reverse statement to Lemma B is true. This is, in our opinion, an intriguing matter and deserves to be stated as an open question.

Question 1. Let $1 and <math>\omega \in \widehat{\mathcal{D}}$, and let ν be a radial weight such that $A_p(\omega, \nu) < \infty$. Is then P_{ω} bounded on L^p_{ν} ?

Despite the fact that we are not able to answer Question 1 in its full generality, we provide an affirmative answer in an important particular case.

Theorem 2. Let $1 , <math>-1 < \alpha < \infty$, $\omega \in \widehat{\mathcal{D}}$ and $\nu = \omega \widehat{\omega_1}^{\alpha}$. Then $P_{\omega} : L^p_{\nu} \to L^p_{\nu}$ is bounded if and only if $A_p(\omega, \nu) < \infty$. Moreover, $P^+_{\omega} : L^p_{\nu} \to L^p_{\nu}$ is bounded if and only if $A_p(\omega, \nu) < \infty$ and $\omega \in \mathcal{D}$.

The classical result [23, Theorem 4.24] for the standard weights, attributed to Forelli and Rudin [8], states that $P_{\gamma}: L^p_{\beta} \to L^p_{\beta}$ is bounded if and only $p(\gamma+1) > \beta+1$. This result is just a particular case of Theorem 2. Namely, if $\omega(s) = (1-s^2)^{\gamma}$ and $\nu(s) = (1-s^2)^{\beta}$, where $\gamma, \beta > -1$, then $\nu = (2(\gamma+1))^{-\alpha}\omega\widehat{\omega_1}^{\alpha}$ for $\alpha = \frac{\beta-\gamma}{\gamma+1} > -1$, and $p(\gamma+1) > \beta+1$ if and only if $A_p(\omega,\nu) < \infty$. Since $\widehat{\mathcal{D}}$ is a very rich class of radial weights, the first statement in Theorem 2 is a far reaching extension of [23, Theorem 4.24]. The second statement is just a consequence of [18, Proposition 20] and the first statement. It is well-known among experts working on operator theory on spaces of analytic functions that the aforementioned result due to Forelli and Rudin is an extremely useful tool with a plethora of applications within this area of research [23]. Therefore we expect that Theorem 2 boosts the development of the operator theory in Bergman spaces A^p_{ω} induced by $\omega \in \widehat{\mathcal{D}}$.

It is very important to realize the distinction between \mathcal{D} and $\widehat{\mathcal{D}} \backslash \mathcal{D}$ when proving Theorem 2. Namely, since P_{ω} and P_{ω}^+ are not always simultaneously bounded when $\widehat{\mathcal{D}} \backslash \mathcal{D}$, any technique, where $|P_{\omega}(f)|$ is dominated by $P_{\omega}^+(|f|)$, is bound to fail all together to prove Theorem 2. In particular, this shows that arguments employed in the proofs of [16, Theorem 11] and [18, Theorem 13] do not serve us here. The techniques we will use to prove Theorem 2 are by no means standard. Our first step towards the theorem is the following reformulation of (1.1) in our setting via a duality relation between weighted Bergman spaces.

Theorem 3. Let $1 and <math>\omega \in \widehat{\mathcal{D}}$, and let ν be a radial weight. Then $P_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded if and only if $\left(A^{p'}_{\sigma}\right)^* \simeq A^p_{\nu}$ via the A^2_{ω} -pairing with equivalence of norms.

The next step in the proof of Theorem 2 is to observe that the dual of $A_{\sigma}^{p'}$ can be identified with the space of coefficient multipliers from $A_{\sigma}^{p'}$ to H^{∞} under the H^2 -pairing (the Cauchy pairing). As usual, for $0 , we write <math>H^p$ for the classical Hardy space. This observation becomes practical once we endow A_{ω}^p with a suitable equivalent norm. The one that works here is an ℓ^p -type norm of the Hardy norms of blocks of the Maclaurin series whose size depend on the weight ω , see [17, Theorem 3.4] and also [14, Theorem 4]. These arguments allow us to show that there exists a space $X_p(\omega,\nu)$ of analytic functions such that $\left(A_{\sigma}^{p'}\right)^{\star} \simeq X_p(\omega,\nu)$ via the H^2 -pairing. Finally, to get the desired pairing and space, we define the operator $I^{\omega}(g)(z) = \sum_{k=0}^{\infty} \widehat{g}(k)\omega_{2k+1}z^k$, and we prove that $I^{\omega}: A_{\nu}^p \to X_p(\omega,\nu)$ is isomorphic and onto. This establishes Theorem 2.

We now turn back to study relationships between the boundedness of the radial averaging operator H_{ω} on L^p_{ν} , and the condition $A_p(\omega,\nu)<\infty$. Recall that $M_p(\omega,\nu)<\infty$ describes the boundedness of H_{ω} on L^p_{ν} by [11, Theorem 2], and $A_p(\omega,\nu)<\infty$ always implies $M_p(\omega,\nu)<\infty$. We proceed with necessary definitions.

The adjoint of H_{ω} , with respect to $\langle \cdot, \cdot \rangle_{L^2_{\omega}}$, is the operator

$$H_{\omega}^{\star}(f)(z) = \int_{0}^{|z|} f\left(s \frac{z}{|z|}\right) \frac{\omega_{1}(s)}{\widehat{\omega}_{1}(s)} ds, \quad z \in \mathbb{D} \setminus \{0\}.$$

For a radial weight η , the associated Hörmander maximal function [9] and Stieltjes transfrom [22] are defined by

$$M_{\eta, \text{rad}}(f)(z) = \sup_{b < |z|} \frac{\int_b^1 \left| f\left(s \frac{z}{|z|}\right) \right| \eta(s) \, ds}{\widehat{\eta}(b)}, \quad z \in \mathbb{D} \setminus \{0\}, \tag{1.4}$$

and

$$S_{\eta, \text{rad}}(f)(z) = \int_0^1 \frac{f\left(s\frac{z}{|z|}\right)\eta(s)}{\widehat{\eta}(s) + \widehat{\eta}(|z|)} ds, \quad z \in \mathbb{D} \setminus \{0\},$$
 (1.5)

respectively. Our findings concerning the averaging radial operators and the condition (1.3) are gathered in the next result.

Theorem 4. Let $1 and let <math>\omega$ and ν be radial weights. Then the following statements are equivalent:

- (i) $H_{\omega} + H_{\omega}^{\star} : L_{\nu}^{p} \to L_{\nu}^{p}$ is bounded;
- (ii) $S_{\omega_1,\text{rad}}: L^p_{\nu} \to L^p_{\nu}$ is bounded;
- (iii) $M_{\omega_1, \text{rad}}: L^p_{\nu} \to L^p_{\nu}$ is bounded;
- (iv) $A_p(\omega, \nu) < \infty$.

Moreover,

$$\|H_{\omega} + H_{\omega}^{\star}\|_{L^{p}_{\omega} \to L^{p}_{\omega}} \simeq \|S_{\omega_{1}, \operatorname{rad}}\|_{L^{p}_{\omega} \to L^{p}_{\omega}} \simeq \|M_{\omega_{1}, \operatorname{rad}}\|_{L^{p}_{\omega} \to L^{p}_{\omega}} \simeq A_{p}(\omega, \nu).$$

As for the proof of Theorem 4, the inequality $\|M_{\omega_1, \text{rad}}\|_{L^p_{\nu} \to L^p_{\nu}} \lesssim A_p(\omega, \nu)$ boils down to a decomposition of the interval [0,1) in level sets for the maximal function $M_{\omega_1, \text{rad}} f_{\theta}$, where $f_{\theta}(r) = f(re^{i\theta})$, together with some properties of the maximal function. Since (iii) \Rightarrow (iv) is deduced by standard techniques, we have the equivalence between (iii) and (iv). After that, we observe the pointwise inequalities

$$M_{\omega_1, \text{rad}}(f)(z) \leq 2S_{\omega_1, \text{rad}}(f)(z) \leq H_{\omega}(f)(z) + H_{\omega}^{\star}(f)(z), \quad z \in \mathbb{D}, \quad f \geqslant 0,$$

which yield (i) \Rightarrow (ii) \Rightarrow (iii). Finally, the implication (iv) \Rightarrow (i) follows from the inequalities $\|H_{\omega}\|_{L^p_{\nu}} \leq \|M_{\omega_1, \text{rad}}\|_{L^p_{\nu} \to L^p_{\nu}} \lesssim A_p(\omega, \nu)$ and duality relations.

The remaining part of the paper is organized as follows. Section 2 contains the proofs of Lemma B, Theorem 3 and Theorem 2 in the said order. Theorem 4 is proved in Section 3.

Throughout the paper we repeatedly use the fact that $\omega \in \mathcal{D}$ (resp. $\omega \in \mathcal{D}$) if and only if $\omega_1 \in \mathcal{D}$ (resp. $\omega_1 \in \mathcal{D}$). Further, as usual, for two non-negative functions A and B, the notation $A \lesssim B$ (or $B \gtrsim A$) means that there exists a constant C > 0, independent of the variables involved, such that $A \leqslant CB$. Furthermore, we will write $A \approx B$ when $A \lesssim B \lesssim A$.

2. Bergman Projection

We begin with a detailed proof of the necessary condition (1.3) for $P_{\omega}: L_{\nu}^{p} \to L_{\nu}^{p}$ to be bounded.

Proof of Lemma B. Assume that $P_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded. Then $P_{\omega}(f)$ is well defined almost everywhere on \mathbb{D} for each $f \in L^p_{\nu}$. It follows that

$$\int_{\mathbb{D}} |f(\zeta)| |B_z^{\omega}(\zeta)| \omega(\zeta) \, dA(\zeta) < \infty, \quad f \in L_{\nu}^p, \tag{2.1}$$

for almost every $z \in \mathbb{D}$. Now that $B_z^{\omega}(\zeta) = \sum_{n=0}^{\infty} \frac{(\overline{z}\zeta)^n}{2\omega_{2n+1}}$, with $\omega_{2n+1} = \int_0^1 s^{2n+1}\omega(s)\,ds$, we have $B_0^{\omega}(\zeta) = \frac{1}{2\omega_1}$. Therefore there exists $\delta = \delta(\omega) > 0$ such that $|B_z^{\omega}(\zeta)| \geqslant \frac{1}{4\omega_1}$ for all $\zeta \in \mathbb{D}$ and z such that $|z| < \delta$. This together with (2.1) yields $L_{\nu}^p \subset L_{\omega}^1$. Let us show that the embedding is continuous. If this were not the case, then for each $n \in \mathbb{N}$ there would exist $f_n \in L_{\nu}^p$ such that $\|f_n\|_{L_{\nu}^p} = 1$ and $\|f_n\|_{L_{\omega}^1} \geqslant n^3$. Consider the function $F = \sum_{n=1}^{\infty} \frac{|f_n|}{n^2}$. It satisfies

$$||F||_{L^p_{\nu}} \le \sum_{n=1}^{\infty} \frac{||f_n||_{L^p_{\nu}}}{n^2} = \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$$
 (2.2)

and $F \geqslant \frac{|f_n|}{n^2}$ for each $n \in \mathbb{N}$. Therefore $||F||_{L^1_{\omega}} \geqslant \frac{||f_n||_{L^1_{\omega}}}{n^2} \geqslant n$ for each $n \in \mathbb{N}$, and thus $F \notin L^1_{\omega}$. This together with (2.2) contradicts $L^p_{\nu} \subset L^1_{\omega}$. Consequently, there exists a constant $C = C(\omega, \nu, p) > 0$ such that

$$||f||_{L^1_{co}} \le C||f||_{L^p_{co}}, \quad f \in L^p_{\nu}.$$
 (2.3)

This inequality implies that ω is absolutely continuous with respect to ν , and hence the function $f_n = \min\{n, \left(\frac{\omega}{\nu}\right)^{\frac{1}{p-1}}\}$ is well-defined almost everywhere on $\mathbb D$ for all $n \in \mathbb N$. Moreover,

 $f_n^p \nu \leqslant f_n \omega$ almost everywhere, and hence (2.3) yields $\sup_n \|f_n\|_{L^1_\omega} \leqslant C^{p'}$. The monotone convergence theorem now shows that

$$2\int_{0}^{1} s \left(\frac{\omega(s)}{\nu(s)^{\frac{1}{p}}}\right)^{p'} ds = \int_{\mathbb{D}} \lim_{n \to \infty} |f_{n}(z)| \omega(z) dA(z) = \lim_{n \to \infty} ||f_{n}||_{L_{\omega}^{1}} < \infty.$$
 (2.4)

Recall that the adjoint operator P_{ω}^* is defined via the identity

$$\langle P_{\omega}(f), g \rangle_{L^{2}_{\omega}} = \langle f, P_{\omega}^{*}(g) \rangle_{L^{2}_{\omega}}, \quad f \in L^{p}_{\nu}, \quad g \in L^{p'}_{\nu}. \tag{2.5}$$

We next establish an integral formula for the image of the monomial $m_n(z) = z^n$ under P_{ω}^* . First observe that for each fixed $z \in \mathbb{D}$ the series $\sum_{k=0}^{\infty} \frac{(z\overline{\zeta})^k}{2\omega_{2k+1}}$ converges uniformly on $\overline{\mathbb{D}}$. By using this and Fubini's theorem we deduce that each bounded function f satisfies

$$\langle P_{\omega}(f), m_{n} \rangle_{L_{\nu}^{2}} = \int_{\mathbb{D}} \left(\int_{\mathbb{D}} f(\zeta) B_{\zeta}^{\omega}(z) \omega(\zeta) \, dA(\zeta) \right) \overline{m_{n}(z)} \nu(z) \, dA(z)$$

$$= \int_{\mathbb{D}} \left(\sum_{k=0}^{\infty} \frac{\left(\int_{\mathbb{D}} f(\zeta) \overline{\zeta}^{k} \omega(\zeta) \, dA(\zeta) \right) z^{k}}{2\omega_{2k+1}} \right) \overline{m_{n}(z)} \nu(z) \, dA(z)$$

$$= \int_{\mathbb{D}} f(\zeta) \frac{\left(\int_{\mathbb{D}} \left(z \overline{\zeta} \right)^{n} \overline{m_{n}(z)} \nu(z) \, dA(z) \right)}{2\omega_{2n+1}} \omega(\zeta) \, dA(\zeta)$$

$$= \int_{\mathbb{D}} f(\zeta) \overline{\left(\frac{\omega(\zeta)}{\nu(\zeta)} \int_{\mathbb{D}} m_{n}(z) B_{z}^{\omega}(\zeta) \nu(z) \, dA(z) \right)} \nu(\zeta) \, dA(\zeta).$$

$$(2.6)$$

Define

$$L_{\omega}(g)(\zeta) = \frac{\omega(\zeta)}{\nu(\zeta)} \int_{\mathbb{D}} g(z) B_z^{\omega}(\zeta) \nu(z) \, dA(z), \quad \zeta \in \mathbb{D}, \quad g \in L_{\nu}^1.$$

Observe that $L_{\omega}(m_n)(\zeta) = \frac{\omega(\zeta)\nu_{2n+1}}{\nu(\zeta)\omega_{2n+1}}\zeta^n$ and $L_{\omega}(m_n) \in L_{\nu}^{p'}$ for all $n \in \mathbb{N}$ by (2.4). For each $f \in L_{\nu}^p$, choose a sequence of simple functions $\{s_k\}$ which converges to f in L_{ν}^p . Then, since $P_{\omega}: L_{\nu}^p \to L_{\nu}^p$ is bounded by the hypothesis, (2.6) yields

$$\langle P_{\omega}(f), m_n \rangle_{L^2_{\nu}} = \lim_{k \to \infty} \langle P_{\omega}(s_k), m_n \rangle_{L^2_{\nu}} = \lim_{k \to \infty} \langle s_k, L_{\omega}(m_n) \rangle_{L^2_{\nu}} = \langle f, L_{\omega}(m_n) \rangle_{L^2_{\nu}},$$

and hence $P_{\omega}^{\star}(m_n) = L_{\omega}(m_n)$ almost everywhere on \mathbb{D} .

Since $\omega \in \widehat{\mathcal{D}}$ by the hypothesis of the theorem and $P_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded, [18, Proposition 19(i)] implies $\nu_1 \in \widehat{\mathcal{D}}$. Therefore [13, Lemma 2.1] yields $||m_n||_{L^{p'}_{\nu}}^{p'} = 2\nu_{np'+1} \asymp \widehat{\nu_1} \left(1 - \frac{1}{n}\right) \asymp \nu_{2n+1}$ and $\omega_{2n+1} \asymp \widehat{\omega_1} \left(1 - \frac{1}{n}\right)$ for all $n \in \mathbb{N}$. Now that $P^*_{\omega}: L^{p'}_{\nu} \to L^{p'}_{\nu}$ is bounded, we deduce

$$\widehat{\nu_1} \left(1 - \frac{1}{n} \right) \approx \|m_n\|_{L^{p'}_{\nu}}^{p'} \gtrsim \|P_{\omega}^{\star}(m_n)\|_{L^{p'}_{\nu}}^{p'} = 2 \left(\frac{\nu_{2n+1}}{\omega_{2n+1}} \right)^{p'} \int_0^1 s^{np'+1} \left(\frac{\omega(s)}{\nu(s)^{\frac{1}{p}}} \right)^{p'} ds$$

$$\gtrsim \left(\frac{\widehat{\nu_1} \left(1 - \frac{1}{n} \right)}{\widehat{\omega_1} \left(1 - \frac{1}{n} \right)} \right)^{p'} \widehat{\sigma_1} \left(1 - \frac{1}{n} \right), \quad n \in \mathbb{N} \setminus \{1\}.$$

which implies

$$\sup_{1/2\leqslant r<1}\frac{\widehat{\nu_1}(r)^{\frac{1}{p}}\widehat{\sigma_1}(r)^{\frac{1}{p'}}}{\widehat{\omega_1}(r)}<\infty$$

since $\omega_1 \in \widehat{\mathcal{D}}$. Then bearing in mind (2.4), it follows that $A_n(\omega, \nu) < \infty$.

Now we proceed with the proof of the convenient reformulation of (1.1) in terms of an appropriate duality.

Proof of Theorem 3. Assume first that $(A_{\sigma}^{p'})^{\star} \simeq A_{\nu}^{p}$ via the A_{ω}^{2} -pairing with equivalence of norms. Let $H \in L_{\nu}^{p}$ and define $h = H\left(\frac{\omega}{\nu}\right)^{-p'/p}$ outside of the set on which ω vanishes. Then

 $\|h\|_{L^p_{\sigma}} = \|H\|_{L^p_{\nu}}$. Consider the bounded linear functional $T_h(f) = \langle f, h \rangle_{L^2_{\sigma}}$ on $L^{p'}_{\sigma}$ that satisfies $\|T_h\| = \|h\|_{L^p_{\sigma}}$. Then $T_h(f) = \langle f, H \rangle_{L^2_{\omega}}$ and $\|T_h\| = \|h\|_{L^p_{\sigma}} = \|H\|_{L^p_{\nu}}$. For each polynomial p, with the Maclaurin coefficients $\{\hat{p}(n)\}$, Fubini's theorem yields

$$T_{h}(p) = \langle P_{\omega}(p), H \rangle_{L_{\omega}^{2}} = \lim_{r \to 1^{-}} \int_{D(0,r)} P_{\omega}(p)(z) \overline{H(z)} \omega(z) \, dA(z)$$

$$= \lim_{r \to 1^{-}} \int_{\mathbb{D}} p(\zeta) \left(\sum_{n=0}^{\infty} \frac{\int_{D(0,r)} \overline{H(z)} z^{n} \omega(z) \, dA(z)}{2\omega_{2n+1}} \overline{\zeta}^{n} \right) \omega(\zeta) \, dA(\zeta)$$

$$= \lim_{r \to 1^{-}} \sum_{n=0}^{\infty} \widehat{p}(n) \int_{D(0,r)} \overline{H(z)} z^{n} \omega(z) \, dA(z) = \langle p, P_{\omega}(H) \rangle_{L_{\omega}^{2}}.$$

$$(2.7)$$

By the hypothesis, there exists $g \in A^p_{\nu}$ such that $T_h(f) = \langle f, g \rangle_{A^2_{\omega}}$ for all $f \in A^{p'}_{\sigma}$, and $||T_h|| \approx ||g||_{A^p_{\nu}}$. In particular, $T_h(m_n) = \langle m_n, g \rangle_{A^2_{\omega}} = \langle m_n, P_{\omega}(H) \rangle_{A^2_{\omega}}$ for each monomial $m_n(z) = z^n$ with $n \in \mathbb{N} \cup \{0\}$. It follows that $P_{\omega}(H) = g$, and hence $||P_{\omega}(H)||_{A^p_{\nu}} \approx ||T_h|| = ||H||_{L^p_{\nu}}$. Since $H \in L^p_{\nu}$ was arbitrarily taken, this shows that $P_{\omega} : L^p_{\nu} \to L^p_{\nu}$ is bounded.

Reciprocally, assume that $P_{\omega}: L^p_{\nu} \to L^p_{\nu}$ is bounded, then (1.3) holds by Lemma B, and in particular σ is a weight. Let $T \in (A^{p'}_{\sigma})^*$, then T can be extended to a bounded linear functional \widetilde{T} on $L^{p'}_{\sigma}$ with $\|T\| = \|\widetilde{T}\|$ by the Hahn-Banach theorem. By arguing as in the first part of the proof, we see that $(L^{p'}_{\sigma})^*$ is isometrically isomorphic to L^p_{ν} via the L^2_{ω} -pairing. Hence there exists $H \in L^p_{\nu}$ such that $\widetilde{T}(f) = \langle f, H \rangle_{L^2_{\omega}}$ for all $f \in L^{p'}_{\sigma}$, and $\|\widetilde{T}\| = \|H\|_{L^p_{\nu}}$. Now, for each polynomial p, with Maclaurin coefficients $\{\widehat{p}(n)\}$, (2.7) yields

$$T(p) = \langle P_{\omega}(p), H \rangle_{L^{2}_{\omega}} = \langle p, P_{\omega}(H) \rangle_{L^{2}_{\omega}} = \langle p, g \rangle_{L^{2}_{\omega}},$$

where $g=P_{\omega}(H)\in A^p_{\nu}$ satisfies $\|g\|_{A^p_{\nu}}\leqslant \|P_{\omega}\|_{L^p_{\nu}\to L^p_{\nu}}\|H\|_{L^p_{\nu}}=\|P_{\omega}\|_{L^p_{\nu}\to L^p_{\nu}}\|T\|<\infty$ by the hypothesis. Bearing in mind that the polynomials are dense in A^p_{σ} , because σ is radial, and using Hölder's inequality, it follows that $T(f)=\langle f,g\rangle_{A^2_{\omega}}$ for all $f\in A^{p'}_{\sigma}$ with $\|g\|_{A^p_{\nu}}\lesssim \|T\|$. Since in addition, each functional $T_g(f)=\langle f,g\rangle_{A^2_{\omega}}$ induced by $g\in A^p_{\nu}$, belongs to $(A^{p'}_{\sigma})^*$ with $\|T_g\|\leqslant \|g\|_{A^p_{\nu}}$ by Hölder's inequality, we deduce $(A^{p'}_{\sigma})^*\simeq A^p_{\nu}$ via the A^2_{ω} -pairing with equivalence of norms.

The proof of Theorem 2 requires some preparations. In particular, we will use a suitable norm in the weighted Bergman space induced by a weight in $\widehat{\mathcal{D}}$. For that purpose some more notation are in order. The Hadamard product of $f(z) = \sum_{k=0}^{\infty} \widehat{f}(k) z^k \in \mathcal{H}(\mathbb{D})$ and $g(z) = \sum_{k=0}^{\infty} \widehat{g}(k) z^k \in \mathcal{H}(\mathbb{D})$ is

$$(f * g)(z) = \sum_{k=0}^{\infty} \widehat{f}(k)\widehat{g}(k)z^k, \quad z \in \mathbb{D}.$$

For a Banach space $X \subset \mathcal{H}(\mathbb{D}), s \in \mathbb{R}, 0 < q \leq \infty$ and a sequence of polynomials $\{P_n\}_{n=0}^{\infty}$, let

$$\ell_s^q(X, \{P_n\}) = \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{\ell_s^q(X, \{P_n\})}^q = \sum_{n=0}^{\infty} \left(2^{-ns} \|P_n * f\|_X \right)^q < \infty \right\}, \quad 0 < q < \infty, \quad (2.8)$$

and

$$\ell_s^{\infty}(X, \{P_n\}) = \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{\ell_s^{\infty}(X, \{P_n\})} = \sup_n \left(2^{-ns} \|P_n * f\|_X \right) < \infty \right\}.$$
 (2.9)

Further, for quasi-normed spaces $X,Y \subset \mathcal{H}(\mathbb{D})$, let $[X,Y] = \{g \in \mathcal{H}(\mathbb{D}) : f * g \in Y \text{ for all } f \in X\}$ denote the space of coefficient multipliers from X to Y equipped with the norm

$$\|g\|_{\lceil X,Y\rceil} = \sup \left\{ \|f * g\|_Y : f \in X, \, \|f\|_X \leqslant 1 \right\}.$$

With these preparations we are ready for the proof.

Proof of Theorem 2. Bearing in mind Lemma B and Theorem 3 we only have to prove that $\left(A_{\sigma}^{p'}\right)^{\star} \simeq A_{\nu}^{p}$ via the A_{ω}^{2} -pairing with equivalence of norms, in order to ensure that P_{ω} is

bounded on L^p_{ν} . Observe that $\sigma = \omega \widehat{\omega_1}^{-\alpha \frac{p'}{p}}$, and hence the condition $A_p(\omega, \nu) < \infty$ is equivalent to the inequality $-1 < \alpha < \frac{p}{p'}$. Assume, without loss of generality, that $\widehat{\omega_1}(0) = 1$, and consider a sequence $\{\rho_n\}_{n=0}^{\infty} = \{\rho_n(\omega)\}_{n=0}^{\infty}$ such that $\widehat{\omega}_1(\rho_n) = 2^{-n}$ for all n. For $n \in \mathbb{N}$, define $M_n = M_n(\omega) = E\left(\frac{1}{1-\rho_n}\right)$, where $E(x) \in \mathbb{N}$ such that $E(x) \leq x < E(x) + 1$. Write

$$I(0) = I_{\omega}(0) = \{k \in \mathbb{N} \cup \{0\} : k < M_1\}$$

and

$$I(n) = I_{\omega}(n) = \{k \in \mathbb{N} : M_n \leqslant k < M_{n+1}\}, \quad n \in \mathbb{N}.$$

For $f \in \mathcal{H}(\mathbb{D})$, with the Maclaurin series $f(z) = \sum_{n=0}^{\infty} \hat{f}(n)z^n$, define the polynomials $\Delta_n^{\omega} f$ by

$$\Delta_n^{\omega} f(z) = \sum_{k \in I_{\omega}(n)} \widehat{f}(k) z^k, \quad n \in \mathbb{N} \cup \{0\}.$$

Then, obviously, $f = \sum_{n=0}^{\infty} \Delta_n^{\omega} f$. Observe that $\nu, \sigma \in \widehat{\mathcal{D}}$ because $\omega \in \widehat{\mathcal{D}}$ and moreover

$$\widehat{\nu}_1(\rho_n) = \frac{2^{-n(1+\alpha)}}{1+\alpha} \quad \text{and} \quad \widehat{\sigma}_1(\rho_n) = \frac{2^{-n\left(1-\alpha\frac{p'}{p}\right)}}{1-\alpha\frac{p'}{p}}, \quad n \in \mathbb{N} \cup \{0\}.$$

Therefore, [17, Theorem 3.4], see also [12, Theorem 4.1] and [14, Theorem 4] for similar results, shows that

$$||f||_{A_{\nu}^{p}}^{p} \approx \sum_{n=0}^{\infty} \hat{\nu}_{1}(\rho_{n}) ||\Delta_{n}^{\omega} f||_{H^{p}}^{p} \approx \sum_{n=0}^{\infty} 2^{-n(1+\alpha)} ||\Delta_{n}^{\omega} f||_{H^{p}}^{p}, \quad f \in \mathcal{H}(\mathbb{D}),$$
 (2.10)

and

$$||f||_{A_{\sigma}^{p'}}^{p'} \approx \sum_{n=0}^{\infty} \widehat{\sigma_1}(\rho_n) ||\Delta_n^{\omega} f||_{H^{p'}}^{p'} \approx \sum_{n=0}^{\infty} 2^{-n\left(1-\alpha\frac{p'}{p}\right)} ||\Delta_n^{\omega} f||_{H^{p'}}^{p'}, \quad f \in \mathcal{H}(\mathbb{D}).$$
 (2.11)

Thus $A^p_{\nu} = \ell^p_{\frac{\alpha+1}{2}}(H^p, \{\Delta^{\omega}_n\})$ and $A^{p'}_{\sigma} = \ell^{p'}_{\frac{1}{2'} - \frac{\alpha}{2}}(H^{p'}, \{\Delta^{\omega}_n\})$ with equivalent norms.

In addition, $\left(A_{\sigma}^{p'}\right)^{\star}$ can be identified with $\left[A_{\sigma}^{p'}, H^{\infty}\right]$ via the H^2 -pairing with equivalence of norms by [12, Proposition 1.3]. That is, for each $L \in \left(A_{\sigma}^{p'}\right)^{\star}$ there exists a unique $g \in \left[A_{\sigma}^{p'}, H^{\infty}\right]$ such that $||L|| = ||g||_{[A^{p'}_{\sigma}, H^{\infty}]}$ and $L(f) = \langle f, g \rangle_{H^2}$ for all $f \in A^{p'}_{\sigma}$, and conversely, each functional $L(f) = \langle f, g \rangle_{H^2}$, induced by $g \in [A^{p'}_{\sigma}, H^{\infty}]$, belongs to $(A^{p'}_{\sigma})^{\star}$ and satisfies $||L|| = ||g||_{[A^{p'}_{\sigma}, H^{\infty}]}$. Now use [12, Theorem 5.4] and the fact that $(H^{p'})^{\star} \simeq [H^{p'}, H^{\infty}] = H^p$ via the H^2 -pairing to deduce

$$\left[\ell^{p'}_{\frac{1}{p'}-\frac{\alpha}{p}}(H^{p'},\{\Delta^{\omega}_n\}),H^{\infty}\right] = \ell^{p}_{\frac{\alpha}{p}-\frac{1}{p'}}\left(\left[H^{p'},H^{\infty}\right],\{\Delta^{\omega}_n\}\right) = \ell^{p}_{\frac{\alpha}{p}-\frac{1}{p'}}\left(H^{p},\{\Delta^{\omega}_n\}\right) \tag{2.12}$$

with equivalent norms. Since $A^{p'}_{\sigma}=\ell^{p'}_{\frac{1}{n'}-\frac{\alpha}{p}}(H^{p'},\{\Delta^{\omega}_n\})$ with equivalent norms, it follows that

$$\left(A_{\sigma}^{p'}\right)^{\star} \simeq \left[A_{\sigma}^{p'}, H^{\infty}\right] = \ell_{\frac{\alpha}{p} - \frac{1}{p'}}^{p} \left(H^{p}, \{\Delta_{n}^{\omega}\}\right), \tag{2.13}$$

via the H^2 -pairing with equivalence of norms. Finally, define the operator $I^\omega:\mathcal{H}(\mathbb{D})\to\mathcal{H}(\mathbb{D})$ by $I^\omega(g)(z)=\sum_{k=0}^\infty \widehat{g}(k)\omega_{2k+1}z^k$ for all $z\in\mathbb{D}$. The sequence $\{\omega_{2k+1}\}_{k=0}^\infty$ is non-increasing, and hence there exists a constant C>0such that

$$C^{-1}\omega_{1+M_{n+1}} \|\Delta_n^{\omega} g\|_{H^p} \leq \|\Delta_n^{\omega} I^{\omega}(g)\|_{H^p} \leq C\omega_{1+M_n} \|\Delta_n^{\omega} g\|_{H^p}, \quad g \in \mathcal{H}(\mathbb{D}),$$

by [14, Lemma E]. This combined with $(\omega_1)_x = \widehat{\omega_1} \left(1 - \frac{1}{x}\right)$, valid for all $1 \le x < \infty$ by [13, Lemma 2.1(vi)], gives $\|\Delta_n^{\omega} I^{\omega}(g)\|_{H^p} \approx 2^{-n} \|\Delta_n^{\omega} g\|_{H^p}$ for all $g \in \mathcal{H}(\mathbb{D})$. Therefore (2.10) yields

$$\begin{split} \|I^{\omega}(g)\|_{\ell^{p}_{\frac{\alpha}{p}-\frac{1}{p'}}(H^{p},\{\Delta^{\omega}_{n}\})}^{p} &= \sum_{n=0}^{\infty} 2^{-n\left(\alpha-\frac{p}{p'}\right)} \|\Delta^{\omega}_{n}I^{\omega}(g)\|_{H^{p}}^{p} \asymp \sum_{n=0}^{\infty} 2^{-n\left(\alpha+p-\frac{p}{p'}\right)} \|\Delta^{\omega}_{n}g\|_{H^{p}}^{p} \\ &= \sum_{n=0}^{\infty} 2^{-n(\alpha+1)} \|\Delta^{\omega}_{n}g\|_{H^{p}}^{p} \asymp \|g\|_{A^{p}_{\nu}}^{p}, \quad g \in \mathcal{H}(\mathbb{D}), \end{split}$$

and hence $I^{\omega}: A^p_{\nu} \to \ell^p_{\frac{\alpha}{p} - \frac{1}{p'}}([H^p, H^{\infty}], \{\Delta^{\omega}_n\})$ is isomorphic and onto. This converts $\left(A^{p'}_{\sigma}\right)^{\star} \simeq \ell^p_{\frac{\alpha}{p} - \frac{1}{p'}}(H^p, \{\Delta^{\omega}_n\})$ via the H^2 -pairing, valid by (2.13), to $\left(A^{p'}_{\sigma}\right)^{\star} \simeq A^p_{\nu}$ via the A^2_{ω} -pairing, and thus finishes the proof of the first statement.

Now assume that $A_p(\omega, \nu) < \infty$ and $\omega \in \mathcal{D}$. Then a calculation shows that $\nu = \omega \widehat{\omega}^{\alpha} \in \mathcal{D}$, and $P_{\omega}^+ : L_{\nu}^p \to L_{\nu}^p$ is bounded by [18, Theorem 13]. Reciprocally, if $P_{\omega}^+ : L_{\nu}^p \to L_{\nu}^p$ is bounded, then $A_p(\omega, \nu) < \infty$ and $\omega \in \mathcal{D}$ by [18, Theorem 13]. Alternatively, we can use Lemma B and [18, Proposition 20] to get the conditions $A_p(\omega, \nu) < \infty$ and $\omega \in \mathcal{D}$.

3. CALDERÓN AND STIELTJES OPERATORS, AND HÖRMANDER MAXIMAL FUNCTION

We will prove the following reformulation of Theorem 4 using ideas coming from [7].

Theorem 5. Let $1 and let <math>\omega$ be a radial weight. Let $\nu : [0,1) \to \mathbb{R}^+$ such that $\nu \in L^1_{\omega}([0,1))$. Then the following statements are equivalent:

- (i) $H_{\omega} + H_{\omega}^{\star} : L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}$ is bounded;
- (ii) $S_{\omega_1, \text{rad}} : L^p_{\omega \nu} \to L^p_{\omega \nu}$ is bounded;
- (iii) $M_{\omega_1, \text{rad}}: L^p_{\omega\nu} \to L^p_{\omega\nu}$ is bounded;
- (iv) $A_p(\omega, \omega \nu) < \infty$.

Moreover,

$$\|H_{\omega} + H_{\omega}^{\star}\|_{L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}} \approx \|S_{\omega_{1}, \text{rad}}\|_{L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}} \approx \|M_{\omega_{1}, \text{rad}}\|_{L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}} \approx A_{p}(\omega, \omega\nu). \tag{3.1}$$

Before embarking on the proof of Theorem 5, two auxiliary results are in order. For a radial weight η , denote

$$M_{\eta}f(t) = \sup_{b < t} \frac{\int_{b}^{1} |f(s)|\eta(s) \, ds}{\widehat{\eta}(b)}, \quad 0 < t < 1.$$
 (3.2)

By denoting $f_{\theta}(r) = f(re^{i\theta})$ for all $re^{i\theta} \in \mathbb{D}$, (1.4) shows that $M_{\eta,\text{rad}}(f)(z) = M_{\eta}f_{\theta}(r)$. Further, for a given measurable function $f, k \in \mathbb{Z}$ and a radial weight ω , define

$$E_k = E_k(f_\theta, \omega) = \{2^k < M_{\omega_1} f_\theta(r) \le 2^{k+1}\}, \quad 0 \le \theta < 2\pi.$$

For each θ , the function $M_{\omega_1} f_{\theta}$ is nondecreasing, and hence $E_k = (b_k(\theta), b_{k+1}(\theta)]$ for some $b_k(\theta) < b_{k+1}(\theta)$ and

$$\frac{\int_{b_k(\theta)}^1 f_\theta(s)\omega_1(s) ds}{\widehat{\omega}_1(b_k(\theta))} = 2^k, \quad k \in \mathbb{Z}, \quad 0 \leqslant \theta < 2\pi.$$
(3.3)

It follows from (3.3) that

$$\frac{\widehat{\omega_1}(b_{k+1}(\theta))}{\widehat{\omega_1}(b_k(\theta))} \leqslant \frac{1}{2}, \quad k \in \mathbb{Z}, \quad 0 \leqslant \theta < 2\pi, \tag{3.4}$$

and therefore

$$\widehat{\omega_1}(b_k(\theta)) = \int_{b_k(\theta)}^{b_{k+1}(\theta)} \omega_1(s) \, ds + \widehat{\omega_1}(b_{k+1}(\theta)) \leqslant \int_{b_k(\theta)}^{b_{k+1}(\theta)} \omega_1(s) \, ds + \frac{1}{2} \widehat{\omega_1}(b_k(\theta)).$$

Thus

$$\widehat{\omega}_1(b_k(\theta)) \leqslant 2 \int_{b_k(\theta)}^{b_{k+1}(\theta)} \omega_1(s) \, ds, \quad k \in \mathbb{Z}, \quad 0 \leqslant \theta < 2\pi.$$
(3.5)

Lemma 6. Let $1 and let <math>\omega$ be a radial weight. Let $\nu : [0,1) \to \mathbb{R}^+$ such that $A_p(\omega,\omega\nu)<\infty$. Then there exists a constant $C=C(\omega,\nu,p)>0$ such that

$$\int_{b_k(\theta)}^1 \nu(s)^{-\frac{p'}{p}} \omega(s) s \, ds \leqslant C \int_{b_k(\theta)}^{b_{k+1}(\theta)} \nu(s)^{-\frac{p'}{p}} \omega(s) s \, ds, \quad 0 \leqslant \theta < 2\pi.$$

Proof. The definition of $A_p(\omega,\omega\nu)$, (3.5) and Hölder's inequality yield

$$\left(\int_{b_{k}(\theta)}^{1} \nu(s)\omega(s)s \, ds\right)^{\frac{1}{p}} \left(\int_{b_{k}(\theta)}^{1} \nu(s)^{-\frac{p'}{p}}\omega(s)s \, ds\right)^{\frac{1}{p'}}
\lesssim \widehat{\omega_{1}}(b_{k}(\theta)) \lesssim \int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \omega_{1}(s) \, ds
\leqslant \left(\int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \nu(s)\omega(s)s \, ds\right)^{\frac{1}{p}} \left(\int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \nu(s)^{-\frac{p'}{p}}\omega(s)s \, ds\right)^{\frac{1}{p'}}
\leqslant \left(\int_{b_{k}(\theta)}^{1} \nu(s)\omega(s)s \, ds\right)^{\frac{1}{p}} \left(\int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \nu(s)^{-\frac{p'}{p}}\omega(s)s \, ds\right)^{\frac{1}{p'}},$$

and the assertion follows.

Lemma 7. Let $1 and let <math>\eta$ be radial weight. Then $M_{\eta}: L^{1}([0,1),\eta) \to L^{1,\infty}([0,1),\eta)$ is bounded with norm at most one. In particular, for each $1 , <math>M_{\eta}: L^{p}([0,1), \eta) \rightarrow$ $L^p([0,1),\eta)$ is bounded with norm depending only on p.

Proof. Since $M_{\eta}f$ is an increasing function, for each $\lambda > 0$ and $f \in L^1([0,1),\eta)$, the level set $\{t \in (0,1): M_{\eta}f(t) > \lambda\}$ coincides with the interval (b,1) for some $b = b(f,\eta,\lambda) \in (0,1)$ such that $\int_{b}^{1} |f(s)| \eta(s) ds = \widehat{\eta}(b) \lambda$. Therefore

$$\int_{\{s \in (0,1): M_n f(s) > \lambda\}} \eta(s) \, ds = \widehat{\eta}(b) = \frac{\int_b^1 |f(s)| \eta(s) \, ds}{\lambda} \leqslant \frac{\int_0^1 |f(t)| \eta(s) \, ds}{\lambda},$$

and thus $M_{\eta}: L^{1}([0,1),\eta) \to L^{1,\infty}([0,1),\eta)$ is bounded with norm at most one. Now that M_{η} is clearly bounded from $L^{\infty}([0,1),\eta)$ into itself with norm at most one, the statement for 1 follows from the Marcinkiewicz interpolation theorem.

Proof of Theorem 5. We begin with showing that (iii) and (iv) are equivalent. Assume first (iv), that is, $A_p(\omega,\omega\nu) < \infty$. Then, for each $0 \le \theta < 2\pi$, by the definition of E_k , (3.3) and the hypothesis (iv), we have

$$I(f_{\theta}) = \int_{0}^{1} (M_{\omega_{1}} f_{\theta}(r))^{p} \nu(r) \omega_{1}(r) dr = \sum_{k \in \mathbb{Z}} \int_{E_{k}} (M_{\omega_{1}} f_{\theta}(r))^{p} \nu(r) \omega_{1}(r) dr$$

$$\leqslant 2^{p} \sum_{k \in \mathbb{Z}} 2^{kp} \int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \nu(r) \omega_{1}(r) dr$$

$$= 2^{p} \sum_{k \in \mathbb{Z}} \frac{\left(\int_{b_{k}(\theta)}^{1} f_{\theta}(r) \omega_{1}(r) dr\right)^{p} \int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \nu(r) \omega_{1}(r) dr}{\widehat{\omega_{1}}(b_{k}(\theta))^{p}}$$

$$\leqslant 2^{p} \sum_{k \in \mathbb{Z}} \frac{\left(\int_{b_{k}(\theta)}^{1} f_{\theta}(r) \omega_{1}(r) dr\right)^{p} \int_{b_{k}(\theta)}^{1} \nu(r) \omega_{1}(r) dr}{\widehat{\omega_{1}}(b_{k}(\theta))^{p}}$$

$$\leqslant 2^{p} A_{p}^{p}(\omega, \omega \nu) \sum_{k \in \mathbb{Z}} \left(\frac{\int_{b_{k}(\theta)}^{1} f_{\theta}(r) \omega_{1}(r) dr}{\int_{b_{k}(\theta)}^{1} \nu^{-\frac{p'}{p}}(r) \omega_{1}(r) dr}\right)^{p} \int_{b_{k}(\theta)}^{1} \nu^{-\frac{p'}{p}}(r) \omega_{1}(r) dr.$$

By applying now Lemmas 6 and 7, we deduce

$$I(f_{\theta}) \lesssim A_{p}^{p}(\omega, \omega \nu) \sum_{k \in \mathbb{Z}} \left(\frac{\int_{b_{k}(\theta)}^{1} f_{\theta}(r) \omega_{1}(r) dr}{\int_{b_{k}(\theta)}^{1} \nu^{-\frac{p'}{p}}(r) \omega_{1}(r) dr} \right)^{p} \int_{b_{k}(\theta)}^{b_{k+1}(\theta)} \nu^{-\frac{p'}{p}}(r) \omega_{1}(r) dr$$

$$\leqslant A_{p}^{p}(\omega, \omega \nu) \int_{0}^{1} \left(M_{\nu^{-\frac{p'}{p}} \omega_{1}}(f_{\theta} \nu^{\frac{p'}{p}})(r) \right)^{p} \nu^{-\frac{p'}{p}}(r) \omega_{1}(r) dr$$

$$\lesssim A_{p}^{p}(\omega, \omega \nu) \int_{0}^{1} \left(|f_{\theta}(r)| \nu^{\frac{p'}{p}}(r) \right)^{p} \nu^{-\frac{p'}{p}}(r) \omega_{1}(r) dr$$

$$= A_{p}^{p}(\omega, \omega \nu) \int_{0}^{1} |f_{\theta}(r)|^{p} \nu(r) \omega_{1}(r) dr, \quad 0 \leqslant \theta < 2\pi.$$

Since

$$\int_{\mathbb{D}} \left(M_{\omega_1, \text{rad}}(f)(z) \right)^p \nu(z) \omega(z) \, dA(z) = \frac{1}{\pi} \int_0^{2\pi} I(f_\theta) \, d\theta,$$

this shows that $M_{\omega_1,\mathrm{rad}}:L^p_{\omega\nu}\to L^p_{\omega\nu}$ is bounded with

$$||M_{\omega_1, \text{rad}}||_{L^p_{\omega\nu} \to L^p_{\omega\nu}} \lesssim A_p(\omega, \omega\nu). \tag{3.6}$$

Conversely, assume that (iii) holds, that is, $M_{\omega_1,\text{rad}}:L^p_{\omega\nu}\to L^p_{\omega\nu}$ is bounded. Then it follows that

$$\int_{0}^{1} (M_{\omega_{1}} f(r))^{p} \nu(r) \omega_{1}(r) dr \leq \|M_{\omega_{1}, \text{rad}}\|_{L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}}^{p} \int_{0}^{1} f(r)^{p} \omega_{1}(r) \nu(r) dr$$

for all non-negative radial functions f. By choosing $f(r) = \nu^{-\frac{p'}{p}}(r)\chi_{[a,1)}(r)$, where $a \in [0,1)$, and using standard arguments, we obtain

$$A_p(\omega, \omega \nu) \le \|M_{\omega_1, \text{rad}}\|_{L^p_{\omega \nu} \to L^p_{\omega \nu}}.$$
(3.7)

The pointwise inequality

$$S_{\omega_1, \text{rad}}(f)(z) \leqslant \frac{\int_{|z|}^1 f\left(s\frac{z}{|z|}\right) s\omega(s) \, ds}{\widehat{\omega}_1(z)} + \int_0^{|z|} f\left(s\frac{z}{|z|}\right) \frac{\omega_1(s)}{\widehat{\omega}_1(s)} \, ds$$
$$= H_{\omega}(f)(z) + H_{\omega}^{\star}(f)(z), \quad z \in \mathbb{D}, \quad f \geqslant 0,$$

yields

$$||S_{\omega_1, \text{rad}}||_{L^p_{\omega\nu} \to L^p_{\omega\nu}} \leq ||H_\omega + H^*_\omega||_{L^p_{\omega\nu} \to L^p_{\omega\nu}}, \tag{3.8}$$

and thus (i) implies (ii).

If b < |z|, then we have

$$\frac{\int_{b}^{1} f\left(s\frac{z}{|z|}\right) \omega(s) s \, ds}{\widehat{\omega}_{1}(b)} \leqslant 2 \int_{b}^{1} \frac{f\left(s\frac{z}{|z|}\right)}{\widehat{\omega}_{1}(b) + \widehat{\omega}_{1}(z)} \omega(s) s \, ds \leqslant 2S_{\omega_{1}, \mathrm{rad}}(f)(z), \quad z \in \mathbb{D}, \quad f \geqslant 0,$$

which implies $M_{\omega_1, \text{rad}}(f) \leq 2S_{\omega_1, \text{rad}}(f)$ for all $f \geq 0$. Therefore

$$||M_{\omega_1, \text{rad}}||_{L^p_{\omega_1} \to L^p_{\omega_1}} \le ||S_{\omega_1, \text{rad}}||_{L^p_{\omega_1} \to L^p_{\omega_1}},$$
 (3.9)

and thus (ii) implies (iii).

To complete the proof, it remains to show that (iv) implies (i) and

$$||H_{\omega} + H_{\omega}^{\star}||_{L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}} \lesssim A_{p}(\omega, \omega\nu).$$

Assume (iv), that is, $A_p(\omega, \omega \nu) < \infty$. Then $\omega \nu^{-\frac{p'}{p}}$ is a weight, and therefore $(L^p_{\omega \nu})^* \simeq L^{p'}_{\omega \nu^{-\frac{p'}{p}}}$ and $(L_{\omega\nu}^{p'})^* \simeq L_{\omega\nu}^p$ via the L_{ω}^2 -pairing. Therefore each linear operator T is bounded from $L^p_{\omega\nu}$ into itself if and only if its adjoint T^* , defined by the identity $\langle Tf, g \rangle_{L^2_{\omega}} = \langle f, T^*g \rangle_{L^2_{\omega}}$, is bounded from $L_{\omega\nu}^{p'}$ into itself, and the operator norms are comparable. Since $A_p(\omega,\omega\nu) < \infty$ by the assumption, (3.6) yields $\|H_{\omega_1,\mathrm{rad}}\|_{L_{\omega\nu}^p \to L_{\omega\nu}^p} \lesssim A_p(\omega,\omega\nu) < \infty$. Moreover, $A_p(\omega,\omega\nu) = 1$ $A_{p'}(\omega,\omega\nu^{-\frac{p'}{p}})$, and hence (3.6) shows that $H_{\omega_1,\mathrm{rad}}:L^{p'}_{\omega\nu^{-\frac{p'}{p}}}\to L^p_{\omega\nu^{-\frac{p'}{p}}}$ is bounded and the operator norm is bounded by a constant times $A_p(\omega,\omega\nu)$. It follows that $H^{\star}_{\omega_1,\mathrm{rad}}:L^p_{\omega\nu}\to L^p_{\omega\nu}$ is bounded, and $\|H_{\omega_1,\mathrm{rad}}^{\star}\|_{L_{\omega\nu}^p\to L_{\omega\nu}^p} \lesssim A_p(\omega,\omega\nu)$. Therefore

$$||H_{\omega} + H_{\omega}^{\star}||_{L_{\omega\nu}^{p} \to L_{\omega\nu}^{p}} \lesssim A_{p}(\omega, \omega\nu), \tag{3.10}$$

and thus (i) is satisfied. The estimates (3.1) for the operator norms follows from the inequalities (3.6)-(3.10).

References

- [1] A. Aleman and O. Constantin, The Bergman projection on vector-valued L^2 -spaces with operator-valued weights, J. Funct. Anal. 262, (2012) no. 5, 2359-2378.
- [2] A. Aleman, S. Pott and M.C. Reguera, Sarason conjecture on the Bergman space, Int. Math. Res. Not. IMRN 2017, no. 14, 4320-4349.
- [3] D. Bekollé, Inégalités á poids pour le projecteur de Bergman dans la boule unité de \mathbb{C}^n [Weighted inequalities for the Bergman projection in the unit ball of \mathbb{C}^n], Studia Math. 71 (1981/82), no. 3, 305–323.
- [4] D. Bekollé and A. Bonami, Inégalités á poids pour le noyau de Bergman, (French) C. R. Acad. Sci. Paris Sér. A-B 286 (1978), no. 18, 775–778.
- [5] B. J. Boudreaux, Equivalent Bergman spaces with inequivalent weights, J. Geom. Anal. 29 (2019), no. 1, 217 - 223.
- [6] M. Dostanić, Unboundedness of the Bergman projections on L^p spaces with exponential weights, Proc. Edinb. Math. Soc. (2) 47 (2004), no. 1, 111–117.
- [7] J. Duoandikoetxea, F. J. MartÃn-Reyes and S. Ombrosi, Calderón weights as Muckenhoupt weights, Indiana Univ. Math. J. 62 (2013), no. 3, 891–910.
- [8] F. Forelli and W. Rudin, Projections on spaces of holomorphic functions in balls, Indiana Univ. Math. J. 24 (1974/75), 593-602.
- [9] L. Hörmander, L^p estimates for (pluri-)subharmonic functions, Math. Scand. (20) (1967), 65–78.
- [10] T. Korhonen, J. A. Peláez and J. Rättyä, Radial averaging operator acting on Bergman and Lebesgue spaces, Forum Math. 31 (2019), no. 4, 1051-1068.
- [11] B. Muckenhoupt, Hardy's inequality with weights, Studia Math.64 (1972), 31–38.
- [12] M. Pavlović, Mixed norm spaces of analytic and harmonic functions. I, Publ. Inst. Math. (Beograd) (N.S.) 40(54) (1986), 117–141.
- [13] J. A. Peláez, Small weighted Bergman spaces, Proceedings of the Summer School in Complex and Harmonic Analysis, and Related Topics, 29–98, Publ. Univ. East. Finl. Rep. Stud. For. Nat. Sci., 22, Univ. East. Finl., Fac. Sci. For., Joensuu, 2016.
- [14] J. A. Peláez and J. Rättyä, Generalized Hilbert operators on weighted Bergman spaces, Adv. Math. 240 (2013), 227–267.

- [15] J. A. Peláez and J. Rättyä, Weighted Bergman spaces induced by rapidly increasing weights, Mem. Amer. Math. Soc. 227 (2014), no. 1066.
- [16] J. A. Peláez and J. Rättyä, Two weight inequality for Bergman projection, J. Math. Pures Appl. (9) 105 (2016), no. 1, 102–130.
- [17] J. A. Peláez and J. Rättyä, On the boundedness of Bergman projection, Advanced courses of mathematical analysis VI, 113–132, World Sci. Publ., Hackensack, NJ, 2017.
- [18] J. A. Peláez and J. Rättyä, Bergman projection induced by radial weight, https://arxiv.org/abs/1902.09837, preprint (submitted).
- [19] J. A. Peláez, J. Rättyä and B. D. Wick, Bergman projection induced by kernel with integral representation, J. Anal. Math. 138 (2019), no. 1, 325–360.
- [20] F. Pérez-González and J. Rättyä, Derivatives of inner functions in weighted Bergman spaces and the Schwarz-Pick lemma, Proc. Amer. Math. Soc. 145 (2017), no. 5, 2155–2166.
- [21] S. Pott and M. C. Reguera, Sharp Bekollé estimate for the Bergman projection, J. Funct. Anal. 265 (2013), 3233–3244.
- [22] D. V. Widder, The Laplace Transform, Princeton Mathematical Series, vol. 6. Princeton University Press, Princeton, N. J., 1941. x+406 pp.
- [23] K. Zhu, Operator Theory in Function Spaces, American Mathematical Society, Providence, RI. 2007.

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, 29071 MÁLAGA, SPAIN

Email address: martin_reyes@uma.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, 29071 MÁLAGA, SPAIN

Email address: portega@uma.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, 29071 MÁLAGA, SPAIN

Email address: japelaez@uma.es

UNIVERSITY OF EASTERN FINLAND, P.O.BOX 111, 80101 JOENSUU, FINLAND *Email address*: jouni.rattya@uef.fi