

Type III Spacetime with Closed Timelike Curves

Faizuddin Ahmed¹

Hindustani Kendriya Vidyalaya, Guwahati-05, Assam, India

**Current Affiliation : National Academy Gauripur, Assam, India,
783331**

Abstract

We present a cyclic symmetric space-time, admitting closed time-like curves (CTCs) which appear after a certain instant of time, *i. e.*, a time-machine space-time. These closed time-like curves evolve from an initial spacelike hypersurface on the plane $z = \text{constant}$ in a causally well-behaved manner. The space-time discussed here is free from curvature singularities and a 4D generalization of the Misner space in curved space-time. The matter field is of pure radiation with a negative Cosmological constant.

1 Introduction

One of the most intriguing aspects of Einstein's theory of gravitation is that solutions of Field Equations admit closed time-like curves (CTC). Presence of CTC in a space-time lead to time-travel which violate the causality condition. The first one being the Gödel's space-time [1] which admit closed time-like curves (CTC) everywhere and an eternal time-machine space-time. There are a considerable number of space-times in literature that admitting closed time-like curves have been constructed. A small sample would be [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21]. One way of classifying such causality violating space-times would be to categorize the metrics as either eternal time-machine in which CTC always exist (in this class would be [1, 2]), or as time-machine space-times in which CTC appear after a

¹faizuddinahmed15@gmail.com ; faiz4U.enter@rediffmail.com

certain instant of time. In the latter category would be the one discussed in [18, 19, 20]. Many of the models, however, suffer from one or more severe drawbacks. For instance, in some of these solutions, for example [13, 14, 20] the weak energy condition (WEC) is violated indicating unrealistic matter-energy content and some other solutions have singularities.

Among the time-machine space-times, we mention two: the first being Ori's compact core [17] which is represented by a vacuum metric locally isometric to pp waves and second, which is more relevant to the present work, the Misner space [22] in 2D. This is essentially a two dimensional metric (hence flat) with peculiar identifications. The Misner space is interesting in the context of CTC as it is a prime example of a space-time where CTC evolve from causally well-behaved initial conditions.

The metric for the Misner space [22]

$$ds_{Misn}^2 = -2 dt dx - t dx^2 \quad . \quad (1)$$

where $-\infty < t < \infty$ but the co-ordinate x periodic. The metric (1) is regular everywhere as $\det g = -1$ including at $t = 0$. The curves $t = t_0$, where t_0 is a constant, are closed since x is periodic. The curves $t < 0$ are spacelike, but $t > 0$ are time-like and the null curves $t = t_0 = 0$ form the chronology horizon. The second type of curves, namely, $t = t_0 > 0$ are closed time-like curves (CTC). This metric has been the subject of intense study and quite recently, D. Levanony *et al* [23] have studied the motion of extended bodies in the 2D Misner space and its flat 4D generalizations. A non-flat 4D space-time, satisfying all the energy conditions, but with causality violating properties of the Misner space, primarily that CTC evolve smoothly from an initially causally well-behaved stage, would be physically more acceptable as a time-machine space-time.

In this paper, we shall attempt to show that causality violating curves appear in non-vacuum space-time with comparatively simple structure. In *section 2*, we analyze the space-time, in *section 3*, the matter distribution and

the energy condition, in *section 4*, the space-time is classified and discussed its kinematical properties and concluding one in *section 5*.

2 Analysis of the space-time

Consider the following metric

$$ds^2 = 4r^2 dr^2 + e^{2\alpha r^2} (dz^2 - t d\phi^2 - 2 dt d\phi) + 4\beta z r e^{-\alpha r^2} dr d\phi, \quad (2)$$

where ϕ coordinate is assumed periodic $0 \leq \phi \leq \phi_0$, with α is an integer and $\beta > 0$ is real number. We have used co-ordinates $x^1 = r$, $x^2 = \phi$, $x^3 = z$ and $x^4 = t$. The ranges of the other co-ordinates are $t, z \in (-\infty, \infty)$ and $0 \leq r < \infty$. The metric has signature $(+, +, +, -)$ and the determinant of the corresponding metric tensor $g_{\mu\nu}$, $\det g = -4r^2 e^{6\alpha r^2}$. The non-zero components of the Einstein tensor are

$$\begin{aligned} G^\mu_\mu &= 3\alpha^2 \\ G^t_\phi &= -\frac{1}{2} e^{-6\alpha r^2} \beta^2 \end{aligned} \quad (3)$$

Consider an azimuthal curves γ defined by $r = r_0$, $z = z_0$ and $t = t_0$, where r_0, z_0, t_0 are constants, then we have from the metric (2)

$$ds^2 = -t e^{2\alpha r^2} d\phi^2 \quad (4)$$

These curves are null for $t = 0$, spacelike throughout for $t = t_0 < 0$, but become time-like for $t = t_0 > 0$, which indicates the presence of closed timelike curves (CTC). Hence CTC form at a definite instant of time satisfy $t = t_0 > 0$.

It is crucial to have analysis that the above CTC evolve from an spacelike $t = \text{constant}$ hypersurface (and thus t is a time coordinate)[17]. This can be ascertained by calculating the norm of the vector $\nabla_\mu t$ (or by determining the sign of the component g^{tt} in the inverse metric tensor $g^{\mu\nu}$ [17]). We find from (2) that

$$g^{tt} = t e^{-2\alpha r^2} + \beta^2 z^2 e^{-6\alpha r^2} \quad (5)$$

A hypersurface $t = \text{constant}$ is spacelike provided $g^{tt} < 0$ for $t = t_0 < 0$, but become time-like provided $g^{tt} > 0$ for $t = t_0 > 0$. Here we choose the $z - \text{planes}$ defined by $z = z_0$, (z_0 , a constant equal to zero) such that the above condition is satisfied. Thus the spacelike $t = \text{constant} < 0$ hypersurface can be chosen as initial conditions over which the initial may be specified. There is a Cauchy horizon for $t = t_0 = 0$ called Chronology horizon which separates the causal and non-causal of the space-time. Hence the space-time evolves from a partial Cauchy hypersurface (initial spacelike hypersurface) in a causally well-behaved manner, upto a moment, i.e., a null hypersurface $t = 0$ and CTC form at a definite instant of time on $z = \text{constant}$ plane.

Consider the Killing vector $\eta = \partial_\phi$ for metric (2) which has the normal form

$$\eta^\mu = (0, 1, 0, 0) \quad (6)$$

Its co-vector is

$$\eta_\mu = \left(2\beta z r e^{-\alpha r^2}, -t e^{2\alpha r^2}, 0, -e^{2\alpha r^2} \right) \quad (7)$$

(6) satisfies the Killing equation $\eta_{\mu;\nu} + \eta_{\nu;\mu} = 0$. For cyclicly symmetric metric, the norm $\eta_\mu \eta^\mu$ of the Killing vector is spacelike, closed orbits [24, 25, 26, 27, 28]. We note that

$$\eta^\mu \eta_\mu = -t e^{2\alpha r^2} \quad (8)$$

which is spacelike for $t < 0$, closed orbits (ϕ co-ordinate being periodic).

An important note is that the Riemann tensor $R_{\mu\nu\rho\sigma}$ can be expressed in terms of metric tensor $g_{\mu\nu}$ as

$$R_{\mu\nu\rho\sigma} = k (g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho}) \quad (9)$$

where $k = -\alpha^2$ for the space-time (2).

Another important note is that if we take $\beta = 0$, then the space-time represented by (2) is maximally symmetric vacuum space-time and locally isometric anti-de Sitter space in four-dimension. One can easily show by

a number of transformations the standard form of locally isometric AdS_4 metric [29]

$$ds^2 = \frac{3}{(-\Lambda) x^2} (-dt^2 + dx^2 + d\phi^2 + dz^2) \quad (10)$$

where one of the co-ordinate ϕ being periodic.

3 Matter distribution of the space-time and the energy condition

The Einstein's Field Equations taking into account the cosmological constant

$$G^{\mu\nu} + \Lambda g^{\mu\nu} = T^{\mu\nu} \quad , \quad \mu, \nu = 1, 2, 3, 4 \quad (11)$$

Consider the energy-momentum tensor that of pure radiation field [30]

$$T^{\mu\nu} = \rho n^\mu n^\nu \quad (12)$$

where n^μ is the null vector defined by

$$n^\mu = (0, 0, 0, 1) \quad (13)$$

The non-zero component of the energy-momentum tensor

$$T_\phi^t = -\rho e^{2\alpha r^2} \quad (14)$$

Equating Field Equations (11) using (3) and (14) we get

$$\Lambda = -3\alpha^2 \quad (15)$$

$$\rho = \frac{1}{2} \beta^2 e^{-8\alpha r^2} \quad , \quad 0 \leq r < \infty \quad (16)$$

The energy-density of pure radiation or null dust decreases exponentially with r and vanish at $r \rightarrow \pm\infty$. The matter field pure radiation satisfy the energy condition and the energy density ρ is always positive.

4 Classification and kinematical properties of the space-time

For classification of the spacetime (2), we can construct the following set of null tetrads (k, l, m, \bar{m}) as

$$k_\mu = (0, 1, 0, 0) \quad , \quad (17)$$

$$l_\mu = \left(-2\beta z r e^{-\alpha r^2}, \frac{t}{2} e^{2\alpha r^2}, 0, e^{2\alpha r^2} \right) \quad , \quad (18)$$

$$m_\mu = \frac{1}{\sqrt{2}} (2r, 0, i e^{\alpha r^2}, 0) \quad , \quad (19)$$

$$\bar{m}_\mu = \frac{1}{\sqrt{2}} (2r, 0, -i e^{\alpha r^2}, 0) \quad , \quad (20)$$

where $i = \sqrt{-1}$. The set of null tetrads above are such that the metric tensor for the line element (2) can be expressed as

$$g_{\mu\nu} = -k_\mu l_\nu - l_\mu k_\nu + m_\mu \bar{m}_\nu + \bar{m}_\mu m_\nu \quad . \quad (21)$$

The vectors (17)–(20) are null vectors and are orthogonal except for $k_\mu l^\mu = -1$ and $m_\mu \bar{m}^\mu = 1$. Using this null tetrad above we have calculated the five Weyl scalars

$$\begin{aligned} \Psi_3 &= -\frac{i\alpha\beta e^{-2\alpha r^2}}{2\sqrt{2}} \\ \Psi_4 &= -\frac{1}{4}\beta e^{-2\alpha r^2} \left(i + 2\alpha z e^{\alpha r^2} \right) \end{aligned} \quad (22)$$

are non-vanishing, while $\Psi_0 = \Psi_1 = \Psi_2 = 0$. The spacetime represented by (2) is of type III in the Petrov classification scheme. Note that the non-zero Weyl scalars Ψ_3 and Ψ_4 are finite at $r \rightarrow 0$ and vanish as $r \rightarrow \pm\infty$ indicating asymptotic flatness of the spacetime (2). The metric (2) is free from curvature singularities. The curvature invariant known as Kretschmann scalar is given by

$$R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} = 24\alpha^4 \quad (23)$$

and the curvature scalar

$$R = -12\alpha^2 \quad (24)$$

are constant being non-zero.

Using the null tetrad (17) we have calculated the *Optical* scalars [30] the *expansion*, the *twist* and the *shear* and they are

$$\begin{aligned} \Theta &= \frac{1}{2} k^\mu_{;\mu} = 0 \quad , \\ \omega^2 &= \frac{1}{2} k_{[\mu;\nu]} k^{\mu;\nu} = 0 \quad , \\ \sigma \bar{\sigma} &= \frac{1}{2} k_{(\mu;\nu)} k^{\mu;\nu} - \Theta^2 = 0 \end{aligned} \quad (25)$$

and the null vector (17) satisfy the geodesics equation

$$k_{\mu;\nu} k^\nu = 0 \quad . \quad (26)$$

Thus the spacetime represented by (2) is non-diverging, have shear-free null geodesics congruence. One can easily show that for constant r and z , the metric (2) reduces to conformal Misner space in 2D

$$ds^2_{conf} = \Omega ds^2_{Misn} \quad (27)$$

where $\Omega = e^{-2\alpha r^2}$ is a constant.

5 Conclusion

Our primary motivation in this paper is to write down a metric for a spacetime that incorporates the Misner space and its causality violating properties and to classify it. The solution presented here is non-vacuum, cyclicly symmetric metric (2) and serves as a model of time-machine spacetime in the sense that CTC appear at a definite instant of time on the $z - plane$. Most of the CTC spacetimes violate one or more energy conditions or unrealistic matter source and are unphysical. The model discussed here is free from all these problems and matter distribution is of pure radiation field with negative cosmological constant satisfying the energy condition.

Appendix

The above metric (2) can be transform to a simple form as follows. Let us perform the following transformation

$$r \rightarrow \sqrt{r'}, \quad (28)$$

into the metric (2), one will get

$$ds^2 = dr'^2 + e^{2\alpha r'} (dz^2 - t d\phi^2 - 2 dt d\phi) + 2\beta z e^{-\alpha r'} dr' d\phi. \quad (29)$$

Finally doing another transformation

$$r' \rightarrow \frac{1}{\alpha} \ln \varrho \quad (30)$$

into the metric (29), we get the following line element

$$ds^2 = \frac{d\varrho^2}{\alpha^2 \varrho^2} + \varrho^2 dz^2 + \varrho^2 (-t d\phi^2 - 2 dt d\phi) + \frac{2\beta z}{\alpha \varrho^2} d\varrho d\phi. \quad (31)$$

Or if one does the following transformation

$$r' \rightarrow \frac{1}{\alpha} \ln(\alpha \varrho) \quad (32)$$

into the metric (29), we get the following line element

$$ds^2 = \frac{d\varrho^2}{\alpha^2 \varrho^2} + \alpha^2 \varrho^2 dz^2 + \alpha^2 \varrho^2 (-t d\phi^2 - 2 dt d\phi) + \frac{2\beta z}{\alpha^2 \varrho^2} d\varrho d\phi. \quad (33)$$

Or if one does the following transformation

$$r' \rightarrow -\frac{1}{\alpha} \ln(\alpha \varrho) \quad (34)$$

into the metric (29), we get the following line element

$$ds^2 = \frac{1}{\alpha^2 \varrho^2} (d\varrho^2 - t d\phi^2 - 2 dt d\phi + dz^2) - 2\beta z d\varrho d\phi. \quad (35)$$

References

- [1] K. Gödel, Rev. Mod. Phys. **21**, 447 (1949).

- [2] W. J. van Stockum, Proc. R. Soc. Edin. **57**, 135 (1937).
- [3] F. J. Tipler, Phys. Rev. **D 9**, 2203 (1974).
- [4] J. R. Gott, Phys. Rev. Lett. **66**, 1126 (1991).
- [5] W. B. Bonnor, Class. Quantum Grav. **19**, 5951 (2002).
- [6] W. B. Bonnor, Int. J. Mod. Phys. **D 12**, 1705 (2003).
- [7] W. B. Bonnor, Gen. Relativ. Grav. **37**, 1833 (2005).
- [8] R. P. Kerr, Phys. Rev. Lett. **11**, 237 (1963).
- [9] B. Carter, Phys. Rev. **D 174**, 1559 (1968).
- [10] M. Gürses, A. Karasu and Ö. Sarioğlu, Class. Quantum Grav. **22**, 1527 (2005).
- [11] M. Alcubierre, Class. Quantum Grav. **11**, L73 (1994).
- [12] F. S. N. Lobo and P. Crawford, NATO Science Series II **volume 95**, 289 (2003).
- [13] M. S. Morris, K. S. Thorne and U. Yurtsever, Phys. Rev. Lett. **61**, 1446 (1988).
- [14] M. S. Morris and K. S. Thorne, Amer. J. Phys. **56**, 395 (1988).
- [15] S. V. Krasnikov, Phys. Rev. **D 57**, 4760 (1998).
- [16] A. E. Everett and T. A. Roman, Phys. Rev. **D 56**, 2100 (1997).
- [17] A. Ori, Phys. Rev. Lett. **95**, 021101 (2005).
- [18] A. Ori, Phys. Rev. **D 76**, 044002 (2007).
- [19] A. Ori, Phys. Rev. **D 54**, 4858 (1996).

- [20] A. Ori, Phys. Rev. **D 49**, 3990 (1994).
- [21] D. Sarma, M. Patgiri and F. U. Ahmed, Gen. Relativ. Grav. **46**, 1633 (2014).
- [22] C. W. Misner, *Relativity Theory and Astrophysics I : Relativity and Cosmology* J. Ehlers (Ed.), Lecture in Applied Mathematics **8**, American Mathematical Society (1967).
- [23] D. Levanony and A. Ori, Phys. Rev. **D 83**, 044043 (2011).
- [24] M. Mars and J. M. M. Senovilla, Class. Quantum Grav. **10**, 1633 (1993).
- [25] B. Carter, Commun. Math. Phys. **17**, 233 (1970).
- [26] A. Branes, Class. Quantum Grav. **18**, 5511 (2001).
- [27] J. Bičák and B. D. Schmidt, J. Math. Phys. **25**, 600 (1984).
- [28] R. M. Wald, *General Relativity*, Cambridge University Press, Cambridge (1984).
- [29] M. Zofka and J. Bičák, Class. Quantum Grav. **25**, 015011 (2008).
- [30] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Herlt, *Exact Solutions to Einstein's Field Equations*, Cambridge University Press, Cambridge (2005).