

Classical string correspondence in 2 and 3 spacetime dimensions

G. Aminov^{*a,b,c}

^aC. N. Yang Institute for Theoretical Physics, State University of New York, Stony Brook, NY 11794-3840, USA

^bSimons Center for Geometry and Physics, State University of New York, Stony Brook, NY 11794-3636, USA

^cITEP NRC KI, Moscow 117218, Russia

Abstract

We present an example of the string correspondence: a relation between two classical string solutions - one in Minkowski spacetime and another one in the AdS spacetime. The first solution describes a 2d motion of the string with massive ends and was derived in [1]. The second solution is the accelerating string solution in the AdS₃ spacetime and describes a heavy quark-antiquark pair [2].

1 Introduction

A notion of duality plays a prominent role in the modern theoretical physics. As the number of theoretical models grows larger and larger, one finds the need to connect them, since they sometimes describe the same physical phenomena. Some of the most famous connections are the AdS/CFT correspondence [3, 4, 5], different string dualities [6, 7, 8, 9, 10, 11, 12] and the AGT correspondence [13, 14]. In this paper we want to describe an example of the correspondence, that has something to do with the first two. From the pure geometrical point of view our example looks like a relation between an open string with massive ends in 2d Minkowski spacetime and a closed string in 3d AdS spacetime (with some specific identifications on the boundary of AdS₃). The time and space coordinates on the string worldsheet are flipped between the two solutions and thus the relation closely resembles T-duality. On the other hand, by cutting the closed string in half on the AdS side one finds a motion of the quark-antiquark pair as proposed by the AdS/CFT correspondence. Parallels with the AdS/CFT correspondence go deeper due to the fact, that the 2d solution exhibits some form of the $SL(2, R)$ symmetry. This could potentially allow one to define a boundary CFT and thus leads to a manifestation of AdS/CFT called AdS/BCFT [15, 16]. Another interesting direction would be the relation between 2d strings and matrix models [17, 18, 19]. In particular, string solutions called short strings and long strings were obtained in [20, 21, 22] as classical solutions of the $SL(2, R)$ WZW model [23, 24, 25, 26, 27]. These string solutions were discussed in different contexts [28, 29, 30] and they are possibly related to the classical solution [1].

The main goal of this paper is to connect two different classical solutions [1] and [2], which have been extensively studied throughout the years. Despite similar effects taking place in either setting, the direct relation has not been established before. The most recent solution [2] was motivated by the study of an energy loss of a heavy quark moving through a hot plasma [31, 32, 33, 34, 35, 36, 37]. It describes a heavy quark-antiquark pair moving with constant acceleration and connected by the string embedded in the AdS₃ space. The string worldsheet develops an event horizon and the ensuing physical implications were studied in a number of subsequent works [38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48]. The first string solution [1] was derived 45 years ago and describes a motion of the string with massive ends in 2d Minkowski spacetime. A particular example of this solution is called a yo-yo string and it corresponds to the case of vanishing masses at the ends of the string or, equivalently, to the limit of strong tension:

$$\text{yo-yo limit: } m \rightarrow 0 \quad \text{or} \quad \gamma \rightarrow \infty. \quad (1.1)$$

In [49] the motivation to study yo-yo strings was the energy loss of gluons and light quarks [50, 51, 52, 53, 54, 55]. The yo-yo string was discussed later in the context of the AdS/BCFT correspondence in [56], but the relation to the AdS solution [2] was not noticed. This is probably due to the fact that the relation appears in a slightly more complicated limit than just (1.1). Recently the general solution [1] was studied under the name of the

*gleb.aminov@stonybrook.edu

periodic zigzag solution [57, 58, 59]. It was observed that the zigzag solution exhibits some properties of a quantum black hole [57] and in the present paper we are going to provide further classical evidence to support this claim.

The paper is organized as follows. In sec. 2 we review the original solution [1] for the string with massive ends. Motivated by [60] in sec. 3 we rewrite the same solution in the conformal gauge, where the string equations of motion look like simple wave equations. Even though we are working with 2 spacetime dimensions, after choosing the conformal gauge the solution is unique due the boundary conditions. This rewriting of the known solution proves to be very insightful. First, we discover a form of $SL(2, R)$ symmetry (3.20), which governs the periodicity properties of the full solution (without taking the yo-yo limit). Second, the change of coordinates on the worldsheet of the string is reminiscent of the coordinate change near the horizon. The latter statement can be unfolded as follows. The solution in the timelike gauge is defined on the infinite set of finite time intervals of the form $[(2n-1)\tau_0, (2n+1)\tau_0]$, $n \in \mathbb{Z}$. At the beginning and at the end of each interval the string is collapsed to a point. When we go to the conformal gauge, each one of these finite time intervals is mapped to the infinite time interval in the corresponding coordinate system and thus the folding points are removed to infinity.

In sec. 4 this black hole like behavior is further explained by introducing two new coordinate systems on the worldsheet related to the accelerated and stationary observers. In the first coordinate system the time " η " is the proper time of the accelerated observer sitting at the end of the string. One could think about these coordinates as a string analog of Rindler coordinates. Indeed, if we take the limit of a large string length and zoom in on one of the endpoints, we would recover the original Rindler coordinates (see [61] for the details). In the second coordinate system the time " t " is the proper time of the stationary observer sitting in the middle of the string. In both coordinate systems the horizon is seen in the induced metric on the worldsheet of the string. In coordinates (t, σ) the induced metric is

$$ds_{ws}^2 = -\frac{dt^2}{(1-t^2/t_0^2)^2} + \alpha^2 t_0^2 d\sigma^2, \quad -t_0 \leq t \leq t_0, \quad (1.2)$$

and the horizon $t = t_0 \equiv \tau_0$ is situated at the folding points. In these coordinates the solution itself is somewhat reminiscent of the short and long string solutions of the $SL(2, R)$ WZW model [62, 63, 20]. From this point of view the existence of black holes on the worldsheets of strings was discovered long time ago [64, 65].

In sec. 5 we establish the relation between two classical solutions [1] and [2]. The main idea is to identify the worldsheet coordinates from both solutions based on the form of the induced metrics. Indeed, the induced metric in the second solution [2]

$$ds_{ws}^2 = -\left(1 - \frac{r^2}{b^2}\right) d\theta^2 + \frac{dr^2}{1 - r^2/b^2} \quad (1.3)$$

is equivalent to (1.2) upon a Weyl transformation and the following identification of coordinates:

$$t = r, \quad \sigma = \frac{\theta}{\alpha b}, \quad (1.4)$$

where we should also assume $b = t_0$. However, the time coordinate θ runs from $-\infty$ to $+\infty$, whereas σ only belongs to the interval $[0, \pi]$. This discrepancy is resolved by taking the limit $\alpha \rightarrow +\infty$ and restricting to the positive time interval $\theta \geq 0$. As it was briefly mentioned earlier, this limit is a bit more complicated than the yo-yo limit (1.1) and can be written in terms of worldsheet coordinates (t, σ) and spacetime Minkowski coordinates (X^0, X^1) as

$$\sigma = \frac{\theta}{\alpha b}, \quad X^0 = \frac{2m}{\gamma} Z_+^0, \quad X^1 - x_0 = \frac{2m}{\gamma} Z_+^1, \quad \frac{\gamma}{m} \rightarrow +\infty, \quad (1.5)$$

where

$$\sinh(\alpha \pi) = \frac{\gamma}{m} t_0 \quad (1.6)$$

and $(-2x_0)$ is the maximum length of the string [1] in the Minkowski spacetime. The end result is that both classical solutions describe the embedding of the same string worldsheet in two different spacetimes: one is 2d Minkowski spacetime and another one is 3d AdS spacetime. The transformation between coordinates in Minkowski and coordinates on the boundary of AdS will be written in the form of the special conformal transformation similar to the ones studied earlier in [66].

2 Periodic zigzag solution in the timelike gauge

Following [1], we consider a pair of point masses joined by a massless string:

$$S = \int_{\tau_1}^{\tau_2} d\tau \left(-m \sqrt{-(\partial_\tau X(\tau, 0))^2} - m \sqrt{-(\partial_\tau X(\tau, \pi))^2} - \gamma \int_0^\pi d\sigma \sqrt{-G} \right). \quad (2.1)$$

The string is described by the usual Nambu-Goto action [67, 68] with

$$G = (\partial_\tau X)^2 (\partial_\sigma X)^2 - (\partial_\tau X^\mu \partial_\sigma X_\mu)^2 \quad (2.2)$$

and a D -component field $X^\mu(\tau, \sigma)$. There are two types of equations of motion: string equations at $0 < \sigma < \pi$ and boundary equations at $\sigma = 0, \pi$. The conserved total momentum is a sum of point masses momenta $p^\mu(\tau, 0)$, $p^\mu(\tau, \pi)$ and the integral of the string momentum density $\gamma K^\mu(\tau, \sigma)$. Definitions we are going to use are:

$$p^\mu(\tau, \sigma) = \frac{m \partial_\tau X^\mu}{\sqrt{-(\partial_\tau X)^2}}, \quad K^\mu(\tau, \sigma) = \frac{1}{\sqrt{-G}} \left(\partial_\tau X^\mu (\partial_\sigma X)^2 - \partial_\sigma X^\mu (\partial_\tau X^\nu \partial_\sigma X_\nu) \right), \quad (2.3)$$

$$N^\mu(\tau, \sigma) = \frac{1}{\sqrt{-G}} \left(\partial_\sigma X^\mu (\partial_\tau X)^2 - \partial_\tau X^\mu (\partial_\tau X^\nu \partial_\sigma X_\nu) \right). \quad (2.4)$$

Then classical equations of motion can be written as

$$0 < \sigma < \pi : \quad \partial_\tau K^\mu + \partial_\sigma N^\mu = 0, \quad (2.5)$$

$$\partial_\tau p^\mu(\tau, 0) + \gamma N^\mu(\tau, 0) = 0, \quad (2.6)$$

$$\partial_\tau p^\mu(\tau, \pi) - \gamma N^\mu(\tau, \pi) = 0, \quad (2.7)$$

and the total momentum is

$$P^\mu = p^\mu(0) + p^\mu(\pi) + \gamma \int_0^\pi d\sigma K^\mu. \quad (2.8)$$

Before solving the equations of motion, we need to fix some gauge choices. First, we choose the timelike gauge $X^0 = \tau$. Second, we can use the σ reparametrization invariance to make $\partial_\sigma X^1$ independent of σ in the interval $(0, \pi)$ [1]:

$$X^1(\tau, \sigma) = \frac{\sigma}{\pi} (X^1(\tau, \pi) - X^1(\tau, 0)) + X^1(\tau, 0). \quad (2.9)$$

The periodic zigzag solution describes a 2d motion, so that only coordinates X^0 and X^1 are involved. In that case the Hamiltonian is particularly simple:

$$H^{2d} = \sqrt{p(0)^2 + m^2} + \sqrt{p(\pi)^2 + m^2} + \gamma |X^1(\pi) - X^1(0)|. \quad (2.10)$$

In the rest frame we can choose $X^1(\tau, 0) = -X^1(\tau, \pi)$ and the solution derived in [1] is

$$X^1(\tau, \sigma) = \left(1 - \frac{2\sigma}{\pi} \right) X^1(\tau, 0) \quad (2.11)$$

with the motion of the endpoints described by

$$X^1(\tau, 0) = \begin{cases} x_0 - \frac{m}{\gamma} + \sqrt{\frac{m^2}{\gamma^2} + \tau^2}, & \tau \in (-\tau_0, \tau_0), \\ -x_0 + \frac{m}{\gamma} - \sqrt{\frac{m^2}{\gamma^2} + (\tau - 2\tau_0)^2}, & \tau \in (\tau_0, 3\tau_0), \end{cases} \quad (2.12)$$

where we introduced $x_0 < 0$ and

$$\tau_0^2 \equiv x_0^2 - \frac{2m}{\gamma} x_0. \quad (2.13)$$

The movement is periodic with period $T = 4\tau_0$. In what follows we are going to be mostly interested in the time interval $(-\tau_0, \tau_0)$. At $\tau = 0$ the endpoints are at maximum separation of $2x_0$ and start moving towards each other (see Fig. 1).




Figure 1: Periodic zigzag solution, $-\tau_0 \leq \tau \leq 3\tau_0$

3 Periodic zigzag solution in the conformal gauge

The goal of this section is to rewrite the 2d solution in another gauge, where the string equations of motion become simple wave equations. In order to achieve this, the following constraints should be imposed [60]:

$$(\partial_\tau X \pm \partial_\sigma X)^2 = 0, \quad (\partial_\tau^2 X \pm \partial_\tau \partial_\sigma X)^2 = \text{const.} \quad (3.1)$$

The first constraint leads to the simplified expressions for G , K^μ and N^μ :

$$G = -(\partial_\tau X^\mu \partial_\tau X_\mu)^2, \quad K^\mu = -\partial_\tau X^\mu \frac{(\partial_\tau X)^2}{|(\partial_\tau X)^2|}, \quad N^\mu = \partial_\sigma X^\mu \frac{(\partial_\tau X)^2}{|(\partial_\tau X)^2|}. \quad (3.2)$$

Taking into the account that $(\partial_\tau X)^2 < 0$, we can rewrite the equations of motion in the form

$$0 < \sigma < \pi: \quad \partial_\tau^2 X^\mu - \partial_\sigma^2 X^\mu = 0, \quad (3.3)$$

$$\partial_\tau p^\mu(\tau, 0) = \gamma \partial_\sigma X^\mu(\tau, 0), \quad (3.4)$$

$$\partial_\tau p^\mu(\tau, \pi) = -\gamma \partial_\sigma X^\mu(\tau, \pi). \quad (3.5)$$

Using the notations introduced in [60], we can solve the boundary equation at $\sigma = 0$ with the help of a new function $\xi(\tau)$ and a vector $e^\mu(\tau)$:

$$\partial_\tau X^\mu|_{\sigma=0} = \xi(\tau) e^\mu(\tau), \quad \partial_\sigma X^\mu|_{\sigma=0} = \frac{m}{\gamma} \dot{e}^\mu(\tau) \text{sign}(\xi), \quad (3.6)$$

where $e^\mu e_\mu = -1$ and $\xi^2 = m^2 (\dot{e})^2 / \gamma^2$. For the 2d solution we can choose e^μ and ξ to be

$$e^\mu(\tau) = \left(\sqrt{1 + f(\tau)^2}, f(\tau) \right), \quad \xi(\tau) = \frac{m}{\gamma} \frac{\dot{f}(\tau)}{\sqrt{1 + f(\tau)^2}}. \quad (3.7)$$

With this choice the second constraint from (3.1) is satisfied and the constant is exactly zero. Therefore, the second constraint does not provide any new information in the 2d case.

The remaining problem is to find such function $f(\tau)$, that the boundary equation at $\sigma = \pi$ is satisfied. This could be done by demanding that the derivatives of X^μ at $\sigma = \pi$ satisfy similar equations to the ones in (3.6), except one of the equations should have an additional minus sign on the right hand side and the new functions $\tilde{\xi}$ and \tilde{e}^μ should be used. This results in a peculiar behaviour of a function $f(\tau)$ under the 2π -shifts of the argument:

$$f(\tau + \pi) + \sqrt{1 + f(\tau + \pi)^2} = g(\tau + \pi) - c_1, \quad (3.8)$$

$$f(\tau - \pi) - \sqrt{1 + f(\tau - \pi)^2} = -\frac{1}{g(\tau + \pi)} - c_2, \quad (3.9)$$

where we introduced two new constants $c_{1,2}$ and a new function $g(\tau)$ with the following properties:

$$c_1 < 0, \quad c_2 > 0, \quad c_1 < g(\tau) < 0, \quad g'(\tau) > 0. \quad (3.10)$$

Before we move on with the analysis, let's discuss the change of variables, which occurs when we go from the timelike gauge of the previous section to the conformal gauge in the current one. If we rename the coordinates on the string worldsheet in the timelike gauge as $(\tilde{\tau}, \tilde{\sigma})$, then the change of coordinates reads

$$X^0 = \tilde{\tau} = \omega^0 + \frac{m}{2\gamma} \left(f + \sqrt{1 + f^2} \right) \Big|_{(\tau+\sigma)} + \frac{m}{2\gamma} \left(f - \sqrt{1 + f^2} \right) \Big|_{(\tau-\sigma)}, \quad (3.11)$$

$$X^1 = \left(1 - \frac{2\tilde{\sigma}}{\pi} \right) \left(x_0 - \frac{m}{\gamma} + \sqrt{\frac{m^2}{\gamma^2} + \tilde{\tau}^2} \right) = \omega^1 + \frac{m}{2\gamma} \left(f + \sqrt{1 + f^2} \right) \Big|_{(\tau+\sigma)} - \frac{m}{2\gamma} \left(f - \sqrt{1 + f^2} \right) \Big|_{(\tau-\sigma)}, \quad (3.12)$$

where we explicitly used the solution for the time interval of $-\tau_0 \leq \tilde{\tau} \leq \tau_0$. Now, the string action is invariant under the reparametrizations of $\tilde{\tau} = \tilde{\tau}(\tau, \sigma)$ and $\tilde{\sigma} = \tilde{\sigma}(\tau, \sigma)$, but the endpoints of the string demand that the reparametrizations of $\tilde{\sigma}$ obey

$$\tilde{\sigma}(\tau, 0) = 0, \quad \tilde{\sigma}(\tau, \pi) = \pi. \quad (3.13)$$

The latter provides us with the consistency check on the behaviour of the function $f(\tau)$ as well as the relations between the various constants. At $\sigma = 0$ we have

$$\tilde{\tau}(\tau, 0) = \omega^0 + \frac{m}{\gamma} f(\tau), \quad (3.14)$$

$$\left(1 - \frac{2\tilde{\sigma}(\tau, 0)}{\pi} \right) \left(x_0 - \frac{m}{\gamma} + \sqrt{\frac{m^2}{\gamma^2} + \tilde{\tau}(\tau, 0)^2} \right) = \omega^1 + \frac{m}{\gamma} \sqrt{1 + f(\tau)^2}. \quad (3.15)$$

Thus, the first condition in (3.13) is satisfied if

$$\omega^0 = 0, \quad \omega^1 = x_0 - \frac{m}{\gamma}. \quad (3.16)$$

And for the time intervals different from $(-\tau_0, \tau_0)$ the ω^0 constant would be non-zero. Next, we are going to look at $\sigma = \pi$ and use the equations (3.8)–(3.9):

$$\tilde{\tau}(\tau, \pi) = \frac{m}{2\gamma} \left(g(\tau + \pi) - c_1 - \frac{1}{g(\tau + \pi)} - c_2 \right), \quad (3.17)$$

$$\left(1 - \frac{2\tilde{\sigma}(\tau, \pi)}{\pi} \right) \left(x_0 - \frac{m}{\gamma} + \sqrt{\frac{m^2}{\gamma^2} + \tilde{\tau}(\tau, \pi)^2} \right) = \omega^1 + \frac{m}{2\gamma} \left(g(\tau + \pi) - c_1 + \frac{1}{g(\tau + \pi)} + c_2 \right). \quad (3.18)$$

We conclude that to satisfy the second condition in (3.13) the following relations should hold:

$$c_2 = -c_1, \quad c_1 = \frac{2\gamma}{m} \omega^1 = 2 \left(\frac{\gamma}{m} x_0 - 1 \right). \quad (3.19)$$

This leaves us with the task of determining the function $g(\tau)$. First, we want to rewrite (3.8)–(3.9) in terms of the g -function alone:

$$g(\tau + 2\pi) = \frac{g(\tau) - c_1}{1 + c_1 c_2 - c_2 g(\tau)}. \quad (3.20)$$

As one can see, $g(\tau)$ and $g(\tau + 2\pi)$ are connected by the $SL(2, R)$ transformation. To find a suitable candidate for the solution, it is useful to take a step back and think about the equations of motion. In the timelike gauge we had a sign function involved in the boundary equations, which made the time derivatives of the point particles momenta to be discontinuous. In the conformal gauge, on the other hand, we do not see any discontinuities neither in the boundary equations (3.4)–(3.5) nor in the string equations of motion (3.3). Moreover, if we look at the change of variables (3.11)–(3.12), we see that the same points of discontinuity $\tilde{\tau} = \pm \tau_0$ make impossible to determine $\tilde{\sigma}(\pm \tau_0, \sigma)$ as a function of σ . These mismatches between the two coordinate systems on the string

worldsheet could be resolved, if we map the finite time interval $(-\tau_0, \tau_0)$ in the timelike gauge to the infinite one in the conformal gauge. This can also be seen in (3.10), where the g -function is described as being bounded from above and below. A good example of a real-valued bounded function, which also has a positive derivative everywhere, is a hyperbolic tangent. Now we are ready to solve (3.20). By implementing the addition formula for the hyperbolic tangent

$$\tanh(\alpha + \beta) = \frac{\tanh(\alpha) + \tanh(\beta)}{1 + \tanh(\alpha)\tanh(\beta)} \quad (3.21)$$

and assuming the general form of the g -function to be $g(\tau) = a \tanh(\alpha \tau) + b$, we can solve for the constants a , α and b to find the solution:

$$g(\tau) = \frac{1}{2} \left(\sqrt{c_1^2 - 4} \tanh(\alpha \tau) + c_1 \right), \quad \tanh(\alpha \pi) = \sqrt{1 - \frac{4}{c_1^2}}. \quad (3.22)$$

Here we would like to emphasize, that we were simply solving the equation (3.20) and not using any information about the map between the time intervals in different gauges. So, the last consistency check is to verify that the time interval $-\tau_0 \leq \tilde{\tau} \leq \tau_0$ in the timelike gauge is mapped to the infinite time interval in the conformal gauge at any value of σ on the string. Using the definition of $\tilde{\tau}(\tau, \sigma)$ from (3.11) and taking the limits $\tau \rightarrow \pm\infty$, we get

$$\tilde{\tau}(\pm\infty, \sigma) = \frac{m}{\gamma} f(\pm\infty). \quad (3.23)$$

We can already see, that the dependence on σ disappears in these limits. Rewriting the function $f(\tau)$ in terms of the g -function

$$f(\tau) = \frac{1}{2} \left(g(\tau) - c_1 - \frac{1}{g(\tau) - c_1} \right) \quad (3.24)$$

and calculating the limits for $g(\tau)$

$$g(\pm\infty) = \frac{1}{2} \left(c_1 \pm \sqrt{c_1^2 - 4} \right), \quad (3.25)$$

we arrive at the positive conclusion

$$\tilde{\tau}(\pm\infty, \sigma) = \pm \frac{m}{2\gamma} \sqrt{c_1^2 - 4} = \pm\tau_0. \quad (3.26)$$

To summarize, we recall that the 2d solution in the timelike gauge is defined on the infinite set of the finite time intervals of the form $[(2n-1)\tau_0, (2n+1)\tau_0]$, $n \in \mathbb{Z}$. When we go to the conformal gauge, each one of these finite time intervals is mapped to the infinite time interval in the corresponding coordinate system. In other words, each finite patch of worldsheet coordinates in the timelike gauge, described by

$$(2n-1)\tau_0 \leq \tilde{\tau} \leq (2n+1)\tau_0 \quad \text{and} \quad 0 \leq \tilde{\sigma} \leq \pi, \quad (3.27)$$

is mapped to the corresponding infinite patch in the conformal gauge:

$$-\infty \leq \tau_{(n)} \leq +\infty \quad \text{and} \quad 0 \leq \sigma \leq \pi. \quad (3.28)$$

This kind of behavior is indeed reminiscent of the black hole solution and the coordinate change near the horizon [57].

4 Polyakov action and the worldsheet metric

In this section we are going to introduce two new coordinate systems on the worldsheet of the string related to the accelerated and stationary observers. This will help us further understand the black hole like properties of the periodic zigzag solution. In the first coordinate system the time " η " is the proper time of the accelerated observer sitting at the end of the string. One could think about these coordinates as a string analog of Rindler coordinates. In the second coordinate system the time " t " is the proper time of the stationary observer sitting in the middle of the string. In both coordinate systems the horizon is seen in the induced metric on the worldsheet.

4.1 Metric for the accelerated observer

In this subsection we want to derive an analog of Rindler coordinates. First, we rewrite the string action in terms of the worldsheet metric $h_{\alpha\beta}$:

$$S_{\text{string}} = -\frac{\gamma}{2} \int d\tau d\sigma \sqrt{-h} h^{\alpha\beta} (\partial_\alpha X^\mu \partial_\beta X_\mu). \quad (4.1)$$

In general, string equations of motion then take the following form:

$$\frac{1}{\sqrt{-h}} \partial_\alpha \left(\sqrt{-h} h^{\alpha\beta} \partial_\beta X^\mu \right) = 0. \quad (4.2)$$

Since the Polyakov action is Weyl invariant, we conclude that the metric in coordinates (τ, σ) is flat:

$$ds^2 = h_{\alpha\beta} d\sigma^\alpha d\sigma^\beta = -d\tau^2 + d\sigma^2. \quad (4.3)$$

However, in the limit $\tau \rightarrow \pm\infty$ the metric exhibits some $\frac{0}{0}$ ambiguity, as seen from (3.2) and the fact that $(\partial_\tau X)^2 \rightarrow 0$ in this limit.

In order to find the string analog of Rindler coordinates, we introduce the following gauge fixing in terms of coordinates (η, σ) :

$$(\partial_\eta X^\mu \partial_\sigma X_\mu) = 0, \quad (\partial_\sigma X)^2 = -\rho(\eta)^2 (\partial_\eta X)^2, \quad (\partial_\eta X)^2 \Big|_{\sigma=0, \pi} = -1, \quad (4.4)$$

where the last condition indicates that η is a proper time for the endpoints of the string and the function $\rho(\eta)$ is fixed by the equations of motion. In these coordinates the string equations of motion take the following form:

$$\partial_\eta (|\rho(\eta)| \partial_\eta X^\mu) - \partial_\sigma \left(\frac{1}{|\rho(\eta)|} \partial_\sigma X^\mu \right) = 0 \quad (4.5)$$

and the metric is

$$ds^2 = -\frac{1}{\rho(\eta)^2} d\eta^2 + d\sigma^2. \quad (4.6)$$

The 2d solution can be obtained with the help of the coordinate change

$$g(\tau) = c_1 + e^{\frac{\gamma}{m}\tau}, \quad (4.7)$$

which gives

$$X^0 = \frac{m}{\gamma} \frac{\eta_0 \sinh\left(\frac{\gamma}{m}\eta_0\right) \sinh\left(\frac{\gamma}{m}\eta\right)}{\eta_0 \sinh\left(\frac{\gamma}{m}\eta_0\right) - \pi\rho(\eta) \sinh(\alpha\sigma) \sinh(\alpha(\pi-\sigma))}, \quad (4.8)$$

$$X^1 = x_0 - \frac{m}{\gamma} + \frac{m}{\gamma} \frac{\eta_0 \sinh\left(\frac{\gamma}{m}\eta_0\right) \cosh\left(\frac{\gamma}{m}\eta\right) + \pi\rho(\eta) \sinh(\alpha\sigma)^2}{\eta_0 \sinh\left(\frac{\gamma}{m}\eta_0\right) - \pi\rho(\eta) \sinh(\alpha\sigma) \sinh(\alpha(\pi-\sigma))}. \quad (4.9)$$

And for the function ρ we have:

$$\rho(\eta) = -\frac{4\eta_0}{\pi} \frac{\sinh\left(\frac{\gamma}{2m}(\eta + \eta_0)\right) \sinh\left(\frac{\gamma}{2m}(\eta - \eta_0)\right)}{\sinh\left(\frac{\gamma}{m}\eta_0\right)}. \quad (4.10)$$

Here we also introduced the constant η_0 , which corresponds to τ_0 in $(\tilde{\tau}, \tilde{\sigma})$ coordinates:

$$\tau_0 = \frac{m}{\gamma} \sinh\left(\frac{\gamma}{m}\eta_0\right), \quad X^1(\pm\eta_0, \sigma) = 0. \quad (4.11)$$

Another useful relation between the constants is

$$\pi\alpha = \frac{\gamma}{m}\eta_0. \quad (4.12)$$

Now we can see, that the solution in coordinates (η, σ) is very similar to the solution described in (2.12). To be more precise, we only wrote the solution for the time interval $(-\eta_0, \eta_0)$ and the generalization to other time intervals is straightforward.

4.2 Metric for the stationary observer

Now that we determined the string analog of the Rindler metric, it is natural to turn to the stationary observer. The obvious placement for such an observer is the middle of the string at $\tilde{\sigma} = \sigma = \pi/2$. Indeed, at $\sigma = \pi/2$ we have

$$X^0\left(\eta, \frac{\pi}{2}\right) = \frac{2m}{\gamma} \cosh\left(\frac{\gamma}{2m}\eta_0\right)^2 \tanh\left(\frac{\gamma}{2m}\eta\right), \quad X^1\left(\eta, \frac{\pi}{2}\right) = 0. \quad (4.13)$$

Thus, we can introduce the proper time t for the stationary observer as

$$t(\eta) = \frac{2m}{\gamma} \cosh\left(\frac{\gamma}{2m}\eta_0\right)^2 \tanh\left(\frac{\gamma}{2m}\eta\right). \quad (4.14)$$

Then the gauge fixing in coordinates (t, σ) is

$$(\partial_t X^\mu \partial_\sigma X_\mu) = 0, \quad (\partial_\sigma X)^2 = -r(t)^2 (\partial_t X)^2, \quad (\partial_t X)^2 \Big|_{\sigma=\pi/2} = -1, \quad (4.15)$$

where

$$r(t) = \frac{\alpha}{t_0} (t_0^2 - t^2), \quad t_0 \equiv t(\eta_0). \quad (4.16)$$

Going back to the coordinates $(\tilde{\tau}, \tilde{\sigma})$, we can see that the time coordinate $\tilde{\tau}$ is also a good candidate for the proper time of the stationary observer in the following sense:

$$(\partial_{\tilde{\tau}} X)^2 \Big|_{\tilde{\sigma}=\pi/2} = -1. \quad (4.17)$$

So, there is no surprise that from (4.14) we get

$$t_0 = \frac{m}{\gamma} \sinh\left(\frac{\gamma}{m}\eta_0\right) = \tau_0. \quad (4.18)$$

However, the induced metric in coordinates $(\tilde{\tau}, \tilde{\sigma})$ is non-diagonal and cumbersome. In coordinates (t, σ) , on the other hand, the metric reads

$$ds^2 = -\frac{dt^2}{(1 - t^2/t_0^2)^2} + \alpha^2 t_0^2 d\sigma^2, \quad -t_0 \leq t \leq t_0. \quad (4.19)$$

The latter is directly related to the worldsheet metric, which appears in the exact string solution in the AdS_3 space [2].

5 2d/3d string correspondence

In this section we are going to establish the relation between two classical solutions [1] and [2]. The main idea is to identify the worldsheet coordinates from both solutions based on the form of the induced metrics.

5.1 The exact string solution in the AdS_3 space

We start by writing down the Nambu-Goto action in the AdS_3 space:

$$S = -T_0 \int dt du \sqrt{-G}, \quad (5.1)$$

where

$$G = (\partial_t Y)^2 (\partial_u Y)^2 - (\partial_t Y^\mu \partial_u Y_\mu)^2. \quad (5.2)$$

The 3-component field Y^μ is a vector in the AdS_3 spacetime with the metric

$$ds^2 = R^2 \left[- (Y^1)^2 (dY^0)^2 + \frac{(dY^1)^2}{(Y^1)^2} + (Y^1)^2 (dY^2)^2 \right]. \quad (5.3)$$

The solution found in [2] describes a motion of the string in the static gauge

$$Y^\mu = (t, u, y(t, u)) \quad (5.4)$$

and can be written as

$$y = \pm \sqrt{t^2 + b^2 - \frac{1}{u^2}}. \quad (5.5)$$

The constant b is the reciprocal of the acceleration of the string and the two branches of the square root are glued together at $y = 0$. Since the solution describes a string with constant proper acceleration, one can go from the AdS space to a generalized Rindler space. The transformation for the positive branch $y > 0$ is

$$y = \sqrt{b^2 - r^2} \exp\left(\frac{\alpha}{b}\right) \cosh\left(\frac{\theta}{b}\right), \quad (5.6)$$

$$t = \sqrt{b^2 - r^2} \exp\left(\frac{\alpha}{b}\right) \sinh\left(\frac{\theta}{b}\right), \quad (5.7)$$

$$\frac{1}{u} = r \exp\left(\frac{\alpha}{b}\right). \quad (5.8)$$

The same can be done for the negative branch $y < 0$. The new coordinates then cover the two wedges $|y| \geq |t|$. Both α and θ run from $-\infty$ to $+\infty$, but r only covers the interval $(0, b)$. The spacetime metric can be rewritten in the new coordinates as

$$ds^2 = \frac{R^2}{r^2} \left[\frac{dr^2}{1 - r^2/b^2} - \left(1 - \frac{r^2}{b^2}\right) d\theta^2 + d\alpha^2 \right]. \quad (5.9)$$

The solution (5.5) corresponds to $\alpha = 0$ and can be viewed as a part of a one-sheet hyperboloid:

$$y^2 + r^2 - t^2 = b^2. \quad (5.10)$$

In what follows we will consider coordinates (r, θ) as new coordinates on the string worldsheet, which correspond to a different gauge choice for Y^μ :

$$Y^\mu = \left(t(r, \theta), \frac{1}{r}, y(r, \theta) \right). \quad (5.11)$$

This identification between Rindler spacetime coordinates and a particular choice of worldsheet coordinates also appears in the Minkowski setting in [61]. The induced metric on the worldsheet is then

$$ds_{ws}^2 = - \left(1 - \frac{r^2}{b^2}\right) d\theta^2 + \frac{dr^2}{1 - r^2/b^2}. \quad (5.12)$$

5.2 Mapping between the solutions

First, we observe that the worldsheet metric in (5.12) coincides with the worldsheet metric in (4.19) upon a Weyl transformation and the following identification of coordinates:

$$t_{(z)} = r, \quad \sigma = \frac{\theta}{\alpha b}, \quad (5.13)$$

where z in the $t_{(z)}$ notation stands for the zigzag solution. Here we see the main difference between the two solutions: θ runs from $-\infty$ to $+\infty$, whereas σ only belongs to the interval $[0, \pi]$. To overcome this obstacle one needs to take a limit $\alpha \rightarrow +\infty$ and consider only the positive time interval $\theta \geq 0$ or, equivalently, $t \geq 0$ in the AdS solution. The same conclusion can be reached by noting that the zigzag solution has two constant parameters $x_0, \frac{m}{\gamma}$ and one needs to match them to the single parameter b in the AdS solution. The limit $\alpha \rightarrow +\infty$ corresponds to the strong tension limit as it can be seen from the following relations:

$$\sinh(\pi \alpha) = \frac{\gamma}{m} \tau_0 \rightarrow +\infty, \quad \tau_0 \rightarrow b. \quad (5.14)$$

Using (2.13), we get the full matching between the parameters:

$$b = -x_0 > 0, \quad \frac{\gamma}{m} \rightarrow +\infty. \quad (5.15)$$

Thus, the reciprocal of the acceleration in the AdS solution corresponds to the half of the maximum string length in the zigzag solution.




Figure 2: Zigzag solution embedded in the AdS space, $b = 1$

To complete the correspondence we want to write down the coordinate transformation between the Minkowski coordinates X^μ in the zigzag solution and coordinates (t, y) on the boundary of AdS. But first, let's introduce a clear geometrical picture of what is going on. The AdS solution can be described as a top part of the hyperboloid (5.10) when it is cut by the plane $r = 0$ (the boundary of the AdS space), where the quark and antiquark are moving. The zigzag solution, on the other hand, is described by the part of the hyperboloid with $t \geq 0$ and $-b \leq r \leq b$. Going from $r > 0$ to $r = 0$ and then to $r < 0$ can be explained by the specific identification on the boundary of AdS, which we will discuss in the next subsection 5.3. Due to the event horizons at $r = \pm b$, we should also cut the hyperboloid by the planes $r = -b$ and $r = b$ (see Figure 2). The zigzag solution then corresponds to the two branches of the following square root:

$$r = \pm \sqrt{b^2 + t^2 - y^2}. \quad (5.16)$$

The positive branch represents $X^0 > 0$ and the negative one represents $X^0 < 0$. Since r is the proper time of the stationary observer in the zigzag solution, the $r = 0$ plane is exactly the $X^0 = 0$ line in Minkowski. Thus, the worldlines of the quark and antiquark are mapped to the string embedded in Minkowski space at $X^0 = 0$. The same is true in reverse. The worldlines of the endpoints of the string in the zigzag solution are mapped to the closed string embedded in the AdS space at $t = 0$.

Now we are ready to derive the coordinate transformation $X^\mu(t, y)$. First, we consider the positive branch of (5.5):

$$y = \sqrt{b^2 - r^2} \cosh\left(\frac{\theta_+}{b}\right), \quad (5.17)$$

$$t = \sqrt{b^2 - r^2} \sinh\left(\frac{\theta_+}{b}\right). \quad (5.18)$$

Substituting the $\eta(t_{(z)})$ in (4.8)–(4.9) and taking the limit $\gamma/m \rightarrow \infty$ we get

$$y > 0: \quad X^0 = \frac{2m}{\gamma} \frac{(\pm)b\sqrt{b^2 + t^2 - y^2}}{(y - t)^2}, \quad X^1 - x_0 = \frac{2m}{\gamma} \frac{b^2 - y(y - t)}{(y - t)^2}, \quad (5.19)$$

where (\pm) in the expression for X^0 corresponds to the two branches of the square root in (5.16). Here we see that the strong tension limit is accompanied by the rescaling of the X^μ coordinates near one of the endpoints of the string. In particular, since our limit describes zooming in on the left endpoint with $\sigma = 0$, the action introduced in (2.1) becomes an action for the semi-infinite string with one massive particle at $\sigma = 0$. To be more precise, we introduce the following limit in terms of the coordinates $(t_{(z)} \equiv r, \sigma)$ and (X^0, X^1) :

$$y > 0: \quad \sigma = \frac{\theta_+}{\alpha b}, \quad X^0 = \frac{2m}{\gamma} Z_+^0, \quad X^1 - x_0 = \frac{2m}{\gamma} Z_+^1, \quad \frac{\gamma}{m} \rightarrow +\infty. \quad (5.20)$$

Then the functions $Z_+^\mu(r, \theta)$ given by (5.19) describe a classical solution to the following string action:

$$S = \frac{2m^2}{\gamma} \int dr \left(-\sqrt{-(\partial_r Z_+(r, 0))^2} - 2 \int_0^\infty d\theta \sqrt{-G} \right), \quad G = (\partial_r Z_+)^2 (\partial_\theta Z_+)^2 - (\partial_r Z_+^\mu \partial_\theta Z_\mu^+)^2. \quad (5.21)$$

As one can see, the action has an overall scale m^2/γ and the equations of motion are independent of the ratio m/γ . The coordinate transformation from (t, y) to (Z_+^0, Z_+^1) can be written in the form of the special conformal transformation:

$$Z_+^\mu = \frac{\hat{y}^\mu - \beta^\mu \hat{y}^2}{1 - 2\beta^\mu \hat{y}_\mu + \beta^2 \hat{y}^2} \quad (5.22)$$

with

$$y > 0: \quad \hat{y}^\mu = \left(\frac{br}{b^2 + t^2}, \frac{ty}{b^2 + t^2} \right), \quad \beta^\mu = (0, 1). \quad (5.23)$$

So far we described the "blue" part of the surface with $y > 0$ in Figure 2, which corresponds to the worldline of the quark and the attached string in the AdS. To describe the "red" part we consider the negative branch of (5.5):

$$y = -\sqrt{b^2 - r^2} \cosh\left(\frac{\theta_-}{b}\right), \quad (5.24)$$

$$t = \sqrt{b^2 - r^2} \sinh\left(\frac{\theta_-}{b}\right), \quad (5.25)$$

and zoom in on the right endpoint of the string with $\sigma = \pi$:

$$t_{(z)} = r, \quad \sigma = \pi - \frac{\theta_-}{\alpha b}, \quad \alpha \rightarrow \infty. \quad (5.26)$$

The corresponding coordinate transformation $X^\mu(t, y)$ is then

$$y < 0: \quad X^0 = \frac{2m(\pm)b\sqrt{b^2 + t^2 - y^2}}{\gamma(y+t)^2}, \quad X^1 + x_0 = \frac{2m(-b^2 + y(y+t))}{\gamma(y+t)^2}, \quad (5.27)$$

Combining this with (5.19), we get the answer for all allowed values of y :

$$|y| \geq t \geq 0: \quad X^0 = \frac{2m(\pm)b\sqrt{b^2 + t^2 - y^2}}{\gamma(|y| - t)^2}, \quad X^1 = \text{sgn}(y) \left(-b + \frac{2mb^2 - |y|(|y| - t)}{\gamma(|y| - t)^2} \right). \quad (5.28)$$

By introducing local Minkowski variables Z_-^μ near the left endpoint of the string with $\sigma = \pi$, we can rewrite (5.28) in the form of the special conformal transformation as

$$\mu = 0, 1: \quad Z_\pm^\mu = (\pm 1)^\mu \frac{\hat{y}^\mu - \beta^\mu \hat{y}^2}{1 - 2\beta^\mu \hat{y}_\mu + \beta^2 \hat{y}^2}, \quad (5.29)$$

where $(+)$ corresponds to $y > 0$, $(-)$ corresponds to $y < 0$ and vectors \hat{y}^μ, β^μ are given by

$$\hat{y}^\mu = \left(\frac{br}{b^2 + t^2}, \frac{t|y|}{b^2 + t^2} \right), \quad \beta^\mu = (0, 1). \quad (5.30)$$

It is helpful to picture the domain in (y, t) variables, which is mapped to the zigzag solution. As it can be seen from Figure 2, the domain is $t \leq |y| \leq \sqrt{b^2 + t^2}$ (shaded regions in Fig. 3). The boundaries of this region are the worldlines of the quark-antiquark pair $|y| = \sqrt{b^2 + t^2}$ and the event horizons $|y| = t$. Since for $r \neq 0$ the hyperboloid covers the domain twice, we have two horizons $Z^0 = \pm\infty$ at $y = t$ and at $y = -t$.

The final step in the correspondence is to associate the two lines $|y| = t$ with the worldline of the middle of the string at $\sigma = \pi/2$ in the zigzag solution. To illustrate this point, we are going to project worldlines of different points on the string on the half-planes $y = t \geq 0$ and $y = -t \leq 0$. As we derived earlier, different points on the string in the zigzag solution are parametrized by the proper time θ of the accelerated observer in the AdS space. To be more precise, there are two different coordinates θ_+ and θ_- describing positive and negative branches of the square root in (5.5). In Figure 4 we see how the different points of the zigzag string are embedded in the AdS space. The θ_+ coordinate describes the points on the string close to $\sigma = 0$ and θ_- describes the points close to $\sigma = \pi$. We know from the full zigzag solution, that as both $\theta_\pm \rightarrow +\infty$ we are getting closer to the middle of the string. Thus, by gluing θ_+ and θ_- at infinity we are able to continuously move from the left endpoint of the string with $\sigma = 0$ to the right one with $\sigma = \pi$. The gluing point $\theta_\pm = +\infty$ corresponds to the middle of the string in the zigzag solution.

5.3 Identifications on the boundary of AdS_3

The solution (5.10) is written in terms of the Poincare coordinates (t, r, y) . Usually, one considers $r > 0$ and in the limit $r \rightarrow +0$ one approaches the boundary of AdS spacetime. However, the solution itself allows for both positive and negative values of r as well as $r = 0$. This can be explained by adding the boundary $r \rightarrow \pm 0$ and introducing specific identifications between different points on the boundary. To figure out the exact form of these identifications, we consider global coordinates $(\hat{\tau}, \hat{\rho}, \hat{\theta})$ on the AdS_3 spacetime:

$$ds^2 = R^2 \left(-\cosh^2 \hat{\rho} d\hat{\tau}^2 + d\hat{\rho}^2 + \sinh^2 \hat{\rho} d\hat{\theta}^2 \right), \quad (5.31)$$

where $\hat{\rho} \geq 0$, $\hat{\tau} \in [0, 2\pi]$ and $\hat{\theta} \in [0, 2\pi]$. The relation to the Poincare coordinates can be written in the following form:

$$\begin{aligned} R \cosh \hat{\rho} \cos \hat{\tau} &= r/2 + (R^2 + y^2 - t^2) / (2r), \\ \cosh \hat{\rho} \sin \hat{\tau} &= t/r, \\ \sinh \hat{\rho} \sin \hat{\theta} &= y/r, \\ R \sinh \hat{\rho} \cos \hat{\theta} &= r/2 - (R^2 - y^2 + t^2) / (2r). \end{aligned} \quad (5.32)$$

We can see, that Poincare coordinates with $r > 0$ cover only "half" of the AdS spacetime. Changing the sign of r corresponds to a simultaneous shift of $\hat{\tau}$ and $\hat{\theta}$ by π :

$$r \rightarrow -r : \quad \hat{\tau} \rightarrow \hat{\tau} + \pi \quad \text{and} \quad \hat{\theta} \rightarrow \hat{\theta} + \pi. \quad (5.33)$$

Thus, two limits $r \rightarrow +0$ and $r \rightarrow -0$ describe two different regions on the AdS boundary. To continuously go from $r > 0$ to $r < 0$, one needs to identify these boundary regions. In our particular example we consider $t \geq 0$, which leaves us with two cases: $t \geq 0, y \geq 0$ and $t \geq 0, y \leq 0$. In each case $r > 0$ and $r < 0$ cover different quadrants in $(\hat{\tau}, \hat{\theta})$ coordinates (see Figure 5). In the limit $r \rightarrow \pm 0$ we identify the points on the boundary according to the following rule (see Figure 6):

$$(\hat{\tau}, \hat{\theta}) \longleftrightarrow (\hat{\tau} + \pi, \hat{\theta} + \pi). \quad (5.34)$$

In this way we get an unconventional Poincare patch with $t \geq 0$ and $-\infty < r, y < +\infty$, which still covers "half" of the AdS_3 . This patch can be visualized in global coordinates $(\hat{\tau}, \hat{\rho}, \hat{\theta})$ as in Figure 7.

6 Conclusion

In this paper we introduced a new relation between two classical string solutions [1] and [2]. The first solution describes two point masses joined by a massless string in the 2d Minkowski spacetime and the second solution describes a quark-antiquark pair connected by a string embedded in the 3d AdS spacetime. As such, the relation takes place in the lowest possible number of spacetime dimensions and requires a particular form of identification on the boundary of AdS_3 . It was shown that the latter identification leads to an alternative definition of the Poincare patch. The following physical interpretation of the established relation could be given. The first solution [1] provides a model for the motion of two quarks subjected to a strong interaction. The second solution is conjectured to describe a motion of the quark-antiquark pair moving in the external electric field. The external electric field not only balances out the attracting force between the quark and antiquark, but also accelerates them with constant proper acceleration in different directions. If the physical interpretation of the second solution [2] proves to be correct, the result of the present paper suggests a non-trivial relation between these two different physical systems. Since we covered classical aspects of the said relation, a straightforward continuation would be to introduce a quantum version of the correspondence. Another interesting direction is to study physical effects like the Unruh effect on both sides of the correspondence. In the case of the second solution the Unruh effect was discussed in the original paper [2] and the corresponding temperature was compared to the Hawking temperature determined by the behavior of the metric near the horizon. In the case of the first solution it is possible to derive an analog of the Unruh effect by adding small transversal excitations and taking the limit of a large string length. This will be discussed in a subsequent paper [61].

Acknowledgements

The author would like to thank Martin Roček and Zohar Komargodski for many valuable discussions and comments. The work was supported by the "BASIS" Foundation grant 18-1-1-50-3 and in part by RFBR grant 19-51-18006-Bolg_a.

A Additional figures




Figure 3: (y, t) domain, $b = 1$




Figure 4: Worldlines of different points on the zigzag string, $y = \pm t$ and $b = 1$




Figure 5: Regions with $r > 0$ and $r < 0$ in $(\hat{t}, \hat{\theta})$ coordinates




Figure 6: Identification between points on the boundary of AdS_3




Figure 7: Unconventional Poincare patch

References

- [1] W. A. Bardeen, Itzhak Bars, Andrew J. Hanson, and R. D. Peccei. Study of the longitudinal kink modes of the string. *Phys. Rev. D*, 13:2364–2382, 1976. doi.org/10.1103.
- [2] Bo-Wen Xiao. On the exact solution of the accelerating string in AdS_5 space. *Physics Letters B*, 665(4):173–177, 2008. arXiv:0804.13435 [hep-th].
- [3] Juan Maldacena. The Large-N Limit of Superconformal Field Theories and Supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998. arXiv:hep-th/9711200.
- [4] Edward Witten. Anti-de Sitter Space And Holography. *Adv. Theor. Math. Phys.*, 2:253–291, 1998. arXiv:hep-th/9802150.
- [5] S.S. Gubser, I.R. Klebanov, and A.M. Polyakov. Gauge theory correlators from non-critical string theory. *Physics Letters B*, 428(1):105–114, 1998. arXiv:hep-th/9802109.
- [6] E. S. Fradkin and Arkady A. Tseytlin. Quantum Equivalence of Dual Field Theories. *Annals Phys.*, 162:31, 1985. doi.org/10.1016.
- [7] Bala Sathiapalan. Duality in statistical mechanics and string theory. *Phys. Rev. Lett.*, 58:1597–1599, 1987. 10.1103/PhysRevLett.58.1597.
- [8] T. H. Buscher. A Symmetry of the String Background Field Equations. *Phys. Lett. B*, 194:59–62, 1987. doi.org/10.1016.
- [9] Philip Candelas, Xenia C. De La Ossa, Paul S. Green, and Linda Parkes. A Pair of Calabi-Yau manifolds as an exactly soluble superconformal theory. *Nucl. Phys. B*, 359:21–74, 1991. doi.org/10.1016.
- [10] Ashoke Sen. Strong - weak coupling duality in four-dimensional string theory. *Int. J. Mod. Phys. A*, 9(21):3707–3750, 1994. arXiv:9402002[hep-th].
- [11] John H. Schwarz. Evidence for nonperturbative string symmetries. *Letters in Mathematical Physics*, 34:309–317, 1995. arXiv:9411178[hep-th].
- [12] C. M. Hull and P. K. Townsend. Unity of superstring dualities. *Nucl. Phys. B*, 438:109–137, 1995. arXiv:9410167[hep-th].
- [13] Luis F. Alday, Davide Gaiotto, and Yuji Tachikawa. Liouville Correlation Functions from Four-Dimensional Gauge Theories. *Letters in Mathematical Physics*, 91(2):167–197, 2010. arXiv:0906.3219[hep-th].
- [14] Niclas Wyllard. A_{N-1} conformal Toda field theory correlation functions from conformal $N = 2$ $SU(N)$ quiver gauge theories. *Journal of High Energy Physics*, 2009(11), 2009. arXiv:0907.2189[hep-th].
- [15] Tadashi Takayanagi. Holographic Dual of a Boundary Conformal Field Theory. *Phys. Rev. Lett.*, 107, 2011. arXiv:1105.5165[hep-th].
- [16] Mitsutoshi Fujita, Tadashi Takayanagi, and Erik Tonni. Aspects of AdS/BCFT. *Journal of High Energy Physics*, 2011, 2011. arXiv:1108.5152[hep-th].
- [17] I. Bars and K. Sfetsos. Global analysis of new gravitational singularities in string and particle theories. *Phys. Rev. D*, 46:4495–4509, 1992. doi.org/10.1103.
- [18] I. Bars and J. Schulze. Folded strings falling into a black hole. *Phys. Rev. D*, 51:1854–1868, 1995. doi.org/10.1103.
- [19] Juan Maldacena. Long strings in two dimensional string theory and non-singlets in the matrix model. *Journal of High Energy Physics*, 2005(09):078, 2005. arXiv:hep-th/0503112.
- [20] Juan Maldacena and Hirosi Ooguri. Strings in AdS_3 and the $SL(2, R)$ WZW model. I: The spectrum. *J. Math. Phys.*, 42(7), 2001. arXiv:hep-th/0001053.
- [21] Juan Maldacena, Hirosi Ooguri, and John Son. Strings in AdS_3 and the $SL(2, R)$ WZW model. II: Euclidean black hole. *J. Math. Phys.*, 42(7), 2001. arXiv:hep-th/0005183.
- [22] Juan Maldacena and Hirosi Ooguri. Strings in AdS_3 and the $SL(2, R)$ WZW model. III: Correlation functions. *Phys. Rev. D*, 65(10):106006, 2002. arXiv:hep-th/0111180.
- [23] J. Wess and B. Zumino. Consequences of anomalous ward identities. *Physics Letters B*, 37(1):95–97, 1971. doi.org/10.1016.
- [24] Edward Witten. Global aspects of current algebra. *Nuclear Physics B*, 223(2):422–432, 1983. doi.org/10.1016.

- [25] Edward Witten. Non-abelian bosonization in two dimensions. *Communications in Mathematical Physics*, 92:455–472, 1984. doi.org/10.1007.
- [26] S.P. Novikov. Multivalued functions and functionals. An analogue of the Morse theory. *Dokl. Akad. Nauk SSSR*, 260(1):31–35, 1981. www.mi-ras.ru/68.pdf.
- [27] S.P. Novikov. The Hamiltonian formalism and a many-valued analogue of Morse theory. *Uspekhi Mat. Nauk*, 37:3–49, 1982. doi.org/10.1070.
- [28] Ofer Aharony and Matan Field. On the effective theory of long open strings. *Journal of High Energy Physics*, 2011, 2011. arXiv:1008.2636[hep-th].
- [29] Ofer Aharony and Zohar Komargodski. The effective theory of long strings. *Journal of High Energy Physics*, 2013, 2013. arXiv:1302.6257[hep-th].
- [30] V. V. Bazhanov, G. A. Kotousov, and S. L. Lukyanov. Equilibrium density matrices for the 2D black hole sigma models from an integrable spin chain. *Journal of High Energy Physics*, 2021(169), 2021. arXiv:2010.10603[hep-th].
- [31] Andrei Mikhailov. Nonlinear waves in AdS/CFT correspondence. 2003. arXiv:hep-th/0305196.
- [32] C.P. Herzog, A. Karch, P. Kovtun, C. Kozcaz, and L.G. Yaffe. Energy loss of a heavy quark moving through $N = 4$ supersymmetric Yang-Mills plasma. *Journal of High Energy Physics*, 2006(7), 2006. arXiv:0605158[hep-th].
- [33] Steven S. Gubser. Drag force in AdS/CFT. *Phys. Rev. D*, 74(12), 2006. arXiv:hep-th/0605182.
- [34] Fabio Dominguez, C. Marquet, A.H. Mueller, Bin Wu, and Bo-Wen Xiao. Comparing energy loss and p_{\perp} -broadening in perturbative QCD with strong coupling $N = 4$ SYM theory. *Nuclear Physics A*, 811(1):197–222, 2008. arXiv:0803.3234[nucl-th].
- [35] Mariano Chernicoff and Alberto Güijosa. Acceleration, energy loss and screening in strongly-coupled gauge theories. *Journal of High Energy Physics*, 2008(06), 2008. arXiv:0803.3070[hep-th].
- [36] Jorge Casalderrey-Solana and Derek Teaney. Heavy quark diffusion in strongly coupled $\mathcal{N} = 4$ Yang-Mills theory. *Phys. Rev. D*, 74, 2006. arXiv:0605199[hep-th].
- [37] Jorge Casalderrey-Solana and Derek Teaney. Transverse momentum broadening of a fast quark in a Script $\mathcal{N} = 4$ Yang-Mills plasma. *Journal of High Energy Physics*, 2007(04), 2007. arXiv:0701123[hep-th].
- [38] Elena Cáceres, Mariano Chernicoff, Alberto Güijosa, and Juan F. Pedraza. Quantum Fluctuations and the Unruh effect in strongly-coupled conformal field theories. *Journal of High Energy Physics*, 2010(6), 2010. arXiv:1003.5332[hep-th].
- [39] Mariano Chernicoff and Angel Paredes. Accelerated detectors and worldsheet horizons in AdS/CFT. *Journal of High Energy Physics*, 2011(3), 2011. arXiv:1011.4206[hep-th].
- [40] Kazuo Ghoroku, Masafumi Ishihara, Kouki Kubo, and Tomoki Taminato. Accelerated quark and holography for confining gauge theory. *Phys. Rev. D*, 83(2):024020, 2011. arXiv:1010.4396[hep-th].
- [41] Alberto Güijosa and Juan F. Pedraza. Early-time energy loss in a strongly-coupled SYM plasma. *Journal of High Energy Physics*, 2011(5), 2011. arXiv:1102.4893[hep-th].
- [42] Takayuki Hirayama, Pei-Wen Kao, Shoichi Kawamoto, and Feng-Li Lin. Unruh effect and holography. *Nuclear Physics B*, 844(1):1–25, 2011. arXiv:1001.1289[hep-th].
- [43] Mariano Chernicoff, J. Antonio García, Alberto Güijosa, and Juan F. Pedraza. Holographic lessons for quark dynamics. *Journal of Physics G: Nuclear and Particle Physics*, 39(5), 2012. arXiv:1111.0872[hep-th].
- [44] Kristan Jensen and Andreas Karch. Holographic Dual of an Einstein-Podolsky-Rosen Pair has a Wormhole. *Phys. Rev. Lett.*, 111(21):211602, 2013. arXiv:1307.1132[hep-th].
- [45] Mariano Chernicoff, Alberto Güijosa, and Juan F. Pedraza. Holographic EPR pairs, wormholes and radiation. *Journal of High Energy Physics*, 2013(10), 2013. arXiv:1308.3695[hep-th].
- [46] Julian Sonner. Holographic Schwinger Effect and the Geometry of Entanglement. *Phys. Rev. Lett.*, 111(21):211603, 2013. arXiv:1307.6850[hep-th].
- [47] Veronika E. Hubeny and Gordon W. Semenoff. String worldsheet for accelerating quark. *Journal of High Energy Physics*, 2015(10), 2015. arXiv:1410.1171[hep-th].
- [48] Mehmet Emre Tasgin. Entanglement, violation of Kramers-Kronig relation and curvature in spacetime. 2019. arXiv:1912.04399[physics.gen-ph].

- [49] Andrej Ficnar and Steven S. Gubser. Finite momentum at string endpoints. *Phys. Rev. D*, 89, 2014. arXiv:1306.6648[hep-th].
- [50] Steven S. Gubser, Daniel R. Gulotta, Silviu S. Pufu, and Fábio D. Rocha. Gluon energy loss in the gauge-string duality. *Journal of High Energy Physics*, 2008(10), 2008. arXiv:0803.1470[hep-th].
- [51] Y. Hatta, E. Iancu, and A.H. Mueller. Jet evolution in the $\mathcal{N} = 4$ SYM plasma at strong coupling. *Journal of High Energy Physics*, 2008(05), 2008. arXiv:0803.2481[hep-th].
- [52] Paul M. Chesler, Kristan Jensen, and Andreas Karch. Jets in strongly coupled $\mathcal{N} = 4$ super Yang-Mills theory. *Phys. Rev. D*, 79, 2009. arXiv:0804.3110[hep-th].
- [53] Paul M. Chesler, Kristan Jensen, Andreas Karch, and Laurence G. Yaffe. Light quark energy loss in strongly coupled $\mathcal{N} = 4$ supersymmetric Yang-Mills plasma. *Phys. Rev. D*, 79, 2009. arXiv:0810.1985[hep-th].
- [54] Andrej Ficnar, Jorge Noronha, and Miklos Gyulassy. Falling Strings and Light Quark Jet Quenching at LHC. *Nuclear Physics A*, 910-911, 2013. arXiv:1208.0305[hep-th].
- [55] Andrej Ficnar. AdS/CFT energy loss in time-dependent string configurations. *Phys. Rev. D*, 86, 2012. arXiv:1201.1780[hep-th].
- [56] Amin Faraji Astaneh and Amir Esmail Mosaffa. Quantum local quench, AdS/BCFT and Yo-Yo string. *Journal of High Energy Physics*, 2015, 2015. arXiv:1405.5469.
- [57] S. Dubovsky. A simple worldsheet black hole. *Journal of High Energy Physics*, 2018(11), 2018. arXiv:1803.00577 [hep-th].
- [58] John C. Donahue and Sergei Dubovsky. Confining strings, infinite statistics, and integrability. *Phys. Rev. D*, 101:081901, 2020. arXiv:1907.07799 [hep-th].
- [59] John C. Donahue and Sergei Dubovsky. Classical integrability of the zigzag model. *Phys. Rev. D*, 102:026005, 2020. arXiv:1912.08885 [hep-th].
- [60] B. M. Barbashov and A. L. Koshkarov. Some solutions of boundary conditions for relativistic string with massive ends. *Letters in Mathematical Physics*, 3:39–46, 1979. .
- [61] Gleb Aminov. On the Unruh effect for the string with massive ends. *To appear*, 2021.
- [62] H. J. de Vega, A. L. Larsen, and N. Sánchez. Quantum string dynamics in the conformal invariant $SL(2, R)$ WZW background: Anti-de Sitter space with torsion. *Phys. Rev. D*, 58(2):026001, 1998. arXiv:hep-th/9803038.
- [63] A. L. Larsen and N. Sánchez. From the WZW model to the Liouville equation: Exact string dynamics in a conformally invariant AdS background. *Phys. Rev. D*, 58(12):126002, 1998. arXiv:hep-th/9805173.
- [64] C. Ahn, M. Rocek, K. Schoutens, and A. Sevrin. Superspace WZW models and black holes. 1991. arXiv:9110035[hep-th].
- [65] Edward Witten. String theory and black holes. *Phys. Rev. D*, 44:314–324, 1991. doi.org/10.1103.
- [66] Shijong Ryang. Conformal transformations and strings for an accelerating quark-antiquark pair in AdS3. *Journal of High Energy Physics*, 2015(2), 2015. arXiv:1412.2428 [hep-th].
- [67] Y. Nambu. Lectures at the Copenhagen symposium. 1970.
- [68] T. Goto. Relativistic Quantum Mechanics of One-Dimensional Mechanical Continuum and Subsidiary Condition of Dual Resonance Model. *Progress of Theoretical Physics*, 46(5):1560–1569, 1971. 10.1143/PTP.46.1560.