

SHARP TRACE AND KORN INEQUALITIES FOR DIFFERENTIAL OPERATORS

LARS DIENING AND FRANZ GMEINER

ABSTRACT. We establish sharp trace- and Korn-type inequalities that involve vectorial differential operators, the focus being on situations where global singular integral estimates are not available. Starting from a novel approach to sharp Besov boundary traces by Riesz potentials and oscillations that equally applies to $p = 1$, a case difficult to be handled by harmonic analysis techniques, we then classify boundary trace- and Korn-type inequalities. For $p = 1$ and so despite the failure of the Calderón-Zygmund theory, we prove that sharp trace estimates can be systematically reduced to full k -th order gradient estimates. Moreover, for $1 < p < \infty$, where sharp trace- yield Korn-type inequalities on smooth domains, we show for the basically optimal class of John domains that Korn-type inequalities persist – even though the reduction to global Calderón-Zygmund estimates by extension operators might not be possible.

CONTENTS

1.	Introduction	1
2.	Domains	5
3.	The sharp representation theorem and Poincaré-type inequalities	7
4.	Proof of Theorem 1.1: The trace estimate via Riesz potentials	14
5.	Proof of Theorem 1.2: The trace estimate via reduction to $W^{k,1}(\Omega; V)$	23
6.	Proof of Theorem 1.3: Korn without global singular integral estimates	31
7.	Appendix	43
	References	46

1. INTRODUCTION

1.1. **Calderón-Zygmund and L^1 -estimates.** A variety of partial differential equations or problems from the calculus of variations require to bound the L^p -norms of full higher order gradients by those of given differential operators. To be more precise, let $k \in \mathbb{N}$, $k \geq 1$ and suppose that \mathbb{A} is a homogeneous, k -th order linear, constant-coefficient differential operator on \mathbb{R}^n with $n \geq 2$ between the two finite dimensional real vector spaces V and W . Then \mathbb{A} has a representation

$$(1.1) \quad \mathbb{A} = \sum_{\substack{\alpha \in \mathbb{N}_0^n \\ |\alpha|=k}} \mathbb{A}_\alpha \partial^\alpha,$$

Keywords: Trace theorem, Uspenskiĭ’s theorem, L^1 -estimates, Korn inequality, \mathbb{C} -ellipticity, Riesz potentials, inverse estimates, NTA domains, John domains, weighted inequalities.

Acknowledgments. The authors are grateful to Herbert Koch for general discussions on the theme of the paper, and to David Cruz-Urbe for pointing out the reference [20] towards extrapolation in Lorentz spaces. F.G. acknowledges financial support by the Hausdorff Centre for Mathematics, Bonn, and the University of Bielefeld. The research of L.D. was supported by the DFG through the CRC 1283.

where for each $\alpha \in \mathbb{N}_0^n$ with $|\alpha| = k$, $\mathbb{A}_\alpha: V \rightarrow W$ is a fixed linear map. Following the foundational work of CALDERÓN & ZYGMUND [15], if $1 < p < \infty$, there exists a constant $c = c(\mathbb{A}, p) > 0$ such that there holds

$$(1.2) \quad \|D^k u\|_{L^p(\mathbb{R}^n)} \leq c \|\mathbb{A}u\|_{L^p(\mathbb{R}^n)} \quad \text{for all } u \in C_c^\infty(\mathbb{R}^n; V)$$

if and only if \mathbb{A} is elliptic; (1.2) is also referred to as *Korn-type inequality* [46]. Here, in the most general sense, we call \mathbb{A} *elliptic* provided the Fourier symbol

$$(1.3) \quad \mathbb{A}[\xi] = \sum_{|\alpha|=k} \xi^\alpha \mathbb{A}_\alpha: V \rightarrow W \quad \text{is injective for any } \xi \in \mathbb{R}^n \setminus \{0\},$$

cf. HÖRMANDER [34] and SPENCER [57]. Due to a celebrated counterexample of ORNSTEIN [53] – colloquially termed *Ornstein’s Non-Inequality* – inequality (1.2) does not persist for $p = 1$ in general. Despite the failure of (1.2) for $p = 1$, one might still hope to find conditions on \mathbb{A} such that lower order quantities depending on u can be controlled in terms of $\|\mathbb{A}u\|_{L^1}$. In the present paper we shall give a resolution of this matter for the boundary traces in the case of k -th order differential operators and *all* $1 \leq p < \infty$, cf. Theorem 1.1 for $p = 1$ (and Theorem 4.4 for $1 < p < \infty$). We then classify this result within situations where *global* Calderón-Zygmund estimates are not available, might it be because of $p = 1$ (Theorem 1.2) or the irregularity of the underlying domains for $1 < p < \infty$ (Theorem 1.3).

1.2. Limiting L^1 -trace estimates via Riesz potentials. The general objective of limiting L^1 -estimates is to classify those operators \mathbb{A} such that well-known inequalities for the full k -th order gradients hold with D^k replaced by \mathbb{A} , too. Starting with BOURGAIN & BREZIS [7, 8], the problem of bounding lower order norms against L^1 -norms of differential expressions $\mathbb{A}u$ has been intensively studied from various perspectives in recent years, cf. VAN SCHAFTINGEN et al. [66, 67, 68, 9, 31] for a systematic treatment of Sobolev-type inequalities; also see [54]. By ORNSTEIN’s Non-Inequality [53, 45, 44], none of these results can be directly inferred from the full k -th order gradient estimates. This equally applies to trace estimates, dealt with in the first order case by BREIT and the authors [11], and in turn gives rise to Sobolev embeddings on domains, cf. RAITA and the second author [30].

The instrumental condition making boundary trace inequalities work is that of *\mathbb{C} -ellipticity*. Following [56, 11, 30], we say that \mathbb{A} of the form (1.1) is *\mathbb{C} -elliptic* provided

$$(1.4) \quad \mathbb{A}[\xi]: V + iV \rightarrow W + iW \text{ is injective for each } \xi \in \mathbb{C}^n \setminus \{0\}.$$

\mathbb{C} -ellipticity is a stronger condition than VAN SCHAFTINGEN’s cancellation condition [67]. So for instance the generalisation of GAGLIARDO’s trace inequality,

$$(1.5) \quad \|u\|_{L^1(\partial\Omega)} \leq c(\|u\|_{L^1(\Omega)} + \|\mathbb{A}u\|_{L^1(\Omega)}) \quad \text{for all } u \in C^\infty(\overline{\Omega}; V)$$

for first order \mathbb{C} -elliptic operators holds if and only if \mathbb{A} is \mathbb{C} -elliptic, see [11]; here $\Omega \subset \mathbb{R}^n$ is a bounded smooth domain. Since the trace operator $\text{Tr}_{\partial\Omega}: W^{1,1}(\Omega; V) \rightarrow L^1(\Omega; V)$ is surjective, (1.5) is optimal. However, if $k \geq 2$, by a result of USPENSKIĬ [64] there exists $c = c(n, k, V) > 0$ such that for any open halfspace $\mathbb{H} \subset \mathbb{R}^n$ there holds

$$(1.6) \quad \|u\|_{\dot{B}_{1,1}^{k-1}(\partial\mathbb{H})} \leq c\|D^k u\|_{L^1(\mathbb{H})} \quad \text{for all } u \in C_c^\infty(\overline{\mathbb{H}}; V)$$

with the homogeneous Besov space $\dot{B}_{1,1}^{k-1}(\partial\mathbb{H}; V) (\subsetneq \dot{W}^{1,1}(\partial\mathbb{H}; V))$ (cf. Section 4.1 for the definition), resulting in a *surjective trace operator*. One may thus wonder whether \mathbb{C} -ellipticity is equally necessary and sufficient to yield the corresponding variant of the trace estimate (1.6), also see [30, Open Problem 4.8]. As our first result, we answer this by

Theorem 1.1 (Higher order traces I). *Let $k \in \mathbb{N}$, $k \geq 2$. Then the following are equivalent for a k -th order differential operator \mathbb{A} of the form (1.1):*

- (a) \mathbb{A} is \mathbb{C} -elliptic in the sense of (1.4).
- (b) There exists $c = c(\mathbb{A}) > 0$ such that for any open halfspace \mathbb{H} there holds

$$(1.7) \quad \|u\|_{\dot{B}_{1,1}^{k-1}(\partial\mathbb{H})} \leq c\|\mathbb{A}u\|_{L^1(\mathbb{H})} \quad \text{for all } u \in C_c^\infty(\overline{\mathbb{H}}; V).$$

The previous theorem particularly implies that, for open and bounded domains Ω with smooth boundary $\partial\Omega$ and \mathbb{C} -elliptic \mathbb{A} , there exists a surjective boundary trace operator $\text{Tr}_{\partial\Omega}: \text{BV}^{\mathbb{A}}(\Omega) \rightarrow B_{1,1}^{k-1}(\partial\Omega; V)$ for *functions of bounded \mathbb{A} -variation* (cf. Theorem 4.7).

Even though a lowering of the smoothness of the boundary is conceivable (and hereafter requires to work with higher order smoothness spaces on less regular sets), our main focus in the previous theorem is not a possibly involved geometry or low regularity of boundaries, but the estimate (1.7) based on Riesz potential estimates; more general domains are addressed in Section 1.3 and 1.4. Note that a weaker estimate, replacing the $\dot{B}_{1,1}^{k-1}$ -norm in (1.7) by the $\dot{W}^{k-1,1}$ -norm has been obtained by RAITA, VAN SCHAFTINGEN and the second author in [31]; again note that $\dot{B}_{1,1}^{k-1} \subsetneq \dot{W}^{k-1,1}$ (cf. BREZIS & PONCE [13, Rem. A.1, p. 1238]).

As pointed out by LEONI & TICE [47], the only available approaches to the classical USPENSKIĬ trace estimate (1.6) for $p = 1$ rely on the finite difference characterisation of Besov spaces; also see MIRONESCU & RUSS [52]. By ORNSTEIN'S Non-Inequality, controlling the requisite finite differences in terms of $\mathbb{A}u$ and *not* $D^k u$ is far from clear, and so (1.7) must be approached differently. Toward Theorem 1.1, our line of action instead is to combine an oscillation characterisation of Besov spaces (cf. Lemma 4.1) and a novel sharp Riesz potential representation formula for \mathbb{C} -elliptic operators (cf. Proposition 3.3). This approach equally seems to be new in the case $1 < p < \infty$ and even for $\mathbb{A} = D^k$, see Section 4.4.

By ARONSZAJN [4] and SMITH [56] (also see KAŁAMAJSKA [41, 42]), it is well-known that for \mathbb{C} -elliptic operators \mathbb{A} and $1 < p < \infty$ one has the norm equivalence

$$(1.8) \quad \|u\|_{W^{k,p}(\Omega)} \simeq \|u\|_{W^{k-1,p}(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)}, \quad u \in W^{k,p}(\Omega; V)$$

for smoothly bounded $\Omega \subset \mathbb{R}^n$. As asserted in Theorem 4.4, for such domains a corresponding sharp variant of the trace estimate (1.7) for $1 < p < \infty$ does not only follow from but *is equivalent* to (1.8), thereby providing a self-contained proof of (1.8). Here, the respective trace estimates yield the existence of a suitable extension operator and so allow to reduce (1.8) to *global* singular integral estimates on \mathbb{R}^n . In particular, the sharp trace estimates obtained by our Riesz potential approach imply (1.8) and then vice versa by the sharp trace theorem for $W^{k,p}$.

In view of the failure of Calderón-Zygmund estimates on L^1 , inequality (1.8) cannot be obtained for $p = 1$ in general and, if $1 < p < \infty$, the above approach only allows to conclude (1.8) for suitable extension domains. We thus proceed by classifying the underlying mechanisms that allow to arrive at Theorem 1.1 and the Korn-type estimate (1.8) in situations where the above approaches are not necessarily available.

1.3. Reduction to $W^{k,1}$ -estimates despite failure of L^1 -CZ theory. Working from Theorem 1.1, \mathbb{C} -ellipticity yields the same trace estimates as known from the full k -th order gradient case. Thus one might wonder whether – *despite the failure of the Calderón-Zygmund estimates for $p = 1$* – \mathbb{C} -ellipticity directly allows to

$$(1.9) \quad \boxed{\text{reduce limiting } L^1\text{-trace estimates to those available for } W^{k,1}.$$

As our second main result, Theorem 1.2 below and, more importantly, its proof, this is the case indeed. It is singled out as its proof seems to be *the first instance in the literature which uses the reduction metaprinciple (1.9)* to arrive at limiting L^1 -estimates for differential operators in the face of ORNSTEIN's Non-Inequality. The proof is naturally set up for NTA_{n-1} -domains (cf. Section 2 for this terminology), leading to the following

Theorem 1.2 (Higher order traces II). *Let $k \in \mathbb{N}_{\geq 1}$. Then the following are equivalent for an operator \mathbb{A} of the form (1.1):*

- (a) \mathbb{A} is \mathbb{C} -elliptic in the sense of (1.4).
- (b) For every open, bounded, NTA_{n-1} -domain $\Omega \subset \mathbb{R}^n$, the space $W^{\mathbb{A},1}(\Omega) := \{u \in L^1(\Omega; V) : \|u\|_{L^1(\Omega)} + \|\mathbb{A}u\|_{L^1(\Omega)} < \infty\}$ has the same trace space on $\partial\Omega$ as $W^{k,1}(\Omega; V)$.

Here, given a Banach space $(\mathcal{X}(\partial\Omega), \|\cdot\|_{\mathcal{X}(\partial\Omega)})$ with $\mathcal{X}(\partial\Omega) \subset L^1_{\text{loc}}(\partial\Omega; V)$, we say that $W^{k,1}(\Omega; V)$ has trace space $\mathcal{X}(\partial\Omega; V)$ provided there exists a linear and bounded surjective trace operator $\text{Tr}_{\partial\Omega} : W^{k,1}(\Omega; V) \rightarrow \mathcal{X}(\partial\Omega; V)$ (and so satisfies $\text{Tr}_{\partial\Omega}(u) = u$ \mathcal{H}^{n-1} -a.e. on $\partial\Omega$ for all $u \in C(\overline{\Omega}; V) \cap W^{k,1}(\Omega; V)$). Complementing Theorem 1.1, Theorem 1.2 does not specify the trace space of $W^{\mathbb{A},1}(\Omega)$ but rather asserts that it equals that of $W^{k,1}(\Omega; V)$. Also note that we could replace $W^{\mathbb{A},1}$ by $BV^{\mathbb{A}}$ in (b).

Theorem 1.2 is approached by an advancement and strengthening of a method employed in the first order case by the authors and BREIT [11]. The underlying key novelty of the proof of Theorem 1.2 is that for a \mathbb{C} -elliptic operator \mathbb{A} , the nullspace of \mathbb{A} consists of polynomials of a fixed degree and so, by the equivalence of all norms on finite dimensional spaces, ORNSTEIN's Non-Inequality becomes invisible on $\ker(\mathbb{A})$. Section 5 is devoted to the implementation of this strategy.

1.4. Korn's inequality without global CZ-estimates. Returning to the Korn-type estimate (1.8) for $1 < p < \infty$, the approach sketched in Section 1.2 works by extensions and applying global Calderón-Zygmund estimates on \mathbb{R}^n . Following the discussions in [23, 39] for the symmetric gradient case $\mathbb{A}u = \frac{1}{2}(Du + Du^\top)$, the natural geometric setup for such inequalities is given by *John domains* (see Section 2 for this terminology). However, John domains need not even be extension domains for $W^{k,p}$. Hence, in this situation, estimate (1.8) cannot be established by means of global Calderón-Zygmund estimates on \mathbb{R}^n . Interestingly, (1.8) still persists for John domains and \mathbb{C} -elliptic operators:

Theorem 1.3 (Korn for John). *Let $1 < p < \infty$ and \mathbb{A} be a differential operator of the form (1.1) with $k \in \mathbb{N}_{\geq 1}$. Then the following are equivalent:*

- (a) \mathbb{A} is \mathbb{C} -elliptic.
- (b) For all open and bounded John domains $\Omega \subset \mathbb{R}^n$ we have

$$\|u\|_{W^{k,p}(\Omega)} \simeq \|u\|_{L^p(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)}, \quad u \in W^{k,p}(\Omega; V).$$

Theorem 1.3 is established by a generalisation of a decomposition method introduced by RŮŽIČKA, SCHUMACHER and the first author [23] to the sharp class of operators for which the above Korn-type inequalities can hold at all. Theorem 1.3 is in the spirit of Theorem 1.2, however, note that John domains need not allow for a boundary trace operator; but if they do, Theorem 1.3 immediately implies the equality of the trace space of $W^{k,p}(\Omega; V)$ and that of the \mathbb{A} -Sobolev space $W^{\mathbb{A},p}(\Omega) := \{u \in L^p(\Omega; V) : \|u\|_{L^p(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)} < \infty\}$. Motivated by recent interest in Korn-type inequalities on more general space scales (cf. [10, 16, 23]), Theorem 1.3 is obtained in Section 6 as a special case of an A_p -weighted version,

cf. Theorem 6.9. By RUBIO DE FRANCIA extrapolation, the latter implies a variety of Korn-type inequalities, so e.g. on Orlicz- or Lorentz spaces; see Section 6.5.

1.5. Organisation of the paper. Section 2 collects the different notions of domains considered in the main part of the paper. As a technical novelty, Section 3 provides an improved representation formula a lá SMITH and KALAMAJSKA. This results in a family of Poincaré-type inequalities for John domains and Riesz potential-type inequalities which should be of independent interest but also display a crucial ingredient for the sequel. In Section 4, after gathering background facts on Besov spaces, we establish Theorem 1.1. Sections 5 and 6 are devoted to the proofs of Theorems 1.2 and 1.3, respectively. The appendix, Section 7 gathers the proofs of auxiliary results.

GENERAL NOTATION

By $\Omega \subset \mathbb{R}^n$ we understand an open set throughout, and the open ball of radius r centered at x_0 is denoted $B(x_0, r) := \{y \in \mathbb{R}^n : |x_0 - y| < r\}$. For $B = B(x_0, r)$ and $\sigma > 0$, we set $\sigma B := B(x_0, \sigma r)$. The n -dimensional Lebesgue and $(n - 1)$ -dimensional Hausdorff measures are denoted \mathcal{L}^n and \mathcal{H}^{n-1} , respectively; for brevity, we sometimes use $|A| := \mathcal{L}^n(A)$ and $d^{n-1} := d\mathcal{H}^{n-1}$. Given a finite dimensional real vector space E , we denote the finite positive Radon measures on Ω by $\mathcal{M}(\Omega)$, the finite E -valued Radon measures on Ω by $\mathcal{M}(\Omega; E)$ and, for $\mu \in \mathcal{M}(\Omega; E)$, $|\mu|(\Omega)$ its total variation. Given $\mu \in \mathcal{M}(\Omega)$ and a Borel subset U of Ω with $\mu(U) > 0$, we put for a μ -integrable map $f : \Omega \rightarrow E$

$$\int_U f \, d\mu := \frac{1}{\mu(U)} \int_U f \, d\mu,$$

and the choice of μ will be clear from the context. If $\mu = \mathcal{L}^n$, we also use the shorthand $(f)_U := \int_U f \, d\mathcal{L}^n$. The space of E -valued polynomials on \mathbb{R}^n of degree at most $m \in \mathbb{N}_0$ is denoted $\mathcal{P}_m(\mathbb{R}^n; E)$, and the homogeneous E -valued polynomials of degree m by $\mathcal{P}_m^h(\mathbb{R}^n; E)$; we also set $\mathcal{P}_{-m}(\mathbb{R}^n; E) = \{0\}$ for $m \in \mathbb{N}$ and $\mathcal{P}(\mathbb{R}^n; E) := \bigcup_{m \in \mathbb{N}_0} \mathcal{P}_m(\mathbb{R}^n; E)$. The symbol $\odot^m(\mathbb{R}^n; E)$ denotes the symmetric m -multilinear maps from \mathbb{R}^n to E . By $c, C > 0$ we denote generic constants which might change from line to line and shall only be specified provided their precise value is required. As such, we write $a \simeq b$ if there exist $c, C > 0$ such that $ca \leq b \leq Ca$ and both c, C do not depend in any essential way on a and b . We also use $X \simeq Y$ for normed spaces to indicate that $X = Y$ with equivalence of norms, but no ambiguities will arise from this.

2. DOMAINS

In this section we collect the various geometric assumptions on domains Ω that arise throughout the main part of the paper.

Let $\Omega \subset \mathbb{R}^n$ be given. We say that Ω satisfies the *interior corkscrew condition* provided there exist $R > 0$ and $M > 1$ such that for all $x \in \partial\Omega$ and all $0 < r < R$ there exists $y \in \Omega$ such that

$$|x - y| < r \quad \text{and} \quad B\left(y, \frac{r}{M}\right) \subset \Omega.$$

Likewise, Ω satisfies the *exterior corkscrew condition* provided $\mathbb{R}^n \setminus \Omega$ satisfies the interior corkscrew condition.

Next, for $x_1, x_2 \in \Omega$, we say that a sequence of balls $B_1, \dots, B_K \subset \Omega$ is a *Harnack chain from x_1 to x_2 of length $K \in \mathbb{N}$* provided $x_1 \in B_1$, $x_2 \in B_K$ and for all $1 \leq j \leq K - 1$ there holds $B_j \cap B_{j+1} \neq \emptyset$ and their radii satisfy $M^{-1}r(B_j) < \text{dist}(B_j, \partial\Omega) < Mr(B_j)$ for all $1 \leq j \leq K$. The set Ω is then said to satisfy the

interior Harnack chain condition if, whenever $\varepsilon > 0$ and $x_1, x_2 \in \Omega \cap B(\xi, \frac{r}{4})$ (for some $\xi \in \partial\Omega$ and $0 < r < R$) satisfy $\text{dist}(x_j, \partial\Omega) > \varepsilon$, $j \in \{1, 2\}$, together with $|x_1 - x_2| < 2^l \varepsilon$ for some $l \in \mathbb{N}$, then there exists a Harnack chain B_1, \dots, B_K from x_1 to x_2 of length $K \leq Ml$ such that $\text{diam}(B_j) \geq \frac{1}{M} \min\{\text{dist}(x_1, \partial\Omega), \text{dist}(x_2, \partial\Omega)\}$ for all $j = 1, \dots, K$.

Definition 2.1 (NTA and NTA_{n-1} -domains). *An open and bounded set $\Omega \subset \mathbb{R}^n$ is non-tangentially accessible (or NTA for brevity) provided Ω satisfies the interior, the exterior corkscrew and the interior Harnack chain condition, we refer to $R > 0$ and $M > 1$ from above as the NTA-parameters of Ω . Moreover, Ω is said to have $(n-1)$ -Ahlfors regular boundary $\partial\Omega$ if there exist $R > 0$ and $L > 0$ such that for all $x \in \partial\Omega$ and $0 < r < R$ there holds*

$$\frac{1}{L} r^{n-1} \leq \mathcal{H}^{n-1}(\partial\Omega \cap B(x, r)) \leq L r^{n-1}.$$

If $\Omega \subset \mathbb{R}^n$ is NTA and has $(n-1)$ -Ahlfors regular boundary, we say that Ω is NTA_{n-1} .

See [2, 33, 38] for more information on NTA domains. Now let $\gamma \subset \mathbb{R}^n$ be a rectifiable path with endpoints a and b and length $|\gamma|$. Assuming that $\gamma : [0, |\gamma|] \rightarrow \mathbb{R}^n$ is parametrised by arclength, we define the α -cigar with core γ and parameter $\alpha > 0$ by

$$\text{cig}(\gamma, \alpha) := \bigcup_{t \in [0, |\gamma|]} B\left(\gamma(t), \frac{1}{\alpha} \min\{t, |\gamma| - t\}\right).$$

Definition 2.2 (John domains). *An open and bounded set $\Omega \subset \mathbb{R}^n$ is called an α -John domain, $\alpha > 0$, if every pair of distinct points $a, b \in \Omega$ can be joined by a rectifiable path γ such that $\text{cig}(\gamma, \alpha) \subset \Omega$. If the constant α is not important, we just say that Ω is a John domain.*

Introduced by JOHN [40] and named after him by MARTIO & SARVAS [48], John domains include sets with fractal boundary such as the Koch snowflake or slit domains. Moreover, they can be decomposed into a suitable set of balls or cubes that satisfy a certain chain condition. In particular, by [23, Thm. 3.8] every bounded John domain also satisfies the *emanating chain condition* in the sense of the following definition.

Definition 2.3 (Emanating chain condition). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded set and let $\sigma_1 > 1$, $\sigma_2 \geq 1$. Then we say that Ω satisfies the emanating chain condition with constants σ_1 and σ_2 if there exists a covering $\mathcal{W} = \{W_i : i \in \mathbb{N}_0\}$ of Ω consisting of open balls (or cubes) such that:*

- (C1) *We have $\sigma_1 W \subset \Omega$ for all $W \in \mathcal{W}$ and $\sum_{W \in \mathcal{W}} \mathbb{1}_{\sigma_1 W} \leq \sigma_2 \mathbb{1}_\Omega$ on \mathbb{R}^n .*
- (C2) *For every $W_i \in \mathcal{W}$ there exists a chain of $W_{i,0}, W_{i,1}, \dots, W_{i,m_i}$ (pairwise different) from \mathcal{W} such that $W_{i,0} = W_i$, $W_{i,m_i} = W_0$, and $W_{i,l_1} \subset \sigma_2 W_{i,l_2}$ for $0 \leq l_1 \leq l_2 \leq m_i$. Moreover, $W_{i,l} \cap W_{i,l+1}$, $0 \leq l < m_i$, contains a ball $B_{i,l}$ such that $W_{i,l} \cup W_{i,l+1} \subset \sigma_2 B_{i,l}$. The chain $W_{i,0}, \dots, W_{i,m_i}$ is called chain emanating from W_i . The number $m_i \in \mathbb{N}_0$ is called the length of this chain.*
- (C3) *The set $\{i \in \mathbb{N}_0 : W_i \cap K \neq \emptyset\}$ is finite for every compact subset $K \subset \Omega$.*

The family \mathcal{W} is called the *chain-covering* of Ω . The ball (or cube) W_0 is called the *central ball* (or cube), since every chain ends in W_0 .

Clearly, all sets satisfying the emanating chain condition are connected. Moreover, if Ω satisfies the emanating chain condition with $\sigma_1 > 1$ and $\sigma_2 \geq 1$, we




FIGURE 1. The emanating chain condition for an exemplary non-extension domain $\Omega \subset \mathbb{R}^2$ with slits, illustrating the key property in (C2): In any chain $W_{i,0} = W_i, \dots, W_{i,m_i} = W_0$, any $W_{i,l}$ might be blown up by σ_2 such that $\sigma_2 W_{i,l}$ contains all $W_{i,j}$, $0 \leq j \leq l$.

have

$$(2.1) \quad \text{diam}(\Omega) \leq \sigma_2 \text{diam}(W_0),$$

since every $x \in \Omega$ is contained in some $W \in \mathcal{W}$ that can be connected with W_0 by a chain satisfying (C2) and so $W \subset \sigma_2 W_0$. Moreover, note that whenever an open and bounded set satisfies the emanating chain condition with $\sigma_1, \sigma_2 \geq 1$, then by [23, Thm. 3.8] it automatically satisfies the emanating chain condition with $\sigma_1 > 1$ and $\sigma_2 \geq 1$ (for some possibly different chain covering); this is why we directly restrict ourselves to $\sigma_1 > 1$ in Definition 2.3.

Remark 2.4 (Choice of overlap balls). By [23, Rem. 3.15], if $\Omega \subset \mathbb{R}^n$ satisfies the requirements of Definition 2.3, the balls $B_{i,l}$ as in (C2) can be chosen to belong to a family \mathcal{B} of balls such that $\sum_{B \in \mathcal{B}} \mathbb{1}_B \leq \sigma_2 \mathbb{1}_\Omega$ as an estimate on \mathbb{R}^n .

Following BOMAN [6] and IWANIEC & NOLDER [37], domains merely satisfying (C1) and (C2) are referred to as *Boman (chain) domains*; also see HURRI [35]. By a result due to BUCKLEY, KOSKELA & LU [14], an open and bounded domain is a Boman chain domain if and only if it is John; moreover, by [23], any open and bounded John domain satisfies the emanating chain condition. In total, for open and bounded sets $\Omega \subset \mathbb{R}^n$ these notions of sets are connected as follows (see, e.g., [2, Sec. 3.7] and [23, Thm. 3.8]):

$$(2.2) \quad \boxed{\text{NTA}_{n-1} \implies \text{NTA} \implies \text{John} = \text{Emanating Chain Condition} = \text{Boman}}$$

3. THE SHARP REPRESENTATION THEOREM AND POINCARÉ-TYPE INEQUALITIES

In this section we derive a new representation formula in terms of $\mathbb{A}u$. From this we conclude Poincaré-type estimates for balls and John domains which should be of independent interest but also play a pivotal role in the proof of Theorems 1.1–1.3.

3.1. The representation formula. The natural framework for the results of this and the subsequent sections is given by the following variants of Sobolev- and BV-spaces adapted to differential operators \mathbb{A} of the form (1.1): Given $1 \leq p < \infty$ and an open set $\Omega \subset \mathbb{R}^n$, we define

$$(3.1) \quad \begin{aligned} W^{\mathbb{A},p}(\Omega) &:= \{u \in L^p(\Omega; V) : \|u\|_{W^{\mathbb{A},p}(\Omega)} := \|u\|_{L^p(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)} < \infty\}, \\ BV^{\mathbb{A}}(\Omega) &:= \{u \in L^1(\Omega; V) : \|u\|_{BV^{\mathbb{A}}(\Omega)} := \|u\|_{L^1(\Omega)} + |\mathbb{A}u|(\Omega) < \infty\}, \end{aligned}$$

where $\|\mathbb{A}u\|_{L^p(\Omega)} < \infty$ or $|\mathbb{A}u|(\Omega) < \infty$ have to be interpreted in the sense that the distributional differential $\mathbb{A}u$ can be represented by an $L^p(\Omega; W)$ -map or a finite W -valued Radon measure, respectively. Moreover, we set $\widetilde{W}^{\mathbb{A},p}(\Omega) := W^{\mathbb{A},p}(\Omega) \cap W^{k-1,p}(\Omega; V)$ (in [30] also denoted $V^{\mathbb{A},p}$) to be endowed with norm $\|u\|_{\widetilde{W}^{\mathbb{A},p}(\Omega)} := \|u\|_{W^{k-1,p}(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)}$. All of these spaces are Banach, and it is routine to check that $C^\infty(\Omega; V) \cap \widetilde{W}^{\mathbb{A},p}(\Omega)$ is dense in $\widetilde{W}^{\mathbb{A},p}(\Omega)$ and, if $\partial\Omega$ is sufficiently regular, we even have density of $C^\infty(\overline{\Omega}; V) \cap \widetilde{W}^{\mathbb{A},p}(\Omega)$. For future reference, note that such density properties cannot be expected for the norm on $BV^{\mathbb{A}}$ (cf. AMBROSIO [3, Chpt. 3] et al. for $\mathbb{A} = D$), and the correct substitute here is the \mathbb{A} -strict metric defined by

$$(3.2) \quad d_{\mathbb{A}}(u, v) := \|u - v\|_{L^1(\Omega)} + \left| |\mathbb{A}u|(\Omega) - |\mathbb{A}v|(\Omega) \right|, \quad u, v \in BV^{\mathbb{A}}(\Omega).$$

(Non-)examples of \mathbb{C} -elliptic operators are as follows; other relevant instances are gathered in [67, Sec. 6.4]:

Example 3.1. For each $k, n \in \mathbb{N}$ and $N \in \mathbb{N}$, the k -th order gradients D^k acting on $u: \mathbb{R}^n \rightarrow \mathbb{R}^N$ are \mathbb{C} -elliptic. For $n \in \mathbb{N}$ and maps $u: \mathbb{R}^n \rightarrow \mathbb{R}^n$, the *symmetric gradient operator* $\varepsilon(u) := \frac{1}{2}(Du + Du^\top)$ is \mathbb{C} -elliptic for all $n \geq 1$, see [11, Sec. 2.2]. For $n \geq 3$, the *trace-free symmetric gradient* $\varepsilon^D u := \varepsilon(u) - \frac{1}{n} \operatorname{div}(u) E_{n \times n}$ is equally \mathbb{C} -elliptic (again, see [11, Sec. 2.2]). It is easy to see that the composition of two \mathbb{C} -elliptic operators is again \mathbb{C} -elliptic; an example that occurs frequently (so, e.g., in the regularity theory for elasticity and plasticity, is $\mathbb{A} = D\varepsilon^D$, see FUCHS & SEREGIN [28]).

Example 3.2. The trace-free symmetric gradient is not \mathbb{C} -elliptic for $n = 2$. This can be seen by the fact (following from Proposition 3.3 below) that \mathbb{C} -elliptic operators have finite dimensional nullspace on connected sets. Elements of $\ker(\varepsilon^D)$ in $n = 2$ have a one-to-one-correspondence with holomorphic functions on \mathbb{C} , hence ε^D is not \mathbb{C} -elliptic, see [11, Sec. 2.2]. On the other hand, if $n \geq 2$ and \mathbb{A} is an elliptic operator on \mathbb{R}^n from V to W , then the \mathbb{A} -Laplacean $\Delta_{\mathbb{A}} := \mathbb{A}^* \mathbb{A}$ is *not* \mathbb{C} -elliptic. Indeed, in this situation $\Delta_{\mathbb{A}}$ is elliptic as an operator on \mathbb{R}^n from V to V , and hence does not even satisfy VAN SCHAFTINGEN's cancellation condition:

$$\bigcap_{\xi \in \mathbb{R}^n \setminus \{0\}} \Delta_{\mathbb{A}}[\xi](V) = V \neq \{0\}.$$

As \mathbb{C} -ellipticity implies ellipticity and cancellation (cf. [30, Lem. 3.2], [31, Prop. 3.1]) for $n \geq 2$, the \mathbb{A} -Laplacean $\Delta_{\mathbb{A}}$ is not \mathbb{C} -elliptic.

Towards Proposition 3.3, we need to revise a decomposition as established in the pioneering works of SMITH [56] and KAŁAMAJSKA [42], cf. (3.7) below. Let $\Omega \subset \mathbb{R}^n$ be a domain that is star-shaped with respect to an open ball B and satisfies $\operatorname{diam}(\Omega) \leq c_0 \operatorname{diam}(B)$. The averaged Taylor polynomial \mathbb{T}_m^B of order m then is defined as in [49, Chpt. 1.1.10] and [12, Chpt. 4]: Choose $\omega \in C_c^\infty(B)$ with $\int_B \omega(y) dy = 1$ and then define for $u \in L^1_{\text{loc}}(\Omega; V)$:

$$(3.3) \quad \mathbb{T}_m^B u(x) := \int_B u(y) \sum_{|\alpha| \leq m} \frac{(-1)^{|\alpha|}}{\alpha!} \partial_y^\alpha (\omega(y)(x - y)^\alpha) dy.$$

The averaged Taylor polynomial has good approximation properties and commutes in some sense with derivatives. In particular, we have

$$(3.4) \quad \mathbb{A} \mathbb{T}_{m-1}^B u = \mathbb{T}_{m-k-1}^B \mathbb{A} u.$$

Moreover, it follows from [49, Chpt. 1.1.10, Thm. 1] that for all $x \in \Omega$ there holds

$$(3.5) \quad u(x) = \mathbb{T}_{m-1}^B u(x) + \sum_{|\alpha|=m} \int_{\Omega} K_{\alpha,B}(x,y) \partial^{\alpha} u(y) \, dy, \quad \text{where}$$

$$K_{\alpha,B}(x,y) = \frac{(-1)^m m}{\alpha!} \frac{(y-x)^{\alpha}}{|x-y|^n} \int_{|x-y|}^{\infty} \omega\left(x + t \frac{y-x}{|y-x|}\right) t^{n-1} \, dt.$$

As established in [42], the \mathbb{C} -ellipticity in combination with the Hilbert Nullstellensatz [42, Lem. 4] implies the existence of $m \in \mathbb{N}$ and a $\odot^m(\mathbb{R}^n; V)$ -valued differential operator \mathbb{L} of order $(m-k)$ such that $D^m = \mathbb{L}\mathbb{A}$. The constant m is not given explicitly, but at least $\ker(\mathbb{A}) \subset \mathcal{P}_{m-1}(\mathbb{R}^n; V)$, so the *polynomial degree* $\deg_{\mathcal{P}}(\mathbb{A})$ of \mathbb{A} satisfies

$$(3.6) \quad m \geq \deg_{\mathcal{P}}(\mathbb{A}) := \min\{j \in \mathbb{N}_0 : \ker(\mathbb{A}) \subset \mathcal{P}_{j-1}(\mathbb{R}^n; V)\} \geq k.$$

For future reference, we note that $\deg_{\mathcal{P}}(\mathbb{A}) < \infty$ not only follows but is in fact equivalent to \mathbb{C} -ellipticity, cf. [30, Prop. 3.1] and [56]. Now, $D^m = \mathbb{L}\mathbb{A}$ and integrating by parts $(m-k)$ -times yields for $u \in C^{\infty}(\bar{\Omega}; V)$

$$(3.7) \quad u(x) = \mathbb{T}_{m-1}^B u(x) + \int_{\Omega} \tilde{\mathfrak{K}}_{\mathbb{A}}^B(x,y) \mathbb{A}u(y) \, dy \quad \text{for all } x \in \Omega,$$

cf. [42, Theorem 4 (v)], where the kernel $\tilde{\mathfrak{K}}_{\mathbb{A}}^B : (\Omega \times \Omega) \setminus \Delta \rightarrow \mathcal{L}(W; V)$ is smooth off the diagonal Δ , satisfies $\tilde{\mathfrak{K}}_{\mathbb{A}}^B(x, \cdot) = 0$ near $\partial\Omega$ for each $x \in \Omega$ and, with $C = C(c_0, \mathbb{A}) > 0$

$$(3.8) \quad |\partial_x^{\alpha} \partial_y^{\beta} \tilde{\mathfrak{K}}_{\mathbb{A}}^B(x,y)| \leq C |x-y|^{k-n-|\alpha|-|\beta|} \quad \text{for all } x, y \in \Omega, x \neq y,$$

for all $\alpha, \beta \in \mathbb{N}_0^n$ with $|\alpha| + |\beta| \leq k$.

In general, the polynomial $\mathbb{T}_{m-1}^B u$ in (3.7) does not belong to the nullspace of \mathbb{A} . As a simple, yet effective improvement, we now modify this representation formula in such a way that we can replace \mathbb{T}_{m-1}^B by a projection $\Pi_{\mathbb{A}}^B$ to $\ker(\mathbb{A})$, thereby yielding a quick argument for Riesz- and Poincaré-type estimates in Sections 3.2 and 4.

Proposition 3.3 (Representation formula). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain which is star-shaped with respect to an open ball $B \subset \Omega$ with $\text{diam } \Omega \leq c_0 \text{ diam } B$. Let \mathbb{A} be a k -th order \mathbb{C} -elliptic differential operator of the form (1.1). Then there exists an integral kernel $\mathfrak{K}_{\mathbb{A}}^B : (\Omega \times \Omega) \setminus \Delta \rightarrow \mathcal{L}(W; V)$ and a linear projection $\Pi_{\mathbb{A}}^B : L^1(\Omega; V) \rightarrow \ker(\mathbb{A})$ such that the following hold:*

(K1) $\mathfrak{K}_{\mathbb{A}}^B$ is smooth off the diagonal, and for $x, y \in \Omega$, $x \neq y$ and $|\alpha| + |\beta| \leq k$ we have

$$|\partial_x^{\alpha} \partial_y^{\beta} \mathfrak{K}_{\mathbb{A}}^B(x,y)| \leq C |x-y|^{k-n-|\alpha|-|\beta|}$$

with a constant $C = C(\mathbb{A}, c_0) > 0$.

(K2) $\mathfrak{K}_{\mathbb{A}}^B(x, \cdot)$ is zero near $\partial\Omega$ for all $x \in \Omega$, i.e., $\text{spt}(\mathfrak{K}_{\mathbb{A}}^B(x, \cdot)) \Subset \Omega$.

(K3) For any $u \in C^{\infty}(\Omega; V)$ and $x \in \Omega$ there holds

$$(3.9) \quad u(x) = \Pi_{\mathbb{A}}^B u(x) + \int_{\Omega} \mathfrak{K}_{\mathbb{A}}^B(x,y) \mathbb{A}u(y) \, dy.$$

(K4) $\Pi_{\mathbb{A}}^B : \mathcal{P}_{\ell}(\mathbb{R}^n; V) \rightarrow \mathcal{P}_{\ell}(\mathbb{R}^n; V) \cap \ker(\mathbb{A})$ for every ℓ with $0 \leq \ell < \deg_{\mathcal{P}}(\mathbb{A})$.

Proof. Using translation and dilation we can assume without loss of generality that B is the unit ball $B(0, 1)$.

Let $m \in \mathbb{N}_0$ and $\tilde{\mathfrak{K}}_{\mathbb{A}}^B$ be such that (3.7) holds. In particular, we have $m \geq \deg_{\mathcal{P}}(\mathbb{A})$ and $\ker(\mathbb{A}) \subset \mathcal{P}_{m-1}(\mathbb{R}^n; V)$. Define $\mathcal{Z}_\ell := \ker(\mathbb{A}) \cap \mathcal{P}_\ell^h(\mathbb{R}^n; V)$ for $\ell = 0, \dots, m-1$. We claim that

$$(3.10) \quad \ker(\mathbb{A}) = \bigoplus_{\ell=0}^{m-1} \mathcal{Z}_\ell$$

as a direct sum. For this, let $q \in \ker(\mathbb{A})$. Then $q = \sum_{\ell=0}^{m-1} q_\ell$ with $q_\ell \in \mathcal{P}_\ell^h(\mathbb{R}^n; V)$. Since $0 = \mathbb{A}q = \sum_{\ell=0}^{m-1} \mathbb{A}q_\ell$ and $\mathbb{A}q_\ell \in \mathcal{P}_{\ell-k}^h(\mathbb{R}^n; W)$ by the homogeneity of \mathbb{A} , we obtain $q_\ell \in \ker(\mathbb{A})$ for $\ell = 0, \dots, m-1$. This proves the decomposition (3.10).

For $\ell = 0, \dots, m-1$, we then choose a linear subspace $\mathcal{W}_\ell \subset \mathcal{P}_\ell^h(\mathbb{R}^n; V)$ such that $\mathcal{P}_\ell^h(\mathbb{R}^n; V) = \mathcal{Z}_\ell \oplus \mathcal{W}_\ell$. Then we have

$$(3.11) \quad \mathcal{P}_{m-1}(\mathbb{R}^n; V) = \bigoplus_{\ell=0}^{m-1} \mathcal{Z}_\ell \oplus \bigoplus_{\ell=0}^{m-1} \mathcal{W}_\ell =: \ker(\mathbb{A}) + \mathcal{W}.$$

Let ψ_1, \dots, ψ_M denote a basis of $\mathcal{P}_{m-1}(\mathbb{R}^n; V)$ such that for each $\ell = 0, \dots, m-1$, $\psi_{j_\ell}, \dots, \psi_{j_{\ell+1}-1}$ is a basis of \mathcal{Z}_ℓ and $\psi_{j_m}, \dots, \psi_M$ is a basis of \mathcal{W} ; here, $j_0 = 1$. Now, let $\psi_1^*, \dots, \psi_M^*$ be a dual basis with respect to the $L^2(B; V)$ -inner product $\langle \cdot, \cdot \rangle_{L^2(B)}$, i.e., $\langle \psi_l, \psi_j^* \rangle_{L^2(B)} = \delta_{l,j}$ for all $j, l = 1, \dots, M$.

Based on (3.11), we define the projection $\tilde{\Pi}_{\mathbb{A}}: \mathcal{P}_{m-1}(\mathbb{R}^n; V) \rightarrow \ker(\mathbb{A})$ by

$$\tilde{\Pi}_{\mathbb{A}}(q) := \sum_{j=1}^{j_m-1} \langle q, \psi_j^* \rangle_{L^2(B)} \psi_j.$$

We define $\Pi_{\mathbb{A}}^B := \tilde{\Pi}_{\mathbb{A}} \mathbb{T}_{m-1}^B$; since \mathbb{T}_{m-1}^B is a projection to $\mathcal{P}_{m-1}(\mathbb{R}^n; V)$, $\Pi_{\mathbb{A}}^B$ is a projection to $\ker(\mathbb{A})$. Moreover, by construction, $\tilde{\Pi}_{\mathbb{A}}$ maps $\mathcal{P}_\ell^h(\mathbb{R}^n; V)$ onto $\mathcal{P}_\ell^h(\mathbb{R}^n; V)$ for $\ell = 0, \dots, m-1$, thus $\Pi_{\mathbb{A}}^B$ also maps $\mathcal{P}_\ell^h(\mathbb{R}^n; V)$ onto $\mathcal{P}_\ell^h(\mathbb{R}^n; V)$ for $\ell = 0, \dots, m-1$. This proves (K4). Moreover, for all $u \in C^\infty(\bar{\Omega}; V)$ we have

$$(3.12) \quad \begin{aligned} u - \Pi_{\mathbb{A}}^B u &= u - \tilde{\Pi}_{\mathbb{A}} \mathbb{T}_{m-1}^B u \\ &= (u - \mathbb{T}_{m-1}^B u) + (\text{Id} - \tilde{\Pi}_{\mathbb{A}}) \mathbb{T}_{m-1}^B u. \end{aligned}$$

Note that $\langle \mathbb{A} \cdot, \mathbb{A} \cdot \rangle_{L^2(B)}$ is an inner product on \mathcal{W} as $\mathcal{P}_{m-1}(\mathbb{R}^n; V) = \ker(\mathbb{A}) \oplus \mathcal{W}$. Thus, for each ψ_ℓ^* with $j_m \leq \ell \leq M$ we find $\xi_\ell \in \mathcal{W}$ such that

$$\langle w, \psi_\ell^* \rangle_{L^2(B)} = \langle \mathbb{A}w, \mathbb{A}\xi_\ell \rangle_{L^2(B)} \quad \text{for all } w \in \mathcal{W} \text{ and } j_m \leq \ell \leq M.$$

If $z \in \ker(\mathbb{A})$, then $\langle z, \psi_\ell^* \rangle_{L^2(B)} = 0 = \langle \mathbb{A}z, \mathbb{A}\xi_\ell \rangle_{L^2(B)}$ for all $j_m \leq \ell \leq M$. Thus,

$$(3.13) \quad \langle q, \psi_\ell^* \rangle_{L^2(B)} = \langle \mathbb{A}q, \mathbb{A}\xi_\ell \rangle_{L^2(B)} \quad \text{for all } q \in \mathcal{P}_{m-1}(\mathbb{R}^n; V) \text{ and } j_m \leq \ell \leq M.$$

In conclusion,

$$(3.14) \quad \begin{aligned} (\text{Id} - \tilde{\Pi}_{\mathbb{A}}) \mathbb{T}_{m-1}^B u &= \sum_{j=j_m}^M \langle \mathbb{T}_{m-1}^B u, \psi_j^* \rangle_{L^2(B)} \psi_j \\ &= \sum_{j=j_m}^M \langle \mathbb{A} \mathbb{T}_{m-1}^B u, \mathbb{A} \xi_j \rangle_{L^2(B)} \psi_j \stackrel{(3.4)}{=} \sum_{j=j_m}^M \langle \mathbb{T}_{m-1-k}^B \mathbb{A}u, \mathbb{A} \xi_j \rangle_{L^2(B)} \psi_j. \end{aligned}$$

Hence, by the definition (3.3) of \mathbb{T}_{m-1}^B , we obtain for all $x \in \Omega$

$$\begin{aligned}
(3.15) \quad & ((\text{Id} - \tilde{\Pi}_{\mathbb{A}})\mathbb{T}_{m-1}^B u)(x) \\
&= \int_B \int_B \sum_{\substack{|\alpha| \leq m-k-1 \\ j_m \leq j \leq M}} \frac{(-1)^{|\alpha|}}{\alpha!} \partial_z^\alpha (\omega(z)(y-z)^\alpha) \psi_j(x) (\mathbb{A} \xi_j(y))^\top dy \mathbb{A} u(z) dz \\
&=: \int_B \bar{\mathfrak{K}}_{\mathbb{A}}^B(x, z) \mathbb{A} u(z) dz,
\end{aligned}$$

where $\bar{\mathfrak{K}}_{\mathbb{A}}^B$ is defined in the obvious manner. Recalling (3.7), put $\mathfrak{K}_{\mathbb{A}}^B := \bar{\mathfrak{K}}_{\mathbb{A}}^B + \tilde{\mathfrak{K}}_{\mathbb{A}}^B$. Then by (3.7), (3.12) and (3.15), (3.9) follows. Note that $\bar{\mathfrak{K}}_{\mathbb{A}}^B(x, \cdot)$ is supported on $\text{spt}(\omega) \Subset B \subset \Omega$ and is therefore zero near $\partial\Omega$. On the other hand, by (3.7)ff. we obtain that $\tilde{\mathfrak{K}}_{\mathbb{A}}^B(x, \cdot)$ is zero near $\partial\Omega$. This proves that $\mathfrak{K}_{\mathbb{A}}^B(x, \cdot)$ is zero near $\partial\Omega$ and so (K2) follows. It remains to establish the estimates for $\mathfrak{K}_{\mathbb{A}}^B$ in (K1); however, the estimates for $\tilde{\mathfrak{K}}_{\mathbb{A}}^B$ are much less singular and follow easily using $\text{diam}(\Omega) \leq c_0 \text{diam}(B)$ and our simplifying assumption $0 \in B$. The proof is complete. \square

Remark 3.4. It is possible to extend (K4) of Proposition 3.3 to all $0 \leq \ell < m$ for any fixed $m \geq \deg_{\varphi}(\mathbb{A})$. For this one has to start with (3.7) with a possible enlarged m . Moreover, if B in Proposition 3.3 is centered at zero or Ω at least contains zero, then it is possible to construct $\Pi_{\mathbb{A}}^B$ such $\Pi_{\mathbb{A}}^B: \mathcal{P}_{\ell}^h(\mathbb{R}^n; V) \rightarrow \mathcal{P}_{\ell}^h(\mathbb{R}^n; V) \cap \ker(\mathbb{A})$ for every ℓ with $0 \leq \ell < \deg_{\varphi} \mathbb{A}$, where \mathcal{P}_{ℓ}^h is the linear space of homogeneous polynomials of order ℓ . Note that not increasing the degree of polynomials when projecting *cannot* be improved to the commuting-type relation $\partial^\alpha \text{proj}_{\ker(\mathbb{A})} u = \text{proj}_{\partial^\alpha \ker(\mathbb{A})}(\partial^\alpha u)$ for $u \in \mathcal{P}_{m-1}(\mathbb{R}^n; V)$ for suitable projections onto $\ker(\mathbb{A})$ or $\partial^\alpha \ker(\mathbb{A})$, respectively.

This can be seen by means of the symmetric gradient $\mathbb{A} = \varepsilon$ (cf. Example 3.1) in $n = 2$ dimensions. In this case, the nullspace of \mathbb{A} is given by the three-dimensional space of *rigid deformations*

$$\ker(\mathbb{A}) = \left\{ (x, y) \mapsto \begin{pmatrix} -ay + b \\ ax + c \end{pmatrix} : a, b, c \in \mathbb{R} \right\},$$

cf. RESHETNYAK [55]. Denote $p(x, y) := (0, x)^\top$. Should the commuting relation $\partial^\alpha \text{proj}_{\ker(\mathbb{A})} p = \text{proj}_{\partial^\alpha \ker(\mathbb{A})}(\partial^\alpha p)$ hold, then we have $\partial_x \text{proj}_{\ker(\mathbb{A})} p(x, y) = (0, 1)^\top$ and $\partial_y \text{proj}_{\ker(\mathbb{A})} p(x, y) = (0, 0)^\top$. Thus, $\text{proj}_{\ker(\mathbb{A})} p(x, y) = (b, x + c)^\top$ for certain $b, c \in \mathbb{R}$, but since $(x, y) \mapsto (b, x + c)^\top$ does not belong to $\ker(\mathbb{A})$, such a projection $\text{proj}_{\ker(\mathbb{A})}$ onto $\ker(\mathbb{A})$ cannot exist.

Lemma 3.5. *Under the assumptions of Proposition 3.3, $\Pi_{\mathbb{A}}^B$ extends to a bounded linear operator $\Pi_{\mathbb{A}}^B: L^1(\Omega; V) \rightarrow \ker(\mathbb{A})$. Moreover, for all $1 \leq p < \infty$ and all $u \in L^1(\Omega; V)$ we have, with $r(B)$ denoting the radius of B ,*

$$\begin{aligned}
(3.16) \quad & \left(\int_{\Omega} |\Pi_{\mathbb{A}}^B u|^p dx \right)^{\frac{1}{p}} \leq \|\Pi_{\mathbb{A}}^B u\|_{L^\infty(\Omega)} \leq C \int_B |u| dx \leq C \left(\int_B |u|^p dx \right)^{\frac{1}{p}}, \\
& \frac{1}{C} \left(\sum_{|\alpha| \leq k} r(B)^{|\alpha|p} \int_B |\partial^\alpha \Pi_{\mathbb{A}}^B u|^p dx \right)^{\frac{1}{p}} \leq \int_B |\Pi_{\mathbb{A}}^B u| dx \leq C \int_B |u| dx,
\end{aligned}$$

with a constant $C = C(\mathbb{A}, c_0) > 0$.

Proof. Since $\Pi_{\mathbb{A}}^B = \tilde{\Pi}_{\mathbb{A}} \mathbb{T}_{m-1}^B$ is a projection to a finite dimensional vector space, this follows immediately by inverse estimates for polynomials using $\text{diam}(\Omega) \leq c_0 \text{diam}(B)$ and scaling. \square

By the support and growth properties of the integral kernel in Proposition 3.3, convolving $u \in W^{\mathbb{A},p}(\Omega)$ or $u \in BV^{\mathbb{A}}(\Omega)$ with smooth bumps and passing to the limit yields

Corollary 3.6. *Under the assumptions of Proposition 3.3, let $u \in W^{\mathbb{A},p}(\Omega)$ for some $1 \leq p \leq \infty$. Then there holds for all $|\alpha| < k$*

$$(3.17) \quad \partial^\alpha u = \partial^\alpha \Pi_{\mathbb{A}}^B u + \int_{\Omega} (\partial_x^\alpha \mathfrak{K}_{\mathbb{A}}^B)(\cdot, y) \mathbb{A}u(y) \, dy \quad \mathcal{L}^n\text{-almost everywhere.}$$

Moreover, for all ℓ with $0 \leq \ell < k$ we have

$$(3.18) \quad |D^\ell(u - \Pi_{\mathbb{A}}^B u)| \leq C \int_{\Omega} |\cdot - y|^{-n+k-\ell} |\mathbb{A}u(y)| \, dy \quad \mathcal{L}^n\text{-almost everywhere.}$$

For $u \in BV^{\mathbb{A}}(\Omega)$, (3.17) and (3.18) remain valid upon replacing $\mathbb{A}u(y) \, dy$ by $d\mathbb{A}u(y)$ and $|\mathbb{A}u(y)| \, dy$ by $d|\mathbb{A}u|(y)$, respectively.

3.2. Poincaré-type inequalities. Throughout this section, let \mathbb{A} be a k -th order \mathbb{C} -elliptic differential operator of the form (1.1). The present section is devoted to the proof of the following Poincaré-type inequality for the vast class of John domains, announced in [21, Rem. 2.4], which shall turn out a crucial tool for the following sections.

Theorem 3.7 (Poincaré for John). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded α -John domain or an open and bounded domain satisfying the emanating chain condition with constants σ_1 and σ_2 and central ball B , respectively. Then for all $u \in W^{\mathbb{A},p}(\Omega)$ with $1 \leq p \leq \infty$ and $\ell \in \{0, \dots, k-1\}$ there holds*

$$(3.19) \quad \left(\int_{\Omega} |D^\ell(u - \Pi_{\mathbb{A}}^B u)|^p \, dx \right)^{\frac{1}{p}} \leq C \operatorname{diam}(\Omega)^{k-\ell} \left(\int_{\Omega} |\mathbb{A}u|^p \, dx \right)^{\frac{1}{p}}$$

and

$$(3.20) \quad \operatorname{diam}(\Omega)^\ell \left(\int_{\Omega} |D^\ell u|^p \, dx \right)^{\frac{1}{p}} \leq C \left(\int_{\Omega} |u|^p \, dx \right)^{\frac{1}{p}} + C \operatorname{diam}(\Omega)^k \left(\int_{\Omega} |\mathbb{A}u|^p \, dx \right)^{\frac{1}{p}},$$

where $C = C(p, \mathbb{A}, \sigma_1, \sigma_2)$. For $p = \infty$, one has to exchange $(\int_{\Omega} |\cdot|^p \, dx)^{\frac{1}{p}}$ by $\operatorname{ess\,sup}_{\Omega}$. For $p = 1$ and $u \in BV^{\mathbb{A}}(\Omega)$, one replaces $|\mathbb{A}u(y)| \, dy$ by $d|\mathbb{A}u|(y)$.

In view of Corollary 3.6, Theorem 3.7 is already available for domains which are star-shaped with respect to a ball; in particular, if Ω is a ball, we may choose $B = \Omega$, giving us back the Poincaré inequalities from [11, 30]. Moreover, the quantity on the left-hand side of (3.19) enjoys a best approximation property, a fact that we shall return to in slightly higher generality in Section 6.2.

Theorem 3.7 is a direct consequence of the following Riesz-type estimates, improving Corollary 3.6 to John domains. The proof is slightly inspired by [22, Section 8.2].

Proposition 3.8. *In the situation of Theorem 3.7, for all $\ell \in \{0, \dots, k-1\}$ and all $u \in W^{\mathbb{A},1}(\Omega)$ there holds with a constant $C = C(\mathbb{A}, \sigma_1, \sigma_2) > 0$*

$$(3.21) \quad |D^\ell(u - \Pi_{\mathbb{A}}^B u)| \leq C \int_{\Omega} |\cdot - y|^{-n+k-\ell} |\mathbb{A}u(y)| \, dy \quad \mathcal{L}^n\text{-almost everywhere.}$$

For $u \in BV^{\mathbb{A}}(\Omega)$, (3.21) remains valid upon replacing $|\mathbb{A}u(y)| \, dy$ by $d|\mathbb{A}u|(y)$.

Proof. Let $x \in \Omega$. Slightly abusing notation, let $m \in \mathbb{N}$ and let $(W_j)_{j=0}^m$ be an emanating chain of balls connecting x with the central ball B such that $W_0 = B$ and $x \in W_m$ in the sense of Definition 2.3, and we remark that each $\sigma_1 W_j$ is still contained in Ω . For the following argument, we may assume that $x \notin \sigma_1 W_j$ for all $j \in \{0, \dots, m-1\}$ as otherwise we may take the minimal j_0 with $x \in \sigma_1 W_{j_0}$, $x \notin \sigma_1 W_i$ for $0 \leq i < j_0$ and redefine m to be j_0 ; we will only use that $x \in \sigma_1 W_m$ (and not that $x \in W_m$) in the sequel.

Now consider the chain $\mathbb{W}_0 := W_0, \dots, \mathbb{W}_{m-1} := W_{m-1}$ and $\mathbb{W}_m := \sigma_1 W_m$. It is then easy to see that this chain satisfies the following modification of (C2): For any $0 \leq j \leq j' \leq m$ we have $\mathbb{W}_{j'} \subset \sigma_1 \sigma_2 \mathbb{W}_j$ and each $W_j \cap W_{j+1}$, $j \in \{0, \dots, m-1\}$ contains an open ball \mathbb{B}_j with $W_j \cup W_{j+1} \subset \sigma_1 \sigma_2 \mathbb{B}_j$. As a main consequence of this slightly modified construction, every $y \in \mathbb{W}_j$, $j \in \{0, \dots, m-1\}$ is at least $(\sigma_1 - 1)r(\mathbb{W}_j)$ distant apart from x , which follows from $x \in \sigma_1 W_m$ but $x \notin \sigma_1 W_j$ for any $0 \leq j \leq m-1$ by the above minimality of m . On the other hand, since $x \in \mathbb{W}_m \subset \sigma_1 \sigma_2 \mathbb{W}_j$ for any $j \in \{0, \dots, m\}$, we have $|x - y| \leq 2\sigma_1 \sigma_2 r(\mathbb{W}_j)$ for all $y \in W_j$. In conclusion, we have

(3.22)

$$(\sigma_1 - 1)r(\mathbb{W}_j) \leq |x - y| \leq 2\sigma_1 \sigma_2 r(\mathbb{W}_j) \quad \text{for all } 1 \leq j \leq m-1 \text{ and } y \in \mathbb{W}_j.$$

Define $\Pi_{\mathbb{A}}^{\mathbb{W}_j}$ and $\Pi_{\mathbb{A}}^{\mathbb{B}_j}$ as indicated after Theorem 3.7. Then, since $\Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} \Pi_{\mathbb{A}}^{\mathbb{B}_i} = \Pi_{\mathbb{A}}^{\mathbb{B}_i}$, $i \in \{j+1, j\}$,

$$\begin{aligned} |D^\ell(u - \Pi_{\mathbb{A}}^{\mathbb{W}_0} u)(x)| &\leq |D^\ell(u - \Pi_{\mathbb{A}}^{\mathbb{B}^{m-1}} u)(x)| + |D^\ell(\Pi_{\mathbb{A}}^{\mathbb{W}_0} - \Pi_{\mathbb{A}}^{\mathbb{B}_0} u)(x)| \\ &\quad + \sum_{j=0}^{m-2} (|D^\ell(\Pi_{\mathbb{A}}^{\mathbb{B}_{j+1}} u - \Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} u)(x)| + |D^\ell(\Pi_{\mathbb{A}}^{\mathbb{B}_j} u - \Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} u)(x)|) \\ &\leq |D^\ell(u - \Pi_{\mathbb{A}}^{\mathbb{B}^{m-1}} u)(x)| + |D^\ell(\Pi_{\mathbb{A}}^{\mathbb{W}_0} - \Pi_{\mathbb{A}}^{\mathbb{B}_0} u)(x)| \\ &\quad + \sum_{j=0}^{m-2} (|D^\ell \Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} (\Pi_{\mathbb{A}}^{\mathbb{B}_{j+1}} u - u)(x)| + |D^\ell \Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} (u - \Pi_{\mathbb{A}}^{\mathbb{B}_j} u)(x)|) \\ &=: \text{I} + \text{II} + \text{III}. \end{aligned}$$

Then, since $x \in \mathbb{W}_m \subset \sigma_1 \sigma_2 \mathbb{W}_{j+1}$ for all $0 \leq j \leq m-2$, we obtain by inverse estimates

$$\begin{aligned} \text{III} &\leq \sum_{j=0}^{m-2} \sum_{i \in \{j, j+1\}} \|D^\ell \Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} (\Pi_{\mathbb{A}}^{\mathbb{B}_i} u - u)\|_{L^\infty(\sigma_1 \sigma_2 \mathbb{W}_{j+1})} \\ &\leq c \sum_{j=0}^{m-2} \sum_{i \in \{j, j+1\}} r(\sigma_1 \sigma_2 \mathbb{W}_{j+1})^{-\ell} \times \\ &\quad \times \left(r(\sigma_1 \sigma_2 \mathbb{W}_{j+1})^\ell \int_{\sigma_1 \sigma_2 \mathbb{W}_{j+1}} |D^\ell \Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} (\Pi_{\mathbb{A}}^{\mathbb{B}_i} u - u)| \, dy \right) \\ &\stackrel{(3.16)_2}{\leq} c \sum_{j=0}^{m-2} \sum_{i \in \{j, j+1\}} r(\mathbb{W}_{j+1})^{-\ell} \left(\int_{\sigma_1 \sigma_2 \mathbb{W}_{j+1}} |\Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} (\Pi_{\mathbb{A}}^{\mathbb{B}_i} u - u)| \, dy \right) \\ &\leq c \sum_{j=0}^{m-2} \sum_{i \in \{j, j+1\}} r(\mathbb{W}_{j+1})^{-\ell} \left(\int_{\mathbb{W}_{j+1}} |\Pi_{\mathbb{A}}^{\mathbb{W}_{j+1}} (\Pi_{\mathbb{A}}^{\mathbb{B}_i} u - u)| \, dy \right) \\ &\leq c \sum_{j=0}^{m-2} \sum_{i \in \{j, j+1\}} r(\mathbb{W}_{j+1})^{-\ell} \left(\int_{\mathbb{W}_{j+1}} |u - \Pi_{\mathbb{A}}^{\mathbb{B}_i} u| \, dy \right) =: c \sum_{j=0}^{m-2} \text{III}_j, \end{aligned}$$

where $c = c(\sigma_1, \sigma_2, \mathbb{A}) > 0$ is a constant. Since each \mathbb{W}_{j+1} , $j \in \{0, \dots, m-2\}$, is starshaped with respect to both \mathbb{B}_{j+1} and \mathbb{B}_j , we obtain by use of Proposition 3.3 with a constant $c = c(\sigma_1, \sigma_2, \mathbb{A}) > 0$ and Fubini's theorem for $i \in \{j, j+1\}$

$$\begin{aligned} \text{III}_j &\leq c \sum_{i \in \{j, j+1\}} r(\mathbb{W}_{j+1})^{k-\ell-n} \int_{\mathbb{W}_{j+1}} |\mathbb{A}u(z)| \, dz \\ &\leq cr(\mathbb{W}_{j+1})^{k-n-\ell} \int_{\mathbb{W}_{j+1}} |\mathbb{A}u(z)| \, dz \stackrel{(3.22)}{\leq} c \int_{\mathbb{W}_{j+1}} |x-z|^{k-n-\ell} |\mathbb{A}u(z)| \, dz =: \text{IV}_j. \end{aligned}$$

On the other hand, the term I can clearly be bounded against IV_j with $j = m-1$, whereas the term II is estimated along the same lines as III, now using that \mathbb{W}_0 is starshaped with respect to \mathbb{B}_0 . In consequence, by the uniformly finite overlap of the balls W_0, \dots, W_m implied by (C2), we conclude (3.21). The proof is hereby complete. \square

4. PROOF OF THEOREM 1.1: THE TRACE ESTIMATE VIA RIESZ POTENTIALS

In this section we establish Theorem 1.1. To this end, we recall in Section 4.1 the scale of Besov spaces and prove Theorem 1.1 in Sections 4.2–4.4.

4.1. Besov spaces. For Theorem 1.1, we briefly recall the scale of Besov spaces and provide an instrumental characterisation by means of oscillations. Besov spaces can be introduced by different means, and we refer to TRIEBEL [61, 63] for comprehensive overviews. In what follows, let $s > 0$, $1 \leq p < \infty$, $1 \leq q < \infty$. Let $\varphi \in C_c^\infty(B(0, 2))$ satisfy $\mathbb{1}_{B(0,1)} \leq \varphi \leq \mathbb{1}_{B(0,2)}$ and put, for $j \in \mathbb{Z}$, $\varphi_j(\xi) := \varphi(2^{-j}\xi) - \varphi(2^{-j+1}\xi)$. We recall that the Besov space $\mathbb{B}_{p,q}^s(\mathbb{R}^n; V)$ consists of all $f \in \mathcal{S}'(\mathbb{R}^n; V)$ such that

$$\|f\|_{\mathbb{B}_{p,q}^s(\mathbb{R}^n)} := \left(\|(\varphi \widehat{f})^\vee\|_{L^p(\mathbb{R}^n)}^q + \sum_{j=1}^{\infty} 2^{jsq} \|(\varphi_j \widehat{f})^\vee\|_{L^p(\mathbb{R}^n)}^q \right)^{\frac{1}{q}} < \infty.$$

The homogeneous Besov space $\dot{\mathbb{B}}_{p,q}^s(\mathbb{R}^n; V)$ consists of all $f \in (\mathcal{S}'/\mathcal{P})(\mathbb{R}^n; V)$ with

$$(4.1) \quad \|f\|_{\dot{\mathbb{B}}_{p,q}^s(\mathbb{R}^n)} := \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|(\varphi_j \widehat{f})^\vee\|_{L^p(\mathbb{R}^n)}^q \right)^{\frac{1}{q}} < \infty.$$

Note that, whereas (4.1) does not give rise to a norm on $\mathcal{S}'(\mathbb{R}^n; V)$ but only on the quotient $(\mathcal{S}'/\mathcal{P})(\mathbb{R}^n; V)$, it does define a norm on $\mathcal{S}(\mathbb{R}^n; V)$. By BERGH & LÖFSTRÖM [5, p. 148] we have for all $s > 0$ and $1 \leq p, q < \infty$

$$(4.2) \quad \mathbb{B}_{p,q}^s(\mathbb{R}^n; V) \simeq (L^p \cap \dot{\mathbb{B}}_{p,q}^s)(\mathbb{R}^n; V).$$

Next, for $1 \leq p < \infty$ and $M \in \mathbb{N}_0$, we define the (p, M) -oscillation of a measurable map $f: \mathbb{R}^n \rightarrow V$ at $x_0 \in \mathbb{R}^n$ by

$$\text{osc}_p^M f(x_0, r) := \inf \left\{ \left(\int_{B(x_0, r)} |f(x) - \pi(x)|^p \, dx \right)^{\frac{1}{p}} : \pi \in \mathcal{P}_M(\mathbb{R}^n; V) \right\}.$$

The next lemma is certainly clear to the experts, but we have been unable to find a precise reference; see, however, DORRONSORO [24] for a related result. In our arguments below, we will merely require inequality ' \lesssim ' in (4.3), and a quick argument for this is sketched in the Appendix, Section 7.2.

Lemma 4.1. *Let $1 \leq p, q < \infty$ and $s > 0$. If $M \in \mathbb{N}$ satisfies $M > \lfloor s \rfloor$, then for all $1 \leq u \leq p$ we have the equivalence of norms*

$$(4.3) \quad \|f\|_{\dot{\mathbb{B}}_{p,q}^s(\mathbb{R}^n)} \simeq \left(\int_0^\infty \|\text{osc}_u^M f(\cdot, t)\|_{L^p(\mathbb{R}^n)}^q \frac{dt}{t^{1+sq}} \right)^{\frac{1}{q}}, \quad f \in \mathcal{S}(\mathbb{R}^n; V),$$

where the constants implicit in \simeq only depend on n, M, s, p, q, u and V .

Let $\Omega \subset \mathbb{R}^n$ be open and bounded with C^∞ -boundary $\partial\Omega$ and outer unit normal $\nu = (\nu_1, \dots, \nu_n)$. By definition, there exists $N \in \mathbb{N}$ and open balls W_1, \dots, W_N with $\partial\Omega \subset \bigcup_{j=1}^N W_j$ such that the following holds for all $j = 1, \dots, N$: We have $\partial\Omega \cap W_j \neq \emptyset$, and there exists $F^{(j)} \in C^\infty(\overline{W_j}; \mathbb{R}^n)$ with $F^{(j)}: W_j \rightarrow \mathbb{R}^n$ injective and having open, bounded image, $F^{(j)}(\partial\Omega \cap W_j) \subset \{x \in \mathbb{R}^n: x_n = 0\}$ and $F^{(j)}(\Omega \cap W_j) \subset \mathbb{R}^{n-1} \times (0, \infty)$ is open, simply connected and satisfies $\det(\nabla F^{(j)}) \neq 0$ on $\overline{W_j}$. Let $W_0 \Subset \Omega$ be open and bounded with C^∞ -boundary ∂W_0 such that $\overline{\Omega} \subset \bigcup_{j=0}^N W_j$ and $(\psi_j)_{j=0}^N$ be a partition of unity subject to $(W_j)_{j=0}^N$. Following TRIEBEL [61, Chpt. 3.2.2, Def. 2], for $s > 0$, $1 \leq p, q < \infty$, we define the Besov norm

$$(4.4) \quad \|f\|_{\mathbb{B}_{p,q}^s(\partial\Omega)} := \sum_{j=1}^N \|(\psi_j f)(F^{(j)})^{-1}(\cdot, 0)\|_{\mathbb{B}_{p,q}^s(\mathbb{R}^{n-1})}, \quad f \in \mathcal{D}'(\partial\Omega; V).$$

In view of a convenient characterisation of trace spaces for smooth domains, we follow the slightly more general approach of MAZ'YA, MITREA & SHAPOSHNIKOVA [50] and define for $m \in \mathbb{N}$, $1 < p < \infty$ and $0 < s < 1$

$$(4.5) \quad \dot{\mathbb{B}}_p^{m-1+s}(\partial\Omega; V) := \overline{\{(\partial^\alpha v|_{\partial\Omega})_{|\alpha| \leq m-1}: v \in C_c^\infty(\mathbb{R}^n; V)\}}^{\left(\mathbb{B}_{p,p}^s(\partial\Omega; V)\right)^\mathbb{M}},$$

where $\mathbb{M} := \#\{\alpha \in \mathbb{N}_0^n: |\alpha| \leq m-1\}$. By [50, Prop. 6.9], $(f^\alpha)_{|\alpha| \leq m-1}$ belongs to $\dot{\mathbb{B}}_p^{m-1+s}(\partial\Omega; V)$ if and only if

$$(4.6) \quad \begin{aligned} f^\alpha &\in \mathbb{B}_{p,p}^s(\partial\Omega; V) \quad \text{for all } |\alpha| \leq m-1, \quad \text{and} \\ (\nu_j \partial_k - \nu_k \partial_j) f^\alpha &= \nu_j f^{\alpha+e_k} - \nu_k f^{\alpha+e_j} \quad \text{for all } |\alpha| \leq m-2, j, k \in \{1, \dots, n\}. \end{aligned}$$

The natural compatibility condition (4.6)₂ on the tangential derivatives is a higher order variant of the Whitney array compatibility conditions considered by ADOLFSSON & PIPHER [1]. As in [1], the equations of (4.6)₂ are understood in the sense of being fulfilled by an arbitrary extension \tilde{f}^α of f^α to a neighbourhood of $\partial\Omega$; since the operator $\nu_j \partial_k - \nu_k \partial_j$ is a tangential derivative on $\partial\Omega$, $(\nu_j \partial_k - \nu_k \partial_j) \tilde{f}^\alpha$ will be independent of the particular extension and hence is well-defined.

Lemma 4.2 ([50, Prop. 6.5]). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded domain with C^∞ -boundary $\partial\Omega$. Let $1 < p < \infty$, $m \in \mathbb{N}$ and define $\text{tr}_{m-1, \partial\Omega}$ on $W^{m,p}(\Omega; V)$ by*

$$(4.7) \quad \text{tr}_{m-1, \partial\Omega} f := (\text{Tr}_{\partial\Omega}(\partial^\alpha f))_{|\alpha| \leq m-1}, \quad f \in W^{m,p}(\Omega; V),$$

where $\text{Tr}_{\partial\Omega}: W^{1,p}(\Omega; V) \rightarrow \mathbb{B}_{p,p}^{1-1/p}(\partial\Omega; V)$ is the trace operator on $W^{1,p}(\Omega; V)$ as usual. Then

$$(4.8) \quad \text{tr}_{m-1, \partial\Omega}: W^{m,p}(\Omega; V) \rightarrow \dot{\mathbb{B}}_p^{m-1/p}(\partial\Omega; V)$$

is a well-defined, linear, bounded operator which is onto. Moreover, there exists a bounded, linear right-inverse $\mathcal{L}: \dot{\mathbb{B}}_p^{m-1/p}(\partial\Omega; V) \rightarrow W^{m,p}(\Omega; V)$ of $\text{tr}_{m-1, \partial\Omega}$ in the sense that, whenever $\dot{f} = (f^\alpha)_{|\alpha| \leq m-1} \in \dot{\mathbb{B}}_p^{m-1/p}(\partial\Omega; V)$, then

$$(4.9) \quad \text{Tr}_{\partial\Omega}(\partial^\alpha(\mathcal{L}\dot{f})) = f^\alpha \quad \text{for all } |\alpha| \leq m-1.$$

4.2. Geometric setup and potential estimates. In this and the following subsection, we suppose that \mathbb{A} is a \mathbb{C} -elliptic differential operator of the form (1.1). Moreover, we shall tacitly work with the particular halfspace $\mathbb{H} := \mathbb{R}^{n-1} \times (0, \infty)$, but our arguments are easily seen to generalise to any halfspace of \mathbb{R}^n .

We start by fixing notation: For some given $z' \in \mathbb{R}^{n-1}$ and $t > 0$ we denote

$$(4.10) \quad K_{z',t} := \{(y', y_n) \in \mathbb{R}^n: 0 \leq y_n \leq t, |y' - z'|^2 < 9y_n^2\}$$




FIGURE 2. Not-to-scale construction of the the umbra shadow estimate underlying the proof of Theorem 1.1. To control the oscillation of $u(\cdot, 0)$, the Riesz potential estimate from Lemma 4.3 is applied to shifted cones emanating from points in the red area, where all cones are star-shaped with respect to a joint ball B .

the single-sided cone emanating from $(z', 0)$ in direction e_n , of height t and base radius $3t$. To distinguish between n - and $(n - 1)$ -dimensional balls, we put for $x' \in \mathbb{R}^{n-1}$, $x \in \mathbb{R}^n$ and $r > 0$

$$B_n(x, r) := B(x, r) \quad \text{and} \quad B_{n-1}(x', r) := \partial\mathbb{H} \cap B((x', 0), r).$$

Adopting this notation, let us note that in this situation there holds

$$(4.11) \quad B_n((x', t), \frac{t}{2}) \subset K_{y', 2t} \quad \text{for all } y' \in \mathbb{R}^{n-1} \text{ with } |x' - y'| < t.$$

In fact, let $(\xi', \xi_n) \in B_n((x', t), \frac{t}{2})$ so that, firstly, $\frac{1}{2}t \leq \xi_n \leq \frac{3}{2}t$ and hence $\frac{3}{2}t \leq 3\xi_n$. As a consequence,

$$|\xi' - y'| \leq |\xi' - x'| + |x' - y'| < \frac{3}{2}t \leq 3\xi_n$$

and so (4.11) follows. This situation is depicted in Figure 2. Combining this setup with Proposition 3.3, we arrive at the following potential estimates on cones:

Lemma 4.3 (Conical Riesz potential estimate). *There exists $c = c(\mathbb{A}) > 0$ with the following property: If $x' \in \mathbb{R}^{n-1}$ and $t > 0$, then for every $y' \in \mathbb{R}^{n-1}$ with $|x' - y'| < t$ there holds for all $u \in C^\infty(\overline{\mathbb{H}}; V)$ and all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq k - 1$*

$$(4.12) \quad \begin{aligned} & |(\partial^\alpha u)(y', 0) - (\partial^\alpha \Pi_{\mathbb{A}}^{B_n((x', t), \frac{t}{2})} u)(y', 0)| \\ & \leq c \int_{K_{y', 2t}} \frac{|\mathbb{A}u(z', z_n)|}{(|z' - y'| + |z_n|)^{n+|\alpha|-k}} d(z', z_n). \end{aligned}$$

Proof. Let x', y' and t be as in the lemma. By (4.11), $B_n((x', t), \frac{t}{2}) \subset K_{y', 2t}$ and clearly $K_{y', 2t}$ is star-shaped with respect to $B_n((x', t), \frac{t}{2})$. Thus we may apply Proposition 3.3 to $\Omega = K_{y', 2t}$ and $B = B_n((x', t), \frac{t}{2})$. Note that $\Pi_{\mathbb{A}}^B u := \tilde{\Pi}_{\mathbb{A}} \mathbb{T}_{m-1}^B u$ is independent of y' and the diameters of $K_{y', 2t}$ and $B_n((x', t), \frac{t}{2})$ are uniformly comparable. Hence (4.12) follows from (K1) via

$$\left| \int_{K_{y', 2t}} (\partial_y^\alpha \mathfrak{K}_{\mathbb{A}}^B)((y', 0), z) \mathbb{A}u(z) dz \right| \leq C \int_{K_{y', 2t}} \frac{|\mathbb{A}u(z', z_n)|}{(|y' - z'| + |z_n|)^{n+|\alpha|-k}} d(z', z_n).$$

The proof is complete. \square

4.3. The trace estimate for smooth maps. After the preparations of the preceding subsection, we can now proceed to showing that \mathbb{C} -ellipticity suffices for the trace estimate of Theorem 1.1 (b). We provide a slightly stronger variant that is also applicable to the $1 < p < \infty$ -scenario of Theorem 4.4 below.

Proof of sufficiency of \mathbb{C} -ellipticity for (1.7). Let $1 \leq p < \infty$ and let \mathbb{A} be a \mathbb{C} -elliptic differential operator of the form (1.1) with $k \geq 2$. Moreover, let $u \in C_c^\infty(\overline{\mathbb{H}}; V)$, $x' \in \partial\mathbb{H}$ and $t > 0$. We distinguish the cases $|\alpha| \leq k-2$ and $|\alpha| = k-1$. In the first case, we will establish a $\dot{B}_{p,p}^{k-|\alpha|-1/p}(\partial\mathbb{H})$ -bound on $(\partial^\alpha u)(\cdot, 0)$ for all $1 \leq p < \infty$, whereas in the second case we will only strive for such a bound provided $1 < p < \infty$.

Let $|\alpha| \leq k-2$ and $1 \leq p < \infty$. Since \mathbb{A} is \mathbb{C} -elliptic, we may choose $m := \deg_{\mathcal{P}}(\mathbb{A}) \in \mathbb{N}$ (cf. (3.6)) so that $\partial^\alpha \ker(\mathbb{A}) \subset \mathcal{P}_m(\mathbb{R}^n; V)$ and, with $s := k - |\alpha| - 1/p > 0$,

$$m > \deg_{\mathcal{P}}(\mathbb{A}) - 1 \geq k - |\alpha| - 1 = \lfloor k - |\alpha| - 1/p \rfloor = \lfloor s \rfloor.$$

Since p and m are fixed, we put for a \mathcal{H}^{n-1} -locally integrable map $v: \partial\mathbb{H} \rightarrow V$

$$(4.13) \quad \text{osc}_{p, \partial\mathbb{H}}^m v(x', t) := \inf \left\{ \left(\int_{B_{n-1}(x', t)} |v - \pi|^p d^{n-1}y \right)^{\frac{1}{p}} : \pi \in \mathcal{P}_m(\mathbb{R}^n; V) \right\}$$

In this situation, Lemma 4.1 implies with $U^\alpha(x') := (\partial^\alpha u)(x', 0)$

$$(4.14) \quad \|U^\alpha\|_{\dot{B}_{p,p}^{k-|\alpha|-1/p}(\partial\mathbb{H})}^p \leq c \int_0^\infty \frac{\|\text{osc}_{p, \partial\mathbb{H}}^m U^\alpha(\cdot, t)\|_{L^p(\partial\mathbb{H})}^p}{t^{(k-|\alpha|)p}} dt.$$

To bound the right-hand side, we use Lemma 4.3 and $\partial^\alpha \ker(\mathbb{A}) \subset \mathcal{P}_m(\mathbb{R}^n; V)$ to obtain

$$(4.15) \quad \begin{aligned} |\text{osc}_{p, \partial\mathbb{H}}^m U^\alpha(x', t)|^p &\leq \int_{B_{n-1}(x', t)} |U^\alpha(y') - \partial^\alpha \Pi_{\mathbb{A}}^{B_n((x', t), \frac{t}{2})} u(y', 0)|^p d^{n-1}y' \\ &\leq c \int_{B_{n-1}(x', t)} \left| \int_0^{2t} \int_{\partial\mathbb{H}} \mathbb{1}_{\{|z'-y'| < 3\tau\}} \frac{|\mathbb{A}u(z', \tau)|}{(|z'-y'| + \tau)^{n-k+|\alpha|}} d^{n-1}z' d\tau \right|^p d^{n-1}y' \\ &\leq c \int_{B_{n-1}(x', t)} t^{p-1} \int_0^{2t} \tau^{(n-1)(p-1)} \times \\ &\quad \times \left(\int_{\partial\mathbb{H}} \mathbb{1}_{\{|z'-y'| < 3\tau\}} \frac{|\mathbb{A}u(z', \tau)|^p}{(|z'-y'| + \tau)^{p(n-k+|\alpha|)}} d^{n-1}z' \right) d\tau d^{n-1}y'. \end{aligned}$$

Let us note that the very first of the previous chain of inequalities is where the \mathbb{C} -ellipticity enters as otherwise the nullspace does not consist of polynomials of a fixed maximal degree. In combination with (4.14), we thus obtain

$$\begin{aligned} \|U^\alpha\|_{\dot{B}_{p,p}^{k-|\alpha|-1/p}(\partial\mathbb{H})}^p &\leq c \int_0^\infty \int_{\partial\mathbb{H}} \int_{B_{n-1}(x', t)} t^{p-1} \int_0^{2t} \tau^{(n-1)(p-1)} \times \\ &\quad \times \left(\int_{\partial\mathbb{H}} \mathbb{1}_{\{|z'-y'| < 3\tau\}} \frac{|\mathbb{A}u(z', \tau)|^p}{(|z'-y'| + \tau)^{p(n-k+|\alpha|)}} d^{n-1}z' \right) d\tau d^{n-1}y' d^{n-1}x' \frac{dt}{t^{(k-|\alpha|)p}} \\ &\leq c \int_0^\infty \int_{\partial\mathbb{H}} \int_{\partial\mathbb{H}} \int_0^{2t} \tau^{(n-1)(p-1)} \mathbb{1}_{\{|x'-y'| < t\}} \times \\ &\quad \times \left(\int_{\partial\mathbb{H}} \frac{\mathbb{1}_{\{|z'-y'| < 3\tau\}} |\mathbb{A}u(z', \tau)|^p}{(|z'-y'| + \tau)^{p(n-k+|\alpha|)}} d^{n-1}z' \right) d\tau d^{n-1}y' d^{n-1}x' \frac{dt}{t^{(k-|\alpha|-1)p+n}} = (*) \end{aligned}$$

Now, if $\mathbb{1}_{\{|z'-y'|<3\tau\}} = 1$, then $\tau \leq |z' - y'| + \tau \leq 4\tau$. Therefore,

$$\begin{aligned}
(*) &\leq c \int_0^\infty \int_{\partial\mathbb{H}} \int_{\partial\mathbb{H}} \int_{\partial\mathbb{H}} \int_0^{2t} \tau^{(n-1)(p-1)} \mathbb{1}_{\{|z'-y'|<3\tau\}} \mathbb{1}_{\{|x'-y'|<t\}} \times \\
&\quad \times \frac{|\mathbb{A}u(z', \tau)|^p}{\tau^{p(n-k+|\alpha|)}} d\tau d^{n-1}z' d^{n-1}y' d^{n-1}x' \frac{dt}{t^{(k-|\alpha|-1)p+n}} \\
&\leq c \int_0^\infty \int_{\partial\mathbb{H}} \int_{\partial\mathbb{H}} \int_0^{2t} \mathbb{1}_{\{|z'-y'|<3\tau\}} \times \\
&\quad \times \frac{|\mathbb{A}u(z', \tau)|^p}{\tau^{n-1+p(1+|\alpha|-k)}} d\tau d^{n-1}z' d^{n-1}y' \frac{dt}{t^{(k-|\alpha|-1)p+1}} \\
&\leq c \int_0^\infty \int_{\partial\mathbb{H}} \int_0^{2t} \frac{|\mathbb{A}u(z', \tau)|^p}{\tau^{p(1+|\alpha|-k)}} d\tau d^{n-1}z' \frac{dt}{t^{(k-|\alpha|-1)p+1}} \\
&= c \int_0^\infty \int_{\partial\mathbb{H}} \int_0^\infty \mathbb{1}_{(0,2t)}(\tau) |\mathbb{A}u(z', \tau)|^p \frac{d\tau}{\tau^{p(1+|\alpha|-k)}} d^{n-1}z' \frac{dt}{t^{(k-|\alpha|-1)p+1}} \\
&= c \int_0^\infty \int_{\partial\mathbb{H}} \int_0^\infty \mathbb{1}_{(0,2t)}(\tau) |\mathbb{A}u(z', \tau)|^p \frac{dt}{t^{(k-1-|\alpha|)p+1}} d^{n-1}z' \frac{d\tau}{\tau^{p(1+|\alpha|-k)}} \\
&\leq c \int_0^\infty \int_{\partial\mathbb{H}} \left(\int_{\frac{\tau}{2}}^\infty \frac{dt}{t^{(k-1-|\alpha|)p+1}} \right) |\mathbb{A}u(z', \tau)|^p d^{n-1}z' \frac{d\tau}{\tau^{p(1+|\alpha|-k)}} \\
&\stackrel{k-|\alpha|-1>0}{\leq} c \int_0^\infty \int_{\partial\mathbb{H}} |\mathbb{A}u(z', \tau)|^p d^{n-1}z' d\tau \\
&= c \|\mathbb{A}u\|_{L^p(\mathbb{H})}^p.
\end{aligned}$$

The reader will notice that this argument requires refinement for $|\alpha| = k - 1$ since in this case, the penultimate estimation cannot be accomplished.

Let $1 < p < \infty$ and $|\alpha| = k - 1$; in this case, the reader will notice that we may even allow for $k = 1$. Defining $\text{osc}_{1, \partial\mathbb{H}}^m$ in the analogous manner as in (4.13), we imitate (4.15) to find by changing variables $z' = x' - \xi'$ and $\eta' = x' - y'$ in the third step

$$\begin{aligned}
|\text{osc}_{1, \partial\mathbb{H}}^m U^\alpha(x', t)| &\leq \frac{c}{t^{n-1}} \int_0^{2t} \int_{\partial\mathbb{H}} \int_{\partial\mathbb{H}} \mathbb{1}_{\{|x'-y'|<t\}} \mathbb{1}_{\{|z'-y'|<3\tau\}} \times \\
&\quad \times \frac{|\mathbb{A}u(z', \tau)|}{(|z' - y'| + \tau)^{n-1}} d^{n-1}y' d^{n-1}z' d\tau \\
&= \frac{c}{t^{n-1}} \int_0^{2t} \int_{\partial\mathbb{H}} |\mathbb{A}u(z', \tau)| \left(\int_{\partial\mathbb{H}} \mathbb{1}_{\{|x'-y'|<t\}} \mathbb{1}_{\{|z'-y'|<3\tau\}} \times \right. \\
&\quad \times \left. \frac{1}{(|z' - y'| + \tau)^{n-1}} d^{n-1}y' \right) d^{n-1}z' d\tau \\
&= \frac{c}{t^{n-1}} \int_0^{2t} \int_{\partial\mathbb{H}} |\mathbb{A}u(x' - \xi', \tau)| \mathbb{1}_{\{|\xi'| \leq 7t\}} \left(\int_{\partial\mathbb{H}} \mathbb{1}_{\{|\eta'|<t\}} \mathbb{1}_{\{|\eta'-\xi'|<3\tau\}} \times \right. \\
&\quad \times \left. \frac{1}{(|\eta' - \xi'| + \tau)^{n-1}} d^{n-1}\eta' \right) d^{n-1}\xi' d\tau = (**),
\end{aligned}$$

where we have used that if $|\xi'| > 7t$, then $|\eta'| < t$ yields

$$3\tau \stackrel{\tau \leq 2t}{\leq} 6t = 7t - t < |\xi'| - |\eta'| \leq |\xi' - \eta'|$$

and so $\mathbb{1}_{\{|\eta'|<t\}} \mathbb{1}_{\{|\eta'-\xi'|<3\tau\}} = \mathbb{1}_{\{|\eta'|<t\}} \mathbb{1}_{\{|\eta'-\xi'|<3\tau\}} \mathbb{1}_{\{|\xi'| \leq 7t\}}$. Since we have

$$c \int_{B_{n-1}(0, 3\tau)} \frac{d^{n-1}\theta}{(|\theta| + \tau)^{n-1}} \leq c \frac{\mathcal{H}^{n-1}(B_{n-1}(0, 3\tau))}{\tau^{n-1}} \leq c$$

as a bound for the very inner integral in (**), we thus conclude

$$|\operatorname{osc}_{1,\partial\mathbb{H}}^m U^\alpha(x', t)| \leq (**) \leq \frac{c}{t^{n-1}} \int_0^{2t} \int_{\partial\mathbb{H}} |\mathbb{A}u(x' - \xi', \tau)| \mathbb{1}_{\{|\xi'| \leq \tau t\}} d^{n-1}\xi' d\tau.$$

Integrating the previous inequality with respect to $x' \in \partial\mathbb{H}$, we choose $0 < \delta < p-1$ to find by Hölder's inequality and a change of variables

$$(4.16) \quad \begin{aligned} & \|\operatorname{osc}_{1,\partial\mathbb{H}}^m U^\alpha(\cdot, t)\|_{L^p(\partial\mathbb{H})}^p \\ & \leq \frac{c}{t^{p(n-1)}} \int_{\partial\mathbb{H}} \left| \int_0^{2t} \frac{1}{\tau^{\frac{\delta}{p}}} \int_{\partial\mathbb{H}} \tau^{\frac{\delta}{p}} \mathbb{1}_{\{|\xi'| \leq \tau t\}} |\mathbb{A}u(x' - \xi', \tau)| d^{n-1}\xi' d\tau \right|^p d^{n-1}x' \\ & \leq \frac{ct^{(p-1)(n-1)}}{t^{p(n-1)}} \int_{\partial\mathbb{H}} \left(\int_0^{2t} \frac{d\tau}{\tau^{\frac{\delta}{p-1}}} \right)^{p-1} \times \\ & \quad \times \left(\int_0^{2t} \tau^\delta \int_{\partial\mathbb{H}} \mathbb{1}_{\{|\xi'| \leq \tau t\}} |\mathbb{A}u(x' - \xi', \tau)|^p d^{n-1}\xi' d\tau \right) d^{n-1}x' \\ & \leq \frac{ct^{(p-1)(n-1)}}{t^{p(n-1)}} t^{p+n-2-\delta} \left(\int_0^{2t} \tau^\delta \int_{\partial\mathbb{H}} |\mathbb{A}u(\xi', \tau)|^p d^{n-1}\xi' d\tau \right) \\ & \leq ct^{p-1-\delta} \int_0^{2t} \tau^\delta \|\mathbb{A}u(\cdot, \tau)\|_{L^p(\partial\mathbb{H})}^p d\tau. \end{aligned}$$

Working from (4.16), we then obtain

$$\begin{aligned} \|U^\alpha\|_{\dot{B}_{p,p}^{1-1/p}(\partial\mathbb{H})}^p & \leq c \int_0^\infty t^{p-1-\delta} \int_0^{2t} \tau^\delta \|\mathbb{A}u(\cdot, \tau)\|_{L^p(\partial\mathbb{H})}^p d\tau \frac{dt}{t^p} \\ & = c \int_0^\infty \left(\int_{\frac{\tau}{2}}^\infty \frac{dt}{t^{1+\delta}} \right) \tau^\delta \|\mathbb{A}u(\cdot, \tau)\|_{L^p(\partial\mathbb{H})}^p d\tau \\ & \leq c \|\mathbb{A}u\|_{L^p(\mathbb{H})}^p. \end{aligned}$$

The proof is complete. \square

Let us note that for $p = 1$ and $|\alpha| = 0$, the above proof works *and is bound to work* for $k \geq 2$ only. Indeed, for $k = 1$, $L^1(\partial\mathbb{H}; V)$ -trace estimates are optimal [29, 11]. However,

$$(4.17) \quad \dot{B}_{1,1}^0(\partial\mathbb{H}) \hookrightarrow \dot{F}_{1,2}^0(\partial\mathbb{H}) = \mathcal{H}^1(\partial\mathbb{H}) \subsetneq L^1(\partial\mathbb{H}),$$

with the homogeneous Hardy space \mathcal{H}^1 , cf. TRIEBEL [61, Chpt. 5.2.4], so that maps with a finite $\dot{B}_{1,1}^0(\partial\mathbb{H})$ -norm need to have integral zero (cf. [52, Prop. 5.8]). Also note that the oscillation characterisation in the zero order case does not hold, cf. [63, Chpt. 3.5.1].

4.4. Proof of Theorem 1.1 and its counterpart for $1 < p < \infty$. In this section we prove Theorem 1.1 and state and prove its counterpart for $1 < p < \infty$, Theorem 4.4 below, where stronger statements are available.

Proof of Theorem 1.1. Implication (a) \Rightarrow (b) has been established in Section 4.3. On the other hand, since $\dot{B}_{1,1}^{k-1}(\partial\mathbb{H}; V) \hookrightarrow \dot{W}^{k-1,1}(\partial\mathbb{H}; V)$ for any open halfspace \mathbb{H} , (b) yields the inequality $\|D^{k-1}u\|_{L^1(\partial\mathbb{H})} \leq c\|\mathbb{A}u\|_{L^1(\mathbb{H})}$ for all $u \in C_c^\infty(\overline{\mathbb{H}}; V)$. By RAITA, VAN SCHAFTINGEN and the second author [31, Thm. 5.2], this already implies that \mathbb{A} is \mathbb{C} -elliptic, and the proof is complete. \square

The rest of the section is devoted to a self-contained proof of the following theorem in the spirit of [4, 56, 42], where our main focus now is on the equivalence of sharp trace estimates and Korn-type inequalities for smooth domains. Here, sharp trace estimates give access to global Calderón-Zygmund estimates, which is

fundamentally different from the non-smooth context of John domains as discussed in Section 6.

Theorem 4.4 (Higher order traces, $1 < p < \infty$). *Let $k \in \mathbb{N}$, $k \geq 1$ and let $1 < p < \infty$. Then the following are equivalent for a k -th order differential operator \mathbb{A} of the form (1.1):*

- (a) \mathbb{A} is \mathbb{C} -elliptic.
- (b) There exists a constant $c = c(p, \mathbb{A}) > 0$ such that for any open halfspace \mathbb{H} and all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq k-1$ there holds

$$(4.18) \quad \|\partial^\alpha u\|_{\dot{B}_{p,p}^{k-|\alpha|-1/p}(\partial\mathbb{H})} \leq c \|\mathbb{A}u\|_{L^p(\mathbb{H})} \quad \text{for all } u \in C_c^\infty(\overline{\mathbb{H}}; V).$$

- (c) For every open and bounded domain Ω with boundary of class C^∞ , there exists a constant $c = c(p, \mathbb{A}, \Omega) > 0$ such that for all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq k-1$ there holds

$$(4.19) \quad \|\partial^\alpha u\|_{\dot{B}_{p,p}^{k-|\alpha|-1/p}(\partial\Omega)} \leq c(\|u\|_{W^{k-1,p}(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)}) \quad \text{for all } u \in C^\infty(\overline{\Omega}; V).$$

- (d) For every open and bounded domain Ω with boundary of class C^∞ and all $1 < p < \infty$, there exists a constant $c = c(p, \mathbb{A}, \Omega) > 0$ such that

$$(4.20) \quad \sum_{j=0}^k \|\nabla^j u\|_{L^p(\Omega)} \leq c(\|u\|_{W^{k-1,p}(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)}) \quad \text{for all } u \in C^\infty(\overline{\Omega}; V),$$

In particular, if any of (a)–(d) holds, then we have (with $W^{\mathbb{A},p}$ and $\widetilde{W}^{\mathbb{A},p}$ as in Section 3.1) for any open and bounded domain with smooth boundary $\partial\Omega$

$$(4.21) \quad W^{k,p}(\Omega; V) \simeq W^{\mathbb{A},p}(\Omega) \simeq \widetilde{W}^{\mathbb{A},p}(\Omega).$$

It will be clear from the proof that some implications also persist for less regular domains (so, e.g., (a) \Rightarrow (c) would also follow for domains satisfying a cone condition by a slight variation of the arguments of Section 4.3), but we stick to smooth domains for ease of exposition. For the proof of Theorem 4.4, we require two preparatory lemmas:

Lemma 4.5 (Peetre-Tartar lemma, [60, Lem. 11.1]). *Let $(E_i, \|\cdot\|_{E_i})$, $i \in \{1, 2, 3\}$, be three Banach spaces, $A: E_1 \rightarrow E_2$ a linear and bounded operator and $B: E_1 \rightarrow E_3$ be a linear and compact operator. If $x \mapsto \|Ax\|_{E_2} + \|Bx\|_{E_3}$ is a norm on E_1 that is equivalent to $\|\cdot\|_{E_1}$, then $\dim(\ker(A)) < \infty$.*

Lemma 4.6 (Traces and Gauß-Green). *Suppose that assertion (c) of Theorem 4.4 holds. Then for every $1 < p < \infty$ and every open and bounded domain $\Omega \subset \mathbb{R}^n$ with boundary of class C^∞ there exists a linear and bounded boundary trace operator*

$$(4.22) \quad \text{tr}_{k-1, \partial\Omega}^{\mathbb{A}}: \widetilde{W}^{\mathbb{A},p}(\Omega) \ni u \mapsto (\text{tr}_{\alpha, \partial\Omega}^{\mathbb{A}}(u))_{|\alpha| \leq k-1} \in \dot{B}_p^{k-1/p}(\partial\Omega; V)$$

such that, in particular, $\text{tr}_{k-1, \partial\Omega}^{\mathbb{A}}(u) = ((\partial^\alpha u)|_{\partial\Omega})_{|\alpha| \leq k-1}$ holds for all $u \in C^\infty(\overline{\Omega}; V)$ and so $\text{tr}_{k-1, \partial\Omega}^{\mathbb{A}} = \text{tr}_{k-1, \partial\Omega}$ on $W^{k,p}(\Omega; V)$ with $\text{tr}_{k-1, \partial\Omega}$ as in Lemma 4.2.

For $\ell \in \{0, \dots, k-1\}$, put $\text{tr}_{\partial\Omega}^{\mathbb{A}, \ell}(u) := (\text{tr}_{\alpha, \partial\Omega}^{\mathbb{A}}(u))_{|\alpha| = \ell}$. For each $l \in \{1, \dots, n\}$ and each $s \in \{1, \dots, k\}$ there exist continuous bilinear forms $\mathcal{B}_{l,s}^{\mathbb{A}}: \odot^{k-s}(\mathbb{R}^n; V) \times \odot^{s-1}(\mathbb{R}^n; W) \rightarrow \mathbb{R}$ such that for all $u \in \widetilde{W}^{\mathbb{A},p}(\Omega)$ and all $\varphi \in C^\infty(\overline{\Omega}; W)$ we have the Gauß-Green identity (with $\nu = (\nu_1, \dots, \nu_n)$ denoting the outer unit normal to

$\partial\Omega$)

$$(4.23) \quad \int_{\Omega} \mathbb{A}u \cdot \varphi \, dx = (-1)^k \int_{\Omega} u \cdot \mathbb{A}^* \varphi \, dx \\ + \sum_{l=1}^n \sum_{s=1}^k \int_{\partial\Omega} \mathcal{B}_{l,s}^{\mathbb{A}} \left(\text{tr}_{\partial\Omega}^{\mathbb{A},k-s}(u), \nabla^{s-1} \varphi \right) \nu_l \, d\mathcal{H}^{n-1},$$

where $\mathbb{A}^* = \sum_{|\alpha|=k} \mathbb{A}_{\alpha}^{\top} \partial^{\alpha}$ is the formal L^2 -adjoint of \mathbb{A} .

Proof. Since Ω has C^{∞} -boundary, an argument similar to that in, e.g., [26, Chpt. 5.3.3, Thm. 5.3] yields that for every $u \in \widetilde{W}^{\mathbb{A},p}(\Omega)$ there exists $(u_j) \subset C^{\infty}(\overline{\Omega}; V)$ such that $\|u - u_j\|_{\widetilde{W}^{\mathbb{A},p}(\Omega)} \rightarrow 0$ as $j \rightarrow \infty$. Let $\alpha \in \mathbb{N}_0^n$ satisfy $|\alpha| \leq k-1$. By (4.19), $((\partial^{\alpha} u_j)|_{\partial\Omega})$ is Cauchy in $B_{p,p}^{k-|\alpha|-1/p}(\partial\Omega; V)$ and therefore converges to some $u^{\alpha} =: \text{tr}_{\alpha,\partial\Omega}^{\mathbb{A}}(u) \in B_{p,p}^{k-|\alpha|-1/p}(\partial\Omega; V)$, which is clearly independent of the approximating sequence and hereafter well-defined. To conclude (4.22), we note that $\{u^{\alpha}: |\alpha| \leq k-1\}$ matches the Besov-Whitney array condition (4.6)₂ \mathcal{H}^{n-1} -a.e. on $\partial\Omega$; indeed, for any $|\alpha| \leq k-2$ and all $i, l \in \{1, \dots, n\}$ we have

$$\begin{aligned} & \|\nu_i \partial_l u^{\alpha} - \nu_l \partial_i u^{\alpha} - (\nu_i u^{\alpha+e_l} - \nu_l u^{\alpha+e_i})\|_{L^p(\partial\Omega)} \\ & \leq \|(\nu_i \partial_l u^{\alpha} - \nu_l \partial_i u^{\alpha}) - (\nu_i \partial_l \partial^{\alpha} u_j - \nu_l \partial_i \partial^{\alpha} u_j)\|_{L^p(\partial\Omega)} \\ & \quad + \|\nu_i u^{\alpha+e_l} - \nu_i \partial^{\alpha+e_l} u_j\|_{L^p(\partial\Omega)} + \|\nu_l u^{\alpha+e_i} - \nu_l \partial^{\alpha+e_i} u_j\|_{L^p(\partial\Omega)} \\ & \quad + \underbrace{\|\nu_i \partial_l \partial^{\alpha} u_j - \nu_l \partial_i \partial^{\alpha} u_j - (\nu_i \partial^{\alpha+e_l} u_j - \nu_l \partial^{\alpha+e_i} u_j)\|_{L^p(\partial\Omega)}}_{=0} =: \text{I}_j + \dots + \text{IV}_j. \end{aligned}$$

Since $(\nu_i \partial_l - \nu_l \partial_i)$ is a tangential differential operator on the smooth boundary $\partial\Omega$, we have with the full tangential gradient ∇_{τ} along $\partial\Omega$

$$\text{I}_j \leq \|\nabla_{\tau}(u^{\alpha} - \partial^{\alpha} u_j)\|_{L^p(\partial\Omega)} \leq \|u^{\alpha} - \partial^{\alpha} u\|_{B_{p,p}^{k-|\alpha|-1/p}(\partial\Omega)} \rightarrow 0, \quad j \rightarrow \infty,$$

as $k - |\alpha| - 1/p \geq 1$ by our assumption on α and p . The terms $\text{II}_j, \text{III}_j$ vanish in the limit by construction, and $\text{IV}_j = 0$ for all $j \in \mathbb{N}$; thus, (4.22) follows. Equation (4.23) follows for maps u and φ which are smooth up to $\partial\Omega$ by repeated use of the usual Gauß-Green theorem, and then inherits to $u \in \widetilde{W}^{\mathbb{A},p}(\Omega)$ by means of smooth approximation. \square

Proof of Theorem 4.4. Let $1 < p < \infty$. Implication (a) \Rightarrow (b) has already been proved in Section 4.3. For implication (b) \Rightarrow (c), we assume that Ω is connected, open and bounded with C^{∞} -boundary; if it is not connected, we apply the following to each of the finitely many connected components of Ω . Let $u \in C^{\infty}(\overline{\Omega}; V)$. Adopting the notation from (4.4), for $j \in \{1, \dots, N\}$, (b) and (4.2) yield for any $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq k-1$ and all $\beta \in \mathbb{N}_0^n$ for the compactly supported, smooth map $\text{I}_{j,\beta} := ((\partial^{\beta} \psi_j)u)(F^{(j)})^{-1}(\cdot, 0)$

$$\begin{aligned} \|\partial^{\alpha} \text{I}_{j,\beta}\|_{B_{p,p}^{k-|\alpha|-1/p}(\mathbb{R}^{n-1})} & \stackrel{(4.2)}{\leq} c \|\partial^{\alpha} \text{I}_{j,\beta}\|_{L^p(\mathbb{R}^{n-1})} + c \|\partial^{\alpha} \text{I}_{j,\beta}\|_{\dot{B}_{p,p}^{k-|\alpha|-1/p}(\mathbb{R}^{n-1})} \\ & \leq c \|\partial^{\alpha} \text{I}_{j,\beta}\|_{\dot{B}_{p,p}^{k-|\alpha|-1/p}(\mathbb{R}^{n-1})} \stackrel{(b)}{\leq} c \|\mathbb{A}(((\partial^{\beta} \psi_j)u)(F^{(j)})^{-1}))\|_{L^p(\mathbb{H})}, \end{aligned}$$

where $\mathbb{H} = \mathbb{R}^{n-1} \times (0, \infty)$ and c also depends on $\text{diam}(\text{spt}(\psi_j))$. Thus, by the Leibniz rule and the properties of ψ_j and $F^{(j)}$,

$$(4.24) \quad \|\partial^{\alpha} \text{I}_{j,\beta}\|_{B_{p,p}^{k-|\alpha|-1/p}(\mathbb{R}^{n-1})} \leq c(\|u\|_{W^{k-1,p}(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)})$$

where $c = c(\alpha, \beta, \psi_j, F^{(j)}, p, \mathbb{A}, \Omega) > 0$. Based on (4.24) and again by the Leibniz rule, we conclude inductively for suitable choices of β and by the properties of $\psi_j, F^{(j)}$,

$$\|(\psi_j \partial^\alpha u)(F^{(j)})^{-1}(\cdot, 0)\|_{\mathbb{B}_{p,p}^{k-|\alpha|-1/p}(\mathbb{R}^{n-1})} \leq c(\|u\|_{\mathbb{W}^{k-1,p}(\Omega)} + \|\mathbb{A}u\|_{L^p(\Omega)})$$

and (c) follows. Ad (c) \Rightarrow (d). Let $B \subset \mathbb{R}^n$ be an open ball with $\Omega \Subset B$. Because of (c), we may consider the bounded, linear operator $\text{tr}_{k-1, \partial\Omega}^{\mathbb{A}}: \widetilde{\mathbb{W}}^{\mathbb{A},p}(\Omega) \rightarrow \dot{\mathbb{B}}_p^{k-1/p}(\partial\Omega; V)$ from Lemma 4.6. We subsequently pick the bounded and linear operator $\mathcal{L}: \dot{\mathbb{B}}_p^{k-1/p}(\partial(B \setminus \overline{\Omega}); V) \rightarrow \mathbb{W}^{k,p}(B \setminus \overline{\Omega}; V)$ from Lemma 4.2 and fix some $\rho \in C_c^\infty(\mathbb{R}^n; [0, 1])$ with $\mathbb{1}_\Omega \leq \rho \leq \mathbb{1}_B$. Then we define for $u \in \widetilde{\mathbb{W}}^{\mathbb{A},p}(\Omega)$

$$(4.25) \quad \mathcal{E}u := \begin{cases} u & \text{in } \Omega, \\ \rho \mathcal{L}(\mathbb{1}_{\partial\Omega} \text{tr}_{k-1, \partial\Omega}^{\mathbb{A}}(u)) & \text{in } \mathbb{R}^n \setminus \overline{\Omega}. \end{cases}$$

By construction of \mathcal{E} , the Gauß-Green formula (4.23) then implies

$$(4.26) \quad \int_{\mathbb{R}^n} \mathbb{A} \mathcal{E}u \cdot \varphi \, dx = (-1)^k \int_{\mathbb{R}^n} \mathcal{E}u \cdot \mathbb{A}^* \varphi \, dx \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^n; W)$$

and so we obtain $\mathcal{E}u \in \widetilde{\mathbb{W}}^{\mathbb{A},p}(\mathbb{R}^n)$ together with $\|\mathcal{E}u\|_{\widetilde{\mathbb{W}}^{\mathbb{A},p}(\mathbb{R}^n)} \leq c\|u\|_{\widetilde{\mathbb{W}}^{\mathbb{A},p}(\Omega)}$ for all $u \in \widetilde{\mathbb{W}}^{\mathbb{A},p}(\Omega)$ with $c > 0$ solely depending on \mathbb{A}, p and Ω .

Now note that, if (c) holds, then the differential operator \mathbb{A} is necessarily elliptic in the sense of (1.3). This can be seen by a modification of a standard construction (see, e.g., [67] or [19]): If it were not elliptic, we would find $\xi \in \mathbb{R}^n \setminus \{0\}$ and $v \in V \setminus \{0\}$ such that $\mathbb{A}[\xi]v = 0$. For any $h \in C_c^\infty(\mathbb{R})$ and any open and bounded set $\Omega \subset \mathbb{R}^n$, the function $u_h(x) := h(x \cdot \xi)v$ belongs to $\mathbb{W}^{k-1,p}(\Omega; V)$ and satisfies both $\partial^\alpha u_h(x) = h^{(|\alpha|)}(x \cdot \xi)\xi^\alpha v$ and $\mathbb{A}u_h = 0$. We extend $\{\xi\}$ to an orthonormal basis $\{\xi, \eta_2, \dots, \eta_n\}$ and choose an open and bounded set $\Omega \subset \mathbb{R}^n$ with smooth boundary such that $\Gamma := (0, 1)\xi \times (0, 1)\eta_2 \times \dots \times (0, 1)\eta_{n-1} \times \{0\} \subset \partial\Omega$ and $\Omega \subset (-2, 2)\xi \times (-2, 2)\eta_2 \times \dots \times (-2, 2)\eta_n$. Choose a sequence $(h_j) \subset C_c^\infty((-2, 2))$ such that $\sup_j \|h_j\|_{\mathbb{W}^{k-1,p}((-2, 2))} < \infty$ but $\limsup_{j \rightarrow \infty} \|h_j^{(k-1)}\|_{L^q((0, 1))} = \infty$ for $q > p$. Let $\alpha \in \mathbb{N}_0^n$ with $|\alpha| = k-1$ be such that $\xi^\alpha \neq 0$. By construction and Fubini's theorem, $\sup_j \|u_{h_j}\|_{\widetilde{\mathbb{W}}^{\mathbb{A},p}(\Omega)} < \infty$ but, with a constant $c = c(n, \xi, \alpha, v) > 0$, we have for $q > p$

$$(4.27) \quad \int_{\partial\Omega} |\partial^\alpha u_{h_j}|^q \, d^{n-1}x \geq \int_\Gamma |\partial^\alpha u_{h_j}|^q \, d^{n-1}x = c \int_0^1 |h_j^{(k-1)}(t)|^q \, dt \rightarrow \infty$$

at least for a suitable subsequence. Now, if $1 < p < n$, choose $q = \frac{p(n-1)}{n-p}$ and let $q > p$ be arbitrary provided $p \geq n$. Since (c) implies that $(\partial^\alpha u_{h_j}|_{\partial\Omega})$ is bounded in $\mathbb{B}_{p,p}^{1-1/p}(\partial\Omega; V)$ and so, by embedding theorems for Besov spaces, in $L^q(\partial\Omega; V)$, (4.27) yields a contradiction. Therefore, \mathbb{A} must be elliptic.

Now, (4.26)ff. and an argument as in [31, Lem. 6.1] or [19, Prop. 4.1] allow to conclude (d): Let $\alpha \in \mathbb{N}_0^n$ satisfy $|\alpha| = k$. Consider for $\psi \in C_c^\infty(\mathbb{R}^n; W)$ the Fourier multiplication operators $\psi \mapsto \Phi_\alpha(\psi)$, where

$$(4.28) \quad \Phi_\alpha(\psi)(x) := \mathcal{F}^{-1}[\xi^\alpha (\mathbb{A}^*[\xi]\mathbb{A}[\xi])^{-1} \mathbb{A}^*[\xi] \widehat{\psi}(\xi)](x), \quad x \in \mathbb{R}^n.$$

Then we have $\Phi_\alpha(\mathbb{A}\varphi) = \partial^\alpha \varphi$ for $\varphi \in C_c^\infty(\mathbb{R}^n; V)$. Since \mathbb{A} is elliptic, $\mathbb{A}[\xi]: V \rightarrow W$ is injective and hence $\mathbb{A}^*[\xi]\mathbb{A}[\xi]: V \rightarrow V$ is bijective for any $\xi \in \mathbb{R}^n \setminus \{0\}$; in particular, it is bijective for any $\xi \in \mathbb{R}^n \setminus \{0\}$, and so (4.28) is well-defined. The Fourier multiplier

$$(4.29) \quad \mathbb{R}^n \setminus \{0\} \ni \xi \mapsto \mathbf{m}_\alpha^\mathbb{A}(\xi) := \xi^\alpha (\mathbb{A}^*[\xi]\mathbb{A}[\xi])^{-1} \mathbb{A}^*[\xi] \in \mathcal{L}(W; W)$$

is of class $C^\infty(\mathbb{R}^n \setminus \{0\}; \mathcal{L}(W; V))$, homogeneous of degree zero, and so [25, Thm. 4.13] yields that Φ_α extends to a bounded operator $\Phi_\alpha: L^p(\mathbb{R}^n; W) \rightarrow L^p(\mathbb{R}^n; V)$. Thus, (4.30)

$$\begin{aligned} \|u\|_{W^{k,p}(\Omega)} &\leq \|\mathcal{E}u\|_{W^{k-1,p}(\mathbb{R}^n)} + \|\nabla^k \mathcal{E}u\|_{L^p(\mathbb{R}^n)} \\ &\leq \|\mathcal{E}u\|_{W^{k-1,p}(\mathbb{R}^n)} + c\|\mathbb{A}\mathcal{E}u\|_{L^p(\mathbb{R}^n)} \leq c\|\mathcal{E}u\|_{\widetilde{W}^{\mathbb{A},p}(\mathbb{R}^n)} \stackrel{(4.26)ff.}{\leq} c\|u\|_{\widetilde{W}^{\mathbb{A},p}(\Omega)} \end{aligned}$$

for all $u \in W^{\mathbb{A},p}(\Omega)$ with some constant $c = c(\mathbb{A}, p, \rho, \Omega) > 0$. This is (d). Now, if (d) holds, we apply Lemma 4.5 to $E_1 = W^{k,p}(\Omega; V)$, $E_2 = L^p(\Omega; W)$, $E_3 = W^{k-1,p}(\Omega; V)$, $A = \mathbb{A}$ and B being the compact embedding $W^{k,p}(\Omega; V) \hookrightarrow W^{k-1,p}(\Omega; V)$. The equivalence of the $W^{k,p}$ - and $\widetilde{W}^{\mathbb{A},p}$ -norms implied by (d) now yields $\dim(\ker(A)) < \infty$ by virtue of Lemma 4.5, and by (3.6)ff., this implies that \mathbb{A} is \mathbb{C} -elliptic. Hence (a) follows, and so all of (a)–(d) are equivalent.

Lastly, by equivalence of (a) and (d), (4.21) follows if we can establish $W^{\mathbb{A},p}(\Omega) \hookrightarrow \widetilde{W}^{\mathbb{A},p}(\Omega)$ provided any of the mutually equivalent conditions (a)–(d) holds. We thus assume that \mathbb{A} is \mathbb{C} -elliptic and assume without loss of generality that Ω is an open, connected and bounded domain with C^∞ -boundary. Since every such domain is John, the Poincaré-type inequality (3.20) now yields $W^{\mathbb{A},p}(\Omega) \hookrightarrow \widetilde{W}^{\mathbb{A},p}(\Omega)$, and the proof is complete. \square

By Theorem 4.4 and Lemma 4.6, we obtain the following theorem for $1 < p < \infty$; for $p = 1$ and $BV^{\mathbb{A}}$, $C^\infty \cap BV^{\mathbb{A}}$ in $BV^{\mathbb{A}}$ is only dense in $BV^{\mathbb{A}}$ for the \mathbb{A} -strict metric and not the norm topology, and so the underlying approximation must be approached differently. For the reader's convenience, the precise argument (hinging on a multiplicative inequality) is given in the Appendix, Section 7.1.

Theorem 4.7 (Trace operator). *Let $\Omega \subset \mathbb{R}^n$ be open and bounded with boundary $\partial\Omega$ of class C^∞ . Then, for any k -th order \mathbb{C} -elliptic differential operator \mathbb{A} of the form (1.1) and $1 \leq p < \infty$, there exists a bounded and linear trace operator*

$$\mathrm{Tr}_{\partial\Omega}: W^{\mathbb{A},p}(\Omega) \rightarrow B_{p,p}^{k-1/p}(\partial\Omega; V)$$

which is onto. Moreover, for $p = 1$, there exists a linear trace operator

$$\mathrm{Tr}_{\partial\Omega}: BV^{\mathbb{A}}(\Omega) \rightarrow B_{1,1}^{k-1}(\partial\Omega; V)$$

which is onto and continuous for the \mathbb{A} -strict metric defined by (3.2).

5. PROOF OF THEOREM 1.2: THE TRACE ESTIMATE VIA REDUCTION TO $W^{k,1}(\Omega; V)$

Let $p = 1$. In comparison with the previous section, we now establish the novel reduction principle (1.9) which leads to the trace estimates underlying Theorem 1.2. This manifests the metaprinciple of \mathbb{C} -ellipticity helping to overcome ORNSTEIN's Non-Inequality by passing to *inverse estimates for polynomials*. Throughout this section, we hereafter suppose that \mathbb{A} is a \mathbb{C} -elliptic differential operator of the form (1.1) with $k \geq 2$.

5.1. Geometric setup. Throughout this section, we suppose that $\Omega \subset \mathbb{R}^n$ is an open and bounded NTA_{n-1} domain, cf. Definition 2.1. To arrive at the conclusion of Theorem 1.2, we borrow and adapt some technology from the precursor [11] to the present paper. Given $j \in \mathbb{Z}$, pick a countable covering $(B_{j,i})_{i \in \mathbb{N}}$ of \mathbb{R}^n by balls, each having diameter $\mathrm{diam}(B_{j,i}) = 2r(B_{j,i})$ and satisfying each of the following properties:

- (C1) For all $j \in \mathbb{Z}, i \in \mathbb{N}$, $2^{-j-4} \leq \ell(B_{j,i}) \leq 2^{-j-3}$,
- (C2) For each $j \in \mathbb{Z}$, $\bigcup_{i \in \mathbb{N}} \frac{7}{8}B_{j,i} = \mathbb{R}^n$,

(C3) There exists $c > 0$ such that $\sup_{j \in \mathbb{Z}} \sum_{i \in \mathbb{N}} \mathbb{1}_{B_{j,i}} \leq c$.

In the sequel, we let $\Omega_j := \{x \in \Omega : \text{dist}(x, \partial\Omega) < 2^{-j}\}$. Since Ω is a non-tangentially accessible domain and thereby satisfies the interior corkscrew as well as the Harnack chain condition, we may record the following further properties, see [11]:

(C4) There exists an index $j_0 \in \mathbb{Z}$ such that for any ball $B_{j,i}$ with $j \geq j_0$, $i \in \mathbb{N}$ and $B_{j,i} \cap \Omega_j \neq \emptyset$, there exists a *reflected ball* $B_{j,i}^\# \subset \Omega$ satisfying

$$\text{diam}(B_{j,i}) \simeq \text{diam}(B_{j,i}^\#) \simeq \text{dist}(B_{j,i}^\#, \partial\Omega), \quad \text{dist}(B_{j,i}, B_{j,i}^\#) \lesssim \text{diam}(B_{j,i}).$$

Here, the constants implicit in ' \simeq ' or ' \lesssim ' are independent of j and i .

(C5) With j_0 as in (C4), if $B_{j,i} \subset \Omega$ and $B_{j,i} \cap \Omega_j \neq \emptyset$ for some $j \geq j_0$, then there exists a chain of balls $\mathbb{B}_1, \dots, \mathbb{B}_\gamma \subset \Omega$ with $\gamma \in \mathbb{N}$ independent of j, i such that

(a) $\mathbb{B}_1 = B_{j,i}, \mathbb{B}_\gamma = B_{j,i}^\#$,

(b) $\mathcal{L}^n(\mathbb{B}_\beta \cap \mathbb{B}_{\beta+1}) \simeq \mathcal{L}^n(\mathbb{B}_\beta) \simeq \mathcal{L}^n(\mathbb{B}_{\beta+1}) \simeq \mathcal{L}^n(B_{j,i})$ for all $\beta \in \{1, \dots, \gamma-1\}$,

(c) $\text{diam}(\mathbb{B}_\beta) \simeq \text{diam}(B_{j,i})$ for all $\beta \in \{1, \dots, \gamma\}$.

Here, the constants implicit in ' \simeq ' are independent of j, i and β . In this situation, we define the *chain set* $\text{ch}(B_{j,i}, B_{j,i}^\#) = \bigcup_{\beta=1}^\gamma \mathbb{B}_\beta$.

(C6) If j_0 is as in (C4) and $j, l \in \mathbb{Z}$ are such that $j, l \geq j_0$ and $|j-l| \leq 1$, then for all balls $B_{j,i}$ and $B_{l,m}$ with $B_{j,i} \cap B_{l,m} \neq \emptyset$, $B_{j,i} \cap \Omega_j \neq \emptyset$, $B_{l,m} \cap \Omega_l \neq \emptyset$ there exists a chain of balls $\mathbb{B}_1, \dots, \mathbb{B}_\gamma \subset \Omega$ with γ independent of j, l, m, i such that

(a) $\mathbb{B}_1 = B_{j,i}, \mathbb{B}_\gamma = B_{l,m}$,

(b) $\mathcal{L}^n(\mathbb{B}_\beta \cap \mathbb{B}_{\beta+1}) \simeq \mathcal{L}^n(\mathbb{B}_\beta) \simeq \text{diam}(B_{j,i})$ for all $\beta \in \{1, \dots, \gamma-1\}$,

(c) $\text{dist}(\mathbb{B}_\beta, \partial\Omega) \simeq \text{diam}(\mathbb{B}_\beta) \simeq \text{diam}(B_{j,i})$ for all $\beta \in \{1, \dots, \gamma\}$.

Here, the constants implicit in ' \simeq ' are independent of j, i and β , and we define similarly as above the respective chain set $\text{ch}(B_{j,i}, B_{l,m}) := \bigcup_{\beta=1}^\gamma \mathbb{B}_\beta$.

Working from (C4)–(C6), we further obtain the existence of an $i_0 \in \mathbb{N}_{\geq 2}$ such that for all $j \geq j_0$ there holds with a universal constant $c > 0$

$$(5.1) \quad \begin{aligned} \sum_{\substack{m: \\ B_{j,m} \cap \Omega_j \neq \emptyset}} \mathbb{1}_{B_{j,m}^\#} &\leq c \mathbb{1}_{\Omega_{j-i_0} \setminus \Omega_{j+i_0}}, \\ \sum_{\substack{m: \\ B_{j,m} \cap \Omega_j \neq \emptyset}} \sum_{\substack{i: \\ B_{j+1,i} \cap B_{j,m} \neq \emptyset \\ B_{j+1,i} \cap \Omega_{j+1} \neq \emptyset}} \mathbb{1}_{\text{ch}(B_{j,m}^\#, B_{j+1,i}^\#)} &\leq c \mathbb{1}_{\Omega_{j-i_0} \setminus \Omega_{j+i_0}}, \\ \sum_{\substack{m: \\ B_{j,m} \cap (\Omega_j \setminus \Omega_{j+2}) \neq \emptyset}} \mathbb{1}_{\text{ch}(B_{j,m}, B_{j,m}^\#)} &\leq c \mathbb{1}_{\Omega_{j-i_0} \setminus \Omega_{j+i_0}}. \end{aligned}$$

For (5.1)₃, note that if $B_{j,m} \cap (\Omega_j \setminus \Omega_{j+2}) \neq \emptyset$, then we have $B_{j,m} \subset \Omega$ by construction. Given $u \in W^{\mathbb{A},1}(\Omega)$ and $j \geq j_0$, $i \in \mathbb{N}$, we introduce the shorthand notation

$$(5.2) \quad \Pi_{j,i} u := \Pi_{\mathbb{A}}^{B_{j,i}^\#} u,$$

with the projection $\Pi_{\mathbb{A}}^{B_{j,i}^\#}$ from Proposition 3.3. The proof of the next lemma is accomplished by an argument similar to that underlying Proposition 3.8 and is therefore omitted.

Lemma 5.1. *There exists a constant $c > 0$ that only depends on \mathbb{A} and the NTA-parameters of Ω such that the following hold for all $j \geq j_0$ and all $u \in W^{\mathbb{A},1}(\Omega)$:*




FIGURE 3. Construction of T_j and idea of the (trace) estimates of Proposition 5.2 and Corollary 5.3. With the dashed balls indicating the covering $(B_{j,i})_i$, the approximation $T_j u$ given by (5.5) splits into two parts: At a distance at least 2^{-j} from $\partial\Omega$ (i.e., in the red area), $T_j u$ leaves u unchanged, whereas close to $\partial\Omega$ (i.e., on $\Omega_j := \{\text{dist}(\cdot, \partial\Omega) < 2^{-j}\}$ represented by the grey area), $T_j u$ essentially replaces u by its projections onto the nullspace of \mathbb{A} on the reflected balls $B_{j,i}^\sharp$.

(a) For all $B_{j,i}$ with $B_{j,i} \cap \Omega_j \neq \emptyset$ and all $\alpha \in \mathbb{N}_0^n$ there holds

$$\|\partial^\alpha \Pi_{j,i} u\|_{L^\infty(B_{j,i})} \leq c \text{diam}(B_{j,i})^{-|\alpha|} \int_{B_{j,i}^\sharp} |u| \, dx.$$

(b) Whenever $B_{j,m} \cap (\Omega_j \setminus \Omega_{j+2}) \neq \emptyset$, then $B_{j,m} \subset \Omega$ and we have, for all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq k-1$,

$$\|\partial^\alpha (u - \Pi_{j,m} u)\|_{L^1(B_{j,m})} \leq c \text{diam}(B_{j,m})^{k-|\alpha|} \int_{\text{ch}(B_{j,m}, B_{j,m}^\sharp)} |\mathbb{A}u| \, dx.$$

(c) Chain control: Whenever $B_{j,m} \cap \Omega_j \neq \emptyset$, $B_{j+1,i} \cap \Omega_{j+1} \neq \emptyset$ and $B_{j+1,i} \cap B_{j,m} \neq \emptyset$, then we have for all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq k-1$

$$\int_{B_{j,m}} |\partial^\alpha (\Pi_{j+1,i} u - \Pi_{j,m} u)| \, dx \leq c \text{diam}(B_{j,m})^{k-|\alpha|} \int_{\text{ch}(B_{j+1,i}^\sharp, B_{j,m}^\sharp)} |\mathbb{A}u| \, dx.$$

5.2. Proof of Theorem 1.2. In view of the requisite trace estimates, we now introduce a suitable approximation $T_j u$ of a given function $u \in W^{\mathbb{A},1}(\Omega)$. For each $j \in \mathbb{Z}$, pick the covering $(B_{j,i})_i$ from Section 5.1 and let $(\rho_{j,i})_i$ be a smooth partition

of unity subject to $(B_{j,i})_i$ such that, with a universal constant $c > 0$,

$$(5.3) \quad \sum_{|\alpha| \leq k} \text{diam}(B_{j,i})^{|\alpha|} \|\partial^\alpha \rho_{j,i}\|_{L^\infty(\Omega)} \leq c \quad \text{for all } i \in \mathbb{N}, j \in \mathbb{Z}.$$

Moreover, let $\rho_j \in C^\infty(\overline{\Omega}; [0, 1])$ be such that $\mathbb{1}_{\Omega_{j+1}} \leq \rho_j \leq \mathbb{1}_{\Omega_j}$ and

$$(5.4) \quad \|\partial^\alpha \rho_j\|_{L^\infty(\Omega)} \leq c 2^{j|\alpha|} \quad \text{for all } \alpha \in \mathbb{N}_0^n, |\alpha| \leq k.$$

We then define the *replacement sequence* $(T_j u)_{j \geq j_0}$ with $j_0 \in \mathbb{Z}$ as in (C4) from above by

$$(5.5) \quad T_j u := T_j^{(1)} u + T_j^{(2)} u := (1 - \rho_j)u + \rho_j \sum_{i \in \mathbb{N}} \rho_{j,i} \Pi_{j,i} u$$

in Ω . Note that, as a locally finite sum and by construction, $T_j^{(2)} u \in C^\infty(\overline{\Omega}; V)$. Thus, in passing from u to $T_j u$ we essentially leave the map u unchanged away from $\partial\Omega$ while close to $\partial\Omega$, we replace it by its projections onto $\ker(\mathbb{A})$ on balls nearby, see Figure 3. The proof of Theorem 1.2 crucially hinges on the decomposition

$$(5.6) \quad \begin{aligned} T_{j+1} u - T_j u &= (\rho_j - \rho_{j+1}) \sum_{m \in \mathbb{N}} \rho_{j,m} (u - \Pi_{j,m} u) \\ &+ \rho_{j+1} \sum_{i, m \in \mathbb{N}} \rho_{j,m} \rho_{j+1,i} (\Pi_{j+1,i} u - \Pi_{j,m} u) =: \text{I}_j[u] + \text{II}_j[u], \end{aligned}$$

which is implied by $\sum_m \rho_{j,m} \equiv 1$ for all j . In comparison with the precursor [11], the key observation is now given by Proposition 5.2 below: The terms $\text{I}_j[u]$ will have trace zero along $\partial\Omega$ and $\text{II}_j[u]$ uniformly belongs to $W^{k,1}(\Omega; V)$. For the reduction argument underlying the proof of Theorem 1.2, it will then suffice to apply the trace estimate for $W^{k,1}$ to $\text{II}_j[u]$, cf. Corollary 5.3.

Proposition 5.2. *Let $u \in W^{\mathbb{A},1}(\Omega)$ and define $T_j u$, $\text{I}_j[u]$ and $\text{II}_j[u]$ for $j \geq j_0$ by (5.5) or (5.6), respectively, and let i_0 be as in (5.1). Then the following holds:*

- (a) *There exists $c > 0$ such that $\|\text{I}_j[u]\|_{W^{\mathbb{A},1}(\Omega)} \leq c \|\mathbb{A}u\|_{L^1(\Omega_{j-i_0} \setminus \Omega_{j+i_0})}$.*
- (b) *There exists $c > 0$ such that $\|\text{II}_j[u]\|_{W^{k,1}(\Omega)} \leq c \|\mathbb{A}u\|_{L^1(\Omega_{j-i_0} \setminus \Omega_{j+i_0})}$.*
- (c) *We have $T_j u \rightarrow u$ with respect to the $W^{\mathbb{A},1}$ -norm.*
- (d) *There exists $(u_j) \subset C^\infty(\overline{\Omega}; V)$ such that $u_j \rightarrow u$ with respect to the $W^{\mathbb{A},1}$ -norm.*
- (e) *If $u \in C(\overline{\Omega}; V) \cap W^{\mathbb{A},1}(\Omega)$, then $T_j u \rightarrow u$ uniformly in Ω .*

In both (a) and (b), the constant $c > 0$ is independent of u .

Proof. Let $u \in W^{\mathbb{A},1}(\Omega)$. Ad (a). For $j \geq j_0$, denote \mathcal{I}_j the set of all indices $m \in \mathbb{N}$ such that $(\rho_j - \rho_{j+1})\rho_{j,m} \neq 0$. Since $\text{spt}(\rho_j - \rho_{j+1}) \subset \Omega_j \setminus \Omega_{j+2}$, $m \in \mathcal{I}_j$ implies that $B_{j,m} \cap (\Omega_j \setminus \Omega_{j+2}) \neq \emptyset$. For such indices, we may use Lemma 5.1 (b). In consequence,

$$\begin{aligned} \|\text{I}_j[u]\|_{W^{\mathbb{A},1}(\Omega)} &\leq \left(\sum_{m \in \mathcal{I}_j} \sum_{l=0}^{k-1} \|D^l((\rho_j - \rho_{j+1})\rho_{j,m}(u - \Pi_{j,m} u))\|_{L^1(\Omega)} \right) \\ &+ \left(\sum_{m \in \mathcal{I}_j} \|\mathbb{A}((\rho_j - \rho_{j+1})\rho_{j,m}(u - \Pi_{j,m} u))\|_{L^1(\Omega)} \right) \\ &\stackrel{(*)}{\leq} c \left(\sum_{m \in \mathcal{I}_j} \sum_{l=0}^{k-1} \sum_{|\alpha|+|\beta|=l} \|\partial^\alpha((\rho_j - \rho_{j+1})\rho_{j,m})\|_{L^\infty(\Omega)} \|\partial^\beta(u - \Pi_{j,m} u)\|_{L^1(B_{j,m})} \right) \end{aligned}$$

$$\begin{aligned}
& + c \left(\sum_{m \in \mathcal{I}_j} \sum_{\substack{|\alpha|+|\beta|=k \\ |\beta| \leq k-1}} \|\partial^\alpha((\rho_j - \rho_{j+1})\rho_{j,m})\|_{L^\infty(\Omega)} \|\partial^\beta(u - \Pi_{j,m}u)\|_{L^1(B_{j,m})} \right. \\
& \quad \left. + \sum_{m \in \mathcal{I}_j} \|(\rho_j - \rho_{j+1})\rho_{j,m}\mathbb{A}u\|_{L^1(\Omega)} \right) \\
\stackrel{\text{Lem. 5.1(b),(5.3),(5.4)}}{\leq} & c \left(\sum_{m \in \mathcal{I}_j} \sum_{\ell=0}^{k-1} \sum_{|\alpha|+|\beta|=\ell} 2^{j|\alpha|} 2^{-j(k-|\beta|)} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j,m}, B_{j,m}^\#))} \right) \\
& + c \left(\sum_{m \in \mathcal{I}_j} \sum_{\substack{|\alpha|+|\beta|=k \\ |\beta| \leq k-1}} 2^{j|\alpha|} 2^{-j(k-|\beta|)} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j,m}, B_{j,m}^\#))} + \sum_{m \in \mathcal{I}_j} \|\mathbb{A}u\|_{L^1(B_{j,m})} \right) \\
& \leq c \sum_{m \in \mathcal{I}_j} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j,m}, B_{j,m}^\#))} \\
& \stackrel{(5.1)_3}{\leq} c \|\mathbb{A}u\|_{L^1(\Omega_{j-i_0} \setminus \Omega_{j+i_0})}.
\end{aligned}$$

Here we used at (*) that, if we expand $\mathbb{A}(\varphi v)$ for sufficiently regular functions $\varphi: \Omega \rightarrow \mathbb{R}$ and $v: \Omega \rightarrow V$ by the Leibniz rule, then the only summand that involves k -th order derivatives of v is of the form $\varphi \mathbb{A}v$. Hence, (a) follows. Ad (b). Let \mathcal{J}_j^m be the set of all indices $i \in \mathbb{N}$ such that $\rho_{j+1}\rho_{j,m}\rho_{j+1,i} \neq 0$. Then, by (C3), $\#\mathcal{J}_j^m$ is uniformly bounded in j and m . If $i \in \mathcal{J}_j^m$, then $\Omega_{j+1} \cap B_{j,m} \neq \emptyset$, $\Omega_{j+1} \cap B_{j+1,i} \neq \emptyset$ and $B_{j,m} \cap B_{j+1,i} \neq \emptyset$. As such, (5.1)₂ is available for such indices. Let $|\beta| \leq k$. Since $\ker(\mathbb{A})$ is a finite dimensional space by the \mathbb{C} -ellipticity of \mathbb{A} , we may invoke the inverse estimates (3.16)₂ on elements of the nullspace of \mathbb{A} to obtain for all $j \geq j_0$, $m \in \mathbb{N}$ and $i \in \mathcal{J}_j^m$:

$$\begin{aligned}
& \|\partial^\beta(\Pi_{j+1,i}u - \Pi_{j,m}u)\|_{L^1(B_{j,m})} \leq \|D^{|\beta|}(\Pi_{j+1,i}u - \Pi_{j,m}u)\|_{L^1(B_{j,m})} \\
& \leq C \text{diam}(B_{j,m})^{n-|\beta|} \text{diam}(B_{j,m})^{|\beta|} \int_{B_{j,m}} |D^{|\beta|}(\Pi_{j+1,i}u - \Pi_{j,m}u)| \, dx \\
& \leq C \text{diam}(B_{j,m})^{n-|\beta|} \times \\
& \quad \times \left(\sum_{\ell=0}^k \text{diam}(B_{j,m})^\ell \int_{B_{j,m}} |D^\ell(\Pi_{j+1,i}u - \Pi_{j,m}u)| \, dx \right) \\
& \stackrel{(3.16)_2}{\leq} C \text{diam}(B_{j,m})^{n-|\beta|} \int_{B_{j,m}} |\Pi_{j+1,i}u - \Pi_{j,m}u| \, dx \\
& \stackrel{\text{Lem. 5.1(c)}}{\leq} C \text{diam}(B_{j,m})^{(k-|\beta|)} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j+1,i}^\#, B_{j,m}^\#))} \\
& \leq C 2^{-j(k-|\beta|)} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j+1,i}^\#, B_{j,m}^\#))}.
\end{aligned}$$

As \mathcal{J}_j^m is non-empty only if $m \in \mathcal{K}_j := \{l \in \mathbb{N} : \Omega_{j+1} \cap B_{j,l} \neq \emptyset\}$, we obtain by (5.3), (5.4) and the preceding inequality

$$\begin{aligned}
\|\Pi_j[u]\|_{W^{k,1}(\Omega)} & \leq c \sum_{m \in \mathcal{K}_j} \sum_{i \in \mathcal{J}_j^m} \sum_{\ell=0}^k \sum_{|\alpha|+|\beta|=\ell} \left(\|\partial^\alpha(\rho_{j+1}\rho_{j,m}\rho_{j+1,i})\|_{L^\infty(\Omega)} \times \right. \\
& \quad \left. \times \|\partial^\beta(\Pi_{j+1,i}u - \Pi_{j,m}u)\|_{L^1(B_{j,m})} \right) \\
& \leq c \sum_{m \in \mathcal{K}_j} \sum_{i \in \mathcal{J}_j^m} \sum_{\ell=0}^k \sum_{|\alpha|+|\beta|=\ell} 2^{j|\alpha|} 2^{-j(k-|\beta|)} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j+1,i}^\#, B_{j,m}^\#))}
\end{aligned}$$

$$\begin{aligned}
&\leq c \sum_{m \in \mathcal{K}_j} \sum_{i \in \mathcal{J}_j^m} \sum_{\ell=0}^k 2^{(\ell-k)j} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j+1,i}^\#, B_{j,m}^\#))} \\
&\leq c \sum_{m \in \mathcal{K}_j} \sum_{i \in \mathcal{J}_j^m} \|\mathbb{A}u\|_{L^1(\text{ch}(B_{j+1,i}^\#, B_{j,m}^\#))} \\
&\stackrel{(5.1)_2}{\leq} c \|\mathbb{A}u\|_{L^1(\Omega_{j-i_0} \setminus \Omega_{j+i_0})}
\end{aligned}$$

This establishes (b). Ad (c). By construction of the sets Ω_j , $\sum_{j \geq l} \mathbb{1}_{\Omega_{j-i_0} \setminus \Omega_{j+i_0}} \leq c(i_0) \mathbb{1}_{\Omega_{l-i_0}}$ for all $l \in \mathbb{N}$. Since $\|\cdot\|_{W^{\mathbb{A},1}}$ always can be dominated by $\|\cdot\|_{W^{k,1}}$, we obtain by (a) and (b) for all $l' \geq l \geq j_0$ by virtue of a telescope sum argument

$$\|T_{l'}u - T_lu\|_{W^{\mathbb{A},1}(\Omega)} \leq \sum_{j=l}^{\infty} \|T_{j+1}u - T_ju\|_{W^{\mathbb{A},1}(\Omega)} \leq c \|\mathbb{A}u\|_{L^1(\Omega_{l-i_0})} \rightarrow 0$$

as $l, l' \rightarrow \infty$. Thus (T_lu) is Cauchy in $W^{\mathbb{A},1}(\Omega)$ and thus converges to some $v \in W^{\mathbb{A},1}(\Omega)$. On the other hand, since by construction there holds $\text{spt}(u - T_lu) \subset \Omega_l$, we have $\|u - T_lu\|_{W^{\mathbb{A},1}(\omega)} \rightarrow 0$ for any open $\omega \Subset \Omega$ as $l \rightarrow \infty$, and so we must have $v = u$. Therefore, $T_ju \rightarrow u$ in $W^{\mathbb{A},1}(\Omega)$ as $j \rightarrow \infty$, which is (c).

Ad (d). We recall the decomposition (5.5). Since $T_j^{(2)}u$ is a locally finite sum of C^∞ -maps, $T_j^{(2)}u \in C^\infty(\overline{\Omega}; V)$. Moreover, since $T_j^{(1)}u \in W^{\mathbb{A},1}(\Omega)$ is compactly supported in Ω , standard mollification yields some $v_j \in C_c^\infty(\Omega; V)$ such that $\|T_j^{(1)}u - v_j\|_{W^{\mathbb{A},1}(\Omega)} < \frac{1}{j}$. Put $u_j := v_j + T_j^{(2)}u$ so that $u_j \in C^\infty(\overline{\Omega}; V)$, and then (c) implies $\|u - u_j\|_{W^{\mathbb{A},1}(\Omega)} \rightarrow 0$ as $j \rightarrow \infty$.

To see (e), let $\varepsilon > 0$. By uniform continuity of u , we find $\delta > 0$ such that $|x - y| < \delta$ implies $|u(x) - u(y)| < \varepsilon$. Let $B_{j,i}$ be such that $B_{j,i} \cap \Omega_j \neq \emptyset$. By (C4) and assuming that $j \geq j_0$, there exists a constant $C > 0$ independent of j and i such that $|x - y| < C2^{-j}$ holds for all $x \in B_{j,i}$ and $y \in B_{j,i}^\#$; moreover, we may assume that $\text{diam}(B_{j,i}^\#) < C2^{-j}$ as well. We thus find $J_0 \geq j_0$ such that for all $j \geq J_0$ there holds $C2^{-j} \leq \delta$. Thus, invoking Lemma 5.1(a) and because of $(u)_{B_{j,i}^\#} = \Pi_{j,i}(u)_{B_{j,i}^\#}$,

$$\begin{aligned}
\|\rho_j \rho_{j,i}(u - \Pi_{j,i}u)\|_{L^\infty(\Omega)} &\leq \|u - (u)_{B_{j,i}^\#}\|_{L^\infty(B_{j,i} \cap \Omega_j)} + \|\Pi_{j,i}(u - (u)_{B_{j,i}^\#})\|_{L^\infty(B_{j,i} \cap \Omega_j)} \\
&\leq \sup_{x \in B_{j,i} \cap \Omega_j} \int_{B_{j,i}^\#} |u(x) - u(y)| \, dy + c \int_{B_{j,i}^\#} |u - (u)_{B_{j,i}^\#}| \, dy \\
&\leq (1 + c)\varepsilon.
\end{aligned}$$

Since $u - T_ju = \sum_i \rho_j \rho_{j,i}(u - \Pi_{j,i}u)$ and the $B_{j,i}$'s have mutual uniformly finite overlap (in i for each fixed j), we thus obtain $T_ju \rightarrow u$ in $L^\infty(\Omega; V)$ and hence (e) follows. The proof is complete. \square

For the following, let $(\mathcal{X}(\partial\Omega; V); \|\cdot\|_{\mathcal{X}(\partial\Omega)})$ be a Banach space with $\mathcal{X}(\partial\Omega; V) \subset L_{\text{loc}}^1(\partial\Omega; V)$ such that $W^{k,1}(\Omega; V)$ has trace space $\mathcal{X}(\partial\Omega; V)$ (cf. Theorem 1.2 ff. for this terminology); we denote $\text{Tr}_{\partial\Omega}: W^{k,1}(\Omega; V) \rightarrow \mathcal{X}(\partial\Omega; V)$ the underlying trace operator.

Corollary 5.3. *There exists a constant $c = c(\Omega, \mathbb{A}) > 0$ such that the following holds: Whenever $u \in C^\infty(\overline{\Omega}; V)$ and $j \geq j_0$ (with $j_0 \in \mathbb{Z}$ as in (C4)), then there holds*

$$(5.7) \quad \sum_{j=j_0}^{\infty} \|\text{Tr}_{\partial\Omega}(T_{j+1}u) - \text{Tr}_{\partial\Omega}(T_ju)\|_{\mathcal{X}(\partial\Omega)} \leq c \|\mathbb{A}u\|_{L^1(\Omega)}.$$

Proof. Let $u \in C^\infty(\overline{\Omega}; V)$ and $j \geq j_0$. Then (5.6) implies that $T_{j+1}u - T_ju$ is a locally finite sum of $C^\infty(\overline{\Omega}; V)$ -maps and so is of class $C^\infty(\overline{\Omega}; V)$, too. Since $\text{spt}(I_j[u]) \subset \{x \in \Omega: \text{dist}(x, \partial\Omega) > 2^{-j-2}\}$ because of $\text{spt}(\rho_j - \rho_{j+1}) \subset \Omega_j \setminus \Omega_{j+2}$, we have $\text{Tr}_{\partial\Omega}(I_j[u]) = 0$. Since $W^{k,1}(\Omega; V)$ has trace space $\mathcal{X}(\partial\Omega; V)$, there exists a constant $c > 0$ such that $\|\text{Tr}_{\partial\Omega}(v)\|_{\mathcal{X}(\partial\Omega)} \leq c\|v\|_{W^{k,1}(\Omega)}$ for all $v \in W^{k,1}(\Omega; V)$. Now, by (5.6) and Proposition 5.2 (b), this entails

$$\begin{aligned}
(5.8) \quad \sum_{j=j_0}^{\infty} \|\text{Tr}_{\partial\Omega}(T_{j+1}u) - \text{Tr}_{\partial\Omega}(T_ju)\|_{\mathcal{X}(\partial\Omega)} &= \sum_{j=j_0}^{\infty} \|\text{Tr}_{\partial\Omega}(\Pi_j[u])\|_{\mathcal{X}(\partial\Omega)} \\
&\leq c \sum_{j=j_0}^{\infty} \|\Pi_j[u]\|_{W^{k,1}(\Omega)} \\
&\stackrel{\text{Prop. 5.2 (b)}}{\leq} c \sum_{j=j_0}^{\infty} \|\mathbb{A}u\|_{L^1(\Omega_{j-i_0} \setminus \Omega_{j+i_0})} \\
&\leq c\|\mathbb{A}u\|_{L^1(\Omega_{j_0-i_0})},
\end{aligned}$$

where in the ultimate step we used that the sets $\Omega_{j-i_0} \setminus \Omega_{j+i_0}$ only have a uniformly finite mutual overlap. This is (5.7), and the proof is complete. \square

Before we proceed, we briefly pause to comment on the preceding proof in

Remark 5.4. Based on Proposition 5.2 (b), in the second line of (5.8) we have employed *full k -th order gradient L^1 -estimates* to obtain the requisite $\mathcal{X}(\partial\Omega; V)$ -estimates for the traces of $T_{j+1}u - T_ju$. We wish to emphasize that a similar strategy *does not* yield any uniform L^1 -bounds on the sequence $(D^k(T_{j+1}u - T_ju))_j$, and so this is in line with ORNSTEIN'S Non-Inequality. Indeed, even for $u \in C^\infty(\overline{\Omega}; V)$ the term $I_j[u]$ in (5.6), for which Proposition 5.2 only provides $W^{\mathbb{A},1}$ - but not $W^{k,1}$ -bounds, might have diverging $W^{k,1}$ -norm as $j \rightarrow \infty$: Precisely by ORNSTEIN'S Non-Inequality, when D^k acts on u in the term $I_j[u]$ in (5.6), one has $\limsup_{j \rightarrow \infty} \|I_j[u]\|_{W^{k,1}(\Omega)} = \infty$ in general. By the inverse estimates on polynomials of a fixed degree, this obstruction becomes invisible for $\Pi_j[u]$, and only $\Pi_j[u]$ matters for the requisite trace estimate (5.7).

Towards the proof of Theorem 1.2, let us note that it is only the sufficiency part that requires proof; similarly as for Theorem 1.1, the necessity part can be established by localising the construction in [31, Proof of Theorem 5.2]. Indeed, if \mathbb{A} is not \mathbb{C} -elliptic, the aforementioned construction leads to some smoothly bounded $\Omega \subset \mathbb{R}^n$ and a $W^{\mathbb{A},1}$ -bounded sequence $(u_j) \subset C^\infty(\overline{\Omega}; V)$ for which $(D^{k-1}u_j)$ is unbounded in $L^1(\partial\Omega; \odot^{k-1}(\mathbb{R}^n; V))$. Since $W^{k,1}(\Omega; V)$ has trace space $B_{1,1}^{k-1}(\Omega; V)$, $W^{\mathbb{A},1}(\Omega)$ cannot have the same trace space as $W^{k,1}(\Omega)$. It thus suffices to give the

Proof of the sufficiency part of Theorem 1.2. Let \mathbb{A} be a \mathbb{C} -elliptic operator and let $u \in C^\infty(\overline{\Omega}; V)$ be given. Since $u \in C^\infty(\overline{\Omega}; V)$, Lemma 5.2 (e) implies by a telescope sum argument

$$(5.9) \quad \text{Tr}_{\partial\Omega}(u) - \text{Tr}_{\partial\Omega}(T_{j_0}u) = \sum_{j=j_0}^{\infty} \text{Tr}_{\partial\Omega}(T_{j+1}u) - \text{Tr}_{\partial\Omega}(T_ju)$$

everywhere on $\partial\Omega$. Based on the decomposition (5.5), we recall that for $u \in C^\infty(\overline{\Omega}; V)$ there holds $\|\text{Tr}_{\partial\Omega}(T_{j_0}^{(1)}u)\|_{\mathcal{X}(\partial\Omega)} = 0$, and so we proceed by proving that the function $T_{j_0}^{(2)}u$ belongs to $W^{k,1}(\Omega; V)$ together with the requisite estimates. Since $T_{j_0}^{(2)}u \in C^\infty(\overline{\Omega}; V)$, it possesses classical traces and so it suffices to




FIGURE 4. Idea of the trace estimate of Theorem 1.2. In the telescope sum (5.9), the trace of $u \in C^\infty(\bar{\Omega}; V)$ is decomposed into a contribution from $T_{j_0}u$ and a sum over $\text{Tr}_{\partial\Omega}(T_{j+1}u - T_ju) = \text{Tr}_{\partial\Omega}(\Pi_j[u])$. By construction, $\Pi_j[u]$ is determined by the projections of u onto $\ker(\mathbb{A})$ on reflected balls contained in the strips $\Omega_{j-i_0} \setminus \Omega_{j+i_0}$; the latter have uniformly finite mutual overlap and yield the entire Ω when united.

give an estimate of the $W^{k,1}$ -norm of $T_{j_0}^{(2)}u$. Denoting \mathcal{L}_{j_0} the set of indices $i \in \mathbb{N}$ such that $\Omega_{j_0} \cap B_{j_0,i} \neq \emptyset$,

$$\begin{aligned}
\|T_{j_0}^{(2)}u\|_{W^{k,1}(\Omega)} &\leq c \sum_{\ell=0}^k \sum_{|\alpha|+|\beta|=\ell} \sum_{i \in \mathcal{L}_{j_0}} \|(\partial^\alpha(\rho_{j_0}\rho_{j_0,i}))(\partial^\beta \Pi_{j_0,i}u)\|_{L^1(\Omega)} \\
&\stackrel{(5.3), (5.4)}{\leq} c \sum_{\ell=0}^k \sum_{|\alpha|+|\beta|=\ell} \sum_{i \in \mathcal{L}_{j_0}} 2^{j_0|\alpha|} \times \\
(5.10) \quad &\quad \times 2^{-j_0(n-|\beta|)} 2^{-j_0|\beta|} \int_{B_{j_0,i}} |\partial^\beta \Pi_{j_0,i}u| \, dx \\
&\stackrel{(3.16)_2, \text{Lem. 5.1(b)}}{\leq} c \sum_{\ell=0}^k 2^{j_0\ell} 2^{-j_0n} \sum_{i \in \mathcal{L}_{j_0}} \int_{B_{j_0,i}^\#} |u| \, dx \\
&\stackrel{(5.1)_1}{\leq} c 2^{j_0k} \|u\|_{L^1(\Omega_{j_0-i_0} \setminus \Omega_{j_0+i_0})},
\end{aligned}$$

where we additionally used the uniform finite overlap property of the $B_{j_0,i}$'s. Since, by assumption, $W^{k,1}(\Omega; V)$ has trace space $\mathcal{X}(\partial\Omega; V)$, we consequently obtain

$$(5.11) \quad \|\text{Tr}_{\partial\Omega}(T_{j_0}u)\|_{\mathcal{X}(\partial\Omega)} = \|\text{Tr}_{\partial\Omega}(T_{j_0}^{(2)}u)\|_{\mathcal{X}(\partial\Omega)} \leq c 2^{j_0k} \|u\|_{L^1(\Omega)}.$$

Combining (5.9), (5.11) and (5.7), we then obtain

$$(5.12) \quad \|\text{Tr}_{\partial\Omega}(u)\|_{\mathcal{X}(\partial\Omega)} \leq c 2^{j_0k} \|u\|_{L^1(\Omega)} + c \|\mathbb{A}u\|_{L^1(\Omega)} \quad \text{for all } u \in C^\infty(\bar{\Omega}; V).$$

If $u \in W^{\mathbb{A},1}(\Omega)$, we invoke Lemma 5.2 (d) to find $(u_j) \subset C^\infty(\bar{\Omega}; V)$ such that $u_j \rightarrow u$ in $W^{\mathbb{A},1}(\Omega)$. As in the proof of Lemma 4.6 and since $(\mathcal{X}(\partial\Omega), \|\cdot\|_{\mathcal{X}(\partial\Omega)})$

is assumed Banach, we may define as a limit in $\mathcal{X}(\partial\Omega; V)$

$$(5.13) \quad \widetilde{\text{Tr}}_{\partial\Omega}(u) := \lim_{j \rightarrow \infty} \text{Tr}_{\partial\Omega}(u_j),$$

being independent of the approximating sequence $(u_j) \subset C^\infty(\overline{\Omega}; V)$. It is then straightforward to verify that $\widetilde{\text{Tr}}_{\partial\Omega}: W^{\mathbb{A},1}(\Omega) \rightarrow \mathcal{X}(\partial\Omega; V)$ is bounded and linear. Clearly $\widetilde{\text{Tr}}_{\partial\Omega}$ and $\text{Tr}_{\partial\Omega}$ coincide on $W^{k,1}(\Omega; V)$, and since $\text{Tr}_{\partial\Omega}: W^{k,1}(\Omega; V) \rightarrow \mathcal{X}(\partial\Omega; V)$ is surjective, $\widetilde{\text{Tr}}_{\partial\Omega}: W^{\mathbb{A},1}(\Omega) \rightarrow \mathcal{X}(\partial\Omega; V)$ is onto, too. The proof is complete. \square

We conclude this section with the following

Remark 5.5 (Homogeneous trace inequalities). To connect with the results of Section 4, let $\Omega = \mathbb{H}$ be an open halfspace and recall that $\dot{W}^{k,1}(\mathbb{H}; V)$ has trace space $\dot{B}_{1,1}^{k-1}(\partial\mathbb{H}; V)$. Given $u \in C_c^\infty(\overline{\mathbb{H}}; V)$, we define $T_j u$ as in (5.5). Note that in this particular geometric situation, the number as it appears in (C4) can be chosen to be arbitrarily negative. Proposition 5.2 directly inherits to the homogeneous situation and hence one obtains (5.7) with $j_0 = -\infty$. As a substitute of (5.10) we then obtain

$$(5.14) \quad \|T_{j_0}^{(2)} u\|_{\dot{W}^{k,1}(\Omega)} \leq c 2^{j_0 k} \|u\|_{L^1(\Omega_{j_0 - i_0} \setminus \Omega_{j_0 + i_0})}.$$

Since $u \in C_c^\infty(\overline{\mathbb{H}}; V)$, the term on the right-hand side of (5.14) will vanish as $j_0 \rightarrow -\infty$. Based on the USPENSKIĬ estimate (1.6), an analogous argument as that for (5.12) then equally yields $\|u\|_{\dot{B}_{1,1}^{k-1}(\partial\mathbb{H})} \leq c \|\mathbb{A}u\|_{L^1(\mathbb{H})}$ for all $u \in C_c^\infty(\overline{\mathbb{H}}; V)$.

6. PROOF OF THEOREM 1.3: KORN WITHOUT GLOBAL SINGULAR INTEGRAL ESTIMATES

We finally turn to the Korn-type inequality of Theorem 1.3 for John domains, which form the canonical class of domains for such inequalities. Since such domains in general need not be extension domains for $W^{\mathbb{A},p}$, the global Calderón-Zygmund estimates from the proof of Theorem 4.4 cannot be utilised and a local argument is required. In Section 6.1, which should be of independent interest, we provide a generalisation of a decomposition for maps on John domains by RŮŽIČKA, SCHUMACHER and the first author [23], the latter being only applicable to maps with zero mean. Building on a best approximation result and an inequality of FEFERMAN-STEIN-type (Sections 6.2 and 6.3), we obtain even more general Korn-type inequalities by means of extrapolation (so e.g. involving Orlicz- or Lorentz norms) once Korn-type inequalities are provided for weighted Lebesgue spaces, see Sections 6.4 and 6.5. Hence, even though Theorem 1.3 is only stated for the usual L^p -spaces, we directly deal with the weighted setting throughout.

6.1. Weights and a decomposition theorem for John domains. As usual, a weight $w : \mathbb{R}^n \rightarrow \mathbb{R}$ with $w > 0$ almost everywhere is said to be a A_q -Muckenhoupt weight with $1 \leq q < \infty$, in short $w \in A_q$, if

$$(6.1) \quad \begin{aligned} [w]_{A_q} &:= \sup_Q \int_Q w \, dx \left(\int_Q w^{-\frac{1}{q-1}} \, dx \right)^{q-1} < \infty, & \text{if } 1 < q < \infty, \\ [w]_{A_1} &:= \sup_Q \left[\int_Q w \, dx \sup_{x \in Q} \frac{1}{w(x)} \right] < \infty, & \text{if } q = 1. \end{aligned}$$

The class A_∞ is defined by $A_\infty = \bigcup_{q>1} A_q$ and endowed with $[w]_{A_\infty} := \lim_{p \rightarrow \infty} [w]_{A_p}$. Now let $\Omega \subset \mathbb{R}^n$ be open and bounded. For $w \in A_\infty$, $L_w^q(\Omega; V)$ is defined in

the obvious manner. For a finite dimensional real vector space E , a subspace $\mathcal{N} \subset \mathcal{P}_m(\mathbb{R}^n; E)$ and a space $X(\Omega; E) \subset L^1(\Omega; E)$, we define

$$(6.2) \quad X_{\mathcal{N}}(\Omega; E) := \left\{ f \in X(\Omega; E) : \int_{\Omega} f \cdot \pi \, dx = 0 \text{ for all } \pi \in \mathcal{N} \right\}.$$

If $w \in A_q$ with $1 \leq q < \infty$ and $f \in L_w^q(\Omega; E)$, then Hölder's inequality and (6.1) imply that $f \in L^1(\Omega; E)$. In particular, for such weights w , $L_{w, \mathcal{N}}^q(\Omega; E)$ is well-defined. We can now state the decomposition theorem that shall later be applied to $\mathcal{N} = \nabla^k \ker(\mathbb{A})$; as usual, \mathcal{M} denotes the Hardy-Littlewood maximal operator.

Theorem 6.1 (Decomposition theorem). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded John domain or an open and bounded domain satisfying the emanating chain condition with constants σ_1, σ_2 and chain-covering $\mathcal{W} = \{W_i : i \in \mathbb{N}_0\}$. Then there exists a family of linear operators $T_i : C_{c, \mathcal{N}}^\infty(\Omega; E) \rightarrow C_{c, \mathcal{N}}^\infty(W_i; E)$, $i \in \mathbb{N}_0$, such that for all $1 < q < \infty$ and all $w \in A_q$ the following holds:*

(a) *For each $i \in \mathbb{N}_0$ and all $f \in L_{w, \mathcal{N}}^q(\Omega; E)$ there holds*

$$(6.3) \quad |T_i f| \leq c(\sigma_2, \mathcal{N}) \mathbb{1}_{W_i} \mathcal{M}(\mathbb{1}_{\Omega} f) \quad \mathcal{L}^n\text{-almost everywhere.}$$

(b) *For each $i \in \mathbb{N}_0$ the operator T_i maps $L_{w, \mathcal{N}}^q(\Omega; E)$ boundedly to $L_{w, \mathcal{N}}^q(W_i; E)$.*

(c) *The family $\{T_i f : i \in \mathbb{N}_0\}$ is a decomposition of f in $L_{w, \mathcal{N}}^q(\Omega; E)$, i.e.,*

$$(6.4) \quad f = \sum_{i \in \mathbb{N}_0} T_i f \quad \text{in } L_{w, \mathcal{N}}^q(\Omega; E)$$

for all $f \in L_{w, \mathcal{N}}^q(\Omega; E)$. The convergence is unconditionally.

(d) *The mapping $f \mapsto (\|T_i f\|_{L_w^q(W_i)})_i$ from $L_{w, \mathcal{N}}^q(\Omega; E)$ to $l^q(\mathbb{N}_0)$ is bounded and*

$$(6.5) \quad \frac{1}{c} \|f\|_{L_w^q(\Omega)} \leq \left(\sum_{i \in \mathbb{N}_0} \|T_i f\|_{L_w^q(W_i)}^q \right)^{\frac{1}{q}} \leq c \|f\|_{L_w^q(\Omega)}$$

with a constant $c = c(\sigma_1, \sigma_2, q, [w]_{A_q}, \mathcal{N}) > 0$.

(e) *If $f \in C_{c, \mathcal{N}}^\infty(\Omega; E)$, then $\{i \in \mathbb{N}_0 : T_i f \neq 0\}$ is finite.*

Proof. The proof is an adaptation of the proof of [23, Thm. 4.2] by RŮŽIČKA, SCHUMACHER and the first author, and so we confine ourselves to the points that need refinement in the setting considered here.

The idea in [23], where \mathcal{N} is given by the constants, is that each f is first decomposed into local functions $S_i f := \xi_i f$ by means of a partition of unity ξ_i subject to the covering $\mathcal{W} = \{W_i : i \in \mathbb{N}_0\}$. Then each $S_i f$ is corrected in [23] such that it has vanishing integral. To compensate this change, the corrections are transported along the chains to the central ball W_0 . In the much more general setting as considered here we have to correct the $S_i f$'s by local projections to \mathcal{N} and transport those corrections along the chains to the central ball W_0 . Throughout, we tacitly suppose f to be extended to \mathbb{R}^n by zero.

Adopting the notation of Definition 2.3, we may suppose that the balls $B_{i,l}$ as in (C2) belong to a family \mathcal{B} such that $\sum_{B \in \mathcal{B}} \mathbb{1}_B \leq \sigma_2 \mathbb{1}_{\Omega}$; see Remark 2.4. For $B \in \mathcal{B}$ we find, by translating and scaling a fixed non-negative $\eta \in C_c^\infty(B(0,1))$ with $\int_{B(0,1)} \eta \, dx = 1$, a non-negative function $\eta_B \in C_c^\infty(B)$ with $\int_B \eta_B \, dx = 1$ and $|\eta_B| \leq c/\mathcal{L}^n(B)$, where $c = c(n) > 0$. Setting $\eta_{i,l} := \eta_{B_{i,l}}$, we may record that

$$(6.6) \quad \begin{aligned} & \eta_{i,l} \in C_c^\infty(W_{i,l}) \cap C_c^\infty(W_{i,l+1}), \\ & \|\eta_{i,l}\|_{L^\infty} \leq c \min \left\{ \frac{1}{\mathcal{L}^n(W_{i,l})}, \frac{1}{\mathcal{L}^n(W_{i,l+1})} \right\} \end{aligned}$$




FIGURE 5. Construction of the decomposition blocks T_i , cf. (6.9). In (6.9), one sums over all chains $W_{j,0} = W_j, \dots, W_{j,m_j} = W_0$ that contain W_i ; here, two such chains and the location of the relevant balls that matter in the definition of T_i are depicted.

for all $i, l \geq 0$ with $0 \leq l \leq m_i - 1$. Equally, we pick a non-negative $\eta_0 \in C_c^\infty(W_0)$ with $\int_{W_0} \eta_0 \, dx = 1$ and $|\eta_0| \leq c/\mathcal{L}^n(W_0)$; since every chain ends in $W_{i,m_i} = W_0$, we put $\eta_{i,m_i} := \eta_0$ for all $i \geq 0$.

Now define an operator $\Pi_{j,l} : L^1(\sigma_2 W_{j,l}; E) \rightarrow \mathcal{N}$ by

$$(6.7) \quad \int_{W_{j,l}} \eta_{j,l}(\Pi_{j,l}f) \, \pi \, dx = \int_{\sigma_2 W_{j,l}} f \, \pi \, dx \quad \text{for all } \pi \in \mathcal{N}.$$

Inserting $\pi = \Pi_{j,l}f$, inverse estimates for polynomials and using that all $\eta_{j,l}$'s are translated and scaled versions of some η , we obtain

$$\begin{aligned} \|\Pi_{j,l}f\|_{L^\infty(\sigma_2 W_{j,l})} &\leq c \left(\int_{\sigma_2 W_{j,l}} \eta_{j,l} |\Pi_{j,l}f|^2 \, dx \right)^{\frac{1}{2}} \\ &\leq c \left(\int_{\sigma_2 W_{j,l}} f \cdot \Pi_{j,l}f \, dx \right)^{\frac{1}{2}} \leq c \|\Pi_{j,l}f\|_{L^\infty(\sigma_2 W_{j,l})}^{\frac{1}{2}} \left(\int_{\sigma_2 W_{j,l}} |f| \, dx \right)^{\frac{1}{2}}, \end{aligned}$$

which implies

$$(6.8) \quad \|\Pi_{j,l}f\|_{L^\infty(\sigma_2 W_{j,l})} \leq c \int_{\sigma_2 W_{j,l}} |f| \, dx.$$

We now modify the decomposition operators of [23] and, adopting the notation of Definition 2.3, set for $f \in L_{w,\mathcal{N}}^q(\Omega; E)$

$$(6.9) \quad \begin{aligned} T_i f &:= (S_i f - \eta_{i,0} \Pi_{i,0}(S_i f)) \\ &+ \sum_{\substack{j \geq 0 \\ j \neq i}} \sum_{\substack{l: 0 < l \leq m_j \\ W_{j,l} = W_i}} (\eta_{j,l-1} \Pi_{j,l-1}(S_j f) - \eta_{j,l} \Pi_{j,l}(S_j f)). \end{aligned}$$

In the ultimate double sum, j effectively ranges over all numbers for which the chain $W_{j,m_j} = W_0, \dots, W_{j,0} = W_j$ from (C2) connects W_j with W_0 where, in addition, $W_{j,l} = W_i$ for some $0 < l \leq m_j$ asserts that W_i must be contained in this chain. If $j \neq i$ and $0 < l \leq m_j$ are such that $W_{j,l} = W_i$, then (6.6) implies that $\text{spt}(\eta_{j,l-1}) \cup \text{spt}(\eta_{j,l}) \subset W_{j,l} = W_i$. We hence obtain $\text{spt}(T_i f) \subset W_i$ and, by (6.6), $\eta_{j,l-1} + \eta_{j,l} \leq c \mathbb{1}_{W_i}/\mathcal{L}^n(W_i)$. Also note that the second sum over $0 < l \leq m_j$ such that $W_{j,l} = W_i$ contains only one summand.

We begin by proving that $T_i f$ is well-defined. Similarly to [23], we define the majorant

$$(6.10) \quad U_i := \sum_{\substack{j \geq 0 \\ j \neq i}} \sum_{\substack{l: 0 < l \leq m_j \\ W_{j,l} = W_i}} (\eta_{j,l-1} |\Pi_{j,l-1}(S_j f)| + \eta_{j,l} |\Pi_{j,l}(S_j f)|).$$

Next, if $W_{j,0} = W_j, \dots, W_{j,m_j} = W_0$ is a chain with $W_{j,l} = W_i$ for some $0 < l \leq m_j$, then we have by (6.6), $B_{j,l-1} \subset \sigma_2 W_{j,l-1} \subset \sigma_2^2 W_{j,l}$, $B_{j,l} \subset W_{j,l} \subset \sigma_2 W_{j,l}$ and the uniform comparability $\mathcal{L}^n(B_{j,l-1}) \simeq \mathcal{L}^n(B_{j,l}) \simeq \mathcal{L}^n(W_{j,l})$ the pointwise estimate

$$\begin{aligned} U_i &\leq c \frac{\mathbb{1}_{W_i}}{\mathcal{L}^n(W_i)} \sum_{\substack{j \geq 0 \\ j \neq i}} \sum_{\substack{l: 0 < l \leq m_j \\ W_{j,l} = W_i}} (\|\Pi_{j,l-1}(S_j f)\|_{L^\infty(B_{j,l-1})} + \|\Pi_{j,l}(S_j f)\|_{L^\infty(B_{j,l})}) \\ &\stackrel{(6.8)}{\leq} c \frac{\mathbb{1}_{W_i}}{\mathcal{L}^n(W_i)} \sum_{\substack{j \geq 0 \\ j \neq i}} \sum_{\substack{l: 0 < l \leq m_j \\ W_{j,l} = W_i}} \|S_j f\|_{L^1(\sigma_2^2 W_{j,l})} \\ &\leq c \frac{\mathbb{1}_{W_i}}{\mathcal{L}^n(W_i)} \sum_{\substack{j \geq 0 \\ j \neq i}} \|S_j f\|_{L^1(\sigma_2 W_i)} \\ &\leq c \mathbb{1}_{W_i} \int_{\sigma_2 W_i} |f| \, dx \end{aligned}$$

where we used in the ultimate step that the W_j 's have uniformly finite overlap, cf. (C1) and, by (C2), there holds $\text{spt}(S_j f) \subset W_j \subset \sigma_2 W_i$. Since $f \in L^1_{\text{loc}}(\Omega; E)$, the sum in $T_i f$ converges absolutely \mathcal{L}^n and so $T_i f$ is well-defined \mathcal{L}^n -a.e.. The term $S_i f - \eta_{i,0} \Pi_{i,0}(S_i f)$ can be estimated analogously, and so (a) follows.

Since \mathcal{M} is bounded on $L^q_w(\mathbb{R}^n)$, we obtain that T_i maps $L^q_w(\Omega; E) \rightarrow L^q_w(W_i; E)$ boundedly. In combination with (a) and recalling that $L^q_w(W_i; E) \hookrightarrow L^1(W_i; E)$, we obtain by Lebesgue's theorem on dominated convergence that the sum defining $T_i f$ also converges in $L^1(W_i; E)$. For the full statement of (b), let $f \in L^q_{w,\mathcal{N}}(\Omega; E)$ and consider some $j \neq i$ such that W_i is contained in the chain connecting W_j and W_0 , so $W_i = W_{j,l}$ for some $0 < l \leq m_j$. By (C2), we then have $W_j \subset \sigma_2 W_{j,l-1} \cap \sigma_2 W_{j,l}$. Therefore, $\text{spt}(S_j f) \subset W_j$ yields for all $\pi \in \mathcal{N}$

$$(6.11) \quad \int_{\sigma_2 W_{j,l-1}} S_j f \cdot \pi \, dx = \int_{\sigma_2 W_{j,l}} S_j f \cdot \pi \, dx.$$

The support properties of the $S_j f$'s and the $\eta_{j,l}$'s, the definition of $\Pi_{j,l}$ and (6.11) imply that each summand in the definition of T_i has integral zero when being tested against any $\pi \in \mathcal{N}$. Since the sum defining $T_i f$ converges in $L^1(\Omega; E)$, we conclude (b). We now establish that $f = \sum_{i \in \mathbb{N}_0} T_i f$ holds pointwisely \mathcal{L}^n -a.e. for $f \in L^q_{w,\mathcal{N}}(\Omega; E)$; working from here, the proof evolves exactly as in [23]. By pointwise absolute convergence of $\sum_{i \in \mathbb{N}_0} T_i f$ \mathcal{L}^n -a.e., we may change the order of summation in the following to find

$$\begin{aligned} \sum_{i \in \mathbb{N}_0} T_i f &= \sum_{i \in \mathbb{N}_0} S_i f - \sum_{i \in \mathbb{N}_0} \eta_{i,0} \Pi_{i,0}(S_i f) \\ &\quad + \sum_{j \geq 0} \sum_{l: 0 < l \leq m_j} \sum_{\substack{i \in \mathbb{N}_0 \setminus \{j\} \\ W_i = W_{j,l}}} (\eta_{j,l-1} \Pi_{j,l-1}(S_j f) - \eta_{j,l} \Pi_{j,l}(S_j f)) \\ &\stackrel{(*)}{=} \sum_{i \in \mathbb{N}_0} S_i f - \sum_{i \in \mathbb{N}_0} \eta_{i,0} \Pi_{i,0}(S_i f) + \sum_{j \geq 0} (\eta_{j,0} \Pi_{j,0}(S_j f) - \eta_{j,m_j} \Pi_{j,m_j}(S_j f)) \end{aligned}$$

$$\begin{aligned}
&= \sum_{i \in \mathbb{N}_0} S_i f - \sum_{j \geq 0} \eta_{j, m_j} \Pi_{j, m_j}(S_j f) \\
&\stackrel{\eta_{j, m_j} = \eta_0}{=} \sum_{i \in \mathbb{N}_0} S_i f - \eta_0 \sum_{j \geq 0} \Pi_{j, m_j}(S_j f) \\
&= f - \eta_0 \sum_{j \geq 0} \Pi_{W_0}(S_j f) \\
&= f - \eta_0 \Pi_{W_0} f = f,
\end{aligned}$$

since $\Pi_{W_0} := \Pi_{j, m_j}$ is independent of j and we have $\Pi_{W_0} f = 0$ as $f \in L^q_{w, \mathcal{N}}(\Omega; E)$. For (*) note that, for fixed $j \geq 0$ and $0 < l \leq m_j$, $\sum_{i \neq j, W_i = W_{j, l}} 1 = 1$, cf. [23, Eq. (4.18)ff.]. Working from here, the rest of the proof is exactly as in [23]. We thus omit the details. \square

Remark 6.2 (Unbounded domains). Theorem 6.1 also holds for unbounded John domains and unbounded domains satisfying the emanating chain condition. The proof requires no change. We refer to the modified definitions for unbounded domains to [23].

6.2. A best approximation property. For our objectives in Section 6.3 and 6.4, we require the following best approximation property of the projections underlying Theorem 3.7:

Proposition 6.3 (Best approximation property). *Let \mathbb{A} be a k -th order \mathbb{C} -elliptic operator of the form (1.1). Let $\Omega \subset \mathbb{R}^n$ be an open and bounded domain satisfying the emanating chain condition with constants σ_1, σ_2 and central ball B . Moreover, let $1 < p < \infty$ and $w \in A_p$. Then for all $\ell \in \{0, \dots, k\}$ and all $u \in W^{\ell, p}(\Omega; V)$ we have*

$$\left(\int_{\Omega} |D^{\ell}(u - \Pi_{\mathbb{A}}^B u)|^p w \, dx \right)^{\frac{1}{p}} \leq C \inf_{q \in \ker(\mathbb{A})} \left(\int_{\Omega} |D^{\ell}(u - q)|^p w \, dx \right)^{\frac{1}{p}}.$$

where $C = C(\mathbb{A}, \sigma_2, p, [w]_{A_p}) > 0$.

Proof. We start by noting that, if $B \subset \Omega$ is an open ball with $\text{diam}(\Omega) \leq c_0 \text{diam}(B)$, then for any $m \in \mathbb{N}_0$ there exists a constant $c_1 = c_1(c_0, m, n, V) > 0$ such that for all $\pi \in \mathcal{P}_m(\mathbb{R}^n; V)$ and all $\ell \in \mathbb{N}_0$ with $\ell \leq m + 1$ there holds

$$\begin{aligned}
(6.12) \quad \left(\int_{\Omega} |D^{\ell} \pi|^p w \, dx \right)^{\frac{1}{p}} &\leq (\sup_{\Omega} |D^{\ell} \pi|) \left(\frac{w(\Omega)}{\mathcal{L}^n(\Omega)} \right)^{\frac{1}{p}} \\
&\leq c_1 r(B)^{-\ell} \int_B |\pi| \, dx \left(\frac{w(\Omega)}{\mathcal{L}^n(\Omega)} \right)^{\frac{1}{p}} \\
&\leq c_1 [w]_{A_p}^{\frac{1}{p}} r(B)^{-\ell} \left(\int_B |\pi|^p w \, dx \right)^{\frac{1}{p}},
\end{aligned}$$

where the ultimate line follows by Hölder's inequality and because w is doubling.

Now let $\ell \in \{0, \dots, k\}$ and let $q \in \ker(\mathbb{A})$ be arbitrary, so that $\Pi_{\mathbb{A}}^B q = q$. Moreover, let $q_1 \in \mathcal{P}_{\ell-1} \subset \ker(\mathbb{A})$ be arbitrary. Then $\Pi_{\mathbb{A}}^B q_1 = q_1$ and therefore $D^{\ell} \Pi_{\mathbb{A}}^B q_1 = 0$. Now, $D^{\ell}(u - \Pi_{\mathbb{A}}^B u) = D^{\ell}(u - q) - D^{\ell} \Pi_{\mathbb{A}}^B(u - q - q_1)$, inverse estimates for polynomials in conjunction with $\text{diam}(\Omega) \leq c_0(\sigma_2) \text{diam}(B)$ (cf. (2.1)) and (6.12) imply

$$\begin{aligned}
&\left(\int_{\Omega} |D^{\ell}(u - \Pi_{\mathbb{A}}^B u)|^p w \, dx \right)^{\frac{1}{p}} \\
&\leq C \left(\int_{\Omega} |D^{\ell}(u - q)|^p w \, dx \right)^{\frac{1}{p}} + C \left(\int_{\Omega} |D^{\ell} \Pi_{\mathbb{A}}^B(u - q - q_1)|^p w \, dx \right)^{\frac{1}{p}}
\end{aligned}$$

$$(6.12) \quad \leq C \left(\int_{\Omega} |D^{\ell}(u - q)|^p w \, dx \right)^{\frac{1}{p}} + C \operatorname{diam}(\Omega)^{-\ell} \left(\int_B |u - q - q_1|^p w \, dx \right)^{\frac{1}{p}}.$$

Now we may fix $q_1 \in \mathcal{P}_{\ell-1}(\mathbb{R}^n; V)$ such that the standard A_p -weighted Poincaré inequality for balls (see [32, Chpt. 1.4]) yields

$$\operatorname{diam}(\Omega)^{-\ell} \left(\int_B |u - q - q_1|^p w \, dx \right)^{\frac{1}{p}} \leq C \left(\int_B |D^{\ell}(u - q)|^p w \, dx \right)^{\frac{1}{p}}.$$

This, the previous estimate and $\operatorname{diam}(\Omega) \leq c_0(\sigma_2) \operatorname{diam}(B)$ prove the claim. \square

6.3. An inequality of Fefferman-Stein-type. In this section we derive an inequality which allows to control the information of a function by its \mathcal{N} -sharp maximal operator. This serves as an instrumental tool in the proof of Theorem 1.3 in Section 6.4, but should be of independent interest.

We require the following restricted variant of maximal operators: Given an open subset $\Omega \subset \mathbb{R}^n$, $\sigma \geq 1$ and, for some finite dimensional real vector space E , a subset $\mathcal{N} \subset \mathcal{P}_m(\mathbb{R}^n; E)$ for some $m \in \mathbb{N}$ and $1 \leq p < \infty$, we put

$$(6.13) \quad \mathcal{M}_{\operatorname{res}, \Omega, \sigma, p} f(x) := \begin{cases} \sup_{Q \ni x: \sigma Q \subset \Omega} \left(\int_Q |f|^p \, dy \right)^{\frac{1}{p}}, & \text{if } x \in \Omega \\ 0 & \text{otherwise} \end{cases}$$

$$(6.14) \quad \mathcal{M}_{\operatorname{res}, \Omega, \sigma, p, \mathcal{N}}^{\sharp} f(x) := \begin{cases} \sup_{Q \ni x: \sigma Q \subset \Omega} \inf_{\pi \in \mathcal{N}} \left(\int_Q |f - \pi|^p \, dy \right)^{\frac{1}{p}}, & \text{if } x \in \Omega \\ 0 & \text{otherwise} \end{cases}$$

for $f \in L_{\operatorname{loc}}^p(\Omega; E)$ and non-degenerate cubes Q . For brevity, we also set $\mathcal{M}_{\operatorname{res}, \Omega, \sigma} := \mathcal{M}_{\operatorname{res}, \Omega, \sigma, 1}$ and $\mathcal{M}_{\operatorname{res}, \Omega, \sigma, \mathcal{N}}^{\sharp} := \mathcal{M}_{\operatorname{res}, \Omega, \sigma, 1, \mathcal{N}}^{\sharp}$.

The main result of this section is a generalisation of [23, Theorem 5.23] for $\mathcal{N} = \mathbb{R}$ and its unweighted version [36, Lemma 4]. We present the proof later in this section.

Theorem 6.4 (of Fefferman-Stein-type). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded domain satisfying the emanating chain condition with constants $\sigma_1 > 1, \sigma_2 \geq 1$ and central ball B . Let $1 < q < \infty$, $w \in A_q$ and $f \in L_{\operatorname{loc}}^1(\Omega; E)$. If $\mathcal{M}_{\operatorname{res}, \Omega, \sigma_1, \mathcal{N}}^{\sharp} f \in L_w^q(\Omega)$, then $f \in L_w^q(\Omega; E)$ and*

$$(6.15) \quad \inf_{\pi \in \mathcal{N}} \|f - \pi\|_{L_w^q(\Omega)} \leq c \|\mathcal{M}_{\operatorname{res}, \Omega, \sigma_1, \mathcal{N}}^{\sharp} f\|_{L_w^q(\Omega)}.$$

The constant c depends only on $\sigma_1, \sigma_2, \mathcal{N}, q$, and $[w]_{A_q}$.

For a cube Q (or ball) we define $\Pi_{\mathcal{N}}^Q$ as the $L^2(Q; E)$ -projection to \mathcal{N} . It follows easily by inverse estimates on \mathcal{N} that $\Pi_{\mathcal{N}}^Q$ is $L^1(Q; E)$ -stable, i.e.

$$(6.16) \quad \int_Q |\Pi_{\mathcal{N}}^Q f| \, dx \leq C_1 \int_Q |f| \, dx,$$

and, similarly to Proposition 6.3, has the approximation property

$$(6.17) \quad \int_Q |f - \Pi_{\mathcal{N}}^Q f| \, dx \leq C_2 \inf_{q \in \mathcal{N}} \int_Q |f - q| \, dx.$$

We need the following version of the Calderón-Zygmund decomposition by IWANIEC:

Lemma 6.5 ([36, Lemma 3]). *Let $Q_0 \subset \mathbb{R}^n$ be an open cube and let $f \in L^1(Q_0; E)$. For every $\alpha \geq (|f|)_{Q_0}$ there exist pairwise disjoint cubes $Q_j^{\alpha} \subset Q_0$ indexed by $j \in \mathbb{N}$ such that*

$$(a) \quad \alpha < \int_{Q_j^{\alpha}} |f| \, dx \leq 2^n \alpha,$$

- (b) If $\alpha \geq \beta \geq (|f|)_{Q_0}$, then each cube Q_j^α is contained in an element of $\{Q_j^\beta : j \in \mathbb{N}\}$,
- (c) $|f| \leq \alpha$ on $Q_0 \setminus \bigcup_j Q_j^\alpha$,
- (d) $\bigcup_j Q_j^\alpha \subset \{\mathcal{M}_{\text{res}, Q_0, 1} f > \alpha\}$ for all $\alpha \geq (|f|)_{Q_0}$,
- (e) $\{\mathcal{M}_{\text{res}, Q_0, 1} f > 5^n \alpha\} \subset \bigcup_j 5Q_j^\alpha$ for all $\alpha \geq (|f|)_{Q_0}$.

We start with a version of Theorem 6.4 for the case where Ω is just a cube.

Lemma 6.6. *Let $Q_0 \subset \mathbb{R}^n$ be an open cube, $1 \leq q < \infty$, $w \in A_\infty$, and let $f \in L^1(Q_0; E)$. If $\mathcal{M}_{\text{res}, Q_0, 1}^\# f \in L_w^q(Q_0)$, then $\mathcal{M}_{\text{res}, Q_0, 1} f \in L_w^q(Q_0)$ and there holds*

$$\int_{Q_0} |\mathcal{M}_{\text{res}, Q_0, 1} f|^q w \, dx \leq c \int_{Q_0} |\mathcal{M}_{\text{res}, Q_0, 1}^\# f|^q w \, dx + c w(Q_0) \left(\int_{Q_0} |f| \, dx \right)^q,$$

where $c = c(n, \mathcal{N}, q, [w]_{A_\infty}) > 0$.

Proof. Since $w \in A_\infty$, there exists a constant $c_0 > 0$, only depending on $[w]_{A_\infty}$, such that $w(5Q) \leq c_0 w(Q)$ for any cube $Q \subset \mathbb{R}^n$, see [59, V 1.6, p. 196]. By the usual inverse estimates and (6.16), there exists $\mathbf{C} = \mathbf{C}(\mathcal{N}, n) > 0$ such that, for any open cube $Q \subset \mathbb{R}^n$,

$$(6.18) \quad \|\Pi_{\mathcal{N}}^Q f\|_{L^\infty(Q)} \leq \mathbf{C} \int_Q |f| \, dx.$$

Then we fix a number $m \in \mathbb{N}$ such that $\mathbf{C}2^{n-m} < \frac{1}{2}$. We claim that for every $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon, n, m, \mathcal{N}, [w]_{A_\infty}) > 0$ such that

$$(6.19) \quad \begin{aligned} w(\{\mathcal{M}_{\text{res}, Q_0, 1} f > 5^n \alpha\}) &\leq c_0 w(\{\mathcal{M}_{\text{res}, Q_0, 1}^\# f > \delta \alpha\}) \\ &\quad + c_0 \varepsilon w(\{\mathcal{M}_{\text{res}, Q_0, 1} f > 2^{-m} \alpha\}) \end{aligned}$$

for all $f \in L^1(Q_0; E)$ and all $\alpha \geq 2^m (|f|)_{Q_0}$. Let $\varepsilon > 0$. As $\alpha \geq 2^{-m} \alpha \geq (|f|)_{Q_0}$, we may pick $(Q_j^\alpha)_{j \in \mathbb{N}}$ as in Lemma 6.5. Toward (6.19), it suffices to show by virtue of Lemma 6.5 (e) and $w(5Q) \leq c_0 w(Q)$ for all cubes Q that

$$(6.20) \quad \sum_j w(Q_j^\alpha) \leq w(\{\mathcal{M}_{\text{res}, Q_0, 1}^\# f > \delta \alpha\}) + \varepsilon w(\{\mathcal{M}_{\text{res}, Q_0, 1} f > 2^{-m} \alpha\}).$$

By Lemma 6.5 (b), every Q_j^α is contained in some Q , $Q \in \{Q_i^{2^{-m} \alpha} : i \in \mathbb{N}\}$. Then

$$\int_{Q_j^\alpha} |\Pi_{\mathcal{N}}^Q f| \, dx \leq \|\Pi_{\mathcal{N}}^Q f\|_{L^\infty(Q)} \stackrel{(6.18)}{\leq} \mathbf{C} \int_Q |f| \, dx \stackrel{\text{Lemma 6.5 (a)}}{\leq} \mathbf{C} 2^{n-m} \alpha \stackrel{(6.18)\text{ff.}}{<} \frac{\alpha}{2}.$$

Thus, by (a) of Lemma 6.5 we obtain for all $j \in \mathbb{N}$

$$\int_{Q_j^\alpha} |f - \Pi_{\mathcal{N}}^Q f| \, dx \geq \int_{Q_j^\alpha} |f| \, dx - \int_{Q_j^\alpha} |\Pi_{\mathcal{N}}^Q f| \, dx \geq \alpha - \frac{\alpha}{2} = \frac{\alpha}{2}.$$

Multiplying the previous inequality by $|Q_j^\alpha|$ and summing over all $Q_j^\alpha \subset Q$, we obtain

$$\sum_{Q_j^\alpha \subset Q} |Q_j^\alpha| \leq \frac{2}{\alpha} \int_Q |f - \Pi_{\mathcal{N}}^Q f| \, dx.$$

Since $w \in A_\infty$, there exists an $\varepsilon_2 > 0$ such that if $\mathcal{E} \subset Q$ with $|\mathcal{E}| \leq \varepsilon_2 |Q|$, then $w(\mathcal{E}) \leq \varepsilon w(Q)$, see [59, V 1.7, p. 196]; here, ε_2 only depends on ε and the A_∞ -constant of w but is independent of Q and \mathcal{E} . With C_2 as in (6.17), we define $\delta := \frac{1}{2C_2} \varepsilon_2$ and then consider two cases:

• If $\int_Q |f - \Pi_{\mathcal{N}}^Q f| dx \leq \frac{\alpha}{2} \varepsilon_2$, then $\sum_{Q_j^\alpha \subset Q} |Q_j^\alpha| \leq \varepsilon_2 |Q|$. By our choice of ε_2 , $\sum_{Q_j^\alpha \subset Q} w(Q_j^\alpha) \leq \varepsilon w(Q)$, where we have used that the Q_j^α , $j \in \mathbb{N}$, are pairwise disjoint.

• If $\int_Q |f - \Pi_{\mathcal{N}}^Q f| dx > \frac{\alpha}{2} \varepsilon_2$, then we have, with C_2 as in (6.17), $\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f(x) > \frac{\alpha}{2C_2} \varepsilon_2 = \delta\alpha$ for all points $x \in Q$. In particular, $Q \subset \{\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f > \delta\alpha\}$ and

$$\sum_{Q_j^\alpha \subset Q} w(Q_j^\alpha) \leq w(Q) = w(Q \cap \{\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f > \delta\alpha\}).$$

Combining both alternatives, we obtain

$$(6.21) \quad \sum_{Q_j^\alpha \subset Q} w(Q_j^\alpha) \leq w(Q \cap \{\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f > \delta\alpha\}) + \varepsilon w(Q).$$

By Lemma 6.5 (b) and since $\alpha \geq 2^m (|f|)_{Q_0}$, summing (6.21) over the pairwise disjoint cubes $Q \in \{Q_i^{2^{-m}\alpha} : i \in \mathbb{N}\}$, all cubes from $\{Q_j^\alpha : j \in \mathbb{N}\}$ are counted exactly once in the overall sum. We thus obtain (6.20) and hereby (6.19). Once (6.19) is established, we may exactly follow the layer cake argument of IWANIEC [36, Lemma 4] to conclude. \square

Corollary 6.7. *Let $Q_0 \subset \mathbb{R}^n$ be an open cube, $1 \leq q < \infty$, $w \in A_q$, and $f \in L_{\mathcal{N}}^1(Q_0; E)$. If $\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f \in L_w^q(Q_0; E)$, then $f \in L_{w, \mathcal{N}}^q(Q_0; E)$ and*

$$\|f\|_{L_w^q(Q_0)} \leq c \|\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f\|_{L_w^q(Q_0)},$$

where the constant c only depends on \mathcal{N} , q and $[w]_{A_q}$.

Proof. Let $f \in L_{\mathcal{N}}^1(Q_0; E)$ so that $\Pi_{\mathcal{N}}^{Q_0} f = 0$. In conjunction with Lemma 6.6 and (6.17), this yields

$$\begin{aligned} \|f\|_{L_w^q(Q_0)}^q &\leq \|\mathcal{M}_{\text{res}, Q_0, 1} f\|_{L_w^q(Q_0)}^q \\ &\leq c \|\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f\|_{L_w^q(Q_0)}^q + c w(Q_0) \left(\int_{Q_0} |f - \Pi_{\mathcal{N}}^{Q_0} f| dx \right)^q \\ &\leq c \|\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f\|_{L_w^q(Q_0)}^q + c \int_{Q_0} |(\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f)(y)|^q w dy \\ &\leq c \|\mathcal{M}_{\text{res}, Q_0, 1, \mathcal{N}}^\# f\|_{L_w^q(Q_0)}^q, \end{aligned}$$

and the proof is complete. \square

Before we come to the proof of Theorem 6.4, the underlying duality argument requires

Lemma 6.8. *Let $\Omega \subset \mathbb{R}^n$ be as in Theorem 6.4. Let $1 < q < \infty$ and $w \in A_q$. Then $C_{c, \mathcal{N}}^\infty(\Omega; E)$ is dense in $L_{w, \mathcal{N}}^q(\Omega; E)$ for the norm topology.*

Proof. Let $\eta_0 \in C_c^\infty(B)$ with $\eta_0 \geq 0$ and $\int_B \eta_0 dx = 1$. Similarly as in (6.7) we define a linear operator $\Pi_0 : L^1(\Omega; E) \rightarrow \mathcal{N}$ by

$$\int_B \eta_0(\Pi_0 f) \xi dx = \int_\Omega f \xi dx \quad \text{for all } \xi \in \mathcal{N}.$$

By inverse estimates on \mathcal{N} and Hölder's inequality we obtain $\|\Pi_0 f\|_{L^\infty(\Omega)} \leq c \int_\Omega |f| dx$. This and $w \in A_q$ imply $\|\eta_0 \Pi_0 f\|_{L_w^q(\Omega)} \leq c \|f\|_{L_w^q(\Omega)}$. Now let $f \in L_{w, \mathcal{N}}^q(\Omega; E)$ so that, in particular, $\Pi_0 f = 0$. Then we find $(g_j) \subset C_c^\infty(\Omega; E)$ such that $g_j \rightarrow f$ in $L_w^q(\Omega; E)$, cf. MILLER [51, Lem. 2.1]. Put $\tilde{\Pi}_0 := |\Omega| \Pi_0$ and define $f_j := g_j - \eta_0 \tilde{\Pi}_0 g_j \in C_{c, \mathcal{N}}^\infty(\Omega; E)$. Since $\Pi_0 f = 0$ we have $f_j = g_j - \eta_0 \tilde{\Pi}_0 (g_j - f)$.

Since $g_j \rightarrow f$ in $L_w^q(\Omega; E)$ it follows with $\|\eta_0 \Pi_0(g_j - f)\|_{L_w^q(\Omega)} \leq c \|g_j - f\|_{L_w^q(\Omega)}$ that $f - f_j \rightarrow 0$ in $L_w^q(\Omega; E)$, as desired. \square

Proof of Theorem 6.4. Let $1 < q < \infty$, $w \in A_q$, and $f \in L_{\text{loc}}^1(\Omega; E)$ with $\mathcal{M}_{\text{res}, \Omega, \sigma_1, \mathcal{N}}^\# f \in L_w^q(\Omega)$. Since $w \in A_q$, we have $w' := w^{\frac{1}{1-q}} \in A_{q'}$.

Let $h \in C_{c, \mathcal{N}}^\infty(\Omega; E)$. By Theorem 6.1 we may then decompose h into the sum of functions $T_i h \in C_{c, \mathcal{N}}^\infty(W_i; E) \subset L_{w', \mathcal{N}}^{q'}(W_i; E)$ such that

$$(6.22) \quad \left(\sum_{i \geq 0} \|T_i h\|_{L_{w', \mathcal{N}}^{q'}(W_i)}^{q'} \right)^{\frac{1}{q'}} \leq c \|h\|_{L_{w'}^{q'}(\Omega)}.$$

Moreover, by Theorem 6.1 (e), only finitely many summands of $h = \sum_{i \geq 0} T_i h$ are non-zero. Since $f \in L_{\text{loc}}^1(\Omega; E)$ and $h \in C_{c, \mathcal{N}}^\infty(\Omega; E)$, $\langle f, h \rangle := \int_\Omega f h \, dx$ is well-defined, and for all $i \in \mathbb{N}$, $\xi \in \mathcal{N}$ we have $\int_{W_i} \xi T_i h \, dx = 0$. Therefore, with $\Pi_{\mathcal{N}}^{W_i}$ as in (6.16)ff.,

$$\langle f, h \rangle = \sum_i \langle f, T_i h \rangle = \sum_i \langle f - \Pi_{\mathcal{N}}^{W_i} f, T_i h \rangle,$$

where we also used that $f \in L^1(W_i; E)$, since $\sigma_1 > 1$ and consequently $\overline{W}_1 \Subset \sigma_1 W_i \subset \Omega$. We estimate with Corollary 6.7 in the second step

$$(6.23) \quad \begin{aligned} \langle f, h \rangle &\leq \sum_i \left\| f - \Pi_{\mathcal{N}}^{W_i} f \right\|_{L_w^q(W_i)} \|T_i h\|_{L_{w'}^{q'}(W_i)} \\ &\leq c \sum_i \left\| \mathcal{M}_{\text{res}, W_i, 1, \mathcal{N}}^\# f \right\|_{L_w^q(W_i)} \|T_i h\|_{L_{w'}^{q'}(W_i)} \\ &\leq c \left(\sum_i \left\| \mathbb{1}_{W_i} \mathcal{M}_{\text{res}, \Omega, \sigma_1, \mathcal{N}}^\# f \right\|_{L_w^q(\Omega)}^q \right)^{\frac{1}{q}} \left(\sum_i \|T_i h\|_{L_{w'}^{q'}(W_i)}^{q'} \right)^{\frac{1}{q'}} \\ &\leq c \left\| \mathcal{M}_{\text{res}, \Omega, \sigma_1, \mathcal{N}}^\# f \right\|_{L_w^q(\Omega)} \|h\|_{L_{w'}^{q'}(\Omega)} \end{aligned}$$

for all $h \in C_{c, \mathcal{N}}^\infty(\Omega; E)$, where we have used in the last step $\sum_{W \in \mathcal{W}} \mathbb{1}_{\sigma_1 W} \leq \sigma_2 \mathbb{1}_\Omega$ and (6.22). Therefore, invoking the smooth approximation from Lemma 6.8,

$$(6.24) \quad \sup_{\substack{h \in L_{w', \mathcal{N}}^{q'}(\Omega; E) \\ \|h\|_{L_{w'}^{q'}(\Omega)} \leq 1}} \langle f, h \rangle \leq c \left\| \mathcal{M}_{\text{res}, \Omega, \sigma_1, \mathcal{N}}^\# f \right\|_{L_w^q(\Omega)}.$$

By an analogous argument to that of Proposition 6.3 and because $\Pi_{\mathcal{N}}^B$ is L_w^q -stable,

$$\begin{aligned} \inf_{\pi \in \mathcal{N}} \|f - \pi\|_{L_w^q(\Omega)} &\leq c \|f - \Pi_{\mathcal{N}}^B f\|_{L_w^q(\Omega)} \\ &= c \sup_{\substack{h \in L_{w', \mathcal{N}}^{q'}(\Omega; E) \\ \|h\|_{L_{w'}^{q'}(\Omega)} \leq 1}} \langle f - \Pi_{\mathcal{N}}^B f, h - \Pi_{\mathcal{N}}^B h \rangle + \underbrace{\langle f - \Pi_{\mathcal{N}}^B f, \Pi_{\mathcal{N}}^B h \rangle}_{=0} \\ &\leq c \sup_{\substack{\tilde{h} \in L_{w', \mathcal{N}}^{q'}(\Omega; E) \\ \|\tilde{h}\|_{L_{w'}^{q'}(\Omega)} \leq 1}} \langle f, \tilde{h} \rangle \stackrel{(6.24)}{\leq} c \left\| \mathcal{M}_{\text{res}, \Omega, \sigma_1, \mathcal{N}}^\# f \right\|_{L_w^q(\Omega)}. \end{aligned}$$

This completes the proof. \square

6.4. Korn-type inequalities and the proof of Theorem 1.3. We now turn to Theorem 6.9 below, a special case of which is Theorem 1.3. Different from Theorem 4.4, where the focus was on the equivalence of trace and Korn-type inequalities on *smooth* domains, and previous contributions (cf. KALAMAJSKA [41, 42]), the previous sections now allow us to cover the natural yet vast class of John domains which, in turn, might be very irregular:

Theorem 6.9 (Weighted Korn inequality). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded domain that satisfies the emanating chain condition with constants $\sigma_1 > 1$ and $\sigma_2 \geq 1$ and central ball B . Moreover, let $k \in \mathbb{N}$, $1 < p < \infty$ and $w \in A_p$. Then for any k -th order \mathbb{C} -elliptic operator \mathbb{A} of the form (1.1) there exists a constant $c = c(\sigma_1, \sigma_2, p, [w]_{A_p}, \mathbb{A}) > 0$ such that*

$$(6.25) \quad \begin{aligned} \|D^k(u - \Pi_{\mathbb{A}}^B u)\|_{L_w^p(\Omega)} &\leq c \|\mathbb{A}u\|_{L_w^p(\Omega)}, \\ \|D^k u\|_{L_w^p(\Omega)} &\leq c(\text{diam}(\Omega))^{-k} \|u\|_{L_w^p(\Omega)} + \|\mathbb{A}u\|_{L_w^p(\Omega)} \end{aligned}$$

hold for all $u \in W^{\mathbb{A}} L_w^p(\Omega) := \{v \in L_w^p(\Omega; V) : \mathbb{A}v \in L_w^p(\Omega; W)\}$.

Proof. Since $w \in A_p$, by the open-endedness property of the Muckenhoupt classes [18], there exists $\varepsilon = \varepsilon(n, p, [w]_{A_p}) > 0$ such that $w \in A_{p-\varepsilon}$, too. We then pick $1 < q < p$ such that $p - \varepsilon < \frac{p}{q} < p$ and record in advance that $w \in A_{p/q}$. Let $\sigma_1 > 1$ and let $Q \subset \Omega$ be an open cube such that $\sigma_1 Q \subset \Omega$. Choose a cut-off function $\rho \in C_c^\infty(\Omega; [0, 1])$ such that $\mathbb{1}_Q \leq \rho \leq \mathbb{1}_{\sigma_1 Q}$ and $|\nabla^\ell \rho| \leq c(\sigma_1 - 1)^{-l} \ell(Q)^{-l}$ for all $l \in \{0, \dots, k\}$. Then $\varphi := \rho(u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u)$ (with $\Pi_{\mathbb{A}}^{\sigma_1 Q}$ being as in Theorem 3.7) is compactly supported in \mathbb{R}^n . Since \mathbb{A} in particular is elliptic, the Fourier multiplier argument underlying (4.28)ff. yields the existence of a constant $c = c(q, \mathbb{A}) > 0$ such that

$$(6.26) \quad \|D^k \varphi\|_{L^q(\mathbb{R}^n)} \leq c \|\mathbb{A}\varphi\|_{L^q(\mathbb{R}^n)}.$$

Moreover, note that $\Pi_{\mathbb{A}}^Q \Pi_{\mathbb{A}}^{\sigma_1 Q} u = \Pi_{\mathbb{A}}^{\sigma_1 Q} u$ on Q and therefore, using the L^q -stability of the projections $\Pi_{\mathbb{A}}^Q$ in the second and the Poincaré-type inequalities from Theorem 3.7 in the fourth step,

$$(6.27) \quad \begin{aligned} \left(\int_Q |\Pi_{\mathbb{A}}^{\sigma_1 Q} u - \Pi_{\mathbb{A}}^Q u|^q dx \right)^{\frac{1}{q}} &= \left(\int_Q |\Pi_{\mathbb{A}}^Q (u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u)|^q dx \right)^{\frac{1}{q}} \\ &\leq c \left(\int_Q |u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u|^q dx \right)^{\frac{1}{q}} \\ &\leq c \left(\int_{\sigma_1 Q} |u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u|^q dx \right)^{\frac{1}{q}} \stackrel{(3.19)}{\leq} c \ell(Q)^k \left(\int_{\sigma_1 Q} |\mathbb{A}u|^q dx \right)^{\frac{1}{q}}, \end{aligned}$$

where $c = c(\sigma_1, q, \mathbb{A}) > 0$. Now, using the Poincaré-type inequalities from Theorem 3.7,

$$\begin{aligned} \left(\int_Q |D^k(u - \Pi_{\mathbb{A}}^Q u)|^q dx \right)^{\frac{1}{q}} &\leq c \left(\int_{\sigma_1 Q} |D^k(\rho(u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u))|^q dx \right)^{\frac{1}{q}} \\ &\quad + c \left(\int_Q |D^k(\Pi_{\mathbb{A}}^{\sigma_1 Q} u - \Pi_{\mathbb{A}}^Q u)|^q dx \right)^{\frac{1}{q}} \\ &\stackrel{(6.26)}{\leq} c \left(\int_{\sigma_1 Q} |\mathbb{A}(\rho(u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u))|^q dx \right)^{\frac{1}{q}} + \left(\int_Q |D^k(\Pi_{\mathbb{A}}^{\sigma_1 Q} u - \Pi_{\mathbb{A}}^Q u)|^q dx \right)^{\frac{1}{q}} \\ &\leq c \left(\sum_{l=1}^k \ell(Q)^{-l} \left(\int_{\sigma_1 Q} |D^{k-l}(u - \Pi_{\mathbb{A}}^{\sigma_1 Q} u)|^q dx \right)^{\frac{1}{q}} \right) + c \left(\int_{\sigma_1 Q} |\mathbb{A}u|^q dx \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
& + c\ell(Q)^{-k} \sum_{l=0}^k \ell(Q)^l \left(\int_Q |D^l(\Pi_{\mathbb{A}}^{\sigma_1 Q} u - \Pi_{\mathbb{A}}^Q u)|^q dx \right)^{\frac{1}{q}} \\
& \stackrel{(3.19), (3.16)_2}{\leq} c \left(\int_{\sigma_1 Q} |\mathbb{A}u|^q dx \right)^{\frac{1}{q}} + c\ell(Q)^{-k} \left(\int_Q |\Pi_{\mathbb{A}}^{\sigma_1 Q} u - \Pi_{\mathbb{A}}^Q u|^q dx \right)^{\frac{1}{q}} \\
& \stackrel{(6.27)}{\leq} c \left(\int_{\sigma_1 Q} |\mathbb{A}u|^q dx \right)^{\frac{1}{q}},
\end{aligned}$$

where again $c = c(\sigma_1, q, \mathbb{A}) > 0$. We define $\mathcal{N} := D^k \ker(\mathbb{A})$. Then, using that $D^k \Pi_{\mathbb{A}}^Q u \in \mathcal{N}$ for any open cube Q ,

$$(6.28) \quad \mathcal{M}_{\text{res}, \Omega, \sigma_1, q, \mathcal{N}}^{\sharp}(D^k u) \leq c \mathcal{M}_{\text{res}, \Omega, 1, q}(\mathbb{A}u).$$

In the sequel, denote B the central ball of Ω , cf. Definition 2.3. We now invoke the best approximation property from Proposition 6.3 with respect to the weight w and $\ell = k$ in the first and the Fefferman-Stein-type inequality from Theorem 6.4 with $f = D^k u$, $\mathcal{N} = D^k \ker(\mathbb{A})$ in the second step to obtain

$$\begin{aligned}
(6.29) \quad \|D^k u - D^k \Pi_{\mathbb{A}}^B u\|_{L_w^p(\Omega)} & \stackrel{\text{Proposition 6.3}}{\leq} c \inf_{\pi \in \mathcal{N}} \|f - \pi\|_{L_w^p(\Omega)} \\
& \stackrel{\text{Theorem 6.4}}{\leq} c \|\mathcal{M}_{\text{res}, \Omega, \sigma_1, q, \mathcal{N}}^{\sharp}(D^k u)\|_{L_w^p(\Omega)} \\
& \stackrel{(6.28)}{\leq} c \|\mathcal{M}_{\text{res}, \Omega, 1, q}(\mathbb{A}u)\|_{L_w^p(\Omega)} \\
& \leq c \|\mathcal{M}_{\text{res}, \Omega, 1, 1}(|\mathbb{A}u|^q)\|_{L_w^{p/q}(\Omega)}^{1/q} \\
& \stackrel{1 < q < p}{\leq} c \|\mathbb{A}u\|_{L_w^p(\Omega)},
\end{aligned}$$

where we used that $\mathcal{M}_{\text{res}, \Omega, 1, 1}: L_w^{p/q}(\Omega) \rightarrow L_w^{p/q}(\Omega)$ is bounded provided $1 < p/q < \infty$ and $w \in A_{p/q}$. Tracking the dependencies of the constants in (6.29), we obtain (6.25)₁. To obtain (6.25)₂, the weighted inverse and stability estimates ((6.12) ff.) yield

$$\|D^k \Pi_{\mathbb{A}}^B u\|_{L_w^p(\Omega)} \leq c \text{diam}(\Omega)^{-k} \|\Pi_{\mathbb{A}}^B u\|_{L_w^p(\Omega)} \leq c \text{diam}(\Omega)^{-k} \|u\|_{L_w^p(\Omega)},$$

where $c = c(n, p, [w]_{A_p}, \sigma_1, \sigma_2, \mathbb{A}) > 0$. Combining this inequality with (6.29), (6.25) follows and the proof is complete. \square

Proof of Theorem 1.3. For a \mathbb{C} -elliptic k -th order operator, Theorems 6.9 and 3.7 immediately imply Theorem 1.3, direction (a) \Rightarrow (b), by taking $w \equiv 1$. For direction (b) \Rightarrow (a) of Theorem 1.3 it suffices to note that Theorem 4.4 remains valid when only requiring (d) to hold for all *connected*, open and bounded domains $\Omega \subset \mathbb{R}^n$; since every such domain is John, we thereby obtain (b) \Rightarrow (a) in Theorem 1.3. \square

Remark 6.10 (Korn for unbounded John domains). Theorem 6.9 also persists for unbounded John domains which are defined in analogy with Definition 2.2. Indeed, by a result due to VÄISÄLÄ [65, Thm. 4.6], if $\Omega \subset \mathbb{R}^n$ is α -John, then there exists a sequence of relatively compact $c\alpha$ -John domains $\Omega_j \subset \Omega$ with $\Omega_j \nearrow \Omega$. Since any open and bounded John domain satisfies the emanating chain condition, we obtain by (6.25)

$$\|D^k u\|_{L_w^p(\Omega_j)} \leq C(\text{diam}(\Omega_j))^{-k} \|u\|_{L_w^p(\Omega)} + \|\mathbb{A}u\|_{L_w^p(\Omega)} \quad \text{for all } u \in W^{\mathbb{A}} L_w^p(\Omega)$$

for any $1 < p < \infty$ and $w \in A_p$, with $C = C(c\alpha, p, [w]_{A_p}, \mathbb{A}) > 0$. Sending $j \rightarrow \infty$ in the preceding inequality then yields the Korn inequality

$$\|D^k u\|_{L_w^p(\Omega)} \leq C \|\mathbb{A}u\|_{L_w^p(\Omega)} \quad \text{for all } u \in W^{\mathbb{A}} L_w^p(\Omega).$$

6.5. Korn-type inequalities for other space scales. Theorem 6.9 easily allows for variants of Korn-type inequalities by means of extrapolation. For this, we recall a far-reaching variant of the RUBIO DE FRANCIA-extrapolation theorem due to CRUZ-URIBE, MARTELL & PÉREZ [20] that requires the following terminology: Given a Banach function space $(\mathbb{X}, \|\cdot\|_{\mathbb{X}})$ on \mathbb{R}^n with respect to \mathcal{L}^n , we call $(\mathbb{X}, \|\cdot\|_{\mathbb{X}})$ *rearrangement invariant* provided for each $f \in \mathbb{X}$ the norm $\|f\|_{\mathbb{X}}$ exclusively depends on its distribution function $d_f(\lambda) := \mathcal{L}^n(\{x \in \mathbb{R}^n : |f(x)| > \lambda\})$. Defining for $f \in \mathbb{X}$ its *decreasing rearrangement* $f^* : [0, \infty) \rightarrow \mathbb{R}$ by $f^*(t) := \inf\{\lambda \geq 0 : d_f(\lambda) \leq t\}$, there exists a unique rearrangement invariant Banach function space $(\overline{\mathbb{X}}, \|\cdot\|_{\overline{\mathbb{X}}})$ on $[0, \infty)$ with respect to \mathcal{L}^1 such that $f \in \mathbb{X}$ precisely if $f^* \in \overline{\mathbb{X}}$. In this situation, defining for $0 < t < \infty$ the dilation operator $D_t : \overline{\mathbb{X}} \ni g \mapsto g(\cdot/t) \in \overline{\mathbb{X}}$ and denoting $\|D_t\|_{\overline{\mathbb{X}}}$ its operator norm, the *lower* or *upper Boyd indices* of \mathbb{X} are given by

$$(6.30) \quad p_{\mathbb{X}} := \lim_{t \rightarrow \infty} \frac{\log(t)}{\log(\|D_t\|_{\overline{\mathbb{X}}})} \quad \text{and} \quad q_{\mathbb{X}} := \lim_{t \searrow 0} \frac{\log(t)}{\log(\|D_t\|_{\overline{\mathbb{X}}})},$$

respectively.

On the other hand, a differentiable function $\varphi : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ with right-continuous, non-decreasing derivative φ' such that $\varphi'(0) = 0$ and $\varphi'(t) > 0$ for $t > 0$ is called an *N-function*. If there exists $c_1 > 0$ such that $\varphi(2t) \leq c_1\varphi(t)$ holds for all $t > 0$, we say that φ satisfies the Δ_2 -condition; in this situation, the infimum over all such possible constants c_1 is denoted $\Delta_2(\varphi)$. If the convex conjugate of φ , $\tilde{\varphi}(t) := \sup_{s>0}(st - \varphi(s))$, satisfies the Δ_2 -condition, we say that φ satisfies the ∇_2 -condition and put $\nabla_2(\varphi) := \Delta_2(\tilde{\varphi})$.

Lemma 6.11 ([20, Thm. 4.10, 4.15], [23, Prop. 6.1]). *Let $1 < p_0 < \infty$ and let $\mathcal{F} \subset L^1_{\text{loc}}(\mathbb{R}^n) \times L^1_{\text{loc}}(\mathbb{R}^n)$ be a family of pairs of non-negative functions not identically zero, such that for any $w \in A_{p_0}$ there exists $c = c(p_0, [w]_{A_{p_0}}) > 0$ with*

$$(6.31) \quad \int_{\mathbb{R}^n} f(x)^{p_0} w(x) \, dx \leq c \int_{\mathbb{R}^n} g(x)^{p_0} w(x) \, dx \quad \text{for all } (f, g) \in \mathcal{F},$$

the inequality tacitly implying finiteness of the left-hand side. Then the following hold:

- (a) *If $(\mathbb{X}, \|\cdot\|_{\mathbb{X}})$ is a rearrangement invariant function space whose lower and upper Boyd indices $p_{\mathbb{X}}, q_{\mathbb{X}}$ satisfy $1 < p_{\mathbb{X}} \leq q_{\mathbb{X}} < \infty$, then there holds*

$$(6.32) \quad \|f\|_{\mathbb{X}} \leq c \|g\|_{\mathbb{X}} \quad \text{for all } (f, g) \in \mathcal{F}.$$

- (b) *For any N-function $\varphi : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ with $\Delta_2(\varphi), \nabla_2(\varphi) < \infty$ there exists a constant $C = C(\Delta_2(\varphi), \nabla_2(\varphi), p_0) > 0$ such that there holds*

$$(6.33) \quad \int_{\mathbb{R}^n} \varphi(f) \, dx \leq C \int_{\mathbb{R}^n} \varphi(cg) \, dx \quad \text{for all } (f, g) \in \mathcal{F}.$$

By virtue of Theorem 6.9, Lemma 6.11 implies a whole family of Korn-type inequalities for \mathbb{C} -elliptic differential operators on John domains, out of which we point out two:

Corollary 6.12 (Korn for Lorentz). *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded domain that satisfies the emanating chain condition with constants $\sigma_1 > 1$ and $\sigma_2 \geq 1$ and central ball B . Let $1 < p < \infty$, $1 \leq q < \infty$. Then for any \mathbb{C} -elliptic operator \mathbb{A} of order k there exists a constant $c = c(\sigma_1, \sigma_2, p, q, \mathbb{A}) > 0$ such that for all $u \in W^{\mathbb{A}} L^{p,q}(\Omega)$ there holds*

$$(6.34) \quad \begin{aligned} \|D^k(u - \Pi_{\mathbb{A}}^B u)\|_{L^{p,q}(\Omega)} &\leq c \|\mathbb{A}u\|_{L^{p,q}(\Omega)}, \\ \|D^k u\|_{L^{p,q}(\Omega)} &\leq c \text{diam}(\Omega)^{-k} \|u\|_{L^{p,q}(\Omega)} + c \|\mathbb{A}u\|_{L^{p,q}(\Omega)}. \end{aligned}$$

Proof. For $p_0 := \frac{1}{2}(p+1)$, put

$$\mathcal{F} := \{(\mathbb{1}_\Omega \mathbb{1}_{\{|D^k(u - \Pi_{\mathbb{A}}^B u)| \leq j\}} |D^k(u - \Pi_{\mathbb{A}}^B u)|, \mathbb{1}_\Omega |\mathbb{A}u|) : u \in W^{\mathbb{A}, p_0}(\Omega), j \in \mathbb{N}\}$$

so that \mathcal{F} meets the requirements of Lemma 6.11 by virtue of Theorem 6.9. The Lorentz space $L^{p,q}(\mathbb{R}^n)$ is rearrangement invariant and by our assumptions on (p, q) , $p_{\mathbb{X}} = q_{\mathbb{X}} = p \in (1, \infty)$ (cf. [20, p. 70]). Therefore, (6.32) and Fatou's lemma yield (6.34) for all $u \in W^{\mathbb{A}, p_0}(\Omega)$. Since $W^{\mathbb{A}} L^{p,q}(\Omega) \subset W^{\mathbb{A}, p_0}(\Omega)$ as $1 < p_0 < p$ and Ω is bounded, this implies (6.34)₁ for all $u \in W^{\mathbb{A}} L^{p,q}(\Omega)$, and (6.34)₂ follows by inverse estimates. \square

Remark 6.13 (L^∞ -estimates). If $\Omega \subset \mathbb{R}^n$ is as in Corollary 6.12 and moreover is such that the embedding $W^1 L^{n,1}(\Omega; V) \hookrightarrow C(\overline{\Omega}; V)$ holds (cf. [17, 43, 58]), then Corollary 6.12 immediately yields the stronger embedding $W^{\mathbb{A}} L^{n,1}(\Omega) \hookrightarrow C(\overline{\Omega}; V)$ for any first order \mathbb{C} -elliptic differential operator \mathbb{A} .

In an analogous way as for Corollary 6.12, we obtain the following

Corollary 6.14 (Korn for Orlicz). *Let $\Omega \subset \mathbb{R}^n$ and \mathbb{A} be as in Corollary 6.12 and let $\varphi: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be an N -function that satisfies the Δ_2 - and ∇_2 -conditions. Then there exists $c = c(\sigma_1, \sigma_2, \Delta_2(\varphi), \nabla_2(\varphi), \mathbb{A}) > 0$ such that for all $u \in W^{\mathbb{A}} L^\varphi(\Omega)$ there holds*

$$(6.35) \quad \begin{aligned} \int_{\Omega} \varphi(|D^k(u - \Pi_{\mathbb{A}}^B u)|) dx &\leq c \int_{\Omega} \varphi(|\mathbb{A}u|) dx, \\ \int_{\Omega} \varphi(|D^k u|) dx &\leq c \text{diam}(\Omega)^n \varphi\left(\text{diam}(\Omega)^{-k} \int_{\Omega} |u| dx\right) + c \int_{\Omega} \varphi(|\mathbb{A}u|) dx. \end{aligned}$$

Remark 6.15. In Theorem 6.9 and Corollaries 6.12, 6.14, the main focus is on functions that do not vanish close to $\partial\Omega$, as otherwise the weaker ellipticity allows the use of global singular integral estimates (cf. [19, 41]). We also wish to point out that, whereas (6.25)₂, (6.34)₂ and (6.35)₂ are in fact equivalent to the \mathbb{C} -ellipticity of \mathbb{A} , this is *not* the case for inequalities (6.25)₁, (6.34)₁ and (6.35)₁, cf. FUCHS [27]. This can be seen by the operator ε^D in $n = 2$ dimensions, cf. Example 3.2, and the classical Cauchy-Pompeiu-formula.

7. APPENDIX

7.1. \mathbb{A} -strict continuity of the trace operator on $BV^{\mathbb{A}}$. The approximation argument underlying the proof of Lemma 4.6 to conclude the existence of a continuous linear trace operator $\text{Tr}_{\partial\Omega}: W^{\mathbb{A}, p}(\Omega) \rightarrow B_{p,p}^{k-1/p}(\partial\Omega; V)$ cannot be applied to $BV^{\mathbb{A}}(\Omega)$. This is due to the non-density of $C^\infty(\Omega; V) \cap BV^{\mathbb{A}}(\Omega)$ in $BV^{\mathbb{A}}(\Omega)$ for the norm topology. In the well-known case of $BV(\Omega)$ (cf. AMBROSIO et al. [3]), this issue is resolved by passing to the *strict metric*. The suitable substitute here is the \mathbb{A} -strict metric, see Section 3.1, but now we have to keep track of the intermediate derivatives as well. To this end, we require a multiplicative inequality as follows:

Lemma 7.1 (Intermediate derivatives). *Let $k \geq 1$ and let \mathbb{A} be a k -th order \mathbb{C} -elliptic differential operator of the form (1.1). Then there exists a constant $c = c(\mathbb{A}) > 0$ such that for all $l \in \{1, \dots, k-1\}$ there holds*

$$(7.1) \quad \|D^l u\|_{L^1(\mathbb{R}^n)} \leq c \|u\|_{L^1(\mathbb{R}^n)}^{1-\frac{l}{k}} \|\mathbb{A}u\|_{L^1(\mathbb{R}^n)}^{\frac{l}{k}}$$

for all $u \in W^{\mathbb{A}, 1}(\mathbb{R}^n)$ with compact support.

Proof. The proof is an adaptation of an argument to be found, e.g., in MAZ'YA [49, Thm. 1.4.7]. Let $u \in C_c^\infty(\mathbb{R}^n; V) \setminus \{0\}$ be supported in $B(x_0, R_0)$ and $l \in \{1, \dots, k-1\}$

be given. Let $x \in \text{spt}(u)$, so that because of $u \in C^\infty(\mathbb{R}^n; V)$, $\|u\|_{L^1(B(x,R))} > 0$ for all $R > 0$. Fix $r_0 > 0$. There are two options: If $\|u\|_{L^1(B(x,r_0))} < r_0^k \|\mathbb{A}u\|_{L^1(B(x,r_0))}$, then we have

$$(7.2) \quad r_0^{-l} < \left(\frac{\|\mathbb{A}u\|_{L^1(B(x,r_0))}}{\|u\|_{L^1(B(x,r_0))}} \right)^{\frac{1}{k}},$$

and define $r_x = r_0$ in this case. Then, by virtue of the Poincaré inequality from Theorem 3.7,

$$(7.3) \quad \begin{aligned} \|D^l u\|_{L^1(B(x,r_x))} &\stackrel{(3.20)}{\leq} cr_x^{-l} \|u\|_{L^1(B(x,r_x))} + cr_x^{k-l} \|\mathbb{A}u\|_{L^1(B(x,r_x))} \\ &\stackrel{(7.2)}{\leq} c \|u\|_{L^1(B(x,r_x))}^{1-\frac{1}{k}} \|\mathbb{A}u\|_{L^1(B(x,r_x))}^{\frac{1}{k}} + cr_0^{k-l} \|\mathbb{A}u\|_{L^1(B(x,r_x))}. \end{aligned}$$

If $\|u\|_{L^1(B(x,r_0))} \geq r_0^k \|\mathbb{A}u\|_{L^1(B(x,r_0))}$, we then find $r_x \geq r_0$ such that $\|u\|_{L^1(B(x,r_x))} = r_x^k \|\mathbb{A}u\|_{L^1(B(x,r_x))}$. This follows from u being compactly supported and the fact that $\|u\|_{L^1(B(x,R))}$ will eventually be constant for large values of R . Fix this r_x , so that

$$(7.4) \quad \|D^l u\|_{L^1(B(x,r_x))} \stackrel{(3.20)}{\leq} c \|u\|_{L^1(B(x,r_x))}^{1-\frac{1}{k}} \|\mathbb{A}u\|_{L^1(B(x,r_x))}^{\frac{1}{k}}$$

Adding (7.3) and (7.4), in each of the cases there holds

$$(7.5) \quad \|D^l u\|_{L^1(B(x,r_x))} \leq c(r_0^{k-l} \|\mathbb{A}u\|_{L^1(B(x,r_x))} + \|u\|_{L^1(B(x,r_x))}^{1-\frac{1}{k}} \|\mathbb{A}u\|_{L^1(B(x,r_x))}^{\frac{1}{k}}).$$

Now, $\mathcal{B} := (B(x, r_x))_{x \in \text{spt}(u)}$ is a covering of $\text{spt}(u)$ by open balls such that $r_x \leq C < \infty$ and the sequence of radii of any sequence of mutually disjoint balls in \mathcal{B} tends to zero. By the Besicovitch-type covering lemma in the form as given in MAZ'YA [49, Chpt. 1.2.1, Thm. 1], there exists a sequence of balls $B_m \in \mathcal{B}$ such that $\text{spt}(u) \subset \bigcup_m B_m$, the balls $\frac{1}{3}B_m$ are mutually disjoint and $\bigcup_{B \in \mathcal{B}} B \subset \bigcup_m 4B_m$. Moreover, there exists a constant $N = N(n) \in \mathbb{N}$ such that every point belongs to at most N balls contained in \mathcal{B} . We then conclude by (7.5) that

$$\begin{aligned} \|D^l u\|_{L^1(\text{spt}(u))} &\leq \sum_m \|D^l u\|_{L^1(B_m)} \\ &\leq c \sum_m \left(r_0^{k-l} \|\mathbb{A}u\|_{L^1(B_m)} + \|u\|_{L^1(B_m)}^{1-\frac{1}{k}} \|\mathbb{A}u\|_{L^1(B_m)}^{\frac{1}{k}} \right) \\ &\leq cr_0^{k-l} \|\mathbb{A}u\|_{L^1(\mathbb{R}^n)} + c \|u\|_{L^1(\mathbb{R}^n)}^{1-\frac{1}{k}} \|\mathbb{A}u\|_{L^1(\mathbb{R}^n)}^{\frac{1}{k}}, \end{aligned}$$

where $c = c(n, \mathbb{A}) > 0$. Now it suffices to send $r_0 \searrow 0$ to conclude (7.1) for $u \in C_c^\infty(\mathbb{R}^n; V)$; the general case of compactly supported $u \in W^{\mathbb{A},1}(\mathbb{R}^n)$ follows by smooth approximation. The proof is complete. \square

We now turn to the remaining part of the

Proof of Theorem 4.7. We start by noting that the existence of a norm-continuous trace operator $\text{Tr}_{\partial\Omega}: W^{\mathbb{A},1}(\Omega) \rightarrow B_{1,1}^{k-1}(\partial\Omega; V)$ follows by smooth approximation along the same lines as for the case $1 < p < \infty$ (cf. Theorem 4.4, direction '(a) \Rightarrow (b) \Rightarrow (c)' and Lemma 4.6, now using Theorem 1.1), the reason being the density of $\widetilde{W}^{\mathbb{A},1}(\Omega) \cap C^\infty(\Omega; V)$ in $W^{\mathbb{A},1}(\Omega)$ for the norm topology on $W^{\mathbb{A},1}(\Omega)$; also note that $W^{\mathbb{A},1}(\Omega) = \widetilde{W}^{\mathbb{A},1}(\Omega)$ by Theorem 3.7.

By Theorem 3.7, $BV^{\mathbb{A}}(\Omega) = W^{k-1,1}(\Omega; V) \cap BV^{\mathbb{A}}(\Omega)$. To define an \mathbb{A} -strictly continuous trace operator by means of smooth approximation, the localisation argument below requires $d(u, u_j) := \|u_j - u\|_{W^{k-1,1}(\Omega)} + \|\mathbb{A}u_j|(\Omega) - \mathbb{A}u|(\Omega)\| \rightarrow 0$ whenever $d_{\mathbb{A}}(u, u_j) = \|u_j - u\|_{L^1(\Omega)} + \|\mathbb{A}u_j|(\Omega) + \mathbb{A}u|(\Omega)\| \rightarrow 0$ as $j \rightarrow \infty$. The Poincaré inequality of Theorem 3.7 does not directly allow for this conclusion. Also

note that, if we define a trace operator as the continuous extension for the metric d as above, then it is not clear that this trace operator will be continuous for $d_{\mathbb{A}}$. It is here where we require the multiplicative inequality of Lemma 7.1.

Let $u \in \text{BV}^{\mathbb{A}}(\Omega) = \text{W}^{k-1,1}(\Omega; V) \cap \text{BV}^{\mathbb{A}}(\Omega)$ and choose $(u_j) \subset C^\infty(\Omega; V) \cap \text{BV}^{\mathbb{A}}(\Omega)$ such that $d_{\mathbb{A}}(u_j, u) \rightarrow 0$ as $j \rightarrow \infty$. We then have $L := \sup_j \|u_j\|_{\text{W}^{\mathbb{A},1}(\Omega)} < \infty$. Since \mathbb{A} is \mathbb{C} -elliptic, the Jones-type extension operator \tilde{E} from [30, Sec. 4.1] maps $\tilde{E}: \widetilde{\text{W}}^{\mathbb{A},1}(\Omega) \rightarrow \widetilde{\text{W}}^{\mathbb{A},1}(\mathbb{R}^n)$ boundedly; clearly, we may assume that all maps $\tilde{E}u$ are compactly supported in a fixed ball $B(0, R)$. This operator moreover satisfies $\|\tilde{E}u\|_{L^1(\mathbb{R}^n)} \leq c\|u\|_{L^1(\Omega)}$. Thus, for all $\ell \in \{0, \dots, k-1\}$,

$$\begin{aligned}
(7.6) \quad \|D^\ell(u_j - u_m)\|_{L^1(\Omega)} &\leq \|D^\ell(\tilde{E}(u_j - u_m))\|_{L^1(\mathbb{R}^n)} \\
&\stackrel{(7.1)}{\leq} c\|\tilde{E}(u_j - u_m)\|_{L^1(\mathbb{R}^n)}^{1-\frac{\ell}{k}} \|\mathbb{A}(\tilde{E}(u_j - u_m))\|_{L^1(\mathbb{R}^n)}^{\frac{\ell}{k}} \\
&\leq c\|u_j - u_m\|_{L^1(\Omega)}^{1-\frac{\ell}{k}} \|u_j - u_m\|_{\text{W}^{\mathbb{A},1}(\Omega)}^{\frac{\ell}{k}} \\
&\leq cL^{\frac{\ell}{k}} \|u_j - u_m\|_{L^1(\Omega)}^{1-\frac{\ell}{k}} \rightarrow 0
\end{aligned}$$

as $j, m \rightarrow \infty$, and so with d as defined above, $d(u_j, u) \rightarrow 0$. We pick a sequence $(\rho_l) \subset C^\infty(\bar{\Omega}; [0, 1])$ such that $\rho_l = 1$ in a $\frac{1}{l}$ -neighbourhood of $\partial\Omega$ and $\rho_l = 0$ in $\Omega \setminus \Omega'_l := \{x \in \Omega: \text{dist}(x, \partial\Omega) > \frac{2}{l}\}$. We may then assume that $|\nabla^\ell \rho_l| \leq cl^\ell$ for some $c > 0$ and all $\ell \in \{1, \dots, k\}$. For each $l \in \mathbb{N}$, we find a set Ω_l with $\text{spt}(\rho_l) \subset \Omega_l \subset \Omega'_l$ and of smooth boundary such that $|\mathbb{A}u|(\partial\Omega_l) = 0$. Denoting $\text{Tr}_{\partial\Omega}$ the trace operator on $\text{W}^{\mathbb{A},1}(\Omega)$ that is already our disposal, we then estimate

$$\begin{aligned}
\|\text{Tr}_{\partial\Omega}(u_j) - \text{Tr}_{\partial\Omega}(u_m)\|_{\text{B}_{1,1}^{k-1}(\partial\Omega)} &\leq c(\|\rho_l(u_j - u_m)\|_{L^1(\Omega)} + \|\mathbb{A}(\rho_l(u_j - u_m))\|_{L^1(\Omega)}) \\
&\leq c\left(\sum_{\substack{|\alpha|+|\beta|\leq k, \\ |\beta|\leq k-1}} \|\partial^\alpha \rho_l \partial^\beta(u_j - u_m)\|_{L^1(\Omega)}\right) \\
&\quad + c\|\rho_l(\mathbb{A}u_j - \mathbb{A}u_m)\|_{L^1(\Omega)} =: \text{I}_{l,j,m} + \text{II}_{l,j,m}.
\end{aligned}$$

For an arbitrary $l \in \mathbb{N}$, (7.6) implies that $\text{I}_{l,j,m} \rightarrow 0$ as $j, m \rightarrow \infty$. Therefore, sending $j, m \rightarrow \infty$ first and then letting $l \rightarrow \infty$,

$$\lim_{j,m \rightarrow \infty} \|\text{Tr}_{\partial\Omega}(u_j) - \text{Tr}_{\partial\Omega}(u_m)\|_{\text{B}_{1,1}^{k-1}(\partial\Omega)} \leq c|\mathbb{A}u|(\Omega_l) \rightarrow 0$$

as $l \rightarrow \infty$; note that $|\mathbb{A}u|$ is a finite Radon measure and $\bigcap_l \Omega_l = \emptyset$. Hence $(\text{Tr}_{\partial\Omega}(u_j))$ is Cauchy in $\text{B}_{1,1}^{k-1}(\partial\Omega; V)$, too. Working from here, the claimed existence of a trace operator with the requisite properties is routine, and the proof thus is complete. \square

7.2. Oscillation characterisation of $\dot{\text{B}}_{p,q}^s$. To the best of our knowledge, the oscillation characterisation of Besov spaces is only available in the inhomogeneous situation, cf. TRIEBEL [63, Chpt. 3.5]. For completeness, we briefly explain how to arrive at the part of Lemma 4.1 that is required for Section 4.3.

Let $M > \lfloor s \rfloor$ and choose $2N > M$. Let $k \in C_c^\infty(\mathbb{R}^n)$ be such that $\text{spt}(k) \subset B(0, 1)$ and consider, for $f \in \mathcal{S}(\mathbb{R}^n)$, the means

$$k(t, f)(x) := \int_{\mathbb{R}^n} k(y)f(x + ty) dy, \quad t > 0, x \in \mathbb{R}^n.$$

Given some $k^0 \in C_c^\infty(\mathbb{R}^n)$ with $\text{spt}(k^0) \subset B(0, 1)$ and $\widehat{k^0}(0) \neq 0$, we put $k := \Delta^N k^0$, where Δ^N stands for the N -fold application of the Laplacean. We now define $\varphi := k^\vee$. It is not too difficult to verify that φ satisfies nonsubstantial modifications

of [62, (62),(78),(79)] (also see [63, Proof of Thm. 2.4.6]) and so, by a combination of [62, Rem. 17, Rem. 18],

$$(7.7) \quad \|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} \approx \left(\int_0^\infty \|\mathcal{F}^{-1}[\varphi(t\cdot)\mathcal{F}f]\|_{L^p(\mathbb{R}^n)}^q \frac{dt}{t^{1+sq}} \right)^{\frac{1}{q}}$$

is an equivalent norm on $\dot{B}_{p,q}^s(\mathbb{R}^n)$. By definition of φ , we have

$$(7.8) \quad \|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} = c(n) \left(\int_0^\infty \|k(t, f)\|_{L^p(\mathbb{R}^n)}^q \frac{dt}{t^{1+sq}} \right)^{\frac{1}{q}}.$$

Since $2N > M$, we have for any $\pi \in \mathcal{P}_M(\mathbb{R}^n)$ that $\Delta^N \pi = 0$ and so, integrating by parts and Jensen's inequality imply

$$|k(t, f)(x)| \leq \int_{\mathbb{R}^n} |k(y)(f - \pi)(x + ty)| dy \leq c(n, N, k^0) \left(\int_{B(x,t)} |f - \pi|^u dy \right)^{\frac{1}{u}}$$

and infimising the very right hand side over $\pi \in \mathcal{P}_M(\mathbb{R}^n)$ yields

$$(7.9) \quad |k(t, f)(x)| \leq c(n) \operatorname{osc}_u^M f(x, t).$$

Combining (7.9) with (7.7) and (7.8) consequently yields the existence of a constant $c = c(n, N, s, p, q, k^0) > 0$ such that

$$\|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} \leq c \left(\int_0^\infty \|\operatorname{osc}_u^M f(\cdot, t)\|_{L^p(\mathbb{R}^n)}^q \frac{dt}{t^{1+sq}} \right)^{\frac{1}{q}} \quad \text{for all } f \in \mathcal{S}(\mathbb{R}^n).$$

The statement for $\dot{B}_{p,q}^s(\mathbb{R}^n; V)$ then follows by componentwise application.

REFERENCES

- [1] V. Adolfsson and J. Pipher. The inhomogeneous Dirichlet problem for Δ^2 in Lipschitz domains. *J. Funct. Anal.*, 159(1):137–190, 1998.
- [2] H. Aikawa. Potential analysis on nonsmooth domains—Martin boundary and boundary Harnack principle. In *Complex analysis and potential theory*, volume 55 of *CRM Proc. Lecture Notes*, pages 235–253. Amer. Math. Soc., Providence, RI, 2012.
- [3] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, 2000.
- [4] N. Aronszajn. On coercive integro-differential quadratic forms. *Conference on Partial Differential Equations, University of Kansas, Technical Report*, (No. 14):94–106, 1954.
- [5] J. Bergh and J. Löfström. *Interpolation spaces. An introduction*. Springer-Verlag, Berlin, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.
- [6] J. Boman. L^p -estimates for very strongly elliptic systems. *Technical Report 29, Technical Report 29, Department of Mathematics, University of Stockholm, Sweden, 1982*.
- [7] J. Bourgain and H. Brezis. New estimates for the Laplacian, the div-curl, and related Hodge systems. *C. R. Math. Acad. Sci. Paris*, 338(7):539–543, 2004.
- [8] J. Bourgain and H. Brezis. New estimates for elliptic equations and Hodge type systems. *J. Eur. Math. Soc. (JEMS)*, 9(2):277–315, 2007.
- [9] P. Bousquet and J. Van Schaftingen. Hardy–Sobolev inequalities for vector fields and canceling linear differential operators. *Indiana Univ. Math.*, 63(5):1419–1445, 2014.
- [10] D. Breit and L. Diening. Sharp conditions for Korn inequalities in Orlicz spaces. *J. Math. Fluid Mech.*, 14:565–573, 2012.
- [11] D. Breit, L. Diening, and F. Gmeiner. On the trace operator for functions of bounded A-variation. *Anal. PDE*, 13(2):559–594, 2020.
- [12] S.C. Brenner and L.R. Scott. *The mathematical theory of finite element methods*, volume 15 of *Texts in Applied Mathematics*. Springer, New York, third edition, 2008.
- [13] H. Brezis and A.C. Ponce. Kato's inequality up to the boundary. *Commun. Contemp. Math.*, 210(6):1217–1241, 2008.
- [14] S. Buckley, P. Koskela, and G. Lu. Boman equals John. *XVIIth Rolf Nevanlinna Colloquium (Joensuu, 1995), de Gruyter, Berlin*, pages 91–99, 1996.
- [15] A.P. Calderón and A. Zygmund. On singular integrals. *Amer. J. Math.*, 78:289–309, 1956.
- [16] A. Cianchi. Korn type inequalities in Orlicz spaces. *J. Funct. Anal.*, 267:2313–2352, 2014.

- [17] A. Cianchi and L. Pick. Sobolev embeddings into BMO, VMO, and L^∞ . *Ark. Mat.*, 36:317–340, 1998.
- [18] R. Coifman and C. Fefferman. Weighted norm inequalities for maximal functions and singular integrals. *Studia Math.*, 51(3):241–250, 1974.
- [19] S. Conti and F. Gmeineder. \mathcal{A} -quasiconvexity and partial regularity. <https://arxiv.org/abs/2009.13820>.
- [20] D.V. Cruz-Uribe, J.M. Martell, and C. Pérez. *Weights, extrapolation and the theory of Rubio de Francia*, volume 215 of *Operator Theory: Advances and Applications*. Birkhäuser/Springer Basel AG, Basel, 2011.
- [21] L. Diening and F. Gmeineder. Continuity points via riesz potentials for \mathbb{C} -elliptic operators. *The Quarterly Journal of Mathematics*, 71(4):1201–1218, 2020.
- [22] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička. *Lebesgue and Sobolev spaces with variable exponents*, volume 2017 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2011.
- [23] L. Diening, M. Růžička, and K. Schumacher. A decomposition technique for John domains. *Ann. Acad. Sci. Fenn. Math.*, 35(1):87–114, 2010.
- [24] J.R. Dorronsoro. Mean oscillation and Besov spaces. *Canad. Math. Bull.*, 28(3), 1985.
- [25] J. Duoandikoetxea. *Fourier Analysis* (translated and revised by David Cruz-Uribe, SFO), volume 29 of *Graduate Studies in Mathematics*. American Mathematical Society, 2001.
- [26] L.C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, 1998.
- [27] M. Fuchs. An estimate for the distance of a complex valued Sobolev function defined on the unit disc to the class of holomorphic functions. *J. Applied Analysis*, 17(1):131–135, 2011.
- [28] M. Fuchs and G. Seregin. Variational methods for problems from plasticity theory and for generalized Newtonian fluids. *Ann. Univ. Sarav. Ser. Math.*, 10(1):iv+283, 1999.
- [29] E. Gagliardo. Caratterizzazione delle sulla frontiera relative ad alcune tracce classi di funzioni in n variabili. *Rend. Sem. Mat. Univ. Padova*, 27:284–305, 1957.
- [30] F. Gmeineder and B. Raita. Embeddings for \mathbb{A} -weakly differentiable functions on domains. *J. Func. Anal.*, 277(12):108278, 2019.
- [31] F. Gmeineder, B. Raita, and J. Van Schaftingen. On limiting trace inequalities for vectorial differential operators. *To appear at Indiana Math. Univ. J.*
- [32] J. Heinonen, T. Kilpeläinen, and O. Martio. *Nonlinear Potential Theory of Degenerate Elliptic Equations*. Dover Books on Mathematics. Dover Publications, 2006. Reprint of the 1993 edition published by Oxford University Press.
- [33] S. Hofmann, M. Mitrea, and M. Taylor. Singular integrals and elliptic boundary problems on regular Semmes-Kenig-Toro domains. *Int. Math. Res. Not. IMRN*, (14):2567–2865, 2010.
- [34] L. Hörmander. Differentiability properties of solutions of systems of differential equations. *Ark. Mat.*, 3:527–535, 1958.
- [35] R. Hurri. Poincaré domains in \mathbb{R}^n . *Ann. Acad. Sci. Fenn. Ser. A I*, 71:1–42, 1988.
- [36] T. Iwaniec. On L^p -integrability in PDEs and quasiregular mappings for large exponents. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 7(2):301–322, 1982.
- [37] T. Iwaniec and C. A. Nolder. Hardy-Littlewood inequality for quasiregular mappings in certain domains in \mathbb{R}^n . *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 10:267–282, 1985.
- [38] D. Jerison and C. Kenig. Boundary behaviour of harmonic functions in non-tangentially accessible domains. *Adv. Math.*, 47(1):80–147, 1982.
- [39] R. Jiang and A. Kauranen. Korn’s inequality and John domains. *J. Calc. Var.*, 56:109, 2017.
- [40] F. John. Rotation and strain. *Comm. Pure Appl. Math.*, 14:391–413, 1961.
- [41] A. Kałamajska. Coercive inequalities on weighted Sobolev spaces. *Colloq. Math.*, LXVI(2):309–318, 1993.
- [42] A. Kałamajska. Pointwise multiplicative inequalities and Nirenberg type estimates in weighted Sobolev spaces. *Studia Math.*, 108(3):275–290, 1994.
- [43] J. Kauhanen, P. Koskela, and J. Malý. On functions with derivatives in a Lorentz space. *Manuscripta Math.*, 100:87–101, 1999.
- [44] Stolyarov D.M. Kazaniecki, K. and M. Wojciechowski. Anisotropic Ornstein non- inequalities. *Analysis & PDE*, 10(2), 2017.
- [45] B. Kirchheim and J. Kristensen. On rank one convex functions that are homogeneous of degree one. *Arch. Ration. Mech. Anal.*, 221(1), 1985.
- [46] A. Korn. Über einige Ungleichungen, welche in der Theorie der elastischen und elektrischen Schwingungen eine Rolle spielen. *Classe des Sciences Mathématiques et Naturels*, (9. Novembre):705–724, 1909.

- [47] G. Leoni and I. Tice. Traces for homogeneous Sobolev spaces in infinite strip-like domains. *J. Funct. Anal.*, 277(7):2288–2380, 2019.
- [48] O. Martio and J. Sarvas. Injectivity theorems in plane and space. *Ann. Acad. Sci. Fenn. Ser. A I Mth.*, 4:383–401, 1978–79.
- [49] V.G. Maz'ya. *Sobolev spaces*. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1985. Translated from the Russian by T. O. Shaposhnikova.
- [50] V.G. Maz'ya, M. Mitrea, and T. Shaposhnikova. The Dirichlet problem in Lipschitz domains with boundary data in Besov spaces for higher order elliptic systems with rough coefficients. *arXiv.math.AP/0505372*.
- [51] N. Miller. Weighted Sobolev spaces and pseudodifferential operators with smooth symbols. *Trans. Amer. Math. Soc.*, 269(1):91–109, 1982.
- [52] P. Mironescu and E. Russ. Traces of weighted Sobolev spaces. Old and new. *Nonlinear Analysis TMA*, 119:354–381, 2015.
- [53] D. Ornstein. A non-equality for differential operators in the L_1 -norm. *Arch. Rational Mech. Anal.*, 11:40–49, 1962.
- [54] B. Raita. Critical L^p -differentiability of BV^A -maps and canceling operators. *Transactions of the American Mathematical Society*, 372(10):7297–7326, 2019.
- [55] Yu. G. Reshetnyak. Estimates for certain differential operators with finite-dimensional kernel. (Russian). *Sibirsk. Mat. Z.*, 11:414–428, 1970.
- [56] K. T. Smith. Formulas to represent functions by their derivatives. *Math. Ann.*, 188:53–77, 1970.
- [57] D. C. Spencer. Overdetermined systems of linear partial differential equations. *Bull. Amer. Math. Soc.*, 75:179–239, 1969.
- [58] E.M. Stein. Editor's note: The differentiability of functions in \mathbb{R}^n . *Annals of Mathematics*, 113:383–385, 1981.
- [59] E.M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.
- [60] L. Tartar. *An Introduction to Sobolev Spaces and Interpolation Spaces*, volume 3 of *Lecture Notes of the Unione Matematica Italiana*. Springer Science & Business Media, 2007.
- [61] H. Triebel. *Theory of function spaces*, volume 78 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1983.
- [62] H. Triebel. Characterizations of Besov-Hardy-Sobolev spaces: A unified approach. *J. Approx. Theory*, 52:162–203, 1988.
- [63] H. Triebel. *Theory of function spaces. II*, volume 84 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1992.
- [64] S.V. Uspenskii. Imbedding theorems for classes with weights. *Tr. Mat. Inst. Steklova*, 60:282–303, 1961. Am. Math. Soc. Transl. 87 (1970) 121–145. English translation.
- [65] J. Väisälä. Exhaustion of John domains. *Ann. Acad. Sci. Fenn., Series A. I. Mathematica*, 19:47–57, 1994.
- [66] J. Van Schaftingen. Estimates for L^1 -vector fields. *C. R. Math. Acad. Sci. Paris*, 339, 2004.
- [67] J. Van Schaftingen. Limiting Sobolev inequalities for vector fields and canceling linear differential operators. *J. Eur. Math. Soc. (JEMS)*, 15(3):877–921, 2013.
- [68] J. Van Schaftingen. Limiting Bourgain–Brezis estimates for systems of linear differential equations: Theme and variations. *J. Fixed Point Theory Appl.*, 15, 2014.

(L. Diening) FACULTY OF MATHEMATICS, UNIVERSITY OF BIELEFELD, UNIVERSITÄTSSTRASSE 25, 33615 BIELEFELD, GERMANY

Email address: lars.diening@uni-bielefeld.de

(F. Gmeiner) MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY (*expiring address*),

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF KONSTANZ, UNIVERSITÄTSSTRASSE 10, 78457 KONSTANZ, GERMANY

Email address: fgmeined@math.uni-bonn.de, franz.gmeined@uni-konstanz.de