On mutually inverse transforms of functions on a half-line

V.Yu.Protasov^{*}, M.E.Shirokov[†]

Abstract

Two transforms of functions on a half-line are considered. It is proved that their composition gives a concave majorant for every nonnegative function. In particular, this composition is the identity transform on the class of nonnegative concave functions. Applications of this result to some problems of mathematical physics are indicated. Several open questions are formulated.

The main result

The following transforms of functions on the positive half-line \mathbb{R} naturally arise in problems of mathematical physics (see Section 3):

$$F[f](x) = \sup_{t>0} \frac{f(xt)}{t+1}, \qquad G[f](x) = \inf_{t>0} f(xt) \left(1 + \frac{1}{t}\right).$$

Throughout this paper, we assume that these transforms are applied to functions on which they are well defined, i.e., F[f](x) and G[f](x) are finite for any x. For example, for F[f], it suffices that the function f(t) be at most linearly growing, i.e.,

$$\sup_{t \in [0,a]} f(t) < C(a+1), \, a > 0.$$

The composition is denoted by G[F[f]] = GF[f]. For concave functions f of a certain class, it was shown in [2] that the composition is an identity transform.

^{*}Lomonosov Moscow State University, National Research University High School of Economics, e-mail: v-protassov@yandex.ru

[†]Steklov Mathematical Institute, e-mail: msh@mi-ras.ru

We will prove that GF[f] = f if and only if f is any monotone concave function satisfying some conditions. Moreover, for an arbitrary nonnegative function f, the composition GF[f] is the smallest concave majorant of f, i.e., it coincides with the function $\tilde{f}(t) = \inf \{\varphi(t) \mid \varphi \ge f, \varphi \text{ concave}\}$. This property of F and G resembles the Legendre–Young duality transform [1], but no relations to it have been found.

Consider two classes of concave functions on \mathbb{R}_+ . The class \mathcal{A} consists of concave nondecreasing nonnegative functions. The graph of each function $\varphi \in \mathcal{A}$ consists of three parts (from left to right): the segment from the point (0,0) to some point A_1 , the arc of the concave function from A_1 to A_2 , and the horizontal ray from A_2 to $+\infty$. Some of these parts may be degenerate, which corresponds to the cases $A_1 = (0, \varphi(0)), A_2 = A_1$ or $A_2 = +\infty$. For example, the function $f \equiv 1$ belongs to \mathcal{A} and for it $A_1 = A_2 = (0, 1)$. Any concave increasing function satisfying the condition $\varphi(0) > 0$ belongs to \mathcal{A} , and, for it $A_1 = (0, \varphi(0)), A_2 = +\infty$. The class \mathcal{B} consists of concave nonincreasing nonpositive functions with the asymptote $y = kt, t \to +\infty$. The graph of such a function consists of three parts (from left to right): a horizontal segment to some point B_1 , the arc of the concave function from B_1 to B_2 , and the ray from B_1 to $+\infty$, whose extension beyond the point \mathcal{B}_2 passes through zero. Some of these parts may be degenerate. For example, for $f \equiv -1$ we have $B_1 = B_2 = +\infty$; while for the function $\varphi(t) = -\frac{1}{t+1} - t$ we have $B_1 = (0, -1), B_2 = +\infty$.

The classes \mathcal{A} and \mathcal{B} are convex and closed in the topology of uniform convergence on each compact set. They intersect only on the zero function. The \mathcal{A} -hull of a function f is the smallest majorant by functions from \mathcal{A} , i.e. $f_{\mathcal{A}}(t) = \inf \{\varphi(t) \mid \varphi \geq f, \varphi \in \mathcal{A}\}$. Similarly we define the \mathcal{B} -hull: $f_{\mathcal{B}}(t) = \inf \{\varphi(t) \mid \varphi \geq f, \varphi \in \mathcal{B}\}$. Note that the \mathcal{A} -hull is defined for any function f with at most linear growth, while the \mathcal{B} -hull is defined only for non-positive f.

Theorem 1 If a function f takes at least one positive value then $GF[f] = f_A$, otherwise $GF[f] = f_B$.

Corollary 1 GF[f] = f if and only if $f \in \mathcal{A} \cup \mathcal{B}$.

Corollary 2 If f is a nonnegative function then GF[f] is the concave majorant of f. For nonnegative functions, the equality GF[f] = f holds if and only if f is concave.

The proof of Theorem 1 is geometric. We begin with an auxiliary assertion. Suppose that the function $f : \mathbb{R}_+ \to \mathbb{R}$ grows at most linearly. Through a given point A = (-a, 0), a > 0 we draw a minimum-slope straight line ℓ lying above the graph of f, i.e. $\ell(t) \geq f(t), t \in \mathbb{R}_+$ (here and below, a linear function is denoted the same as the corresponding line). We call a left supporting line of the graph of f. In contrast to a usual supporting line ℓ , for which $\inf_{t \in \mathbb{R}_+} (\ell(t) - f(t)) = 0$, the left supporting line must intersect the negative horizontal half-line $\{(-a, 0) \mid a \geq 0\}$.

Lemma 1 For any x > 0, the straight line passing through the points (-x, 0) and (0, F[f](x)) is a left supporting line to the graph of the function f.

Proof. At the point with abscissa z, this line has the ordinate

$$y = \frac{(x+z)F[f](x)}{x} = (1+t)F[f](x)$$

where t = z/x. Hence,

$$\inf_{z \ge 0} (y(z) - f(z)) = \inf_{t \ge 0} ((1+t)F[f](x) - f(xt)) \ge 0.$$

Therefore, the line is above the graph of f and its slope cannot be reduced, since $\inf_{t\geq 0} \left(F[f](x) - \frac{1}{1+t} f(xt) \right) = 0.$

Proof of the theorem 1. Denote z = xt. Then $G[f](x) = \inf_{z>0} f(z) \frac{x+z}{z}$. Hence,

$$GF[f](x) = \inf_{z>0} F[f](z) \frac{x+z}{z}.$$
 (1)

From the point A = (-z, 0), we draw a supporting line to the graph of f. It intersects the vertical axis at the point M = (0, F[f](z)). Let also B = (x, 0). Drop a perpendicular to the horizontal axis at the point B; it crosses the supporting line at the point N. Since the triangles AMO and ANB, where O is the origin, are similar, we obtain. BN/OM = AB/AO = (z + x)/z. Thus, BN = F[f](z)(z + x)/z. Combining this with (1), we conclude that GF[f](x) is equal to the smallest value of $\ell(x)$ over all left supporting lines ℓ to the graph of f. Let denote \tilde{f} the smallest concave majorant of f. It is the pointwise minimum of all supporting lines to the graph of f.

Assume that f(t) > 0 at least at one point t. In the graph of f there are points A_1, A_2 (which may coincide or grow to $+\infty$) such that the supporting line to the graph of \tilde{f} at the point A_1 passes through the origin (we use the rightmost of these points), while the supporting line at the point A_2 is horizontal (we use the leftmost point). Thus, A_2 is a maximizer of \tilde{f} and, by assumption, this maximum is positive. For any point between A_1 and A_2 , the supporting line to \tilde{f} intersects the negative horizontal half-line; hence, this is a left supporting line. Therefore, the graph of GF[f] coincides with the graph of f between the points A_1 and A_2 , with the graph of the left supporting line through the origin to the left of A_1 , and with the horizontal line through to the right of A_2 . Thus, $GF[f] = f_{\mathcal{A}}$. The second case is treated in a similar manner: $f \leq 0$ then $GF[f] = f_{\mathcal{B}}$. \Box

2. Comments

Remark 1 After interchanging the factors in the composition, the resulting operator FG is not an identity one in the classes \mathcal{A} and \mathcal{B} . We do not know, for which functions f, it holds that FG[f] = f. One condition can be obtained by applying the following argument. Let denote ℓ_z the line passing through the points (-z, 0) and (0, f(z)). It is easy to see that $G[f](t) = \inf_{z>0} \ell_z(t)$. Then the equality FG[f] = f is equivalent to the fact that each line of the family $\{\ell_z \mid z > 0\}$ is a right supporting line to the graph of G[f]. If f is nonnegative, this condition is equivalent to this requirement for f. Denote by \mathcal{H} the family of functions $\varphi(t) = \frac{at}{t+b}$, a, b > 0. The equality FG[f] = f holds if and only if G[f] is not identically zero and, for any pair of points in the graph of f, the arc of a function from the family \mathcal{H} that connects these points lies above the graph of f. Finding more adequate conditions for the equality FG[f] = f remains an open problem.

3. Some applications

Denote by \mathbb{T} the convex cone in a linear space. Let α, β and γ be nonnegative linear functions on \mathbb{T} such that $\alpha(t) > 0$ for all t from \mathbb{T} not equal to zero. Consider the nonnegative functions

$$f(x) = \sup\{\gamma(t) \mid \alpha(t) \le 1, \beta(t) \le x\}, \quad g(x) = \sup\{\gamma(t) \mid \alpha(t) + x^{-1}\beta(t) \le 1\}$$

on \mathbb{R}_+ . Corollary 1 allows to prove the following

Proposition 1 The above functions are connected by the relations g = F[f]and f = G[g], where F and G are the transforms defined in Section 1.

Proof. It is easy to see that the first function is concave. Therefore, by Corollary 1, it suffices to show that

$$g(x) = \sup_{r \in (0,1)} rf\left(x\frac{1-r}{r}\right) = \sup_{t>0} \frac{f(xt)}{1+t}.$$
 (2)

To prove (2) note first that $g(x) \ge rf\left(x\frac{1-r}{r}\right)$ for all $r \in (0,1]$. For any $\varepsilon > 0$ there is $t_{\varepsilon} \in \mathbb{T}$ such that $\alpha(t_{\varepsilon}) \le 1, \beta(t_{\varepsilon}) \le x\frac{1-r}{r}$ and $f\left(x\frac{1-r}{r}\right) \le \varepsilon$

 $\gamma(t_{\varepsilon}) + \varepsilon$. Then $rt_{\varepsilon} \in \mathbb{T}$ and $\alpha(rt_{\varepsilon}) + x^{-1}\beta(rt_{\varepsilon}) \leq r + (1-r) \leq 1$. Hence $g(x) \geq \gamma(rt_{\varepsilon}) \geq rf\left(x\frac{1-r}{r}\right) - r\varepsilon$. Since ε is arbitrary, we obtain the required inequality.

For any $\varepsilon > 0$ there is $t_{\varepsilon} \in \mathbb{T} \setminus \{0\}$ such that $\alpha(t_{\varepsilon}) + x^{-1}\beta(t_{\varepsilon}) \leq 1$ and $g(x) \leq \gamma(t_{\varepsilon}) + \varepsilon$. Since $r = \alpha(t_{\varepsilon}) \in (0, 1], \ \alpha(t_{\varepsilon}/r) = 1$. Then $\beta(t_{\varepsilon}/r) \leq x(1-r)/r$ and hence $\gamma(t_{\varepsilon}/r) \leq f\left(x\frac{1-r}{r}\right)$. Thus,

$$g(x) \le \gamma(t_{\varepsilon}) + \varepsilon \le rf\left(x\frac{1-r}{r}\right) + r\varepsilon.$$

It suffices to note that the condition $r \in (0, 1)$ in (2) can be replaced by the condition $r \in (0, 1]$ by the concavity of f.

Example 1. Let \mathbb{T} be the cone of positive operators on a Hilbert space H with the inner product $\langle \cdot, \cdot \rangle$ and the norm $|\cdot|$. To avoid the technical difficulties, we assume that H is finite-dimensional.

Given any positive operator G on H with a zero lower spectral bound and any E > 0, the space H can be equipped with the inner product

$$\langle \varphi, \psi \rangle_E^G = \langle \varphi, \psi \rangle + \langle \varphi, G\psi \rangle / E, \quad \varphi, \psi \in H$$

and the corresponding norm $|\varphi|_E^G = \sqrt{\langle \varphi, \varphi \rangle_E^G}$ of any vector $\varphi \in H$. Then, for any linear operator A on H, the quantity

$$|||A|||_E^G = \sup\{|A\varphi| \mid \varphi \in H, \ |\varphi|_E^G \le 1\}$$

is the operator norm of A regarded as an operator from the space H with the inner product $\langle \cdot, \cdot \rangle_E^G$ to the space H with the inner product $\langle \cdot, \cdot \rangle$. One can show (see [2]) that

$$|||A|||_E^G = \sup\{\sqrt{\mathrm{Tr}AXA^*} \,|\, X \in \mathbb{T}, \, \mathrm{Tr}X + E^{-1}\mathrm{Tr}GX \le 1\}$$

For any linear operator A on H, we can define the family of norms

$$||A||_E^G = \sup\{|A\varphi| \,|\, \varphi \in H, \, |\varphi| \le 1, \, \langle \varphi, G\varphi \rangle \le E\}, \quad E > 0,$$

which are useful for a number of problems of mathematical physics [2].¹ It is proved in [2] that

$$||A||_E^G = \sup\{\sqrt{\operatorname{Tr} AXA^*} \,|\, X \in \mathbb{T}, \, \operatorname{Tr} X \le 1, \, \operatorname{Tr} GX \le E\}.$$

¹Specific properties of these norms are exhibited in the case of an unbounded operator G and an infinite-dimensional Hilbert space H, in which these norms define topologies weaker than the topology of the operator norm.

This representation shows that the function $E \to [||A||_E^G]^2$ is concave on \mathbb{R}_+ and tends to the ordinary operator norm of A as $E \to +\infty$. By Proposition 1 for each operator A on H the nonnegative functions $f_A(E) = [||A||_E^G]^2$ and $g_A(E) = [|||A|||_E^G]^2$ are related by the transforms F and G:

$$g_A = F[f_A], \qquad f_A = G[g_A]. \tag{3}$$

These relations were obtained in [2, Theorem 3] by applying a rather complicated method involving special properties of the functions f_A and g_A . Proposition 1 shows that these relations are a special case of the more general property presented in Corollary 1.

The norms $\|\cdot\|_{E}^{G}$ (naturally defined for operators on separable Hilbert spaces in the case of an unbounded operator G) were found useful for some problems in the theory of open quantum systems. They were used to prove a generalization of the Kretschmann– Schlingemann–Werner theorem [2, Theorem 2], which made it possible to obtain a Stinespring representation for strongly converging sequences of quantum channels. These norms also arise in describing strongly continuous quantum dynamic semigroups in the case of energy constraints imposed on the states of the quantum system [3].

The norms $||| \cdot |||_E^G$ naturally arise in the theory of relatively bounded operators on infinite-dimensional Hilbert spaces [4, 5].

Relation (3) defines a one-to-one correspondence between the norms $\|\cdot\|_{E}^{G}$ and $\|\cdot\|_{E}^{G}$ as functions on \mathbb{R}_{+} , which is of interest for both theoretical considerations and physical applications (see [2], the example after Theorem 3).

References

- G.G.Magaril-Il'yaev, V.M.Tikhomirov, "Convex Analysis: Theory and Applications", Am. Math. Soc. Providence, R.I., 2003.
- M. E. Shirokov, "Operator E-norms and their use", Sb. Math., 211:9 (2020), 1323–1353; arXiv:1806.05668.
- [3] A.S.Holevo, M.E.Shirokov, "Energy-constrained diamond norms and quantum dynamical semigroups", Lobachevskii J. Math., 40:10 (2019), 1569–1586; arXiv:1812.07447.
- [4] T.Kato, "Perturbation Theory for Linear Operators". Springer-Verlag, New York-Heidelberg-Berlin, 1980.

[5] B.Simon, "Operator Theory: A Comprehensive Course in Analysis, Part IV", American Mathematical Society, 2015.