

# Multiscalar-metric gravity: cosmological constant screening and emergence of massive-graviton dark components of the Universe

Yury F. Pirogov\*

Theory Division, Logunov Institute for High Energy Physics of NRC “Kurchatov Institute”,  
 Protvino, 142281 Moscow Region, Russia

## Abstract

In the multiscalar-metric frameworks, the issues of the vacuum energy/cosmological constant (CC) screening due to the Weyl-scale enhancement of the Diff gauge symmetry, along with emergence of the massive dark gravity components through the gravitational Higgs mechanism are considered. A generic dark gravity model is developed, with the two extreme versions of the model of particular interest based on General Relativity (GR) and its classically equivalent Weyl transverse alternative being compared and argued to be, generally, inequivalent. The so constructed spontaneously broken Weyl Transverse Relativity (WTR) is proposed as a viable beyond-GR effective field theory of gravity, with screening of the Lagrangian CC, superseded by the induced one, and emergence of the massive tensor and scalar gravitons as the dark gravity components. A basic concept with the spontaneously broken Diff gauge symmetry/relativity – in particular, WTR vs. GR – as a principle source of the emergent dark gravity components of the Universe is put forward.

**Keywords:** modified gravity, Weyl transverse relativity, Higgs mechanism for gravity, vacuum energy, cosmological constant, dark energy, dark matter.

## 1 Introduction

The issue of the vacuum energy/cosmological constant (CC) in General Relativity (GR) due to the so-called zero-point quantum fluctuations is well-known already for a long time.<sup>1</sup> But only recently, in the wake of the appearance of the convincing observational evidences for (an effective) CC responsible for the late-time accelerated expansion of the Universe, as expressed by the cosmological Standard Model (SM), or, otherwise, the  $\Lambda$ CDM model, the problem of CC (or, more generally, the vacuum energy) seems to become the most crucial one in the realm of the contemporary fundamental physics.<sup>2</sup> Though not meaning any explicit discrepancy with observations, this problem signifies nevertheless at least a tension between the two present-day basic physical theories: the theory of gravity – GR and the theory of matter and its interactions — the quantum field theory (QFT) — expressed more particularly by the particle SM. Namely, in the effective field theory (EFT) framework the CC problem is, in fact, at least threefold:<sup>3</sup>

(I) *The (un)naturalness problem:* What forbids the huge Lagrangian CC, expected in the EFT framework, from the explicit manifestations, with a non-zero tiny observational CC being (technically) “unnatural”?

(II) *The coincidence problem:* What defines such a tiny value of the observational CC, so that the latter manifests itself in the accelerated expansion of the Universe just “lately” on the cosmological time scale?

(III) *The quantum stability problem:* What prevents the tiny observational CC from being drastically renormalized by the radiative corrections, a priori expected in the EFT framework?

---

\*E-mail: pirogov@ihep.ru

<sup>1</sup>For an original treatment of the vacuum energy/CC problem, see, [1].

<sup>2</sup>For the observational status of CC, see, e.g., [2]. For the modern view of the vacuum energy/CC problem, see, e.g., [3, 4].

<sup>3</sup>For the importance of the EFT approach to the CC problem, cf, e.g., [5].

Solving these and related problems presents a challenge to the modern fundamental physics and could ultimately imply either a revision of QFT, or a GR modification, with a variety of the latter ones proposed to this end up to now.<sup>4</sup> More particularly, to solve the CC problem the modified gravity should, conceivably, be treated as EFT valid at the relatively low energy scales (compared to the fundamental one given by the Planck mass). At that, GR as EFT is well-known, first, to be constructed from a symmetric second-rank tensor field – the metric  $g_{\mu\nu}$ , with its determinant  $g$  defining a four-volume element  $\sqrt{-g}d^4x$  (in the spacetime dimension  $n = 4$ ), with the spacetime measure  $\sqrt{-g}$ , and, second, to be based on the gauge symmetry of the (full) diffeomorphisms (Diff's) incorporating, in particular, the longitudinal ones. The fact that the spacetime measure  $\sqrt{-g}$  transforms under the longitudinal diffeomorphisms proves to be in GR (and its siblings) an ultimate reason of the emergent CC problem. By this token, we take the issue of  $g$  as a guide for choosing a proper route towards solving the CC problems, with some principle steps leading to this goal shortly indicated below.

(i) *Unimodular relativity.* First of all, long before the CC problem had become so acute, for solving such a problem there was proposed in [11] (with the numerous subsequent elaborations) to substitute GR based on the full Diff symmetry by Unimodular Relativity (UR), known also as Unimodular Gravity, based on the transverse diffeomorphism (TDiff  $\subset$  Diff) symmetry.<sup>5</sup> By construction, instead of the conventional ten-component (at  $n = 4$ ) metric  $g_{\mu\nu}$ , UR/TDiff uses a reduced nine-component one  $\hat{g}_{\mu\nu}$ , with the determinant  $\hat{g} = \kappa$  given by a fixed absolute/non-dynamical scalar density  $\kappa$  (for simplicity, say,  $\kappa = -1$ ).<sup>6</sup> The latter defines simultaneously the spacetime volume element  $\sqrt{-\kappa}d^4x$ . The Lagrangian CC within UR proves to be irrelevant being superseded by an integration constant having yet no clear-cut physical meaning. Due to this, UR remains, in a sense, dynamically incomplete. Nevertheless, solving the first part of the CC problem, UR definitely presents a step in the right direction, though hardly sufficient to completely solve the problem, and implies further elaboration.

(ii) *Weyl transverse relativity.* In the latter respect, in [13] there was proposed (cf. also [14]) a ten-component modification of UR constructed from a relative tensor  $\hat{g}_{\mu\nu} \equiv g_{\mu\nu}/(-g)^{1/4}$  (at  $n = 4$ ), with a fixed determinant (put for definiteness  $\hat{g} = -1$ ), the former being liable to be converted into the true tensor through multiplication by  $(-\kappa)^{1/4}$ . By construction, the respective EFT of gravity — to be called Weyl Transverse Relativity (WTR) — satisfies a gauge symmetry, WTDiff, consisting of TDiff enhanced by the Weyl scale transformations, and provides ultimately an improved, compared to UR, solution of the CC problem.<sup>7</sup> Likewise UR/TDiff, its modification WTR/WTDiff may naturally give a justification of the disappearance of the (huge) Lagrangian CC. At the classical level (under the covariant conservation of the energy-momentum tensor) WTR/WTDiff proves to be equivalent to GR/Diff, with an arbitrary induced CC to be treated as a global degree of freedom. Moreover at the quantum level, WTR/WTDiff could ensure the quantum stability of the residual classical CC against the radiative corrections [17, 18]. Besides, on the field-theoretic side, WTR may be obtained as a viable alternative to GR proceeding in a self-consistent manner, in the complete analogy to GR, from a theory of the precisely two-d.o.f. self-interacting massless graviton [19].<sup>8</sup> Altogether, solving (at least partially) the first and the third parts of the CC problem and possessing a solid field-theoretic status WTR/WTDiff could thus constitute a viable alternative to GR/Diff, realizing one more step in the right direction, with the next one(s) still needed. More particularly, within WTR per se (likewise UR) there appears no indication on a infrared mass scale for the tiny residual CC, with the second part of the CC problem remaining to be resolved, too.

(iii) *Higgs mechanism for gravity.* The looked-for infrared mass scale could naturally be provided by a massive graviton, the latter serving as the putative dark energy (DE) mimicking the observed CC. At that, the original tensor-graviton mass term due to Fierz and Pauli [20] explicitly violates the Diff gauge symmetry, and was later shown to result in the massive-massless perturbative discontinuity [21, 22], to remedy which there was proposed [23] a nonperturbative mechanism for recovering the

<sup>4</sup>For the modifications of gravity, see, e.g., [6]–[8] and, in particular, [9, 10] for the massive gravity.

<sup>5</sup>For the relevance of the TDiff gauge symmetry as a substitute of the full Diff one for the consistent description of the massless tensor graviton, see [12].

<sup>6</sup>The notation in Introduction is in accord with what follows.

<sup>7</sup>For a development of WTR, known also as Weyl Transverse Gravity, cf., e.g., [15, 16].

<sup>8</sup>This fact alone could, conceivably, justify WTR (and its siblings) to be further explored at least on par with GR (and its siblings).

transition. It was also found [24] that such an explicit Diff gauge symmetry violation results in the appearance of a ghost, with a violation of unitarity. Thus, with the massless GR treated as a Diff gauge theory, the consistent ghost-free theory of massive gravity would, conceivably, imply not the explicit violation but the spontaneous breaking of such a gauge symmetry by a counterpart of the Higgs mechanism for gravity. An attempt to approach such a goal, modifying GR through the Higgs mechanism in terms of a quartet of the scalar fields, was undertaken in [25] (see also [26, 27]), in the wake of which there was proposed in [28]–[30] a simple solution to the problem elaborated further in [31, 32]. In this respect, GR augmented by the Higgs mechanism for gravity could naturally provide a required scale for the late exponential expansion of the Universe, associating such a scale with the tiny tensor-graviton mass. The latter, on the other hand, should naturally be treated as a parameter of the theory (given still ad hoc) likewise any other fundamental parameter of Nature. Out of the four (at  $n = 4$ ) scalar fields used in such a Higgs mechanism within GR, the three components can, at a proper choice of the (quasi-)Higgs potential, be absorbed as the additional components of the massive tensor graviton. At that, one more component may be stated to become a ghost independently of the potential, though in a sufficiently high order of the perturbation theory [32]. Besides, in the massive-modified GR there still remains unclear the reason of absence of the (huge) Lagrangian CC, as well as the radiative stability of the Higgs mechanism (in the absence of a protecting symmetry). So, the Higgs mechanism in GR is on its own hardly sufficient to exhaust the CC problem, too.

(iv) *Quartet-metric gravity.* Altogether, merging the above mentioned directions of the gravity modification to reconcile the various parts of the CC problem, with the two principle ingredients of the properly modified gravity — an effective metric  $\hat{g}_{\mu\nu}$ ,  $\hat{g} = \kappa$ , and a (quasi-)Higgs field — to stem from the same source — the scalar quartet — may be obligatory.<sup>9</sup> A new concept should, conceivably, be invoked to this end, with a change of the spacetime paradigm [35, 36]. According to the latter, among the arbitrary kinematic observer’s coordinates there exist some distinct dynamical ones, associated ultimately with the vacuum and given by a quartet (at  $n = 4$ ) of the peculiar (invertible) scalar fields. Such a scalar quartet is to be considered on par with the metric as a basic field variable to construct EFT of the so-called quartet-metric gravity. Introducing in such a framework, on the one hand, a proper effective metric, without adding the (quasi-)Higgs potential for the massive tensor graviton, one can eliminate the (huge) Lagrangian CC as in WTR. On the other hand, introducing the proper (quasi-)Higgs potential without modifying metric, one can produce, as in GR, the massive tensor graviton serving, conceivably, as DE. Superimposing these two routes of the gravity modification within the quartet-modified WTR may, conceivably, solve both the problem of screening of the (huge) Lagrangian CC and emergence of the (tiny) observable one supplemented by the associated dark gravity components – the massive tensor and scalar gravitons. This is a goal of the present paper.

In section 2, the basics of the quartet-metric (dark) gravity are exposed. In section 3, a sufficiently general prototype dark gravity model is constructed to realize the concept of the spontaneously broken Diff gauge symmetry/relativity as a source of the emergent dark gravity components. The two extreme versions of such a generic model of the particular interest corresponding, respectively, to the spontaneous breaking of GR vs. its classically equivalent alternative — WTR — are worked out in sections 4 and 5. These versions are compared in the context of the CC screening and emergence of the dark gravity components through the gravitational Higgs mechanism, and are argued to be, generally, non-equivalent. The consistency of the so constructed spontaneously broken WTR as a theory of the massive tensor and scalar gravitons is demonstrated in the weak-field limit. In Conclusion, the viability of the spontaneously broken WTR as a beyond-GR EFT of gravity solving the CC problem through screening the Lagrangian CC, superseding it by the induced one and supplementing by the massive tensor and scalar gravitons as the dark gravity components is argued. The importance of further developing the basic concept of the spontaneously broken Diff gauge symmetry/relativity — in particular, WTR vs. GR — as a principle reason of emergence of the dark gravity components of the Universe is stressed.<sup>10</sup>

---

<sup>9</sup>This, in turn, ensures one more important ingredient for solving the CC problem – the dynamical non-gravitating spacetime measure  $\sqrt{-\kappa}$ . For the relevance of such a measure built of the so-called “measure fields” within the modified GR still with the conventional metric  $g_{\mu\nu}$ , cf., e.g., [33, 34].

<sup>10</sup>Though adhering primarily to the spacetime dimension  $n = 4$ , not to be formally confined by this value, the term “scalar quartet” valid at  $n = 4$  is substituted in what follows by a more general one — the “multiscalar” — and,

## 2 Multiscalar-metric dark gravity: basics

### 2.1 Multiscalar-extended field set

Under the term “dark gravity” there will be understood a merger of the (conventional) gravity with the emergent dark gravitational components. The EFT of the multiscalar-metric dark gravity may basically be defined through the generally covariant action functional<sup>11</sup>

$$S[g_{\mu\nu}, X^a] = \int \mathcal{L}(g_{\mu\nu}, X^a) d^n x \quad (1)$$

in terms of a Lagrangian density  $\mathcal{L}$  dependent on a tensor field — the basic metric —  $g_{\mu\nu}$  (possessing by an inverse  $g^{-1\mu\nu}$ ) extended through a set of the scalar fields  $X^a$ , with  $a, b, \dots = 0, 1, \dots, n-1$  being the global Lorentz indices and  $n$  a spacetime dimension. The scalar multiplet  $X^a = X^a(x)$  — the multiscalar — is peculiar by the fact of being (in a patch-wise manner) invertible,  $x^\mu = x^\mu(X)$ . This allows to use it as some distinct dynamical spacetime coordinates  $X^\alpha \equiv \delta_a^\alpha X^a(x)$  among the arbitrary kinematic ones  $x^\mu$ , as well as to introduce the respective dynamical frame  $X_\mu^\alpha \equiv \partial_\mu X^\alpha$ ,  $\det(X_\mu^\alpha) \neq 0$ , possessing by an inverse  $X^{-1\alpha}_\mu \equiv X^\mu_\alpha$ . Stress that such a multiscalar is not an auxiliary (non-dynamical) variable introduced just to restore the general covariance, but a faithful dynamical variable on par with the basic metric. Ultimately, the dynamical coordinates  $X^\alpha$  may be treated as those associated with the (dynamical) vacuum. The action  $S$  is moreover postulated to be invariant under the reparametrizations of  $X^a$  given by the Lorentz transformations and the finite constant shifts,  $X^a \rightarrow X^a + C^a$ . By this token,  $X^a$  should, in fact, enter through a derivative term  $\kappa_{\mu\nu}$  constructed as

$$\kappa_{\mu\nu} \equiv \partial_\mu X^a \partial_\nu X^b \eta_{ab}, \quad (2)$$

with  $\eta_{ab}$  being the Lorentz-invariant Minkowski symbol. By default,  $\kappa_{\mu\nu}$ , serving as a (quasi-)metric, possesses by an inverse  $\kappa^{-1\mu\nu}$  (at least patch-wise with the proper matching), with

$$\kappa \equiv \det(\kappa_{\mu\nu}) = (\det(\partial_\mu X^a))^2 \det(\eta_{ab}) < 0 \quad (3)$$

to ensure the non-degeneracy and invertibility of such a (quasi-)metric.<sup>12</sup> In terms of the independent field variables, the multiscalar-metric action, in fact, looks like

$$S[g_{\mu\nu}, X^a] = \int \mathcal{L}(g_{\mu\nu}, \kappa_{\mu\nu}) d^n x. \quad (4)$$

Further, in terms of  $\kappa$  and  $g = \det(g_{\mu\nu}) < 0$  one can define an effective scalar field

$$\sigma = \ln(\sqrt{-g}/\sqrt{-\kappa}), \quad (5)$$

to be called the scalar graviton. By this token, in terms of the fields  $g_{\mu\nu}$  and  $\kappa_{\mu\nu}$  one can introduce the effective ones as a conformal metric  $\bar{g}_{\mu\nu}$  and a tensor field  $\bar{\alpha}^\mu_\nu$  given by a correlator (partial contraction) of the two metrics as follows:

$$\bar{g}_{\mu\nu} \equiv e^{\bar{w}} g_{\mu\nu}, \quad \bar{\alpha}^\mu_\nu \equiv \bar{g}^{\mu\lambda} \kappa_{\lambda\nu}, \quad (6)$$

where  $\bar{w} = \bar{w}(\sigma)$  is a gravitational scale factor,<sup>13</sup>  $\bar{g}^{\mu\nu} \equiv \bar{g}^{-1\mu\nu} = e^{-\bar{w}} g^{-1\mu\nu}$  is an inverse of  $\bar{g}_{\mu\nu}$  and  $\det(\bar{\alpha}^\mu_\nu) = \kappa/\bar{g} = e^{-2(2\bar{w}+\sigma)}$ ,  $\bar{g} \equiv \det(\bar{g}_{\mu\nu}) < 0$ .<sup>14</sup> The tensor field  $\bar{\alpha}^\mu_\nu$  may be treated as a realization of

---

accordingly, “quartet-metric” by “multiscalar-metric”, etc.

<sup>11</sup>Here we start directly from the EFT level. For the justification of EFT of the multiscalar-metric gravity as an affine-Goldstone nonlinear model, see [37].

<sup>12</sup>To this end, we assume the (patch-wise) Lorentz-preservation in the field space of  $X^a$ , up to the physically equivalent affine redefinitions of the fields  $X^a$  and the metric  $\eta_{ab}$ .

<sup>13</sup>The dependence  $\bar{w}(\sigma)$  presents the simplest case. More generally, one could envisage within EFT the dependence of the effective metric scale  $\bar{w}$  on the matter fields, environment, etc, with the ultraviolet and infrared behavior of the effective metric being a priori quite different.

<sup>14</sup>By default, the spacetime indices are now lowered (raised) through the effective metric  $\bar{g}_{\mu\nu}$  ( $\bar{g}^{\mu\nu}$ ) if not stated explicitly otherwise.

the (a priori loose) DE. By means of these effective fields, merging gravity and its dark components, the effective action of the purely multiscalar-metric dark gravity may be presented quite generally as

$$S[g_{\mu\nu}, X^a] = \int \bar{L}(\bar{g}_{\mu\nu}, \bar{\alpha}_\nu^\mu, \sigma) \sqrt{-\bar{g}} d^n x, \quad (7)$$

with  $\bar{L}$  being an effective Lagrangian defining, in the line with  $\bar{w}$ , a particular dark gravity model. Due to not all of the variables in such a form of the effective action being independent, the action is ultimately still a function of the independent basic variables  $g_{\mu\nu}$  and  $X^a$ . At last, by means of the frame  $X_\mu^a \equiv \partial_\mu X^a$  and its inverse  $X_a^\mu$ , the DE tensor  $\bar{\alpha}_\nu^\mu$  may equivalently be converted into the scalar form as  $\bar{H}_b^a \equiv X_\mu^a \bar{\alpha}_\nu^\mu X_b^\nu$ , with  $\det(\bar{\alpha}_\nu^\mu) = \det(\bar{H}_b^a)$ , where

$$\bar{H}_c^a \eta^{cb} \equiv \bar{H}^{ab} = \bar{g}^{\kappa\lambda} \partial_\kappa X^a \partial_\lambda X^b \quad (8)$$

is a modification of the original (quasi-)Higgs field for gravity [28]–[30]. Ultimately, the effective action in the form (7), with the dependence on the DE field  $\bar{\alpha}_\nu^\mu$  expressed through a scalar potential  $\bar{V}_\alpha(\bar{\alpha}_\nu^\mu) = \bar{V}_H(\bar{H}^{ab})$ , is aimed at the spontaneous breaking of the Diff gauge symmetry/relativity, with producing the dark gravity components ultimately due to the multiscalar.

## 2.2 Weyl-scale enhanced Diff gauge symmetry

Besides a set of fields, any EFT is basically characterized by a pattern of its symmetries. For the generic multiscalar-metric dark gravity, the effective action (7) is assumed to be generally covariant and gauge invariant under the (full) Diff symmetry given by a Lie derivative  $D_\xi$ . The latter depends on a (infinitesimal) vector parameter  $\xi^\mu(x)$  and may by default be constructed in terms of a covariant derivative  $\nabla_\lambda(g_{\mu\nu})$  defined by the basic metric  $g_{\mu\nu}$ , subsequently expressed through the effective fields as  $g_{\mu\nu} = e^{-\bar{w}(\sigma)} \bar{g}_{\mu\nu}$ . Under  $D_\xi$ , all the basic fields transform conventionally in accord with their covariant tensor structure still irrespective of their nature. To account for the different origin of the basic fields, consider the (local) Weyl scale transformations distinguishing such fields as follows:<sup>15</sup>

$$\Delta_\zeta g_{\mu\nu} = \zeta g_{\mu\nu}, \quad \Delta_\zeta X^a = 0, \quad (9)$$

where  $\zeta(x)$  is an arbitrary (infinitesimal) scalar parameter, so that<sup>16</sup>

$$\begin{aligned} \Delta_\zeta g &= 4\zeta g, \\ \Delta_\zeta \left( g_{\mu\nu} / (-g)^{1/4} \right) &= 0, \\ \Delta_\zeta \kappa_{\mu\nu} = \Delta_\zeta \kappa &= 0. \end{aligned} \quad (10)$$

For the effective fields, this implies in turn

$$\begin{aligned} \Delta_\zeta \sigma &= 2\zeta, \\ \Delta_\zeta \bar{g}_{\mu\nu} &= \zeta(1 + 2\bar{w}') \bar{g}_{\mu\nu}, \\ \Delta_\zeta \bar{g} &= 4\zeta(1 + 2\bar{w}') \bar{g}, \\ \Delta_\zeta \bar{\alpha}_\nu^\mu &= -\zeta(1 + 2\bar{w}') \bar{\alpha}_\nu^\mu, \end{aligned} \quad (11)$$

where  $\bar{w}' = d\bar{w}/d\sigma$ . Generally, this reflects the explicit Weyl scale violation in the multiscalar-metric dark gravity.<sup>17</sup> In an exceptional case  $\bar{w}' = -1/2$ , one independent component of  $g_{\mu\nu}$ , namely, the determinant  $g$  is missing from  $\bar{g}_{\mu\nu}$  and  $\bar{\alpha}_\nu^\mu$ . At that, the residual dependence of the effective action (7) on  $g$  is mediated only by  $\sigma$ . Under  $g$  being a principle source of the CC problem, such a Weyl scale symmetry, signifying otherwise the  $g$ -independence, is of the particular importance for the CC screening in an exceptional case of a generic dark gravity model to be exposed in what follows.

<sup>15</sup>A priori, the Weyl scale symmetry is assumed to be the specifically metric one, with the conventional matter fields, likewise  $X^a$ , to be inert under the Weyl rescalings, unless stated otherwise.

<sup>16</sup>For definiteness, we restrict consideration here and in what follows by the spacetime dimension  $n = 4$ .

<sup>17</sup>Note though, that the exclusively derivative couplings of  $\sigma$  still admit the residual global Weyl scale symmetry  $g_{\mu\nu} \rightarrow e^{\zeta_0} g_{\mu\nu}$ , with the constant shifts  $\sigma \rightarrow \sigma + 2\zeta_0$ .

### 3 Spontaneous Diff gauge symmetry/relativity breaking

#### 3.1 Generic dark gravity model

To be more particular with the (multiscalar) spontaneous breaking of the Diff gauge symmetry (or, otherwise, relativity) consider a prototype dark gravity model merging tensor gravity and the associated dark gravity components, with the effective Lagrangian in a restricted but still rather general partitioned form

$$\bar{L} = \bar{L}_G(\bar{g}_{\mu\nu}, \bar{\alpha}_\nu^\mu) + \bar{L}_M(\bar{g}_{\mu\nu}, \sigma, \phi_I) + \Delta\bar{L}, \quad (12)$$

consisting of an extended gravity Lagrangian  $\bar{L}_G$ , an extended matter Lagrangian  $\bar{L}_M$  and a rest,  $\Delta\bar{L}$ . By default,  $\bar{L}_G$  may depend a priori arbitrarily on the tensor fields  $\bar{g}_{\mu\nu}$  and  $\bar{\alpha}_\nu^\mu$  (and thus on  $\kappa_{\mu\nu}$ ), while  $\bar{L}_M$  is assumed to be independent of  $\bar{\alpha}_\nu^\mu$  and to depend only minimally on  $\bar{g}_{\mu\nu}$ , but a priori arbitrarily on the scalar graviton  $\sigma$  and the generic matter fields  $\phi_I$ ,  $I = 1, \dots$ . The mixing part  $\Delta\bar{L}$ , dependent, generally, on all the variables, is assumed to be negligible in the main approximation.<sup>18</sup> More particularly, adopt for  $\bar{L}_G$  the partitioned form:

$$\bar{L}_G = \bar{L}_\Lambda + \bar{L}_g(\bar{g}_{\mu\nu}) - \bar{V}_\alpha(\bar{\alpha}_\nu^\mu), \quad (13)$$

where the conventional gravity part,  $\bar{L}_g$ , is taken for simplicity in the GR-like form for  $\bar{g}_{\mu\nu}$ :

$$\bar{L}_g = -\frac{1}{2}M_{\text{Pl}}^2 R(\bar{g}_{\mu\nu}), \quad (14)$$

with  $M_{\text{Pl}} = 1/(8\pi G_N)^{1/2}$  being the Planck mass and  $\bar{R} = R(\bar{g}_{\mu\nu})$  the Ricci scalar curvature.<sup>19</sup> This term is supplemented by a constant one

$$\bar{L}_\Lambda \equiv -M_{\text{Pl}}^2 \bar{\Lambda}, \quad (15)$$

with  $\bar{\Lambda}$  being an effective Lagrangian CC. At last,  $\bar{V}_\alpha$  is a scalar potential dependent on the traces of the products of the DE tensor field  $\bar{\alpha}_\nu^\mu$ :<sup>20</sup>

$$\bar{V}_\alpha = M_{\text{Pl}}^2 \sum \bar{v}_{n_1 n_2 \dots} \left( \text{tr}(\bar{\alpha}_\nu^\mu)^{n_1} \text{tr}(\bar{\alpha}_\nu^\mu)^{n_2} \dots \right), \quad (16)$$

with a priori arbitrary coefficients  $\bar{v}_{n_1 n_2 \dots}$ . The powers  $n_i = 0, 1, 2, \dots$  are to be positive finite for the potential to be analytic near  $\bar{\alpha}_\nu^\mu = 0$ , allowing in such a case a smooth limit to GR. The potential may be normalized as  $\bar{V}_\alpha|_\star = 0$  in a priori arbitrary background point in the unbroken or spontaneously broken phases, say, at  $\bar{\alpha}_{\star\nu}^\mu = 0$  or  $\bar{\alpha}_{\star\nu}^\mu = \delta_\nu^\mu$ , respectively, defining, in turn, the proper normalization of  $\bar{\Lambda}$ . In what follows, we restrict consideration by the simplest quartic potential which proves to be sufficient in the leading approximation (see, section 5). The consistency of the theory in the higher-order approximations may require a further elaboration of the potential including, in particular, the fractional powers of  $\bar{\alpha}_\nu^\mu$ , as well as the negative ones corresponding to  $(\bar{\alpha}_\nu^\mu)^{-1} \equiv \bar{\alpha}^{-1\mu}_\nu = \kappa^{-1\mu\lambda} \bar{g}_{\lambda\nu}$ , etc.<sup>21</sup> Finally, the extended matter Lagrangian  $\bar{L}_M$  (incorporating, conceivably, some kinds of the particle DM in addition to the scalar-graviton dark component) remains a priori arbitrary, elaborating further the purely-tensor dark gravity due to  $\bar{L}_G$ .

#### 3.2 Generic field equations

A straightforward way of deriving the dynamical field equations (FEs) for EFT of the multiscalar-metric dark gravity is to extremize its effective action with respect to the independent variations of the primary metric  $g_{\mu\nu}$  (or, rather, its inverse  $g^{-1\mu\nu}$ ) and the multiscalar  $X^a$ , expressing afterwards

<sup>18</sup>In the spirit of EFT,  $\bar{L}_G$  is assumed to be of the order of the Plank mass squared,  $M_{\text{Pl}}^2$ , while  $\bar{L}_M$  and  $\Delta\bar{L}$  to be relatively suppressed in the scale of  $M_{\text{Pl}}$ .

<sup>19</sup>In principle, any modification of  $\bar{L}_g$ , like  $f(\bar{R})$ , etc, is conceivable, too.

<sup>20</sup>At that,  $\det(\bar{\alpha}_\nu^\mu) = \kappa/\bar{g} = e^{-2(2\bar{w}(\sigma)+\sigma)}$  is explicitly missed being expressed, in principle, through  $\sigma$  accounted in  $\bar{L}_M$ .

<sup>21</sup>A priori, one may admit another mode of the spontaneous braking, with the potential  $\bar{V}_\alpha$  dependent instead of  $\bar{\alpha}_\nu^\mu$  on its inverse  $\bar{\alpha}^{-1\mu}_\nu$ , subject to the analyticity requirement near  $\bar{\alpha}^{-1\mu}_\nu = 0$  and possessing by the flat background at the symmetric point  $\bar{\alpha}^{-1\mu}_\nu|_\star = \bar{\alpha}|_\star^\mu = \delta_\nu^\mu$ .

the coefficients at the independent variations  $\delta g^{-1\mu\nu}$  and  $\delta X^a$  through the effective fields  $\bar{g}_{\mu\nu}$  (and its inverse  $\bar{g}^{\mu\nu}$ ), as well as  $\sigma$  (in particular,  $\bar{w}(\sigma)$ ). By this token, the effective variations look like

$$\begin{aligned}\delta\bar{g}^{\mu\nu} &= e^{-\bar{w}}\delta g^{-1\mu\nu} - \bar{w}'\bar{g}^{\mu\nu}\delta\sigma, \\ \frac{\delta\sqrt{-\bar{g}}}{\sqrt{-\bar{g}}} &= -\frac{1}{2}\bar{g}_{\kappa\lambda}\delta\bar{g}^{\kappa\lambda} = -\frac{1}{2}e^{-\bar{w}}\bar{g}_{\kappa\lambda}\delta g^{-1\kappa\lambda} + 2\bar{w}'\delta\sigma,\end{aligned}\quad (17)$$

with a prime meaning the derivative with respect to  $\sigma$ , as well as

$$\delta\bar{\alpha}_\nu^\mu = \bar{g}^{\mu\lambda}\delta\kappa_{\lambda\nu} + \kappa_{\nu\lambda}\delta\bar{g}^{\lambda\mu}.\quad (18)$$

In the above, one should also account for the constraints (5) and (2) (implying  $\sigma$  and  $\kappa_{\mu\nu}$  to be the effective composite fields), resulting due to

$$\frac{\delta\sqrt{-g}}{\sqrt{-g}} = -\frac{1}{2}e^{-\bar{w}}\bar{g}_{\kappa\lambda}\delta g^{-1\kappa\lambda}, \quad \frac{\delta\sqrt{-\kappa}}{\sqrt{-\kappa}} = \frac{1}{2}\kappa^{-1\kappa\lambda}\delta\kappa_{\kappa\lambda},\quad (19)$$

in the respective variation of  $\sigma$

$$\delta\sigma = \frac{\delta\sqrt{-g}}{\sqrt{-g}} - \frac{\delta\sqrt{-\kappa}}{\sqrt{-\kappa}} = -\frac{1}{2}(e^{-\bar{w}}\bar{g}_{\kappa\lambda}\delta g^{-1\kappa\lambda} + \kappa^{-1\kappa\lambda}\delta\kappa_{\kappa\lambda}),\quad (20)$$

where

$$\delta\kappa_{\mu\nu} = (X_\mu^a\delta X_\nu^b + X_\nu^a\delta X_\mu^b)\eta_{ab}, \quad \delta X_\mu^a \equiv \partial_\mu\delta X^a.\quad (21)$$

Extremizing then the effective action (7), given by the effective Lagrangian (12)–(16), with respect to the independent variations  $\delta g^{-1\mu\nu}$  and  $\delta\kappa_{\mu\nu}$  (the latter expressed finally through  $\delta X^a$ ), we get the generic (under an arbitrary  $\bar{w}$ ) FEs:

$$\begin{aligned}\bar{R}_{\mu\nu} - \frac{1}{4}\bar{R}\bar{g}_{\mu\nu} - \frac{1}{M_{\text{Pl}}^2}(\bar{T}_{\mu\nu} - \frac{1}{4}\bar{T}\bar{g}_{\mu\nu}) &= 0, \\ -\frac{1}{4}(1 + 2\bar{w}')(\bar{R} + 4\bar{\Lambda} + \frac{1}{M_{\text{Pl}}^2}\bar{T}) + \frac{1}{M_{\text{Pl}}^2}\frac{\delta\bar{L}_M}{\delta\sigma} &= 0, \\ \bar{\nabla}_\lambda\left(\frac{1}{M_{\text{Pl}}^2}\bar{g}^{\lambda\rho}\frac{\partial\bar{V}_{\bar{\alpha}}}{\partial\bar{\alpha}_\kappa^\rho}X_{\kappa a} + \frac{1}{8}(\bar{R} + 4\bar{\Lambda} + \frac{1}{M_{\text{Pl}}^2}\bar{T})X_a^\lambda\right) &= 0, \\ \frac{\delta\bar{L}_M}{\delta\phi_I} &= 0\end{aligned}\quad (22)$$

for the three sectors of the multiscalar-metric dark gravity — respectively, tensor-traceless, tensor-trace and multiscalar — supplemented by FE for matter. Here and in what follows,  $\partial/\partial$  and  $\delta/\delta$  mean, respectively, the partial and total (incorporating the derivatives with respect to the derivatives of the proper fields) variational derivatives.<sup>22</sup> Besides,  $X_{\kappa a} \equiv X_\kappa^b\eta_{ba}$  and

$$\bar{T}_{\mu\nu} \equiv \bar{T}_{\bar{\alpha}\mu\nu} + \bar{T}_{M\mu\nu}, \quad \bar{T} \equiv \bar{g}^{\kappa\lambda}\bar{T}_{\kappa\lambda} = \bar{T}_{\bar{\alpha}} + \bar{T}_M,\quad (23)$$

is the total energy-momentum tensor for the DE field  $\bar{\alpha}_\nu^\mu$  and the extended matter  $M = (\sigma, \phi_I)$ , with the partial contributions, respectively, as follows:

$$\begin{aligned}\bar{T}_{\bar{\alpha}\mu\nu} &\equiv -\left(\frac{\partial\bar{V}_{\bar{\alpha}}}{\partial\bar{\alpha}_\lambda^\mu}\bar{g}_{\rho\nu} + \frac{\partial\bar{V}_{\bar{\alpha}}}{\partial\bar{\alpha}_\lambda^\nu}\bar{g}_{\rho\mu}\right)\bar{\alpha}_\lambda^\rho + \bar{V}_{\bar{\alpha}}\bar{g}_{\mu\nu}, \\ \bar{T}_{M\mu\nu} &\equiv \frac{2}{\sqrt{-\bar{g}}}\frac{\partial(\sqrt{-\bar{g}}\bar{L}_M)}{\partial\bar{g}^{\mu\nu}} = 2\frac{\partial\bar{L}_M}{\partial\bar{g}^{\mu\nu}} - \bar{L}_M\bar{g}_{\mu\nu}.\end{aligned}\quad (24)$$

A priori, the derived generic FEs do not automatically imply the covariant conservation of the total energy-momentum tensor, to be discussed below.

<sup>22</sup>At that, the scalar-graviton FE as such is absent, with the composite  $\sigma$  being, generally, off-mass-shell,  $\delta\bar{L}_M/\delta\sigma \neq 0$ .

### 3.3 Geometry consistency condition

To explicitly account for the Riemannian structure of the spacetime in terms of the effective metric  $\bar{g}_{\mu\nu}$ , apply the proper covariant derivative  $\bar{\nabla}_\mu$  to the sum of the trace-free and trace (times  $\bar{g}_{\mu\nu}$ ) parts of the tensor-gravity FEs. Due to the reduced Bianchi identity,  $\bar{\nabla}^\lambda(\bar{R}_{\lambda\nu} - 1/2 \bar{R}\bar{g}_{\lambda\nu}) = 0$ , this gives a geometrical consistency condition

$$-\frac{1}{2}\partial_\mu\left(\bar{w}'(\bar{R} + 4\bar{\Lambda} + \frac{1}{M_{\text{Pl}}^2}\bar{T})\right) = \frac{1}{M_{\text{Pl}}^2}\bar{\nabla}^\lambda\bar{\Theta}_{\lambda\mu}, \quad (25)$$

to be satisfied for all the solutions to the multiscalar-metric FEs. In the above, the tensor

$$\bar{\Theta}_{\mu\nu} \equiv \bar{T}_{\mu\nu} + \bar{\theta}_{s\mu\nu} = \bar{T}_{\mathfrak{e}\mu\nu} + \bar{T}_{M\mu\nu} + \bar{\theta}_{s\mu\nu} \quad (26)$$

is an extension of the total energy-momentum tensor accounting also for the off-mass-shell contribution

$$\bar{\theta}_{s\mu\nu} \equiv -\delta\bar{L}_M/\delta\sigma\bar{g}_{\mu\nu} \quad (27)$$

of the (composite) scalar graviton  $\sigma$ .<sup>23</sup> A priori,  $\bar{\Theta}_{\mu\nu}$  is not bound to covariantly conserve, likewise any its part (26) separately, in particular,  $\bar{\theta}_{s\mu\nu}$ . The covariant conservation of  $\bar{\Theta}_{\mu\nu}$  can still be imposed at an arbitrary  $\bar{w}$  as an additional restriction on the solutions. The two extreme versions of the so constructed generic dark gravity model, with  $\bar{w}' = 0$  and  $-1/2$ , presenting the special interest — respectively, as the spontaneously broken GR vs. WTR — are worked out in more detail below.

### 3.4 Linear approximation

To illustrate the spontaneous breaking of the Diff gauge symmetry/relativity expand the basic fields  $g_{\mu\nu}$  and  $X^a$  on the background ones  $g_{\star\mu\nu}$  and  $X_\star^a$  supplemented by some perturbations:

$$g_{\mu\nu} \equiv g_{\star\mu\nu}(x) + h_{\mu\nu}(x), \quad X^a \equiv X_\star^a(x) + \chi^a(x). \quad (28)$$

Choose then the proper background coordinates  $x_\star^\alpha \equiv \delta_\alpha^\alpha X_\star^\alpha(x)$ , with the flat background looking in such coordinates as follows:

$$g_{\star\alpha\beta} = \bar{g}_{\star\alpha\beta} = \kappa_{\star\alpha\beta} = \eta_{\alpha\beta}, \quad \bar{\mathfrak{a}}_\star^{\pm 1\alpha} = \delta_\beta^\alpha, \quad \sigma_\star = 0, \quad (29)$$

where  $\eta_{\alpha\beta}$  is the Minkowski symbol. Identifying the global Lorentz indices  $a, b, \dots$  with the flat background ones  $\alpha, \beta, \dots$  results then for the basic fields in the linear approximation (LA) in the relations:

$$\begin{aligned} g_{\alpha\beta}^{\pm 1} &= \eta_{\alpha\beta} \pm h_{\alpha\beta}(x_\star), \\ X_\alpha^{\pm 1\beta} &= \delta_\alpha^\beta \pm \partial_\alpha\chi^\beta(x_\star), \\ \kappa_{\alpha\beta}^{\pm 1} &= \eta_{\alpha\beta} \pm k_{\alpha\beta}(x_\star), \\ k_{\alpha\beta} &\equiv \partial_\alpha\chi_\beta(x_\star) + \partial_\beta\chi_\alpha(x_\star), \end{aligned} \quad (30)$$

with  $\partial_\alpha \equiv \partial/\partial x_\star^\alpha$  and  $\chi_\alpha \equiv \eta_{\alpha\beta}\chi^\beta$ , etc. Here and in what follows, the tensors are explicitly marked vs. their inverse values by the superscripts  $\pm 1$ . For the effective fields, this in turn implies in LA:

$$\begin{aligned} \bar{g}_{\alpha\beta}^{\pm 1} &= \eta_{\alpha\beta} \pm \bar{h}_{\alpha\beta}, \\ \bar{\mathfrak{a}}_\beta^{\pm 1\alpha} &= \delta_\beta^\alpha \mp \bar{f}_\beta^\alpha \end{aligned} \quad (31)$$

and  $\sigma = s$ . Designating in LA  $\bar{w} = \bar{\alpha}\sigma$ , with  $\bar{\alpha}$  a constant interpolating between  $\bar{\alpha} = 0$  for the modified GR and  $\bar{\alpha} = -1/2$  for the modified WTR, we get in the same approximation:

$$\begin{aligned} \bar{h}_{\alpha\beta} &\equiv h_{\alpha\beta} + \bar{\alpha}/2(h - 2\partial\chi)\eta_{\alpha\beta}, \\ \bar{f}_{\alpha\beta} \equiv \eta_{\alpha\gamma}\bar{f}_\beta^\gamma &= \bar{h}_{\alpha\beta} - k_{\alpha\beta}, \quad \bar{f} = (1 + 2\bar{\alpha})(h - 2\partial\chi), \\ s &= 1/2(h - 2\partial\chi), \end{aligned} \quad (32)$$

<sup>23</sup>Note that the value  $\bar{\theta}_{s\mu\nu} \neq 0$  additionally closes or opens a spacetime region, i.e., acts in this region as an effective dark matter (DM) or DE, depending on  $\delta\bar{L}_M/\delta\sigma$  being positive or negative, respectively.

with  $\partial\chi \equiv \partial_\gamma\chi^\gamma$ ,  $h = h_{\alpha\beta}\eta^{\alpha\beta}$ ,  $\bar{f} = \bar{f}_{\alpha\beta}\eta^{\alpha\beta}$ , etc. By construction, under the Diff gauge symmetry enhanced by the Weyl rescalings, for the basic fields there takes place:

$$\begin{aligned}\chi^\alpha &\rightarrow \chi^\alpha + \xi^\alpha, \\ k_{\alpha\beta} &\rightarrow k_{\alpha\beta} + (\partial_\alpha\xi_\beta + \partial_\beta\xi_\alpha) = \partial_\alpha(\chi_\beta + \xi_\beta) + \partial_\beta(\chi_\alpha + \xi_\alpha), \\ h_{\alpha\beta} &\rightarrow h_{\alpha\beta} + (\partial_\alpha\xi_\beta + \partial_\beta\xi_\alpha) + \zeta\eta_{\alpha\beta}, \\ h &\rightarrow h + 2\partial\xi + 4\zeta,\end{aligned}\tag{33}$$

with the arbitrary functions  $\xi^\alpha(x_\star)$  and  $\zeta(x_\star)$  as the gauge parameters. For the effective fields, this results in LA in transformations:

$$\begin{aligned}\bar{h}_{\alpha\beta} &\rightarrow \bar{h}_{\alpha\beta} + (\partial_\alpha\xi_\beta + \partial_\beta\xi_\alpha) + (1 + 2\bar{\alpha})\zeta\eta_{\alpha\beta}, \\ \bar{f}_{\alpha\beta} &\rightarrow \bar{f}_{\alpha\beta} + (1 + 2\bar{\alpha})\zeta\eta_{\alpha\beta}, \\ s &\rightarrow s + 2\zeta.\end{aligned}\tag{34}$$

Otherwise,  $\bar{h}_{\alpha\beta}$  is a tensor under Diff, coinciding in the ‘‘unitary’’ gauge  $\chi^\alpha = 0$  (and thus  $k_{\alpha\beta} = 0$ ) according to (32) with the Diff gauge-invariant  $\bar{f}_{\alpha\beta}$ , whereas  $s$  is a scalar under Diff. Hence, the fields  $\bar{f}_{\alpha\beta}$  and  $s$  may be used to describe the spontaneous Diff gauge symmetry/relativity breaking in the dark gravity in the explicitly Diff gauge-invariant terms. At that,  $\bar{f}_{\alpha\beta}$  at  $\bar{\alpha} \neq -1/2$  and  $s$  explicitly violate the Weyl-scale gauge symmetry, with  $s$  serving, in effect, as a Weyl scale in disguise.<sup>24</sup>

## 4 Spontaneously broken General Relativity

Let first  $\bar{w}' = 0$ , so that without loss of generality  $\bar{w} = w = 0$  and  $\bar{g}_{\mu\nu} = g_{\mu\nu}$ , with the bar-sign to be dropped-off everywhere. In the Lagrangian  $L = L_G + L_M$ , both  $L_G$  and  $L_M$  are Diff gauge invariant, with the explicitly violated Weyl scale symmetry. The FEs (22) can now be combined as follows:

$$\begin{aligned}R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} - \Lambda g_{\mu\nu} - \frac{1}{M_{\text{Pl}}^2}\Theta_{\mu\nu} &= 0, \\ \nabla_\lambda\left(g^{\lambda\rho}\frac{\partial V_{\text{ae}}}{\partial\text{ae}^\rho}X_{\kappa a} + \frac{1}{2}\frac{\delta L_M}{\delta\sigma}X_a^\lambda\right) &= 0, \\ \frac{\delta L_M}{\delta\phi_I} &= 0,\end{aligned}\tag{35}$$

with the scalar-graviton FE  $\delta L_M/\delta\sigma = 0$  as such being, generally, missing. Due to the reduced Bianchi identity, the extended total energy-momentum tensor  $\Theta_{\mu\nu}$  should for the geometrical consistency be covariantly conserved for all the solutions to FEs,  $\nabla^\lambda\Theta_{\lambda\nu} \equiv 0$ , though separately each of its three contributions to the total sum (26) may, generally, not satisfy this requirement. In general, FEs (35) correspond to the multiscalar-modified/spontaneously broken GR, with the Lagrangian CC  $\Lambda$  supplemented by the emergent tensor- and scalar-graviton dark components contributing through  $\Theta_{\mu\nu}$ . Under the conventional FE for the scalar graviton  $\sigma$ ,  $\delta L_M/\delta\sigma = 0$ , and the absence of the tensor DE,  $V_{\text{ae}} \equiv 0$ , the multiscalar part of FEs (35) is clearly trivial, with the tensor FE reducing due to  $T_{\text{ae}\mu\nu} = 0$  to that of the unbroken GR supplemented by a scalar field  $\sigma$ .

In LA at  $\bar{\alpha} = \alpha = 0$ , one has  $\bar{h}_{\alpha\beta} = h_{\alpha\beta}$ , with the tensor and scalar dark gravity fields

$$\begin{aligned}f_{\alpha\beta} &= h_{\alpha\beta} - k_{\alpha\beta}, \quad f = h - 2\partial\chi, \\ s &= 1/2(h - 2\partial\chi)\end{aligned}\tag{36}$$

transforming under the enhanced Diff gauge symmetry as follows:

$$\begin{aligned}h_{\alpha\beta} &\rightarrow h_{\alpha\beta} + (\partial_\alpha\xi_\beta + \partial_\beta\xi_\alpha) + \zeta\eta_{\alpha\beta}, \\ f_{\alpha\beta} &\rightarrow f_{\alpha\beta} + \zeta\eta_{\alpha\beta}, \quad s \rightarrow s + 2\zeta,\end{aligned}\tag{37}$$

<sup>24</sup>To be more concise, the scalar graviton being a measure of the (Weyl) scale transformations may be called the *systolon*, with the tensor graviton being conventionally the graviton.

resulting in the Weyl scale symmetry to be explicitly violated. At that, the trace metric part  $h$  enters both the Diff gauge-invariant tensor field  $f_{\alpha\beta}$  and scalar  $s$ , and, generally, results in a ghost. In the case with the missing  $\sigma$ ,  $\delta L_M/\delta\sigma \equiv 0$ , under the proper choice of the potential  $V_{\mathfrak{a}}$ , there appears in the broken phase the massive tensor graviton, with  $h$  as a ghost missing. But the ghost inevitably reappears in a sufficiently high order of the perturbation theory [28]–[32]. This may, conceivably, be traced back to the absence of a protecting symmetry. On the other hand, in the reduced case  $V_{\mathfrak{a}} = 0$  the tensor graviton remains massless, with  $h$  in the transverse gauge  $\partial\chi = 0$  becoming the physical massive scalar graviton.<sup>25</sup> The general tensor-scalar mixed case beyond LA, with both  $V_{\mathfrak{a}} \neq 0$  and  $\delta L_M/\delta\sigma \neq 0$ , is to be investigated.

Altogether, though being able to produce the dark gravity components, the spontaneously broken GR at its face value can hardly help in solving the entire CC problem, with the latter still to be resolved by some other means. Nevertheless, an alternative to GR classically equivalent to the latter up to CC — WTR — proves a priori to be well suited to this end, as is discussed in more detail below, with the case of GR reserved as the canonical reference one.

## 5 Spontaneously broken Weyl Transverse Relativity

### 5.1 Full nonlinear theory

#### 5.1.1 General field equations

Let then  $\bar{w}' = -1/2$ , with the effective gravity scale factor  $\bar{w} = \hat{w} \equiv -\sigma/2$ . Replacing in this case everywhere the bar-sign by the hat, one has for the effective metric

$$\hat{g}_{\mu\nu} = e^{-\sigma/2} g_{\mu\nu} = (\kappa/g)^{1/4} g_{\mu\nu}, \quad \hat{g} = \kappa, \quad (38)$$

with  $\det(\hat{\mathfrak{a}}_{\nu}^{\mu}) = \kappa/\hat{g} = 1$ . In the effective Lagrangian  $\hat{L} = \hat{L}_G + \hat{L}_M$ , the purely tensor-gravity part  $\hat{L}_G$  possesses by the Diff gauge symmetry enhanced by the “hidden” local Weyl scale symmetry. Due to this, one independent component in  $\hat{L}_G$  is, in fact, missing. With account for  $\hat{L}_M$ , the missing component reappears again as  $\sigma$  signifying the explicit violation of the local Weyl scale symmetry. The FEs (22) now read as follows:

$$\begin{aligned} \hat{R}_{\mu\nu} - \frac{1}{4}\hat{R}\hat{g}_{\mu\nu} - \frac{1}{M_{\text{Pl}}^2}(\hat{T}_{\mu\nu} - \frac{1}{4}\hat{T}\hat{g}_{\mu\nu}) &= 0, \\ \hat{\nabla}_{\lambda}\left(\frac{1}{M_{\text{Pl}}^2}\hat{g}^{\lambda\rho}\frac{\partial\hat{V}_{\mathfrak{a}}}{\partial\hat{\mathfrak{a}}_{\kappa}^{\rho}}X_{\kappa a} + \frac{1}{8}(\hat{R} + 4\hat{\Lambda} + \frac{1}{M_{\text{Pl}}^2}\hat{T})X_a^{\lambda}\right) &= 0, \\ \frac{\delta\hat{L}_M}{\delta\sigma} = \frac{\delta\hat{L}_M}{\delta\phi_I} &= 0, \end{aligned} \quad (39)$$

with  $\hat{T}_{\mu\nu} = \hat{T}_{\mathfrak{a}\mu\nu} + \hat{T}_{M\mu\nu}$  given by eq. (24). The tensor-gravity FE (39) is trace-free, being principle in the context of the CC problem.<sup>26</sup> In the multiscalar-metric framework, such a FE alone is, in fact, incomplete, to be supplemented by the multiscalar one. At that, the Lagrangian CC  $\hat{\Lambda}$  clearly influences the tensor gravity just indirectly through the multiscalar FE. A crucial point here is that the dynamical measure  $\sqrt{-\kappa}$ , becoming, in a sense, non-gravitating, depends entirely on the multiscalar  $X^a$ , but not on the basic metric  $g_{\mu\nu}$ . The scalar graviton behaves now likewise a conventional scalar particle, with  $\hat{\theta}_{s\mu\nu} \equiv -\delta\hat{L}_M/\delta\sigma \hat{g}_{\mu\nu} = 0$ . Due to the general covariance and the conventional FEs for  $\sigma$  and  $\phi_I$ , the extended matter energy-momentum tensor  $\hat{T}_{M\mu\nu}$  should be covariantly conserved,  $\hat{\nabla}^{\lambda}\hat{T}_{M\lambda\nu} = 0$ , whereas the DE  $\hat{T}_{\mathfrak{a}\mu\nu}$  and thus the total  $\hat{T}_{\mu\nu}$  may, generally, not satisfy this requirement.<sup>27</sup>

<sup>25</sup>For the reduced case  $V_{\mathfrak{a}} \equiv 0$ , with the (ultralight) physical  $\sigma$  serving as a scalar DE superseding the Lagrangian CC, see [38]. For the similar case, with  $\sigma$  as a scalar DM, cf. [39].

<sup>26</sup>For the viability of the trace-free tensor-gravity FE in cosmology, cf., e.g., [40, 41].

<sup>27</sup>In addition to the scalar graviton  $\sigma$ , the extended matter Lagrangian  $\hat{L}_M = \hat{L}_M(\hat{g}_{\mu\nu}, \sigma, \phi_I)$  could, in principle, account for the (more conventional) DM components. In the absence of matter, the purely scalar-graviton Lagrangian  $\hat{L}_M = \hat{L}_s(\hat{g}_{\mu\nu}, \sigma)$  could be looked-for, e.g., in a most general form for the scalar field  $\sigma$  satisfying the second-order FE to avoid the explicit Ostrogradsky instabilities [42, 43].

### 5.1.2 Covariant conservation condition

Impose additionally on a class of the solutions to FEs (39) the condition of the covariant conservation of the total energy-momentum tensor  $\hat{\Theta}_{\mu\nu} = \hat{T}_{\mu\nu}$  in eq. (25), resulting upon integration in the relation

$$\hat{R} + 4\hat{\Lambda}_0 + \frac{1}{M_{\text{Pl}}^2}\hat{T} = 0, \quad (40)$$

where  $\hat{\Lambda}_0$  is an integration constant, in particular,  $\hat{\Lambda}_0 = 0$  or the Lagrangian CC  $\hat{\Lambda}$ . Incorporating this relation refines, in turn, the tensor-gravity and multiscalar FEs as follow:

$$\begin{aligned} \hat{R}_{\mu\nu} - \frac{1}{2}\hat{R}\hat{g}_{\mu\nu} - \hat{\Lambda}_0\hat{g}_{\mu\nu} - \frac{1}{M_{\text{Pl}}^2}\hat{T}_{\mu\nu} &= 0, \\ \hat{\nabla}_\lambda \left( \frac{1}{M_{\text{Pl}}^2}\hat{g}^{\lambda\rho}\frac{\partial\hat{V}_{\mathfrak{a}}}{\partial\hat{x}^\rho}X_{\kappa\alpha} + \frac{1}{2}(\hat{\Lambda} - \hat{\Lambda}_0)X_a^\lambda \right) &= 0, \\ \frac{\delta\hat{L}_M}{\delta\sigma} = \frac{\delta\hat{L}_M}{\delta\phi_I} &= 0, \end{aligned} \quad (41)$$

with the Lagrangian CC  $\hat{\Lambda}$  clearly screened from the tensor-gravity FE, and  $\hat{T}_{\mu\nu}$  explicitly covariantly conserved due to the reduced Bianchi identity. Moreover, due to  $\hat{\nabla}^\lambda\hat{T}_{M\lambda\nu} = 0$  there should now separately fulfill  $\hat{\nabla}^\lambda\hat{T}_{\mathfrak{a}\lambda\nu} = 0$ . Under an arbitrary  $\hat{V}_{\mathfrak{a}}$ , FEs (41) describe the multiscalar-modified/spontaneously broken WTR, with DE manifesting itself through the covariantly conserved  $\hat{T}_{\mathfrak{a}\mu\nu}$ , as well as the induced and Lagrangian CCs  $\hat{\Lambda}_0$  and  $\hat{\Lambda}$ , respectively. In the case  $\hat{V}_{\mathfrak{a}} \equiv 0$ , the multiscalar part of FEs (41) factorizes due to  $\hat{g} = \kappa$  as  $\partial_\lambda(\sqrt{-\kappa}X_a^\lambda) = 0$ . By this token,  $\kappa$  can be chosen arbitrary, looking formally as non-dynamical (similarly to UR). With account for  $\hat{T}_{\mathfrak{a}\mu\nu} = 0$ , FEs in this case reduce to those of UR, equivalent classically to GR up to CC, with the matter set (containing, conceivably, DM) extended through the scalar graviton  $\sigma$  as a conventional scalar particle.<sup>28</sup>

## 5.2 Weak-field limit

### 5.2.1 Linear approximation

At  $\bar{\alpha} = \hat{\alpha} = -1/2$ , in the flat background one has in LA

$$\begin{aligned} \hat{h}_{\alpha\beta} &\equiv h_{\alpha\beta} - 1/4(h - 2\partial\chi)\eta_{\alpha\beta}, \quad \hat{h} = 2\partial\chi, \\ \hat{f}_{\alpha\beta} &= \hat{h}_{\alpha\beta} - k_{\alpha\beta}, \quad \hat{f} = 0, \\ s &= 1/2(h - 2\partial\chi), \end{aligned} \quad (42)$$

Under the Weyl-scale enhanced Diff gauge transformations one gets:

$$\begin{aligned} \hat{h}_{\alpha\beta} &\rightarrow \hat{h}_{\alpha\beta} + (\partial_\alpha\xi_\beta + \partial_\beta\xi_\alpha), \\ \hat{f}_{\alpha\beta} &\rightarrow \hat{f}_{\alpha\beta}, \quad s \rightarrow s + 2\zeta. \end{aligned} \quad (43)$$

This means that  $\hat{h}_{\alpha\beta}$ , being a tensor under Diff, is Weyl-scale invariant,  $\hat{f}_{\alpha\beta}$  is completely invariant under the Weyl-scale enhanced Diff gauge symmetry, while  $s$ , being a scalar under Diff, transforms inhomogeneously under the Weyl rescalings. Hence, the fields  $\hat{f}_{\alpha\beta}$  and  $s$  may be used to describe the spontaneously broken Diff gauge symmetry in the explicitly Diff-invariant terms realizing the unitary gauge ( $\chi^\alpha = 0$ ,  $\hat{h}_{\alpha\beta} = \hat{f}_{\alpha\beta}$ ) for the dark gravity. At that, one independent DE component in the tensor  $\hat{f}_{\alpha\beta}$  is missing due to  $\hat{f} = 0$  as a manifestation of the Weyl scale symmetry. In the end, such a component reappears again through the scalar graviton  $\sigma$  as a conventional particle, reflecting ultimately the explicit local Weyl scale violation.<sup>29</sup> The account beyond LA for the higher orders in the presence of the protecting Weyl scale symmetry, as well as the local Weyl-scale violation due to  $\sigma$ , are to be investigated.

<sup>28</sup>Otherwise, the multiscalar-modified UR could be obtained as a reduction of the multiscalar-modified WTR under imposing ab initio the constraint  $g = \kappa$ , consistent with  $\hat{g} = \kappa$ . This would imply, in particular, the explicit absence of the scalar graviton  $\sigma$ , which is to be added ad hock as an additional scalar particle.

<sup>29</sup>Note that imposing ab initio the UR restriction  $g = \kappa$  eliminating, in fact, one independent component would result in LA in  $h = \hat{h} = 2\partial\chi$  (still preserving  $\hat{f} = 0$ ) and  $\hat{h}_{\alpha\beta} = h_{\alpha\beta}$ , as well as  $s = 0$ , with the lost of the Weyl scale symmetry.

### 5.2.2 Massive tensor graviton

To clarify the viability of the multiscalar-modified/spontaneously broken WTR as a theory of gravitons retain first only the Weyl-scale invariant purely tensor-gravity Lagrangian  $\hat{L}_G$  putting  $\hat{L}_M(\sigma, \phi_I) = 0$  in neglect by matter and the explicit scalar-graviton contribution. Choosing then as independent variables for the DE tensor field  $\hat{\mathfrak{a}}_\nu^\mu$  its trace  $\hat{\mathfrak{a}}$  and the traceless part  $\hat{\hat{\mathfrak{a}}}_\nu^\mu$ , respectively, as

$$\hat{\mathfrak{a}} \equiv \hat{\mathfrak{a}}_\lambda^\kappa \delta_\kappa^\lambda, \quad \hat{\hat{\mathfrak{a}}}_\nu^\mu \equiv \hat{\mathfrak{a}}_\nu^\mu - \frac{1}{4} \hat{\mathfrak{a}} \delta_\nu^\mu, \quad (44)$$

consider the even quartic potential  $\hat{V}_{\mathfrak{a}}(\hat{\hat{\mathfrak{a}}}_\nu^\mu, \hat{\mathfrak{a}})$  as follows:

$$\hat{V}_{\mathfrak{a}} = \frac{1}{2} M_{\text{Pl}}^2 \hat{v}_2 \left( \hat{\hat{\mathfrak{a}}}_\nu^\mu \hat{\hat{\mathfrak{a}}}_\mu^\nu + \hat{\beta}_4 \left( \left( \frac{1}{4} \hat{\mathfrak{a}} \right)^2 - 1 \right)^2 \right), \quad (45)$$

where  $\hat{v}_2 > 0$  and  $\hat{\beta}_4$  are some constant parameters.<sup>30</sup> For definiteness,  $\hat{V}_{\mathfrak{a}}$  is normalized as  $\hat{V}_{\mathfrak{a}}|_\star = 0$  in the flat background at  $\hat{\mathfrak{a}}_\star^\mu = \delta_\nu^\mu$ ,  $\hat{\mathfrak{a}}_\star = 4$ , with an additional constant contribution assigned by default to the Lagrangian CC  $\hat{\Lambda}$ . Such a potential is peculiar by possessing by the two extremums under  $\hat{\mathfrak{a}}$ : at  $\hat{\mathfrak{a}} = 0$  and  $|\hat{\mathfrak{a}}| = 4$ , which may be ascribed at  $\hat{\beta}_4 > 0$  to the unbroken trivial and broken flat backgrounds, respectively. In the trivial background at  $\hat{\mathfrak{a}}_\star^\mu = 0$ ,  $\hat{\mathfrak{a}}_\star = 0$ , the potential  $\hat{V}_{\mathfrak{a}}|_\star$  at  $\hat{\beta}_4 > 0$  lies higher, being positive and unstable near  $\hat{\mathfrak{a}}_\star = 0$ . In a vicinity of the flat background at  $\hat{\mathfrak{a}}_\star^\alpha = \delta_\beta^\alpha$ , the potential  $\hat{V}_{\mathfrak{a}}$  in the quadratic approximation is as follows:<sup>31</sup>

$$\hat{V}_{\mathfrak{a}} = \frac{1}{2} M_{\text{Pl}}^2 \hat{v}_2 \hat{f}_\beta^\alpha \hat{f}_\alpha^\beta + \frac{1}{4} (\hat{\beta}_4 - 1) \hat{f}^2, \quad (46)$$

being in LA due to  $\hat{f} = 0$ , in fact, independent of  $\hat{f}$  irrespective of  $\hat{\beta}_4$ . This gives in turn:

$$\hat{T}_{\mathfrak{a}\alpha\beta} = 2M_{\text{Pl}}^2 \hat{v}_2 \hat{f}_{\alpha\beta}, \quad \hat{T}_{\mathfrak{a}} \equiv \hat{T}_{\mathfrak{a}\alpha\beta} \eta^{\alpha\beta} = 0. \quad (47)$$

Due to the Diff gauge symmetry, accounting under the unitary gauge  $\chi^\alpha = 0$  (and thus  $k_{\alpha\beta} = 0$ ) for (42), one can substitute in  $\hat{R}_{\alpha\beta}$  the tensor-graviton field  $\hat{h}_{\alpha\beta}$ , with  $\hat{h} = 0$ , by the DE one  $\hat{f}_{\alpha\beta}$ , with  $\hat{f} = 0$ , getting in LA

$$\hat{R}_{\alpha\beta} = \frac{1}{2} (\partial_\alpha \partial_\gamma \hat{f}_\beta^\gamma + \partial_\beta \partial_\gamma \hat{f}_\alpha^\gamma - \partial^2 \hat{f}_{\alpha\beta}), \quad \hat{R} = \partial_\gamma \partial_\delta \hat{f}^{\gamma\delta}. \quad (48)$$

This results in LA in the tensor and multiscalar FEs (39) as follows:

$$\begin{aligned} \partial_\alpha \partial_\gamma \hat{f}_\beta^\gamma + \partial_\beta \partial_\gamma \hat{f}_\alpha^\gamma - \partial^2 \hat{f}_{\alpha\beta} - \frac{1}{2} \partial_\gamma \partial_\delta \hat{f}^{\gamma\delta} \eta_{\alpha\beta} &= m_g^2 \hat{f}_{\alpha\beta}, \\ \partial_\alpha (m_g^2 \hat{f}_\beta^\alpha - \frac{1}{2} \partial_\gamma \partial_\delta \hat{f}^{\gamma\delta} \delta_\beta^\alpha) &= 0, \end{aligned} \quad (49)$$

clearly independent of  $\hat{\Lambda}$ . The latter property can be explicitly confirmed under an arbitrary  $\chi^\alpha$  due to the Diff gauge symmetry. In the above, we have put  $\hat{v}_2 \equiv m_g^2/4$ , with  $m_g$  to be associated with the tensor-graviton mass. It proves that the multiscalar FE follows, in fact, from the tensor one, with all the nine dark-gravity components remaining thus independent. At that, one missing component corresponds to the neglected scalar graviton, the inclusion of which would explicitly violate the hidden Weyl scale symmetry. To reduce the number of the independent components, impose additionally the covariant conservation condition (25) resulting in LA in the transversality condition

$$\partial_\gamma \hat{f}^{\gamma\alpha} = 0 \quad (50)$$

<sup>30</sup>In particular, a similar potential  $V_{\mathfrak{a}}$  at a proper  $\beta_4$  is used for the consistent description of the massive tensor graviton within the spontaneously broken GR [28].

<sup>31</sup>In fact, due to  $\hat{f} = 0$  in LA the second term in (46) in the quadratic approximation is irrelevant, with the potential being (quasi-)stable irrespective of  $\hat{\beta}_4$ , admitting, in particular,  $\hat{\beta}_4 = 0$  and the purely-quadratic  $\hat{V}_{\mathfrak{a}}(\hat{\hat{\mathfrak{a}}}_\nu^\mu)$ .

followed by  $\hat{\Lambda}_0 = 0$ . This finally results in FE

$$(\partial^2 + m_g^2)\hat{f}_{\alpha\beta} = 0 \quad (51)$$

for the gauge-invariant  $\hat{f}_{\alpha\beta}$  coinciding with the gauge-variant  $\hat{h}_{\alpha\beta} = h_{\alpha\beta} - 1/4 h \eta_{\alpha\beta}$  under the unitary gauge  $\chi_\alpha = 0$ . The FEs (50) and (51) describe precisely the five independent DE degrees of freedom without ghosts, similarly to the Fierz-Pauli case for the massive tensor graviton within the spontaneously broken GR [28]. The covariant conservation condition in the spontaneously broken WTR, likewise GR, proves thus to be of principle to ensure the consistency of the massive tensor graviton description. Still, the conceivable violation of the covariant conservation condition in the context of DE is of particular interest.

### 5.2.3 Massless limit

At last, in the massless limit  $m_g \rightarrow 0$ , in FEs (49) there appears a three-parameter residual gauge symmetry

$$\hat{f}_{\alpha\beta} \rightarrow \hat{f}_{\alpha\beta} - (\partial_\alpha \hat{\varphi}_\beta + \partial_\beta \hat{\varphi}_\alpha), \quad \partial \hat{\varphi} = 0, \quad \partial^2 \hat{\varphi}_\alpha = 0, \quad (52)$$

satisfied also by the constraint (50). This reduces the number of the independent tensor-graviton components on-mass-shell  $m_g = 0$  to two, making thus the massless limit in LA consistent. In fact, in this limit  $\hat{f}_{\alpha\beta}$  reduces to the conventional massless unimodular  $\hat{h}_{\alpha\beta}$ , with  $\hat{h} = 0$ . The presence of the residual gauge symmetry ensuring in the spontaneously broken WTR the smooth massless limit to UR is thus crucial for the explicit consistency in LA of the massive tensor-graviton description.

### 5.2.4 Scalar graviton

Finally, retaining the tensor-graviton description through the Weyl-scale invariant Lagrangian  $\hat{L}_G$ , with the three (at  $n = 4$ ) gauge components converted into the additional physical ones for the massive tensor graviton, one could add the explicitly Weyl-scale violating  $\hat{L}_M(\sigma)$  to consistently realize the last component contained in the multiscalar  $X^a$ . In the neglect by matter, this would result in LA in FE for the scalar graviton  $s$  as a conventional particle:

$$(\partial^2 + m_s^2)s = 0, \quad (53)$$

with a scalar-graviton mass  $m_s$ . In the above, the gauge-invariant scalar field  $s$  coincides with the gauge-variant metric component  $h/2$  under the transverse gauge  $\partial\chi = 0$  (embodying, in particular, the unitary one  $\chi^\alpha = 0$ ). Going beyond LA and accounting for the mixing of the scalar graviton  $\sigma$  with matter remain to be studied.

## 6 Conclusion

The generic multiscalar-metric framework provides a promising route for the modification of gravity, with the spontaneous breaking of the Diff gauge symmetry/relativity as a basic reason of the dark gravity – a merger of gravity with the emergent dark components. The multiscalar-modified/spontaneously broken WTR, built on this route, incorporating the Weyl scale symmetry and the Higgs mechanism for gravity, may well serve as a viable beyond-GR EFT of gravity. It is able to screen the Lagrangian CC  $\hat{\Lambda}$ , with emergence of the induced one  $\hat{\Lambda}_0$  supplemented by the massive tensor and scalar gravitons as the dark gravity components. As EFT, the so constructed dark gravity model, being applied to the (extensions of the) particle SM, allows, in principle, to account also for the spontaneous breaking of the internal symmetries. This should be followed by the step-wise modifications of the Lagrangian CC and matching of the emergent descriptions at the phase transition scales. In the leading approximation, the spontaneously broken WTR proves to be well-behaving as the theory of the massive tensor and scalar gravitons without ghosts. Further developing the basic concept of the spontaneously broken Diff gauge symmetry/relativity — in particular, WTR vs. GR — as a principle source of the emergent dark gravity components of the Universe is urgent.

**Acknowledgment** The author is sincerely grateful to S. S. Gershtein for encouraging discussions.

## References

- [1] Y. B. Zel'dovich, "The cosmological constant and elementary particles", *JETP Lett.* **6**, 316 (1967).
- [2] S. M. Carroll, "The cosmological constant", *Living Rev. Rel.* **4**, 1 (2001); arXiv:astro-ph/0004075.
- [3] S. Weinberg, "The cosmological constant problem", *Rev. Mod. Phys.* **61**, 1 (1989).
- [4] J. Martin, "Everything you always wanted to know about the cosmological constant problem (but were afraid to ask)", *Comptes Rendus Physique* **13**, 566 (2012); arXiv:1205.3365.
- [5] C. P. Burgess, "The cosmological constant problem: Why it's hard to get dark energy from micro-physics", *The Les Houches Summer School *Post-Planck Cosmology**; arXiv:1309.4133.
- [6] S. Capozziello and M. De Laurentis, "Extended theories of gravity", *Phys. Rept.* **509**, 167 (2011); [arXiv:1108.6266].
- [7] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, "Modified gravity and cosmology", *Phys. Rept.* **513**, 1 (2012); arXiv:1106.2476.
- [8] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, "Modified gravity theories on a nutshell: Inflation, bounce and late-time evolution", *Phys. Rept.* **692**, 1 (2017); arXiv:1705.11098v3.
- [9] V. A. Rubakov and P. G. Tinyakov, "Infrared-modified gravities and massive gravitons", *Phys. Usp.* **51**, 759 (2008); arXiv:0802.4379.
- [10] C. de Rham, "Massive gravity", *Living Rev. Rel.* **17**, 7 (2014); arXiv:1401.4173.
- [11] J. L. Anderson and D. Finkelstein, "Cosmological constant and fundamental length", *Am. J. Phys.* **39**, 901 (1971).
- [12] J. J. van der Bij, H. van Dam, and Y. J. Ng, "The exchange of massless spin-two particles", *Physica A* **116**, 307 (1982).
- [13] K.-I. Izawa, "Derivative expansion in quantum theory of gravitation", *Prog. Theor. Phys.* **93**, 615 (1995); arXiv:hep-th/9410111.
- [14] E. Alvarez, D. Blas, J. Garriga, and E. Verdaguer, "Transverse Fierz-Pauli symmetry", *Nucl. Phys. B* **756**, 148 (2006); arXiv:hep-th/0606019.
- [15] E. Alvarez and R. Vidal, "Weyl transverse gravity (WTDiff) and the cosmological constant", *Phys. Rev. D* **81**, 084057 (2010); arXiv:1001.4458.
- [16] I. Oda, "Classical Weyl transverse gravity", *Eur. Phys. J. C* **77**, 284 (2017); arXiv:1610.05441.
- [17] R. Carballo-Rubio, "Longitudinal diffeomorphisms obstruct the protection of vacuum energy", *Phys. Rev. D* **91**, 124071 (2015); arXiv:1502.05278.
- [18] E. Alvarez, S. Gonzalez-Martin, M. Herrero-Valea, and C. P. Martin, "Unimodular gravity redux", *Phys. Rev. D* **92**, 061502 (2015); arXiv:1505.00022.
- [19] C. Barcelo, R. Carballo-Rubio, and L. J. Garay, "Absence of cosmological constant problem in special relativistic field theory of gravity", *Ann. Phys.* **398**, 9 (2018); arXiv:1406.7713.
- [20] M. Fierz and W. Pauli, "On relativistic wave equations for particles of arbitrary spin in an electromagnetic field", *Proc. Roy. Soc. Lond. A* **173**, 211 (1939).
- [21] H. van Dam and M. J. G. Veltman, "Massive and massless Yang-Mills and gravitational fields", *Nucl. Phys. D* **22**, 397 (1970).

- [22] V. I. Zakharov, “Linearized gravitaion theory and the graviton mass”, JETP Lett. **12**, 312 (1970).
- [23] A. I. Vainshtein, “To the problem of nonvanishing graviton mass”, Phys. Lett. B **39**, 393 (1972).
- [24] D. G. Boulware and S. Deser, “Can graviton have a finite range?”, Phys. Rev. D **6**, 3368 (1972).
- [25] G. ’t Hooft, “Unitarity in the Brout-Englert-Higgs mechanism for gravity”, arXiv:0708.3184.
- [26] Z. Kakushadze, “Gravitational Higgs mechanism and massive gravity”, Int. J. Mod. Phys. A **23**, 1581 (2008); arXiv:0709.1673.
- [27] D. A. Demir and N. K. Pak, “General tensor Lagrangians from gravitational Higgs mechanism”, Class. Quant. Grav. **26**, 105018 (2009); arXiv:0904.0089.
- [28] A. H. Chamseddine and V. Mukhanov, “Higgs for graviton: simple and elegant solution”, JHEP **1008**, 011 (2010); arXiv:1002.3877.
- [29] I. Oda, “Higgs mechanism for gravitons”, Mod. Phys. Lett. **A 25**, 2411 (2010); arXiv:1003.1437.
- [30] I. Oda, “Remarks on Higgs mechanism for gravitons”, Phys. Lett. B **690**, 322 (2010); arXiv:1004.3078.
- [31] L. Alberte, A. H. Chamseddine, and V. Mukhanov, “Massive gravity: resolving the puzzles”, JHEP **1012**, 023 (2010); arXiv:1008.5132.
- [32] L. Alberte, A. H. Chamseddine, and V. Mukhanov, “Massive gravity: exorcising the ghost”, JHEP **1104**, 004 (2011); arXiv:1011.0183.
- [33] E. I. Guendelman and A. B. Kaganovich, “The principle of non-gravitating vacuum energy and some of its consequences”, Phys. Rev. D **53**, 7020 (1996); arXiv:gr-qc/9605026.
- [34] E. I. Guendelman and A. B. Kaganovich, “Gravitational theory without the cosmological constant problem”, Phys. Rev. D **55**, 5970 (1997); arXiv:gr-qc/9611046.
- [35] Y. F. Pirogov, “Quartet-metric general relativity: scalar graviton, dark matter and dark energy”, Eur. Phys. J. C **76**, 215 (2016); arXiv:1511.04742.
- [36] Y. F. Pirogov, “Quartet-metric gravity and dark components of the Universe”, Int. J. Mod. Phys.: Conf. Series **47**, 1860101 (2018); arXiv:1712.00612.
- [37] Y. F. Pirogov, “Affine-Goldstone/quartet-metric gravity and beyond”, Phys. Atom. Nucl. **82**, 503 (2019); arXiv:1807.02160v2.
- [38] Y. F. Pirogov, “Quartet-metric/multi-component gravity: scalar graviton as emergent dark substance”, JCAP **01**, 055 (2019); arXiv:1811.12923.
- [39] Y. F. Pirogov, “Unimodular bimode gravity and the coherent scalar-graviton field as galaxy dark matter”, Eur. Phys. J. C **72**, 2017 (2012); arXiv:1111.1437.
- [40] G. F. R. Ellis, H. van Elst, J. Murugan, and J.-P. Uzan, “On the trace-free Einstein equations as a viable alternative to General Relativity”, Class. Quant. Grav. **28**, 225007 (2011); arXiv:1008.1196.
- [41] G. F. R. Ellis, “The trace-free Einstein equations and inflation”, Gen. Rel. Grav. **46**, 1619 (2014); arXiv:1306.3021.
- [42] P. G. Bergmann, “Comments on the scalar-tensor theory”, Int. J. Theor. Phys. **1**, 25 (1968).
- [43] G. W. Horndeski, “Second-order scalar-tensor field equations in a four-dimensional space”, Int. J. Theor. Phys. **10**, 363 (1974).