

Addendum to “Hasenöhrl and the Equivalence of Mass and Energy”

Tony Rothman*

NYU, Dept. of Applied Physics, (retired)

L^AT_EX-ed September 18, 2021

Abstract

This addendum addresses several objections regarding accelerating cavities in the paper by Boughn and Rothman, “Hasenöhrl and the Equivalence of Mass and Energy,” arXiv:1108.2250.

PACS: 03.30.+p, 01.65.+g, 03.50.De

Keywords: Hasenöhrl, Poincaré, Abraham, Einstein, Fermi, mass-energy equivalence, blackbody radiation, special relativity

The paper “Hasenöhrl and the Equivalence of Mass and Energy,” by Stephen Boughn and Tony Rothman (BR)[1] concerned the early attempts by Fritz Hasenöhrl to derive the relationship between the energy of black body radiation in a cavity and its equivalent mass. Section 5.2 of that paper was an attempt to give a relativistically correct derivation for slowly accelerating cavities. The derivation, however, contained one mistake and also brought criticisms about a further point. This note attempts to correct the error and also addresses the second criticism.

The external force density (force per unit volume) on a radiation-fluid element is given by $\mathbf{f}_\mu = T_{\mu\nu}{}^{,\nu}$, or as in BR (5.14),

$$\mathbf{f}_\mu = \frac{d(\rho_o + p_o)}{c^2 d\tau} u_\mu + \frac{(\rho_o + p_o)}{c^2} \left[\frac{du_\mu}{d\tau} + u_\mu \frac{\partial u_\nu}{\partial x_\nu} \right] + \frac{\partial p_o}{\partial x_\nu} \eta_{\mu\nu}. \quad (1)$$

The 4-force density, including heat transfer, can be written as

$$\vec{\mathbf{f}} = [(1/c)(\mathcal{F} \cdot \mathbf{v} + q_{,t}), \mathcal{F}] \quad (2)$$

*tonyrothman@gmail.com

where \mathcal{F} is the Newtonian three-force density q is the heat density. In that case, one gets BR (5.16), the fundamental equation of motion for the radiation:

$$\mathfrak{f}_\mu = \frac{(\rho_o + p_o)}{c^2} \frac{du_\mu}{d\tau} + \frac{u_\mu}{c^2} \frac{dp_o}{d\tau} + \frac{\partial p_o}{\partial x_\nu} \eta_{\mu\nu} + \frac{\dot{q}_o u_\mu}{c^2}. \quad (3)$$

However, from BR (3.32) the heat term is $\sim E\beta^4$, and so when $ax/c^2 \sim v^2/c^2 \ll 1$, which is the approximation assumed in BR, it can be neglected. Also in that case one can use the equilibrium values of ρ_o and p_o ($= \rho_o/3$). For motion in the x -direction and retaining terms only of order $a \sim v^2/c^2$, we have

$$\mathfrak{f}' = \frac{4}{3c^2} \rho_o a + \frac{\partial p_o}{\partial x} (1 + \beta^2), \quad (4)$$

The total force on the fluid in the lab frame is then:

$$F = \int_{V'} \mathfrak{f}' dV' = \int_{V_o} \left[\frac{4}{3c^2} \rho_o a + \frac{\partial p_o}{\partial x} (1 + \beta^2) \right] \gamma^{-1} dV_o. \quad (5)$$

BR took γ to be the general time-dilation factor

$$\gamma = \frac{1}{(1 + \frac{2ax}{c^2} - \frac{v^2}{c^2})^{1/2}}. \quad (6)$$

However, if following Rindler[2], we define γ to be the ratio of the proper cavity length to its length in the lab frame, there can be only one γ , which we take to be $\gamma^{-1} = (1 - \beta^2)^{1/2}$. (A "Born rigid" cavity will have different proper accelerations at each point. Nevertheless, for such a cavity, boosting to the instantaneous co-moving frame of any fiducial point provides an instantaneous rest frame for the entire cavity, if it is small enough. Each point in the cavity is undergoing hyperbolic motion, and because at fixed time in the lab frame the different cavity points must be moving with different velocities, the value of γ might appear to be ambiguous. However, γ^{-1} at opposite ends differ only at $\mathcal{O}(\beta^4)$, and so which value of β one chooses is immaterial for this calculation.) To order β^2 Eq. (5) gives

$$F = \int_{V_o} \left[\frac{4}{3c^2} \rho_o a + \frac{\partial p_o}{\partial x} (1 + \frac{1}{2}\beta^2) \right] dV_o, \quad (7)$$

or in terms of a

$$F = \int_{V_o} \left[\frac{4}{3c^2} \rho_o a + \frac{\partial p_o}{\partial x} (1 + ax) \right] dV_o. \quad (8)$$

The remainder of the derivation goes through as in BR: To integrate the second term we use the divergence theorem $\int_V (\nabla p_o) dV = \oint_A p_o \hat{\mathbf{n}} dA$ for outwardly directed normal $\hat{\mathbf{n}}$. We integrate the third term by parts and get

$$\int_{V_o} \frac{\partial p_o}{\partial x} (1 + ax) dV_o = \oint_A p_o \hat{\mathbf{n}} dA + \frac{a}{3c^2} (\rho_o V_o)|_{boundary} - \frac{a}{3c^2} \int_{V_o} \rho_o \frac{\partial x}{\partial x} dV_o. \quad (9)$$

The first term in this equation is the negative of the external force on the volume element of a fluid. Inserting that term into Eq.(8) and ignoring the last term gives the relativistic equation of hydrostatic equilibrium when $F = 0$, in agreement with Boughn[3]. That is,

$$\oint_A p_o \hat{\mathbf{n}} dA = \frac{4}{3} \frac{E_o}{c^2} a. \quad (10)$$

Since the left-hand-side represents a force $= ma$, one immediately has $E_o = 3/4 mc^2$, also in accord with Boughn.

However, in this case, if the above assumptions are correct, then the last term in Eq. (8) apparently provides another relativistic correction to the effective mass of the system, lowering the total external force. That is, if the system is spatially bounded and we extend the integration volume to infinity where the density and pressure is zero, we can drop the boundary terms. Then the total external force becomes

$$F = \frac{E_o}{c^2} a, \quad (11)$$

and we immediately have $E_o = mc^2$, as desired.

Although the neglect of the boundary terms may seem like a risky move, that procedure is employed in the standard proofs of von Laue's and Klein's theorem (see BR §4 and Ohanian[4]). If one takes the point of view that in order for the fluid to be spatially bounded, it must be contained in a physical cavity (which has energy in its own mass density and stresses), one can proceed as follows.

If the rest energy density of the shell is $T_{00} = \mu$, the stresses are $T_{xx} = \sigma$ and $T_{0x} = 0$, then the stress-energy tensor for the cavity can be written in perfect-fluid form[5]:

$$(T_{\mu\nu})_{shell} = (\mu + \sigma) u_\mu u_\nu + \eta_{\mu\nu} \sigma \quad (12)$$

We then require that the divergence of the combined stress-energy tensor be the force external to the entire system: $f_x = (T_{x\nu}{}^{,\nu})_{fluid} + (T_{x\nu}{}^{,\nu})_{shell}$. Hence, in analogy to above,

$$F_x = \int_{cav} \left[\frac{\rho + p}{c^2} a + \frac{\partial p}{\partial x} (1 + ax) \right] dV_{cav} + \int_{shell} \left[\frac{\mu + \sigma}{c^2} a + \frac{\partial \sigma}{\partial x} (1 + ax) \right] dV_{shell}. \quad (13)$$




Figure 1: The fluid is assumed to be held in a cavity of length D , radius R and wall thickness h .

Referring to Figure 1, we see that for the gas to be contained by the shell, we must have $\pi R^2 p = -2\pi R h \sigma$, where h is the thickness of the shell walls and R is the cavity radius. Thus $\sigma = -pR/2h$. However, $dV_{cav} = \pi R^2 dx$, while $dV_{shell} = 2\pi R h dx = (2h/R)dV_{cav}$. Thus the integrals involving p cancel with those involving σ , leaving

$$F_{ext} = \int_{cav} \frac{\rho a}{c^2} dV_{cav} + \int_{shell} \frac{\mu a}{c^2} dV_{shell} = E \frac{a}{c^2}. \quad (14)$$

Once again, if $F = ma$, we can immediately conclude that $E = mc^2$ for the entire system. Consequently, if one ignores the boundary terms in Eq. (9) it apparently becomes unnecessary to introduce elastic stresses in order to achieve the correct answer for the radiation alone. In a sense, this appears to be a vindication of Hasenöhl's approach, although because he effectively neglected the relativistic corrections in the fluid's equation of motion, he achieved an incorrect result.

Acknowledgement

Many thanks to Steve Boughn for his continued interest in this problem and many helpful remarks.

References

- [1] S. Boughn and T. Rothman, "Hasenöhl and the Equivalence of Mass and Energy" (arXiv:1108.2250).

- [2] W. Rindler, *Essential Relativity* (Springer Verlag: New York, 1977).
- [3] S. Boughn, Eur. Phys. J. H. **38**, 262-78 (2013).
- [4] H. Ohanian, Am. J. Phys. **80**, 1067 (2012).
- [5] C. Møller, *The Theory of Relativity* (Oxford University Press: Oxford, 1962).