

Positive least energy solutions for k -coupled Schrödinger system with critical exponent: the higher dimension and cooperative case*

Xin Yin, Wenming Zou

Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China.

Abstract

In this paper, we study the following k -coupled nonlinear Schrödinger system with Sobolev critical exponent:

$$\begin{cases} -\Delta u_i + \lambda_i u_i = \mu_i u_i^{2^*-1} + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{\frac{2^*}{2}-1} u_j^{\frac{2^*}{2}} & \text{in } \Omega, \\ u_i > 0 & \text{in } \Omega \quad \text{and} \quad u_i = 0 \quad \text{on } \partial\Omega, \quad i = 1, 2, \dots, k. \end{cases}$$

Here $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, $2^* = \frac{2N}{N-2}$ is the Sobolev critical exponent, $-\lambda_1(\Omega) < \lambda_i < 0, \mu_i > 0$ and $\beta_{ij} = \beta_{ji} \neq 0$, where $\lambda_1(\Omega)$ is the first eigenvalue of $-\Delta$ with the Dirichlet boundary condition. We characterize the positive least energy solution of the k -coupled system for the purely cooperative case $\beta_{ij} > 0$, in higher dimension $N \geq 5$. Since the k -coupled case is much more delicate, we shall introduce *the idea of induction*. We point out that *the key idea* is to give a more accurate upper bound of the least energy. It's interesting to see that the least energy of the k -coupled system decreases as k grows. Moreover, we establish the existence of positive least energy solution of the limit system in \mathbb{R}^N , as well as classification results.

Key words: k -Coupled Schrödinger System; Positive least energy solution; Existence; Uniqueness.

2010 Mathematics Subject Classification: 35J50, 35J15, 35J60.

*Supported by NSFC(20171301826, 20181301532).
E-mails: yinx16@mails.tsinghua.edu.cn (Yin); zou-wm@mail.tsinghua.edu.cn (Zou)

1 Introduction

In this paper we consider the following k -coupled nonlinear Schrödinger system:

$$\begin{cases} -\Delta u_i + \lambda_i u_i = \mu_i u_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{p-1} u_j^p & \text{in } \Omega, \\ u_i > 0 & \text{in } \Omega \quad \text{and} \quad u_i = 0 \quad \text{on } \partial\Omega, \quad i = 1, 2, \dots, k, \end{cases} \quad (1.1)$$

where $\Omega = \mathbb{R}^N$ or $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, $p > 1$ and $p \leq 2^*/2$ if $N \geq 3$, $\mu_i > 0$ and $\beta_{ij} = \beta_{ji} \neq 0$ is a coupling constant.

System (1.1) arises when we consider the standing wave solutions to the following time-depending coupled nonlinear Schrödinger system:

$$\begin{cases} -i \frac{\partial}{\partial t} \Phi_1 = \Delta \Phi_1 + \mu_1 |\Phi_1|^2 \Phi_1 + \beta |\Phi_2|^2 \Phi_1, \\ -i \frac{\partial}{\partial t} \Phi_2 = \Delta \Phi_2 + \mu_2 |\Phi_2|^2 \Phi_2 + \beta |\Phi_1|^2 \Phi_2, \\ \Phi_j = \Phi_j(x, t) \in \mathbb{C}, \quad j = 1, 2, \quad N \leq 3, \end{cases} \quad (x, t) \in \mathbb{R}^N \times \mathbb{R}, \quad (1.2)$$

where i is the imaginary unit, $\mu_1, \mu_2 > 0$ and $\beta \neq 0$ is a coupling constant. The system (1.2) has many applications in physics, such as the occurrence of phase separation in Bose–Einstein condensates with multiple states, or the propagation of mutuary in coherent wave packets in nonlinear optics, see e.g. [2, 25, 26, 44].

From the physical aspect, the solution Φ_j denotes the j th component of the beam in Kerr-like photorefractive media (see [2]), μ_j represents self-focusing in the j th component, and the coupling constant β represents the interaction between the two components of the beam.

To study the solitary wave solutions of system (1.2), we set $\Phi_j(x, t) = e^{i\lambda_j t} u_j(x)$ for $j = 1, 2$. Then, it is reduced to system (1.1) with $k = 2$ and $p = 2$.

For the subcritical case, that is, $N \leq 3$, the existence of least energy solutions were studied in [3, 8, 9, 18, 21, 22, 23, 29, 35, 38, 40, 43, 45] and the references therein. The existence and multiplicity of positive and sign-changing solutions were studied by [3, 7, 9, 14, 15, 18, 29, 30, 32, 33, 34, 38, 41] and the references therein.

For the critical case $p = 2$ and $N = 4$, Chen and Zou [17] proved that there exists $0 < \beta_1 < \beta_2$ (β_1 and β_2 depend on λ_i and β), such that (1.1) has a positive least energy solution if $\beta < \beta_1$ or $\beta > \beta_2$, while (1.1) has no positive least energy solution if $\beta_1 \leq \beta \leq \beta_2$.

Later in [19], Chen and Zou considered the critical case $2p = 2^* := \frac{2N}{N-2}$ of system (1.1) in higher dimension $N \geq 5$. It turns out that different phenomena happen from the special case $N = 4$. Indeed, the authors showed that system (1.1) has a positive least energy solution for any $\beta \neq 0$. When $\beta = 0$, the system is reduced to the well-known Brézis–Nirenberg problem[11]

$$-\Delta u + \lambda_i u = \mu_i |u|^{2^*-2} u, \quad u \in H_0^1(\Omega). \quad (1.3)$$

The Brézis-Nirenberg problem (1.3) has been studied intensively, and we refer the reader to [13, 16, 24, 39] and references therein.

From then on, many results corresponding to the coupled system with critical Sobolev exponent have been studied. See [31, 37, 50, 52, 51] and references therein.

For the more general case where $k \geq 2$, physically the coefficient $\beta_{ii} = \mu_i$ represents self-focusing within the i th component, while β_{ij} ($i \neq j$) illustrates interaction between two different components. If $\beta_{ij} > 0$, the interaction between the i th and j th components is of cooperative type; while $\beta_{ij} < 0$ means that the interaction is of competitive type. The relation $\beta_{ij} = \beta_{ji}$ illustrates the symmetry in the interaction between different components.

Let $\lambda_1(\Omega)$ be the first eigenvalue of $-\Delta$ with the Dirichlet boundary condition.

For the purely cooperative case, where $\beta_{ij} > 0$ for $i, j = 1, 2, \dots, k$, Guo, Luo and Zou [27] obtained the existence and classification of positive least energy solutions to (1.1) under $-\lambda_1(\Omega) < \lambda_1 = \dots = \lambda_k < 0$ in case of a bounded smooth domain $\Omega \subset \mathbb{R}^4$ with some additional conditions on the coefficients. In the same paper, the authors treated the system with $\lambda_1 = \dots = \lambda_k = 0$ and $\Omega = \mathbb{R}^4$, which can be seen as the limit problem corresponding to the previous problem. Wu [49] also obtained the existence of positive least energy solution in \mathbb{R}^N with $N \geq 4$ and $\beta_{ij} = \beta$, $i \neq j$ via variational arguments.

Meanwhile, for the purely competitive case $\beta_{ij} < 0$, [20, 42, 48] studied the existence of positive least energy solutions and phase separation phenomena respectively for $N = 4$ and $N \geq 5$. And some results are obtained for mixed cooperative and competitive case when $N = 4$. See [42, 46] and references therein.

To the best of our knowledge, there is no paper considering (1.1) with arbitrary purely cooperative coefficients $\beta_{ij} > 0$ in higher dimension $N \geq 5$. It is natural to ask *whether the more general k -coupled system (1.1) admits a positive least energy solution similarly as the 2-coupled system proved in [19]*. In the sequel we assume that

$$2p = 2^*. \tag{1.4}$$

We shall see that the k -coupled case are more delicate than the 2-coupled case. *The key reason* is that $k \geq 2$ and $1 < p < 2$ when $N \geq 5$. The fact that $k \geq 2$ and $1 < p < 2$ make the problem more complicated comparing to the cases $p = 2, k \geq 2$ and $1 < p < 2, k = 2$. Some more ideas and techniques are needed.

First, we obtained the following results.

Theorem 1.1. *Assume that $N \geq 3$ and (1.4). Then system (1.1) has no positive solutions if one of the following conditions holds.*

- (1) $\lambda_i \leq -\lambda_1(\Omega)$, $\mu_i > 0$ and $\beta_{ij} > 0$ for $i, j = 1, 2, \dots, k$.
- (2) $\mu_i, \beta_{ij} \in \mathbb{R}$, $\lambda_i \geq 0$ for $i = 1, 2, \dots, k$, and Ω is star-shaped.

If $\Omega = \mathbb{R}^N$, and (u_1, \dots, u_k) is any a solution of (1.1), by the Pohozaev identity (2.1), it is easy to get that $\int_{\mathbb{R}^N} \sum \lambda_i u_i^2 dx = 0$, so $(u_1, \dots, u_k) \equiv (0, \dots, 0)$ if $\lambda_1, \dots, \lambda_k$ are of the same sign. Therefore, in the sequel we assume that $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain. Some non-existence results also have been attained in Theorem 1.1. In order to study the existence of positive least energy solution of (1.1), we assume that $-\lambda_1(\Omega) < \lambda_i < 0$, $\mu_i > 0$ and $\beta_{ij} > 0$ for $i, j = 1, 2, \dots, k$.

We call a solution (u_1, \dots, u_k) is nontrivial if $u_i \not\equiv 0$ for all $i = 1, 2, \dots, k$, while a solution (u_1, \dots, u_k) is semi-trivial if $u_i \equiv 0$ for some $i = 1, 2, \dots, k$. We only study the nontrivial solutions of (1.1) in this paper.

Denote $H := (H_0^1(\Omega))^k$ and $\vec{u} = (u_1, u_2, \dots, u_k) \in H$ is a vector function. It is well known that solutions of (1.1) correspond to the critical points of C^1 functional $E : H \rightarrow \mathbb{R}$ given by

$$\begin{aligned} E(\vec{u}) &= \frac{1}{2} \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) \\ &\quad - \frac{1}{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right). \end{aligned} \quad (1.5)$$

We say a solution \vec{u} of (1.1) is a least energy solution, if \vec{u} is nontrivial and

$$E(\vec{u}) \leq E(\vec{v})$$

for any other nontrivial solution \vec{v} of (1.1). As in [29], we define a Nehari type manifold

$$\mathcal{M} = \{\vec{u} \in H \mid u_i \not\equiv 0, E'(\vec{u})(0, \dots, u_i, \dots, 0) = 0 \text{ for } i = 1, 2, \dots, k\}.$$

Then any nontrivial solutions of (1.1) belong to \mathcal{M} . Take $\varphi_1, \varphi_2, \dots, \varphi_k \in C_0^\infty(\Omega)$ with $\varphi_i \not\equiv 0$ and $\text{supp}(\varphi_i) \cap \text{supp}(\varphi_j) = \emptyset$ for $\forall i, j = 1, 2, \dots, k$; then there exists $c_1, c_2, \dots, c_k > 0$ such that $(\sqrt{c_1}\varphi_1, \sqrt{c_2}\varphi_2, \dots, \sqrt{c_k}\varphi_k) \in \mathcal{M}$, so $\mathcal{M} \neq \emptyset$.

Define the least energy

$$B := \inf_{\vec{u} \in \mathcal{M}} E(\vec{u}) = \inf_{\vec{u} \in \mathcal{M}} \frac{1}{N} \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2). \quad (1.6)$$

Now we consider the special case $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \dots = \lambda_k = \lambda < 0$. It is well known that the Brézis-Nirenberg problem [11]

$$-\Delta u + \lambda u = |u|^{2^*-2}u, \quad u \in H_0^1(\Omega) \quad (1.7)$$

has a positive least energy solution ω with energy

$$B_1 := \frac{1}{N} \int_{\Omega} (|\nabla \omega|^2 + \lambda \omega^2) dx = \frac{1}{N} \int_{\Omega} \omega^{2^*} dx. \quad (1.8)$$

Consider the following nonlinear algebraic problem

$$\begin{cases} \mu_i t_i^{2p-2} + \sum_{j=1, j \neq i}^k \beta_{ij} t_i^{p-2} t_j^p = 1, & \text{for } i = 1, 2, \dots, k, \\ t_i > 0, & \text{for } i = 1, 2, \dots, k. \end{cases} \quad (1.9)$$

In [17, 19], Chen and Zou proved the existence of solution to (1.9) in the case $k = 2$. Recently, Wu [49] proved that (1.9) has a solution for $k \geq 2$ with $\beta_{ij} = \beta$, by applying the variational argument to the related system. We generalize the existence result of the algebraic system (1.9) to arbitrary $\beta_{ij} > 0$ and give a more minute and geometric proof in Lemma 2.2. Moreover, we will prove in Lemma 2.2 that there exists $\vec{\mathbf{t}} = (\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$ such that

$$\vec{\mathbf{t}} \text{ satisfies (1.9) and } \vec{\mathbf{t}} \text{ attains } d_k. \quad (1.10)$$

Here, $d_k := \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \mathcal{G}_k(\vec{\mathbf{t}})$, where

$$\mathcal{G}_k(\vec{\mathbf{t}}) := \sum_{i=1}^k \left(\frac{t_i^2}{2} - \frac{\mu_i |t_i|^{2p}}{2p} \right) - \frac{1}{2p} \left(\sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p \right),$$

$$\mathcal{P}_k := \{ \vec{\mathbf{t}} \in \mathbb{R}^k \setminus \{ \vec{\mathbf{0}} \} \mid P_k(\vec{\mathbf{t}}) := \sum_{i=1}^k (t_i^2 - \mu_i |t_i|^{2p}) - \sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p = 0 \}.$$

We will see that d_k has a similar form as the minimizing problem (1.23) and d_k plays an important role in this paper.

Our second result also deals with the special case $\lambda_1 = \lambda_2 = \dots = \lambda_k := \lambda$.

Theorem 1.2. *Assume that $N \geq 5$ and (1.4), $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \dots = \lambda_k := \lambda < 0$ and $\mu_i, \beta_{ij} > 0$. Then system (1.1) has a positive solution of the synchronized form*

$$(t_1 \omega, t_2 \omega, \dots, t_k \omega), \quad (1.11)$$

where ω is the positive least energy solution of (1.7) and (t_1, t_2, \dots, t_k) solves (1.9).

Remark 1.1.

- (1) Denote $\mathcal{R} = (\beta_{ij})_{k \times k}$ as the coupling matrix, where $\beta_{ij} = \beta_{ji}$ and $\beta_{ii} = \mu_i$. For the special case $N = 4$, Guo, Luo and Zou [27] proved that the least energy B is attained by a symmetrized form solution (1.11), under the assumptions that

$$\mathcal{R} = (\beta_{ij})_{k \times k} \text{ is invertible and } \sum_{j=1}^k (\mathcal{R}^{-1})_{ij} > 0, \quad i = 1, 2, \dots, k. \quad (1.12)$$

Actually, under these assumptions, the algebraic system has pairs of solutions $(\pm t_1, \pm t_2, \dots, \pm t_k)$, then the signed solutions $(\pm t_1 \omega, \dots, \pm t_k \omega)$ are nontrivial signed solutions of (1.1).

(2) If $\lambda \leq -\lambda_1(\Omega)$, we can obtain sign-changing solutions of synchronized form.

We shall give a classification result of the positive least energy solution of the (1.1).

Theorem 1.3. *Assume that $N \geq 5$, $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \dots = \lambda_k := \lambda < 0$ and $\mu_i, \beta_{ij} > 0$. Then the positive least energy solution of the system (1.1) must be of the synchronized form*

$$(\tilde{t}_1\omega, \tilde{t}_2\omega, \dots, \tilde{t}_k\omega),$$

where ω is a positive least energy solution of (1.7) and $(\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$ is defined in (1.10).

Moreover, assume additionally that $\beta_{ij} := \beta > 0$ and $\Omega \subset \mathbb{R}^N$ is a ball, then there exists $\beta_k > 0$, such that the positive least energy solution of the system (1.1) is unique for $\beta > \beta_k$.

Remark 1.2.

- (1) For the special case $N = 4$, Guo, Luo and Zou [27] classified the positive least energy solution of the synchronized form (1.11), under the assumptions (1.12) and \mathcal{R} is positively or negatively definite. In particular, the positive least energy solution is unique if Ω is a ball in \mathbb{R}^4 .
- (2) Recently, Tavares and You [42] also obtained the classification result for the special case $N = 4$, under the same assumption as our result. But the uniqueness problem was not covered in [42].

Now let us consider the general case $-\lambda_1(\Omega) < \lambda_1, \lambda_2, \dots, \lambda_k < 0$.

Theorem 1.4. *Assume that $N \geq 5$ and (1.4). Suppose that $-\lambda_1(\Omega) < \lambda_1, \lambda_2, \dots, \lambda_k < 0$ and $\mu_i > 0$. Then system (1.1) has a positive least energy solution \vec{u} with $E(\vec{u}) = B$ for any $\beta_{ij} > 0$.*

Remark 1.3. *In [20, 48], the authors obtained the existence of positive least energy solutions of the system (1.1) for the purely competitive case $\beta_{ij} < 0$.*

For each $\Theta \subsetneq \{1, \dots, k\}$, we obtain the following $(k - |\Theta|)$ -coupled system

$$\begin{cases} -\Delta u_i + \lambda_i u_i = \mu_i u_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{p-1} u_j^p & \text{in } \Omega, \\ u_i > 0 \text{ in } \Omega \text{ and } u_i = 0 \text{ on } \partial\Omega, \quad i, j \in \{1, 2, \dots, k\} \setminus \Theta, \end{cases} \quad (1.13)$$

by replacing $\lambda_i, \mu_i, \beta_{ij}, \beta_{ji}$ with 0 if $i \in \Theta$, where $|\Theta|$ is the cardinality of Θ .

Note that the nontrivial solutions of (1.13) can be extended to the semi-trivial solutions of k -coupled system (1.1).

Denote $\tau = k - |\Theta|$. Indeed, we can establish C_k^τ different τ -coupled systems from (1.1). For every fixed Θ , we denote $\{i_1, \dots, i_\tau\} = \{1, 2, \dots, k\} \setminus \Theta$ and $B_{\mu_{i_1} \dots \mu_{i_\tau}}$ as

the least energy of the τ -coupled system (4.3) with coefficients $(\mu_{i_1}, \dots, \mu_{i_\tau})$, that is,

$$B_{\mu_{i_1} \dots \mu_{i_\tau}} := \inf \{ E(\vec{u}) : \vec{u} = (u_1, \dots, u_k) \text{ solves (1.1) and } u_i = 0 \text{ iff } i \in \Theta \}. \quad (1.14)$$

Denote

$$\bar{B}_\tau := \min_{\forall \{l_1, \dots, l_\tau\} \subsetneq \{1, 2, \dots, k\}} \{ B_{\mu_{l_1} \dots \mu_{l_\tau}} \}. \quad (1.15)$$

Next we need to obtain a more refined estimates on the least energy of the k -coupled system.

Theorem 1.5. *Assume that $N \geq 5$ and (1.4). Suppose that $-\lambda_1(\Omega) < \lambda_1, \lambda_2, \dots, \lambda_k < 0$ and $\mu_i, \beta_{ij} > 0$. Then the least energy of (1.1) has an upper bound*

$$B < \min \{ \bar{B}_1, \bar{B}_2, \dots, \bar{B}_{k-1} \}. \quad (1.16)$$

To be more specific,

$$B < \bar{B}_{k-1} < \dots < \bar{B}_1.$$

Moreover,

$$B := B_{\mu_1 \mu_2 \dots \mu_k} < B_{\mu_1 \mu_2 \dots \mu_{k-1}} < B_{\mu_1 \mu_2 \dots \mu_{k-2}} < \dots < B_{\mu_1 \mu_2} < B_{\mu_1}.$$

By Theorem 1.5, it's interesting to see that the least energy of the k -coupled system may decrease as k grows.

Since the nonlinearity and the coupling term are both critical in (1.1), the existence of nontrivial solutions of (1.1) depends heavily on the existence of the least energy solution of the following limit problem

$$\begin{cases} -\Delta u_i = \mu_i u_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{p-1} u_j^p, & x \in \mathbb{R}^N, \\ u_i \in D^{1,2}(\mathbb{R}^N), & i = 1, 2, \dots, k, \end{cases} \quad (1.17)$$

where $D^{1,2}(\mathbb{R}^N) := \{u \in L^{2^*}(\mathbb{R}^N) : |\nabla u| \in L^2(\mathbb{R}^N)\}$ with norm $\|u\|_{D^{1,2}} := (\int_{\mathbb{R}^N} |\nabla u|^2 dx)^{1/2}$. Let S be the sharp constant of $D^{1,2}(\mathbb{R}^N) \hookrightarrow L^{2^*}(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} |\nabla u|^2 dx \geq S \left(\int_{\mathbb{R}^N} |u|^{2^*} dx \right)^{\frac{2}{2^*}}. \quad (1.18)$$

For $\varepsilon > 0$ and $y \in \mathbb{R}^N$, we consider the Aubin-Talenti instanton [6, 47] $U_{\varepsilon, y} \in D^{1,2}(\mathbb{R}^N)$ defined by

$$U_{\varepsilon, y}(x) := [N(N-2)]^{\frac{N-2}{4}} \left(\frac{\varepsilon}{\varepsilon^2 + |x-y|^2} \right)^{\frac{N-2}{2}}. \quad (1.19)$$

Then $U_{\varepsilon, y}$ satisfies $-\Delta u = |u|^{2^*-2}u$ in \mathbb{R}^N and

$$\int_{\mathbb{R}^N} |\nabla U_{\varepsilon, y}|^2 dx = \int_{\mathbb{R}^N} |U_{\varepsilon, y}|^{2^*} dx = S^{N/2}. \quad (1.20)$$

Clearly (1.17) has semi-trivial solutions

$$(\mu_1^{\frac{2-N}{4}} U_{\varepsilon,y}, 0, \dots, 0), (0, \mu_2^{\frac{2-N}{4}} U_{\varepsilon,y}, 0, \dots, 0), \dots, (0, \dots, 0, \mu_k^{\frac{2-N}{4}} U_{\varepsilon,y}).$$

Here, we are only concerned with nontrivial solutions of (1.17). Define $D := (D^{1,2}(\mathbb{R}^N))^k$ and a C^1 functional $I : D \rightarrow \mathbb{R}$ given by

$$I(\vec{\mathbf{u}}) := \frac{1}{2} \int_{\mathbb{R}^N} \sum_{i=1}^k |\nabla u_i|^2 - \frac{1}{2p} \int_{\mathbb{R}^N} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right). \quad (1.21)$$

We consider the following set as in [29]:

$$\mathcal{N} = \{ \vec{\mathbf{u}} \in D : u_i \neq 0, I'(\vec{\mathbf{u}})(0, \dots, 0, u_i, 0, \dots, 0) = 0, i = 1, 2, \dots, k \}. \quad (1.22)$$

Then any nontrivial solution of (1.17) belongs to \mathcal{N} . Similarly $\mathcal{N} \neq \emptyset$. We set

$$A := \inf_{\vec{\mathbf{u}} \in \mathcal{N}} I(\vec{\mathbf{u}}) = \inf_{\vec{\mathbf{u}} \in \mathcal{N}} \frac{1}{N} \int_{\mathbb{R}^N} \sum_{i=1}^k |\nabla u_i|^2. \quad (1.23)$$

Then we have the following theorem, which plays an important role in the proof of Theorem 1.4.

Theorem 1.6. *Assume that $N \geq 5$ and (1.4). If $\mu_i > 0$ and $\beta_{ij} > 0$, then system (1.17) has a positive least energy solution $\vec{\mathbf{U}}$ with $I(\vec{\mathbf{U}}) = A$, which is radially symmetric decreasing. Moreover,*

- (1) *the positive least energy solution of (1.17) must be the least energy synchronized type solution of the form $(\tilde{t}_1 U_{\varepsilon,y}, \tilde{t}_2 U_{\varepsilon,y}, \dots, \tilde{t}_k U_{\varepsilon,y})$, where $(\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$ is defined in (1.10).*
- (2) *there exists $\beta_1 > 0$, and for any $0 < \beta_{ij} = \beta < \beta_1$, there exists a solution $(t_1(\beta), t_2(\beta), \dots, t_k(\beta))$ of (1.9), such that*

$$I(t_1(\beta) U_{\varepsilon,y}, t_2(\beta) U_{\varepsilon,y}, \dots, t_k(\beta) U_{\varepsilon,y}) > A = I(\vec{\mathbf{U}}).$$

That is, $(t_1(\beta) U_{\varepsilon,y}, t_2(\beta) U_{\varepsilon,y}, \dots, t_k(\beta) U_{\varepsilon,y})$ is a different positive solution of (1.17) with respect to $\vec{\mathbf{U}}$.

Remark 1.4.

- (1) *The multiplicity of solution of algebraic system (1.9) arises much more difficulty when dealing with the classification problem.*

For the special case $\beta_{ij} = \beta > 0$, Wu proved that the positive least energy solution of (1.17) must be of synchronized type $(t_1 U_{\varepsilon,y}, t_2 U_{\varepsilon,y}, \dots, t_k U_{\varepsilon,y})$ where (t_1, t_2, \dots, t_k) is solves (1.9). (See [49, Theorem 1.2, Proposition 3.2].)

We point out that the solution (t_1, t_2, \dots, t_k) of (1.9) might not attain d_k , then $(t_1 U_{\varepsilon,y}, t_2 U_{\varepsilon,y}, \dots, t_k U_{\varepsilon,y})$ might not be of the least energy, since the method of Lagrange multiplier yields only a necessary condition for constrained problems.

It is interesting that Theorem 1.6 (2) coincides with the multiplicity of solution of algebraic system (1.9), and admits the existence of another positive solution of (1.17) which is not of the least energy.

- (2) Theorem 1.6 (1) generalizes [19, Theorem 1.6] in the sense that it asserts that the positive least energy of (1.17) must be the synchronized type solution for the purely cooperative case $\beta_{ij} > 0$.
- (3) We note that a similar existence result of ground state (which seems probably to be semi-trivial) is obtained by He and Yang [28] via concentration-compactness lemma. We remark that our method is quite different from that. We illustrate the positive least energy of synchronized type and any other result concerning (1.17) cannot be found in [28].

In rest of this paper we prove these theorems. In Sect. 2, we obtain the fundamental Lemma 2.2 on the existence of positive solution of the algebraic system (1.9) with arbitrary $\beta_{ij} > 0$, which generalizes [49, Proposition 3.2] and gives out a detailed and geometric explanation. Theorem 1.1 and 1.2 are proved subsequently. Since the k -coupled case is much more delicatated than 2-couple case, we shall introduce *the idea of induction* in the proof of Theorem 1.4 and 1.6, respectively in Sect. 3 and 4. *The key idea* is to give a more accurate upper bound of the least energy, which is inspired by [19, 29]. See Lemma 3.2 and 4.2. We mention that the idea of induction is also introduced in [20, 42, 48] to obtain the positive least energy solution of the purely competitive system. Theorem 1.5 is a byproduct of Lemma 4.2 and Theorem 1.4. Theorem 1.3 is proved in Sect. 4.

2 Preliminaries

In this section, we introduce some fundamental results and prove Theorem 1.1 and Theorem 1.2.

Lemma 2.1. *(the Pohozaev identity)*

If Ω is star-shaped with respect to some point $y_0 \in \mathbb{R}^N$ and $\vec{u} \in H$ be a solution of (1.1), then \vec{u} satisfies

$$\begin{aligned} & \frac{1}{2p} \int_{\Omega} \sum_{i=1}^k |\nabla u_i|^2 dx + \frac{1}{2N} \int_{\partial\Omega} \sum_{i=1}^k \left| \frac{\partial u_i}{\partial \vec{n}} \right|^2 (x - y_0) \cdot \vec{n} dS \\ &= \frac{1}{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) dx - \frac{1}{2} \int_{\Omega} \sum_{i=1}^k \lambda_i u_i^2 dx, \end{aligned} \quad (2.1)$$

where \vec{n} denotes the unit outward normal.

Proof. The Pohozaev identity (2.1) follows by multiplying i th equation

$$-\Delta u_i + \lambda_i u_i = \mu_i u_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{p-1} u_j^p$$

by $x \cdot \nabla u_i$ and integrating by parts. \square

First we prove Theorem 1.1, which shows the nonexistence of positive solution of system (1.1) under some assumptions.

Proof of Theorem 1.1 (1) Let ϕ_1 be the first eigenfunction of $-\Delta$ with the Dirichlet boundary condition with respect to $\lambda_1(\Omega)$.

By multiplying each equations in (1.1) with ϕ_1 and integrating over Ω , we obtain

$$(\lambda_1(\Omega) + \lambda_i) \int_{\Omega} u_i \phi_1 dx = \int_{\Omega} (\mu_i u_i^{2p-1} \phi_1 + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{p-1} u_j^p \phi_1) dx.$$

It is easy to see that when $\lambda_i \leq -\lambda_1(\Omega)$, $\mu_i > 0$ and $\beta_{ij} > 0$ for $i, j = 1, 2, \dots, k$, system (1.1) has no positive solutions.

(2) Assume that Ω is star-shaped with respect to some point $y_0 \in \mathbb{R}^N$ and $\vec{u} = (u_1, u_2, \dots, u_k)$ be a solution of (1.1), then \vec{u} satisfies

$$\int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) dx = \int_{\Omega} (\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i,j=1, i < j}^k 2\beta_{ij} |u_i|^p |u_j|^p) dx.$$

If $\lambda_i \geq 0$, combined with the Pohozaev identity (2.1) we have

$$0 \leq \frac{1}{2N} \int_{\partial\Omega} \sum_{i=1}^k \left| \frac{\partial u_i}{\partial \vec{n}} \right|^2 (x - y_0) \cdot \vec{n} dS = \left(\frac{1}{2p} - \frac{1}{2} \right) \int_{\Omega} \sum_{i=1}^k \lambda_i u_i^2 dx \leq 0,$$

then system (1.1) has no positive solutions. \square

Now we turn to study the existence of solution to the algebraic system (1.9), which may play a fundamental role in the subsequent research. We generalize [49, Proposition 3.2] to arbitrary $\beta_{ij} > 0$. Our proof is more specific and gives out a geometric explanation.

Lemma 2.2. *Assume $\mu_i > 0$ and $\beta_{ij} > 0$, then algebraic system*

$$\begin{cases} \mu_i t_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} t_i^{p-1} t_j^p = t_i, & \text{for } i = 1, 2, \dots, k, \\ t_i \in \mathbb{R}, & \text{for } i = 1, 2, \dots, k \end{cases}$$

has a nontrivial positive solution.

Proof. Let

$$d_k := \inf_{\vec{t} \in \mathcal{P}_k} \mathcal{G}_k(\vec{t}), \quad (2.2)$$

where

$$\mathcal{G}_k(\vec{t}) = \sum_{i=1}^k \left(\frac{t_i^2}{2} - \frac{\mu_i |t_i|^{2p}}{2p} \right) - \frac{1}{2p} \left(\sum_{i=1, j > i}^k 2\beta_{ij} |t_i|^p |t_j|^p \right),$$

$$\mathcal{P}_k = \{\vec{\mathbf{t}} \in \mathbb{R}^k \setminus \{\vec{\mathbf{0}}\} \mid P_k(\vec{\mathbf{t}}) := \sum_{i=1}^k (t_i^2 - \mu_i |t_i|^{2p}) - \sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p = 0\}.$$

Denote $\vec{\mathbf{e}}_i = (0, \dots, 0, \mu_i^{-\frac{1}{2p-2}}, 0, \dots, 0)$, it is easy to check that $\vec{\mathbf{e}}_i \in \mathcal{P}_k$ for all $i = 1, 2, \dots, k$, which implies $\mathcal{P}_k \neq \emptyset$ and the minimizing problem (2.2) is well defined. Notice that

$$\begin{aligned} d_k &= \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \mathcal{G}_k(\vec{\mathbf{t}}) = \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \frac{1}{N} \sum_{i=1}^k t_i^2 \\ &= \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \frac{1}{N} \left(\text{dist}(\vec{\mathbf{0}}, \vec{\mathbf{t}}) \right)^2. \end{aligned}$$

Since $\mathcal{P}_k \neq \emptyset$ and symmetric, the minimizing problem (2.2) is reduced to a geometric problem, that is, to find the closest point to original point $\vec{\mathbf{0}}$ in a nonempty set \mathcal{P}_k . Then d_k can be attained by some $\vec{\mathbf{t}}$ with $\tilde{t}_i \geq 0$ for all $i = 1, 2, \dots, k$, and $\tilde{t}_i > 0$ for some i .

By the method of Lagrange's multiplier, there exists a Lagrange multiplier $\gamma \in \mathbb{R}$ such that

$$\nabla \mathcal{G}_k(\vec{\mathbf{t}}) - \gamma \nabla P_k(\vec{\mathbf{t}}) = 0, \quad (2.3)$$

then $\vec{\mathbf{t}}$ satisfies the system

$$\begin{cases} \mu_i \tilde{t}_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} \tilde{t}_i^{p-1} \tilde{t}_j^p = \tilde{t}_i, & \text{for } i = 1, 2, \dots, k, \\ \tilde{t}_i \geq 0, \quad \text{and} \quad \sum_{i=1}^k \tilde{t}_i > 0, & \text{for } i = 1, 2, \dots, k. \end{cases} \quad (2.4)$$

In fact, for any nonzero $\vec{\mathbf{t}} \in \mathcal{P}_k$, we assume that $\vec{\mathbf{t}} = (t_1, \dots, t_k)$ with $t_s \neq 0$ for $1 \leq s \leq k$, we see that

$$t_i \partial_i P_k(\vec{\mathbf{t}}) = 2t_i^2 - 2p\mu_i t_i^{2p} - p \sum_{j=1, j \neq i}^k 2\beta_{ij} t_i^p t_j^p,$$

then we have

$$\begin{aligned} \sum_{i=1}^s t_i \partial_i P_k(\vec{\mathbf{t}}) &= 2 \sum_{i=1}^s t_i^2 - 2p \sum_{i=1}^s \mu_i t_i^{2p} - p \sum_{i=1}^s \sum_{j=1, j \neq i}^k 2\beta_{ij} t_i^p t_j^p \\ &= 2 \sum_{i=1}^s t_i^2 - 2p \sum_{i=1}^s \mu_i t_i^{2p} - 2p \sum_{i=1, j>i}^s 2\beta_{ij} t_i^p t_j^p \\ &= (2 - 2p) \sum_{i=1}^s t_i^2 \neq 0, \end{aligned}$$

which implies that if $\vec{\mathbf{t}} \in \mathcal{P}_k$ then $\nabla P_k(\vec{\mathbf{t}}) \neq \vec{\mathbf{0}}$. Then by (2.3), we have

$$\partial_i \mathcal{G}_k(\vec{\mathbf{t}}) - \gamma \partial_i P_k(\vec{\mathbf{t}}) = 0.$$

Thus $\partial_i \mathcal{G}_k(\vec{\mathbf{t}}) = 0$ if $\tilde{t}_i = 0$, while

$$0 = P_k(\vec{\mathbf{t}}) = \sum_{i=1}^s \tilde{t}_i \partial_i \mathcal{G}_k(\vec{\mathbf{t}}) = \gamma \sum_{i=1}^s \tilde{t}_i \partial_i P_k(\vec{\mathbf{t}}),$$

if $\tilde{t}_1, \dots, \tilde{t}_s \neq 0$, that is

$$(2 - 2p)\gamma \sum_{i=1}^s \tilde{t}_i^2 = 0,$$

which implies that $\gamma = 0$ and $\partial_i \mathcal{G}_k(\vec{\mathbf{t}}) = 0$ for $\tilde{t}_i \neq 0$. Hence, $\vec{\mathbf{t}}$ satisfies (2.4).

We claim that if $\beta_{ij} > 0$, then $d_k < \min\{d_1, d_2, \dots, d_{k-1}\}$.

Our proof is inspired by [1]. By the method of induction, it suffices to prove that $d_k < d_{k-1}$. Assume that d_{k-1} is attained by (t_1, \dots, t_{k-1}) , then

$$d_{k-1} = \frac{1}{N} \sum_{i=1}^{k-1} t_i^2 = \frac{1}{N} \left(\sum_{i=1}^{k-1} \mu_i |t_i|^{2p} + \sum_{i=1, j>i}^{k-1} 2\beta_{ij} |t_i|^p |t_j|^p \right).$$

Note that for any $s \in \mathbb{R}$, there exists $f(s) > 0$ such that $f(s)(t_1, \dots, t_{k-1}, sl) \in \mathcal{P}_k$, where $l \in \mathbb{R} \setminus \{0\}$. In fact,

$$f(s)^{2p-2} = \frac{Nd_{k-1} + s^2 l^2}{Nd_{k-1} + \mu_k |sl|^{2p} + \sum_{i=1}^{k-1} 2\beta_{ik} |t_i|^p |sl|^p}.$$

By direct computations we have

$$\lim_{s \rightarrow 0} \frac{f'(s)}{|s|^{p-2} s} = -\frac{p}{(p-1)Nd_{k-1}} \sum_{i=1}^{k-1} \beta_{ik} |t_i|^p |l|^p,$$

that is,

$$f'(s) = -\frac{p}{(p-1)Nd_{k-1}} \left(\sum_{i=1}^{k-1} \beta_{ik} |t_i|^p |l|^p \right) |s|^{p-2} s (1 + o(1)), \quad \text{as } s \rightarrow 0.$$

Note that $f(0) = 1$, then

$$f(s) = 1 - \frac{1}{(p-1)Nd_{k-1}} \left(\sum_{i=1}^{k-1} \beta_{ik} |t_i|^p |l|^p \right) |s|^p (1 + o(1)), \quad \text{as } s \rightarrow 0.$$

This implies that

$$\begin{aligned} f(s)^{2p} &= 1 - \frac{2p}{(p-1)Nd_{k-1}} \left(\sum_{i=1}^{k-1} \beta_{ik} |t_i|^p |l|^p \right) |s|^p (1 + o(1)) \\ &= 1 - \frac{1}{d_{k-1}} \left(\sum_{i=1}^{k-1} \beta_{ik} |t_i|^p |l|^p \right) |s|^p (1 + o(1)) \quad \text{as } s \rightarrow 0, \end{aligned}$$

then we see that

$$\begin{aligned} d_k &\leq \mathcal{G}_k(f(s)t_1, \dots, f(s)t_{k-1}, f(s)sl) \\ &= \frac{1}{N} f(s)^{2p} \left(Nd_{k-1} + \mu_k |sl|^{2p} + \sum_{i=1}^{k-1} 2\beta_{ik} |t_i|^p |sl|^p \right) \\ &= d_{k-1} - \left(\frac{1}{2} - \frac{1}{N} \right) \left(\sum_{i=1}^{k-1} 2\beta_{ik} |t_i|^p |l|^p \right) |s|^p + o(|s|^p) \\ &< d_{k-1} \quad \text{as } |s| > 0 \text{ small enough.} \end{aligned}$$

By the idea of induction, we see that $d_k < \min\{d_1, d_2, \dots, d_{k-1}\}$. Next we prove that if $\beta_{ij} > 0$, then d_k is attained by $(\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$, where $\tilde{t}_i > 0$ for all $i = 1, 2, \dots, k$. Without loss of generality, we assume by contradiction that $\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_{k-1} > 0$, and $\tilde{t}_k = 0$. Denote

$$\vec{\mathbf{t}}_k = (\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_{k-1}, 0), \quad \vec{\mathbf{t}}_{k-1} = (\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_{k-1}),$$

then $\vec{\mathbf{t}}_{k-1} \in \mathcal{P}_{k-1}$. Notice that

$$\begin{aligned} d_k &= \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \mathcal{G}_k(\vec{\mathbf{t}}) = \mathcal{G}_k(\vec{\mathbf{t}}_k) \\ &= \mathcal{G}_{k-1}(\vec{\mathbf{t}}_{k-1}) \geq \inf_{\vec{\mathbf{t}} \in \mathcal{P}_{k-1}} \mathcal{G}_{k-1}(\vec{\mathbf{t}}) = d_{k-1}, \end{aligned} \tag{2.5}$$

while we see that

$$d_k < \min\{d_1, d_2, \dots, d_{k-1}\},$$

a contradiction. Hence we conclude that all $\tilde{t}_i > 0$. \square

Now assume that $N \geq 4$, $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \dots = \lambda_k := \lambda < 0$. We shall start to prove Theorem 1.2.

Proof of Theorem 1.2 In the special case $N = 4$, where $2p = 2^* = 4$, the nonlinear algebraic problem (1.9) reduces to

$$\mu_i t_i^2 + \sum_{j=1, j \neq i}^k \beta_{ij} t_j^2 = 1, \quad \text{for } i = 1, 2, \dots, k,$$

which can be seen as

$$\mathcal{R} \begin{pmatrix} t_1^2 \\ t_2^2 \\ \vdots \\ t_k^2 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix}, \quad (2.6)$$

where $\mathcal{R} = (\beta_{ij})_{k \times k}$ is the coupling matrix.

It is easy to check that if \mathcal{R} is invertible and the sum of each column of \mathcal{R}^{-1} is greater than 0, then (2.6) has solutions $(\pm t_1, \pm t_2, \dots, \pm t_k)$.

We can construct nontrivial solutions of (1.1) as

$$(\pm t_1 \omega, \pm t_2 \omega, \dots, \pm t_k \omega), \quad (2.7)$$

where ω is the positive least energy solution of (1.7). Remark that these solutions are signed solutions and $(t_1 \omega, t_2 \omega, \dots, t_k \omega)$ is a positive solution of (1.1). Moreover, it's proved in [27, Theorem 1.1] that $(t_1 \omega, t_2 \omega, \dots, t_k \omega)$ is a positive least energy solution.

In the general case $N \geq 5$, Lemma 2.2 admits a positive solution of the nonlinear algebraic problem (1.9), we can also construct a positive solution of (1.1) as

$$(t_1 \omega, t_2 \omega, \dots, t_k \omega),$$

where ω is the positive least energy solution of (1.7) and (t_1, t_2, \dots, t_k) solves (1.9).

Moreover, we recall that in [5, 12], the authors obtained a nontrivial solution for the Brézis-Nirenberg problem (1.7) if $N \geq 4$ and $\lambda < 0$. Note that (1.7) has a positive least energy solution if $-\lambda_1(\Omega) < \lambda < 0$. While any nontrivial solution of (1.7) is sign-changing if $\lambda \leq -\lambda_1(\Omega)$ (See [39].) Thus, we observe that if $\lambda \leq -\lambda_1(\Omega)$, we can also construct a sign-changing solution of (1.1) as

$$(t_1 \bar{u}, t_2 \bar{u}, \dots, t_k \bar{u}),$$

where \bar{u} is a nontrivial solution of (1.7) and (t_1, t_2, \dots, t_k) solves (1.9). \square

3 Proof of Theorem 1.6

Recall that

$$A := \inf_{\vec{\mathbf{u}} \in \mathcal{N}} I(\vec{\mathbf{u}}) = \inf_{\vec{\mathbf{u}} \in \mathcal{N}} \frac{1}{N} \int_{\mathbb{R}^N} \sum_{i=1}^k |\nabla u_i|^2,$$

where

$$\mathcal{N} = \{\vec{\mathbf{u}} \in D : u_i \not\equiv 0, I'(\vec{\mathbf{u}})(0, \dots, 0, u_i, 0, \dots, 0) = 0, i = 1, 2, \dots, k\}.$$

Define

$$A' := \inf_{\vec{\mathbf{u}} \in \mathcal{N}'} I(\vec{\mathbf{u}}), \quad (3.1)$$

where

$$\mathcal{N}' = \{\vec{\mathbf{u}} \in D \setminus \{\vec{\mathbf{0}}\} : I'(\vec{\mathbf{u}}) \vec{\mathbf{u}} = 0\}. \quad (3.2)$$

Note that $\mathcal{N} \subset \mathcal{N}'$, one has that $A' \leq A$. By Sobolev inequality (1.18), we have $A, A' > 0$.

Define $B(0, R) := \{x \in \mathbb{R}^N : |x| < R\}$ and $H(0, R) := (H_0^1(B(0, R)))^k$. Consider

$$\begin{cases} -\Delta u_i = \mu_i |u_i|^{2p-2} u_i + \sum_{j=1, j \neq i}^k \beta_{ij} |u_i|^{p-2} u_i |u_j|^p, & x \in B(0, R), \\ u_i \in H_0^1(B(0, R)), & i = 1, 2, \dots, k, \end{cases} \quad (3.3)$$

and define

$$A'(R) := \inf_{\vec{u} \in \mathcal{N}'(R)} I(\vec{u}), \quad (3.4)$$

where

$$\begin{aligned} \mathcal{N}'(R) = \left\{ \vec{u} \in H(0, R) \setminus \{\vec{0}\} : \int_{B(0, R)} \sum_{i=1}^k |\nabla u_i|^2 dx \right. \\ \left. - \int_{B(0, R)} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i, j=1, i < j}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) dx = 0 \right\}. \end{aligned} \quad (3.5)$$

We need the following lemma from [19], which still holds in our case.

Lemma 3.1. $A'(R) \equiv A'$ for all $R > 0$.

Proof. Take any $R_1 > R_2$. By $\mathcal{N}'(R_2) \subset \mathcal{N}'(R_1)$, we have $A'(R_1) \leq A'(R_2)$. On the other hand, for any $(u_1, u_2, \dots, u_k) \in \mathcal{N}'(R_1)$, we define

$$u'_i(x) := \left(\frac{R_1}{R_2}\right)^{\frac{N-2}{2}} u_i\left(\frac{R_1}{R_2}x\right),$$

then it is standard to see that $(u'_1, u'_2, \dots, u'_k) \in \mathcal{N}'(R_2)$, and we have

$$A'(R_2) \leq I(u'_1, u'_2, \dots, u'_k) = I(u_1, u_2, \dots, u_k), \quad \forall (u_1, u_2, \dots, u_k) \in \mathcal{N}'(R_1).$$

That is, $A'(R_2) \leq A'(R_1)$ and so $A'(R_1) = A'(R_2)$.

Clearly $A' \leq A'(R)$. Let $(u_{1n}, u_{2n}, \dots, u_{kn})$ be a minimizing sequence of A' . We may assume that $u_{1n}, u_{2n}, \dots, u_{kn} \in H_0^1(B(0, R_n))$ for some $R_n > 0$. Then $u_{1n}, u_{2n}, \dots, u_{kn} \in \mathcal{N}'(R_n)$, and

$$A' = \lim_{n \rightarrow \infty} I(u_{1n}, u_{2n}, \dots, u_{kn}) \geq \lim_{n \rightarrow \infty} A'(R_n) \equiv A'(R).$$

Therefore, $A'(R) \equiv A'$ for all $R > 0$. \square

Let $0 \leq \varepsilon < p - 1$. Consider

$$\begin{cases} -\Delta u_i = \mu_i |u_i|^{2p-2-2\varepsilon} u_i + \sum_{j=1, j \neq i}^k \beta_{ij} |u_i|^{p-2-\varepsilon} u_i |u_j|^{p-\varepsilon}, & x \in B(0, 1), \\ u_i \in H_0^1(B(0, 1)), & i = 1, 2, \dots, k, \end{cases} \quad (3.6)$$

and define

$$A_\varepsilon := \inf_{\vec{\mathbf{u}} \in \mathcal{N}'_\varepsilon} I_\varepsilon(\vec{\mathbf{u}}), \quad (3.7)$$

where

$$I_\varepsilon(\vec{\mathbf{u}}) := \frac{1}{2} \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i|^2 dx - \frac{1}{2p-2\varepsilon} \int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i|^{2p-2\varepsilon} + \sum_{i,j=1, i < j}^k 2\beta_{ij} |u_i|^{p-\varepsilon} |u_j|^{p-\varepsilon} \right) dx, \quad (3.8)$$

$$\mathcal{N}'_\varepsilon = \{ \vec{\mathbf{u}} \in H(0,1) \setminus \{ \vec{\mathbf{0}} \} : H_\varepsilon(\vec{\mathbf{u}}) := I'_\varepsilon(\vec{\mathbf{u}}) \vec{\mathbf{u}} = 0 \}. \quad (3.9)$$

We shall introduce **the idea of induction**. We notice that the proof of Lemma 3.2 depends on the existence result Theorem 3.1 for the $(k-1)$ -coupled system case.

From now on, we assume that Lemma 3.2 and Theorem 3.1 hold true for $(k-1)$ -coupled system. We shall give out the proof of Lemma 3.2 and Theorem 3.1 for k -coupled system in the sequel. Then by idea of induction, these results hold true for arbitrary k -coupled system.

Denote $\hat{\mathbf{u}}_i \in H(0,1)$ as an vector, where the number of nonzero components of $\hat{\mathbf{u}}_i$ is i for $i = 1, 2, \dots, k$. Denote

$$\begin{aligned} C_\varepsilon(i) &:= \min \{ \inf_{\hat{\mathbf{u}}_i \in \mathcal{N}'_\varepsilon} I_\varepsilon(\hat{\mathbf{u}}_i) \} \\ &= \min \left\{ \inf_{(u_1, \dots, u_i, 0, \dots, 0) \in \mathcal{N}'_\varepsilon} I_\varepsilon(u_1, \dots, u_i, 0, \dots, 0), \dots, \right. \\ &\quad \left. \inf_{(0, \dots, 0, u_1, \dots, u_i) \in \mathcal{N}'_\varepsilon} I_\varepsilon(0, \dots, 0, u_1, \dots, u_i) \right\}. \end{aligned}$$

For the case $k = 2$, $C_\varepsilon(1) = \min \{ \inf_{(u,0) \in \mathcal{N}'_\varepsilon} I_\varepsilon(u, 0), \inf_{(0,v) \in \mathcal{N}'_\varepsilon} I_\varepsilon(0, v) \}$, $C_\varepsilon(2) = A_\varepsilon$.

Lemma 3.2. *For any $0 < \varepsilon < p - 1$, there holds*

$$A_\varepsilon < \min \{ C_\varepsilon(1), C_\varepsilon(2), \dots, C_\varepsilon(k-1) \}. \quad (3.10)$$

Proof. Fix any $0 < \varepsilon < p - 1$. By the idea of induction, we only need to prove

$$A_\varepsilon := C_\varepsilon(k) < C_\varepsilon(k-1).$$

Recall that $2 < 2p - 2\varepsilon < 2^*$, we may let $(u_1, u_2, \dots, u_{k-1})$ be a least energy solution of the $(k-1)$ -coupled system (3.6) with energy

$$c_{k-1} := \inf_{(u_1, u_2, \dots, u_{k-1}, 0) \in \mathcal{N}'_\varepsilon} I_\varepsilon(u_1, u_2, \dots, u_{k-1}, 0).$$

We note that for any $s \in \mathbb{R}$, there exists a unique $t(s) > 0$ such that

$$(t(s)u_1, t(s)u_2, \dots, t(s)u_{k-1}, t(s)s\phi) \in \mathcal{N}'_\varepsilon.$$

where $\phi \in H_0^1(\Omega) \setminus \{0\}$. In fact,

$$\begin{aligned} t(s)^{2p-2\varepsilon-2} & \left(\int_{B(0,1)} \left(\sum_{i=1}^{k-1} \mu_i |u_i|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^{k-1} 2\beta_{ij} |u_i|^{p-\varepsilon} |u_j|^{p-\varepsilon} \right. \right. \\ & \left. \left. + \mu_k |s\phi|^{2p-2\varepsilon} + \sum_{i=1}^{k-1} 2\beta_{ik} |u_i|^{p-\varepsilon} |s\phi|^{p-\varepsilon} \right) \right) \\ & = \int_{B(0,1)} \left(\sum_{i=1}^{k-1} |\nabla u_i|^2 + s^2 |\nabla \phi|^2 \right), \end{aligned}$$

Note that $t(0) = 1$. Recall that $0 < p - \varepsilon < 2$, by direct computations we have

$$\lim_{s \rightarrow 0} \frac{t'(s)}{|s|^{p-2-\varepsilon} s} = -\frac{p-\varepsilon}{(p-\varepsilon-1)p'c_{k-1}} \int_{B(0,1)} \sum_{i=1}^{k-1} \beta_{ik} |u_i|^{p-\varepsilon} |\phi|^{p-\varepsilon},$$

where $p' = \frac{2p-2\varepsilon}{p-1-\varepsilon}$. That is,

$$t'(s) = -\frac{p-\varepsilon}{(p-\varepsilon-1)p'c_{k-1}} \left(\int_{B(0,1)} \sum_{i=1}^{k-1} \beta_{ik} |u_i|^{p-\varepsilon} |\phi|^{p-\varepsilon} \right) |s|^{p-\varepsilon-2} s (1 + o(1)),$$

as $s \rightarrow 0$ and so

$$t(s) = 1 - \frac{1}{(p-\varepsilon-1)p'c_{k-1}} \left(\int_{B(0,1)} \sum_{i=1}^{k-1} \beta_{ik} |u_i|^{p-\varepsilon} |\phi|^{p-\varepsilon} \right) |s|^{p-\varepsilon} (1 + o(1)), \quad \text{as } s \rightarrow 0.$$

This implies that

$$\begin{aligned} t(s)^{2p-2\varepsilon} & = 1 - \frac{2p-2\varepsilon}{(p-\varepsilon-1)p'c_{k-1}} \left(\int_{B(0,1)} \sum_{i=1}^{k-1} \beta_{ik} |u_i|^{p-\varepsilon} |\phi|^{p-\varepsilon} \right) |s|^{p-\varepsilon} (1 + o(1)), \\ & = 1 - \frac{1}{c_{k-1}} \left(\int_{B(0,1)} \sum_{i=1}^{k-1} \beta_{ik} |u_i|^{p-\varepsilon} |\phi|^{p-\varepsilon} \right) |s|^{p-\varepsilon} (1 + o(1)) \quad \text{as } s \rightarrow 0, \end{aligned}$$

then we see that

$$\begin{aligned}
A_\varepsilon &\leq I_\varepsilon(t(s)u_1, \dots, t(s)u_{k-1}, t(s)s\phi) \\
&= \frac{1}{p'} t(s)^{2p-2\varepsilon} \left(p' c_{k-1} + \int_{B(0,1)} (\mu_k |s\phi|^{2p-2\varepsilon} + \sum_{i=1}^{k-1} 2\beta_{ik} |u_i|^{p-\varepsilon} |s\phi|^{p-\varepsilon}) \right) \\
&= c_{k-1} - \left(\frac{1}{2} - \frac{1}{p'} \right) |s|^{p-\varepsilon} \int_{B(0,1)} \sum_{i=1}^{k-1} 2\beta_{ik} |u_i|^{p-\varepsilon} |\phi|^{p-\varepsilon} + o(|s|^{p-\varepsilon}) \\
&< c_{k-1} \quad \text{as } |s| > 0 \text{ small enough.}
\end{aligned}$$

Then by similar arguments, we have

$$A_\varepsilon < C_\varepsilon(k-1).$$

By the idea of induction, we have $C_\varepsilon(k-1) < C_\varepsilon(k-2) < \dots < C_\varepsilon(1)$. This completes the proof. \square

Denote

$$\begin{aligned}
C(i) &:= \min\{ \inf_{\hat{\mathbf{u}}_i \in \mathcal{N}'} I(\hat{\mathbf{u}}_i) \} \\
&= \min \left\{ \inf_{(u_1, \dots, u_i, 0, \dots, 0) \in \mathcal{N}'} I(u_1, \dots, u_i, 0, \dots, 0), \dots, \right. \\
&\quad \left. \inf_{(0, \dots, 0, u_1, \dots, u_i) \in \mathcal{N}'} I(0, \dots, 0, u_1, \dots, u_i) \right\}.
\end{aligned}$$

Similarly as Lemma 3.2, we have

$$\begin{aligned}
A' &< \min\{C(1), C(2), \dots, C(k-1)\} \\
&< C(1) \\
&= \min\{I(\omega_{\mu_1}, 0, \dots, 0), \dots, I(0, \dots, 0, \omega_{\mu_k})\} \\
&= \min\left\{ \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2}, \dots, \frac{1}{N} \mu_k^{-\frac{N-2}{2}} S^{N/2} \right\}.
\end{aligned} \tag{3.11}$$

Theorem 3.1. *For any $0 < \varepsilon < p-1$, (3.6) has a classical least energy solution $\vec{u}_\varepsilon = (u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon)$, and $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon \in C^2(B(0,1))$ are all positive radially symmetric decreasing.*

Proof. Fix any $0 < \varepsilon < p-1$, it is easy to see that $A_\varepsilon > 0$. For $(u_1, u_2, \dots, u_k) \in \mathcal{N}'_\varepsilon$ with $u_1, u_2, \dots, u_k \geq 0$, we denote by $(u_1^*, u_2^*, \dots, u_k^*)$ as its Schwarz symmetrization. Then by the properties of Schwarz symmetrization and $\beta_{ij} > 0$, we have

$$\begin{aligned}
\int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^*|^2 dx &\leq \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i|^2 dx \\
&= \int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_i|^{p-\varepsilon} |u_j|^{p-\varepsilon} \right) dx \\
&\leq \int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i^*|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_i^*|^{p-\varepsilon} |u_j^*|^{p-\varepsilon} \right) dx.
\end{aligned} \tag{3.12}$$

Recall (3.9), there exists $t^* > 0$ such that $(t^* u_1^*, t^* u_2^*, \dots, t^* u_k^*) \in \mathcal{N}'_\varepsilon$ with

$$(t^*)^{2p-2\varepsilon-2} = \frac{\int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^*|^2 dx}{\int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i^*|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_i^*|^{p-\varepsilon} |u_j^*|^{p-\varepsilon} \right) dx} \leq 1,$$

and then

$$\begin{aligned}
I_\varepsilon(t^* u_1^*, t^* u_2^*, \dots, t^* u_k^*) &= \left(\frac{1}{2} - \frac{1}{2p-2\varepsilon} \right) (t^*)^2 \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^*|^2 dx \\
&\leq \left(\frac{1}{2} - \frac{1}{2p-2\varepsilon} \right) \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i|^2 dx \\
&= I_\varepsilon(u_1, u_2, \dots, u_k).
\end{aligned} \tag{3.13}$$

Therefore, we may take a minimizing sequence $\vec{\mathbf{u}}_n = (u_{1n}, u_{2n}, \dots, u_{kn}) \in \mathcal{N}'_\varepsilon$ such that $(u_{1n}, u_{2n}, \dots, u_{kn}) = (u_{1n}^*, u_{2n}^*, \dots, u_{kn}^*)$ and $I_\varepsilon(\vec{\mathbf{u}}_n) \rightarrow A_\varepsilon$. We see from (3.13) $u_{1n}, u_{2n}, \dots, u_{kn}$ are uniformly bounded in $H_0^1(B(0,1))$. Passing to a subsequence, we may assume that $u_{in} \rightharpoonup u_i^\varepsilon$ weakly in $H_0^1(B(0,1))$. By the compactness of the embedding $H_0^1(B(0,1)) \hookrightarrow L^{2p-2\varepsilon}(B(0,1))$ we have

$$\begin{aligned}
&\int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i^\varepsilon|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_i^\varepsilon|^{p-\varepsilon} |u_j^\varepsilon|^{p-\varepsilon} \right) dx \\
&= \lim_{n \rightarrow \infty} \int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_{in}|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_{in}|^{p-\varepsilon} |u_{jn}|^{p-\varepsilon} \right) dx \\
&= \frac{2p-2\varepsilon}{p-1-\varepsilon} \lim_{n \rightarrow \infty} I_\varepsilon(\vec{\mathbf{u}}_n) = \frac{2p-2\varepsilon}{p-1-\varepsilon} A_\varepsilon > 0,
\end{aligned}$$

which implies $(u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon) \neq (0, 0, \dots, 0)$. Moreover, $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon \geq 0$ are radially symmetric. Meanwhile,

$$\int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^\varepsilon|^2 dx \leq \lim_{n \rightarrow \infty} \int_{B(0,1)} \sum_{i=1}^k |\nabla u_{in}|^2 dx,$$

then

$$\int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^\varepsilon|^2 dx \leq \int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i^\varepsilon|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_i^\varepsilon|^{p-\varepsilon} |u_j^\varepsilon|^{p-\varepsilon} \right) dx.$$

Therefore, there exists $0 < t_\varepsilon \leq 1$ such that $t_\varepsilon \vec{\mathbf{u}}_\varepsilon = (t_\varepsilon u_1^\varepsilon, t_\varepsilon u_2^\varepsilon, \dots, t_\varepsilon u_k^\varepsilon) \in \mathcal{N}'_\varepsilon$, and then

$$\begin{aligned} A_\varepsilon &\leq I_\varepsilon(t_\varepsilon \vec{\mathbf{u}}_\varepsilon) \\ &= \left(\frac{1}{2} - \frac{1}{2p-2\varepsilon} \right) (t_\varepsilon)^2 \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^\varepsilon|^2 dx \\ &\leq \lim_{n \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{2p-2\varepsilon} \right) \int_{B(0,1)} \sum_{i=1}^k |\nabla u_{in}|^2 dx \\ &= \lim_{n \rightarrow \infty} I_\varepsilon(\vec{\mathbf{u}}_n) = A_\varepsilon. \end{aligned}$$

Therefore, $t_\varepsilon = 1$ and $\vec{\mathbf{u}}_\varepsilon \in \mathcal{N}'_\varepsilon$ with $I(\vec{\mathbf{u}}_\varepsilon) = A_\varepsilon$. Moreover,

$$\int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^\varepsilon|^2 dx = \lim_{n \rightarrow \infty} \int_{B(0,1)} \sum_{i=1}^k |\nabla u_{in}|^2 dx,$$

that is, $u_{in} \rightarrow u_i^\varepsilon$ strongly in $H_0^1(B(0,1))$. There exists a Lagrange multiplier $\gamma \in \mathbb{R}$ such that

$$I'_\varepsilon(\vec{\mathbf{u}}_\varepsilon) - \gamma H'_\varepsilon(\vec{\mathbf{u}}_\varepsilon) = 0.$$

Since $I'_\varepsilon(\vec{\mathbf{u}}_\varepsilon) \vec{\mathbf{u}}_\varepsilon = H'_\varepsilon(\vec{\mathbf{u}}_\varepsilon) = 0$ and

$$H'_\varepsilon(\vec{\mathbf{u}}_\varepsilon) \vec{\mathbf{u}}_\varepsilon = (2+2\varepsilon-2p) \int_{B(0,1)} \left(\sum_{i=1}^k \mu_i |u_i^\varepsilon|^{2p-2\varepsilon} + \sum_{i,j=1, i<j}^k 2\beta_{ij} |u_i^\varepsilon|^{p-\varepsilon} |u_j^\varepsilon|^{p-\varepsilon} \right) dx < 0,$$

we get that $\gamma = 0$ and so $I'_\varepsilon(\vec{\mathbf{u}}_\varepsilon) = 0$. By Lemma 3.2, we see that $u_i^\varepsilon \not\equiv 0$, otherwise $\vec{\mathbf{u}}_\varepsilon$ cannot minimize $I_\varepsilon(\vec{\mathbf{u}})$. This means that $\vec{\mathbf{u}}_\varepsilon$ is a least energy solution of (3.6). Recall that $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon \geq 0$ are radially symmetric non-increasing. By regularity theory and the maximum principle, we see that $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon > 0$ in $B(0,1)$ and $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon \in C^2(B(0,1))$ are radially symmetric decreasing. \square

Now we start to prove Theorem 1.6.

Proof of Theorem 1.6 Recall the definition of $\mathcal{N}'(R)$, for any $\vec{\mathbf{u}} \in \mathcal{N}'(1)$, there exists $t_\varepsilon > 0$ such that $t_\varepsilon \vec{\mathbf{u}} = (t_\varepsilon u_1, t_\varepsilon u_2, \dots, t_\varepsilon u_k) \in \mathcal{N}'_\varepsilon$ with $t_\varepsilon \rightarrow 1$ as $\varepsilon \rightarrow 0$. Then we have

$$\limsup_{\varepsilon \rightarrow 0} A_\varepsilon \leq \limsup_{\varepsilon \rightarrow 0} I_\varepsilon(t_\varepsilon \vec{\mathbf{u}}) = I(\vec{\mathbf{u}}), \quad \forall \vec{\mathbf{u}} \in \mathcal{N}'(1).$$

By Lemma 3.1 we have

$$\limsup_{\varepsilon \rightarrow 0} A_\varepsilon \leq A'(1) = A'. \quad (3.14)$$

By Theorem 3.1, let $\vec{u}_\varepsilon = (u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon)$ be a positive least energy solution of (3.6), which is radially symmetric decreasing.

Fix $0 < \varepsilon \leq \frac{p-1}{2}$. Then by $I'_\varepsilon(\vec{u}_\varepsilon) = 0$ and Sobolev inequality, we see that

$$\frac{2p-2\varepsilon}{p-\varepsilon-1} A_\varepsilon = \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^\varepsilon|^2 dx \geq C_0, \quad \forall 0 < \varepsilon \leq \frac{p-1}{2}, \quad (3.15)$$

where C_0 is a positive constant independent of ε . Then $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon$ are uniformly bounded in $H_0^1(B(0,1))$. Passing to a subsequence, we assume that $u_i^\varepsilon \rightharpoonup u_i^0$ weakly in $H_0^1(B(0,1))$ for $i = 1, 2, \dots, k$. Then $\vec{u}_0 = (u_1^0, u_2^0, \dots, u_k^0)$ is a solution of

$$\begin{cases} -\Delta u_i = \mu_i |u_i|^{2p-2} u_i + \sum_{j=1, j \neq i}^k \beta_{ij} |u_i|^{p-2} u_i |u_j|^p, & x \in B(0,1), \\ u_i \in H_0^1(B(0,1)), & i = 1, 2, \dots, k. \end{cases} \quad (3.16)$$

Assume by contradiction that $\sum_{i=1}^k \|u_i^\varepsilon\|_\infty$ is uniformly bounded, then by the Dominated Convergence Theorem, we get that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{B(0,1)} |u_i^\varepsilon|^{2p-2\varepsilon} dx &= \int_{B(0,1)} |u_i^0|^{2p} dx, \\ \lim_{\varepsilon \rightarrow 0} \int_{B(0,1)} |u_i^\varepsilon|^{p-\varepsilon} |u_j^\varepsilon|^{p-\varepsilon} dx &= \int_{B(0,1)} |u_i^0|^p |u_j^0|^p dx. \end{aligned}$$

Combining these with $I'_\varepsilon(\vec{u}_\varepsilon) = I'(\vec{u}_0) = 0$, we have that $u_i^\varepsilon \rightarrow u_i^0$ strongly in $H_0^1(B(0,1))$ for $i = 1, 2, \dots, k$.

Then by (3.15), we see that $\vec{u}_0 \neq \vec{0}$. Moreover $u_1^0, u_2^0, \dots, u_k^0 \geq 0$. We may assume $u_1^0 \not\equiv 0$. By the strong maximum principle, $u_1^0 > 0$ in $B(0,1)$. Note that $2p = 2^*$. Combining these with the Pohozaev identity (2.1), we have

$$0 < \int_{\partial B(0,1)} \sum_{i=1}^k |\nabla u_i|^2 (x \cdot \vec{n}) dS = 0,$$

a contradiction. Here, \vec{n} denotes the outward unit normal vector on $\partial B(0,1)$. Therefore,

we deduce that $\sum_{i=1}^k \|u_i^\varepsilon\|_\infty \rightarrow +\infty$ as $\varepsilon \rightarrow 0$.

Next we will use a blow up analysis.

Note that $u_1^\varepsilon, u_2^\varepsilon, \dots, u_k^\varepsilon$ are radially symmetric decreasing, then $u_i^\varepsilon(0) = \max_{B(0,1)} u_i^\varepsilon(x)$.

We define $K_\varepsilon := \max_i \{u_i^\varepsilon(0)\}$, then $K_\varepsilon \rightarrow +\infty$ as $\varepsilon \rightarrow 0$. Define

$$U_i^\varepsilon(x) = K_\varepsilon^{-1} u_i^\varepsilon(K_\varepsilon^{-\alpha_\varepsilon} x),$$

where $\alpha_\varepsilon = p - 1 - \varepsilon$. Then

$$1 = \max_i \{U_i^\varepsilon(0)\} = \max_i \left\{ \max_{x \in B(0, K_\varepsilon^{\alpha_\varepsilon})} U_i^\varepsilon(x) \right\}, \quad (3.17)$$

and $U_1^\varepsilon, U_2^\varepsilon, \dots, U_k^\varepsilon$ satisfy the following system:

$$-\Delta U_i^\varepsilon = \mu_i (U_i^\varepsilon)^{2p-2\varepsilon-1} + \sum_{j=1, j \neq i}^k \beta_{ij} (U_i^\varepsilon)^{p-1-\varepsilon} (U_j^\varepsilon)^{p-\varepsilon}, \quad x \in B(0, K_\varepsilon^{\alpha_\varepsilon}). \quad (3.18)$$

Note that $0 < \varepsilon \leq \frac{p-1}{2}$, then

$$\int_{\mathbb{R}^N} |\nabla U_i^\varepsilon|^2 dx = K_\varepsilon^{-(N-2)\varepsilon} \int_{\mathbb{R}^N} |\nabla u_i^\varepsilon|^2 dx \leq \int_{\mathbb{R}^N} |\nabla u_i^\varepsilon|^2 dx,$$

then $\{(U_1^\varepsilon, U_2^\varepsilon, \dots, U_k^\varepsilon)\}_{n \geq 1}$ is bounded in D . By elliptic estimates, for a subsequence we have $(U_1^\varepsilon, U_2^\varepsilon, \dots, U_k^\varepsilon) \rightarrow (U_1, U_2, \dots, U_k) \in D$ uniformly in every compact subset of \mathbb{R}^N as $\varepsilon \rightarrow 0$, and (U_1, U_2, \dots, U_k) satisfies (1.17). Then we see that $I'(U_1, U_2, \dots, U_k) = 0$. Moreover, $U_1, U_2, \dots, U_k \geq 0$ are radially symmetric decreasing.

By (3.17) we have $(U_1, U_2, \dots, U_k) \neq (0, \dots, 0)$, and so $(U_1, U_2, \dots, U_k) \in \mathcal{N}'$. Then we deduce from (3.14) that

$$\begin{aligned} A' &\leq I(U_1, U_2, \dots, U_k) = \left(\frac{1}{2} - \frac{1}{2p}\right) \int_{\mathbb{R}^N} \sum_{i=1}^k |\nabla U_i|^2 dx \\ &\leq \liminf_{\varepsilon \rightarrow 0} \left(\frac{1}{2} - \frac{1}{2p-2\varepsilon}\right) \int_{B(0, K_\varepsilon^{\alpha_\varepsilon})} \sum_{i=1}^k |\nabla U_i^\varepsilon|^2 dx \\ &\leq \liminf_{\varepsilon \rightarrow 0} \left(\frac{1}{2} - \frac{1}{2p-2\varepsilon}\right) \int_{B(0,1)} \sum_{i=1}^k |\nabla u_i^\varepsilon|^2 dx \\ &= \liminf_{\varepsilon \rightarrow 0} A_\varepsilon \leq A'. \end{aligned}$$

This implies that $I(U_1, U_2, \dots, U_k) = A'$. By (3.11) we have that $U_i \not\equiv 0$ for all $i = 1, 2, \dots, k$. By strong maximum principle, $U_1, U_2, \dots, U_k > 0$ are radially symmetric decreasing. Notice that $(U_1, U_2, \dots, U_k) \in \mathcal{N}$ and so $I(U_1, U_2, \dots, U_k) \geq A \geq A'$, that is,

$$I(U_1, U_2, \dots, U_k) = A' = A, \quad (3.19)$$

and (U_1, U_2, \dots, U_k) is a positive least energy solution of (1.17), which is radially symmetric decreasing.

Now we shall prove that the positive least energy solution of (1.17) must be the least energy synchronized type solution of the form $(t_1\omega, t_2\omega, \dots, t_k\omega)$. Recall the minimizing problem

$$A' := \inf_{\vec{\mathbf{u}} \in \mathcal{N}'} I(\vec{\mathbf{u}}),$$

where

$$\mathcal{N}' = \{\vec{\mathbf{u}} \in D \setminus \{\vec{\mathbf{0}}\} : I'(\vec{\mathbf{u}}) \vec{\mathbf{u}} = 0\},$$

and the minimizing problem introduced in the proof of Lemma 2.2:

$$d_k := \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \mathcal{G}_k(\vec{\mathbf{t}}),$$

where

$$\mathcal{G}_k(\vec{\mathbf{t}}) = \sum_{i=1}^k \left(\frac{t_i^2}{2} - \frac{\mu_i |t_i|^{2p}}{2p} \right) - \frac{1}{2p} \left(\sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p \right),$$

$$\mathcal{P}_k = \{\vec{\mathbf{t}} \in \mathbb{R}^k \setminus \{\vec{\mathbf{0}}\} \mid P_k(\vec{\mathbf{t}}) := \sum_{i=1}^k (t_i^2 - \mu_i |t_i|^{2p}) - \sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p = 0\}.$$

By standard argument (cf. [36]), we see that

$$d_k = \inf_{\vec{\mathbf{t}} \in \mathbb{R}^k \setminus \{\vec{\mathbf{0}}\}} \frac{\left(\sum_{i=1}^k t_i^2 \right)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i |t_i|^{2p} - \sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p \right)^{\frac{N-2}{2}}},$$

and by (3.19) we have

$$A = A' = \inf_{\vec{\mathbf{u}} \in D \setminus \{\vec{\mathbf{0}}\}} \frac{\left(\sum_{i=1}^k \|\nabla u_i\|_2^2 \right)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i \|u_i\|_{2p}^{2p} + \sum_{i,j=1, i<j}^k 2\beta_{ij} \|u_i\|_{2p}^p \|u_j\|_{2p}^p \right)^{\frac{N-2}{2}}}, \quad (3.20)$$

then by Sobolev inequality (1.18), we have

$$A' \geq \inf_{\vec{\mathbf{u}} \in D \setminus \{\vec{\mathbf{0}}\}} \frac{\left(\sum_{i=1}^k \|u_i\|_{2p}^2 \right)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i \|u_i\|_{2p}^{2p} + \sum_{i,j=1, i<j}^k 2\beta_{ij} \|u_i\|_{2p}^p \|u_j\|_{2p}^p \right)^{\frac{N-2}{2}}} S^{\frac{N}{2}},$$

where $\|\cdot\|_q$ denotes the norm on $L^q(\mathbb{R}^N)$ and $2p = 2^* = \frac{2N}{N-2}$. Recall Lemma 2.2, then we see that

$$A' = d_k S^{\frac{N}{2}}$$

is attained by

$$(t_1 U_{\varepsilon, y}, t_2 U_{\varepsilon, y}, \dots, t_k U_{\varepsilon, y}),$$

where (t_1, t_2, \dots, t_k) is the positive solution of (1.9) that attains d_k , and $U_{\varepsilon, y}$ is the Aubin-Talenti instanton given by (1.19).

In fact, we deduce that

$$A' = \inf_{\vec{\mathbf{u}} \in D \setminus \{\vec{\mathbf{0}}\}} \frac{\left(\sum_{i=1}^k \|u_i\|_{2p}^2 \right)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i \|u_i\|_{2p}^{2p} + \sum_{i,j=1, i<j}^k 2\beta_{ij} \|u_i\|_{2p}^p \|u_j\|_{2p}^p \right)^{\frac{N-2}{2}}} S^{\frac{N}{2}}. \quad (3.21)$$

Suppose that A' is attained by some nonzero $\vec{v} = (v_1, v_2, \dots, v_k)$. Then by Hölder inequality and Sobolev inequality, we see from (3.20) and (3.21) that

$$\|\nabla v_i\|_2^2 = S\|v_i\|_{2p}^2$$

for all $i = 1, 2, \dots, k$, which implies $v_i = c_i U_{\varepsilon, y}$ for some $c_i \neq 0, \varepsilon > 0$ and $y \in \mathbb{R}^N$, or $v_i = 0$. Redenote all $v_i = c_i U_{\varepsilon, y}$ for all i where $c_i = 0$ or $c_i \neq 0$.

By (3.21), we have that $(\|v_1\|_{2p}, \|v_2\|_{2p}, \dots, \|v_k\|_{2p})$ attains d_k . It is easy to see that (c_1, c_2, \dots, c_k) attains d_k . Recall Lemma 2.2 that d_k is attained by a positive solution if $\beta_{ij} > 0$. Then $c_i > 0$ for all $i = 1, 2, \dots, k$.

That is, A' is attained by \vec{u} if and only if

$$\vec{u} = (\tilde{t}_1 U_{\varepsilon, y}, \tilde{t}_2 U_{\varepsilon, y}, \dots, \tilde{t}_k U_{\varepsilon, y}),$$

where $(\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$ is defined in (1.10). We see that, if $\vec{v} \in \mathcal{N}$ attains A , then $\vec{v} \in \mathcal{N}'$ and by (3.19) we have

$$I(\vec{v}) = A = A',$$

which means \vec{v} attains A' . Since A' is attained only by $\vec{u} = (\tilde{t}_1 U_{\varepsilon, y}, \tilde{t}_2 U_{\varepsilon, y}, \dots, \tilde{t}_k U_{\varepsilon, y})$, which solves (1.17), then $\vec{u} \in \mathcal{N}$, by

$$I(\vec{u}) = A = A',$$

we see that \vec{u} also attains A . Hence, the positive least energy solution of (1.17) must be the least energy synchronized type $(\tilde{t}_1 U_{\varepsilon, y}, \tilde{t}_2 U_{\varepsilon, y}, \dots, \tilde{t}_k U_{\varepsilon, y})$, where $(\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$ is defined in (1.10).

Finally, we show the existence of $(t_1(\beta), t_2(\beta), \dots, t_k(\beta))$ for $\beta_{ij} = \beta > 0$ small. Denote functions

$$\begin{aligned} f_i(t_1, t_2, \dots, t_k, \beta) &:= \mu_i t_i^{2p-2} \\ &+ \sum_{j=1, j \neq i}^k \beta t_i^{p-2} t_j^p - 1, \quad t_i > 0, \text{ and } t_j \geq 0 \text{ for } j \neq i, \end{aligned} \quad (3.22)$$

and define $t_i(0) = \mu_i^{-\frac{1}{2p-2}}$, then $f_i(t_1(0), t_2(0), \dots, t_k(0), 0) = 0$ for all $i = 1, 2, \dots, k$. Note that

$$\begin{aligned} \partial_i f_i(t_1(0), t_2(0), \dots, t_k(0), 0) &= (2p-2)\mu_i(t_i(0))^{2p-3} > 0, \\ \partial_j f_i(t_1(0), t_2(0), \dots, t_k(0), 0) &= 0, \quad \text{if } j \neq i \text{ and } j = 1, 2, \dots, k, \end{aligned}$$

which implies that

$$\det(F_{ij})_{k \times k} := \det(\partial_j f_i(t_1(0), t_2(0), \dots, t_k(0), 0))_{k \times k} > 0.$$

Therefore, by the implicit function theorem, $(t_1(\beta), t_2(\beta), \dots, t_k(\beta))$ are well defined and class C^1 on $(-\beta_2, \beta_2)$ for some $\beta_2 > 0$, and

$$f_i(t_1(\beta), t_2(\beta), \dots, t_k(\beta), \beta) \equiv 0, \quad i = 1, 2, \dots, k.$$

This implies that $(t_1(\beta), t_2(\beta), \dots, t_k(\beta))$ solves (1.9). Moreover, we notice that $(t_1(\beta)U_{\varepsilon,y}, t_2(\beta)U_{\varepsilon,y}, \dots, t_k(\beta)U_{\varepsilon,y})$ is a positive solution of (1.17) and

$$\lim_{\beta \rightarrow 0} \sum_{i=1}^k t_i^2(\beta) = \sum_{i=1}^k t_i^2(0) = \sum_{i=1}^k \mu_i^{-\frac{1}{p-1}},$$

that is, there exists $0 < \beta_1 \leq \beta_2$, such that

$$\sum_{i=1}^k t_i^2(\beta) > \min_i \{\mu_i^{-\frac{1}{p-1}}\}, \quad \forall \beta \in (0, \beta_1).$$

Combining this with (3.11), we have

$$I(\vec{U}) = A' = A < I(t_1(\beta)U_{\varepsilon,y}, t_2(\beta)U_{\varepsilon,y}, \dots, t_k(\beta)U_{\varepsilon,y}), \quad \forall \beta \in (0, \beta_1),$$

that is, $(t_1(\beta)U_{\varepsilon,y}, t_2(\beta)U_{\varepsilon,y}, \dots, t_k(\beta)U_{\varepsilon,y})$ is a different positive solution of (1.17) with respect to \vec{U} . This completes the proof. \square

Similarly to [19, Proposition 2.1], the following properties still hold in our case.

Proposition 3.1. *Assume that $\mu_i > 0$ and $\beta_{ij} > 0$. Let $\vec{U} = (U_1, U_2, \dots, U_k)$ be a positive radially symmetric least energy solution of (1.17) obtained in Theorem 1.6. Then there exists $C > 0$ such that*

$$\sum_{i=1}^k U_i(x) \leq C(1 + |x|)^{2-N}, \quad \sum_{i=1}^k |\nabla U_i(x)| \leq C(1 + |x|)^{2-N}.$$

Proof. Define the Kelvin transformation:

$$U_i^*(x) := |x|^{2-N} U_i(x^*), \quad x^* = \frac{x}{|x|^2},$$

then $U_1^*, \dots, U_k^* \in D^{1,2}(\mathbb{R}^N)$ and (U_1^*, \dots, U_k^*) satisfies the same system (1.17). By a standard Brézis-Kato argument [10], we see that $U_1^*, \dots, U_k^* \in L^\infty(\mathbb{R}^N)$. Therefore, there exists $C > 0$ such that

$$U_i^*(x^*) = |x|^{N-2} U_i(x) \leq C,$$

then

$$\sum_{i=1}^k U_i(x) \leq C|x|^{2-N}. \quad (3.23)$$

On the other hand, we note that U_1, U_2, \dots, U_k are radially symmetric decreasing. We also have $U_1, U_2, \dots, U_k \in L^\infty(\mathbb{R}^N)$, and then

$$\sum_{i=1}^k U_i(x) \leq C(1 + |x|)^{2-N}.$$

Moreover, by standard elliptic regularity theory, we have that $U_1, U_2, \dots, U_k \in C^2(\mathbb{R}^N)$. We write $U_i(|x|) = U_i(x)$ for convenience. Then

$$(r^{N-1}(U_i)_r)_r = -r^{N-1}(\mu_i U_i^{2^*-1} + \sum_{j=1, j \neq i}^k \beta_{ij} U_i^{\frac{2^*}{2}-1} U_j^{\frac{2^*}{2}}),$$

and so for any $R \geq 1$, we see from (3.23) that

$$\begin{aligned} R^{N-1} |(U_i)_r(R)| &\leq |(U_i)_r(1)| + \int_1^R r^{N-1} (\mu_i U_i^{2^*-1} + \sum_{j=1, j \neq i}^k \beta_{ij} U_i^{\frac{2^*}{2}-1} U_j^{\frac{2^*}{2}}) dr \\ &\leq C + C \int_1^{+\infty} r^{N-1} r^{-N-2} dr \leq C. \end{aligned}$$

Therefore, it is easy to see that $|\nabla U_i(x)| \leq C(1 + |x|)^{2-N}$ for some $C > 0$. Then there exists $C > 0$ such that

$$\sum_{i=1}^k |\nabla U_i(x)| \leq C(1 + |x|)^{2-N}.$$

□

4 Proof of Theorem 1.4

In this section, without loss of generality, we assume that

$$-\lambda_1(\Omega) < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k < 0.$$

Recall the definition of B in (1.6), since

$$\int_{\Omega} (|\nabla u|^2 + \lambda_i u^2) dx \geq (1 + \frac{\lambda_i}{\lambda_1(\Omega)}) \int_{\Omega} |\nabla u|^2 dx, \quad i = 1, 2, \dots, k,$$

it is standard to see that $B > 0$. Recall [11] that the Brézis-Nirenberg problem

$$-\Delta u + \lambda_i u = \mu_i |u|^{2^*-2} u, \quad u \in H_0^1(\Omega) \quad (4.1)$$

has a positive least energy solution $u_{\mu_i} \in C^2(\Omega) \cap C(\bar{\Omega})$ with energy

$$\begin{aligned} B_{\mu_i} &:= \frac{1}{2} \int_{\Omega} (|\nabla u_{\mu_i}|^2 + \lambda_i u_{\mu_i}^2) dx - \frac{\mu_i}{2^*} \int_{\Omega} u_{\mu_i}^{2^*} dx, \\ &< \frac{1}{N} \mu_i^{-\frac{N-2}{2}} S^{N/2}, \quad i = 1, 2, \dots, k. \end{aligned} \quad (4.2)$$

We need the following lemma from [19].

Lemma 4.1. *Let $u_n \rightharpoonup u, v_n \rightharpoonup v$ in $H_0^1(\Omega)$ as $n \rightarrow \infty$, then passing to a subsequence, there holds*

$$\lim_{n \rightarrow \infty} \int_{\Omega} (|u_n|^p |v_n|^p - |u_n - u|^p |v_n - v|^p - |u|^p |v|^p) dx = 0.$$

Remark 4.1. *Similar to the proof of Theorem 1.6 in Sect. 3, before starting the proof, we shall introduce **the idea of induction** again. From now on, we assume that Theorem 1.4 hold true for $(k-1)$ -coupled system, that is,*

the $(k-1)$ -coupled system has a positive least energy solution.

We shall give out the proof of these results for k -coupled system in the sequel. Then by the idea of induction, these theorems hold true for arbitrary k -coupled system.

For convinience, here are some notations. Consider the k -coupled system (1.1), where the coefficients $\lambda_i, \mu_i, \beta_{ij}$ are fixed. For each $\Theta \subsetneq \{1, \dots, k\}$, we obtain the following $(k - |\Theta|)$ -coupled system

$$\begin{cases} -\Delta u_i + \lambda_i u_i = \mu_i u_i^{2p-1} + \sum_{j=1, j \neq i}^k \beta_{ij} u_i^{p-1} u_j^p & \text{in } \Omega, \\ u_i > 0 \text{ in } \Omega \quad \text{and} \quad u_i = 0 \text{ on } \partial\Omega, \quad i, j \in \{1, 2, \dots, k\} \setminus \Theta, \end{cases} \quad (4.3)$$

by replacing $\lambda_i, \mu_i, \beta_{ij}, \beta_{ji}$ with 0 if $i \in \Theta$, where $|\Theta|$ is the cardinality of Θ . Note that the nontrivial solutions of (4.3) can be extended to the semi-trivial solutions of k -coupled system (1.1).

Denote $\tau = k - |\Theta|$. Indeed, we can establish C_k^τ different τ -coupled systems from (1.1). For every fixed Θ , we denote $\{i_1, \dots, i_\tau\} = \{1, 2, \dots, k\} \setminus \Theta$ and $B_{\mu_{i_1} \dots \mu_{i_\tau}}$ as the least energy of the τ -coupled system (4.3) with coefficients $(\mu_{i_1}, \dots, \mu_{i_\tau})$, that is,

$$B_{\mu_{i_1} \dots \mu_{i_\tau}} := \inf \{E(\vec{u}) : \vec{u} = (u_1, \dots, u_k) \text{ solves (1.1) and } u_i = 0 \text{ iff } i \in \Theta\}. \quad (4.4)$$

Denote

$$\bar{B}_\tau := \min_{\forall \{l_1, \dots, l_\tau\} \subsetneq \{1, 2, \dots, k\}} \{B_{\mu_{l_1} \dots \mu_{l_\tau}}\}. \quad (4.5)$$

Define

$$\mathcal{B} := \inf_{h \in \Gamma} \max_{t \in [0, 1]} E(h(t)), \quad (4.6)$$

where $\Gamma = \{h \in C([0, 1], H) : h(0) = \vec{\mathbf{0}}, E(h(1)) < 0\}$. By (1.5), we see that for any $\vec{u} \in H, \vec{u} \neq \vec{\mathbf{0}}$,

$$\begin{aligned} \max_{t > 0} E(t\vec{u}) &= E(t\vec{u}) \\ &= \frac{1}{N} t_{\vec{u}}^2 \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) \\ &= \frac{1}{N} t_{\vec{u}}^{2^*} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j > i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right), \end{aligned} \quad (4.7)$$

where $t_{\vec{\mathbf{u}}} > 0$ satisfies

$$t_{\vec{\mathbf{u}}}^{2p-2} = \frac{\int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2)}{\int_{\Omega} (\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p)}. \quad (4.8)$$

Note that $t_{\vec{\mathbf{u}}} \vec{\mathbf{u}} \in \mathcal{M}'$, where

$$\mathcal{M}' := \left\{ \vec{\mathbf{u}} \in H \setminus \{\vec{\mathbf{0}}\} : G(\vec{\mathbf{u}}) := \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) - \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) = 0 \right\}. \quad (4.9)$$

Note that $\mathcal{M} \subset \mathcal{M}'$, one has that $0 < \mathcal{B} \leq B$. It is easy to check that

$$\mathcal{B} = \inf_{\vec{\mathbf{0}} \neq \vec{\mathbf{u}} \in H} \max_{t>0} E(t\vec{\mathbf{u}}) = \inf_{\vec{\mathbf{u}} \in \mathcal{M}'} E(\vec{\mathbf{u}}). \quad (4.10)$$

In fact, by (4.7) we have that

$$\inf_{\vec{\mathbf{0}} \neq \vec{\mathbf{u}} \in H} \max_{t>0} E(t\vec{\mathbf{u}}) = \inf_{\vec{\mathbf{u}} \in \mathcal{M}'} E(\vec{\mathbf{u}}),$$

and

$$\begin{aligned} E'(t\vec{\mathbf{u}})(t\vec{\mathbf{u}}) &= t^2 \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) - t^{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) \\ &\geq 0 \quad \text{if } 0 \leq t \leq t_{\vec{\mathbf{u}}}. \end{aligned}$$

On the one hand, we see that $E(t\vec{\mathbf{u}}) < 0$ for $\vec{\mathbf{u}} \in H \setminus \{\vec{\mathbf{0}}\}$ and $t > 0$ large enough. Then by the definition of \mathcal{B} , we have

$$\mathcal{B} \leq \inf_{\vec{\mathbf{0}} \neq \vec{\mathbf{u}} \in H} \max_{t>0} E(t\vec{\mathbf{u}}).$$

On the other hand, \mathcal{M}' separates H into two components

$$\begin{aligned} \mathcal{M}'_1 &= \{\vec{\mathbf{u}} \in H \setminus \{\vec{\mathbf{0}}\} : E'(\vec{\mathbf{u}})\vec{\mathbf{u}} > 0\} \cup \{\vec{\mathbf{0}}\}, \\ \mathcal{M}'_2 &= \{\vec{\mathbf{u}} \in H \setminus \{\vec{\mathbf{0}}\} : E'(\vec{\mathbf{u}})\vec{\mathbf{u}} < 0\}. \end{aligned}$$

Since $E'(t\vec{\mathbf{u}})(t\vec{\mathbf{u}}) \geq 0$ if $0 \leq t \leq t_{\vec{\mathbf{u}}}$, we have

$$\mathcal{M}'_1 = \{t\vec{\mathbf{u}} : \vec{\mathbf{u}} \in H \setminus \{\vec{\mathbf{0}}\}, 0 \leq t < t_{\vec{\mathbf{u}}}\}, \quad \mathcal{M}'_2 = \{t\vec{\mathbf{u}} : \vec{\mathbf{u}} \in H \setminus \{\vec{\mathbf{0}}\}, t > t_{\vec{\mathbf{u}}}\}.$$

Note that for any $\vec{\mathbf{0}} \neq t\vec{\mathbf{u}} \in \mathcal{M}'_1$,

$$\begin{aligned} E(t\vec{\mathbf{u}}) &= \frac{t^2}{2} \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) - \frac{t^{2p}}{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) \\ &> \frac{1}{2p} \left[t^2 \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda_i u_i^2) - t^{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) \right] \\ &= \frac{1}{2p} E'(t\vec{\mathbf{u}})(t\vec{\mathbf{u}}) > 0, \quad \text{if } 0 < t < t_{\vec{\mathbf{u}}}. \end{aligned}$$

then every $h \in \Gamma$ has to cross \mathcal{M}' and $\inf_{\vec{u} \in \mathcal{M}'} E(\vec{u}) \leq \mathcal{B}$. It follows that (4.10) holds.

Next we introduce the key lemma in the proof of Theorem 1.4 which gives out an estimate of \mathcal{B} .

Lemma 4.2. (the comparison lemma) *Let $\beta_{ij} > 0$, then*

$$\mathcal{B} < \min\{\bar{B}_1, \bar{B}_2, \dots, \bar{B}_{k-1}, A\}.$$

Proof. We will prove the lemma in three steps.

Step 1 ($\mathcal{B} < A$)

Without loss of generality, we may assume that $0 \in \Omega$. Then there exists $\rho > 0$ such that $B(0, 2\rho) := \{x : |x| \leq 2\rho\} \subset \Omega$. Let $\psi \in C_0^1(B(0, 2\rho))$ be a nonnegative function with $0 \leq \psi \leq 1$ and $\psi \equiv 1$ for $|x| \leq \rho$. Recall that $\vec{U} = (U_1, U_2, \dots, U_k)$ in Theorem 1.6. We define

$$U_i^\varepsilon(x) := \varepsilon^{-\frac{N-2}{2}} U_i(x/\varepsilon), \quad u_i^\varepsilon := \psi U_i^\varepsilon, \quad i = 1, 2, \dots, k.$$

Then it is easy to see that

$$\int_{\mathbb{R}^N} |\nabla U_i^\varepsilon|^2 = \int_{\mathbb{R}^N} |\nabla U_i|^2, \quad \int_{\mathbb{R}^N} |U_i^\varepsilon|^{2^*} = \int_{\mathbb{R}^N} |U_i|^{2^*}, \quad i = 1, 2, \dots, k.$$

It's proved in [19] that

$$\int_{\Omega} |\nabla u_i^\varepsilon|^2 \leq \int_{\mathbb{R}^N} |\nabla U_i|^2 + O(\varepsilon^{N-2}), \quad (4.11)$$

$$\int_{\Omega} |u_i^\varepsilon|^{2^*} \geq \int_{\mathbb{R}^N} |U_i|^{2^*} + O(\varepsilon^N), \quad (4.12)$$

$$\int_{\Omega} |u_i^\varepsilon|^{\frac{2^*}{2}} |u_j^\varepsilon|^{\frac{2^*}{2}} \geq \int_{\mathbb{R}^N} |U_i|^{\frac{2^*}{2}} |U_j|^{\frac{2^*}{2}} + O(\varepsilon^N), \quad (4.13)$$

$$\int_{\Omega} |u_i^\varepsilon|^2 \geq C\varepsilon^2 + O(\varepsilon^{N-2}), \quad (4.14)$$

where C is a positive constant. Recall that $I(U_1, U_2, \dots, U_k) = A$, we have

$$NA = \int_{\mathbb{R}^N} \sum_{i=1}^k |\nabla U_i|^2 = \int_{\mathbb{R}^N} \left(\sum_{i=1}^k \mu_i |U_i|^{2^*} + \sum_{i=1, j>i}^k 2\beta_{ij} |U_i|^{\frac{2^*}{2}} |U_j|^{\frac{2^*}{2}} \right).$$

Combining this with (4.11)-(4.14) and recalling that $\lambda_i < 0$, $2p = 2^*$, $N \geq 5$, we have for any $t > 0$ that

$$\begin{aligned}
E(tu_1^\varepsilon, \dots, tu_k^\varepsilon) &= \frac{1}{2}t^2 \int_{\Omega} \sum_{i=1}^k (|\nabla u_i^\varepsilon|^2 + \lambda_i (u_i^\varepsilon)^2) \\
&\quad - \frac{1}{2p}t^{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i^\varepsilon|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i^\varepsilon|^p |u_j^\varepsilon|^p \right) \\
&\leq \frac{1}{2} \left(\int_{\mathbb{R}^N} \sum_{i=1}^k |\nabla U_i|^2 - C\varepsilon^2 + O(\varepsilon^{N-2}) \right) t^2 \\
&\quad - \frac{1}{2^*} \left(\int_{\mathbb{R}^N} \sum_{i=1}^k \mu_i |U_i|^{2^*} + \sum_{i=1, j>i}^k 2\beta_{ij} |U_i|^{\frac{2^*}{2}} |U_j|^{\frac{2^*}{2}} \right) + O(\varepsilon^N) t^{2^*} \\
&= \frac{1}{2} \left(NA - C\varepsilon^2 + O(\varepsilon^{N-2}) \right) t^2 - \frac{1}{2^*} \left(NA + O(\varepsilon^N) \right) t^{2^*} \\
&\leq \frac{1}{N} \left(NA - C\varepsilon^2 + O(\varepsilon^{N-2}) \right) \left(\frac{NA - C\varepsilon^2 + O(\varepsilon^{N-2})}{NA + O(\varepsilon^N)} \right)^{\frac{N-2}{2}} \\
&< A \quad \text{for } \varepsilon > 0 \text{ small enough.}
\end{aligned}$$

Hence, for $\varepsilon > 0$ small enough, there holds

$$\mathcal{B} \leq \max_{t>0} E(tu_1^\varepsilon, \dots, tu_k^\varepsilon) < A. \quad (4.15)$$

From now on, we redenote $\mathcal{B}_k := \mathcal{B}$ as the minimax value defined in (4.6) for the k -coupled system.

Step 2 ($k = 3$ and $\mathcal{B}_3 < \min\{\bar{B}_1, \bar{B}_2\}$)

In [19], it's proved that the 2-coupled system

$$\begin{cases} -\Delta u + \lambda_i u = \mu_i u^{2p-1} + \beta_{ij} u^{p-1} v^p, & x \in \Omega, \\ -\Delta v + \lambda_j v = \mu_j v^{2p-1} + \beta_{ij} v^{p-1} u^p, & x \in \Omega, \\ u, v \geq 0 \text{ in } \Omega \quad \text{and} \quad u = v = 0 \text{ on } \partial\Omega, \end{cases} \quad (4.16)$$

has a positive least energy solution (u_{ij}, v_{ij}) with energy $B_{\mu_i \mu_j}$ and

$$B_{\mu_i \mu_j} < \min\{B_{\mu_i}, B_{\mu_j}\}, \quad (4.17)$$

then $\bar{B}_2 = \min\{B_{\mu_1 \mu_2}, B_{\mu_1 \mu_3}, B_{\mu_2 \mu_3}\} < \bar{B}_1$, it remains to prove $\mathcal{B}_3 < \bar{B}_2$.

We shall prove $\mathcal{B}_3 < B_{\mu_1 \mu_2}$.

Recall (4.8), we note that for any $s \in \mathbb{R}$, there exists a unique $t(s) := t_{u_{12}, v_{12}, s\phi}$

such that $(t(s)u_{12}, t(s)v_{12}, t(s)s\phi) \in \mathcal{M}'$ and

$$\begin{aligned} & t(s)^{2p-2} \left(\int_{\Omega} (\mu_1 u_{12}^{2p} + \mu_2 v_{12}^{2p} + \mu_3 |s|^{2p} \phi^{2p}) \right. \\ & \quad \left. + 2\beta_{12} u_{12}^p v_{12}^p + 2\beta_{13} |s|^p u_{12}^p \phi^p + 2\beta_{23} |s|^p v_{12}^p \phi^p \right) \\ & = \int_{\Omega} (|\nabla u_{12}|^2 + \lambda_1 u_{12}^2 + |\nabla v_{12}|^2 + \lambda_2 v_{12}^2 + s^2 |\nabla \phi|^2 + \lambda_3 s^2 \phi^2), \end{aligned}$$

where (u_{12}, v_{12}) is a positive least energy solution of (4.16) and $\phi \in H_0^1(\Omega)$ is a positive function. Recall that

$$\begin{aligned} NB_{\mu_1 \mu_2} & = \int_{\Omega} (\mu_1 u_{12}^{2p} + 2\beta_{12} u_{12}^p v_{12}^p + \mu_2 v_{12}^{2p}) \\ & = \int_{\Omega} (|\nabla u_{12}|^2 + \lambda_1 u_{12}^2 + |\nabla v_{12}|^2 + \lambda_2 v_{12}^2) := c_{12}, \end{aligned}$$

and we denote

$$\begin{aligned} g(s) & := t(s)^{2p-2} \\ & = \frac{c_{12} + s^2 \int_{\Omega} (|\nabla \phi|^2 + \lambda_3 \phi^2)}{c_{12} + \int_{\Omega} (\mu_3 |s|^{2p} \phi^{2p} + 2\beta_{13} |s|^p u_{12}^p \phi^p + 2\beta_{23} |s|^p v_{12}^p \phi^p)} := \frac{A(s)}{B(s)}. \end{aligned} \quad (4.18)$$

Note that $t'(s) = \frac{1}{2p-2} g(s)^{\frac{3-2p}{2p-2}} g'(s)$ and

$$\lim_{s \rightarrow 0} g(s) = g(0) = t(0) = 1, \quad \lim_{s \rightarrow 0} A(s) = \lim_{s \rightarrow 0} B(s) = c_{12}. \quad (4.19)$$

By direct computations we have

$$g'(s) = \frac{A'(s)}{B(s)} - \frac{A(s)B'(s)}{B^2(s)} := F_1(s) - F_2(s).$$

Note that $1 < p < 2$, then

$$\lim_{s \rightarrow 0} \frac{F_1(s)}{|s|^{p-2}s} = \lim_{s \rightarrow 0} \frac{2s^{2-p} \int_{\Omega} (|\nabla \phi|^2 + \lambda_3 \phi^2)}{c_{12} + \int_{\Omega} (\mu_3 |s|^{2p} \phi^{2p} + 2\beta_{13} |s|^p u_{12}^p \phi^p + 2\beta_{23} |s|^p v_{12}^p \phi^p)} = 0. \quad (4.20)$$

On the other hand, we see that

$$\begin{aligned} F_2(s) & = \frac{2p|s|^{p-2}s}{B^2(s)} \left[c_{12} + s^2 \left(\int_{\Omega} |\nabla \phi|^2 + \lambda_3 \phi^2 \right) \right] \\ & \quad \cdot \left(|s|^p \int_{\Omega} \mu_3 \phi^{2p} + \int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p \right), \end{aligned}$$

then

$$\lim_{s \rightarrow 0} \frac{F_2(s)}{|s|^{p-2}s} = \frac{2p}{c_{12}} \int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p. \quad (4.21)$$

Then we have

$$\lim_{s \rightarrow 0} \frac{t'(s)}{|s|^{p-2}s} = -\frac{p}{(p-1)c_{12}} \int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p,$$

that is,

$$t'(s) = -\frac{p}{(p-1)c_{12}} \left(\int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p \right) |s|^{p-2}s (1 + o(1)), \quad \text{as } s \rightarrow 0,$$

and so

$$t(s) = 1 - \frac{1}{(p-1)c_{12}} \left(\int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p \right) |s|^p (1 + o(1)), \quad \text{as } s \rightarrow 0.$$

This implies that

$$t(s)^{2p} = 1 - \frac{2p}{(p-1)c_{12}} \left(\int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p \right) |s|^p (1 + o(1)), \quad \text{as } s \rightarrow 0.$$

Therefore, we deduce from (4.7) that

$$\begin{aligned} \mathcal{B}_3 &\leq E(t(s)u_{12}, t(s)v_{12}, t(s)s\phi) \\ &= \frac{1}{N} t(s)^{2p} \int_{\Omega} (\mu_1 u_{12}^{2p} + \mu_2 v_{12}^{2p} + \mu_3 |s|^{2p} \phi^{2p} \\ &\quad + 2\beta_{12} u_{12}^p v_{12}^p + 2\beta_{13} |s|^p u_{12}^p \phi^p + 2\beta_{23} |s|^p v_{12}^p \phi^p) \\ &= t(s)^{2p} \left[B_{\mu_1 \mu_2} + \frac{1}{N} \int_{\Omega} \mu_3 |s|^{2p} \phi^{2p} + 2\beta_{13} |s|^p u_{12}^p \phi^p + 2\beta_{23} |s|^p v_{12}^p \phi^p \right] \\ &= B_{\mu_1 \mu_2} + \frac{1}{N} \int_{\Omega} \mu_3 |s|^{2p} \phi^{2p} + 2\beta_{13} |s|^p u_{12}^p \phi^p + 2\beta_{23} |s|^p v_{12}^p \phi^p + o(|s|^p) \\ &\quad - \frac{1}{N} \frac{2p}{p-1} |s|^p \int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p \\ &\quad - \frac{1}{N} \frac{2p}{(p-1)c_{12}} |s|^{2p} \left(\int_{\Omega} \beta_{13} u_{12}^p \phi^p + \beta_{23} v_{12}^p \phi^p \right) \left(\int_{\Omega} \mu_3 |s|^p \phi^{2p} + 2\beta_{13} u_{12}^p \phi^p + 2\beta_{23} v_{12}^p \phi^p \right) \\ &= B_{\mu_1 \mu_2} - \left(\frac{1}{2} - \frac{1}{N} \right) |s|^p \int_{\Omega} 2\beta_{13} u_{12}^p \phi^p + 2\beta_{23} v_{12}^p \phi^p + o(|s|^p) \\ &< B_{\mu_1 \mu_2} \quad \text{as } |s| > 0 \text{ small enough.} \end{aligned}$$

That is $\mathcal{B}_3 < B_{\mu_1\mu_2}$. By a similar argument, we can prove that $\mathcal{B}_3 < B_{\mu_i\mu_j}$, then $\mathcal{B}_3 < \bar{B}_2$, which implies that

$$\mathcal{B}_3 < \min\{\bar{B}_1, \bar{B}_2\}.$$

This completes the proof of the case $k = 3$.

Step 3 (for general $k \geq 3$)

Suppose that the $(k-i)$ -coupled system (4.3) with coefficients $(\mu_1, \mu_2, \dots, \mu_{k-i})$ has a positive least energy solution $\mathbf{w}_{k-i} = (\omega_1, \omega_2, \dots, \omega_{k-i})$ with energy $B_{\mu_1 \dots \mu_{k-i}}$. Denote

$$\mathcal{W} := (\omega_1, \omega_2, \dots, \omega_{k-i}, s\phi, s^2\phi, \dots, s^i\phi),$$

where $\phi \in H_0^1(\Omega)$ and $\phi \not\equiv 0$.

Recall (4.8), we note that for any $s \in \mathbb{R}$, there exists a unique $t(s) := t_{\mathcal{W}}$ such that $t(s)(\mathcal{W}) \in \mathcal{M}'$. Then by similar argument as in **Step 2**, we have

$$\begin{aligned} \mathcal{B}_k &\leq E(t(s)(\omega_1, \dots, \omega_{k-i}, s\phi, s^2\phi, \dots, s^i\phi)) \\ &= B_{\mu_1 \dots \mu_{k-i}} - \left(\frac{1}{2} - \frac{1}{N}\right) |s|^p \int_{\Omega} \sum_{j=1}^{k-i} 2\beta_{j, k-i+1} |\omega_j|^p |\phi|^p + o(|s|^p) \\ &< B_{\mu_1 \dots \mu_{k-i}}, \end{aligned}$$

which implies that $\mathcal{B}_k < \bar{B}_{k-i}$. This completes the proof. \square

Let's begin to prove Theorem 1.4.

Proof of Theorem 1.4. Assume that $\beta_{ij} > 0$. Since the functional E has a mountain pass structure, by the mountain pass theorem [4, 47] there exists $\{\bar{\mathbf{u}}_n\} \in H$, such that

$$\lim_{n \rightarrow +\infty} E(\bar{\mathbf{u}}_n) = \mathcal{B}, \quad \lim_{n \rightarrow +\infty} E'(\bar{\mathbf{u}}_n) = 0,$$

where $\bar{\mathbf{u}}_n = (u_{1n}, u_{2n}, \dots, u_{kn})$. By standard argument it is easy to see that $\{\bar{\mathbf{u}}_n\}$ is bounded in H , and so we may assume that $(u_{1n}, \dots, u_{kn}) \rightharpoonup (u_1, \dots, u_k)$ weakly in H . Passing to a subsequence, we may assume that $u_{in} \rightharpoonup u_i$ weakly in $L^{2p}(\Omega)$, $|u_{in}|^{q-1} u_{in} \rightharpoonup |u_i|^{q-1} u_i$ weakly in $L^{2p/q}(\Omega)$ if $1 < q < 2p$, and $u_{in} \rightarrow u_i$ strongly in $L^2(\Omega)$. Since $E'(\bar{\mathbf{u}}_n) \rightarrow 0$, then we have $E'(u_1, \dots, u_k) = 0$.

Set $\sigma_{in} = u_{in} - u_i$, then by Brézis-Lieb Lemma, there holds

$$|u_{in}|_{2p}^{2p} = |u_i|_{2p}^{2p} + |\sigma_{in}|_{2p}^{2p} + o(1), \quad i = 1, 2, \dots, k. \quad (4.22)$$

Note that $\bar{\mathbf{u}}_n \in \mathcal{M}$ and $E'(\bar{\mathbf{u}}_n) = 0$. Then combined with Lemma 4.1, we have

$$\int_{\Omega} |\nabla \sigma_{in}|^2 - \int_{\Omega} (\mu_i |\sigma_{in}|^{2p} + \sum_{j=1, j \neq i}^k \beta_{ij} |\sigma_{in}|^p |\sigma_{jn}|^p) = o(1), \quad (4.23)$$

$$E(u_{1n}, \dots, u_{kn}) = E(u_1, \dots, u_k) + I(\sigma_{1n}, \dots, \sigma_{kn}) + o(1), \quad (4.24)$$

for $i = 1, 2, \dots, k$.

Passing to a subsequence, we may assume that

$$\lim_{n \rightarrow +\infty} \int_{\Omega} |\nabla \sigma_{in}|^2 = b_i, \quad i = 1, 2, \dots, k,$$

then by (4.23) we have

$$I(\sigma_{1n}, \dots, \sigma_{kn}) = \frac{1}{N}(b_1 + \dots + b_k) + o(1).$$

Letting $n \rightarrow +\infty$ in (4.24), we get that

$$\begin{aligned} 0 \leq E(u_1, \dots, u_k) &\leq E(u_1, \dots, u_k) + \frac{1}{N}(b_1 + \dots + b_k) \\ &= \lim_{n \rightarrow +\infty} E(\vec{u}_n) = \mathcal{B}. \end{aligned} \quad (4.25)$$

Case 1. $u_1, u_2, \dots, u_k \equiv 0$.

By (4.25), we have $b_1 + b_2 + \dots + b_k > 0$. Then we assume that $(\sigma_{1n}, \dots, \sigma_{kn}) \neq (0, \dots, 0)$ for n large. Recall the definition of \mathcal{N}' , and by (4.23), it is easy to check that there exists t_n such that $(t_n \sigma_{1n}, \dots, t_n \sigma_{kn}) \in \mathcal{N}'$ and $t_n \rightarrow 1$ as $n \rightarrow +\infty$. Then by (3.19) and (4.25), we have

$$\begin{aligned} \mathcal{B} &= \frac{1}{N} \sum_{i=1}^k b_i = \lim_{n \rightarrow +\infty} I(\sigma_{1n}, \dots, \sigma_{kn}) \\ &= \lim_{n \rightarrow +\infty} I(t_n \sigma_{1n}, \dots, t_n \sigma_{kn}) \\ &\geq A' = A, \end{aligned}$$

a contradiction with Lemma 4.2. Therefore, Case 1 is impossible.

Case 2. $u_i \not\equiv 0$ and $u_j \equiv 0$ for $i \in I \subsetneq \{1, 2, \dots, k\}$ and $j \in \{1, 2, \dots, k\} \setminus I$.

Without loss of generality, we may assume that $u_1, \dots, u_\tau \not\equiv 0, u_{\tau+1}, \dots, u_k \equiv 0$. Then (u_1, \dots, u_τ) is a nontrivial solution of the τ -coupled system, and so

$$\mathcal{B} \geq E(u_1, \dots, u_\tau, 0, \dots, 0) \geq B_{\mu_1 \dots \mu_\tau} \geq \bar{B}_\tau,$$

a contradiction with Lemma 4.2. Therefore, Case 2 is also impossible.

Therefore, $u_1, \dots, u_k \not\equiv 0$.

Since $E'(u_1, \dots, u_k) = 0$, we have $(u_1, \dots, u_k) \in \mathcal{M}$, by $\mathcal{B} \leq B$ and (4.25) we have

$$E(u_1, \dots, u_k) = \mathcal{B} = B. \quad (4.26)$$

This means $(|u_1|, \dots, |u_k|) \in \mathcal{M} \subset \mathcal{M}'$ and $E(|u_1|, \dots, |u_k|) = \mathcal{B} = B$. By (4.9) and (4.10), there exists a Lagrange multiplier $\gamma \in \mathbb{R}$ such that

$$E'(|u_1|, \dots, |u_k|) - \gamma G'(|u_1|, \dots, |u_k|) = 0.$$

Since $E'(|u_1|, \dots, |u_k|)(|u_1|, \dots, |u_k|) = G(|u_1|, \dots, |u_k|) = 0$ and

$$\begin{aligned} & G'(|u_1|, \dots, |u_k|)(|u_1|, \dots, |u_k|) \\ &= (2 - 2p) \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right) < 0, \end{aligned}$$

we get that $\gamma = 0$ and $E'(|u_1|, \dots, |u_k|) = 0$. This means $(|u_1|, \dots, |u_k|)$ is a least energy solution of (1.1). Then by the maximum principle, we see that $|u_1|, \dots, |u_k| > 0$ in Ω . Therefore, $(|u_1|, \dots, |u_k|)$ is a positive least energy solution of (1.1). \square

Proof of Theorem 1.3 Assume that $N \geq 5$, $-\lambda_1(\Omega) < \lambda_1 = \dots = \lambda_k := \lambda < 0$ and $\mu_i, \beta_{ij} > 0$. The following proof is similar to the proof of Theorem 1.6 (1). It is well known that the Brézis-Nirenberg problem [11]

$$-\Delta u + \lambda u = |u|^{2^*-2}u, \quad u \in H_0^1(\Omega)$$

has a positive least energy solution ω that attains

$$S_{\lambda} := \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\int (|\nabla u|^2 + \lambda u^2) dx}{\left(\int |u|^{2^*} dx \right)^{\frac{2}{2^*}}}.$$

By (4.10) and (4.26), we have

$$B := \inf_{\vec{\mathbf{u}} \in \mathcal{M}} E(\vec{\mathbf{u}}) = \inf_{\vec{\mathbf{u}} \in \mathcal{M}'} E(\vec{\mathbf{u}}),$$

where

$$E(\vec{\mathbf{u}}) = \frac{1}{2} \int_{\Omega} \sum_{i=1}^k (|\nabla u_i|^2 + \lambda u_i^2) - \frac{1}{2p} \int_{\Omega} \left(\sum_{i=1}^k \mu_i |u_i|^{2p} + \sum_{i=1, j>i}^k 2\beta_{ij} |u_i|^p |u_j|^p \right),$$

$$\mathcal{M} = \{ \vec{\mathbf{u}} \in H \mid u_i \not\equiv 0, E'(\vec{\mathbf{u}})(0, \dots, u_i, \dots, 0) = 0 \text{ for } i = 1, 2, \dots, k \},$$

$$\mathcal{M}' = \{ \vec{\mathbf{u}} \in H \setminus \{ \vec{\mathbf{0}} \} \mid E'(\vec{\mathbf{u}}) \vec{\mathbf{u}} = 0 \text{ for } i = 1, 2, \dots, k \}.$$

Then by standard argument(cf. [36]), we have

$$B = \inf_{\vec{\mathbf{u}} \in H \setminus \{ \vec{\mathbf{0}} \}} \frac{\left(\sum_{i=1}^k \int (|\nabla u_i|^2 + \lambda u_i^2) \right)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i \int |u_i|^{2p} + \sum_{i,j=1, i<j}^k 2\beta_{ij} \int |u_i u_j|^p \right)^{\frac{N-2}{2}}}. \quad (4.27)$$

Recall the minimizing problem introduced in the proof of Lemma 2.2:

$$d_k := \inf_{\vec{\mathbf{t}} \in \mathcal{P}_k} \mathcal{G}_k(\vec{\mathbf{t}}),$$

where

$$\mathcal{G}_k(\vec{\mathbf{t}}) = \sum_{i=1}^k \left(\frac{t_i^2}{2} - \frac{\mu_i |t_i|^{2p}}{2p} \right) - \frac{1}{2p} \left(\sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p \right),$$

$$\mathcal{P}_k = \{\vec{t} \in \mathbb{R}^k \setminus \{\vec{0}\} \mid P_k(\vec{t}) := \sum_{i=1}^k (t_i^2 - \mu_i |t_i|^{2p}) - \sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p = 0\}.$$

By standard argument (cf. [36]), we see that

$$d_k = \inf_{\vec{t} \in \mathbb{R}^k \setminus \{\vec{0}\}} \frac{(\sum_{i=1}^k t_i^2)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i |t_i|^{2p} - \sum_{i=1, j>i}^k 2\beta_{ij} |t_i|^p |t_j|^p \right)^{\frac{N-2}{2}}},$$

then by Sobolev inequality

$$\int_{\Omega} (|\nabla u|^2 + \lambda u^2) dx \geq S_{\lambda} \left(\int_{\Omega} |u|^{2p} dx \right)^{\frac{2}{2p}},$$

we have

$$B \geq \inf_{\vec{u} \in H \setminus \{\vec{0}\}} \frac{(\sum_{i=1}^k \|u_i\|_{2p}^2)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i \|u_i\|_{2p}^{2p} + \sum_{i,j=1, i<j}^k 2\beta_{ij} \|u_i\|_{2p}^p \|u_j\|_{2p}^p \right)^{\frac{N-2}{2}}} S_{\lambda}^{\frac{N}{2}},$$

where $\|\cdot\|_q$ denotes the norm on $L^q(\Omega)$ and $2p = 2^* = \frac{2N}{N-2}$. Recall Lemma 2.2, then we see that

$$B = d_k S_{\lambda}^{\frac{N}{2}}$$

is attained by

$$(t_1 \omega, t_2 \omega, \dots, t_k \omega),$$

where (t_1, t_2, \dots, t_k) is the positive solution of (1.9) that attains d_k . In fact, we deduce that

$$B = \inf_{\vec{u} \in D \setminus \{\vec{0}\}} \frac{(\sum_{i=1}^k \|u_i\|_{2p}^2)^{\frac{N}{2}}}{N \left(\sum_{i=1}^k \mu_i \|u_i\|_{2p}^{2p} + \sum_{i,j=1, i<j}^k 2\beta_{ij} \|u_i\|_{2p}^p \|u_j\|_{2p}^p \right)^{\frac{N-2}{2}}} S_{\lambda}^{\frac{N}{2}}. \quad (4.28)$$

Suppose that B is attained by some nonzero $\vec{v} = (v_1, v_2, \dots, v_k)$. Then by Hölder inequality and Sobolev inequality, we see from (4.27) and (4.28) that

$$\begin{aligned} \int_{\Omega} (|\nabla v_i|^2 + \lambda v_i^2) dx &= S_{\lambda} \left(\int_{\Omega} |v_i|^{2p} dx \right)^{\frac{2}{2p}}, \\ \left(\int_{\Omega} |v_i v_j|^p dx \right)^2 &= \int_{\Omega} |v_i|^{2p} dx \int_{\Omega} |v_j|^{2p} dx \end{aligned}$$

for all $i, j = 1, 2, \dots, k$, which implies $v_i = 0$ or $v_i = c_i \omega_i$ for some $c_i \neq 0$, where ω_i is a positive least energy solution of Brézis-Nirenberg problem (1.7).

Moreover, for $v_i = c_i \omega_i \neq 0$, Hölder's inequality implies that there exists $k_i, k_j > 0$, such that $k_i (c_i \omega_i)^2 = k_j (c_j \omega_j)^2$. Thus we can redenote all $v_i = c_i \omega$ for all i where $c_i = 0$ or $c_i \neq 0$.

By (4.28), we have that $(\|v_1\|_{2p}, \|v_2\|_{2p}, \dots, \|v_k\|_{2p})$ attains d_k . It is easy to see that (c_1, c_2, \dots, c_k) attains d_k . Recall Lemma 2.2 that d_k is attained by a positive solution if $\beta_{ij} > 0$. Then $c_i > 0$ for all $i = 1, 2, \dots, k$.

That is, B is attained by $\vec{\mathbf{u}}$ if and only if

$$\vec{\mathbf{u}} = (\tilde{t}_1\omega, \tilde{t}_2\omega, \dots, \tilde{t}_k\omega),$$

where $(\tilde{t}_1, \tilde{t}_2, \dots, \tilde{t}_k)$ is defined in (1.10). Then the positive least energy solution of the system (1.1) must be of the synchronized form $(\tilde{t}_1\omega, \tilde{t}_2\omega, \dots, \tilde{t}_k\omega)$.

Now we assume that $\beta_{ij} := \beta > 0, i \neq j, i, j = 1, 2, \dots, k$. Assume that $\Omega \subset \mathbb{R}^N$ is a ball, then the positive least energy solution of the Brézis-Nirenberg problem (1.7) is unique (cf. [11]). Recall [49, Proposition 3.3] that there exists $\beta_k > 0$, such that the algebraic system (1.9) has a unique solution that attains d_k for $\beta > \beta_k$. Therefore, the positive least energy solution of (1.1) is unique under these assumptions. \square

Acknowledgements The authors wish to thank Dr. Y. Wu very much for his kind suggestion and valuable comments on the proof of Theorem 1.3.

References

- [1] B. Abdellaoui, V. Felli and I. Peral. Some remarks on systems of elliptic equations doubly critical in the whole \mathbb{R}^N . *Calc. Var. PDE*, 34:97–137 (2009)
- [2] N. Akhmediev and A. Ankiewicz. Partially coherent solitons on a finite background. *Phys. Rev. Letters*, 82(13):2661 (1999)
- [3] A. Ambrosetti and E. Colorado. Standing waves of some coupled nonlinear Schrödinger equations. *J. London Math. Soc.*, 75(1):67–82 (2007)
- [4] A. Ambrosetti and P. H. Rabinowitz. Dual variational methods in critical point theory and applications. *J. Func. Anal.*, 14(4):349–381 (1973)
- [5] A. Ambrosetti and M. Struwe. A note on the problem $-\Delta u = \lambda u + |u|^{2^*-2}u$. *Manuscripta Math.*, 54:373–379 (1986)
- [6] T. Aubin. Problèmes isopérimétriques et espaces de Sobolev. *J. Diff. Geom.*, 11(4):573–598 (1976)
- [7] T. Bartsch, N. Dancer, and Z.-Q. Wang. A Liouville theorem, a-priori bounds, and bifurcating branches of positive solutions for a nonlinear elliptic system. *Calc. Var. PDE.*, 37(3-4):345–361 (2010)
- [8] T. Bartsch and Z.-Q. Wang. Note on ground states of nonlinear Schrödinger systems. *J. Partial Diff. Eq.*, 19:200–207 (2006)
- [9] T. Bartsch, Z.-Q. Wang, and J. Wei. Bound states for a coupled Schrödinger system. *J. Fixed Point Theory A.*, 2(2):353–367 (2007)

- [10] H. Brézis and T. Kato. Remarks on the Schrödinger operator with singular complex potentials. *J. Math. Pures Appl.*, 58:137–151 (1979)
- [11] H. Brézis and L. Nirenberg. Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. *Commun. Pure Appl. Math.*, 36(4):437–477 (1983)
- [12] A. Capozzi, D. Fortunato and G. Palmieri. An existence result for nonlinear elliptic problems involving critical Sobolev exponent. *Ann. Inst. H. Poincaré-Analyse, Nonlinear*, 2:463–470 (1985)
- [13] G. Cerami, D. Fortunato, and M. Struwe. Bifurcation and multiplicity results for nonlinear elliptic problems involving critical Sobolev exponents. *Ann. Inst. H. Poincaré-Analyse, Nonlinear*, 1(5):341–350 (1984)
- [14] Z. Chen, C.-S. Lin, and W. Zou. Multiple sign-changing and semi-nodal solutions for coupled Schrödinger equations. *J. Diff. Equ.*, 255(11):4289–4311 (2013)
- [15] Z. Chen, C.-S. Lin, and W. Zou. Infinitely many sign-changing and semi-nodal solutions for a nonlinear Schrödinger system. *Annali della Scuola Normale Superiore di Pisa. Classe di scienze*, 15(1):859–897 (2016)
- [16] Z. Chen and W. Zou. On the Brézis-Nirenberg problem in a ball. *Differential and Integral Equations*, 25(5/6):527–542 (2012)
- [17] Z. Chen and W. Zou. Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent. *Arch. Ration. Mech. Anal.*, 205(2):515–551 (2012)
- [18] Z. Chen and W. Zou. An optimal constant for the existence of least energy solutions of a coupled Schrödinger system. *Calc. Var. PDE.*, 48(3-4):695–711 (2013)
- [19] Z. Chen and W. Zou. Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent: Higher dimensional case. *Calc. Var. PDE.*, 52(1-2):423–467 (2015)
- [20] M. Clapp and A. Szulkin. A simple variational approach to weakly coupled competitive elliptic systems. *Nonl. Diff. Eq. Appl. NoDEA*, 26(4):26 (2019)
- [21] M. Conti, S. Terracini, and G. Verzini. Nehari’s problem and competing species systems. *Ann. Inst. H. Poincaré-Analyse, Nonlinear*, 19(6):871–888 (2002)
- [22] M. Conti, S. Terracini, and G. Verzini. Asymptotic estimates for the spatial segregation of competitive systems. *Advances in Mathematics*, 195(2):524–560 (2005)
- [23] E. Dancer, J. Wei, and T. Weth. A priori bounds versus multiple existence of positive solutions for a nonlinear Schrödinger system. *Annales de l’IHP Analyse non linéaire*, 27(3):953–969 (2010)
- [24] G. Devillanova and S. Solimini. Concentration estimates and multiple solutions to elliptic problems at critical growth. *Adv. Diff. Eq.*, 7(10):1257–1280 (2002)

- [25] B. Esry, C. H. Greene, J. P. Burke Jr, and J. L. Bohn. Hartree-fock theory for double condensates. *Phys. Rev. Letters*, 78(19):3594 (1997)
- [26] D. Frantzeskakis. Dark solitons in atomic Bose–Einstein condensates: from theory to experiments. *Journal of Physics A: Mathematical and Theoretical*, 43(21):213001 (2010)
- [27] Y. Guo, S. Luo, and W. Zou. The existence, uniqueness and nonexistence of the ground state to the N-coupled Schrödinger systems in \mathbb{R}^N ($N \leq 4$). *Nonlinearity*, 31(1):314 (2018)
- [28] Q. He and J. Yang. Quantitative properties of ground-states to an M-coupled system with critical exponent in \mathbb{R}^n . *Sci. China Math.*, 61(4):709–726 (2018)
- [29] T.-C. Lin and J. Wei. Ground state of N coupled nonlinear Schrödinger equations in \mathbb{R}^N , $N \leq 3$. *Commun. Math. Phys.*, 255(3):629–653 (2005)
- [30] T.-C. Lin and J. Wei. Spikes in two coupled nonlinear Schrödinger equations. *Annales de l’IHP Analyse non linéaire*, 22(4):403–439 (2005)
- [31] C. Liu and Z. Q. Wang. A complete classification of ground-states for a coupled nonlinear Schrödinger system. *Commun. Pure Appl. Anal.*, 16(1):115–130 (2017)
- [32] J. Liu, X. Liu, and Z.-Q. Wang. Multiple mixed states of nodal solutions for nonlinear Schrödinger systems. *Calc. Var. PDE.*, 52(3-4):565–586 (2015)
- [33] L. Maia, E. Montefusco, and B. Pellacci. Infinitely many nodal solutions for a weakly coupled nonlinear Schrödinger system. *Commun. Contemp. Math.*, 10(05):651–669 (2008)
- [34] B. Noris and M. Ramos. Existence and bounds of positive solutions for a nonlinear Schrödinger system. *Proc. Amer. Math. Soc.*, 138(5):1681–1692 (2010)
- [35] B. Noris, S. Terracini, H. Tavares, and G. Verzini. Uniform Hölder bounds for nonlinear Schrödinger systems with strong competition. *Comm. Pure and Applied Mathematics*, 63(3):267–302 (2010)
- [36] S. Peng, Y.-f. Peng, and Z.-Q. Wang. On elliptic systems with Sobolev critical growth. *Calc. Var. PDE.*, 55(6):142 (2016)
- [37] Y. F. Peng and H. Y. Ye. Positive solutions for coupled Schrödinger system with critical exponent in \mathbb{R}^N ($N \geq 5$). *Bound. Value Probl.*, 2017(1) (2017)
- [38] Y. Sato and Z.-Q. Wang. On the least energy sign-changing solutions for a nonlinear elliptic system. *Discrete Contin. Dyn. Syst.*, 35:2151–2164 (2015)
- [39] M. Schechter and W. Zou. On the Brézis-Nirenberg problem. *Arch. Ration. Mech. Anal.*, 197(1):337–356 (2010)
- [40] B. Sirakov. Least energy solitary waves for a system of nonlinear Schrödinger equations in \mathbb{R}^N . *Comm. Math. Phys.*, 271(1):199–221 (2007)

- [41] H. Tavares and S. Terracini. Sign-changing solutions of competition–diffusion elliptic systems and optimal partition problems. *Annales de l’IHP Analyse non linéaire*, 29(2):279–300 (2012)
- [42] H. Tavares and S. You. Existence of least energy positive solutions to Schrödinger systems with mixed competition and cooperation terms: the critical case. *Calc. Var. PDE.*, 59(1):26 (2020)
- [43] S. Terracini and G. Verzini. Multipulse phases in k-mixtures of Bose–Einstein condensates. *Arch. Ration. Mech. Anal.*, 194(3):717–741 (2009)
- [44] E. Timmermans. Phase separation of Bose–Einstein condensates. *Phys. Rev. Letters*, 81(26):5718 (1998)
- [45] J. Wei and T. Weth. Radial solutions and phase separation in a system of two coupled Schrödinger equations. *Arch. Ration. Mech. Anal.*, 190(1):83–106 (2008)
- [46] J. Wei and Y. Wu. Ground states of Nonlinear Schrödinger System with Mixed Couplings. *preprint.*, <https://arxiv.org/abs/1903.05340>; (2019)
- [47] M. Willem. *Minimax theorems*. Springer Science & Business Media (1997)
- [48] Y. Wu. On a K-component elliptic system with the Sobolev critical exponent in high dimensions: the repulsive case. *Calc. Var. PDE.*, 56(5) (2017)
- [49] Y. Wu. Ground States of a K-Component Critical System with Linear and Nonlinear Couplings: The Attractive Case. *Adv. Nonlinear Stud.*, 19(3):595–623 (2019)
- [50] Y. Wu. On finding the ground state solution to the linearly coupled Brézis–Nirenberg system in high dimensions: The cooperative case. *Topol. Methods Nonlinear Anal.*, 53(2):697–729 (2019)
- [51] H. Yang and W. Zou. Stable and finite morse index solutions of a nonlinear elliptic system. *J. Math. Anal. Appl.*, 471(1-2):147–169 (2019)
- [52] H. Ye and Y. Peng. Positive least energy solutions for a coupled Schrödinger system with critical exponent. *J. Math. Anal. Appl.*, 417(1):308–326 (2014)