

# A VARIANT OF THE PRIME NUMBER THEOREM

KUI LIU, JIE WU & ZHISHAN YANG

ABSTRACT. Let  $\Lambda(n)$  be the von Mangoldt function, and let  $[t]$  be the integral part of real number  $t$ . In this note, we prove that for any  $\varepsilon > 0$  the asymptotic formula

$$\sum_{n \leq x} \Lambda\left(\left[\frac{x}{n}\right]\right) = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + O_\varepsilon(x^{9/19+\varepsilon}) \quad (x \rightarrow \infty)$$

holds. This improves a recent result of Bordellès, which requires  $\frac{97}{203}$  in place of  $\frac{9}{19}$ .

## 1. INTRODUCTION

The prime number theorem is a basic result in number theory and has many applications. Denoting by  $\Lambda(n)$  the von Mangoldt function, the prime number theorem states, in strong form, as follows: there is a constant  $c > 0$  such that for  $x \rightarrow \infty$ , we have

$$\sum_{n \leq x} \Lambda(n) = x + O(x \exp\{-c(\log x)^{3/5}(\log \log x)^{-1/5}\})$$

and

$$\sum_{n \leq x} \Lambda(n) = x + O_\varepsilon(x^{1/2+\varepsilon}) \Leftrightarrow \text{Riemann Hypothesis,}$$

where  $\varepsilon$  is an arbitrarily small positive constant. Clearly it is also interesting to study the distribution of prime numbers in different sequences of integers such as the arithmetic progressions, the Beatty sequence  $([\alpha n + \beta])_{n \geq 1}$ , the Piatetski-Shapiro sequence  $([n^c])_{n \geq 1}$ , etc, where  $[t]$  denotes the integral part of the real number. For example, Banks and Shparlinski [2, Corollary 5.6] proved the following result: Let  $\alpha$  and  $\beta$  be fixed real numbers with  $\alpha > 0$ , irrational and of finite type, then there is a positive constant  $c = c(\alpha, \beta)$  such that

$$\sum_{n \leq x} \Lambda([\alpha n + \beta]) = x + O(x \exp\{-c(\log x)^{3/5}(\log \log x)^{-1/5}\})$$

as  $x \rightarrow \infty$ . About works related to the Piatetski-Shapiro prime number theorem, we infer the reader to see [12, 8, 13]. On the other hand, Bordellès-Dai-Heyman-Pan-Shparlinski [4] established an asymptotic formula of

$$(1.1) \quad S_f(x) := \sum_{n \leq x} f\left(\left[\frac{x}{n}\right]\right),$$

under some simple assumptions of  $f$ . Subsequently, Wu [16] and Zhai [17] improved their results independently. In particular, applying [16, Theorem 1.2(i)] or [17, Theorem 1] to the von Mangoldt function  $\Lambda(n)$ , we have

$$S_\Lambda(x) = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + O_\varepsilon(x^{1/2+\varepsilon})$$

---

*Date:* December 10, 2021.

*2010 Mathematics Subject Classification.* 11N37, 11L07.

*Key words and phrases.* The prime number theorem, Exponential sums, Vaughan's identity.

for  $x \geq 1$ . With the help of the Vaughan identity and the method of one-dimensional exponential sum, Ma and Wu [11] broke the  $\frac{1}{2}$ -barrier:

$$S_{\Lambda}(x) = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + O_{\varepsilon}(x^{35/71+\varepsilon}) \quad (x \geq 1).$$

Very recently Bordellès [3, Corollary 1.3] sharpened the exponent  $\frac{35}{71}$  to  $\frac{97}{203}$  by using a result of Baker on 2-dimensional exponential sums [1, Theorem 6].

The aim of this short note is to propose a better exponent by establishing an estimate on 3-dimensional exponential sums (see Proposition 3.1 below).

**Theorem 1.** *For any  $\varepsilon > 0$ , we have*

$$(1.2) \quad S_{\Lambda}(x) = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + O_{\varepsilon}(x^{9/19+\varepsilon}) \quad \text{as } x \rightarrow \infty.$$

For comparison, we have  $\frac{97}{203} \approx 0.4778$  and  $\frac{9}{19} \approx 0.4736$ . In fact, Bordellès established a more general result (see [3, Theorem 1.1]):

$$(1.3) \quad S_{\Lambda}(x) = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + O_{\varepsilon}(x^{\frac{14(\kappa+1)}{29\kappa-\lambda+30}+\varepsilon}) \quad (x \rightarrow \infty),$$

where  $(\kappa, \lambda)$  is an exponent pair satisfying  $\kappa \leq \frac{1}{6}$  and  $\lambda^2 + \lambda + 3 - \kappa(5 + 9\kappa - \lambda) > 0$ . The exponent  $\frac{97}{203}$  comes from the choice of  $(\kappa, \lambda) = (\frac{13}{84} + \varepsilon, \frac{55}{84} + \varepsilon)$  — Bourgain's new exponent pair [5, Theorem 6]. We note that under the exponent pair hypothesis (i.e.  $(\varepsilon, \frac{1}{2} + \varepsilon)$  is an exponent pair, see [7]), Bordellès' (1.3) only gives the exponent  $\frac{28}{59} \approx 0.4745$ , which is larger than our constant  $\frac{9}{19} \approx 0.4736$ .

Some related works on the quantity (1.1) can be found in [10, 15].

## 2. PRELIMINARY LEMMAS

In this section, we shall cite three lemmas, which will be needed in the next section. The first one is [6, Lemma 1].

**Lemma 2.1.** *Let  $\alpha \in \mathbb{R}^*$  and  $\beta \in \mathbb{R}^*$ . For  $M \geq 1$ ,  $N \geq 1$  and  $\Delta > 0$ , define*

$$\mathcal{D}(M, N; \Delta) := \left| \left\{ (m, \tilde{m}, n, \tilde{n}) \in \mathbb{N}^4 : m, \tilde{m} \sim M; n, \tilde{n} \sim N; \left| \left( \frac{m}{\tilde{m}} \right)^{\alpha} - \left( \frac{n}{\tilde{n}} \right)^{\beta} \right| \leq \Delta \right\} \right|,$$

where  $m \sim M$  means that  $M < m \leq 2M$ . Then we have

$$\mathcal{D}(M, N) \ll_{\alpha, \beta} MN \log(2MN) + \Delta(MN)^2$$

uniformly for  $M \geq 1$ ,  $N \geq 1$  and  $\Delta > 0$ , where the implied constant depends on  $\alpha$  and  $\beta$ .

The second one is the Vaughan identity [14, formula (3)].

**Lemma 2.2.** *There are six real arithmetical functions  $\alpha_k(n)$  verifying  $|\alpha_k(n)| \ll_{\varepsilon} n^{\varepsilon}$  for  $(n \geq 1, 1 \leq k \leq 6)$  such that, for all  $D \geq 100$  and any arithmetical function  $g$ , we have*

$$(2.1) \quad \sum_{D < d \leq 2D} \Lambda(d)g(d) = S_1 + S_2 + S_3 + S_4,$$

where

$$\begin{aligned}
S_1 &:= \sum_{m \leq D^{1/3}} \alpha_1(m) \sum_{D < mn \leq 2D} g(mn), \\
S_2 &:= \sum_{m \leq D^{1/3}} \alpha_2(m) \sum_{D < mn \leq 2D} g(mn) \log n, \\
S_3 &:= \sum_{\substack{D^{1/3} < m, n \leq D^{2/3} \\ D < mn \leq 2D}} \alpha_3(m) \alpha_4(n) g(mn), \\
S_4 &:= \sum_{\substack{D^{1/3} < m, n \leq D^{2/3} \\ D < mn \leq 2D}} \alpha_5(m) \alpha_6(n) g(mn).
\end{aligned}$$

The sums  $S_1$  and  $S_2$  are called as type I,  $S_3$  and  $S_4$  are called as type II.

The third one is due to Vaaler (see [7, Theorem A.6]).

**Lemma 2.3.** Let  $\psi(t) := \{t\} - \frac{1}{2}$ , where  $\{t\}$  means the fractional part of real number  $t$ . For  $x \geq 1$  and  $H \geq 1$ , we have

$$\psi(x) = - \sum_{1 \leq |h| \leq H} \Phi\left(\frac{h}{H+1}\right) \frac{e(hx)}{2\pi i h} + R_H(x),$$

where  $e(t) := e^{2\pi i t}$ ,  $\Phi(t) := \pi t(1 - |t|) \cot(\pi t) + |t|$  and the error term  $R_H(x)$  satisfies

$$(2.2) \quad |R_H(x)| \leq \frac{1}{2H+2} \sum_{0 \leq |h| \leq H} \left(1 - \frac{|h|}{H+1}\right) e(hx).$$

### 3. MULTIPLE EXPONENTIAL SUMS

Let  $\alpha > 0$ ,  $\beta > 0$ ,  $\gamma > 0$  and  $\delta \in \mathbb{R}$  be some constants. For  $X > 0$ ,  $H \geq 1$ ,  $M \geq 1$  and  $N \geq 1$ , define

$$(3.1) \quad S_\delta = S_\delta(H, M, N) := \sum_{h \sim H} \sum_{m \sim M} \sum_{n \sim N} a_{h,m} b_n e\left(X \frac{M^\beta N^\gamma}{H^\alpha} \frac{h^\alpha}{m^\beta n^\gamma + \delta}\right),$$

where the  $a_{h,m}$  and  $b_n$  are complex numbers such that  $|a_{h,m}| \leq 1$  and  $|b_n| \leq 1$ . When  $\delta = 0$ , this sum has been studied by Heath-Brown [8] and Fouvry-Iwaniec [6]. The aim of this section is to prove an estimate of  $S_\delta$  by adapting and refining Heath-Brown's approach (see also [9]). The following proposition will play a key role in the proof of Theorem 1.

**Proposition 3.1.** Under the previous notation, for any  $\varepsilon > 0$  we have

$$(3.2) \quad S_\delta \ll ((XHMN)^{1/2} + (HM)^{1/2}N + HMN^{1/2} + X^{-1/2}HMN)X^\varepsilon,$$

$$(3.3) \quad S_\delta \ll ((X^\kappa H^{2+\kappa} M^{2+\kappa} N^{1+\kappa+\lambda})^{1/(2+2\kappa)} + HMN^{1/2}$$

$$(3.4) \quad + (HM)^{1/2}N + X^{-1/2}HMN)X^\varepsilon$$

uniformly for  $M \geq 1$ ,  $N \geq 1$ ,  $H \leq M^{\beta-1}N^\gamma$  and  $|\delta| \leq 1/\varepsilon$ , where  $(\kappa, \lambda)$  is an exponent pair and the implied constant depends on  $(\alpha, \beta, \gamma, \varepsilon)$  only.

*Proof.* Obviously we have

$$h^\alpha m^{-\beta} \leq C_1 H^\alpha M^{-\beta} =: \Xi \quad (h \sim H, m \sim M),$$

where the  $C_j = C_j(\alpha, \beta, \gamma)$  denotes some positive constant depending on  $(\alpha, \beta, \gamma)$  at most. Let  $K \geq 1$  be a parameter to be chosen later. Introducing the following set

$$T_k := \{(h, m) : h \sim H, m \sim M, \Xi(k-1) < h^\alpha m^{-\beta} K \leq \Xi k\},$$

we can write

$$S_\delta = \sum_{n \sim N} b_n \sum_{k \leq K} \sum_{(h, m) \in T_k} a_{h, m} e \left( X \frac{M^\beta N^\gamma}{H^\alpha} \frac{h^\alpha}{m^\beta n^\gamma + \delta} \right).$$

By the Cauchy-Schwarz inequality, we derive

$$|S_\delta|^2 \leq NK \sum_{n \sim N} \sum_{k \leq K} \sum_{(h, m) \in T_k} \sum_{(h', m') \in T_k} a_{h, m} \overline{a_{h', m'}} e \left( X \frac{M^\beta N^\gamma}{H^\alpha} \left( \frac{h^\alpha}{m^\beta n^\gamma + \delta} - \frac{h'^\alpha}{m'^\beta n'^\gamma + \delta} \right) \right).$$

Inverting the order of summations, it follows that

$$|S_\delta|^2 \leq NK \sum_{k \leq K} \sum_{(h, m) \in T_k} \sum_{(h', m') \in T_k} |\mathcal{S}(h, h', m, m')|,$$

where

$$(3.5) \quad \mathcal{S}(h, h', m, m') := \sum_{n \sim N} e \left( X \frac{M^\beta N^\gamma}{H^\alpha} \left( \frac{h^\alpha}{m^\beta n^\gamma + \delta} - \frac{h'^\alpha}{m'^\beta n'^\gamma + \delta} \right) \right).$$

Noticing that  $(h, m), (h', m') \in T_k$  imply  $|h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq \Xi K^{-1}$ , we can write

$$(3.6) \quad |S_\delta|^2 \leq NK \sum_{\substack{h, h' \sim H \\ |h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq \Xi K^{-1}}} \sum_{\substack{m, m' \sim M}} |\mathcal{S}(h, h', m, m')| \leq NK (S_\delta^\dagger + S_\delta^\sharp),$$

where

$$\begin{aligned} S_\delta^\dagger &:= \sum_{\substack{h, h' \sim H \\ |h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq \Xi(HM)^{-1}}} \sum_{\substack{m, m' \sim M}} |\mathcal{S}(h, h', m, m')|, \\ S_\delta^\sharp &:= \sum_{\substack{h, h' \sim H \\ \Xi(HM)^{-1} < |h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq \Xi K^{-1}}} \sum_{\substack{m, m' \sim M}} |\mathcal{S}(h, h', m, m')|. \end{aligned}$$

[We have made the convention that  $S_\delta^\sharp = 0$  if  $K \leq HM$ .] Since

$$\left| \frac{h^\alpha}{m^\beta} - \frac{h'^\alpha}{m'^\beta} \right| \leq \frac{\Xi}{HM} \Rightarrow \left| \frac{h^\alpha}{h'^\alpha} - \frac{m^\beta}{m'^\beta} \right| \leq \frac{C_2}{\Xi} \left| \frac{h^\alpha}{m^\beta} - \frac{h'^\alpha}{m'^\beta} \right| \leq \frac{C_2}{HM},$$

Lemma 2.1 implies that the number of  $(h, h', m, m')$  verifying  $|h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq \Xi(HM)^{-1}$  is  $\ll \mathcal{D}(H, M; C_2/HM) \ll_\varepsilon HMX^\varepsilon$ . Thus we have trivially

$$(3.7) \quad S_\delta^\dagger \ll_\varepsilon HMNX^\varepsilon.$$

Next we bound  $S_\delta^\sharp$ . We write

$$\begin{aligned} (3.8) \quad & \frac{h^\alpha}{m^\beta n^\gamma + \delta} - \frac{h'^\alpha}{m'^\beta n'^\gamma + \delta} \\ &= \frac{1}{n^\gamma} \left( \frac{h^\alpha}{m^\beta} - \frac{h'^\alpha}{m'^\beta} \right) - \frac{\delta}{n^\gamma} \left( \frac{h^\alpha}{m^\beta(m^\beta n^\gamma + \delta)} - \frac{h'^\alpha}{m'^\beta(m'^\beta n'^\gamma + \delta)} \right) =: f(n). \end{aligned}$$

Since  $H \leq M^{\beta-1}N^\gamma$ , we have

$$\left| \frac{h^\alpha}{m^\beta(m^\beta n^\gamma + \delta)} - \frac{h'^\alpha}{m'^\beta(m'^\beta n'^\gamma + \delta)} \right| \leq \frac{C_3 \Xi}{M^\beta N^\gamma} \ll \frac{\Xi}{HM} \quad (h, h' \sim H; m, m' \sim M),$$

Therefore for  $(h, h', m, m')$  verifying  $\Xi(HM)^{-1} < |h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq \Xi K^{-1}$ , the first member on the right-hand side of (3.8) dominates the second one. Split  $(\Xi(HM)^{-1}, \Xi K^{-1}]$  into dyadic intervals  $(\Delta\Xi, 2\Delta\Xi]$  with  $1/HM \leq \Delta \leq 1/K$ . Take  $K = \max\{2C_4X/N, 1\}$  such that for  $(h, h', m, m')$  verifying  $\Xi\Delta < |h^\alpha m^{-\beta} - h'^\alpha m'^{-\beta}| \leq 2\Xi\Delta$  we have

$$\max_{n \sim N} |f'(n)| = C_4 X \Delta N^{-1} \leq \frac{1}{2}.$$

By Kusmin-Landau's inequality and Lemma 2.1, we have

$$(3.9) \quad S_\delta^\sharp \ll_\varepsilon X^\varepsilon \max_{4/HM \leq \Delta \leq 1/K} \Delta (HM)^2 (X \Delta N^{-1})^{-1} \ll_\varepsilon X^{-1+\varepsilon} (HM)^2 N.$$

Combining (3.7) and (3.9) with (3.6) gives us

$$\begin{aligned} |S_\delta|^2 &\ll_\varepsilon (HMN^2K + X^{-1}(HMN)^2K)X^\varepsilon \\ &\ll_\varepsilon (XHMN + HMN^2 + (HM)^2N + X^{-1}(HMN)^2)X^\varepsilon, \end{aligned}$$

which is equivalent to (3.2).

Next we prove (3.4). If  $X \leq HM$ , then (3.2) implies that

$$(3.10) \quad S_\delta \ll_\varepsilon ((HM)^{1/2}N + HMN^{1/2} + X^{-1/2}HMN)X^\varepsilon.$$

Now we can suppose that  $X \geq HM$ . Applying the exponent pair  $(\kappa, \lambda)$  to  $\mathcal{S}(h, h', m, m')$  and using Lemma 2.1, we can derive that

$$\begin{aligned} (3.11) \quad S_\delta^\sharp &\ll_\varepsilon X^\varepsilon \max_{4/HM \leq \Delta \leq 1/K} \Delta (HM)^2 ((X \Delta N^{-1})^\kappa N^\lambda + (X \Delta N^{-1})^{-1}) \\ &\ll_\varepsilon X^\varepsilon (X^\kappa H^2 M^2 N^{-\kappa+\lambda} K^{-1-\kappa} + X^{-1} H^2 M^2 N). \end{aligned}$$

Combining (3.7) and (3.11) with (3.6) gives us

$$|S_\delta|^2 \ll_\varepsilon (X^\kappa H^2 M^2 N^{1-\kappa+\lambda} K^{-\kappa} + HMN^2 K) X^\varepsilon$$

for all  $K \in [1, HM]$ , where we have removed the term  $X^{-1} H^2 M^2 N^2 K$  ( $\leq HMN^2 K$  since we have suppose that  $X \geq HM$ ). Noticing that this estimate is trivial if  $K \geq HM$ , we can optimise the parameter  $K$  over  $[1, \infty)$  to get

$$|S_\delta|^2 \ll_\varepsilon ((X^\kappa H^{2+\kappa} M^{2+\kappa} N^{1+\kappa+\lambda})^{1/(1+\kappa)} + HMN^2) X^\varepsilon.$$

Combining this with (3.10), we obtain (3.4).  $\square$

#### 4. A KEY INEQUALITY

The aim of this section is to prove the following proposition, which will play a key role for the proof of Theorem 1. Define

$$(4.1) \quad \mathfrak{S}_\delta(x, D) := \sum_{d \sim D} \Lambda(d) \psi\left(\frac{x}{d + \delta}\right).$$

**Proposition 4.1.** *Let  $\delta \notin -\mathbb{N}$  be a fixed constant. Then we have*

$$(4.2) \quad \begin{aligned} \mathfrak{S}_\delta(x, D) &\ll ((x^{2\kappa} D^{3+\lambda})^{1/(4\kappa+4)} + D^{5/6} \\ &\quad + (x^{3\kappa'} D^{-2\kappa'+2\lambda'+1})^{1/(3\kappa'+3)} + (x^{3\kappa'} D^{-5\kappa'+2\lambda'+1})^{1/3}) x^\varepsilon. \end{aligned}$$

uniformly for  $x \geq 3$  and  $1 \leq D \leq x^{2/3}$ , where  $(\kappa, \lambda)$  and  $(\kappa', \lambda')$  are exponent pairs. In particular, uniformly for  $x^{6/13} \leq D \leq x^{2/3}$  we have

$$(4.3) \quad \mathfrak{S}_\delta(x, D) \ll_\varepsilon (x^2 D^7)^{1/12} x^\varepsilon.$$

*Proof.* We apply the Vaughan identity (2.1) with  $g(d) = \psi\left(\frac{x}{d+\delta}\right)$  to write

$$(4.4) \quad \mathfrak{S}_\delta(x, D) = \mathfrak{S}_{\delta,1} + \mathfrak{S}_{\delta,2} + \mathfrak{S}_{\delta,3} + \mathfrak{S}_{\delta,4},$$

where

$$\begin{aligned} \mathfrak{S}_{\delta,1} &:= \sum_{m \leq D^{1/3}} \alpha_1(m) \sum_{D < mn \leq 2D} \psi\left(\frac{x}{mn + \delta}\right), \\ \mathfrak{S}_{\delta,2} &:= \sum_{m \leq D^{1/3}} \alpha_2(m) \sum_{D < mn \leq 2D} \psi\left(\frac{x}{mn + \delta}\right) \log n, \\ \mathfrak{S}_{\delta,3} &:= \sum_{\substack{D^{1/3} < m, n \leq D^{2/3} \\ D < mn \leq 2D}} \alpha_3(m) \alpha_4(n) \psi\left(\frac{x}{mn + \delta}\right), \\ \mathfrak{S}_{\delta,4} &:= \sum_{\substack{D^{1/3} < m, n \leq D^{2/3} \\ D < mn \leq 2D}} \alpha_5(m) \alpha_6(n) \psi\left(\frac{x}{mn + \delta}\right). \end{aligned}$$

Firstly we estimate  $\mathfrak{S}_{\delta,3}$ . In view of Lemma 2.3, we can write

$$(4.5) \quad \mathfrak{S}_{\delta,3} = -\frac{1}{2\pi i} \sum_{H'} \sum_M \sum_N (\mathfrak{S}_{\delta,3}^b(H', M, N) + \overline{\mathfrak{S}_{\delta,3}^b(H', M, N)}) + \sum_M \sum_N \mathfrak{S}_{\delta,3}^\dagger(M, N).$$

where  $MN \asymp D$  (i.e.  $D \ll MN \ll D$ ),  $\alpha(h) := \frac{H'}{h} \Phi\left(\frac{h}{H+1}\right) \ll 1$  and

$$\begin{aligned} \mathfrak{S}_{\delta,3}^b(H', M, N) &:= \frac{1}{H'} \sum_{h \sim H'} \sum_{m \sim M} \sum_{\substack{n \sim N \\ D < mn \leq 2D}} \alpha(h) \alpha_3(m) \alpha_4(n) e\left(\frac{hx}{mn + \delta}\right), \\ \mathfrak{S}_{\delta,3}^\dagger(M, N) &:= \sum_{\substack{m \sim M \\ D < mn \leq 2D}} \sum_{n \sim N} \alpha_3(m) \alpha_4(n) R_H\left(\frac{x}{mn + \delta}\right). \end{aligned}$$

Firstly we bound  $\mathfrak{S}_{\delta,3}^b(H', M, N)$ . We remove the extra multiplicative condition  $D < mn \leq 2D$  at the cost of a factor  $\log x$ . On the other hand, in view of the symmetry of the variables  $m$  and  $n$ , we can suppose that

$$(4.6) \quad D^{1/3} \leq M \leq D^{1/2} \leq N \leq D^{2/3}.$$

By Proposition 3.1 with  $\alpha = \beta = \gamma = 1$  and  $(X, H, M, N) = (xH'/MN, H', M, N)$ , we have

$$(4.7) \quad \mathfrak{S}_{\delta,3}^b(H', M, N) \ll_\varepsilon \left( (x^\kappa M^2 N^{1+\lambda})^{1/(2+2\kappa)} + MN^{1/2} + M^{1/2}N + (x^{-1}DH')^{1/2} \right) x^\varepsilon,$$

provided  $H' \leq H \leq N$ .

Secondly we bound  $\mathfrak{S}_{\delta,3}^\dagger(M, N)$ . Using (2.2), we have

$$\begin{aligned} \mathfrak{S}_{\delta,3}^\dagger(M, N) &\ll x^\varepsilon \sum_{m \sim M} \sum_{n \sim N} \left| R_H\left(\frac{x}{mn + \delta}\right) \right| \\ &\ll \frac{x^\varepsilon}{H} \sum_{m \sim M} \sum_{n \sim N} \sum_{0 \leq |h| \leq H} \left( 1 - \frac{|h|}{H+1} \right) e\left(\frac{hx}{mn + \delta}\right) \\ &\ll x^\varepsilon (DH^{-1} + \max_{1 \leq H' \leq H} |\tilde{\mathfrak{S}}_{\delta,3}^\dagger(H', M, N)|), \end{aligned}$$

where

$$\tilde{\mathfrak{S}}_{\delta,3}^\dagger(H', M, N) := \frac{1}{H} \sum_{m \sim M} \sum_{n \sim N} \sum_{h \sim H'} \left( 1 - \frac{|h|}{H+1} \right) e\left(\frac{hx}{mn + \delta}\right).$$

Clearly we can bound  $\tilde{\mathfrak{S}}_{\delta,3}^\dagger(H', M, N)$  in the same way as  $\mathfrak{S}_{\delta,3}^\flat(H', M, N)$  and obtain

$$(4.8) \quad \mathfrak{S}_{\delta,3}^\dagger(M, N) \ll_\varepsilon \left( DH^{-1} + (x^\kappa M^2 N^{1+\lambda})^{1/(2+2\kappa)} \right. \\ \left. + MN^{1/2} + M^{1/2}N + (x^{-1}DH)^{1/2} \right) x^\varepsilon,$$

provided  $H \leq N$ . Combining (4.7) and (4.8) with (4.5) and using (4.6), we find that

$$\mathfrak{S}_{\delta,3} \ll_\varepsilon \left( DH^{-1} + (x^\kappa M^2 N^{1+\lambda})^{1/(2+2\kappa)} + MN^{1/2} + M^{1/2}N + (x^{-1}DH)^{1/2} \right) x^\varepsilon \\ \ll_\varepsilon \left( DH^{-1} + (x^{2\kappa} D^{3+\lambda})^{1/(4+4\kappa)} + D^{5/6} + (x^{-1}DH)^{1/2} \right) x^\varepsilon,$$

provided  $H \leq D^{1/2} (\leq N)$ . Optimising  $H$  over  $[1, D^{1/2}]$ , it follows that

$$(4.9) \quad \mathfrak{S}_{\delta,3} \ll_\varepsilon \left( (x^{2\kappa} D^{3+\lambda})^{1/(4+4\kappa)} + D^{5/6} + (x^{-1}D^2)^{1/3} \right) x^\varepsilon.$$

Clearly the same estimate also holds for  $\mathfrak{S}_{\delta,4}$ .

On the other hand, let  $(\kappa', \lambda')$  be an exponent pair, then [11, (5.16)] gives us

$$(4.10) \quad \mathfrak{S}_{\delta,j} \ll \left( (x^{3\kappa'} D^{-2\kappa'+2\lambda'+1})^{1/(3\kappa'+3)} + x^{\kappa'} D^{(-5\kappa'+2\lambda'+1)/3} + x^{-1}D^2 \right) x^\varepsilon.$$

for  $j = 1, 2$ . Inserting (4.10) and (4.9) into (4.4) and using the fact that

$$\max\{(x^{-1}D^2)^{1/3}, x^{-1}D^2\} \leq D^{5/6}$$

for  $1 \leq D \leq x^{2/3}$ , we get (4.2).

Taking  $(\kappa, \lambda) = (\kappa', \lambda') = (\frac{1}{2}, \frac{1}{2})$  in (4.2), we find that

$$\mathfrak{S}_\delta(x, D) \ll \left( (x^2 D^7)^{1/12} + D^{5/6} + (x^3 D^2)^{1/9} + (x^3 D^{-1})^{1/6} \right) x^\varepsilon,$$

which implies (4.3), since the last three terms can be absorbed by the first one provided  $x^{6/13} \leq D \leq x^{2/3}$ .  $\square$

## 5. PROOF OF THEOREM 1

Let  $N \in [x^{6/13}, x^{1/2})$  be a parameter which can be chosen later. First we write

$$(5.1) \quad \sum_{n \leq x} \Lambda\left(\left[\frac{x}{n}\right]\right) := S_1(x) + S_2(x)$$

with

$$S_1(x) := \sum_{n \leq N} \Lambda\left(\left[\frac{x}{n}\right]\right), \quad S_2(x) := \sum_{N < n \leq x} \Lambda\left(\left[\frac{x}{n}\right]\right).$$

We have trivially

$$(5.2) \quad S_1(x) \ll_\varepsilon Nx^\varepsilon.$$

Next we bound  $S_2(x)$ . Putting  $d = [x/n]$ , then  $x/n - 1 < d \leq x/n \Leftrightarrow x/(d+1) < n \leq x/d$ . Thus we can write

$$(5.3) \quad S_2(x) = \sum_{d \leq x/N} \Lambda(d) \sum_{x/(d+1) < n \leq x/d} 1 \\ = \sum_{d \leq x/N} \Lambda(d) \left( \frac{x}{d} - \psi\left(\frac{x}{d}\right) - \frac{x}{d+1} + \psi\left(\frac{x}{d+1}\right) \right) \\ = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + \mathcal{R}_1(x) - \mathcal{R}_0(x) + O(N),$$

where we have used the following bounds

$$x \sum_{d > x/N} \frac{\Lambda(d)}{d(d+1)} \ll_{\varepsilon} Nx^{\varepsilon}, \quad \sum_{d \leq N} \Lambda(d) \left( \psi\left(\frac{x}{d+1}\right) - \psi\left(\frac{x}{d}\right) \right) \ll_{\varepsilon} Nx^{\varepsilon}$$

and defined

$$\mathcal{R}_{\delta}(x) = \sum_{N < d \leq x/N} \Lambda(d) \psi\left(\frac{x}{d+\delta}\right).$$

Writing  $D_j := x/(2^j N)$ , we have  $x^{6/13} \leq N \leq D_j \leq x/N \leq x^{7/13}$  for  $0 \leq j \leq \frac{\log(x/N^2)}{\log 2}$  since  $x^{6/13} \leq N \leq x^{1/2}$ . Thus we can apply (4.3) of Proposition 4.1 to get

$$\begin{aligned} |\mathcal{R}_{\delta}(x)| &\leq \sum_{0 \leq j \leq \log(x/N^2)/\log 2} |\mathfrak{S}_{\delta}(x, D_j)| \\ &\ll \sum_{0 \leq j \leq \log(x/N^2)/\log 2} (x^2 D_j^7)^{1/12} x^{\varepsilon} \\ &\ll (x^9 N^{-7})^{1/12} x^{\varepsilon}. \end{aligned}$$

Putting this into (5.3) and taking  $N = x^{9/19}$ , we find that

$$(5.4) \quad S_2(x) = x \sum_{d \geq 1} \frac{\Lambda(d)}{d(d+1)} + O_{\varepsilon}(x^{9/19+\varepsilon}).$$

Inserting (5.2) with  $N = x^{9/19}$  and (5.4) into (5.1), we get the required result.

**Acknowledgement.** This work is supported in part by the National Natural Science Foundation of China (Grant Nos. 12071238, 11771252, 11971370 and 12071375) and by the NSF of Chongqing (Grant No. cstc2019jcyj-msxm1651).

## REFERENCES

- [1] R. C. Baker, *Sums of two relatively prime cubes*, Acta Arith. **129** (2007), 103–146.
- [2] William D. Banks and Igor E. Shparlinski, *Prime numbers with Beatty sequences*, Colloquium Math. **115** (2009), No. 2, 147–157.
- [3] O. Bordellès, *On certain sums of number theory*, arXiv:2009.05751v2 [math.NT] 25 Nov 2020.
- [4] O. Bordellès, L. Dai, R. Heyman, H. Pan and I. E. Shparlinski, *On a sum involving the Euler function*, J. Number Theory **202** (2019), 278–297.
- [5] J. Bourgain, *Decoupling, exponential sums and the Riemann zeta function*, J. Amer. Math. Soc. **30** (2017), 205–224.
- [6] E. Fouvry and H. Iwaniec, *Exponential sums with monomials*, J. Number Theory **33** (1989), 331–333.
- [7] S. W. Graham and G. Kolesnik, *Van der Corput's Method of Exponential Sums*, Cambridge University Press, Cambridge, 1991.
- [8] D. R. Heath-Brown, *The Pjateckiĭ-Šapiro Prime Number Theorem*, J. Number Theory **16** (1983), 242–266.
- [9] H.-Q. Liu & J. Wu, *Numbers with a large prime factor*, Acta Arith. **89** (1999), 163–187.
- [10] J. Ma & H.-Y. Sun, *On a sum involving the divisor function*, Period. Math. Hung. , to appear.
- [11] J. Ma & J. Wu, *On a sum involving the von Mangoldt function*, Period. Math. Hung. **83** (2021), Number 1, 39–48.
- [12] I. I. Pjateckiĭ-Šapiro, *On the distribution of prime numbers in sequences of the form  $[f(n)]$* , Mat. Sb. **33** (1953), 559–566.
- [13] J. Rivat & J. Wu, *Prime numbers of the forms  $[n^c]$* , Glasg. Math. **43** (2001), no. 2, 237–254.
- [14] R. C. Vaughan, *An elementary method in prime number theory*, Acta Arith. **37** (1980), 111–115.
- [15] J. Wu, *On a sum involving the Euler totient function*, Indag. Mathematicae **30** (2019), 536–541.



- [16] J. Wu, *Note on a paper by Bordellès, Dai, Heyman, Pan and Shparlinski*, Period. Math. Hung. **80** (2020), Number 1, 95–102.
- [17] W. Zhai, *On a sum involving the Euler function*, J. Number Theory **211** (2020), 199–219.

KUI LIU, SCHOOL OF MATHEMATICS AND STATISTICS, QINGDAO UNIVERSITY, 308 NINGXIA ROAD,  
QINGDAO, SHANDONG 266071, CHINA  
*Email address:* liukui@qdu.edu.cn

JIE WU, CNRS UMR 8050, LABORATOIRE D'ANALYSE ET DE MATHÉMATIQUES APPLIQUÉES,  
UNIVERSITÉ PARIS-EST CRÉTEIL, 94010 CRÉTEIL CEDEX, FRANCE  
*Email address:* jie.wu@u-pec.fr

ZHISHAN YANG, SCHOOL OF MATHEMATICS AND STATISTICS, QINGDAO UNIVERSITY, 308 NINGXIA  
ROAD, QINGDAO, SHANDONG 266071, CHINA  
*Email address:* zsyang@qdu.edu.cn