

Existence result for fractional problems with logarithmic and critical exponential nonlinearities ^{*}

Yuanyuan Zhang

School of Science

Jiangnan university

Wuxi, Jiangsu 214122, China

E-mail: 6191204010@stu.jiangnan.edu.cn

Yang Yang[†]

School of Science

Jiangnan University

Wuxi, Jiangsu 214122, China

E-mail: yynjnu@126.com

Abstract

We study the existence of nontrivial solutions for a nonlinear fractional elliptic equation in presence of logarithmic and critical exponential nonlinearities. This problem extends [5] to fractional p -Laplacian equations with logarithmic nonlinearity. We overcome the lack of compactness due to the critical exponential nonlinearity by using the fractional Trudinger-Moser inequality. The existence result is established via critical point theory.

1 Introduction and main results

Let $N \geq 2$, $s \in (0, 1)$ and $\Omega \subset \mathbb{R}^N$ be a bounded domain with Lipschitz boundary. For $\mu > 0$ and $\frac{N}{s} < q < \frac{N\sigma}{s}$, we consider the existence of solutions for the following fractional problem

$$\begin{cases} (-\Delta)_{N/s}^s u = |u|^{q-2}u \ln |u|^2 + \mu |u|^{(N\sigma-2s)/s} u e^{|u|^{N/(N-s)}}, & \text{in } \Omega, \\ u = 0, & \text{in } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (1.1)$$

^{*}MSC2020: Primary 35J35 Secondary 35J60, 35D30

Key Words and Phrases: Fractional Trudinger-Moser embedding, Logarithmic nonlinearity, Exponential nonlinearity, Existence of solutions

[†]Corresponding author. Project supported by NSFC(No. 11501252, No. 11571176).

Here, $(-\Delta)_{N/s}^s$ is the nonlinear nonlocal operator defined on smooth functions by

$$(-\Delta)_{N/s}^s \phi(x) := 2 \lim_{\epsilon \searrow 0} \int_{\mathbb{R}^N \setminus B_\epsilon(x)} \frac{|\phi(x) - \phi(y)|^{\frac{N}{s}-2} (\phi(x) - \phi(y))}{|x - y|^{2N}} dy, \quad x \in \mathbb{R}^N,$$

for all $\phi \in C_0^\infty(\Omega)$, where $B_\epsilon(x) \subset \mathbb{R}^N$ denotes an open ball centered at $x \in \mathbb{R}^N$ with radius $\epsilon > 0$.

Since the fractional Trudinger-Moser inequality (see Lemma 2.1) was established, the existence and multiplicity of solutions for various nonlinear nonlocal problems with exponential nonlinearity were investigated. We refer to [1, 2, 6, 9, 10, 12] and references therein. In particular, Perera and Squassina [8] obtained the bifurcation results of the following problems

$$\begin{cases} (-\Delta)_{N/s}^s u = \lambda |u|^{(N-2s)/s} u \exp(|u|^{N/(N-s)}), & \text{in } \Omega, \\ u = 0, & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases}$$

for appropriate $\lambda > 0$. On the other hand, there have been several papers that deal with fractional problems involving logarithmic nonlinearities. Xiang, Hu and Yang [11] considered the following Kirchhoff problems:

$$\begin{cases} M([u]_{s,p}^p) (-\Delta)_p^s u = h(x) |u|^{\theta p-2} u \ln |u| + |u|^{q-2} u, & \text{in } \Omega, \\ u = 0, & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases}$$

for $1 < p < N/s$, $\theta \in (1, p_s^*/p)$, $q \in (1, \theta p)$ and appropriate $h \in C(\bar{\Omega})$, where $p_s^* = \frac{Np}{N-sp}$ is the fractional critical Sobolev exponent. Liang et al [5], subsequently, studied the case $q = p_s^*$. However, to our best knowledge, the existence of solutions for fractional N/s -Laplacian equations with logarithmic and critical exponential nonlinearities was not tackled in the literature, which is precisely the main goal of this manuscript.

Here is our main existence result.

Theorem 1.1. *Assume that $N \geq 2$, $0 < s < 1$ and $\frac{N}{s} < q < \frac{N\sigma}{s}$. Then there exists $\mu^* > 0$ such that for any $\mu \geq \mu^*$ problem (1.1) admits a nontrivial solution.*

The outline of the paper is as follows: Section 2 contains the variational framework and some auxiliary results. And we also show that the corresponding energy functional satisfies the Palais-Smale condition below appropriate critical level. In Section 3, we prove the existence of at least one solution for problem (1.1) using some asymptotic estimates and the Mountain Pass Theorem.

2 Auxiliary results

We start this section with the fractional Sobolev space and some information about the weak formulation of problem (1.1). Throughout this paper, we work in the fractional

Sobolev space $W_0^{s,N/s}(\Omega)$, defined as the completion of $C_0^\infty(\Omega)$ with respect to the Gagliardo seminorm

$$\|u\| = [u]_{s,N/s} = \left(\int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{N/s}}{|x - y|^{2N}} dx dy \right)^{s/N}.$$

The well-known Sobolev embedding theorem states that for $\nu \in [1, \infty)$, $W_0^{s,N/s}(\Omega) \hookrightarrow L^\nu(\Omega)$ is continuous and compact. One can refer to [3] for more detailed account of the properties of $W_0^{s,N/s}(\Omega)$. To study problems involving exponential growth in the fractional Sobolev space, the main tool is the fractional Trudinger-Moser inequality due to [7] (see also [8, 13]).

Lemma 2.1. [7, Theorem 1.1] *Let Ω be a bounded, open domain of \mathbb{R}^N ($N \geq 2$) with Lipschitz boundary, and let $s \in (0, 1)$. Then there exists $\alpha_{s,N}$ such that*

$$\sup \left\{ \int_{\Omega} \exp(\alpha |u|^{N/(N-s)}) dx : u \in W_0^{s,N/s}(\Omega), [u]_{s,N/s} \leq 1 \right\} < \infty,$$

for $\alpha \in [0, \alpha_{s,N})$. Moreover,

$$\sup \left\{ \int_{\Omega} \exp(\alpha |u|^{N/(N-s)}) dx : u \in W_0^{s,N/s}(\Omega), [u]_{s,N/s} \leq 1 \right\} = \infty,$$

for $\alpha \in (\alpha_{s,N}^*, \infty)$, where

$$\alpha_{s,N}^* := N \left(\frac{2(N\omega_N)^2 \Gamma(N/s + 1)}{N!} \sum_{k=0}^{\infty} \frac{(N+k-1)!}{k!} \frac{1}{(N+2k)^{N/s}} \right)^{\frac{s}{N-s}}.$$

and ω_N is the volume of the N -dimensional unit ball.

Lemma 2.2. [7, Proposition 3.2] *If $u \in W_0^{s,N/s}(\Omega)$, then for any $\alpha \geq 0$, it holds*

$$\int_{\Omega} \exp(\alpha |u|^{N/(N-s)}) dx < \infty.$$

Let $f(t) = |t|^{(N\sigma-2s)/s} t e^{|t|^{N/(N-s)}}$ for $t \in \mathbb{R}$ and $F(t) = \int_0^t f(\tau) d\tau$. For each $\mu > 0$, a function $u \in W_0^{s,N/s}(\Omega)$ is a weak solution of problem (1.1) if

$$\begin{aligned} & \int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{(N-2s)/s} (u(x) - u(y))(v(x) - v(y))}{|x - y|^{2N}} dx dy \\ &= \int_{\Omega} |u|^{q-2} uv \ln |u|^2 dx + \mu \int_{\Omega} f(u)v dx, \quad \forall v \in W_0^{s,N/s}(\Omega). \end{aligned}$$

The last integral is well-defined in view of Lemma 2.2 and the Hölder inequality. The weak solutions of problem (1.1) coincide with the critical points of the C^1 functional

$$I_\mu(u) = \frac{s}{N} \|u\|^{N/s} + \frac{2}{q^2} \int_{\Omega} |u|^q dx - \frac{1}{q} \int_{\Omega} |u|^q \ln |u|^2 dx - \mu \int_{\Omega} F(u) dx, \quad u \in W_0^{s,N/s}(\Omega).$$

It follows from $q > N/s$ that $\lim_{t \rightarrow 0} \frac{|t|^{q-1} \ln |t|^2}{|t|^{\theta_1-1}} = 0$ and $\lim_{t \rightarrow \infty} \frac{|t|^{q-1} \ln |t|^2}{|t|^{\theta_2-1}} = 0$ for all $\theta_1 \in (N/s, q)$ and $\theta_2 \in (q, \infty)$. Then for any $\epsilon > 0$, there exists $C_\epsilon > 0$ such that

$$|t|^{q-1} \ln |t|^2 \leq \epsilon |t|^{\theta_1-1} + C_\epsilon |t|^{\theta_2-1}. \quad (2.1)$$

Besides, we observe that

$$2t^q - qt^q \ln |t|^2 \leq 2, \text{ for all } t \in (0, \infty), \quad (2.2)$$

by simple calculation.

Next, we will collect some elementary estimates of the exponential nonlinearity.

Lemma 2.3. *For all $t \in \mathbb{R}$,*

$$(i) \quad F(t) \leq \frac{1}{q} f(t)t;$$

$$(ii) \quad F(t) \leq \frac{s}{N\sigma} |t|^{\frac{N\sigma}{s}} + |t|^\theta e^{|t|^{N/(N-s)}};$$

$$(iii) \quad F(t) \geq \frac{s}{N\sigma} |t|^{\frac{N\sigma}{s}} + \frac{1}{\theta} |t|^\theta, \text{ where } \theta = \frac{N\sigma}{s} + \frac{N}{N-s}.$$

Proof. Since f is odd, $F(t) = \int_0^t f(\tau) d\tau = \int_0^{|t|} \tau^{(N\sigma-s)/s} e^{\tau^{N/(N-s)}} d\tau$.

(i) Integrating by parts,

$$F(t) = \frac{s}{N\sigma} t f(t) - \frac{s}{\sigma(N-s)} \int_0^{|t|} \tau^{\theta-1} e^{\tau^{N/(N-s)}} d\tau \leq \frac{1}{q} t f(t).$$

(ii) Since $e^\tau \leq 1 + \tau e^\tau$ for all $\tau \geq 0$,

$$F(t) \leq \int_0^{|t|} \tau^{(N\sigma-s)/s} d\tau + \int_0^{|t|} |t|^{\theta-1} e^{|t|^{N/(N-s)}} d\tau \leq \frac{s}{N\sigma} |t|^{\frac{N\sigma}{s}} + |t|^\theta e^{|t|^{N/(N-s)}}.$$

(iii) Since $e^\tau \geq 1 + \tau$ for all $\tau \geq 0$,

$$F(t) \geq \int_0^{|t|} \tau^{(N\sigma-s)/s} (1 + \tau^{N/(N-s)}) d\tau = \frac{s}{N\sigma} |t|^{\frac{N\sigma}{s}} + \frac{1}{\theta} |t|^\theta.$$

□

The next result is concerned in the $(PS)_c$ condition.

Lemma 2.4. *I_μ satisfies the $(PS)_c$ condition for all $0 < c < (\frac{s}{N} - \frac{1}{q})\alpha_{s,N}^{(N-s)/s}$.*

Proof. Let $(u_j) \subset W_0^{s,N/s}(\Omega)$ satisfy $I_\mu(u_j) \rightarrow c$ and $I'_\mu(u_j) \rightarrow 0$ for $0 < c < (\frac{s}{N} - \frac{1}{q})\alpha_{s,N}^{(N-s)/s}$. If $\inf_j \|u_j\| = 0$, then there exists a subsequence of (u_j) , still denoted by (u_j) , such that $u_j \rightarrow 0$ in $W_0^{s,N/s}(\Omega)$ as $j \rightarrow \infty$, which is impossible for $c > 0$. Thus we assume that $\inf_j \|u_j\| > 0$.

It follows from Lemma 2.3 (i) that

$$\begin{aligned}
c + o(1)\|u_j\| &= I_\mu(u_j) - \frac{1}{q}\langle I'_\mu(u_j), u_j \rangle \\
&= \left(\frac{s}{N} - \frac{1}{q}\right)\|u_j\|^{N/s} + \frac{2}{q^2} \int_\Omega |u_j|^q dx + \mu \int_\Omega \left[\frac{1}{q}f(u_j)u_j - F(u_j)\right] dx \\
&\geq \left(\frac{s}{N} - \frac{1}{q}\right)\|u_j\|^{N/s}.
\end{aligned} \tag{2.3}$$

By $\frac{N}{s} < q$, we deduce that the sequence (u_j) is bounded in $W_0^{s, N/s}(\Omega)$. Therefore, there exists some $u \in W_0^{s, N/s}(\Omega)$ such that (u_j) converges to u weakly in $W_0^{s, N/s}(\Omega)$, strongly in $L^\nu(\Omega)$ for $\nu \in [1, \infty)$ and a.e. in Ω . Then we deduce from (2.1), the Sobolev inequality and $\langle I'_\mu(u_j), u_j \rangle \rightarrow 0$ that $\int_\Omega |u_j|^q \ln |u_j|^2 dx$ and $\int_\Omega f(u_j)u_j dx$ are bounded.

Since $I'_\mu(u_j) \rightarrow 0$,

$$\begin{aligned}
&\int_{\mathbb{R}^{2N}} \frac{|u_j(x) - u_j(y)|^{(N-2s)/s} (u_j(x) - u_j(y))(v(x) - v(y))}{|x - y|^{2N}} dx dy \\
&\quad - \int_\Omega |u_j|^{q-2} u_j v \ln |u_j|^2 dx - \mu \int_\Omega f(u_j) v dx \rightarrow 0,
\end{aligned} \tag{2.4}$$

for all $v \in W_0^{s, N/s}(\Omega)$. For $v \in C_0^\infty(\Omega)$ and $M > 0$,

$$\begin{aligned}
\int_\Omega f(u_j) v dx &= \int_{\{|u_j| < M\}} f(u_j) v dx + \int_{\{|u_j| \geq M\}} f(u_j) v dx \\
&\leq \int_{\{|u_j| < M\}} f(u_j) v dx + \frac{\sup |v|}{M} \int_\Omega f(u_j) u_j dx \\
&= \int_{\{|u_j| < M\}} f(u_j) v dx + \mathcal{O}\left(\frac{1}{M}\right),
\end{aligned}$$

which shows that

$$\int_\Omega f(u_j) v dx \rightarrow \int_\Omega f(u) v dx, \tag{2.5}$$

by letting $j \rightarrow \infty$ first and then $M \rightarrow \infty$. Similarly, we also have that

$$\int_\Omega |u_j|^{q-2} u_j v \ln |u_j|^2 dx \rightarrow \int_\Omega |u|^{q-2} u v \ln |u|^2 dx, \tag{2.6}$$

as $j \rightarrow \infty$. It follows from (2.4)-(2.6) that

$$\begin{aligned}
&\int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{(N-2s)/s} (u(x) - u(y))(v(x) - v(y))}{|x - y|^{2N}} dx dy \\
&= \int_\Omega |u|^{q-2} u v \ln |u|^2 dx + \mu \int_\Omega f(u) v dx.
\end{aligned}$$

Then the last equality holds for all $v \in W_0^{s, N/s}(\Omega)$ by density and

$$\|u\|^{N/s} = \int_{\Omega} |u|^q \ln |u|^2 dx + \mu \int_{\Omega} f(u)u dx \quad (2.7)$$

holds by taking $v = u$.

Next, we claim that

$$\int_{\Omega} f(u_j)u_j dx \rightarrow \int_{\Omega} f(u)u dx. \quad (2.8)$$

On one hand, by (2.3), Lemma 2.1 and choosing $t > 1$ such that $t\|u_j\|^{N/(N-s)} < \alpha_{s,N}$, we can conclude that $\int_{\Omega} e^{t|u_j|^{N/(N-s)}} dx \leq M$. Since $e^{t|u_j|^{N/(N-s)}} \rightarrow e^{t|u|^{N/(N-s)}}$ a.e. in Ω , we use [4, Lemma 4.8] and conclude that

$$e^{|u_j|^{N/(N-s)}} \rightharpoonup e^{|u|^{N/(N-s)}} \text{ in } L^t(\Omega). \quad (2.9)$$

On the other hand, note that

$$|u_j|^{N\sigma/s} \rightarrow |u|^{N\sigma/s} \text{ in } L^{t'}(\Omega), \quad (2.10)$$

for $t' > 1$ with $\frac{1}{t} + \frac{1}{t'} = 1$.

Then it follows from the Hölder inequality, (2.9), (2.10) and Lemma 2.1 that

$$\begin{aligned} \int_{\Omega} [f(u_j)u_j - f(u)u] dx &= \int_{\Omega} [|u_j|^{N\sigma/s} - |u|^{N\sigma/s}] e^{|u_j|^{N/(N-s)}} dx \\ &\quad + \int_{\Omega} |u|^{N\sigma/s} [e^{|u_j|^{N/(N-s)}} - e^{|u|^{N/(N-s)}}] dx \\ &\leq \left(\int_{\Omega} [|u_j|^{N\sigma/s} - |u|^{N\sigma/s}]^{t'} dx \right)^{1/t'} \left(\int_{\Omega} e^{t|u_j|^{N/(N-s)}} dx \right)^{1/t} \\ &\quad + \int_{\Omega} |u|^{N\sigma/s} [e^{|u_j|^{N/(N-s)}} - e^{|u|^{N/(N-s)}}] dx \\ &\leq M \left(\int_{\Omega} [|u_j|^{N\sigma/s} - |u|^{N\sigma/s}]^{t'} dx \right)^{1/t'} \\ &\quad + \int_{\Omega} |u|^{N\sigma/s} [e^{|u_j|^{N/(N-s)}} - e^{|u|^{N/(N-s)}}] dx \\ &\rightarrow 0 \text{ as } j \rightarrow \infty, \end{aligned}$$

which complete the proof of (2.8).

Now, we are ready to show that the sequence (u_j) converges strongly to u in $W_0^{s, N/s}(\Omega)$. Indeed, by the weak lower semicontinuity of the norm, the Lebesgue dominated convergence

theorem, (2.1), (2.7) and (2.8), we obtain that

$$\begin{aligned}
0 &= \lim_{j \rightarrow \infty} \langle I'_\mu(u_j), u_j \rangle \\
&= \lim_{j \rightarrow \infty} [\|u_j\|^{N/s} - \int_\Omega |u_j|^q \ln |u_j|^2 dx - \mu \int_\Omega f(u_j) u_j dx] \\
&\geq \|u\|^{N/s} - \int_\Omega |u|^q \ln |u|^2 dx - \mu \int_\Omega f(u) u dx \\
&= 0,
\end{aligned}$$

then (u_j) converges strongly to u in $W_0^{s, N/s}(\Omega)$. This completes the proof. \square

3 Proof of Theorem 1.1

Lemma 3.1. *The functional I_μ has a mountain pass geometry.*

Proof. From (2.1), Lemma 2.3 (ii), the Hölder inequality, Theorem 2.1 and Sobolev embedding inequality, we have that

$$\begin{aligned}
I_\mu(u) &= \frac{s}{N} \|u\|^{N/s} + \frac{2}{q^2} \int_\Omega |u|^q dx - \frac{1}{q} \int_\Omega |u|^q \ln |u|^2 dx - \mu \int_\Omega F(u) dx \\
&\geq \frac{s}{N} \|u\|^{N/s} - \frac{\epsilon}{q} \int_\Omega |u|^{\theta_1} dx - \frac{C_\epsilon}{q} \int_\Omega |u|^{\theta_2} dx - \frac{\mu s}{N\sigma} \int_\Omega |u|^{N\sigma/s} dx \\
&\quad - \mu \int_\Omega |u|^\theta e^{|u|^{N/(N-s)}} dx \\
&\geq \frac{s}{N} \|u\|^{N/s} - \frac{\epsilon}{q} \int_\Omega |u|^{\theta_1} dx - \frac{C_\epsilon}{q} \int_\Omega |u|^{\theta_2} dx - \frac{\mu s}{N\sigma} \int_\Omega |u|^{N\sigma/s} dx \\
&\quad - \left(\int_\Omega |u|^{t'\theta} dx \right)^{1/t'} \left(\int_\Omega e^{t|u|^{N/(N-s)}} dx \right)^{1/t} \\
&\geq \frac{s}{N} \|u\|^{N/s} - \frac{\epsilon}{q} C_1 \|u\|^{\theta_1} - \frac{C_\epsilon}{q} C_2 \|u\|^{\theta_2} - \frac{\mu s}{N\sigma} C_3 \|u\|^{N\sigma/s} - C_4 \|u\|^\theta,
\end{aligned}$$

where $C_i, i = 1, 2, 3, 4$ are some positive constants and $t, t' > 1$ with $\frac{1}{t} + \frac{1}{t'} = 1$. Since $q, \theta_1, \theta_2, \theta > \frac{N}{s}$, there exist $\rho, \alpha > 0$ such that $I_\mu(u) \geq \alpha$ for $\|u\| = \rho$.

Let $u_0 \in W_0^{s, N/s}(\Omega)$ with $\|u_0\| = 1$. Thus it follows from Lemma 2.3 (iii) and (2.2).

$$\begin{aligned}
I_\mu(tu_0) &= \frac{s}{N} \|tu_0\|^{N/s} + \frac{1}{q^2} \int_\Omega [2|tu_0|^q - q|tu_0|^q \ln |tu_0|^2] dx - \mu \int_\Omega F(tu_0) dx \\
&\leq \frac{s}{N} \|tu_0\|^{N/s} + \frac{2|\Omega|}{q^2} - \frac{\mu s}{N\sigma} \int_\Omega |tu_0|^{N\sigma/s} dx - \frac{\mu}{\theta} \int_\Omega |tu_0|^\theta dx \\
&\rightarrow -\infty, \text{ as } t \rightarrow \infty.
\end{aligned}$$

Then we deduce that $I_\mu(t_0 u_0) < 0$ and $\|t_0 u_0\| > \rho$ for $t_0 > 0$ large enough, which completes the proof. \square

Proof of Theorem 1.1. Let

$$\Gamma = \{\gamma \in C([0, 1], W_0^{s, N/s}(\Omega)) : \gamma(0) = 0 \text{ and } \gamma(1) = t_0 u_0\}$$

be the class of joining 0 and $t_0 u_0$. Next, we claim that

$$c_\mu := \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} I_\mu(\gamma(t)) < \left(\frac{s}{N} - \frac{1}{q}\right) \alpha_{s, N}^{(N-s)/s}, \quad (3.1)$$

for large μ . Assuming that (3.1) holds true, we can deduce from Lemma 3.1, Lemma 2.4 and the Mountain Pass Theorem that there exists at least one nontrivial critical point of I_μ .

Note that $\lim_{t \rightarrow \infty} I_\mu(tu_0) = -\infty$ for $u_0 \in W_0^{s, N/s}(\Omega)$ with $\|u_0\| = 1$, from which there holds $\sup_{t \geq 0} I_\mu(tu_0) = I_\mu(t_\mu u_0)$ for some $t_\mu > 0$. Therefore, by Lemma 2.3 (i), (iii), t_μ satisfies that

$$\begin{aligned} t_\mu^{N/s} &= \int_\Omega |t_\mu u_0|^q \ln |t_\mu u_0|^2 dx + \mu \int_\Omega f(t_\mu u_0) t_\mu u_0 dx \\ &\geq \int_\Omega |t_\mu u_0|^q \ln |t_\mu u_0|^2 dx + \frac{sq}{N\sigma} \mu \int_\Omega |t_\mu u_0|^{N\sigma/s} dx + \frac{q}{\theta} \mu \int_\Omega |t_\mu u_0|^\theta dx. \end{aligned} \quad (3.2)$$

The sequence (t_μ) is bounded since $\theta > N/s$. Now we claim that $t_\mu \rightarrow 0$ as $\mu \rightarrow \infty$. Suppose, by contradiction, that there exist $t_{\mu_0} > 0$ and a sequence $\mu_n \rightarrow \infty$ as $n \rightarrow \infty$ such that $t_{\mu_n} \rightarrow t_{\mu_0}$ as $n \rightarrow \infty$. By the Lebesgue's convergence theorem and (2.1), we have that $\int_\Omega |t_{\mu_n} u_0|^q \ln |t_{\mu_n} u_0|^2 dx \rightarrow \int_\Omega |t_{\mu_0} u_0|^q \ln |t_{\mu_0} u_0|^2 dx$ as $n \rightarrow \infty$. It follows that $t_{\mu_0}^{N/s} \geq \infty$, which is absurd. Therefore, $t_\mu \rightarrow 0$ as $\mu \rightarrow \infty$.

We deduce from (3.2) that $\lim_{\mu \rightarrow \infty} \mu \int_\Omega f(t_\mu u_0) t_\mu u_0 dx = 0$ and

$$\lim_{\mu \rightarrow \infty} [\sup_{t \geq 0} I_\mu(tu_0)] = \lim_{\mu \rightarrow \infty} I_\mu(t_\mu u_0) = 0.$$

Then there exists $\mu_* > 0$ such that $c_\mu \leq \sup_{t \geq 0} I_\mu(tu_0) < \left(\frac{s}{N} - \frac{1}{q}\right) \alpha_{s, N}^{(N-s)/s}$. The proof of Theorem 1.1 is now complete. \square

References

- [1] W. J. Chen and F. Yu. On a fractional Kirchhoff type problem with critical exponential growth nonlinearity. *Applied Mathematics Letters*, 105:106279, 2020. [2](#)
- [2] M. de Souza and Y. L. Araújo. Semilinear elliptic equations for the fractional Laplacian involving critical exponential growth. *Mathematical Methods in the Applied Sciences*, 40(5):1757–1772, 2017. [2](#)

- [3] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bulletin Des Sciences Mathématiques*, 136(5):521–573, 2012. [3](#)
- [4] O. Kavian. *Introduction à la Théorie des Points Critiques*. Springer-Verlag, Berlin, 1991. [6](#)
- [5] S. Liang, H. Pu, and V. D. Rădulescu. High perturbations of critical fractional Kirchhoff equations with logarithmic nonlinearity. *Applied Mathematics Letters*, 116:107027, 2021. [1](#), [2](#)
- [6] O. H. Miyagaki and P. Pucci. Nonlocal Kirchhoff problems with Trudinger-Moser critical nonlinearities. *Nonsmooth Nonlinear Differential Equations and Applications*, 26(4):27, 2019. [2](#)
- [7] E. Parini and B. Ruf. On the Moser-Trudinger inequality in fractional Sobolev-Slobodeckij spaces. *Journal D Analyse Mathématique*, 138(1):281–300, 2019. [3](#)
- [8] K. Perera and M. Squassina. Bifurcation results for problems with fractional Trudinger-Moser nonlinearity. *Discrete and Continuous Dynamical Systems-Series S*, 11(3):561–576, 2018. [2](#), [3](#)
- [9] M. d. Souza, U. B. Severo, and T. L. d. Rêgo. Nodal solutions for fractional elliptic equations involving exponential critical growth. *Mathematical Methods in the Applied Sciences*, 43(6):3650–3672, 2020. [2](#)
- [10] M. d. Souza, U. B. Severo, and T. L. d. Rêgo. On solutions for a class of fractional Kirchhoff-type problems with Trudinger–Moser nonlinearity. *Communications in Contemporary Mathematics*, 2021. [2](#)
- [11] M. Xiang, D. Hu, and D. Yang. Least energy solutions for fractional Kirchhoff problems with logarithmic nonlinearity. *Nonlinear Analysis-Theory Methods & Applications*, 198:111899, 2020. [2](#)
- [12] M. Q. Xiang, V. D. Rădulescu, and B. L. Zhang. Fractional Kirchhoff problems with critical Trudinger-Moser nonlinearity. *Calculus of Variations and Partial Differential Equations*, 58(2):57, 2019. [2](#)
- [13] C. Zhang. Trudinger-Moser inequalities in Fractional Sobolev-Slobodeckij spaces and multiplicity of weak solutions to the Fractional-Laplacian equation. *Advanced Nonlinear Studies*, 19(1):197–217, 2019. [3](#)