# FAST REACTION LIMIT AND FORWARD-BACKWARD DIFFUSION: A RADON-NIKODYM APPROACH

### JAKUB SKRZECZKOWSKI

ABSTRACT. We consider two singular limits: fast reaction limit with nonmonotone nonlinearity and regularization of forward-backward diffusion equation. It was proved by Plotnikov that for cubic-type (nondegenerate) nonlinearities, the limit oscillates between at most three states. In this paper we make his argument more optimal and we sharpen the previous result: we use Radon-Nikodym theorem to obtain a pointwise identity characterizing the Young measure. As a consequence, we establish a simpler condition which implies Plotnikov result for piecewise affine functions. We also prove that the result is true if the Young measure is not supported in the so-called unstable zone, the fact observed in numerical simulations.

# Contents

1.	Introduction and main results	2
2.	Properties of the fast-reaction system (1.1)	8
3.	Proof of Theorem 1 for fast-reaction system $(1.1)$	9
4.	Proofs of Theorems 2, 3 and 4 for fast-reaction system $(1.1)$	12
5.	Proof of Theorems 1–3 to the forward-backward diffusion system $(1.2)$	15
Ap	ppendix A. Useful notions and results	17
References		10

 $<sup>2020\</sup> Mathematics\ Subject\ Classification.\ 35K57,\ 35B25,\ 35B36.$ 

Key words and phrases. reaction-diffusion, fast reaction limit, forward-backward diffusion, Young measures, cross-diffusion, oscillations, compensated compactness.

Jakub Skrzeczkowski was supported by National Science Center, Poland through project no. 2017/27/B/ST1/01569. He is grateful to Benoît Perthame for fruitful discussions and helpful suggestions.




FIGURE 1. Plot of a typical function F. It is strictly increasing in the intervals  $I_1 := (-\infty, \alpha_+], I_3 := [\beta_-, \infty)$  and strictly decreasing in  $I_2 := (\alpha_+, \beta_-)$ . For  $r \in [f_-, f_+]$ , the function F is not invertible and equation F(u) = r has three roots  $u = S_1(r) \le S_2(r) \le S_3(r)$ .

#### 1. Introduction and main results

1.1. **Presentation of the problem.** In this paper we focus on two interesting limit problems: fast-reaction limit in the reaction-diffusion system

(1.1) 
$$\partial_t u^{\varepsilon} = \frac{F(u^{\varepsilon}) - v^{\varepsilon}}{\varepsilon},$$
$$\partial_t v^{\varepsilon} = \Delta v^{\varepsilon} + \frac{v^{\varepsilon} - F(u^{\varepsilon})}{\varepsilon}$$

and regularization of the forward-backward parabolic equation  $\partial_t u = \Delta F(u)$ 

(1.2) 
$$\partial_t u^{\varepsilon} = \Delta v^{\varepsilon},$$
$$v^{\varepsilon} = F(u^{\varepsilon}) + \varepsilon \, \partial_t u^{\varepsilon},$$

where F is a nonmonotone function, for simplicity assumed to look like as in Fig. 1. Notice that due to nonmonotone character of F, it has three inverses  $S_1$ ,  $S_2$  and  $S_3$ . Both problems are posed on some bounded domain  $\Omega \subset \mathbb{R}^d$  and are equipped with initial conditions and usual Neumann boundary conditions.

System (1.1) is an interesting toy model for studying oscillations in reaction-diffusion systems as they are known to occur in its steady states [28]. For monotonone F the problem is fairly classical

and has been studied for a great variety of reaction-diffusion systems, also with more than two components [5,6,14,29] or reaction-diffusion equation coupled with an ODE [21]. In the limit  $\varepsilon \to 0$  one obtains widely studied cross-diffusion systems [9,10,15,16,22,24] where the gradient of one quantity induces a flux of another one. A slightly different yet connected type of problem deals with the fast-reaction limit for irreversible reactions which leads to free boundary problems [11,17,20]. Finally, for nonmonotone F as in this paper, the only available result was established very recently in [33] (see below). We also refer to the recent stability analysis of problems of the type (1.1) [12,13,25].

System (1.2) was extensively studied by Plotnikov [34,35] who identified limits as  $\varepsilon \to 0$  in terms of Young measures (see below) and by Novick-Cohen and Pego who studied its asymptotics with  $\varepsilon > 0$  fixed [31]. Regularization term in (1.2) was also generalized in [3,4,40]. Recently, so called nonstandard analysis was used to study the limit problem in the space of grid functions [7,8].

It is known [33, 35] that both systems exhibit the following surprising phenomenon: as  $\varepsilon \to 0$ ,  $F(u^{\varepsilon}) \to v$  and  $v^{\varepsilon} \to v$  converge strongly without any known a priori estimates allowing to conclude so. As a consequence,  $u^{\varepsilon}$  converges weakly to

$$u(t,x) = \lambda_1(t,x) S_1(v(t,x)) + \lambda_2(t,x) S_2(v(t,x)) + \lambda_3(t,x) S_3(v(t,x))$$

where  $\sum_{i=1}^{3} \lambda_i(t,x) = 1$ . More precisely, if  $\mu_{t,x}$  is a Young measure generated by  $\{u^{\varepsilon}\}_{{\varepsilon}>0}$  we have

$$\mu_{t,x} = \lambda_1(t,x) \, \delta_{S_2(v(t,x))} + \lambda_2(t,x) \, \delta_{S_2(v(t,x))} + \lambda_3(t,x) \, \delta_{S_2(v(t,x))}$$

which represents oscillations between phases  $S_1(v(t,x))$ ,  $S_2(v(t,x))$  and  $S_3(v(t,x))$ . The proof exploits a family of energies as well as analysis of related Young measures in the spirit of Murat and Tartar work on conservation laws and compensated compactness [30,41]. The numerical simulations suggests that the middle state, referred to as an unstable phase, is not present [19] which motivates research on two-phase solutions to such problems [23,26,39,42] with a result of nonuniqueness when the unstable phase is present [43].

So far, the main assumption on F that allows to deduce strong convergence is the so-called nondegeneracy condition: for (1.1) it reads

(1.3) for all intervals 
$$R \subset (f_-, f_+)$$
:  $\sum_{i=1}^3 a_i \left( S_1'(r) + 1 \right) = 0$  for  $r \in R \implies a_1 + a_2 + a_3 = 0$ 

while for (1.2) it reads

(1.4) for all intervals 
$$R \subset (f_-, f_+)$$
:  $\sum_{i=1}^3 a_i S_1'(r) = 0$  for  $r \in R \implies a_1 + a_2 + a_3 = 0$ .

While it is fairly classical for this type of problems [1,31,35], it is hard to be verified for a given nonlinearity F. Moreover, the nondegeneracy condition excludes piecewise affine functions which allows for more explicit computations as in [26].

1.2. Main results and outline of the paper. In this paper, we take a slightly different approach to study strong convergence. Although we use family of energy identities to characterize Young measure as Plotnikov [35], we aim at pointwise identities to obtain optimal amount of information from these energy identities, in particular new results. To achieve this, we use Radon-Nikodym Theorem as explained below.

Let  $\{\mu_{t,x}\}_{t,x}$  be Young measure generated by sequence  $\{u^{\varepsilon}\}_{\varepsilon\in(0,1)}$  solving either (1.1) or (1.2), i.e. for any bounded function  $G: \mathbb{R} \to \mathbb{R}$  we have (up to a subsequence and for a.e.  $(t,x) \in (0,T) \times \Omega$ )

$$G(u^{\varepsilon}) \stackrel{*}{\rightharpoonup} \int_{\mathbb{R}} G(\lambda) \, \mathrm{d}\mu_{t,x}(\lambda),$$

see Appendix A.3 if necessary. To analyze amount of  $\mu_{t,x}$  on the intervals  $I_1$ ,  $I_2$  and  $I_3$ , see Figure 1, we introduce restrictions

$$\mu_{t,x}^{(1)} := \mu_{t,x} \, \mathbb{1}_{I_1}, \qquad \qquad \mu_{t,x}^{(2)} := \mu_{t,x} \, \mathbb{1}_{I_2}, \qquad \qquad \mu_{t,x}^{(3)} := \mu_{t,x} \, \mathbb{1}_{I_3}.$$

The reason we introduce these measures is that in the sequel, we will gain information only about measure  $F^{\#}\mu_{t,x}$ , i.e. push-forward (image) of  $\mu_{t,x}$  along F defined as

$$F^{\#}\mu_{t,x} = \mu_{t,x}(F^{-1}(A)), \qquad A \subset \mathbb{R}^+.$$

Observe that for all i = 1, 2, 3 measures  $F^{\#}\mu_{t,x}^{(i)}$  are absolutely continuous with respect to  $F^{\#}\mu_{t,x}$ . Therefore, Radon-Nikodym theorem implies that there exist densities  $g^{(1)}(\lambda)$ ,  $g^{(2)}(\lambda)$  and  $g^{(3)}(\lambda)$  such that

(1.5) 
$$F^{\#}\mu_{t,x}^{(i)}(A) = \int_{A} g^{(i)}(\lambda) \, \mathrm{d}F^{\#}\mu_{t,x}(\lambda), \qquad i = 1, 2, 3.$$

We also note that for all  $A \subset \mathbb{R}^+$ 

(1.6) 
$$\sum_{i=1}^{3} F^{\#} \mu_{t,x}^{(i)}(A) = \sum_{i=1}^{3} \mu_{t,x}(F^{-1}(A) \cap I_i) = \mu_{t,x}(F^{-1}(A)) = F^{\#} \mu_{t,x}(A).$$

In particular, from (1.5) and (1.6) we deduce that for  $F^{\#}\mu_{t,x}$ -a.e.  $\lambda$  we have

(1.7) 
$$\sum_{i=1}^{3} g_i(\lambda) = 1.$$

The main result of this paper reads:

**Theorem 1.** (A) Let  $\{\mu_{t,x}\}_{t,x}$  be Young measure generated by sequence  $\{u^{\varepsilon}\}_{\varepsilon\in(0,1)}$  solving (1.1). Then, for almost all  $\lambda_0$  (with respect to  $F^{\#}\mu_{t,x}$ ) and all  $\tau_0 \neq f_-, f_+$  we have

$$\sum_{i=1}^{3} (S_i'(\tau_0) + 1) \left[ \mathbb{1}_{\lambda_0 > \tau_0} g_i(\lambda_0) - F^{\#} \mu_{t,x}^{(i)}(\tau_0, \infty) \right] + \left( S_1'(\tau_0) - S_2'(\tau_0) \right) \left( F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_0) \right) = 0.$$

where  $S_i$  are inverses of F as in Notation 1.1 and  $g_i$  are Radon-Nikodym densities as in (1.5). Moreover, if  $\lambda_0 \neq f_-, f_+$  it holds

(1.8) 
$$\left(1 - F^{\#} \mu_{t,x} \left\{\lambda_0\right\}\right) \sum_{i=1}^{3} \left(S_i'(\lambda_0) + 1\right) g_i(\lambda_0) = 0.$$

(B) If  $\{\mu_{t,x}\}_{t,x}$  is the Young measure generated by sequence  $\{u^{\varepsilon}\}_{\varepsilon\in(0,1)}$  solving (1.2) the equalities above holds with functions  $S'_i$  instead of  $S'_i+1$ .

As  $F^{\#}\mu_{t,x}$  turns out to be the Young measure generated by  $\{v^{\varepsilon}\}_{\varepsilon>0}$  cf. Corollary 2.3, strong convergence  $v^{\varepsilon} \to v$  follows from proving that  $F^{\#}\mu_{t,x}$  is a Dirac mass cf. Lemma A.5 (A). Equation (1.8) shows that it is sufficient to find  $\lambda_0$  in the support such that the sum  $\sum_{i=1}^{3} (S'_i(\lambda_0) + 1) g_i(\lambda_0)$  does not vanish (some additional care is needed when  $\lambda_0 = f_-, f_+$ , cf. Lemma 4.1).

We remark that similar forms of entropy equality as in Theorem 1 are well-known however they are not so easily formulated and they are usually stated without explicitly identified coefficients standing next to  $(S'_i(\tau_0) + 1)$ . First, we show that the form presented in Theorem 1 can be used to recover already known result due to Plotnikov [35] as well as Perthame and Skrzeczkowski [33].

**Theorem 2.** Suppose that nondegeneracy condition (1.3) or (1.4) is satisfied. Then,  $v^{\varepsilon} \to v$  strongly in  $L^2((0,T) \times \Omega)$ . Moreover, there are nonnegative numbers  $\lambda_1(t,x)$ ,  $\lambda_2(t,x)$ ,  $\lambda_3(t,x)$  such that  $\sum_{i=1}^3 \lambda_i(t,x) = 1$  and

$$\mu_{t,x} = \lambda_1(t,x) \, \delta_{S_1(v(t,x))} + \lambda_2(t,x) \, \delta_{S_2(v(t,x))} + \lambda_3(t,x) \, \delta_{S_3(v(t,x))}.$$

Now, we move to the new results that easily follow from Theorem 1. The first one asserts that if one knows a priori that the Young measure  $\mu_{t,x}$  is not supported in the interval  $I_2$  where F is

decreasing, the strong convergence occurs. The fact concerning support of  $\mu_{t,x}$  was observed in numerical simulations [19] and so, the next theorem may serve as a tool to prove strong convergence without nondegeneracy condition.

# **Theorem 3.** Suppose that:

- there exists  $\tau_0 \in (f_-, f_+)$  such that  $S'_1(\tau_0) S'_3(\tau_0) \neq 0$ ,
- the Young measure  $\{\mu_{t,x}\}_{t,x}$  is not supported in the interval  $I_2$  (see Figure 1).

Then,  $v^{\varepsilon} \to v$  strongly in  $L^{2}((0,T) \times \Omega)$ . Moreover, there are nonnegative numbers  $\lambda_{1}(t,x)$ ,  $\lambda_{3}(t,x)$  such that  $\lambda_{1}(t,x) + \lambda_{3}(t,x) = 1$  and

$$\mu_{t,x} = \lambda_1(t,x) \, \delta_{S_1(v(t,x))} + \lambda_3(t,x) \, \delta_{S_3(v(t,x))}.$$

The next result shows that the systems (1.1) and (1.2) are not exactly the same in view of the strong convergence. Indeed, for the first one, we can establish a simple condition on F implying strong convergence of  $v^{\varepsilon} \to v$  that does not exclude piecewise affine functions as in the case of nondegeneracy condition (1.3).

**Theorem 4.** Let  $\{\mu_{t,x}\}_{t,x}$  be a Young measure generated by sequence  $\{u^{\varepsilon}\}_{\varepsilon\in(0,1)}$  solving (1.1). Suppose that:

- there exists  $\tau_0 \in (f_-, f_+)$  such that  $S_1'(\tau_0) S_3'(\tau_0) \neq 0$ ,
- $S'_2(\lambda) + 1 > 0$  for all  $\lambda \in (f_-, f_+)$ .

Then,  $v^{\varepsilon} \to v$  strongly in  $L^{2}((0,T) \times \Omega)$ . Moreover, there are nonnegative numbers  $\lambda_{1}(t,x)$ ,  $\lambda_{2}(t,x)$ ,  $\lambda_{3}(t,x)$  such that  $\sum_{i=1}^{3} \lambda_{i}(t,x) = 1$  and

$$\mu_{t,x} = \lambda_1(t,x) \, \delta_{S_1(v(t,x))} + \lambda_2(t,x) \, \delta_{S_2(v(t,x))} + \lambda_3(t,x) \, \delta_{S_3(v(t,x))}.$$

As an example of function F satisfying assumptions of Theorem 4 consider

$$F(\lambda) = \begin{cases} 2\lambda & \text{if } \lambda \in [0, 1], \\ 3 - 2\lambda & \text{if } \lambda \in \left[1, \frac{5}{4}\right], \\ 4\lambda - \frac{9}{2} & \text{if } \lambda \in \left[\frac{5}{4}, \infty\right) \end{cases}$$

Then,  $S_1'(\lambda) = \frac{1}{2}$ ,  $S_2'(\lambda) = -\frac{1}{2}$  and  $S_3'(\lambda) = \frac{1}{4}$  so that  $S_1'(\lambda) - S_3'(\lambda) = \frac{1}{4} \neq 0$  and  $S_2'(\lambda) + 1 = \frac{1}{2} > 0$ . Note that F does not satisfy nondegeneracy condition (1.3) that was used in the previous paper on fast reaction limit with nonmonotone reaction function [33]. Proofs of Theorem 3 and 4 are based on equation (1.8), namely one uses  $g_1(\lambda_0) + g_2(\lambda_0) + g_3(\lambda_0) = 1$  to show that for  $\lambda_0 \in \text{supp } F$  we have  $F^{\#}\mu_{t,x}\{\lambda_0\} = 1$ . Note however that (1.8) is not valid for  $\lambda_0 = f_-, f_+$  so that some additional care is needed if the support of measure  $F^{\#}\mu_{t,x}$  accumulates only in these points. This is studied in Lemma 4.1 and it requires an additional assumption that  $S'_1(\tau) - S'_3(\tau)$  does not vanish at least for one value of  $\tau$ , see also Remark 4.2.

The structure of the paper is as follows. In Section 2 we review (well-known) properties of the fast-reaction system (1.1). Then, in Section 3 we use compensated compactness approach to prove Theorem 1. Section 4 is devoted to the simple proofs of Theorems 2, 3 and 4 while in Section 5 we show how to easily adapt Theorems 1–3 to the case of system (1.2). Finally, Appendix A provides necessary background on Young measures, supports of measures and compensated compactness results.

1.3. **Technical assumptions and notation.** For the sake of completeness, we list here assumptions of technical nature.

**Notation 1.1.** Let  $S_1(\lambda) \leq S_2(\lambda) \leq S_3(\lambda)$  be the solutions of equation  $F(S_i(\lambda)) = \lambda$  as already introduced in Figure 1. These are inverses of F satisfying

$$S_1: (-\infty, f_+] \to (-\infty, \alpha_+], \qquad S_2: (f_-, f_+) \to (\alpha_+, \beta_-), \qquad S_3: [f_-, \infty) \to [\beta_-, \infty).$$

Their role is too focus analysis on parts of the plot of F where the monotonicity of F does not change. By a small abuse of notation, we extend functions  $S_i$  by a constant value to the whole of  $\mathbb{R}$ . We usually write, for images of functions  $S_1$ ,  $S_2$ ,  $S_3$  and for their domains

$$I_1 = (-\infty, \alpha_+],$$
  $I_2 = (\alpha_+, \beta_-),$   $I_3 = [\beta_-, \infty),$ 

$$J_1 = (-\infty, f_+],$$
  $J_2 = (f_-, f_+),$   $J_3 = [f_-, \infty).$ 

**Assumption 1.2** (Initial data for (1.1)). Functions  $u^{\varepsilon}(0,x) = u_0(x)$ ,  $v^{\varepsilon}(0,x) = v_0(x)$  satisfy

- (1) Nonnegativity:  $u_0, v_0 \ge 0$ .
- (2) Regularity:  $u_0, v_0 \in C^{2+\alpha}(\overline{\Omega})$  for some  $\alpha \in (0,1)$ .
- (3) Boundary condition:  $u_0, v_0$  satisfy the Neumann boundary condition.

**Assumption 1.3** (Reaction function F). We assume that the function F(u) satisfies:

(1) Regularity, nonnegativity: F is Lipschitz continuous with F(0) = 0 and  $F \ge 0$ .

- (2) Piecewise monotonicity of F: there are  $\alpha_{-} < \alpha_{+} < \beta_{-} < \beta_{+}$  such that  $F(\beta_{-}) = F(\alpha_{-})$ ,  $F(\alpha_{+}) = F(\beta_{+})$ , F is strictly increasing on  $(-\infty, \alpha_{+}) \cup (\beta_{-}, \infty)$  and strictly decreasing on  $(\alpha_{+}, \beta_{-})$  (see Fig. 1). Moreover,  $\lim_{u \to \infty} F(u) = \infty$ .
- (3) Regularity of inverses: functions  $\lambda \mapsto S'_i(\lambda)$  are continuous except  $\lambda = f_-, f_+$ .

# 2. Properties of the fast-reaction system (1.1)

We begin from recalling energy equality and well-posedness result from [33] which we prove below for the sake of completeness.

**Lemma 2.1** (energy equality). Given a smooth test function  $\phi : \mathbb{R} \to \mathbb{R}$ , we define

(2.1) 
$$\Psi(\lambda) := \int_0^{\lambda} \phi(F(\tau)) d\tau, \qquad \Phi(\lambda) := \int_0^{\lambda} \phi(\tau) d\tau.$$

Then, if  $(u^{\varepsilon}, v^{\varepsilon})$  solve (1.1), it holds

(2.2) 
$$\partial_t \Psi(u^{\varepsilon}) + \partial_t \Phi(v^{\varepsilon}) = \Delta \Phi(v^{\varepsilon}) - \phi'(v^{\varepsilon}) |\nabla v^{\varepsilon}|^2 - \frac{\left(v^{\varepsilon} - F(u^{\varepsilon})\right) \left(\phi(v^{\varepsilon}) - \phi(F(u^{\varepsilon}))\right)}{\varepsilon}$$

*Proof.* Multiplying equation for  $u^{\varepsilon}$  in (1.1) with  $\phi(F(u^{\varepsilon}))$  and equation for  $v^{\varepsilon}$  in (1.1) with  $\phi(v^{\varepsilon})$  we obtain

$$\partial_t \Psi(u^{\varepsilon}) = \frac{v^{\varepsilon} - F(u^{\varepsilon})}{\varepsilon} \phi(F(u^{\varepsilon})),$$

$$\partial_t \Phi(v^{\varepsilon}) = \Delta \Phi(v^{\varepsilon}) - \phi'(v^{\varepsilon}) |\nabla v^{\varepsilon}|^2 + \frac{F(u^{\varepsilon}) - v^{\varepsilon}}{\varepsilon} \phi(v^{\varepsilon}).$$

Summing up these equations we deduce (2.2).

**Lemma 2.2.** There exists the unique classical solution  $u^{\varepsilon}, v^{\varepsilon} : [0, \infty) \times \Omega \to \mathbb{R}$  of (1.1) which is nonnegative and has regularity

$$u^{\varepsilon} \in C^{\alpha,1+\alpha/2}\left([0,\infty)\times\overline{\Omega}\right), \qquad v^{\varepsilon} \in C^{2+\alpha,1+\alpha/2}\left([0,\infty)\times\overline{\Omega}\right).$$

Moreover, we have

- (1)  $0 \le u^{\varepsilon} \le M$ ,  $0 \le v^{\varepsilon} \le M$  with  $M = \max(\|F(u_0)\|_{\infty}, \|u_0\|_{\infty}, \|v_0\|_{\infty}, f_+, \beta_+)$ ,
- (2)  $\{\nabla v^{\varepsilon}\}_{\varepsilon\in(0,1)}$  is uniformly bounded in  $L^{2}((0,\infty)\times\Omega)$ ,
- (3)  $\left\{\frac{F(u^{\varepsilon})-v^{\varepsilon}}{\sqrt{\varepsilon}}\right\}_{\varepsilon\in(0,1)}$  and  $\left\{\sqrt{\varepsilon}\,\Delta v^{\varepsilon}\right\}_{\varepsilon\in(0,1)}$  are uniformly bounded in  $L^{2}((0,\infty)\times\Omega)$ ,
- (4)  $\{\partial_t u^{\varepsilon} + \partial_t v^{\varepsilon}\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $L^2(0,T;H^{-1}(\Omega))$ ,
- (5) for all smooth  $\varphi : \mathbb{R} \to \mathbb{R}$ ,  $\{\nabla \varphi(v^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $L^{2}((0,\infty) \times \Omega)$ ,
- (6) for all smooth  $\phi : \mathbb{R} \to \mathbb{R}$ ,  $\{\partial_t \Psi(u^{\varepsilon}) + \partial_t \Phi(v^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $(C(0,T;H^k(\Omega)))^*$  for sufficiently large  $k \in \mathbb{N}$ .

*Proof.* Existence and uniqueness of global solution as well as points (1)–(3) were proven in [33, Theorem 3.1] so we only sketch the argument. First, local well-posedness and nonnegativity follows from classical theory [37]. To extend existence and uniqueness to an arbitrary interval of time, we need to prove a priori estimates as in (1). To this end, we note that thanks to (2.2), the nonnegative map

$$t \mapsto \int_{\Omega} \left[ \Psi(u^{\varepsilon}(t,x)) + \Phi(v^{\varepsilon}(t,x)) \right] \mathrm{d}x$$

is nonincreasing whenever  $\phi' \geq 0$ . Choosing  $\phi$  vanishing on (0, M) and strictly increasing for  $(M, \infty)$  we obtain (1) and global well-posedness. Then, (2) and (3) follows from (2.2) with  $\phi(v) = v$ . Furthermore, (4) follows from the equality  $\partial_t u^{\varepsilon} + \partial_t v^{\varepsilon} = \Delta v^{\varepsilon}$  and property (2) while (5) follows from the chain rule for Sobolev functions, boundedness of  $v^{\varepsilon}$  from (1) and (2). Finally, to see (6) we choose  $k \geq d$  so that  $H^k(\Omega)$  embedds continuously into  $L^{\infty}(\Omega)$ . Let  $\varphi \in C(0,T;H^k(\Omega))$ . Note that there is a constant C such that

Thanks to (2.2) we have

$$\int_{(0,T)\times\Omega} (\partial_t \Psi(u^{\varepsilon}) + \partial_t \Phi(v^{\varepsilon})) \varphi \, dt \, dx - \int_{(0,T)\times\Omega} \nabla \Phi(v^{\varepsilon}) \cdot \nabla \varphi \, dt \, dx =$$

$$= -\int_{(0,T)\times\Omega} \phi'(v^{\varepsilon}) |\nabla v^{\varepsilon}|^2 \varphi \, dt \, dx - \int_{(0,T)\times\Omega} \frac{(v^{\varepsilon} - F(u^{\varepsilon})) (\phi(v^{\varepsilon}) - \phi(F(u^{\varepsilon})))}{\varepsilon} \varphi \, dt \, dx.$$

As  $|\phi'(v^{\varepsilon})| \leq C$  and  $|\phi(v^{\varepsilon}) - \phi(F(u^{\varepsilon}))| \leq C|v^{\varepsilon} - F(u^{\varepsilon})|$  we use bounds (2.3) together with points (2) and (3) to deduce for some possibly larger constant C (independent of  $\varepsilon$ )

$$\left| \int_{(0,T)\times\Omega} \left( \partial_t \Psi(u^{\varepsilon}) + \partial_t \Phi(v^{\varepsilon}) \right) \varphi \, \mathrm{d}t \, \mathrm{d}x \right| \le C \, \|\varphi\|_{C(0,T;H^k(\Omega))}.$$

Corollary 2.3. Let  $\{\mu_{t,x}\}_{t,x}$  and  $\{\nu_{t,x}\}_{t,x}$  be Young measures generated by sequences  $\{u^{\varepsilon}\}_{\varepsilon>0}$  and  $\{v^{\varepsilon}\}_{\varepsilon>0}$  respectively. Combining Lemma 2.2 (3) and Lemma A.5 (B, C) we obtain that  $F^{\#}\mu_{t,x} = \nu_{t,x}$ .

#### 3. Proof of Theorem 1 for fast-reaction system (1.1)

We begin with an entropy equality.

**Lemma 3.1** (entropy equality). Let  $\Psi$  and  $\Phi$  be defined with (2.1). Let  $g_i$  be densities given by (1.5). Then, for almost all  $\lambda_0$  (with respect to  $F^{\#}\mu_{t,x}$ ) we have

(3.1) 
$$\sum_{i=1}^{3} (\Psi(S_i(\lambda_0)) + \Phi(\lambda_0)) g_i(\lambda_0) = \sum_{i=1}^{3} \int_{\mathbb{R}^+} (\Psi(S_i(\lambda)) + \Phi(\lambda)) g_i(\lambda) dF^{\#} \mu_{t,x}(\lambda),$$

where  $S_i$  are inverses of F as in Notation 1.1.

Proof. Thanks to Lemma 2.2 (6), for all smooth  $\phi : \mathbb{R} \to \mathbb{R}$ ,  $\{\partial_t \Psi(u^{\varepsilon}) + \partial_t \Phi(v^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $(C(0,T;H^k(\Omega)))^*$ . Similarly, for all smooth  $\varphi : \mathbb{R} \to \mathbb{R}$ ,  $\{\nabla \varphi(v^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $L^2((0,\infty) \times \Omega)$ . Hence, Lemma A.1 implies

$$\underset{\varepsilon \to 0}{\operatorname{w}^*-\lim} \left(\Psi(u^{\varepsilon}) + \Phi(v^{\varepsilon})\right) \varphi(v^{\varepsilon}) = \underset{\varepsilon \to 0}{\operatorname{w}^*-\lim} \left(\Psi(u^{\varepsilon}) + \Phi(v^{\varepsilon})\right) \underset{\varepsilon \to 0}{\operatorname{w}^*-\lim} \varphi(v^{\varepsilon}).$$

As  $v^{\varepsilon} - F(u^{\varepsilon}) \to 0$  cf. Lemma 2.2 (3), we may replace  $v^{\varepsilon}$  with  $F(u^{\varepsilon})$  in the identity above to obtain

$$\underset{\varepsilon \to 0}{\operatorname{w}^*\text{-}\mathrm{lim}} \left( \Psi(u^\varepsilon) + \Phi(F(u^\varepsilon)) \right) \varphi(F(u^\varepsilon)) = \underset{\varepsilon \to 0}{\operatorname{w}^*\text{-}\mathrm{lim}} \left( \Psi(u^\varepsilon) + \Phi(F(u^\varepsilon)) \right) \ \underset{\varepsilon \to 0}{\operatorname{w}^*\text{-}\mathrm{lim}} \ \varphi(F(u^\varepsilon)).$$

In the language of Young measures, this identity reads

$$\int_{\mathbb{R}^+} (\Psi(\lambda) + \Phi(F(\lambda))) \varphi(F(\lambda)) d\mu_{t,x}(\lambda) = \int_{\mathbb{R}^+} (\Psi(\lambda) + \Phi(F(\lambda))) d\mu_{t,x}(\lambda) \int_{\mathbb{R}^+} \varphi(F(\lambda)) d\mu_{t,x}(\lambda).$$

We observe that  $\lambda = \sum_{i=1}^{3} S_i(F(\lambda)) \mathbb{1}_{\lambda \in I_i}$ . Hence, we may use push-forward measure to write

$$\sum_{i=1}^{3} \int_{\mathbb{R}^{+}} (\Psi(S_{i}(\lambda)) + \Phi(\lambda)) \varphi(\lambda) dF^{\#} \mu_{t,x}^{(i)}(\lambda) =$$

$$= \sum_{i=1}^{3} \int_{\mathbb{R}^{+}} (\Psi(S_{i}(\lambda)) + \Phi(\lambda)) dF^{\#} \mu_{t,x}^{(i)}(\lambda) \int_{\mathbb{R}^{+}} \varphi(\lambda) dF^{\#} \mu_{t,x}(\lambda).$$

Using (1.5) with densities  $g_1(\lambda)$ ,  $g_2(\lambda)$  and  $g_3(\lambda)$  we obtain

$$\sum_{i=1}^{3} \int_{\mathbb{R}^{+}} (\Psi(S_{i}(\lambda)) + \Phi(\lambda)) \varphi(\lambda) g_{i}(\lambda) dF^{\#} \mu_{t,x}(\lambda) =$$

$$= \sum_{i=1}^{3} \int_{\mathbb{R}^{+}} (\Psi(S_{i}(\lambda)) + \Phi(\lambda)) g_{i}(\lambda) dF^{\#} \mu_{t,x}(\lambda) \int_{\mathbb{R}^{+}} \varphi(\lambda) dF^{\#} \mu_{t,x}(\lambda).$$

Hence, when  $\lambda_0$  belongs to the support of the measure  $F^{\#}\mu_{t,x}$ , we obtain

$$\sum_{i=1}^{3} (\Psi(S_i(\lambda_0)) + \Phi(\lambda_0)) g_i(\lambda_0) = \sum_{i=1}^{3} \int_{\mathbb{R}^+} (\Psi(S_i(\lambda)) + \Phi(\lambda)) g_i(\lambda) dF^{\#} \mu_{t,x}(\lambda).$$

To analyze entropy inequality, we need to deal with integrals of the form  $\int_0^{S_i(\lambda)} \phi(F(\tau)) d\tau$ . This is the content of the next lemma.

# Lemma 3.2. We have

$$\Psi(S_i(\lambda_0)) = \int_0^{S_i(\lambda_0)} \phi(F(\tau)) d\tau = \int_0^{\lambda_0} \phi(\tau) S_i'(\tau) d\tau + C_i(\phi)$$

where  $C_1(\phi) = 0$  and  $C_2(\phi) = C_3(\phi) = \int_0^{f_+} \phi(\tau) \left( S_1'(\tau) - S_2'(\tau) \right) d\tau$ .

*Proof.* For i=1 we note that F is invertible on  $(0,S_1(\lambda))$  so that simple change of variables implies

$$\Psi(S_1(\lambda_0)) = \int_0^{S_1(\lambda_0)} \phi(F(\tau)) d\tau = \int_0^{\lambda_0} \phi(\tau) S_1'(\tau) d\tau.$$

For i = 2 we first split the integral for two intervals  $(0, \alpha_+)$ ,  $(\alpha_+, \lambda_0)$  cf. Notation 1.1. On each of them F is invertible so we can apply change of variables again:

$$\Psi(S_{2}(\lambda_{0})) = \int_{0}^{\alpha_{+}} \phi(F(\tau)) d\tau + \int_{\alpha_{+}}^{S_{2}(\lambda_{0})} \phi(F(\tau)) d\tau =$$

$$= \int_{0}^{f_{+}} \phi(\tau) S'_{1}(\tau) d\tau - \int_{\lambda_{0}}^{f_{+}} \phi(\tau) S'_{2}(\tau) d\tau = C_{2}(\phi) + \int_{0}^{\lambda_{0}} \phi(\tau) S'_{2}(\tau) d\tau.$$

For i=3 we split the integral for three intervals and apply change of variables again:

$$\Psi(S_3(\lambda_0)) = \int_0^{\alpha_+} \phi(F(\tau)) d\tau + \int_{\alpha_+}^{\beta_-} \phi(F(\tau)) d\tau + \int_{\beta_-}^{S_3(\lambda_0)} \phi(F(\tau)) d\tau =$$

$$= \int_0^{f_+} \phi(\tau) S_1'(\tau) d\tau - \int_{f_-}^{f_+} \phi(\tau) S_2'(\tau) d\tau + \int_{f_-}^{\lambda_0} \phi(\tau) S_3'(\tau) d\tau.$$

As  $S_2'(\tau) = 0$  and  $S_3'(\tau) = 0$  for  $\tau \in (0, f_-)$ , the proof is concluded.

### Lemma 3.3. Consider function

$$\mathcal{F}(\tau_0) = \sum_{i=1}^{3} (S_i'(\tau_0) + 1) F^{\#} \mu_{t,x}^{(i)}((\tau_0, \infty)) + (S_1'(\tau_0) - S_2'(\tau_0)) (1 - F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+)).$$

Then, for almost all  $\lambda_0$  (with respect to  $F^{\#}\mu_{t,x}$ ) and  $\tau_0 \neq f_-, f_+$  we have

$$\mathbb{1}_{\lambda_0 > \tau_0} \sum_{i=1}^{3} (S_i'(\tau_0) + 1) g_i(\lambda_0) + (S_1'(\tau_0) - S_2'(\tau_0)) (1 - g_1(\lambda_0)) = \mathcal{F}(\tau_0).$$

Proof. We consider  $\phi(\tau) = \phi^{\delta}(\tau) = \frac{1}{\delta} \mathbb{1}_{[\tau_0, \tau_0 + \delta]}$  and send  $\delta \to 0$  so that  $\Phi(\lambda_0) = \int_0^{\lambda_0} \phi^{\delta}(\tau) d\tau \to \mathbb{1}_{\lambda > \tau_0}$ . Moreover,  $\int_0^{\lambda_0} \phi^{\delta}(\tau) S_i'(\tau) d\tau \to S_i'(\tau_0) \mathbb{1}_{\lambda_0 > \tau_0}$ . Therefore, from Lemmas 3.1 and 3.2 we

deduce

$$\sum_{i=1}^{3} \left( \mathbb{1}_{\lambda_0 > \tau_0} \left( S_i'(\tau_0) + 1 \right) + \left( S_1'(\tau_0) - S_2'(\tau_0) \right) \mathbb{1}_{i=2,3} \right) g_i(\lambda_0) = 
= \sum_{i=1}^{3} \int_{\mathbb{R}^+} \left( \mathbb{1}_{\lambda > \tau_0} \left( S_i'(\tau_0) + 1 \right) + \left( S_1'(\tau_0) - S_2'(\tau_0) \right) \mathbb{1}_{i=2,3} \right) g_i(\lambda) \, \mathrm{d}F^{\#} \mu_{t,x}(\lambda).$$

Using identities from (1.6) and (1.7)

$$1 - g_1(\lambda_0) = g_2(\lambda_0) + g_3(\lambda_0), \qquad 1 - F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) = F^{\#} \mu_{t,x}^{(2)}(\mathbb{R}^+) + F^{\#} \mu_{t,x}^{(3)}(\mathbb{R}^+)$$

we conclude the proof.

Proof of Theorem 1. The first part of Theorem 1 is proved in Lemma 3.3. To see the second one, fix  $\lambda_0 \neq f_-, f_+$ . For  $\tau_0 := \eta > \lambda_0$  we obtain

$$\sum_{i=1}^{3} (S_i'(\eta) + 1) F^{\#} \mu_{t,x}^{(i)}((\eta, \infty)) + (S_1'(\eta) - S_2'(\eta)) (F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_0)) = 0$$

while for  $\tau_0 := \xi < \lambda_0$  we deduce

$$\sum_{i=1}^{3} \left(S_i'(\xi) + 1\right) \left(g_i(\lambda_0) - F^{\#} \mu_{t,x}^{(i)}((\xi, \infty))\right) + \left(S_1'(\xi) - S_2'(\xi)\right) \left(F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_0)\right) = 0.$$

Sending  $\xi, \eta \to \lambda_0$  and using continuity of  $\lambda \mapsto S'_i(\lambda)$  at  $\lambda \neq f_-, f_+$  we obtain

$$\sum_{i=1}^{3} (S_i'(\lambda_0) + 1) g_i(\lambda_0) = \sum_{i=1}^{3} (S_i'(\lambda_0) + 1) F^{\#} \mu_{t,x}^{(i)} \{\lambda_0\}.$$

Finally, we note that for almost all  $\lambda_0$  (with respect to  $F^{\#}\mu_{t,x}$ )  $F^{\#}\mu_{t,x}^{(i)}\{\lambda_0\} = g_i(\lambda_0) F^{\#}\mu_{t,x}\{\lambda_0\}$  and this concludes the proof.

4. Proofs of Theorems 2, 3 and 4 for fast-reaction system (1.1)

Proof of Theorem 2. If supp  $F^{\#}\mu_{t,x} \cap (0, f_{-})$  is nonempty, we take any  $\lambda_0 \in \text{supp } F^{\#}\mu_{t,x} \cap (0, f_{-})$ . Note that  $S'_2(\lambda_0) = S'_3(\lambda_0) = 0$  and (1.8) in Theorem 1 implies

$$(1 - F^{\#}\mu_{t,x} \{\lambda_0\}) (S_1'(\lambda_0) + 1) g_1(\lambda_0) = 0.$$

For almost all  $\lambda_0 \in (0, f_-)$  we have  $g_1(\lambda_0) = 1$  so we conclude  $F^{\#}\mu_{t,x}\{\lambda_0\} = 1$ . Similar argument works in the case  $\lambda_0 \in (f_+, \infty)$ .

Now, let  $\lambda_0 \in [f_-, f_+] \cap \operatorname{supp} F^{\#} \mu_{t,x}$ . If  $\operatorname{supp} F^{\#} \mu_{t,x} = \{\lambda_0\}$ , we conclude  $F^{\#} \mu_{t,x} = \delta_{\lambda_0}$ . Otherwise, there are  $\lambda_1, \lambda_2 \in \operatorname{supp} F^{\#} \mu_{t,x}$  such that  $f_- \leq \lambda_1 < \lambda_2 \leq f_+$ . For any  $\tau_0 \in (\lambda_1, \lambda_2)$  we use Theorem 1 with  $\lambda_0 = \lambda_1, \lambda_2$  to obtain two equations:

$$\sum_{i=1}^{3} (S_i'(\tau_0) + 1) \left[ g_i(\lambda_2) - F^{\#} \mu_{t,x}^{(i)}(\tau_0, \infty) \right] + \left( S_1'(\tau_0) - S_2'(\tau_0) \right) \left( F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_2) \right) = 0,$$

$$-\sum_{i=1}^{3} (S_i'(\tau_0) + 1) F^{\#} \mu_{t,x}^{(i)}(\tau_0, \infty) + (S_1'(\tau_0) - S_2'(\tau_0)) (F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_1)) = 0.$$

Hence,  $\sum_{i=1}^{3} (S_i'(\tau_0) + 1) g_i(\lambda_2) + (S_1'(\tau_0) - S_2'(\tau_0)) (g_1(\lambda_1) - g_1(\lambda_2)) = 0$ . But then, nondegeneracy condition (1.3) implies that the sum  $\sum_{i=1}^{3} g_i(\lambda_2) = 0 \neq 1$  raising contradiction.

It follows that  $F^{\#}\mu_{t,x}$  is a Dirac mass. From Corollary 2.3 we deduce that the Young measure  $\{\nu_{t,x}\}_{t,x}$  generated by  $\{v^{\varepsilon}\}_{\varepsilon}$  is also a Dirac mass so  $v^{\varepsilon} \to v$  strongly and  $\nu_{t,x} = \delta_{v(t,x)}$ , cf. Lemma A.5. The representation formula for  $\mu_{t,x}$  follows from  $F^{\#}\mu_{t,x} = \delta_{v(t,x)}$ .

Before proceeding to the proofs of Theorems 3 and 4, we will state a simple lemma concerning the case when  $F^{\#}\mu_{t,x}$  is supported only at  $f_{-}$  and  $f_{+}$ . This needs some care as functions  $S'_{1}$ ,  $S'_{2}$  and  $S'_{3}$  are not continuous at these points.

**Lemma 4.1** (Accumulation at the interface). Suppose that there exists  $\tau_0 \in (f_-, f_+)$  such that  $S_1'(\tau_0) - S_3'(\tau_0) \neq 0$ . Assume that supp  $F^{\#}\mu_{t,x} \subset \{f_-, f_+\}$ . Then,  $F^{\#}\mu_{t,x} = \delta_{f_-}$  or  $F^{\#}\mu_{t,x} = \delta_{f_+}$ .

*Proof.* Aiming at contradiction, we assume that  $F^{\#}\mu_{t,x}\{f_+\}>0$  and  $F^{\#}\mu_{t,x}\{f_-\}>0$ . Note that

$$0 = \mu_{t,x}^{(2)}(F^{-1}(f_+) \cap I_2) = F^{\#}\mu_{t,x}^{(2)}\{f_+\} = g_2(f_+)F^{\#}\mu_{t,x}\{f_+\}$$

so that  $g_2(f_+) = 0$  and similarly  $g_2(f_-) = 0$ . Applying Theorem 1 with  $\tau_0 \in (f_-, f_+)$  and  $\lambda_0 \in \{f_-, f_+\}$  we obtain

$$\sum_{i=1}^{3} (S_i'(\tau_0) + 1) \left[ \mathbb{1}_{\lambda_0 > \tau_0} g_i(\lambda_0) - F^{\#} \mu_{t,x}^{(i)}(\tau_0, \infty) \right] + \left( S_1'(\tau_0) - S_2'(\tau_0) \right) \left( F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_0) \right) = 0.$$

As  $\tau_0 \in (f_-, f_+)$ , we have

$$F^{\#}\mu_{t,x}^{(i)}(\tau_0,\infty) = F^{\#}\mu_{t,x}^{(i)}\{f_+\} = g_i(f_+) F^{\#}\mu_{t,x}\{f_+\}.$$

But this implies

$$\left(\mathbb{1}_{\lambda_0 > \tau_0} - F^{\#} \mu_{t,x} \{f_+\}\right) \sum_{i=1,3} \left(S_i'(\tau_0) + 1\right) g_i(\lambda_0) + \left(S_1'(\tau_0) - S_2'(\tau_0)\right) \left(F^{\#} \mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(\lambda_0)\right) = 0.$$

Considering  $\lambda_0 = f_+$ ,  $f_-$  and using  $1 - F^{\#}\mu_{t,x}\{f_+\} = F^{\#}\mu_{t,x}\{f_-\}$  we obtain two equations:

$$(4.1) F^{\#}\mu_{t,x}\{f_{-}\} \sum_{i=1,3} (S'_{i}(\tau_{0}) + 1) g_{i}(f_{+}) + (S'_{1}(\tau_{0}) - S'_{2}(\tau_{0})) (F^{\#}\mu_{t,x}^{(1)}(\mathbb{R}^{+}) - g_{1}(f_{+})) = 0,$$

$$(4.2) -F^{\#}\mu_{t,x}\{f_{+}\} \sum_{i=1,3} (S'_{i}(\tau_{0})+1) g_{i}(f_{-}) + (S'_{1}(\tau_{0})-S'_{2}(\tau_{0})) (F^{\#}\mu_{t,x}^{(1)}(\mathbb{R}^{+})-g_{1}(f_{-})) = 0.$$

Using  $1 - F^{\#}\mu_{t,x}\{f_+\} = F^{\#}\mu_{t,x}\{f_-\}$  once again we obtain that

$$F^{\#}\mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(f_+) = g_1(f_+) F^{\#}\mu_{t,x}\{f_+\} + g_1(f_-) F^{\#}\mu_{t,x}\{f_-\} - g_1(f_+) =$$

$$= (g_1(f_-) - g_1(f_+)) F^{\#}\mu_{t,x}\{f_-\}$$

$$= (g_1(f_-) - g_1(f_+)) F^{\#}\mu_{t,x}\{f_-\}$$

and similarly for  $F^{\#}\mu_{t,x}^{(1)}(\mathbb{R}^+) - g_1(f_-)$ . As we assume that  $F^{\#}\mu_{t,x}\{f_-\}, F^{\#}\mu_{t,x}\{f_+\} > 0$  we may simplify (4.1)–(4.2) to obtain

(4.3) 
$$\sum_{i=1,3} \left( S_i'(\tau_0) + 1 \right) g_i(f_+) + \left( S_1'(\tau_0) - S_2'(\tau_0) \right) \left( g_1(f_-) - g_1(f_+) \right) = 0,$$

$$(4.4) -\sum_{i=1,3} \left(S_i'(\tau_0) + 1\right) g_i(f_-) + \left(S_1'(\tau_0) - S_2'(\tau_0)\right) \left(g_1(f_+) - g_1(f_-)\right) = 0.$$

We observe further that  $g_1(\lambda_0) + g_3(\lambda_0) = 1$ , cf. (1.7), so that

$$\sum_{i=1,3} (S_i'(\tau_0) + 1) g_i(\lambda_0) = (S_1'(\tau_0) - S_3'(\tau_0)) g_1(\lambda_0) + (S_3'(\tau_0) + 1).$$

Hence, we may further simplify (4.3)–(4.4) to get

$$(4.5) (S_1'(\tau_0) - S_3'(\tau_0)) g_1(f_+) + (S_3'(\tau_0) + 1) + (S_1'(\tau_0) - S_2'(\tau_0)) (g_1(f_-) - g_1(f_+)) = 0,$$

$$(4.6) -(S_1'(\tau_0) - S_3'(\tau_0))g_1(f_-) - (S_3'(\tau_0) + 1) + (S_1'(\tau_0) - S_2'(\tau_0))(g_1(f_+) - g_1(f_-)) = 0.$$

By assumption, there is  $\tau_0 \in (f_-, f_+)$  such that  $S'_1(\tau_0) - S'_3(\tau_0) \neq 0$ . Using (4.5)–(4.6) for such  $\tau_0$  we see that  $g_1(f_+) = g_1(f_-)$ . But then, coming back to (4.5)–(4.6), we deduce that

$$\sum_{i=1,3} (S_i'(\tau_0) + 1) g_i(f_-) = 0, \qquad \sum_{i=1,3} (S_i'(\tau_0) + 1) g_i(f_-) = 0.$$

As  $S_1$ ,  $S_3$  are increasing, this implies  $g_1(f_-) = g_3(f_-) = g_1(f_+) = g_3(f_+) = 0$  raising contradiction with  $g_1(f_-) + g_3(f_-) = 1$  and  $g_1(f_+) + g_3(f_+) = 1$ .

**Remark 4.2.** Without assumption that there is  $\tau_0 \in (f_-, f_+)$  such that  $S'_1(\tau_0) - S'_3(\tau_0) \neq 0$  we observe that (4.5)–(4.6) degenerate to the same equation:

$$g_1(f_+) - g_1(f_-) = \frac{1 + S_3'(\tau_0)}{S_1'(\tau_0) - S_2'(\tau_0)}$$

valid for all  $\tau_0 \in (f_-, f_+)$ . Hence, it the function  $\tau_0 \mapsto \frac{1+S_3'(\tau_0)}{S_1'(\tau_0)-S_2'(\tau_0)}$  is not constant, we may also obtain contradiction. But we believe that assumption on  $S_1'(\tau_0) - S_3'(\tau_0)$  is easier to formulate. It also allows for piecewise affine nonlinearities.

Proof of Theorem 3. As in the proof of Theorem 2 we may assume that supp  $F^{\#}\mu_{t,x} \subset [f_-, f_+]$  (this did not use nondegeneracy condition!). By assumption, for any set  $A \subset \mathbb{R}^+$ 

$$0 = \mu_{t,x}(F^{-1}(A) \cap I_2) = F^{\#}\mu_{t,x}^{(2)}(A) = \int_A g_2(\lambda) \, dF^{\#}\mu_{t,x}(\lambda)$$

so  $g_2(\lambda) = 0$  for almost all  $\lambda$ . Hence, when  $\lambda_0 \in \operatorname{supp} F^{\#}\mu_{t,x} \cap (f_-, f_+)$ , the sum

$$\sum_{i=1}^{3} (S_i'(\lambda_0) + 1) g_i(\lambda_0) \ge \min(S_1'(\lambda_0) + 1, S_3'(\lambda_0) + 1) > 0$$

because  $g_1(\lambda_0) + g_3(\lambda_0) = 1$  and  $S_1$ ,  $S_3$  are strictly increasing. It follows from Theorem 1 that  $F^{\#}\mu_{t,x}\{\lambda_0\} = 1$ , i.e.  $F^{\#}\mu_{t,x} = \delta_{\lambda_0}$ . Finally, if there is no such  $\lambda_0 \in \text{supp } F^{\#}\mu_{t,x} \cap (f_-, f_+)$ , we apply Lemma 4.1.

It follows that  $F^{\#}\mu_{t,x}$  is a Dirac mass and now, we can conclude as in Theorem 2.

Proof of Theorem 4. Mimicking the proof of Theorem 3, we let  $\lambda_0 \in \operatorname{supp} F^{\#} \mu_{t,x} \cap (f_-, f_+)$  and we observe that the sum

$$\sum_{i=1}^{3} (S_i'(\lambda_0) + 1) g_i(\lambda_0) \ge \min(1, \delta(\lambda_0)) \sum_{i=1}^{3} g_i(\lambda_0) = \min(1, \delta(\lambda_0)) > 0$$

where  $\delta(\lambda_0)$  is such that  $S_2'(\lambda_0) + 1 > \delta(\lambda_0) > 0$ . We conclude as in the proof of Theorem 3.

5. Proof of Theorems 1-3 to the forward-backward diffusion system (1.2)

We first formulate basic well-posedness result for (1.2). This comes mostly from [31,35] but the compactness estimates are simplified.

**Lemma 5.1.** Let  $u_0 \in L^{\infty}(\Omega)$ . Then, there exists the unique solution  $u^{\varepsilon} : [0, \infty) \times \Omega \to \mathbb{R}$  of (1.2) which is nonnegative and has regularity  $C^1([0,T];L^2(\Omega)) \cap L^{\infty}(\Omega)$ . Moreover, we have

(1) for 
$$M = \max(\|F(u_0)\|_{\infty}, f_+)$$
 we have  $0 < u^{\varepsilon} < M$ .

- (2)  $\{\nabla v^{\varepsilon}\}_{\varepsilon\in(0,1)}$  is uniformly bounded in  $L^{2}((0,T)\times\Omega)$ ,
- $(3) \ \left\{ \frac{v^{\varepsilon} F(u^{\varepsilon})}{\sqrt{\varepsilon}} \right\}_{\varepsilon \in (0,1)} = \left\{ \sqrt{\varepsilon} \ u_t^{\varepsilon} \ \right\}_{\varepsilon \in (0,1)} \ are \ uniformly \ bounded \ in \ L^2((0,T) \times \Omega),$
- (4) for all smooth  $\varphi: \mathbb{R} \to \mathbb{R}$ ,  $\{\nabla \varphi(v^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $L^{2}((0,T) \times \Omega)$ .
- (5) for all smooth  $\phi : \mathbb{R} \to \mathbb{R}$ ,  $\{\partial_t \Psi(u^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $(C(0,T;H^k(\Omega)))^*$  for sufficiently large  $k \in \mathbb{N}$ .

*Proof.* We observe that equation is equivalent to the following ODE:

$$\partial_t u^{\varepsilon} = (I - \varepsilon \Delta)^{-1} \Delta F(u^{\varepsilon}).$$

As long as  $\varepsilon > 0$ , the (RHS) is Lipschitz continuous, say on  $L^2(\Omega)$ , so the local well-posedness follows. To obtain global well-posedness, we consider functions  $\Psi$ ,  $\Phi$  defined in (2.1). We have

$$(5.1) \quad \partial_t \Psi(u^{\varepsilon}) = \phi(F(u^{\varepsilon})) \, u_t^{\varepsilon} = (\phi(F(u^{\varepsilon})) - \phi(v^{\varepsilon})) \, u_t^{\varepsilon} + \phi(v^{\varepsilon}) \, \Delta v^{\varepsilon} = \\ = (\phi(F(u^{\varepsilon})) - \phi(v^{\varepsilon})) \, u_t^{\varepsilon} + \Delta \Phi(v^{\varepsilon}) - \phi'(v^{\varepsilon}) \, |\nabla v^{\varepsilon}|^2.$$

If  $\phi$  is nondecreasing, we have

$$\left(\phi(F(u^{\varepsilon})) - \phi(v^{\varepsilon})\right)u_t^{\varepsilon} = \left(\phi(F(u^{\varepsilon})) - \phi(v^{\varepsilon})\right)\frac{v^{\varepsilon} - F(u^{\varepsilon})}{\varepsilon} \le 0$$

so after integration in space, the (RHS) of (5.1) is nonnegative. Hence,  $\partial_t \int_{\Omega} \Psi(u^{\varepsilon}) \leq 0$ . Choosing  $\phi = 0$  for [0, M] and  $\phi'(x) > 0$  for  $x \notin [0, M]$  we prove (1) and conclude the proof of global well-posedness. To see (2) and (3) we take  $\phi(x) = x$  and integrate (5.1) in time and space. Part (4) easily follows from chain rule and (2). Finally, (5) follows from (5.1) and exactly the same computations as in Lemma 2.2.

Now, we formulate an analog of Lemma 3.1.

**Lemma 5.2** (entropy equality). Let  $\Psi$  be defined with (2.1). Let  $g_i$  be densities given by (1.5). Then, for almost all  $\lambda_0$  (with respect to  $F^{\#}\mu_{t,x}$ ) we have

(5.2) 
$$\sum_{i=1}^{3} \Psi(S_i(\lambda_0)) g_i(\lambda_0) = \sum_{i=1}^{3} \int_{\mathbb{R}^+} \Psi(S_i(\lambda)) g_i(\lambda) dF^{\#} \mu_{t,x}(\lambda).$$

*Proof.* Thanks to Lemma 5.1 (5), for all smooth  $\phi: \mathbb{R} \to \mathbb{R}$ ,  $\{\partial_t \Psi(u^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $(C(0,T;H^k(\Omega)))^*$ . Similarly, for all smooth and bounded  $\varphi: \mathbb{R} \to \mathbb{R}$ ,  $\{\nabla \varphi(v^{\varepsilon})\}_{\varepsilon \in (0,1)}$  is uniformly bounded in  $L^2((0,\infty) \times \Omega)$ . Hence, Lemma A.1 implies

$$\mathbf{w}^*_{\varepsilon \to 0} \cdot \Psi(u^{\varepsilon}) \varphi(v^{\varepsilon}) = \mathbf{w}^*_{\varepsilon \to 0} \cdot \Psi(u^{\varepsilon}) \cdot \mathbf{w}^*_{\varepsilon \to 0} \cdot \varphi(v^{\varepsilon}).$$

As  $v^{\varepsilon} - F(u^{\varepsilon}) = \varepsilon u_t^{\varepsilon} \to 0$  cf. Lemma 5.1 (3), we may replace  $v^{\varepsilon}$  with  $F(u^{\varepsilon})$  in the identity above to obtain

$$\mathbf{w}^*_{\varepsilon \to 0} \cdot \Psi(u^{\varepsilon}) \, \varphi(F(u^{\varepsilon}))) = \mathbf{w}^*_{\varepsilon \to 0} \cdot \Psi(u^{\varepsilon}) \, \mathbf{w}^*_{\varepsilon \to 0} \cdot \varphi(F(u^{\varepsilon}))).$$

In the language of Young measures, this identity reads

$$\int_{\mathbb{R}^+} \Psi(\lambda) \, \varphi(F(\lambda)) \, \mathrm{d}\mu_{t,x}(\lambda) = \int_{\mathbb{R}^+} \Psi(\lambda) \, \mathrm{d}\mu_{t,x}(\lambda) \, \int_{\mathbb{R}^+} \varphi(F(\lambda)) \, \mathrm{d}\mu_{t,x}(\lambda).$$

We observe that  $\lambda = \sum_{i=1}^{3} S_i(F(\lambda)) \mathbb{1}_{\lambda \in I_i}$ . Hence, we may use push-forward measure to write

$$\sum_{i=1}^{3} \int_{\mathbb{R}^{+}} \Psi(S_{i}(\lambda)) \varphi(\lambda) \, \mathrm{d}F^{\#} \mu_{t,x}^{(i)}(\lambda) = \sum_{i=1}^{3} \int_{\mathbb{R}^{+}} \Psi(S_{i}(\lambda)) \, \mathrm{d}F^{\#} \mu_{t,x}^{(i)}(\lambda) \int_{\mathbb{R}^{+}} \varphi(\lambda) \, \mathrm{d}F^{\#} \mu_{t,x}(\lambda).$$

Using densities  $g_1(\lambda)$ ,  $g_2(\lambda)$  and  $g_3(\lambda)$  we obtain

$$\sum_{i=1}^{3} \int_{\mathbb{R}^{+}} \Psi(S_{i}(\lambda)) \varphi(\lambda) g_{i}(\lambda) dF^{\#} \mu_{t,x}(\lambda) =$$

$$= \sum_{i=1}^{3} \int_{\mathbb{R}^{+}} \Psi(S_{i}(\lambda)) g_{i}(\lambda) dF^{\#} \mu_{t,x}(\lambda) \int_{\mathbb{R}^{+}} \varphi(\lambda) dF^{\#} \mu_{t,x}(\lambda).$$

Hence, if  $\lambda_0$  belongs to the support of the measure  $F^{\#}\mu_{t,x}$ , we obtain (5.2).

Proof of Theorems 1–3. Comparing formulations of Lemma 3.1 and 5.2 we see that it is sufficient to modify proofs in Sections 3-4 by replacing  $S'_1+1$ ,  $S'_2+1$  and  $S'_3+1$  with  $S'_1$ ,  $S'_2$  and  $S'_3$  respectively.  $\square$ 

Note that Theorem 4 is only true for fast-reaction limit (1.1) because its proof exploits presence of  $S'_2 + 1$  in the entropy formulations.

# APPENDIX A. USEFUL NOTIONS AND RESULTS

A.1. Compensated compactness lemma. We formulate lemma used in the proof of Theorem 1, more precisely in Lemma 3.1. For the proof see [27, Proposition 1].

**Lemma A.1.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded domain. Suppose that  $\{a_n\}_{n\in\mathbb{N}}$  is uniformly bounded in  $L^2(0,T;H^1(\Omega))$  and  $\{b_n\}_{n\in\mathbb{N}}$  is uniformly bounded in  $L^2(0,T;L^2(\Omega))$ . Moreover, assume that the sequence of distributional time derivatives  $\{\partial_t b_n\}_{n\in\mathbb{N}}$  is uniformly bounded in the dual space  $C(0,T;H^m(\Omega))^*$  for some  $m\in\mathbb{N}$ . Then, if  $a_n\rightharpoonup a$  and  $b_n\rightharpoonup b$  we have  $a_n\,b_n\to a\,b$  in the sense of distributions.

In our case, the considered sequences are also in  $L^{\infty}((0,T)\times\Omega)$  so the resulting convergence is true in the weak\* sense.

A.2. Support of the measure. We recall definition of the support of measure on  $\mathbb{R}^n$  [38, Definition 1.14]. For this, let B(x,r) denote a ball of radius r > 0 centered at  $x \in \mathbb{R}^n$ .

**Definition A.2.** Let  $\mu$  be a nonnegative measure on  $\mathbb{R}^n$ . We say that  $x \in \text{supp } \mu$  if and only if  $\mu(B(x,r)) > 0$  for all r > 0.

Remark A.3. When given property (like equation) is satisfied for almost every x (with respect to  $\mu$ ) one may worry that it is not true for the particularly chosen value of x. This is not the problem if one takes  $x \in \text{supp } \mu$  because in each neighbourhood of x there is  $y \in \text{supp } \mu$  such that the property has to be satisfied because the measure of each neighbourhood is nonzero.

A.3. Young measures. Finally, we recall the theory of Young measures introduced by Young [44,45] and recalled in the seminal paper of Ball [2]. Reader interested in modern presentation may consult [18], [32, Chapter 6] or [36, Chapter 4]. For simplicity, we formulate it for sequences of functions  $\{u_n\}_{n\in\mathbb{N}}$  uniformly bounded in  $L^p(\Omega)$  with some  $1 \leq p \leq \infty$  and  $\Omega \subset \mathbb{R}^n$  being a bounded domain. We start with the most important result that we cite from [32, Theorem 6.2]:

**Theorem A.4** (Fundamental Theorem of Young Measures). Let  $\Omega \subset \mathbb{R}^n$  be a a bounded domain and let  $\{u_n\}_{n\in\mathbb{N}}$  be a sequence bounded in  $L^p(\Omega)$  with  $1 \leq p \leq \infty$ . Then, there exists a subsequence (not relabeled) and a weakly-\* measurable family of probability measures  $\{\mu_x\}_{x\in\Omega}$  such that for all bounded and smooth  $G: \mathbb{R} \to \mathbb{R}$ , we have

(A.1) 
$$G(u_n(x)) \stackrel{*}{\rightharpoonup} \int_{\mathbb{R}} G(\lambda) d\mu_x(\lambda) \quad in L^{\infty}(\Omega).$$

We say that the sequence  $\{u_n\}_{n\in\mathbb{N}}$  generates the family of Young measures  $\{\mu_x\}_{x\in\Omega}$ .

Now we list properties of Young measures used in the paper.

**Lemma A.5.** Under notation of Theorem A.4, the following hold true.

- (A) We have  $u_n \to u$  a.e. (up to a subsequence) if and only if  $\mu_{t,x} = \delta_{u(t,x)}$ .
- (B) If  $\{w_n\}_{n\in\mathbb{N}}$  is another bounded sequence such that  $u_n-w_n\to 0$  a.e. then Young measures generated by  $\{u_n\}_{n\in\mathbb{N}}$  and  $\{w_n\}_{n\in\mathbb{N}}$  coincide.
- (C) If  $F : \mathbb{R} \to \mathbb{R}$ , the sequence  $\{F(u_n)\}_{n \in \mathbb{N}}$  generates Young measure  $F^{\#}\mu_{t,x}$  (i.e. push-forward  $\mu_{t,x} \circ F^{-1}$ ).

Sketch of the proof. For (A) we consider G(u) = u and  $G(u) = u^2$  to deduce that  $u_n \to u$  in  $L^2(\Omega)$  so that up to a subsequence also a.e. The opposite direction is clear because  $G(u_n(x)) \to G(u(x))$ 

a.e. For (B) we note that for all bounded and smooth G, weak limits of  $G(u_n(x))$  and  $G(w_n(x))$  coincide. For (C) we write

$$G(F(u_n)) = \int_{\mathbb{R}} G(F(\lambda)) d\mu_{t,x}(\lambda) = \int_{\mathbb{R}} G(\lambda) d(\mu_{t,x} \circ F^{-1})(\lambda).$$

#### References

- G. Andrews and J. M. Ball. Asymptotic behaviour and changes of phase in one-dimensional nonlinear viscoelasticity. J. Differential Equations, 44(2):306-341, 1982. Special issue dedicated to J. P. LaSalle, https://doi.org/10.1016/0022-0396(82)90019-5.
- [2] J. M. Ball. A version of the fundamental theorem for Young measures. In PDEs and continuum models of phase transitions (Nice, 1988), volume 344 of Lecture Notes in Phys., pages 207-215. Springer, Berlin, 1989. URL: https://doi.org/10.1007/BFb0024945.
- [3] M. Bertsch, F. Smarrazzo, and A. Tesei. Pseudoparabolic regularization of forward-backward parabolic equations: a logarithmic nonlinearity. *Anal. PDE*, 6(7):1719–1754, 2013. URL: https://doi.org/10.2140/apde.2013.6.1719.
- [4] M. Bertsch, F. Smarrazzo, and A. Tesei. Pseudo-parabolic regularization of forward-backward parabolic equations: power-type nonlinearities. J. Reine Angew. Math., 712:51-80, 2016. URL: https://doi.org/10.1515/crelle-2013-0123.
- [5] D. Bothe and D. Hilhorst. A reaction-diffusion system with fast reversible reaction. J. Math. Anal. Appl., 286(1):125-135, 2003. URL: https://doi.org/10.1016/S0022-247X(03)00457-8.
- [6] D. Bothe, M. Pierre, and G. Rolland. Cross-diffusion limit for a reaction-diffusion system with fast reversible reaction. Comm. Partial Differential Equations, 37(11):1940–1966, 2012. URL: https://doi.org/10.1080/03605302.2012.715706.
- [7] E. Bottazzi. Grid functions of nonstandard analysis in the theory of distributions and in partial differential equations. Adv. Math., 345:429-482, 2019. URL: https://doi.org/10.1016/j.aim.2019.01.024.
- [8] E. Bottazzi. A grid function formulation of a class of ill-posed parabolic equations. *J. Differential Equations*, 271:39-75, 2021. URL: https://doi.org/10.1016/j.jde.2020.08.002.
- [9] F. Bubba, B. Perthame, C. Pouchol, and M. Schmidtchen. Hele-Shaw Limit for a System of Two Reaction-(Cross-)Diffusion Equations for Living Tissues. Arch. Ration. Mech. Anal., 236(2):735-766, 2020. URL: https://doi.org/10.1007/s00205-019-01479-1.
- [10] J. A. Carrillo, F. Filbet, and M. Schmidtchen. Convergence of a finite volume scheme for a system of interacting species with cross-diffusion. *Numer. Math.*, 145(3):473–511, 2020. URL: https://doi.org/10.1007/s00211-020-01121-3.
- [11] E. C. M. Crooks and D. Hilhorst. Self-similar fast-reaction limits for reaction-diffusion systems on unbounded domains. J. Differential Equations, 261(3):2210-2250, 2016. URL: https://doi.org/10.1016/j.jde.2016.04.028.

- [12] S. Cygan, A. Marciniak-Czochra, G. Karch, and K. Suzuki. Instability of all regular stationary solutions to reaction-diffusion-ode systems. arXiv preprint arXiv:2105.05023, 2021. URL: https://arxiv.org/abs/2105.05023.
- [13] S. Cygan, A. Marciniak-Czochra, G. Karch, and K. Suzuki. Stable discontinuous stationary solutions to reactiondiffusion-ode systems. In preparation, 2021.
- [14] E. S. Daus, L. Desvillettes, and A. Jüngel. Cross-diffusion systems and fast-reaction limits. Bull. Sci. Math., 159:102824, 29, 2020. URL: https://doi.org/10.1016/j.bulsci.2019.102824.
- [15] L. Desvillettes, T. Lepoutre, and A. Moussa. Entropy, duality, and cross diffusion. SIAM J. Math. Anal., 46(1):820–853, 2014. URL: https://doi.org/10.1137/130908701.
- [16] L. Desvillettes, T. Lepoutre, A. Moussa, and A. Trescases. On the entropic structure of reaction-cross diffusion systems. Comm. Partial Differential Equations, 40(9):1705–1747, 2015. URL: https://doi.org/10.1080/03605302.2014.998837.
- [17] L. C. Evans. A convergence theorem for a chemical diffusion-reaction system. Houston J. Math., 6(2):259–267, 1980.
- [18] L. C. Evans. Weak convergence methods for nonlinear partial differential equations, volume 74 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1990. URL: https://doi.org/10.1090/cbms/074.
- [19] L. C. Evans and M. Portilheiro. Irreversibility and hysteresis for a forward-backward diffusion equation. Math. Models Methods Appl. Sci., 14(11):1599–1620, 2004. URL: https://doi.org/10.1142/S0218202504003763.
- [20] D. Hilhorst, M. Mimura, and H. Ninomiya. Fast reaction limit of competition-diffusion systems. In *Handbook of differential equations: evolutionary equations. Vol. V*, Handb. Differ. Equ., pages 105–168. Elsevier/North-Holland, Amsterdam, 2009. URL: https://doi.org/10.1016/S1874-5717(08)00209-0.
- [21] M. Iida, H. Monobe, H. Murakawa, and H. Ninomiya. Vanishing, moving and immovable interfaces in fast reaction limits. J. Differential Equations, 263(5):2715-2735, 2017. URL: https://doi.org/10.1016/j.jde.2017.04.009.
- [22] A. Jüngel. The boundedness-by-entropy method for cross-diffusion systems. *Nonlinearity*, 28(6):1963–2001, 2015.
   URL: https://doi.org/10.1088/0951-7715/28/6/1963.
- [23] P. Lafitte and C. Mascia. Numerical exploration of a forward-backward diffusion equation. *Math. Models Methods Appl. Sci.*, 22(6):1250004, 33, 2012. URL: https://doi.org/10.1142/S0218202512500042.
- [24] Y. Lou and W.-M. Ni. Diffusion, self-diffusion and cross-diffusion. J. Differential Equations, 131(1):79–131, 1996.
  URL: https://doi.org/10.1006/jdeq.1996.0157.
- [25] A. Marciniak-Czochra, G. Karch, and K. Suzuki. Unstable patterns in reaction-diffusion model of early carcinogenesis. J. Math. Pures Appl. (9), 99(5):509-543, 2013. URL: https://doi.org/10.1016/j.matpur.2012.09.011.
- [26] C. Mascia, A. Terracina, and A. Tesei. Two-phase entropy solutions of a forward-backward parabolic equation. Arch. Ration. Mech. Anal., 194(3):887–925, 2009. URL: https://doi.org/10.1007/s00205-008-0185-6.
- [27] A. Moussa. Some variants of the classical Aubin-Lions lemma. J. Evol. Equ., 16(1):65-93, 2016. URL: https://doi.org/10.1007/s00028-015-0293-3.
- [28] A. Moussa, B. Perthame, and D. Salort. Backward parabolicity, cross-diffusion and Turing instability. *J. Nonlinear Sci.*, 29(1):139–162, 2019. URL: https://doi.org/10.1007/s00332-018-9480-z.

- [29] H. Murakawa and H. Ninomiya. Fast reaction limit of a three-component reaction-diffusion system. J. Math. Anal. Appl., 379(1):150-170, 2011. URL: https://doi.org/10.1016/j.jmaa.2010.12.040.
- [30] F. Murat. A survey on compensated compactness. In Contributions to modern calculus of variations (Bologna, 1985), volume 148 of Pitman Res. Notes Math. Ser., pages 145–183. Longman Sci. Tech., Harlow, 1987.
- [31] A. Novick-Cohen and R. L. Pego. Stable patterns in a viscous diffusion equation. Trans. Amer. Math. Soc., 324(1):331–351, 1991. URL: https://doi.org/10.2307/2001511.
- [32] P. Pedregal. Parametrized measures and variational principles, volume 30 of Progress in Non-linear Differential Equations and their Applications. Birkhäuser Verlag, Basel, 1997. URL: https://doi.org/10.1007/978-3-0348-8886-8.
- [33] B. Perthame and J. Skrzeczkowski. Fast reaction limit with nonmonotone reaction function. To appear in Communications on Pure and Applied Mathematics. arXiv preprint arXiv:2008.11086, 2020. URL: https://arxiv.org/abs/2008.11086.
- [34] P. I. Plotnikov. Equations with a variable direction of parabolicity and the hysteresis effect. *Dokl. Akad. Nauk*, 330(6):691–693, 1993.
- [35] P. I. Plotnikov. Passage to the limit with respect to viscosity in an equation with a variable direction of parabolicity. *Differentsial' nye Uravneniya*, 30(4):665–674, 734, 1994. URL: http://mi.mathnet.ru/de8352.
- [36] F. Rindler. Calculus of variations. Universitext. Springer, Cham, 2018. URL: https://doi.org/10.1007/978-3-319-77637-8.
- [37] F. Rothe. Global solutions of reaction-diffusion systems, volume 1072 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1984. URL: https://doi.org/10.1007/BFb0099278.
- [38] F. Santambrogio. Optimal transport for applied mathematicians, volume 87 of Progress in Nonlinear Differential Equations and their Applications. Birkhäuser/Springer, Cham, 2015. Calculus of variations, PDEs, and modeling, https://doi.org/10.1007/978-3-319-20828-2.
- [39] F. Smarrazzo and A. Tesei. Long-time behavior of solutions to a class of forward-backward parabolic equations. SIAM J. Math. Anal., 42(3):1046-1093, 2010. URL: https://doi.org/10.1137/090763561.
- [40] F. Smarrazzo and A. Tesei. Degenerate regularization of forward-backward parabolic equations: the vanishing viscosity limit. *Math. Ann.*, 355(2):551–584, 2013. URL: https://doi.org/10.1007/s00208-012-0798-9.
- [41] L. Tartar. Compensated compactness and applications to partial differential equations. In Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV, volume 39 of Res. Notes in Math., pages 136–212. Pitman, Boston, Mass.-London, 1979.
- [42] A. Terracina. Qualitative behavior of the two-phase entropy solution of a forward-backward parabolic problem. SIAM J. Math. Anal., 43(1):228–252, 2011. URL: https://doi.org/10.1137/090778833.
- [43] A. Terracina. Non-uniqueness results for entropy two-phase solutions of forward-backward parabolic problems with unstable phase. J. Math. Anal. Appl., 413(2):963-975, 2014. URL: https://doi.org/10.1016/j.jmaa.2013.12.045.
- [44] L. C. Young. Generalized surfaces in the calculus of variations. Ann. of Math. (2), 43:84–103, 1942. URL: https://doi.org/10.2307/1968882.
- [45] L. C. Young. Generalized surfaces in the calculus of variations. II. Ann. of Math. (2), 43:530–544, 1942. URL: https://doi.org/10.2307/1968809.

Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Stefana Banacha 2, 02-097 Warsaw, Poland

 $Email\ address: \verb"jakub.skrzeczkowski@student.uw.edu.pl"$