

Asymptotic Stability of Phase Separation States for Compressible Immiscible Two-Phase Flow with Periodic Boundary Condition in 3D

Yazhou Chen* Hakho Hong[†] Xiaoding Shi[‡]

Abstract

This paper is concerned with a diffuse interface model called as Navier-Stokes/Cahn-Hilliard system. This model is usually used to describe the motion of immiscible two-phase flow with diffusion interface. For the periodic boundary value problem of this system in torus \mathbb{T}^3 , we prove that there exists a global unique strong solution near the phase separation state, which means no vacuum, shock wave, mass concentration, interface collision and rupture will be developed in finite time. Furthermore, we established the large time behavior of these global strong solution of this system. In particular, we find that the phase field decays algebraically to the phase separation state.

MSC 2020: 35B40, 35B65, 35L65, 76N05, 76N10, 76T10.

Keywords: Navier-Stokes/Cahn-Hilliard system, existence, uniqueness, large-time behavior.

1 Introduction

Two-phase flows or multi-phase flows are important in many industrial applications, for instance, in aerospace, chemical engineering, micro-technology and so on. They have attracted studies from many engineers, geophysicists and astrophysicists. We focus on diffuse interface model which describes the motion of a mixture of two compressible viscous fluids with different densities. Macroscopically immiscible two-phase fluids are assumed to be separated by a sharp interface. However, in order to describe topological transitions, such as droplet formation, coalescence of several droplet or droplet breakup, we need to take into account a partial mixing on a small length scale in the model. As a result, the sharp interface of the two fluids is replaced by a narrow transition layer, and an order parameter related to the concentration difference of both fluids is introduced.

In this paper, we consider the following barotropic compressible Navier-Stokes/Cahn-Hilliard system (see Heida-Málek-Rajagopal [15] and Abels-Feireisl [1]) describing immis-

*Department of Mathematics, College of Mathematics and Physics, Beijing University of Chemical Technology, Beijing 100029, P R China (chenyz@mail.buct.edu.cn).

[†]Institute of Mathematics, State Academy of Sciences, Pyongyang, D P R Korea (hhhong@star-co.net.kp and hhong@amss.ac.cn).

[‡]Corresponding author. Department of Mathematics, College of Mathematics and Physics, Beijing University of Chemical Technology, Beijing 100029, P R China (shixd@mail.buct.edu.cn)

cible two-phase flow with diffuse interface:

$$\begin{cases} \rho_t + \operatorname{div}(\rho \mathbf{u}) = 0, \\ (\rho \mathbf{u})_t + \operatorname{div}(\rho \mathbf{u} \otimes \mathbf{u}) + \nabla p(\rho) = \operatorname{div} \mathbf{S} - \operatorname{div} \left(\nabla \phi \otimes \nabla \phi - \frac{|\nabla \phi|^2}{2} \mathbb{I} \right), \\ (\rho \phi)_t + \operatorname{div}(\rho \phi \mathbf{u}) = \Delta \mu, \\ \mu = -\frac{\epsilon}{\rho} \Delta \phi + \frac{1}{\epsilon} \partial_\phi f(\phi), \end{cases} \quad (1.1)$$

where $\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{T}^3$, \mathbb{T}^3 is the torus in \mathbb{R}^3 , $t > 0$ the time variable. The unknown functions ρ , \mathbf{u} and ϕ represent the density, velocity and phase field respectively. Notice that, here $\{\mathbf{x} : \phi(\mathbf{x}, t) = 1\}$ is occupied by fluid 1 and $\{\mathbf{x} : \phi(\mathbf{x}, t) = -1\}$ by fluid 2 in the area of immiscible two-phase flow. The phase function ϕ is introduced to distinguish between different fluids. In terms of simple, taking any volume element V in the flow, M_i is assumed to be the mass of the components in the representative material volume V , $\phi_i = \frac{\rho_i}{\rho}$ the mass concentration, $\rho_i = \frac{M_i}{V}$ the apparent mass density of the fluid i ($i = 1, 2$). $\rho = \rho_1 + \rho_2$ the total density, and the phase function $\phi = \phi_1 - \phi_2$. Obviously, We can determine the location of the spread interface by the phase function ϕ .

The constant $\epsilon > 0$ is the thickness of the interface between the phases, and the deformation tensor \mathbf{S} is given by

$$\mathbf{S} = 2\nu(\phi)\mathbf{D}(\mathbf{u}) + \lambda(\phi)\operatorname{div}\mathbf{u}\mathbb{I}, \quad (1.2)$$

where $\mathbf{D}(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + \nabla^\top \mathbf{u})$, \top represents the transpose of a matrix, \mathbb{I} is the identity matrix, $\nu(\phi)$ and $\lambda(\phi)$ are two viscosity coefficients. In this paper, we suppose that $\nu(\cdot), \lambda(\cdot) \in C^3(\mathbb{R})$ and there exists positive constant ν_0 such that

$$\nu(s) \geq \nu_0 > 0, \quad \lambda(s) \geq 0. \quad (1.3)$$

Also, $p = p(\rho)$ is the pressure, satisfies

$$p'(\rho) > 0 \quad \text{for all } \rho > 0. \quad (1.4)$$

Here f is the potential function. It is reasonable to take a potential function f as follows (see Heida-Málek-Rajagopal [15])

$$f(\phi) = \frac{\phi^4}{4} - \frac{\phi^2}{2}. \quad (1.5)$$

Remark 1.1 *In the theory of the Cahn-Hilliard equation, double-well structural potential is often considered. A typical example of such potential is of logarithmic type*

$$f(\phi) = \frac{\alpha_1}{2} \left((1 - \phi) \ln\left(\frac{1 - \phi}{2}\right) + (1 + \phi) \ln\left(\frac{1 + \phi}{2}\right) \right) - \frac{\alpha_2}{2} \phi^2,$$

where α_1 and α_2 are positive constants, which is suggested by Cahn-Hilliard [2]. However, this potential is usually replaced by a polynomial approximation of the type $\beta_1 \phi^4 - \beta_2 \phi^2$, where β_1 and β_2 are positive constants (for example, see [28], [7]). Therefore, without loss of generality, the approximate expression (1.5) is usually adopted.

The main purpose of this paper is to give a rigorous analysis of the asymptotic stability of the phase separation state of immiscible two-phase flow. We consider the initial boundary problem of the system (1.1) in torus $\mathbb{T}^3 = (0, 1)^3$ with the periodic boundary condition

$$(\rho, \mathbf{u}, \phi)(\mathbf{x} + \mathbf{L}, t) = (\rho, \mathbf{u}, \phi)(\mathbf{x}, t), \quad t \geq 0, \quad \mathbf{L} = (1, 1, 1), \quad (1.6)$$

and the initial condition

$$(\rho, \mathbf{u}, \phi)(\mathbf{x}, 0) = (\rho_0, \mathbf{u}_0, \phi_0)(\mathbf{x}). \quad \mathbf{x} \in \mathbb{T}^3. \quad (1.7)$$

For the last few decades, the mathematical study of diffusive interface models for compressible immiscible two-phase flow has been attracted by many researchers. For the system (1.1) which can be regarded as a simple variant of the model derived originally by Heida-Málek-Rajagopal [15], Abels-Feireisl [1] proved existence of global weak solutions to the initial boundary value problem in bounded domain $\Omega \subset \mathbb{R}^3$. For the more general system, Kotschote-Zacher [20] established a local existence and uniqueness of strong solutions for the initial boundary value problem in bounded domain $\Omega \subset \mathbb{R}^n (n = 2, 3)$. Recently, in the case of one-dimensional initial boundary value problem, Chen-He-Mei-Shi [4] studied the global existence and the large time behavior of the strong solutions, even for the large initial disturbance of the density and the large velocity data. For the full compressible Navier-Stokes/Cahn-Hilliard system with heat-conductivity, a case more closer to the physical reality, there exist some recent progresses to derive a thermodynamically consistent model (see [15, 19, 10, 11] and so on).

It should be noted that Navier-Stokes/Allen-Cahn system is another commonly used immiscible two-phase flow model for diffusion interfaces. The essential difference between the two models is that, the phase field equation in the Navier-Stokes/Cahn-Hilliard system is a fourth order equation, which is conserved with respect to the $\rho\phi$. But the phase field equation in the Navier-Stokes/Allen-Cahn system is a second order equation, and it is not conserved with respect to the $\rho\phi$. Therefore, the Navier-Stokes/Cahn-Hilliard system is more suitable to describe the phase separation phenomenon, while the Navier-Stokes/Allen-Cahn is more suitable to describe the phase transition phenomenon. There are also many recent studies for the compressible Navier-Stokes/Allen-Cahn system recently, see [9, 18, 7, 3, 22, 14, 6, 8, 27] and references therein.

Notation: In this paper, $L^p(\mathbb{T}^3)$ and $W^{k,p}(\mathbb{T}^3)$ denote the usual Lebesgue and Sobolev spaces on \mathbb{T}^3 , with norms $\|\cdot\|_{L^p}$ and $\|\cdot\|_{W^{k,p}}$, respectively. When $p = 2$, we denote $W^{k,p}(\mathbb{T}^3)$ by $H^k(\mathbb{T}^3)$ with the norm $\|\cdot\|_{H^k}$ and $\|\cdot\|_{H^0} = \|\cdot\|$ will be used to denote the usual L^2 -norm. The notation $\|(A_1, A_2, \dots, A_l)\|_{H^k}$ means the summation of $\|A_i\|_{H^k}$ from $i = 1$ to $i = l$. For an integer m , the symbol ∇^m denotes the summation of all terms D^α with the multi-index α satisfying $|\alpha| = m$. We use C, c to denote the constants which are independent of \mathbf{x}, t and may change from line to line. For 3×3 -matrices F, H , denote $F : H = \sum_{i,j=1}^3 F_{ij}H_{ij}$, $|F| \equiv (F : F)^{1/2}$. For vectors a and b , we denote their tensor product by $a \otimes b := (a_i b_j)_{3 \times 3}$. The integral mean is denoted by the following symbol

$$\int_{\mathbb{T}^3} \cdot d\mathbf{x} \stackrel{\text{def}}{=} \frac{1}{|\mathbb{T}^3|} \int_{\mathbb{T}^3} \cdot d\mathbf{x}. \quad (1.8)$$

We will employ the notation $a \lesssim b$ to mean that $a \leq Cb$ for a universal constant $C > 0$ that only depends on the parameters coming from the problem. Next, in order to establish

the negative Sobolev estimates, we should review the following useful results. But let us first introduce the following necessary definition.

Definition 1.1 For $s \in \mathbb{R}$, $\dot{H}_{\text{per}}^s(\mathbb{T}^3)$ is defined as the homogeneous Sobolev space of f , with the periodic condition and following norm:

$$\|f\|_{\dot{H}^s} \stackrel{\text{def}}{=} \|\Lambda^s f\|,$$

where Λ^s is defined by

$$(\Lambda^s f)(\mathbf{x}) \stackrel{\text{def}}{=} \sum_{\mathbf{k} \in \mathbb{Z}^3} (2\pi\mathbf{k})^s \hat{f}(\mathbf{k}) e^{2\pi i \mathbf{x} \cdot \mathbf{k}}, \quad \mathbf{k} = (k_1, \dots, k_3),$$

where $\hat{f}(\mathbf{k})$ is the Fourier transform of f defined by

$$\hat{f}(\mathbf{k}) = \int_{\mathbb{T}^3} f(\mathbf{x}) e^{-2\pi i \mathbf{x} \cdot \mathbf{k}} d\mathbf{x}.$$

Before giving the main results, for convenience, we present several important Lemmas commonly used in this paper. We set

$$W_{\text{per}}^{k,q}(\mathbb{T}^3) \stackrel{\text{def}}{=} \left\{ w \in W^{k,q}(\mathbb{T}^3) \mid w(\mathbf{x} + \mathbf{L}) = w(\mathbf{x}) \right\}. \quad (1.9)$$

Then, we need the following result which is a direct consequence of Gagliardo-Nirenberg inequality ([23]):

Lemma 1.1 Let l, s and k be any real numbers satisfying $0 \leq l, s < k$, and let $p, r, q \in [1, \infty]$ and $\frac{l}{k} \leq \theta \leq 1$ such that

$$\frac{l}{3} - \frac{1}{p} = \left(\frac{s}{3} - \frac{1}{r} \right) (1 - \theta) + \left(\frac{k}{3} - \frac{1}{q} \right) \theta.$$

Then, we have

$$\|\nabla^l w\|_{L^p} \lesssim \|\nabla^s w\|_{L^r}^{1-\theta} \|\nabla^k w\|_{L^q}^\theta, \quad (1.10)$$

for any $w \in W_{\text{per}}^{k,q}(\mathbb{T}^3)$.

Next, we need the following result:

Lemma 1.2 Let $f(\sigma)$ and $f(\sigma, w)$ be smooth functions of σ and (σ, w) , respectively, with bounded derivatives of any order, and $\|\sigma\|_{L^\infty(\mathbb{T}^3)} \lesssim 1$. Then for any integer $m \geq 1$, we have

$$\|\nabla^m f(\sigma)\|_{L^p} \leq C \|\nabla^m \sigma\|_{L^p}, \quad \|\nabla^m f(\sigma, w)\|_{L^p} \leq C \|\nabla^m(\sigma, w)\|_{L^p}, \quad \forall 1 \leq p \leq \infty, \quad (1.11)$$

where C may depend f and m .

Proof: The (1.11) can be obtained directly making use of the Gagliardo-Nirenberg inequality (Lemma 1.1), we therefore omitted here for the sake of brevity. \blacksquare

Next, we recall the following Moser-type calculus inequalities ([17]):

Lemma 1.3 *Let Ω be a domain of \mathbb{R}^3 . Then, for $f, g \in H^s(\Omega) \cap L^\infty(\Omega)$ and $|\alpha| \leq s$, $s > \frac{3}{2}$, it holds that*

$$\|D^\alpha(fg)\| \lesssim \|f\|_{L^\infty} \|\nabla^s g\| + \|g\|_{L^\infty} \|\nabla^s f\|. \quad (1.12)$$

By the Parseval theorem and Hölder's inequality, it is easy to check the following result (see [13]).

Lemma 1.4 *Let $s \geq 0$ and $l \geq 0$. Then, we have*

$$\|\nabla^l f\| \lesssim \|\nabla^{l+1} f\|^{1-\theta} \|f\|_{\dot{H}^{-s}}^\theta, \quad \text{with } \theta = \frac{1}{l+s+1}. \quad (1.13)$$

If $s \in (0, 3)$, $\Lambda^{-s}g$ is the Riesz potential. Then, we have the following L^p type inequality by the discrete Hardy-Littlewood-Sobolev inequality (see [5],[16],[24]):

Lemma 1.5 *Let $0 < s < 3$, $1 < p < q < \infty$ and $\frac{1}{q} + \frac{s}{3} = \frac{1}{p}$. Then, we have*

$$\|\Lambda^{-s} f\|_{L^q} \lesssim \|f\|_{L^p}. \quad (1.14)$$

Without loss of generality, we assume the thickness of the interface $\epsilon = 1$ here, moreover, noticing that $\text{div}(\nabla\phi \otimes \nabla\phi) = \nabla\left(\frac{|\nabla\phi|^2}{2}\right) + \nabla\phi\Delta\phi$, combining with (1.5)-(1.7), (1.1) is simplified to

$$\begin{cases} \rho_t + \text{div}(\rho\mathbf{u}) = 0, \\ (\rho\mathbf{u})_t + \text{div}(\rho\mathbf{u} \otimes \mathbf{u}) + \nabla p(\rho) + \nabla\phi\Delta\phi = 2\text{div}[\nu(\phi)\mathbf{D}(\mathbf{u})] + \nabla[\lambda(\phi)\text{div}\mathbf{u}], \\ \rho\phi_t + \rho\mathbf{u} \cdot \nabla\phi = \Delta\mu, \\ \rho\mu = -\Delta\phi + \rho(\phi^3 - \phi), \\ (\rho, \mathbf{u}, \phi)(\mathbf{x} + \mathbf{L}, t) = (\rho, \mathbf{u}, \phi)(\mathbf{x}, t), \\ (\rho, \mathbf{u}, \phi)(\mathbf{x}, 0) = (\rho_0, \mathbf{u}_0, \phi_0)(\mathbf{x}). \end{cases} \quad (1.15)$$

Now, the main results is as following below:

Theorem 1.1 *Assume that (1.3)-(1.5) and*

$$\rho_0, \mathbf{u}_0, \phi_0 \in H_{\text{per}}^3(\mathbb{T}^3), \quad \inf_{\mathbf{x} \in \mathbb{T}^3} \rho_0(\mathbf{x}) > 0. \quad (1.16)$$

Then, there exists a positive constant $\delta > 0$ such that if

$$\|\rho_0 - \bar{\rho}\|_{H^3} + \|\mathbf{u}_0\|_{H^3} + \|\nabla\phi_0\|_{H^2} + \|\phi_0^2 - 1\|_{L^2} \leq \delta, \quad (1.17)$$

where $\bar{\rho} = \int_{\mathbb{T}^3} \rho_0(\mathbf{x}) d\mathbf{x}$, then the system (1.15) admits a unique solution (ρ, \mathbf{u}, ϕ) on $[0, \infty)$ satisfying

$$\begin{aligned} \rho, \mathbf{u}, \phi &\in C([0, \infty); H_{\text{per}}^3(\mathbb{T}^3)), \quad \rho \in L^2(0, \infty; H_{\text{per}}^3(\mathbb{T}^3)), \\ \nabla\mathbf{u} &\in L^2(0, \infty; H_{\text{per}}^3(\mathbb{T}^3)), \quad \nabla\phi \in L^2(0, \infty; H_{\text{per}}^4(\mathbb{T}^3)). \end{aligned}$$

and

$$\begin{aligned}
& \|(\rho - \bar{\rho}, \mathbf{u})(t)\|_{H^3}^2 + \|\nabla\phi(t)\|_{H^2}^2 + \|\phi^2(t) - 1\|_{L^2}^2 \\
& + \int_0^t \|\rho(\tau) - \bar{\rho}\|_{H^3}^2 d\tau + \int_0^t \|\nabla\mathbf{u}(\tau)\|_{H^3}^2 d\tau + \int_0^t \|\nabla\phi(\tau)\|_{H^4}^2 d\tau \\
& \leq C (\|(\rho_0 - \bar{\rho}, \mathbf{u}_0)\|_{H^3}^2 + \|\nabla\phi_0\|_{H^2}^2 + \|\phi_0^2 - 1\|_{L^2}^2), \quad \forall t \geq 0,
\end{aligned} \tag{1.18}$$

where C is the positive constant independent of \mathbf{x}, t and δ .

Moreover, if $(\rho_0 - \bar{\rho}, \mathbf{u}_0, \nabla\phi_0, \phi_0^2 - 1) \in \dot{H}^{-s}$ for some $s \in [0, \frac{3}{2})$, we have the following algebraic decay estimates:

$$\|(\rho - \bar{\rho}, \mathbf{u}, \nabla\phi, \phi^2 - 1)(t)\|_{\dot{H}^{-s}} \leq C_0, \tag{1.19}$$

$$\|\mathbf{u}(t)\|_{L^2}^2 + \|\phi^2(t) - 1\|_{L^2} \leq C_0(1+t)^{-s}, \tag{1.20}$$

and

$$\|\rho(t) - \bar{\rho}\|_{H^3}^2 + \|\nabla\mathbf{u}(t)\|_{H^2}^2 + \|\nabla\phi(t)\|_{H^2}^2 \leq C_0(1+t)^{-(2+s)}, \tag{1.21}$$

where \dot{H}^{-s} denotes the homogeneous negative Sobolev space with periodic condition (see Definition 1.1).

By Lemma 1.5, we obtain that for $p \in (1, 2]$, $L^p(\mathbb{T}^3) \subset \dot{H}^{-s}(\mathbb{T}^3)$ with $s = 3\left(\frac{1}{p} - \frac{1}{2}\right) \in [0, \frac{3}{2})$. Then by Theorem 1.1, we have the following corollary of the usual $L^p - L^2$ type of decay results:

Corollary 1.1 *Under the assumptions of Theorem 1.1, if $(\rho_0 - \bar{\rho}, \mathbf{u}_0, \nabla\phi_0, \phi_0^2 - 1) \in L^p_{\text{per}}(\mathbb{T}^3)$ for some $p \in (1, 2]$, then the following decay results hold:*

$$\|\mathbf{u}(t)\|_{L^2}^2 + \|\phi^2(t) - 1\|_{L^2}^2 \lesssim (1+t)^{-3\left(\frac{1}{p} - \frac{1}{2}\right)},$$

and

$$\|\rho(t) - \bar{\rho}\|_{H^3}^2 + \|\nabla\mathbf{u}(t)\|_{H^2}^2 + \|\nabla\phi(t)\|_{H^2}^2 \lesssim (1+t)^{-3+1-3\left(\frac{1}{p} - \frac{1}{2}\right)}.$$

Remark 1.2 *The constraint $s < \frac{3}{2}$ in Theorem 1.1 comes from applying Lemma 1.5 to estimate the nonlinear terms when doing the negative Sobolev estimates via Λ^{-s} . This in turn restricts $p > 1$ in Corollary 1.1 by our method. Note that the nonlinear estimates would not work for $s \geq \frac{3}{2}$.*

Remark 1.3 *A particular feature of the system (1.15) is that the viscosity coefficients ν, λ depend on the mass concentration difference ϕ , which makes our arguments to be difficult. In general, the viscosity coefficients ν, λ may depend not only on the mass concentration difference ϕ , but also the density ρ . Here, we'd like to emphasis that the result of Theorem 1.1 holds by the same lines in this case. Moreover, when the dimension of the space is 2 dimensions, the conclusion of the theorem 1.1 still holds.*

Remark 1.4 *The conclusion of the Theorem 1.1 shows that, in the small perturbation condition near the phase separation state, discontinuous phenomena such as shock wave, vacuum, interface rupture for compressible immiscible two-phase flow will not occur in finite time. Moreover, velocity, density and phase field of immiscible two-phase flow decay algebraically with time. In particular, the interface of two-phase flow algebra decays to sharp interface limit.*

Corollary 1.2 Assume that (1.3)-(1.5), d represents the dimension of space, and for an integer $N \geq 3$,

$$\rho_0, \mathbf{u}_0, \phi_0 \in H^N(\mathbb{T}^d), \quad \inf_{x \in \mathbb{T}^d} \rho_0(x) > 0.$$

So, there exists a positive constant $\delta_0 > 0$ such that if

$$\|\rho_0 - \bar{\rho}\|_{H^N} + \|\mathbf{u}_0\|_{H^N} + \|\nabla \phi_0\|_{H^{N-1}} + \|\phi_0^2 - 1\| \leq \delta_0, \quad (1.22)$$

then the system (1.15) admits a unique solution (ρ, \mathbf{u}, ϕ) on $[0, \infty)$ satisfying

$$\begin{aligned} \rho, \mathbf{u}, \phi &\in C([0, \infty); H^N(\mathbb{T}^d)), \quad \rho \in L^2(0, \infty; H^N(\mathbb{T}^d)), \\ \nabla \mathbf{u} &\in L^2(0, \infty; H^N(\mathbb{T}^d)), \quad \nabla \phi \in L^2(0, \infty; H^{N+1}(\mathbb{T}^d)). \end{aligned}$$

and

$$\begin{aligned} &\|(\rho - \bar{\rho}, \mathbf{u})(t)\|_{H^N}^2 + \|\nabla \phi(t)\|_{H^{N-1}}^2 + \|\phi^2(t) - 1\|^2 \\ &+ \int_0^t \|\rho(\tau) - \bar{\rho}\|_{H^N}^2 d\tau + \int_0^t \|\nabla \mathbf{u}(\tau)\|_{H^N}^2 d\tau + \int_0^t \|\nabla \phi(\tau)\|_{H^{N+1}}^2 d\tau \\ &\leq C (\|(\rho_0 - \bar{\rho}, \mathbf{u}_0)\|_{H^N}^2 + \|\nabla \phi_0\|_{H^{N-1}}^2 + \|\phi_0^2 - 1\|^2), \end{aligned} \quad (1.23)$$

for any $t \geq 0$, where C is the positive constant independent of x, t and η_0 . Moreover, we have the following algebraic decay estimates: if $(\rho_0 - \bar{\rho}, \mathbf{u}_0, \nabla \phi_0, \phi_0^2 - 1) \in \dot{H}_{per}^{-s}(\mathbb{T}^d)$ for some $s \in [0, \frac{d}{2})$, then both (1.19) and (1.20) still hold, and

$$\|(\rho(t) - \bar{\rho}, \mathbf{u}(t))\|_{H^{N-1}}^2 + \|\nabla \phi(t)\|_{H^{N-1}}^2 \leq C_0(1+t)^{-(N-1+s)}. \quad (1.24)$$

To prove Theorem 1.1, we will use the energy method developed by [13], which relies essentially on the following two main steps:

Step 1. Energy estimates at l -th level.

$$\frac{d}{dt} \mathcal{E}_l^m(t) + \|\nabla^l(\rho - \bar{\rho})(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \mathbf{u}(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \phi(t)\|_{H^{m+1-l}}^2 \leq 0, \quad (1.25)$$

for any $0 \leq l < m \leq 3$, where

$$\mathcal{E}_l^m(t) \simeq \|\nabla^l(\rho - \bar{\rho}, \mathbf{u})(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \phi(t)\|_{H^{m-1-l}}^2 + \|\phi^2 - 1\|_{L^2}^2,$$

where $A \simeq B$ means $CA \leq B \leq \frac{1}{C}A$ for a generic constant $C > 0$.

Step 2. Negative Sobolev norm estimate.

$$\frac{d}{dt} \mathcal{E}_{-s}(t) \lesssim (\|\nabla(\rho, \mathbf{u})\|_{H^2}^2 + \|\nabla \phi\|_{H^2}^2) \mathcal{E}_{-s}(t), \quad 0 < s \leq \frac{3}{2}, \quad (1.26)$$

where

$$\mathcal{E}_{-s} = \|\Lambda^{-s}(\rho - \bar{\rho})\|_{L^2}^2 + \|\Lambda^{-s} \mathbf{u}\|_{L^2}^2 + \|\Lambda^{-s} \nabla \phi\|_{L^2}^2 + \|\Lambda^{-s}(\phi^2 - 1)\|_{L^2}^2.$$

If we prove (1.25), then it is easy to show that there exists a solution of the system (1.15) satisfying (1.18) by the continuation argument of local solution (Subsection 3.1). Also, by using (1.25) and the Poincaré inequality

$$\left\| w - \int_{\mathbb{T}^3} w(\mathbf{x}) d\mathbf{x} \right\|_{L^p} \lesssim \|\nabla w\|_{L^p}, \quad \text{for all } w \in W_{per}^{1,p}(\mathbb{T}^3), \quad 1 < p < \infty, \quad (1.27)$$

combining (1.26), Lemma 1.4, the differential inequality

$$\frac{df(t)}{dt} + c_0(f(t))^{1+\frac{1}{l+s}} \leq 0 \Rightarrow f(t) \leq \left(f(0)^{-\frac{1}{l+s}} + \frac{c_0 t}{l+s} \right)^{-(l+s)} \lesssim (1+t)^{-(l+s)}, \quad (1.28)$$

and

$$\|\nabla^k w\|_{L^2} \lesssim \|\nabla^{k+1} w\|_{L^2}, \quad k = 1, 2, \quad \text{for all } w \in H_{\text{per}}^3(\mathbb{T}^3), \quad (1.29)$$

we can get the algebraic decay (1.20) and (1.21) (Subsection 3.2). Therefore, the estimates (1.25) and (1.26) are essential in the proof of Theorem 1.1.

Here, we briefly review some difficulties and key analytical techniques in deriving (1.25) and (1.26), compared with previous works in [26, 25, 12]. The main difficulty comes from the Cahn-Hilliard equation (1.15)_{3,4}, rewritten it as a fourth order nonlinear PDE

$$\phi_t + \frac{1}{\rho} \Delta \left(\frac{\Delta \phi}{\rho} \right) - \frac{2}{\rho} \Delta \phi + \frac{3}{\rho} (1 - \phi^2) \Delta \phi = -\mathbf{u} \cdot \nabla \phi + \frac{6\phi}{\rho} (\nabla \phi)^2, \quad (1.30)$$

where the strong nonlinear term $(1 - \phi^2) \Delta \phi$ makes a trouble for desired estimates because both of $\|\phi(t)\|_{L^2}$ and $\|\phi(t)\|_{L^\infty}$ are not small. On the other hand, the coupling between the Navier-Stokes equations (1.15)_{1,2} and the Cahn-Hilliard equation (1.30) also bring trouble to us. In order to overcome this difficulties, we first find that $\|\phi^2 - 1\|_{L^2}$ is small for small initial data and then $\|\phi(t)\|_{L^\infty}$ is bounded (Lemma 2.1). This makes for us to assume that not only $\|(\sigma, \mathbf{u})\|_{H^3} + \|\nabla \phi\|_{H^2}$ is small, but also $\|\phi^2 - 1\|_{L^2}$ is small in a priori estimates. Based on these facts, we complete the estimate (2.17) for $\nabla \phi$ at k -th level, where a new L^p -estimate (1.11) for nonlinear functions is used essentially together with Gagliardo-Nirenberg inequality (1.10) (Lemma 2.2). Next, we derive the estimate (2.21) and (2.27) for $(\rho - \bar{\rho}, \mathbf{u})$ at k -th level, where the more delicate estimates are needed because of the dependence on ϕ of viscosity coefficients ν and λ (Lemma 2.3 and Lemma 2.4). Combining Lemmas 2.1-2.4, we get (1.25) (see Subsection 3.1). Another main point in this paper is how to get the negative Sobolev norm estimate (1.26) in the case of periodic boundary problem, which is quite different with Cauchy problem in \mathbb{R}^3 of previous works [26, 25, 12] due to they are based essentially on the following inequality

$$\|w\|_{L^{\frac{3}{s}}} \lesssim \begin{cases} \|\nabla w\|^{\frac{1}{2}+s} \|\nabla^2 w\|^{\frac{1}{2}-s}, & s \in (0, \frac{1}{2}], \\ \|w\|^{s-\frac{1}{2}} \|\nabla w\|^{\frac{3}{2}-s}, & s \in (\frac{1}{2}, \frac{3}{2}), \end{cases}$$

which does not hold for periodic boundary problem. Therefore, in order to estimate $\|\Lambda^{-s}(\rho - \bar{\rho}, \mathbf{u}, \nabla \phi)\|^2$, we must rely on Lemma 1.5 and the Poincaré inequality (1.27) (see Lemma 2.5). For example, we have

$$\begin{aligned} & \int_{\mathbb{T}^3} \Lambda^{-s} [(\mathbf{u}, \nabla) \mathbf{u}] \Lambda^{-s} \mathbf{u} dx = \int_{\mathbb{T}^3} (\mathbf{u}', \nabla) \Lambda^{-s} \mathbf{u} \cdot \Lambda^{-s} \mathbf{u} dx \\ & + \int_{\mathbb{T}^3} \Lambda^{-s} [(\mathbf{u} - \mathbf{u}', \nabla) \mathbf{u}] \cdot \Lambda^{-s} \mathbf{u} dx = \int_{\mathbb{T}^3} \Lambda^{-s} [(\mathbf{u} - \mathbf{u}', \nabla) \mathbf{u}] \cdot \Lambda^{-s} \mathbf{u} dx \\ & \stackrel{(1.14)}{\lesssim} \|(\mathbf{u} - \mathbf{u}', \nabla) \mathbf{u}\|_{L^{\frac{1}{\frac{1}{2}+\frac{s}{3}}}} \|\Lambda^{-s} \mathbf{u}\| \lesssim \|\mathbf{u} - \mathbf{u}'\|_{L^{\frac{3}{s}}} \|\nabla \mathbf{u}\| \|\Lambda^{-s} \mathbf{u}\| \\ & \stackrel{(1.27)}{\lesssim} \|\nabla \mathbf{u}\|_{L^{\frac{3}{s}}} \|\nabla \mathbf{u}\| \|\Lambda^{-s} \mathbf{u}\| \lesssim \|\nabla \mathbf{u}\|_{H^2}^2 \|\Lambda^{-s} \mathbf{u}\|, \end{aligned}$$

where $\mathbf{u}' = \int_{\mathbb{T}^3} \mathbf{u} dx$. Also, for the estimate on $\|\Lambda^{-s}(\phi^2 - 1)\|^2$, multiplying (1.30) by 2ϕ and applying Λ^{-s} to the resulting equality (2.36), we could obtain the following type of inequality

$$\frac{d}{dt} \|\Lambda^{-s}(\phi^2 - 1)\|^2 + \|\Lambda^{-s} \nabla(\phi^2 - 1)\|^2 + \|\Lambda^{-s} \Delta(\phi^2 - 1)\|^2 \lesssim \dots, ,$$

which is another key point (see Lemma 2.6).

2 The local existence and some energy estimates

In this section, a priori estimate for (1.15) are established. First, we will give the solution space. For any interval $I \subset [0, \infty)$, and $\forall M > 0$, we suppose that $(\sigma, \mathbf{u}, \phi) \in X_M(I)$ is the solution to the system (1.15), where the solution space $X_M(I)$ is defined as follows

$$\begin{aligned} X_M(I) \stackrel{\text{def}}{=} \left\{ (\sigma, \mathbf{u}, \phi) \mid (\sigma, \mathbf{u}) \in C(I; H_{\text{per}}^3(\mathbb{T}^3)), \nabla \phi \in C(I; H_{\text{per}}^2(\mathbb{T}^3)), \right. \\ \phi^2 - 1 \in C(I; L_{\text{per}}^2(\mathbb{T}^3)), \nabla \sigma \in L^2(I; H_{\text{per}}^2(\mathbb{T}^3)), \\ \nabla \phi \in L^2(I; H_{\text{per}}^4(\mathbb{T}^3)), \nabla \mathbf{u} \in L^2(I; H_{\text{per}}^3(\mathbb{T}^3)), \\ \left. \sup_{t \in I} (\|(\sigma, \mathbf{u})\|_{H^3} + \|\nabla \phi\|_{H^2} + \|\phi^2(t) - 1\|) \leq M, \inf_{\mathbf{x} \in \mathbb{T}^3, t \in I} \rho(\mathbf{x}, t) > 0 \right\}. \end{aligned} \quad (2.1)$$

Now the local existence for (1.15) is established below. The proof process is classical, and it can be easily obtained by using traditional methods such as linearization techniques for equations and fixed point theorems. We will not give the details.

Proposition 2.1 (local existence). *Suppose that (1.3)-(1.5) and (1.16) are satisfied, $\bar{\rho} = \int_{\mathbb{T}^3} \rho_0(\mathbf{x}) dx$. Let $\|(\rho_0 - \bar{\rho}, \mathbf{u}_0)\|_{H^3} + \|\nabla \phi_0\|_{H^2} + \|\phi_0^2 - 1\| \leq M$, $\inf_{\mathbf{x} \in \mathbb{T}^3} \rho_0(\mathbf{x}) > 0$, then there exists T^* small enough, such that, there exists a unique solution $(\rho, \mathbf{u}, \phi) \in X_{2M}([0, T^*])$ to the system (1.15), satisfying*

$$\begin{aligned} (\rho, \mathbf{u}, \phi) \in C([0, T^*]; H^3(\mathbb{T}^3)), \rho \in L^2([0, T^*]; H^3(\mathbb{T}^3)), \\ \nabla \phi \in L^2([0, T^*]; H^4(\mathbb{T}^3)), \nabla \mathbf{u} \in L^2([0, T^*]; H^3(\mathbb{T}^3)). \end{aligned} \quad (2.2)$$

On the basis of the existence of local solutions, We will now extend the local solution to the global solution. For this purpose, we need to give some related energy inequalities. Setting

$$\sigma = \rho - \bar{\rho}, \quad (2.3)$$

we rewrite (1.15)₁ - (1.15)₂ and (1.15)_{3,4} as following

$$\begin{cases} \sigma_t + \bar{\rho} \operatorname{div} \mathbf{u} = g_1, \\ \mathbf{u}_t - \frac{2}{\bar{\rho}} \operatorname{div} [\nu(\phi) \mathbf{D}(\mathbf{u})] - \frac{1}{\bar{\rho}} \nabla [(\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}] + \frac{p'(\bar{\rho})}{\bar{\rho}} \nabla \sigma + \frac{1}{\bar{\rho}} \nabla \phi \Delta \phi = \mathbf{g}_2, \end{cases} \quad (2.4)$$

and

$$\phi_t + \mathbf{u} \cdot \nabla \phi = -\frac{1}{\rho} \Delta \left(\frac{\Delta \phi}{\rho} \right) + \frac{3\phi^2 - 1}{\rho} \Delta \phi + \frac{6\phi}{\rho} (\nabla \phi)^2, \quad (2.5)$$

where

$$g_1 = -\operatorname{div}(\sigma \mathbf{u}), \quad (2.6)$$

$$\begin{aligned} \mathbf{g}_2 = & -(\mathbf{u}, \nabla) \mathbf{u} + h_1(\sigma) \nabla \sigma - 2h_2(\sigma) \operatorname{div} [\nu(\phi) \mathbf{D}(\mathbf{u})] \\ & - h_2(\sigma) \nabla [(\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u} - \nabla \phi \Delta \phi], \end{aligned} \quad (2.7)$$

and

$$h_1(\sigma) = \frac{p'(\bar{\rho})}{\bar{\rho}} - \frac{p'(\rho)}{\rho}, \quad h_2(\sigma) = \frac{1}{\bar{\rho}} - \frac{1}{\rho}.$$

By using the solution space (2.1), combining with the Sobolev embedding theorem, we can choose $M_0 > 0$, such that, $\forall 0 < M < M_0$,

$$0 < \frac{\bar{\rho}}{2} \leq \rho(\mathbf{x}, t) \leq 2\bar{\rho}. \quad (2.8)$$

We first give the following energy estimate:

Lemma 2.1 *Under the assumption (2.1), it holds that*

$$\frac{d}{dt} \int_{\mathbb{T}^3} \left(\frac{\rho}{2} \mathbf{u}^2 + G(\rho) + \frac{1}{2} |\nabla \phi|^2 + \frac{\rho}{4} (\phi^2 - 1)^2 \right) d\mathbf{x} + \frac{\nu_0}{2} \|\nabla \mathbf{u}\|^2 + \|\nabla \mu\|^2 \leq 0, \quad (2.9)$$

$$\|\phi(t)\| \lesssim 1 \quad \text{and} \quad \|\phi(t)\|_{L^\infty} \lesssim 1, \quad (2.10)$$

where

$$G(\rho) = \rho \int_{\bar{\rho}}^{\rho} \frac{p(z) - p(\bar{\rho})}{z^2} dz, \quad \rho > 0. \quad (2.11)$$

Proof: Using (2.11) and (1.15)₁, it is easy to see

$$\begin{aligned} \rho G'(\rho) &= G(\rho) + (p(\rho) - p(\bar{\rho})), \quad \rho G''(\rho) = p'(\rho), \\ G(\rho)_t + \operatorname{div}(G(\rho) \mathbf{u}) + (p(\rho) - p(\bar{\rho})) \operatorname{div} \mathbf{u} &= 0. \end{aligned} \quad (2.12)$$

Using (1.15)₂ and (2.12) yields that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{T}^3} \left(\frac{1}{2} \rho \mathbf{u}^2 + G(\rho) \right) d\mathbf{x} + \frac{1}{2} \int_{\mathbb{T}^3} \nu(\phi) |\nabla \mathbf{u} + \nabla \mathbf{u}^T|^2 d\mathbf{x} \\ + \int_{\mathbb{T}^3} \lambda(\phi) |\operatorname{div} \mathbf{u}|^2 d\mathbf{x} = - \int_{\mathbb{T}^3} \mathbf{u} \cdot \nabla \phi \Delta \phi d\mathbf{x}. \end{aligned} \quad (2.13)$$

Multiplying (1.15)₃ by μ and using (1.15)₄, we have

$$\frac{1}{2} \frac{d}{dt} \int_{\mathbb{T}^3} |\nabla \phi|^2 d\mathbf{x} + \frac{1}{4} \frac{d}{dt} \int_{\mathbb{T}^3} \rho (\phi^2 - 1)^2 d\mathbf{x} + \int_{\mathbb{T}^3} |\nabla \mu|^2 d\mathbf{x} = - \int_{\mathbb{T}^3} \nabla(\mathbf{u} \cdot \nabla \phi) \cdot \nabla \phi d\mathbf{x}, \quad (2.14)$$

where we used

$$\frac{1}{4} \int_{\mathbb{T}^3} (\phi^2 - 1)^2 \operatorname{div}(\rho \mathbf{u}) d\mathbf{x} = - \int_{\mathbb{T}^3} \rho \mathbf{u} \cdot \nabla \phi (\phi^3 - \phi) d\mathbf{x}.$$

Adding (2.13) and (2.14), and using (1.3), we get (2.9). By (2.11), (1.4) and (2.8), we have $G(\bar{\rho}) = G'(\bar{\rho}) = 0$ and

$$c_{\bar{\rho}}(\rho - \bar{\rho})^2 \leq G(\rho) \leq C_{\bar{\rho}}(\rho - \bar{\rho})^2. \quad (2.15)$$

Integrating (2.9) for t , and using (2.8) and (2.15), we obtain

$$\begin{aligned} & \|(\rho - \bar{\rho}, \mathbf{u}, \nabla\phi)(t)\|^2 + \|\phi^2(t) - 1\|^2 + \int_0^t \|(\nabla\mathbf{u}, \nabla\mu)(\tau)\|^2 d\tau \\ & \lesssim \|(\rho_0 - \bar{\rho}, \mathbf{u}_0, \nabla\phi_0)\|^2 + \|\phi_0^2 - 1\|^2. \end{aligned} \quad (2.16)$$

Also, by (2.16), we have

$$\|\phi(t)\|^2 = \int_{\mathbb{T}^3} (\phi^2(t) - 1 + 1) d\mathbf{x} \leq |\mathbb{T}^3|^{\frac{1}{2}} \left(\|\phi^2(t) - 1\| + |\mathbb{T}^3|^{\frac{1}{2}} \right) \leq C,$$

and by Gagliardo-Nirenberg inequality, we get

$$\|\phi(t)\|_{L^\infty} \leq C_0 \|\nabla^2\phi(t)\|^{\frac{3}{4}} \|\phi(t)\|^{\frac{1}{4}} + C_0 \|\phi(t)\| \leq C.$$

The proof of Lemma 2.1 is completed. ■

Next, we derive the following estimate on $\nabla\phi$, which plays an essential role in the proof of Theorem 1.1.

Lemma 2.2 *Under the assumption (2.1), it holds that*

$$\frac{d}{dt} \|\nabla^{k+1}\phi\|^2 + \frac{1}{4\rho^2} \|\nabla^{k+3}\phi\|^2 \lesssim \eta_1 \|\nabla^{k+1}\sigma\|^2, \quad \text{for } k = 0, 1, 2. \quad (2.17)$$

Proof: Applying ∇^k to (2.5) and multiplying it by $-\Delta\nabla^k\phi$ yields that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{T}^3} |\nabla^{k+1}\phi|^2 d\mathbf{x} + \int_{\mathbb{T}^3} \left| \nabla \left(\frac{\nabla^k \Delta\phi}{\rho} \right) \right|^2 d\mathbf{x} - \int_{\mathbb{T}^3} \nabla^k(\mathbf{u} \cdot \nabla\phi) \Delta\nabla^k\phi d\mathbf{x} \\ & = \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{1}{\rho} \right) \nabla^{k-l} \Delta \left(\frac{\Delta\phi}{\rho} \right) \nabla^k \Delta\phi d\mathbf{x} \\ & + \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \Delta \left[\nabla^l \left(\frac{1}{\rho} \right) \nabla^{k-l} \Delta\phi \right] \frac{\nabla^k \Delta\phi}{\rho} d\mathbf{x} - \int_{\mathbb{T}^3} \frac{3\phi^2 - 1}{\rho} |\nabla^k \Delta\phi|^2 d\mathbf{x} \\ & - \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{3\phi^2 - 1}{\rho} \right) \nabla^{k-l} \Delta\phi \nabla^k \Delta\phi d\mathbf{x} - \int_{\mathbb{T}^3} \nabla^k \left(\frac{6\phi}{\rho} (\nabla\phi)^2 \right) \Delta\nabla^k\phi d\mathbf{x}. \end{aligned} \quad (2.18)$$

Noticing that

$$\begin{aligned} & \int_{\mathbb{T}^3} \left| \nabla \left(\frac{\nabla^k \Delta\phi}{\rho} \right) \right|^2 d\mathbf{x} \\ & = \int_{\mathbb{T}^3} \frac{1}{\rho^2} |\nabla^{k+1}\Delta\phi|^2 d\mathbf{x} + \int_{\mathbb{T}^3} \frac{|\nabla\rho|^2}{\rho^4} |\nabla^k \Delta\phi|^2 d\mathbf{x} - \int_{\mathbb{T}^3} \frac{3\phi^2 - 1}{\rho} |\nabla^k \Delta\phi|^2 d\mathbf{x} \\ & \leq - \int_{\mathbb{T}^3} \frac{\phi^2 - 1}{\rho} |\nabla^k \Delta\phi|^2 d\mathbf{x}, \end{aligned}$$

and using (2.8), we obtain from (2.18)

$$\frac{1}{2} \frac{d}{dt} \|\nabla^{k+1}\phi\|^2 + \frac{1}{4\rho^2} \|\nabla^{k+1}\Delta\phi\|^2 \leq I_1, \quad (2.19)$$

where

$$\begin{aligned}
I_1 &= - \underbrace{\int_{\mathbb{T}^3} \frac{\phi^2 - 1}{\rho} |\nabla^k \Delta \phi|^2 dx}_{I_1^1} + \underbrace{\int_{\mathbb{T}^3} \nabla^k (\mathbf{u} \cdot \nabla \phi) \Delta \nabla^k \phi dx}_{I_1^2} \\
&+ \underbrace{\sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{1}{\rho} \right) \nabla^{k-l} \Delta \left(\frac{\Delta \phi}{\rho} \right) \nabla^k \Delta \phi dx}_{I_1^3} \\
&+ \underbrace{\sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \Delta \left[\nabla^l \left(\frac{1}{\rho} \right) \nabla^{k-l} \Delta \phi \right] \frac{\nabla^k \Delta \phi}{\rho} dx}_{I_1^4} \\
&- \underbrace{\sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{3\phi^2 - 1}{\rho} \right) \nabla^{k-l} \Delta \phi \nabla^k \Delta \phi dx}_{I_1^5} \\
&- \underbrace{\sum_{0 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{6\phi}{\rho} \right) \nabla^{k-l} (\nabla \phi)^2 \Delta \nabla^k \phi dx}_{I_1^6} .
\end{aligned} \tag{2.20}$$

We estimate I_1 . For I_1^1 and I_1^2 , by using Höder inequality and (2.1), we have

$$\begin{aligned}
I_1^1 &\lesssim \|\phi^2 - 1\| \|\Delta \nabla^k \phi\|_{L^4}^2 \lesssim \eta_1 \|\nabla^{k+2} \phi\|_{L^4}^2 \\
&\stackrel{(1.10)}{\lesssim} \eta_1 \|\nabla^{k+3} \phi\|^{2-\frac{3}{2}} \|\nabla^{k+2} \phi\|^{\frac{3}{2}} \lesssim \eta_1 \|\nabla^{k+3} \phi\|^2, \\
I_1^2 &\lesssim \|\nabla^k (\mathbf{u} \cdot \nabla \phi)\| \|\Delta \nabla^k \phi\| \\
&\stackrel{(1.12)}{\lesssim} \left(\|\mathbf{u}\|_{L^\infty} \|\nabla^{k+1} \phi\| + \|\nabla \phi\|_{L^\infty} \|\nabla^k \mathbf{u}\| \right) \|\nabla^{k+2} \phi\| \\
&\lesssim \eta_1 \|\nabla^{k+3} \phi\|^2.
\end{aligned}$$

Also, for I_1^3 , by using Höder inequality and (2.1), we have

$$\begin{aligned}
I_1^3 &= - \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla \left[\nabla^l \left(\frac{1}{\rho} \right) \nabla^k \Delta \phi \right] \cdot \nabla^{k-l} \nabla \left[\operatorname{div} \left(\frac{\nabla \phi}{\rho} \right) - \nabla \left(\frac{1}{\rho} \right) \cdot \nabla \phi \right] dx \\
&\lesssim \sum_{1 \leq l \leq k} \left(\left\| \nabla^l \left(\frac{1}{\rho} \right) \right\|_{L^\infty} \|\nabla^{k+3} \phi\| + \left\| \nabla^{l+1} \left(\frac{1}{\rho} \right) \right\|_{L^3} \|\nabla^{k+2} \phi\|_{L^6} \right) \times \\
&\quad \times \left(\left\| \nabla^{k-l+2} \left(\frac{\nabla \phi}{\rho} \right) \right\| + \left\| \nabla^{k-l+1} \left[\nabla \left(\frac{1}{\rho} \right) \cdot \nabla \phi \right] \right\| \right) \\
&\stackrel{(1.11), (1.12)}{\lesssim} \sum_{1 \leq l \leq k} \left(\|\nabla^l \sigma\|_{L^\infty} \|\nabla^{k+3} \phi\| + \|\nabla^{l+1} \sigma\|_{L^3} \|\nabla^{k+2} \phi\|_{L^6} \right) \times \\
&\quad \times \left(\|\nabla^{k-l+3} \phi\| + \|\nabla \phi\|_{L^\infty} \|\nabla^{k-l+2} \sigma\| + \|\nabla^{k-l+2} \phi\| \right) \\
&\stackrel{(1.10), (2.1)}{\lesssim} \eta_1 \sum_{1 \leq l \leq k} \|\nabla^{k+3} \phi\| \left(\|\nabla^{k-l+3} \phi\| + \|\nabla^{k-l+2} \sigma\| + \|\nabla^{k-l+2} \phi\| \right) \\
&\stackrel{(1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+3} \phi\|^2 \right).
\end{aligned}$$

By the similar arguments, we have

$$\begin{aligned}
I_1^4 &= - \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla \left[\nabla^l \left(\frac{1}{\rho} \right) \nabla^{k-l} \Delta \phi \right] \cdot \nabla \left[\frac{\nabla^k \Delta \phi}{\rho} \right] d\mathbf{x} \\
&\lesssim \sum_{1 \leq l \leq k} \left(\left\| \nabla^l \left(\frac{1}{\rho} \right) \right\|_{L^\infty} \|\nabla^{k-l+3} \phi\| + \left\| \nabla^{l+1} \left(\frac{1}{\rho} \right) \right\|_{L^3} \|\nabla^{k-l+2} \phi\|_{L^6} \right) \times \\
&\quad \times \left(\|\nabla^{k+3} \phi\| + \|\nabla^{k+2} \phi\| \left\| \nabla \left(\frac{1}{\rho} \right) \right\|_{L^\infty} \right) \\
&\stackrel{(1.11)}{\lesssim} \sum_{1 \leq l \leq k} \left(\|\nabla^l \sigma\|_{L^\infty} \|\nabla^{k-l+3} \phi\| + \|\nabla^{l+1} \sigma\|_{L^3} \|\nabla^{k-l+2} \phi\|_{L^6} \right) \times \\
&\quad \times \left(\|\nabla^{k+3} \phi\| + \|\nabla^{k+2} \sigma\| \|\nabla \sigma\|_{L^\infty} \right) \\
&\stackrel{(1.10), (2.1)}{\lesssim} \eta_1 \sum_{1 \leq l \leq k} \|\nabla^{k-l+3} \phi\| \left(\|\nabla^{k+3} \phi\| + \|\nabla^{k+2} \sigma\| \right) \stackrel{(1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+3} \phi\|^2 \right),
\end{aligned}$$

$$\begin{aligned}
I_1^5 &= \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{3\phi^2 - 1}{\rho} \right) \nabla^{k-l} \Delta \phi \nabla^k \Delta \phi d\mathbf{x} \\
&\lesssim \sum_{1 \leq l \leq k} \left(\left\| \nabla^l \left(\frac{\phi^2}{\rho} \right) \right\| + \left\| \nabla^l \left(\frac{1}{\rho} \right) \right\| \right) \|\nabla^{k-l+2} \phi\|_{L^3} \|\nabla^{k+2} \phi\|_{L^6} \\
&\stackrel{(1.11)}{\lesssim} \sum_{1 \leq l \leq k} \left(\|\nabla^l \sigma\| + \|\nabla^l \phi\| \right) \|\nabla^{k-l+2} \phi\|_{L^3} \|\nabla^{k+2} \phi\|_{L^6} \\
&\stackrel{(1.10), (2.1)}{\lesssim} \eta_1 \sum_{1 \leq l \leq k} \left(\|\nabla^l \sigma\| + \|\nabla^l \phi\| \right) \|\nabla^{k+3} \phi\| \stackrel{(1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+3} \phi\|^2 \right),
\end{aligned}$$

and

$$\begin{aligned}
I_1^6 &= \sum_{0 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \left(\frac{6\phi}{\rho} \right) \nabla^{k-l} (\nabla \phi)^2 \Delta \nabla^k \phi d\mathbf{x} \\
&\lesssim \sum_{1 \leq l \leq k} \left\| \nabla^l \left(\frac{\phi}{\rho} \right) \right\|_{L^3} \|\nabla^{k-l} (\nabla \phi)^2\| \|\nabla^{k+2} \phi\|_{L^6} \\
&\stackrel{(1.11), (1.12)}{\lesssim} \sum_{1 \leq l \leq k} \left(\|\nabla^l \sigma\|_{L^3} + \|\nabla^l \phi\|_{L^3} \right) \|\nabla \phi\|_{L^\infty} \|\nabla^{k-l+1} \phi\| \|\nabla^{k+2} \phi\|_{L^6} \\
&\stackrel{(1.10), (2.1)}{\lesssim} \eta_1 \sum_{1 \leq l \leq k} \|\nabla^{k-l+1} \phi\| \|\nabla^{k+3} \phi\| \stackrel{(1.29)}{\lesssim} \eta_1 \|\nabla^{k+3} \phi\|^2.
\end{aligned}$$

Substituting the estimates for I_1^j ($j = 1, \dots, 6$) into (2.20), we have

$$I_1 \lesssim \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+3} \phi\|^2 \right),$$

and then, we obtain (2.17) from (2.19). The proof of Lemma 2.2 is completed. \blacksquare

In the following two lemmas, we will give the estimates for (σ, \mathbf{u}) .

Lemma 2.3 *Under the assumption (2.1), it holds that*

$$\frac{d}{dt} \left(\|\nabla^k \mathbf{u}\|^2 + \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\nabla^k \sigma\|^2 \right) + \frac{\nu_0}{\bar{\rho}} \|\nabla^{k+1} \mathbf{u}\|^2 \lesssim \eta_1 \left(\|\nabla^k \sigma\|^2 + \|\nabla^{k+2} \phi\|^2 \right), \quad (2.21)$$

for $k = 0, \dots, 3$.

Proof: We first derive (2.21) for $k = 0$. Multiplying (2.4)₂ by \mathbf{u} , and using (2.4)₁ and (1.3), we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|\mathbf{u}\|^2 + \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\sigma\|^2 \right) + \frac{\nu_0}{\bar{\rho}} \int_{\mathbb{T}^3} |\mathbf{D}(\mathbf{u})|^2 d\mathbf{x} \\ & \leq \underbrace{\int_{\mathbb{T}^3} \left(\mathbf{g}_2 \cdot \mathbf{u} - \frac{p'(\bar{\rho})}{\bar{\rho}^2} g_1 \sigma \right) d\mathbf{x}}_{I_2} - \underbrace{\frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \nabla \phi \Delta \phi \cdot \mathbf{u} d\mathbf{x}}_{I_3}. \end{aligned} \quad (2.22)$$

Using (2.6) and (2.7), it is easy to check by the same lines as in [26, Lemma 2.1] that

$$I_2 \lesssim \eta_1 (\|\sigma\|^2 + \|\nabla \mathbf{u}\|^2).$$

Also, for I_3 , we have

$$I_3 \lesssim \|\nabla \phi\|_{L^3} \|\Delta \phi\| \|\mathbf{u}\|_{L^6} \stackrel{(1.10), (2.1)}{\lesssim} \eta_1 (\|\nabla^2 \phi\|^2 + \|\nabla \mathbf{u}\|^2).$$

Therefore, we obtain (2.21) for $k = 0$ from (2.22).

Next, we derive (2.21) for $k = 1, \dots, 3$. Applying ∇^k to (2.4), we have

$$\begin{aligned} & \nabla^k \sigma_t + \bar{\rho} \operatorname{div} \nabla^k \mathbf{u} = \nabla^k g_1, \\ & \nabla^k \mathbf{u}_t - \frac{2}{\bar{\rho}} \operatorname{div} \left[\nu(\phi) \mathbf{D}(\nabla^k \mathbf{u}) \right] - \frac{1}{\bar{\rho}} \nabla \left[(\nu(\phi) + \lambda(\phi)) \operatorname{div} \nabla^k \mathbf{u} \right] + \frac{p'(\bar{\rho})}{\bar{\rho}} \nabla^k \nabla \sigma \\ & \quad + \frac{1}{\bar{\rho}} \nabla^k [\nabla \phi \Delta \phi] = \nabla^k \mathbf{g}_2 + \frac{2}{\bar{\rho}} \operatorname{div} \left[\nabla^k (\nu(\phi) \mathbf{D}(\mathbf{u})) - \nu(\phi) \mathbf{D}(\nabla^k \mathbf{u}) \right] \\ & \quad + \frac{1}{\bar{\rho}} \nabla \left[\nabla^k ((\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}) - (\nu(\phi) + \lambda(\phi)) \operatorname{div} \nabla^k \mathbf{u} \right]. \end{aligned} \quad (2.23)$$

Multiplying (2.23)₂ by $\nabla^k \mathbf{u}$, and using (2.23)₁, (2.6) and (2.7), we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|\nabla^k \mathbf{u}\|^2 + \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\nabla^k \sigma\|^2 \right) + \frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \nu(\phi) |\mathbf{D}(\nabla^k \mathbf{u})|^2 d\mathbf{x} \\ & \quad + \frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} (\nu(\phi) + \lambda(\phi)) |\operatorname{div} \nabla^k \mathbf{u}|^2 d\mathbf{x} = I_4, \end{aligned} \quad (2.24)$$

where

$$\begin{aligned} I_4 = & \underbrace{\frac{P'(\bar{\rho})}{\bar{\rho}^2} \int_{\mathbb{T}^3} \nabla^k \sigma \nabla^k \operatorname{div}(\sigma \mathbf{u}) d\mathbf{x}}_{I_4^1} + \underbrace{\int_{\mathbb{T}^3} \nabla^k [-(\mathbf{u}, \nabla) \mathbf{u} + h_1(\sigma) \nabla \sigma] \cdot \nabla^k \mathbf{u} d\mathbf{x}}_{I_4^2} \\ & - \underbrace{\int_{\mathbb{T}^3} \nabla^k [2h_2(\sigma) \operatorname{div}(\nu(\phi) \mathbf{D}(\mathbf{u})) + h_2(\sigma) \nabla((\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u})] \cdot \nabla^k \mathbf{u} d\mathbf{x}}_{I_4^3} \end{aligned}$$

$$\begin{aligned}
& \underbrace{-\frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \nabla^k [\nabla \phi \Delta \phi] \cdot \nabla^k \mathbf{u} d\mathbf{x}}_{I_4^4} + \underbrace{\frac{2}{\bar{\rho}} \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \nu(\phi) \mathbf{D}(\nabla^{k-l} \mathbf{u}) : \nabla \nabla^k \mathbf{u} d\mathbf{x}}_{I_4^5} \\
& + \underbrace{\frac{1}{\bar{\rho}} \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l (\nu(\phi) + \lambda(\phi)) \operatorname{div} \nabla^{k-l} \mathbf{u} \operatorname{div} \nabla^k \mathbf{u} d\mathbf{x}}_{I_4^6} \\
& + \underbrace{\int_{\mathbb{T}^3} \nabla^k [h_2(\sigma) \nabla \phi \Delta \phi] \cdot \nabla^k \mathbf{u} d\mathbf{x}}_{I_4^7} .
\end{aligned} \tag{2.25}$$

We estimate I_4 . Noticing that

$$I_4^1 = \frac{1}{2} \int_{\mathbb{T}^3} \operatorname{div} \mathbf{u} (\nabla^k \sigma)^2 d\mathbf{x} + \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \operatorname{div} (\nabla^l \mathbf{u} \nabla^{k-l} \sigma) \nabla^k \sigma d\mathbf{x},$$

we have

$$\begin{aligned}
I_4^1 & \lesssim \|\operatorname{div} \mathbf{u}\|_{L^\infty} \|\nabla^k \sigma\|^2 + \left(\|\nabla^2 \mathbf{u}\|_{L^6} \|\nabla^{k-1} \sigma\|_{L^3} + \|\nabla \mathbf{u}\|_{L^\infty} \|\nabla^k \sigma\| \right) \|\nabla^k \sigma\| \\
& + \sum_{2 \leq l \leq k} \left(\|\nabla^{l+1} \mathbf{u}\| \|\nabla^{k-l} \sigma\|_{L^\infty} + \|\nabla^l \mathbf{u}\|_{L^6} \|\nabla^{k-l+1} \sigma\|_{L^3} \right) \|\nabla^k \sigma\| \\
& \stackrel{(2.1), (1.10)}{\lesssim} \eta_1 \left(\|\nabla^k \sigma\|^2 + \|\nabla^{k-1} \sigma\|^2 \right) + \eta_1 \sum_{2 \leq l \leq k} \|\nabla^{l+1} \mathbf{u}\|^2 \\
& \stackrel{(1.29)}{\lesssim} \eta_1 \left(\|\nabla^k \sigma\|^2 + \|\nabla^{k+1} \mathbf{u}\|^2 \right).
\end{aligned}$$

Next, for I_4^2 , we have

$$\begin{aligned}
I_4^2 & = - \int_{\mathbb{T}^3} \nabla^{k-1} [-(\mathbf{u}, \nabla) \mathbf{u} + h_1(\sigma) \nabla \sigma] \cdot \nabla^{k+1} \mathbf{u} d\mathbf{x} \\
& \stackrel{(1.12)}{\lesssim} \left(\|\mathbf{u}\|_{L^\infty} \|\nabla^k \mathbf{u}\| + \|\nabla^{k-1} \mathbf{u}\| \|\nabla \mathbf{u}\|_{L^\infty} \right) \|\nabla^{k+1} \mathbf{u}\| \\
& + \left(\|h_1(\sigma)\|_{L^\infty} \|\nabla^k \sigma\| + \|\nabla^{k-1} h_1(\sigma)\| \|\nabla \sigma\|_{L^\infty} \right) \|\nabla^{k+1} \mathbf{u}\| \\
& \stackrel{(2.1), (1.11)}{\lesssim} \eta_1 \left(\|\nabla^k \mathbf{u}\| + \|\nabla^{k-1} \mathbf{u}\| + \|\nabla^k \sigma\| + \|\nabla^{k-1} \sigma\| \right) \|\nabla^{k+1} \mathbf{u}\| \\
& \stackrel{(1.29)}{\lesssim} \eta_1 \left(\|\nabla^k \sigma\|^2 + \|\nabla^{k+1} \mathbf{u}\|^2 \right).
\end{aligned}$$

For the term I_4^3 , using Leibniz formula and Höder inequality yields that

$$\begin{aligned}
I_4^3 & = \int_{\mathbb{T}^3} \nabla^{k-1} [2h_2(\sigma) \operatorname{div} (\nu(\phi) \mathbf{D}(\mathbf{u})) + h_2(\sigma) \nabla ((\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u})] \cdot \nabla^{k+1} \mathbf{u} d\mathbf{x} \\
& = \underbrace{\sum_{0 \leq l \leq k-1} C_{k-1}^l \int_{\mathbb{T}^3} \nabla^l h_2(\sigma) \operatorname{div} \nabla^{k-l-1} (\nu(\phi) \mathbf{D}(\mathbf{u})) \cdot \nabla^{k+1} \mathbf{u} d\mathbf{x}}_{J_1} \\
& + \underbrace{\sum_{0 \leq l \leq k-1} C_{k-1}^l \int_{\mathbb{T}^3} \nabla^l h_2(\sigma) \nabla^{k-l} ((\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}) \cdot \nabla^{k+1} \mathbf{u} d\mathbf{x}}_{J_2} .
\end{aligned} \tag{2.26}$$

By using Höder inequality, (2.1) and Sobolev embedding, we have

$$\begin{aligned}
J_1 &\lesssim \|h_2(\sigma)\|_{L^\infty} \|\nabla(\nu(\phi)\mathbf{D}(\mathbf{u}))\| \|\nabla^2\mathbf{u}\| \\
&\lesssim \eta_1 (\|\nu(\phi)\|_{L^\infty} \|\nabla^2\mathbf{u}\| + \|\nu'(\phi)\|_{L^\infty} \|\nabla\phi\| \|\nabla\mathbf{u}\|) \|\nabla^2\mathbf{u}\| \\
&\stackrel{(2.10)}{\lesssim} \eta_1 (\|\nabla^2\mathbf{u}\| + \|\nabla\mathbf{u}\|) \|\nabla^2\mathbf{u}\| \stackrel{(1.29)}{\lesssim} \eta_1 \|\nabla^2\mathbf{u}\|^2
\end{aligned}$$

for $k = 1$, and

$$\begin{aligned}
J_1 &\lesssim \sum_{0 \leq l \leq k-1} \|\nabla^l h_2(\sigma) \nabla^{k-l}(\nu(\phi)\mathbf{D}(\mathbf{u}))\| \|\nabla^{k+1}\mathbf{u}\| \\
&\lesssim \|h_2(\sigma)\|_{L^\infty} \|\nabla^k(\nu(\phi)\mathbf{D}(\mathbf{u}))\| \|\nabla^{k+1}\mathbf{u}\| \\
&+ \sum_{1 \leq l \leq k-1} \|\nabla^l h_2(\sigma)\|_{L^3} \|\nabla^{k-l}(\nu(\phi)\mathbf{D}(\mathbf{u}))\| \|\nabla^{k+1}\mathbf{u}\| \\
&\stackrel{(1.12),(1.11)}{\lesssim} \eta_1 \left(\|\nu(\phi)\|_{L^\infty} \|\nabla^{k+1}\mathbf{u}\| + \|\nabla^k\phi\| \|\nabla\mathbf{u}\|_{L^\infty} \right) \|\nabla^{k+1}\mathbf{u}\| \\
&+ \sum_{1 \leq l \leq k-1} \|\nabla^l \sigma\|_{L^3} \left(\|\nu(\phi)\|_{L^\infty} \|\nabla^{k-l+1}\mathbf{u}\| + \|\nabla^{k-l}\phi\| \|\nabla\mathbf{u}\|_{L^\infty} \right) \|\nabla^{k+1}\mathbf{u}\| \\
&+ \sum_{1 \leq l \leq k-1} \|\nabla^l \sigma\|_{L^3} \left(\|\nu(\phi)\|_{L^\infty} \|\nabla^{k-l+2}\mathbf{u}\| + \|\nabla^{k-l+1}\phi\| \|\nabla\mathbf{u}\|_{L^\infty} \right) \|\nabla^{k+1}\mathbf{u}\| \\
&\stackrel{(2.1),(1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+2}\phi\|^2 + \|\nabla^{k+1}\mathbf{u}\|^2 \right)
\end{aligned}$$

for $k = 2, \dots, 3$. Therefore, by following the same argument on J_2 , we obtain from (2.26) that

$$I_4^3 \lesssim \eta_1 \left(\|\nabla^{k+2}\phi\|^2 + \|\nabla^{k+1}\mathbf{u}\|^2 \right).$$

Also, for I_4^4 , we have

$$\begin{aligned}
I_4^4 &= - \sum_{0 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^{l+1}\phi \nabla^{k-l}\Delta\phi \cdot \nabla^k \mathbf{u} dx \\
&\lesssim \|\nabla\phi\|_{L^\infty} \|\nabla^{k+2}\phi\| \|\nabla^k\mathbf{u}\| + \sum_{1 \leq l \leq k} \|\nabla^{l+1}\phi\|_{L^6} \|\nabla^{k-l+2}\phi\|_{L^3} \|\nabla^k\mathbf{u}\| \\
&\stackrel{(2.1),(1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+2}\phi\|^2 + \|\nabla^{k+1}\mathbf{u}\|^2 \right).
\end{aligned}$$

For I_4^5 , we have

$$\begin{aligned}
I_4^5 &= \sum_{1 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l \nu(\phi) \mathbf{D}(\nabla^{k-l}\mathbf{u}) : \nabla \nabla^k \mathbf{u} dx \\
&\stackrel{(1.11)}{\lesssim} \sum_{1 \leq l \leq k} \|\nabla^l \phi\|_{L^3} \|\nabla^{k-l+1}\mathbf{u}\|_{L^6} \|\nabla^{k+1}\mathbf{u}\| \stackrel{(2.1),(1.29)}{\lesssim} \eta_1 \|\nabla^{k+1}\mathbf{u}\|^2.
\end{aligned}$$

By the same lines as in I_4^5 and I_4^3 , we have

$$I_4^6 \lesssim \eta_1 \|\nabla^{k+1}\mathbf{u}\|^2, \quad I_4^7 \lesssim \eta_1 (\|\nabla^{k+2}\phi\|^2 + \|\nabla^{k+1}\mathbf{u}\|^2).$$

Using (2.25) and the estimates for $I_4^j (j = 1, \dots, 6)$, we obtain (2.21) for $k = 1, \dots, 3$ from (2.24). The proof of Lemma 2.3 is completed. \blacksquare

Lemma 2.4 *Under the assumption (2.1), it holds that*

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{T}^3} \nabla^k \mathbf{u} \cdot \nabla^{k+1} \sigma d\mathbf{x} + \frac{p'(\bar{\rho})}{2\bar{\rho}} \|\nabla^{k+1} \sigma\|^2 \\ & \lesssim \eta_1 \left(\|\nabla^{k+2} \mathbf{u}\|^2 + \|\nabla^{k+3} \phi\|^2 \right) + \|\nabla^{k+1} \mathbf{u}\|^2, \quad \text{for } k = 0, \dots, 2. \end{aligned} \quad (2.27)$$

Proof: Multiplying (2.23)₂ by $\nabla^{k+1} \sigma$, and using (2.23)₁, (2.6) and (2.7), we have

$$\frac{d}{dt} \int_{\mathbb{T}^3} \nabla^k \mathbf{u} \cdot \nabla^{k+1} \sigma d\mathbf{x} + \frac{p'(\bar{\rho})}{\bar{\rho}} \|\nabla^{k+1} \sigma\|^2 - \bar{\rho} \int_{\mathbb{T}^3} (\operatorname{div} \nabla^k \mathbf{u})^2 d\mathbf{x} = I_5, \quad (2.28)$$

where

$$\begin{aligned} I_5 &= \underbrace{\int_{\mathbb{T}^3} \nabla^k \operatorname{div}(\sigma \mathbf{u}) \operatorname{div} \nabla^k \mathbf{u} d\mathbf{x}}_{I_5^1} + \underbrace{\int_{\mathbb{T}^3} \nabla^k [-(\mathbf{u}, \nabla) \mathbf{u} + h_1(\sigma) \nabla \sigma] \cdot \nabla^{k+1} \sigma d\mathbf{x}}_{I_5^2} \\ & - \underbrace{\int_{\mathbb{T}^3} \nabla^k [2h_2(\sigma) \operatorname{div}(\nu(\phi) \mathbf{D}(\mathbf{u})) + h_2(\sigma) \nabla((\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u})] \cdot \nabla^{k+1} \sigma d\mathbf{x}}_{I_5^3} \\ & - \underbrace{\frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \nabla^k [\nabla \phi \Delta \phi] \cdot \nabla^{k+1} \sigma d\mathbf{x}}_{I_5^4} + \underbrace{\frac{2}{\bar{\rho}} \int_{\mathbb{T}^3} \operatorname{div} \nabla^k [\nu(\phi) \mathbf{D}(\mathbf{u})] \cdot \nabla^{k+1} \sigma d\mathbf{x}}_{I_5^5} \\ & + \underbrace{\frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \nabla^{k+1} [(\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}] \cdot \nabla^{k+1} \sigma d\mathbf{x}}_{I_5^6} \\ & + \underbrace{\int_{\mathbb{T}^3} \nabla^k [h_2(\sigma) \nabla \phi \Delta \phi] \cdot \nabla^{k+1} \sigma d\mathbf{x}}_{I_5^7}. \end{aligned} \quad (2.29)$$

We estimate I_5 . For the term I_5^1 , we get

$$\begin{aligned} I_5^1 & \lesssim \|\nabla^{k+1}(\sigma \mathbf{u})\| \|\nabla^{k+1} \mathbf{u}\| \\ & \stackrel{(1.12)}{\lesssim} \left(\|\sigma\|_{L^\infty} \|\nabla^{k+1} \mathbf{u}\| + \|\mathbf{u}\|_{L^\infty} \|\nabla^{k+1} \sigma\| \right) \|\nabla^{k+1} \mathbf{u}\| \\ & \stackrel{(2.1), (1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+2} \mathbf{u}\|^2 + \|\nabla^{k+1} \sigma\|^2 \right). \end{aligned}$$

Next, for I_5^2 , we have

$$\begin{aligned} I_5^2 & \lesssim \|\nabla^k [-(\mathbf{u}, \nabla) \mathbf{u} + h_1(\sigma) \nabla \sigma]\| \|\nabla^{k+1} \sigma\| \\ & \stackrel{(1.12)}{\lesssim} \left(\|\mathbf{u}\|_{L^\infty} \|\nabla^{k+1} \mathbf{u}\| + \|\nabla^k \mathbf{u}\| \|\nabla \mathbf{u}\|_{L^\infty} \right) \|\nabla^{k+1} \sigma\| \\ & + \left(\|h_1(\sigma)\|_{L^\infty} \|\nabla^{k+1} \sigma\| + \|\nabla^k h_1(\sigma)\| \|\nabla \sigma\|_{L^\infty} \right) \|\nabla^{k+1} \sigma\| \\ & \stackrel{(2.1), (1.11)}{\lesssim} \eta_1 \left(\|\nabla^{k+1} \mathbf{u}\| + \|\nabla^k \mathbf{u}\| + \|\nabla^{k+1} \sigma\| + \|\nabla^k \sigma\| \right) \|\nabla^{k+1} \sigma\| \\ & \stackrel{(1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+2} \mathbf{u}\|^2 \right). \end{aligned}$$

By the same lines as above, noticing that

$$\begin{aligned} I_5^3 &= - \sum_{0 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^l h_2(\sigma) \operatorname{div} \nabla^{k-l-1} (\nu(\phi) \mathbf{D}(\mathbf{u})) \cdot \nabla^{k+1} \sigma d\mathbf{x} \\ &\quad - \sum_{0 \leq l \leq k} C_{k-1}^l \int_{\mathbb{T}^3} \nabla^l h_2(\sigma) \nabla^{k-l} ((\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}) \cdot \nabla^{k+1} \sigma d\mathbf{x}, \end{aligned}$$

we have

$$I_5^3 \lesssim \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+2} \mathbf{u}\|^2 + \|\nabla^{k+3} \phi\|^2 \right).$$

Also, for I_5^4 , we have

$$\begin{aligned} I_5^4 &= - \sum_{0 \leq l \leq k} C_k^l \int_{\mathbb{T}^3} \nabla^{l+1} \phi \nabla^{k-l} \Delta \phi \cdot \nabla^{k+1} \sigma d\mathbf{x} \\ &\lesssim \|\nabla \phi\|_{L^\infty} \|\nabla^{k+2} \phi\| \|\nabla^{k+1} \sigma\| + \sum_{1 \leq l \leq k} \|\nabla^{l+1} \phi\|_{L^6} \|\nabla^{k-l+2} \phi\|_{L^3} \|\nabla^{k+1} \sigma\| \\ &\stackrel{(2.1), (1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+3} \phi\|^2 + \|\nabla^{k+1} \sigma\|^2 \right). \end{aligned}$$

For I_5^5 , we have

$$\begin{aligned} I_5^5 &= \int_{\mathbb{T}^3} \operatorname{div} \nabla^k [\nu(\phi) \mathbf{D}(\mathbf{u})] \cdot \nabla^{k+1} \sigma d\mathbf{x} \\ &= \sum_{0 \leq l \leq k} C_{k+1}^l \int_{\mathbb{T}^3} \left(\nabla^l \nu(\phi) \operatorname{div} \mathbf{D}(\nabla^{k-l} \mathbf{u}) + \nabla^{l+1} \nu(\phi) \mathbf{D}(\nabla^{k-l} \mathbf{u}) \right) \cdot \nabla^{k+1} \sigma d\mathbf{x} \\ &\stackrel{(1.11)}{\lesssim} \sum_{1 \leq l \leq k} \left(\|\nabla^l \phi\|_{L^3} \|\nabla^{k-l+2} \mathbf{u}\|_{L^6} + \|\nabla^{l+1} \phi\|_{L^3} \|\nabla^{k-l+1} \mathbf{u}\|_{L^6} \right) \|\nabla^{k+1} \sigma\| \\ &\quad + \left(\|\nu(\phi)\|_{L^\infty} \|\nabla^{k+2} \mathbf{u}\| + \|\nabla \nu(\phi)\|_{L^3} \|\nabla^{k+1} \mathbf{u}\|_{L^6} \right) \|\nabla^{k+1} \sigma\| \\ &\stackrel{(2.1), (1.29)}{\lesssim} \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+2} \mathbf{u}\|^2 \right). \end{aligned}$$

By the same lines as in I_5^5 and I_5^4 , we have

$$I_5^6 \lesssim \eta_1 \left(\|\nabla^{k+1} \sigma\|^2 + \|\nabla^{k+2} \mathbf{u}\|^2 \right), \quad I_5^7 \lesssim \eta_1 \left(\|\nabla^{k+3} \phi\|^2 + \|\nabla^{k+1} \sigma\|^2 \right).$$

Using (2.29) and the estimates for I_5^j ($j = 1, \dots, 7$), we obtain (2.27) from (2.28). The proof of Lemma 2.4 is completed. \blacksquare

To get the decay of the solution, we need the following evolution of the negative Sobolev norms of the solution to the system (1.15).

Lemma 2.5 *Under the assumption (2.1), it holds that*

$$\begin{aligned} &\frac{d}{dt} \left(\frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\Lambda^{-s} \sigma\|^2 + \|\Lambda^{-s} \mathbf{u}\|^2 + \|\Lambda^{-s} \nabla \phi\|^2 \right) \\ &\lesssim \left(\|\nabla(\sigma, \mathbf{u})\|_{H^2}^2 + \|\nabla \phi\|_{H^3}^2 \right) \left(\|\Lambda^{-s} \sigma\| + \|\Lambda^{-s} \mathbf{u}\| + \|\Lambda^{-s} \nabla \phi\| \right), \quad \text{for } s \in \left(0, \frac{3}{2}\right). \end{aligned} \tag{2.30}$$

Proof: Applying Λ^{-s} to (2.4)₁, (2.4)₂ and (2.5), multiplying the resulting identities by $\frac{p'(\bar{\rho})}{\bar{\rho}^2}\Lambda^{-s}\sigma$, $\Lambda^{-s}\mathbf{u}$ and $-\Lambda^{-s}\Delta\phi$, respectively, summing up and then integrating by parts, we deduce that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \left(\frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\Lambda^{-s}\sigma\|^2 + \|\Lambda^{-s}\mathbf{u}\|^2 + \|\Lambda^{-s}\nabla\phi\|^2 \right) + \frac{\nu'}{\bar{\rho}} \|\Lambda^{-s}\nabla\mathbf{u}\|^2 \\
& + \frac{\nu' + \lambda'}{\bar{\rho}} \|\Lambda^{-s}\operatorname{div}\mathbf{u}\|^2 + \frac{1}{\bar{\rho}^2} \|\Lambda^{-s}\nabla\Delta\phi\|^2 \\
& = - \frac{2}{\bar{\rho}} \int_{\mathbb{T}^3} \Lambda^{-s} \operatorname{div} [(\nu(\phi) - \nu')\mathbf{D}(\mathbf{u})] \Lambda^{-s}\mathbf{u} d\mathbf{x} \quad \underline{I_6} \\
& - \frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \Lambda^{-s} \nabla [(\nu(\phi) - \nu') + (\lambda(\phi) - \lambda')\operatorname{div}\mathbf{u}] \Lambda^{-s}\mathbf{u} d\mathbf{x} \quad \underline{I_7} \\
& - \frac{p'(\bar{\rho})}{\bar{\rho}^2} \int_{\mathbb{T}^3} \Lambda^{-s} (\sigma \operatorname{div}\mathbf{u} + \nabla\sigma \cdot \mathbf{u}) \Lambda^{-s}\sigma d\mathbf{x} \quad \underline{I_8} - \int_{\mathbb{T}^3} \Lambda^{-s} [(\mathbf{u} \cdot \nabla)\mathbf{u} - h_1(\sigma)\nabla\sigma] \cdot \Lambda^{-s}\mathbf{u} d\mathbf{x} \quad \underline{I_9} \\
& - \int_{\mathbb{T}^3} \Lambda^{-s} [2h_2(\sigma)\operatorname{div}[\nu(\phi)\mathbf{D}(\mathbf{u})] + h_2(\sigma)\nabla[(\nu(\phi) + \lambda(\phi))\operatorname{div}\mathbf{u}]] \cdot \Lambda^{-s}\mathbf{u} d\mathbf{x} \quad \underline{I_{10}} \\
& - \frac{1}{\bar{\rho}} \int_{\mathbb{T}^3} \Lambda^{-s} (\nabla\phi\Delta\phi) \cdot \Lambda^{-s}\mathbf{u} d\mathbf{x} \quad \underline{I_{11}} + \int_{\mathbb{T}^3} \Lambda^{-s} \nabla \left(\frac{3\phi^2 - 1}{\rho} \Delta\phi + \frac{6\phi}{\rho} (\nabla\phi)^2 \right) \Lambda^{-s}\nabla\phi d\mathbf{x} \quad \underline{I_{12}} \\
& + \int_{\mathbb{T}^3} \Lambda^{-s} \nabla \left(-\mathbf{u} \cdot \nabla\phi + h_3(\sigma)\Delta^2\phi + \frac{2}{\rho^3} \Delta\nabla\phi \cdot \nabla\sigma + \frac{\Delta\phi}{\rho} \operatorname{div}\left(\frac{\nabla\sigma}{\rho^2}\right) \right) \Lambda^{-s}\nabla\phi d\mathbf{x} \quad \underline{I_{13}} \\
& + \int_{\mathbb{T}^3} \Lambda^{-s} (h_2(\sigma)\nabla\phi\Delta\phi) \cdot \Lambda^{-s}\mathbf{u} d\mathbf{x} \quad \underline{I_{14}},
\end{aligned} \tag{2.31}$$

where

$$\nu' = \nu \left(\int_{\mathbb{T}^3} \phi d\mathbf{x} \right), \quad \lambda' = \lambda \left(\int_{\mathbb{T}^3} \phi d\mathbf{x} \right) \quad \text{and} \quad h_3(\sigma) = \frac{1}{\bar{\rho}^2} - \frac{1}{\rho^2}.$$

In order to estimate the nonlinear terms in the right-hand side of (2.31), we shall use the estimate (1.14). If $s \in (0, \frac{3}{2})$, then $\frac{1}{2} + \frac{s}{3} < 1$ and we have

$$\begin{aligned}
I_6 & \lesssim \|\Lambda^{-s} \operatorname{div} [(\nu(\phi) - \nu')\mathbf{D}(\mathbf{u})]\| \|\Lambda^{-s}\mathbf{u}\| \\
& \stackrel{(1.14)}{\lesssim} \|\operatorname{div} [(\nu(\phi) - \nu')\mathbf{D}(\mathbf{u})]\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}} \|\Lambda^{-s}\mathbf{u}\| \\
& \lesssim \|\operatorname{div}[\nu(\phi) - \nu']\|_{L^{\frac{3}{s}}} \|\nabla\mathbf{u}\| \|\Lambda^{-s}\mathbf{u}\| \\
& \lesssim \left(\|\nabla\nu(\phi)\|_{L^{\frac{3}{s}}} \|\nabla\mathbf{u}\| + \left\| \phi - \int_{\mathbb{T}^3} \phi d\mathbf{x} \right\|_{L^{\frac{3}{s}}} \|\nabla^2\mathbf{u}\| \right) \|\Lambda^{-s}\mathbf{u}\| \\
& \stackrel{(2.10), (1.27)}{\lesssim} \|\nabla\phi\|_{L^{\frac{3}{s}}} \|\nabla\mathbf{u}\| \|\Lambda^{-s}\mathbf{u}\| \stackrel{(2.32)}{\lesssim} \|\nabla\phi\|_{H^2} \|\nabla\mathbf{u}\| \|\Lambda^{-s}\mathbf{u}\| \\
& \lesssim (\|\nabla\phi\|_{H^2}^2 + \|\nabla\mathbf{u}\|^2) \|\Lambda^{-s}\mathbf{u}\|,
\end{aligned}$$

where we used the Sobolev inequality

$$\|w\|_{L^p} \lesssim \|w\|_{H^2} \quad \text{for all } w \in H^2(\mathbb{T}^3), \quad 1 \leq p \leq \infty. \tag{2.32}$$

Similarly, we can prove that

$$I_7 \lesssim (\|\nabla\phi\|_{H^2}^2 + \|\nabla\mathbf{u}\|^2) \|\Lambda^{-s}\mathbf{u}\|.$$

To estimate I_8 and I_9 , we rewrite them respectively as

$$I_8 = \int_{\mathbb{T}^3} \Lambda^{-s} [\sigma \operatorname{div} \mathbf{u} + \nabla \sigma \cdot (\mathbf{u} - \mathbf{u}')] \Lambda^{-s} \sigma \, d\mathbf{x}$$

and

$$I_9 = \int_{\mathbb{T}^3} \Lambda^{-s} [(\mathbf{u} - \mathbf{u}', \nabla) \mathbf{u} - h_1(\sigma) \nabla \sigma] \cdot \Lambda^{-s} \mathbf{u} \, d\mathbf{x},$$

where $\mathbf{u}' = \int_{\mathbb{T}^3} \mathbf{u} \, d\mathbf{x}$ and we used

$$\int_{\mathbb{T}^3} \mathbf{u}' \cdot \nabla \Lambda^{-s} \sigma \Lambda^{-s} \sigma \, d\mathbf{x} = 0, \quad \int_{\mathbb{T}^3} (\mathbf{u}', \nabla) \Lambda^{-s} \mathbf{u} \cdot \Lambda^{-s} \mathbf{u} \, d\mathbf{x} = 0.$$

Then, we have

$$\begin{aligned} I_8 &\lesssim \|\Lambda^{-s} [\sigma \operatorname{div} \mathbf{u} + \nabla \sigma \cdot (\mathbf{u} - \mathbf{u}')] \| \|\Lambda^{-s} \sigma\| \\ &\stackrel{(1.14)}{\lesssim} \|\sigma \operatorname{div} \mathbf{u} + \nabla \sigma \cdot (\mathbf{u} - \mathbf{u}')\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}}} \|\Lambda^{-s} \sigma\| \\ &\lesssim \left(\|\sigma\|_{L^{\frac{3}{s}}} \|\nabla \mathbf{u}\| + \|\nabla \sigma\|_{L^{\frac{3}{s}}} \|\mathbf{u} - \mathbf{u}'\| \right) \|\Lambda^{-s} \sigma\| \\ &\stackrel{(2.32), (1.27)}{\lesssim} \|\sigma\|_{H^3} \|\nabla \mathbf{u}\| \|\Lambda^{-s} \sigma\| \stackrel{(1.27)}{\lesssim} (\|\nabla \sigma\|_{H^2}^2 + \|\nabla \mathbf{u}\|^2) \|\Lambda^{-s} \sigma\|, \end{aligned}$$

and

$$\begin{aligned} I_9 &\lesssim \|\Lambda^{-s} [(\mathbf{u} - \mathbf{u}', \nabla) \mathbf{u} - h_1(\sigma) \nabla \sigma] \| \|\Lambda^{-s} \mathbf{u}\| \\ &\stackrel{(1.14)}{\lesssim} \|(\mathbf{u} - \mathbf{u}', \nabla) \mathbf{u} - h_1(\sigma) \nabla \sigma\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}}} \|\Lambda^{-s} \mathbf{u}\| \\ &\lesssim \left(\|\mathbf{u} - \mathbf{u}'\|_{L^{\frac{3}{s}}} \|\nabla \mathbf{u}\| + \|\sigma\|_{L^{\frac{3}{s}}} \|\nabla \sigma\| \right) \|\Lambda^{-s} \mathbf{u}\| \\ &\stackrel{(1.27), (2.32)}{\lesssim} (\|\nabla \mathbf{u}\|_{H^2}^2 + \|\sigma\|_{H^2}^2) \|\Lambda^{-s} \mathbf{u}\| \\ &\stackrel{(1.27)}{\lesssim} (\|\nabla \mathbf{u}\|_{H^2}^2 + \|\nabla \sigma\|_{H^1}^2) \|\Lambda^{-s} \mathbf{u}\|. \end{aligned}$$

Also, we have

$$\begin{aligned} I_{10} &\lesssim (\|\Lambda^{-s} [h_2(\sigma) \operatorname{div} [\nu(\phi) \mathbf{D}(\mathbf{u})]] \| + \|\Lambda^{-s} [h_2(\sigma) \nabla [(\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}]] \|) \|\Lambda^{-s} \mathbf{u}\| \\ &\stackrel{(1.14)}{\lesssim} \left(\|h_2(\sigma) \operatorname{div} [\nu(\phi) \mathbf{D}(\mathbf{u})]\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}}} + \|h_2(\sigma) \nabla [(\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}]\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}}} \right) \|\Lambda^{-s} \mathbf{u}\| \\ &\lesssim \|h_2(\sigma)\|_{L^{\frac{3}{s}}} (\|\operatorname{div} [\nu(\phi) \mathbf{D}(\mathbf{u})]\| + \|\nabla [(\nu(\phi) + \lambda(\phi)) \operatorname{div} \mathbf{u}]\|) \|\Lambda^{-s} \mathbf{u}\| \\ &\stackrel{(1.11), (2.1)}{\lesssim} \|\sigma\|_{L^{\frac{3}{s}}} \|\nabla \mathbf{u}\|_{H^1} \|\Lambda^{-s} \mathbf{u}\| \\ &\stackrel{(2.32), (1.27)}{\lesssim} (\|\nabla \sigma\|_{H^1}^2 + \|\nabla \mathbf{u}\|_{H^1}^2) \|\Lambda^{-s} \mathbf{u}\| \end{aligned}$$

and

$$\begin{aligned}
I_{11} &\lesssim \|\Lambda^{-s}(\nabla\phi\Delta\phi)\| \|\Lambda^{-s}\mathbf{u}\| \\
&\stackrel{(1.14)}{\lesssim} \|\nabla\phi\Delta\phi\|_{L^{\frac{1}{\frac{1}{2}+\frac{s}{3}}}} \|\Lambda^{-s}\mathbf{u}\| \lesssim \|\nabla\phi\|_{L^{\frac{3}{s}}} \|\nabla^2\phi\| \|\Lambda^{-s}\mathbf{u}\| \\
&\stackrel{(2.32)}{\lesssim} \|\nabla\phi\|_{H^2}^2 \|\Lambda^{-s}\mathbf{u}\|.
\end{aligned}$$

The estimate on I_{12} is more subtle. To this end, we rewrite it as

$$\begin{aligned}
I_{12} &= \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s}\nabla \left[(3\phi^2 - 1) \left(\frac{1}{\rho} - \frac{1}{\bar{\rho}} \right) \Delta\phi \right] \Lambda^{-s}\nabla\phi d\mathbf{x}}_{I_{12}^1} \\
&\quad + \underbrace{\frac{3}{\bar{\rho}} \int_{\mathbb{T}^3} \Lambda^{-s}\nabla \left((\phi^2 - \bar{\phi}^2) \Delta\phi \right) \Lambda^{-s}\nabla\phi d\mathbf{x}}_{I_{12}^2} \\
&\quad - \underbrace{\frac{3\bar{\phi}^2 - 1}{\bar{\rho}} \int_{\mathbb{T}^3} |\Lambda^{-s}\nabla\phi|^2 d\mathbf{x}}_{I_{12}^3} + 6 \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s}\nabla \left(\frac{\phi}{\rho} (\nabla\phi)^2 \right) \Lambda^{-s}\nabla\phi d\mathbf{x}}_{I_{12}^4},
\end{aligned} \tag{2.33}$$

where $\bar{\phi}^2 = \int_{\mathbb{T}^3} \phi^2 d\mathbf{x}$. Then, we have

$$\begin{aligned}
I_{12}^1 &\lesssim \left\| \Lambda^{-s}\nabla \left[(3\phi^2 - 1) \left(\frac{1}{\rho} - \frac{1}{\bar{\rho}} \right) \Delta\phi \right] \right\| \|\Lambda^{-s}\nabla\phi\| \\
&\stackrel{(1.14)}{\lesssim} \left\| \nabla \left[(3\phi^2 - 1) \left(\frac{1}{\rho} - \frac{1}{\bar{\rho}} \right) \Delta\phi \right] \right\|_{L^{\frac{1}{\frac{1}{2}+\frac{s}{3}}}} \|\Lambda^{-s}\nabla\phi\| \\
&\lesssim \left(\|\sigma\|_{L^{\frac{3}{s}}} \|\nabla^3\phi\| + \|\nabla\phi\|_{L^{\frac{3}{s}}} \|\nabla^2\phi\| + \|\nabla\sigma\|_{L^{\frac{3}{s}}} \|\nabla^2\phi\| \right) \|\Lambda^{-s}\nabla\phi\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} (\|\nabla\sigma\|_{H^2}^2 + \|\nabla\phi\|_{H^2}^2) \|\Lambda^{-s}\nabla\phi\|,
\end{aligned}$$

and

$$\begin{aligned}
I_{12}^2 &\lesssim \left\| \Lambda^{-s}\nabla \left((\phi^2 - \bar{\phi}^2) \Delta\phi \right) \right\| \|\Lambda^{-s}\nabla\phi\| \\
&\stackrel{(1.14)}{\lesssim} \left\| \nabla \left((\phi^2 - \bar{\phi}^2) \Delta\phi \right) \right\|_{L^{\frac{1}{\frac{1}{2}+\frac{s}{3}}}} \|\Lambda^{-s}\nabla\phi\| \\
&\lesssim \left(\|\phi^2 - \bar{\phi}^2\|_{L^{\frac{3}{s}}} \|\nabla^3\phi\| + \|\nabla\phi\|_{L^{\frac{3}{s}}} \|\nabla^2\phi\| \right) \|\Lambda^{-s}\nabla\phi\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} \|\nabla\phi\|_{H^2}^2 \|\Lambda^{-s}\nabla\phi\|,
\end{aligned}$$

moreover,

$$\begin{aligned}
&I_{12}^3 + \frac{2}{\bar{\rho}} \int_{\mathbb{T}^3} |\Lambda^{-s}\nabla\phi|^2 d\mathbf{x} \\
&\leq \frac{3}{\bar{\rho}} (1 - \bar{\phi}^2) \int_{\mathbb{T}^3} |\Lambda^{-s}\nabla\phi|^2 d\mathbf{x} \\
&\lesssim \eta_1 \int_{\mathbb{T}^3} |\Lambda^{-s}\nabla\phi|^2 d\mathbf{x},
\end{aligned}$$

where we used

$$1 - \bar{\phi}^2 = \int_{\mathbb{T}^3} (1 - \phi^2) d\mathbf{x} \lesssim \|1 - \phi^2\| \lesssim \eta_1, \tag{2.34}$$

and

$$\begin{aligned}
I_{12}^4 &\lesssim \left\| \Lambda^{-s} \nabla \left(\frac{\phi}{\rho} (\nabla \phi)^2 \right) \right\| \left\| \Lambda^{-s} \nabla \phi \right\| \stackrel{(1.14)}{\lesssim} \left\| \nabla \left(\frac{\phi}{\rho} (\nabla \phi)^2 \right) \right\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}} \left\| \Lambda^{-s} \nabla \phi \right\| \\
&\lesssim \left\| \nabla \phi \right\|_{L^{\frac{3}{s}}} (\left\| \nabla \phi \right\| + \left\| \nabla^2 \phi \right\|) \left\| \Lambda^{-s} \nabla \phi \right\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} \left\| \nabla \phi \right\|_{H^2}^2 \left\| \Lambda^{-s} \nabla \phi \right\|.
\end{aligned}$$

Therefore, we obtain from (2.33) that

$$I_{12} \lesssim (\left\| \nabla \sigma \right\|_{H^2}^2 + \left\| \nabla \phi \right\|_{H^2}^2) \left\| \Lambda^{-s} \nabla \phi \right\|.$$

Last, for I_{13} , noticing that $\int_{\mathbb{T}^3} \mathbf{u}' \cdot \Lambda^{-s} \nabla \nabla \phi \Lambda^{-s} \nabla \phi d\mathbf{x} = 0$, we have

$$\begin{aligned}
I_{13} &\lesssim \left\| \Lambda^{-s} \nabla [(\mathbf{u}' - \mathbf{u}) \cdot \nabla \phi + h_3(\sigma) \Delta^2 \phi] \right\| \left\| \Lambda^{-s} \nabla \phi \right\| \\
&+ \left\| \Lambda^{-s} \nabla \left(\frac{2}{\rho^3} \Delta \nabla \phi \cdot \nabla \sigma + \frac{\Delta \phi}{\rho} \operatorname{div} \left(\frac{\nabla \sigma}{\rho^2} \right) \right) \right\| \left\| \Lambda^{-s} \nabla \phi \right\| \\
&\stackrel{(1.14)}{\lesssim} \left\| \nabla [(\mathbf{u}' - \mathbf{u}) \cdot \nabla \phi + h_3(\sigma) \Delta^2 \phi] \right\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}} \left\| \Lambda^{-s} \nabla \phi \right\| \\
&+ \left\| \nabla \left(\frac{2}{\rho^3} \Delta \nabla \phi \cdot \nabla \sigma + \frac{\Delta \phi}{\rho} \operatorname{div} \left(\frac{\nabla \sigma}{\rho^2} \right) \right) \right\|_{L^{\frac{1}{\frac{1}{2} + \frac{s}{3}}}} \left\| \Lambda^{-s} \nabla \phi \right\| \\
&\lesssim \left[(\left\| \mathbf{u} - \mathbf{u}' \right\|_{L^{\frac{3}{s}}} + \left\| \nabla \sigma \right\|_{L^{\frac{3}{s}}}) \left\| \nabla^2 \phi \right\| + \left\| \nabla \mathbf{u} \right\|_{L^{\frac{3}{s}}} \left\| \nabla \phi \right\| + \left\| \sigma \right\|_{L^{\frac{3}{s}}} \left\| \nabla^3 \phi \right\| \right] \left\| \Lambda^{-s} \nabla \phi \right\| \\
&+ \left[\left\| \nabla \sigma \right\|_{L^{\frac{3}{s}}} \left\| \Delta \nabla \phi \right\|_{H^1} + \left\| \nabla \phi \right\|_{W^{1, \frac{3}{s}}} \left\| \Delta \sigma \right\|_{H^1} \right] \left\| \Lambda^{-s} \nabla \phi \right\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} (\left\| \nabla \mathbf{u} \right\|_{H^2}^2 + \left\| \nabla \sigma \right\|_{H^2}^2 + \left\| \nabla \phi \right\|_{H^3}^2) \left\| \Lambda^{-s} \nabla \phi \right\|.
\end{aligned}$$

By the same lines as in I_{11} , we obtain

$$I_{14} \lesssim \left\| \nabla \phi \right\|_{H^2}^2 \left\| \Lambda^{-s} \mathbf{u} \right\|.$$

Substituting the estimates on $I_i (i = 6, \dots, 14)$ into (2.31) yields (2.30). The proof of Lemma 2.5 is completed. \blacksquare

Lemma 2.6 *Under the assumption (2.1), it holds that*

$$\frac{d}{dt} \left\| \Lambda^{-s} (\phi^2 - 1) \right\|^2 \lesssim (\left\| \nabla(\sigma, \mathbf{u}) \right\|_{H^2}^2 + \left\| \nabla \phi \right\|_{H^3}^2) \left\| \Lambda^{-s} (\phi^2 - 1) \right\|, \quad \text{for } s \in (0, \frac{3}{2}). \quad (2.35)$$

Proof: We rewrite (2.5) as

$$\begin{aligned}
&(\phi^2 - 1)_t + \mathbf{u} \cdot \nabla (\phi^2 - 1) - \frac{2}{\rho} \Delta (\phi^2 - 1) + \frac{1}{\rho^2} \Delta^2 (\phi^2 - 1) \\
&= \frac{12\phi^3}{\rho} |\nabla \phi|^2 + \frac{6\phi}{\rho} (\phi^2 - 1) \Delta \phi + 4\phi \left(\frac{1}{\rho} - \frac{1}{\bar{\rho}} \right) \Delta \phi - \frac{2}{\rho} |\nabla \phi|^2 \\
&+ \frac{2}{\bar{\rho}^2} [|\nabla \phi|^2 + 2\nabla \phi \cdot \nabla \Delta \phi + \Delta(|\nabla \phi|^2)] \\
&+ 2\phi \left(\frac{1}{\rho} - \frac{1}{\bar{\rho}} \right) \Delta \left(\frac{\Delta \phi}{\rho} \right) + \frac{2\phi}{\bar{\rho}} \Delta \left[\left(\frac{1}{\rho} - \frac{1}{\bar{\rho}} \right) \Delta \phi \right].
\end{aligned} \tag{2.36}$$

Then, applying Λ^{-s} to (2.36) and multiplying the resulting identity by $\Lambda^{-s}(\phi^2 - 1)$, we deduce that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\Lambda^{-s}(\phi^2 - 1)\|^2 + \frac{2}{\bar{\rho}} \|\Lambda^{-s} \nabla(\phi^2 - 1)\|^2 + \frac{1}{\bar{\rho}^2} \|\Lambda^{-s} \Delta(\phi^2 - 1)\|^2 \\
&= \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s} \left[\left(\frac{12\phi^3}{\rho} - \frac{2}{\bar{\rho}} \right) |\nabla\phi|^2 + 4\phi h_2(\sigma) \Delta\phi \right] \Lambda^{-s}(\phi^2 - 1) \, d\mathbf{x}}_{I_{15}} \\
&+ \underbrace{\frac{2}{\bar{\rho}^2} \int_{\mathbb{T}^3} \Lambda^{-s} [|\nabla\phi|^2 + 2\nabla\phi \cdot \nabla\Delta\phi + \Delta(|\nabla\phi|^2)] \Lambda^{-s}(\phi^2 - 1) \, d\mathbf{x}}_{I_{16}} \\
&- 2 \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s} \left[\phi h_2(\sigma) \Delta \left(\frac{\Delta\phi}{\rho} \right) + \frac{\phi}{\bar{\rho}} \Delta(h_2(\sigma) \Delta\phi) \right] \Lambda^{-s}(\phi^2 - 1) \, d\mathbf{x}}_{I_{17}} \\
&- \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s} [\mathbf{u} \cdot \nabla(\phi^2 - 1)] \Lambda^{-s}(\phi^2 - 1) \, d\mathbf{x}}_{I_{18}} \\
&+ 6 \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s} \left[\frac{\phi}{\rho} (\phi^2 - 1) \Delta\phi \right] \Lambda^{-s}(\phi^2 - 1) \, d\mathbf{x}}_{I_{19}}, \tag{2.37}
\end{aligned}$$

where $h_2(\sigma) = \frac{1}{\bar{\rho}} - \frac{1}{\rho}$ (see (2.7)). In order to estimate the nonlinear terms in the right-hand side of (2.37), we shall use the estimate (1.14). If $s \in (0, \frac{3}{2})$, then $\frac{1}{2} + \frac{s}{3} < 1$ and we have

$$\begin{aligned}
I_{15} &\lesssim \left\| \Lambda^{-s} \left[\left(\frac{12\phi^3}{\rho} - \frac{2}{\bar{\rho}} \right) |\nabla\phi|^2 + 4\phi h_2(\sigma) \Delta\phi \right] \right\| \|\Lambda^{-s}(\phi^2 - 1)\| \\
&\stackrel{(1.14), (2.10), (2.8)}{\lesssim} \left(\|\nabla\phi\|_{L^{\frac{3}{s}}} \|\nabla\phi\| + \|\Delta\phi\| \|\sigma\|_{L^{\frac{3}{s}}} \right) \|\Lambda^{-s}(\phi^2 - 1)\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} (\|\nabla\phi\|_{H^3}^2 + \|\nabla\sigma\|_{H^2}^2) \|\Lambda^{-s}(\phi^2 - 1)\|,
\end{aligned}$$

$$\begin{aligned}
I_{16} &\lesssim \|\Lambda^{-s} [|\nabla\phi|^2 + 2\nabla\phi \cdot \nabla\Delta\phi + \Delta(|\nabla\phi|^2)]\| \|\Lambda^{-s}(\phi^2 - 1)\| \\
&\stackrel{(1.14)}{\lesssim} \left[\|\nabla\phi\|_{L^{\frac{3}{s}}} (\|\nabla\phi\| + \|\nabla\Delta\phi\|) + \|\Delta\phi\|_{L^{\frac{3}{s}}} \|\Delta\phi\| \right] \|\Lambda^{-s}(\phi^2 - 1)\| \\
&\stackrel{(1.27)}{\lesssim} \|\nabla\phi\|_{H^3}^2 \|\Lambda^{-s}(\phi^2 - 1)\|,
\end{aligned}$$

and

$$\begin{aligned}
I_{17} &\lesssim \left\| \Lambda^{-s} \left[\phi h_2(\sigma) \Delta \left(\frac{\Delta\phi}{\rho} \right) + \frac{\phi}{\bar{\rho}} \Delta(h_2(\sigma) \Delta\phi) \right] \right\| \|\Lambda^{-s}(\phi^2 - 1)\| \\
&\stackrel{(1.14), (2.10), (2.8)}{\lesssim} \|\sigma\|_{L^{\frac{3}{s}}} \left(\left\| \Delta \left(\frac{\Delta\phi}{\rho} \right) \right\| + \|\Delta^2\phi\| \right) \|\Lambda^{-s}(\phi^2 - 1)\| \\
&+ \left(\|\nabla\sigma\|_{L^{\frac{3}{s}}} \|\nabla\Delta\phi\| + \|\Delta\sigma\| \|\Delta\phi\|_{L^{\frac{3}{s}}} \right) \|\Lambda^{-s}(\phi^2 - 1)\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} (\|\nabla\phi\|_{H^3}^2 + \|\nabla\sigma\|_{H^2}^2) \|\Lambda^{-s}(\phi^2 - 1)\|.
\end{aligned}$$

For I_{17} , noticing that

$$\int_{\mathbb{T}^3} \mathbf{u}' \cdot \Lambda^{-s} \nabla(\phi^2 - 1) \Lambda^{-s}(\phi^2 - 1) \, d\mathbf{x} = 0,$$

where $\mathbf{u}' = \int_{\mathbb{T}^3} \mathbf{u} d\mathbf{x}$, we have

$$\begin{aligned}
I_{18} &= \int_{\mathbb{T}^3} \Lambda^{-s} [(\mathbf{u} - \mathbf{u}') \cdot \nabla (\phi^2 - 1)] \Lambda^{-s} (\phi^2 - 1) d\mathbf{x} \\
&\lesssim \|\Lambda^{-s} [(\mathbf{u} - \mathbf{u}') \cdot \nabla (\phi^2 - 1)]\| \|\Lambda^{-s} (\phi^2 - 1)\| \\
&\stackrel{(1.14)}{\lesssim} \|\mathbf{u} - \mathbf{u}'\|_{L^{\frac{3}{s}}} \|\nabla (\phi^2 - 1)\| \|\Lambda^{-s} (\phi^2 - 1)\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} (\|\nabla \mathbf{u}\|_{H^2}^2 + \|\nabla \phi\|^2) \|\Lambda^{-s} (\phi^2 - 1)\|.
\end{aligned}$$

Last, noticing that

$$\phi \Delta \phi = \frac{1}{2} \Delta (\phi^2 - 1) + |\nabla \phi|^2,$$

we rewrite I_{19} as

$$\begin{aligned}
I_{19} &= \underbrace{-\frac{(\overline{\phi^2} - 1)}{2\bar{\rho}} \int_{\mathbb{T}^3} |\Lambda^{-s} \nabla (\phi^2 - 1)|^2 d\mathbf{x}}_{I_{19}^1} \\
&\quad + \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s} \left[\left(\frac{(\phi^2 - 1)}{2\rho} - \frac{(\overline{\phi^2} - 1)}{2\bar{\rho}} \right) \Delta (\phi^2 - 1) \right] \Lambda^{-s} (\phi^2 - 1)}_{I_{19}^2} \\
&\quad + \underbrace{\int_{\mathbb{T}^3} \Lambda^{-s} \left[\frac{(\phi^2 - 1)}{\rho} |\nabla \phi|^2 \right] \Lambda^{-s} (\phi^2 - 1) d\mathbf{x}}_{I_{19}^3}, \tag{2.38}
\end{aligned}$$

where $\overline{\phi^2} = \int_{\mathbb{T}^3} \phi^2 d\mathbf{x}$. Then we have

$$\begin{aligned}
I_{19}^1 &\stackrel{(2.34)}{\lesssim} \eta_1 \|\Lambda^{-s} \nabla (\phi^2 - 1)\|^2, \\
I_{19}^2 &\lesssim \left\| \Lambda^{-s} \left[\left(\frac{(\phi^2 - 1)}{2\rho} - \frac{(\overline{\phi^2} - 1)}{2\bar{\rho}} \right) \Delta (\phi^2 - 1) \right] \right\| \|\Lambda^{-s} (\phi^2 - 1)\| \\
&\stackrel{(1.14)}{\lesssim} \left(\|\rho - \bar{\rho}\|_{L^{\frac{3}{s}}} + \|\phi^2 - \overline{\phi^2}\|_{L^{\frac{3}{s}}} \right) \|\Delta (\phi^2 - 1)\| \|\Lambda^{-s} (\phi^2 - 1)\| \\
&\stackrel{(2.32), (1.27)}{\lesssim} (\|\nabla \sigma\|_{H^2}^2 + \|\nabla \phi\|_{H^2}^2) \|\Lambda^{-s} (\phi^2 - 1)\|,
\end{aligned}$$

and

$$\begin{aligned}
I_{19}^3 &\lesssim \left\| \Lambda^{-s} \left[\frac{(\phi^2 - 1)}{\rho} |\nabla \phi|^2 \right] \right\| \|\Lambda^{-s} (\phi^2 - 1)\| \\
&\stackrel{(1.14), (2.10), (2.8)}{\lesssim} \|\nabla \phi\|_{L^{\frac{3}{s}}} \|\nabla \phi\| \|\Lambda^{-s} (\phi^2 - 1)\| \\
&\stackrel{(2.32)}{\lesssim} \|\nabla \phi\|_{H^2}^2 \|\Lambda^{-s} (\phi^2 - 1)\|.
\end{aligned}$$

Therefore, we obtain from (2.38) that

$$I_{19} \lesssim \eta_1 \|\Lambda^{-s} \nabla (\phi^2 - 1)\|^2 + (\|\nabla \sigma\|_{H^2}^2 + \|\nabla \phi\|_{H^2}^2) \|\Lambda^{-s} (\phi^2 - 1)\|.$$

Substituting the estimates on $I_i (i = 15, \dots, 19)$ into (2.37) yields (2.35). The proof of Lemma 2.6 is completed. \blacksquare

3 The proof of Theorem 1.1

In this section, we shall combine all the energy estimates that we have derived in the previous section and the Sobolev interpolation to prove Theorem 1.1.

3.1 The existence of global solutions

Proposition 3.1 (a priori estimate). *There exists a positive constant and $M_1 > 0$, if $(\rho, \mathbf{u}, \phi) \in X_{M_1}([0, T^*])$, then*

$$\begin{aligned} & \sup_{t \in [0, T]} \left\{ \|(\rho - \bar{\rho}, \mathbf{u})(t)\|_{H^3}^2 + \|\nabla \phi(t)\|_{H^2}^2 + \|\phi^2(t) - 1\|^2 \right\} \\ & + \int_0^T \|\nabla \rho\|_{H^2}^2 d\tau + \int_0^T \|\nabla \mathbf{u}\|_{H^3} d\tau + \int_0^T \|\nabla \phi\|_{H^4}^2 d\tau \\ & \lesssim \|\rho_0 - \bar{\rho}\|_{H^3}^2 + \|\mathbf{u}_0\|_{H^3}^2 + \|\nabla \phi_0\|_{H^2}^2 + \|\phi_0^2 - 1\|^2. \end{aligned} \quad (3.1)$$

Proof: We first close the energy estimates at each l -th level in our weak sense to prove (1.18). Let $0 \leq l \leq m$ with $1 \leq m \leq 3$. Summing up the estimate (2.17) of Lemma 2.2 for from $k = l$ to $m - 1$, we obtain

$$\begin{aligned} & \frac{d}{dt} \sum_{l \leq k \leq m-1} \|\nabla^{k+1} \phi\|^2 + \frac{1}{4\bar{\rho}^2} \sum_{l+1 \leq k \leq m} \|\nabla^{k+3} \phi\|^2 \\ & \lesssim \eta_1 \sum_{l+1 \leq k \leq m} \left(\|\nabla^k \sigma\|^2 + \|\nabla^{k+1} \mathbf{u}\|^2 \right). \end{aligned} \quad (3.2)$$

Also, summing up the estimate (2.21) of Lemma 2.3 for from $k = l$ to m , we obtain

$$\begin{aligned} & \frac{d}{dt} \sum_{l \leq k \leq m} \left(\|\nabla^k \mathbf{u}\|^2 + \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\nabla^k \sigma\|^2 \right) + \frac{\nu_0}{\bar{\rho}} \sum_{l \leq k \leq m} \|\nabla^{k+1} \mathbf{u}\|^2 \\ & \lesssim \eta_1 \sum_{l \leq k \leq m} \left(\|\nabla^k \sigma\|^2 + \|\nabla^{k+2} \phi\|^2 \right). \end{aligned} \quad (3.3)$$

Last, summing up the estimate (2.27) of Lemma 2.4 for from $k = l$ to $m - 1$, we obtain

$$\begin{aligned} & \frac{d}{dt} \sum_{l \leq k \leq m-1} \int_{\mathbb{T}^3} \nabla^k \mathbf{u} \cdot \nabla^{k+1} \sigma d\mathbf{x} + \frac{p'(\bar{\rho})}{2\bar{\rho}} \sum_{l+1 \leq k \leq m} \|\nabla^k \sigma\|^2 \\ & \lesssim \eta_1 \sum_{l+1 \leq k \leq m} \left(\|\nabla^{k+1} \mathbf{u}\|^2 + \|\nabla^{k+3} \phi\|^2 \right) + \sum_{l \leq k \leq m-1} \|\nabla^{k+1} \mathbf{u}\|^2. \end{aligned} \quad (3.4)$$

Let $\epsilon \in (0, 1]$ be suitably small. Then, summing (2.9), (3.2), (3.3) and $\epsilon \times (3.4)$, and choosing $\eta_1 > 0$ to be small, we obtain

$$\frac{d}{dt} \mathcal{E}_l^m(t) + \frac{1}{2} \Lambda_l^m(t) \leq 0 \quad (3.5)$$

for any $0 \leq l < m \leq 3$, where

$$\begin{aligned} \mathcal{E}_l^m(t) &:= \sum_{l \leq k \leq m} \left(\|\nabla^k \mathbf{u}\|^2 + \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\nabla^k \sigma\|^2 \right) + \epsilon \sum_{l \leq k \leq m-1} \int_{\mathbb{T}^3} \nabla^k \mathbf{u} \cdot \nabla^{k+1} \sigma \, dx \\ &\quad + \sum_{l \leq k \leq m-1} \|\nabla^{k+1} \phi\|^2 + \int_{\mathbb{T}^3} \left(\frac{\rho}{2} \mathbf{u}^2 + G(\rho) + \frac{1}{2} |\nabla \phi|^2 + \frac{\rho}{4} (\phi^2 - 1)^2 \right) dx, \\ \Lambda_l^m(t) &:= \epsilon \frac{p'(\bar{\rho})}{2\bar{\rho}} \sum_{l+1 \leq k \leq m} \|\nabla^k \sigma\|^2 + \frac{1}{4\bar{\rho}^2} \sum_{l+1 \leq k \leq m} \|\nabla^{k+2} \phi\|^2 \\ &\quad + \frac{\nu_0}{\bar{\rho}} \sum_{l \leq k \leq m} \|\nabla^{k+1} \mathbf{u}\|^2 - \epsilon C_0 \sum_{l \leq k \leq m-1} \|\nabla^{k+1} \mathbf{u}\|^2. \end{aligned} \quad (3.6)$$

Notice that since $\epsilon \in (0, 1]$ be suitably small, we obtain from (3.6) that

$$\begin{aligned} \mathcal{E}_l^m(t) &\simeq \|\nabla^l(\sigma, \mathbf{u})(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \phi(t)\|_{H^{m-1-l}}^2 + \|\phi^2 - 1\|^2, \\ \Lambda_l^m(t) &\simeq \|\nabla^l \sigma(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \mathbf{u}(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \phi(t)\|_{H^{m+1-l}}^2 \end{aligned} \quad (3.7)$$

uniformly for all $t \geq 0$. Now taking $l = 0$ and $m = 3$ in (3.5), and then using (3.7) and (3.5), we get

$$\begin{aligned} \|\sigma(t)\|_{H^3}^2 + \|\mathbf{u}(t)\|_{H^3}^2 + \|\nabla \phi(t)\|_{H^{3-1}}^2 + \|\phi^2 - 1\|^2 &\lesssim \mathcal{E}_0^3(t) \\ &\lesssim \mathcal{E}_0^3(0) \lesssim \|\rho_0 - \bar{\rho}\|_{H^3}^2 + \|\mathbf{u}_0\|_{H^3}^2 + \|\nabla \phi_0\|_{H^{3-1}}^2 + \|\phi_0^2 - 1\|^2. \end{aligned} \quad (3.8)$$

Using (1.22) and (3.8), by a standard continuity argument, we can close the a priori estimate (3.1). This in turn allows us to take $l = 0$ and $m = 3$ in (3.5), and then integrate it directly in time to obtain (3.1). \blacksquare

By using the Proposition 2.1, a solution for $t \in [0, T^*]$ can be obtained which satisfies (2.8) in $\mathbb{T}^3 \times [0, T^*]$, T^* only depends on the initial data of the periodic boundary value problem (1.15). From and Proposition 3.1, one can start again from T^* , by the same way, one can find a solution in $[0, 2T^*]$, and so on. Thus the existence and uniqueness of the global solution for system (2.8) is obtained.

3.2 Algebraic decay

We prove (1.19)-(1.21). Define

$$\mathcal{E}_{-s} := \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\Lambda^{-s} \sigma\|^2 + \|\Lambda^{-s} \mathbf{u}\|^2 + \|\Lambda^{-s} \nabla \phi\|^2 + \|\Lambda^{-s} (\phi^2 - 1)\|^2$$

for $s \in (0, \frac{3}{2})$. Then, integrating in time (2.30) and (2.35), and using (1.18), we get

$$\begin{aligned} \mathcal{E}_{-s}(t) &\lesssim \mathcal{E}_{-s}(0) + \int_0^t (\|\nabla(\sigma, \mathbf{u})\|_{H^2}^2 + \|\nabla \phi\|_{H^3}^2) \sqrt{\mathcal{E}_{-s}(\tau)} \, d\tau \\ &\lesssim 1 + \sup_{0 \leq \tau \leq t} \sqrt{\mathcal{E}_{-s}(\tau)}, \end{aligned}$$

which implies (1.19). If $l = 0, \dots, 2$, we may use (1.13) to have

$$\|\nabla^{l+1} f\| \gtrsim \|\Lambda^{-s} f\|^{-\frac{1}{l+s}} \|\nabla^l f\|^{1+\frac{1}{l+s}}. \quad (3.9)$$

By (3.9) and (1.19), we get

$$\|\nabla \phi\| \stackrel{(2.10)}{\gtrsim} \|\nabla(\phi^2 - 1)\| \gtrsim \|\phi^2 - 1\|^{1+\frac{1}{s}}, \quad (3.10)$$

and

$$\|\nabla^{l+1} \sigma\|^2 + \|\nabla^{l+1} \mathbf{u}\|^2 + \|\nabla^{l+1} \nabla \phi\|^2 \gtrsim \left(\|\nabla^l \sigma\|^2 + \|\nabla^l \mathbf{u}\|^2 + \|\nabla^l \nabla \phi\|^2 \right)^{1+\frac{1}{l+s}},$$

which implies

$$\begin{aligned} & \|\nabla^{l+1} \sigma(t)\|_{H^{m-l-1}}^2 + \|\nabla^{l+1} \mathbf{u}(t)\|_{H^{m-l}}^2 + \|\nabla^{l+2} \phi(t)\|_{H^{m-l-1}}^2 \\ & \gtrsim \left(\|\nabla^l \sigma(t)\|_{H^{m-l-1}}^2 + \|\nabla^l \mathbf{u}(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1} \phi(t)\|_{H^{m-l-1}}^2 \right)^{1+\frac{1}{l+s}}. \end{aligned} \quad (3.11)$$

Also, using $\|\sigma(t)\|_{H^3} \lesssim 1$ due to (1.18), we have

$$\|\nabla^m \sigma\| \gtrsim \|\nabla^m \sigma\|^{1+\frac{1}{l+s}}. \quad (3.12)$$

By (3.10)-(3.12) and (1.29), we obtain from (3.7) that

$$\Lambda_0^1(t) \gtrsim (\mathcal{E}_0^1(t))^{1+\frac{1}{s}}. \quad (3.13)$$

Using (3.13), we deduce from (3.5) with $l = 0$ and $m = 1$ that

$$\frac{d}{dt} \mathcal{E}_0^1(t) + C_0 (\mathcal{E}_0^1(t))^{1+\frac{1}{s}} \leq 0.$$

Solving this inequality directly gives (see (1.28))

$$\mathcal{E}_0^1(t) \lesssim (1+t)^{-s},$$

which implies (1.20), that is,

$$\|(\sigma, \mathbf{u})(t)\|_{H^1}^2 + \|\nabla \phi(t)\|^2 + \|\phi^2(t) - 1\|^2 \lesssim (1+t)^{-s},$$

due to (3.7)₁. We prove (1.21). For the small $\epsilon \in (0, 1]$, summing (3.2), (3.3) and $\epsilon \times (3.4)$, and choosing $\eta_1 > 0$ to be small, we have

$$\frac{d}{dt} \mathcal{F}_l^m(t) + G_l^m(t) \leq 0, \quad \text{for any } 0 \leq l < m \leq 3, \quad (3.14)$$

where

$$\begin{aligned} \mathcal{F}_l^m(t) &:= \sum_{l \leq k \leq m} \left(\|\nabla^k \mathbf{u}\|^2 + \frac{p'(\bar{\rho})}{\bar{\rho}^2} \|\nabla^k \sigma\|^2 \right) \\ &+ \epsilon \sum_{l \leq k \leq m-1} \int_{\mathbb{T}^3} \nabla^k \mathbf{u} \cdot \nabla^{k+1} \sigma \, dx + \sum_{l \leq k \leq m-1} \|\nabla^{k+1} \phi\|^2, \\ G_l^m(t) &:= \epsilon \frac{p'(\bar{\rho})}{2\bar{\rho}} \sum_{l+1 \leq k \leq m} \|\nabla^k \sigma\|^2 + \frac{1}{4\bar{\rho}^2} \sum_{l+1 \leq k \leq m} \|\nabla^{k+2} \phi\|^2 \\ &+ \left(\frac{\nu_0}{\bar{\rho}} - \epsilon C_0 \right) \sum_{l \leq k \leq m} \|\nabla^{k+1} \mathbf{u}\|^2. \end{aligned} \quad (3.15)$$

Notice that since $\epsilon \in (0, 1]$ be suitably small, we obtain from (3.15) that

$$\begin{aligned}\mathcal{F}_l^m(t) &\simeq \|\nabla^l(\sigma, \mathbf{u})(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1}\phi(t)\|_{H^{m-1-l}}^2, \\ G_l^m(t) &\simeq \|\nabla^l\sigma(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1}\mathbf{u}(t)\|_{H^{m-l}}^2 + \|\nabla^{l+1}\phi(t)\|_{H^{m+1-l}}^2,\end{aligned}\tag{3.16}$$

uniformly for all $t \geq 0$. By (3.11) and (3.12), we obtain from (3.16) that

$$G_l^m(t) \gtrsim (\mathcal{F}_l^m(t))^{1+\frac{1}{l+s}}.\tag{3.17}$$

Using (3.17), we deduce from (3.14) with $l = 2$ and $m = 3$ that

$$\frac{d}{dt}\mathcal{F}_2^3(t) + C_0 (\mathcal{F}_2^3(t))^{1+\frac{1}{3-1+s}} \leq 0.$$

Solving this inequality directly gives (see (1.28))

$$\mathcal{F}_2^3(t) \lesssim (1+t)^{-(2+s)},$$

which implies

$$\|\nabla^2(\sigma, \mathbf{u})(t)\|_{H^1}^2 + \|\nabla^3\phi(t)\|^2 \lesssim (2+t)^{-(l+s)},\tag{3.18}$$

due to (3.16)₁. Using (1.29), we obtain (1.21) from (3.18), and so far, combining with Subsection 3.1, Theorem 1.1 is obtained.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- [1] H. Abels, E. Feireisl, On a diffuse interface model for a two-phase flow of compressible viscous fluids, *Indiana Univ. Math J.*, 57 (2) (2008) 569-578.
- [2] J. W. Cahn, J. E. Hilliard, Free energy of a nonuniform system, I. Interfacial free energy, *J. Chem. Phys.*, 28 (1958) 258-267.
- [3] M. Chen, X. Guo, Global large solutions for a coupled compressible Navier-Stokes/Allen-Cahn system with initial vacuum, *Nonlinear Analysis: Real World Applications*, 37 (2017) 350-373.
- [4] Y. Chen, Q. He, M. Mei, X. Shi, Asymptotic stability of solutions for 1-D compressible Navier-Stokes-Cahn-Hilliard system, *J. Math. Anal. Appl.*, 467 (2018) 185-206.
- [5] Z.Chen, C. Li, An extended discrete Hardy-Littlewood-Sobolev inequality, *Discrete and Continuous Dynamical Systems*, 34(5) (2014) 1951-1959.
- [6] S. Chen, H. Wen, C. Zhu, Golbal existence of weak solution to compressible Navier-Stokes/Allen-Cahn system in three dimensions. *J. Math. Anal. Appl.*, 477 (2019) 1265-1295.

- [7] S. Ding, Y. Li, W. Luo, Global solutions for a coupled compressible Navier-Stokes/Allen-Cahn system in 1D, *J. Math. Fluid Mech.*, 15 (2013) 335-360.
- [8] S. Ding, Y. Li, Y. Tang, Strong solutions to 1D compressible Navier-Stokes/Allen-Cahn system with free boundary, *Math. Methods Appl. Sci.*, 42 (2019) 4780-4794.
- [9] E. Feireisl, H. Petzeltová, E. Rocca, G. Schimperna, Analysis of a phase-field model for two-phase compressible fluids, *math. Models Meth. Appl. Sci.*, 20 (7) (2010) 1129-1160.
- [10] H. Freistühler, M. Kotschote, Phase-field and Korteweg-type models for the time-dependent flow of compressible two-phase fluids, *Arch. Rational Mech. Anal.*, 224 (2017) 1-20.
- [11] H. Freistühler, M. Kotschote, The Blesgen and Lowengrub-Truskinovsky Descriptions of Two-Phase Compressible Fluid Flow: Interstitial Working and a Reduction to Korteweg Theory, arXiv:1711.10041
- [12] J. Gao, Q. Tao, Z.-A. Yao, Long-time behavior of solution for the compressible nematic liquid crystal flows in \mathbb{R}^3 , *J. Differential Equations*, 261 (2016) 2334-2383.
- [13] Y. Guo, Y.J. Wang, Decay of dissipative equations and negative Sobolev spaces, preprint, *Commun. P.D.E.*, 37 (2012) 2165-2208.
- [14] Q. He, C. Liu, X. Shi, Numerical study of phase transition in van der Waals fluid, *Discret. Contin.Dyn. Syst. Ser. B*, 23 (2018) 4519-4540.
- [15] M. Heida, J. Málek, K.R. Rajagopal, On the development and generalizations of Cahn-Hilliard equations within a thermodynamic framework, *Z. Angew. Math. Phys.*, 63 (1) (2012) 145-169.
- [16] G. Huang, C. Li, X. Yin, Existence of the maximizing pair for the discrete Hardy-Littlewood-Sobolev inequality, *Discret. Contin.Dyn. Syst.*, 35(3) (2015) 935-942.
- [17] S. Klainerman, A. Majda, Singular limits of quasilinear hyperbolic systems with large parameters and the incompressible limit of compressible fluids, *Comm. Pure Appl. Math.*, 34 (1981) 481-524.
- [18] M. Kotschote, Strong solutions of the Navier-Stokes equations for a compressible fluid of Allen-Cahn type, *Arch. Rational Mech. Anal.*, 206 (2012) 489-514.
- [19] M. Kotschote, Mixing rules and the Navier-Stokes-Cahn-Hilliard equations for compressible heat-conductive fluids, *Bull. Braz. Math. Soc. (3.S.)*, 47 (2) (2016) 457-471.
- [20] M. Kotschote, R. Zacher, Strong solutions in the dynamical theory of compressible fluid mixtures, *Math. Models Meth. Appl. Sciences*, 25 (7) (2015) 1217-1256.
- [21] C. Liu, J. Shen, A phase field model for the mixture of two incompressible fluids and its approximation by a Fourier-spectral method. *Phys. D*, 179 (3-4) (2003) 211-228.
- [22] T. Luo, H. Yin, Stability of the rarefaction wave for a coupled compressible Navier-Stokes/Allen-Cahn system, *Math. Methods Appl. Sci.*, 41 (12) (2018) 4724-4736.

- [23] L. Nirenberg, On elliptic partial differential equations, *Ann. Scuola Norm. Sup. Pisa*, 13 (1959) 115-162.
- [24] E. M. Stein, *Singular integrals and differentiability properties of functions*. Princeton University Press, Princeton, 1970.
- [25] Z. Tan, R. Zhang, Optimal decay rates of the compressible fluid models of Korteweg type, *Z. Angew. Math. Phys.*, 65 (2014) 279-300.
- [26] Y. Wang, Decay of the Navier-Stokes-Poisson equations, *J. Differential Equations*, 253 (2012) 273-297.
- [27] H. Yin, C. Zhu, Asymptotic stability of superposition of stationary solutions and rarefaction waves for 1D Navier-Stokes/Allen-Cahn system, *J. Differential Equations*, 266 (2019) 7291-7326.
- [28] J. Yin, On the Cahn-Hilliard equation with nonlinear principal part, *Journal of Partial Differential Equations*, 7 (1) (1994) 77-96.