# CONCENTRATION PHENOMENA FOR A FRACTIONAL RELATIVISTIC SCHRÖDINGER EQUATION WITH CRITICAL GROWTH

VINCENZO AMBROSIO

ABSTRACT. In this paper, we are concerned with the following fractional relativistic Schrödinger equation with critical growth:

$$\left\{\begin{array}{ll} (-\Delta+m^2)^s u+V(\varepsilon\,x)u=f(u)+u^{2^*_s-1} \text{ in } \mathbb{R}^N,\\ u\in H^s(\mathbb{R}^N), \quad u>0 \text{ in } \mathbb{R}^N, \end{array}\right.$$

where  $\varepsilon > 0$  is a small parameter,  $s \in (0, 1)$ , m > 0, N > 2s,  $2_s^* = \frac{2N}{N-2s}$  is the fractional critical exponent,  $(-\Delta + m^2)^s$  is the fractional relativistic Schrödinger operator,  $V : \mathbb{R}^N \to \mathbb{R}$  is a continuous potential, and  $f : \mathbb{R} \to \mathbb{R}$  is a superlinear continuous nonlinearity with subcritical growth at infinity. Under suitable assumptions on the potential V, we construct a family of positive solutions  $u_{\varepsilon} \in H^s(\mathbb{R}^N)$ , with exponential decay, which concentrates around a local minimum of V as  $\varepsilon \to 0$ .

### 1. INTRODUCTION

In this paper, we continue the study started in [6] concerning the concentration phenomena for a class of fractional relativistic Schrödinger equations. More precisely, we focus on the following nonlinear fractional elliptic equation with critical growth:

$$\begin{cases} (-\Delta + m^2)^s u + V(\varepsilon x)u = f(u) + u^{2^*_s - 1} \text{ in } \mathbb{R}^N, \\ u \in H^s(\mathbb{R}^N), \quad u > 0 \text{ in } \mathbb{R}^N, \end{cases}$$
(1.1)

where  $\varepsilon > 0$  is a small parameter, m > 0,  $s \in (0, 1)$ , N > 2s,  $2_s^* := \frac{2N}{N-2s}$  is the fractional critical exponent, and  $V : \mathbb{R}^N \to \mathbb{R}$  and  $f : \mathbb{R} \to \mathbb{R}$  are continuous functions. The operator  $(-\Delta + m^2)^s$  is defined in Fourier space as multiplication by the symbol  $(|k|^2 + m^2)^s$  (see [28,29]), i.e., for each function  $u : \mathbb{R}^N \to \mathbb{R}$  that belongs to the Schwartz space  $\mathcal{S}(\mathbb{R}^N)$  of rapidly decreasing functions, we have

$$\mathcal{F}((-\Delta + m^2)^s u)(k) := (|k|^2 + m^2)^s \mathcal{F}u(k), \quad k \in \mathbb{R}^N,$$

where we denoted by

$$\mathcal{F}u(k) := (2\pi)^{-\frac{N}{2}} \int_{\mathbb{R}^N} e^{-\imath k \cdot x} u(x) \, dx, \quad k \in \mathbb{R}^N$$

the Fourier transform of u. We also recall the following alternative representation of  $(-\Delta + m^2)^s$  in terms of singular integrals (see [24, 29]):

$$(-\Delta + m^2)^s u(x) := m^{2s} u(x) + C(N,s) m^{\frac{N+2s}{2}} P.V. \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x - y|^{\frac{N+2s}{2}}} K_{\frac{N+2s}{2}}(m|x - y|) \, dy, \quad x \in \mathbb{R}^N, \quad (1.2)$$

where P.V. indicates the Cauchy principal value,  $K_{\nu}$  is the modified Bessel function of the third kind of index  $\nu$  (see [9,23]), and

$$C(N,s) := 2^{-\frac{N+2s}{2}+1} \pi^{-\frac{N}{2}} 2^{2s} \frac{s(1-s)}{\Gamma(2-s)}$$

When  $s = \frac{1}{2}$ , the operator  $\sqrt{-\Delta + m^2}$  was considered in [43,44] for spectral problems and has a clear meaning in relativistic quantum mechanics. Indeed, the energy for the motion of a free relativistic particle of mass m and momentum p is given by:

$$\mathcal{E} := \sqrt{p^2 c^2 + m^2 c^4},$$

<sup>2010</sup> Mathematics Subject Classification. 35R11, 35J10, 35J20, 35J60, 35B09, 35B33.

Key words and phrases. fractional relativistic Schrödinger operator; critical exponent; extension method; variational methods.

where c is the speed of the light. With the usual quantization rule  $p \mapsto -i\hbar \nabla$ , where  $\hbar$  is Planck's constant, we obtain the so-called relativistic Hamiltonian operator:

$$\mathcal{H} := \sqrt{-\hbar^2 c^2 \Delta + m^2 c^4} - mc^2.$$

The point of the subtraction of the constant  $mc^2$  is to make sure that the spectrum of the operator  $\mathcal{H}$  is  $[0, \infty)$ , and this explains the terminology of relativistic Schrödinger operators for the operators of the form  $\mathcal{H} + V(x)$ , where V(x) is a potential (see [16]). Equations involving  $\mathcal{H}$  arise in the study of time-dependent Schrödinger equations of the type:

$$i\hbar \frac{\partial \Phi}{\partial t} = \mathcal{H}\Phi - f(x, |\Phi|^2)\Phi, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^N,$$

where  $\Phi : \mathbb{R} \times \mathbb{R}^N \to \mathbb{C}$  is a wave function and  $f : \mathbb{R}^N \times [0, \infty) \to \mathbb{R}$  is a nonlinear function, which describe the dynamics of systems consisting of identical spin-0 bosons whose motions are relativistic, for instance, boson stars. Physical models related to  $\mathcal{H}$  have been widely analyzed over the past 30 years, and there exists an important literature on the spectral properties of relativistic Hamiltonians; most of it has been strongly influenced by the works of Lieb on the stability of relativistic matter (see [22,27,30,31] and references therein). On the other hand, from a probabilistic point of view,  $m^{2s} - (-\Delta + m^2)^s$  is the infinitesimal generator of a Lévy process  $X_t^{2s,m}$  called 2s-stable relativistic process having the following characteristic function:

$$E^{0}e^{\imath k\cdot X_{t}^{2s,m}} = e^{-t[(|k|^{2}+m^{2})^{s}-m^{2s}]}, \quad k \in \mathbb{R}^{N};$$

(see, for example, [16, 36]). For a more detailed discussion on  $(-\Delta + m^2)^s$ , we refer the interested reader to [8].

When  $m \to 0$ ,  $(-\Delta + m^2)^s$  reduces to the well-known fractional Laplacian  $(-\Delta)^s$  defined via Fourier transform by:

$$\mathcal{F}((-\Delta)^s u)(k) := |k|^{2s} \mathcal{F}u(k), \quad k \in \mathbb{R}^N$$

or through singular integrals by:

$$(-\Delta)^{s}u(x) := C_{N,s} P.V. \int_{\mathbb{R}^{N}} \frac{u(x) - u(y)}{|x - y|^{N + 2s}} \, dy, \quad x \in \mathbb{R}^{N}, \quad C_{N,s} := \pi^{-\frac{N}{2}} 2^{2s} \frac{\Gamma(\frac{N + 2s}{2})}{\Gamma(2 - s)} s(1 - s). \tag{1.3}$$

This operator has gained tremendous popularity during the last two decades thanks to its applications in different fields, such as, among others, phase transition phenomena, crystal dislocation, population dynamics, anomalous diffusion, flame propagation, chemical reactions of liquids, conservation laws, quasi-geostrophic flows, and water waves. Moreover, the fractional Laplacian is the infinitesimal generator of a (rotationally) symmetric 2s-stable Lévy process. For a very nice introduction to  $(-\Delta)^s$  and its applications, consult [13,20]. Note that the most striking difference between the operators  $(-\Delta)^s$  and  $(-\Delta + m^2)^s$  is that the first one is homogeneous in scaling, whereas the second one is inhomogeneous as should be clear from the presence of the Bessel function  $K_{\nu}$  in (1.2).

We emphasize that in these years, several authors dealt with the existence and multiplicity of solutions for the following fractional Schrödinger equation:

$$\begin{cases} \varepsilon^{2s}(-\Delta)^{s}u + V(x)u = f(u) + \gamma |u|^{2^{s}_{s}-2}u \text{ in } \mathbb{R}^{N}, \\ u \in H^{s}(\mathbb{R}^{N}), \quad u > 0 \text{ in } \mathbb{R}^{N}, \end{cases}$$
(1.4)

where  $\varepsilon > 0, \gamma \in \{0, 1\}, V : \mathbb{R}^N \to \mathbb{R}$  and  $f : \mathbb{R} \to \mathbb{R}$  satisfy suitable conditions (see, for instance, [7] and references therein).

As  $s \to 1$ , equation (1.4) boils down to the classical nonlinear Schrödinger equation of the form:

$$\begin{cases} -\varepsilon^2 \Delta u + V(x)u = g(u) \text{ in } \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N), \quad u > 0 \text{ in } \mathbb{R}^N. \end{cases}$$
(1.5)

Since we cannot review the huge bibliography on this topic, we refer to [2,3,19,25,26,35,42] for some results on the existence, multiplicity, and concentration of positive solutions to (1.5) for small  $\varepsilon > 0$ . We recall that a positive solution  $u_{\varepsilon}$  of (1.5) is said to concentrate at  $x_0 \in \mathbb{R}^N$  as  $\varepsilon \to 0$  if

$$\forall \delta > 0, \quad \exists \, \varepsilon_0 > 0, R > 0: \quad u_{\varepsilon}(x) \leq \delta, \quad \forall |x - x_0| \geq \varepsilon \, R, \, \varepsilon < \varepsilon_0 \, .$$

The interest in studying semiclassical solutions of (1.5), i.e., solutions of (1.5) with small  $\varepsilon > 0$ , is justified by the well-known fact that the transition from quantum mechanics to classical mechanics can be described

by letting  $\varepsilon \to 0$ . A typical feature of semiclassical solutions is that they tend to concentrate as  $\varepsilon \to 0$  around critical points of the potential V.

On the other hand, several existence and multiplicity results for fractional equations driven by  $(-\Delta + m^2)^s$ , with m > 0, have been established in [5, 6, 14, 17, 37, 39]. In particular, in [6], the author investigated (1.1) without the presence of the critical term  $u^{2_s^*-1}$  and obtained the existence of solutions concentrating in a given set of local minima of V as  $\varepsilon \to 0$ . He also related the number of positive solutions to the topology of the set where V attains its minimum value. We point out that, in all the aforementioned articles, only equations with subcritical nonlinearities are considered.

Motivated by the previous facts, in this paper we examine the existence of concentrating solutions to (1.1), by assuming that the potential  $V: \mathbb{R}^N \to \mathbb{R}$  is a continuous function fulfilling the following conditions: by assuming that the potential V is  $V_1 = \inf_{x \in \mathbb{R}^N} V(x)$ ,  $(V_1)$  there exists  $V_1 \in (0, m^{2s})$  such that  $-V_1 := \inf_{x \in \mathbb{R}^N} V(x)$ ,

$$(V_2)$$
 there exists a bounded open set  $\Lambda \subset \mathbb{R}^N$  such that

$$-V_0 := \inf_{x \in \Lambda} V(x) < \min_{x \in \partial \Lambda} V(x),$$

with  $V_0 > 0$ , and  $0 \in M := \{x \in \Lambda : V(x) = -V_0\},\$ 

and that the nonlinearity  $f : \mathbb{R} \to \mathbb{R}$  is a continuous, f(t) = 0 for  $t \leq 0$ , and satisfies the following hypotheses:  $(f_1) \lim_{t \to 0} \frac{f(t)}{t} = 0,$ 

 $(f_2)$  there exist  $p,q\in(2,2^*_s)$  and  $\lambda>0$  such that

$$f(t) \ge \lambda t^{p-1}$$
, for all  $t \ge 0$ , and  $\lim_{t \to \infty} \frac{f(t)}{t^{q-1}} = 0$ ,

where  $\lambda > 0$  is such that

- $\lambda > 0$  if either  $N \ge 4s$ , or 2s < N < 4s and  $2_s^* 2 ,$
- $\lambda > 0$  is sufficiently large if 2s < N < 4s and 2 ,
- $(f_3)$  there exists  $\theta \in (2,q)$  such that  $0 < \theta F(t) \le tf(t)$  for all t > 0, where  $F(t) := \int_0^t f(\tau) d\tau$ ,

 $(f_4)$  the function  $t \mapsto \frac{f(t)}{t}$  is increasing in  $(0, \infty)$ . The main result of this paper can be stated as follows:

**Theorem 1.1.** Assume that  $(V_1)$ - $(V_2)$  and  $(f_1)$ - $(f_4)$  hold. Then, for every small  $\varepsilon > 0$ , there exists a solution  $u_{\varepsilon}$  to (1.1) such that  $u_{\varepsilon}$  has a maximum point  $x_{\varepsilon}$  satisfying

$$\lim_{\varepsilon \to 0} \operatorname{dist}(\varepsilon \, x_{\varepsilon}, M) = 0$$

and for which

$$0 < u_{\varepsilon}(x) \le C_1 e^{-C_2 |x - x_{\varepsilon}|}, \quad \text{for all } x \in \mathbb{R}^N.$$

for suitable constants  $C_1, C_2 > 0$ . Moreover, for each sequence  $(\varepsilon_n)$  with  $\varepsilon_n \to 0$ , there exists a subsequence, still denoted by itself, such that there exist a point  $x_0 \in M$  with  $\varepsilon_n x_{\varepsilon_n} \to x_0$  and a positive ground state solution  $u \in H^{s}(\mathbb{R}^{N})$  of the limiting problem:

$$(-\Delta + m^2)^s u - V_0 u = f(u) + u^{2^*_s - 1}$$
 in  $\mathbb{R}^N$ 

for which we have

$$u_{\varepsilon_n}(x) = u(x - x_{\varepsilon_n}) + \mathcal{R}_n(x),$$

where  $\lim_{n \to \infty} \|\mathcal{R}_n\|_{H^s(\mathbb{R}^N)} = 0.$ 

The proof of Theorem 1.1 relies on appropriate variational techniques. Since the operator  $(-\Delta + m^2)^s$  is nonlocal, we transform (1.1) into a degenerate elliptic equation in a half-space with a nonlinear Neumann boundary condition via a variant of the extension method [15] (see [17, 24, 41]). Then, we adapt the penalization approach in [19], the so-called *local mountain pass*, by building a convenient modification of the energy functional associated with the extended problem in such a way that the corresponding modified energy functional  $J_{\varepsilon}$  satisfies the hypotheses of the mountain pass theorem [4], and then we prove that, for  $\varepsilon > 0$ sufficiently small, the trace of the associated mountain pass solution is, indeed, a solution to the original equation with the stated properties. The modification of the functional corresponds to a penalization outside  $\Lambda$ , and this is why no other global assumptions are required. With respect to [6], it is more difficult to obtain compactness for  $J_{\varepsilon}$  due to the presence of the critical exponent. To overcome this obstacle, we first estimate

from above the mountain pass level  $c_{\varepsilon}$  of  $J_{\varepsilon}$ , by constructing a suitable cut-off function. Roughly speaking, we choose a function, appropriately rescaled, of the type  $v_{\epsilon}(x, y) = \vartheta(my)\phi(x, y)w_{\epsilon}(x, y)$ , with  $\epsilon > 0$ , where  $\vartheta$  is expressed via the Bessel function  $K_s$ ,  $\phi$  is a smooth cut-off function, and  $w_{\epsilon}$  is the s-harmonic extension of the extremal function  $u_{\epsilon}$  for the fractional Sobolev inequality (see [18]), in such a way that the control of the quadratic term

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v_{\epsilon}|^2 + m^2 v_{\epsilon}^2) \, dx \, dy - m^{2s} \int_{\mathbb{R}^N} v_{\epsilon}^2(x,0) \, dx$$

is, in some sense, reduced to the control of the term:

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla(\phi w_{\epsilon})|^2 \, dx dy,$$

and thus, we are able to verify that  $c_{\varepsilon} < c_*$ , where the threshold value  $c_*$  depends on the best constant  $S_*$  for the critical Sobolev trace inequality (see [11]), and the constants  $V_1$ ,  $m^{2s}$ , and  $\theta$  (see Lemma 3.2). In view of this bound and by establishing a concentration-compactness principle in the spirit of Lions [32, 33], we show that the modified energy functional satisfies the Palais-Smale condition in the range  $(0, c_*)$  (see Lemmas 3.1 and 3.5). Finally, we prove that, for  $\varepsilon > 0$  small enough, the solution of the auxiliary problem is, indeed, solution of the original one by combining a Moser iteration scheme [34], a comparison argument, and some crucial properties of the Bessel kernel [9, 40] (see Lemmas 5.3 and 5.4). As far as we know, this is the first time that the penalization method is used to study the concentration phenomena for a fractional relativistic Schrödinger equation with critical growth.

The structure of the paper is the following. In section 2, we define some function spaces. In section 3, we focus on the modified problem. In section 4, we deal with the autonomous critical problems related to the extended modified problem. Section 5 is devoted to the proof of Theorem 1.1. In section 6, we discuss a multiplicity result for (1.1).

**Notations:** We denote the upper half-space in  $\mathbb{R}^N$  by  $\mathbb{R}^{N+1}_+ := \{(x,y) \in \mathbb{R}^{N+1} : y > 0\}$ . For  $(x,y) \in \mathbb{R}^{N+1}_+$ , we set  $|(x,y)| := \sqrt{|x|^2 + y^2}$ . The letters c, C, C', and  $C_i$  will be repeatedly used to denote various positive constants whose exact values are irrelevant and can change from line to line. For  $x \in \mathbb{R}^N$  and R > 0, we will denote by  $B_R(x)$  the ball in  $\mathbb{R}^N$  centered at  $x \in \mathbb{R}^N$  with radius r > 0. When x = 0, we set  $B_R := B_R(0)$ . For  $(x_0, y_0) \in \mathbb{R}^{N+1}_+$  and R > 0, we put  $B_R^+(x_0, y_0) := \{(x, y) \in \mathbb{R}^{N+1}_+ : |(x, y) - (x_0, y_0)| < R\}$  and  $B_R^+ := B_R^+(0, 0)$ . Let  $p \in [1, \infty]$  and  $A \subset \mathbb{R}^N$  be a measurable set. The notation  $A^c := \mathbb{R}^N \setminus A$  stands for the complement of A in  $\mathbb{R}^N$ . We will use  $|u|_{L^p(A)}$  for the  $L^p$ -norm of  $u : \mathbb{R}^N \to \mathbb{R}$ . If  $A = \mathbb{R}^N$ , we simply write  $|u|_p$  instead of  $|u|_{L^p(\mathbb{R}^N)}$ . With  $||v||_{L^p(\mathbb{R}^{N+1})}$  we denote the norm of  $v \in L^p(\mathbb{R}^{N+1}_+)$ . For a generic real-valued function w, we set  $w^+ := \max\{w, 0\}$  and  $w^- := \min\{w, 0\}$ .

# 2. Function spaces

Let  $H^s(\mathbb{R}^N)$  be the fractional Sobolev space defined as the completion of  $C_c^{\infty}(\mathbb{R}^N)$  with respect to the norm

$$|u|_{H^{s}(\mathbb{R}^{N})} := \left( \int_{\mathbb{R}^{N}} (|k|^{2} + m^{2})^{s} |\mathcal{F}u(k)|^{2} dk \right)^{\frac{1}{2}}.$$

Then,  $H^s(\mathbb{R}^N)$  is continuously embedded in  $L^p(\mathbb{R}^N)$  for all  $p \in [2, 2_s^*]$  and compactly in  $L^p_{loc}(\mathbb{R}^N)$  for all  $p \in [1, 2_s^*)$ ; see [1, 7, 8, 20, 29]. We denote by  $X^s(\mathbb{R}^{N+1}_+)$  the completion of  $C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  with respect to the norm:

$$\|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})} := \left(\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s}(|\nabla v|^{2} + m^{2}v^{2}) \, dx dy\right)^{\frac{1}{2}}.$$

By [24, Lemma 3.1] (see also [21, Proposition 3.1.1]), it follows that  $X^s(\mathbb{R}^{N+1}_+)$  is continuously embedded in  $L^{2\gamma}(\mathbb{R}^{N+1}_+, y^{1-2s})$ , i.e.,

$$\|v\|_{L^{2\gamma}(\mathbb{R}^{N+1}_+, y^{1-2s})} \le \hat{S} \|v\|_{X^s(\mathbb{R}^{N+1}_+)}, \quad \text{for all } v \in X^s(\mathbb{R}^{N+1}_+),$$
(2.1)

for some  $\hat{S} > 0$ , where  $\gamma := 1 + \frac{2}{N-2s}$ , and  $L^r(\mathbb{R}^{N+1}_+, y^{1-2s})$  is the weighted Lebesgue space, with  $r \in (1, \infty)$ , endowed with the norm:

$$\|v\|_{L^{r}(\mathbb{R}^{N+1}_{+},y^{1-2s})} := \left(\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |v|^{r} \, dx dy\right)^{\frac{1}{r}}.$$

In light of [21, Lemma 3.1.2], we also know that  $X^{s}(\mathbb{R}^{N+1}_{+})$  is compactly embedded in  $L^{2}(B^{+}_{R}, y^{1-2s})$  for all R > 0. By [24, Proposition 5], there exists a (unique) linear trace operator  $\text{Tr} : X^{s}(\mathbb{R}^{N+1}_{+}) \to H^{s}(\mathbb{R}^{N})$  such that

$$\sqrt{\sigma_s} |\operatorname{Tr}(v)|_{H^s(\mathbb{R}^N)} \le ||v||_{X^s(\mathbb{R}^{N+1}_+)}, \quad \text{for all } v \in X^s(\mathbb{R}^{N+1}_+).$$
(2.2)

where  $\sigma_s := 2^{1-2s}\Gamma(1-s)/\Gamma(s)$ ; see [11,41]. In order to lighten the notation, we will denote  $\operatorname{Tr}(v)$  by  $v(\cdot, 0)$ . We note that (2.2) yields

$$\sigma_s m^{2s} \int_{\mathbb{R}^N} v^2(x,0) \, dx \le \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v|^2 + m^2 v^2) \, dx dy, \tag{2.3}$$

for all  $v \in X^{s}(\mathbb{R}^{N+1}_{+})$ , which can be also written as:

$$\sigma_s \int_{\mathbb{R}^N} v^2(x,0) \, dx \le m^{-2s} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla v|^2 \, dx \, dy + m^{2-2s} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v^2 \, dx \, dy. \tag{2.4}$$

From the previous facts, we deduce the following fundamental embeddings.

**Theorem 2.1.**  $\operatorname{Tr}(X^s(\mathbb{R}^{N+1}_+))$  is continuously embedded in  $L^r(\mathbb{R}^N)$  for all  $r \in [2, 2^*_s]$  and compactly embedded in  $L^r_{loc}(\mathbb{R}^N)$  for all  $r \in [1, 2^*_s]$ .

To deal with (1.1) via variational methods, we use a variant of the extension method [15] given in [17,24,41]. More precisely, for each  $u \in H^s(\mathbb{R}^N)$ , there exists a unique function  $U \in X^s(\mathbb{R}^{N+1}_+)$  solving the problem

$$\left\{ \begin{array}{ll} -\operatorname{div}(y^{1-2s}\nabla U)+m^2y^{1-2s}U=0 & \text{ in } \mathbb{R}^{N+1}_+, \\ U=u & \text{ on } \partial \mathbb{R}^{N+1}_+ \end{array} \right.$$

The function U is called the extension of u and fulfills the following properties: (E1)

$$\frac{\partial U}{\partial \nu^{1-2s}} := -\lim_{y \to 0} y^{1-2s} \frac{\partial U}{\partial y}(x,y) = \sigma_s(-\Delta + m^2)^s u(x) \quad \text{in } H^{-s}(\mathbb{R}^N),$$

where  $H^{-s}(\mathbb{R}^N)$  denotes the dual of  $H^s(\mathbb{R}^N)$ .

(E2)  $\sqrt{\sigma_s}|u|_{H^s(\mathbb{R}^N)} = \|U\|_{X^s(\mathbb{R}^{N+1}_+)} \le \|V\|_{X^s(\mathbb{R}^{N+1}_+)}$  for all  $V \in X^s(\mathbb{R}^{N+1}_+)$  such that  $V(\cdot, 0) = u$ , (E3) if  $u \in \mathcal{S}(\mathbb{R}^N)$ , then  $U \in C^\infty(\mathbb{R}^{N+1}_+) \cap C(\mathbb{R}^{N+1}_+)$ , and it can be expressed as:

$$U(x,y) = \int_{\mathbb{R}^N} P_{s,m}(x-z,y)u(z) \, dz,$$

with

$$P_{s,m}(x,y) := c'_{N,s} y^{2s} m^{\frac{N+2s}{2}} |(x,y)|^{-\frac{N+2s}{2}} K_{\frac{N+2s}{2}}(m|(x,y)|)$$

and

$$c'_{N,s} := p_{N,s} \frac{2^{\frac{N+2s}{2}-1}}{\Gamma(\frac{N+2s}{2})}$$

where  $p_{N,s} := \pi^{-\frac{N}{2}} \frac{\Gamma(\frac{N+2s}{2})}{\Gamma(s)}$  is the constant for the (normalized) Poisson kernel with m = 0 (see [41]).

**Remark 2.1.** We recall (see [24]) that  $P_{s,m}$  is the Fourier transform of  $k \mapsto \vartheta(\sqrt{|k|^2 + m^2})$  and that

$$\int_{\mathbb{R}^N} P_{s,m}(x,y) \, dx = \vartheta(my), \tag{2.5}$$

where

$$\vartheta(r) := \frac{2}{\Gamma(s)} \left(\frac{r}{2}\right)^s K_s(r) \tag{2.6}$$

belongs to  $H^1(\mathbb{R}_+, y^{1-2s})$  and solves the following ordinary differential equation:

$$\begin{cases} \vartheta'' + \frac{1-2s}{y}\vartheta' - \vartheta = 0 \quad in \mathbb{R}_+,\\ \vartheta(0) = 1, \quad \lim_{y \to \infty} \vartheta(y) = 0. \end{cases}$$
(2.7)

We also have

$$\int_0^\infty y^{1-2s} (|\vartheta'(y)|^2 + |\vartheta(y)|^2) \, dy = -\lim_{y \to 0} y^{1-2s} \vartheta'(y) = \kappa_s.$$
(2.8)

Consequently, (1.1) can be realized in a local manner through the following nonlinear boundary value problem:

$$\begin{cases} -\operatorname{div}(y^{1-2s}\nabla v) + m^2 y^{1-2s} v = 0 & \text{in } \mathbb{R}^{N+1}_+, \\ \frac{\partial v}{\partial \nu^{1-2s}} = \sigma_s [-V_{\varepsilon} v(\cdot, 0) + f(v(\cdot, 0)) + (v^+(\cdot, 0))^{2^*_s - 1}] & \text{on } \mathbb{R}^N, \end{cases}$$
(2.9)

where  $V_{\varepsilon}(x) := V(\varepsilon x)$ . For simplicity of notation, we will drop the constant  $\sigma_s$  from the second equation in (2.9). In order to examine (2.9), for  $\varepsilon > 0$ , we introduce the space

$$X_{\varepsilon} := \left\{ v \in X^{s}(\mathbb{R}^{N+1}_{+}) : \int_{\mathbb{R}^{N}} V_{\varepsilon}(x) v^{2}(x,0) \, dx < \infty \right\}$$

equipped with the norm:

$$\|v\|_{\varepsilon} := \left(\|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} + \int_{\mathbb{R}^{N}} V_{\varepsilon}(x)v^{2}(x,0) \, dx\right)^{\frac{1}{2}}$$

Clearly,  $X_{\varepsilon} \subset X^{s}(\mathbb{R}^{N+1}_{+})$ , and using (2.3) and  $(V_{1})$ , we see that

$$\|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} \leq \left(\frac{m^{2s}}{m^{2s} - V_{1}}\right) \|v\|_{\varepsilon}^{2}, \quad \text{for all } v \in X_{\varepsilon}.$$
(2.10)

Furthermore,  $X_{\varepsilon}$  is a Hilbert space endowed with the inner product:

$$\langle v, w \rangle_{\varepsilon} := \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla v \cdot \nabla w + m^2 v w) \, dx dy + \int_{\mathbb{R}^N} V_{\varepsilon}(x) v(x, 0) w(x, 0) \, dx, \quad \text{ for all } v, w \in X_{\varepsilon}.$$

Henceforth, with  $X_{\varepsilon}^*$ , we will denote the dual space of  $X_{\varepsilon}$ .

# 3. PENALIZATION ARGUMENT

To study (2.9), we adapt the penalization approach in [19] (see also [6]). Fix  $\kappa > \max\{\frac{V_1}{m^{2s}-V_1}, \frac{\theta}{\theta-2}\} > 1$ and a > 0 such that  $f(a) + a^{2^*_s - 1} = \frac{V_1}{\kappa}a$ . Define

$$\tilde{f}(t) := \begin{cases} f(t) + (t^+)^{2_s^* - 1}, & \text{for } t < a \\ \frac{V_1}{\kappa} t, & \text{for } t \ge a \end{cases}$$

and

$$g(x,t) := \chi_{\Lambda}(x)f(t) + (1 - \chi_{\Lambda}(x))\tilde{f}(t), \quad \text{for } (x,t) \in \mathbb{R}^N \times \mathbb{R},$$

where  $\chi_{\Lambda}$  denotes the characteristic function of  $\Lambda$ . Set  $G(x,t) := \int_0^t g(x,\tau) d\tau$ . By assumptions  $(f_1)$ - $(f_4)$ , it is easy to prove that g is a Carathéodory function satisfying the following properties:

- (g1)  $\lim_{t\to 0} \frac{g(x,t)}{t} = 0$  uniformly in  $x \in \mathbb{R}^N$ , (g2)  $g(x,t) \leq f(t) + t^{2^*_s 1}$  for all  $x \in \mathbb{R}^N$ , t > 0, (g3) (i)  $0 < \theta G(x,t) \leq tg(x,t)$  for all  $x \in \Lambda$  and t > 0, or,  $x \in \Lambda^c$  and  $0 < t \leq a$ , (ii)  $0 \leq 2G(x,t) \leq tg(x,t) \leq \frac{V_1}{\kappa}t^2$  for all  $x \in \Lambda^c$  and t > 0, (g4) for each  $x \in \Lambda$ , the function  $t \mapsto \frac{g(x,t)}{t}$  is increasing in  $(0,\infty)$ , and for each  $x \in \Lambda^c$ , the function  $t \mapsto \frac{g(x,t)}{t}$ is increasing in (0, a).

Let us introduce the following auxiliary problem:

$$\begin{cases} -\operatorname{div}(y^{1-2s}\nabla v) + m^2 y^{1-2s} v = 0 & \text{in } \mathbb{R}^{N+1}_+, \\ \frac{\partial v}{\partial \nu^{1-2s}} = -V_{\varepsilon} v(\cdot, 0) + g_{\varepsilon}(\cdot, v(\cdot, 0)) & \text{on } \mathbb{R}^N, \end{cases}$$
(3.1)

where  $g_{\varepsilon}(x,t) := g(\varepsilon x,t)$ . It is clear that if  $v_{\varepsilon}$  is a positive solution of (3.1) satisfying  $v_{\varepsilon}(x,0) < a$  for all  $x \in \Lambda_{\varepsilon}^{c}$ , where  $\Lambda_{\varepsilon} := \{x \in \mathbb{R}^{N} : \varepsilon x \in \Lambda\}$ , then  $v_{\varepsilon}$  is a positive solution of (2.9). The energy functional associated with (3.1) is defined by:

$$J_{\varepsilon}(v) := \frac{1}{2} \|v\|_{\varepsilon}^2 - \int_{\mathbb{R}^N} G_{\varepsilon}(x, v(x, 0)) \, dx, \quad \text{ for all } v \in X_{\varepsilon}.$$

It is standard to check that  $J_{\varepsilon} \in C^1(X_{\varepsilon}, \mathbb{R})$  and that its differential is given by

$$\langle J'_{\varepsilon}(v), w \rangle = \langle v, w \rangle_{\varepsilon} - \int_{\mathbb{R}^N} g_{\varepsilon}(x, v(x, 0)) w(x, 0) \, dx \quad \text{ for all } v, w \in X_{\varepsilon}.$$

Hence, the critical points of  $J_{\varepsilon}$  correspond to the weak solutions of (3.1). To seek these critical points, we will apply suitable variational arguments. First, we show that  $J_{\varepsilon}$  possesses the geometric assumptions of the mountain pass theorem [4].

**Lemma 3.1.** The functional  $J_{\varepsilon}$  satisfies the following properties:

(i)  $J_{\varepsilon}(0) = 0$ ,

- (ii) there exist  $\alpha, \rho > 0$  such that  $J_{\varepsilon}(v) \ge \alpha$  for all  $v \in X_{\varepsilon}$  such that  $||v||_{\varepsilon} = \rho$ ,
- (iii) there exists  $\bar{v} \in X_{\varepsilon}$  such that  $\|\bar{v}\|_{\varepsilon} > \rho$  and  $J_{\varepsilon}(\bar{v}) < 0$ .

*Proof.* Condition (i) is obvious. By  $(f_1)$ ,  $(f_2)$ ,  $(g_1)$ , and  $(g_2)$ , we see that for all  $\eta > 0$ , there exists  $C_{\eta} > 0$  such that

$$|g_{\varepsilon}(x,t)| \le \eta |t| + C_{\eta} |t|^{2_{s}^{*}-1}, \quad \text{for } (x,t) \in \mathbb{R}^{N} \times \mathbb{R},$$
(3.2)

and

$$G_{\varepsilon}(x,t)| \leq \frac{\eta}{2}|t|^2 + \frac{C_{\eta}}{2_s^*}|t|^{2_s^*}, \quad \text{for } (x,t) \in \mathbb{R}^N \times \mathbb{R}.$$

$$(3.3)$$

Pick  $\eta \in (0, m^{2s} - V_1)$ . By (3.3), (2.3), (2.10), and using Theorem 2.1, we have

$$J_{\varepsilon}(v) \geq \frac{1}{2} \|v\|_{\varepsilon}^{2} - \frac{\eta}{2} |v(\cdot, 0)|_{2}^{2} - \frac{C_{\eta}}{2_{s}^{*}} |v(\cdot, 0)|_{2_{s}^{*}}^{2_{s}^{*}}$$

$$= \frac{1}{2} \|v\|_{\varepsilon}^{2} - \frac{\eta}{2m^{2s}} m^{2s} |v(\cdot, 0)|_{2}^{2} - \frac{C_{\eta}}{2_{s}^{*}} |v(\cdot, 0)|_{2_{s}^{*}}^{2_{s}^{*}}$$

$$\geq \frac{1}{2} \|v\|_{\varepsilon}^{2} - \frac{\eta}{2m^{2s}} \|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} - C_{\eta}C \|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2_{s}^{*}}$$

$$\geq \left(\frac{1}{2} - \frac{\eta}{2(m^{2s} - V_{1})}\right) \|v\|_{\varepsilon}^{2} - C_{\eta}C' \|v\|_{\varepsilon}^{2_{s}^{*}},$$

from which we deduce that (*ii*) is fulfilled. Finally, take  $v_0 \in C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  such that  $v_0 \ge 0$ ,  $v_0 \ne 0$ , and  $\sup (v_0(\cdot, 0)) \subset \Lambda_{\varepsilon}$ . Then, for all t > 0,

$$\begin{aligned} J_{\varepsilon}(tv_0) &\leq \frac{t^2}{2} \|v_0\|_{\varepsilon}^2 - \int_{\mathbb{R}^N} F(tv_0(x,0)) \, dx \\ &\leq \frac{t^2}{2} \|v_0\|_{\varepsilon}^2 - C_1 t^{\theta} \int_{\Lambda_{\varepsilon}} (v_0(x,0))^{\theta} \, dx + C_2 |\Lambda_{\varepsilon}| \end{aligned}$$

where we have used  $(g_3)$ . Since  $\theta \in (2, 2_s^*)$ , we obtain  $J_{\varepsilon}(tv_0) \to -\infty$  as  $t \to \infty$ .

By Lemma 3.1 and invoking a variant of the mountain pass theorem without the Palais-Smale condition (see [45, Theorem 2.9]), we can find a Palais-Smale sequence  $(v_n) \subset X_{\varepsilon}$  at the mountain pass level  $c_{\varepsilon}$ , i.e.,

$$J_{\varepsilon}(v_n) \to c_{\varepsilon}$$
 and  $J'_{\varepsilon}(v_n) \to 0$  in  $X^*_{\varepsilon}$ ,

as  $n \to \infty$ , where

$$c_{\varepsilon} := \inf_{\gamma \in \Gamma_{\varepsilon}} \max_{t \in [0,1]} J_{\varepsilon}(\gamma(t))$$

and

$$\Gamma_{\varepsilon} := \{ \gamma \in C([0,1], X_{\varepsilon}) : \gamma(0) = 0, J_{\varepsilon}(\gamma(1)) < 0 \}$$

In view of the properties of g, it is easy to verify (see [35, 45]) that

$$c_{\varepsilon} = \inf_{v \in \mathcal{N}_{\varepsilon}} J_{\varepsilon}(v) = \inf_{v \in X_{\varepsilon} \setminus \{0\}} \max_{t \ge 0} J_{\varepsilon}(tv),$$

where

$$\mathcal{N}_{\varepsilon} := \{ v \in X_{\varepsilon} : \langle J'_{\varepsilon}(v), v \rangle = 0 \}$$

is the Nehari manifold associated with  $J_{\varepsilon}$ .

Next, we provide an important upper bound for the minimax level  $c_{\varepsilon}$ . For this purpose, we remember the following trace inequality (see [11, Theorem 2.1]):

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla v|^2 \, dx dy \ge S_* \left( \int_{\mathbb{R}^N} |v(x,0)|^{2^*_s} \, dx \right)^{\frac{2}{2^*_s}},\tag{3.4}$$

for all  $v \in X_0^s(\mathbb{R}^{N+1}_+)$ , where  $X_0^s(\mathbb{R}^{N+1}_+)$  is the completion of  $C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  under the norm:

$$\left(\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla v|^2 \, dx dy\right)^{\frac{1}{2}}$$

and the exact value of the best constant  $S_* = S(N, s) > 0$  is

$$S_* := \frac{2\pi^s \Gamma(1-s) \Gamma(\frac{N+2s}{2}) \Gamma(\frac{N}{2})^{\frac{2s}{N}}}{\Gamma(s) \Gamma(\frac{N-2s}{2}) \Gamma(N)^{\frac{2s}{N}}}.$$

This constant is achieved on the family of functions  $w_{\epsilon} = \mathcal{E}_s(u_{\epsilon})$ , where  $\mathcal{E}_s$  denotes the s-harmonic extension [15], and

$$u_{\epsilon}(x) := \frac{\epsilon^{\frac{N-2s}{2}}}{\left(|x|^2 + \epsilon^2\right)^{\frac{N-2s}{2}}}, \quad \epsilon > 0;$$

see [11, 18, 38] for more details. Hence,

$$w_{\epsilon}(x,y) := (P_s(\cdot,y) * u_{\epsilon})(x) = \int_{\mathbb{R}^N} P_s(x-\xi,y) u_{\epsilon}(\xi) \, d\xi,$$

where

$$P_s(x,y) := \frac{p_{N,s} y^{2s}}{(|x|^2 + y^2)^{\frac{N+2s}{2}}}$$

is the Poisson kernel for the extension problem in the half-space  $\mathbb{R}^{N+1}_+$ . Note that  $w_{\epsilon}(x,y) = \epsilon^{\frac{2s-N}{2}} w_1(\frac{x}{\epsilon},\frac{y}{\epsilon})$ .

**Lemma 3.2.** It holds  $0 < c_{\varepsilon} < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$ , where  $\zeta := 1 - \frac{V_1}{m^{2s}} \left(1 + \frac{1}{\kappa}\right) \in (0, 1)$ .

*Proof.* For simplicity, we assume that  $\varepsilon = 1$ . Let  $\phi(x, y) := \phi_0(|(x, y)|)$ , where  $\phi_0 \in C^{\infty}([0, \infty))$  is a non-increasing function such that

$$\phi_0(t) = 1$$
 if  $t \in [0, 1]$ ,  $\phi_0(t) = 0$  if  $t \ge 2$ ,

and suppose that  $B_2 \subset \Lambda$ . Then, we consider

$$\eta_{\epsilon}(x,y) := \frac{(\phi w_{\epsilon})(x,y)}{|(\phi w_{\epsilon})(\cdot,0)|_{2_{s}^{*}}}, \quad \epsilon > 0$$

Let us recall the following useful estimates (see [7, 10, 38]):

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \eta_{\epsilon}(x,y)|^2 \, dx dy = S_* + O(\epsilon^{N-2s}), \tag{3.5}$$

$$|\eta_{\epsilon}(\cdot,0)|_{2}^{2} = \begin{cases} O(\epsilon^{2s}), & \text{if } N > 4s, \\ O(\epsilon^{2s} \log(1/\epsilon)), & \text{if } N = 4s, \\ O(\epsilon^{N-2s}), & \text{if } 2s < N < 4s, \end{cases}$$
(3.6)

A FRACTIONAL RELATIVISTIC SCHRÖDINGER EQUATION WITH CRITICAL GROWTH

$$|\eta_{\epsilon}(\cdot,0)|_{q}^{q} \geq \begin{cases} O(\epsilon^{\frac{2N-(N-2s)q}{2}}), & \text{if } q > \frac{N}{N-2s}, \\ O(\epsilon^{\frac{N}{2}}|\log(\epsilon)|), & \text{if } q = \frac{N}{N-2s}, \\ O(\epsilon^{\frac{(N-2s)q}{2}}), & \text{if } q < \frac{N}{N-2s}. \end{cases}$$
(3.7)

Now, we define  $v_{\epsilon}(x,y) := \vartheta(my)\eta_{\epsilon,\beta}(x,y)$ , where  $\eta_{\epsilon,\beta}(x,y) = \eta_{\epsilon}(\beta x,\beta y)$ ,  $\beta := \zeta^{-\frac{2}{N-2s}}$ , and  $\vartheta$  is given in (2.6). Evidently,  $v_{\epsilon}(x,0) = \eta_{\epsilon,\beta}(x,0)$  and

$$\partial_{x_j} v_{\epsilon}(x,y) = \beta \vartheta(my) \partial_{x_j} \eta_{\epsilon,\beta}(x,y), \text{ for all } j = 1, \dots, N,$$

$$\partial_y v_{\epsilon}(x,y) = m\vartheta'(my)\eta_{\epsilon,\beta}(x,y) + \beta\vartheta(my)\partial_y\eta_{\epsilon,\beta}(x,y)$$

Therefore,

$$\begin{split} \|v_{\epsilon}\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} &= \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} \left[ \vartheta^{2}(my)(\beta^{2}|\nabla\eta_{\epsilon,\beta}|^{2} + m^{2}\eta_{\epsilon,\beta}^{2}) + 2m\beta\vartheta'(my)\vartheta(my)\eta_{\epsilon,\beta}\partial_{y}\eta_{\epsilon,\beta} + m^{2}(\vartheta'(my))^{2}\eta_{\epsilon,\beta}^{2} \right] dxdy \\ &= \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s}\vartheta^{2}(my)\beta^{2}|\nabla\eta_{\epsilon,\beta}|^{2} dxdy + \iint_{\mathbb{R}^{N+1}_{+}} 2m\beta y^{1-2s}\vartheta'(my)\vartheta(my)\eta_{\epsilon,\beta}\partial_{y}\eta_{\epsilon,\beta} dxdy \\ &+ \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s}m^{2} \left[ (\vartheta'(my))^{2} + (\vartheta(my))^{2} \right] \eta_{\epsilon,\beta}^{2} dxdy \\ &= \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s}\vartheta^{2}(my)\beta^{2}|\nabla\eta_{\epsilon,\beta}|^{2} dxdy + m^{2s} \int_{\mathbb{R}^{N}} \eta_{\epsilon,\beta}^{2}(x,0) dx, \end{split}$$
(3.8)

where we have used integration by parts,  $\kappa_s = 1$ , (2.7), and (2.8) to deduce that

$$\begin{split} &\iint_{\mathbb{R}^{N+1}_+} 2m\beta y^{1-2s}\vartheta'(my)\vartheta(my)\eta_{\epsilon,\beta}\partial_y\eta_{\epsilon,\beta}\,dxdy = \iint_{\mathbb{R}^{N+1}_+} my^{1-2s}\vartheta'(my)\vartheta(my)\partial_y(\eta^2_{\epsilon,\beta})\,dxdy \\ &= -\int_{\mathbb{R}^N} \left[\lim_{y\to 0} \vartheta(my)\vartheta'(my)my^{1-2s}\eta^2_{\epsilon,\beta}(x,y)\right]\,dx - \iint_{\mathbb{R}^{N+1}_+} m\eta^2_{\epsilon,\beta}\partial_y[y^{1-2s}\vartheta(my)\vartheta'(my)]\,dxdy \\ &= m^{2s}\kappa_s \int_{\mathbb{R}^N} \eta^2_{\epsilon,\beta}(x,0)\,dx - \iint_{\mathbb{R}^{N+1}_+} y^{1-2s}m^2\left[(\vartheta'(my))^2 + (\vartheta(my))^2\right]\eta^2_{\epsilon,\beta}\,dxdy \\ &= m^{2s} \int_{\mathbb{R}^N} \eta^2_{\epsilon,\beta}(x,0)\,dx - \iint_{\mathbb{R}^{N+1}_+} y^{1-2s}m^2\left[(\vartheta'(my))^2 + (\vartheta(my))^2\right]\eta^2_{\epsilon,\beta}\,dxdy. \end{split}$$

Consequently, using (3.8), a change of variable theorem,  $|\eta_{\varepsilon}(\cdot, 0)|_{2_s^*} = 1$ ,  $(g_2)$ , and  $0 \leq \vartheta(my) \leq 1$ , for all t > 0,

$$J_{1}(tv_{\epsilon}) \leq \frac{t^{2}}{2} \|v_{\epsilon}\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} + \frac{1}{2} \int_{\mathbb{R}^{N}} V(x)(tv_{\epsilon}(x,0))^{2} dx - \frac{\lambda t^{p}}{p} \int_{\mathbb{R}^{N}} |v_{\epsilon}(x,0)|^{p} dx - \frac{t^{2}_{s}^{*}}{2_{s}^{*}} \int_{\mathbb{R}^{N}} |v_{\epsilon}(x,0)|^{2_{s}^{*}} dx$$

$$\leq \frac{t^{2}}{2} \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} \beta^{2} |\nabla \eta_{\epsilon,\beta}|^{2} dx dy + (\bar{V} + m^{2s}) \frac{t^{2}}{2} |\eta_{\epsilon,\beta}(\cdot,0)|^{2}_{2} - \frac{\lambda t^{p}}{p} |\eta_{\epsilon,\beta}(\cdot,0)|^{p}_{p} - \frac{t^{2_{s}^{*}}}{2_{s}^{*}} |\eta_{\epsilon,\beta}(\cdot,0)|^{2_{s}^{*}}_{2_{s}^{*}}$$

$$\leq \frac{t^{2}}{2} \left[ \beta^{2s-N} \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla \eta_{\epsilon}|^{2} dx dy + \beta^{-N} (\bar{V} + m^{2s}) |\eta_{\epsilon}(\cdot,0)|^{2}_{2} \right] - \frac{\lambda t^{p}}{p} \beta^{-N} |\eta_{\epsilon}(\cdot,0)|^{p}_{p} - \frac{t^{2_{s}^{*}}}{2_{s}^{*}} \beta^{-N} =: \alpha(t).$$

$$(3.9)$$

where  $\bar{V} := \max_{\bar{\Lambda}} V > -m^{2s}$ . Clearly,  $\alpha(t) > 0$  for t > 0 small and  $\alpha(t) \to -\infty$  as  $t \to \infty$ , so  $\alpha(t)$  attains its maximum at some  $t_{\epsilon} > 0$  with  $\alpha'(t_{\epsilon}) = 0$ . This fact combined with (3.5)-(3.7) implies that there exist  $\delta_1, \delta_2 > 0$ , independent of  $\epsilon > 0$ , such that

$$\delta_1 \le t_\epsilon \le \delta_2.$$

Then, observing that the function

$$t \mapsto \frac{t^2}{2} \left[ \beta^{2s-N} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \eta_{\epsilon}|^2 \, dx \, dy + \beta^{-N} (\bar{V} + m^{2s}) |\eta_{\epsilon}(\cdot, 0)|_2^2 \right] - \frac{t^{2s}}{2s} \beta^{-N} dx \, dy + \beta^{-N} (\bar{V} + m^{2s}) |\eta_{\epsilon}(\cdot, 0)|_2^2 \right] - \frac{t^{2s}}{2s} \beta^{-N} dx \, dy + \beta^{-N} (\bar{V} + m^{2s}) |\eta_{\epsilon}(\cdot, 0)|_2^2 = 0$$

is increasing in the interval

$$\left[0, \left(\beta^{2s-N} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \eta_{\epsilon}|^2 \, dx dy + \beta^{-N} (\bar{V}+m^{2s}) |\eta_{\epsilon}(\cdot,0)|_2^2\right)^{\frac{1}{2s-2}} \beta^{\frac{N}{2s-2}}\right],$$

and exploiting the well-known inequality

$$(a+b)^{\alpha} \le a^{\alpha} + \alpha(a+b)^{\alpha-1}, \quad \text{for } a, b > 0, \ \alpha \ge 1,$$

and the estimates (3.5)-(3.7), we can deduce that

$$\begin{aligned} \alpha(t_{\epsilon}) &\leq \frac{s}{N} \left( \beta^{2s-N} \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla \eta_{\epsilon}|^{2} \, dx dy + \beta^{-N} (\bar{V} + m^{2s}) |\eta_{\epsilon}(\cdot, 0)|_{2}^{2} \right)^{\frac{N}{2s}} \beta^{\frac{N(N-2s)}{4s}} - \lambda C |\eta_{\epsilon}(\cdot, 0)|_{p}^{p} \\ &\leq \frac{s}{N} (\zeta S_{*})^{\frac{N}{2s}} + O(\epsilon^{N-2s}) + C |\eta_{\epsilon}(\cdot, 0)|_{2}^{2} - \lambda C |\eta_{\epsilon}(\cdot, 0)|_{p}^{p}, \end{aligned}$$

where we have used

$$\beta^{(2s-N)\frac{N}{2s} + \frac{N(N-2s)}{4s}} = \beta^{-\frac{N(N-2s)}{4s}} = \left(\zeta^{-\frac{2}{N-2s}}\right)^{-\frac{N(N-2s)}{4s}} = \zeta^{\frac{N}{2s}}$$

Next, we show that, for  $\epsilon > 0$  small enough,

$$\alpha(t_{\epsilon}) < \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}}.$$
(3.10)

Assume that N > 4s. Thus,  $p > 2 > \frac{N}{N-2s}$ , and using (3.7), we find

$$\alpha(t_{\epsilon}) \leq \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}} + O(\epsilon^{N-2s}) + O(\epsilon^{2s}) - O(\epsilon^{N-\frac{(N-2s)p}{2}}).$$

Since

$$N - \frac{(N-2s)p}{2} < 2s < N - 2s,$$

we infer that (3.10) holds as long as  $\epsilon > 0$  is sufficiently small. If N = 4s, then  $p > 2 = \frac{N}{N-2s}$ , and in view of (3.7), we obtain

$$\alpha(t_{\epsilon}) \leq \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}} + O(\epsilon^{2s} (1 + |\log \epsilon|)) - O(\epsilon^{4s - sp}).$$

Observing that p > 2 yields

$$\lim_{\epsilon \to 0} \frac{\epsilon^{4s-sp}}{\epsilon^{2s}(1+|\log \epsilon \,|)} = \infty,$$

we arrive at the assertion for  $\epsilon > 0$  small enough.

Now, let us consider the case 2s < N < 4s. First, we suppose that  $2_s^* - 2 . Hence,$ 

$$p > 2_s^* - 2 > \frac{N}{N - 2s},$$

which combined with (3.7) gives

$$\alpha(t_{\epsilon}) \leq \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}} + O(\epsilon^{N-2s}) - O(\epsilon^{N-\frac{(N-2s)p}{2}}).$$

Because

$$N - \frac{(N-2s)p}{2} < N - 2s < 2s,$$

we conclude that (3.10) is satisfied for  $\epsilon > 0$  small enough. Second, we deal with the case 2 .We distinguish the following subcases:

$$2 ,  $p = \frac{N}{N-2s}$ , and  $\frac{N}{N-2s} .$$$

If 2 , by (3.7), we see that

$$\alpha(t_{\epsilon}) \leq \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}} + O(\epsilon^{N-2s}) - \lambda O(\epsilon^{\frac{(N-2s)p}{2}})$$

and noting that

$$N-2s < \frac{(N-2s)p}{2},$$

we can put  $\lambda = e^{-\nu}$ , with  $\nu > \frac{(N-2s)(p-2)}{2}$ , to reach the required estimate. If  $p = \frac{N}{N-2s}$ , then, thanks to (3.7), we have

$$\alpha(t_{\epsilon}) \leq \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}} + O(\epsilon^{N-2s}) - \lambda O(\epsilon^{\frac{N}{2}} |\log \epsilon|),$$

and taking  $\lambda = \epsilon^{-\nu}$ , with  $\nu > 2s - \frac{N}{2}$ , we deduce the assertion for  $\epsilon > 0$  small enough. Finally, when  $\frac{N}{N-2s} , it follows from (3.7) that$ 

$$\alpha(t_{\epsilon}) \leq \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}} + O(\epsilon^{N-2s}) - \lambda O(\epsilon^{N-\frac{(N-2s)p}{2}}),$$

and choosing  $\lambda = \epsilon^{-\nu}$ , with  $\nu > 2s - \frac{(N-2s)p}{2}$ , we obtain the claim for  $\epsilon > 0$  sufficiently small. Consequently, (3.9) and (3.10) yield

$$\max_{t\geq 0} J_1(tv_{\epsilon}) < \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}},$$

which together with  $c_1 \leq \max_{t\geq 0} J_1(tv_{\epsilon})$  implies the desired conclusion.

In what follows, we show that  $J_{\varepsilon}$  satisfies a local compactness condition. First, we prove the boundedness of Palais-Smale sequences of  $J_{\varepsilon}$ .

**Lemma 3.3.** Let  $0 < c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$  and  $(v_n) \subset X_{\varepsilon}$  be a Palais-Smale sequence of  $J_{\varepsilon}$  at the level c. Then,  $(v_n)$  is bounded in  $X_{\varepsilon}$ .

*Proof.* By assumptions, we know that

$$J_{\varepsilon}(v_n) \to c \quad \text{and} \quad J'_{\varepsilon}(v_n) \to 0 \text{ in } X^*_{\varepsilon},$$

$$(3.11)$$

as  $n \to \infty$ . Using (3.11), (g<sub>3</sub>), (2.3), and (2.10), we see that, for n big enough,

$$\begin{split} c+1+\|v_n\|_{\varepsilon} &\geq J_{\varepsilon}(v_n) - \frac{1}{\theta} \langle J_{\varepsilon}'(v_n), v_n \rangle \\ &= \left(\frac{1}{2} - \frac{1}{\theta}\right) \|v_n\|_{\varepsilon}^2 + \frac{1}{\theta} \int_{\mathbb{R}^N} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) - \theta G_{\varepsilon}(x, v_n(x, 0)) \, dx \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta}\right) \|v_n\|_{\varepsilon}^2 - \left(\frac{1}{2} - \frac{1}{\theta}\right) \frac{V_1}{\kappa} \int_{\mathbb{R}^N} v_n^2(x, 0) \, dx \\ &= \left(\frac{1}{2} - \frac{1}{\theta}\right) \|v_n\|_{\varepsilon}^2 - \left(\frac{1}{2} - \frac{1}{\theta}\right) \frac{V_1}{\kappa m^{2s}} m^{2s} \int_{\mathbb{R}^N} v_n^2(x, 0) \, dx \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta}\right) \|v_n\|_{\varepsilon}^2 - \left(\frac{1}{2} - \frac{1}{\theta}\right) \frac{V_1}{\kappa m^{2s}} \|v_n\|_{X^s(\mathbb{R}^{N+1}_+)}^2 \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta}\right) \left(1 - \frac{V_1}{\kappa (m^{2s} - V_1)}\right) \|v_n\|_{\varepsilon}^2. \end{split}$$

Since  $\theta > 2$  and  $\kappa > \frac{V_1}{m^{2s} - V_1}$ , we conclude that  $(v_n)$  is bounded in  $X_{\varepsilon}$ .

**Remark 3.1.** If  $0 < c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$  and  $(v_n) \subset X_{\varepsilon}$  is a Palais-Smale sequence of  $J_{\varepsilon}$  at the level c, then we may always assume that  $(v_n)$  is nonnegative. In fact, Lemma 3.3 implies that also  $(v_n^-)$  is bounded in  $X_{\varepsilon}$ . Then, we have  $\langle J'_{\varepsilon}(v_n), v_n^- \rangle = o_n(1)$ , which combined with  $g(\cdot, t) = 0$  for  $t \leq 0$  yields  $||v_n^-||_{\varepsilon} = o_n(1)$ . Furthermore, it is easy to check that  $J_{\varepsilon}(v_n) = J_{\varepsilon}(v_n^+) + o_n(1)$  and  $J'_{\varepsilon}(v_n) = J'_{\varepsilon}(v_n^+) + o_n(1)$ .

The next concentration-compactness principle in the spirit of Lions [32, 33] will be used in the proof of the local compactness of  $J_{\varepsilon}$ . We start by recalling some useful definitions. A sequence  $(u_n) \subset L^1(\mathbb{R}^N)$  is tight if for every  $\xi > 0$ , there exists R > 0 such that

$$\int_{B_R^c} |u_n| \, dx < \xi, \quad \text{ for all } n \in \mathbb{N}$$

A sequence  $(v_n) \subset X^s(\mathbb{R}^{N+1})$  is tight if for every  $\xi > 0$ , there exists R > 0 such that

$$\iint_{\mathbb{R}^{N+1}_+ \setminus B^+_R} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \, dx dy < \xi, \quad \text{ for all } n \in \mathbb{N}.$$

**Remark 3.2.** Let us observe that if  $(v_n)$  is a bounded tight sequence in  $X^s(\mathbb{R}^{N+1}_+)$ , then  $(|v_n(\cdot, 0)|^{2^*_s})$  is a (bounded) tight sequence in  $L^1(\mathbb{R}^N)$ . To prove this, let R > 0 and consider  $\eta_R \in C^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  given by:

$$\eta_R(x,y) := \begin{cases} 0, & \text{if } (x,y) \in \frac{B_{R/2}^+}{1}, \\ 1, & \text{if } (x,y) \in \overline{\mathbb{R}^{N+1}_+} \setminus B_R^+ \end{cases}$$

with  $0 \leq \eta_R \leq 1$  and  $\|\nabla \eta_R\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \leq C/R$ , for some C > 0 independent of R > 0. Using the definition and the properties of  $\eta_R$ , Theorem 2.1, the boundedness of  $(v_n)$  in  $X^s(\mathbb{R}^{N+1}_+)$ , and  $|x+y|^2 \leq 2(|x|^2+|y|^2)$ for all  $x, y \in \mathbb{R}^N$ , we have that, for all  $n \in \mathbb{N}$ ,

$$\begin{split} &\left(\int_{B_{R}^{c}}|v_{n}(x,0)|^{2_{s}^{*}}\,dx\right)^{\frac{2}{2_{s}^{*}}} \leq \left(\int_{\mathbb{R}^{N}}|(v_{n}\eta_{R})(x,0)|^{2_{s}^{*}}\,dx\right)^{\frac{2}{2_{s}^{*}}} \\ \leq & C\left(\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla(v_{n}\eta_{R})|^{2}+m^{2}v_{n}^{2}\eta_{R}^{2})\,dxdy\right) \\ \leq & C\left(\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla v_{n}|^{2}+m^{2}v_{n}^{2})\eta_{R}^{2}\,dxdy+\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}|\nabla\eta_{R}|^{2}v_{n}^{2}\,dxdy\right) \\ \leq & C\left(\iint_{\mathbb{R}^{N+1}_{+}\setminus B_{R/2}^{+}}y^{1-2s}(|\nabla v_{n}|^{2}+m^{2}v_{n}^{2})\,dxdy+\frac{C}{R^{2}}\right). \end{split}$$

From the aforementioned estimate and the tightness of  $(v_n)$  in  $X^s(\mathbb{R}^{N+1}_+)$ , we derive that  $(|v_n(\cdot, 0)|^{2^s_s})$  is tight in  $L^1(\mathbb{R}^N)$ , as desired.

**Proposition 3.1.** Let  $(v_n)$  be a bounded tight sequence in  $X^s(\mathbb{R}^{N+1}_+)$  such that  $v_n \rightharpoonup v$  in  $X^s(\mathbb{R}^{N+1}_+)$ . Let  $\mu$  and  $\nu$  be two bounded nonnegative measures on  $\mathbb{R}^{N+1}_+$  and  $\mathbb{R}^N$ , respectively, and such that

$$y^{1-2s}(|\nabla v_n|^2 + m^2 v_n^2) \rightharpoonup \mu \quad weakly in the sense of measures$$
 (3.12)

and

$$|v_n(\cdot, 0)|^{2^*_s} \rightharpoonup \nu$$
 weakly in the sense of measures. (3.13)

Then, there exist an at most countable set I and three families  $(x_i)_{i\in I} \subset \mathbb{R}^N$ ,  $(\mu_i)_{i\in I} \subset (0,\infty)$ ,  $(\nu_i)_{i\in I} \subset (0,\infty)$  such that

$$\nu = |v(\cdot, 0)|^{2^*_s} + \sum_{i \in I} \nu_i \delta_{x_i}, \tag{3.14}$$

$$\mu \ge y^{1-2s}(|\nabla v|^2 + m^2 v^2) + \sum_{i \in I} \mu_i \delta_{(x_i,0)}, \qquad (3.15)$$

$$\mu_i \ge S_* \nu_i^{\frac{2}{2s}}, \quad \text{for all } i \in I.$$
(3.16)

*Proof.* We follow the strategy used in the proof of [32, Lemma I.1] (see also [33, Lemma 2.3], [21, Proposition 3.2.1], and [10, Theorem 5.1]). We first suppose that  $v \equiv 0$ . We claim that, for all  $\varphi \in C_c^{\infty}(\mathbb{R}^{N+1}_+)$ , it holds

$$\left(\int_{\mathbb{R}^N} |\varphi(x,0)|^{2^*_s} d\nu\right)^{\frac{1}{2^*_s}} \le C_0 \left(\iint_{\mathbb{R}^{N+1}_+} \varphi^2 d\mu\right)^{\frac{1}{2}},\tag{3.17}$$

for some constant  $C_0 > 0$ . For this purpose, we fix  $\varphi \in C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  and let  $K := \operatorname{supp}(\varphi)$ . By (3.4), we deduce that

$$\left(\int_{\mathbb{R}^{N}} |(\varphi v_{n})(x,0)|^{2^{*}_{s}} dx\right)^{\frac{1}{2^{*}_{s}}} \leq S^{-\frac{1}{2}}_{*} \left(\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} [|\nabla(\varphi v_{n})|^{2} + m^{2}(\varphi v_{n})^{2}] \, dx dy\right)^{\frac{1}{2}}.$$
(3.18)

Now, we note that (3.13) implies that

$$\int_{\mathbb{R}^N} |(\varphi v_n)(x,0)|^{2^*_s} dx \to \int_{\mathbb{R}^N} |\varphi(x,0)|^{2^*_s} d\nu.$$
(3.19)

On the other hand,

$$\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} [|\nabla(\varphi v_n)|^2 + m^2 (\varphi v_n)^2] \, dx \, dy$$
  
= 
$$\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} \varphi^2 [|\nabla v_n|^2 + m^2 v_n^2] \, dx \, dy + \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} v_n^2 |\nabla \varphi|^2 \, dx \, dy$$
  
+ 
$$2 \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} v_n \varphi \nabla \varphi \cdot \nabla v_n \, dx \, dy.$$
(3.20)

Since  $H^1(K, y^{1-2s})$  is compactly embedded in  $L^2(K, y^{1-2s})$  (see [21, Lemma 3.1.2]), we have that  $v_n \to 0$  in  $L^2(K, y^{1-2s})$ , which yields

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n^2 |\nabla \varphi|^2 \, dx dy \le C \iint_K y^{1-2s} v_n^2 \, dx dy \to 0. \tag{3.21}$$

Furthermore, the Hölder inequality, the boundedness of  $(v_n)$  in  $X^s(\mathbb{R}^{N+1}_+)$ , and (3.21) lead to

$$\begin{aligned} \left| \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} v_n \varphi \nabla \varphi \cdot \nabla v_n \, dx dy \right| \\ &\leq \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} v_n^2 |\nabla \varphi|^2 \, dx dy \right)^{\frac{1}{2}} \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} \varphi^2 |\nabla v_n|^2 \, dx dy \right)^{\frac{1}{2}} \\ &\leq C \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} v_n^2 \, dx dy \right)^{\frac{1}{2}} \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla v_n|^2 \, dx dy \right)^{\frac{1}{2}} \\ &\leq C \left( \iint_{K} y^{1-2s} v_n^2 \, dx dy \right)^{\frac{1}{2}} \to 0. \end{aligned}$$
(3.22)

Finally, taking (3.12) into account, we see that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} \varphi^2(|\nabla v_n|^2 + m^2 v_n^2) \, dx dy \to \iint_{\mathbb{R}^{N+1}_+} \varphi^2 \, d\mu. \tag{3.23}$$

Putting together (3.18)-(3.23), we can infer that (3.17) holds with  $C_0 = S_*^{-\frac{1}{2}}$ . Now, assume that v is not necessarily 0. Set  $w_n := v_n - v$ . Clearly,  $(w_n)$  is a bounded tight sequence in  $X^s(\mathbb{R}^{N+1}_+)$  such that  $w_n \to 0$  in  $X^s(\mathbb{R}^{N+1}_+)$ . Moreover, there exist two bounded nonnegative measures  $\tilde{\mu}$  and  $\tilde{\nu}$  on  $\mathbb{R}^{N+1}_+$  and  $\mathbb{R}^N$ , respectively, such that

 $y^{1-2s}(|\nabla w_n|^2 + m^2 w_n^2) \rightharpoonup \tilde{\mu}$  weakly in the sense of measures, (3.24)

$$|w_n(\cdot, 0)|^{2^*_s} \to \tilde{\nu}$$
 weakly in the sense of measures. (3.25)

Then, we are in the previous case, and we can use (3.17) to deduce that

$$\left(\int_{\mathbb{R}^N} |\varphi(x,0)|^{2^*_s} d\tilde{\nu}\right)^{\frac{1}{2^*_s}} \le S^{-\frac{1}{2}}_* \left(\iint_{\mathbb{R}^{N+1}_+} \varphi^2 d\tilde{\mu}\right)^{\frac{1}{2}}, \quad \text{for all } \varphi \in C^\infty_c(\overline{\mathbb{R}^{N+1}_+}).$$

Hence, as in [32, Lemma 1.2], we can find an at most countable set I, a family of distinct points  $(x_i)_{i \in I} \subset \mathbb{R}^N$ and  $(\nu_i)_{i \in I} \subset (0, \infty)$  such that

$$\tilde{\nu} = \sum_{i \in I} \nu_i \delta_{x_i}.$$
(3.26)

Pick  $\varphi \in C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$ . By the Brezis-Lieb lemma [12], we know that

$$|(\varphi w_n)(\cdot,0)|_{2_s^*}^{2_s^*} = |(\varphi v_n)(\cdot,0)|_{2_s^*}^{2_s^*} - |(\varphi v)(\cdot,0)|_{2_s^*}^{2_s^*} + o_n(1)$$

The aforementioned fact combined with (3.13), (3.25), and (3.26), the boundedness of  $(v_n)$  in  $X^s(\mathbb{R}^{N+1}_+)$ , and the tightness of  $(|v_n(\cdot, 0)|^{2^*_s})$  implies that (3.14) holds. Now, take  $\psi \in C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  such that  $0 \le \psi \le 1$ ,  $\psi = 1$  in  $B_{\frac{1}{2}}^+, \psi = 0$  in  $(B_1^+)^c$  and  $\|\nabla \psi\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \le 2$ . Fix  $i \in I$ . For  $\rho > 0$ , we define  $\psi_{\rho}(x, y) := \psi(\frac{x-x_i}{\rho}, \frac{y}{\rho})$ . Applying (3.4) to  $\psi_{\rho}v_n$ , we have that

$$S_*^{\frac{1}{2}} \left( \int_{\mathbb{R}^N} |\psi_{\rho}(x,0)v_n(x,0)|^{2^*_s} \, dx \right)^{\frac{1}{2^*_s}} \le \left( \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla(\psi_{\rho}v_n)|^2 \, dx dy \right)^{\frac{1}{2}}$$

from which

$$\begin{split} S^{\frac{1}{2}}_{*} \left( \int_{\mathbb{R}^{N}} |\psi_{\rho}(x,0)v_{n}(x,0)|^{2^{*}_{s}} dx \right)^{\frac{1}{2^{*}_{s}}} \\ &\leq \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla\psi_{\rho}|^{2} v_{n}^{2} dx dy \right)^{\frac{1}{2}} + \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla v_{n}|^{2} \psi_{\rho}^{2} dx dy \right)^{\frac{1}{2}} \\ &\leq \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla\psi_{\rho}|^{2} v_{n}^{2} dx dy \right)^{\frac{1}{2}} + \left( \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} (|\nabla v_{n}|^{2} + m^{2} v_{n}^{2}) \psi_{\rho}^{2} dx dy \right)^{\frac{1}{2}}. \end{split}$$

Letting  $n \to \infty$  and exploiting the fact that  $\psi_{\rho}$  has compact support, we obtain

$$S_*^{\frac{1}{2}} \left( \int_{\mathbb{R}^N} |\psi_{\rho}(x,0)|^{2^*_s} \, d\nu \right)^{\frac{1}{2^*_s}} \le \left( \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \psi_{\rho}|^2 v^2 \, dx dy \right)^{\frac{1}{2}} + \left( \iint_{\mathbb{R}^{N+1}_+} \psi_{\rho}^2 \, d\mu \right)^{\frac{1}{2}}. \tag{3.27}$$

Using the Hölder inequality with exponents  $\gamma$  and  $\frac{\gamma}{\gamma-1}$  and recalling (2.1), we see that

$$\begin{split} \left(\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla\psi_{\rho}|^{2} v^{2} \, dx dy\right)^{\frac{1}{2}} &\leq \frac{C}{\rho} \left(\iint_{B^{+}_{\rho}(x_{i},0)} y^{1-2s} v^{2} \, dx dy\right)^{\frac{1}{2}} \\ &\leq \frac{C}{\rho} \left(\iint_{B^{+}_{\rho}(x_{i},0)} y^{1-2s} |v|^{2\gamma} \, dx dy\right)^{\frac{1}{2\gamma}} \left(\iint_{B^{+}_{\rho}(x_{i},0)} y^{1-2s} \, dx dy\right)^{\frac{\gamma-1}{2\gamma}} \\ &\leq C \left(\iint_{B^{+}_{\rho}(x_{i},0)} y^{1-2s} |v|^{2\gamma} \, dx dy\right)^{\frac{1}{2\gamma}} \to 0 \quad \text{as } \rho \to 0. \end{split}$$

Then, passing to the limit as  $\rho \to 0$  in (3.27), we find

$$S_* \nu_i^{\frac{2}{2s}} \le \mu_i := \lim_{\rho \to 0} \mu(B_\rho^+(x_i, 0)),$$

and so (3.16) is true. Since  $\mu \geq \sum_{i \in I} \mu_i \delta_{(x_i,0)}, \ \mu \geq y^{1-2s}(|\nabla v|^2 + m^2 v^2)$  (by the weak convergence), and  $y^{1-2s}(|\nabla v|^2 + m^2 v^2)$  and  $\sum_{i \in I} \mu_i \delta_{(x_i,0)}$  are orthogonal, we deduce that (3.15) hold.

Next, we prove the tightness of the Palais-Smale sequences of  $J_{\varepsilon}$ . More precisely, we establish the following result.

**Lemma 3.4.** Let  $0 < c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$  and  $(v_n) \subset X_{\varepsilon}$  be a Palais-Smale sequence of  $J_{\varepsilon}$  at the level c. Then, for all  $\xi > 0$ , there exists  $R = R(\xi) > 0$  such that

$$\limsup_{n \to \infty} \left[ \iint_{\mathbb{R}^{N+1} \setminus B_R^+} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \, dx \, dy + \int_{\mathbb{R}^N \setminus B_R} (V_{\varepsilon}(x) + V_1) v_n^2(x, 0) \, dx \right] < \xi. \tag{3.28}$$

*Proof.* For R > 0, let  $\eta_R \in C^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  be a function such that

$$\eta_R(x,y) := \begin{cases} 0 & \text{if } (x,y) \in \frac{B_{R/2}^+}{1}, \\ 1 & \text{if } (x,y) \in \overline{\mathbb{R}^{N+1}_+} \setminus B_R^+, \end{cases}$$

with  $0 \leq \eta_R \leq 1$  and  $\|\nabla \eta_R\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \leq C/R$ , for some C > 0 independent of R > 0. Since  $(v_n)$  is a bounded Palais-Smale sequence in  $X_{\varepsilon}$ , we see that  $\langle J'_{\varepsilon}(v_n), v_n \eta_R^2 \rangle = o_n(1)$ , i.e.,

$$\begin{aligned} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \eta_R^2 \, dx dy + \int_{\mathbb{R}^N} V_{\varepsilon}(x) v_n^2(x,0) \eta_R^2(x,0) \, dx \\ &= -2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \eta_R \nabla v_n \cdot \nabla \eta_R \, dx dy + \int_{\mathbb{R}^N} g_{\varepsilon}(x,v_n(x,0)) v_n(x,0) \eta_R^2(x,0) \, dx + o_n(1), \end{aligned}$$

which can be rewritten as:

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \eta_R^2 \, dx \, dy + \int_{\mathbb{R}^N} (V_{\varepsilon}(x) + V_1) v_n^2(x, 0) \eta_R^2(x, 0) \, dx$$
$$= \int_{\mathbb{R}^N} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \eta_R^2(x, 0) \, dx + V_1 \int_{\mathbb{R}^N} v_n^2(x, 0) \eta_R^2(x, 0) \, dx$$
$$- 2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \eta_R \nabla v_n \cdot \nabla \eta_R \, dx \, dy + o_n(1).$$
(3.29)

Choose R > 0 such that  $\Lambda_{\varepsilon} \subset B_{R/2}$ . Thus, thanks to  $(g_3)$ -(ii),

$$\int_{\mathbb{R}^N} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \eta_R^2(x, 0) \, dx + V_1 \int_{\mathbb{R}^N} v_n^2(x, 0) \eta_R^2(x, 0) \, dx \le V_1 \left(1 + \frac{1}{\kappa}\right) \int_{\mathbb{R}^N} v_n^2(x, 0) \eta_R^2(x, 0) \, dx.$$
(3.30)

On the other hand, by Hölder's inequality,  $0 \le \eta_R \le 1$ ,  $\|\nabla \eta_R\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \le C/R$ , and the boundedness of  $(v_n)$  in  $X_{\varepsilon}$ , we have

$$\left| 2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \eta_R \nabla v_n \cdot \nabla \eta_R \, dx dy \right| \leq \frac{C}{R} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n |\nabla v_n| \, dx dy$$
$$\leq \frac{C}{R} \left( \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n^2 \, dx dy \right)^{\frac{1}{2}} \left( \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla v_n|^2 \, dx dy \right)^{\frac{1}{2}}$$
$$\leq \frac{C}{R}. \tag{3.31}$$

Then, applying (2.4) to  $v_n\eta_R$ , and using  $\|\nabla\eta_R\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \leq C/R$  and (3.31), we infer that

$$\begin{split} V_1\left(1+\frac{1}{\kappa}\right) &\int_{\mathbb{R}^N} v_n^2(x,0)\eta_R^2(x,0) \, dx \\ &\leq V_1 m^{-2s} \left(1+\frac{1}{\kappa}\right) \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla(v_n\eta_R)|^2 \, dx dy + V_1 m^{2-2s} \left(1+\frac{1}{\kappa}\right) \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n^2 \eta_R^2 \, dx dy \\ &= V_1 m^{-2s} \left(1+\frac{1}{\kappa}\right) \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla v_n|^2 \eta_R^2 \, dx dy + V_1 m^{-2s} \left(1+\frac{1}{\kappa}\right) \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \eta_R|^2 v_n^2 \, dx dy \end{split}$$

$$+ 2V_{1}m^{-2s}\left(1+\frac{1}{\kappa}\right)\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}v_{n}\eta_{R}\nabla v_{n}\cdot\nabla\eta_{R}\,dxdy + V_{1}m^{2-2s}\left(1+\frac{1}{\kappa}\right)\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}v_{n}^{2}\eta_{R}^{2}\,dxdy$$

$$\leq V_{1}m^{-2s}\left(1+\frac{1}{\kappa}\right)\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla v_{n}|^{2}+m^{2}v_{n}^{2})\eta_{R}^{2}\,dxdy + \frac{C}{R^{2}}\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}v_{n}^{2}\,dxdy + \frac{C}{R}$$

$$\leq V_{1}m^{-2s}\left(1+\frac{1}{\kappa}\right)\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla v_{n}|^{2}+m^{2}v_{n}^{2})\eta_{R}^{2}\,dxdy + \frac{C}{R^{2}}+\frac{C}{R}.$$
(3.32)

Putting together (3.29)-(3.32), we arrive at

$$\left[1 - V_1 m^{-2s} \left(1 + \frac{1}{\kappa}\right)\right] \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \eta_R^2 \, dx \, dy + \int_{\mathbb{R}^N} (V_\varepsilon(x) + V_1) v_n^2(x, 0) \eta_R^2(x, 0) \, dx$$
  
$$\leq \frac{C}{R} + \frac{C}{R^2} + o_n(1).$$
(3.33)

By means of  $\kappa > \frac{V_1}{m^{2s}-V_1}$ ,  $(V_1)$  and the definition of  $\eta_R$ , we deduce that (3.33) implies the assertion.

**Remark 3.3.** Differently from [3, 7, 19], we use  $\langle J'_{\varepsilon}(v_n), v_n \eta_R^2 \rangle = o_n(1)$  instead of  $\langle J'_{\varepsilon}(v_n), v_n \eta_R \rangle = o_n(1)$  to obtain (3.28). This is motivated by the fact that to estimate the quadratic terms, it is needed to apply in a careful way Inequality (2.4).

**Remark 3.4.** Let  $r \in [2, 2_s^*]$ . Exploiting  $\eta_R(\cdot, 0) = 1$  in  $B_R^c$ , Theorem 2.1,  $|x + y|^2 \leq 2(|x|^2 + |y|^2)$  for all  $x, y \in \mathbb{R}^N$ ,  $\|\nabla \eta_R\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \leq C/R$  and the boundedness of  $(v_n)$  in  $X_{\varepsilon}$ , we can see that

$$\begin{split} &\left(\int_{B_{R}^{c}}|v_{n}(x,0)|^{r}\,dx\right)^{\frac{2}{r}} \leq \left(\int_{\mathbb{R}^{N}}|(v_{n}\eta_{R})(x,0)|^{r}\,dx\right)^{\frac{2}{r}} \\ &\leq C\left(\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla(v_{n}\eta_{R})|^{2}+m^{2}v_{n}^{2}\eta_{R}^{2})\,dxdy\right) \\ &\leq C\left(\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla v_{n}|^{2}+m^{2}v_{n}^{2})\eta_{R}^{2}\,dxdy+\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}|\nabla\eta_{R}|^{2}v_{n}^{2}\,dxdy\right) \\ &\leq C\left(\iint_{\mathbb{R}^{N+1}_{+}}y^{1-2s}(|\nabla v_{n}|^{2}+m^{2}v_{n}^{2})\eta_{R}^{2}\,dxdy+\frac{C}{R^{2}}\right), \end{split}$$

which combined with (3.33) gives

$$\left(\int_{B_R^c} |v_n(x,0)|^r \, dx\right)^{\frac{2}{r}} \le \frac{C}{R} + \frac{C}{R^2} + o_n(1).$$

Thus,

$$\lim_{R \to \infty} \limsup_{n \to \infty} \int_{B_R^c} |v_n(x,0)|^r \, dx = 0, \quad \text{for all } r \in [2, 2_s^*].$$
(3.34)

At this point, we can show that the modified functional fulfills a local compactness condition.

**Lemma 3.5.** Let  $0 < c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$ . Then,  $J_{\varepsilon}$  satisfies the Palais-Smale condition at the level c.

*Proof.* Let  $0 < c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$  and  $(v_n) \subset X_{\varepsilon}$  be a Palais-Smale sequence at the level c, namely,

$$J_{\varepsilon}(v_n) \to c$$
 and  $J'_{\varepsilon}(v_n) \to 0$  in  $X^*_{\varepsilon}$ ,

as  $n \to \infty$ . By Lemma 3.3, we know that  $(v_n)$  is bounded in  $X_{\varepsilon}$ . In view of Remark 3.1, we can suppose that  $(v_n)$  is nonnegative. Thanks to the reflexivity of  $X_{\varepsilon}$  and Theorem 2.1, up to a subsequence, we may assume that

$$\begin{cases} v_n \rightharpoonup v & \text{in } X_{\varepsilon}, \\ v_n(\cdot, 0) \rightarrow v(\cdot, 0) & \text{in } L^r_{loc}(\mathbb{R}^N), \text{ for all } r \in [1, 2^*_s), \\ v_n(\cdot, 0) \rightarrow v(\cdot, 0) & \text{a.e. in } \mathbb{R}^N. \end{cases}$$
(3.35)

We are going to demonstrate that  $v_n \to v$  in  $X_{\varepsilon}$ . Using the density of  $C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  in  $X_{\varepsilon}$ ,  $(g_1)$ ,  $(g_2)$ ,  $(f_2)$ , and (3.35), it is easy to check that  $\langle J'_{\varepsilon}(v), \varphi \rangle = 0$  for all  $\varphi \in X_{\varepsilon}$ . In particular,

$$\|v\|_{\varepsilon}^{2} = \int_{\mathbb{R}^{N}} g_{\varepsilon}(x, v(x, 0))v(x, 0) \, dx.$$

$$(3.36)$$

On the other hand,  $\langle J'_{\varepsilon}(v_n), v_n \rangle = o_n(1)$ , i.e.,

$$\|v_n\|_{\varepsilon}^2 = \int_{\mathbb{R}^N} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \, dx + o_n(1).$$
(3.37)

In light of (3.36) and (3.37), if we prove that

$$\int_{\mathbb{R}^N} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \, dx = \int_{\mathbb{R}^N} g_{\varepsilon}(x, v(x, 0)) v(x, 0) \, dx + o_n(1), \tag{3.38}$$

then we deduce that  $||v_n||_{\varepsilon} \to ||v||_{\varepsilon}$  as  $n \to \infty$ , and recalling that  $X_{\varepsilon}$  is a Hilbert space, we conclude that  $v_n \to v$  in  $X_{\varepsilon}$  as  $n \to \infty$ . Next, we verify that (3.38) is valid. By virtue of Lemma 3.4, fixed  $\xi > 0$ , there exists  $R = R(\xi) > 0$  such that (3.28) is true. By  $(g_2)$ ,  $(f_1)$ ,  $(f_2)$ , and (3.34), we see that

$$\limsup_{n \to \infty} \left| \int_{B_R^c} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \, dx \right| \leq C \limsup_{n \to \infty} \int_{B_R^c} (|v_n(x, 0)|^2 + |v_n(x, 0)|^q + |v_n(x, 0)|^{2^*_s}) \, dx \\ \leq C\xi. \tag{3.39}$$

On the other hand, because  $g_{\varepsilon}(\cdot, v(\cdot, 0))v(\cdot, 0) \in L^1(\mathbb{R}^N)$ , we can take R > 0 large enough so that

$$\int_{B_R^c} g_{\varepsilon}(x, v(x, 0)) v(x, 0) \, dx \le \xi.$$
(3.40)

Thus, (3.39) and (3.40) yield

$$\lim_{n \to \infty} \sup_{n \to \infty} \left| \int_{B_R^c} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \, dx - \int_{B_R^c} g_{\varepsilon}(x, v(x, 0)) v(x, 0) \, dx \right| \le C\xi.$$
(3.41)

Now, it follows from the definition of g that

$$g_{\varepsilon}(x, v_n(x, 0))v_n(x, 0) \le f(v_n(x, 0))v_n(x, 0) + a^{2^*_s} + \frac{V_1}{\kappa}v_n^2(x, 0), \quad \text{ for a.e. } x \in \Lambda_{\varepsilon}^c.$$

Since  $B_R \cap \Lambda_{\varepsilon}^c$  is bounded, we can use the aforementioned estimate,  $(f_1)$ ,  $(f_2)$ , (3.35), and the dominated convergence theorem to infer that, as  $n \to \infty$ ,

$$\int_{B_R \cap \Lambda_{\varepsilon}^c} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \, dx = \int_{B_R \cap \Lambda_{\varepsilon}^c} g_{\varepsilon}(x, v(x, 0)) v(x, 0) \, dx + o_n(1). \tag{3.42}$$

At this point, we aim to show that

$$\int_{\Lambda_{\varepsilon}} v_n^{2^*_s}(x,0) \, dx = \int_{\Lambda_{\varepsilon}} v^{2^*_s}(x,0) \, dx + o_n(1). \tag{3.43}$$

In fact, if we assume that (3.43) holds, then we can exploit  $(g_2)$ ,  $(f_1)$ ,  $(f_2)$ , (3.35), and the dominated convergence theorem again to obtain

$$\int_{B_R \cap \Lambda_{\varepsilon}} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) \, dx = \int_{B_R \cap \Lambda_{\varepsilon}} g_{\varepsilon}(x, v(x, 0)) v(x, 0) \, dx + o_n(1),$$

which combined with (3.41) and (3.42) gives (3.38). Therefore, we shall prove that (3.43) is satisfied. Taking into account that  $(v_n)$  is a bounded tight sequence in  $X^s(\mathbb{R}^{N+1}_+)$ , we may suppose that

$$y^{1-2s}(|\nabla v_n|^2 + m^2 v_n^2) \rightharpoonup \mu \quad \text{weakly in the sense of measures,}$$
$$v_n^{2^*_s}(\cdot, 0) \rightharpoonup \nu \quad \text{weakly in the sense of measures,}$$
(3.44)

where  $\mu$  and  $\nu$  are two bounded nonnegative measures on  $\mathbb{R}^{N+1}_+$  and  $\mathbb{R}^N$ , respectively. Thus, applying Proposition 3.1, we can find an at most countable index set I and sequences  $(x_i)_{i \in I} \subset \mathbb{R}^N$ ,  $(\mu_i)_{i \in I} \subset (0, \infty)$ , and  $(\nu_i)_{i \in I} \subset (0, \infty)$  such that

$$\nu = v^{2^{s}_{s}}(\cdot, 0) + \sum_{i \in I} \nu_{i} \delta_{x_{i}},$$

$$\mu \ge y^{1-2s}(|\nabla v|^{2} + m^{2}v^{2}) + \sum_{i \in I} \mu_{i} \delta_{(x_{i},0)},$$

$$\mu_{i} \ge S_{*} \nu_{i}^{\frac{2^{*}}{2^{s}}}, \quad \text{for all } i \in I.$$
(3.45)

Let us show that  $(x_i)_{i\in I} \cap \Lambda_{\varepsilon} = \emptyset$ . Assume, by contradiction, that  $x_i \in \Lambda_{\varepsilon}$  for some  $i \in I$ . Fixed  $\rho > 0$ , we define  $\psi_{\rho}(x, y) := \psi(\frac{x-x_i}{\rho}, \frac{y}{\rho})$ , where  $\psi \in C_c^{\infty}(\overline{\mathbb{R}^{N+1}_+})$  is such that  $\psi = 1$  in  $B_{\frac{1}{2}}^+$  and  $\psi = 0$  in  $(B_1^+)^c$ ,  $0 \le \psi \le 1$ , and  $\|\nabla \psi\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \le 2$ . We suppose that  $\rho > 0$  is such that  $\operatorname{supp}(\psi_{\rho}(\cdot, 0)) \subset \Lambda_{\varepsilon}$ . Because  $(\psi_{\rho}v_n)$ is bounded in  $X_{\varepsilon}$ , we have that  $\langle J'_{\varepsilon}(v_n), \psi_{\rho}v_n \rangle = o_n(1)$ , and so

$$\begin{aligned} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \psi_\rho \, dx dy + \int_{\mathbb{R}^N} V_\varepsilon(x) v_n^2(x,0) \psi_\rho(x,0) \, dx \\ &= -\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \nabla v_n \cdot \nabla \psi_\rho \, dx dy + \int_{\mathbb{R}^N} f(v_n(x,0)) \psi_\rho(x,0) v_n(x,0) \, dx + \int_{\mathbb{R}^N} \psi_\rho(x,0) v_n^{2^*_s}(x,0) \, dx + o_n(1), \end{aligned}$$

or equivalently,

$$\begin{aligned} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (|\nabla v_n|^2 + m^2 v_n^2) \psi_\rho \, dx dy + \int_{\mathbb{R}^N} (V_\varepsilon(x) + V_1) v_n^2(x, 0) \psi_\rho(x, 0) \, dx \\ &= V_1 \int_{\mathbb{R}^N} v_n^2(x, 0) \psi_\rho(x, 0) \, dx + \int_{\mathbb{R}^N} f(v_n(x, 0)) \psi_\rho(x, 0) v_n(x, 0) \, dx + \int_{\mathbb{R}^N} \psi_\rho(x, 0) v_n^{2s}(x, 0) \, dx \\ &- \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \nabla v_n \cdot \nabla \psi_\rho \, dx dy + o_n(1). \end{aligned}$$
(3.46)

Since f has subcritical growth and  $\psi_{\rho}(\cdot, 0)$  has compact support, we can use (3.35) to see that

$$\lim_{\rho \to 0} \lim_{n \to \infty} \int_{\mathbb{R}^N} f(v_n(x,0))\psi_\rho(x,0)v_n(x,0)\,dx = \lim_{\rho \to 0} \int_{\mathbb{R}^N} f(v(x,0))\psi_\rho(x,0)v(x,0)\,dx = 0 \tag{3.47}$$

and

$$\lim_{\rho \to 0} \lim_{n \to \infty} \int_{\mathbb{R}^N} v_n^2(x,0) \psi_\rho(x,0) \, dx = \lim_{\rho \to 0} \int_{\mathbb{R}^N} v^2(x,0) \psi_\rho(x,0) \, dx = 0.$$
(3.48)

Now, we prove that

$$\lim_{\rho \to 0} \limsup_{n \to \infty} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \nabla v_n \cdot \nabla \psi_\rho \, dx dy = 0.$$
(3.49)

From the Hölder inequality,  $(v_n)$  is bounded in  $X^s(\mathbb{R}^{N+1}_+)$ ,  $X^s(\mathbb{R}^{N+1}_+)$  is compactly embedded in  $L^2(B^+_{\rho}(x_i, 0), y^{1-2s})$ , and  $\|\nabla \psi_{\rho}\|_{L^{\infty}(\mathbb{R}^{N+1}_+)} \leq \frac{C}{\rho}$ , we obtain that

$$\begin{split} & \limsup_{n \to \infty} \left| \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} v_n \nabla v_n \cdot \nabla \psi_\rho \, dx dy \right| \\ & \leq \limsup_{n \to \infty} \left( \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla v_n|^2 \, dx dy \right)^{\frac{1}{2}} \left( \iint_{B^+_\rho(x_i,0)} y^{1-2s} v_n^2 |\nabla \psi_\rho|^2 \, dx dy \right)^{\frac{1}{2}} \\ & \leq \frac{C}{\rho} \left( \iint_{B^+_\rho(x_i,0)} y^{1-2s} v^2 \, dx dy \right)^{\frac{1}{2}}. \end{split}$$

Applying the Hölder inequality with exponents  $\gamma$  and  $\frac{\gamma}{\gamma-1}$  and bearing in mind (2.1), we have that

$$\begin{split} & \frac{C}{\rho} \left( \iint_{B_{\rho}^{+}(x_{i},0)} y^{1-2s} v^{2} \, dx dy \right)^{\frac{1}{2}} \\ & \leq \frac{C}{\rho} \left( \iint_{B_{\rho}^{+}(x_{i},0)} y^{1-2s} v^{2\gamma} \, dx dy \right)^{\frac{1}{2\gamma}} \left( \iint_{B_{\rho}^{+}(x_{i},0)} y^{1-2s} \, dx dy \right)^{\frac{\gamma-1}{2\gamma}} \\ & \leq C \left( \iint_{B_{\rho}^{+}(x_{i},0)} y^{1-2s} v^{2\gamma} \, dx dy \right)^{\frac{1}{2\gamma}} \to 0 \quad \text{as } \rho \to 0. \end{split}$$

The aforementioned estimates show that (3.49) is satisfied. Therefore, from (3.45) and taking the limit as  $n \to \infty$  and  $\rho \to 0$  in (3.46), we deduce that (3.47), (3.48), (3.49) and ( $V_1$ ) yield  $\mu_i \leq \nu_i$ . This fact combined with the last statement in (3.45) implies that

$$\nu_i \ge S_*^{\frac{N}{2s}}.$$
(3.50)

Hence, exploiting  $(f_4)$ ,  $(g_3)$ ,  $\operatorname{supp}(\psi_{\rho}(\cdot, 0)) \subset \Lambda_{\varepsilon}$ , and  $0 \leq \psi_{\rho} \leq 1$ , we arrive at

$$\begin{split} c &= J_{\varepsilon}(v_n) - \frac{1}{2} \langle J'_{\varepsilon}(v_n), v_n \rangle + o_n(1) \\ &= \int_{\Lambda_{\varepsilon}^{\varepsilon}} \left[ \frac{1}{2} g_{\varepsilon}(x, v_n(x, 0)) v_n(x, 0) - G_{\varepsilon}(x, v_n(x, 0)) \right] \, dx + \int_{\Lambda_{\varepsilon}} \left[ \frac{1}{2} f(v_n(x, 0)) v_n(x, 0) - F(v_n(x, 0)) \right] \, dx \\ &+ \frac{s}{N} \int_{\Lambda_{\varepsilon}} v_n^{2^*_s}(x, 0) \, dx + o_n(1) \\ &\geq \frac{s}{N} \int_{\Lambda_{\varepsilon}} v_n^{2^*_s}(x, 0) \, dx + o_n(1) \\ &\geq \frac{s}{N} \int_{\Lambda_{\varepsilon}} \psi_{\rho}(x, 0) v_n^{2^*_s}(x, 0) \, dx + o_n(1). \end{split}$$

Letting  $n \to \infty$  and using (3.45) and (3.50), we obtain

$$c \ge \frac{s}{N} \sum_{\{i \in I: x_i \in \Lambda_{\varepsilon}\}} \psi_{\rho}(x_i, 0) \nu_i \ge \frac{s}{N} \nu_i \ge \frac{s}{N} S_*^{\frac{N}{2s}},$$

which leads to a contradiction because  $c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$  and  $\zeta \in (0,1)$ . Consequently, (3.43) holds, and we can conclude that  $v_n \to v$  in  $X_{\varepsilon}$ .

In light of Lemmas 3.1 and 3.5, we can apply the mountain pass theorem [4] to infer the next existence result for (3.1).

**Theorem 3.1.** For all  $\varepsilon > 0$ , there exists a nonnegative function  $v_{\varepsilon} \in X_{\varepsilon} \setminus \{0\}$  such that

$$J_{\varepsilon}(v_{\varepsilon}) = c_{\varepsilon}$$
 and  $J'_{\varepsilon}(v_{\varepsilon}) = 0.$  (3.51)

**Remark 3.5.** It is possible to give an alternative proof of Theorem 3.1 without using Proposition 3.1. We outline the details. Let  $(v_n) \subset X_{\varepsilon}$  be a Palais-Smale sequence at the mountain pass level  $c_{\varepsilon}$ . By Lemma 3.3, we know that  $(v_n)$  is bounded in  $X_{\varepsilon}$ . Exploiting  $J_{\varepsilon}(v_n) \to c_{\varepsilon} > 0$  and  $\langle J'_{\varepsilon}(v_n), v_n \rangle = o_n(1)$ , we can argue as in the proof of Lemma 5.1 to prove that there exist  $(z_n) \subset \mathbb{R}^N$  and  $r, \beta > 0$  such that

$$\liminf_{n \to \infty} \int_{B_r(z_n)} v_n^2(x,0) \, dx \ge \beta$$

On the other hand, reasoning as in Remark 3.4, we have that for all R > 0 such that  $\Lambda_{\varepsilon} \subset B_{R/2}$ ,

$$\int_{B_R^c} v_n^2(x,0) \, dx \le \frac{C}{R} + \frac{C}{R^2} + o_n(1).$$

In view of the aforementioned estimates, we deduce that the sequence  $(z_n)$  is bounded in  $\mathbb{R}^N$ . Now, because  $(v_n)$  is bounded in  $X_{\varepsilon}$ , up to a subsequence, we may assume that there exists  $v_{\varepsilon} \in X_{\varepsilon}$  such that  $v_n \rightharpoonup v_{\varepsilon}$  in

 $X_{\varepsilon}, v_n(\cdot, 0) \to v_{\varepsilon}(\cdot, 0)$  in  $L^r_{loc}(\mathbb{R}^N)$  for all  $r \in [1, 2^*_s)$ , and  $v_n(\cdot, 0) \to v_{\varepsilon}(\cdot, 0)$  a.e. in  $\mathbb{R}^N$ . Then, it is easy to check that  $v_{\varepsilon}$  is a critical point of  $J_{\varepsilon}$ . Moreover, we can see that  $v_{\varepsilon} \neq 0$ . In fact, due to the boundedness of  $(z_n)$ , we can find k > 0 such that  $B_r(z_n) \subset B_k$  for all  $n \in \mathbb{N}$ . Consequently,

$$\int_{B_k} v_{\varepsilon}^2(x,0) \, dx = \liminf_{n \to \infty} \int_{B_k} v_n^2(x,0) \, dx \ge \liminf_{n \to \infty} \int_{B_r(z_n)} v_n^2(x,0) \, dx \ge \beta$$

which implies that  $v_{\varepsilon} \neq 0$ . Finally, to verify that  $J_{\varepsilon}(v_{\varepsilon}) = c_{\varepsilon}$ , it suffices to use  $(g_3)$  and Fatou's lemma to get

$$c_{\varepsilon} \leq J_{\varepsilon}(v_{\varepsilon}) - \frac{1}{2} \langle J_{\varepsilon}'(v_{\varepsilon}), v_{\varepsilon} \rangle \leq \liminf_{n \to \infty} \left[ J_{\varepsilon}(v_n) - \frac{1}{2} \langle J_{\varepsilon}'(v_n), v_n \rangle \right] = c_{\varepsilon}.$$

# 4. Autonomous critical problems

Let  $\mu > -V_1(1+\frac{1}{\kappa})$  and introduce the following autonomous problem related to (1.1):

$$\begin{cases} (-\Delta + m^2)^s u + \mu u = f(u) + u^{2^*_s - 1} \text{ in } \mathbb{R}^N, \\ u \in H^s(\mathbb{R}^N), \quad u > 0 \text{ in } \mathbb{R}^N. \end{cases}$$
(4.1)

The extended problem associated with (4.1) is given by:

$$\begin{cases} -\operatorname{div}(y^{1-2s}\nabla v) + m^2 y^{1-2s} v = 0 & \text{in } \mathbb{R}^{N+1}_+, \\ \frac{\partial v}{\partial \nu^{1-2s}} = -\mu v(\cdot, 0) + f(v(\cdot, 0)) + (v^+(\cdot, 0))^{2^*_s - 1} & \text{on } \mathbb{R}^N, \end{cases}$$
(4.2)

and the corresponding energy functional

$$L_{\mu}(v) := \frac{1}{2} \|v\|_{Y_{\mu}}^{2} - \int_{\mathbb{R}^{N}} \left[ F(v(x,0)) + \frac{1}{2_{s}^{*}} (v^{+}(x,0))^{2_{s}^{*}} \right] dx$$

is well defined on  $Y_{\mu} := X^s(\mathbb{R}^{N+1}_+)$  endowed with the norm:

$$\|v\|_{Y_{\mu}} := \left(\|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} + \mu|v(\cdot,0)|_{2}^{2}\right)^{\frac{1}{2}}.$$

To check that  $\|\cdot\|_{Y_{\mu}}$  is a norm equivalent to  $\|\cdot\|_{X^{s}(\mathbb{R}^{N+1}_{+})}$ , one can argue as at pag. 5671 in [6] (observe that  $\mu > -V_{1}(1+\frac{1}{\kappa}) > -m^{2s}$ ). We also note that, by (2.3), for all  $v \in Y_{\mu}$ ,

$$\|v\|_{Y_{\mu}}^{2} \ge \zeta \|v\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2}.$$
(4.3)

Obviously,  $Y_{\mu}$  is a Hilbert space with the inner product

$$\langle v, w \rangle_{Y_{\mu}} := \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla v \cdot \nabla w + m^2 v w) \, dx dy + \mu \int_{\mathbb{R}^N} v(x, 0) w(x, 0) \, dx, \quad \text{ for all } v, w \in Y_{\mu}$$

Denote by  $\mathcal{M}_{\mu}$  the Nehari manifold associated with  $L_{\mu}$ , i.e.,

$$\mathcal{M}_{\mu} := \{ v \in Y_{\mu} : \langle L'_{\mu}(v), v \rangle = 0 \}.$$

As in the previous section, it is easy to verify that  $L_{\mu}$  has a mountain pass geometry [4]. Thus, invoking a variant of the mountain pass theorem without the Palais-Smale condition (see [45, Theorem 2.9]), we can find a Palais-Smale sequence  $(v_n) \subset Y_{\mu}$  at the mountain pass level  $d_{\mu}$  of  $L_{\mu}$  given by:

$$d_{\mu} := \inf_{\gamma \in \Gamma_{\mu}} \max_{t \in [0,1]} L_{\mu}(\gamma(t)),$$

where

$$\Gamma_{\mu} := \{ \gamma \in C([0,1], Y_{\mu}) : \gamma(0) = 0, \, L_{\mu}(\gamma(1)) < 0 \}$$

We stress that  $(v_n)$  is bounded in  $Y_{\mu}$ . In fact, by  $(f_3)$ ,

$$C(1 + \|v_n\|_{Y_{\mu}}) \ge L_{\mu}(v_n) - \frac{1}{\theta} \langle L'_{\mu}(v_n), v_n \rangle$$
  
=  $\left(\frac{1}{2} - \frac{1}{\theta}\right) \|v_n\|_{Y_{\mu}}^2 + \frac{1}{\theta} \int_{\mathbb{R}^N} [f(v_n(x,0))v_n(x,0) - \theta F(v_n(x,0))] \, dx + \left(\frac{1}{\theta} - \frac{1}{2_s^*}\right) |v_n^+(\cdot,0)|_{2_s^*}^2$   
$$\ge \left(\frac{1}{2} - \frac{1}{\theta}\right) \|v_n\|_{Y_{\mu}}^2,$$

which implies the boundedness of  $(v_n)$  in  $Y_{\mu}$ . As in [35, 45], by assumptions on f, we can see that

$$< d_{\mu} = \inf_{v \in \mathcal{M}_{\mu}} L_{\mu}(v) = \inf_{v \in Y_{\mu} \setminus \{0\}} \max_{t \ge 0} L_{\mu}(tv).$$

Arguing as in the proof of Lemma 3.2, it is easy to prove that

$$0 < d_{\mu} < \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}}.$$
 (4.4)

Our claim is to establish the existence of a ground state solution for (4.2). We start by recalling a vanishing Lions-type result.

**Lemma 4.1.** [6, Lemma 3.3] Let  $t \in [2, 2_s^*)$  and R > 0. If  $(v_n) \subset X^s(\mathbb{R}^{N+1}_+)$  is a bounded sequence such that

$$\lim_{n \to \infty} \sup_{z \in \mathbb{R}^N} \int_{B_R(z)} |v_n(x,0)|^t \, dx = 0,$$

then  $v_n(\cdot, 0) \to 0$  in  $L^r(\mathbb{R}^N)$  for all  $r \in (2, 2^*_s)$ .

The next lemma is a critical version of [6, Lemma 3.4].

**Lemma 4.2.** Let  $(v_n) \subset Y_{\mu}$  be a Palais-Smale sequence for  $L_{\mu}$  at the level  $c < \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$  and such that  $v_n \rightharpoonup 0$  in  $Y_{\mu}$ . Then, we have either

(a)  $v_n \to 0$  in  $Y_{\mu}$ , or

(b) there exist a sequence  $(z_n) \subset \mathbb{R}^N$  and constants  $R, \beta > 0$  such that

$$\liminf_{n \to \infty} \int_{B_R(z_n)} v_n^2(x,0) \, dx \ge \beta$$

*Proof.* Assume that (b) does not occur. Therefore, for all R > 0, it holds

$$\lim_{n \to \infty} \sup_{z \in \mathbb{R}^N} \int_{B_R(z)} v_n^2(x,0) \, dx = 0$$

By Lemma 4.1, we know that  $v_n(\cdot, 0) \to 0$  in  $L^r(\mathbb{R}^N)$  for all  $r \in (2, 2^*_s)$ . This fact and  $(f_1)$ - $(f_2)$  imply that

$$\int_{\mathbb{R}^N} f(v_n(x,0))v_n(x,0) \, dx = \int_{\mathbb{R}^N} F(v_n(x,0)) \, dx = o_n(1). \tag{4.5}$$

Exploiting  $\langle L'_{\mu}(v_n), v_n \rangle = o_n(1)$  and (4.5), we see that

$$||v_n||_{Y_{\mu}}^2 = |v_n^+(\cdot, 0)|_{2_s^*}^{2_s^*} + o_n(1)$$

Because  $(v_n)$  is bounded in  $Y_{\mu}$ , we may assume that there exists  $\ell \geq 0$  such that

$$||v_n||_{Y_{\mu}}^2 \to \ell \quad \text{and} \quad |v_n^+(\cdot, 0)|_{2_s^s}^{2_s^s} \to \ell.$$
 (4.6)

Suppose by contradiction that  $\ell > 0$ . By virtue of  $L_{\mu}(v_n) = d_{\mu} + o_n(1)$ , (4.5), and (4.6), we have

$$d_{\mu} + o_{n}(1) = L_{\mu}(v_{n}) = \frac{1}{2} ||v_{n}||_{Y_{\mu}}^{2} - \int_{\mathbb{R}^{N}} F(v_{n}(x,0)) dx - \frac{1}{2_{s}^{*}} |v_{n}^{+}(\cdot,0)|_{2_{s}^{*}}^{2_{s}^{*}}$$
$$= \frac{\ell}{2} - \frac{\ell}{2_{s}^{*}} + o_{n}(1) = \frac{s}{N}\ell + o_{n}(1),$$

i.e.,

$$\ell = \frac{N}{s} d_{\mu}. \tag{4.7}$$

On the other hand, by Theorem 2.1 and (4.3),

$$\zeta S_* |v_n^+(\cdot, 0)|_{2_s^*}^2 \le \zeta S_* |v_n(\cdot, 0)|_{2_s^*}^2 \le \zeta ||v_n||_{X^s(\mathbb{R}^{N+1}_+)}^2 \le ||v_n||_{Y_\mu}^2,$$

and passing to the limit as  $n \to \infty$ , we arrive at

$$\zeta S_* \ell^{\frac{2}{2^*_s}} \le \ell.$$

Taking (4.7) into account, we obtain that  $d_{\mu} \geq \frac{s}{N}(\zeta S_*)^{\frac{N}{2s}}$ , which is impossible in view of (4.4). Then,  $\ell = 0$ , and this completes the proof.

Now we are ready to provide the main result of this section.

**Theorem 4.1.** Let  $\mu > -V_1(1+\frac{1}{\kappa})$ . Then, (4.1) has a ground state solution.

Proof. Since  $L_{\mu}$  has a mountain pass geometry [4], we can find a Palais-Smale sequence  $(v_n) \subset Y_{\mu}$  at the level  $d_{\mu}$ . Hence,  $(v_n)$  is bounded in  $Y_{\mu}$ , and so, up to a subsequence, we may suppose that there exists  $v \in Y_{\mu}$  such that  $v_n \rightharpoonup v$  in  $Y_{\mu}$ . Using the growth assumptions on f and the density of  $C_c^{\infty}(\mathbb{R}^{N+1}_+)$  in  $X^s(\mathbb{R}^{N+1}_+)$ , it is standard to verify that  $\langle L'_{\mu}(v), \varphi \rangle = 0$  for all  $\varphi \in Y_{\mu}$ . If  $v \equiv 0$ , we can apply Lemma 4.2 to deduce that for some sequence  $(z_n) \subset \mathbb{R}^N$ ,  $\bar{v}_n(x,y) := v_n(x + z_n, y)$  is a bounded Palais-Smale sequence at the level  $d_{\mu}$  and  $\bar{v}_n \rightharpoonup \bar{v} \neq 0$  in  $Y_{\mu}$ . Thus,  $\bar{v} \in \mathcal{M}_{\mu}$ . Moreover, by  $(f_3)$  and Fatou's lemma,

$$\begin{split} d_{\mu} &\leq L_{\mu}(\bar{v}) \\ &= L_{\mu}(\bar{v}) - \frac{1}{\theta} \langle L'_{\mu}(\bar{v}), \bar{v} \rangle \\ &= \left(\frac{1}{2} - \frac{1}{\theta}\right) \|\bar{v}\|_{Y_{\mu}}^{2} + \frac{1}{\theta} \int_{\mathbb{R}^{N}} [\bar{v}(x,0)f(\bar{v}(x,0)) - \theta F(\bar{v}(x,0))] \, dx + \left(\frac{1}{\theta} - \frac{1}{2_{s}^{*}}\right) |\bar{v}^{+}(\cdot,0)|_{2_{s}^{*}}^{2_{s}^{*}} \\ &\leq \liminf_{n \to \infty} \left[ \left(\frac{1}{2} - \frac{1}{\theta}\right) \|\bar{v}_{n}\|_{Y_{\mu}}^{2} + \frac{1}{\theta} \int_{\mathbb{R}^{N}} [\bar{v}_{n}(x,0)f(\bar{v}_{n}(x,0)) - \theta F(\bar{v}_{n}(x,0))] \, dx + \left(\frac{1}{\theta} - \frac{1}{2_{s}^{*}}\right) |\bar{v}_{n}^{+}(\cdot,0)|_{2_{s}^{*}}^{2_{s}^{*}} \\ &= \liminf_{n \to \infty} \left[ L_{\mu}(\bar{v}_{n}) - \frac{1}{\theta} \langle L'_{\mu}(\bar{v}_{n}), \bar{v}_{n} \rangle \right] = d_{\mu}, \end{split}$$

and so  $L_{\mu}(\bar{v}) = d_{\mu}$ . When  $v \neq 0$ , as before, we can prove that v is a ground state solution to (4.2). Consequently, for each  $\mu > -V_1(1 + \frac{1}{\kappa})$ , there exists a ground state solution  $w = w_{\mu} \in Y_{\mu} \setminus \{0\}$  such that

$$L_{\mu}(w) = d_{\mu}$$
 and  $L'_{\mu}(w) = 0.$ 

As f(t) = 0 for  $t \leq 0$ , it follows from  $\langle L'_{\mu}(w), w^{-} \rangle = 0$  that  $w \geq 0$  in  $\mathbb{R}^{N+1}_{+}$  and  $w \neq 0$ . A standard Moser iteration argument (see, for instance, [7, Lemma 4.4.1], [8, Lemma 4.1] or Lemma 5.3) shows that  $w(\cdot, 0) \in L^{r}(\mathbb{R}^{N})$  for all  $r \in [2, \infty]$ . According to [6, Corollary 3], we know that  $w(\cdot, 0) \in C^{0,\alpha}(\mathbb{R}^{N})$  for some  $\alpha \in (0, 1)$ . By the weak Harnack inequality [24, Proposition 2], we obtain that  $w(\cdot, 0) > 0$  in  $\mathbb{R}^{N}$ .

We conclude this section by establishing an important relation between  $c_{\varepsilon}$  and  $d_{V(0)} = d_{-V_0}$  (note that  $V(0) = -V_0 > -V_1(1 + \frac{1}{\kappa})$  thanks to  $0 \in M$  and  $V_1 - V_0 \ge 0 > -\frac{V_1}{\kappa}$ ).

**Lemma 4.3.** The numbers  $c_{\varepsilon}$  and  $d_{V(0)}$  verify the following inequality:

$$\limsup_{\varepsilon \to 0} c_{\varepsilon} \le d_{V(0)} < \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}}.$$

Proof. In light of Theorem 4.1, there exists a ground state solution w to (4.2) with  $\mu = V(0)$ . Take  $\eta \in C_c^{\infty}(\mathbb{R})$  such that  $0 \leq \eta \leq 1$ ,  $\eta = 1$  in [-1, 1] and  $\eta = 0$  in  $\mathbb{R} \setminus (-2, 2)$ . Suppose that  $B_2 \subset \Lambda$ . Define  $w_{\varepsilon}(x, y) := \eta(\varepsilon | (x, y) |) w(x, y)$  and note that  $\sup(w_{\varepsilon}(\cdot, 0)) \subset \Lambda_{\varepsilon}$ . It is easy to see that  $w_{\varepsilon} \to w$  in  $X^s(\mathbb{R}^{N+1}_+)$  and that  $L_{V(0)}(w_{\varepsilon}) \to L_{V(0)}(w)$  as  $\varepsilon \to 0$ . On the other hand, by the definition of  $c_{\varepsilon}$ , we have

$$c_{\varepsilon} \leq \max_{t \geq 0} J_{\varepsilon}(tw_{\varepsilon}) = J_{\varepsilon}(t_{\varepsilon}w_{\varepsilon}) = \frac{t_{\varepsilon}^{2}}{2} \|w_{\varepsilon}\|_{\varepsilon}^{2} - \int_{\mathbb{R}^{N}} \left[ F(t_{\varepsilon}w_{\varepsilon}(x,0)) + \frac{1}{2_{s}^{*}}(t_{\varepsilon}w_{\varepsilon}(x,0))^{2_{s}^{*}} \right] dx,$$
(4.8)

for some  $t_{\varepsilon} > 0$ . Using  $w \in \mathcal{M}_{V(0)}$  and  $(f_4)$ , we deduce that  $t_{\varepsilon} \to 1$  as  $\varepsilon \to 0$ . Observe that

$$J_{\varepsilon}(t_{\varepsilon}w_{\varepsilon}) = L_{V(0)}(t_{\varepsilon}w_{\varepsilon}) + \frac{t_{\varepsilon}^2}{2} \int_{\mathbb{R}^N} (V_{\varepsilon}(x) - V(0))w_{\varepsilon}^2(x,0) \, dx.$$

$$\tag{4.9}$$

Then, since  $V_{\varepsilon}(x)$  is bounded on the support of  $w_{\varepsilon}(\cdot, 0)$  and  $V_{\varepsilon}(x) \to V(0)$  as  $\varepsilon \to 0$ , we can exploit the dominated convergence theorem, (4.4), (4.8), and (4.9) to reach the desired conclusion.

**Remark 4.1.** From  $(V_1)$  and  $(g_2)$ , we derive that  $c_{\varepsilon} \geq d_{-V_1} > 0$  for all  $\varepsilon > 0$ .

#### 5. Proof of Theorem 1.1

This section is devoted to the proof of Theorem 1.1. Let us recall that, by Theorem 3.1, for all  $\varepsilon > 0$  there exists a nonnegative mountain pass solution  $v_{\varepsilon}$  to (3.1). We begin with a useful result.

**Lemma 5.1.** There exist  $r, \beta, \varepsilon^* > 0$  and  $(y_{\varepsilon}) \subset \mathbb{R}^N$  such that

$$\int_{B_r(y_{\varepsilon})} v_{\varepsilon}^2(x,0) \, dx \ge \beta, \quad \text{for all } \varepsilon \in (0,\varepsilon^*).$$

*Proof.* On account of (3.51) and the growth conditions on f, we can find  $\alpha > 0$ , independent of  $\varepsilon > 0$ , such that

$$\|v_{\varepsilon}\|_{\varepsilon}^{2} \ge \alpha, \quad \text{for all } \varepsilon > 0.$$
 (5.1)

Let  $(\varepsilon_n) \subset (0,\infty)$  be such that  $\varepsilon_n \to 0$ . Assume, by contradiction, that there exists r > 0 such that

$$\lim_{n \to \infty} \sup_{y \in \mathbb{R}^N} \int_{B_r(y)} v_{\varepsilon_n}^2(x,0) \, dx = 0.$$

By Lemma 4.1, we know that  $v_{\varepsilon_n}(\cdot, 0) \to 0$  in  $L^r(\mathbb{R}^N)$  for all  $r \in (2, 2^*_s)$ . Hence, (3.51) and the growth assumptions on f yield

$$\int_{\mathbb{R}^N} F(v_{\varepsilon_n}(x,0)) \, dx = \int_{\mathbb{R}^N} f(v_{\varepsilon_n}(x,0)) v_{\varepsilon_n}(x,0) \, dx = o_n(1).$$

This implies that

$$\int_{\mathbb{R}^N} G_{\varepsilon_n}(x, v_{\varepsilon_n}(x, 0)) \, dx \le \frac{1}{2^*_s} \int_{\Lambda_{\varepsilon_n} \cup \{v_{\varepsilon_n}(\cdot, 0) \le a\}} v_{\varepsilon_n}^{2^*_s}(x, 0) \, dx + \frac{V_1}{2\kappa} \int_{\Lambda^c_{\varepsilon_n} \cap \{v_{\varepsilon_n}(\cdot, 0) > a\}} v_{\varepsilon_n}^2(x, 0) \, dx + o_n(1) \quad (5.2)$$

and

$$\int_{\mathbb{R}^N} g_{\varepsilon_n}(x, v_{\varepsilon_n}(x, 0)) v_{\varepsilon_n}(x, 0) \, dx = \int_{\Lambda_{\varepsilon_n} \cup \{v_{\varepsilon_n}(\cdot, 0) \le a\}} v_{\varepsilon_n}^{2^*}(x, 0) \, dx + \frac{V_1}{\kappa} \int_{\Lambda_{\varepsilon_n}^c \cap \{v_{\varepsilon_n}(\cdot, 0) > a\}} v_{\varepsilon_n}^2(x, 0) \, dx + o_n(1).$$

$$(5.3)$$

Because of  $\langle J'_{\varepsilon_n}(v_{\varepsilon_n}), v_{\varepsilon_n} \rangle = 0$  and (5.3), we obtain

$$\|v_{\varepsilon_n}\|_{\varepsilon_n}^2 - \frac{V_1}{\kappa} \int_{\Lambda_{\varepsilon_n}^c \cap \{v_{\varepsilon_n}(\cdot,0) > a\}} v_{\varepsilon_n}^2(x,0) \, dx = \int_{\Lambda_{\varepsilon_n} \cup \{v_{\varepsilon_n}(\cdot,0) \le a\}} v_{\varepsilon_n}^{2^*_s}(x,0) \, dx + o_n(1). \tag{5.4}$$

Let  $\ell \geq 0$  be such that

$$\|v_{\varepsilon_n}\|_{\varepsilon_n}^2 - \frac{V_1}{\kappa} \int_{\Lambda_{\varepsilon_n}^c \cap \{v_{\varepsilon_n}(\cdot, 0) > a\}} v_{\varepsilon_n}^2(x, 0) \, dx \to \ell.$$

It is clear that  $\ell > 0$ ; otherwise,  $\|v_{\varepsilon_n}\|_{\varepsilon_n} \to 0$ , and this is impossible due to (5.1) (alternatively, one can observe that  $\|v_{\varepsilon_n}\|_{\varepsilon_n} \to 0$  yields  $c_{\varepsilon_n} = J_{\varepsilon_n}(v_{\varepsilon_n}) \to 0$ , which is a contradiction since Remark 4.1 ensures that  $c_{\varepsilon_n} \ge d_{-V_1} > 0$  for all  $n \in \mathbb{N}$ ). From (5.4), we derive that

$$\int_{\Lambda_{\varepsilon_n} \cup \{v_{\varepsilon_n}(\cdot, 0) \le a\}} v_{\varepsilon_n}^{2^*_s}(x, 0) dx \to \ell.$$

Using  $J_{\varepsilon_n}(v_{\varepsilon_n}) - \frac{1}{2_s^*} \langle J'_{\varepsilon_n}(v_{\varepsilon_n}), v_{\varepsilon_n} \rangle = c_{\varepsilon_n}$ , (5.2), and (5.3), we arrive at

$$\frac{s}{N}\ell \le \liminf_{n \to \infty} c_{\varepsilon_n}.$$
(5.5)

On the other hand, noting that

$$\int_{\mathbb{R}^N} V_{\varepsilon_n}(x) v_{\varepsilon_n}^2(x,0) \, dx - \frac{V_1}{\kappa} \int_{\Lambda_{\varepsilon_n}^c \cap \{v_{\varepsilon_n}(\cdot,0) > a\}} v_{\varepsilon_n}^2(x,0) \, dx \ge -\left(1 + \frac{1}{\kappa}\right) V_1 \int_{\mathbb{R}^N} v_{\varepsilon_n}^2(x,0) \, dx$$
$$\ge -\left(1 + \frac{1}{\kappa}\right) \frac{V_1}{m^{2s}} \|v_{\varepsilon_n}\|_{X^s(\mathbb{R}^{N+1}_+)}^2,$$

by the definitions of  $S_*$  and  $\zeta$ , we see that

$$\|v_{\varepsilon_n}\|_{\varepsilon_n}^2 - \frac{V_1}{\kappa} \int_{\Lambda_{\varepsilon_n}^c \cap \{v_{\varepsilon_n}(\cdot, 0) > a\}} v_{\varepsilon_n}^2(x, 0) \, dx \ge \zeta S_* \left( \int_{\Lambda_{\varepsilon_n} \cup \{v_{\varepsilon_n}(\cdot, 0) \le a\}} v_{\varepsilon_n}^{2^*_s}(x, 0) \, dx \right)^{\frac{2^*}{2^*_s}},$$

and letting  $n \to \infty$ , we infer that

 $\ell \ge \zeta S_* \ell^{\frac{2}{2_s^*}}.\tag{5.6}$ 

Combining (5.5) and (5.6) with the fact that  $\ell > 0$ , we obtain

$$\liminf_{n \to \infty} c_{\varepsilon_n} \ge \frac{s}{N} (\zeta S_*)^{\frac{N}{2s}},$$

which contradicts Lemma 4.3.

**Lemma 5.2.** For each sequence  $(\varepsilon_n)$  such that  $\varepsilon_n \to 0$ , consider the sequence  $(y_{\varepsilon_n}) \subset \mathbb{R}^N$  given in Lemma 5.1. Set  $w_n(x,y) := v_{\varepsilon_n}(x+y_{\varepsilon_n},y)$ . Then, there exist a subsequence of  $(w_n)$ , still denoted by itself, and  $w \in X^s(\mathbb{R}^{N+1}_+) \setminus \{0\}$  such that

$$w_n \to w$$
 in  $X^s(\mathbb{R}^{N+1}_+)$ .

Moreover, there exists  $x_0 \in \Lambda$  such that

$$\varepsilon_n y_{\varepsilon_n} \to x_0 \quad and \quad V(x_0) = -V_0.$$

*Proof.* Hereafter, we denote by  $(y_n)$  and  $(v_n)$ , the sequences  $(y_{\varepsilon_n})$  and  $(v_{\varepsilon_n})$ , respectively. Exploiting (3.51), Lemma 4.3 and (2.10), we can argue as in the proof of Lemma 3.3 to deduce that  $(w_n)$  is bounded in  $X^s(\mathbb{R}^{N+1}_+)$ . Hence, up to a subsequence, there exists  $w \in X^s(\mathbb{R}^{N+1}_+) \setminus \{0\}$  such that, as  $n \to \infty$ ,

$$\begin{cases} w_n \rightharpoonup w & \text{in } X^s(\mathbb{R}^{N+1}_+), \\ w_n(\cdot, 0) \rightarrow w(\cdot, 0) & \text{in } L^r_{loc}(\mathbb{R}^N), \text{ for all } r \in [1, 2^*_s), \\ w_n(\cdot, 0) \rightarrow w(\cdot, 0) & \text{a.e. in } \mathbb{R}^N, \end{cases}$$
(5.7)

and

$$\int_{B_r} w^2(x,0) \, dx \ge \beta > 0, \tag{5.8}$$

where we have used Lemma 5.1. Next, we will show that  $(\varepsilon_n y_n)$  is bounded in  $\mathbb{R}^N$ . To this end, it suffices to prove that

dist
$$(\varepsilon_n y_n, \overline{\Lambda}) \to 0$$
 as  $n \to \infty$ . (5.9)

In fact, if (5.9) does not hold, there exist  $\delta > 0$  and a subsequence of  $(\varepsilon_n y_n)$ , still denoted by itself, such that

 $\operatorname{dist}(\varepsilon_n y_n, \overline{\Lambda}) \geq \delta$ , for all  $n \in \mathbb{N}$ .

Consequently, we can find R > 0 such that  $B_R(\varepsilon_n y_n) \subset \Lambda^c$  for all  $n \in \mathbb{N}$ . Since  $w \ge 0$ , it follows from the definition of  $X^s(\mathbb{R}^{N+1}_+)$  that there exists a nonnegative sequence  $(\psi_j) \subset X^s(\mathbb{R}^{N+1}_+)$  such that  $\psi_j$  has compact support in  $\mathbb{R}^{N+1}_+$  and  $\psi_j \to w$  in  $X^s(\mathbb{R}^{N+1}_+)$  as  $j \to \infty$ . Fix  $j \in \mathbb{N}$ . Inserting  $\psi_j$  into the relation  $\langle J'_{\varepsilon_n}(v_n), \phi \rangle = 0$  for all  $\phi \in X_{\varepsilon_n}$ , we can write

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w_n \cdot \nabla \psi_j + m^2 w_n \psi_j) \, dx dy + \int_{\mathbb{R}^N} V_{\varepsilon_n}(x+y_n) w_n(x,0) \psi_j(x,0) \, dx$$
$$= \int_{\mathbb{R}^N} g_{\varepsilon_n}(x+y_n, w_n(x,0)) \psi_j(x,0) \, dx. \tag{5.10}$$

Note that, by the properties of  $g_{\varepsilon}$ ,

$$\begin{split} &\int_{\mathbb{R}^N} g_{\varepsilon_n}(x+y_n, w_n(x,0))\psi_j(x,0)\,dx \\ &= \int_{B_{\frac{R}{\varepsilon_n}}} g_{\varepsilon_n}(x+y_n, w_n(x,0))\psi_j(x,0)\,dx + \int_{B_{\frac{R}{\varepsilon_n}}} g_{\varepsilon_n}(x+y_n, w_n(x,0))\psi_j(x,0)\,dx \end{split}$$

A FRACTIONAL RELATIVISTIC SCHRÖDINGER EQUATION WITH CRITICAL GROWTH

$$\leq \frac{V_1}{\kappa} \int_{B_{\frac{R}{\varepsilon_n}}} w_n(x,0)\psi_j(x,0) \, dx + \int_{B_{\frac{R}{\varepsilon_n}}^c} [f(w_n(x,0)) + w_n^{2^*_s - 1}(x,0)]\psi_j(x,0) \, dx,$$

which combined with  $(V_1)$  and (5.10) yields

$$\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} (\nabla w_n \cdot \nabla \psi_j + m^2 w_n \psi_j) \, dx \, dy - V_1 \left(1 + \frac{1}{\kappa}\right) \int_{\mathbb{R}^N} w_n(x, 0) \psi_j(x, 0) \, dx$$
  
$$\leq \int_{B_{\frac{R}{\varepsilon_n}}} [f(w_n(x, 0)) + w_n^{2^*_s - 1}(x, 0)] \psi_j(x, 0) \, dx.$$

Recalling that  $\psi_j$  has compact support and exploiting  $\varepsilon_n \to 0$ , (5.7), Theorem 2.1, the growth assumptions on f, we have that, as  $n \to \infty$ ,

$$\int_{B_{\frac{R}{\varepsilon_n}}^c} [f(w_n(x,0)) + w_n^{2^*_s - 1}(x,0)]\psi_j(x,0) \, dx \to 0$$

and

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w_n \cdot \nabla \psi_j + m^2 w_n \psi_j) \, dx dy - V_1 \left(1 + \frac{1}{\kappa}\right) \int_{\mathbb{R}^N} w_n(x,0) \psi_j(x,0) \, dx dx \\ \rightarrow \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w \cdot \nabla \psi_j + m^2 w \psi_j) \, dx dy - V_1 \left(1 + \frac{1}{\kappa}\right) \int_{\mathbb{R}^N} w(x,0) \psi_j(x,0) \, dx.$$

Hence, for all  $j \in \mathbb{N}$ ,

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w \cdot \nabla \psi_j + m^2 w \psi_j) \, dx \, dy - V_1 \left(1 + \frac{1}{\kappa}\right) \int_{\mathbb{R}^N} w(x,0) \psi_j(x,0) \, dx \le 0.$$

and letting  $j \to \infty$ , we obtain

$$\|w\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} - V_{1}\left(1 + \frac{1}{\kappa}\right)|w(\cdot, 0)|_{2}^{2}dx \le 0.$$

Using (2.3) and  $\kappa > \frac{V_1}{m^{2s} - V_1}$ , we arrive at

$$0 \le \left(1 - \frac{V_1}{m^{2s}} \left(1 + \frac{1}{\kappa}\right)\right) \|w\|_{X^s(\mathbb{R}^{N+1}_+)}^2 \le 0,$$

which contradicts (5.8). By virtue of (5.9), there exist a subsequence of  $(\varepsilon_n y_n)$ , still denoted by itself, and  $x_0 \in \overline{\Lambda}$  such that  $\varepsilon_n y_n \to x_0$  as  $n \to \infty$ . Next, we claim that  $x_0 \in \Lambda$ .

By  $(g_2)$  and (5.10), we have that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w_n \cdot \nabla \psi_j + m^2 w_n \psi_j) \, dx \, dy + \int_{\mathbb{R}^N} V_{\varepsilon_n}(x+y_n) w_n(x,0) \psi_j(x,0) \, dx$$
  
$$\leq \int_{\mathbb{R}^N} [f(w_n(x,0)) + w_n^{2^*_s - 1}(x,0)] \psi_j(x,0) \, dx,$$

and taking the limit as  $n \to \infty$ , it follows from (5.7), Theorem 2.1 and the continuity of V that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w \cdot \nabla \psi_j + m^2 w \psi_j) \, dx \, dy + \int_{\mathbb{R}^N} V(x_0) w(x,0) \psi_j(x,0) \, dx$$
  
$$\leq \int_{\mathbb{R}^N} [f(w(x,0)) + w^{2^*_s - 1}(x,0)] \psi_j(x,0) \, dx.$$

Letting  $j \to \infty$ , we find

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s}(|\nabla w|^2 + m^2 w^2) \, dx \, dy + \int_{\mathbb{R}^N} V(x_0) w^2(x,0) \, dx \le \int_{\mathbb{R}^N} \left[ f(w(x,0)) w(x,0) + w^{2s}(x,0) \right] \, dx.$$

Hence, there is  $t_1 \in (0,1)$  such that  $t_1 w \in \mathcal{M}_{V(x_0)}$ . Thus, by Lemma 4.3, we see that

$$d_{V(x_0)} \le L_{V(x_0)}(t_1 w) \le \liminf_{n \to \infty} J_{\varepsilon_n}(v_n) = \liminf_{n \to \infty} c_{\varepsilon_n} \le d_{V(0)}.$$

Therefore,  $d_{V(x_0)} \leq d_{V(0)}$ , which implies that  $V(x_0) \leq V(0) = -V_0$ . Since  $-V_0 = \inf_{x \in \overline{\Lambda}} V(x)$ , we deduce that  $V(x_0) = -V_0$ . Moreover, by  $(V_2)$ ,  $x_0 \notin \partial \Lambda$ , and we can infer that  $x_0 \in \Lambda$ . Now, we aim to show that  $w_n \to w$  in  $X^s(\mathbb{R}^{N+1}_+)$  as  $n \to \infty$ . For this purpose, for all  $n \in \mathbb{N}$  and  $x \in \mathbb{R}^N$ ,

we set

$$\tilde{\Lambda}_n := \frac{\Lambda - \varepsilon_n \, \tilde{y}_n}{\varepsilon_n}$$

and

$$\begin{split} \tilde{\chi}_n^1(x) &:= \begin{cases} 1, & \text{if } x \in \tilde{\Lambda}_n, \\ 0, & \text{if } x \in \tilde{\Lambda}_n^c, \end{cases} \\ \tilde{\chi}_n^2(x) &:= 1 - \tilde{\chi}_n^1(x). \end{split}$$

Let us define the following functions for  $x \in \mathbb{R}^N$  and  $n \in \mathbb{N}$ :

$$\begin{split} h_n^1(x) &:= \left(\frac{1}{2} - \frac{1}{\theta}\right) \left(V_{\varepsilon_n}(x+y_n) + V_1\right) w_n^2(x,0) \tilde{\chi}_n^1(x), \\ h^1(x) &:= \left(\frac{1}{2} - \frac{1}{\theta}\right) \left(V(x_0) + V_1\right) w^2(x,0), \\ h_n^2(x) &:= \left[\left(\frac{1}{2} - \frac{1}{\theta}\right) \left(V_{\varepsilon_n}(x+y_n) + V_1\right) w_n^2(x,0) + \frac{1}{\theta} g_{\varepsilon_n}(x+y_n, w_n(x,0)) w_n(x,0) - G_{\varepsilon_n}(x+y_n, w_n(x,0))\right] \tilde{\chi}_n^2(x), \\ h_n^3(x) &:= \left(\frac{1}{\theta} g_{\varepsilon_n}(x+y_n, w_n(x,0)) w_n(x,0) - G_{\varepsilon_n}(x+y_n, w_n(x,0))\right) \tilde{\chi}_n^1(x) \\ &= \left[\frac{1}{\theta} \left( \left(f(w_n(x,0)) w_n(x,0) + (w_n(x,0))^{2_s^*}\right) - \left(F(w_n(x,0)) + \frac{1}{2_s^*}(w_n(x,0))^{2_s^*}\right)\right) \right] \tilde{\chi}_n^1(x), \\ h^3(x) &:= \frac{1}{\theta} \left( f(w(x,0)) w(x,0) + (w(x,0))^{2_s^*} \right) - \left(F(w(x,0)) + \frac{1}{2_s^*}(w(x,0))^{2_s^*}\right). \end{split}$$

In view of  $(f_3)$ ,  $(g_3)$ ,  $(V_1)$ , and our choice of  $\kappa$ , the aforementioned functions are nonnegative in  $\mathbb{R}^N$ . Furthermore, using the following relations of limits, as  $n \to \infty$ ,

$$w_n(x,0) \to w(x,0)$$
 for a.e.  $x \in \mathbb{R}^N$ ,  
 $\varepsilon_n y_n \to x_0 \in \Lambda$ ,

we deduce that, as  $n \to \infty$ ,

$$\tilde{\chi}_n^1(x) \to 1, \ h_n^1(x) \to h^1(x), \ h_n^2(x) \to 0, \ \text{ and } \ h_n^3(x) \to h^3(x) \ \text{ for a.e. } x \in \mathbb{R}^N.$$

Then, by a direct computation,

$$\begin{aligned} d_{V(0)} &\geq \limsup_{n \to \infty} c_{\varepsilon_n} = \limsup_{n \to \infty} \left( J_{\varepsilon_n}(v_n) - \frac{1}{\theta} \langle J_{\varepsilon_n}'(v_n), v_n \rangle \right) \\ &\geq \limsup_{n \to \infty} \left\{ \left( \frac{1}{2} - \frac{1}{\theta} \right) \left[ \|w_n\|_{X^s(\mathbb{R}^{N+1}_+)}^2 - V_1 |w_n(\cdot, 0)|_2^2 \right] + \int_{\mathbb{R}^N} (h_n^1 + h_n^2 + h_n^3) \, dx \right\} \\ &\geq \liminf_{n \to \infty} \left\{ \left( \frac{1}{2} - \frac{1}{\theta} \right) \left[ \|w_n\|_{X^s(\mathbb{R}^{N+1}_+)}^2 - V_1 |w_n(\cdot, 0)|_2^2 \right] + \int_{\mathbb{R}^N} (h_n^1 + h_n^2 + h_n^3) \, dx \right\} \\ &\geq \left( \frac{1}{2} - \frac{1}{\theta} \right) \left[ \|w\|_{X^s(\mathbb{R}^{N+1}_+)}^2 - V_1 |w(\cdot, 0)|_2^2 \right] + \int_{\mathbb{R}^N} (h^1 + h^3) \, dx = d_{V(0)}. \end{aligned}$$

The aforementioned inequalities yield

$$\lim_{n \to \infty} \|w_n\|_{X^s(\mathbb{R}^{N+1}_+)}^2 - V_1 |w_n(\cdot, 0)|_2^2 = \|w\|_{X^s(\mathbb{R}^{N+1}_+)}^2 - V_1 |w(\cdot, 0)|_2^2$$
(5.11)

and

$$h_n^1 \to h^1, h_n^2 \to 0 \text{ and } h_n^3 \to h^3 \text{ in } L^1(\mathbb{R}^N).$$

Hence,

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} (V_{\varepsilon_n}(x+y_n) + V_1) w_n^2(x,0) \, dx = \int_{\mathbb{R}^N} (V(x_0) + V_1) w^2(x,0) \, dx,$$

and we obtain

$$\lim_{n \to \infty} |w_n(\cdot, 0)|_2^2 = |w(\cdot, 0)|_2^2.$$
(5.12)

Combining (5.11) and (5.12), and recalling that  $X^{s}(\mathbb{R}^{N+1}_{+})$  is a Hilbert space, we conclude that

$$\|w_n - w\|_{X^s(\mathbb{R}^{N+1}_+)} \to 0 \quad \text{as } n \to \infty.$$

Now we use a Moser iteration argument [34] to establish a fundamental  $L^{\infty}$ -estimate.

**Lemma 5.3.** Let  $(w_n)$  be the sequence defined as in Lemma 5.2. Then,  $(w_n(\cdot, 0)) \subset L^{\infty}(\mathbb{R}^N)$ , and there exists C > 0 such that

$$|w_n(\cdot, 0)|_{\infty} \le C, \quad \text{for all } n \in \mathbb{N}.$$

*Proof.* It suffices to argue as in the proof of [6, Lemma 4.1]. However, for the reader's convenience, we provide a different proof here. First, we observe that  $w_n$  is a weak solution to

$$\begin{cases} -\operatorname{div}(y^{1-2s}\nabla w_n) + m^2 y^{1-2s} w_n = 0 & \text{in } \mathbb{R}^{N+1}_+, \\ \frac{\partial w_n}{\partial \nu^{1-2s}} = -V_{\varepsilon_n}(\cdot + y_n) w_n(\cdot, 0) + g_{\varepsilon_n}(\cdot + y_n, w_n(\cdot, 0)) & \text{on } \mathbb{R}^N. \end{cases}$$
(5.13)

For  $\beta > 1$  and T > 0, we consider the following function:

$$H(t) := \begin{cases} 0, & \text{if } t \le 0, \\ t^{\beta}, & \text{if } 0 < t < T, \\ \beta T^{\beta - 1}(t - T) + T^{\beta}, & \text{if } t \ge T. \end{cases}$$

Note that  $H : \mathbb{R} \to \mathbb{R}$  is convex, nondecreasing, and Lipschitz continuous with Lipschitz constant  $\beta T^{\beta-1}$ . Define  $\mathcal{L}(t) := \int_0^t (H'(\tau))^2 d\tau$  for all  $t \in \mathbb{R}$ . Clearly,  $\mathcal{L} \in C^1(\mathbb{R}) \cap W^{1,\infty}(\mathbb{R})$  and  $\mathcal{L}(0) = 0$ . Set

$$\varphi_n(x,y) := \mathcal{L}(w_n(x,y)) = \int_0^{w_n(x,y)} (H'(\tau))^2 d\tau$$

Testing (5.13) with  $\varphi_n$ , we can write

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w_n \cdot \nabla \varphi_n + m^2 w_n \varphi_n) \, dx dy$$
  
=  $-\int_{\mathbb{R}^N} V_{\varepsilon_n}(x+y_n) w_n(x,0) \varphi_n(x,0) \, dx + \int_{\mathbb{R}^N} g_{\varepsilon_n}(x+y_n, w_n(x,0)) \varphi_n(x,0) \, dx.$  (5.14)

By  $(V_1)$ , (3.2), and  $\varphi_n \ge 0$ , we see that

$$\begin{aligned} \left[-V_{\varepsilon_n}(x+y_n)w_n(x,0) + g_{\varepsilon_n}(x+y_n,w_n(\cdot,0))\right]\varphi_n(x,0) &\leq \left[(V_1+\delta)w_n(x,0) + C_{\delta}w_n^{2^*_s-1}(x,0)\right]\varphi_n(x,0) \\ &\leq C(1+w_n^{2^*_s-1}(x,0))\varphi_n(x,0) \\ &\leq C(1+w_n^{2^*_s-1}(x,0))w_n(x,0)(H'(w_n(x,0)))^2, \end{aligned}$$

where we have used  $\varphi_n(x,0) \leq w_n(x,0)(H'(w_n(x,0)))^2$ . Then, from (5.14), we derive that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} (\nabla w_n \cdot \nabla \varphi_n + m^2 w_n \varphi_n) \, dx \, dy \le C \int_{\mathbb{R}^N} (1 + w_n^{2^*_s - 1}(x, 0)) w_n(x, 0) (H'(w_n(x, 0)))^2 \, dx.$$

Since  $H(w_n(x,0))H'(w_n(x,0)) \le \beta^2 w_n^{2\beta-1}(x,0)$  and  $w_n(x,0)H'(w_n(x,0)) \le \beta H(w_n(x,0))$ , we obtain

$$\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} (\nabla w_n \cdot \nabla \varphi_n + m^2 w_n \varphi_n) \, dx \, dy$$
  
$$\leq C\beta \int_{\mathbb{R}^N} (1 + w_n^{2^*_s - 1}(x, 0)) H(w_n(x, 0)) H'(w_n(x, 0)) \, dx$$
  
$$\leq C\beta \int_{\mathbb{R}^N} \left[ \beta^2 w_n^{2\beta - 1}(x, 0) + \beta w_n^{2^*_s - 2}(x, 0) H^2(w_n(x, 0)) \right] \, dx$$

$$\leq C\beta^3 \int_{\mathbb{R}^N} \left[ w_n^{2\beta-1}(x,0) + w_n^{2^*_s - 2}(x,0) H^2(w_n(x,0)) \right] dx.$$
(5.15)

On the other hand, using Theorem 2.1 and  $w_n, \varphi_n \ge 0$ , we obtain

$$\begin{aligned} \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} (\nabla w_n \cdot \nabla \varphi_n + m^2 w_n \varphi_n) \, dx dy &= \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} [|\nabla w_n|^2 (H'(w_n))^2 + m^2 w_n \varphi_n] \, dx dy \\ &\geq \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla w_n|^2 (H'(w_n))^2 \, dx dy \\ &= \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla H(w_n)|^2 \, dx dy \\ &\geq C \left( \int_{\mathbb{R}^N} |H(w_n(x,0))|^{2^*_s} \, dx \right)^{\frac{2}{2^*_s}}. \end{aligned}$$
(5.16)

Combining (5.15) and (5.16), we find

$$\left(\int_{\mathbb{R}^N} |H(w_n(x,0))|^{2^*_s} \, dx\right)^{\frac{2^*}{2^*_s}} \le C\beta^3 \int_{\mathbb{R}^N} \left[ w_n^{2\beta-1}(x,0) + w_n^{2^*_s-2}(x,0)H^2(w_n(x,0)) \right] \, dx,\tag{5.17}$$

where C > 0 is independent of  $\beta$  and T. We stress that the last integral in (5.17) is well defined for every T > 0 in the definition of H. Now we choose  $\beta$  in (5.17) such that  $2\beta - 1 = 2_s^*$ , and we name it  $\beta_1$ , i.e.,

$$\beta_1 := \frac{2_s^* + 1}{2}.\tag{5.18}$$

Let R > 0 to be fixed later. Concerning the last integral in (5.17), applying the Hölder inequality with exponents  $r := \frac{2_s^*}{2}$  and  $r' := \frac{2_s^*}{2_s^*-2}$ , we see that

$$\int_{\mathbb{R}^{N}} w_{n}^{2^{*}_{s}-2}(x,0) H^{2}(w_{n}(x,0)) dx \\
= \int_{\{w_{n}(\cdot,0) \leq R\}} w_{n}^{2^{*}_{s}-2}(x,0) H^{2}(w_{n}(x,0)) dx + \int_{\{w_{n}(\cdot,0) > R\}} w_{n}^{2^{*}_{s}-2}(x,0) H^{2}(w_{n}(x,0)) dx \\
\leq R^{2^{*}_{s}-1} \int_{\{w_{n}(\cdot,0) \leq R\}} \frac{H^{2}(w_{n}(x,0))}{w_{n}(x,0)} dx + \left(\int_{\mathbb{R}^{N}} |H(w_{n}(x,0))|^{2^{*}_{s}} dx\right)^{\frac{2^{*}_{s}}{2^{*}_{s}}} \left(\int_{\{w_{n}(\cdot,0) > R\}} w_{n}^{2^{*}_{s}}(x,0) dx\right)^{\frac{2^{*}_{s}-2}{2^{*}_{s}}}.$$
(5.19)

Because  $(w_n(\cdot, 0))$  strongly converges in  $L^{2^*}(\mathbb{R}^N)$  (by Lemma 5.2), we can take R sufficiently large such that

$$\left(\int_{\{w_n(\cdot,0)>R\}} w_n^{2^*}(x,0) \, dx\right)^{\frac{2^*_s-2}{2^*_s}} \le \frac{1}{2C\beta_1^3},$$

where C is the constant appearing in (5.17). This together with (5.17), (5.18), and (5.19) yields

$$\left(\int_{\mathbb{R}^N} |H(w_n(x,0))|^{2^*_s} dx\right)^{\frac{2^*}{2^*_s}} \le 2C\beta_1^3 \left(\int_{\mathbb{R}^N} w_n^{2^*_s}(x,0) dx + R^{2^*_s - 1} \int_{\mathbb{R}^N} \frac{H^2(w_n(x,0))}{w_n(x,0)} dx\right).$$
(5.20)

In view of  $H(w_n(x,0)) \leq w_n^{\beta_1}(x,0)$  and (5.18), and letting  $T \to \infty$  in (5.20), we obtain

$$\left(\int_{\mathbb{R}^N} w_n^{2^*_s\beta_1}(x,0) \, dx\right)^{\frac{2}{2^*_s}} \le 2C\beta_1^3 \left(\int_{\mathbb{R}^N} w_n^{2^*_s}(x,0) \, dx + R^{2^*_s-1} \int_{\mathbb{R}^N} w_n^{2^*_s}(x,0) \, dx\right),$$

which combined the boundedness of  $(w_n(\cdot, 0))$  in  $L^{2^*_s}(\mathbb{R}^N)$  implies

 $|w_n(\cdot,0)|_{2^*_s\beta_1} \le C', \quad \text{for all } n \in \mathbb{N}.$ (5.21)

28

Now, we suppose  $\beta > \beta_1$ . Thus, using  $H(w_n(x,0)) \le w_n^\beta(x,0)$  on the right-hand side of (5.17) and passing to the limit as  $T \to \infty$ , we deduce that

$$\left(\int_{\mathbb{R}^N} w_n^{2^*_s\beta}(x,0) \, dx\right)^{\frac{2}{2^*_s}} \le C\beta^3 \left(\int_{\mathbb{R}^N} w_n^{2\beta-1}(x,0) \, dx + \int_{\mathbb{R}^N} w_n^{2\beta+2^*_s-2}(x,0) \, dx\right).$$
(5.22)

Put

$$a_1 := \frac{2^*_s(2^*_s - 1)}{2(\beta - 1)}$$
 and  $a_2 := 2\beta - 1 - a_1$ .

Note that  $0 < a_1 < 2_s^*$  and  $a_2 > 0$  (since  $\beta > \beta_1$ ). Applying the Young inequality with exponents  $r := \frac{2_s^*}{a_1}$  and  $r' := \frac{2_s^*}{2_s^* - a_1}$ , we have that

$$\begin{split} \int_{\mathbb{R}^{N}} w_{n}^{2\beta-1}(x,0) \, dx &\leq \frac{a_{1}}{2_{s}^{*}} \int_{\mathbb{R}^{N}} w_{n}^{2_{s}^{*}}(x,0) \, dx + \frac{2_{s}^{*}-a_{1}}{2_{s}^{*}} \int_{\mathbb{R}^{N}} w_{n}^{\frac{2_{s}^{*}-a_{1}}{2_{s}^{*}-a_{1}}}(x,0) \, dx \\ &\leq \int_{\mathbb{R}^{N}} w_{n}^{2_{s}^{*}}(x,0) \, dx + \int_{\mathbb{R}^{N}} w_{n}^{2\beta+2_{s}^{*}-2}(x,0) \, dx \\ &\leq C \left(1 + \int_{\mathbb{R}^{N}} w_{n}^{2\beta+2_{s}^{*}-2}(x,0) \, dx\right), \end{split}$$
(5.23)

with C > 0 independent of  $\beta$  and  $n \in \mathbb{N}$ . Combining (5.22) and (5.23), we get

$$\left(\int_{\mathbb{R}^N} w_n^{2^*_s\beta}(x,0) \, dx\right)^{\frac{2^*}{2^*_s}} \le C\beta^3 \left(1 + \int_{\mathbb{R}^N} w_n^{2\beta+2^*_s-2}(x,0) \, dx\right)$$

with C > 0 changing from line to line, but remaining independent of  $\beta$  and  $n \in \mathbb{N}$ . Therefore,

$$\left(1 + \int_{\mathbb{R}^N} w_n^{2_s^*\beta}(x,0) \, dx\right)^{\frac{1}{2_s^*(\beta-1)}} \le (C\beta^3)^{\frac{1}{2(\beta-1)}} \left(1 + \int_{\mathbb{R}^N} w_n^{2\beta+2_s^*-2}(x,0) \, dx\right)^{\frac{1}{2(\beta-1)}}.$$
(5.24)

For  $k \in \mathbb{N}$ , we define  $\beta_k$  inductively so that  $2\beta_{k+1} + 2_s^* - 2 = 2_s^*\beta_k$ , i.e.,

$$\beta_{k+1} := \left(\frac{2^*_s}{2}\right)^k (\beta_1 - 1) + 1$$

Hence, from (5.24), we obtain

$$\left(1+\int_{\mathbb{R}^N} w_n^{2^*_s\beta_{k+1}}(x,0)\,dx\right)^{\frac{1}{2^*_s(\beta_{k+1}-1)}} \le (C\beta_{k+1}^3)^{\frac{1}{2(\beta_{k+1}-1)}} \left(1+\int_{\mathbb{R}^N} w_n^{2^*_s\beta_k}(x,0)\,dx\right)^{\frac{1}{2^*_s(\beta_k-1)}}$$

Setting

$$A_{k,n} := \left(1 + \int_{\mathbb{R}^N} w_n^{2^*_s \beta_k}(x,0) \, dx\right)^{\frac{1}{2^*_s (\beta_k - 1)}}$$

and

$$C_{k+1} := C\beta_{k+1}^3,$$

we can find a constant  $C_0 > 0$  independent of k such that

$$A_{k+1,n} \le \prod_{j=2}^{k+1} C_j^{\frac{1}{2(\beta_j-1)}} A_{1,n} \le C_0 A_{1,n}, \quad \text{for all } k, n \in \mathbb{N}.$$

Since (5.21) implies that, for some  $A_0 > 0$ ,  $A_{1,n} \leq A_0$  for all  $n \in \mathbb{N}$ , we infer that

 $A_{k+1,n} \leq C_0 A_0$ , for all  $k, n \in \mathbb{N}$ .

Consequently, letting  $k \to \infty$ ,

 $|w_n(\cdot, 0)|_{\infty} \le C$ , for all  $n \in \mathbb{N}$ .

The proof of the lemma is now complete.

**Remark 5.1.** According to [24, Proposition 3], we have that  $w_n \in C^{0,\alpha}_{loc}(\overline{\mathbb{R}^{N+1}_+})$  for all  $n \in \mathbb{N}$ .

Now, we observe that  $w_n(\cdot, 0)$  is a weak solution to

$$(-\Delta + m^2)^s w_n(\cdot, 0) = -V_{\varepsilon_n}(\cdot + y_n) w_n(\cdot, 0) + g_{\varepsilon_n}(\cdot + y_n, w_n(\cdot, 0)) \quad \text{in } \mathbb{R}^N.$$

Fix  $\eta \in (0, m^{2s} - V_1)$ . Using  $(V_1)$  and (3.2), we can deduce that  $w_n(\cdot, 0)$  is a weak subsolution to

$$-\Delta + m^2)^s w_n(\cdot, 0) = (V_1 + \eta) w_n(\cdot, 0) + C_\eta w_n^{2_s^* - 1}(\cdot, 0) =: \mu_n \quad \text{in } \mathbb{R}^N,$$
(5.25)

for some  $C_{\eta} > 0$ . Note that  $\mu_n \ge 0$  in  $\mathbb{R}^N$ . By Lemma 5.3 and interpolation in  $L^r$  spaces, we know that, for all  $r \in [2, \infty)$ ,

$$\mu_n \to \mu := (V_1 + \eta)w(\cdot, 0) + C_\eta w^{2^*_s - 1}(\cdot, 0) \quad \text{in } L^r(\mathbb{R}^N),$$

and  $|\mu_n|_{\infty} \leq C$  for all  $n \in \mathbb{N}$ . Let  $z_n \in H^s(\mathbb{R}^N)$  be the unique solution to

$$(-\Delta + m^2)^s z_n = \mu_n \quad \text{in } \mathbb{R}^N.$$
(5.26)

Then,  $z_n = \mathcal{G}_{2s,m} * \mu_n$ , where  $\mathcal{G}_{2s,m}(x) := (2\pi)^{-\frac{N}{2}} \mathcal{F}^{-1}((|k|^2 + m^2)^{-s})(x)$  is the Bessel kernel with parameter m and  $\mathcal{F}^{-1}$  denotes the inverse Fourier transform. By the scaling property of the Fourier transform, it follows that  $\mathcal{G}_{2s,m}(x) = m^{N-2s} \mathcal{G}_{2s,1}(mx)$ . Exploiting formula (4.1) at pag. 416 in [9] (where  $\mathcal{G}_{2s,1}$  is denoted by  $G_{2s}$ ), we can see that

$$\mathcal{G}_{2s,m}(x) = \frac{1}{2^{\frac{N+2s-2}{2}}\pi^{\frac{N}{2}}\Gamma(s)} m^{\frac{N-2s}{2}} K_{\frac{N-2s}{2}}(m|x|)|x|^{\frac{2s-N}{2}},$$

and it satisfies the following properties (see pag. 416-417 in [9] and pag. 132 in [40], with  $\alpha = 2s$  and m = 1): (G1)  $\mathcal{G}_{2s,m}$  is positive, radially symmetric, and smooth in  $\mathbb{R}^N \setminus \{0\}$ , (G2)  $\mathcal{G}_{2s,m}(x) \leq C(\chi_{B_2}(x)|x|^{2s-N} + \chi_{B_2^c}(x)e^{-c|x|})$  for all  $x \in \mathbb{R}^N$ , for some C, c > 0,

$$(\mathcal{G}3) \ \mathcal{G}_{2s,m} \in L^r(\mathbb{R}^N) \text{ for all } r \in [1, \frac{N}{N-2s})$$

In view of the aforementioned facts, we can prove the next crucial result.

**Lemma 5.4.** The sequence  $(w_n)$  satisfies  $w_n(\cdot, 0) \to 0$  as  $|x| \to \infty$  uniformly in  $n \in \mathbb{N}$ .

*Proof.* We start by showing that  $z_n(x) \to 0$  as  $|x| \to \infty$  uniformly in  $n \in \mathbb{N}$ . Note that, fixed  $\delta \in (0, \frac{1}{2})$ , it holds

$$z_n(x) = (\mathcal{G}_{2s,m} * \mu_n)(x) = \int_{B_{\frac{1}{\delta}}^c(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \,d\xi + \int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \,d\xi.$$
(5.27)

From  $(\mathcal{G}_1)$  and  $(\mathcal{G}_2)$ , we derive that the first integral in (5.27) can be estimated as follows:

$$0 \leq \int_{B_{\frac{1}{\delta}}^{c}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_{n}(\xi) d\xi \leq C|\mu_{n}|_{\infty} \int_{B_{\frac{1}{\delta}}^{c}(x)} e^{-c|x-\xi|} d\xi$$
$$\leq C \int_{\frac{1}{\delta}}^{\infty} e^{-cr} r^{N-1} dr =: CA(\delta) \to 0 \quad \text{as } \delta \to 0.$$
(5.28)

Concerning the second integral in (5.27), we observe that

$$0 \le \int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \, d\xi = \int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)(\mu_n(\xi)-\mu(\xi)) \, d\xi + \int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu(\xi) \, d\xi.$$

Fix  $q \in (1, \min\{\frac{N}{N-2s}, 2\})$  so that q' > 2, where q' is the conjugate exponent of q, i.e.  $\frac{1}{q} + \frac{1}{q'} = 1$ . By means of  $(\mathcal{G}3)$  and Hölder's inequality, we have that

$$\int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \, d\xi \le |\mathcal{G}_{2s,m}|_q |\mu_n - \mu|_{q'} + |\mathcal{G}_{2s,m}|_q |\mu|_{L^{q'}(B_{\frac{1}{\delta}}(x))}$$

Because  $|\mu_n - \mu|_{q'} \to 0$  as  $n \to \infty$  and  $|\mu|_{L^{q'}(B_{\frac{1}{4}}(x))} \to 0$  as  $|x| \to \infty$ , there exist R > 0 and  $n_0 \in \mathbb{N}$  such that

$$\int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \,d\xi \le C\delta \tag{5.29}$$

for all  $n \ge n_0$  and  $|x| \ge R$ . Putting together (5.28) and (5.29), we obtain that

$$\int_{\mathbb{R}^N} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \, d\xi \le C(A(\delta)+\delta) \tag{5.30}$$

for all  $n \ge n_0$  and  $|x| \ge R$ . On the other hand, for each  $n \in \{1, \ldots, n_0 - 1\}$ , there exists  $R_n > 0$  such that  $|\mu_n|_{L^{q'}(B_1(x))} < \delta$  as  $|x| \ge R_n$ . Thus, for  $|x| \ge R_n$ ,

$$\int_{\mathbb{R}^{N}} \mathcal{G}_{2s,m}(x-\xi)\mu_{n}(\xi) d\xi \leq CA(\delta) + \int_{B_{\frac{1}{\delta}}(x)} \mathcal{G}_{2s,m}(x-\xi)\mu_{n}(\xi) d\xi$$

$$\leq CA(\delta) + |\mathcal{G}_{2s,m}|_{q}|\mu_{n}|_{L^{q'}(B_{\frac{1}{\delta}}(x))}$$

$$\leq C(A(\delta) + \delta).$$
(5.31)

Hence, taking  $\bar{R} := \max\{R, R_1, \dots, R_{n_0-1}\}, (5.30), \text{ and } (5.31), \text{ ensure that}$ 

$$\int_{\mathbb{R}^N} \mathcal{G}_{2s,m}(x-\xi)\mu_n(\xi) \,d\xi \le C(A(\delta)+\delta)$$

for  $|x| \geq \overline{R}$ , uniformly in  $n \in \mathbb{N}$ . Letting  $\delta \to 0$ , we reach the desired result for  $z_n$ . In light of (5.25) and (5.26), a simple comparison argument (see [6, Theorem 4.3] with  $\Omega = \mathbb{R}^N$ ) shows that  $0 \leq w_n(\cdot, 0) \leq z_n$  in  $\mathbb{R}^N$ . Consequently,  $w_n(\cdot, 0) \to 0$  as  $|x| \to \infty$  uniformly in  $n \in \mathbb{N}$ .

**Remark 5.2.** An alternative proof of Lemma 5.4 can be established by using the elliptic estimates in [24] (see [6, Lemma 4.2]). For completeness, we give the details. By Remark 5.1, we know that  $w_n \in C_{loc}^{0,\alpha}(\mathbb{R}^{N+1}_+)$ for some  $\alpha \in (0,1)$  independent of n. Thanks to (2.1) and  $w_n \to w$  in  $X^s(\mathbb{R}^{N+1}_+)$ , we deduce that  $w_n \to w$ in  $L^{2\gamma}(\mathbb{R}^{N+1}_+, y^{1-2s})$ , where  $\gamma = 1 + \frac{2}{N-2s}$ . Since  $w_n(\cdot, 0) \to w(\cdot, 0)$  in  $H^s(\mathbb{R}^N)$ , it follows from Lemma 5.3 that  $w_n(\cdot, 0) \to w(\cdot, 0)$  in  $L^r(\mathbb{R}^N)$  for all  $r \in [2, \infty)$ . Now, let  $\bar{x} \in \mathbb{R}^N$  be fixed. Using  $(V_1)$  and the growth assumptions on g, we see that  $w_n$  is a weak subsolution to

$$\begin{cases} -\operatorname{div}(y^{1-2s}\nabla w_n) + m^2 y^{1-2s} w_n = 0 & \text{ in } Q_1(\bar{x},0) := B_1(\bar{x}) \times (0,1) \\ \frac{\partial w_n}{\partial \nu^{1-2s}} = (V_1 + \eta) w_n(\cdot,0) + C_\eta w_n^{2^*_s - 1}(\cdot,0) & \text{ on } B_1(\bar{x}), \end{cases}$$

where  $\eta \in (0, m^{2s} - V_1)$  is fixed. By [24, Proposition 1], we obtain

$$\sup_{Q_{\frac{1}{2}}(\bar{x},0)} w_n(\cdot,0) \le C\left( \|w_n\|_{L^{2\gamma}(Q_1(\bar{x},0),y^{1-2s})} + |w_n^{2s^{-1}}(\cdot,0)|_{L^{q_0}(B_1(\bar{x}))} \right), \text{ for all } n \in \mathbb{N},$$

where  $q_0 > \frac{N}{2s}$  is fixed and C > 0 is a constant independent of  $n \in \mathbb{N}$  and  $\bar{x}$ . Exploiting the strong convergence of  $(w_n)$  in  $L^{2\gamma}(\mathbb{R}^{N+1}_+, y^{1-2s})$  and of  $(w_n(\cdot, 0))$  in  $L^{q_0(2^*_s - 1)}(\mathbb{R}^N)$ , respectively, we infer that  $w_n(\bar{x}, 0) \to 0$  as  $|\bar{x}| \to \infty$  uniformly in  $n \in \mathbb{N}$ , and thus, the claim is proved. In this paper we prefer to give a proof based on the properties of the Bessel kernel  $\mathcal{G}_{2s,m}$  that we believe to be useful for future references.

We also have the following lemma.

**Lemma 5.5.** There exists  $\delta > 0$  such that

$$|w_n(\cdot, 0)|_{\infty} \ge \delta, \quad \text{for all } n \in \mathbb{N}.$$

*Proof.* By Lemma 5.1, there exist  $r, \beta > 0$  and  $n_0 \in \mathbb{N}$  such that

$$\int_{B_r} w_n^2(x,0) \, dx \ge \beta, \quad \text{ for all } n \ge n_0.$$

If by contradiction  $|w_n(\cdot, 0)|_{\infty} \to 0$  as  $n \to \infty$ , then

$$0 < \beta \le \int_{B_r} w_n^2(x,0) \, dx \le |B_r| |w_n(\cdot,0)|_{\infty}^2 \to 0 \quad \text{as } n \to \infty,$$

which of course is absurd.

We are now ready to give the proof of the main result of this paper.

Proof of Theorem 1.1. By Lemma 5.2, up to a subsequence, there exist  $(y_n) \subset \mathbb{R}^N$  and  $w \in X^s(\mathbb{R}^{N+1}_+) \setminus \{0\}$ such that  $w_n(x, y) := v_n(x+y_n, y) \to w$  in  $X^s(\mathbb{R}^{N+1}_+)$  and  $\varepsilon_n y_n \to x_0$  for some  $x_0 \in \Lambda$  such that  $V(x_0) = -V_0$ . From the last limit, we can find r > 0 such that for some subsequence, still denoted by itself, it holds

 $B_r(\varepsilon_n y_n) \subset \Lambda$ , for all  $n \in \mathbb{N}$ .

Hence,

$$B_{\frac{r}{\varepsilon_n}}(y_n) \subset \Lambda_{\varepsilon_n}, \quad \text{for all } n \in \mathbb{N}$$

or equivalently,

$$\Lambda_{\varepsilon_n}^c \subset B_{\frac{r}{\varepsilon_n}}^c(y_n), \quad \text{for all } n \in \mathbb{N}.$$
(5.32)

Now, by Lemma 5.4, there exists R > 0 such that

$$v_n(x,0) < a$$
, for all  $|x| \ge R$  and  $n \in \mathbb{N}$ .

Therefore,  $v_n(x,0) = w_n(x-y_n,0) < a$  for all  $x \in B_R^c(y_n)$  and  $n \in \mathbb{N}$ . On the other hand, there exists  $n_0 \in \mathbb{N}$  such that

$$\Lambda_{\varepsilon_n}^c \subset B_{\varepsilon_n}^c(y_n) \subset B_R^c(y_n), \quad \text{for all } n \ge n_0.$$

Consequently,

$$v_n(x,0) < a$$
, for all  $x \in \Lambda_{\varepsilon_n}^c$  and  $n \ge n_0$ .

Then, there exists  $\varepsilon_0 > 0$  such that, for all  $\varepsilon \in (0, \varepsilon_0)$ , Problem (2.9) admits a solution  $v_{\varepsilon}$ . Invoking the weak Harnack inequality [24, Proposition 2], we conclude that  $v_{\varepsilon}(\cdot, 0) > 0$  in  $\mathbb{R}^N$ .

Next, we study the behavior of the maximum points of solutions to Problem (1.1). Take  $\varepsilon_n \to 0$  and let  $(v_n) \subset X_{\varepsilon_n}$  be a sequence of mountain pass solutions to (3.1). By Lemma 5.2, up to a subsequence, there exist  $(y_n) \subset \mathbb{R}^N$  and  $w \in X^s(\mathbb{R}^{N+1}_+) \setminus \{0\}$  such that  $w_n(x,y) := v_n(x+y_n,y)$  strongly converges to w in  $X^s(\mathbb{R}^{N+1}_+)$  as  $n \to \infty$ . Moreover, there exists  $x_0 \in M$  such that  $\varepsilon_n y_n \to x_0$ . Let now  $q_n$  be a global maximum point of  $w_n(\cdot, 0)$ . Lemma 5.4 and Lemma 5.5 guarantee that there exists  $\overline{R} > 0$  such that  $|q_n| \leq \overline{R}$ for all  $n \in \mathbb{N}$ . Thus,  $x_n := q_n + y_n$  is a global maximum point of  $v_n(\cdot, 0)$ , and  $\varepsilon_n x_n \to x_0 \in M$ . This fact together with the continuity of V produces

$$\lim_{n \to \infty} V(\varepsilon_n \, x_n) = V(x_0) = -V_0$$

Finally, we prove a decay estimate for  $v_n(\cdot, 0)$ . Using Lemma 5.4,  $(f_1)$ , and the definition of g, there exists  $R_0 > 0$  sufficiently large such that

$$g_{\varepsilon_n}(x+y_n, w_n(x,0)) \le \delta w_n(x,0), \quad \text{for all } |x| > R_0 \text{ and } n \in \mathbb{N},$$
(5.33)

where  $\delta \in (0, m^{2s} - V_1)$  is fixed. Arguing as in [6] (see formulas (57) and (58) in [6]), we can find a positive continuous function  $\bar{w} \in H^s(\mathbb{R}^N)$  and  $R_1 > 0$  such that

$$(-\Delta + m^2)^s \bar{w} - (V_1 + \delta) \bar{w} = 0 \quad \text{in } \bar{B}^c_{R_1}$$
(5.34)

and

$$0 < \bar{w}(x) \le Ce^{-c|x|}, \quad \text{for all } x \in \mathbb{R}^N, \tag{5.35}$$

for some C, c > 0. Put  $R_2 := \max\{R_0, R_1\}$ . Thus, by  $(V_1)$  and (5.33), we have that

$$(-\Delta + m^2)^s w_n(\cdot, 0) - (V_1 + \delta) w_n(\cdot, 0) \le 0 \quad \text{in } \bar{B}^c_{R_2}.$$
 (5.36)

Set  $b := \min_{\bar{B}_{R_2}} \bar{w} > 0$  and  $z_n := (\ell + 1)\bar{w} - bw_n(\cdot, 0)$ , where  $\ell := \sup_{n \in \mathbb{N}} |w_n(\cdot, 0)|_{\infty} < \infty$ . Let us observe that  $z_n \ge 0$  in  $\bar{B}_{R_2}$  and that (5.34) and (5.36) yield

$$(-\Delta + m^2)^s z_n - (V_1 + \delta) z_n \ge 0$$
 in  $\bar{B}_{R_2}^c$ 

Because  $V_1 + \delta < m^{2s}$ , we can use a comparison argument (see [6, Theorem 4.3] with  $\Omega = \bar{B}_{R_2}^c$ ) to deduce that  $z_n \geq 0$  in  $\mathbb{R}^N$ . In view of (5.35), there exists  $C_0 > 0$  such that

$$0 \le w_n(x,0) \le C_0 e^{-c|x|}$$
, for all  $x \in \mathbb{R}^N$  and  $n \in \mathbb{N}$ .

Recalling that  $v_n(x,0) = w_n(x-y_n,0)$ , we arrive at

$$v_n(x,0) = w_n(x-y_n,0) \le C_0 e^{-c|x-y_n|}, \quad \text{for all } x \in \mathbb{R}^N \text{ and } n \in \mathbb{N}.$$

This completes the proof of Theorem 1.1.

# 6. Final comments: multiple concentrating solutions to (1.1)

As in [6], if we suppose that the continuous potential  $V : \mathbb{R}^N \to \mathbb{R}$  satisfies the following conditions:  $(V'_1)$  there exists  $V_0 \in (0, m^{2s})$  such that  $-V_0 := \inf_{x \in \mathbb{R}^N} V(x)$ ,  $(V'_1)$  there exists a base of a group of  $A \subseteq \mathbb{R}^N$  such that  $-V_0 := \inf_{x \in \mathbb{R}^N} V(x)$ ,

 $(V_2')$  there exists a bounded open set  $\Lambda \subset \mathbb{R}^N$  such that

$$-V_0 < \min_{x \in \partial \Lambda} V(x) \text{ and } 0 \in M := \{x \in \Lambda : V(x) = -V_0\},\$$

then we obtain the next multiplicity result:

**Theorem 6.1.** Assume that  $(V'_1)$ - $(V'_2)$  and  $(f_1)$ - $(f_4)$  hold. Then, for each  $\delta > 0$  such that

$$M_{\delta} := \{ x \in \mathbb{R}^N : \operatorname{dist}(x, M) \leq \delta \} \subset \Lambda,$$

there exists  $\varepsilon_{\delta} > 0$  such that, for each  $\varepsilon \in (0, \varepsilon_{\delta})$ , Problem (1.1) has at least  $\operatorname{cat}_{M_{\delta}}(M)$  positive solutions. Moreover, if  $u_{\varepsilon}$  denotes one of these solutions and  $x_{\varepsilon}$  is a global maximum point of  $u_{\varepsilon}$ , then we have

$$\lim_{\varepsilon \to 0} V(\varepsilon \, x_{\varepsilon}) = -V_0$$

Since the proof of Theorem 6.1 is similar to the one of [6, Theorem 1.2], we only point out the main differences. For the proof of the critical version of [6, Lemma 5.1], we only need to replace in formula (67) in [6] the term

$$\int_{\Lambda_{\varepsilon}} F(tu_n(x,0)) \, dx$$

by the term

$$\int_{\Lambda_{\varepsilon}} \left[ F(tu_n(x,0)) + \frac{t^{2^*_s}}{2^*_s} (u_n^+(x,0))^{2^*_s} \right] dx,$$

and use the same estimates given in the proof of [6, Lemma 5.1-(iv)]. For the analog of [6, Corollary 1], we exploit Lemma 3.5 instead of [6, Lemma 3.2]. For the proof of the critical version of [6, Lemma 5.4], it suffices to replace in formula (71) in [6] the term

$$\int_{\mathbb{R}^N} F(t_{\varepsilon_n} \eta(|(\varepsilon_n x', 0)|) w(x', 0)) \, dx'$$

by the term

$$\int_{\mathbb{R}^N} F(t_{\varepsilon_n} \eta(|(\varepsilon_n \, x', 0)|) w(x', 0)) + \frac{t_{\varepsilon_n}^{2^*_s}}{2^*_s} (\eta(|(\varepsilon_n \, x', 0)|) w(x', 0))^{2^*_s} \, dx',$$

in formula (72) in [6] we substitute the term

$$\int_{\mathbb{R}^N} f(t_{\varepsilon_n} \eta(|(\varepsilon_n x', 0)|) w(x', 0)) t_{\varepsilon_n} \eta(|(\varepsilon_n x', 0)|) w(x', 0) \, dx$$

with

$$\int_{\mathbb{R}^{N}} [f(t_{\varepsilon_{n}}\eta(|(\varepsilon_{n} x', 0)|)w(x', 0))t_{\varepsilon_{n}}\eta(|(\varepsilon_{n} x', 0)|)w(x', 0) + (t_{\varepsilon_{n}}\eta(|(\varepsilon_{n} x', 0)|)w(x', 0))^{2^{*}_{s}}] dx',$$

and we replace formula (73) in [6] by

$$\begin{split} \|\Psi_{\varepsilon_{n},z_{n}}\|_{\varepsilon_{n}}^{2} &\geq \int_{B_{\frac{\delta}{2}}} \frac{[f(t_{\varepsilon_{n}}w(x',0)) + (t_{\varepsilon_{n}}w(x',0))^{2^{*}_{s}-1}]}{t_{\varepsilon_{n}}w(x',0)} w^{2}(x',0) \, dx' \\ &\geq t_{\varepsilon_{n}}^{2^{*}_{s}-2} \int_{B_{\frac{\delta}{2}}} w^{2^{*}_{s}}(x',0) \, dx' \geq t_{\varepsilon_{n}}^{2^{*}_{s}-2} |B_{\frac{\delta}{2}}| w^{2^{*}_{s}}(\hat{x},0). \end{split}$$

Finally, for the proof of the critical version of [6, Proposition 4], we consider Lemma 4.2 instead of [6, Lemma 3.4], and replace the terms

$$\int_{\mathbb{R}^N \setminus B_{R/\varepsilon_n}} f(v_n(x,0))v_n(x,0) \, dx,$$
$$\int_{\mathbb{R}^N} F(\tilde{v}_n(x,0)) \, dx,$$
$$\int_{\mathbb{R}^N} F(t_n u_n(x,0)) \, dx,$$

by

$$\int_{\mathbb{R}^N \setminus B_{R/\varepsilon_n}} \left[ f(v_n(x,0))v_n(x,0) + (v_n^+(x,0))^{2^*_s} \right] dx,$$
$$\int_{\mathbb{R}^N} \left[ F(\tilde{v}_n(x,0)) + \frac{t_n^{2^*_s}}{2^*_s} (\tilde{v}_n^+(x,0))^{2^*_s} \right] dx,$$
$$\int_{\mathbb{R}^N} \left[ F(t_n u_n(x,0)) + \frac{t_n^{2^*_s}}{2^*_s} (u_n^+(x,0))^{2^*_s} \right] dx,$$

respectively. No additional substantial modifications are necessary to deduce the required multiplicity result.

### References

- R. A. Adams, Sobolev spaces, Pure and Applied Mathematics, Vol. 65. Academic Press, New York-London, 1975. xviii+268 pp.
- [2] C.O. Alves, J.M. do Ó, M.A.S. Souto, Local mountain-pass for a class of elliptic problems in R<sup>N</sup> involving critical growth, Nonlinear Analysis 46 (2001) 495–510.
- [3] C.O. Alves, G.M. Figueiredo, Multiplicity of positive solutions for a quasilinear problem in ℝ<sup>N</sup> via penalization method, Adv. Nonlinear Stud. 5 (2005), no. 4, 551–572.
- [4] A. Ambrosetti, P. H. Rabinowitz, Dual variational methods in critical point theory and applications, J. Functional Analysis 14 (1973), 349–381.
- [5] V. Ambrosio, Ground states solutions for a non-linear equation involving a pseudo-relativistic Schrödinger operator, J. Math. Phys. 57 (2016), no. 5, 051502, 18 pp.
- [6] V. Ambrosio, The nonlinear fractional relativistic Schrödinger equation: existence, multiplicity, decay and concentration results, Discrete Contin. Dyn. Syst. 41 (2021), no. 12, 5659–5705.
- [7] V. Ambrosio, Nonlinear fractional Schrödinger equations in  $\mathbb{R}^N$ , Birkhäuser, 2021.
- [8] V. Ambrosio, On the fractional relativistic Schrödinger operator, J. Differential Equations 308 (2022), 327–368.
- [9] N. Aronszajn, K. T. Smith, Theory of Bessel potentials. I, Ann. Inst. Fourier (Grenoble) 11 (1961), 385-475.
- [10] B. Barrios, E. Colorado, A. de Pablo, U. Sánchez, On some critical problems for the fractional Laplacian operator, J. Differential Equations 252 (2012), no. 11, 6133–6162.
- [11] C. Brändle, E. Colorado, A. de Pablo, U. Sánchez, A concave-convex elliptic problem involving the fractional Laplacian Proc. Roy. Soc. Edinburgh Sect. A 143 (2013), no. 1, 39–71.
- [12] H. Brézis, E. Lieb, A relation between pointwise convergence of functions and convergence of functionals, Proc. Amer. Math. Soc. 88 (1983), no. 3, 486–490.
- [13] C. Bucur, E. Valdinoci, Nonlocal diffusion and applications. Lecture Notes of the Unione Matematica Italiana, 20. Springer, Unione Matematica Italiana, Bologna, 2016. xii+155 pp.
- [14] H. Bueno, O.H. Miyagaki, G.A. Pereira, Remarks about a generalized pseudo-relativistic Hartree equation, J. Differential Equations 266 (2019), no. 1, 876–909.
- [15] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian, Comm. Partial Differential Equations 32 (2007), no. 7-9, 1245–1260
- [16] R. Carmona, W.C. Masters, B. Simon, Relativistic Schrödinger operators: Asymptotic behavior of the eigenfunctions, J. Func. Anal 91 (1990), 117–142.
- [17] V. Coti Zelati, M. Nolasco, Existence of ground states for nonlinear, pseudo-relativistic Schrödinger equations, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. 22 (2011), 51–72.
- [18] A. Cotsiolis, N.K. Tavoularis, Best constants for Sobolev inequalities for higher order fractional derivatives, J. Math. Anal. Appl. 295 (2004), no. 1, 225–236.
- [19] M. del Pino, P.L. Felmer, Local mountain passes for semilinear elliptic problems in unbounded domains, Calc. Var. Partial Differential Equations, 4 (1996), 121–137.

- [20] E. Di Nezza, G. Palatucci, E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. 136 (2012), no. 5, 521–573.
- [21] S. Dipierro, M. Medina, E. Valdinoci, Fractional elliptic problems with critical growth in the whole of R<sup>n</sup>, Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)], 15. Edizioni della Normale, Pisa, 2017. viii+152 pp.
- [22] A. Elgart, B. Schlein, Mean field dynamics of Boson stars, Comm. Pure Appl. Math. 60 (2007), no. 4, 500-545.
- [23] A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, *Higher transcendental functions. Vol. II*, Based on notes left by Harry Bateman. Reprint of the 1953 original. Robert E. Krieger Publishing Co., Inc., Melbourne, Fla., 1981. xviii+396 pp.
- [24] M. M. Fall, V. Felli, Unique continuation properties for relativistic Schrödinger operators with a singular potential, Discrete Contin. Dyn. Syst. 35 (2015), no. 12, 5827–5867.
- [25] G.M. Figueiredo, M. F. Furtado, Positive solutions for a quasilinear Schrödinger equation with critical growth, J. Dynam. Differential Equations 24 (2012), no. 1, 13–28.
- [26] A. Floer, A. Weinstein, Nonspreading wave packets for the cubic Schrödinger equation with a bounded potential, J. Funct. Anal. 69 (1986), no. 3, 397–408.
- [27] J. Fröhlich, B.L.G. Jonsson, E. Lenzmann, Boson stars as solitary waves, Comm. Math. Phys. 274 (2007), no. 1, 1–30.
- [28] L. Hörmander, The analysis of linear partial differential operators. III. Pseudo-differential operators, Reprint of the 1994 edition. Classics in Mathematics. Springer, Berlin, 2007. viii+525 pp.
- [29] E. H. Lieb, M. Loss, Analysis, Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, RI, 1997. xviii+278 pp.
- [30] E. H. Lieb, R. Seiringer, The stability of matter in quantum mechanics, Cambridge University Press, Cambridge, 2010. xvi+293 pp.
- [31] E. H. Lieb, H. T. Yau, The Chandrasekhar theory of stellar collapse as the limit of quantum mechanics, Comm. Math. Phys. 112 (1987), no. 1, 147–174.
- [32] P.L. Lions, The concentration-compactness principle in the calculus of variations. The limit case. I, Rev. Mat. Iberoamericana 1 (1985), 145–201.
- [33] P.L. Lions, The concentration-compactness principle in the calculus of variations. The limit case. II, Rev. Mat. Iberoamericana 1 (1985), 45–121.
- [34] J. Moser, A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations, Comm. Pure Appl. Math., 13 (1960), 457–468.
- [35] P. Rabinowitz, On a class of nonlinear Schrödinger equations Z. Angew. Math. Phys. 43 (1992), no. 2, 270–291.
- [36] M. Ryznar, Estimate of Green function for relativistic  $\alpha$ -stable processes, Potential Analysis, **17**, (2002), 1–23.
- [37] S. Secchi, On some nonlinear fractional equations involving the Bessel potential, J. Dynam. Differential Equations 29 (2017), no. 3, 1173–1193.
- [38] R. Servadei, E. Valdinoci, The Brezis-Nirenberg result for the fractional Laplacian, Trans. Amer. Math. Soc. 367 (2015), no. 1, 67–102.
- [39] L. Shen, Existence result for fractional Schrödinger-Poisson systems involving a Bessel operator without Ambrosetti-Rabinowitz condition, Comput. Math. Appl. 75 (2018), no. 1, 296–306.
- [40] E. Stein, Singular integrals and differentiability properties of functions, Princeton Mathematical Series, No. 30 Princeton University Press, Princeton, N.J. 1970 xiv+290 pp.
- [41] P. R. Stinga, J. L. Torrea, Extension problem and Harnack's inequality for some fractional operators, Comm. Partial Differential Equations 35 (2010), no. 11, 2092–2122.
- [42] X. Wang, On concentration of positive bound states of nonlinear Schrödinger equations, Comm. Math. Phys. 53 (1993), 229–244.
- [43] R. A. Weder, Spectral properties of one-body relativistic spin-zero Hamiltonians, Ann. Inst. H. Poincaré Sect. A (N.S.) 20 (1974), 211–220.
- [44] R. A. Weder, Spectral analysis of pseudodifferential operators, J. Functional Analysis 20 (1975), no. 4, 319–337.
- [45] M. Willem, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications, 24. Birkhäuser Boston, Inc., Boston, MA, 1996. x+162 pp.

Vincenzo Ambrosio

Dipartimento di Ingegneria Industriale e Scienze Matematiche

Università Politecnica delle Marche

VIA BRECCE BIANCHE, 12

60131 Ancona (Italy)

Email address: v.ambrosio@staff.univpm.it