Phenomenological implications of modified loop cosmologies: an overview

Bao-Fei Li ^{1,2*}, Parampreet Singh^{1†}, and Anzhong Wang^{3‡}

¹ Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803, USA

Zhejiang University of Technology, Hangzhou, 310032, China

³ GCAP-CASPER, Department of Physics, Baylor University, Waco, TX, 76798-7316, USA

(Dated: December 15, 2021)

In this paper, we first provide a brief review of the effective dynamics of two recently well-studied models of modified loop quantum cosmologies (mLQCs), which arise from different regularizations of the Hamiltonian constraint and show the robustness of a generic resolution of the big bang singularity, replaced by a quantum bounce due to non-perturbative Planck scale effects. As in loop quantum cosmology (LQC), in these modified models the slow-roll inflation happens generically. We consider the cosmological perturbations following the dressed and hybrid approaches and clarify some subtle issues regarding the ambiguity of the extension of the effective potential of the scalar perturbations across the quantum bounce, and the choice of initial conditions. Both of the modified regularizations yield primordial power spectra that are consistent with current observations for the Starobinsky potential within the framework of either the dressed or the hybrid approach. But differences in primordial power spectra are also identified among the mLQCs and LQC. In particular, for mLQC-I, striking differences arise between the dressed and hybrid approaches in the infrared and oscillatory regimes. While the differences between the two modified models can be attributed to differences in the Planck scale physics, the permissible choices of the initial conditions and the differences between the two perturbation approaches have been reported for the first time. All these differences, due to either the different regularizations or the different perturbation approaches, in principle can be observed in terms of non-Gaussianities.

I. INTRODUCTION

Despite offering a solution to several fundamental and conceptual problems of the standard big bang cosmology, including the flatness, horizon, and exotic-relic problems, the cosmic inflation in the early universe also provides a mechanism to produce density perturbations and primordial gravitational waves (PGWs) [1–3]. The latter arise from quantum fluctuations of spacetimes and produce not only a temperature anisotropy, but also polarizations in the cosmic microwave background (CMB), a smoking gun of PGWs. However, the inflationary paradigm is incomplete without the knowledge of key elements from quantum gravity. First, it is well-known that the cosmic inflation is sensitive to the ultraviolet (UV) physics, and its successes are tightly contingent on the understanding of this UV physics [4–6]. In particular, if the inflationary phase lasts somewhat longer than the minimal period required to solve the above mentioned problems, the length scales we observe today will originate from modes that are smaller than the Planck length during inflation [4]. Then, the treatment of the underlying quantum field theory on a classical spacetime becomes questionable, as now the quantum geometric effects are expected to be large, and the space and time cannot be treated classically any more. This is often referred to as the trans-Planckian problem of cosmological fluctuations 1 .

The second problem of the inflationary paradigm is more severe. It is well known that inflationary spacetimes are past-incomplete because of the big bang singularity [10], with which it is not clear how to impose the initial conditions. This problem gets aggravated for low energy inflation in spatially-closed models which are slightly favored by current observations where the universe encounters a big crunch singularity and lasts only for a few Planck seconds [11].

Another problematic feature of inflation is that one often ignores the pre-inflationary dynamics and sets the Bunch-Davies (BD) vacuum at a very early time. But, it is not clear how such a vacuum state can be realized dynamically in the framework of quantum cosmology [6], considering the fact that a pre-inflationary phase always exists between the Plank and inflation scales, which are about $10^3 \sim 10^{12}$ orders of magnitude difference [2, 6]. While these problems of inflationary paradigm demand a completion from quantum theory of spacetimes, they also open an avenue to overcome one of the main obstacles in the development of quantum gravity, which concerns with the lack of experimental evidences. Thus, understanding inflation in the framework of quantum gravity could offer valuable guidances to the construction of the underlying theory [13–15].

 $^{^2}$ Institute for Theoretical Physics & Cosmology,

^{*}E-mail address: baofeili1@lsu.edu

 $^{^{\}dagger}\textsc{E-mail}$ address: psingh@lsu.edu

 $^{^{\}ddagger}\text{E-mail}$ address: Anzhong_Wang@baylor.edu

¹ It has been conjectured using models in string theory that the trans-Planckian problem might never arise [7], which results on severe constraints on various cosmological models (See [8, 9] for more details).

In particular, when applying the techniques of loop quantum gravity (LQG) to homogeneous and isotropic universe, namely loop quantum cosmology (LQC) [16], it was shown that, purely due to quantum geometric effects, the big bang singularity is generically resolved and replaced by a quantum bounce at which the spacetime curvature becomes Planckian [17–20]. The robustness of the singularity resolution has been shown for a varietv of isotropic and anisotropic spacetimes [21]. Interestingly, there exists a reliable effective spacetime description, which has been used to confirm a generic resolution of all strong curvature singularities [22]. Various phenomenological implications have been studied using this effective spacetime description, whose validity has been verified for isotropic and anisotropic spacetimes [23–27]. For low energy inflation models with a positive spatial curvature, the singularity resolution and a successful onset of inflation for classically inadmissible initial conditions have been demonstrated [28–30].

In the last couple of years, several approaches have been proposed, in order to address the impacts of the quantum geometry on the primordial power spectra. These include the approaches of the deformed algebra [31–33], dressed metric [34–36], and hybrid [37–41] (For a recent discussion about similar ideas in anisotropic Bianchi I LQC spacetimes see Refs. [42–45] and references therein.). In particular, the last two approaches have been widely studied and found that they are all consistent with current cosmological observations [46–59]. In addition, within the framework of the dressed metric approach recently it has been also shown that some anomalies from the CMB data [60–62] can be reconciled purely due to the quantum geometric effects [63, 64].

In addition to the standard LQC, in which the Lorentzian term of the classical Hamiltonian constraint is first expressed in terms of the Euclidean term in the spatially flat Friedmann-Lemaître-Robertson-Walker (FLRW) universe, and then only the quantization of the Euclidean term is considered, the robustness of the singularity resolution with respect to different quantizations of the classical Hamiltonian constraint in the symmetry reduced spacetimes have been extensively studied. Two notable examples are the so-called modified LOC-I (mLOC-I) and modified LQC-II (mLQC-II) models, which were first proposed by Yang, Ding and Ma more than a decade ago [65]. In a recent study, Dapor and Liegener (DL) [66, 67] obtained the expectation values of the Hamiltonian operator in LQG using complexifier coherent states [68], adapted to the spatially flat FLRW universe. Using the non-graph changing regularization of the Hamiltonian advocated by Thiemann [69], DL obtained an effective Hamiltonian constraint, which, to the leading order, agrees with the mLQC-I model first obtained in [65]. Sometimes, this model has also been referred to as the DL model or Thiemann regularized LQC. Strictly speaking, when constructing loops in [66] DL treated the edge length μ as a free parameter, but in [65] it was considered as a specific triad dependent function, the so-called

 $\bar{\mu}$ scheme [19], which is known to be the only possible choice in LQC, and results in physics that is independent from underlying fiducial structures used during quantization, and meanwhile yields a consistent infrared behavior for all matter obeying the weak energy condition [70]. Lately, the studies of [66] have been extended to the $\bar{\mu}$ scheme [71].

In the two modified LQC models, mLQC-I and mLQC-II, since different regularizations of the Lorentzian term were used, the resulting equations become the fourthorder and non-singular quantum difference equations, instead of the second-order difference ones obtained in LQC. In these two models the big bang singularity is also generically resolved and replaced by a quantum bounce. In addition, the inflationary phase can naturally take place with a very high probability [72–76]. In addition, the dynamics in LQC and mLQC-II is qualitatively similar in the whole evolution of the universe, while the one in mLQC-I becomes significantly different from LQC (as well as mLQC-II) in the contracting phase, in which an emergent quasi de Sitter space is present. This implies that the contracting phase in mLQC-I is purely a quantum regime without any classical limit 2 .

An important question now is what are the effects of these models and approaches on the CMB observations. The answer to this question requires the knowledge of how the quantum fluctuations propagate on a quantum spacetime in LQC and modified loop cosmological models. In particular, in the framework of the dressed metric approach the power spectra of the cosmological perturbations for both mLQC-I and mLQC-II models were investigated [78]. In the same framework but restricted only to the mLQC-I model, the power spectra of the cosmological perturbations were studied in [79]. More recently, the hybrid approach was applied to mLQC-I [80–82], for which the time-dependent mass of the perturbations was studied in detail [83]. The primordial scalar power spectra obtained in the three models, LQC, mLQC-I and mLQC-II, were also investigated in the hybrid approach [84], and found that the relative differences in the amplitudes of the power spectra among the three models could be as large as 2% in the UV regime of the spectra, which is relevant to the current observations. Interestingly, in the above work, differences in primordial power spectra were found between the hybrid and dressed metric approaches in the infra-red and oscillatory regimes in mLQC-I.

In this brief review, we shall focus mainly on the states that are sharply peaked along the classical trajectories, so that the description of the "effective" dynamics of the universe becomes available [16], and the questions raised recently in [85] are avoided. This includes the studies of the "effective" dynamics of the homogeneous and isotropic mLQC-I and mLQC-II models, and their cos-

² A similar contracting branch is found in certain anisotropic models in the standard regularization of LQC (see for eg. [77]).

mological perturbations in the framework of the dressed metric and hybrid approaches. We shall first clarify the issue regarding the ambiguities in the extension of the effective potential for the scalar perturbations across the quantum bounce, and then pay particular attention to the differences among the three models, LQC, mLQC-I and mLQC-II, and possible observational signals. It is important to note that initial conditions are another subtle and important issue not only in LOC but also in mLQCs. This includes two parts: (i) when to impose the initial conditions, and (ii) which kind of initial conditions one can impose *consistently*. To clarify this issue, we discuss it at length by showing the (generalized) comoving Hubble radius in each model and in each of the dressed and hybrid approaches. From this analysis, one can see clearly what initial conditions can and cannot be imposed at a chosen initial time.

The outline of this brief overview is as follows. In Sec. II we consider the effective dynamics of mLQC-I and mLOC-II, and discuss some universal features of their dynamics such as the resolution of big bang singularity. In addition, in this section we also show that for states such that the evolution of the homogeneous universe was dominated initially at the bounce by the kinetic energy of the inflaton, that is, $\dot{\phi}_B^2 \gg V(\phi_B)$, the post-bounce evolution between the bounce and the reheating can be always divided universally into three different phases: the bouncing, transition, and slow-roll inflation [cf. Fig. 1]. During each of these phases the expansion factor a(t) and the scalar field $\phi(t)$ can be given analytically. In particular, they are given by Eqs.(2.54)-(2.55) and (2.56)-(2.57)during the bouncing phase for mLQC-I and mLQC-II, respectively. In this same section, the probabilities of the slow-roll inflation is considered, and shown that it occurs generically. This particular consideration is restricted to the quadratic potential, but is expected to be also true for other cases.

In Sec. III, the cosmological perturbations of mLQC-I and mLQC-II are studied. We discuss initial conditions and the subtle issue of the ambiguity in the choice of the variables π_a^{-2} and π_a^{-1} (present in the effective potential), which correspond to the quadratic and linear inverse of the momentum conjugate to the scale factor. In addition, to understand the issue of initial conditions properly, we first introduce the comoving Hubble radius λ_H^2 and then state clearly how this is resolved in GR [cf. Fig. 2], and which are the relevant questions in mLQC-I [cf. Fig. 5] and mLQC-II [cf. Fig. 6]. From these figures it is clear that the BD vacuum cannot be consistently imposed at the bounce ³, as now some modes are inside the (comoving) Hubble radius while others not. However, the fourth-order adiabatic vacuum may be imposed at this moment for both of these two modified LQC models, as that adopted in LQC [36]. In addition, in mLQC-I the de Sitter state given by Eq.(3.7)⁴ can be imposed in the contracting phase as long as t_0 is sufficiently early, so the universe is well inside the de Sitter phase. On the other hand, in mLQC-II and LQC, the BD vacuum can be imposed in the contracting phase as long as t_0 is sufficiently early, so the universe becomes so large that the spacetime curvature is very small, and particle creation is negligible. With these in mind, the power spectra obtained in the three models, mLQC-I, mLQC-II and LQC, within the framework of the dressed metric approach were calculated and compared by imposing the initial conditions in the contracting phase. In particular, the spectra can be universally divided into three regimes, the infrared, intermediate and UV. In the infrared and intermediate regimes, the relative differences in the amplitudes of the spectra can be as large as 100% between mLQC-I and mLQC-II (the same is also true between mLQC-I and LQC), but in the UV regime such differences get dramatically reduced, which is no larger than 0.1%. Since the modes in the UV regime are the relevant ones to the current observations and also their corresponding power spectra are scale-invariant, so these three models are all consistent with observations.

In Sec. IV, the cosmological perturbations of mLQC-I and mLQC-II are studied within the hybrid approach, and the subtleties of the initial conditions are shown in Figs. 8, 9 and 10, where Figs. 8 and 9 are respectively for the quadratic and Starobinsky potentials in mLQC-I. while Fig. 10 is for the Starobinsky potential in mLQC-II. The case with the quadratic potential in mLQC-II is similar to that of mLQC-I, given by Fig. 8. From these figures it is clear that imposing the initial conditions now becomes a more delicate issue, and sensitively depends on the potential $V(\phi)$ of the inflaton field. First, in the cases described by Figs. 8 and 9, all the modes are oscillating during the time $t_i^p < t < t_i$, so one might intend to impose the BD vacuum at the bounce. However, for $t < t_i^p$ the quantity $\Omega_{\rm tot}$ defined in Eq.(3.9) experiences a period during which it is very large and negative. As a result, particle creation is expected not to be negligible during this period. Then, imposing the BD vacuum at the bounce will not account for these effects, and the resulting power spectra shall be quite different from the case, in which in the deep contracting phase $(t \ll t_B)$ the BD vacuum is imposed for mLQC-II and LQC, and the de Sitter state for mLQC-I. On the other hand, in the case described by Fig. 10, even if the BD vacuum is

³ It should be noted that anisotropies rise during the contracting phase and generically dominate the earliest stages of the postbounce of the homogeneous universe [42–45]. So, cautions must be taken, when imposing initial conditions at the bounce.

⁴ To be distinguished from the BD vacuum described by Eq.(3.4) we refer to the state described by Eq.(3.7) as the de Sitter state. The difference between them is due to the term $i/(k\eta)$, which is not negligible in the deep contracting phase of the de Sitter background, as now $|k\eta|$ could be very small. For more details, see the discussions given in Sec. III.A, especially the paragraph after Eq.(3.9).

chosen at the bounce, it may not be quite different from the one imposed in the deep contracting phase, as now in the whole contracting phase all the modes are oscillating, and particle creation is not expected to be important up to the bounce. To compare the results from the three different models, in this section the second-order adiabatic vacuum conditions are chosen in the contracting phase, which is expected not to be much different from the de Sitter state for mLOC-I and the BD vacuum for mLQC-II and LQC, as long as $t_0 \ll t_B$ in all the cases described by Figs. 8 - 10. The ambiguities of the choice of π_a^{-2} and π_a^{-1} also occur in this approach, but as far as the power spectra are concerned, different choices lead to similar conclusions [82]. So, in this section only the so-called prescription A is considered. Then, similar conclusions are obtained in this approach regarding the differences among the amplitudes of the power spectra in the three different models. In particular, the relative differences can be as large as 100% between mLQC-I and mLQC-II/LQC, but in the UV regime such differences are reduced to about 2%. A remarkable feature between the two different approaches is also identified: in the infrared and oscillatory regimes, the power spectrum in mLQC-I is suppressed as compared with its counterpart in LQC in the hybrid approach. On the other hand, in the dressed metric approach, the power spectrum in mLQC-I is largely amplified in the infrared regime where its magnitude is as large as of the Planck scale [78, 79]. The main reason for such differences is that the effective mass in the hybrid approach is strictly positive near the bounce, while it is strictly negative in the dressed metric approach for states that are initially dominated by the kinetic energy of the inflaton [78, 79, 82].

The review is concluded in Sec. V, in which we summarize the main conclusions and point out some open questions for future studies.

II. EFFECTIVE QUANTUM DYNAMICS IN MODIFIED LQCS

To facilitate our following discussions, let us first briefly review the standard process of quantization carried out in LQC, from which one can see clearly the similarities and differences among the three models, LQC, mLQC-I and mLQC-II.

A. Quantum Dynamics of LQC

The key idea of LQC is to use the fundamental variables and quantization techniques of LQG to cosmological spacetimes, by taking advantage of the simplifications that arise from the symmetries of these spacetimes. In the spatially-flat FLRW spacetime,

$$ds^{2} = -N^{2}(t)dt^{2} + q_{ab}(t)dx^{a}dx^{b} \equiv -N^{2}(t)dt^{2} + a^{2}(t)\delta_{ab}dx^{a}dx^{b}, \qquad (2.1)$$

there exists only one degree of freedom, the scale factor a(t), where N(t) is the lapse function and can be freely chosen, given the freedom in reparametrizing t, and $q_{ab}(t)$ denotes the 3-dimensional (3D) spatial metric of the hypersurface t = Constant. In this paper, we shall use the indices a, b, c, \ldots to denote spatial coordinates and i, j, k, \ldots to denote the internal su(2) indices. Repeated indices will represent sum, unless otherwise specified.

In full GR, the gravitational phase space consists of the connection A_a^i and density weighted triad E_i^a . In the present case, the 3D spatial space M has a \mathbb{R}^3 topology, from which we can introduce a fiducial cell \mathcal{V} and restrict all integrations to this cell, in order to avoid some artificial divergences and have a well-defined symplectic structure. Within this cell, we introduce a fiducial flat metric \mathring{q}_{ab} via the relation $q_{ab}(t) = a^2(t) \mathring{q}_{ab}$, and then an associated constant orthogonal triad \mathring{e}_i^a and a cotriad $\mathring{\omega}_a^i$. Then, after symmetry reduction A_a^i and E_i^a are given by,

$$A_a^i = c \, v_o^{-1/3} \, \mathring{\omega}_a^i, \ E_i^a = |p| \, v_o^{-2/3} \sqrt{\mathring{q}} \, \mathring{e}_i^a, \qquad (2.2)$$

where $|p| = v_o^{2/3}a^2$, $\kappa = 8\pi G/c^4$, v_o denotes the volume of the fiducial cell measured by \mathring{q}_{ab} , \mathring{q} is the determinant of \mathring{q}_{ab} , and γ is the Barbero-Immirzi parameter whose value can be set to $\gamma \approx 0.2375$ using black hole thermodynamics in LQG [86]. For classical solutions, symmetry reduced connection c is related to time derivative of scale factor as $c = \gamma \dot{a}$, where an over dot denotes a derivative with respect to t for the choice N = 1.

The physical triad and cotriad are given by $e_i^a = (\text{sgn p}) |p|^{-1/2} v_o^{1/3} \mathring{e}_i^a$ and $\omega_a^i = (\text{sgn p}) |p|^{1/2} v_o^{-1/3} \mathring{\omega}_a^i$, where (sgn p) arises because in connection dynamics the phase space contains triads with both orientations. In the following we choose this orientation to be positive and volume of the fiducial cell to be $v_o = 1$. The variables c and p satisfy the communication relation,

$$\{c, p\} = \frac{\kappa\gamma}{3}.\tag{2.3}$$

Then, the gravitational part of the Hamiltonian is a sum of the Euclidean and Lorentzian terms,

$$\mathcal{H}_{\text{grav}} = \mathcal{H}_{\text{grav}}^{(E)} - (1 + \gamma^2) \mathcal{H}_{\text{grav}}^{(L)}, \qquad (2.4)$$

where, with the choice N = 1, these two terms are given, respectively, by

$$\mathcal{H}_{\text{grav}}^{(E)} = \frac{1}{2\kappa} \int d^3x \,\epsilon_{ijk} F_{ab}^i \frac{E^{aj} E^{bk}}{\sqrt{q}}, \qquad (2.5)$$

$$\mathcal{H}_{\rm grav}^{(L)} = \frac{1}{\kappa} \int \mathrm{d}^3 x \, K_{[a}^j K_{b]}^k \frac{E^{aj} E^{bk}}{\sqrt{q}}, \qquad (2.6)$$

where $q = \det(q_{ab}) = a^6 \ \mathring{q}$, F^k_{ab} is the field strength of the connection A^i_a , and K^i_a is the extrinsic curvature, given, respectively, by

$$F_{ab}^{k} \equiv 2\partial_{[a}A_{b]}^{k} + \epsilon_{ij}{}^{k}A_{a}^{i}A_{b}^{j} = c^{2}\epsilon_{ij}{}^{k}\overset{a}{\omega}_{a}^{i}\overset{b}{\omega}_{b}^{j},$$

$$K_{a}^{i} \equiv K_{ab}e_{i}^{b} = \frac{e_{i}^{b}}{2N}\left(\dot{q}_{ab} - 2D_{(a}N_{b)}\right) = \pm \dot{a}\overset{a}{\omega}_{a}^{i}. \quad (2.7)$$

Upon quantization, ambiguities can arise due to different treatments of the Euclidean and Lorentzian terms in the Hamiltonian constraint. In particular, LQC takes the advantage that in the spatially-flat FLRW universe the Lorentzian part is proportional to the Euclidean part,

$$\mathcal{H}_{\rm grav}^{(L)} = \gamma^{-2} \mathcal{H}_{\rm grav}^{(E)}, \qquad (2.8)$$

so that, when coupled to a massless scalar field, the classical Hamiltonian can be rewritten as [19, 87]

$$\mathcal{H} \equiv \mathcal{H}_{\text{grav}} + \mathcal{H}_{M}$$
$$= -\frac{1}{2\kappa\gamma^{2}} \int d^{3}x \,\epsilon_{ijk} F^{i}_{ab} \frac{E^{aj}E^{bk}}{\sqrt{q}} + \mathcal{H}_{M}, \quad (2.9)$$

where $\mathcal{H}_M = p_{\phi}^2 / (2\sqrt{q})$, with p_{ϕ} being the momentum conjugate of ϕ .

The elementary operators in the standard LQC are the triads⁵ p and elements of the holonomies given by $\widehat{e^{i\mu c/2}}$ of c, where $\overline{\mu} = \sqrt{\Delta l_{pl}^2/|p|}$ with $\Delta \equiv 4\sqrt{3}\pi\gamma$, and Δl_{pl}^2 being the minimum non-zero eigenvalue of the area operator, and the Planck length l_{pl} is defined as $l_{pl} \equiv \sqrt{\hbar G}$. However, it is found that, instead of using the eigenket $|p\rangle$ of the area operator p as the basis, it is more convenient to use the eigenket $|v\rangle$ of the volume operator $\hat{v} (\equiv \operatorname{sgn}(p)|p|^{3/2})$, where

$$\hat{v} |v\rangle = \left(\frac{8\pi\gamma}{6}\right)^{3/2} \frac{|v|}{K} l_{pl}^3 |v\rangle, \quad \widehat{e^{i\bar{\mu}c/2}} |v\rangle = |v+1\rangle,$$
(2.10)

with $K \equiv 2\sqrt{2} \left(3\sqrt{3\sqrt{3}}\right)^{-1}$. Let $\Psi(v, \phi)$ denote the wavefunction in the kinematical Hilbert space of the gravitational field coupled with the scalar field ϕ , we have

$$\hat{\phi}\Psi(v,\phi) = \phi\Psi(v,\phi),$$

$$\hat{p}_{\phi}\Psi(v,\phi) = -i\hbar\frac{\partial}{\partial\phi}\Psi(v,\phi),$$

$$\widehat{p|^{-3/2}}\Psi(v,\phi) = \mathcal{B}(v)\Psi(v,\phi),$$
(2.11)

where

$$\mathcal{B}(v) \equiv \left(\frac{6}{8\pi\gamma l_{pl}^2}\right)^{3/2} B(v),$$

$$B(v) \equiv \left(\frac{3}{2}\right)^3 K|v| \left| |v+1|^{1/3} - |v-1|^{1/3} \right|^3. (2.12)$$

Then, the equation satisfied by selecting the physical states $\hat{\mathcal{H}}\Psi(v,\phi) = 0$ can be cast in the form,

$$\partial_{\phi}^{2}\Psi(v,\phi) = \frac{1}{B(v)} \Big[C^{+}(v)\Psi(v+4,\phi) - C^{o}(v)\Psi(v,\phi) + C^{-}(v)\Psi(v-4,\phi) \Big], \qquad (2.13)$$

where

$$C^{+}(v) \equiv \frac{3\pi KG}{8} |v+2| ||v+1| - |v+3||,$$

$$C^{-}(v) \equiv C^{+}(v-4), \quad C^{o}(v) \equiv C^{+}(v) + C^{-}(v). \quad (2.14)$$

This is the main result of LQC [19], which shows that: (i) It is a second order quantum difference equation with *uniform discreteness in volume*, rather than a simple differential equation, a direct consequence of the discrete nature of loop quantum geometry. (ii) It provides the evolution of the quantum cosmological wavefunction $\Psi(v,\phi)$, in which the scalar field serves as a clock. Thus, once an initial state $\Psi(v, \phi_0)$ is given at the initial moment ϕ_0 , the study of the quantum dynamics of LQC can be carried out. It is found that, instead of a big bang singularity, a quantum bounce is generically produced, a result confirmed through extensive numerical simulations [23–27] and an exactly solvable model [20]. Using this model, one can compute the probability for the quantum bounce which turns out to be unity for an arbitrary superposition of wavefunctions [89].

For the states sharply peaked around a classical solution, we can obtain "effective" Friedmann and Raychaudhuri (FR) equations, by using the geometric quantum mechanics in terms of the expectation values of the operators $(\hat{b}, \hat{v}, \hat{\phi}, \widehat{p_{\phi}})$,

$$\dot{b} = \{b, \mathcal{H}\}, \ \dot{v} = \{v, \mathcal{H}\},$$
 (2.15)

$$\dot{\phi} = \{\phi, \mathcal{H}\}, \ \dot{p}_{\phi} = \{p_{\phi}, \mathcal{H}\},$$
 (2.16)

which take the same forms as their classical ones, but all the quantities now represent their expectation values, $A_I \equiv \left\langle \widehat{A_I} \right\rangle$. Then, it was found that the effective Hamiltonian is given by [19],

$$\mathcal{H}_{\text{eff.}} = -\frac{3}{8\pi\gamma^2\bar{\mu}^2G}|p|^{1/2}\sin^2(\bar{\mu}c) + \frac{1}{2}|p|^{3/2}p_{\phi}^2, \ (2.17)$$

which can also be expressed in terms of v and b via the relations $v = |p|^{3/2}$ and $b = c/\sqrt{|p|}$. Then, from Eqs. (2.15) and (2.16) one can find that the "effective" FR equations are given by,

$$H^2 = \frac{8\pi G}{3} \rho \left(1 - \frac{\rho}{\rho_c}\right), \qquad (2.18)$$

$$\dot{H} = -4\pi G \left(\rho + P\right) \left(1 - \frac{\rho}{\rho_c}\right), \qquad (2.19)$$

where

$$H \equiv \frac{\dot{v}}{3v} = \frac{\dot{a}}{a}, \quad \rho_c \equiv \frac{3}{8\pi\lambda a^2\gamma^2 G},$$

$$\rho \equiv \frac{\mathcal{H}_M}{v}, \quad P \equiv -\frac{\partial\mathcal{H}_M}{\partial v}, \quad (2.20)$$

and $v = v_o a^3$. Since H^2 cannot be negative, from Eq.(2.18) we can see that we must have $\rho \leq \rho_c$, and at $\rho = \rho_c$ we have $H^2 = 0$, that is, a quantum bounce

⁵ For a modification of LQC based on using gauge-covariant fluxes, see [88].

occurs at this moment. When $\rho \ll \rho_c$, the quantum gravity effects are negligible, whereby the classical relativistic limit is obtained.

For a scalar field ϕ with its potential $V(\phi)$, we have

$$\mathcal{H}_M \equiv \mathcal{H}_\phi = v \left[\frac{p_\phi^2}{2v^2} + V(\phi) \right]. \tag{2.21}$$

Then, from Eq.(2.16) we find that

$$\dot{\phi} = \frac{p_{\phi}}{v}, \qquad (2.22)$$

$$\dot{p}_{\phi} = -vV_{,\phi}(\phi).$$
 (2.23)

In the rest of this review, we shall consider only the states that are sharply peaked around a classical solution, so the above "effective" descriptions are valid, and the questions raised recently in [85] are avoided.

B. Effective Dynamics of mLQC-I

As mentioned in the introduction, an important open issue in LQC is its connection with LQG [90]. In particular, in LQC the spacetime symmetry is first imposed (in the classical level), before the quantization process is carried out. However, it is well-known that this is different from the general process of LQG [15], and as a result, different Hamiltonian constraints could be resulted, hence resulting in different Planck scale physics. Though the question of ambiguities in obtaining the Hamiltonian in LQG is still open, based on some rigorous proposals by Thiemann [69], various attempts have been carried out, in order to obtain deeper insights into the question.

One of the first attempts to understand this issue was made in [65], in which the Euclidean and Lorentzian terms given by Eqs.(2.5) and (2.6) are treated differently, by closely following the actual construction of LQG. To be more specific, in the full theory [15], the extrinsic curvature in the Lorentzian term (2.6) can be expressed in terms of the connection and the volume as

$$K_{a}^{i} = \frac{1}{\kappa \gamma^{3}} \{ A_{a}^{i}, \{ \mathcal{H}_{\text{grav}}^{(E)}, V \} \}, \qquad (2.24)$$

which once substituted back into Eq. (2.6) lead to an expression of $\mathcal{H}_{\text{grav}}^{(L)}$ different from that of $\mathcal{H}_{\text{grav}}^{(E)}$ in the standard LQC (see [65] for more details). Correspondingly, one is able to obtain the following "effective" Hamiltonian [65, 72],

$$\mathcal{H}_{\text{eff.}}^{\text{I}} = \frac{3v}{8\pi G\lambda^2} \Big\{ \sin^2(\lambda b) - \frac{(\gamma^2 + 1)\sin^2(2\lambda b)}{4\gamma^2} \Big\} + \mathcal{H}_M.$$
(2.25)

Hence, the Hamilton's equations take the form,

$$\dot{v} = \frac{3v\sin(2\lambda b)}{2\gamma\lambda} \Big\{ (\gamma^2 + 1)\cos(2\lambda b) - \gamma^2 \Big\}, \quad (2.26)$$
$$\dot{b} = \frac{3\sin^2(\lambda b)}{2\gamma\lambda^2} \Big\{ \gamma^2 \sin^2(\lambda b) - \cos^2(\lambda b) \Big\}$$
$$-4\pi G\gamma P, \quad (2.27)$$

where P represents the pressure defined in Eq.(2.20). Once the matter Hamiltonian \mathcal{H}_M is specified, together with the Hamiltonian constraint,

$$\mathcal{H} \approx 0,$$
 (2.28)

Eqs.(2.26) and (2.27) uniquely determine the evolution of the universe. Using the non-graph changing regularization of the Hamiltonian [69], the expectation values of the Hamiltonian operator yield the same "effective" Hamiltonian of Eq.(2.25) to the leading order [66].

It has been shown in detail that the big bang singularity is generically replaced by a quantum bounce when the energy reaches its maximum $\rho_c^{\rm I}$ [65, 66, 72–74], where

$$\rho_c^{\rm I} \equiv \frac{\rho_c}{4(1+\gamma^2)},\tag{2.29}$$

and the universe is asymmetric with respect to the bounce, in contrast to LQC.

To write Eqs.(2.26)-(2.28) in terms of H, ρ and P, it was found that one must distinguish the pre- and postbounce phases [72]. In particular, before the bounce, the modified FR equations take the form [72],

$$H^{2} = \frac{8\pi G \alpha \rho_{\Lambda}}{3} \left(1 - \frac{\rho}{\rho_{c}^{\mathrm{I}}}\right) \left[1 + \left(\frac{1 - 2\gamma^{2} + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}}{4\gamma^{2} \left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right)}\right) \frac{\rho}{\rho_{c}^{\mathrm{I}}}\right], \qquad (2.30)$$
$$\frac{\ddot{a}}{a} = -\frac{4\pi \alpha G}{3} \left(\rho + 3P - 2\rho_{\Lambda}\right) + 4\pi G \alpha P \left(\frac{2 - 3\gamma^{2} + 2\sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}}{\left(1 - 5\gamma^{2}\right) \left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right)}\right) \frac{\rho}{\rho_{c}^{\mathrm{I}}}$$
$$-\frac{4\pi \alpha G \rho}{3} \left[\frac{2\gamma^{2} + 5\gamma^{2} \left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right) - 4 \left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right)^{2}}{\left(1 - 5\gamma^{2}\right) \left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right)^{2}}\right] \frac{\rho}{\rho_{c}^{\mathrm{I}}}, \qquad (2.31)$$

where

$$\alpha \equiv \frac{1-5\gamma^2}{\gamma^2+1}, \quad \rho_{\Lambda} \equiv \frac{\gamma^2 \rho_c}{(1+\gamma^2)(1-5\gamma^2)}.$$
(2.32)

As $\rho \ll \rho_c^{\scriptscriptstyle \rm I}$, Eqs.(2.30) and (2.31) reduce, respectively, to

$$H^2 \approx \frac{8\pi\alpha G}{3} \left(\rho + \rho_{\Lambda}\right),$$
 (2.33)

$$\frac{\ddot{a}}{a} \approx -\frac{4\pi\alpha G}{3} \left(\rho + 3P - 2\rho_{\Lambda}\right). \tag{2.34}$$

These are exactly the FR equations in GR for an ordinary matter field coupled with a positive cosmological constant ρ_{Λ} , and a modified Newton's constant, $G_{\alpha} \equiv \alpha G$. For $\gamma \approx 0.2375$, we have $\rho_{\Lambda} \approx 0.03 \rho_{pl}$, which is of the same order as the one deduced conventionally in quantum field theory for the vacuum energy in our universe. In addition, we also have

$$\left|\frac{G_{\alpha}}{G} - 1\right|_{\gamma \approx 0.2375} \simeq 0.32 > \frac{1}{8}.$$
 (2.35)

Finally, we want to emphasize that the minimal energy density of this branch, for which the Hubble rate vanishes, turns out to be negative which can be shown as $\rho_{\min} = -\frac{3}{8\pi G \lambda^2} \approx -0.023$. As a result, the necessary condition to generate a cyclic universe in mLQC-I is the violation of the weak energy condition which is in contrast to the cyclic universes in LQC where the energy density is always non-negative [91].

In the post-bounce phase $(t > t_B)$, from Eqs.(2.26)-(2.28) we find that [72],

$$H^{2} = \frac{8\pi G\rho}{3} \left(1 - \frac{\rho}{\rho_{c}^{\mathrm{I}}}\right) \left[1 + \frac{\gamma^{2}}{\gamma^{2} + 1} \left(\frac{\sqrt{\rho/\rho_{c}^{\mathrm{I}}}}{1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}}\right)^{2}\right], \qquad (2.36)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + 3P\right) + \frac{4\pi G\rho}{3} \left[\frac{(7\gamma^{2} + 8) - 4\rho/\rho_{c}^{\mathrm{I}} + (5\gamma^{2}14 + 8)\sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}}{(\gamma^{2} + 1)\left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right)^{2}}\right] \frac{\rho}{\rho_{c}^{\mathrm{I}}} + 4\pi GP \left[\frac{3\gamma^{2} + 2 + 2\sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}}{(\gamma^{2} + 1)\left(1 + \sqrt{1 - \rho/\rho_{c}^{\mathrm{I}}}\right)}\right] \frac{\rho}{\rho_{c}^{\mathrm{I}}}, \qquad (2.37)$$

from which we obtain

$$\dot{H} = \frac{4G\pi(P+\rho)}{(1+\gamma^2)} \left(2\gamma^2 + 2\frac{\rho}{\rho_c^{\rm I}} - 3\gamma^2 \sqrt{1-\frac{\rho}{\rho_c^{\rm I}}} - 1\right).$$
(2.38)

Therefore, regardless of the matter content, the superinflation (starting at the bounce) will always end at $\rho = \rho_s$, where

$$\rho_s = \frac{\rho_c^{\rm I}}{8} \left(4 - 8\gamma^2 - 9\gamma^4 + 3\gamma^2 \sqrt{8 + 16\gamma^2 + 9\gamma^4} \right),$$
(2.39)

for which we have $\dot{H}(\rho_s) = 0$.

In the classical limit $\rho/\rho_c^{\rm I} \ll 1$, Eqs. (2.36) and (2.37) reduce, respectively, to

$$H^2 \approx \frac{8\pi G}{3}\rho, \qquad (2.40)$$

$$\frac{\ddot{a}}{a} \approx -\frac{4\pi G}{3} \left(\rho + 3P\right), \qquad (2.41)$$

whereby the standard relativistic cosmology is recovered.

It is remarkable to note that in the pre-bounce phase the limit $\rho/\rho_c^{\rm I} \ll 1$ leads to Eqs.(2.33) and (2.34) with a modified Newtonian constant G_{α} , while in the postbounce the same limits leads to Eqs.(2.40) and (2.41) but now with the precise Newtonian constant G.

C. Effective Dynamics of mLQC-II

In LQG, the fundamental variables for the gravitational sector are the su(2) Ashtekar-Barbero connection A_a^i and the conjugate triad E_i^a . When the Gauss and spatial diffeomorphism constraints are fixed, in the homogeneous and isotropic universe the only relevant constraint is the Hamiltonian constraint, from which we obtain the FR equations, as shown in the previous section. The Hamiltonian in mLQC-II arises from the substitution

$$K_a^i = \frac{A_a^i}{\gamma},\tag{2.42}$$

in the Lorentzian term (2.6). Then, the following effective Hamiltonian is resulted [65],

$$\mathcal{H}_{\text{eff.}}^{\text{II}} = -\frac{3v}{2\pi G\lambda^2 \gamma^2} \sin^2\left(\frac{\lambda b}{2}\right) \left\{1 + \gamma^2 \sin^2\left(\frac{\lambda b}{2}\right)\right\} + \mathcal{H}_M, \qquad (2.43)$$

from which we find that the corresponding Hamilton's equations are given by,

$$\dot{v} = \frac{3v\sin(\lambda b)}{\gamma\lambda} \Big\{ 1 + \gamma^2 - \gamma^2\cos(\lambda b) \Big\}, \qquad (2.44)$$

$$\dot{b} = -\frac{6\sin^2\left(\frac{\lambda D}{2}\right)}{\gamma\lambda^2} \left\{ 1 + \gamma^2 \sin^2\left(\frac{\lambda b}{2}\right) \right\} - 4\pi G\gamma P$$

= $-4\pi G\gamma(\rho + P).$ (2.45)

It can be shown that the corresponding (modified) FR equations now read [73],

$$H^{2} = \frac{8\pi G\rho}{3} \left(1 + \gamma^{2} \frac{\rho}{\rho_{c}} \right) \left(1 - \frac{(\gamma^{2} + 1)\rho}{\Delta^{2}\rho_{c}} \right), \quad (2.46)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + 3P \right) - \frac{4\pi GP\rho}{\Delta\rho_{c}} \left[3 \left(\gamma^{2} + 1 \right) - 2\Delta \right]$$

$$-\frac{4\pi G\rho^{2}}{3\Delta^{2}\rho_{c}} \left[7\gamma^{2} - 1 + \left(5\gamma^{2} - 3 \right) \left(\Delta - 1 \right) - \frac{4\gamma^{2}\rho}{\rho_{c}} \right], \quad (2.47)$$

where $\Delta \equiv 1 + \sqrt{1 + \gamma^2 \rho / \rho_c}$. From these equations we can see that now the quantum bounce occurs when $\rho = \rho_c^{II}$, at which we have H = 0 and $\ddot{a} > 0$, where

$$\rho_c^{\text{II}} = 4(\gamma^2 + 1)\rho_c, \qquad (2.48)$$

which is different from the critical density ρ_c in LQC as well as the one $\rho_c^{\rm I}$ in mLQC-I. Therefore, the big bang singularity is also resolved in this model, and replaced by a quantum bounce at $\rho = \rho_c^{\rm II}$, similar to LQC and mLQC-I, despite the fact that the bounce in each of these models occurs at a different energy density. However, in contrast to mLQC-I, the evolution of the universe is symmetric with respect to the bounce, which is quite similar to the standard LQC model.

In addition, similar to the other two cases, now the bounce is accompanied by a phase of super-inflation, i.e. $\dot{H} > 0$, which ends at $\dot{H}(\rho_s) = 0$, but now ρ_s is given by,

$$\rho_s = \frac{\rho_c}{8\gamma^2} \left(3(\gamma^2 + 1)\sqrt{1 + 2\gamma^2 + 9\gamma^4} + 9\gamma^4 + 10\gamma^2 - 3 \right).$$
(2.49)

For $\gamma = 0.2375$, we find $\rho_s = 0.5132 \rho_c^{\text{II}}$.

When $\rho \ll \rho_c^{\scriptscriptstyle \rm II},$ the modified FR equations (2.46)-(2.47) reduce to,

$$H^2 \approx \frac{8\pi G}{3}\rho, \qquad (2.50)$$

$$\frac{\ddot{a}}{a} \; \approx \; -\frac{4\pi G}{3} \left(\rho + 3P \right), \tag{2.51}$$

which are identical to those given in GR. Therefore, in this model, the classical limit is obtained in both preand post-bounce when the energy density ρ is much lower than the critical one ρ_c^{II} .

D. Universal Properties of mLQC-I/II Models

To study further the evolution of the universe, it is necessary to specify the matter content \mathcal{H}_M . For a single scalar field with its potential $V(\phi)$, the corresponding Hamiltonian takes the form (2.21). As a result, the Hamilton's equations of the matter sector are given by Eqs.(2.22) and (2.23).

The effective quantum dynamics of LQC, mLQC-I, and mLQC-II were studied in detail recently in [73] for a single scalar field with various potentials, including the chaotic inflation, Starobinsky inflation, fractional monodromy inflation, non-minimal Higgs inflation, and inflation with an exponential potential, by using dynamical system analysis. It was found that, while several features of LQC were shared by the mLQC-I and mLQC-II models, others belong to particular models. In particular, in the pre-bounce phase, the qualitative dynamics of LQC and mLQC-II are quite similar, but are strikingly different from that of mLQC-I. In all the three models, the non-perturbative quantum gravitational effects always result in a non-singular post-bounce phase, in which a short period of super-inflation always exists right after the bounce, and is succeeded by the conventional inflation. The latter is an attractor in the phase space for all the three models.

Moreover, similar to LQC [51, 52]⁶, for the initially kinetic energy dominated conditions,

$$\frac{1}{2}\dot{\phi}_B^2 \gg V(\phi_B),\tag{2.52}$$

it was found that the evolution of the universe before the reheating is universal. In particular, in the post-bounce phase (between the quantum bounce and the reheating), the evolution can be uniquely divided into three phases: *bouncing, transition and slow-roll inflation*, as shown in Fig. 1 for the Starobinsky potential,

$$V(\phi) = \frac{3m^2}{32\pi G} \left(1 - e^{-\sqrt{16\pi G/3}\phi}\right)^2.$$
 (2.53)

For other potentials, similar results can be obtained, as long as at the bounce the evolution of the universe is dominated by the kinetic energy of the inflaton $w(\phi_B) \simeq$ 1 [74, 102].

In each of these three phases, the evolutions of a(t) and $\phi(t)$ can be well approximated by analytical solutions. In

⁶ In LQC, this universality was first found for the quadratic and Starobinsky potentials [51, 52] (see also [92]), and later was shown that they are also true for other potentials, including the power-law potentials [93, 94], α-attractor potentials [95, 96], Monodromy potentials [97], warm inflation [98], Tachyonic inflation [99] and even in Brans-Dicke LQC [100, 101].



FIG. 1: The evolution of the scale factor a(t), the scalar field $\phi(t)$, and the equation w_{ϕ} of state of the scalar field (from top to bottom) in the post-bounce phase are depicted and compared among the three modes, LQC (red solid curves), mLQC-I (blue dotted curves) and mLQC-II (green dot-dashed curves), with the Starobinsky potential. In the last panel, w_{ϕ} is defined via $w_{\phi} \equiv P(\phi)/\rho(\phi) = [\dot{\phi}^2 - 2V(\phi)]/[\dot{\phi}^2 + 2V(\phi)]$. The initial condition for the simulation is chosen at the bounce with $\phi_B = -1.6 \ m_{pl}, \dot{\phi}_B > 0$ [74].

particular, during the bouncing phase, they are given by

$$a(t) = \left[1 + 24\pi G \rho_{c}^{\mathrm{I}} \left(1 + \frac{A\gamma^{2}}{1 + Bt}\right) t^{2}\right]^{1/6},$$

$$\phi(t) = \phi_{\mathrm{B}} \pm \frac{m_{pl} \operatorname{arcsinh} \left(\sqrt{24\pi G \rho_{c}^{\mathrm{I}} \left(1 + \frac{C\gamma^{2}}{1 + Dt}\right)}t\right)}{\sqrt{12\pi G \left(1 + \frac{C\gamma^{2}}{1 + Dt}\right)}},$$
(2.54)

for mLQC-I model, where the parameters A, B, C and D are fixed through numerical simulations. It was found

that the best fitting is provided by [74],

$$A = C = 1.2, \quad B = 6, \quad D = 2.$$
 (2.55)

For the mLQC-II model, during the bouncing phase a(t) and $\phi(t)$ are given by

$$a(t) = \left[1 + 24\pi G\rho_c^{\text{II}} \left(1 + \frac{A\gamma^2}{1+Bt}\right) t^2\right]^{1/6},$$

$$\phi(t) = \phi_{\text{B}} \pm \frac{m_{pl} \operatorname{arcsinh}\left(\sqrt{24\pi G\rho_c^{\text{II}} \left(1 + \frac{C\gamma^2}{1+Dt}\right)}t\right)}{\sqrt{12\pi G\left(1 + \frac{C\gamma^2}{1+Dt}\right)}},$$
(2.56)

but now with,

$$A = 2.5, \quad B = 7, \quad C = D = 2.$$
 (2.57)

In the transition and slow-roll inflationary phases, the functions a(t) and $\phi(t)$ were given explicitly in [74].

For the initially potential energy dominated cases,

$$\frac{1}{2}\dot{\phi}_B^2 \ll V(\phi_B),\tag{2.58}$$

it was found that such universalities are lost. In particular, for the Starobinsky potential, the potential energy dominated bounce cannot give rise to any period of inflation for both mLQC-I and mLQC-II models, quite similar to what happens in LQC [47, 48].

E. Probabilities of the Slow-roll Inflation in mLQC-I/II Models

To consider the probability of the slow-roll inflation in the modified LQC models, let us start with the phase space S of the modified Friedmann and Klein-Gordon equations, which now is four-dimensional (4D), and consists of the four variables, (v, b) and (ϕ, p_{ϕ}) , from the gravitational and matter sectors, respectively. Using the canonical commutation relations, the symplectic form on the 4D phase space is given by [103–110],

$$\Omega = dp_{\phi} \wedge d\phi + \frac{dv \wedge db}{4\pi G\gamma}.$$
(2.59)

However, after taking the effective Hamiltonian constraint into account,

$$\mathcal{C} = 16\pi G \left\{ \mathcal{H}_{grav}(v,b) + \frac{p_{\phi}^2}{2v} + vV(\phi) \right\} \simeq 0, \quad (2.60)$$

where " \simeq " means that the equality holds only on $\overline{\Gamma}$, we can see that the 4D phase space \mathbb{S} reduces to a threedimensional (3D) hypersurface $\overline{\Gamma}$.

On the other hand, the phase space S is isomorphic to a 2-dimensional (2D) gauge-fixed surface $\hat{\Gamma}$ of $\bar{\Gamma}$, which is intersected by each dynamical trajectory once and only once [105]. From the FR equations, it can be shown that for both mLQC-I and mLQC-II the variable b satisfies the equation [72–74],

$$\dot{b} = -4\pi G\gamma(\rho + P). \tag{2.61}$$

For any given matter field that satisfies the weak energy condition [111], we have $\rho + P > 0$, so the function b is monotonically decreasing. Then, a natural parameterization of this 2D surface is b = constant, say, b_0 . Hence, using the Hamiltonian constraint (2.60) we find

$$p_{\phi}^{\mathcal{A}} = v \left\{ -2 \left[\hat{\mathcal{H}}_{grav}^{\mathcal{A}} + V(\phi) \right] \right\}^{1/2}, \qquad (2.62)$$

where A = I, II, and

$$\hat{\mathcal{H}}_{grav}^{A} \equiv v^{-1} \mathcal{H}_{grav}^{A}(v, b_0).$$
(2.63)

On the other hand, from Eqs.(2.25) and (2.43) we find that $\hat{\mathcal{H}}_{grav}^{A} = \hat{\mathcal{H}}_{grav}^{A}(b_0) = \text{constant on }\hat{\Gamma}$. Thus, we find

$$dp_{\phi}^{A}|_{\hat{\Gamma}} = \frac{p_{\phi}^{A}}{v}dv - \frac{v^{2}V_{,\phi}}{p_{\phi}}d\phi.$$
 (2.64)

Inserting this expression into Eq.(2.59), we find that the pulled-back symplectic structure $\hat{\Omega}$ reads

$$\hat{\Omega}^{\mathcal{A}}\Big|_{\hat{\Gamma}} = \left\{-2\left[\hat{\mathcal{H}}_{grav}^{\mathcal{A}}(b_0) + V(\phi)\right]\right\}^{1/2} d\phi \wedge dv, \ (2.65)$$

from which we find that the Liouville measure $d\hat{\mu}_L$ on $\hat{\Gamma}$ is given by

$$d\hat{\mu}_L^{\mathcal{A}} = \left\{ -2 \left[\hat{\mathcal{H}}_{grav}^{\mathcal{A}}(b_0) + V(\phi) \right] \right\}^{1/2} d\phi dv. \quad (2.66)$$

Note that $d\hat{\mu}_L^A$ does not depend on v, so that the integral with respect to dv is infinite. However, this divergency shall be cancelled when calculating the probability, as it will appear in both denominator and numerator. Therefore, the measure for the space of physically distinct solutions can be finally taken as

$$d\omega^{A} = \left\{-2\left[\hat{\mathcal{H}}_{grav}^{A}(b_{0}) + V(\phi)\right]\right\}^{1/2} d\phi, \qquad (2.67)$$

so that the 2D phase space $\hat{\Gamma}$ is further reduced to an interval $\phi \in (\phi_{\min}, \phi_{\max})$. It should be noted that such a defined measure depends explicitly on b_0 , and its choice in principle is arbitrary. However, in loop cosmology there exists a preferred choice, which is its value at the quantum bounce $b_0 = b(t_B)$ [105]. With such a choice, the probability of the occurrence of an event E becomes

$$P(E) = \frac{1}{\mathcal{D}} \int_{\mathcal{I}(E)} \left\{ -2 \left[\hat{\mathcal{H}}_{grav}^{\mathrm{A}}(b_B) + V(\phi) \right] \right\}^{1/2} d\phi,$$
(2.68)

where $\mathcal{I}(E)$ is the interval on the ϕ_B -axis, which corresponds to the physically distinct initial conditions in which the event E happens, and \mathcal{D} is the total measure

$$\mathcal{D} \equiv \int_{\phi_{\min}}^{\phi_{\max}} \left\{ -2 \left[\hat{\mathcal{H}}_{grav}^{A}(b_B) + V(\phi) \right] \right\}^{1/2} d\phi. \quad (2.69)$$

Once the probability is properly defined, we can calculate it in different models of the modified LQCs. In LQC [105], the calculations were carried out for the quadratic potential. In order to compare the results obtained in different models, let us consider the same potential. Then, for the mLQC-I model it was found that [74]

$$\sin(\lambda b_B^{\rm I}) = \sqrt{\frac{1}{2\gamma^2 + 2}}, \quad \sin(2\lambda b_B^{\rm I}) = \frac{\sqrt{2\gamma^2 + 1}}{\gamma^2 + 1},$$
$$p_{\phi}^{\rm I} = v \left(2\rho_c^{\rm I} - 2V\right)^{\frac{1}{2}}, \quad d\omega^{\rm I} = \left(2\rho_c^{\rm I} - 2V\right)^{\frac{1}{2}} d\phi, \quad (2.70)$$

so that the probability for the desired slow-roll not to happen is,

$$P^{\rm I}(\text{not realized}) \lesssim \frac{\int_{-5.158}^{0.917} d\omega^{\rm I}}{\int_{-\phi_{\rm max}}^{\phi_{\rm max}^{\rm I}} d\omega^{\rm I}} \simeq 1.12 \times 10^{-5}, \quad (2.71)$$

where $\phi_{\max}^{I} = 3.49 \times 10^{5} m_{pl}$.

In mLQC-II, following a similar analysis, it can be shown that the probability for the desired slow-roll not to happen is [74],

$$P^{\text{II}}(\text{not realized}) \lesssim 2.62 \times 10^{-6}.$$
 (2.72)

Note that in LQC the probability for the desired slow roll inflation not to occur is [105],

$$P^{\text{LQC}}(\text{not realized}) \lesssim 2.74 \times 10^{-6},$$
 (2.73)

which is smaller than that for mLQC-I and slightly larger than the one for mLQC-II. However, it is clear that the desired slow-roll inflation is very likely to occur in all the models, including the two modified LQC ones.

III. PRIMORDIAL POWER SPECTRA OF MODIFIED LQCS IN DRESSED METRIC APPROACH

As mentioned above, in the literature there exist several approaches to investigate the inhomogeneities of the universe. Such approaches can be generalized to the modified LQC models, including mLQC-I and mLQC-II. In this section we shall focus ourselves on cosmological perturbations in the framework of mLQCs following the dressed metric approach [34–36], while in the next section we will be following the hybrid approach [37–41]. We shall restrict ourselves to the effective dynamics, as we did with the homogeneous background in the last section. Such investigations in general include two parts: (a) the initial conditions; and (b) the dynamical evolutions of perturbations. In the framework of effective dynamics, the latter is a second-order ordinary differential equation in the momentum space, so in principle once the initial conditions are given, it uniquely determines the cosmological (scalar and tensor) perturbations.

However, the initial conditions are a subtle issue, which is true not only in LQC but also in mLQCs. This is mainly because that in general there does not exist a preferred initial time and state for a quantum field in an arbitrarily curved space-time [12]. If the universe is sufficiently smooth and its evolution is sufficiently slow, so that the characteristic scale of perturbations is much larger than the wavelengths of all the relevant modes, a well justified initial state can be defined: *the BD vacuum*. This is precisely the initial state commonly adopted in GR at the beginning of the slow-roll inflation, in which all the relevant perturbation modes are well inside *the comoving Hubble radius* [112] [cf. Fig.2].

However, in LQC/mLQCs, especially near the bounce, the evolution of the background is far from "slow", and its geometry is also far from the de Sitter. In particular, for the perturbations during the bouncing phase, the wavelengths could be larger, equal, or smaller than the corresponding characteristic scale, as it can be seen, for example, from Fig. 5. Thus, it is in general impossible to assume that the universe is in the BD vacuum at the bounce [36, 49–52]. Therefore, in the following let us first elaborate in more details about the subtle issues regarding the initial conditions.

A. Initial Conditions for Cosmological Perturbations

The initial conditions for cosmological perturbations in fact consists of two parts: when and which? However, both questions are related to each other. In LQC literature, for cosmological perturbations, two different moments have been chosen so far in the dressed and hybrid approaches: (i) the remote past in the contracting phase [78] and (ii) the bounce [36, 49, 50]. To see which conditions we need to impose at a given moment, let us first recall how to impose the initial conditions in GR, in which the scalar perturbations are governed by the equation,

$$v_k'' + \left(k^2 - \frac{z''}{z}\right)v_k = 0, \qquad (3.1)$$

where k denotes the comoving wave number, and $z \equiv a \dot{\delta \phi}/H$, with $\delta \phi$ being the scalar field perturbations, $\phi = \bar{\phi}(t) + \delta \phi(t, x)$. A prime denotes a derivative with respect to the conformal time η , while an over dot denotes a derivative with respect to the cosmic time t, where $d\eta = dt/a(t)$. The standard choice of the initial sate is the Minkowski vacuum of an incoming observer in the far past, $k \gg aH$ [cf. Fig. 2]. In this limit, Eq.(3.1) becomes

 $v_k'' + k^2 v_k = 0$, which has the solution,

$$v_k \simeq \frac{\alpha_k}{\sqrt{2k}} e^{-ik\eta} + \frac{\beta_k}{\sqrt{2k}} e^{ik\eta}, \qquad (3.2)$$

where α_k and β_k are two integration constants, and must satisfy the normalized condition,

$$v_k^* v_k' - v_k^{*'} v_k = -i. ag{3.3}$$

If we further require the vacuum to be the minimum energy state, a unique solution exists, which is given by $\alpha_k = 1, \ \beta_k = 0$, that is,

$$\lim_{k\gg aH} v_k \simeq \frac{1}{\sqrt{2k}} e^{-ik\eta},\tag{3.4}$$

which is often referred to as the BD vacuum [112].



FIG. 2: The evolution of the comoving Hubble radius $\ln(L_H)$ vs $\ln(a)$ in GR, where t_i denotes the moment of the onset of the slow-roll inflation, t_* the horizon crossing time of a mode with the wavenumber k, and $t_{\rm end}$ the moment that the slow-roll inflation ends.

Consider the de Sitter space as the background, we have $a(\eta) = 1/(-\eta H)$, and $z''/z = a''/a = 2/L_H^2$, where $L_H \equiv 1/(aH) = -\eta$ is the corresponding comoving Hubble radius. Then, Eq.(3.1) reads,

$$v_k'' + \left(\frac{1}{\lambda^2} - \frac{2}{L_H^2}\right)v_k = 0,$$
 (3.5)

where $\lambda \equiv 1/k$ denotes the comoving wavelength. The above equation has the following exact solutions,

$$v_k = \frac{\alpha_k}{\sqrt{2k}} e^{-ik\eta} \left(1 - \frac{i}{k\eta}\right) + \frac{\beta_k}{\sqrt{2k}} e^{ik\eta} \left(1 + \frac{i}{k\eta}\right). \quad (3.6)$$

It is clear that on scales much smaller than the comoving Hubble radius ($\lambda \ll L_H$), v_k is oscillating with frequency k and constant amplitude, given by Eq.(3.2). Then, setting (α_k , β_k) = (1,0) we find that Eq. (3.6) reduces to

$$v_k = \frac{1}{\sqrt{2k}} e^{-ik\eta} \left(1 - \frac{i}{k\eta} \right). \tag{3.7}$$

Note that if the initial time t_i is chosen sufficiently small, i.e., $t_i \ll t_{\text{end}}$ or $|k\eta| \gg 1$, all the modes are inside the comoving Hubble radius L_H [cf. Fig. 2], and the BD vacuum (3.4) becomes a natural choice.

However, on the scales much larger than the comoving Hubble radius $(\lambda \gg L_H)$, the k^2 term is negligible compared to the squeezing term, z''/z, and as a result, the fluctuations will stop oscillating and the amplitude of v_k starts to increase, yielding

$$v_k \simeq z(\eta). \tag{3.8}$$

As shown in Fig. 2, if the initial time t_i is chosen to be sufficiently early, all the currently observed modes $k_{\rm ph} \in (0.1, 1000) \times k_0^*$ will be well inside the comoving Hubble radius at $t = t_i$, so the mode function v_k can be well approximately given by Eq.(3.4), which is the wellknown zeroth order adiabatic state, where $k_0^* = 0.002$ Mpc⁻¹ and $k_{\rm ph}(t) \equiv k/a(t)$ [113].

In modified LQC models, the mode function v_k satisfies the following modified equation,

$$v_k'' + \Omega_{\text{tot}}^2(\eta, k) v_k = 0, \qquad (3.9)$$

where $\Omega^2_{\text{tot}}(\eta, k)$ depends on the homogeneous background and the inflation potential $V(\phi)$, so it is modeldependent. Therefore, the choice of the initial conditions will depend on not only the modified LQC models to be considered but also the moment at which the initial conditions are imposed.

One of the main reasons to choose the remote past in the contracting phase as the initial time for perturbations is that at such time either the background is well described by the de Sitter space (mLQC-I) or the expansion factor a(t) becomes so large that the curvature of the background is negligible (mLQC-II and LQC), so imposing the BD vacuum for mLQC-II and LQC and the de Sitter state given by Eq.(3.7) for mLQC-I at this moment is well justified. It should be noted that the reason to refer to the state described by Eq.(3.7) as the de Sitter state is the following: In the slow-roll inflation, the homogeneous and isotropic universe is almost de Sitter, as the Hubble parameter $H \equiv \dot{a}/a$ is almost a constant, so we have $a(\eta) \simeq 1/(-H\eta)$. For $t_i \ll t_{end}$ we have $a(\eta) \ll 1$, and $|\eta k| \simeq |H\eta| \gg 1$, so the choice $(\alpha_k \ \beta_k) = (1,0)$ will lead Eq.(3.7) directly to Eq.(3.4) at the onset of the slow-roll inflation [cf. Fig. 2]. However, in the deep contracting phase of the same de Sitter space, now the universe is very large, that is, $a(\eta) \gg 1$, so we must have $|H\eta| \ll 1$ and $|\eta k| \simeq |k/H_{\Lambda}|a^{-1}(\eta)$, which can be very small in sufficiently early times of the contracting phase, so the terms $i/(k\eta)$ in Eq.(3.6) now cannot be neglected. To distinguish this case from the one described by Eq.(3.4), In this review we refer the state described by Eq.(3.7) with the term $i/(k\eta)$ not being negligible as the de Sitter state, while the state described by Eq.(3.4)is still called the BD vacuum state, or simply the BD vacuum.

On the other hand, if the initial time is chosen to be at the bounce, cautions must be taken on what initial conditions can be imposed *consistently*. In particular, if at this moment some modes are inside the comoving Hubble radius and others are not, it is clear that in this case imposing the BD vacuum at the bounce will lead to inconsistencies. In addition, there also exist the cases in which particle creation in the contracting phase is not negligible, then it is unclear how a BD vacuum can be imposed at the bounce, after the universe is contracting for such a long time before the bounce. Thus, in these cases other initial conditions need to be considered, such as the fourth-order adiabatic vacuum [36, 49–52].

With the above in mind, in the following we turn to consider power spectra of the cosmological perturbations.

B. Power Spectra of Cosmological Perturbations

Since the evolutions of the effective dynamics of the homogeneous backgrounds for mLQC-I and mLQC-II are different, in this subsection let us first consider the case of mLQC-I and then mLQC-II. To compare the results with those obtained in LQC, at the end of this subsection, we also discuss the LQC case.

1. mLQC-I

For mLQC-I, the power spectrum of the cosmological scalar perturbations was first studied in [79] for the quadratic ϕ^2 potential, and re-examined in [78]. In the terminology used in [79], it was found that the corresponding mode function $v_k (\equiv q_k/a)$ satisfies Eq.(3.9) with

$$\Omega_{\text{tot}}^{2} = k^{2} - \frac{a''}{a} + \mathfrak{A}_{-}, \quad r \equiv \frac{24\pi G \dot{\phi}^{2}}{\rho}, \\ \mathfrak{A}_{-} \equiv a^{2} \left[V(\phi)r - 2V_{,\phi}(\phi)\sqrt{r} + V_{,\phi\phi}(\phi) \right], (3.10)$$

where $r = 24\pi G \dot{\phi}^2 / \rho$ and $V(\phi)$ denotes the scalar field potential.

It should be noted that, when generalizing the classical expression of the function \mathfrak{A}_{-} defined in Eq.(3.10) to its corresponding quantum mechanics operator, there exists ambiguities. In fact, classically \mathfrak{A}_{-} only coincides with Ω_Q^2 in the expanding phase. The latter is a function of the phase space variables which is explicitly given by [36, 59, 78, 79],

$$\Omega_Q^2 = 3\kappa \frac{p_\phi^2}{a^4} - 18 \frac{p_\phi^4}{a^6 \pi_a^2} - 12a \frac{p_\phi}{\pi_a} V_{,\phi} + a^2 V_{,\phi\phi}, \quad (3.11)$$

where π_a is the moment conjugate to a, and given by one of Hamilton's dynamical equations,

$$\pi_a = -\frac{3a^2}{4\pi G}H,\tag{3.12}$$

with the choice of the lapse function N = 1. Therefore, $\pi_a < 0 \ (\pi_a > 0)$ corresponds to $H > 0 \ (H < 0)$. At the quantum bounce we have $H(t_B) = 0$, so that $\pi_a(t_B) = 0$. Then, Ω_Q^2 defined by Eq.(3.11) diverges at the bounce. Hence, from the Friedmann equation, $H^2 = (8\pi G/3)\rho$, we find that

$$\frac{1}{\pi_a^2} = \frac{2\pi G}{3a^4\rho}, \quad \frac{1}{\pi_a} = \pm \sqrt{\frac{2\pi G}{3a^4\rho}}, \quad (3.13)$$

where "-" corresponds to H > 0, and "+" to H < 0. Then, a direct generalization leads to [78],

$$\Omega_Q^2 = \begin{cases} \mathfrak{A}_-, & H \ge 0, \\ \mathfrak{A}_+, & H \le 0, \end{cases}$$
(3.14)

where

$$\mathfrak{A}_{\pm} \equiv a^2 \left[V(\phi)r \pm 2V_{,\phi}(\phi)\sqrt{r} + V_{,\phi\phi}(\phi) \right].$$
(3.15)

It should be noted that in [79] only the function \mathfrak{A}_{-} was chosen over the whole process of the evolution of the universe. The same choice was also adopted in [59, 64].

In addition, \mathfrak{A} defined by Eq.(3.13) is not continuous across the bounce, as the coefficient $2V_{,\phi}(\phi)\sqrt{r}$ in general does not vanish at the bounce. In [51, 52, 55] \mathfrak{A}_{-} appearing in Eq.(3.10) was replaced by $U(\phi) (\equiv \Omega_{+}^2)$ over the whole process of the evolution of the homogeneous universe, where

$$\Omega_{\pm}^2 \equiv a^2 \left[\mathfrak{f}^2 V(\phi) \pm 2\mathfrak{f} V_{,\phi}(\phi) + V_{,\phi\phi}(\phi) \right], \qquad (3.16)$$

and $\mathfrak{f} \equiv (24\pi G/\rho)^{1/2} \dot{\phi}$.

Another choice was introduced in [78], which was motivated from the following considerations. The functions Ω_{\pm}^2 defined above are not continuous across the bounce, quite similar to \mathfrak{A}_{\pm} . However, if we introduce the quantity Ω^2 as,

$$\Omega^{2} = a^{2} \left[f^{2} V(\phi) + 2\Theta(b) f V_{,\phi}(\phi) + V_{,\phi\phi}(\phi) \right], \quad (3.17)$$

to replace \mathfrak{A}_{-} in Eq.(3.10), it could be continuous across the bounce by properly choosing $\Theta(b)$. In particular, the variable b(t) satisfies Eq.(2.61) [72–74]⁷, from which we can see that b(t) is always a monotonically decreasing function for any matter that satisfies the weak energy condition [111]. Moreover, one can construct a step-like function of b with the bounce as its symmetry axis [72– 74]. Therefore, if we define $\Theta(b)$ as

$$\Theta(b) = 1 - 2(1 + \gamma^2) \sin^2(\lambda b),$$
 (3.18)

it behaves precisely as a step function, so that Ω^2 smoothly connects Ω^2_{\pm} across the bounce, as shown in Fig. 3.



FIG. 3: The potential terms Ω^2_+ and Ω^2_- are compared with their smooth extension Ω^2 across the bounce in mLQC-I for the quadratic potential $V(\phi) = \frac{1}{2}m^2\phi^2$ [78].



FIG. 4: The potential terms Ω^2 and Ω_{eff}^2 are compared with the curvature term a''/a in mLQC-I near the bounce and the preinflationary regime for the quadratic potential $V(\phi) = \frac{1}{2}m^2\phi^2$ [78].

In addition to the above choices, motivated by the hybrid approach [37–41], the following replacements for π_a^{-2} and π_a^{-1} in Eq.(3.11) were also introduced in [78],

$$\frac{1}{\pi_a^2} \to \frac{64\pi^2 G^2 \lambda^2 \gamma^2}{9a^4 \mathcal{D}}, \qquad (3.19)$$

$$\frac{1}{\pi_a} \rightarrow -\frac{8\pi G\lambda\gamma\Theta(b)}{3a^2\mathcal{D}^{1/2}},$$
 (3.20)

⁷ It is interesting to note that Eq.(2.61) holds not only in mLQC-I, but also in LQC and mLQC-II.

where

$$\mathcal{D} \equiv \left(1 + \gamma^2\right) \sin^2\left(2\lambda b\right) - 4\gamma^2 \sin^2\left(\lambda b\right). \qquad (3.21)$$

Such obtained Ω_Q^2 was referred to as Ω_{eff}^2 in [78], and in Fig. 4, we show the differences among Ω^2 , Ω_{eff}^2 and the quantity a''/a, from which one can see that the term a''/a dominates the other two terms over the whole range $t/t_{pl} \in (-8, 10^4)$.

To study the effects of the curvature term, let us first introduce the quantity,

$$k_B^{\rm I} = \left. \left(\frac{a''}{a} \right)^{1/2} \right|_{t=t_B} \approx 1.60,$$
 (3.22)

which is much larger than other two terms Ω^2 and Ω_{eff}^2 , where $\Omega^2(t_B) = 1.75 \times 10^{-10}$ and $\Omega_{\text{eff}}^2(t_B) = 0.006$. Therefore, the differences between Ω^2 and Ω_{eff}^2 near the bounce are highly diluted by the background. On the other hand, in the post-bounce phase, Ω^2 and Ω_{eff}^2 coincide after $t/t_{pl} \simeq 10^4$, while near the onset of the inflation their amplitudes first become almost equal to that of the curvature term, and then quickly exceeds it during the slow-roll inflation, as we can see from Fig. 4.

From Fig. 4 we can also see that the difference between Ω^2 and Ω_{eff}^2 lies mainly in the region near the bounce. However, as the curvature term a''/a overwhelmingly dominates in this region, it is usually expected that the impact of the different choices of Ω^2 on the power spectrum might not be very large [36, 59, 79]. However, in [78], it was found that the relative difference in the magnitude of the power spectrum in the IR and oscillating regimes could be as large as 10%, where the relative difference is defined as,

$$\mathcal{E} \equiv 2 \left| \frac{\mathcal{P}_1 - \mathcal{P}_2}{\mathcal{P}_1 + \mathcal{P}_2} \right|. \tag{3.23}$$

However, the power spectra obtained from Ω^2 and Ω_{\pm}^2 are substantially different even in the UV regime due to the (tiny) difference between Ω_{\pm}^2 at the bounce, see Fig. 14 given in [78]. In fact, the difference is so large that the power spectrum calculated from Ω_{\pm}^2 is essentially already ruled out by current observations.

With the clarification of the ambiguities caused by the quantum mechanical generalization of the function \mathfrak{A}_{-} defined in Eq.(3.10), now let us turn to the issue of the initial conditions, for which we consider only two representative potentials, the quadratic and Starobinsky, given explicitly by

$$V = \begin{cases} \frac{1}{2}m^2\phi^2, & \text{quadratic,} \\ \frac{3m^2}{32\pi G}\left(1 - e^{-\sqrt{16\pi G/3}}\phi\right)^2, & \text{Starobinsky.} \end{cases}$$
(3.24)

In mLQC-I, the evolution of the effective (quantum) homogeneous universe is asymmetric with respect to the bounce [66, 72–74]. In particular, before the bounce (t <

 t_B), the universe is asymptotically de Sitter, and only very near the bounce (about several Planck seconds), the Hubble parameter H which is negative in the pre-bounce regime suddenly increases to zero at the bounce. Then, the universe enters a very short super-acceleration phase $\dot{H} > 0$ (super-inflation) right after the bounce, which lasts until $\rho \simeq \rho_c^{\rm I}/2$, where $\dot{H}(t)_{\rho\simeq\frac{1}{2}\rho_c^{\rm I}} = 0$. Afterwards, for a kinetic energy dominated bounce $\dot{\phi}_B^2 \gg V(\phi_B)$, it takes about $10^4 \sim 10^6$ Planck seconds before entering the slow-roll inflation [72–74]. Introducing the quantity,

$$\lambda_H^2 \equiv \frac{a}{a^{\prime\prime} - a\mathfrak{A}_-} = -\frac{1}{m_{\text{eff}}},\qquad(3.25)$$

where m_{eff} is the effective mass of the modes, from Eq.(3.10) we find that

$$\Omega_{\text{tot}}^2 = \frac{1}{\lambda^2} - \frac{1}{\lambda_H^2} = \begin{cases} > 0, & \lambda_H^2 > \lambda^2, \\ < 0, & 0 < \lambda_H^2 < \lambda^2, \\ > 0, & \lambda_H^2 < 0, \end{cases}$$
(3.26)

where $\lambda \ (\equiv k^{-1})$ denotes the comoving wavelength of the mode k, as mentioned above. Note that such a defined quantity λ_H^2 becomes negative when the effective mass is positive. In Fig. 5, we plot λ_H^2 schematically for the quadratic and the Starobinsky potentials with the mass of the inflaton set to $1.21 \times 10^{-6} m_{\rm pl}$ and $2.44 \times 10^{-6} m_{\rm pl}$ respectively. The initial conditions for the background evolution are set as follows: for the quadratic potential, the inflaton starts with a positive velocity on the right wing of the potential and for the Starobinsky potential the inflaton is released from the left wing of the potential with a positive velocity. For both potentials, the initial conditions are set at the bounce which is dominated by the kinetic energy of the inflaton field. The same mass parameters and similar initial conditions are also used in the following figures where the comoving Hubble radius is plotted schematically. In Fig. 5, the moments t_H and t_i are defined, respectively, by $a''(t_H) = a''(t_i) = 0$, so t_i represents the beginning of the inflationary phase, and during the slow-roll inflation (Region III), we have $\lambda_H^2 \approx L_H^2/2 \simeq 1/(2a^2H^2)$, which is exponentially decreasing, and all the modes observed today were inside the comoving Hubble radius at $t = t_i$. Between the times t_H and t_i , λ_H^2 is negative, and $\Omega_{\rm tot}^2$ is strictly positive. Therefore, during this period the mode functions are oscillating, while during the epoch between t_B and t_H , some modes $(k^{-2} > k_B^{-2})$ are inside the comoving Hubble radius, and others $(k^{-2} < k_B^{-2})$ are outside it right after the bounce, where $k_B \equiv \lambda_B^{-1}(t_B)$. In the contracting phase, when $t \ll t_B$, the universe is quasi-de Sitter and $\lambda_H^2 \simeq 1/(2a^2H^2)$ increases exponentially toward the bounce $t \to t_B$, as now a(t) is decreasing exponentially. However, several Planck seconds before the bounce, the universe enters a non-de Sitter state, during which λ_{H}^{2} starts to decrease until the bounce, at which a characteristic Planck scale $k_B \ (\equiv 1/\lambda_H)$ can be well defined. Therefore, for $t \ll t_B$, all the modes are outside the comoving Hubble radius. Then, following our

previous arguments, if the initial moment is chosen at $t_0 \ll t_B$, the de Sitter state seems not to be viable. However, when $t_0 \ll t_B$ we have $a(\eta) \simeq 1/(-\eta |H_\Lambda|)$, where $H_{\Lambda} = -[\lambda(1+\gamma^2)]^{-1}$ and

$$\Omega_{\rm tot}^2 \simeq k^2 - \frac{2}{\eta^2},\tag{3.27}$$

for which Eq.(3.9) has the exact solutions given by Eq.(3.6). Therefore, at sufficient early times, choosing $\alpha_k = 1$, $\beta_k = 0$ leads us to the de Sitter state (3.7). From the above analysis it is clear that this is possible precisely because of the isometry of the de Sitter space, which is sufficient to single out a preferred state, the de Sitter state [79].

With the exact solution (3.7) as the initial conditions imposed at the moment $t_0 \ (\ll t_B)$ in the contracting phase, it was found that the power spectrum of the cosmological scalar perturbations can be divided into three different regimes: (i) the ultraviolet (UV) $(k > k_{mLQC-I})$; (ii) intermediate $(k_i < k < k_{mLQC-I})$; and (iii) infrared $(k < k_i)$, where $k_{mLQC-I} \equiv a_B \sqrt{R_B/6}$ and $k_i = a_i \sqrt{R_i/6}$, and R_B and R_i are the curvatures given at the bounce and the beginning of the slow-roll inflation, respectively [cf. Fig. 5]. During the infrared regime, the power spectrum increases as k increases, while in the intermediate regime it is oscillating very fast and the averaged amplitude of the power spectrum is decreasing as k increases. In the UV regime, the spectrum is almost scale-invariant, which is consistent with the current observations. There exists a narrow band, $0.1 \times k_0^* < k < k_{\rm mLQC-I}$, in which the quantum gravitational effects could be detectable by current or forthcoming cosmological observations [79]. Within the dressed metric approach, one of the most distinctive features of the power spectrum in mLQC-I is that its magnitude in the IR regime is of the Planck scale [78, 79]. This is because those infrared modes are originally outside the Hubble horizon in the contracting phase and thus their magnitudes are frozen as they propagate across the bounce and then into the inflationary phase. Considering that the contracting phase is a quasi de Sitter phase with a Plank-scale Hubble rate, the magnitude of the IR modes is thus also Planckian [78].

It should be noted that if the initial conditions are imposed at the bounce, from Fig. 5 we can see clearly that some modes are inside the comoving Hubble radius, and some are not. In addition, in the neighborhood of the bounce, the background is far from de Sitter. So, it is impossible to impose either the BD vacuum or the de Sitter state at the bounce. In this case, one of the choices of the initial conditions is the fourth-order adiabatic vacuum, similar to that in LQC [36, 49–52].

2. mLQC-II

In mLQC-II, the evolution of the effective homogeneous universe is different from that of mLQC-I. In par-



FIG. 5: Schematic plot of λ_H^2 defined by Eq.(3.25) vs t for mLQC-I in the dressed metric approach for the quadratic and the Starobinsky potentials, where $a''(t_H) = 0$ and $a''(t_i) = 0$ with t_i being the starting time of the inflationary phase. During the slow-roll inflation, we have $\lambda_H^2 \approx L_H^2/2$. In the contracting phase $t < t_B$, the universe is initially de Sitter and we still have $\lambda_H^2 \approx L_H^2/2$, but now it increases exponentially toward bounce $t \to t_B$, as the universe in this phase is exponentially contracting. However, several Planck seconds before the bounce, the universe enters a non-de Sitter state, during which λ_H^2 starts to decrease until the bounce. The qualitative behavior of the comoving Hubble radius is the same for the quadratic and the Starobinsky potentials. Different potentials will change the the values of t_H and t_i correspondingly.

ticular, it is symmetric with respect to the bounce and in the initially kinetic energy dominated case at the bounce the solutions can be well approximated by Eq.(2.56) in the bouncing phase [73, 74], similar to that of LQC [16].

When considering the cosmological perturbations, similar ambiguities in the choices of π_a^{-2} and π_a^{-1} in Eq.(3.11) exist. In particular, for the choice of Eq.(3.17) now the function $\Theta(b)$ is replaced by

$$\Theta(b) = \cos\left(\frac{\lambda b}{2}\right),\tag{3.28}$$

which behaves also like a step function across the bounce and picks up the right sign in both contracting and expanding phases, so it smoothly connects Ω_{\pm} defined by Eq.(3.16).

On the other hand, Ω_{eff}^2 is obtained from Eq.(3.11) by the replacements,

$$\frac{1}{\pi_a^2} \to \frac{4\pi^2 \gamma^2 \lambda^2}{9a^4 \sin^2 \left(\lambda b/2\right) \mathcal{D}},\tag{3.29}$$

$$\frac{1}{\pi_a} \rightarrow \frac{-2\pi\gamma\lambda\cos\left(\lambda b/2\right)}{3a^2\sin\left(\lambda b/2\right)\mathcal{D}^{1/2}},\tag{3.30}$$

but now with

$$\mathcal{D} \equiv 1 + \gamma^2 \sin^2\left(\frac{\lambda b}{2}\right).$$
 (3.31)

Such obtained Ω^2 , Ω^2_{\pm} and Ω^2_{eff} are quite similar to those given by Figs. 3 and 4 in mLQC-I. In particular,

at the bounce, we have

$$\Omega^{2}(t_{B}) = 1.59 \times 10^{-10}, \quad \Omega^{2}_{\text{eff}}(t_{B}) = 0.265,$$
$$k_{B}^{\text{II}} = \left. \left(\frac{a''}{a} \right)^{1/2} \right|_{t=t_{B}} \approx 6.84, \quad (3.32)$$

that is, the curvature term a''/a still dominates the evolution near the bounce.



FIG. 6: Schematic plot of λ_H^2 defined by Eq.(3.25) vs t for mLQC-II in the dressed metric approach for the quadratic and the Starobinsky potentials, where $a''(t_H^p) = a''(t_H) = 0$ and $a''(t_i^p) = a''(t_i) = 0$, and $t = t_i$ denotes the starting time of the inflationary phase, while $t = t_i^p$ is the end time of the deflationary phase in the contracting branch. During the slow-roll inflation, we have $\lambda_H^2 \approx L_H^2/2$. In particular, λ_H^2 is decreasing (increasing) exponentially in Region III (Region III'). The corresponding effective mass near the bounce is always negative. Similar behavior also happens in LQC in the dressed metric approach [51]. The bounce is dominated by the kinetic energy of the scalar field, which leads to $t_H^p \approx -t_H$. However, in general we find that $t_i^p \neq -t_i$ due to the effects of the potential energy of the scalar field far from the bouncing point. The comoving Hubble radius has the same qualitative behavior for the quadratic and the Starobinsky potentials, while the values of t_i (t_i^P) and t_H (t_H^p) depend on the type of the potentials and the initial conditions.

To see how to impose the initial conditions, let us introduce the quantity λ_H^2 defined by Eq.(3.25) but now \mathfrak{A}_- will be replaced either by Ω^2 or Ω_{eff}^2 . The details here are not important, and λ_H^2 is schematically plotted in Fig. 6, from which we can see that if the initial conditions are chosen to be imposed at the bounce, the BD vacuum (as well as the de Sitter state) is still not available, and the fourth-order adiabatic vacuum is one of the possible choices, similar to the LQC case. However, if the initial conditions are imposed in the contracting phase at $t_0 \ll t_i^p$, the universe becomes very large $a(t) \gg 1$ and can be practically considered as flat, then the BD vacuum can be chosen.

Certainly, one can choose different initial conditions. In particular, the fourth-order adiabatic vacuum was chosen even in the contracting phase in [78]. With such a choice, the power spectra from Ω^2 and $\Omega_{\rm eff}^2$ in the region

 $k \in (5 \times 10^{-6}, 50)$ were studied and found that the relative difference in the magnitude of the power spectra is around 30% in the IR regime and less than 10% in the intermediate regime. In the UV regime, the relative difference can be as small as 0.1% or even less.

3. LQC

To consider the effects of the ambiguities in the choice of π_a^{-2} and π_a^{-1} in Eq.(3.11)⁸, power spectra of the cosmological perturbations were also studied in the framework of LQC in [78]. In this case, Ω^2 is obtained from Eq.(3.17) with

$$\Theta(b) = \cos(\lambda b), \tag{3.33}$$

while Ω_{eff}^2 is obtained from Eq.(3.11) by the replacements,

$$\frac{1}{\pi_a^2} \to \frac{16\pi^2 G^2 \gamma^2 \lambda^2}{9a^4 \sin^2(\lambda b)},\tag{3.34}$$

$$\frac{1}{\pi_a} \rightarrow \frac{-4\pi\gamma\lambda\cos\left(\lambda b\right)}{3a^2\sin\left(\lambda b\right)}.$$
(3.35)

As shown explicitly, the term Ω^2_+ is always negligible comparing with the curvature term a''/a in the expression of Ω^2_{tot} defined in Eq.(3.10) by replacing \mathfrak{A}_- with Ω^2_+ . So, from Eq.(3.25) we find that

$$\lambda_H^2 = \frac{1}{a''/a - \Omega_+^2} \simeq \frac{a}{a''},$$
(3.36)

during the bouncing phase $t \in (t_B, t_i)$, and $\lambda_H^2 \simeq a/a''$ was shown schematically by Fig. 18 in [52], which is quite similar to Fig. 6 given above for mLQC-II.

As a result, the initial states of the linear perturbations can be either imposed in the contracting phase at a moment $t_0 \ll t_i^p$ as the BD vacuum, or at the bounce as the fourth-order adiabatic vacuum [36]. However, it was shown analytically that such two conditions lead to the same results [52].

To compare the results obtained from the three different models, in [78] the fourth-order adiabatic vacuum was chosen even in the contracting phase for LQC. Here, we cite some of the results in Fig. 7. In particular, it was found that the relative difference in the amplitudes of the power spectra of the scalar perturbations due to the choice of Ω^2 or Ω_{eff}^2 is about 10% in the infrared regime, about 100% in the intermediate regime, and about 0.1% in the UV regime. Since only modes in the UV regime can be observed currently, clearly this difference is out of

⁸ In the framework of LQC, such effects were also studied in [36, 51, 52, 55, 78]. In particular, in [59, 64, 79] the function \mathfrak{A}_{-} defined in Eq.(3.15) was chosen over the whole process of the evolution of the universe.



FIG. 7: The figure shows the results of the scalar power spectra from three models presented in [78] when the potential term is given by $\Omega_{\rm eff}^2$. The inflationary potential is chosen to be the quadratic potential and the e-foldings of the inflationary phases in all three models are chosen to be 72.8. The first panel shows the scalar power spectrum in mLQC-I which is characterized by its unique infrared regime. In the second panel, we compare the scalar power spectra from LQC and mLQC-II.

the sensitivities of the current and forhcoming observations [114].

However, comparing the power spectra obtained from the three different models, even with the same choice of π_a^{-2} and π_a^{-1} , it was found that the relative difference among LQC, mLQC-I and mQLC-II are significant only in the IR and oscillating regimes, while in the UV regime, all three models give quite similar results. In particular, with the same regularization of π_a the difference can be as large as 100% throughout the IR and oscillating regimes, while in the UV regime it is about 0.1%.

For the tensor perturbations, the potential term Ω_Q vanishes identically, so no ambiguities related to the choice of π_a exist. But, due to different models, the differences of the power spectra of the tensor perturbations can be still very large in the IR and oscillating regimes among the three models, although they are very small in the UV regime, see, for example, Fig. 12 given in [78].

IV. PRIMORDIAL POWER SPECTRA OF MODIFIED LQCS IN HYBRID APPROACH

As in the previous section, in this section we also consider the three different models, LQC, mLQC-I, and mLQC-II, but now in the hybrid approach, and pay particular attention to the differences of the power spectra among these models. Since the scalar perturbations are the most relevant ones in the current CMB observations, in the following we shall mainly focus on them, and such studies can be easily extended to the tensor perturbations.

A. mLQC-I

Power spectra of the cosmological scalar and tensor perturbations for the effective Hamilton in mLQC-I were recently studied in the hybrid approach [82–84]. In particular, the mode function v_k of the scalar perturbations satisfies the differential equation [84],

$$v_k'' + (k^2 + s) v_k = 0, (4.1)$$

where

$$s = \frac{4\pi G p_{\phi}^{2}}{3v^{4/3}} \left(19 - 24\pi G \gamma^{2} \frac{p_{\phi}^{2}}{\pi_{a}^{2}} \right) \\ + v^{2/3} \left(V_{,\phi\phi} + \frac{16\pi G \gamma p_{\phi}}{\pi_{a}} V_{,\phi} - \frac{16\pi G}{3} V \right) \\ = -\frac{4\pi G}{3} a^{2} \left(\rho - 3P \right) + \mathcal{U}, \qquad (4.2)$$

which is the effective mass of the scalar mode, with

$$\mathcal{U} \equiv a^2 \left[V_{,\phi\phi} - 12 \frac{V_{,\phi}}{\pi_a} + \frac{64a^6 V(\phi)}{\pi G} \left(\rho - \frac{3V(\phi)}{4\pi G} \right) \frac{1}{\pi_a^2} \right].$$
(4.3)

Note that in [84], instead of π_a , the symbol Ω was used. In addition, the cosmological tensor perturbations are also given by Eqs.(4.1) and (4.2) but with the vanishing potential $\mathcal{U} = 0$. Then, we immediately realize that in the hybrid approach quantum mechanically there are also ambiguities in the replacements π_a^{-2} and π_a^{-1} , as mentioned in the last section. So far, two possibilities were considered [82, 83]. One is given by the replacements,

$$\frac{1}{\pi_a^2} \to \frac{1}{\Omega_{\rm I}^2}, \qquad \frac{1}{\pi_a} \to \frac{\Lambda_{\rm I}}{\Omega_{\rm I}^2},$$
(4.4)

in Eq.(4.2), where

$$\Omega_{\rm I}^2 \equiv -\frac{v^2 \gamma^2}{\lambda^2} \Big\{ \sin^2 \left(\lambda b\right) - \frac{\gamma^2 + 1}{4\gamma^2} \sin^2 \left(2\lambda b\right) \Big\}, \Lambda_{\rm I} \equiv v \frac{\sin(2\lambda b)}{2\lambda}.$$
(4.5)

This is the case referred to as prescription A in [83].

The other possibility is obtained by the replacement of Eqs.(3.34) and (3.35), which was referred to as Prescription B [83], and showed that the two prescriptions lead to almost the same results. So, in the rest of this section we restrict ourselves only to prescription A.

Then, for the case in which the evolution of the homogeneous universe was dominated by kinetic energy at the bounce,

$$\dot{\phi}_B^2 \gg 2V(\phi_B),$$
(4.6)

it was shown that the effective mass is always positive at the bounce [83]. In fact, near the bounce we have [58],

$$s = -\frac{4\pi G}{3}a^{2}(\rho - 3P) + \mathcal{U}(\eta)$$

\$\approx \frac{8\pi G}{3}a^{2}\rho > 0. \quad (4.7)\$

Note that in writing the above expression, we have used the fact that during the bouncing phase we have $w_{\phi} \equiv P/\rho \simeq 1$, and $|\mathcal{U}(\eta)| \ll 1$. On the other hand, in the pre-bounce phase, when $t \ll t_B$ the background is a contracting de Sitter spacetime, so we have [83],

$$s = -\frac{4\pi G}{3}a^2 \left(\rho - 3P\right) + \mathcal{U}(\eta) \simeq \mathcal{U}(\eta) \simeq 5a^2 V_{,\phi\phi},$$

$$a \simeq a_B e^{H_\Lambda(t-t_B)}, \qquad (4.8)$$

where $H_{\Lambda} \equiv -\sqrt{8\pi\alpha G\rho_{\Lambda}/3}$. Thus, the effective mass remains positive in the pre-bounce phase, as long as $V_{,\phi\phi}(t \ll t_B) > 0$. This is the case for both quadratic and Starobinsky potentials. In fact, from (3.24), we find that

$$V_{,\phi\phi} = \begin{cases} m^2, & \text{quadratic,} \\ m^2 \left(2 - e^4 \sqrt{\pi G/3} \phi\right) e^{-8\sqrt{\pi G/3} \phi}, & \text{Starobinsky.} \end{cases}$$
(4.9)

For the case that satisfies the condition (4.6) initially at the bounce, we find that $\phi(t)$ becomes very negative at $t \ll t_B$ for the Starobinsky potential, so $V_{,\phi\phi}(t \ll t_B)$ is positive even in this case. Then, the quantity defined by

$$\lambda_H^2 \equiv -\frac{1}{s},\tag{4.10}$$

has similar behavior in the post-bounce phases for the case in which the evolution of the homogeneous universe was dominated by kinetic energy at the bounce, but has different behaviors in the pre-bounce phases, depending specifically on the potentials considered.

In Figs. 8 and 9 we show the comoving Hubble radius for the quadratic and Starobinsky potentials, respectively. From these figures it is clear that for $t_i^p < t < t_i$, λ_H^2 is strictly negative, which implies the effective mass *s* is positive in this regime. Hence, all the modes assume the oscillatory behavior as the modes inside the Hubble horizon, and we may impose the BD vacuum at the bounce. In addition, when $t \ll t_i^p$, the background is well described by the de Sitter space, so the de Sitter state can be imposed in the deep contracting phase. However, imposing the BD vacuum at the bounce will clearly lead to different power spectra at the end of the slow-roll inflation from that obtained by imposing the de Sitter state in the deep contracting phase. This is because, when the background is contracting to about the moments $t \simeq t_i^{p-}$, the effective mass becomes so large and negative that the mode function v_k will be modified significantly, in comparison with that given at $t_0 (\ll t_i^p)$, or in other words, particle creation now becomes not negligible during the contracting phase. Then, other initial conditions at the bounce may need to be considered.



FIG. 8: Schematic plot of λ_H^2 defined by Eq.(4.10) in mLQC-I for the quadratic potential in the hybrid approach, where $s(t_i) = s(t_i^p) = 0$, and $t = t_i$ is the starting time of the inflationary phase. During the slow-roll inflation, we have $\lambda_H^2 \approx L_{\rm H}^2/2$ (Region III). In the contracting phase, the background is asymptotically de Sitter. The evolution of the universe is asymmetric with respect to the bounce. In particular, λ_H^2 is strictly negative for $t_i^p < t < t_i$, while for $t \simeq t_i^{p-1}$ the "generalized" comoving Hubble radius λ_H^2 becomes positive and large. However, as t decreases, λ_H^2 becomes negative again. Although the values of t_i and t_H depend on the initial conditions for the background evolution, for example, when $\phi_B = 1.27 \ m_{\rm pl}$ at the bounce, $t_i \approx 7.55 \times 10^4 \ t_{\rm pl}$ and $t_H \approx -21.85 t_{\rm pl}$, the qualitative behavior of the comoving Hubble radius is robust with respect to the choice of the initial conditions as long as the bounce is dominated by the kinetic energy of the scalar field.

B. mLQC-II

Similar to LQC, the homogeneous universe of mLQC-II is symmetric with respect to the bounce, and is well described by the analytical solutions given by Eqs.(2.56) and (2.57) for the states that are dominated by kinetic energy at the bounce.

In this model, the cosmological perturbations are also given by Eqs.(4.1)-(4.3) but now with the replacement



FIG. 9: Schematic plot of λ_H^2 defined by Eq.(4.10) for the Starobinsky potential and mLQC-I in the hybrid approach, where $s(t_i) = s(t_i^p) = s(t_H^p) = 0$, and $t = t_i$ is the starting time of the inflationary phase. During the slow-roll inflation, we have $\lambda_H^2 \approx L_H^2/2$ (Region III). In the contracting phase, the background is asymptotically de Sitter. The evolution of the universe is asymmetric with respect to the bounce. In particular, λ_H^2 is strictly negative for $t_i^p < t < t_i$, while for $t \simeq t_i^{p-}$ it becomes positive and large. However, as t decreases, λ_H^2 becomes negative again. The qualitative behavior of λ_H^2 does not change with the choice of the initial conditions as long as the inflaton initially starts from the left wing of the potential at the kinetic-energy-dominated bounce. However, the exact values of t_i and t_H depend on the initial conditions. For example, when $\phi_B = -1.32 \ m_{\rm pl}$, $t_H^p = -7.88 \ t_{\rm pl}$, $t_i^p = -4.11 \ t_{\rm pl}$ and $t_i = 4.90 \times 10^5 \ t_{\rm pl}$.

[84],

$$\frac{1}{\pi_a^2} \to \frac{1}{\Omega_{\rm II}^2}, \qquad \frac{1}{\pi_a} \to \frac{\Lambda_{\rm II}}{\Omega_{\rm II}^2}, \tag{4.11}$$

where

$$\Omega_{\rm II}^2 \equiv \frac{4v^2}{\lambda^2} \sin^2\left(\frac{\lambda b}{2}\right) \left\{1 + \gamma^2 \sin^2\left(\frac{\lambda b}{2}\right)\right\},$$

$$\Lambda_{\rm II} \equiv v \frac{\sin\left(\lambda b\right)}{\lambda}.$$
 (4.12)

In this case, it can be shown that the effective mass defined by Eq.(4.2) is always positive in the neighborhood of the bounce, but far from the bounce, the properties of λ_H^2 depend on the potential in the pre-bounce phase, similar to mLQC-I.

In Fig. 10, we plot λ_H^2 for the Starobinsky potential, while for the quadratic one, it is quite similar to the corresponding one in mLQC-I, given by Fig. 8. From Fig. 10 we can see that λ_H^2 now is negative not only near the bounce but also in the whole contracting phase, so that all the modes are oscillating for $t < t_i$. Then, one can choose the BD vacuum at the bounce. It is remarkable that for the quadratic potential, this is impossible [cf. Fig. 8].

Moreover, as $t \to -\infty$, the expansion factor becomes very large, and the corresponding curvature is quite low, so to a good approximation, the BD vacuum can also be chosen in the distant past, not only for the Starobinsky potential but also for other potentials. Due to the oscillating behavior of the mode function over the whole contracting phase, imposing the BD vacuum at the bounce is expected not to lead to significant difference in the power spectra from that in which the same condition is imposed in the deep contracting phase.



FIG. 10: Schematic plot of λ_H^2 defined by Eq.(4.10) for the Starobinsky potential and mLQC-II in the hybrid approach, where $s(t_i) = 0$, and $t = t_i$ is the starting time of the inflationary phase. During the slow-roll inflation, we have $\lambda_H^2 \approx L_{\rm H}^2/2$ (Region III) since the contribution from the potential is in general less than a''/a. Again the qualitative behavior of λ_H^2 remains the same as long as the inflaton starts from the left wing of the potential with a positive velocity and the bounce is initially dominated by the kinetic energy of the inflaton field.

C. LQC

The evolution of the homogeneous universe of standard LQC model is also symmetric with respect to the bounce, and is well described by the analytical solutions given in [51, 52] for the states that are dominated by kinetic energy at the bounce.

In this model, the cosmological perturbations are also given by Eqs.(4.1)-(4.3) but now with the replacement [84],

$$\frac{1}{\pi_a^2} \to \frac{1}{\Omega_{LQC}^2}, \qquad \frac{1}{\pi_a} \to \frac{\Lambda_{LQC}}{\Omega_{LQC}^2}, \qquad (4.13)$$

where

$$\Omega_{\text{LQC}} \equiv \frac{v \sin(\lambda b)}{\lambda}, \quad \Lambda_{\text{LQC}} \equiv \frac{v \sin(2\lambda b)}{2\lambda}.$$
 (4.14)

In this case, it can be shown that the effective mass defined by Eq.(4.2) is always positive for the states that are dominated by kinetic energy at the bounce [55, 58], and the quantity λ_H^2 defined by Eq.(4.10) is negative near the bounce. Again, similar to the mLQC-II case, the modes are oscillating near the bounce. However, in the contracting phase the behavior of λ_H^2 sensitively depends

on the inflation potentials. For the Starobinsky one, λ_H^2 behaves similar to that described by Fig. 10, so the BD vacuum can be imposed either in the deep contracting phase or at the bounce, and such resulted power spectra are expected not to be significantly different from one another. But for the quadratic potential the situation is quite different, and a preferred choice is to impose the BD vacuum in the deep contracting phase $(t_0 \ll t_B)$.

D. Primordial Power Spectra

As it can be seen that one of the preferred moments to impose the initial conditions for the cosmological perturbations in all these three models is a moment in the contracting phase $t_0 < t_B$. In this phase, we can impose the BD vacuum state as long as the moment is sufficiently earlier, $t_0 \ll t_B$. Certainly, other initial conditions can also be chosen. In particular, in [84] the second-order adiabatic vacuum conditions were selected, but it was found that the same results can also be obtained even when the BD vacuum or the fourth-order adiabatic vacuum is imposed initially.

The nth-order adiabatic vacuum conditions can be obtained as follows: Let us first consider the solution,

$$\nu_k = \frac{1}{\sqrt{2W_k}} e^{-i\int^{\eta} W_k(\bar{\eta})d\bar{\eta}}.$$
 (4.15)

Then, inserting it into (4.1), one can find an iterative equation for W_k . In particular, it can be shown that the zeroth-order solution is given by $W_k^{(0)} = k$, while the second and fourth order adiabatic solutions are given by,

$$W_k^{(2)} = \sqrt{k^2 + s}, \qquad W_k^{(4)} = \frac{\sqrt{f(s,k)}}{4|k^2 + s|}.$$
 (4.16)

Here $f(s,k) = 5s'^2 + 16k^4(k^2+3s) + 16s^2(3k^2+s) - 4s''(s+k^2)$. It should be noted that, in order to compare directly with observations, it is found convenient to calculate the power spectrum of the comoving curvature perturbation \mathcal{R}_k , which is related to the Mukhanov-Sasaki variable via the relation $\mathcal{R}_k = \nu_k/z$, with $z = a\dot{\phi}/H$. Then, its power spectrum reads

$$\mathcal{P}_{\mathcal{R}_k} = \frac{\mathcal{P}_{\nu_k}}{z^2} = \frac{k^3}{2\pi^2} \frac{|\nu_k|^2}{z^2}.$$
 (4.17)

In addition, the power spectrum is normally evaluated at the end of inflation, at which all the relevant modes are well outside the Hubble horizon [cf. Fig. 2].

It should be also noted that the above formula is only applicable to the case where $W_k^{(2)}$ and/or $W_k^{(4)}$ remains real at the initial time. This is equivalent to require $k^2 + s \ge 0$ for $W_k^{(2)}$ and $f(s,k) \ge 0$ for $W_k^{(4)}$. Since the effective mass s in general depends on t, it is clear that the validity of (4.16) depends not only on the initial states but also on the initial times.



FIG. 11: The primordial power spectra of the cosmological scalar perturbations in the hybrid approach with the Starobinsky potential, respectively, for LQC, mLQC-I, and mLQC-II. The mass of the inflaton field is set to $2.44 \times 10^{-6} m_{\rm pl}$. The background evolution is chosen so that the pivot mode is $k_* = 5.15$ in all three models. The initial states are the second-order adiabatic states imposed in the contracting phase at the moment t_0 with $t_0 \ll t_B$ [84].

In addition, in the following only the Starobinsky potential given in Eq.(3.24) will be considered, as it represents one of the most favorable models by current observations [113]. Let us turn to consider the power spectrum of the scalar perturbations in each of the three models. Similar results can be also obtained for the tensor perturbations. In particular, it was found that the scalar power spectra in these three models can be still divided into three distinctive regimes: the infrared, oscillatory and UV, as shown in Fig. 11.

In the infrared and oscillatory regimes, the relative difference between LQC and mLQC-I can be as large as 100%, while this difference reduces to less than 1% in the UV regime. This is mainly because LQC and mLQC-I have the same classical limit in the post-bounce phase, and as shown in Figs. 5 and 8, the effective masses in both approaches tend to be the same during the inflationary phase.

However, it is interesting to note that in the infrared and oscillatory regimes, the power spectrum in mLQC-I is suppressed in comparison with that of LQC, which has been found only in the hybrid approach. As a matter of fact, in the dressed metric approach, the power spectrum in mLQC-I is largely amplified in the infrared regime, and its magnitude is of the Planck scale as depicted in Fig. 7 [78, 79]. The main reason might root in the distinctive behavior of the effective masses in the two approaches, as shown explicitly in Figs. 5 and 8.

On the other hand, the difference of the power spectra between LQC and mLQC-II is smaller than that between LQC and mLQC-I. In particular, in the infrared regime, it is about 50%. The large relative difference (more than 100%) of the power spectra between mLQC-I and mLQC-II also happens in the infrared and oscillatory regimes, while in the UV regime it reduces to about 2%. To summarize, in the hybrid approach the maximum relative differences of the power spectra among these three different models always happen in the infrared and oscillatory regimes, while in the UV regime, the differences reduce to no larger than 2%, and all the three models predict a scale-invariant power spectrum, and is consistent with the current CMB observations. However, in the hybrid approach, the power spectrum in mLQC-I is suppressed in the infrared and oscillatory regimes. The latter is in a striking contrast to the results obtained from the dressed metric approach, which might be closely related to the fact that the effective masses in these two approaches are significantly different, especially near the bounce and in the prebounce stage.

V. CONCLUSIONS AND OUTLOOK

In the past two decades, LQC has been studied extensively, and several remarkable features have been found [16], including the generic resolution of the big bang singularity (replaced by a quantum bounce) in the Planckian scale, the slow-roll inflation as an attractor in the post-bounce evolution of the universe, and the scaleinvariant power spectra of the cosmological perturbations, which are consistent with the current CMB observations. Even more interestingly, it was shown recently that some anomalies from the CMB data [60–62] can be reconciled purely due to the quantum geometric effects in the framework of LQC [63, 64].

Despite of all these achievements, LQC is still plagued with some ambiguities in the quantization procedure. In particular, its connection with LQG is still not established [90], and the quantization procedure used in LQC owing to symmetry reduction before quantization can result in different Hamiltonian constraints than the one of LQG.

Motivated by the above considerations, recently various modified LQC models have been proposed, see, for example, [115–124] and references therein. In this brief review, we have restricted ourselves only to mLQC-I and mLQC-II [65–67], as they are the ones that have been extensively studied in the literature not only the dynamics of the homogeneous universe [72–76, 80, 81], but also the cosmological perturbations [78, 79, 82–84].

In these two modified LQC models, it was found that the resolution of the big bang singularity is also generic [72–76]. This is closely related to the fact that the area operator in LQG has a minimal but nonzero eigenvalue [15, 16], quite similar to the eigenvalue of the ground state of the energy operator of a simple harmonic oscillator in quantum mechanics. This deep connection also shows that the resolution of the big bang singularity is purely due to the quantum geometric effects. In addition, similar to LQC, the slow-roll inflation also occurs generically in both mLQC-I and mLQC-II [74]. In particular, when the inflaton has a quadratic potential, $V(\phi) = m^2 \phi^2/2$, the probabilities for the desired slow-roll inflation not to occur are $\leq 1.12 \times 10^{-5}$, $\leq 2.62 \times 10^{-6}$, and $\leq 2.74 \times 10^{-6}$ for mLQC-I, mLQC-II and LQC, respectively.

When dealing with perturbations, another ambiguity rises in the replacement of the momentum conjugate π_a of the expansion factor a in the effective potential of the scalar perturbations. This ambiguity occurs not only in the dressed metric approach [cf. Eq.(3.11)] but also in the hybrid approach [cf. Eq.(4.3)], as it is closely related to the quantization strategy used in LQG/LQC, because now only the holonomies (complex exponentials) of π_a are defined as operators. Several choices have been proposed in the literature [36, 59, 78, 79, 82–84, 125]. In Secs. III and IV, we have shown that for some choices the effects on the power spectra are non-trivial, while for others the effects are negligible. However, even with the same choice, the relative differences in the amplitudes of the power spectra among the three different models can be as large as 100% in the infrared and intermediate regimes of the spectra, while in the UV regime the relative differences are no larger than 2%, and the corresponding power spectra are scale-invariant. Since only the modes in the UV regime are relevant to the current observations, the power spectra obtained in all the three models are consistent with current observations [113].

However, the interactions between the infrared and UV modes appearing in non-Gaussianities might provide an excellent window to observe such effects. This was initially done in LQC [57–59], and lately generalized to bouncing cosmologies [64]. It should be noted that in [64], the expansion factor a(t) near the bounce was assumed to take the form,

$$a(t) = a_B \left(1 + bt^2\right)^n,$$

where b and n are two free parameters. For example, for LQC we have n = 1/6 and $b = R_B/2$, where R_B is the Ricci scalar at the bounce [51, 52]. But, it is clear that near the bounce a(t) takes forms different from the above expression for mLQC-I/II, as one can see from Eqs.(2.54)-(2.57). Thus, it would be very interesting to study such effects in mLQC-I/II, and look for some observational signals.

Moreover, initial conditions are another subtle and important issue not only in LQC but also in mLQCs. As a matter of fact, the initial conditions consist of two parts: the initial time, and the initial conditions. Different choices of the initial times lead to different choices of the initial conditions, or vice versa. To clarify these issues, in Sections III and IV we have discussed it at length by showing the (generalized) comoving Hubble radius in each model as well as in each of the two approaches, dressed metric and hybrid. From these analyses, we have shown clearly which initial conditions can and cannot be imposed at a given initial time.

In addition, when the universe changes from contraction to expansion at the bounce, particle and entropy creations are expected to be very large, and it is crucial to keep such creations under control, so that the basic assumptions of the models are valid, including the one that the cosmological perturbations are small and can be treated as test fields propagating on the quantum homogeneous background, as assumed in both the dressed metric and hybrid approaches.

Yet, different initial conditions also affect the amplitudes and shapes of the primordial power spectra, and it would be very interesting to investigate the consistency of such obtained spectra with current observations, in particular the possible explanations to the anomalies found in the CMB data [60–62], and the naturalness of such initial conditions.

On the other hand, bouncing cosmologies, as an alternative to the cosmic inflation paradigm, have been extensively studied in the literature, see, for example, [126] and references therein. However, in such classical bounces, exotic matter fields are required in order to keep the bounce open. This in turn raises the question of instabilities of the models. On the other hand, quantum bounces found in LQC/mLQCs are purely due to

- A. Guth, Inflationary universe: A possible solution to the horizon and flatness problems, Phys. Rev. D23, 347 (1981); A.A. Starobinsky, A new type of isotropic cosmological models without singularity, Phys. Lett. B91, 99 (1980); K. Sato, First-order phase transition of a vacuum and the expansion of the universe, Mon. Not. R. Astron. Soc. 195, 467 (1981).
- S. Dodelson, Modern Cosmology (Academic Press, New York, 2003); V. Mukhanov, Physics Foundations of Cosmology (Cambridge University Press, Cambridge, 2005)); S. Weinberg, Cosmology (Oxford University Press, Oxford, 2008).
- [3] H. Kodama, M. Sasaki, Cosmological Perturbation Theory, Progr. Theoret. Phys. Suppl. 78 (1984) 1; V.F. Mukhanov, H.A. Feldman, R.H. Brandenberger, Theory of Cosmological Perturbations, Phys. Rep. 215 (1992) 203; K.A. Malik, Cosmological Perturbations in an Inflationary Universe, arXiv:astro-ph/0101563; K.A. Malik and D. Wands, Cosmological perturbations, Phys. Rept. 475 (2009) 1; L. Senatore, Lectures on Inflation, in New Frontiers in Fields and Strings, Proceedings of the 2015 Theoretical Advanced Study Institute in Elementary Particle Physics, Boulder, Colorado, 1 26 June 2015, Edited by J. Polchinski, P. Vieira and O. DeWolfe (World Scientific, Singapore, 2017) pp. 447-543.
- [4] R.H. Brandenberger, Inflationary Cosmology: Progress and Problems, arXiv:hep-th/9910410; J. Martin and R.H. Brandenberger, Trans-Planckian problem of inflationary cosmology, Phys. Rev. D63, 123501 (2001); J.C. Niemeyer and R. Parentani, Trans-Planckian dispersion relation and scale invariance of inflationary perturbations, Phys. Rev. D64, 101301(R) (2001); Corley-Jacobson dispersion relation and trans-Planckian inflation, Phys. Rev. D65, 103514 (2002); Dependence of the spectra of fluctuations in inflationary cosmology on trans-Planckian physics, Phys. Rev. D68, 063513 (2003); L. Bergstorm and U.H. Danielsson, Can MAP and Planck map Planck physics?, J. High Energy Phys.

the quantum geometric effects, and the instability problem is automatically out of the question. So, it would be very interesting to study bouncing cosmologies in the framework of LQC/mLQCs. The first step in this direction has already been taken [122], and more detailed and extensive analyses are still needed.

Acknowledgements

We are grateful to Robert Brandenberger and David Wands for valuable comments on the manuscript and helpful discussions. We thank Javier Olmedo and Tao Zhu for various discussions related to works presented in this manuscript. B.-F.L and P.S. are supported by NSF grant PHY-1454832. B.-F.L is also partially supported by the National Natural Science Foundation of China (NNSFC) with the Grants No. 12005186.

038 (2002); R. Easther, W. H. Kinney, and H. Peiris, Observing trans-Planckian signatures in the cosmic microwave background, J. Cosmol. Astropart. Phys.
 05 (2005) 009; A. E. Joras and G. Marozzi, Trans-Planckian physics from a nonlinear dispersion relation, Phys. Rev. D79, 023514 (2009); A. Ashoorioon, D. Chialva and U. Danielsson, Effects of Nonlinear Dispersion Relations on Non-Gaussianities, J. Cosmol. Astropart. Phys. 06, 034 (2011); M.G. Jackson and K. Schalm, Model Independent Signatures of New Physics in the Inflationary Power Spectrum, Phys. Rev. Lett. 108, 111301 (2012); R.H. Brandenberger and J. Martin, Trans-Planckian issues for inflationary cosmology, Class. Quantum. Grav. 30, 113001 (2013).

- [5] C. Kiefer, and M. Kramer, Quantum Gravitational Contributions to the CMB Anisotropy Spectrum, Phys. Rev. Lett. 108, 021301 (2012); L.M. Krauss and F. Wilczek, Using Cosmology to Establish the Quantization of Gravity, Phys. Rev. D89, 047501 (2014); R.P. Woodard, Perturbative Quantum Gravity Comes of Age, arXiv:1407.4748.
- [6] L. McAllister and E. Silverstein, String Cosmology: A Review, Gen. Relativ. Grav. 40, 565 (2007); C.P. Burgess, M. Cicoli, and F. Quevedo, String Inflation After Planck 2013, JCAP 1311 (2013) 003; D.F. Chernoff, S.-H. H. Tye, Inflation, String Theory and Cosmology, arXiv:1412.0579; D. Baumann and L. McAllister, Inflation and String Theory (Cambridge Monographs on Mathematical Physics, Cambridge University Press, 2015); M. Cicoli, Recent developments in string model-building and cosmology, arXiv:1604.00904; E. Silverstein, TASI lectures on cosmological observables and string theory, arXiv:1606.03640.
- [7] A. Bedroya and C. Vafa, Trans-Planckian Censorship and the Swampland, JHEP 2009, 123 (2020) [arXiv:1909.11063].
- [8] A. Bedroya, R. Brandenberger, M. Loverde and C. Vafa, *Trans-Planckian Censorship and Inflation*ary Cosmology, Phys. Rev. D101, 103502 (2020)

[arXiv:1909.11106].

- [9] R. Brandenberger, Trans-Planckian Censorship Conjecture and Early Universe Cosmology, arXiv:2102.09641.
- [10] A. Borde and A. Vilenkin, Eternal inflation and the initial singularity, Phys. Rev. Lett. **72**, 3305 (1994); A. Borde, A.H. Guth, and A. Vilenkin, Inflationary Spacetimes Are Incomplete in Past Directions, Phys. Rev. Lett. **90**, 151301 (2003).
- [11] A. Linde, On the problem of initial conditions for inflation, Found Phys 48, 1246(2018); A. Linde, Inflationary Cosmology after Planck 2013, arXiv: 1402.0526.
- [12] N. D. Birrell and P.C.W. Davies, Quantum fields in curved space (Cambridge University Press, Cambridge, 1982); R.M. Wald, Quantum Field Theory om Curved Spacetime and Black Hole Thermodynamics (The University of Chicago Press, Chicago, 1994); V.F. Mukhanov, and S. Winitzki, Introduction to Quantum Effects in Gravity (Cambridge University Press, Cambridge, 2007); L. Parker and D. Toms, Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity (Cambridge University Press, Cambridge, 2009).
- [13] S. Weinberg, in General Relativity, An Einstein Centenary Survey, edited by S.W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1980); S. Carlip, Quantum Gravity in 2+1 Dimensions, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, 2003); C. Kiefer, Quantum Gravity (Oxford Science Publications, Oxford University Press, 2007).
- [14] M.B. Green, J.H. Schwarz and E. Witten, Superstring Theory: Vol.1 & 2, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, 1999); J. Polchinski, String Theory, Vol. 1 & 2 (Cambridge University Press, Cambridge, 2001); C. V. Johson, D-Branes, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, 2003); K. Becker, M. Becker, and J.H. Schwarz, String Theory and M-Theory (Cambridge University Press, Cambridge, 2007).
- [15] A. Ashtekar and J. Lewandowski, Background independent quantum gravity: A status report, Class. Quantum Grav. 21, R53 (2004); T. Thiemann, Modern Canonical Quantum General Relativity Cambridge University Press, Cambridge, 2007); C. Rovelli, Quantum gravity (Cambridge University Press, Cambridge, 2008); M. Bojowald, Canonical Gravity and Applications: Cosmology, Black Holes, and Quantum Gravity (Cambridge University Press, Cambridge, 2011); R. Gambini and J. Pullin, A First Course in Loop Quantum Gravity (Oxford University Press, Oxford, 2011); A. Ashtekar and J. Pullin, Loop Quantum Gravity the first 30 years (World Scientific, Singapore, 2017).
- [16] A. Ashtekar and P. Singh, Loop quantum cosmology: a status report, Class. Quantum Grav. 28 (2011) 213001;
 A. Ashtekar and A. Barrau, Loop quantum cosmology: From pre-inflationary dynamics to observations, Class. Quantum Grav. 32 (2015) 234001; M. Bojowald, Quantum cosmology: a review, Rep. Prog. Phys. 78 (2015) 023901; I. Agullo and P. Singh, Loop Quantum Cosmology, in Loop Quantum Gravity: The First 30 Years, edited by A. Ashtekar and J. Pullin (Wald Scientific, Singapore, 2017).
- [17] A. Ashtekar, T. Pawlowski and P. Singh, Quantum na-

ture of the big bang, Phys. Rev. Lett. 96, 141301 (2006).

- [18] A. Ashtekar, T. Pawlowski and P. Singh, Quantum Nature of the Big Bang: An Analytical and Numerical Investigation. I., Phys. Rev. D73, 124038 (2006).
- [19] A. Ashtekar, T. Pawlowski and P. Singh, Quantum nature of the big bang: improved dynamics, Phys. Rev. D74, 084003 (2006).
- [20] A. Ashtekar, A. Corichi. and P. Singh, Robustness of key features of loop quantum cosmology, Phys. Rev. D77, 024046 (2010).
- [21] K. Giesel, B.-F. Li and P. Singh, Towards a reduced phase space quantization in loop quantum cosmology with an inflationary potential, Phys. Rev. D102, 126024 (2020).
- [22] P. Singh, Are loop quantum cosmos never singular?, Class. Quant. Grav. 26, 125005 (2009); P. Singh, Loop quantum cosmology and the fate of cosmological singularities, Bull. Astron. Soc. India 42, 121 (2014).
- [23] P. Singh, Glimpses of Space-Time Beyond the Singularities Using Supercomputers, Comput. Sci. Eng. 20, 26 (2018).
- [24] P. Diener, B. Gupt and P. Singh, Numerical simulations of a loop quantum cosmos: robustness of the quantum bounce and the validity of effective dynamics, Class. Quant. Grav. **31**, 105015 (2014).
- [25] P. Diener, B. Gupt, M. Megevand and P. Singh, Numerical evolution of squeezed and non-Gaussian states in loop quantum cosmology, Class. Quant. Grav. 31, 165006 (2014).
- [26] P. Diener, A. Joe, M. Megevand and P. Singh, Numerical simulations of loop quantum Bianchi-I spacetimes, Class. Quant. Grav. 34, 094004 (2017).
- [27] I. Agullo, A. Ashtekar and B. Gupt, *Phenomenology with fluctuating quantum geometries in loop quantum cosmology*, Class. Quantum Grav. 34, 074003 (2017).
- [28] J. L. Dupuy and P. Singh, Hysteresis and beats in loop quantum cosmology, Phys. Rev. D101, 086016 (2020).
- [29] L. Gordon, B.-F. Li and P. Singh, Quantum gravitational onset of Starobinsky inflation in a closed universe, Phys. Rev. D103, 046016 (2021).
- [30] M. Motaharfar and P. Singh, On the role of dissipative effects in the quantum gravitational onset of warm Starobinsky inflation in a closed universe, arXiv:grqc/2102.09578.
- [31] M. Bojowald, G. M. Hossain, M. Kagan, S. Shankaranarayanan, Anomaly freedom in perturbative loop quantum gravity, Phys. Rev. D78, 063547 (2008).
- [32] T. Cailleteau, A. Barrau, J. Grain, F. Vidotto, Consistency of holonomy-corrected scalar, vector and tensor perturbations in Loop Quantum Cosmology, Phys. Rev. D86, 087301 (2012).
- [33] T. Cailleteau, J. Mielczarek, A. Barrau, J. Grain, Anomaly-free scalar perturbations with holonomy corrections in loop quantum cosmology, Class. Quant. Grav. 29, 095010 (2012).
- [34] I. Agullo, A. Ashtekar and W. Nelson, *Quantum gravity extension of the inflationary scenario*, Phys. Rev. Lett. 109, 251301 (2012).
- [35] I. Agullo, A. Ashtekar and W. Nelson, Extension of the quantum theory of cosmological perturbations to the Planck era, Phys. Rev. D87, 043507 (2013).
- [36] I. Agullo, A. Ashtekar and W. Nelson, The preinflationary dynamics of loop quantum cosmology: confronting quantum gravity with observations, Class.

Quantum Grav. **30**, 085014 (2013).

- [37] M. Fernández-Méndez, G. A. Mena Marugán and J. Olmedo, *Hybrid quantization of an inflationary uni*verse, Phys. Rev. D86, 024003 (2012).
- [38] M. Fernández-Méndez, G. A. Mena Marugán and J. Olmedo, *Hybrid quantization of an inflationary model: The flat case*, Phys. Rev. D88, 044013 (2013).
- [39] L. Castelló Gomar, M. Fernández-Méndez, G. A. Mena Marugan and J. Olmedo, Cosmological perturbations in hybrid loop quantum cosmology: Mukhanov-Sasaki variables, Phys. Rev. D90, 064015 (2014).
- [40] L. Castelló Gomar, M. Martín-Benito, G. A. Mena Marugán, Gauge-invariant perturbations in hybrid quantum cosmology, JCAP 1506 (2015) 045.
- [41] F. Benítez Martínez and J. Olmedo, Primordial tensor modes of the early Universe, Phys. Rev. D93, 124008 (2016).
- [42] B. Gupt, P. Singh, Quantum gravitational Kasner transitions in Bianchi-I spacetime, Phys. Rev. D86, 024034 (2012).
- [43] B. Gupt, P. Singh, A quantum gravitational inflationary scenario in Bianchi-I spacetime, Class. Quantum Grav. 30, 145013 (2013).
- [44] I. Agullo, J. Olmedo and V. Sreenath, Predictions for the Cosmic Microwave Background from an Anisotropic Quantum Bounce, Phys. Rev. Lett. 124, 251301 (2020).
- [45] I. Agullo, J. Olmedo and V. Sreenath, Observational consequences of Bianchi I spacetimes in loop quantum cosmology, Phys. Rev. D102, 043523 (2020).
- [46] I. Agullo and N. A. Morris, Detailed analysis of the predictions of loop quantum cosmology for the primordial power spectra, Phys. Rev. D92, 124040 (2015).
- [47] B. Bonga and B. Gupt, Inflation with the Starobinsky potential in loop quantum cosmology, Gen. Relativ. Grav. 48, 1 (2016).
- [48] B. Bonga and B. Gupt, Phenomenological investigation of a quantum gravity extension of inflation with the Starobinsky potential, Phys. Rev. D93, 063513 (2016).
- [49] A. Ashtekar and B. Gupt, Quantum Gravity in the Sky: Interplay between fundamental theory and observations, Class. Quantum Grav. 34, 014002 (2017).
- [50] A. Ashtekar and B. Gupt, Initial conditions for cosmological perturbations, Class. Quant. Grav. 34, 035004 (2017).
- [51] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, Q. Sheng, Universal features of quantum bounce in loop quantum cosmology, Phys. Lett. B773 (2017) 196.
- [52] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, Q. Sheng, Preinflationary universe in loop quantum cosmology, Phys. Rev. D96, 083520 (2017).
- [53] D. Martín de Blas and J. Olmedo, Primordial power spectra for scalar perturbations in loop quantum cosmology, JCAP 1606, 029 (2016).
- [54] L. Castelló Gomar, D. Martín de Blas, G. A. Mena Marugán and J. Olmedo, Hybrid loop quantum cosmology and predictions for the cosmic microwave background, Phys. Rev. D96, 103528 (2017).
- [55] B. E. Navascues, D. M. de Blas and G. A. M. Marugan, Time-dependent mass of cosmological perturbations in the hybrid and dressed metric approaches to loop quantum cosmology, Phys. Rev. D97, 043523 (2018).
- [56] B. Elizaga Navascués, D. Martín de Blas and G. A. Mena Marugán, The vacuum state of primordial fluctuations in hybrid loop quantum cosmology, Universe 4,

98 (2018).

- [57] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, and Q. Sheng, Primoridial non-Gaussianity and power asymmetry with quantum gravitational effects in loop quantum cosmology, Phys. Rev. D97, 043501 (2018).
- [58] Q. Wu, T. Zhu, and A. Wang, Nonadiabatic evolution of primordial perturbations and non-Gaussianity in hybrid approach of loop quantum cosmology Phys. Rev. D98, 103528 (2018).
- [59] I. Agullo, B. Bolliet and V. Sreenath, Non-Gaussianity in loop quantum cosmology, Phys. Rev. D97, 066021 (2018).
- [60] Y. Akrami et al. (Planck collaboration), Planck 2018 results. I. Overview and the cosmological legacy of Planck, A&A, 641 (2020) A1.
- [61] Y. Akrami et al. (Planck collaboration), Planck 2018 results. VII. Isotropy and Statistics of the CMB, A&A, 641 (2020) A7.
- [62] D.J. Schwarz, C.J. Copi, D. Huterer, G.D. Starkman, *CMB Anomalies after Planck*, Class. Quantum Grav. 33 (2016) 184001.
- [63] A. Ashtekar, B. Gupt, D. Jeong, and V. Sreenath, Alleviating the Tension in the Cosmic Microwave Background Using Planck-Scale Physics, Phys. Rev. Lett. 125, 051302 (2020); Cosmic tango between the very small and the very large: Addressing CMB anomalies through Loop Quantum Cosmology, arXiv:2103.14568.
- [64] I. Agullo, D. Kranas, and V. Sreenath, Anomalies in the CMB from a cosmic bounce, Gen. Relativ. Grav. 53 (2021) 17; Large scale anomalies in the CMB and non-Gaussianity in bouncing cosmologies, Class. Quantum Grav. 38, 065010 (2021).
- [65] J. Yang, Y. Ding and Y. Ma, Alternative quantization of the Hamiltonian in loop quantum cosmology II: Including the Lorentz term, Phys. Lett. B682 (2009) 1.
- [66] A. Dapor and K. Liegener, Cosmological Effective Hamiltonian from full Loop Quantum Gravity Dynamics, Phys. Lett. B785 (2018) 506.
- [67] A. Dapor and K. Liegener, Cosmological Coherent State Expectation Values in LQG I. Isotropic Kinematics, Quant. Grav. 35 (2018) 135011.
- [68] T. Thiemann, Gauge Field Theory Coherent States (GCS): I. General Properties, Class. Quantum Grav. 18 (2001) 2025; T. Thiemann, O. Winkler, Gauge Field Theory Coherent States (GCS): II. Peakedness Properties, Class. Quantum Grav. 18 (2001) 2561; T. Thiemann, Complexifier coherent states for quantum general relativity, Class. Quantum Grav. 23 (2006) 2063.
- [69] T. Thiemann, Quantum Spin Dynamics (QSD), Class. Quantum Grav. 15, 839 (1998); Quantum Spin Dynamics (QSD) II, Class. Quantum Grav. 15, 875 (1998);
 K. Giesel, T. Thiemann, Algebraic Quantum Gravity (AQG) I. Conceptual Setup, Class. Quantum Grav. 24, 2465 (2007).
- [70] A. Corichi and P. Singh, Is loop quantization in cosmology unique?, Phys. Rev. D78, 024034 (2008).
- [71] M. Assanioussi, A. Dapor, K. Liegener and T. Pawlowski, Emergent de Sitter epoch of the quantum Cosmos from Loop Quantum Cosmology, Phys. Rev. Lett. 121, 081303 (2018) [arXiv:1801.00768]; Emergent de Sitter epoch of the loop quantum cosmos: A detailed analysis, Phys. Rev. D100, 084003 (2019); Challenges in recovering a consistent cosmology from the effective dynamics of loop quantum gravity, Phys. Rev. D100,

106016 (2019); K. Liegener and P. Singh, Some physical implications of regularization ambiguities in SU(2) gauge-invariant loop quantum cosmology, Phys. Rev. D100, 124049 (2019).

- [72] B.-F. Li, P. Singh, A. Wang, Towards cosmological dynamics from loop quantum gravity, Phys. Rev. D97, 084029 (2018).
- [73] B.-F. Li, P. Singh, A. Wang, Qualitative dynamics and inflationary attractors in loop cosmology, Phys. Rev. D98, 066016 (2018).
- [74] B.-F. Li, P. Singh, A. Wang, Genericness of preinflationary dynamics and probability of the desired slow-roll inflation in modified loop quantum cosmologies, Phys. Rev. D100, 063513 (2019).
- [75] S. Saini, P. Singh, Generic absence of strong singularities and geodesic completeness in modified loop quantum cosmologies, Class. Quantum Grav. 36 (2019) 105014.
- [76] S. Saini, P. Singh, Von Neumann stability of modified loop quantum cosmologies, Class. Quantum Grav. 36 (2019) 105010.
- [77] N. Dadhich, A. Joe and P. Singh, Emergence of product of constant curvature spaces in loop quantum cosmology, Class. Quantum Grav. 32 (2015) 185006.
- [78] B.-F. Li, P. Singh, A. Wang, Primordial power spectrum from the dressed metric approach in loop cosmologies, Phys. Rev. D100, 086004 (2020).
- [79] I. Agullo, Primordial power spectrum from the Dapor-Liegener model of loop quantum cosmology, Gen. Relativ. Grav. 50 (2018) 91.
- [80] A. García-Quismondo and G. A. Mena Marugán, Martin-Benito-Mena Marugan-Olmedo prescription for the Dapor-Liegener model of loop quantum cosmology, Phys. Rev. D99, 083505 (2019).
- [81] A. García-Quismondo and G. A. Mena Marugán, Dapor-Liegener formalism of loop quantum cosmology for Bianchi I spacetimes, Phys. Rev. D101, 023520 (2020).
- [82] L. Castelló Gomar, A. García-Quismondo and G. A. Mena Marugán, Primordial perturbations in the Dapor-Liegener model of hybrid loop quantum cosmology, Phys. Rev. D102, 083524 (2020).
- [83] A. García-Quismondo, G. A. Mena Marugán and G. Sánchez Pérez, The time-dependent mass of cosmological perturbations in loop quantum cosmology: Dapor-Liegener regularization, Class. Quantum Grav. 37, 195003 (2020).
- [84] B.-F. Li, J. Olmedo, P. Singh, and A. Wang, primordial scalar power spectrum from the hybrid approach in loop cosmologies, Phys. Rev. D102, 126025 (2020).
- [85] W. Kamiński, M. Kolanowski, and J. Lewandowski, Dressed metric predictions revisited, Class. Quantum Grav. 37 (2020) 095001.
- [86] K.A. Meissner, Black hole entropy in Loop Quantum Gravity, Class. Quantum Grav. 21 (2004) 5245.
- [87] A. Ashtekar, M. Bojowald, and J. Lewandowski, Mathematical structure of loop quantum cosmology, Adv. Theor. Math. Phys. 7, 233 (2003).
- [88] K. Liegener and P. Singh, Gauge-invariant bounce from loop quantum gravity, Class. Quant. Grav. 37, 085015 (2020); New Loop Quantum Cosmology Modifications from Gauge-covariant Fluxes, Phys. Rev. D 100, 124048 (2019).
- [89] D. A. Craig and P. Singh, Consistent probabilities in loop quantum cosmology, Class. Quant. Grav. 30,

205008 (2013).

- [90] J. Engle, Relating loop quantum cosmology to loop quantum gravity: Symmetric sectors and embeddings, Class. Quant. Grav. 24, 5777 (2007); J. Brunnemann and C. Fleischhack, On the configuration spaces of homogeneous loop quantum cosmology and loop quantum gravity, arXiv:0709.1621; J. Brunnemann and T. A. Koslowski, Symmetry Reduction of Loop Quantum Gravity, Class. Quant. Grav. 28, 245014 (2011).
- [91] B.-F. Li and P. Singh, Singularity resolution may be forbid a cyclic evolution. To appear.
- [92] A. Bhardwaj, E.J. Copeland, J. Louko, Inflation in Loop Quantum Cosmology, Phys. Rev. D99, 063520 (2019).
- [93] M. Shahalam, M. Sharma, Q. Wu, A. Wang, Preinflationary dynamics in loop quantum cosmology: Power-law potentials, Phys. Rev. D96 (2017) 123533.
- [94] M. Shahalam, Preinflationary dynamics of power-law potential in loop quantum cosmology, Universe 4 (2018) 87.
- [95] M. Shahalam, M. Sami, A. Wang, Pre-inflationary dynamics of α-attractor in loop quantum cosmology, Phys. Rev. D98, 043524 (2018).
- [96] M. Shahalam, M.A. Ajmi, R. Myrzakulov, A. Wang, *Revisiting pre-inflationary universe of family of αattractor in loop quantum cosmology*, Class.Quant.Grav. **37**, 195026 (2020).
- [97] M. Sharma, M. Shahalam, W. Qiang, A. Wang, Preinflationary dynamics in loop quantum cosmology: Monodromy Potential, JCAP, 11 (2018) 003.
- [98] K. Xiao and S.-Q. Wang, Pre-inflation dynamical behavior of warm inflation in loop quantum cosmology, Mod. Phys. Lett. A35(2020) 2050293.
- [99] K. Xiao, Tachyonic inflation in loop quantum cosmology, Eur. Phys. J. C79 (2019) 1019.
- [100] M. Sharma, T. Zhu, A. Wang, Background dynamics of pre-inflationary scenario in Brans-Dicke loop quantum cosmology, Commun. Theor. Phys. 71 (2019) 1205.
- [101] W.-J. Jin, Y.-G. Ma, T. Zhu, Pre-inflationary dynamics of Starobinsky inflation and its generalization in Loop Quantum Brans-Dicke Cosmology, JCAP 02 (2019) 010.
- [102] K. Xiao, Tachyon field in loop cosmology, Phys. Lett. B811 (2020) 135859.
- [103] P. Singh, K. Vandersloot and G. V. Vereshchagin, Nonsingular bouncing universes in loop quantum cosmology, Phys. Rev. D74 (2006) 043510.
- [104] X. Zhang and Y. Ling, Inflationary universe in loop quantum cosmology, JCAP, 08 (2007) 012.
- [105] A. Ashtekar and D. Sloan, Probability of Inflation in Loop Quantum Cosmology, Gen. Relativ. Gravit. 43, 3619 (2011); A. Ashtekar and D. Sloan, Loop quantum cosmology and slow roll inflation, Phys. Lett. B 694, 108 (2011)
- [106] A. Corichi and A. Karami, On the measure problem in slow roll inflation and loop quantum cosmology, Phys. Rev. D83 104006 (2011).
- [107] A. Corichi and D. Sloan, Inflationary Attractors and their Measures, Class. Quant. Grav. 31, 062001 (2014)
- [108] S. Bedic and G. Vereshchagin, Probability of inflation in Loop Quantum Cosmology, Phys. Rev. D99, no. 4, 043512 (2019)
- [109] L. Linsefors and A. Barrau, Duration of inflation and conditions at the bounce as a prediction of effective isotropic loop quantum cosmology, Phys. Rev. D87 (2013) 123509.

- [110] L. Chen and J.-Y. Zhu, Loop quantum cosmology: The horizon problem and the probability of inflation, Phys. Rev. D92 (2015) 084063.
- [111] S.W. Hawking, G.F.R. Ellis, *The Large Scale Structure of Spacetime* (Cambridge University Press, Cambridge, 1973).
- [112] D. Baumann, TASI Lectures on Inflation, arXiv:0907.5424.
- [113] C. L. Bennett et al., 4-Year COBE DMR Cosmic Microwave Background Observations: Maps and Basic Results, Astrophys. J. 464, L1 (1996); K. M. Gorski et al., Power Spectrum of Primordial Inhomogeneity Determined from the 4-Year COBE DMR Sky Maps, ibid., 464, L11 (1996); E. Komatsu et al. (WMAP Collaboration), Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation, Astrophys. J. Suppl. Ser. 192, 18 (2011); D. Larson et al. (WMAP Collaboration), Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Power Spectra and WMAP-Derived Parameters, ibid., 192, 16 (2011); P. Ade et al. (PLANCK Collaboration), Planck 2015 results. XX. Constraints on inflation, A&A 594, A20 (2016); N. Aghanim, et al. (PLANCK Collaboration), Planck 2018 results. X. Constraints on inflation, A&A 641, A10 (2020).
- [114] K.N. Abazajian et al., Inflation Physics from the Cosmic Microwave Background and Large Scale Structure, Astropart. Phys. 63, 55 (2015); K. Abazajian, et al., CMB-S4 Decadal Survey APC White Paper, arXiv:1908.01062.
- [115] E. Wilson-Ewing, Testing loop quantum cosmology, Comptes Rendus Physique 18, 207 (2017).
- [116] F. Gerhardt, D. Oriti, and E. Wilson-Ewing, The separate universe framework in group field theory condensate cosmology, Phys. Rev. D98, 066011 (2018): D. Oriti, The universe as a quantum gravity condensate, Comptes Rendus Physique 18, 235 (2017); D. Oriti, L. Sindoni, and E. Wilson-Ewing, Bouncing cosmologies from quantum gravity condensates, Class. Quantum Grav. 34, 04LT01 (2017); B. Baytas, M. Bojowald, S. Crowe, Equivalence of models in loop quantum cosmology and group field theory, Universe 5 (2019) 41.
- [117] J. Olmedo, E. Alesci, Power spectrum of primordial perturbations for an emergent universe in quantum reduced loop gravity, JCAP 04 (2019) 030; E. Alesci, G. Botta, F. Cianfrani, and S. Liberati, Cosmological singularity resolution from quantum gravity: the emergent-bouncing universe, Phys. Rev. D96, 046008 (2017); E. Alesci and

F. Cianfrani, Quantum Reduced Loop Gravity: a realistic Universe, Phys. Rev. D92, 084065 (2015); E. Alesci and F. Cianfrani, Quantum-Reduced Loop Gravity: Cosmology, Phys. Rev. D87, 083521 (2013).

- [118] J. Engle and I. Vilensky, Deriving loop quantum cosmology dynamics from diffeomorphism invariance, Phys. Rev. D98, 023505 (2018); Uniqueness of minimal loop quantum cosmology dynamics, Phys. Rev. D100, 121901 (2019).
- [119] É. Wilson-Ewing, The loop quantum cosmology bounce as a Kasner transition, Class. Quantum Grav. 35, 065005 (2018).
- [120] K. Giesel, B.-F. Li and P. Singh, Towards a reduced phase space quantization in loop quantum cosmology with an inflationary potential, Phys. Rev. D102, 126024 (2020).
- [121] K. Giesel, B.-F. Li and P. Singh, Revisiting the Bardeen and Mukhanov-Sasaki equations in the Brown-Kuchar and Gaussian dust models, arXiv:2012.14443.
- [122] B.-F. Li, S. Saini, and P. Singh, Primordial power spectrum from a matter-Ekpyrotic bounce scenario in loop quantum cosmology, arXiv:2012.10462.
- [123] M. Han and H. Liu, Effective Dynamics from Coherent State Path Integral of Full Loop Quantum Gravity, Phys. Rev. D101, 046003 (2020); Semiclassical limit of new path integral formulation from reduced phase space loop quantum gravity, Phys. Rev. D102, 024083 (2020); M. Han, H. Li, and H. Liu, Manifestly Gauge-Invariant Cosmological Perturbation Theory from Full Loop Quantum Gravity, arXiv:2005.00883.
- [124] S. Schander, T. Thiemann, Quantum Cosmological Backreactions I: Cosmological Space Adiabatic Perturbation Theory, arXiv:1906.08166; J. Neuser, S. Schander, T. Thiemann, Quantum Cosmological Backreactions II: Purely Homogeneous Quantum Cosmology, arXiv:1906.08185; S. Schander, T. Thiemann, Quantum Cosmological Backreactions III: Deparametrised Quantum Cosmological Perturbation Theory, arXiv:1906.08194.
- [125] G. A. Mena Marugán, J. Olmedo, and T. Pawlowski, Prescriptions in loop quantum cosmology: A comparative analysis, Phys. Rev. D 84, 064012 (2011).
- R. Brandenberger and P. Peter, Bouncing Cosmologies: Progress and Problems, Found Phys. 47 (2017) 797; D. Wand, Duality invariance of cosmological perturbation spectra, Phys. Rev. D60, 023507 (1999).