THE FREE m-CONE OF A MATROID AND ITS G-INVARIANT

JOSEPH E. BONIN AND KEVIN LONG

ABSTRACT. For a matroid M, its configuration determines its \mathcal{G} -invariant. Few examples are known of pairs of matroids with the same \mathcal{G} -invariant but different configurations. In order to produce new examples, we introduce the free m-cone $Q_m(M)$ of a loopless matroid M, where m is a positive integer. We show that the \mathcal{G} -invariant of M determines the \mathcal{G} -invariant of $Q_m(M)$, and that the configuration of $Q_m(M)$ determines M; so if Mand N are nonisomorphic and have the same \mathcal{G} -invariant, then $Q_m(M)$ and $Q_m(N)$ have the same \mathcal{G} -invariant but different configurations. We prove analogous results for several variants of the free m-cone. We also define a new matroid invariant of M, and show that it determines the Tutte polynomial of $Q_m(M)$.

1. INTRODUCTION

Of the following three matroid invariants, listed from weakest to strongest, we are mainly interested in the second and third (we recall them below and treat them more fully in Section 2):

- (1) the Tutte polynomial,
- (2) the \mathcal{G} -invariant (introduced by Derksen [6]), and
- (3) the configuration (introduced by Eberhardt [8]).

Having two matroid invariants, one of which can be derived from the other, raises a basic question: how can we construct matroids that share the weaker invariant but not the stronger? We address this question for the \mathcal{G} -invariant and configuration. Eberhardt proved that the configuration of a matroid M determines its Tutte polynomial. Bonin and Kung [3] strengthened this result, proving that the configuration of M determines its \mathcal{G} -invariant. We treat a construction with which we can produce matroids with the same \mathcal{G} -invariant but different configurations.

The natural starting point is the most well known matroid invariant, the Tutte polynomial. Given a matroid M = (E, r), its *Tutte polynomial* is

$$T(M; x, y) = \sum_{A \subseteq E} (x - 1)^{r(E) - r(A)} (y - 1)^{|A| - r(A)}.$$

The data in this polynomial is the multiset $\{(|A|, r(A)) : A \subseteq E\}$. The significance of the Tutte polynomial comes in part from its being a universal invariant for the deletion-contraction rule [5].

Derksen [6] introduced the \mathcal{G} -invariant, which generalizes the Tutte polynomial. Derksen and Fink [7] later showed that the \mathcal{G} -invariant is a universal valuative invariant for subdivisions of matroid base polytopes. For a matroid M = (E, r) where |E| = n, the data in its \mathcal{G} -invariant is the multiset of *n*-tuples of rank increases

$$(r(\{e_1\}), r(\{e_1, e_2\}) - r(\{e_1\}), r(\{e_1, e_2, e_3\}) - r(\{e_1, e_2\}), \dots, r(E) - r(E - \{e_n\}))$$

Date: January 28, 2022.

²⁰²⁰ Mathematics Subject Classification. 05B35.

over all permutations e_1, e_2, \ldots, e_n of E. Bonin and Kung [3] showed that this data is equivalent to the multiset of (r(E) + 1)-tuples

$$(|F_0|, |F_1 - F_0|, |F_2 - F_1|, \dots, |F_{r(E)} - F_{r(E)-1}|)$$

over all flags, that is, chains $(F_0, F_1, \ldots, F_{r(E)})$ of flats of M where $r(F_i) = i$.

Eberhardt [8] showed that, for a matroid with no coloops, the Tutte polynomial can be computed from a small amount of data about the cyclic flats (the flats that are unions of circuits), namely, from the abstract lattice formed by the cyclic flats, along with the size and rank of the cyclic flat corresponding to each element in this lattice. This data is the configuration of the matroid. Eberhardt's result was extended in Bonin and Kung [3], who showed that the configuration of a matroid with no coloops determines its \mathcal{G} -invariant. (Extending this to all matroids by also recording the number of coloops is routine.)

We develop a construction that yields examples that show that the configuration is strictly stronger than the \mathcal{G} -invariant: we show how to construct pairs of matroids with the same \mathcal{G} -invariant and different configurations. (When we started this work, the only other such examples we knew of were those treated in [3], namely, for $k \ge 4$, rank-kDowling matroids based on nonisomorphic groups of the same order. Since then, a very different technique for constructing such examples has been developed; see [1].) Let M be a loopless matroid and m be a positive integer. We will define a matroid $Q_m(M)$ that we call the free m-cone of M. Our main result is that if M and N are nonisomorphic and have the same \mathcal{G} -invariant, then $Q_m(M)$ and $Q_m(N)$ have the same \mathcal{G} -invariant and different configurations.

We use the notation and terminology in Oxley [10]. More specialized background is treated in Section 2. In Section 3, we define the free m-cone and develop some of its properties, which we then use to prove that the \mathcal{G} -invariant of M determines that of its free m-cone, and that the configuration of the free m-cone of M determines M. In Section 4, we treat several variants of the free m-cone and show that, with some exceptions for small m, our main results in Section 3 also hold for them. The Higgs lift of a matroid is a special case of these variants. In Section 5, we introduce a matroid invariant that lies between the Tutte polynomial and the \mathcal{G} -invariant, and show that it determines the Tutte polynomial of the free m-cone.

We use [n] to denote the set $\{1, 2, \ldots, n\}$.

2. BACKGROUND

Given a rank-k matroid M on E = [n] with rank function r and a permutation π on E, the rank sequence $\underline{r}(\pi) = r_1 r_2 \dots r_n$ is given by $r_1 = r(\pi(1))$ and, for i > 1,

$$r_i = r(\{\pi(j) : j \in [i]\}) - r(\{\pi(j) : j \in [i-1]\}).$$

Thus, $\{\pi(i) : r_i = 1\}$ is a basis of M. Each rank sequence is an (n, k)-sequence, that is, a sequence of k ones and n - k zeroes. For each (n, k)-sequence \underline{r} , let $[\underline{r}]$ be a formal symbol, and let $\mathcal{G}(n, k)$ be the vector space over a field of characteristic zero consisting of all formal linear combinations of such symbols. The \mathcal{G} -invariant of M is defined by

$$\mathcal{G}(M) = \sum_{\pi} [\underline{r}(\pi)]$$

where the sum is over all permutations π of E.

Another perspective on $\mathcal{G}(M)$ was developed in [3]. An (n, k)-composition is an integer sequence $(a_i) = (a_0, a_1, \ldots, a_k)$ for which $a_0 + a_1 + \cdots + a_k = n$ where $a_0 \ge 0$ and $a_i > 0$ for $i \in [k]$. Let the matroid M be as above. A flag of M is a sequence




FIGURE 1. Two rank-3 sparse paving matroids.

 $(X_i) = (X_0, X_1, \ldots, X_k)$ where X_i is a rank-*i* flat of M, and $X_i \subset X_{i+1}$ for i < k. The *composition* of a flag (X_i) is (a_i) where $a_0 = |X_0|$ and $a_i = |X_i - X_{i-1}|$ for $i \in [k]$. Thus, (a_i) is an (n, k)-composition. Let $\nu(M; (a_i))$ be the number of flags of M with composition (a_i) . The *catenary data* of M is the $\binom{n}{k}$ -dimensional vector $(\nu(M; (a_i)))$ indexed by (n, k)-compositions.

Bonin and Kung [3] defined a special basis of $\mathcal{G}(n, k)$, the γ -basis, whose vectors $\gamma((a_i))$ are indexed by (n, k)-compositions (a_i) . The change-of-basis result from [3], stated next, connects $\mathcal{G}(M)$ and the catenary data of M.

Theorem 2.1. The catenary data of M determines $\mathcal{G}(M)$ and conversely since

$$\mathcal{G}(M) = \sum_{(a_i)} \nu(M; (a_i))\gamma((a_i)).$$

Given a matroid M = (E, r), a subset A of E is *cyclic* if M|A has no coloops. Just as lines and planes refer to flats of ranks 2 and 3, respectively, if such flats are cyclic, then we call them cyclic lines or cyclic planes. More generally, a cyclic flat is a flat that is cyclic. The set $\mathcal{Z}(M)$ of cyclic flats of M is a lattice under inclusion. The configuration, introduced in Eberhart [8], is the abstract lattice of cyclic flats of M, without the sets but with their size and rank. More precisely, the *configuration* of a matroid M with no coloops is the triple (L, s, ρ) , where L is a lattice and s and ρ are functions with domain L where there is an isomorphism $\phi : L \to \mathcal{Z}(M)$ for which $s(x) = |\phi(x)|$ and $\rho(x) = r(\phi(x))$ for all $x \in L$. Many nonisomorphic matroids can have the same configuration: O. Giménez constructed n! non-paving matroids of rank 2n + 2 on 4n + 5 elements, all with the same configuration [2, see Theorem 5.7].

Bonin and Kung [3] showed that if M has no coloops, then $\mathcal{G}(M)$ can be found from the configuration of M. If M has coloops, $\mathcal{G}(M)$ can be found from $\mathcal{G}(M \setminus X)$ and |X| where X is the set of coloops, so we focus on matroids without coloops.

In Figures 1 and 2, we give an example of nonisomorphic matroids M_1 and M_2 that have the same configuration. (We will refer to this example throughout the paper.) It follows that M_1 and M_2 have the same \mathcal{G} -invariant, which is

$$\mathcal{G}(M_1) = \mathcal{G}(M_2) = 648[111000] + 72[110100].$$

They then have the same catenary data:

$$\nu(M_1; 0, 1, 2, 3) = \nu(M_2; 0, 1, 2, 3) = 6$$
 and $\nu(M_1; 0, 1, 1, 4) = \nu(M_2; 0, 1, 1, 4) = 18$.

Brylawski [4] showed that a matroid is determined by its cyclic flats and their ranks, which is more data than the configuration gives. We will use the following related result of Sims [11] and Bonin and de Mier [2].

Theorem 2.2. For a set Z of subsets of a set E and a function $r : Z \to \mathbb{Z}$, there is a matroid M on E with Z(M) = Z and $r_M(X) = r(X)$ for all $X \in Z$ if and only if

J. BONIN AND K. LONG




FIGURE 2. Parts (a) and (b) show the lattice of cyclic flats of M_1 and M_2 . Replacing each set by just its size and rank gives the configuration of M_1 and M_2 , shown in part (c).

(Z0) (\mathcal{Z}, \subseteq) is a lattice, (Z1) $r(0_{\mathcal{Z}}) = 0$, where $0_{\mathcal{Z}}$ is the least set in \mathcal{Z} , (Z2) 0 < r(Y) - r(X) < |Y - X| for all $X, Y \in \mathcal{Z}$ with $X \subsetneq Y$, and (Z3) for all sets X, Y in \mathcal{Z} (or, equivalently, just incomparable sets in \mathcal{Z}),

(2.1)
$$r(X \lor Y) + r(X \land Y) + |(X \cap Y) - (X \land Y)| \le r(X) + r(Y).$$

3. The Free m-Cone

Let M = (E, r) be a rank-k loopless matroid. For each integer $m \ge 1$, we will define a rank-(k + 1) matroid $Q_m(M)$, which we often shorten to Q, on a set of (m + 1)|E| + 1elements. The construction is illustrated in Figure 3. For each $e \in E$, let T_e be a set of size m that is disjoint from E and from all other sets $T_{e'}$, and let T be the union of these |E|sets. Let a be an element not in $E \cup T$, which we call the *tip* of Q. Let $E(Q) = E \cup T \cup \{a\}$.

Define $q: 2^E \to 2^{E(Q)}$ as follows: for $S \subseteq E$,

$$q(S) = S \cup \{a\} \cup \left(\bigcup_{e \in S} T_e\right).$$

Thus, $q(\emptyset) = \{a\}$. In the examples in Figure 3, if $e \in [6]$, then $q(\{e\})$ is the 3-point line $\{e, \overline{e}, a\}$. Define $p: 2^{E(Q)} \to 2^E$ as follows: for $S \subseteq E(Q)$,

$$p(S) = (S \cap E) \cup \{e : S \cap T_e \neq \emptyset\}.$$

For $e \in E$ and $x \in T_e$, we have $p(\{e\}) = \{e\} = p(\{x\})$. Henceforth we omit the braces when applying p and q to singleton sets. The set q(S) will be the ground set of the free m-cone of M|S, so, for convenience, we refer to q(S) as the *cone of* S.

Let $\mathcal{Z}(Q) = \mathcal{Z}(M) \cup \{q(F) : F \text{ is a nonempty flat of } M\}$. In each example in Figure 3, besides the cyclic flats of the original matroid (\emptyset , two 3-point lines, and E), this set contains the six 3-point lines that contain a, the cone of each 2- and 3-point line of the original matroid (e.g., $\{3, \overline{3}, 6, \overline{6}, a\}$ and $\{1, \overline{1}, 2, \overline{2}, 3, \overline{3}, a\}$), and q(E). Define $r_Q : \mathcal{Z}(Q) \to \mathbb{Z}$ by

(3.1)
$$r_Q(Z) = \begin{cases} r(Z), & \text{if } Z \in \mathcal{Z}(M) \\ r(F) + 1, & \text{if } Z = q(F) \text{ where } F \text{ is a nonempty flat of } M. \end{cases}$$

We next show that $\mathcal{Z}(Q)$ and r_Q define a matroid on E(Q). We denote this matroid by $Q_m(M)$ and call it the *free m-cone* of M.

Theorem 3.1. Let M = (E, r) be a loopless matroid. The set $\mathcal{Z}(Q)$ and the function r_Q defined above satisfy axioms (Z0)–(Z4) in Theorem 2.2 and so define a matroid on E(Q).




FIGURE 3. The free 1-cones of the matroids M_1 and M_2 , on E = [6], in Figure 1. Each of M_1 and M_2 is shown in a face of a tetrahderon. The tip a is at the vertex opposite that face. For each $e \in [6]$, the set T_e is $\{\overline{e}\}$.

Proof. Since $\emptyset \in \mathbb{Z}(M)$, the least set in $\mathbb{Z}(Q)$ is \emptyset , so property (Z1) holds by Equation (3.1). Each of properties (Z0), (Z2), and (Z3) involves two sets in $\mathbb{Z}(Q)$. When both sets are in $\mathbb{Z}(M)$, the properties hold since M is a matroid and $r_Q(X) = r(X)$ for such sets, so we focus on the case having at least one set not in $\mathbb{Z}(M)$. We use \vee and \wedge for the join and meet operations of $\mathbb{Z}(Q)$, and \vee_M and \wedge_M for those of $\mathbb{Z}(M)$. We use cl for the closure operator of M.

Let X and Y be distinct sets in $\mathcal{Z}(Q) - \mathcal{Z}(M)$, so X = q(X') and Y = q(Y') where X' and Y' are nonempty flats of M. Now $X \cap Y = q(X' \cap Y')$, which is in $\mathcal{Z}(Q)$ unless the flat $X' \cap Y'$ of M is empty. So $X \wedge Y = q(X' \cap Y')$ unless $X' \cap Y' = \emptyset$, in which case $X \wedge Y = \emptyset$. Since $X \vee Y$ can only be the cone of a flat of M, and the least flat of M that contains X' and Y' is $cl(X' \cup Y')$, we have $X \vee Y = q(cl(X' \cup Y'))$. For property (Z3), submodularity in M gives $r(cl(X' \cup Y')) + r(X' \cap Y') \leq r(X') + r(Y')$. If $X' \cap Y' \neq \emptyset$, then $(X \cap Y) - (X \wedge Y) = \emptyset$, so Inequality (2.1) follows by Equation (3.1). If $X' \cap Y' = \emptyset$, then $X \cap Y = \{a\}, X' \wedge Y' = \emptyset$, and $r(cl(X' \cup Y')) \leq r(X') + r(Y')$, and Inequality (2.1) follows in this case too by Equation (3.1). For property (Z2), we assume, in addition, that $X \subsetneq Y$. Thus, $X' \subsetneq Y'$. Equation (3.1) gives $r_Q(Y) - r_Q(X) = r(Y') - r(X') > 0$. Since |Y - X| = (m + 1)|Y' - X'|, property (Z2) follows.

Now assume that $X \in \mathcal{Z}(M)$ and $Y \in \mathcal{Z}(Q) - \mathcal{Z}(M)$; let Y = q(Y') where Y' is a nonempty flat of M. Now $X \cap Y = X \cap Y'$, which, as an intersection of flats of M, is a flat of M. Let C be the set of coloops of $M|X \cap Y$. Now $(X \cap Y) - C$ is a cyclic flat of M, and it contains each cyclic flat that is a subset of X and Y, so it is $X \wedge Y$. Also,

$$r_Q(X \wedge Y) = r(X \cap Y) - |C| = r(X \cap Y) - |(X \cap Y) - (X \wedge Y)|.$$

Since $X \vee Y$ can only be the cone of a flat of M, and the least flat of M that contains X and Y' is $cl(X \cup Y')$, we have $X \vee Y = q(cl(X \cup Y'))$. This completes the proof of property (Z0). To finish the proof of property (Z3), Inequality (2.1) follows since

$$\begin{aligned} r_Q(X \lor Y) + r_Q(X \land Y) + |(X \cap Y) - (X \land Y)| &= r(X \cup Y') + 1 + r(X \cap Y') \\ &\leq r(X) + r(Y') + 1 \\ &= r_Q(X) + r_Q(Y). \end{aligned}$$

For property (Z2), assume, in addition, that $X \subsetneq Y$. Property (Z2) holds if Y = q(X)since $r_Q(q(X)) - r_Q(X) = 1 < 1 + m|X| = |q(X) - X|$. If $Y \neq q(X)$, then property (Z2) follows from this and the inequality $0 < r_Q(Y) - r_Q(q(X)) < |Y - q(X)|$ proven in the previous paragraph. \Box

This result justifies extending r_Q , as defined in equation (3.1), so that r_Q is the rank function of the matroid $Q_m(M)$.

If M is binary, graphic, or transversal, $Q_m(M)$ might not share these properties. However, if M is representable over \mathbb{R} , then so is $Q_m(M)$, and if M is representable over some field of characteristic p, then so is $Q_m(M)$, although not necessarily over the same field of characteristic p. This holds since another way to define $Q_m(M)$ (less suited to our purposes) is via iterated principal extensions of the direct sum of M and the rank-1 matroid on the tip, adding points freely to the lines $cl_M(\{a, e\})$ for $e \in E(M)$.

Note that in the examples in Figure 3, and their generalization for any $m \ge 1$, each cyclic line in $Q_m(M_1)$ is in a cyclic plane that contains four cyclic lines, but that fails for $Q_m(M_2)$, so the configurations differ. Our results below imply that $\mathcal{G}(Q_m(M_1))$ and $\mathcal{G}(Q_m(M_2))$ are equal. So, if $m \ge 1$, then $Q_m(M_1)$ and $Q_m(M_2)$ have the same \mathcal{G} -invariant and different configurations. Our main theorem generalizes this example.

Theorem 3.2. Let M and N be nonisomorphic loopless matroids with $\mathcal{G}(M) = \mathcal{G}(N)$. For all $m \ge 1$, $Q_m(M)$ and $Q_m(N)$ have the same \mathcal{G} -invariant and different configurations.

This result can be iterated, producing $Q_{m_1}(Q_m(M))$ and $Q_{m_1}(Q_m(N))$, and so on.

We prove Theorem 3.2 in two parts: in Theorem 3.8, we show that the configuration of $Q_m(M)$ determines M; in Theorem 3.10, we show that $\mathcal{G}(M)$ determines $\mathcal{G}(Q_m(M))$. We first treat some preliminary results that enter into the proofs of both theorems. By Theorem 2.1, $\mathcal{G}(M)$ is equivalent to the catenary data of M. We will show that the catenary data of M determines that of $Q_m(M)$. To do so, we use the next few results to characterize the flats of $Q_m(M)$, which then allows us to characterize its flags.

Lemma 3.3. Fix $m \ge 1$. Let M = (E, r) be a loopless matroid and let $Q = Q_m(M)$.

- (i) If F is a flat of Q and $a \notin F$, then all elements of F E are coloops of Q|F.
- (ii) All flats of M are flats of Q and the restriction Q|E is M.

Proof. For any circuit C with $C \not\subseteq E$, its closure $cl_Q(C)$ is a cyclic flat and $cl_Q(C) \not\subseteq E$, so $a \in cl_Q(C)$. So for any flat F of Q with $a \notin F$, no circuit of Q|F contains elements of F - E, so statement (i) follows.

If $e \in E$ and $x \in T_e$, then $a \in q(e) \subseteq \operatorname{cl}_Q(\{x, e\})$, so the only flat of Q that properly contains E is E(Q). Let Y be the (perhaps empty) set of coloops of M, so E - Y is in $\mathcal{Z}(M)$ and so in $\mathcal{Z}(Q)$. Thus, $r(E) = r(E - Y) + |Y| = r_Q(E - Y) + |Y| \ge r_Q(E)$. Since $\operatorname{cl}_Q(E \cup a) = E(Q)$ and Q has rank r(E) + 1, it follows that $r(E) = r_Q(E)$ and that E is a flat of Q. If F is a flat of M, then $F = q(F) \cap E$ and, as an intersection of flats of Q, it is a flat of Q, so the first part of statement (ii) holds. By considering chains of flats, we now get $r(F) = r_Q(F)$ for all flats F of M.

If the second part failed, then there would be a flat X of Q|E of minimal rank, say *i*, that is not a flat of M. Now i > 0. Let Y be a flat of rank i - 1 with $Y \subsetneq X$. Fix x in X - Y. Only one flat of rank *i* in Q contains $Y \cup x$, and $cl_M(Y \cup x)$ has this property, so $X = cl_M(Y \cup x)$, contrary to X not being a flat of M. This completes the proof. \Box

Having $r(X) = r_Q(X)$ for $X \subseteq E$ allows us to simplify the notation by using r for the rank function of Q.

The next lemma is useful for describing the flats of Q. While we will apply it to the free m-cone of a matroid, the lemma holds for any matroid.

Lemma 3.4. Let L_1, L_2, \ldots, L_j be distinct lines in a matroid N, each containing the rank-1 flat $\{a\}$. Fix sets $X = \{x_1, x_2, \ldots, x_j\}$ and $Y = \{y_1, y_2, \ldots, y_j\}$ where $x_i, y_i \in L_i - a$ for each $i \in [j]$. Then $X \cup a$ is independent if and only if $Y \cup a$ is independent. Thus, if F is a flat of N with $a \notin F$ and $X \cup Y \subseteq F$, then X is independent if and only if Y is independent.

Proof. Let $F = cl(L_1 \cup L_2 \cup \cdots \cup L_j)$. If $X \cup a$ is independent, then it is a basis of N|F. Since $F \subseteq cl(Y \cup a)$ and $|X \cup a| = |Y \cup a|$, it follows that $Y \cup a$ is also a basis of N|F, and so is independent. The result follows by symmetry. \Box

Corollary 3.5. Let $Q = Q_m(M) = (E(Q), r)$ for some $m \ge 1$. For $S \subseteq E(Q)$,

(3.2)
$$r(p(S)) = \begin{cases} r(S), & \text{if } a \notin cl(S), \\ r(S) - 1, & \text{if } a \in cl(S). \end{cases}$$

Proof. Assume that $a \notin cl(S)$. No two elements of S are colinear with a, so by Lemma 3.4, a subset U of S is independent if and only if p(U) is independent. Therefore r(S) = r(p(S)). Now assume that $a \in cl(S)$. Let B be a basis of $Q|S \cup a$ with $a \in B$. So p(B-a) is independent by Lemma 3.4, so $r(p(S)) \ge r(S) - 1$. Now r(p(S)) < r(S) since $p(S) \subseteq cl(S)$ and $a \in cl(S) - cl(p(S))$. Thus, r(p(S)) = r(S) - 1. \Box

The next lemma characterizes the flats of Q. Recall that $T = E(Q) - (E \cup a)$ is the set of elements of Q that are neither the tip, a, nor in M.

Lemma 3.6. Let $Q = Q_m(M)$ for some $m \ge 1$. The flats F of Q that contain a are the cones of flats in M; indeed, $F = q(F \cap E)$. If $a \notin F$, then F is a flat of Q if and only if

- (1) $F \cap E$ is a flat of M,
- (2) $p(x) \neq p(y)$ for all $x, y \in F$ with $x \neq y$, and
- (3) for each basis B of $M|F \cap E$, the set $B \cup p(F \cap T)$ is independent.

Proof. If F is a flat of Q with $a \in F$, then $F \cap E$ (an intersection of flats) is a flat of Q and of M, and F is its cone. The converse holds by the definition of $\mathcal{Z}(Q)$.

Let F be a flat of Q not containing a. Condition (1) holds since $F \cap E$ (an intersection of flats of Q) is a flat of Q, and so of M. If p(x) = p(y) for some $x, y \in F$ with $x \neq y$, then $a \in cl(\{x, y\}) \subseteq F$, which is a contradiction, so condition (2) holds. Let B be a basis of $M|F \cap E$. By Lemma 3.3, since $a \notin F$, elements in $F \cap T$ are coloops of Q|F, so $B \cup (F \cap T)$ is independent. Therefore $B \cup p(F \cap T)$ is independent by Lemma 3.4, so condition (3) holds.

For the converse, take $F \subseteq E(Q)$ with $a \notin F$ satisfying conditions (1)–(3). Assume that F is not a flat of Q. Then there is a circuit C of Q and element $x \in C - F$ with $C - x \subseteq F$. By condition (1), $F \cap E$ is a flat of M, and hence of Q, so $C - x \notin E$. Now cl(C) is a cyclic flat and $cl(C) \notin E$, so $a \in cl(C)$, and so $a \in cl(C - x)$. Corollary 3.5 now gives r(p(C - x)) = |C - x| - 1. This contradicts condition (3) since condition (2) gives |p(C - x)| = |C - x|.

Corollary 3.7. Fix $m \ge 1$. Let M be a rank-k loopless matroid and let $Q = Q_m(M)$. Let $(Y_i) = (Y_0, Y_1, \ldots, Y_{k+1})$ be a flag in Q. Fix $j \in [k+1]$. If $a \in Y_{j-1}$, then $Y_{j-1} = q(X_{j-2})$ and $Y_j = q(X_{j-1})$ where X_{j-2} and X_{j-1} are flats of M of rank j-2 and j-1 respectively with $X_{j-2} \subset X_{j-1}$. If $a \notin Y_{j-1}$, then Y_j satisfies exactly one of the following conditions:

(i)
$$Y_j = \operatorname{cl}(Y_{j-1} \cup a) = q(\operatorname{cl}(p(Y_{j-1}))),$$

(ii) $Y_j - E = Y_{j-1} - E$ and $r(Y_j \cap E) = r(Y_{j-1} \cap E) + 1,$

(iii) $Y_j = Y_{j-1} \cup x$, where $x \in T$ and $p(x) \notin cl(p(Y_{j-1}))$. Case (i) occurs when Y_j is the first flat in (Y_i) that contains a.

Proof. Corollary 3.5 and Lemma 3.6 give the result when $a \in Y_{j-1}$. Now assume that $a \notin Y_{j-1}$. Then $r(Y_{j-1}) = r(p(Y_{j-1}))$ by Corollary 3.5. Option (i) accounts for the unique rank-*j* flat that contains $Y_{j-1} \cup a$. Now assume $a \notin Y_j$. If some $x \in Y_j - Y_{j-1}$ is in *T*, then *x* is a coloop in $Q|Y_j$ by Lemma 3.3, so $Y_j = Y_{j-1} \cup x$ and $p(x) \notin cl(p(Y_{j-1}))$. Finally, if $Y_j - Y_{j-1} \subseteq E$, then $Y_j - E = Y_{j-1} - E$.

We now prove the first result into which we divide the proof of Theorem 3.2.

Theorem 3.8. Fix $m \ge 1$. Let M = (E, r) be a loopless matroid and Q be its free m-cone. The configuration of Q determines M.

Proof. The cases with $r(M) \leq 2$ are easy, so assume that r(M) > 2. Let \mathcal{F}_1 be the set of rank-1 flats of M. Consider sets \mathcal{C} of lines in $\mathcal{Z}(Q)$ that satisfy

- (L1) for each $L \in C$, at most one proper, nonempty subset of L is in $\mathcal{Z}(Q)$,
- (L2) $\bigvee_{L \in \mathcal{C}} L = E(Q)$, and

(L3) if $L_1, L_2 \in \mathcal{C}$ with $L_1 \neq L_2$, then $r(L_1 \vee L_2) = 3$.

By Lemma 3.6, each cyclic line of Q is either a cone q(F) with $F \in \mathcal{F}_1$ (and so contains a) or a cyclic line of M. Let \mathcal{L} be the set of all lines of Q that contain a. From the definition of Q, it follows that \mathcal{L} satisfies properties (L1)–(L3). If L is a line with $L \subseteq E$, then there is an $F \in \mathcal{F}_1$ disjoint from L (since r(M) > 2) and $r(L \lor q(F)) > 3$. So \mathcal{L} is a maximal set with these properties.

Let C be a set of lines satisfying properties (L1)–(L3) with $C \not\subseteq L$. Fix $L \in C$ with $L \subseteq E$. By property (L2), C must contain a line J with $a \in J$. We claim that J is the only such line. To see this, assume that $J' \in C - \{J\}$ and $a \in J'$. Let $J \cap E = P$ and $J' \cap E = P'$. Any line of M that is coplanar with both J and J' must contain $P \cup P'$. Exactly one line of M contains $P \cup P'$, so L is the only line of M that is coplanar with both J and J' must containing a that L is coplanar with are q(cl(x)) for $x \in L$, so C would fail property (L2). So $L' \subseteq E$ for each $L' \in C - \{J\}$.

Each $F \in \mathcal{F}_1$ is contained in a unique line in \mathcal{L} , so $|\mathcal{L}| = |\mathcal{F}_1|$. Since all lines in \mathcal{C} are cyclic, by property (L1) each line in \mathcal{C} contains at least three rank-1 flats. By property (L3), if two lines of \mathcal{C} , say L_1 and L_2 , contain some $F \in \mathcal{F}_1$, then all lines in \mathcal{C} contain F since for any such line L, the planes $\operatorname{cl}(L_1 \cup L)$ and $\operatorname{cl}(L_2 \cup L)$ intersect in the line L and contain the point F. If $F \in \mathcal{F}_1$ is in all lines in \mathcal{C} , then $|\mathcal{C}| \leq \frac{|\mathcal{F}_1| - 1}{2} + 1$, otherwise $|\mathcal{C}| \leq \frac{|\mathcal{F}_1|}{3} + 1$ (the +1 is for J). So $|\mathcal{L}| > |\mathcal{C}|$. So \mathcal{L} is the unique set of largest size satisfying properties (L1)–(L3).

We can detect properties (L1)–(L3) from the configuration, so we can identify \mathcal{L} in the configuration of Q. The configuration does not give the sets in $\mathcal{Z}(Q)$, but for each $X, Y \in \mathcal{Z}(Q)$, it gives us |X|, |Y|, and whether X and Y are comparable. To get M up to isomorphism, first pick pairwise disjoint sets X_L , one for each $L \in \mathcal{L}$, where if there is a (necessarily unique) nonempty cyclic flat Y with Y < L, then $|X_L| = |Y|$, otherwise $|X_L| = 1$. The cyclic flats in $\mathcal{Z}(Q) - \mathcal{Z}(M)$ are the cones of nonempty flats of M, so the flats of M, up to isomorphism, are, for each $F \in \mathcal{Z}(Q)$, the union of the sets X_L such that $L \in \mathcal{L}$ and $L \leq F$ (note that \emptyset is such a union).

To complete the proof of Theorem 3.2, we work mainly with the catenary data, which, by Theorem 2.1, is equivalent to the \mathcal{G} -invariant. We first develop one of the keys to the

8

proof: for a size-n, rank-k loopless matroid M, we define a bijection from a certain set of 4-tuples, one component of which is a flag of M, onto the set of flags of its free m-cone Q.

To motivate the bijection, which appears in the lemma below, we first consider how the flags of Q relate to those of M. If $(Y_i) = (Y_0, Y_1, \ldots, Y_{k+1})$ is a flag of Q, then there is an integer h with $0 \le h \le k$ so that $a \in Y_i$ if and only if i > h. By Lemma 3.6, each Y_i with i > h is the cone $q(X_{i-1})$ of a flat X_{i-1} of M; the flats X_h, \ldots, X_k are the end of a flag of M. Let $|Y_h - E| = b$. The elements of $Y_h - E$ are coloops of $Q|Y_h$ by Lemma 3.3, so the list $Y_0 \cap E, Y_1 \cap E, \ldots, Y_h \cap E$ contains h - b + 1 distinct flats of M, say $X_0, X_1, \ldots, X_{h-b}$; these flats begin a flag of M. To complete $X_0, X_1, \ldots, X_{h-b}$ and X_h, \ldots, X_k to a flag of M, consider the elements of $Y_h - E$, say e_1, e_2, \ldots, e_b , listed in the order in which each first appears in the flag (Y_i) . For each $j \in [b]$, let $X_{h-b+j} = cl(X_{h-b} \cup p(\{e_1, e_2, \ldots, e_j\}))$. From Corollary 3.5, it follows that $(X_i) = (X_0, X_1, \ldots, X_k)$ is a flag of M. Each e_j has $p(e_j) \in X_{h-b+j} - X_{h-b+j-1}$.

Lemma 3.9. Fix $m \ge 1$. Let M be a size-n, rank-k loopless matroid and let Q be $Q_m(M)$. Let \mathcal{T} be the set of all 4-tuples $((X_i), h, C, (e_j))$ where

- (X_i) is a flag of M,
- *h* is an integer with $0 \le h \le k$,
- C is a subset of [h], and
- $(e_i) = (e_1, e_2, \dots, e_{|C|})$ is a |C|-tuple of elements of T where, for i with $1 \le i \le |C|$, we have $p(e_i) \in X_{h-|C|+i} X_{h-|C|+i-1}$.

Write the set C in a 4-tuple in \mathcal{T} as $\{c_1, c_2, \ldots, c_{|C|}\}$ with $c_1 < c_2 < \cdots < c_{|C|}$, and the difference [h] - C as $D = \{d_1, d_2, \ldots, d_{h-|C|}\}$ with $d_1 < d_2 < \cdots < d_{h-|C|}$. The *j*th element of C refers to c_j , and likewise for D. Let \mathcal{F} be the set of flags of Q. Define $f: \mathcal{T} \to \mathcal{F}$ by $f((X_i), h, C, (e_j)) = (Y_i)$ where

$$Y_{i} = \begin{cases} \varnothing, & \text{if } i = 0, \\ Y_{i-1} \cup e_{j}, & \text{if } i \text{ is the } j\text{th element of } C, \\ Y_{i-1} \cup X_{j}, & \text{if } i \text{ is the } j\text{th element of } D, \\ q(X_{i-1}), & \text{if } h < i \leq k+1. \end{cases}$$

Then the map f is a bijection.

Proof. We first show that (Y_i) is indeed a flag of Q. Clearly $Y_i \subsetneq Y_{i+1}$, so we focus on showing that each Y_i is a flat of Q and $r(Y_i) = i$. This is clear for Y_0 as well as for Y_i with $h < i \le k + 1$. For $0 < i \le h$, we show that Y_i satisfies the conditions in Lemma 3.6, and so is a flat. Condition (1) is immediate. Condition (2) holds since (i) $Y_h - E = \{e_1, e_2, \ldots, e_{|C|}\}$, and (ii) $p(e_j)$ is in $X_{h-|C|+j} - X_{h-|C|+j-1}$, and all |C| such differences are disjoint from $X_{h-|C|}$ and each other. For condition (3), the intersection $Y_i \cap E$ is X_l for some l with $0 \le l \le h - |C|$, and each element e_j in Y_i has $p(e_j) \in$ $X_{h-|C|+j} - X_{h-|C|+j-1}$, so for any basis B of $M|Y_i \cap E$, we get $B \cup p(Y_i \cap T)$ by adjoining to B a sequence of elements, none of which is in the closure of B and the earlier elements in the sequence. Thus, $B \cup p(Y_i \cap T)$ is independent, so condition (3) holds. So Y_i is a flat. If i is the j-th element of C, then e_j is a coloop of Y_i ; if i is the j-th element of D, then $Y_i \cap T = Y_{i-1} \cap T$, and $Y_i \cap E = X_j$ and $Y_{i-1} \cap E = X_{j-1}$; both options give $r(Y_i) = r(Y_{i-1}) + 1$, and so we get $r(Y_i) = i$ by induction.

The paragraph before the proof in effect treats the inverse: map a flag (Y_i) of Q to the 4-tuple $((X_i), h, C, (e_j))$ where $(X_i), h$, and (e_j) are as stated in that paragraph and $C = \{i : Y_i - Y_{i-1} = \{e_j\}$ for some $j\}$. So f is a bijection.

We next prove the result that completes the proof of Theorem 3.2.

Theorem 3.10. Fix $m \ge 1$. Let M be a size-n, rank-k loopless matroid and let Q be $Q_m(M)$. Then $\mathcal{G}(Q)$ can be determined from $\mathcal{G}(M)$.

Proof. As stated above, we work mainly with the catenary data, and for that we use the bijection f in Lemma 3.9. More precisely, from the composition of the flag (X_i) of M and the other three entries in a 4-tuple in $((X_i), h, C, (e_j)) \in \mathcal{T}$, we find the composition of the image $f((X_i), h, C, (e_j))$. With this, we express $\mathcal{G}(Q)$ using the catenary data of M.

By Theorem 2.1,

$$\mathcal{G}(Q) = \sum_{(b_i)} \nu(Q; (b_i)) \gamma((b_i))$$

where the sum is over all ((m + 1)n + 1, k + 1)-compositions $(b_i) = (b_0, b_1, \dots, b_{k+1})$, and $\gamma((b_i))$ is the γ -basis vector indexed by (b_i) . Let $f((X_i), h, C, (e_j)) = (Y_i)$, let (a_i) be the composition of (X_i) , and let (b_i) be that of (Y_i) . We next show that (b_i) depends only on (a_i) , h, and C; this justifies letting $\hat{f}((a_i), h, C)$ denote (b_i) . Since Q has no loops, $b_0 = 0$. For $i \in [h]$, if $i \in C$, then $Y_i - Y_{i-1}$ is a singleton subset of T, so $b_i = 1$; if i is the j-th element of D, then $Y_i - Y_{i-1} = X_j - X_{j-1}$, so $b_i = a_j$. If i > h, then $Y_i = q(X_{i-1})$, so $b_{h+1} = 1 + \sum_{j=1}^{h} (a_j(m+1) - b_j)$, and $b_i = (m+1)a_{i-1}$ for i > h+1. The triple $((a_i), h, C)$ determines another key item: there are $m \cdot a_{h-|C|+j}$ options for

The triple $((a_i), h, C)$ determines another key item: there are $m \cdot a_{h-|C|+j}$ options for e_j since $e_j \in (q(X_{h-|C|+j}) - q(X_{h-|C|+j-1})) - E$. With this, we get

$$\mathcal{G}(Q) = \sum_{(a_i)} \nu(M; (a_i)) \sum_{h=0}^k \sum_{C \subseteq [h]} \left(\prod_{j=1}^{|C|} m \cdot a_{h-|C|+j} \right) \gamma(\hat{f}((a_i), h, C)).$$

The sums account for all triples $((a_i), h, C)$. The term $\nu(M; a_i)$ accounts for all flags (X_i) of M that have composition (a_i) . When h = 0, we take $[h] = \emptyset$. The product accounts for the choices of $(e_1, e_2, \ldots, e_{|C|})$, and $\gamma(\hat{f}((a_i), h, C))$ is the γ -basis element that all flags of Q of the form $f((X_i), h, C, (e_j))$, where (X_i) and (e_j) vary over all of their respective options, contribute to $\mathcal{G}(Q)$.

The next corollary better fits our main result.

Corollary 3.11. If M and N are loopless and $\mathcal{G}(M) = \mathcal{G}(N)$, then $\mathcal{G}(Q_m(M)) = \mathcal{G}(Q_m(N))$.

4. VARIANTS OF THE FREE m-cone

As we show in this section, the proof of Theorem 3.10 adapts to give analogous results for the following deletions of the free *m*-cone: the *tipless m*-cone $Q \setminus a$, the *baseless m*-cone $Q \setminus E$, and the *tipless baseless m*-cone $Q \setminus (E \cup a)$.

We first prove a result noted by Kung [9]: these variants include an important, wellstudied construction, the Higgs lift. Recall that the *Higgs lift* of a matroid M = (E, r) is the matroid $L(M) = (E, r_L)$ where $r_L(X) = \min\{r(X) + 1, |X|\}$ for all $X \subseteq E$.

Theorem 4.1. If M has no loops, then the tipless baseless 1-cone $Q \setminus (E \cup a)$ of M is isomorphic to the Higgs lift L(M) of M.

Proof. For a 1-cone, the restriction $p: 2^T \to 2^E$ of the map p defined earlier is a bijection, and p gives a bijection from the ground set of $Q \setminus (E \cup a)$ onto that of L(M). We use p^{-1}

to relabel L(M) so that its ground set is T. With this relabeling, the rank function r_L of L(M) is given by

$$r_L(S) = \begin{cases} r(p(S)), & \text{if } p(S) \text{ is independent in } M, \\ r(p(S)) + 1, & \text{otherwise,} \end{cases}$$

for $S \subseteq T$. By Corollary 3.5, the rank of S in $Q \setminus (E \cup a)$ is given by

$$r(S) = \begin{cases} r(p(S)), & \text{if } a \notin cl(S), \\ r(p(S)) + 1, & \text{otherwise.} \end{cases}$$

When p(S) is independent, Lemma 3.4 applies to the sets S and p(S); now $p(S) \cup a$ is independent since $a \notin cl(p(S))$, so $S \cup a$ is independent, so $a \notin cl(S)$; therefore, $r_L(S) = r(p(S)) = r(S)$. When p(S) is dependent in M, either (i) p(S) contains parallel elements, and so S contains two points that are collinear with a, so $a \in cl(S)$, or (ii) Lemma 3.4 applies and gives that S is dependent, but the closure of any circuit in T contains a, so $a \in cl(S)$; thus, $r_L(S) = r(p(S)) + 1 = r(S)$. \Box

Given a matroid M, a flag (X_i) of M, and $S \subset E(M)$, we say that (X_i) collapses in $M \setminus S$ if $X_i - S = X_{i+1} - S$ for some i. If it is obvious what S is, we just say that (X_i) collapses. The next lemma relates the flags of M to those of $M \setminus S$ when at least one flag of M does not collapse.

Lemma 4.2. Let S be a nonempty subset of E(M). Not all flags of M collapse in $M \setminus S$ if and only if $r(M) = r(M \setminus S)$. In that case, the flags of $M \setminus S$ are exactly the sequences $(Y_i - S)$ where (Y_i) is a flag of M that does not collapse.

Proof. Let (Y_i) be a flag of M that does not collapse. The r(M) + 1 flats in the chain of flats $(Y_i - S)$ in $M \setminus S$ strictly increase in rank, so $r(M \setminus S) = r(M)$ and $(Y_i - S)$ is a flag of $M \setminus S$. Now assume that $r(M \setminus S) = r(M)$ and let (Z_i) be a flag of $M \setminus S$. Since $r(cl_M(Z_i)) = i$ and $r(M \setminus S) = r(M)$, it follows that $(cl_M(Z_i))$ is a flag of M and $(cl_M(Z_i) - S) = (Z_i)$.

Theorem 4.3. Fix $m \ge 1$. Let M be a rank-k loopless matroid and let $Q = Q_m(M)$. The \mathcal{G} -invariants of $Q \setminus a$, $Q \setminus E$, and $Q \setminus (E \cup a)$ can be computed from $\mathcal{G}(M)$.

Proof. A flag (Y_i) of Q collapses in $Q \setminus a$ if and only if $Y_{i+1} - Y_i = \{a\}$ for some i, which, by Corollary 3.7, occurs if and only if $Y_1 = \{a\}$. So, using the bijection f in Lemma 3.9, a flag (Y_i) of Q does not collapse if and only if $(Y_i) = f((X_i), h, C, (e_j))$ for some $((X_i), h, C, (e_j)) \in \mathcal{T}$ with $h \ge 1$. The composition (b_i) of the flag $(Y_i - a)$ of $Q \setminus a$ is the same as that of the flag (Y_i) except that $b_{h+1} = \sum_{j=1}^{h} (a_j(m+1) - b_j)$. Letting $f_1((a_i), h, C)$ denote (b_i) , we get

$$\mathcal{G}(Q \setminus a) = \sum_{(a_i)} \nu(M; (a_i)) \sum_{h=1}^k \sum_{C \subseteq [h]} \left(\prod_{j=1}^{|C|} m \cdot a_{h-|C|+j} \right) \gamma(f_1((a_i), h, C)).$$

A flag (Y_i) of Q collapses in $Q \setminus E$ if and only if $Y_{i+1} - Y_i \subseteq E$ for some i, which, by Corollary 3.7, occurs if and only if $r(Y_{i+1} \cap E) = r(Y_i \cap E) + 1$ for some i such that $a \notin Y_{i+1}$. So, a flag (Y_i) of Q does not collapse if and only if $(Y_i) = f((X_i), h, C, (e_j))$ for some $((X_i), h, C, (e_j)) \in \mathcal{T}$ with C = [h]. The composition (b_i) of the flag $(Y_i - E)$ is the same as that of (Y_i) , except that $b_{h+1} = 1 + \sum_{j=1}^{h} (ma_j - 1)$ and $b_i = ma_{i-1}$ for i > h + 1. Letting $f_2((a_i), h)$ denote (b_i) , since C = [h], we get

$$\mathcal{G}(Q \setminus E) = \sum_{(a_i)} \nu(M; (a_i)) \sum_{h=0}^k \left(\prod_{j=1}^h m \cdot a_j\right) \gamma(f_2((a_i), h)).$$

For $Q \setminus (E \cup a)$, we treat the cases m = 1 and m > 1 separately, with the latter first. A flag (Y_i) of Q collapses in $Q \setminus (E \cup a)$ if and only if $Y_{i+1} - Y_i \subseteq E \cup a$ for some i. By Corollary 3.7, the flag (Y_i) does not collapse if and only if $Y_1 \neq \{a\}$, and for all Y_{i+1} with $a \notin Y_{i+1}$, we have $Y_{i+1} - Y_i = \{x\}$ for some $x \in T$. So, a flag (Y_i) of Q does not collapse if and only if $(Y_i) = f((X_i), h, C, (e_j))$ for some $((X_i), h, C, (e_j)) \in \mathcal{T}$ with C = [h] and $h \ge 1$. The composition (b_i) of the flag $(Y_i - (E \cup a))$ of $Q \setminus (E \cup a)$ is the same as that of (Y_i) , except that $b_{h+1} = \sum_{j=1}^{h} (ma_j - 1)$ and $b_i = ma_{i-1}$ for i > h + 1. Letting $f_3((a_i), h)$ denote (b_i) , since C = [h], we get

$$\mathcal{G}(Q \setminus (E \cup a)) = \sum_{(a_i)} \nu(M; (a_i) \sum_{h=1}^k \left(\prod_{j=1}^h m \cdot a_j\right) \gamma(f_3((a_i), h))$$

Now consider m = 1, which has one more way for the inclusion $Y_{i+1} - Y_i \subseteq E \cup a$ to occur, namely, if i = h, $Y_h \subseteq T$, and $p(Y_h)$ is a (necessarily independent) flat of M, since then $Y_{h+1} = \operatorname{cl}(Y_h \cup a) = Y_h \cup p(Y_h) \cup a$. So a flag (Y_i) of Q does not collapse if and only if $(Y_i) = f((X_i), h, C, (e_j))$ for some $((X_i), h, C, (e_j)) \in \mathcal{T}$ with $C = [h], h \ge 1$, and $|X_i - X_{i-1}| > 1$ for at least one $i \le h$. When such flags exist, the composition (b_i) of the flag $(Y_i - (E \cup a))$ of $Q \setminus (E \cup a)$ is the same as when m > 1. The formula for the \mathcal{G} -invariant is also the same as when m > 1, except that for a composition (a_i) and integer $h \in [k]$ to contribute, we must have some $i \in [h]$ with $a_i > 1$.

Finally, if all flags of Q collapse, then M is the uniform matroid $U_{k,k}$; this can be detected from $\mathcal{G}(M)$, which is k![11...1]. It follows from Theorem 4.1 that $Q \setminus (E \cup a)$ is also $U_{k,k}$, so $\mathcal{G}(Q) = \mathcal{G}(M)$.

The case of $Q \setminus (E \cup a)$ with m = 1, which is isomorphic to the Higgs lift of M, was treated in [3, Proposition 4.2] using rank sequences; the expression above using the γ -basis is new.

We extend Theorem 3.8 to the variants of the free *m*-cone. We use the following observations about cyclic flats in a deletion $N \setminus X$ of a matroid N. If $F \in \mathcal{Z}(N \setminus X)$, then $cl_N(F) \in \mathcal{Z}(N)$. The flats of $N \setminus X$ are the sets F - X as F ranges over the flats of N. So the cyclic flats of $N \setminus X$ are the sets F - X, with $F \in \mathcal{Z}(N)$, that are cyclic.

Theorem 4.4. Let M be a loopless matroid and let $Q = Q_m(M)$.

- (1) If m > 1, then the configuration of $Q \setminus a$ determines M.
- (2) If m > 1, then m and the configuration of $Q \setminus E$ determine M.
- (3) If m > 2, then m and the configuration of $Q \setminus (E \cup a)$ determine M.

Proof. Each line of Q that contains a has at least four points, so, using the observations above, the cyclic flats of $Q \setminus a$ are those of M along with the sets q(F) - a as F ranges over the nonempty flats of M. Let $\mathcal{L} = \{L - a : L \text{ is a line of } Q \text{ with } a \in L\}$. From the proof of Theorem 3.8, we see that \mathcal{L} is the largest set of cyclic lines of $Q \setminus a$ that satisfies properties (L1)–(L3) in that proof (using E(Q) - a in (L2)), and we can deduce, from the configuration of $Q \setminus a$, which of its elements come from lines in \mathcal{L} . The cyclic flats of Q that contain a correspond to the elements F in the configuration of $Q \setminus a$ where $L \leq F$




FIGURE 4. The matroids above have the same Tutte polynomial, but $\mathcal{G}(N_1) \neq \mathcal{G}(N_2)$. Their free 1-cones have different Tutte polynomials.

for at least one $L \in \mathcal{L}$, so we get the configuration of Q by increasing the size assigned to such elements by 1. So, from the configuration of $Q \setminus a$, we get that of Q, from which we get M by Theorem 3.8.

The cyclic flats of $Q \setminus E$ are \emptyset and the sets q(F) - F as F ranges over the nonempty flats of M. Increase the size assigned to each element F in the configuration of $Q \setminus E$ by

$$\sum_{\substack{L \text{ in the configuration,} \\ L \leq F, \text{ and } r(L) = 2}} \frac{|L| - 1}{m}.$$

This gives the data that was used in the proof of Theorem 3.8 to get M.

Similarly, when m > 2, the cyclic flats of $Q \setminus (E \cup a)$ are precisely \emptyset and the sets $q(F) - (F \cup a)$ as F ranges over the nonempty flats of M. For each element of positive rank in the configuration of $Q \setminus (E \cup a)$, increase its assigned size by 1; this gives the configuration of $Q \setminus E$, from which we just showed that we get M.

The matroids M_1 and M_2 in Figure 1 show that the inequalities m > 1 for (1) and (2), and m > 2 for (3), are necessary.

5. SIZE-RANK-COLOOP DATA

In contrast to Theorem 3.10, the Tutte polynomial of M does not determine that of Q: the matroids N_1 and N_2 in Figure 4 have the same Tutte polynomial, but their free 1-cones do not. (The same counterexample settles this question also for the characteristic polynomial, which is $\chi(M; x) = (-1)^{r(M)}T(M; 1 - x, 0)$.) We introduce a matroid invariant that determines the Tutte polynomial of Q.

The Tutte polynomial T(M; x, y) of M is equivalent to the *size-rank data* of M, which is the multiset of pairs (|S|, r(S)) for all $S \subseteq E(M)$. The *size-rank-coloop data* of M is the multiset of triples (|S|, r(S), c(S)), for all $S \subseteq E(M)$, where c(S) is the number of coloops of M|S. We show that we can compute T(Q; x, y) using this stronger invariant.

Theorem 5.1. Let $Q = Q_m(M)$ for some $m \ge 1$. The Tutte polynomial of Q can be computed from the size-rank-coloop data of M.

Proof. We refer to the following information as the *type* of a subset S' of E(Q):

- (1) the set S = p(S'),
- (2) r(S),
- (3) whether $a \in S'$,
- (4) for each $e \in S$, (a) the size $|S' \cap T_e|$, (b) whether $e \in S'$, and (c) whether e is a coloop of M|S.

We call S the underlying set of the type. Below we show that from the type of S', we can deduce |S'| and r(S'), and so the contribution of S' to T(Q; x, y). The proof is then completed by showing that for each $S \subseteq E$ and each possible type of subsets of E(Q) with underlying set S, we can find the number of subsets of E(Q) having that type from the triple (|S|, r(S), c(S)).

Let $s = |S \cap S'| + \sum_{e \in S} |S' \cap T_e|$. If $a \notin S'$, then |S'| = s, otherwise |S'| = s + 1. Thus, we get |S'| from the type of S'.

Now r(S') = r(S) if $a \notin cl(S')$, otherwise r(S') = r(S) + 1, by Corollary 3.5, so to show that we get r(S') from the type of S', it suffices to show that the type of S' determines whether $a \in cl(S')$. We have $a \in cl(S')$ if either $a \in S'$ or $|S' \cap (T_e \cup e)| > 1$ for some $e \in S$, so now assume $a \notin S'$ and $|S' \cap (T_e \cup e)| = 1$ for all $e \in S$.

With those assumptions, we claim that $a \in cl(S')$ if and only if $|S' \cap T_e| > 0$ for some $e \in S$ that is not a coloop in M|S. First assume that $a \in cl(S')$. Now cl(S') = q(F) for some flat F of M by Lemma 3.6. Let $B \subseteq S'$ be a basis of Q|q(F). We have |p(B)| = |B| = r(F) + 1 and r(p(B)) = r(F), so p(B) contains a circuit C. Since $C \not\subseteq B$, there is an $e \in C - S'$. So $e \in S$, it is not a coloop in M|S, and $|S' \cap T_e| > 0$. Now assume that $a \notin cl(S')$. If $e \in S$ and $e' \in S' \cap T_e$, then e' is a coloop of Q|S' by Lemma 3.3. Since r(S) = r(S') and r(S - e) = r(S' - e'), it follows that e is a coloop of M|S. This completes the proof that we get r(S') from the type of S'.

We now show that for each set $S \subseteq E$ and possible type with underlying set S, we get the number of subsets of E(Q) of that type from the triple (|S|, r(S), c(S)). As just shown, r(S') = r(S) if and only if (i) $a \notin S'$, (ii) $|S' \cap (T_e \cup e)| = 1$ for all $e \in S$, and (iii) each $e \in S$ with $|S' \cap T_e| = 1$ is a coloop of M|S; otherwise r(S') = r(S) + 1. For any positive integer h, the number of sets S' with p(S') = S and |S'| = h depends just on m and |S|, so the proof is completed by observing that the number of such sets S' that satisfy conditions (i)–(iii) is determined by |S| and c(S).

The free 1-cones of N_1 and N_2 in Figure 4 have different Tutte polynomials, so N_1 and N_2 have different size-rank-coloop data. Indeed, N_1 has 20 sets of size 4, rank 3, with one coloop, while N_2 has 18.

The proof of Theorem 5.1 applies, with only minor changes in which sets S' are of concern, to prove the following extension to the variants of the free *m*-cone introduced in Section 4.

Corollary 5.2. Let $Q = Q_m(M)$ for some $m \ge 1$. The Tutte polynomial of $Q \setminus a$, of $Q \setminus E$, and of $Q \setminus (E \cup a)$ can be determined from the size-rank-coloop data of M.

We now prove that $\mathcal{G}(M)$ gives the size-rank-coloop data of M.

Theorem 5.3. Let M = (E, r) be a matroid with |E| = n. The size-rank-coloop data of M can be computed from $\mathcal{G}(M)$.

Proof. For a triple of nonnegative integers (s, t, c) with $s \ge t \ge c$, let F(s, t, c) be the set of subsets X of E with |X| = s and r(X) = t for which M|X has c coloops, and let G(s, t, c) be the set of permutations π of E for which $[\underline{r}(\pi)]$ has exactly t ones in the first s places, with each of the last c of those being one. More elements than those corresponding to the last string of ones may be coloops among the first s elements, so if $\pi \in G(s, t, c)$, then $\{\pi(i) : i \in [s]\} \in \bigcup_{c' \ge c} F(s, t, c')$. A set $X \in F(s, t, c')$ gives rise to permutations $\pi \in G(s, t, c)$ by picking c of the coloops, taking a permutation of X with those c coloops

last, and placing the elements of E - X after those c coloops. So,

$$\sum_{c' \ge c} |F(s,t,c')| \binom{c'}{c} c! (s-c)! (n-s)! = |G(s,t,c)|.$$

We get |G(s, t, c)| from $\mathcal{G}(M)$. For a fixed s and t, these equalities form an upper triangular system of equations in |F(s, t, c)|, so, as needed, we can find the terms |F(s, t, c)|. \Box

The results above show that the size-rank-coloop data is strictly stronger than the Tutte polynomial, and the \mathcal{G} -invariant is at least as strong as the size-rank-coloop data. It seems most likely that the \mathcal{G} -invariant is strictly stronger than the size-rank-coloop data, but we do not currently have pairs of matroids with the same size-rank-coloop data and different \mathcal{G} -invariants. In summary, we have the invariants below listed from weakest to strongest, where (2) is strictly stronger than (1), and (4) is strictly stronger than (3), but we do not yet know about (3) and (2):

- (1) the Tutte polynomial,
- (2) the size-rank-coloop data,
- (3) the \mathcal{G} -invariant, and
- (4) the configuration.

ACKNOWLEDGMENTS

We thank Joseph Kung for valuable observations and comments on the topics of this paper. We thank both referees for their careful reading of the manuscript, their valuable comments, and for pointing out a gap in the original proof of Theorem 3.8.

REFERENCES

- [1] J. Bonin, Matroids with different configurations and the same G-invariant (submitted).
- [2] J. Bonin and A. de Mier, The lattice of cyclic flats of a matroid, Ann. Comb. 12 (2008) 155–170.
- [3] J. Bonin and J.P.S. Kung, The *G*-invariant and catenary data of a matroid, *Adv. in Appl. Math.* 94 (2018) 39–70.
- [4] T. Brylawski, An affine representation for transversal geometries, Studies in Appl. Math. 54 (1975) 143-160.
- [5] T. Brylawski and J. Oxley, The Tutte Polynomial and Its Applications, In N. White (ed), *Matroid Applications* (Encyclopedia of Mathematics and its Applications, pp. 123–225). Cambridge: Cambridge University Press (1992).
- [6] H. Derksen, Symmetric and quasi-symmetric functions associated to polymatroids, J. Algebraic Combin. 30 (2009) 43–86.
- [7] H. Derksen and A. Fink, Valuative invariants for polymatroids, Adv. Math. 225 (2010) 1840–1892
- [8] J. Eberhardt, Computing the Tutte polynomial of a matroid from its lattice of cyclic flats, *Electron. J. Combin.* 21 (2014) Paper 3.47, 12 pp.
- [9] J.P.S. Kung, personal communications, May 2021.
- [10] J. Oxley, Matroid Theory, second edition (Oxford University Press, Oxford, 2011).
- [11] J.A. Sims, Some Problems in Matroid Theory, (Ph.D. Dissertation, Linacre College, Oxford University, Oxford, 1980).

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON, D.C. 20052, USA

Email address: jbonin@gwu.edu, kevinlong@gwmail.gwu.edu