

BLOWUP OF SOLUTIONS TO A DAMPED EULER EQUATION WITH HOMOGENEOUS THREE-POINT BOUNDARY CONDITION

IKECHUKWU OBI-OKOYE AND ALEJANDRO SARRIA

ABSTRACT. It has been established that solutions to the inviscid Proudman-Johnson equation subject to a homogeneous three-point boundary condition can develop singularities in finite time. In this paper, we consider the possibility of singularity formation in solutions of the generalized, inviscid Proudman-Johnson equation with damping subject to the same homogeneous three-point boundary condition. In particular, we derive conditions the initial data must satisfy in order for solutions to blowup in finite time with either bounded or unbounded smooth damping term.

1. INTRODUCTION

We study the initial value problem (IVP)

$$\begin{cases} u_{xxt} + uu_{xxx} + \beta u_x u_{xx} + \alpha(t)u_{xx} = 0, & x \in [0, 1], t > 0, \\ u(x, 0) = u_0(x), & x \in [0, 1], \end{cases} \quad (1.1)$$

where $\beta = \frac{n-3}{n-1}$, $n \in \mathbb{Z}^+$, $n \geq 2$, and $u(x, t)$ satisfies the homogeneous three-point boundary condition

$$u(1, t) = u_x(0, t) = u_x(1, t) = 0 \quad (1.2)$$

Equation (1.1)i can be obtained by imposing on the incompressible n -dimensional Euler equations with damping

$$\mathbf{u}_t + (\mathbf{u} \cdot \nabla)\mathbf{u} + \alpha(t)\Delta\mathbf{u} = -\nabla p, \quad \nabla \cdot \mathbf{u} = 0$$

velocities of the form

$$\mathbf{u}(x, \mathbf{x}', t) = (u(x, t), -\frac{\mathbf{x}'}{n-1}u_x(x, t))$$

for $\mathbf{x}' = \{x_2, \dots, x_n\}$, or by using the cylindrical coordinate representation

$$u^r = -\frac{r}{n-1}u_x(x, t), \quad u^\theta = 0, \quad u^x = u(x, t)$$

where $r = |\mathbf{x}'|$ ([4], [22], [20], [14], [6], [16]).

We will refer to (1.1)-(1.2) as the initial boundary value problem for the generalized, inviscid Proudman-Johnson equation with damping.

We remark that the undamped version of (1.1)i, i.e.

$$u_{xxt} + uu_{xxx} + \beta u_x u_{xx} = 0 \quad (1.3)$$

arises in several physical and geometrical contexts. For instance, when $\beta = 3$, (1.3) reduces to the Burgers' equation of gas dynamics. For $\beta = 2$, it becomes the Hunter-Saxton equation

2000 *Mathematics Subject Classification.* 35B44, 35B65, 35Q35.

Key words and phrases. Generalized Proudman-Johnson equation, blow-up, Three-point boundary condition.

(HS) describing the orientation of waves in massive nematic liquid crystals ([9], [3], [5], [24]). From a more geometric point of view, periodic solutions to the HS equation also describe geodesics on the group $\mathcal{D}(\mathbb{S}) \setminus \text{Rot}(\mathbb{S})$ of orientation preserving diffeomorphisms on the unit circle modulo rigid rotations with respect to the right-invariant metric ([10], [3], [21], [11])

$$\langle f, g \rangle = \int_{\mathbb{S}} f_x g_x dx$$

Furthermore, Lenells and Misiolek [12] showed that, for any β , (1.3) arises as the geodesic equation of the affine connection $\nabla^{(\alpha)}$ on the group $\mathcal{D}(\mathbb{S}) \setminus \text{Rot}(\mathbb{S})$. See also [2] for yet another derivation of (1.3) as a geodesic equation.

From a more heuristic point of view, (1.1)i) may serve as a tool to better understand the role that convection and stretching play in the regularity of solutions to one-dimensional fluid evolution equations; it has been argued that the convection term can sometimes cancel some of the nonlinear effects and contribute positively to regularity of solutions ([13], [8], [15]). More particularly, setting $\omega = u_{xx}$ in (1.1)i) yields

$$\omega_t + \underbrace{u\omega_x}_{\text{convection}} + \beta \underbrace{\omega u_x}_{\text{stretching}} + \alpha(t)\omega = 0 \quad (1.4)$$

In the undamped case ($\alpha(t) \equiv 0$) and $\beta = -1$, (1.4) becomes a one-dimensional analogue of the three-dimensional vorticity equation of incompressible inviscid fluids

$$\omega_t + \mathbf{u} \cdot \nabla \omega = \omega \cdot \nabla \mathbf{u}, \quad \omega = \nabla \times \mathbf{u}$$

The nonlinear terms in equation (1.4) represent the competition between nonlinear convection and stretching ([7], [23]). More particularly, the parameter $\beta \in \mathbb{R}$ is related to the ratio of stretching to convection.

2. PREVIOUS REGULARITY RESULTS

The global regularity of solutions of the damped equation (1.1)i) for a particular value of β is discussed in [19] for solutions satisfying periodic boundary conditions. In the undamped case, finite-time singularity formation has been extensively studied for periodic solutions as well as Dirichlet boundary conditions (see e.g. [17], [18] and references therein). However, we are unaware of any results concerning singularity formation in solutions of the damped equation (1.1)i) with the homogeneous three-point boundary condition (1.2). In the undamped case with homogeneous three-point boundary condition (1.2), the only result we are aware of is Theorem (2.1) below established by Yuen ([1]) where conditions on the initial data leading to finite-time blowup are derived for $\beta = -1$. Therefore, it is of interest to study whether finite-time blowup of solutions is still a possibility after incorporating damping into the system as well as determining if varying the value of the parameter β has any effect on the regularity of solutions which, in turn, could lead to a better understanding of the competing effects between nonlinear convection and stretching as discussed at the end of the previous section.

Theorem 2.1. (Yuen [1]) *Consider the C^3 solutions to the IBVP (1.1)-(1.2) for the inviscid undamped Proudman Johnson equation ($n = 2$, $\alpha(t) \equiv 0$). If the initial velocity satisfies*

$$U_0 = \int_0^1 u'_0(x) dx = -u_0(0) > 0$$

then the solutions blowup on or before the finite time $1/(2U_0)$.

The outline of the paper is as follows. In Section 3, we prove finite-time blowup of solutions to (1.1)-(1.2) with smooth and bounded time-dependent damping term, while the formation of singularities in finite time with smooth and unbounded damping term is discussed in Section 4.

3. FINITE-TIME BLOWUP WITH SMOOTH, BOUNDED DAMPING

In this section, we will derive conditions on the initial data $u_0(x)$ which show that the presence of a smooth (bounded or unbounded) time-dependent damping term $\alpha(t)$ is insufficient to arrest finite-time blowup for any value of the parameter $\beta \in \mathbb{R}$.

Theorem 3.1. *Consider the IBVP (1.1)-(1.2) for the generalized inviscid damped Proudman Johnson equation with initial data $u_0(x) \in C^\infty([0, 1])$ and smooth bounded damping term $\alpha(t) : [0, \infty) \rightarrow \mathbb{R}^+$.*

Set

$$H_0 = - \int_0^1 u_0'(x) dx = u_0(0)$$

and

$$M := \sup_{t \in [0, \infty)} \alpha(t) \tag{3.2}$$

for some $M \in \mathbb{R}^+$. If

$$H_0 < 0 \quad \text{and, more particularly,} \quad H_0 < M \left(\frac{1-n}{n} \right) \tag{3.3}$$

for fixed $n \in \mathbb{Z}^+$, $n \geq 2$, then

$$\lim_{t \nearrow t^*} u(0, t) = -\infty \tag{3.4}$$

for positive t^* given by

$$t^* = -\frac{1}{M} \ln \left(1 - \frac{M(1-n)}{nH_0} \right)$$

Proof. Multiplying (1.1)i) by x , integrating over $x \in [0, 1]$ and using (1.2) gives

$$H'(t) + \alpha(t)H(t) + \frac{n}{n-1} \|u_x(\cdot, t)\|_2^2 = 0 \tag{3.5}$$

for

$$H(t) = - \int_0^1 u_x(x, t) dx = u(0, t) \tag{3.6}$$

Now, the Cauchy-Schwarz inequality implies that

$$H^2 \leq \|u_x(\cdot, t)\|_2^2$$

and so equation (3.5) yields the inequality

$$H'(t) + \alpha(t)H(t) + \frac{n}{n-1} H^2 \leq 0 \tag{3.7}$$

Setting $f = H^{-1}$ in (3.7) and using an integrating factor argument we obtain

$$\frac{d}{dt} \left(f(t) e^{-\int_0^t \alpha(s) ds} \right) \geq \frac{n}{n-1} e^{-\int_0^t \alpha(s) ds}$$

which gives, after integrating,

$$\frac{1}{H(t)} \geq \frac{n(1 + H_0 g(t))}{(n-1)H_0 g'(t)} \quad (3.8)$$

for $H_0 = H(0)$ and

$$g(t) = \frac{n}{n-1} \int_0^t e^{-\int_0^s \alpha(z) dz} ds \quad (3.9)$$

Next, from (3.2),

$$\alpha(t) \leq M$$

for all $t \in [0, \infty)$ for some $M \in \mathbb{R}^+$. Consequently,

$$g'(t) \geq \frac{n}{n-1} e^{-Mt}$$

and, after integrating,

$$g(t) \geq \frac{n}{n-1} \left(\frac{1 - e^{-Mt}}{M} \right)$$

Using these inequalities on (3.8) we obtain

$$\frac{1}{H(t)} \geq \Lambda(t) \quad (3.10)$$

for

$$\Lambda(t) = \frac{nN(t)}{(n-1)^2 M H_0 g'(t)} \quad (3.11)$$

and

$$N(t) = M(n-1) + nH_0(1 - e^{-Mt})$$

Now, (3.3)ii) implies that

$$\lim_{t \rightarrow \infty} N(t) = M(n-1) + nH_0 < 0$$

This, along with

$$N(0) = M(n-1) > 0, \quad N'(t) = nMH_0 e^{-Mt} < 0 \quad \text{and} \quad N(t) \in C([0, \infty))$$

implies the existence of a finite time $t^* > 0$ such that

$$\lim_{t \nearrow t^*} N(t) = 0 \quad (3.12)$$

In turn, the above argument implies that $\Lambda(t) < 0$ when $t \in [0, t^*)$ and

$$\lim_{t \nearrow t^*} \Lambda(t) = 0$$

Finally, (3.7) gives

$$H'(t) + \alpha(t)H(t) \leq 0$$

which yields

$$H(t) \leq H_0 e^{-\int_0^t \alpha(s) ds} \quad (3.13)$$

Since $H_0 < 0$, (3.13) implies that $H(t) < 0$ for as long as it exists. Thus

$$0 \geq \frac{1}{H(t)} \geq \Lambda(t) \quad (3.14)$$

and the Theorem follows by letting t approach t^* in (3.14). □

4. FINITE-TIME BLOWUP WITH SMOOTH, UNBOUNDED DAMPING

In Theorem 3.1 of the previous section, finite-time blowup was established for smooth and bounded time-dependent damping. In Theorem 4.1 below we show that finite-time blowup is still possible with smooth but unbounded time-dependent damping.

Theorem 4.1. *Consider the IBVP (1.1)-(1.2) for the generalized inviscid damped Proudman Johnson equation with initial data $u_0(x) \in C^\infty([0, 1])$. Let $E_1(x)$ be the exponential integral*

$$E_1(x) = \int_x^\infty \frac{e^{-t}}{t} dt, \quad x > 0 \quad (4.2)$$

If

$$u_0(0) < -\frac{c(n-1)}{ne^{1/c}E_1(1/c)} \quad (4.3)$$

for $c = \alpha'(0) \in \mathbb{R}^+$ and $n \in \mathbb{Z}^+$, $n \geq 2$, then there exists smooth, unbounded damping $\alpha(t) : [0, \infty) \rightarrow \mathbb{R}^+$ and a finite time $t^* > 0$ such that

$$\lim_{t \nearrow t^*} u(0, t) = -\infty \quad (4.4)$$

Proof. Suppose

$$H_0 = -\int_0^1 u_0'(x) dx = u_0(0) < 0$$

From (3.8) and (3.13) we have that

$$0 > \frac{1}{H(t)} \geq \frac{n(1 + H_0 g(t))}{(n-1)H_0 g'(t)} \quad (4.5)$$

for $g(t)$ as in (3.9) and $n \in \mathbb{Z}^+$, $n \geq 2$. Set

$$\alpha(t) = e^{ct}$$

for some $c \in \mathbb{R}^+$. Then, after some straightforward computations, we can write (4.5) as

$$0 > \frac{1}{H(t)} \geq \frac{1 + H_0 \phi(n) \eta(t)}{e^{\frac{1}{c}} H_0 e^{-\frac{ect}{c}}} \quad (4.6)$$

for

$$\phi(n) = \frac{ne^{\frac{1}{c}}}{c(n-1)}$$

and

$$\eta(t) = \int_{\frac{1}{c}}^{\frac{ect}{c}} \frac{e^{-u}}{u} du$$

Note that $\eta(0) = 0$, while $\eta'(t) > 0$ and $\eta''(t) < 0$ for all $t \in [0, \infty)$. Moreover, for the exponential integral $E_1(x)$ as defined in (4.2), we have that

$$\lim_{t \rightarrow \infty} \eta(t) = E_1(1/c)$$

The above remarks, along with the fact that the denominator of the right hand-side of (4.6) is negative for all $t \in [0, \infty)$, imply that if $H_0 = u_0(0)$ satisfies (4.3), then there exists a finite time $t^* > 0$ such that (4.4) follows. □

Remark 4.7. *Below are some sample upper bounds (4.3) (computed using Mathematica) for the initial data H_0 in Theorem 4.1 for certain values of n and c .*

- If $n = 2$ and $c = 1$, then

$$H_0 < -\frac{1}{2eE_1(1)} \approx -0.84$$

- If $n = 2$ and $c = 0.01$, then

$$H_0 < -\frac{0.01}{2e^{100}E_1(100)} \approx -0.51$$

5. CONCLUSIONS AND OPEN QUESTIONS

In this paper, we derived conditions on the initial data which guarantee the formation of singularities in finite time in solutions of the generalized inviscid Proudman-Johnson when smooth and bounded damping is incorporated into the system. Similarly, we have shown that finite-time blowup is also possible if the damping term is unbounded. It is unknown if solutions will persist for all time (instead of blowing up in finite time) if the initial data satisfies the opposite of inequality (3.3)ii) in Theorem 3.1, namely, if

$$H_0 \geq M \left(\frac{1-n}{n} \right)$$

It is also of interest to investigate how solutions to other fluid-related models such as the Boussinesq equations or the MHD equations behave under a homogeneous three-point boundary condition.

REFERENCES

- [1] M. Yuen, “*Blowup for the Inviscid Proudman-Johnson Equation with the Homogeneous Three-point Boundary Condition*”, Applied Mathematical Sciences (52) **7** (2013), 2591-2597.
- [2] M. Bauer, Y. Lu, C. Maor, A geometric view on the generalized Proudman-Johnson and r -Hunter-Saxton equations, arXiv:2101.03601.
- [3] A. Bressan and A. Constantin, Global solutions of the Hunter-Saxton equation, SIAM J. Math. Anal. **37**(3) (2005), 996-1026.
- [4] S. Childress, G.R. Ierley, E.A. Spiegel and W.R. Young, Blow-up of unsteady two-dimensional Euler and Navier-Stokes equations having stagnation-point form, J. Fluid Mech. **203** (1989), 1-22.
- [5] C. M. Dafermos, Generalized characteristics and the Hunter-Saxton equation, J. Hyperbol. Differ. Eq., Vol. 8, No. 1 (2011), 159-168.
- [6] J. Escher and M. Wunsch, Restrictions on the geometry of the periodic vorticity equation, pre-print, arXiv:1009.1029, 2010.
- [7] D.D. Holm and M.F. Staley, Wave structure and nonlinear balances in a family of evolutionary PDEs, SIAM J. Appl. Dyn. Syst. **2** (2003), 323-380.
- [8] T. Y. Hou and C. Li, Dynamic stability of the three-dimensional axisymmetric Navier-Stokes equations with swirl. Comm. Pure Appl. Math. **LXI** (2008) 661-697
- [9] J.K. Hunter and R. Saxton, Dynamics of director fields, SIAM J. Appl. Math. **51**(6) (1991), 1498-1521.
- [10] B. Khesin, G. Misiolek, Euler equations on homogeneous spaces and Virasoro orbits, Adv. Math. **176** (2003) no. 1, 116-144.

- [11] J. Lenells, Weak geodesic flow and global solutions of the Hunter-Saxton equation, *Discrete Contin. Dyn. Syst.* 18 (4) (2007), 643-656.
- [12] J. Lenells and G. Misiolek, Amari-Chentsov connections and their geodesics on homogeneous spaces of diffeomorphism groups. *J. Math. Sci.* (2014), 196, 144-151.
- [13] H. Okamoto and K. Ohkitani, On the role of the convection term in the equations of motion of incompressible fluid, *J. Phys. Soc. Japan* 74 (2005), 2737-2742.
- [14] H. Okamoto and J. Zhu, Some similarity solutions of the Navier-Stokes equations and related topics, *Taiwanese J. Math.* 4 (2000), 65-103.
- [15] H. Okamoto, T. Sakajo, M. Wunsch, On a generalization of the Constantin-Lax-Majda equation, *Nonlinearity* 21, (2008), 2447-2461.
- [16] I. Proudman and K. Johnson, Boundary-layer growth near a rear stagnation point, *J. Fluid Mech.* 12 (1962), 161-168.
- [17] A. Sarria and R. Saxton, Blow-up of solutions to the generalized, inviscid Proudman-Johnson equation, *SIAM J. Math. Anal.* 4 (2008), 1499-1515.
- [18] A. Sarria and R. Saxton, The role of initial curvature in solutions to the generalized inviscid Proudman-Johnson equation, *Q Appl Math* 73 (1) (2015), 55-91.
- [19] W. Chen and A. Sarria, Damped infinite energy solutions of the 3D Euler and Boussinesq equations, *J Differ Equations* 265 (2018), 3841-3857.
- [20] R. Saxton and F. Tiglay, Global existence of some infinite energy solutions for a perfect incompressible fluid, *SIAM J. Math. Anal.* 4 (2008), 1499-1515.
- [21] F. Tiglay, The periodic Cauchy problem of the modified Hunter-Saxton equation, *J. Evol. Eq.* 5 (4) (2005), 509-527.
- [22] H. Weyl, On the differential equations of the simplest boundary-layer problems, *Ann. Math.* 43 (1942), 381-407.
- [23] M. Wunsch, The generalized Hunter-Saxton system. *SIAM J. Math. Anal.* 42 (3) (2010) 1286-1304.
- [24] Z. Yin, On the Structure of Solutions to the Periodic Hunter-Saxton Equation, *SIAM J. Math. Anal.*, Vol. 36, No. 1, (2004), 272-283.

IKECHUKWU OBI-OKOYE
DEPARTMENT OF PHYSICS
UNIVERSITY OF NORTH GEORGIA
82 COLLEGE CIRCLE
DAHLONEGA, GA 30597
Email address: ivobio3570@ung.edu

ALEJANDRO SARRIA
DEPARTMENT OF MATHEMATICS AND STATISTICS
UNIVERSITY OF NORTH GEORGIA
82 COLLEGE CIRCLE
DAHLONEGA, GA 30597
Email address: alejandro.sarria@ung.edu