

Perfectoid overconvergent Siegel modular forms and the overconvergent Eichler–Shimura morphism

HANSHENG DIAO, GIOVANNI ROSSO AND JU-FENG WU

Abstract. The aim of this paper is twofold. We first present a construction of overconvergent automorphic sheaves for Siegel modular forms by generalising the perfectoid method, originally introduced by Chojecki–Hansen–Johansson for automorphic forms on compact Shimura curves over \mathbf{Q} . These sheaves are then verified to be isomorphic to the ones introduced by Andreatta–Iovita–Pilloni. Secondly, we establish an overconvergent Eichler–Shimura morphism for Siegel modular forms, generalising the result of Andreatta–Iovita–Stevens for elliptic modular forms.

Contents

1	Introduction	2
2	Siegel modular varieties and the Hodge–Tate period map	5
2.1	Algebraic and p -adic groups	5
2.2	Siegel modular varieties	8
2.3	The flag variety	12
2.4	Vector bundles on the flag variety	14
2.5	The Hodge–Tate period map and the w -ordinary locus	15
3	Constructions of the overconvergent automorphic sheaf	18
3.1	The perfectoid construction	18
3.2	Hecke operators	24
3.3	Admissibility	29
3.4	Classical Siegel modular forms	36
3.5	The construction à la Andreatta–Iovita–Pilloni	38
3.6	The w -ordinary loci and the (pseudo-)canonical subgroups	41
3.7	Comparison of the two constructions	47
4	Overconvergent cohomology groups	51
4.1	Analytic functions and analytic distributions	51
4.2	Overconvergent cohomology groups	53
5	The overconvergent Eichler–Shimura morphism	55
5.1	The Kummer étale and the pro-Kummer étale cohomology groups	55
5.2	The overconvergent Eichler–Shimura morphism	57
5.3	The image of the overconvergent Eichler–Shimura morphism at classical weights	62
6	The overconvergent Eichler–Shimura morphism on the cuspidal eigenvariety	67
6.1	Some preliminaries on overconvergent cohomology groups	68
6.2	The cuspidal eigenvariety	69
6.3	Sheaves on the cuspidal eigenvariety	71

A Kummer étale and pro-Kummer étale sites of log adic spaces	73
A.1 Review of log adic spaces	73
A.2 Calculation of $R^i \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$	78
A.3 Banach sheaves and a (generalised) projection formula	82
B Toroidal compactifications of the Siegel modular varieties	92
B.1 Toroidal compactifications and boundary strata	93
B.2 The perfectoid Siegel modular variety	97
References	99

1 Introduction

A classical result, due to M. Eichler and G. Shimura, states that the first cohomology of the complex modular curve with coefficients in $\text{Sym}^{k-2} \mathbf{Q}^2$, after scalar extension to \mathbf{C} , admits a Hecke-equivariant decomposition as the direct sum of the space of weight k holomorphic modular forms and the space of weight k anti-holomorphic cusp forms.

G. Faltings establishes a p -adic analogue of the Eichler–Shimura decomposition. Let p be a prime number and let \mathbf{C}_p be the completion of a fixed algebraic closure $\overline{\mathbf{Q}_p}$ of \mathbf{Q}_p . Suppose X is the modular curve (of some tame level N) over \mathbf{Q}_p and let \overline{X} be its compactification. Let $\pi : E \rightarrow \overline{X}$ be the universal generalised elliptic curve and let $\underline{\omega} := \pi_* \Omega_{E/\overline{X}}^1$. In [Fal87], Faltings constructs a Hecke- and $\text{Gal}(\overline{\mathbf{Q}_p}/\mathbf{Q}_p)$ -equivariant decomposition

$$H_{\text{ét}}^1(X_{\overline{\mathbf{Q}_p}}, \text{Sym}^k \mathbf{Q}_p^2) \otimes_{\mathbf{Q}_p} \mathbf{C}_p(1) \simeq (H^0(\overline{X}, \underline{\omega}^{k+2}) \otimes_{\mathbf{Q}_p} \mathbf{C}_p) \oplus (H^1(\overline{X}, \underline{\omega}^{-k}) \otimes_{\mathbf{Q}_p} \mathbf{C}_p(k+1))$$

where the Galois actions on the coherent cohomology groups are trivial.

This type of results have been generalised to families of overconvergent modular forms by F. Andreatta, A. Iovita and G. Stevens in [AIS15] and later extended to the case of compact Shimura curves over \mathbf{Q} by P. Chojecki, D. Hansen and C. Johansson in [CHJ17]. The novelty of the second work consists of a “perfectoid construction” of overconvergent families of automorphic forms as well as a succinct calculation of overconvergent cohomology groups on the pro-étale site. The same method has also been adapted to the case of elliptic and Hilbert modular forms in [BHW19].

There are two main goals we would like to achieve in this paper. We first present a construction of automorphic sheaves for overconvergent Siegel modular forms (of genus g) using the perfectoid method. Then we follow the spirit of Chojecki–Hansen–Johansson to establish an overconvergent Eichler–Shimura morphism for Siegel modular forms.

One key ingredient for such a perfectoid construction is the (toroidally compactified) perfectoid Siegel modular variety $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ studied in [PS16]. This perfectoid space comes equipped with the *Hodge–Tate period map* $\pi_{\text{HT}} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \mathcal{F}\ell$ where the adic flag variety $\mathcal{F}\ell$ parameterises maximal lagrangian subspaces of a fixed symplectic space of rank $2g$. We also consider the Siegel modular variety $\overline{\mathcal{X}}_{\text{Iw}^+}$ for the finite *strict Iwahori level* (see Definition 2.2.1 for details), viewed as an adic space. Roughly speaking, the strict Iwahori level serves as a deeper level compared with the usual Iwahori level. There is a natural projection $\overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\text{Iw}^+}$ from the Siegel modular variety at infinite level to the one at finite level.

In order to investigate the overconvergent Siegel modular forms, we restrict our attentions to certain open subspaces $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ and $\overline{\mathcal{X}}_{\text{Iw}^+, w}$ of $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ and $\overline{\mathcal{X}}_{\text{Iw}^+}$, respectively. They are referred as the *w-ordinary loci* (see Definition 2.5.3) and they contain the usual ordinary locus.

Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a weight (see Definition 3.1.1) and let $r_{\mathcal{U}}$ be the integer as defined in Definition 3.1.8. For any $w \in \mathbf{Q}_{>0}$ with $w > 1 + r_{\mathcal{U}}$, we construct a sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ over $\overline{\mathcal{X}}_{\text{Iw}^+, w}$ whose global sections are exactly the

w -overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$. Roughly speaking, sections of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ consist of functions f on $\overline{\mathcal{X}}_{\Gamma(p^\infty),w}$ which

- take value in a certain weight- $\kappa_{\mathcal{U}}$ analytic representation $C_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}})$ of the Iwahori subgroup of $\text{GL}_g(\mathbf{Z}_p)$, and
- satisfy the following formula regarding the natural action of the strict Iwahori subgroup $\text{Iw}_{\text{GSp}_{2g}}^+$ of $\text{GSp}_{2g}(\mathbf{Z}_p)$:

$$\gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f \quad \text{for any} \quad \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \text{Iw}_{\text{GSp}_{2g}}^+,$$

where \mathfrak{z} stands for the pullback of the coordinate function on the flag variety and $\rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)$ stands for a certain automorphism on $C_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}})$.

When $p > 2g$, this sheaf coincides with (the pullback to the strict Iwahori level of) the automorphic sheaf constructed in [AIP15]. See §3 for a complete story.

The next step is to look at the overconvergent cohomology groups. Suppose the weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ is *small* in the sense of Definition 3.1.1. A typical example of a small weight is a *wide-open* disc whose ring of functions is isomorphic to $\mathbf{Z}_p[[T_1, \dots, T_g]]$. For such a weight, we consider a space $D_{\kappa}^r(\mathbf{T}_0, R_{\mathcal{U}})$ of analytic distributions on certain p -adic manifold \mathbf{T}_0 . (See §4.1 for the precise definitions.) This space of distributions is equipped with a natural action of the strict Iwahori subgroup and hence induces a sheaf $\mathcal{D}_{\kappa_{\mathcal{U}}}^r$ on the Siegel modular variety $\overline{\mathcal{X}}_{\text{Iw}^+}$. The cohomology groups of $\mathcal{D}_{\kappa_{\mathcal{U}}}^r$ are precisely the *overconvergent modular symbols* studied by A. Ash and G. Stevens, among others (see for example [AS08; Han17; JN19]).

Our goal is to explicitly construct a morphism that relates the automorphic sheaves with the overconvergent cohomology groups. To this end, we will compute the objects on both sides on the *pro-Kummer étale site* $\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}$. On one hand, the overconvergent cohomology groups can be calculated in terms of the cohomology groups of some sheaf of distributions $\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r$ over the pro-Kummer étale site. On the other hand, let $\widehat{\omega}_w^{\kappa_{\mathcal{U}}}$ be the “completed” pull-back of the automorphic sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ to this site. The key step is then Proposition 5.2.5 where we construct a morphism of sheaves

$$\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r \rightarrow \widehat{\omega}_w^{\kappa_{\mathcal{U}}}.$$

This morphism can be thought of as an analytic analogue of the “projection onto the highest weight vector”. We remark that the construction involves taking transposes of matrices while the usual Iwahori subgroup is not closed under taking transposes. This is why we have to work with a smaller subgroup—the strict Iwahori subgroup—as a compromise.

Consequently, we obtain the *overconvergent Eichler-Shimura morphism*. It is a natural Hecke- and Galois-equivariant morphism from the (family of) overconvergent cohomology groups to the (family of) overconvergent Siegel modular forms

$$\text{ES}_{\kappa_{\mathcal{U}}} : H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r) \rightarrow H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}+g+1})(-n_0),$$

where $n_0 = \dim_{\mathbf{C}_p} \overline{\mathcal{X}}_{\text{Iw}^+}$ and “ $(-n_0)$ ” stands for the Tate twist.

We list some properties of this map. First of all, we show that it is compatible with base change on the weights. Secondly, we are able to control its image when specialising to a dominant algebraic weight $k \in \mathbf{Z}_{\geq 0}^g$. In particular, we show in Theorem 5.3.3 that the image of ES_k is contained in the space of classical algebraic Siegel modular forms. The proof uses the fact that when the weight is a dominant algebraic weight, the “highest weight vector” is an algebraic function. Lastly, we conclude the paper by showing that $\text{ES}_{\kappa_{\mathcal{U}}}$ can be glued to a morphism of sheaves on a suitable cuspidal eigenvariety.

There are many things we don't do: we don't control the cokernel of the map $\mathrm{ES}_{\kappa_{\mathcal{U}}}$ in general and we do not investigate the kernel of this map at all. A natural expectation is that one could construct a full filtration of $H_{\mathrm{prok\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{O}_{\kappa_{\mathcal{U}}}^r)$ using higher Coleman theory, recently developed by G. Boxer and V. Pilloni [BP20], in terms of information on suitable strata of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$. When $g = 1$, such a result is recently announced by J. E. Rodríguez Camargo ([RC21]).

We also remark that while preparing this paper, Andreatta and Iovita have announced the article [AI20], in which they upgrade their previous work [AIS15] to an “overconvergent de Rham Eichler–Shimura morphism”, meaning that their Eichler–Shimura map has as source the overconvergent cohomology groups tensored with \mathbb{B}_{dR} . They have also announced upcoming work concerning this type of de Rham Eichler–Shimura morphisms for overconvergent Siegel modular forms of genus 2. This suggests that the results in this paper can be upgraded to investigate finer p -adic Hodge theoretic properties of overconvergent cohomology groups; for example, the construction of de Rham (or even crystalline) periods in p -adic families. We shall leave this to future studies.

The paper is organised as follows. In §2, we define the main geometric objects of interest, including the Siegel modular varieties for various level structures, the perfectoid Siegel modular variety at infinite level, the flag variety, and the Hodge–Tate period map. The next section, §3, contributes to the construction of the overconvergent automorphic sheaves. In particular, when $p > 2g$, we show that our construction coincides with the (pullback to the strict Iwahori level of the) automorphic sheaves of Andreatta–Iovita–Pilloni. We warn the reader that §3.3–3.7 is the most technical part of the paper. These subsections can be skipped on a first reading. In §4, we study the space of analytic distributions and the overconvergent cohomology groups. Finally, in §5 and §6, we construct the Eichler–Shimura morphism as well as its upgraded version on the cuspidal eigenvariety.

In §A, we recall the basics of logarithmic adic spaces and their Kummer pro-étale topology following [Dia+19a]. Also in the same section, we include some technical calculations of the derived functor $R^i\nu_*$ and a generalised projection formula, both of which are used in the main text. In §B, we recall the basics of the toroidal compactifications of Siegel modular varieties. We also review the “modified integral structures” studied in [PS16] as well as the construction of the perfectoid Siegel modular variety at the infinite level.

Acknowledgement. The authors would like to express their gratitude to Przemysław Chojecki, David Hansen, and Christian Johansson for discussions on the perfectoid construction of sheaves of families of Siegel modular forms and for pointing out an inaccuracy in the first version of this paper. G.R. would like to thank Daniel Barrera-Salazar and Riccardo Brasca. J.-F. W. would like to thank Ulrich Görtz for helpful conversations about toroidal compactifications of Siegel modular varieties and p -divisible groups. He would also like to thank Marc Levine for a succinct introduction on 1-motives during a casual conversation.

Conventions and notations. Through out this paper, we fix the following:

- $g \in \mathbb{Z}_{\geq 1}$.
- $p \in \mathbb{Z}_{>0}$ is an odd prime number. Due to certain technicality, we will have to assume $p > 2g$ at some places. Such an assumption shall be clear in the context.
- $N \in \mathbb{Z}_{\geq 3}$ is an integer coprime to p .
- We fix once and forever an algebraic closure $\overline{\mathbb{Q}}_p$ of \mathbb{Q}_p and an algebraic isomorphism $\mathbb{C}_p \simeq \mathbb{C}$, where \mathbb{C}_p is the p -adic completion of $\overline{\mathbb{Q}}_p$. We write $G_{\mathbb{Q}_p}$ for the absolute Galois group $\mathrm{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$. We also fix the p -adic absolute value on \mathbb{C}_p so that $|p| = p^{-1}$.
- For any $w \in \mathbb{Q}_{>0}$, we denote by “ p^w ” an element in \mathbb{C}_p with absolute value p^{-w} . All constructions in the paper will not depend on such choices.

- We adopt the language of almost mathematics. In particular, for an $\mathcal{O}_{\mathbf{C}_p}$ -module M , we denote by M^a for the associated almost $\mathcal{O}_{\mathbf{C}_p}$ -module.
- For $n \in \mathbf{Z}_{\geq 1}$ and any set R , we denote by $M_n(R)$ the set of n by n matrices with coefficients in R .
- The transpose of a matrix α is denoted by ${}^t\alpha$.
- For any $n \in \mathbf{Z}_{\geq 1}$, we denote by $\mathbb{1}_n$ the $n \times n$ identity matrix and denote by $\check{\mathbb{1}}_n$ the $n \times n$ anti-diagonal matrix whose non-zero entries are 1; *i.e.*,

$$\mathbb{1}_n = \begin{pmatrix} 1 & & \\ & \ddots & \\ & & 1 \end{pmatrix} \quad \text{and} \quad \check{\mathbb{1}}_n = \begin{pmatrix} & & 1 \\ & \ddots & \\ 1 & & \end{pmatrix}$$

- We use \cong to denote canonical isomorphisms and \simeq to denote non-canonical ones.
- In principle (except for §A), symbols in Gothic font (*e.g.* $\mathfrak{X}, \mathfrak{Y}, \mathfrak{Z}$) stand for formal schemes; symbols in calligraphic font (*e.g.* $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$) stand for adic spaces; and symbols in script font (*e.g.* $\mathcal{O}, \mathcal{F}, \mathcal{E}$) stand for sheaves (over various geometric objects).

2 Siegel modular varieties and the Hodge–Tate period map

In this section, we introduce the Siegel modular varieties for various level structures, viewed as adic spaces, as well as their toroidal compactifications. We also recall the perfectoid Siegel modular variety introduced in [PS16] together with the Hodge–Tate period map. The notion of perfectoid Siegel modular variety has its root in [Sch15]. However, we point out that the author of *loc. cit.* only considers minimal compactifications while it is important for us to work with the toroidal compactifications.

2.1 Algebraic and p -adic groups

We start with a list of algebraic and p -adic groups that will appear repeatedly throughout the paper.

Let $V := \mathbf{Z}^{2g}$ equipped with an alternating pairing

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbf{Z}, \quad (\vec{v}, \vec{v}') \mapsto {}^t\vec{v} \begin{pmatrix} & -\check{\mathbb{1}}_g \\ \check{\mathbb{1}}_g & \end{pmatrix} \vec{v}',$$

where we view elements in V as column vectors. In particular, if e_1, \dots, e_{2g} is the standard basis for V , then

$$\langle e_i, e_j \rangle = \begin{cases} -1, & \text{if } i < j \text{ and } j = 2g + 1 - i \\ 1, & \text{if } i > j \text{ and } j = 2g + 1 - i \\ 0, & \text{else} \end{cases}.$$

We define the algebraic group GSp_{2g} to be the subgroup of GL_{2g} that preserves this pairing up to a unit. In other words, for any ring R ,

$$\mathrm{GSp}_{2g}(R) := \left\{ \gamma \in \mathrm{GL}_{2g}(R) : {}^t\gamma \begin{pmatrix} & -\check{\mathbb{1}}_g \\ \check{\mathbb{1}}_g & \end{pmatrix} \gamma = \varsigma(\gamma) \begin{pmatrix} & -\check{\mathbb{1}}_g \\ \check{\mathbb{1}}_g & \end{pmatrix} \text{ for some } \varsigma(\gamma) \in R^\times \right\}.$$

Equivalently, for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{GL}_{2g}$, $\gamma \in \mathrm{GSp}_{2g}$ if and only if

$${}^t\gamma_a \check{\mathbb{1}}_g \gamma_c = {}^t\gamma_c \check{\mathbb{1}}_g \gamma_a, \quad {}^t\gamma_b \check{\mathbb{1}}_g \gamma_d = {}^t\gamma_d \check{\mathbb{1}}_g \gamma_b, \quad \text{and} \quad {}^t\gamma_a \check{\mathbb{1}}_g \gamma_d - {}^t\gamma_c \check{\mathbb{1}}_g \gamma_b = \varsigma(\gamma) \check{\mathbb{1}}_g$$

for some $\varsigma(\gamma) \in \mathbb{G}_m$.

Taking base change to \mathbf{Z}_p , we consider $V_p := V \otimes_{\mathbf{Z}} \mathbf{Z}_p$, equipped with the induced alternating pairing

$$\langle \cdot, \cdot \rangle : V_p \times V_p \rightarrow \mathbf{Z}_p, \quad (\tilde{v}, \tilde{v}') \mapsto \mathfrak{t}\tilde{v} \begin{pmatrix} & -\check{\mathbb{I}}_g \\ \check{\mathbb{I}}_g & \end{pmatrix} \tilde{v}'.$$

Let e_1, \dots, e_{2g} be the standard basis for V_p and let $\text{Fil}_{\bullet}^{\text{std}}$ denote the standard increasing filtration on V_p defined by $\text{Fil}_0^{\text{std}} = 0$ and

$$\text{Fil}_i^{\text{std}} = \langle e_1, \dots, e_i \rangle,$$

for $i = 1, \dots, 2g$.

The algebraic and p -adic subgroups of GL_g and GSp_{2g} considered in the present paper are the following:

- For every $m \in \mathbf{Z}_{\geq 1}$, we write

$$\Gamma(p^m) := \ker \left(\text{GSp}_{2g}(\mathbf{Z}_p) \xrightarrow{\text{mod } p^m} \text{GSp}_{2g}(\mathbf{Z}/p^m \mathbf{Z}) \right).$$

- The Borel subgroups are

$$\begin{aligned} B_{\text{GL}_g} &:= \text{the Borel subgroup of upper triangular matrices in } \text{GL}_g, \\ B_{\text{GSp}_{2g}} &:= \text{the Borel subgroup of upper triangular matrices in } \text{GSp}_{2g}. \end{aligned}$$

- The corresponding unipotent radicals are

$$\begin{aligned} U_{\text{GL}_g} &:= \text{the upper triangular } g \times g \text{ matrices whose diagonal entries are all 1,} \\ U_{\text{GSp}_{2g}} &:= \text{the upper triangular } 2g \times 2g \text{ matrices in } \text{GSp}_{2g} \text{ whose diagonal entries are all 1.} \end{aligned}$$

- The corresponding maximal tori for GL_g and GSp_{2g} are the maximal tori of diagonal matrices, denoted by T_{GL_g} and $T_{\text{GSp}_{2g}}$, respectively. The Levi decomposition then yields

$$B_{\text{GL}_g} = T_{\text{GL}_g} U_{\text{GL}_g} \quad \text{and} \quad B_{\text{GSp}_{2g}} = T_{\text{GSp}_{2g}} U_{\text{GSp}_{2g}}.$$

- Let $B_{\text{GL}_g}^{\text{opp}}$ and $B_{\text{GSp}_{2g}}^{\text{opp}}$ be the opposite Borel subgroups of B_{GL_g} and $B_{\text{GSp}_{2g}}$, respectively. They consist of lower triangular matrices of the corresponding algebraic groups. Similarly, $U_{\text{GL}_g}^{\text{opp}}$ and $U_{\text{GSp}_{2g}}^{\text{opp}}$ stand for the opposite unipotent radicals of U_{GL_g} and $U_{\text{GSp}_{2g}}$, respectively.
- To simplify the notations, we write

$$\begin{aligned} T_{\text{GL}_g,0} &= T_{\text{GL}_g}(\mathbf{Z}_p), \quad U_{\text{GL}_g,0} = U_{\text{GL}_g}(\mathbf{Z}_p), \quad B_{\text{GL}_g,0} = B_{\text{GL}_g}(\mathbf{Z}_p), \\ T_{\text{GSp}_{2g},0} &= T_{\text{GSp}_{2g}}(\mathbf{Z}_p), \quad U_{\text{GSp}_{2g},0} = U_{\text{GSp}_{2g}}(\mathbf{Z}_p), \quad B_{\text{GSp}_{2g},0} = B_{\text{GSp}_{2g}}(\mathbf{Z}_p). \end{aligned}$$

The subgroups $B_{\text{GL}_g,0}^{\text{opp}}$, $B_{\text{GSp}_{2g},0}^{\text{opp}}$, $U_{\text{GL}_g,0}^{\text{opp}}$, and $U_{\text{GSp}_{2g},0}^{\text{opp}}$ are defined similarly.

For every $s \in \mathbf{Z}_{\geq 1}$, define

$$\begin{aligned} T_{\text{GL}_g,s} &:= \ker(T_{\text{GL}_g}(\mathbf{Z}_p) \rightarrow T_{\text{GL}_g}(\mathbf{Z}/p^s \mathbf{Z})), & T_{\text{GSp}_{2g},s} &:= \ker(T_{\text{GSp}_{2g}}(\mathbf{Z}_p) \rightarrow T_{\text{GSp}_{2g}}(\mathbf{Z}/p^s \mathbf{Z})), \\ U_{\text{GL}_g,s} &:= \ker(U_{\text{GL}_g}(\mathbf{Z}_p) \rightarrow U_{\text{GL}_g}(\mathbf{Z}/p^s \mathbf{Z})), & U_{\text{GSp}_{2g},s} &:= \ker(U_{\text{GSp}_{2g}}(\mathbf{Z}_p) \rightarrow U_{\text{GSp}_{2g}}(\mathbf{Z}/p^s \mathbf{Z})), \\ B_{\text{GL}_g,s} &:= \ker(B_{\text{GL}_g}(\mathbf{Z}_p) \rightarrow B_{\text{GL}_g}(\mathbf{Z}/p^s \mathbf{Z})), & B_{\text{GSp}_{2g},s} &:= \ker(B_{\text{GSp}_{2g}}(\mathbf{Z}_p) \rightarrow B_{\text{GSp}_{2g}}(\mathbf{Z}/p^s \mathbf{Z})), \end{aligned}$$

where all of the maps are reduction modulo p^s .

The subgroups $B_{\text{GL}_g,s}^{\text{opp}}$, $B_{\text{GSp}_{2g},s}^{\text{opp}}$, $U_{\text{GL}_g,s}^{\text{opp}}$, and $U_{\text{GSp}_{2g},s}^{\text{opp}}$ are defined similarly.

- The Iwahori subgroups of $\mathrm{GL}_g(\mathbf{Z}_p)$ and $\mathrm{GSp}_{2g}(\mathbf{Z}_p)$ are

$$\begin{aligned} \mathrm{Iw}_{\mathrm{GL}_g} &:= \text{the preimage of } B_{\mathrm{GL}_g}(\mathbf{F}_p) \text{ under the reduction map } \mathrm{GL}_g(\mathbf{Z}_p) \rightarrow \mathrm{GL}_g(\mathbf{F}_p), \\ \mathrm{Iw}_{\mathrm{GSp}_{2g}} &:= \text{the preimage of } B_{\mathrm{GSp}_{2g}}(\mathbf{F}_p) \text{ under the reduction map } \mathrm{GSp}_{2g}(\mathbf{Z}_p) \rightarrow \mathrm{GSp}_{2g}(\mathbf{F}_p). \end{aligned}$$

The Iwahori decomposition yields

$$\mathrm{Iw}_{\mathrm{GL}_g} = U_{\mathrm{GL}_g,1}^{\mathrm{opp}} T_{\mathrm{GL}_g,0} U_{\mathrm{GL}_g,0} \quad \text{and} \quad \mathrm{Iw}_{\mathrm{GSp}_{2g}} = U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}} T_{\mathrm{GSp}_{2g},0} U_{\mathrm{GSp}_{2g},0}.$$

- We consider the *strict Iwahori subgroups* of $\mathrm{GL}_g(\mathbf{Z}_p)$ and $\mathrm{GSp}_{2g}(\mathbf{Z}_p)$ defined as

$$\begin{aligned} \mathrm{Iw}_{\mathrm{GL}_g}^+ &:= \text{the preimage of } T_{\mathrm{GL}_g}(\mathbf{F}_p) \text{ under the reduction map } \mathrm{GL}_g(\mathbf{Z}_p) \rightarrow \mathrm{GL}_g(\mathbf{F}_p) \\ \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ &:= \left\{ \gamma \in \mathrm{GSp}_{2g}(\mathbf{Z}_p) : \gamma \equiv \begin{pmatrix} * & & & * & \cdots & * \\ & \ddots & & \vdots & & \vdots \\ & & * & * & \cdots & * \\ \hline & & & * & & \\ & & & & \ddots & \\ & & & & & * \end{pmatrix} \pmod{p} \right\}, \end{aligned}$$

where the blocks in the definition of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ are all $(g \times g)$ -matrices.

Clearly, $\mathrm{Iw}_{\mathrm{GL}_g}^+ \subset \mathrm{Iw}_{\mathrm{GL}_g}$ and $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \subset \mathrm{Iw}_{\mathrm{GSp}_{2g}}$. Also observe that, for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$, we have $\gamma_a \in \mathrm{Iw}_{\mathrm{GL}_g}^+$. Moreover, the Iwahori decomposition induces decompositions

$$\mathrm{Iw}_{\mathrm{GL}_g}^+ = U_{\mathrm{GL}_g,1}^{\mathrm{opp}} T_{\mathrm{GL}_g,0} U_{\mathrm{GL}_g,1} \quad \text{and} \quad \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ = U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}} T_{\mathrm{GSp}_{2g},0} U_{\mathrm{GSp}_{2g},0}^+,$$

where $U_{\mathrm{GSp}_{2g},0}^+ := U_{\mathrm{GSp}_{2g},0} \cap \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$.

- Finally, we introduce the notion of “ w -neighbourhood” of some aforementioned p -adic groups. For any $w \in \mathbf{Q}_{>0}$, define

$$\begin{aligned} T_{\mathrm{GL}_g,0}^{(w)} &:= \left\{ \lambda = (\lambda_{ij})_{i,j} \in T_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in T_{\mathrm{GL}_g,0} \right\}, \\ U_{\mathrm{GL}_g,0}^{(w)} &:= \left\{ \lambda = (\lambda_{ij})_{i,j} \in U_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in U_{\mathrm{GL}_g,0} \right\}, \\ B_{\mathrm{GL}_g,0}^{(w)} &:= \left\{ \lambda = (\lambda_{ij})_{i,j} \in B_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in B_{\mathrm{GL}_g,0} \right\}. \end{aligned}$$

The groups $B_{\mathrm{GL}_g,0}^{\mathrm{opp},(w)}$ and $U_{\mathrm{GL}_g,0}^{\mathrm{opp},(w)}$ are defined similarly.

For every $s \in \mathbf{Z}_{\geq 1}$, define

$$\begin{aligned} T_{\mathrm{GL}_g,s}^{(w)} &:= \left\{ \lambda = (\lambda_{ij})_{i,j} \in T_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in T_{\mathrm{GL}_g,s} \right\}, \\ U_{\mathrm{GL}_g,s}^{(w)} &:= \left\{ \lambda = (\lambda_{ij})_{i,j} \in U_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in U_{\mathrm{GL}_g,s} \right\}, \\ B_{\mathrm{GL}_g,s}^{(w)} &:= \left\{ \lambda = (\lambda_{ij})_{i,j} \in B_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in B_{\mathrm{GL}_g,s} \right\}. \end{aligned}$$

The groups $U_{\mathrm{GL}_g, s}^{\mathrm{opp}, (w)}$ and $B_{\mathrm{GL}_g, s}^{\mathrm{opp}, (w)}$ are defined similarly.

Define

$$\mathrm{Iw}_{\mathrm{GL}_g}^{(w)} := \left\{ \lambda = (\lambda_{ij})_{i,j} \in \mathrm{GL}_g(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in \mathrm{Iw}_{\mathrm{GL}_g} \right\}$$

and

$$\mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)} := \left\{ \lambda = (\lambda_{ij})_{i,j} \in \mathrm{GL}_g(\mathcal{O}_{\mathbf{C}_p}) : |\lambda_{ij} - \lambda'_{ij}| \leq p^{-w} \text{ for some } \lambda' = (\lambda'_{ij})_{i,j} \in \mathrm{Iw}_{\mathrm{GL}_g}^+ \right\}.$$

The Iwahori decomposition induces

$$\mathrm{Iw}_{\mathrm{GL}_g}^{(w)} = U_{\mathrm{GL}_g, 1}^{\mathrm{opp}, (w)} T_{\mathrm{GL}_g, 0}^{(w)} U_{\mathrm{GL}_g, 0}^{(w)}$$

and

$$\mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)} = U_{\mathrm{GL}_g, 1}^{\mathrm{opp}, (w)} T_{\mathrm{GL}_g, 0}^{(w)} U_{\mathrm{GL}_g, 1}^{(w)}.$$

We also define

$$T_w = \ker(T_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) \rightarrow T_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}/p^w)),$$

$$U_w = \ker(U_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) \rightarrow U_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}/p^w)),$$

$$B_w = \ker(B_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}) \rightarrow B_{\mathrm{GL}_g}(\mathcal{O}_{\mathbf{C}_p}/p^w)).$$

The groups U_w^{opp} and B_w^{opp} are defined similarly.

Then we have

$$T_{\mathrm{GL}_g, 0}^{(w)} = T_{\mathrm{GL}_g, 0} T_w, \quad U_{\mathrm{GL}_g, 0}^{(w)} = U_{\mathrm{GL}_g, 0} U_w, \quad B_{\mathrm{GL}_g, 0}^{(w)} = B_{\mathrm{GL}_g, 0} B_w.$$

There are similarly identities for $U_{\mathrm{GL}_g, 0}^{\mathrm{opp}, (w)}$ and $B_{\mathrm{GL}_g, 0}^{\mathrm{opp}, (w)}$.

2.2 Siegel modular varieties

We consider Siegel modular varieties of genus g (of principal tame level N) for various level structures at p .

Definition 2.2.1. (i) The **Siegel modular scheme** is the scheme X_0 over $\mathcal{O}_{\mathbf{C}_p}$ that parameterises triples (A, λ, ψ_N) , where (A, λ) is a principally polarised abelian scheme over $\mathcal{O}_{\mathbf{C}_p}$ and

$$\psi_N : V \otimes_{\mathbf{Z}} (\mathbf{Z}/N\mathbf{Z}) \xrightarrow{\sim} A[N]$$

is a symplectic isomorphism with respect to the pairing induced by $\{\cdot, \cdot\}$ on the left and the Weil pairing on the right. Let X denote the base change of X_0 to \mathbf{C}_p .

(ii) For every $n \in \mathbf{Z}_{\geq 1}$, the **Siegel modular variety of principal p^n -level** is the algebraic variety $X_{\Gamma(p^n)}$ over \mathbf{C}_p that parameterises quadruples $(A, \lambda, \psi_N, \psi_{p^n})$, where (A, λ) is a principally polarised abelian variety over \mathbf{C}_p and

$$\psi_N : V \otimes_{\mathbf{Z}} (\mathbf{Z}/N\mathbf{Z}) \xrightarrow{\sim} A[N]$$

and

$$\psi_{p^n} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n\mathbf{Z}) \xrightarrow{\sim} A[p^n]$$

are symplectic isomorphisms.

(iii) The **Siegel modular variety of Iwahori level** is the algebraic variety X_{Iw} over \mathbf{C}_p that parameterises quadruples $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p])$, where (A, λ, ψ_N) is as in (ii) and $\text{Fil}_\bullet A[p]$ is a full flag of $A[p]$ that satisfies

$$(\text{Fil}_\bullet A[p])^\perp \cong \text{Fil}_{2g-\bullet} A[p]$$

with respect to the Weil pairing.

(iv) The **Siegel modular variety of strict Iwahori level** is the algebraic variety X_{Iw^+} over \mathbf{C}_p that parameterises quintuples $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\})$, where $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p])$ is as in (iii) and $\{C_i : i = 1, \dots, g\}$ is a collection of subgroups $C_i \subset A[p]$ of order p such that

$$\text{Fil}_i A[p] = \langle C_1, \dots, C_i \rangle$$

for all $i = 1, \dots, g$.

For $\Gamma \in \{\Gamma(p^n), \text{Iw}^+ = \text{Iw}_{\text{GSp}_{2g}}^+, \text{Iw} = \text{Iw}_{\text{GSp}_{2g}}, \emptyset\}$, it is well-known that the \mathbf{C} -points of the algebraic variety X_Γ can be identified with the locally symmetric space

$$X_\Gamma(\mathbf{C}) = \text{GSp}_{2g}(\mathbf{Q}) \backslash \text{GSp}_{2g}(\mathbf{A}_f) \times \mathbb{H}_g / \Gamma(N) \cdot \Gamma,$$

where

- \mathbf{A}_f is the ring of finite adèles of \mathbf{Q} ;
- \mathbb{H}_g is the disjoint union of the Siegel upper- and lower-half spaces;
- $\Gamma(N) = \{\gamma \in \text{GSp}_{2g}(\widehat{\mathbf{Z}}) : \gamma \equiv 1 \pmod{N}\}$.

Here, we use the fix isomorphism $\mathbf{C}_p \simeq \mathbf{C}$ to view \mathbf{C} as a \mathbf{C}_p -algebra.

Moreover, we have a chain of forgetful maps

$$X_{\Gamma(p^n)} \rightarrow X_{\Gamma(p)} \rightarrow X_{\text{Iw}^+} \rightarrow X_{\text{Iw}} \rightarrow X$$

with the arrows described as follows:

- The first arrow sends $(A, \lambda, \psi_N, \psi_{p^n})$ to $(A, \lambda, \psi_N, p^{n-1}\psi_{p^n})$.
- The second arrow sends $(A, \lambda, \psi_N, \psi_p)$ to $(A, \lambda, \psi_N, \text{Fil}_\bullet^{\psi_p} A[p], \{\langle \psi_p(e_i) \rangle : i = 1, \dots, g\})$ where $\text{Fil}_\bullet^{\psi_p} A[p]$ stands for the full flag

$$0 \subset \langle \psi_p(e_1) \rangle \subset \langle \psi_p(e_1), \psi_p(e_2) \rangle \subset \dots \subset \langle \psi_p(e_1), \dots, \psi_p(e_{2g}) \rangle.$$

- The third arrow sends $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\})$ to $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p])$.
- The fourth arrow sends $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p])$ to (A, λ, ψ_N) .

For $\Gamma \in \{\Gamma(p^n), \text{Iw}^+, \text{Iw}, \emptyset\}$, let \mathcal{X}_Γ be the adic space over $\text{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ associated with X_Γ . The analytifications of the forgetful maps yield

$$\mathcal{X}_{\Gamma(p^n)} \rightarrow \mathcal{X}_{\Gamma(p)} \rightarrow \mathcal{X}_{\text{Iw}^+} \rightarrow \mathcal{X}_{\text{Iw}} \rightarrow \mathcal{X}. \quad (1)$$

By fixing a $\text{GSp}_{2g}(\mathbf{Z})$ -admissible polyhedral cone decomposition as in §B, we show in §B.1 that the chain (1) extends to a chain of log adic spaces¹

$$\overline{\mathcal{X}}_{\Gamma(p^n)} \rightarrow \overline{\mathcal{X}}_{\Gamma(p)} \rightarrow \overline{\mathcal{X}}_{\text{Iw}^+} \rightarrow \overline{\mathcal{X}}_{\text{Iw}} \rightarrow \overline{\mathcal{X}},$$

where, for each $\Gamma \in \{\Gamma(p^n), \text{Iw}^+, \text{Iw}, \emptyset\}$,

¹For a quick review of log adic spaces and the pro-Kummer étale site, see §A.

- $\overline{\mathcal{X}}_\Gamma$ is the adic space over $\mathrm{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ associated with the toroidal compactification \overline{X}_Γ of X_Γ , determined by the fixed polyhedral cone decomposition;
- the log structure on $\overline{\mathcal{X}}_\Gamma$ is the divisorial log structure associated with the boundary divisor $\mathcal{Z}_\Gamma := \overline{\mathcal{X}}_\Gamma \setminus \mathcal{X}_\Gamma$; namely, the corresponding sheaf of monoids \mathcal{M}_Γ on $\overline{\mathcal{X}}_{\Gamma, \text{ét}}$ consists of sections of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma, \text{ét}}}$ that are invertible on the locus away from the boundary divisor;
- $\overline{\mathcal{X}}_\Gamma$ is finite Kummer étale over $\overline{\mathcal{X}}$ and
 - (i) $\overline{\mathcal{X}}_{\Gamma(p^n)} \rightarrow \overline{\mathcal{X}}$ is Galois with Galois group $\mathrm{GSp}_{2g}(\mathbf{Z}/p^n \mathbf{Z})$;
 - (ii) $\overline{\mathcal{X}}_{\Gamma(p)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}}$ is Galois with Galois group $B_{\mathrm{GSp}_{2g}}(\mathbf{Z}/p \mathbf{Z})$;
 - (iii) $\overline{\mathcal{X}}_{\Gamma(p)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+}$ is Galois with Galois group

$$B_{\mathrm{GSp}_{2g}}^+(\mathbf{Z}/p \mathbf{Z}) := \left\{ \begin{pmatrix} \gamma_a & \gamma_b \\ & \gamma_d \end{pmatrix} \in B_{\mathrm{GSp}_{2g}}(\mathbf{Z}/p \mathbf{Z}) : \gamma_a \text{ is diagonal} \right\}.$$

We call $\overline{\mathcal{X}}_\Gamma$ the **toroidal compactification** of \mathcal{X}_Γ (determined by the fixed polyhedral cone decomposition).

Furthermore, we have the *perfectoid Siegel modular variety of infinite level* constructed in [PS16]. See §B.2 for details.

Theorem 2.2.2 ([PS16, Corollaire 4.14]). *There exists a perfectoid space $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ such that*

$$\overline{\mathcal{X}}_{\Gamma(p^\infty)} \sim \varprojlim_n \overline{\mathcal{X}}_{\Gamma(p^n)},$$

where “ \sim ” is in the sense of [SW13, Definition 2.4.1].

Remark 2.2.3. The perfectoid Siegel modular variety is constructed by introducing certain *modified integral structures* at finite levels. More precisely, consider the toroidal compactification $\overline{X}_0^{\mathrm{tor}}$ of X_0 and let $\overline{\mathfrak{X}}^{\mathrm{tor}}$ denote the formal scheme obtained by taking completion along the special fibre of $\overline{\mathfrak{X}}_0^{\mathrm{tor}}$. Let $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$ denote the normalisation of $\overline{\mathcal{X}}^{\mathrm{tor}}$ inside $\overline{\mathcal{X}}_{\Gamma(p^n)}$. In [PS16], the authors consider the modified formal schemes $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}$ obtained by taking certain admissible formal blowups from $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$ and then consider the projective limit

$$\overline{\mathfrak{X}}_{\Gamma(p^\infty)}^{\mathrm{tor-mod}} = \varprojlim_n \overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}$$

in the category of p -adic formal schemes. Finally, $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ is defined to be the adic generic fibre of $\overline{\mathfrak{X}}_{\Gamma(p^\infty)}^{\mathrm{tor-mod}}$.

We summarise the discussion above in the following commutative diagram

$$\begin{array}{ccc}
\mathcal{X}_{\Gamma(p^\infty)} & \hookrightarrow & \overline{\mathcal{X}}_{\Gamma(p^\infty)} \\
\downarrow & & \downarrow \\
\mathcal{X}_{\Gamma(p^n)} & \hookrightarrow & \overline{\mathcal{X}}_{\Gamma(p^n)} \\
\downarrow & & \downarrow \\
\mathcal{X}_{\Gamma(p)} & \hookrightarrow & \overline{\mathcal{X}}_{\Gamma(p)} \\
\downarrow & & \downarrow \\
\mathcal{X}_{\mathrm{Iw}^+} & \hookrightarrow & \overline{\mathcal{X}}_{\mathrm{Iw}^+} \\
\downarrow & & \downarrow \\
\mathcal{X}_{\mathrm{Iw}} & \hookrightarrow & \overline{\mathcal{X}}_{\mathrm{Iw}} \\
\downarrow & & \downarrow \\
\mathcal{X} & \hookrightarrow & \overline{\mathcal{X}}
\end{array}
.$$

where $\mathcal{X}_{\Gamma(p^\infty)}$ is the part of $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ away from the boundary.

There is a natural $\mathrm{GSp}_{2g}(\mathbf{Z}_p)$ -action on $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ permuting the p -power level structures. In particular, the chain of natural projections

$$\overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\Gamma(p^n)} \rightarrow \overline{\mathcal{X}}_{\Gamma(p)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}} \rightarrow \overline{\mathcal{X}}$$

is $\mathrm{GSp}_{2g}(\mathbf{Z}_p)$ -equivariant. According to Proposition B.2.1 (ii), the projection $h_{\Gamma(p^n)} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\Gamma(p^n)}$ (resp., $h_{\mathrm{Iw}^+} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+}$, $h_{\mathrm{Iw}} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}}$) is a pro-Kummer étale Galois cover with Galois group $\Gamma(p^n)$ (resp., $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$, $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$).²

Lemma 2.2.4. *We have the following identities of sheaves*

$$\begin{aligned}
\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}}}^+ &= \left(h_{\mathrm{Iw},*} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+ \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}}, & \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}}} &= \left(h_{\mathrm{Iw},*} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}} \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}} \\
\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}}^+ &= \left(h_{\mathrm{Iw}^+,*} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+ \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+}, & \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}} &= \left(h_{\mathrm{Iw}^+,*} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}} \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+}.
\end{aligned}$$

Proof. We give the proof of the first pair of identities. The second pair can be proven by the same argument.

It suffices to prove the first identity. For any affinoid open $\mathcal{V} \subset \overline{\mathcal{X}}_{\mathrm{Iw}}$ with preimages $\mathcal{V}_n \subset \overline{\mathcal{X}}_{\Gamma(p^n)}$ for $n \in \mathbf{Z}_{>0} \cup \{\infty\}$ such that

$$\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+(\mathcal{V}_\infty) = \left(\varinjlim_n \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^+(\mathcal{V}_n) \right)^\wedge,$$

we have to show

$$\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}}}^+(\mathcal{V}) = \left(\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+(\mathcal{V}_\infty) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}}.$$

²Here we have abused the notation and identify the perfectoid space $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ with the object $\varprojlim_n \overline{\mathcal{X}}_{\Gamma(p^n)}$ in the pro-Kummer étale site $\overline{\mathcal{X}}_{\mathrm{prok\acute{e}t}}$.

Here, “ \wedge ” stands for the p -adic completion. Consider the object $\tilde{\mathcal{V}}_\infty := \varprojlim_n \mathcal{V}_n$ in the pro-Kummer étale site $\overline{\mathcal{X}}_{\text{Iw,prokét}}$. By Lemma A.1.11, each \mathcal{V}_n is finite Kummer étale over \mathcal{V} with Galois group $G_n := \text{IwGSp}_{2g}/\Gamma(p^n)$. Thus,

$$\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+(\mathcal{V}_\infty) = \left(\varinjlim_n \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^+(\mathcal{V}_n) \right)^\wedge = \left(\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw,prokét}}}^+(\tilde{\mathcal{V}}_\infty) \right)^\wedge.$$

By [Dia+19a, Lemma 4.1.7 & Corollary 4.4.13], we know

$$\left(\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^+(\mathcal{V}_n)/p^m \right)^{\text{IwGSp}_{2g}} = \left(\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^+(\mathcal{V}_n)/p^m \right)^{G_n} = \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}}}^+(\mathcal{V})/p^m$$

for every $m \in \mathbf{Z}_{\geq 1}$. This implies

$$\left(\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw,prokét}}}^+(\tilde{\mathcal{V}}_\infty)/p^m \right)^{\text{IwGSp}_{2g}} = \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}}}^+(\mathcal{V})/p^m.$$

Consequently, we have

$$\begin{aligned} \left(\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+(\mathcal{V}_\infty) \right)^{\text{IwGSp}_{2g}} &= \left(\left(\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw,prokét}}}^+(\tilde{\mathcal{V}}_\infty) \right)^\wedge \right)^{\text{IwGSp}_{2g}} = \left(\varprojlim_m \left(\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw,prokét}}}^+(\tilde{\mathcal{V}}_\infty)/p^m \right) \right)^{\text{IwGSp}_{2g}} \\ &= \varprojlim_m \left(\left(\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw,prokét}}}^+(\tilde{\mathcal{V}}_\infty)/p^m \right)^{\text{IwGSp}_{2g}} \right) = \varprojlim_m \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}}}^+(\mathcal{V})/p^m = \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}}}^+(\mathcal{V}). \end{aligned}$$

□

Remark 2.2.5. We point out that the main geometric object studied in [AIP15] is the (toroidally compactified) Siegel modular variety of Iwahori level while ours is of strict Iwahori level. We introduce this deeper level to deal with certain technical issue involved in the construction of the overconvergent Eichler–Shimura morphism in §5.2.

2.3 The flag variety

The Hodge–Tate period map is a $\text{GSp}_{2g}(\mathbf{Z}_p)$ -equivariant morphism from the perfectoid Siegel modular variety to certain flag variety. In this section, let us first describe the target flag variety (and its variants) carefully.

Recall that $V_p = V_{\mathbf{Z}} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ is the standard symplectic space of rank $2g$ over \mathbf{Z}_p . Let P_{Siegel} be the (opposite) Siegel parabolic subgroup of GSp_{2g} defined by

$$P_{\text{Siegel}} := \begin{pmatrix} \text{GL}_g & \\ M_g & \text{GL}_g \end{pmatrix} \cap \text{GSp}_{2g}.$$

Let $\text{Fl} := P_{\text{Siegel}} \backslash \text{GSp}_{2g}$ be the flag variety over \mathbf{Z}_p , parameterising the maximal lagrangians $W \subset V_p$. There is a natural action of GSp_{2g} on Fl by right multiplication. Let $\mathcal{F}\ell$ be the associated adic space of Fl over $\text{Spa}(\mathbf{Q}_p, \mathbf{Z}_p)$, equipped with the induced right action of $\text{GSp}_{2g}(\mathbf{Q}_p)$. Hence, for any p -adically complete sheafy $(\mathbf{Q}_p, \mathbf{Z}_p)$ -algebra (R, R^+) , $\mathcal{F}\ell(R, R^+)$ parameterises maximal lagrangians $W \subset V_p \otimes_{\mathbf{Z}_p} R$. Consider the open subset $\mathcal{F}\ell^\times \subset \mathcal{F}\ell$ whose (R, R^+) -points are

$$\mathcal{F}\ell^\times(R, R^+) = \left\{ (W \subset V_p \otimes_{\mathbf{Z}_p} R) \in \mathcal{F}\ell(R, R^+) : \begin{array}{l} \text{there exists a basis } \{w_i\} \text{ of } W \text{ such that} \\ \text{the matrix } ((w_i, e_{2g+1-j}))_{1 \leq i, j \leq g} \text{ is invertible} \end{array} \right\}.$$

For any $\mathbf{x}_W = (W \subset V_p \otimes_{\mathbf{Z}_p} R) \in \mathcal{F}\ell^\times(R, R^+)$, there exists a unique basis $\{w_i^\square\}$ of W such that

$$((w_i^\square, e_{2g+1-j}))_{1 \leq i, j \leq g} = \mathbb{1}_g.$$

Therefore, there exist global sections $\mathbf{z}_{i,j} \in \mathcal{O}_{\mathcal{F}^\times}(\mathcal{F}^\times)$ such that for any $\mathbf{x}_W \in \mathcal{F}^\times(R, R^+)$,

$$w_i^\square = e_i + \sum_{j=1}^g \mathbf{z}_{i,j}(\mathbf{x}_W) e_{g+j}.$$

Since $\langle w_i^\square, w_j^\square \rangle = 0$, we have

$$\begin{aligned} 0 &= \langle w_i^\square, w_j^\square \rangle \\ &= \langle e_i, \sum_{k=1}^g \mathbf{z}_{j,k}(\mathbf{x}_W) e_{g+k} \rangle + \langle \sum_{k=1}^g \mathbf{z}_{i,k}(\mathbf{x}_W) e_{g+k}, e_j \rangle \\ &= \mathbf{z}_{j,g+1-i}(\mathbf{x}_W) - \mathbf{z}_{i,g+1-j}(\mathbf{x}_W). \end{aligned}$$

That is, the matrix

$$\mathbf{z} := \begin{pmatrix} \mathbf{z}_{1,1} & \cdots & \mathbf{z}_{1,g} \\ \vdots & & \vdots \\ \mathbf{z}_{g,1} & \cdots & \mathbf{z}_{g,g} \end{pmatrix}$$

is symmetric with respect to the anti-diagonal. Moreover, we may use the matrix $\begin{pmatrix} \mathbb{1}_g & \mathbf{z}(\mathbf{x}_W) \end{pmatrix}$ (or just the matrix $\mathbf{z}(\mathbf{x}_W)$) to represent the element $\mathbf{x}_W \in \mathcal{F}^\times(R, R^+)$ because the basis $\{w_i^\square\}$ is represented by the matrix

$$\begin{pmatrix} 1 & & \mathbf{z}_{1,1}(\mathbf{x}_W) & \cdots & \mathbf{z}_{1,g}(\mathbf{x}_W) \\ & \ddots & \vdots & & \vdots \\ & & 1 & \mathbf{z}_{g,1}(\mathbf{x}_W) & \cdots & \mathbf{z}_{g,g}(\mathbf{x}_W) \end{pmatrix} = \begin{pmatrix} \mathbb{1}_g & \mathbf{z}(\mathbf{x}_W) \end{pmatrix}$$

with respect to the standard basis e_1, \dots, e_{2g} of V_p .

In the rest of the paper, we take base change of the adic spaces $\mathcal{F}\ell$ and $\mathcal{F}\ell^\times$ to $\mathrm{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$.

For every $w \in \mathbf{Q}_{>0}$, consider an open adic subspace $\mathcal{F}\ell_w^\times \subset \mathcal{F}\ell^\times$ defined by

$$\mathcal{F}\ell_w^\times := \left\{ \mathbf{x} \in \mathcal{F}\ell^\times : \max_{i,j} \inf_{h \in \mathbf{Z}_p} \{ |\mathbf{z}_{i,j}(\mathbf{x}) - h| \} \leq p^{-w} \right\}.$$

For any algebraically closed complete nonarchimedean field C containing \mathbf{Q}_p , let

$$\mathrm{GSp}_{2g,w}(C) := \left\{ \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{GSp}_{2g}(C) : \begin{array}{l} \gamma_a \in \mathrm{GL}_g(C), \text{ and} \\ \max_{i,j} \inf_{h \in \mathbf{Z}_p} \{ |(\gamma_a^{-1} \gamma_b)_{ij} - h| \} \leq p^{-w} \end{array} \right\}$$

where $(\gamma_a^{-1} \gamma_b)_{ij}$ is the (i, j) -th entry of the matrix $\gamma_a^{-1} \gamma_b$. Then the (C, \mathcal{O}_C) -points of $\mathcal{F}\ell_w^\times$ can be identified with the quotient

$$\mathcal{F}\ell_w^\times(C, \mathcal{O}_C) = P_{\mathrm{Siegel}}(C) \backslash \mathrm{GSp}_{2g,w}(C)$$

so that the natural inclusion $\mathcal{F}\ell_w^\times(C, \mathcal{O}_C) \subset \mathcal{F}\ell(C, \mathcal{O}_C)$ is induced by

$$\mathcal{F}\ell_w^\times(C, \mathcal{O}_C) = P_{\mathrm{Siegel}}(C) \backslash \mathrm{GSp}_{2g,w}(C) \hookrightarrow P_{\mathrm{Siegel}}(C) \backslash \mathrm{GSp}_{2g}(C) = \mathcal{F}\ell(C, \mathcal{O}_C).$$

Recall that there is a natural right action of $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ on $\mathcal{F}\ell$. The following lemma shows that $\mathcal{F}\ell_w^\times$ is stable under the action of the subgroup $\mathrm{Iw}_{\mathrm{GSp}_{2g}} \subset \mathrm{GSp}_{2g}(\mathbf{Q}_p)$.

Lemma 2.3.1. *The adic space $\mathcal{F}\ell_w^\times$ is stable under the right action of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$. Coordinate-wise, the action is given by*

$$\mathcal{F}\ell_w^\times \times \mathrm{Iw}_{\mathrm{GSp}_{2g}} \rightarrow \mathcal{F}\ell_w^\times, \quad \left(\mathbf{z}, \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \right) \mapsto (\gamma_a + \mathbf{z} \gamma_c)^{-1} (\gamma_b + \mathbf{z} \gamma_d).$$

In particular, $\mathcal{F}\ell_w^\times$ is also stable under the right action of the subgroup $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$.

Proof. It follows from the definition that the right action of $\gamma \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}$ indeed sends $\begin{pmatrix} \mathbb{1}_g & \mathbf{z}(\mathbf{x}_W) \end{pmatrix}$ to

$$\begin{pmatrix} \mathbb{1}_g & (\gamma_a + \mathbf{z}(\mathbf{x}_W)\gamma_c)^{-1}(\gamma_b + \mathbf{z}(\mathbf{x}_W)\gamma_d) \end{pmatrix} = (\gamma_a + \mathbf{z}(\mathbf{x}_W)\gamma_c)^{-1} \begin{pmatrix} \mathbb{1}_g & \mathbf{z}(\mathbf{x}_W) \end{pmatrix} \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix}.$$

It remains to show that, for every $\mathbf{x}_W \in \mathcal{F}_w^\times$, the matrix $(\gamma_a + \mathbf{z}(\mathbf{x}_W)\gamma_c)^{-1}(\gamma_b + \mathbf{z}(\mathbf{x}_W)\gamma_d)$ lands in \mathcal{F}_w^\times . But this is straightforward. \square

2.4 Vector bundles on the flag variety

Let $\mathcal{W}_{\mathrm{Fl}} \subset \mathcal{O}_{\mathrm{Fl}}^{2g}$ be the universal maximal lagrangian over Fl . The total space of $\mathcal{W}_{\mathrm{Fl}}$ can be naturally identified with

$$\mathcal{W}_{\mathrm{Fl}} \simeq P_{\mathrm{Siegel}} \backslash (\mathbb{A}^g \times \mathrm{GSp}_{2g})$$

where

- by viewing elements $\vec{v} \in \mathbb{A}^g$ as column vectors, P_{Siegel} acts on \mathbb{A}^g from the left via $\gamma * \vec{v} = {}^t\gamma_a^{-1} \vec{v}$, for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in P_{\mathrm{Siegel}}$;
- P_{Siegel} acts on GSp_{2g} via the left multiplication.

Similarly, consider the linear dual $\mathcal{W}_{\mathrm{Fl}}^\vee$ of $\mathcal{W}_{\mathrm{Fl}}$. Then the total space of $\mathcal{W}_{\mathrm{Fl}}^\vee$ can be naturally identified with

$$\mathcal{W}_{\mathrm{Fl}}^\vee \simeq P_{\mathrm{Siegel}} \backslash (\mathbb{A}^g \times \mathrm{GSp}_{2g})$$

where, by viewing elements $\vec{v} \in \mathbb{A}^g$ as row vectors, P_{Siegel} acts on \mathbb{A}^g from the left via $\gamma * \vec{v} = \vec{v} {}^t\gamma_a$, for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in P_{\mathrm{Siegel}}$. Under this identification, global sections of $\mathcal{W}_{\mathrm{Fl}}^\vee$ are identified with

$$\left\{ \text{algebraic functions } \phi : \mathrm{GSp}_{2g} \rightarrow \mathbb{A}^g : \phi(\gamma \alpha) = \phi(\alpha) \cdot {}^t\gamma_a, \forall \gamma \in P_{\mathrm{Siegel}}, \alpha \in \mathrm{GSp}_{2g} \right\}.$$

For every $i = 1, \dots, g$, we consider a global section \mathbf{s}_i of $\mathcal{W}_{\mathrm{Fl}}^\vee$ defined by

$$\mathbf{s}_i(\alpha) := \text{the } i\text{-th row of } {}^t\alpha_a$$

for all $\alpha = \begin{pmatrix} \alpha_a & \alpha_b \\ \alpha_c & \alpha_d \end{pmatrix} \in \mathrm{GSp}_{2g}$. If we write

$$\mathbf{s} := \begin{pmatrix} \mathbf{s}_1 \\ \vdots \\ \mathbf{s}_g \end{pmatrix} \in (\mathcal{W}_{\mathrm{Fl}}^\vee)^g$$

then we have $\mathbf{s}(\alpha) = {}^t\alpha_a$.

By passing to the adic space \mathcal{F} and restricting to \mathcal{F}_w^\times , the (algebraic) sheaves $\mathcal{W}_{\mathrm{Fl}}$ and $\mathcal{W}_{\mathrm{Fl}}^\vee$ yield (analytic) sheaves $\mathcal{W}_{\mathcal{F}_w^\times}$ and $\mathcal{W}_{\mathcal{F}_w^\times}^\vee$ on \mathcal{F}_w^\times . We still use \mathbf{s}_i 's to denote the restrictions on \mathcal{F}_w^\times of the corresponding algebraic sections. By definition, the sections \mathbf{s}_i 's are non-vanishing on \mathcal{F}_w^\times and hence \mathbf{s}_i^\vee 's are well-defined sections on $\mathcal{W}_{\mathcal{F}_w^\times}^\vee$. We similarly write

$$\mathbf{s}^\vee := \begin{pmatrix} \mathbf{s}_1^\vee & \cdots & \mathbf{s}_g^\vee \end{pmatrix} \in (\mathcal{W}_{\mathcal{F}_w^\times}^\vee)^g.$$

Moreover, the right action of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$ on \mathcal{F}_w^\times induces a right action of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$ on $\mathcal{W}_{\mathcal{F}_w^\times}^\vee$. For later use, we would like to understand the behavior of \mathbf{s}_i^\vee 's under this action.

Lemma 2.4.1. *For any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}$, we have*

$$\gamma^*(s^\vee) = s^\vee \cdot {}^t(\gamma_a + z\gamma_c)^{-1}.$$

The right-hand side means the right multiplication of matrices, where we view s^\vee as a $(1 \times g)$ -matrix with entries s_i^\vee .

Proof. To prove the identity, it suffices to check on the level of (C, \mathcal{O}_C) -points. Using the identification

$$\mathcal{F}\ell_w^\times(C, \mathcal{O}_C) = P_{\mathrm{Siegel}}(C) \backslash \mathrm{GSp}_{2g,w}(C),$$

the sections of $\mathcal{W}_{\mathcal{F}\ell_w^\times}$ can be identified with

$$\left\{ \text{analytic functions } \phi : \mathrm{GSp}_{2g,w} \rightarrow C^g : \phi(\gamma\alpha) = {}^t\gamma_a^{-1} \cdot \phi(\alpha), \forall \gamma \in P_{\mathrm{Siegel}}(C), \alpha \in \mathrm{GSp}_{2g,w}(C) \right\}.$$

Here, elements in C^g are viewed as column vectors. Under this identification, s^\vee sends $\alpha \in \mathrm{GSp}_{2g,w}(C)$ to ${}^t\alpha_a^{-1}$. Notice that a section $\phi : \mathrm{GSp}_{2g,w}(C) \rightarrow C^g$ of $\mathcal{W}_{\mathcal{F}\ell_w^\times}$ is determined by its restriction on

$$\left\{ \begin{pmatrix} \mathbb{1}_g & z \\ & \mathbb{1}_g \end{pmatrix} : {}^t z \mathbb{1}_g = \mathbb{1}_g z, \max_{i,j} \inf_{h \in \mathbf{Z}_p} \{|z_{i,j}(x) - h|\} \leq p^{-w} \right\}.$$

Let $\alpha = \begin{pmatrix} \mathbb{1}_g & z \\ & \mathbb{1}_g \end{pmatrix}$. Then $s^\vee(\alpha) = \mathbb{1}_g$ and

$$(\gamma^*(s^\vee))(\alpha) = s^\vee(\alpha\gamma) = s^\vee \left(\begin{pmatrix} \gamma_a + z\gamma_c & \gamma_b + z\gamma_d \\ \gamma_c & \gamma_d \end{pmatrix} \right) = {}^t(\gamma_a + z\gamma_c)^{-1} = s^\vee(\alpha) \cdot {}^t(\gamma_a + z\gamma_c)^{-1}$$

as desired. \square

An immediate corollary of the lemma above is the following:

Corollary 2.4.2. *For any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}$, we have*

$$\gamma^*(s) = {}^t(\gamma_a + z\gamma_c) \cdot s.$$

2.5 The Hodge–Tate period map and the w -ordinary locus

We briefly recall the well-known Hodge–Tate period map in the setup of (toroidally compactified) Siegel modular variety.

The Hodge–Tate period map (see [PS16, §1] and §B.2) is a morphism of adic spaces

$$\pi_{\mathrm{HT}} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \mathcal{F}\ell.$$

On the level of points, and away from the boundary, the Hodge–Tate period map has the following explicit description. Suppose C is an algebraically closed and complete extension of \mathbf{Q}_p and (A, λ) is a principally polarised abelian variety over C . The Hodge–Tate sequence of A is

$$0 \rightarrow \mathrm{Lie} A \rightarrow T_p A \otimes_{\mathbf{Z}_p} C \rightarrow \omega_{A^\vee} \rightarrow 0,$$

where ω_{A^\vee} is the dual of the Lie algebra of the dual abelian variety A^\vee and the second last map is induced from the Hodge–Tate map $\mathrm{HT}_A : T_p A \rightarrow \omega_{A^\vee}$. Here, we ignore the Tate twist by fixing a compatible system of p -power roots of unity $(\zeta_{p^n})_{n \in \mathbb{Z}_{\geq 1}}$ in \mathbf{C}_p . Notice that every point $\mathbf{x} \in \mathcal{X}_{\Gamma(p^\infty)}(C, \mathcal{O}_C)$ corresponds to a quadruple $(A, \lambda, \psi_N, \psi)$ where (A, λ, ψ_N) is a principally polarised abelian variety over C with a principal level N structure and ψ is a symplectic isomorphism $\psi : V_p \simeq T_p A$. Then π_{HT} sends \mathbf{x} to the maximal lagrangian

$$\mathrm{Lie} A \subset T_p A \otimes_{\mathbf{Z}_p} C \xrightarrow{\psi^{-1}} V_p \otimes_{\mathbf{Z}_p} C.$$

One can extend such an explicit description to the boundary points as well using the language of 1-motives. The details are left to the interested readers.

Remark 2.5.1. There are right $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -actions on both sides of the Hodge–Tate map. The $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -action on $\mathcal{F}\ell$ is given in §2.3. As for the $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -action on $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$, here we only describe the action away from the boundary. (A similar description applies to the boundary points as well, using the language of 1-motives.) Let $\gamma \in \mathrm{GSp}_{2g}(\mathbf{Q}_p)$ and let $m \in \mathbf{Z}$ such that $p^m \gamma \in M_{2g}(\mathbf{Z}_p)$ and $p^{m-1} \gamma \notin M_{2g}(\mathbf{Z}_p)$. Choose $k \in \mathbf{Z}_{\geq 0}$ sufficiently large such that the kernel of $p^m \gamma : A[p^k] \rightarrow A[p^k]$ stabilises. Let $H \subset A[p^k]$ denote the corresponding kernel. Then γ sends $(A, \lambda, \psi_N, \psi)$ to $(A' = A/H, \lambda', \psi'_N, \psi')$ where

- λ' is the induced polarisation on A' ;
- ψ'_N is induced from ψ_N via the isomorphism $A[N] \simeq A'[N]$;
- ψ' is given by the composition

$$V_p \rightarrow V_p \otimes_{\mathbf{Z}_p} \mathbf{Q}_p \xrightarrow{\psi} T_p A \otimes_{\mathbf{Z}_p} \mathbf{Q}_p \rightarrow T_p A' \otimes_{\mathbf{Z}_p} \mathbf{Q}_p$$

with the first map $V_p \rightarrow V_p \otimes_{\mathbf{Z}_p} \mathbf{Q}_p$ sending \vec{v} to $(p^m \gamma)^{-1} \vec{v}$.

One checks that π_{HT} is equivariant with respect to these $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -actions.

Let $\mathcal{G}^{\mathrm{univ}}$ be the tautological semiabelian variety over $\overline{\mathcal{X}}$ extending the universal abelian variety $\mathcal{A}^{\mathrm{univ}}$ over \mathcal{X} . Let $\pi : \mathcal{G}^{\mathrm{univ}} \rightarrow \overline{\mathcal{X}}$ be the structure morphism and let

$$\underline{\omega} := \pi_* \Omega_{\mathcal{G}^{\mathrm{univ}}/\overline{\mathcal{X}}}^1$$

which is a vector bundle of rank g over $\overline{\mathcal{X}}$. Pulling back along the projection $h : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}$, we obtain a vector bundle

$$\underline{\omega}_{\Gamma(p^\infty)} := h^* \underline{\omega}$$

over $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$.

Proposition 2.5.2. *There is a natural isomorphism*

$$\pi_{\mathrm{HT}}^* \mathcal{W}_{\mathcal{F}\ell}^\vee \cong \underline{\omega}_{\Gamma(p^\infty)}.$$

Proof. Let $\mathcal{A}_{\Gamma(p^\infty)}^{\mathrm{univ}}$ be the pullback of $\mathcal{A}^{\mathrm{univ}}$ to $\mathcal{X}_{\Gamma(p^\infty)}$. Away from the boundary, we have a universal trivialisation $\psi^{\mathrm{univ}} : V_p \cong T_p \mathcal{A}_{\Gamma(p^\infty)}^{\mathrm{univ}}$. Let $\psi^{\mathrm{univ}, \vee} : V_p^\vee \cong T_p^\vee \mathcal{A}_{\Gamma(p^\infty)}^{\mathrm{univ}, \vee}$ be the dual trivialisation. The Hodge–Tate map on the universal abelian variety $\mathcal{A}_{\Gamma(p^\infty)}^{\mathrm{univ}}$ induces a map

$$\mathrm{HT}_{\Gamma(p^\infty)} : V_p^\vee \xrightarrow{\psi^{\mathrm{univ}, \vee}} T_p^\vee \mathcal{A}_{\Gamma(p^\infty)}^{\mathrm{univ}, \vee} \rightarrow \underline{\omega}_{\Gamma(p^\infty)}|_{\mathcal{X}_{\Gamma(p^\infty)}}$$

which induces a surjection

$$\mathrm{HT}_{\Gamma(p^\infty)} \otimes \mathrm{id} : V_p^\vee \otimes_{\mathbf{Z}_p} \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty)}} \rightarrow \underline{\omega}_{\Gamma(p^\infty)}|_{\mathcal{X}_{\Gamma(p^\infty)}}.$$

According to §B.2, this surjection extends to a surjection³

$$\mathrm{HT}_{\Gamma(p^\infty)} \otimes \mathrm{id} : V_p^\vee \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}} \rightarrow \underline{\omega}_{\Gamma(p^\infty)}$$

on the entire perfectoid Siegel modular variety.

Consequently, the sheaf $\pi_{\mathrm{HT}}^* \mathcal{W}_{\mathcal{F}\ell}^\vee$, being the universal maximal Lagrangian quotient of $V_p^\vee \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}$, coincides with $\underline{\omega}_{\Gamma(p^\infty)}$. \square

Recall the sections \mathfrak{s}_i of $\mathcal{W}_{\mathcal{F}\ell}^\vee$ defined in §2.4. We define sections $\mathfrak{s}_i \in \underline{\omega}_{\Gamma(p^\infty)}$ by

$$\mathfrak{s}_i := \pi_{\mathrm{HT}}^* \mathfrak{s}_i. \quad (2)$$

From the construction, one sees that

$$\mathfrak{s}_i = \mathrm{HT}_{\Gamma(p^\infty)}(e_i^\vee) \quad (3)$$

for all $i = 1, \dots, g$. These \mathfrak{s}_i 's are examples of *fake Hasse invariants* studied in [Sch15]. We also write

$$\mathfrak{s} := \begin{pmatrix} \mathfrak{s}_1 \\ \vdots \\ \mathfrak{s}_g \end{pmatrix} = \pi_{\mathrm{HT}}^* \mathfrak{s}.$$

To wrap up the section, we introduce the notion of “ w -ordinary locus” of the perfectoid Siegel modular variety. In particular, it is an open subset of $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ which contains the usual ordinary locus.

Definition 2.5.3. *For every $w \in \mathbf{Q}_{>0}$, define*

$$\overline{\mathcal{X}}_{\Gamma(p^\infty),w} := \pi_{\mathrm{HT}}^{-1}(\mathcal{F}\ell_w^\times).$$

We also define

$$\overline{\mathcal{X}}_{\Gamma(p^n),w} := h_n(\overline{\mathcal{X}}_{\Gamma(p^\infty),w}), \quad \overline{\mathcal{X}}_{\mathrm{Iw}^+,w} := h_{\mathrm{Iw}^+}(\overline{\mathcal{X}}_{\Gamma(p^\infty),w}), \quad \overline{\mathcal{X}}_{\mathrm{Iw},w} := h_{\mathrm{Iw}}(\overline{\mathcal{X}}_{\Gamma(p^\infty),w}), \quad \text{and} \quad \overline{\mathcal{X}}_w := h(\overline{\mathcal{X}}_{\Gamma(p^\infty),w}),$$

*where $h_n : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\Gamma(p^n)}$, $h_{\mathrm{Iw}^+} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+}$, $h_{\mathrm{Iw}} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}}$, and $h : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}$ are the natural projections. The subspaces $\overline{\mathcal{X}}_{\Gamma(p^\infty),w}$, $\overline{\mathcal{X}}_{\Gamma(p^n),w}$, $\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}$, $\overline{\mathcal{X}}_{\mathrm{Iw},w}$, and $\overline{\mathcal{X}}_w$ are called the w -**ordinary loci** of $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$, $\overline{\mathcal{X}}_{\Gamma(p^n)}$, $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$, $\overline{\mathcal{X}}_{\mathrm{Iw}}$, and $\overline{\mathcal{X}}$, respectively.*

We still denote by

$$\pi_{\mathrm{HT}} : \overline{\mathcal{X}}_{\Gamma(p^\infty),w} \rightarrow \mathcal{F}\ell_w^\times$$

the restriction of the Hodge–Tate period map on the w -ordinary locus. It is equivariant under the right $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$ -actions on both sides.

Denote by $\mathfrak{z}_{ij} := \pi_{\mathrm{HT}}^* \mathfrak{z}_{ij}$ and $\mathfrak{z} := (\mathfrak{z}_{i,j})_{1 \leq i,j \leq g} = \pi_{\mathrm{HT}}^* \mathfrak{z}$. Let $\mathfrak{s}_i^\vee := \pi_{\mathrm{HT}}^*(\mathfrak{s}_i^\vee)$ and

$$\mathfrak{s}^\vee := (\mathfrak{s}_1^\vee \quad \cdots \quad \mathfrak{s}_g^\vee) = \pi_{\mathrm{HT}}^*(\mathfrak{s}^\vee).$$

³The map $\mathrm{HT}_{\Gamma(p^\infty)} \otimes \mathrm{id}$ here coincides with the map $\mathrm{HT}_{\Gamma(p^\infty)} \otimes \mathrm{id} : V_p \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}} \rightarrow \underline{\omega}_{\Gamma(p^\infty)}$ in §B.2 via the symplectic isomorphism $V_p \simeq V_p^\vee$ sending e_i to $-e_{2g+1-i}^\vee$, for $i = 1, \dots, g$, and sending e_i to e_{2g+1-i}^\vee , for $i = g+1, \dots, 2g$.

By Lemma 2.4.1 and Corollary 2.4.2, we have

$$\gamma^*(\mathfrak{s}^\vee) = \mathfrak{s}^\vee \cdot {}^t(\gamma_a + \mathfrak{z}\gamma_c)^{-1}$$

and

$$\gamma^*\mathfrak{s} = {}^t(\gamma_a + \mathfrak{z}\gamma_c) \cdot \mathfrak{s} \quad (4)$$

for all $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSP}_{2g}}$. We will need these sections \mathfrak{s}_i 's and \mathfrak{s}_i^\vee 's in §3.3 and §3.7.

3 Constructions of the overconvergent automorphic sheaf

In this section, we construct the overconvergent automorphic sheaves using the geometric objects introduced in the previous section. In particular, we generalise the “perfectoid method” which was originally adopted by Chojecki–Hansen–Johansson in [CHJ17] to handle the compact Shimura curves over \mathbf{Q} . Notice that overconvergent automorphic sheaves are first introduced by Andreatta–Iovita–Pilloni in [AIP15] using a different approach. At the end of the section we shall compare the two constructions (when $p > 2g$).

3.1 The perfectoid construction

Let $\mathbf{ALG}_{(\mathbf{Z}_p, \mathbf{Z}_p)}$ be the category of complete sheafy $(\mathbf{Z}_p, \mathbf{Z}_p)$ -algebras. We consider the functor

$$\mathbf{ALG}_{(\mathbf{Z}_p, \mathbf{Z}_p)} \rightarrow \mathbf{SETS}, \quad (R, R^+) \mapsto \mathrm{Hom}_{\mathbf{GROUP}}^{\mathrm{cts}}(T_{\mathrm{GL}_g, 0}, R^\times),$$

which is represented by the $(\mathbf{Z}_p, \mathbf{Z}_p)$ -algebra $(\mathbf{Z}_p[[T_{\mathrm{GL}_g, 0}]], \mathbf{Z}_p[[T_{\mathrm{GL}_g, 0}]])$. The *weight space* is

$$\mathcal{W} := \mathrm{Spa}(\mathbf{Z}_p[[T_{\mathrm{GL}_g, 0}]], \mathbf{Z}_p[[T_{\mathrm{GL}_g, 0}]])^{\mathrm{rig}},$$

where the superscript “rig” stands for taking the generic fibre. Every continuous group homomorphism $\kappa : T_{\mathrm{GL}_g, 0} \rightarrow R^\times$ can be expressed as $\kappa = (\kappa_1, \dots, \kappa_g)$ where each $\kappa_i : \mathbf{Z}_p^\times \rightarrow R^\times$ is a continuous group homomorphism. We write $\kappa^\vee := (-\kappa_g, \dots, -\kappa_1)$ where $-\kappa_i$ is the inverse of κ_i .

We adapt the terminologies of “small weights” and “affinoid weights” introduced in [CHJ17] to our setting:

Definition 3.1.1. (i) A **small \mathbf{Z}_p -algebra** is a p -torsion free reduced ring which is also a finite $\mathbf{Z}_p[[T_1, \dots, T_d]]$ -algebra for some $d \in \mathbf{Z}_{\geq 0}$. In particular, a small \mathbf{Z}_p -algebra is equipped with a canonical adic profinite topology and is complete with respect to the p -adic topology.

(ii) A **small weight** is a pair $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ where $R_{\mathcal{U}}$ is a small \mathbf{Z}_p -algebra and $\kappa_{\mathcal{U}} : T_{\mathrm{GL}_g, 0} \rightarrow R_{\mathcal{U}}^\times$ is a continuous group homomorphism such that $\kappa_{\mathcal{U}}((1+p)\mathbb{1}_g) - 1$ is topologically nilpotent in $R_{\mathcal{U}}$ with respect to the p -adic topology. By the universal property of the weight space, we obtain a natural morphism

$$\mathrm{Spa}(R_{\mathcal{U}}, R_{\mathcal{U}})^{\mathrm{rig}} \rightarrow \mathcal{W}.$$

Occasionally, we abuse the terminology and call $\mathcal{U} := \mathrm{Spa}(R_{\mathcal{U}}, R_{\mathcal{U}})$ a small weight. For later use, we write $R_{\mathcal{U}}^+ := R_{\mathcal{U}}$ in this situation.

(iii) An **affinoid weight** is a pair $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ where $R_{\mathcal{U}}$ is a reduced Tate algebra topologically of finite type over \mathbf{Q}_p and $\kappa_{\mathcal{U}} : T_{\mathrm{GL}_g, 0} \rightarrow R_{\mathcal{U}}^\times$ is a continuous group homomorphism. By the universal property of weight space, we obtain a natural morphism

$$\mathrm{Spa}(R_{\mathcal{U}}, R_{\mathcal{U}}^\circ) \rightarrow \mathcal{W}.$$

Occasionally, we abuse the terminology and call $\mathcal{U} := \mathrm{Spa}(R_{\mathcal{U}}, R_{\mathcal{U}}^\circ)$ an affinoid weight. For later use, we write $R_{\mathcal{U}}^+ = R_{\mathcal{U}}^\circ$ in this situation.

(iv) By a **weight**, we shall mean either a small weight or an affinoid weight.

Remark 3.1.2. For any $n \in \mathbf{Z}_{\geq 0}$, we view n as a weight by identifying it with the character

$$T_{\mathrm{GL}_g,0} \rightarrow \mathbf{Z}_p^\times, \quad \mathrm{diag}(\tau_1, \dots, \tau_g) \mapsto \prod_{i=1}^g \tau_i^n.$$

Moreover, for any weight $\kappa = (\kappa_1, \dots, \kappa_g)$, we write $\kappa + n$ for the weight $(\kappa_1 + n, \dots, \kappa_g + n)$ defined by

$$\mathrm{diag}(\tau_1, \dots, \tau_g) \mapsto \prod_{i=1}^g \kappa_i(\tau_i) \tau_i^n.$$

We adopt the notation of “mixed completed tensor” used in [CHJ17]:

Definition 3.1.3. Let R be a small \mathbf{Z}_p -algebra.

(i) For any \mathbf{Z}_p -module M , we define ⁴

$$M \widehat{\otimes}' R := \varprojlim_{j \in J} (M \otimes_{\mathbf{Z}_p} R/I_j)$$

where $\{I_j : j \in J\}$ runs through a cofinal system of neighborhood of 0 consisting of \mathbf{Z}_p -submodules of R . If, in addition, M is a \mathbf{Z}_p -algebra, then $M \widehat{\otimes}' R$ is also a \mathbf{Z}_p -algebra.

(ii) Let B be a \mathbf{Q}_p -Banach space and let B_0 be an open and bounded \mathbf{Z}_p -submodule. We define the **mixed completed tensor**

$$B \widehat{\otimes} R := (B_0 \widehat{\otimes}' R)[\frac{1}{p}].$$

which is in fact independent of the choice of B_0 .

Definition 3.1.4. Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a weight.

- (i) For any \mathbf{Z}_p -module M , the term $M \widehat{\otimes} R_{\mathcal{U}}^+$ will either stand for $M \widehat{\otimes}' R_{\mathcal{U}}$ in the case of small weights (notice that $R_{\mathcal{U}} = R_{\mathcal{U}}^+$ in this case), or stand for the p -adically completed tensor over \mathbf{Z}_p in the case of affinoid weights.
- (ii) For any \mathbf{Q}_p -Banach space B , the term $B \widehat{\otimes} R_{\mathcal{U}}$ will either stand for the mixed completed tensor in the case of small weights, or stand for the usual p -adically completed tensor over \mathbf{Q}_p in the case of affinoid weights.

Remark 3.1.5. For a uniform Banach \mathbf{Q}_p -algebra B and any weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$, the tensor product $B \widehat{\otimes} R_{\mathcal{U}}$ also admits a structure of uniform \mathbf{Q}_p -Banach algebra. In particular, if B° is the unit ball of B (with respect to the unique power-multiplicative Banach algebra norm), then the unit ball in $B \widehat{\otimes} R_{\mathcal{U}} = B \widehat{\otimes}_{\mathbf{Q}_p} R_{\mathcal{U}}^+[1/p]$ is given by $B^\circ \widehat{\otimes} R_{\mathcal{U}}^+$. Here, note that $R_{\mathcal{U}}^+[1/p]$ has a structure of a uniform Banach \mathbf{Q}_p -algebra given by the corresponding spectral norm (see [CHJ17, pp. 202]).

Next, we introduce the notion of “ r -analytic functions”.

Definition 3.1.6. Let $r \in \mathbf{Q}_{>0}$ and $n \in \mathbf{Z}_{\geq 1}$. Let B be a uniform \mathbf{C}_p -Banach algebra and let B° be the corresponding unit ball.

⁴Our notation $\widehat{\otimes}'$ corresponds to the notation $\widehat{\otimes}$ in [CHJ17, Definition 6.3]. We make this change to distinguish from the one in Definition 3.1.3 (ii).

- (i) A function $f : \mathbf{Z}_p^n \rightarrow B$ (resp., a function $f : (\mathbf{Z}_p^\times)^n \rightarrow B$) is called **r -analytic** if for every $\underline{a} = (a_1, \dots, a_n) \in \mathbf{Z}_p^n$ (resp., every $\underline{a} = (a_1, \dots, a_n) \in (\mathbf{Z}_p^\times)^n$), there exists a power series $f_{\underline{a}} \in B[[T_1, \dots, T_n]]$ which converges on the n -dimensional closed unit ball $\mathbf{B}^n(0, p^{-r}) \subset \mathbf{C}_p^n$ of radius p^{-r} such that

$$f(x_1 + a_1, \dots, x_n + a_n) = f_{\underline{a}}(x_1, \dots, x_n)$$

for all $x_i \in p^{\lceil r \rceil} \mathbf{Z}_p$, $i = 1, \dots, n$. Here $\lceil r \rceil$ stands for the smallest integer that is greater or equal to r .

- (ii) Let $C^{r\text{-an}}(\mathbf{Z}_p^n, B)$ (resp., $C^{r\text{-an}}((\mathbf{Z}_p^\times)^n, B)$) denote the set of r -analytic functions from \mathbf{Z}_p^n (resp., $(\mathbf{Z}_p^\times)^n$) to B .
- (iii) Let $C^{r\text{-an}}(\mathbf{Z}_p^n, B^\circ)$ (resp., $C^{r\text{-an}}((\mathbf{Z}_p^\times)^n, B^\circ)$) denote the subset of $C^{r\text{-an}}(\mathbf{Z}_p^n, B)$ (resp., $C^{r\text{-an}}((\mathbf{Z}_p^\times)^n, B)$) consisting of those functions with value in B° .

Remark 3.1.7. We claim that $C^{r\text{-an}}(\mathbf{Z}_p^n, B)$ (resp., $C^{r\text{-an}}((\mathbf{Z}_p^\times)^n, B)$) admits a natural structure of uniform \mathbf{C}_p -Banach algebra. Indeed, express \mathbf{Z}_p^n as the disjoint union of $p^{n\lceil r \rceil}$ closed balls of radius $p^{\lceil r \rceil}$, labelled by an index set A of size $p^{n\lceil r \rceil}$. Then, for every $f \in C^{r\text{-an}}(\mathbf{Z}_p^n, B)$, the restriction of f on each closed ball (with label $a \in A$) is given by a power series

$$f_a \in B\left(\frac{T_1}{p^r}, \dots, \frac{T_n}{p^r}\right)$$

where $B\left(\frac{T_1}{p^r}, \dots, \frac{T_n}{p^r}\right)$ stands for the subset of $B[[T_1, \dots, T_n]]$ which converges on the n -dimensional closed unit ball $\mathbf{B}^n(0, p^{-r}) \subset \mathbf{C}_p^n$. Let $|\bullet|_B$ be the unique power-multiplicative norm on B . We can equip $B\left(\frac{T_1}{p^r}, \dots, \frac{T_n}{p^r}\right)$ with the following norm: for every $g = \sum_{\nu \in \mathbf{Z}_{\geq 0}^n} b_\nu T^\nu$, we put

$$|g| := \sup_{\nu \in \mathbf{Z}_{\geq 0}^n} |b_\nu|_B \cdot p^{-r|\nu|}.$$

Finally, if $f \in C^{r\text{-an}}(\mathbf{Z}_p^n, B)$ is represented by $\{f_a\}_{a \in A}$, we put $|f| := \sup_{a \in A} |f_a|$. This is indeed a uniform Banach norm with unit ball $C^{r\text{-an}}(\mathbf{Z}_p^n, B^\circ)$.

Definition 3.1.8. (i) A weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ is called **r -analytic** if it is r -analytic when viewed as a function

$$\kappa_{\mathcal{U}} : (\mathbf{Z}_p^\times)^g \rightarrow R_{\mathcal{U}}^\times \subset \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}$$

via the identification $T_{\text{GL}_g, 0} \cong (\mathbf{Z}_p^\times)^g$.

- (ii) For a weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$, we write $r_{\mathcal{U}}$ for the smallest positive integer r such that the weight is r -analytic.

Remark 3.1.9. (i) It is well-known that every continuous character $\mathbf{Z}_p^\times \rightarrow R_{\mathcal{U}}^\times$ is r -analytic for sufficiently large r . Moreover, if such a character is r -analytic, it necessarily extends to a character

$$\mathbf{Z}_p^\times(1 + p^{r+1} \mathcal{O}_{\mathbf{C}_p}) \rightarrow (\mathcal{O}_{\mathbf{C}_p} \widehat{\otimes} R_{\mathcal{U}}^+)^{\times} \subset \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}.$$

See, for example, [CHJ17, Proposition 2.6].

- (ii) If we write $\kappa_{\mathcal{U}} = (\kappa_{\mathcal{U}, 1}, \dots, \kappa_{\mathcal{U}, g})$ with components $\kappa_{\mathcal{U}, i} : \mathbf{Z}_p^\times \rightarrow R_{\mathcal{U}}^\times$, then $\kappa_{\mathcal{U}}$ is r -analytic if and only if all $\kappa_{\mathcal{U}, i}$'s are r -analytic. In this case, for any $w \in \mathbf{Q}_{>0}$ with $w > 1 + r_{\mathcal{U}}$, $\kappa_{\mathcal{U}}$ extends to a character

$$\kappa_{\mathcal{U}} : T_{\text{GL}_g, 0}^{(w)} \rightarrow (\mathcal{O}_{\mathbf{C}_p} \widehat{\otimes} R_{\mathcal{U}}^+)^{\times} \subset \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}.$$

Definition 3.1.10. Let B be a uniform \mathbf{C}_p -Banach algebra.

(i) A function $\psi : U_{\text{GL}_g,1}^{\text{opp}} \rightarrow B$ is called *r-analytic* if, under the (topological) identification

$$U_{\text{GL}_g,1}^{\text{opp}} = \begin{pmatrix} 1 & & & \\ p\mathbf{Z}_p & 1 & & \\ \vdots & & \ddots & \\ p\mathbf{Z}_p & \dots & p\mathbf{Z}_p & 1 \end{pmatrix} \simeq \mathbf{Z}_p^{\frac{(g-1)g}{2}},$$

the function

$$\psi : \mathbf{Z}_p^{\frac{(g-1)g}{2}} \rightarrow B$$

is *r-analytic*. Let $C^{r\text{-an}}(U_{\text{GL}_g,1}^{\text{opp}}, B)$ denote the space of such functions.

(ii) Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be an *r-analytic weight*. Using the decomposition $B_{\text{GL}_g,0} = T_{\text{GL}_g,0} U_{\text{GL}_g,0}$, we extend $\kappa_{\mathcal{U}}$ to a group homomorphism $\kappa_{\mathcal{U}} : B_{\text{GL}_g,0} \rightarrow R_{\mathcal{U}}^{\times}$ by setting $\kappa_{\mathcal{U}}|_{U_{\text{GL}_g,0}} = 1$. Define

$$C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B) := \left\{ f : \text{Iw}_{\text{GL}_g} \rightarrow B : \begin{array}{l} f(\gamma\beta) = \kappa_{\mathcal{U}}(\beta)f(\gamma), \forall \beta \in B_{\text{GL}_g,0}, \gamma \in \text{Iw}_{\text{GL}_g} \\ f|_{U_{\text{GL}_g,1}^{\text{opp}}} \text{ is } r\text{-analytic} \end{array} \right\}.$$

(iii) Let $C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B^{\circ})$ denote the subset of $C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B)$ consisting of those functions with value in B° .

Remark 3.1.11. According to Remark 3.1.7, $C^{r\text{-an}}(U_{\text{GL}_g,1}^{\text{opp}}, B)$ admits a structure of uniform \mathbf{C}_p -Banach algebra. Notice that an element in $C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B)$ is determined by its restriction on $U_{\text{GL}_g,1}^{\text{opp}}$. Consequently, $C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B)$ admits a structure of uniform \mathbf{C}_p -Banach algebra via the identification

$$C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B) \cong C^{r\text{-an}}(U_{\text{GL}_g,1}^{\text{opp}}, B).$$

In particular, $C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B^{\circ})$ is the corresponding unit ball in $C_{\kappa_{\mathcal{U}}}^{r\text{-an}}(\text{Iw}_{\text{GL}_g}, B)$.

Remark 3.1.12. Let $\kappa_{\mathcal{U}}$ be a weight and let $w \in \mathbf{Q}_{>0}$ with $w > r_{\mathcal{U}} + 1$. Recall that we have a decomposition $B_{\text{GL}_g,0}^{(w)} = T_{\text{GL}_g,0}^{(w)} U_{\text{GL}_g,0}^{(w)}$. Since $w > 1 + r_{\mathcal{U}}$, $\kappa_{\mathcal{U}}$ extends to a character on $T_{\text{GL}_g,0}^{(w)}$, and hence to a character on $B_{\text{GL}_g,0}^{(w)}$ by setting $\kappa_{\mathcal{U}}|_{U_{\text{GL}_g,0}^{(w)}} = 0$.

We claim that every element f in $C_{\kappa_{\mathcal{U}}}^{w\text{-an}}(\text{Iw}_{\text{GL}_g}, B)$ (resp., $C_{\kappa_{\mathcal{U}}}^{w\text{-an}}(\text{Iw}_{\text{GL}_g}, B^{\circ})$) naturally extends to a function

$$f : \text{Iw}_{\text{GL}_g}^{(w)} \rightarrow B \quad (\text{resp., } f : \text{Iw}_{\text{GL}_g}^{(w)} \rightarrow B^{\circ})$$

such that $f(\gamma\beta) = \kappa_{\mathcal{U}}(\beta)f(\gamma)$ for all $\beta \in B_{\text{GL}_g,0}^{(w)}$ and $\gamma \in \text{Iw}_{\text{GL}_g}^{(w)}$. Indeed, we have decomposition

$$\text{Iw}_{\text{GL}_g}^{(w)} = U_{\text{GL}_g,1}^{\text{opp},(w)} T_{\text{GL}_g,0}^{(w)} U_{\text{GL}_g,0}^{(w)}.$$

Then for every $\nu \in U_{\text{GL}_g,1}^{\text{opp},(w)}$, $\tau \in T_{\text{GL}_g,0}^{(w)}$, and $\nu' \in U_{\text{GL}_g,0}^{(w)}$, we put

$$f(\nu\tau\nu') = \kappa_{\mathcal{U}}(\tau)f(\nu).$$

It is straightforward to check that f is well-defined and satisfies the required condition.

Definition 3.1.13. As a consequence of Remark 3.1.12, given $w \in \mathbf{Q}_{>0}$ with $w > 1 + r_{\mathcal{U}}$, there is a natural left action of $\mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)}$ on $C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, B)$ and $C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, B^\circ)$ (hence also a left action of $\mathrm{Iw}_{\mathrm{GL}_g}^+$) given by

$$(\gamma \cdot f)(\gamma') = f({}^t\gamma \gamma')$$

for all $\gamma \in \mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)}$, $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$, and $f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, B)$ (resp., $C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, B^\circ)$). This left action is denoted by $\rho_{\kappa_{\mathcal{U}}} : \mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)} \rightarrow \mathrm{Aut}(C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, B))$ (resp., $\rho_{\kappa_{\mathcal{U}}} : \mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)} \rightarrow \mathrm{Aut}(C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, B^\circ))$).

We are ready to define the sheaf of overconvergent Siegel modular forms.

Definition 3.1.14. Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a weight and let $w \in \mathbf{Q}_{>0}$ with $w > 1 + r_{\mathcal{U}}$.

(i) Let $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}}$ be the sheaf on $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ given by

$$\mathcal{Y} \mapsto \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}(\mathcal{Y}) \widehat{\otimes} R_{\mathcal{U}}$$

for every affinoid open subset $\mathcal{Y} \subset \overline{\mathcal{X}}_{\Gamma(p^\infty), w}$. This is in fact a sheaf of uniform \mathbf{C}_p -Banach algebra; i.e., $(\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})(\mathcal{Y})$ is a uniform \mathbf{C}_p -Banach algebra for every affinoid open \mathcal{Y} .

Similarly, let $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}^+ \widehat{\otimes} R_{\mathcal{U}}^+$ be the sheaf on $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ given by

$$\mathcal{Y} \mapsto \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}^+(\mathcal{Y}) \widehat{\otimes} R_{\mathcal{U}}^+$$

for every affinoid open subset $\mathcal{Y} \subset \overline{\mathcal{X}}_{\Gamma(p^\infty), w}$.

(ii) For any $r \in \mathbf{Q}_{>0}$ with $r > 1 + r_{\mathcal{U}}$, let $\mathcal{C}_{\kappa_{\mathcal{U}}}^{r-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ denote the sheaf on $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ given by

$$\mathcal{Y} \mapsto C_{\kappa_{\mathcal{U}}}^{r-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}(\mathcal{Y}) \widehat{\otimes} R_{\mathcal{U}})$$

for every affinoid open subset $\mathcal{Y} \subset \overline{\mathcal{X}}_{\Gamma(p^\infty), w}$. This is also a sheaf of uniform \mathbf{C}_p -Banach algebra.

The sheaf $\mathcal{C}_{\kappa_{\mathcal{U}}}^{r-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}^+ \widehat{\otimes} R_{\mathcal{U}}^+)$ is defined in the same way.

(iii) The **sheaf of w -overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is a subsheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ of $h_{\mathrm{Iw}^+, *}\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ defined as follows. For every affinoid open subset $\mathcal{V} \subset \overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$ with $\mathcal{V}_\infty = h_{\mathrm{Iw}^+}^{-1}(\mathcal{V})$, we put

$$\underline{\omega}_w^{\kappa_{\mathcal{U}}}(\mathcal{V}) := \left\{ f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}) : \gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f, \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \right\}.$$

Here, $\gamma^* f$ stands for the left action of γ on $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}$ induced by the natural right $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -action on $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ defined in §2.5.

Similarly, the **sheaf of integral w -overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is a subsheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}, +}$ of $h_{\mathrm{Iw}^+, *}\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}^+ \widehat{\otimes} R_{\mathcal{U}}^+)$ defined as follows.

For every affinoid open subset $\mathcal{V} \subset \overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$ with $\mathcal{V}_\infty = h_{\mathrm{Iw}^+}^{-1}(\mathcal{V})$, we put

$$\underline{\omega}_w^{\kappa_{\mathcal{U}}, +}(\mathcal{V}) := \left\{ f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}^+) : \gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f, \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \right\}.$$

- (iv) The **space of w -overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is defined to be

$$M_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}} := H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}).$$

We similarly define the **space of integral w -overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** to be

$$M_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}, +} := H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}, +}).$$

- (v) Taking limit with respect to w , the **space of overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is

$$M_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}} := \lim_{w \rightarrow \infty} M_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}}.$$

Similarly, the **space of integral overconvergent Siegel modular forms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is

$$M_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}, +} := \lim_{w \rightarrow \infty} M_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}, +}.$$

- (vi) Recall that $\mathcal{Z}_{\mathrm{Iw}^+} = \overline{\mathcal{X}}_{\mathrm{Iw}^+} \setminus \mathcal{X}_{\mathrm{Iw}^+}$ is the boundary divisor. The **sheaf of w -overconvergent Siegel cuspforms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is defined to be the subsheaf $\underline{\omega}_{w, \mathrm{cusp}}^{\kappa_{\mathcal{U}}} = \underline{\omega}_w^{\kappa_{\mathcal{U}}}(-\mathcal{Z}_{\mathrm{Iw}^+})$ of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ consisting of sections that vanish along $\mathcal{Z}_{\mathrm{Iw}^+}$.

A w -overconvergent Siegel modular form of strict Iwahori level and weight $\kappa_{\mathcal{U}}$ is called **cuspidal** if it is an element of

$$S_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}} := H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_{w, \mathrm{cusp}}^{\kappa_{\mathcal{U}}}).$$

Moreover, by taking limit with respect to w , the **space of overconvergent Siegel cuspforms of strict Iwahori level and weight $\kappa_{\mathcal{U}}$** is defined to be

$$S_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}} := \lim_{w \rightarrow \infty} S_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}}.$$

Remark 3.1.15. Notice that, in Definition 3.1.14 (iii), for every $\mathbf{x} \in \overline{\mathcal{X}}_{\Gamma(p^\infty), w}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and any $\begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$, we have $\gamma_a + \mathfrak{z}(\mathbf{x})\gamma_c \in \mathrm{Iw}_{\mathrm{GL}_g}^{+, (w)}$. Hence, $\rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z}\gamma_c)$ is well-defined.

Remark 3.1.16. The definition above yields an analogue to the complex theory, which we describe in the following.

Suppose $k = (k_1, \dots, k_g) \in \mathbf{Z}_{\geq 0}^g$ is a dominant weight for GL_g and let $\rho_k : \mathrm{GL}_g(\mathbf{C}) \rightarrow \mathrm{GL}(\mathbf{V}_k)$ be the corresponding irreducible representation of GL_g of highest weight k . Let \mathbb{H}_g^+ be the (complex) Siegel upper-half space. Then a classical Siegel modular form of weight k and level Γ is a holomorphic function $f : \mathbb{H}_g^+ \rightarrow \mathbf{V}_k$ such that

$$f(\gamma \cdot \mathbf{x}) = \rho_k(\gamma_c \mathbf{x} + \gamma_d)f(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{H}_g^+$ and $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma \subset \mathrm{GSp}_{2g}(\mathbf{Z})$.

In our case, a w -overconvergent Siegel modular form $f \in M_{\mathrm{Iw}^+, w}^{\kappa_{\mathcal{U}}}$ can be viewed as a function

$$f : \overline{\mathcal{X}}_{\Gamma(p^\infty), w} \rightarrow C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}})$$

satisfying

$$f(\mathbf{x} \cdot \gamma) = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z}\gamma_c)^{-1}f(\mathbf{x})$$

for all $\mathbf{x} \in \overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ and $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \subset \mathrm{GSp}_{2g}(\mathbf{Z}_p)$. Notice that $C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}})$ is an analytic analogue of the algebraic representation \mathbf{V}_k .

To simplify the notation, we defined a “twisted” left action of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ on $\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ by

$$\gamma \cdot f := \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c) \gamma^* f.$$

Then sections of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ are precisely the $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -invariant sections of $h_{\mathrm{Iw}^+, *}\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ under the twisted action.

Remark 3.1.17. The sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ is functorial in the weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$. Given a map of weights $R_{\mathcal{U}} \rightarrow R_{\mathcal{U}'}$ and $w > \max\{1 + r_{\mathcal{U}}, 1 + r_{\mathcal{U}'}\}$, we obtain a natural map $\underline{\omega}_w^{\kappa_{\mathcal{U}}} \rightarrow \underline{\omega}_w^{\kappa_{\mathcal{U}'}}$ induced from

$$C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}}) \rightarrow C_{\kappa_{\mathcal{U}'}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}'}).$$

3.2 Hecke operators

In this subsection, we spell out how the Hecke operators act on the overconvergent Siegel modular forms. The Hecke operators at the primes dividing the tame level N are not considered in this paper.

Through out this subsection, let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a weight and $w > 1 + r_{\mathcal{U}}$.

Hecke operators outside Np . We define the Hecke operators outside Np using correspondences. Let ℓ be a rational prime that does not divide Np . For every $\gamma \in \mathrm{GSp}_{2g}(\mathbf{Q}_\ell) \cap M_{2g}(\mathbf{Z}_\ell)$, consider the moduli space $X_{\gamma, \mathrm{Iw}^+}$ over X_{Iw^+} parameterising *isogenies of type γ* . More precisely, $X_{\gamma, \mathrm{Iw}^+}$ is the moduli space of sextuples

$$(A, \lambda, \psi_N, \mathrm{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\}, L)$$

where $(A, \lambda, \psi_N, \mathrm{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\}) \in X_{\mathrm{Iw}^+}$ and $L \subset A$ is a subgroup of finite order such that the isogeny $(A, \lambda) \rightarrow (A/L, \lambda')$ is of type γ in the sense of [FC90, Chapter VII, §3], where λ' stands for the induced principal polarisation. According to *loc. cit.*, for every isogeny of type γ , its dual isogeny is also of type γ . In particular, the assignment

$$(A, \lambda, \psi_N, \mathrm{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\}, L) \mapsto (A' = A/L, \lambda', \psi'_N, \mathrm{Fil}_\bullet A'[p], \{C'_i : i = 1, \dots, g\}, L')$$

defines an isomorphism $\Phi_\gamma : X_{\gamma, \mathrm{Iw}^+} \xrightarrow{\sim} X_{\gamma, \mathrm{Iw}^+}$, where

- λ' is the induced polarisation on A' ;
- ψ'_N , $\mathrm{Fil}_\bullet A'[p]$, and C'_i 's are induced from ψ_N , $\mathrm{Fil}_\bullet A[p]$, and C_i 's, respectively, via the isomorphisms $A[N] \simeq A'[N]$ and $A[p] \simeq A'[p]$;
- L' is defined by the dual isogeny of $(A, \lambda) \rightarrow (A', \lambda')$.

There are two finite étale projections

$$\begin{array}{ccc} & X_{\gamma, \mathrm{Iw}^+} & \\ \mathrm{pr}_1 \swarrow & & \searrow \mathrm{pr}_2 \\ X_{\mathrm{Iw}^+} & & X_{\mathrm{Iw}^+} \end{array}$$

where pr_1 is the forgetful map and pr_2 sends the sextuple $(A, \lambda, \psi_N, \text{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\}, L)$ to the quintuple $(A' = A/L, \lambda', \psi'_N, \text{Fil}_\bullet A'[p], \{C'_i : i = 1, \dots, g\})$ described as above. Clearly, we have $\text{pr}_1 = \text{pr}_2 \circ \Phi_\gamma$.

Let $\mathcal{X}_{\gamma, \text{Iw}^+}$ be the adic space associated with X_{γ, Iw^+} by taking analytification. We obtain finite étale morphisms $\text{pr}_1, \text{pr}_2 : \mathcal{X}_{\gamma, \text{Iw}^+} \rightrightarrows \mathcal{X}_{\text{Iw}^+}$ as well as an isomorphism $\Phi_\gamma : \mathcal{X}_{\gamma, \text{Iw}^+} \rightarrow \mathcal{X}_{\gamma, \text{Iw}^+}$. We further pass to the w -ordinary locus. More precisely, let $\mathcal{X}_{\gamma, \text{Iw}^+, w}$ denote the preimage of $\mathcal{X}_{\text{Iw}^+, w}$ under the projection pr_1 . Notice that Φ_γ preserves $\mathcal{X}_{\gamma, \text{Iw}^+, w}$ as the isogeny $(A, \lambda) \rightarrow (A', \lambda')$ induces a symplectic isomorphism $T_p A \cong T_p A'$. Hence, we obtain finite étale morphisms

$$\begin{array}{ccc} & \mathcal{X}_{\gamma, \text{Iw}^+, w} & \\ \text{pr}_1 \swarrow & & \searrow \text{pr}_2 \\ \mathcal{X}_{\text{Iw}^+, w} & & \mathcal{X}_{\text{Iw}^+, w} \end{array} \quad (5)$$

and an isomorphism $\Phi_\gamma : \mathcal{X}_{\gamma, \text{Iw}^+, w} \xrightarrow{\sim} \mathcal{X}_{\gamma, \text{Iw}^+, w}$. We still have $\text{pr}_1 = \text{pr}_2 \circ \Phi_\gamma$.

In order to define the Hecke operator, we shall first construct a natural isomorphism

$$\varphi_\gamma : \text{pr}_2^* \underline{\omega}_w^{\kappa_{\mathcal{U}}} \xrightarrow{\sim} \text{pr}_1^* \underline{\omega}_w^{\kappa_{\mathcal{U}}}.$$

Here we have abused the notation and still write $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ for its restriction to $\mathcal{X}_{\text{Iw}^+, w}$.

Indeed, pulling back the diagram (5) along the projection $h_{\text{Iw}^+} : \mathcal{X}_{\Gamma(p^\infty), w} \rightarrow \mathcal{X}_{\text{Iw}^+, w}$, we obtain finite étale morphisms

$$\begin{array}{ccc} & \mathcal{X}_{\gamma, \Gamma(p^\infty), w} & \\ \text{pr}_{1, \infty} \swarrow & & \searrow \text{pr}_{2, \infty} \\ \mathcal{X}_{\Gamma(p^\infty), w} & & \mathcal{X}_{\Gamma(p^\infty), w} \end{array}$$

between perfectoid spaces and an $\text{Iw}_{\text{GSp}_{2g}}^+$ -equivariant isomorphism $\Phi_{\gamma, \infty} : \mathcal{X}_{\gamma, \Gamma(p^\infty), w} \xrightarrow{\sim} \mathcal{X}_{\gamma, \Gamma(p^\infty), w}$. The isomorphism $\Phi_{\gamma, \infty}$ induces an isomorphism

$$\Phi_{\gamma, \infty}^* : \text{pr}_{2, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \xrightarrow{\sim} \text{pr}_{1, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}}.$$

It then induces an isomorphism

$$\Phi_{\gamma, \infty}^* : \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \text{pr}_{2, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}}) \xrightarrow{\sim} \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \text{pr}_{1, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$$

by taking the identity on $R_{\mathcal{U}}$.

Recall that \mathfrak{z} is the pullback of the coordinate z via the Hodge–Tate period map $\pi_{\text{HT}} : \mathcal{X}_{\Gamma(p^\infty), w} \rightarrow \mathcal{F}\ell_w^\times$. Let $\mathfrak{z}' := \text{pr}_{1, \infty}^* \mathfrak{z}$ and $\mathfrak{z}'' := \text{pr}_{2, \infty}^* \mathfrak{z}$. Since $\Phi_{\gamma, \infty}$ induces an isomorphism on the p -adic Tate module, we have $\mathfrak{z}' = \mathfrak{z}''$. Consequently, a section f of $\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \text{pr}_{2, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ satisfies

$$\gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z}'' \gamma_c)^{-1} f$$

for all $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \text{Iw}_{\text{GSp}_{2g}}^+$, if and only if the section $\Phi_{\gamma, \infty}^*(f)$ of $\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \text{pr}_{1, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ satisfies

$$\gamma^* (\Phi_{\gamma, \infty}^*(f)) = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z}' \gamma_c)^{-1} (\Phi_{\gamma, \infty}^*(f))$$

for all $\gamma \in \text{Iw}_{\text{GSp}_{2g}}^+$. This yields the desired isomorphism

$$\varphi_\gamma : \text{pr}_2^* \underline{\omega}_w^{\kappa_{\mathcal{U}}} \xrightarrow{\sim} \text{pr}_1^* \underline{\omega}_w^{\kappa_{\mathcal{U}}}.$$

Given this, we consider the composition

$$T_\gamma : \quad \begin{array}{ccc} H^0(\mathcal{X}_{Iw^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) & \xrightarrow{\text{pr}_2^*} & H^0(\mathcal{X}_{\gamma, Iw^+,w}, \text{pr}_2^* \underline{\omega}_w^{\kappa_{\mathcal{U}}}) \\ & \searrow \varphi_\gamma & \\ & H^0(\mathcal{X}_{\gamma, Iw^+,w}, \text{pr}_1^* \underline{\omega}_w^{\kappa_{\mathcal{U}}}) & \xrightarrow{\text{Tr pr}_1} H^0(\mathcal{X}_{Iw^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}). \end{array}$$

Finally, we have to extend the construction to the boundary. In fact, we shall prove that the sections of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ on $\overline{\mathcal{X}}_{Iw^+,w}$ are precisely the *bounded* sections of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ over the open part $\mathcal{X}_{Iw^+,w}$. By definition, a section of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ on $\overline{\mathcal{X}}_{Iw^+,w}$ can be viewed a section of $\mathcal{O}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}} \widehat{\otimes} R_{\mathcal{U}})$, which is a sheaf of uniform \mathbf{C}_p -Banach algebra by Definition 3.1.14 (ii). In particular, the notion of bounded section of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ on $\overline{\mathcal{X}}_{Iw^+,w}$ is well-defined.

Lemma 3.2.1. (i) Every bounded section of $\mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty),w}}$ uniquely extends to a section of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}}$.

(ii) Every bounded section of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ on $\mathcal{X}_{Iw^+,w}$ uniquely extends to a section of $\overline{\mathcal{X}}_{Iw^+,w}$.

Proof. (i) The assertion follows from that every bounded section of $\mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty)}}$ uniquely extends to a section of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}$. To show this, let $\mathcal{X}_{\Gamma(p^\infty)}^*$ be the minimal compactification of $\mathcal{X}_{\Gamma(p^\infty)}$ considered in [Sch15] and let

$$\psi_{\Gamma(p^\infty)} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \mathcal{X}_{\Gamma(p^\infty)}^*$$

be the natural map. By the proof of [PS16, Proposition 1.11], we know that $\psi_{\Gamma(p^\infty)}$ is a surjection and the natural map

$$\mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty)}^*} \rightarrow \psi_{\Gamma(p^\infty),*} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}$$

is an injection. Hence, it is enough to show that every bounded section of $\mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty)}}$ extends to a section in $\mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty)}^*}$. However, this follows from the fact that $\mathcal{X}_{\Gamma(p^\infty)}^*$ is covered by translates of the anticanonical locus ([Sch15, Lemma 3.3.11]) and the assertion holds for the anticanonical locus ([op. cit., Theorem 3.2.36]).

(ii) For simplicity, let $m = \frac{g(g-1)}{2}$. According to Remark 3.1.7 and Remark 3.1.11, every section f in $\mathcal{O}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}} \widehat{\otimes} R_{\mathcal{U}})$ is given by an $p^{m[w]}$ -tuple $\{f_a\}_{a \in A}$ where each

$$f_a \in \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}} \widehat{\otimes} R_{\mathcal{U}} \langle \frac{T_1}{p^w}, \dots, \frac{T_m}{p^w} \rangle.$$

By choosing a pseudo-basis $\{e_\sigma : \sigma \in \Sigma\}$ of $R_{\mathcal{U}}$ as in [CHJ17, Proposition 6.2], we can identify $R_{\mathcal{U}} \simeq \prod_{\sigma \in \Sigma} \mathbf{Z}_p e_\sigma$, equipped with the product topology. Under this identification, each f_a can be viewed as an element

$$f_a = \{f_{a,\sigma}\}_{\sigma \in \Sigma} \in \prod_{\sigma \in \Sigma} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}} \langle \frac{T_1}{p^w}, \dots, \frac{T_m}{p^w} \rangle.$$

If we express each $f_{a,\sigma}$ in terms of power series expansion

$$f_{a,\sigma} = \sum_{\nu \in \mathbf{Z}_{\geq 0}^m} g_{a,\sigma,\nu} T^\nu$$

with $g_{a,\sigma,\nu} \in \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty),w}}$, it is clear that f is bounded if and only if each $g_{a,\sigma,\nu}$ is bounded. In this case, by (i), each $g_{a,\sigma,\nu}$ extends uniquely to a section of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}}$. One checks that the resulting extension

of each f_a still satisfies the desired convergence condition. Hence, $f = \{f_a\}_{a \in A}$ extends to a section of $\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IwGL}_g, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$. Finally, by continuity, f satisfies the condition

$$\gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f$$

for all $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \text{Iw}_{\text{GSp}_{2g}}^+$, and hence defines an element of $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ on $\overline{\mathcal{X}}_{\text{Iw}^+, w}$.

□

Thanks to Lemma 3.2.1 (ii), and observe that

$$\Phi_{\gamma, \infty}^* : \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IwGL}_g, \text{pr}_{2, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}}) \xrightarrow{\sim} \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IwGL}_g, \text{pr}_{1, \infty}^* \mathcal{O}_{\mathcal{X}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$$

sends bounded sections to bounded sections, we know that T_γ extends to the boundary. We arrive at the Hecke operator

$$T_\gamma : M_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}}} = H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) = M_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}}}.$$

Hecke operators at p . For $1 \leq i \leq g$, we consider matrices $\mathbf{u}_{p, i} \in \text{GSp}_{2g}(\mathbf{Q}_p) \cap M_{2g}(\mathbf{Z}_p)$ defined by

$$\mathbf{u}_{p, i} := \begin{pmatrix} \mathbb{1}_i & & & & \\ & p \mathbb{1}_{g-i} & & & \\ & & p \mathbb{1}_{g-i} & & \\ & & & p^2 \mathbb{1}_i & \\ & & & & \end{pmatrix}$$

for $1 \leq i \leq g-1$, and

$$\mathbf{u}_{p, g} := \begin{pmatrix} \mathbb{1}_g & \\ & p \mathbb{1}_g \end{pmatrix}.$$

For later use, we write

$$\mathbf{u}_{p, i} = \begin{pmatrix} \mathbf{u}_{p, i}^\square & \\ & \mathbf{u}_{p, i}^\blacksquare \end{pmatrix}$$

where $\mathbf{u}_{p, i}^\square$ and $\mathbf{u}_{p, i}^\blacksquare$ are the corresponding $g \times g$ diagonal matrices.

Notice that the $\mathbf{u}_{p, i}$ -action on $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ preserves $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$. This can be checked at the infinite level via local coordinates; *i.e.*, the action of $\mathbf{u}_{p, i}$ on \mathbf{z} is given by

$$\mathbf{z} \cdot \mathbf{u}_{p, i} = \mathbf{u}_{p, i}^{\square, -1} \mathbf{z} \mathbf{u}_{p, i}^\blacksquare = \begin{cases} \begin{pmatrix} p \mathbf{z}_{1,1} & \cdots & p \mathbf{z}_{1, g-i} & p^2 \mathbf{z}_{1, g+1-i} & \cdots & p^2 \mathbf{z}_{1, g} \\ \vdots & & \vdots & \vdots & & \vdots \\ p \mathbf{z}_{i,1} & \cdots & p \mathbf{z}_{i, g-i} & p^2 \mathbf{z}_{i, g+1-i} & \cdots & p^2 \mathbf{z}_{i, g} \\ \mathbf{z}_{i+1,1} & \cdots & \mathbf{z}_{i+1, g-i} & p \mathbf{z}_{i+1, g+1-i} & \cdots & p \mathbf{z}_{i+1, g} \\ \vdots & & \vdots & \vdots & & \vdots \\ \mathbf{z}_{g,1} & \cdots & \mathbf{z}_{g, g-i} & p \mathbf{z}_{g, g+1-i} & \cdots & p \mathbf{z}_{g, g} \end{pmatrix}, & \text{if } i = 1, \dots, g-1 \\ p \mathbf{z}, & \text{if } i = g \end{cases}.$$

In particular, when $i = g$, the $\mathbf{u}_{p, g}$ -action actually sends $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$ into $\overline{\mathcal{X}}_{\Gamma(p^\infty), w+1}$.

Recall the twisted left action of $\text{Iw}_{\text{GSp}_{2g}}^+$ on $\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IwGL}_g, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$, given by the formula

$$\gamma \cdot f := \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c) \gamma^* f.$$

Definition 3.2.2. (i) For $f \in \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$, we define $\mathbf{u}_{p,i} \cdot f \in \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ by

$$\mathbf{u}_{p,i} \cdot f(\gamma') := \mathbf{u}_{p,i}^* f(\mathbf{u}_{p,i}^\square \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0)$$

where $\gamma' = \gamma'_0 \beta'_0 \in \text{Iw}_{\text{GL}_g}$ with $\gamma'_0 \in U_{\text{GL}_g, 1}^{\text{opp}}$ and $\beta'_0 \in B_{\text{GL}_g, 0}$.

(ii) Suppose $f \in \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ satisfies

$$\gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f$$

for all $\gamma \in \text{Iw}_{\text{GSp}_{2g}}^+$; i.e., $\gamma \cdot f = f$. Pick a decomposition of the double coset

$$\text{Iw}_{\text{GSp}_{2g}}^+ \mathbf{u}_{p,i} \text{Iw}_{\text{GSp}_{2g}}^+ = \bigsqcup_{j=1}^m \delta_{ij} \mathbf{u}_{p,i} \text{Iw}_{\text{GSp}_{2g}}^+$$

with $\delta_{i,j} \in \text{Iw}_{\text{GSp}_{2g}}^+$. Define

$$U_{p,i}(f) := \sum_{j=1}^m \delta_{i,j} \cdot (\mathbf{u}_{p,i} \cdot f) \in \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}}).$$

Of course, we have to verify that $U_{p,i}(f)$ is independent to the choice of the representatives δ_{ij} 's. Suppose $\{\delta'_{i,j}\}_{j=1}^m$ is another set of representatives. Up to re-labelling, we may assume that

$$\delta'_{ij} \mathbf{u}_{p,i} \text{Iw}_{\text{GSp}_{2g}}^+ = \delta_{ij} \mathbf{u}_{p,i} \text{Iw}_{\text{GSp}_{2g}}^+$$

for every $j = 1, \dots, m$. Write $\delta'_{ij} \mathbf{u}_{p,i} = \delta_{ij} \mathbf{u}_{p,i} \gamma_j$ for some $\gamma_j \in \text{Iw}_{\text{GSp}_{2g}}^+$. We have to check that $\delta_{ij} \cdot (\mathbf{u}_{p,i} \cdot f) = \delta'_{ij} \cdot (\mathbf{u}_{p,i} \cdot f)$. Indeed, if we write $\delta_{ij} = \begin{pmatrix} \delta_{ija} & \delta_{ijb} \\ \delta_{ijc} & \delta_{ijd} \end{pmatrix}$, $\delta'_{ij} = \begin{pmatrix} \delta'_{ija} & \delta'_{ijb} \\ \delta'_{ijc} & \delta'_{ijd} \end{pmatrix}$, and $\gamma_j = \begin{pmatrix} \gamma_{ja} & \gamma_{jb} \\ \gamma_{jc} & \gamma_{jd} \end{pmatrix}$, then for every $\gamma' \in \text{Iw}_{\text{GL}_g}$, we have

$$\begin{aligned} \delta'_{ij} \cdot (\mathbf{u}_{p,i} \cdot f)(\gamma') &= \rho_{\kappa_{\mathcal{U}}}(\delta'_{ija} + \mathfrak{z} \delta'_{ijc}) \delta_{ij}^* (\mathbf{u}_{p,i}^* f)(\mathbf{u}_{p,i}^\square \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= (\delta'_{ij} \mathbf{u}_{p,i})^* f(\mathbf{u}_{p,i}^\square {}^t(\delta'_{ija} + \mathfrak{z} \delta'_{ijc}) \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= (\delta'_{ij} \mathbf{u}_{p,i})^* f({}^t(\gamma_{ja} + \mathfrak{z} \gamma_{jc}) \mathbf{u}_{p,i}^\square {}^t(\delta_{ija} + \mathfrak{z} \delta_{ijc}) \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= (\delta'_{ij} \mathbf{u}_{p,i})^* (\rho_{\kappa_{\mathcal{U}}}(\gamma_{ja} + \mathfrak{z} \gamma_{jc}) f)(\mathbf{u}_{p,i}^\square {}^t(\delta_{ija} + \mathfrak{z} \delta_{ijc}) \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= (\delta'_{ij} \mathbf{u}_{p,i})^* (\gamma_j^{-1, *} f)(\mathbf{u}_{p,i}^\square {}^t(\delta_{ija} + \mathfrak{z} \delta_{ijc}) \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= (\delta_{ij} \mathbf{u}_{p,i})^* f(\mathbf{u}_{p,i}^\square {}^t(\delta_{ija} + \mathfrak{z} \delta_{ijc}) \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= \rho_{\kappa_{\mathcal{U}}}(\delta_{ija} + \mathfrak{z} \delta_{ijc}) (\delta_{ij} \mathbf{u}_{p,i})^* f(\mathbf{u}_{p,i}^\square \gamma'_0 \mathbf{u}_{p,i}^{\square, -1} \beta'_0) \\ &= \delta_{ij} \cdot (\mathbf{u}_{p,i} \cdot f)(\gamma') \end{aligned}$$

as desired. Here, the third equality follows from the identity

$$\mathbf{u}_{p,i}^\square {}^t(\delta'_{ija} + \mathfrak{z} \delta'_{ijc}) = {}^t(\gamma_{ja} + \mathfrak{z} \gamma_{jc}) \mathbf{u}_{p,i}^\square {}^t(\delta_{ija} + \mathfrak{z} \delta_{ijc}).$$

Lemma 3.2.3. Suppose $f \in \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ such that $\gamma \cdot f = f$. Then, the section $U_{p,i}(f) \in \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}} \widehat{\otimes} R_{\mathcal{U}})$ satisfies $\gamma \cdot (U_{p,i}(f)) = U_{p,i}(f)$ for all $\gamma \in \text{Iw}_{\text{GSp}_{2g}}^+$.

Proof. We have

$$\gamma \cdot (U_{p,i}(f)) = \sum_{j=1}^m \gamma \cdot (\delta_{ij} \cdot (\mathbf{u}_{p,i}(f))) = \sum_{j=1}^m (\gamma \delta_{ij}) \cdot (\mathbf{u}_{p,i}(f)).$$

The last term indeed computes $U_{p,i}(f)$ because $\{\gamma \delta_{ij} : 1 \leq j \leq m\}$ is also a valid set of representatives. \square

Consequently, we arrive at the Hecke operator

$$U_{p,i} : M_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}} = H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) = M_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}}.$$

Definition 3.2.4. The *Hecke algebra outside Np* is defined to be

$$\mathbb{T}^{Np} := \mathbf{Z}_p [T_{\gamma}; \gamma \in \mathrm{GSp}_{2g}(\mathbf{Q}_{\ell}) \cap M_{2g}(\mathbf{Z}_{\ell}), \ell \nmid Np]$$

and the *total Hecke algebra* is defined to be

$$\mathbb{T} := \mathbb{T}^{Np} \otimes_{\mathbf{Z}_p} \mathbf{Z}_p[U_{p,i}; i = 1, \dots, g].$$

We now define $U_p := \prod_{i=1}^g U_{p,i}$ and conclude with the following proposition

Proposition 3.2.5. The operator U_p is a compact operator on $M_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}}$.

Proof. Note that the action of $\mathbf{u}_{p,g}$ on \mathbf{z} is given by $p\mathbf{z}$ and that, by definition, the action of $\prod_{i=1}^g \mathbf{u}_{p,i}$ on $\mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^{\infty}),w}} \widehat{\otimes} R_{\mathcal{U}})$ factors through the inclusion

$$\mathcal{C}_{\kappa_{\mathcal{U}}}^{(w-1)-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^{\infty}),w}} \widehat{\otimes} R_{\mathcal{U}}) \hookrightarrow \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^{\infty}),w}} \widehat{\otimes} R_{\mathcal{U}}).$$

This means that U_p factors as

$$U_p : H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+,w+1}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+,w+1}, \underline{\omega}_{w-1}^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}),$$

where the first arrow is the natural restriction map.

To show the desired result, note that it is known that restrictions of the structure sheaf of $\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}$ are compact operators. Moreover, by the discussion in [Han17, §2.2], the injection of $(w-1)$ -analytic functions into w -analytic functions is compact. The assertion then follows by combining these two facts. \square

Remark 3.2.6. Note that the subspace $S_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}} \subset M_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}}$ of w -overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ is stable under the action of \mathbb{T} . Moreover, as U_p is a compact operator on $M_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}}$, it is also a compact operator on $S_{\mathrm{Iw}^+,w}^{\kappa_{\mathcal{U}}}$.

3.3 Admissibility

Throughout this subsection, let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a small weight and $w > 1 + r_{\mathcal{U}}$. We fix an ideal $\mathfrak{a}_{\mathcal{U}} \subset R_{\mathcal{U}}$ defining the profinite adic topology on $R_{\mathcal{U}}$. In addition, we assume $p \in \mathfrak{a}_{\mathcal{U}}$.

The purpose of this subsection is to show that the overconvergent Siegel modular sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ can be identified with the G -invariants of an *admissible Kummer étale Banach sheaf* in the sense of Definition A.3.9, where G is a finite group. Such a description allows us to apply Corollary A.3.14 to the sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$. This will be used in §5.2.

Firstly, we introduce the notion of w -compatibility inspired by [AIP15, §4.5].

Definition 3.3.1. Let R be a flat $\mathcal{O}_{\mathbb{C}_p}$ -algebra and suppose M is a free R -module of rank g . We write $R_w := R \otimes_{\mathcal{O}_{\mathbb{C}_p}} \mathcal{O}_{\mathbb{C}_p}/p^w$ and $M_w := M \otimes_R R_w$. Let $\underline{\mathbf{m}} := (\mathbf{m}_1, \dots, \mathbf{m}_g)$ be an R_w -basis for M_w . We denote by $\text{Fil}_{\bullet}^{\underline{\mathbf{m}}}$ the full flag

$$0 \subset \langle \mathbf{m}_1 \rangle \subset \langle \mathbf{m}_1, \mathbf{m}_2 \rangle \subset \dots \subset \langle \mathbf{m}_1, \dots, \mathbf{m}_g \rangle$$

of the free R_w -module M_w . Namely, $\text{Fil}_i^{\underline{\mathbf{m}}} = \langle \mathbf{m}_1, \dots, \mathbf{m}_i \rangle$ for all $i = 1, \dots, g$.

(i) A full flag Fil_{\bullet} of the free R -module M is called *w-compatible with $\underline{\mathbf{m}}$* if

$$\text{Fil}_i \otimes_R R_w = \text{Fil}_i^{\underline{\mathbf{m}}}$$

for all $i = 1, \dots, g$.

(ii) Suppose Fil_{\bullet} is a *w-compatible full flag* as in (i). Consider a collection $\{w_i : i = 1, \dots, g\}$ where each w_i is an R -basis for $\text{Fil}_i / \text{Fil}_{i-1}$. Then $\{w_i : i = 1, \dots, g\}$ is called *w-compatible with $\underline{\mathbf{m}}$* if

$$w_i \mod (p^w M + \text{Fil}_{i-1}) = \mathbf{m}_i \mod \text{Fil}_{i-1}^{\underline{\mathbf{m}}}$$

for all $i = 1, \dots, g$.

Pick a positive integer $n > \sup\{w, \frac{g}{p-1}\}$. Recall from §B.2 the locally free $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^+$ -module $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}$ over $\overline{\mathcal{X}}_{\Gamma(p^n)}$. Also recall the Hodge–Tate map

$$\text{HT}_{\Gamma(p^n)} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \underline{\omega}_{\Gamma(p^n)}^{\text{mod},+} / p^n \underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}$$

over $\overline{\mathcal{X}}_{\Gamma(p^n)}$. Restricting to the w -ordinary locus $\overline{\mathcal{X}}_{\Gamma(p^n),w}$ and composing with a natural projection, we obtain

$$\text{HT}_{\Gamma(p^n),w} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+} / p^n \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+} \twoheadrightarrow \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+} / p^w \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+}$$

where $\underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+}$ is the restriction of $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}$ on $\overline{\mathcal{X}}_{\Gamma(p^n),w}$.

Lemma 3.3.2. The sheaf $\underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+} / p^w \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+}$ is a free $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w}}^+ / p^w$ -module of rank g generated by the basis $\text{HT}_{\Gamma(p^n),w}(e_{g+1}), \dots, \text{HT}_{\Gamma(p^n),w}(e_{2g})$.

Proof. Notice that $\underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+} / p^w \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+}$ is locally free of rank g . It follows from the definition of w -ordinary locus that $\text{HT}_{\Gamma(p^n),w}(e_{g+1}), \dots, \text{HT}_{\Gamma(p^n),w}(e_{2g})$ span $\underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+} / p^w \underline{\omega}_{\Gamma(p^n),w}^{\text{mod},+}$. Hence they must form a set of free generators. \square

We consider an adic space \mathcal{IW}_w^+ over $\overline{\mathcal{X}}_{\Gamma(p^n),w}$ parameterising certain w -compatible objects. More precisely, for every affinoid open subset $\text{Spa}(R, R^+) \subset \overline{\mathcal{X}}_{\Gamma(p^n),w}$, the set $\mathcal{IW}_w^+(R, R^+)$ consists of triples

$$(\sigma, \text{Fil}_{\bullet}, \{w_i : i = 1, \dots, g\})$$

where

- (i) σ is a matrix in $\text{Iw}_{\text{GL}_g}^+(\mathbf{Z}/p^n \mathbf{Z})$ where $\text{Iw}_{\text{GL}_g}^+(\mathbf{Z}/p^n \mathbf{Z})$ is the preimage of $T_{\text{GL}_g}(\mathbf{Z}/p \mathbf{Z})$ under the surjection $\text{GL}_g(\mathbf{Z}/p^n \mathbf{Z}) \xrightarrow{\text{mod } p} \text{GL}_g(\mathbf{Z}/p \mathbf{Z})$;
- (ii) both Fil_{\bullet} and $\{w_i : i = 1, \dots, g\}$ are w -compatible with $(\text{HT}_{\Gamma(p^n),w}(e_{g+1}), \dots, \text{HT}_{\Gamma(p^n),w}(e_{2g})) \cdot \sigma$.

Let $\pi : \mathcal{IW}_w^+ \rightarrow \overline{\mathcal{X}}_{\Gamma(p^n),w}$ denote the natural projection. To further understand \mathcal{IW}_w^+ , we consider the following group objects in adic spaces.

Definition 3.3.3. Let $\mathbf{B}(0,1) = \mathrm{Spa}(\mathbf{C}_p\langle X \rangle, \mathcal{O}_{\mathbf{C}_p}\langle X \rangle)$ be the closed unit ball.

(i) Define

$$\mathcal{B}_w^{\mathrm{opp}} = \begin{pmatrix} 1 + p^w \mathbf{B}(0,1) & & & \\ p^w \mathbf{B}(0,1) & 1 + p^w \mathbf{B}(0,1) & & \\ \vdots & \vdots & \ddots & \\ p^w \mathbf{B}(0,1) & p^w \mathbf{B}(0,1) & \cdots & 1 + p^w \mathbf{B}(0,1) \end{pmatrix}$$

In particular, the underlying adic space is isomorphic to a $\frac{1}{2}g(g+1)$ -dimension ball of radius p^{-w} .

(ii) Define

$$\begin{aligned} \mathcal{T}_{\mathrm{GL}_g,0}^{(w)} &= \begin{pmatrix} \mathbf{Z}_p^\times + p^w \mathbf{B}(0,1) & & & \\ & \mathbf{Z}_p^\times + p^w \mathbf{B}(0,1) & & \\ & & \ddots & \\ & & & \mathbf{Z}_p^\times + p^w \mathbf{B}(0,1) \end{pmatrix} \\ &= \bigsqcup_{h_1, \dots, h_g \in S} \begin{pmatrix} h_1 + p^w \mathbf{B}(0,1) & & & \\ & h_2 + p^w \mathbf{B}(0,1) & & \\ & & \ddots & \\ & & & h_g + p^w \mathbf{B}(0,1) \end{pmatrix} \end{aligned}$$

where $S \subset \mathbf{Z}_p^\times$ is a set of representatives of $\mathbf{Z}_p^\times / (1 + p^n \mathbf{Z}_p)$. In particular, the underlying adic space is isomorphic to the disjoint union of $(p-1)^g p^{g(n-1)}$ copies of g -dimension ball of radius p^{-w} .

(iii) Define

$$\begin{aligned} \mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} &= \begin{pmatrix} 1 & & & \\ p \mathbf{Z}_p + p^w \mathbf{B}(0,1) & 1 & & \\ \vdots & \vdots & \ddots & \\ p \mathbf{Z}_p + p^w \mathbf{B}(0,1) & p \mathbf{Z}_p + p^w \mathbf{B}(0,1) & \cdots & 1 \end{pmatrix} \\ &= \bigsqcup_{\substack{h_{i,j} \in S \\ 1 \leq j < i \leq g}} \begin{pmatrix} 1 & & & \\ h_{2,1} + p^w \mathbf{B}(0,1) & 1 & & \\ \vdots & \vdots & \ddots & \\ h_{g,1} + p^w \mathbf{B}(0,1) & h_{g,2} + p^w \mathbf{B}(0,1) & \cdots & 1 \end{pmatrix} \end{aligned}$$

where $S \subset \mathbf{Z}_p^\times$ is a set of representatives of $p \mathbf{Z}_p / p^n \mathbf{Z}_p$. In particular, the underlying adic space is isomorphic to the disjoint union of $p^{\frac{1}{2}g(g-1)(n-1)}$ copies of $\frac{1}{2}g(g-1)$ -dimension ball of radius p^{-w} .

All of the group structures are given by the matrix multiplications.

Remark 3.3.4. The $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ -points of $\mathcal{B}_w^{\mathrm{opp}}$, $\mathcal{T}_{\mathrm{GL}_g,0}^{(w)}$, and $\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)}$ coincide with the groups B_w^{opp} , $T_{\mathrm{GL}_g,0}^{(w)}$, and $U_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)}$, respectively. This justifies the notations.

Lemma 3.3.5. \mathcal{IW}_w^+ is a $\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times \mathcal{T}_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z})$ -torsor over $\overline{\mathcal{X}}_{\Gamma(p^n),w}$ where $U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z})$ is the kernel of the natural surjection $U_{\mathrm{GL}_g}(\mathbf{Z}/p^n \mathbf{Z}) \xrightarrow{\mathrm{mod } p} U_{\mathrm{GL}_g}(\mathbf{Z}/p \mathbf{Z})$. Namely, locally on $\overline{\mathcal{X}}_{\Gamma(p^n),w}$, we have an identification

$$\mathcal{IW}_w^+ \simeq \overline{\mathcal{X}}_{\Gamma(p^n),w} \times_{\mathrm{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})} \left(\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times \mathcal{T}_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z}) \right)$$

where $\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times \mathcal{T}_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z})$ acts from the right by matrix multiplication.

Proof. This is clear from the construction. \square

Using the adic space \mathcal{IW}_w^+ , we construct two auxiliary sheaves $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}$ and $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$.

Definition 3.3.6. (i) The sheaf $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}$ over $\overline{\mathcal{X}}_{\Gamma(p^n),w}$ is defined to be

$$\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+} := (\pi_* \mathcal{O}_{\mathcal{IW}_w^+}^+ \widehat{\otimes} R_{\mathcal{U}}) [\kappa_{\mathcal{U}}^\vee];$$

i.e., the subsheaf of $\pi_* \mathcal{O}_{\mathcal{IW}_w^+}^+ \widehat{\otimes} R_{\mathcal{U}}$ consisting of those sections on which $T_{\mathrm{GL}_g,0}$ -acts through the character $\kappa_{\mathcal{U}}^\vee$ and $U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z})$ acts trivially.

(ii) The sheaf

$$\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}} := (\pi_* \mathcal{O}_{\mathcal{IW}_w^+} \widehat{\otimes} R_{\mathcal{U}}) [\kappa_{\mathcal{U}}^\vee]$$

is defined similarly.

Remark 3.3.7. Since $\kappa_{\mathcal{U}}$ is w -analytic, the character $\kappa_{\mathcal{U}}^\vee : T_{\mathrm{GL}_g,0} \rightarrow R_{\mathcal{U}}^\times$ extends to a character on $\mathcal{T}_{\mathrm{GL}_g,0}^{(w)}$; namely, a character

$$\kappa_{\mathcal{U}}^\vee : \mathcal{T}_{\mathrm{GL}_g,0}^{(w)}(R, R^+) \rightarrow (R^+ \widehat{\otimes} R_{\mathcal{U}})^\times$$

for every affinoid $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ -algebra (R, R^+) . It turns out, in Definition 3.3.6, there is no difference between taking $\kappa_{\mathcal{U}}^\vee$ -eigenspaces with respect to $T_{\mathrm{GL}_g,0}$ - or $\mathcal{T}_{\mathrm{GL}_g,0}^{(w)}$ -actions.

Lemma 3.3.8. The sheaf $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$ is a projective Banach sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w}} \widehat{\otimes} R_{\mathcal{U}}$ -modules in the sense of Definition A.3.3 (ii). Moreover, $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}$ is an integral model of $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$ in the sense of Definition A.3.3 (iv).

Proof. Let $\{\mathcal{V}_{n,i} : i \in I\}$ be an affinoid open covering of $\overline{\mathcal{X}}_{\Gamma(p^n),w}$ such that $\underline{\omega}_{\Gamma(p^n)}^{\mathrm{mod},+}|_{\mathcal{V}_{n,i}}$ is free, for every $i \in I$. By choosing a basis for $\underline{\omega}_{\Gamma(p^n)}^{\mathrm{mod},+}|_{\mathcal{V}_{n,i}}$, we can identify

$$\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}|_{\mathcal{V}_{n,i}} \simeq \mathcal{O}_{\mathcal{V}_{n,i}}^+ \langle T_{st} : 1 \leq s < t \leq g \rangle \widehat{\otimes} R_{\mathcal{U}}$$

which is the p -adic completion of a free $\mathcal{O}_{\mathcal{V}_{n,i}}^+ \widehat{\otimes} R_{\mathcal{U}}$ -module, as desired. \square

We also consider the associated p -adically completed sheaves on the Kummer étale site.

Definition 3.3.9. Let

$$\underline{\omega}_{n,w,k\acute{e}t}^{\kappa_{\mathcal{U}},+} := \varprojlim_m \left(\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w}}^+} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w,k\acute{e}t}}^+ / p^m \right)$$

and let

$$\underline{\omega}_{n,w,k\acute{e}t}^{\kappa_{\mathcal{U}}} := \underline{\omega}_{n,w,k\acute{e}t}^{\kappa_{\mathcal{U}},+} \left[\frac{1}{p} \right].$$

By Lemma 3.3.8 and Corollary A.3.6, $\underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U}}}$ is a projective Kummer étale Banach sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w,\text{két}}} \widehat{\otimes} R_{\mathcal{U}}$ -modules in the sense of Definition A.3.4 (ii). Moreover, $\underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U},+}$ is an integral model of $\underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U}}}$ in the sense of Definition A.3.4 (iv).

In fact, we show that the Kummer étale Banach sheaf $\underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U}}}$ is *admissible*.

Lemma 3.3.10. *The sheaf $\underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U}}}$ is an admissible Kummer étale Banach sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w,\text{két}}} \widehat{\otimes} R_{\mathcal{U}}$ -modules (in the sense of Definition A.3.9) with integral model $\underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U},+}$.*

Proof. The proof is inspired by the discussion in [AIP15, §8.1]. We provide a sketch of proof.

To simplify the notation, we write $\mathcal{F}^+ = \underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U},+}$ and $\mathcal{F} = \underline{\omega}_{n,w,\text{két}}^{\kappa_{\mathcal{U}}}$. We also write $\mathcal{F}_m^+ := \mathcal{F}^+ / \mathfrak{a}_{\mathcal{U}}^m$, for every $m \in \mathbf{Z}_{\geq 1}$.

Let $\mathcal{U} = \{\mathcal{V}_{n,i} : i \in I\}$ be an open affinoid covering for $\overline{\mathcal{X}}_{\Gamma(p^n),w}$ such that $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}|_{\mathcal{V}_{n,i}}$ is free, for every $i \in I$. We equip each $\mathcal{V}_{n,i}$ the induced log structure from $\overline{\mathcal{X}}_{\Gamma(p^n),w}$. By choosing a basis for $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}|_{\mathcal{V}_{n,i}}$, we can identify

$$\mathcal{F}^+|_{\mathcal{V}_{n,i}} \simeq \mathcal{O}_{\mathcal{V}_{n,i}}^+ \langle T_{st} : 1 \leq s < t \leq g \rangle \widehat{\otimes} R_{\mathcal{U}}$$

which is the p -adic completion of a free $\mathcal{O}_{\mathcal{V}_{n,i}}^+ \widehat{\otimes} R_{\mathcal{U}}$ -module. Modulo $\mathfrak{a}_{\mathcal{U}}^m$, we obtain

$$\mathcal{F}_m^+|_{\mathcal{V}_{n,i}} \simeq (\mathcal{O}_{\mathcal{V}_{n,i}}^+ \otimes_{\mathbf{Z}_p} (R_{\mathcal{U}} / \mathfrak{a}_{\mathcal{U}}^m)) [T_{st} : 1 \leq s < t \leq g].$$

For any $d \in \mathbf{Z}_{\geq 0}$, consider the subsheaf $(\mathcal{F}_m^+|_{\mathcal{V}_{n,i}})^{\leq d} \subset \mathcal{F}_m^+|_{\mathcal{V}_{n,i}}$ consisting of those polynomials of degree $\leq d$, and consider

$$\mathcal{F}_{m,d}^+ := \ker \left(\prod_{i \in I} (\mathcal{F}_m^+|_{\mathcal{V}_{n,i}})^{\leq d} \rightarrow \prod_{i,j \in I} \mathcal{F}_m^+|_{\mathcal{V}_{n,i} \cap \mathcal{V}_{n,j}} \right).$$

Then each $\mathcal{F}_{m,d}^+$ is a coherent $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w,\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R_{\mathcal{U}} / \mathfrak{a}_{\mathcal{U}}^m)$ -module and we have $\mathcal{F}_m^+ = \varinjlim_d \mathcal{F}_{m,d}^+$, as desired. \square

Next, we are going to relate the overconvergent Siegel modular sheaves $\underline{\omega}_w^{\kappa_{\mathcal{U},+}$ and $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ with the auxiliary sheaves $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U},+}$ and $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$. To this end, we need two intermediate sheaves $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U},+}$ and $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$ over $\overline{\mathcal{X}}_{\Gamma(p^n),w}$.

Definition 3.3.11. *Let $h_{\Gamma(p^n)} : \overline{\mathcal{X}}_{\Gamma(p^\infty),w} \rightarrow \overline{\mathcal{X}}_{\Gamma(p^n),w}$ be the natural projection.*

(i) *The subsheaf $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$ of $h_{\Gamma(p^n),*} \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IW}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}} \widehat{\otimes} R_{\mathcal{U}})$ is defined as follows. For every affinoid open subset $\mathcal{V} \subset \overline{\mathcal{X}}_{\Gamma(p^n),w}$ with $\mathcal{V}_\infty = h_{\Gamma(p^n)}^{-1}(\mathcal{V})$, we put*

$$\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}(\mathcal{V}) := \left\{ f \in C_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IW}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}}(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}) : \gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f, \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n) \right\}.$$

(ii) *The subsheaf $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U},+}$ of $h_{\Gamma(p^n),*} \mathcal{C}_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IW}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}}^+ \widehat{\otimes} R_{\mathcal{U}})$ is defined as follows. For every affinoid open subset $\mathcal{V} \subset \overline{\mathcal{X}}_{\Gamma(p^n),w}$ with $\mathcal{V}_\infty = h_{\Gamma(p^n)}^{-1}(\mathcal{V})$, we put*

$$\underline{\omega}_{n,w}^{\kappa_{\mathcal{U},+}}(\mathcal{V}) := \left\{ f \in C_{\kappa_{\mathcal{U}}}^{w-\text{an}}(\text{IW}_{\text{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty),w}}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}) : \gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f, \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n) \right\}.$$

Remark 3.3.12. Let $h_n : \overline{\mathcal{X}}_{\Gamma(p^n),w} \rightarrow \overline{\mathcal{X}}_{\text{Iw}^+,w}$ denote the natural projection. Then the overconvergent Siegel modular sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ can be identified as the $\text{Iw}_{\text{GSp}_{2g}}^+ / \Gamma(p^n)$ -invariants of the sheaf $h_{n,*} \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$ with respect to the “twisted” action $\gamma \cdot f := \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c) \gamma^* f$ for every $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n)$ and $f \in \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$. Similar result holds for the integral sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U},+}$.

Proposition 3.3.13. *There is a natural isomorphism of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w}}^+ \widehat{\otimes} R_{\mathcal{U}}$ -modules $\Psi^+ : \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+} \simeq \widetilde{\underline{\omega}}_{n,w}^{\kappa_{\mathcal{U}},+}$. Inverting p , we obtain a natural isomorphism of $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n),w}} \widehat{\otimes} R_{\mathcal{U}}$ -modules $\Psi : \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}}} \simeq \widetilde{\underline{\omega}}_{n,w}^{\kappa_{\mathcal{U}}}$.*

Proof. As a preparation, consider the pullback diagram

$$\begin{array}{ccc} \mathcal{IW}_{w,\infty}^+ & \longrightarrow & \mathcal{IW}_w^+ \\ \pi_\infty \downarrow & & \downarrow \pi \\ \overline{\mathcal{X}}_{\Gamma(p^\infty),w} & \xrightarrow{h_{\Gamma(p^n)}} & \overline{\mathcal{X}}_{\Gamma(p^n),w} \end{array}$$

in the category of adic spaces. To show the existence of such a pullback, it suffices to check this locally on $\overline{\mathcal{X}}_{\Gamma(p^n),w}$. Recall that \mathcal{IW}_w^+ is a $\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times \mathcal{T}_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n\mathbf{Z})$ -torsor over $\overline{\mathcal{X}}_{\Gamma(p^n),w}$, and $\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times \mathcal{T}_{\mathrm{GL}_g,0}^{(w)}$ is isomorphic to finitely many copies of $\mathbf{B}(0,1)^{\frac{g(g+1)}{2}}$. It remains to show that the fibre product $\overline{\mathcal{X}}_{\Gamma(p^\infty),w} \times_{\mathrm{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})} \mathbf{B}(0,1)^{\frac{g(g+1)}{2}}$ exists. Indeed, by [SW20, Proposition 6.3.3 (3)], such a fibre product exists and is a sousperfectoid space. In addition, we know that the pullback $\mathcal{IW}_{w,\infty}^+$ is likewise a $\mathcal{U}_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times \mathcal{T}_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n\mathbf{Z})$ -torsor over $\overline{\mathcal{X}}_{\Gamma(p^\infty),w}$.

For every affinoid open $\mathcal{V} \subset \overline{\mathcal{X}}_{\Gamma(p^n),w}$ and $\mathcal{V}_\infty := h_{\Gamma(p^n)}^{-1} \mathcal{V}$, the desired isomorphism Ψ^+ will be established via a sequence of isomorphisms

$$\Psi^+ : \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}(\mathcal{V}) \xrightarrow[\Psi_1]{\sim} \omega^{(1)} \xrightarrow[\Psi_2]{\sim} \omega^{(2)} \xrightarrow[\Psi_3]{\sim} \widetilde{\underline{\omega}}_{n,w}^{\kappa_{\mathcal{U}},+}(\mathcal{V}),$$

where

$$\omega^{(1)} := \left\{ f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}) : \gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z} \gamma_c^\dagger) f, \quad \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n) \right\}$$

and

$$\omega^{(2)} := \left\{ f \in \pi_{\infty,*} \mathcal{O}_{\mathcal{IW}_{w,\infty}^+}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}} : \begin{array}{l} \gamma^* f = f, \quad \tau^* f = \kappa_{\mathcal{U}}^\vee(\tau) f, \quad \nu^* f = f \\ \forall (\gamma, \tau, \nu) \in \Gamma(p^n) \times T_{\mathrm{GL}_g,0} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n\mathbf{Z}) \end{array} \right\}.$$

Here, for any $\delta \in M_g$, we write $\delta^\dagger := \check{\mathbb{I}}_g {}^t \delta \check{\mathbb{I}}_g$, which can be viewed as the “transpose with respect to the anti-diagonal”. Notice that $\mathfrak{z}^\dagger = \mathfrak{z}$.

Construction of Ψ_1 . Observe that there is an isomorphism

$$\Psi_1 : C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}) \rightarrow C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}})$$

defined by

$$\Psi_1(f)(\gamma') := f(\check{\mathbb{I}}_g {}^t \gamma'^{-1} \check{\mathbb{I}}_g)$$

for all $f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}})$ and $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$.

We claim that Ψ_1 induces an isomorphism $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}(\mathcal{V}) \simeq \omega^{(1)}$. It suffices to check that if $\gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f$ for every $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n)$, then $\gamma^*(\Psi_1(f)) = \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z} \gamma_c^\dagger) \Psi_1(f)$. Indeed, for any $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$, we

have

$$\begin{aligned}
\gamma^*(\Psi_1(f))(\gamma') &= \rho_{\kappa_{\mathcal{U}}}(\gamma_a + \mathfrak{z}\gamma_c)^{-1} f(\check{\mathbb{I}}_g \mathfrak{t} \gamma'^{-1} \check{\mathbb{I}}_g) \\
&= f(\mathfrak{t}(\gamma_a + \mathfrak{z}\gamma_c)^{-1} \check{\mathbb{I}}_g \mathfrak{t} \gamma'^{-1} \check{\mathbb{I}}_g) \\
&= f(\check{\mathbb{I}}_g \check{\mathbb{I}}_g \mathfrak{t}(\gamma_a + \mathfrak{z}\gamma_c)^{-1} \check{\mathbb{I}}_g \mathfrak{t} \gamma'^{-1} \check{\mathbb{I}}_g) \\
&= f(\check{\mathbb{I}}_g \mathfrak{t}(\check{\mathbb{I}}_g \gamma_a \check{\mathbb{I}}_g + \check{\mathbb{I}}_g \mathfrak{z} \check{\mathbb{I}}_g \check{\mathbb{I}}_g \gamma_c \check{\mathbb{I}}_g)^{-1} \mathfrak{t} \gamma'^{-1} \check{\mathbb{I}}_g) \\
&= f(\check{\mathbb{I}}_g \mathfrak{t}(\mathfrak{t}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger) \gamma')^{-1} \check{\mathbb{I}}_g) \\
&= \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger) \Psi_1(f)(\gamma').
\end{aligned}$$

Construction of Ψ_2 . To construct Ψ_2 , consider $\mathfrak{s}^\dagger = (\mathfrak{s}_g \ \cdots \ \mathfrak{s}_1) \in \underline{\omega}_{\Gamma(p^\infty)}(\mathcal{V}_\infty)^g$. Recall that $\mathfrak{s} = \begin{pmatrix} \mathfrak{s}_1 \\ \vdots \\ \mathfrak{s}_g \end{pmatrix}$

and thus

$$\mathfrak{s}^\dagger = \mathfrak{t} \mathfrak{s} \check{\mathbb{I}}_g.$$

Moreover, for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n)$, we have $\gamma^* \mathfrak{s} = \mathfrak{t}(\gamma_a + \mathfrak{z}\gamma_c) \mathfrak{s}$ by (4). Hence

$$\gamma^* \mathfrak{s}^\dagger = \mathfrak{t}(\gamma^* \mathfrak{s}) \check{\mathbb{I}}_g = \mathfrak{t}(\mathfrak{t}(\gamma_a + \mathfrak{z}\gamma_c) \mathfrak{s}) \check{\mathbb{I}}_g = \mathfrak{t} \mathfrak{s} \check{\mathbb{I}}_g \check{\mathbb{I}}_g (\gamma_a + \mathfrak{z}\gamma_c) \check{\mathbb{I}}_g = (\mathfrak{t} \mathfrak{s} \check{\mathbb{I}}_g) \mathfrak{t}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger) = \mathfrak{s}^\dagger \mathfrak{t}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger).$$

Let $\text{Fil}_\bullet^\dagger$ be the full flag of the free $\mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty)$ -module $\underline{\omega}_{\Gamma(p^\infty)}(\mathcal{V}_\infty)$ given by

$$\text{Fil}_\bullet^\dagger = 0 \subset \langle \mathfrak{s}_g \rangle \subset \langle \mathfrak{s}_g, \mathfrak{s}_{g-1} \rangle \subset \cdots \langle \mathfrak{s}_g, \dots, \mathfrak{s}_1 \rangle$$

and let w_i^\dagger be the image of \mathfrak{s}_{g+1-i} in $\text{Fil}_i^\dagger / \text{Fil}_{i-1}^\dagger$, for all $i = 1, \dots, g$. Then the triple $(\mathbb{I}_g, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\})$ defines a section of the $\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)} \times \mathcal{T}_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g}(\mathbf{Z}/p^n \mathbf{Z})$ -torsor $\pi_\infty^{-1}(\mathcal{V}_\infty) \rightarrow \mathcal{V}_\infty$. Consequently, one obtains an isomorphism

$$\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)} \times \mathcal{T}_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z}) \xrightarrow{\sim} \pi_\infty^{-1}(\mathcal{V}_\infty), \quad \gamma' \mapsto (\mathbb{I}_g, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\}) \cdot \gamma'$$

and thus an isomorphism

$$\begin{aligned}
\Phi : \pi_{\infty,*} \mathcal{O}_{\mathcal{I}\mathcal{W}_{w,\infty}}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}^+ &\xrightarrow{\sim} \left\{ \text{analytic functions } U_{\text{GL}_g,1}^{\text{opp},(w)} \times T_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}} \right\} \\
f &\mapsto (\gamma' \mapsto f((\mathbb{I}_g, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\}) \cdot \gamma')).
\end{aligned}$$

We claim that if $\gamma^* f = f$ for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n)$, then $\gamma^* \Phi(f) = \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger) \Phi(f)$. Indeed, for any $\gamma' \in U_{\text{GL}_g,1}^{\text{opp},(w)} \times T_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g,1}(\mathbf{Z}/p^n \mathbf{Z})$, we have

$$\begin{aligned}
(\gamma^* \Phi(f))(\gamma') &= (\gamma^* f)(\gamma^* (\psi^\dagger, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\}) \cdot \gamma') \\
&= f((\psi^\dagger, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\}) \cdot \mathfrak{t}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger) \cdot \gamma') \\
&= \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger) \Phi(f)(\gamma'),
\end{aligned}$$

where the second equation follows from the identity $\gamma^* \mathfrak{s}^\dagger = \mathfrak{s}^\dagger \mathfrak{t}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger)$.

On the other hand, we can identify $\omega^{(1)}$ with the set of analytic functions

$$f : U_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times T_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g,1}(\mathbf{Z}/p^n\mathbf{Z}) \rightarrow \mathcal{O}_{\mathcal{V}_\infty}^+(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}$$

satisfying

- $f(\mathbf{v}\tau\nu) = \kappa_{\mathcal{U}}^\vee(\tau)f(\mathbf{v})$ for all $(\mathbf{v}, \tau, \nu) \in U_{\mathrm{GL}_g,1}^{\mathrm{opp},(w)} \times T_{\mathrm{GL}_g,0}^{(w)} \times U_{\mathrm{GL}_g}(\mathbf{Z}/p^n\mathbf{Z})$;
- $\gamma^* f = \rho_{\kappa_{\mathcal{U}}^\vee}(\gamma_a^\dagger + \mathfrak{z}\gamma_c^\dagger)f$ for all $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \Gamma(p^n)$.

Therefore, putting $\Psi_2 := \Phi^{-1}$, one obtains the desired isomorphism

$$\Psi_2 : \omega^{(1)} \xrightarrow{\sim} \omega^{(2)}.$$

Construction of Ψ_3 . By the construction of $\underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}$ and Lemma 2.2.4, one immediately obtains an identification of $\omega^{(2)}$ with $\widetilde{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}(\mathcal{V})$. We simply take Ψ_3 to be this identification.

Putting everything together, the composition $\Psi^+ = \Psi_3 \circ \Psi_2 \circ \Psi_1$ yields an isomorphism

$$\Psi^+ : \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}(\mathcal{V}) \xrightarrow{\sim} \widetilde{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}(\mathcal{V}).$$

It is also straightforward to check that the construction is functorial in \mathcal{V} . By gluing, we arrive at an isomorphism

$$\Psi^+ : \underline{\omega}_{n,w}^{\kappa_{\mathcal{U}},+} \xrightarrow{\sim} \widetilde{\omega}_{n,w}^{\kappa_{\mathcal{U}},+}.$$

□

Consider the p -adically completed sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{k\acute{e}t}}}^+$ -modules associated with $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$; namely, let

$$\underline{\omega}_{w,\mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}},+} := \varprojlim_m \left(\underline{\omega}_w^{\kappa_{\mathcal{U}},+} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}}^+} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{k\acute{e}t}}}^+ / p^m \right)$$

and

$$\underline{\omega}_{w,\mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}} := \underline{\omega}_{w,\mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}},+} \left[\frac{1}{p} \right].$$

By Remark 3.3.12 and Proposition 3.3.13, $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ can be identified with the sheaf of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+/\Gamma(p^n)$ -invariants of $h_{n,*}\widetilde{\omega}_{n,w}^{\kappa_{\mathcal{U}}}$. Hence, $\underline{\omega}_{w,\mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}}$ can be identified with the sheaf of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+/\Gamma(p^n)$ -invariants of $h_{n,*}\widetilde{\omega}_{n,w,\mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}}$, later of which is an admissible Kummer étale Banach sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{k\acute{e}t}}} \widehat{\otimes} R_{\mathcal{U}}$ -modules by Lemma 3.3.10 and Lemma A.3.10. Consequently, such a description allows us to apply Corollary A.3.14 to the sheaf $\underline{\omega}_{w,\mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}}$. This will be used in the proof of Lemma 5.2.2.

3.4 Classical Siegel modular forms

In this subsection, we show that the space of w -overconvergent Siegel modular forms does contain all of the classical Siegel modular forms.

Let $k = (k_1, \dots, k_g) \in \mathbf{Z}_{\geq 0}^g$ be a dominant weight and consider $k^\vee = (-k_g, \dots, -k_1)$. Let

$$\mathcal{M} := \mathrm{Isom}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}}(\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}}^g, \underline{\omega}_{\mathrm{Iw}^+})$$

be the GL_g -torsor over $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$ together with the structure morphism $\vartheta : \mathcal{M} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+}$. Then the sheaf $\underline{\omega}_{\mathrm{Iw}^+}^k$ of *classical Siegel modular forms of weight k (of strict Iwahori level)* is defined to be

$$\underline{\omega}_{\mathrm{Iw}^+}^k := \vartheta_* \mathcal{O}_{\mathcal{M}}[k^\vee];$$

namely, the subsheaf of $\vartheta_* \mathcal{O}_{\mathcal{M}}$ on which T_{GL_g} acts through the character k^\vee . The *space of classical Siegel modular forms of weight k (of strict Iwahori level)* is defined to be

$$M_{\mathrm{Iw}^+}^{k, \mathrm{cl}} := H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \underline{\omega}_{\mathrm{Iw}^+}^k)$$

equipped with naturally defined Hecke operators.

Remark 3.4.1. One can also define the sheaf of integral classical Siegel modular forms by

$$\underline{\omega}_{\mathrm{Iw}^+}^{k, +} := \vartheta_* \mathcal{O}_{\mathcal{M}}^+[k^\vee].$$

But we do not need this in the current subsection.

Restricting to the w -ordinary locus, we may consider the sheaf $\underline{\omega}_{\mathrm{Iw}^+}^k|_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}}$. Repeating the strategy as in the proof of Proposition 3.3.13, we arrive at the following explicit description of $\underline{\omega}_{\mathrm{Iw}^+}^k|_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}}$.

Definition 3.4.2. (i) Let $P(\mathrm{GL}_g, \mathbb{A}^1)$ denote the \mathbf{Q}_p -vector space of maps $\mathrm{GL}_g \rightarrow \mathbb{A}^1$ between algebraic varieties over \mathbf{Q}_p .

(ii) For every uniform \mathbf{C}_p -Banach algebra B , define

$$P(\mathrm{GL}_g, B) := P(\mathrm{GL}_g, \mathbb{A}^1) \widehat{\otimes}_{\mathbf{Q}_p} B$$

and let $P_k(\mathrm{GL}_g, B)$ denote the subspace of $P(\mathrm{GL}_g, B)$ consisting of those $f : \mathrm{GL}_g \rightarrow B$ such that $f(\gamma\beta) = k(\beta)f(\gamma)$ for all $\gamma \in \mathrm{GL}_g$ and $\beta \in B_{\mathrm{GL}_g}$.

(iii) There is a natural left action of GL_g on $P_k(\mathrm{GL}_g, B)$ given by

$$(\gamma \cdot f)(\gamma') = f({}^t\gamma \gamma')$$

for all $\gamma, \gamma' \in \mathrm{GL}_g$ and $f \in P_k(\mathrm{GL}_g, B)$. This left action is denoted by

$$\rho_k : \mathrm{GL}_g \rightarrow \mathrm{Aut}(P_k(\mathrm{GL}_g, B)).$$

Proposition 3.4.3. For any affinoid open $\mathcal{V} \subset \overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$ with preimage \mathcal{V}_∞ in $\overline{\mathcal{X}}_{\Gamma(p^\infty), w}$, we have a natural identification

$$\underline{\omega}_{\mathrm{Iw}^+}^k(\mathcal{V}) = \left\{ f \in P_k(\mathrm{GL}_g, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}(\mathcal{V}_\infty)) : \gamma^* f = \rho_k(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f, \quad \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \right\}.$$

In particular, there is a natural injection

$$\underline{\omega}_{\mathrm{Iw}^+}^k|_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}} \hookrightarrow \underline{\omega}_w^k. \quad (6)$$

Proof. For the first statement, the strategy in the proof of Proposition 3.3.13 applies verbatim, except that we consider the torsor \mathcal{M} in place of \mathcal{IW}_w^+ . The details are left to the reader. The inclusion $\underline{\omega}_{\mathrm{Iw}^+}^k|_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}} \hookrightarrow \underline{\omega}_w^k$ follows from the natural inclusion $P_k(\mathrm{GL}_g, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}(\mathcal{V}_\infty)) \hookrightarrow C_k^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty), w}}(\mathcal{V}_\infty))$. \square

The following result shows that the space of classical forms naturally injects into the space of overconvergent modular forms.

Lemma 3.4.4. *The Hecke-equivariant composition of maps*

$$M_{\mathrm{Iw}^+}^{k,\mathrm{cl}} = H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \underline{\omega}_{\mathrm{Iw}^+}^k) \xrightarrow{\mathrm{Res}} H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_{\mathrm{Iw}^+}^k) \hookrightarrow M_{\mathrm{Iw}^+, w}^k$$

is injective.

Proof. It suffices to show that

$$\mathrm{Res} : H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \underline{\omega}_{\mathrm{Iw}^+}^k) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_{\mathrm{Iw}^+}^k)$$

is injective; namely, given any global section f of $\underline{\omega}_{\mathrm{Iw}^+}^k$ that vanishes on $\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$, we have to show that $f = 0$ on every irreducible component of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$.

For every algebraic variety Y over \mathbf{C}_p , we know that Y is irreducible if and only if the associated adic space \mathcal{Y} over $\mathrm{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ is irreducible (see [Con99, Theorem 2.3.1] and [Hub13, §1.1.11.(c)]). In particular, the irreducible components of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$ coincide with the irreducible components of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}^{\mathrm{tor}}$. As $\overline{\mathcal{X}}_{\mathrm{Iw}^+}^{\mathrm{tor}}$ is a compactification of X_{Iw^+} , its irreducible components correspond to the irreducible components of X_{Iw^+} . Under the identification

$$X_{\mathrm{Iw}^+}(\mathbf{C}) = \mathrm{GSp}_{2g}(\mathbf{Q}) \backslash \mathrm{GSp}_{2g}(\mathbf{A}_f) \times \mathbb{H}_g / \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \Gamma(N),$$

[Del71, §2] provides the following description of the irreducible components of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$:

$$\pi_0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}) = \mathbf{Q}_{>0} \backslash \mathbb{G}_m(\mathbf{A}_f) / \varsigma \left(\Gamma(N) \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \right).$$

where ς is the character of similitude involved in the definition of GSp_{2g} . There is a similar description for $\pi_0(\overline{\mathcal{X}})$. Note that $\pi_0(\overline{\mathcal{X}}_{\mathrm{Iw}^+})$ is the same as $\pi_0(\overline{\mathcal{X}})$ because $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ and $\mathrm{GSp}_{2g}(\mathbf{Z}_p)$ have the same image via ς . In particular, since every irreducible component in $\pi_0(\overline{\mathcal{X}})$ contains an ordinary point, every irreducible component of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$ intersects $\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$.

By definition, f can be viewed as a global section of the structure sheaf of \mathcal{M} . Let \mathcal{C} be any irreducible component of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$, it remains to show that f vanishes on $\mathcal{M} \times_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}} \mathcal{C}$. Indeed, observe that $\mathcal{M} \times_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}} \mathcal{C}$ is irreducible and f vanishes on $\mathcal{M} \times_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}} (\mathcal{C} \cap \overline{\mathcal{X}}_{\mathrm{Iw}^+, w})$. Hence, the desired vanishing follows from [Ber96, Proposition 0.1.13] which states that a rigid analytic function vanishing on an open subset of an irreducible rigid analytic variety is identically zero. \square

3.5 The construction à la Andreatta–Iovita–Pilloni

The sheaves $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ constructed in §3.1 are analogues of the overconvergent automorphic sheaves constructed by Andreatta–Iovita–Pilloni in [AIP15]. It is a natural question whether these two constructions coincide. In this subsection, we recall the construction in [AIP15]. Later in §3.7, we will present a comparison result.

Choose $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2})$ and let n be a positive integer such that $v < \frac{1}{2p^{n-1}}$. Consider the open subset

$$\overline{\mathcal{X}}(v) := \{ \mathbf{x} \in \overline{\mathcal{X}} : |\widetilde{\mathrm{Ha}}(\mathbf{x})| \geq p^{-v} \} \subset \overline{\mathcal{X}},$$

where $\widetilde{\mathrm{Ha}}$ is a fixed lift of the Hasse invariant. (We point out that, for those \mathbf{x} at the boundary, the Hasse invariant of \mathbf{x} is defined to be the Hasse invariant of the abelian part of the semiabelian scheme associated with \mathbf{x} .) Thanks to [AIP15, Proposition 4.1.3], for every $1 \leq m \leq n$, there is a universal canonical subgroup \mathcal{H}_m of level m of the tautological semiabelian variety over $\overline{\mathcal{X}}(v)$. Let $\underline{\omega}_v$ denote the restriction of $\underline{\omega}$ on $\overline{\mathcal{X}}(v)$.

We also consider the following finite covers of $\overline{\mathcal{X}}(v)$:

- Let

$$\overline{\mathcal{X}}_1(p^n)(v) := \text{Isom}_{\overline{\mathcal{X}}(v)}((\mathbf{Z}/p^n \mathbf{Z})^g, \mathcal{H}_n^\vee)$$

be the adic space over $\overline{\mathcal{X}}(v)$ which parameterises trivialisations of \mathcal{H}_n^\vee . Notice that the group $\text{GL}_g(\mathbf{Z}/p^n \mathbf{Z})$ naturally acts on $\overline{\mathcal{X}}_1(p^n)(v)$ from the right by permuting the trivialisations.

- Let

$$\overline{\mathcal{X}}_1(v) := \text{Isom}_{\overline{\mathcal{X}}(v)}((\mathbf{Z}/p \mathbf{Z})^g, \mathcal{H}_1^\vee)$$

be the adic space over $\overline{\mathcal{X}}(v)$ which parameterises trivialisations of \mathcal{H}_1^\vee .

- The group $\text{GL}_g(\mathbf{Z}/p \mathbf{Z})$ naturally acts on $\overline{\mathcal{X}}_1(v)$ from the right by permuting the trivialisations. By taking the quotient

$$\overline{\mathcal{X}}_{\text{Iw}}(v) := \overline{\mathcal{X}}_1(v)/B_{\text{GL}_g}(\mathbf{Z}/p \mathbf{Z}),$$

we obtain an adic space $\overline{\mathcal{X}}_{\text{Iw}}(v)$ over $\overline{\mathcal{X}}(v)$ which parameterises full flags $\text{Fil}_\bullet \mathcal{H}_1^\vee$ of \mathcal{H}_1^\vee .

- Let $\underline{\omega}_{n,v}$ be the pullback of $\underline{\omega}_v$ along $\overline{\mathcal{X}}_1(p^n)(v) \rightarrow \overline{\mathcal{X}}(v)$.
- Let $\underline{\omega}_{\text{Iw},v}$ be the pullback of $\underline{\omega}_v$ along $\overline{\mathcal{X}}_{\text{Iw}}(v) \rightarrow \overline{\mathcal{X}}(v)$.

In order to proceed, we need to introduce formal models of aforementioned geometric objects:

- Recall that $\overline{\mathfrak{X}}^{\text{tor}}$ is the formal completion of $\overline{X}_0^{\text{tor}}$ along the special fibre. Let $\widetilde{\mathfrak{X}}^{\text{tor}}(v)$ be the blowup of $\overline{\mathfrak{X}}^{\text{tor}}$ along the ideal $(\widetilde{\text{Ha}}, p^v)$. Let $\overline{\mathfrak{X}}^{\text{tor}}(v)$ be the p -adic completion of the normalisation of the largest open formal subscheme of $\widetilde{\mathfrak{X}}^{\text{tor}}(v)$ where the ideal $(\widetilde{\text{Ha}}, p^v)$ is generated by $\widetilde{\text{Ha}}$. Then $\overline{\mathfrak{X}}^{\text{tor}}(v)$ is a formal model of $\overline{\mathcal{X}}(v)$.
- Let $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$ be the normalisation of $\overline{\mathfrak{X}}^{\text{tor}}(v)$ in $\overline{\mathcal{X}}_1(p^n)(v)$. The group $\text{GL}_g(\mathbf{Z}/p^n \mathbf{Z})$ naturally acts on $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$.
- Let $\overline{\mathfrak{X}}_1^{\text{tor}}(v)$ be the normalisation of $\overline{\mathfrak{X}}^{\text{tor}}(v)$ in $\overline{\mathcal{X}}_1(v)$. The group $\text{GL}_g(\mathbf{Z}/p \mathbf{Z})$ naturally acts on $\overline{\mathfrak{X}}_1^{\text{tor}}(v)$.
- Let $\overline{\mathfrak{X}}_{\text{Iw}}^{\text{tor}}(v)$ be the normalisation of $\overline{\mathfrak{X}}^{\text{tor}}(v)$ in $\overline{\mathcal{X}}_{\text{Iw}}(v)$. We can identify $\overline{\mathfrak{X}}_{\text{Iw}}^{\text{tor}}(v)$ with the quotient $\overline{\mathfrak{X}}_1^{\text{tor}}(v)/B_{\text{GL}_g}(\mathbf{Z}/p \mathbf{Z})$.
- Let $\mathfrak{G}_v^{\text{univ}}$ be the tautological semiabelian scheme over $\overline{\mathfrak{X}}^{\text{tor}}(v)$ with the structure morphism

$$\pi : \mathfrak{G}_v^{\text{univ}} \rightarrow \overline{\mathfrak{X}}^{\text{tor}}(v).$$

Define

$$\underline{\Omega}_v := \pi_* \Omega_{\mathfrak{G}_v^{\text{univ}}/\overline{\mathfrak{X}}^{\text{tor}}(v)}^1.$$

- Let $\underline{\Omega}_{n,v}$ be the pullback of $\underline{\Omega}_v$ along $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v) \rightarrow \overline{\mathfrak{X}}^{\text{tor}}(v)$.
- Let $\underline{\Omega}_{\text{Iw},v}$ be the pullback of $\underline{\Omega}_v$ along $\overline{\mathfrak{X}}_{\text{Iw}}^{\text{tor}}(v) \rightarrow \overline{\mathfrak{X}}^{\text{tor}}(v)$.

Now suppose $w \in \mathbf{Q}_{>0}$ lies in the interval $\left(n - 1 + \frac{v}{p-1}, n - \frac{vp^n}{p-1}\right]$. Let

$$\psi_n^{\text{univ}} : (\mathbf{Z}/p^n \mathbf{Z})^g \cong \mathcal{H}_n^\vee$$

denote the universal trivialisation of \mathcal{H}_n^\vee over $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$. Then [AIP15, Proposition 4.3.1] yields a locally free $\mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)}$ -submodule $\mathcal{F} \subset \underline{\Omega}_{n,v}$ of rank g , equipped with a map

$$\text{HT}_{n,v,w} : (\mathbf{Z}/p^n \mathbf{Z})^g \xrightarrow{\psi_n^{\text{univ}}} \mathcal{H}_n^\vee \rightarrow \mathcal{F} \otimes_{\mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)}} \mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)} / p^w$$

which induces an isomorphism

$$\text{HT}_{n,v,w} \otimes \text{id} : (\mathbf{Z}/p^n \mathbf{Z})^g \otimes_{\mathbf{Z}} \mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)} / p^w \cong \mathcal{F} \otimes_{\mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)}} \mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)} / p^w.$$

More precisely, locally on $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$, consider the family version of the Hodge-Tate map

$$\text{HT}_n : (\mathbf{Z}/p^n \mathbf{Z})^g \xrightarrow{\psi_n^{\text{univ}}} \mathcal{H}_n^\vee \rightarrow \omega_{\mathcal{H}_n}$$

studied in [AIP15, §4]. Let $\epsilon_1, \dots, \epsilon_g$ be the standard $(\mathbf{Z}/p^n \mathbf{Z})$ -basis for $(\mathbf{Z}/p^n \mathbf{Z})^g$ and let $\widetilde{\text{HT}}_n(\epsilon_i)$ be lifts of $\text{HT}_n(\epsilon_i)$ from $\omega_{\mathcal{H}_n}$ to $\underline{\Omega}_{n,v}$. Then \mathcal{F} is generated by $\widetilde{\text{HT}}_n(\epsilon_1), \dots, \widetilde{\text{HT}}_n(\epsilon_g)$. It turns out this local construction glues to a locally free $\mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)}$ -module of rank g .

In [AIS15, §4.5], Andreatta–Iovita–Pilloni constructs a formal scheme $\mathfrak{W}_{w,v}^+$ over $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$ which parameterises certain w -compatible objects. More precisely, $\mathfrak{W}_{w,v}^+$ is the formal schemes over $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$ such that for every affine open subset $\text{Spf } R \subset \overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$ on which \mathcal{F} is free, $\mathfrak{W}_{w,v}^+(R)$ consists of pairs $(\text{Fil}_\bullet, \{w_i : i = 1, \dots, g\})$ where both Fil_\bullet and $\{w_i : i = 1, \dots, g\}$ are w -compatible with $\text{HT}_n(\epsilon_1), \dots, \text{HT}_n(\epsilon_g)$ in the sense of Definition 3.3.1.

Now we go back to the generic fibres. Let $\mathcal{W}_{w,v}^+$ be the adic space associated with the formal scheme $\mathfrak{W}_{w,v}^+$ over $\text{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Then we have a chain of morphisms of adic spaces

$$\pi^{\text{AIP}} : \mathcal{W}_{w,v}^+ \rightarrow \overline{\mathcal{X}}_1(p^n)(v) \rightarrow \overline{\mathcal{X}}_1(v) \rightarrow \overline{\mathcal{X}}_{\text{Iw}}(v).$$

Recall the group adic spaces $\mathcal{B}_w^{\text{opp}}$, $\mathcal{T}_{\text{GL}_g,0}^{(w)}$, and $\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)}$ defined in Definition 3.3.3.

Lemma 3.5.1. *(i) $\mathcal{W}_{w,v}^+$ is a $\mathcal{B}_w^{\text{opp}}$ -torsor over $\overline{\mathcal{X}}_1(p^n)(v)$. Namely, locally on $\overline{\mathcal{X}}_1(p^n)(v)$, we have identification*

$$\mathcal{W}_{w,v}^+ \simeq \overline{\mathcal{X}}_1(p^n)(v) \times_{\text{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})} \mathcal{B}_w^{\text{opp}}$$

where $\mathcal{B}_w^{\text{opp}}$ permutes the points $(\text{Fil}_\bullet, \{w_i\})$ from the right.

(ii) Similarly, $\mathcal{W}_{w,v}^+$ is a $\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)} \times \mathcal{T}_{\text{GL}_g,0}^{(w)}$ -torsor over $\overline{\mathcal{X}}_{\text{Iw}}(v)$.

Proof. These are clear from the construction. □

Finally, we are ready to define the overconvergent automorphic sheaves of Andreatta–Iovita–Pilloni.

Definition 3.5.2. *Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a w -analytic weight.*

- (i) *Andreatta–Iovita–Pilloni’s sheaf of w -analytic v -overconvergent Siegel modular forms of weight $\kappa_{\mathcal{U}}$ (of Iwahori level) is defined to be*⁵

$$\underline{\omega}_{w,v}^{\kappa_{\mathcal{U}},\text{AIP}} := \pi_*^{\text{AIP}} \mathcal{O}_{\mathcal{IW}_{w,v}^+}[\kappa_{\mathcal{U}}^{\vee}],$$

where $\pi_*^{\text{AIP}} \mathcal{O}_{\mathcal{IW}_{w,v}^+}[\kappa_{\mathcal{U}}^{\vee}]$ stands for the subsheaf of $\pi_*^{\text{AIP}}(\mathcal{O}_{\mathcal{IW}_{w,v}^+} \widehat{\otimes} R_{\mathcal{U}})$ consisting of sections on which $T_{\text{GL}_g,0}$ acts via the character $\kappa_{\mathcal{U}}^{\vee}$ and $U_{\text{GL}_g}(\mathbf{Z}/p^n\mathbf{Z})$ acts trivially.

- (ii) *Andreatta–Iovita–Pilloni’s space of w -analytic v -overconvergent Siegel modular forms of weight $\kappa_{\mathcal{U}}$ (of Iwahori level) is*

$$M_{\text{Iw},w,v}^{\kappa_{\mathcal{U}},\text{AIP}} := H^0(\overline{\mathcal{X}}_{\text{Iw}}(v), \underline{\omega}_{w,v}^{\kappa_{\mathcal{U}},\text{AIP}}).$$

- (iii) *The space of locally analytic overconvergent Siegel modular forms of weight $\kappa_{\mathcal{U}}$ (of Iwahori level) is*

$$M_{\text{Iw}}^{\kappa_{\mathcal{U}},\text{AIP}} := \lim_{\substack{v \rightarrow 0 \\ w \rightarrow \infty}} M_{\text{Iw},w,v}^{\kappa_{\mathcal{U}},\text{AIP}}.$$

- (iv) *Recall that $\mathcal{Z}_{\text{Iw}} = \overline{\mathcal{X}}_{\text{Iw}} \setminus \mathcal{X}_{\text{Iw}}$ is the boundary divisor. Andreatta–Iovita–Pilloni’s sheaf of w -analytic v -overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ (of Iwahori level) is defined to be the subsheaf $\underline{\omega}_{w,v,\text{cusp}}^{\kappa_{\mathcal{U}},\text{AIP}} = \underline{\omega}_{w,v}^{\kappa_{\mathcal{U}},\text{AIP}}(-\mathcal{Z}_{\text{Iw}})$ of $\underline{\omega}_{w,v}^{\kappa_{\mathcal{U}},\text{AIP}}$ consisting of sections that vanish along \mathcal{Z}_{Iw} .*

Andreatta–Iovita–Pilloni’s space of w -analytic v -overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ (of Iwahori level) is defined to be

$$S_{\text{Iw},w,v}^{\kappa_{\mathcal{U}},\text{AIP}} := H^0(\overline{\mathcal{X}}_{\text{Iw}}(v), \underline{\omega}_{w,v,\text{cusp}}^{\kappa_{\mathcal{U}},\text{AIP}}),$$

and the space of locally analytic overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ (of Iwahori level) is

$$S_{\text{Iw}}^{\kappa_{\mathcal{U}},\text{AIP}} := \lim_{\substack{v \rightarrow 0 \\ w \rightarrow \infty}} S_{\text{Iw},w,v}^{\kappa_{\mathcal{U}},\text{AIP}}.$$

Remark 3.5.3. Similar to Remark 3.3.7, in Definition 3.5.2 (i), there is no difference between taking $\kappa_{\mathcal{U}}^{\vee}$ -eigenspaces with respect to $T_{\text{GL}_g,0}$ - or $\mathcal{T}_{\text{GL}_g,0}^{(w)}$ -actions.

3.6 The w -ordinary loci and the (pseudo-)canonical subgroups

In §3.7, we will prove the comparison between our perfectoid construction of the overconvergent Siegel modular forms and the construction of Andreatta–Iovita–Pilloni. Immediate from the definitions, one observes the incompatibility of the underlying adic spaces used in the two constructions. That is, we employ the w -ordinary locus in the perfectoid construction while the authors of [AIP15] make use of the “ v -locus” $\overline{\mathcal{X}}_{\text{Iw}}(v)$. Therefore, as a preparation for the comparison result, we have to first compare these two different loci. Due to technical reasons, we assume $p > 2g$ in this subsection.

We start with the definition of the v -locus at the strict Iwahori level. Recall from §3.5 that, for any $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2})$, $\overline{\mathcal{X}}_1(p^n)(v)$ (resp., $\overline{\mathcal{X}}_1(v)$; resp., $\overline{\mathcal{X}}_{\text{Iw}}(v)$) is the adic space over $\overline{\mathcal{X}}(v)$ which parameterises trivialisations of \mathcal{H}_n^{\vee} (resp., trivialisations of \mathcal{H}_1^{\vee} ; resp., full flags of \mathcal{H}_1^{\vee}). In particular, $\overline{\mathcal{X}}_1(v)$ is equipped with a natural right action of $\text{GL}_g(\mathbf{Z}/p\mathbf{Z})$ permuting the trivialisations. Consider the quotient

$$\overline{\mathcal{X}}_{\text{Iw}^+}(v) := \overline{\mathcal{X}}_1(v)/T_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z})$$

⁵The notations $\underline{\omega}_{w,v}^{\kappa_{\mathcal{U}},\text{AIP}}$, $M_{\text{Iw},w,v}^{\kappa_{\mathcal{U}},\text{AIP}}$, and $M_{\text{Iw}}^{\kappa_{\mathcal{U}},\text{AIP}}$ correspond to the notations $\omega_w^{\dagger \kappa_{\mathcal{U}}}$, $M_w^{\dagger \kappa_{\mathcal{U}}}(\mathcal{X}_{\text{Iw}}(p)(v))$, and $M^{\dagger \kappa_{\mathcal{U}}}(\mathcal{X}_{\text{Iw}}(p))$, respectively, in [AIP15].

which is an adic space over $\overline{\mathcal{X}}(v)$ parameterising the “strict Iwahori structures” of \mathcal{H}_1^\vee ; namely, it parameterises full flags $\text{Fil}_\bullet \mathcal{H}_1^\vee$ of \mathcal{H}_1^\vee together with a collection of subgroups $\{D_i : i = 1, \dots, g\}$ of \mathcal{H}_1^\vee of order p such that

$$\text{Fil}_i \mathcal{H}_1^\vee = \langle D_1, \dots, D_i \rangle$$

for all $i = 1, \dots, g$. There is a chain of natural projections among these v -loci

$$\overline{\mathcal{X}}_1(p^n)(v) \rightarrow \overline{\mathcal{X}}_1(v) \rightarrow \overline{\mathcal{X}}_{\text{Iw}^+}(v) \rightarrow \overline{\mathcal{X}}_{\text{Iw}}(v) \rightarrow \overline{\mathcal{X}}(v).$$

One can identify $\overline{\mathcal{X}}_{\text{Iw}}(v)$ as the quotient of $\overline{\mathcal{X}}_{\text{Iw}^+}(v)$ by the finite group $U_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z})$.

The main result of this subsection is the following:

Theorem 3.6.1. *Given $\Gamma \in \{\text{Iw}^+, \text{Iw}\}$, the system of w -ordinary loci $\{\overline{\mathcal{X}}_{\Gamma, w} : w \in \mathbf{Q}_{>0}\}$ and the system of v -loci $\{\overline{\mathcal{X}}_\Gamma(v) : v \in \mathbf{Q}_{>0} \cap [0, 1/2)\}$ are mutually cofinal. More precisely,*

- (i) *For any given $v \in \mathbf{Q}_{>0} \cap [0, 1/2)$, there exists sufficiently large $w \in \mathbf{Q}_{>0}$ such that $\overline{\mathcal{X}}_{\Gamma, w} \subset \overline{\mathcal{X}}_\Gamma(v)$.*
- (ii) *For any given $w \in \mathbf{Q}_{>0}$, there exists sufficiently small $v \in \mathbf{Q}_{>0} \cap [0, 1/2)$ such that $\overline{\mathcal{X}}_\Gamma(v) \subset \overline{\mathcal{X}}_{\Gamma, w}$.*

To prove Theorem 3.6.1, we follow the strategy in [CHJ17, §2.3]. However, we have to generalise their study of pseudocanonical subgroups to the case of semiabelian schemes with constant toric rank.

Let C be an algebraically closed complete nonarchimedean field containing \mathbf{Q}_p and let \mathcal{O}_C be its ring of integers. Suppose the valuation v_p on C is normalised so that $v_p(p) = 1$. Let G be a semiabelian scheme over \mathcal{O}_C of dimension g with constant toric rank $r \leq g$. That is, G sits inside an extension

$$0 \rightarrow T \rightarrow G \rightarrow A \rightarrow 0,$$

where T is a torus of rank r over \mathcal{O}_C and A is an abelian scheme of dimension $g - r$ over \mathcal{O}_C . (We say that G is **principally polarised** if A is principally polarised.) One sees that the p -adic Tate module $T_p G := \varprojlim_n G[p^n](C)$ is isomorphic to \mathbf{Z}_p^{2g-r} .

Recall the Hodge–Tate complex over \mathcal{O}_C

$$0 \rightarrow \text{Lie } G \rightarrow T_p G \otimes_{\mathbf{Z}_p} \mathcal{O}_C \rightarrow \omega_{G^\vee} \rightarrow 0,$$

where ω_{G^\vee} is the dual of the Lie algebra $\text{Lie } G^\vee$ of the dual semiabelian scheme G^\vee , and the second last map is induced from the Hodge–Tate map $\text{HT}_G : T_p G \rightarrow \omega_{G^\vee}$. By [FGL08, Théorème II. 1.1], the cohomology of this complex is killed by $p^{1/(p-1)}$.

Definition 3.6.2. *Recall that $V_p = V \otimes_{\mathbf{Z}} \mathbf{Z}_p \simeq \mathbf{Z}_p^{2g}$ is equipped with the standard basis e_1, \dots, e_{2g} together with a symplectic pairing. For every $0 \leq r \leq g$, let $V_{p,r}$ denote the \mathbf{Z}_p -submodule of V_p spanned by $e_{r+1}, e_{r+2}, \dots, e_{2g-r}$, equipped with the induced symplectic pairing. We also write $V'_{p,r}$ to be \mathbf{Z}_p -submodule of V_p spanned by e_1, \dots, e_{2g-r} and write $W_{p,r}$ to be the one spanned by e_1, \dots, e_r . There is an obvious split exact sequence*

$$0 \rightarrow W_{p,r} \rightarrow V'_{p,r} \rightarrow V_{p,r} \rightarrow 0.$$

Definition 3.6.3. *Let G be a principally polarised semiabelian scheme over \mathcal{O}_C of dimension g with constant toric rank $r \leq g$.*

- (i) *An isomorphism $\alpha : V'_{p,r} \xrightarrow{\sim} T_p G$ is called a **trivialisation** of $T_p G$ if it is part of a commutative diagram*

$$\begin{array}{ccc} V_{p,r} & \xrightarrow{\sim} & T_p A \\ \uparrow & & \uparrow \\ V'_{p,r} & \xrightarrow{\sim} & T_p G \\ \uparrow & & \uparrow \\ W_{p,r} & \xrightarrow{\sim} & T_p T \end{array}$$

where

- the vertical arrows on the left are the ones as in Definition 3.6.2;
- the vertical arrows on the right are induced from the exact sequence $0 \rightarrow T \rightarrow G \rightarrow A \rightarrow 0$;
- the top arrow preserves the symplectic pairings.

(ii) A trivialisation $\alpha : V'_{p,r} \rightarrow T_p G$ is **w-ordinary** if $\text{HT}_G(\alpha(e_i)) \in p^w \omega_{G^\vee}$ for all $i = 1, \dots, g$.

(iii) We say that G is **w-ordinary** if it admits a w-ordinary trivialisation.

Remark 3.6.4. From the definition, if G is w-ordinary, it is w' -ordinary for any $w' > w$. It is also clear that G is ordinary if and only if it is w-ordinary for all $w \in \mathbf{Q}_{>0}$.

Lemma 3.6.5. Let G be a w-ordinary semiabelian scheme (of dimension g with constant toric rank r) over \mathcal{O}_C and let $n \in \mathbf{Z}_{\geq 1}$ such that $n < w + 1$. The Hodge-Tate map HT_G induces a map

$$G[p^n](C) \rightarrow (\text{image } \text{HT}_G) / p^{\min\{n, w\}} (\text{image } \text{HT}_G).$$

Then the schematic closure of the kernel of this map defines a flat subgroup scheme $H_n \subset G[p^n]$ whose generic fibre is isomorphic to $(\mathbf{Z}/p^n \mathbf{Z})^g$. Moreover, if α is a w-ordinary trivialisation of $T_p G$, then $H_n(C)$ is generated by $\alpha(e_1), \dots, \alpha(e_g)$. Here we have abused the notations and still use $\alpha(e_i)$'s to denote their images in $G[p^n](C)$.

Proof. Since the Hodge-Tate complex is exact after inverting p , the image of $\text{Lie } G$ in $T_p G \otimes_{\mathbf{Z}_p} \mathcal{O}_C$ is a rank g sub-lattice in the kernel of $T_p G \otimes_{\mathbf{Z}_p} \mathcal{O}_C \rightarrow \omega_{G^\vee}$. Hence, the kernel of $\text{HT}_G : T_p G \rightarrow \omega_{G^\vee}$ has rank at most g .

On the other hand, there is a commutative diagram

$$\begin{array}{ccc} T_p G & \xrightarrow{\text{HT}_G} & \omega_{G^\vee} \\ \downarrow & & \downarrow \\ G[p^n](C) & \xrightarrow{\text{HT}_{G[p^n]}} & \omega_{G[p^n]^\vee} \end{array},$$

where the right vertical arrow is induced from the natural identification $\omega_{G[p^n]^\vee} = \omega_{G^\vee} / p^n \omega_{G^\vee}$. Consequently, $\ker \text{HT}_{G[p^n]}$ also has rank at most g .

Let α be a w-ordinary trivialisation of $T_p G$. Since $n < w + 1$, the kernel of the composition

$$T_p G \xrightarrow{\text{HT}_G} \omega_{G^\vee} \rightarrow \omega_{G^\vee} / p^n \omega_{G^\vee}$$

necessarily contains $\alpha(e_i)$, for all $i = 1, \dots, g$. Since $\alpha(e_i)$'s are \mathbf{Z}_p -linearly independent, their images in $G[p^n](C)$ are $(\mathbf{Z}/p^n \mathbf{Z})$ -linearly independent and hence generate $\ker \text{HT}_{G[p^n]}$. Consequently, H_n is precisely the schematic closure in $G[p^n]$ of the subgroup of $G[p^n](C)$ generated by $\{\alpha(e_i) : i = 1, \dots, g\}$. Flatness of H_n follows from the flatness of G . \square

Definition 3.6.6. The subgroup scheme H_n defined in Lemma 3.6.5 is called the **pseudocanonical subgroup of level n** . When $n = 1$, we simply call H_1 the **pseudocanonical subgroup** of G .

Lemma 3.6.7. Let $m \leq n$ be positive integers and let $w \in \mathbf{Q}_{>0}$ such that $w > n$. Let G be a semiabelian scheme (of dimension g with constant toric rank r) over \mathcal{O}_C . Suppose G is w-ordinary. Then, G/H_m is $(w - m)$ -ordinary, and for any $m' \in \mathbf{Z}$ with $m < m' \leq n$, we have $H'_{m'-m} = H_{m'}/H_m$, where $H'_{m'-m}$ is the pseudocanonical subgroup of G/H_m of level $m' - m$.

Proof. The proof is the same as in [CHJ17, Lemma 2.11] as long as we use the matrix $\text{diag}(p^m \mathbb{1}_g, \mathbb{1}_{g-r})$ in place of $\text{diag}(1, p^m)$. Notice that the “ p^m ” factor appears at the bottom right corner in *loc. cit.* because they work with a slightly different action of $\text{GL}_2(\mathbf{Q}_p)$. \square

Before stating the next lemma, let us recall the notion of the *degree* of a finite flat group scheme over \mathcal{O}_C studied in [Far11]. If M is a p -power torsion \mathcal{O}_C -module of finite presentation, we can write

$$M \simeq \bigoplus_{i=1}^l \mathcal{O}_C / a_i \mathcal{O}_C$$

for some $a_i \in \mathcal{O}_C$, $i = 1, \dots, l$. Then the degree of M is defined to be $\deg M := \sum_{i=1}^l v_p(a_i)$. Now, if H is a finite flat group scheme over \mathcal{O}_C and let ω_H denote the \mathcal{O}_C -module of invariant differentials on H , then we define the *degree* of H to be $\deg H := \deg \omega_H$.

Lemma 3.6.8. *Let G be a w -ordinary semiabelian scheme (of dimension g with constant toric rank r) over \mathcal{O}_C and let α be a w -ordinary trivialisation. Let ω_{H_1} be the dual of $\text{Lie } H_1$ and let $\omega_{H_1^\vee}$ be the dual of $\text{Lie } H_1^\vee$. For $i = 1, \dots, g$, let $H_{1,i}$ be the schematic closure in H_1 of the subgroup generated by $\alpha(e_i)$. Then*

- (i) *Each $H_{1,i}$ is isomorphic to $\text{Spec}(\mathcal{O}_C[X]/(X^p - a_i X))$ for some $a_i \in \mathcal{O}_C$. The dual $H_{1,i}^\vee$ is isomorphic to $\text{Spec}(\mathcal{O}_C[X]/(X^p - b_i X))$ with $a_i b_i = p$.*
- (ii) *We have isomorphisms $\omega_{H_1} \simeq \bigoplus_{i=1}^g \mathcal{O}_C / a_i \mathcal{O}_C$ and $\omega_{H_1^\vee} \simeq \bigoplus_{i=1}^g \mathcal{O}_C / b_i \mathcal{O}_C$. In particular, we have $\deg H_1 = \sum_{i=1}^g v_p(a_i)$ and $\deg H_1^\vee = \sum_{i=1}^g v_p(b_i) = g - \sum_{i=1}^g v_p(a_i)$.*
- (iii) *Under the identification $\omega_{H_1^\vee} \simeq \bigoplus_{i=1}^g \mathcal{O}_C / b_i \mathcal{O}_C$, the image of the (linearised) Hodge–Tate map*

$$H_1(C) \otimes_{\mathbf{Z}_p} \mathcal{O}_C \rightarrow \omega_{H_1^\vee}$$

is equal to $\bigoplus_{i=1}^g c_i \mathcal{O}_C / b_i \mathcal{O}_C$ for some $c_i \in \mathcal{O}_C$ such that $v_p(c_i) = v_p(a_i)/(p-1)$, $i = 1, \dots, g$.

Proof. Since each $H_{1,i}$ is a finite flat group scheme over \mathcal{O}_C of degree p , the assertion follows from classical Oort–Tate theory. See, for example, [Far11, §6.5, Lemme 9]. \square

Recall from [AIP15, §3.1] that the *Hodge height* of G is defined to be the “truncated” p -adic valuation of the Hasse invariant of G . See *loc. cit.* for details.

Lemma 3.6.9. *Let G be a w -ordinary semiabelian scheme (of dimension g with constant toric rank r) over \mathcal{O}_C . Suppose $\frac{(2g-1)p}{2g(p-1)} < w \leq 1$.⁶ Then H_1 coincides with the canonical subgroup of G . Moreover, the Hodge height of G is smaller than $1/2$.*

Proof. We follow the strategy of the proof of [CHJ17, Lemma 2.14]. Consider the commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & H_1(C) & \longrightarrow & G[p](C) \\ & & \downarrow \text{HT}_{H_1} & & \downarrow \text{HT}_{G[p]} \\ 0 & \longrightarrow & \omega_{H_1^\vee} & \longrightarrow & \omega_{G[p]^\vee} \end{array}$$

with exact rows. Notice that we have an identification $\omega_{G[p]^\vee} = \omega_{G^\vee}/p\omega_{G^\vee}$. Let α be a w -ordinary trivialisation of $T_p G$. According to Lemma 3.6.5, $\alpha(e_1), \dots, \alpha(e_g)$ form a basis for $H_1(C)$. Also, by definition, we have $\text{HT}_{G[p]}(\alpha(e_i)) \in p^w \omega_{G[p]^\vee}$.

⁶The inequalities are valid because of the assumption $p > 2g$ at the beginning of the subsection.

Now, with respect to the generators $\alpha(e_1), \dots, \alpha(e_g)$ of $H_1(C)$, the map $\omega_{H_1^\vee} \rightarrow \omega_{G[p]^\vee}$ can be identified with the inclusion

$$\bigoplus_{i=1}^g \mathcal{O}_C / b_i \mathcal{O}_C \rightarrow (\mathcal{O}_C / p \mathcal{O}_C)^g, \quad (x_1, \dots, x_g) \mapsto (a_1 x_1, \dots, a_g x_g).$$

Therefore, we see that

$$a_i \text{HT}_{H_1}(\alpha(e_i)) = \text{HT}_{G[p]}(\alpha(e_i)) \in p^w \omega_{G[p]^\vee}.$$

By Lemma 3.6.8 (iii), we know that $\text{HT}_{H_1}(\alpha(e_i))$ has valuation $v_p(a_i)/(p-1)$. This implies

$$w \leq v_p(a_i) + \frac{v_p(a_i)}{p-1} = \frac{pv_p(a_i)}{p-1}.$$

Consequently, we have

$$\deg H_1 = \sum_{i=1}^g v_p(a_i) \geq \frac{gw(p-1)}{p} > \frac{g(p-1)}{p} \cdot \frac{(2g-1)p}{2g(p-1)} = \frac{2g-1}{2} = g - \frac{1}{2}.$$

It follows from [AIP15, Proposition 3.1.2] that H_1 is exactly the canonical subgroup of G and the Hodge height of G is less than $\frac{1}{2}$. \square

Remark 3.6.10. The lemma might hold without the assumption $p > 2g$ as long as one can produce finer estimates on the degree and the Hodge height. However, we do not attempt to find these better estimates.

Proposition 3.6.11. *Let G be a w -ordinary semiabelian scheme (of dimension g with constant toric rank r) over \mathcal{O}_C . Suppose $\frac{(2g-1)p}{2g(p-1)} + n - 1 < w \leq n$, then H_n coincides with the canonical subgroup of G of level n . In this case, the Hodge height of G is less than $\frac{1}{2p^{n-1}}$.*

Proof. The proof follows from induction. The case for $n = 1$ is precisely Lemma 3.6.9.

Assume that the statement is affirmative for $n - 1$. By Lemma 3.6.7, G/H_1 is $(w-1)$ -ordinary and we have $\frac{(2g-1)p}{2g(p-1)} + n - 2 < w - 1 \leq n - 1$. The induction hypothesis implies that the pseudocanonical subgroup H_n/H_1 of level $n - 1$ of G/H_1 is the canonical subgroup of level $n - 1$ and that the Hodge height of G/H_1 is less than $\frac{1}{2p^{n-2}}$.

However, H_1 coincides with the canonical subgroup of G by Lemma 3.6.9. Hence, by [Far11, Théorème 6 (4)] (see also [AIP15, Theorem 3.1.1 (5)]), we see that the Hodge height of G is bounded by $\frac{1}{2p^{n-1}}$ and that H_n is the canonical subgroup of level n of G . \square

Corollary 3.6.12. *Let $n \in \mathbf{Z}_{\geq 1}$ and suppose $w \in \mathbf{Q}_{>0}$ such that $\frac{(2g-1)p}{2g(p-1)} < w \leq n$. Then there exists $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2p^{n-1}})$ and a natural inclusion $\overline{\mathcal{X}}_w \hookrightarrow \overline{\mathcal{X}}(v)$.*

Proof. It suffices to work with (C, \mathcal{O}_C) -points for algebraically closed complete nonarchimedean field C containing \mathbf{Q}_p . (Notice that the classical points determine these adic spaces by [Hub13, (1.1.11)]). Let $\mathbf{x} \in \overline{\mathcal{X}}_w(C, \mathcal{O}_C)$. By the properness of $\overline{\mathcal{X}}$, the point \mathbf{x} extends to an \mathcal{O}_C -point $\tilde{\mathbf{x}}$ of $\overline{\mathcal{X}}^{\text{tor}}$. One can associate with $\tilde{\mathbf{x}}$ a 1-motive $\widetilde{M}_{\tilde{\mathbf{x}}} = [Y \rightarrow \widetilde{G}_{\tilde{\mathbf{x}}}]$ where $\widetilde{G}_{\tilde{\mathbf{x}}}$ is a semiabelian scheme (of dimension g with constant toric rank) over \mathcal{O}_C and Y is a free \mathbf{Z} -module of finite rank (see, for example, [Str10]).

From the definition of the Hodge–Tate period map (see §B.2 for a quick review), we see that $\widetilde{G}_{\tilde{\mathbf{x}}}$ is w -ordinary. By Proposition 3.6.11, the Hodge height of $\widetilde{G}_{\tilde{\mathbf{x}}}$ is smaller than $\frac{1}{2p^{n-1}}$. This means $\mathbf{x} \in \overline{\mathcal{X}}(v)(C, \mathcal{O}_C)$ for some $v < \frac{1}{2p^{n-1}}$ and so we are done. \square

Recall that, for any $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2})$, \mathcal{H}_1 is the universal canonical subgroup of the tautological semiabelian variety over $\overline{\mathcal{X}}(v)$. Let $w > \frac{(2g-1)p}{2g(p-1)}$ and pick v so that $\overline{\mathcal{X}}_w \hookrightarrow \overline{\mathcal{X}}(v)$ as in Corollary 3.6.12. We still write \mathcal{H}_1 for its pullback to $\overline{\mathcal{X}}_w$.

In this case, consider

$$\overline{\mathcal{X}}_{1,w} := \text{Isom}_{\overline{\mathcal{X}}_w}((\mathbf{Z}/p\mathbf{Z})^g, \mathcal{H}_1^\vee);$$

namely, the adic space over $\overline{\mathcal{X}}_w$ which parameterises trivialisations of \mathcal{H}_1^\vee . The group $\text{GL}_g(\mathbf{Z}/p\mathbf{Z})$ naturally acts on $\overline{\mathcal{X}}_{1,w}$ by permuting the trivialisations.

Lemma 3.6.13. *For $w > \frac{(2g-1)p}{2g(p-1)}$, there are natural identifications*

$$\overline{\mathcal{X}}_{1,w}/B_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z}) = \overline{\mathcal{X}}_{\text{Iw},w} \quad \text{and} \quad \overline{\mathcal{X}}_{1,w}/T_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z}) = \overline{\mathcal{X}}_{\text{Iw}^+,w}.$$

Proof. We only give the proof for the first identity. The second one is similar and left to the readers.

We first focus on the part away from the boundary. Let $\mathcal{X}_w = \overline{\mathcal{X}}_w \cap \mathcal{X}$ and let $\mathcal{A}_w^{\text{univ}}$ be the universal abelian variety over \mathcal{X}_w .

The key observation is that any trivialisation $\psi : (\mathbf{Z}/p\mathbf{Z})^g \rightarrow \mathcal{H}_1^\vee$ induces a full flag $\text{Fil}_\bullet^\psi \mathcal{A}_w^{\text{univ}}[p]$ on $\mathcal{A}_w^{\text{univ}}[p]$. Indeed, let $\epsilon_1, \dots, \epsilon_g$ denote the standard basis for $(\mathbf{Z}/p\mathbf{Z})^g$ and let $\text{Fil}_\bullet^\psi \mathcal{H}_1^\vee$ be the full flag of \mathcal{H}_1^\vee given by

$$0 \subset \langle \psi(\epsilon_1) \rangle \subset \langle \psi(\epsilon_1), \psi(\epsilon_2) \rangle \subset \dots \subset \langle \psi(\epsilon_1), \dots, \psi(\epsilon_g) \rangle.$$

Consider the natural projection

$$\text{pr} : \mathcal{A}_w^{\text{univ}}[p] \xrightarrow{\sim} \mathcal{A}_w^{\text{univ}}[p]^\vee \twoheadrightarrow \mathcal{H}_1^\vee$$

where the first isomorphism is induced from the principal polarisation. Then the desired full flag $\text{Fil}_\bullet^\psi \mathcal{A}_w^{\text{univ}}[p]$ is given by

$$\text{Fil}_i^\psi \mathcal{A}_w^{\text{univ}}[p] := \begin{cases} \text{pr}^{-1} \text{Fil}_{i-g}^\psi \mathcal{H}_1^\vee, & i > g \\ (\text{pr}^{-1} \text{Fil}_{g-i}^\psi \mathcal{H}_1^\vee)^\perp, & i \leq g \end{cases}.$$

Moreover, if two such ψ 's induce the same $\text{Fil}_\bullet^\psi \mathcal{H}_1^\vee$, then the associated $\text{Fil}_\bullet^\psi \mathcal{A}_w^{\text{univ}}[p]$ coincide. Hence, the assignment $\psi \mapsto \text{Fil}_\bullet^\psi \mathcal{A}_w^{\text{univ}}[p]$ induces a natural inclusion $\mathcal{X}_{1,w}/B_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z}) \subset \mathcal{X}_{\text{Iw},w}$ away from the boundary.

Conversely, using the w -ordinarity, one sees that the universal full flag $\text{Fil}_\bullet \mathcal{A}_w^{\text{univ}}[p]$ on $\mathcal{X}_{\text{Iw},w}$ induces a full flag $\text{Fil}_\bullet \mathcal{H}_1^\vee$ of \mathcal{H}_1^\vee given by

$$\text{Fil}_i \mathcal{H}_1^\vee = \text{pr}((\text{Fil}_{g-i} \mathcal{A}_w^{\text{univ}}[p])^\perp)$$

for $i = 1, \dots, g$. This yields the opposite inclusion away from the boundary.

In order to extend to the boundary, one considers the 1-motives on the boundary strata and same argument as above applies verbatim. The details are left to the reader. \square

Finally, we prove Theorem 3.6.1.

Proof of Theorem 3.6.1. (i) We may assume $v = \frac{1}{2p^{n-1}}$ for some sufficiently large n . In this case, we can take any $\frac{(2g-1)p}{2g(p-1)} + n - 1 < w \leq n$. Indeed, by Corollary 3.6.12, we have a Cartesian diagram

$$\begin{array}{ccc} \overline{\mathcal{X}}_{1,w} & \hookrightarrow & \overline{\mathcal{X}}_1(v) \\ \downarrow & & \downarrow \\ \overline{\mathcal{X}}_w & \hookrightarrow & \overline{\mathcal{X}}(v) \end{array}$$

where the top arrow is equivariant under the action of $\text{GL}_g(\mathbf{Z}/p\mathbf{Z})$. Taking the quotient by either $B_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z})$ or $T_{\text{GL}_g}(\mathbf{Z}/p\mathbf{Z})$, and applying Lemma 3.6.13, we obtain the desired inclusions.

- (ii) We may assume $n - 1 < w < n$ for some sufficiently large n . Pick $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2p^{n-1}})$ such that $w \in (n - 1 + \frac{v}{p-1}, n - \frac{vp^n}{p-1}]$. Applying [AIP15, Proposition 3.2.1], on the level of classical points, we obtain a natural inclusion $\overline{\mathcal{X}}(v)(C, \mathcal{O}_C) \hookrightarrow \overline{\mathcal{X}}_w(C, \mathcal{O}_C)$ and hence an inclusion $\overline{\mathcal{X}}(v) \hookrightarrow \overline{\mathcal{X}}_w$. There is a Cartesian diagram

$$\begin{array}{ccc} \overline{\mathcal{X}}_1(v) & \hookrightarrow & \overline{\mathcal{X}}_{1,w} \\ \downarrow & & \downarrow \\ \overline{\mathcal{X}}(v) & \hookrightarrow & \overline{\mathcal{X}}_w \end{array}$$

Once again, applying Lemma 3.6.13 and taking the corresponding quotients yield the desired inclusions. \square

3.7 Comparison of the two constructions

In this section, we still assume $p > 2g$. The aim of this subsection is to prove the following theorem which compares the overconvergent automorphic sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ constructed in §3.1 and the sheaf $\underline{\omega}_{w,v}^{\kappa_{\mathcal{U}}, \text{AIP}}$ of Andreatta–Iovita–Pilloni.

For any $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2})$, let $h_{\diamond} : \overline{\mathcal{X}}_{\text{Iw}^+}(v) \rightarrow \overline{\mathcal{X}}_{\text{Iw}}(v)$ denote the natural projection.

Theorem 3.7.1. *Suppose $n > \frac{g}{p-1}$ and let $v \in \mathbf{Q}_{>0} \cap [0, \frac{1}{2p^{n-1}})$, $w \in \mathbf{Q}_{>0} \cap (n - 1 + \frac{v}{p-1}, n - \frac{vp^n}{p-1}]$. (In particular, by the proof of Theorem 3.6.1 (ii), there are natural inclusions $\overline{\mathcal{X}}_{\text{Iw}}(v) \hookrightarrow \overline{\mathcal{X}}_{\text{Iw},w}$ and $\overline{\mathcal{X}}_{\text{Iw}^+}(v) \hookrightarrow \overline{\mathcal{X}}_{\text{Iw}^+,w}$.) Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a weight such that $w > 1 + r_{\mathcal{U}}$. Then, over $\overline{\mathcal{X}}_{\text{Iw}^+}(v)$, there is a canonical isomorphism of sheaves*

$$\Psi : \underline{\omega}_w^{\kappa_{\mathcal{U}}} |_{\overline{\mathcal{X}}_{\text{Iw}^+}(v)} \xrightarrow{\sim} h_{\diamond}^* \underline{\omega}_{w,v}^{\kappa_{\mathcal{U}}, \text{AIP}}$$

Recall that the space of overconvergent Siegel modular forms of weight $\kappa_{\mathcal{U}}$ of strict Iwahori level (see Definition 3.1.14 (v)) is defined to be

$$M_{\text{Iw}^+}^{\kappa_{\mathcal{U}}} = \varinjlim_{w \rightarrow \infty} M_{\text{Iw}^+,w}^{\kappa_{\mathcal{U}}}$$

where

$$M_{\text{Iw}^+,w}^{\kappa_{\mathcal{U}}} = H^0(\overline{\mathcal{X}}_{\text{Iw}^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}).$$

We can also extend the notion of overconvergent Siegel modular forms of Andreatta–Iovita–Pilloni to the case of strict Iwahori level.

Definition 3.7.2. *Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a weight.*

- (i) *Let $v \in \mathbf{Q}_{>0} \cap [0, 1/2)$ and $w \in \mathbf{Q}_{>0}$. Suppose $\kappa_{\mathcal{U}}$ is w -analytic. The **space of w -analytic v -overconvergent Siegel modular forms of weight $\kappa_{\mathcal{U}}$ (of strict Iwahori level)** of Andreatta–Iovita–Pilloni is defined to be*

$$M_{\text{Iw}^+,w,v}^{\kappa_{\mathcal{U}}, \text{AIP}} := H^0(\overline{\mathcal{X}}_{\text{Iw}^+}(v), h_{\diamond}^* \underline{\omega}_{w,v}^{\kappa_{\mathcal{U}}, \text{AIP}}).$$

- (ii) *The **space of locally analytic overconvergent Siegel modular forms of weight $\kappa_{\mathcal{U}}$ (of strict Iwahori level)** of Andreatta–Iovita–Pilloni is defined to be*

$$M_{\text{Iw}^+}^{\kappa_{\mathcal{U}}, \text{AIP}} := \varinjlim_{\substack{v \rightarrow 0 \\ w \rightarrow \infty}} M_{\text{Iw}^+,w,v}^{\kappa_{\mathcal{U}}, \text{AIP}}.$$

(iii) Similarly, the **space of w -analytic v -overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ (of strict Iwahori level)** of Andreatta–Iovita–Pilloni is defined to be

$$S_{\mathrm{Iw}^+, w, v}^{\kappa_{\mathcal{U}}, \mathrm{AIP}} := H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}(v), h_{\diamond}^* \underline{\omega}_{w, v, \mathrm{cusp}}^{\kappa_{\mathcal{U}}, \mathrm{AIP}}),$$

and the **space of locally analytic overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ (of strict Iwahori level)** of Andreatta–Iovita–Pilloni is defined to be

$$S_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}, \mathrm{AIP}} := \lim_{\substack{v \rightarrow 0 \\ w \rightarrow \infty}} S_{\mathrm{Iw}^+, w, v}^{\kappa_{\mathcal{U}}, \mathrm{AIP}}.$$

Then we have the following immediate corollary of Theorem 3.7.1 and Theorem 3.6.1.

Corollary 3.7.3. *There are canonical isomorphisms*

$$M_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}} \cong M_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}, \mathrm{AIP}} \quad \text{and} \quad S_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}} \cong S_{\mathrm{Iw}^+}^{\kappa_{\mathcal{U}}, \mathrm{AIP}}.$$

Remark 3.7.4. In fact, it will follow from the construction of Ψ that the isomorphisms in Corollary 3.7.3 is also Hecke-equivariant, except that the U_p -operators are only equivariant up to p -power scalars. More precisely, the authors of [AIP15] normalise their U_p -operators $U_{p, i}^{\mathrm{AIP}}$ by dividing by a certain power of p . Therefore, for all $i = 1, \dots, g-1$, our $U_{p, i}$ acts as $p^{i(g+1)} U_{p, i}^{\mathrm{AIP}}$, and $U_{p, g}$ acts as $p^{g(g+1)/2} U_{p, g}^{\mathrm{AIP}}$.

The rest of the subsection is dedicated to the proof of Theorem 3.7.1.

Let n, v, w , and $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be as in Theorem 3.7.1. Recall that the $\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}(v)}$ -module (resp., $\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}}(v)}$ -module) $\underline{\omega}_{\mathrm{Iw}^+, v}$ (resp., $\underline{\omega}_{\mathrm{Iw}, v}$) is locally free of rank g . Let $\mathcal{V}' \subset \overline{\mathcal{X}}_{\mathrm{Iw}}(v)$ be an affinoid open subset such that $\underline{\omega}_{\mathrm{Iw}, v}|_{\mathcal{V}'}$ is free, and let $\mathcal{V} \subset \overline{\mathcal{X}}_{\mathrm{Iw}^+}(v)$ be the preimage of \mathcal{V}' . To construct Ψ , it suffices to establish a canonical isomorphism

$$\Psi : \underline{\omega}_{w, v}^{\kappa_{\mathcal{U}}}(\mathcal{V}) \xrightarrow{\sim} h_{\diamond}^* \underline{\omega}_{w, v}^{\kappa_{\mathcal{U}}, \mathrm{AIP}}(\mathcal{V})$$

for every such \mathcal{V} , which is also functorial in \mathcal{V} .

As a preparation, consider the pullback diagram

$$\begin{array}{ccc} \mathcal{IW}_{w, v, \infty}^+ & \longrightarrow & \mathcal{IW}_{w, v}^+ \\ \pi_{\infty}^{\mathrm{AIP}} \downarrow & & \downarrow \pi^{\mathrm{AIP}} \\ \overline{\mathcal{X}}_{\Gamma(p^{\infty})}(v) & \xrightarrow{h_{\mathrm{Iw}}} & \overline{\mathcal{X}}_{\mathrm{Iw}}(v) \end{array}$$

where $\overline{\mathcal{X}}_{\Gamma(p^{\infty})}(v)$ is the preimage of $\overline{\mathcal{X}}_{\mathrm{Iw}}(v)$ under the natural morphism $h_{\mathrm{Iw}} : \overline{\mathcal{X}}_{\Gamma(p^{\infty}), w} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}, w}$. The existence of the pullback follows from the same argument as in the proof of Proposition 3.3.13. For later usage, we denote by \mathcal{V}_{∞} (resp., \mathcal{V}_{∞}^+) the preimage of \mathcal{V}' in $\overline{\mathcal{X}}_{\Gamma(p^{\infty})}(v)$ (resp., in $\mathcal{IW}_{w, v, \infty}^+$) under the projection h_{Iw} (resp., $h_{\mathrm{Iw}} \circ \pi_{\infty}^{\mathrm{AIP}}$).

Since $\mathcal{IW}_{w, v}^+$ is a $\mathcal{U}_{\mathrm{GL}_g, 1}^{\mathrm{opp}, (w)} \times \mathcal{T}_{\mathrm{GL}_g, 0}^{(w)} \times U_{\mathrm{GL}_g}(\mathbf{Z}/p^n \mathbf{Z})$ -torsor over $\overline{\mathcal{X}}_{\mathrm{Iw}}$, we know that $\mathcal{IW}_{w, v, \infty}^+$ is likewise a $\mathcal{U}_{\mathrm{GL}_g, 1}^{\mathrm{opp}, (w)} \times \mathcal{T}_{\mathrm{GL}_g, 0}^{(w)} \times U_{\mathrm{GL}_g}(\mathbf{Z}/p^n \mathbf{Z})$ -torsor over $\overline{\mathcal{X}}_{\Gamma(p^{\infty})}(v)$. In what follows, we provide an explicit moduli interpretation of this torsor, in three steps.

Step 1. Observe that the natural projection $h_{\mathrm{Iw}} : \overline{\mathcal{X}}_{\Gamma(p^{\infty})}(v) \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}}(v)$ factors as

$$h_{\mathrm{Iw}} : \overline{\mathcal{X}}_{\Gamma(p^{\infty})}(v) \xrightarrow{h_1} \overline{\mathcal{X}}_1(p^n)(v) \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}}(v).$$

Indeed, away from the boundary, the map h_1 can be described as follows. Let $\mathcal{X}_{\Gamma(p^\infty)}(v)$ be the part of $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$ away from the boundary. For every point $(A, \lambda, \psi_N, \psi_{p^\infty}) \in \mathcal{X}_{\Gamma(p^\infty)}(v)$, consider the dual trivialisation

$$\psi_{p^\infty}^\vee : V_p^\vee \xrightarrow{\sim} T_p A^\vee.$$

Modulo p^n , we obtain a symplectic isomorphism

$$\psi_{p^n}^\vee : V_p^\vee \otimes_{\mathbf{Z}_p} (\mathbf{Z}/p^n \mathbf{Z}) \xrightarrow{\sim} A[p^n]^\vee.$$

Then h_1 sends $(A, \lambda, \psi_N, \psi_{p^\infty})$ to $(A, \lambda, \psi_N, \psi)$ where ψ is the composition

$$\psi : (\mathbf{Z}/p^n \mathbf{Z})^g \hookrightarrow V_p^\vee \otimes_{\mathbf{Z}_p} (\mathbf{Z}/p^n \mathbf{Z}) \xrightarrow{\psi_{p^n}^\vee} A[p^n]^\vee \twoheadrightarrow H_n^\vee$$

with the first arrow sending e_i to $e_{g+1-i}^\vee \otimes 1$, for all $i = 1, \dots, g$, and the last arrow being the natural surjection. From the proof of Lemma 3.6.13, we see that ψ is indeed a trivialisation of H_n^\vee .

Using the language of 1-motives, this description of h_1 also extends to the boundary. The details are left to the readers.

Step 2. Recall that, in §3.5, we defined a locally free $\mathcal{O}_{\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)}$ -submodule $\mathcal{F} \subset \underline{\Omega}_{n,v}$ on $\overline{\mathfrak{X}}_1^{\text{tor}}(p^n)(v)$. Passing to the adic generic fibre, let $\underline{\omega}_{n,v}^+$ denote the sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_1(p^n)(v)}^+$ -module on $\overline{\mathcal{X}}_1(p^n)(v)$ associated with $\underline{\Omega}_{n,v}$. Then \mathcal{F} can be identified with a locally free $\mathcal{O}_{\overline{\mathcal{X}}_1(p^n)(v)}^+$ -submodule of $\underline{\omega}_{n,v}^+$, which is still denoted by \mathcal{F} . Moreover, let \mathcal{F}_∞ be the pullback of \mathcal{F} to $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$ along h_1 .

Recall as well the $\mathcal{O}_{\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}}$ -modules $\underline{\Omega}_{\Gamma(p^n)}^{\text{mod}} \subset \underline{\Omega}_{\Gamma(p^n)}$ constructed in §B.2. Passing to the adic generic fibre, they induce $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}^+}$ -modules $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+} \subset \underline{\omega}_{\Gamma(p^n)}^+$ on $\overline{\mathcal{X}}_{\Gamma(p^n)}$. Let $\underline{\omega}_{\Gamma(p^\infty)}^{\text{mod},+} \subset \underline{\omega}_{\Gamma(p^\infty)}^+$ be their pullbacks to $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ and let $\underline{\omega}_{\Gamma(p^\infty),v}^{\text{mod},+} \subset \underline{\omega}_{\Gamma(p^\infty),v}^+$ be their restrictions on $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$.

We claim that there is a natural inclusion

$$\mathcal{F}_\infty \subset \underline{\omega}_{\Gamma(p^\infty),v}^{\text{mod},+}.$$

Indeed, recall the map

$$\text{HT}_n : (\mathbf{Z}/p^n \mathbf{Z})^g \rightarrow \omega_{\mathcal{H}_n}$$

on $\overline{\mathcal{X}}_1(p^n)(v)$ constructed in §3.5. Pulling back to $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$, we obtain a map

$$\text{HT}_{n,\infty} : (\mathbf{Z}/p^n \mathbf{Z})^g \rightarrow \omega_{\mathcal{H}_{n,\infty}}$$

where $\mathcal{H}_{n,\infty}$ is the pullback of \mathcal{H}_n along the projection $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v) \rightarrow \overline{\mathcal{X}}_1(p^n)(v)$. On the other hand, recall the map $\text{HT}_{\Gamma(p^\infty)}$ on $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ constructed in §B.2. Restricting to $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$ and modulo p^n , we obtain a map

$$\text{HT}_{\Gamma(p^\infty),n,v} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \underline{\omega}_{\Gamma(p^\infty),v}^{\text{mod},+} / p^n \underline{\omega}_{\Gamma(p^\infty),v}^{\text{mod},+}.$$

These maps fit into a commutative diagram

$$\begin{array}{ccccc} & \underline{\omega}_{\Gamma(p^\infty),v}^{\text{mod},+} & \hookrightarrow & \underline{\omega}_{\Gamma(p^\infty),v}^+ & \\ & \downarrow & & \downarrow & \\ (\mathbf{Z}/p^n \mathbf{Z})^g & \xrightarrow{\text{HT}_{n,\infty}} & \omega_{\mathcal{H}_{n,\infty}} & & \\ \downarrow & & \downarrow & & \downarrow \\ V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) & \xrightarrow{\text{HT}_{\Gamma(p^\infty),n,v}} & \underline{\omega}_{\Gamma(p^\infty),v}^{\text{mod},+} / p^n & \hookrightarrow & \underline{\omega}_{\Gamma(p^\infty),v}^+ / p^n = \omega_{\mathcal{H}_{n,\infty}} / p^n \end{array}.$$

where the left inclusion sends ϵ_i to $e_{2g+1-i} \otimes 1$, for all $i = 1, \dots, g$. The equality at the bottom right corner follows from [AIP15, Proposition 3.2.1]. By definition, \mathcal{F}_∞ is generated by the lifts of $\text{HT}_{n,\infty}(\epsilon_i)$'s from $\omega_{\mathcal{H}_{n,\infty}}$ to $\underline{\omega}_{\Gamma(p^\infty),v}^+$ and hence the desired inclusion follows.

Step 3. We are now able to describe the torsor. Recall that there is a universal full flag $\text{Fil}_\bullet^{\text{univ}} \mathcal{H}_1^\vee$ of \mathcal{H}_1^\vee on $\overline{\mathcal{X}}_{\text{Iw}}(v)$. Pulling back to $\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$, we obtain universal full flag $\text{Fil}_\bullet^{\text{univ}} \mathcal{H}_{1,\infty}^\vee$ of $\mathcal{H}_{1,\infty}^\vee$. There is a natural projection $\Theta : \mathcal{H}_{n,\infty}^\vee \rightarrow \mathcal{H}_{1,\infty}^\vee$. Moreover, the Hodge–Tate map on $\mathcal{H}_{n,\infty}^\vee$ induces a map

$$\text{HT}_{\mathcal{H}_{n,\infty}^\vee} : \mathcal{H}_{n,\infty}^\vee \rightarrow \mathcal{F}_\infty \otimes_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)}^+ / p^w.$$

Then, for every affinoid open $\mathcal{Y} = \text{Spa}(R, R^+) \subset \overline{\mathcal{X}}_{\Gamma(p^\infty)}(v)$, the sections $\mathcal{IW}_{w,\infty}^+(\mathcal{Y})$ parametrise triples $(\psi, \text{Fil}_\bullet, \{w_i : i = 1, \dots, g\})$ where

- $\psi : (\mathbf{Z}/p^n \mathbf{Z})^g \xrightarrow{\sim} \mathcal{H}_{n,\infty}^\vee|_{\mathcal{Y}}$ is a trivialisation such that

$$\psi\langle \epsilon_1, \dots, \epsilon_i \rangle = \Theta(\text{Fil}_i^{\text{univ}} \mathcal{H}_{1,\infty}^\vee)$$

for all $i = 1, \dots, g$.

- Fil_\bullet is a full flag of the free R^+ -module $\mathcal{F}_\infty(\mathcal{Y})$, which is w -compatible with $\text{HT}_{\mathcal{H}_{n,\infty}^\vee}(\psi(\epsilon_1)), \dots, \text{HT}_{\mathcal{H}_{n,\infty}^\vee}(\psi(\epsilon_g))$ in the sense of Definition 3.3.1 (i).
- Each w_i is an R^+ -basis for $\text{Fil}_i / \text{Fil}_{i-1}$, which is w -compatible with $\text{HT}_{\mathcal{H}_{n,\infty}^\vee}(\psi(\epsilon_1)), \dots, \text{HT}_{\mathcal{H}_{n,\infty}^\vee}(\psi(\epsilon_g))$ in the sense of Definition 3.3.1 (ii).

Moreover, $\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)} \times \mathcal{T}_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g}(\mathbf{Z}/p^n \mathbf{Z})$ permutes these triples by right multiplication.

We are now ready to prove Theorem 3.7.1.

Proof of Theorem 3.7.1. The construction of Ψ is similar to the proof of Proposition 3.3.13. We only give a sketch of the proof. Indeed, the isomorphism Ψ is established via a sequence of isomorphisms

$$\Psi : \underline{\omega}_w^{\kappa_{\mathcal{U}}}(\mathcal{V}) \xrightarrow[\Psi_1]{\sim} \omega^{(1)} \xrightarrow[\Psi_2]{\sim} \omega^{(2)} \xrightarrow[\Psi_3]{\sim} h_{\diamond}^* \underline{\omega}_{w,v}^{\kappa_{\mathcal{U}}, \text{AIP}}(\mathcal{V}),$$

where

$$\omega^{(1)} := \left\{ f \in C_{\kappa_{\mathcal{U}}}^{w\text{-an}}(\text{Iw}_{\text{GL}_g}, \mathcal{O}_{\mathcal{V}_\infty}(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}}) : \gamma^* f = \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z} \gamma_c^\dagger) f, \quad \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \text{Iw}_{\text{GSp}_{2g}}^+ \right\}$$

and

$$\omega^{(2)} := \left\{ f \in \pi_{\infty,*}^{\text{AIP}} \mathcal{O}_{\mathcal{IW}_{w,\infty}^+}(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}} : \begin{array}{lll} \gamma^* f = f, & \tau^* f = \kappa_{\mathcal{U}}^\vee(\tau) f, & \nu^* f = f \\ \forall (\gamma, \tau, \nu) \in \text{Iw}_{\text{GSp}_{2g}}^+ \times T_{\text{GL}_g,0} \times U_{\text{GL}_g}(\mathbf{Z}/p^n \mathbf{Z}) \end{array} \right\}.$$

The construction of Ψ_1 and Ψ_3 follows verbatim as in Proposition 3.3.13. To construct Ψ_2 , consider $\mathfrak{s}^\dagger = (\mathfrak{s}_g \ \cdots \ \mathfrak{s}_1) \in \mathcal{F}_\infty(\mathcal{V}_\infty)^g$. Let $\text{Fil}_\bullet^\dagger$ be the full flag of the free $\mathcal{O}_{\mathcal{V}_\infty}^+$ -module $\mathcal{F}_\infty(\mathcal{V}_\infty)$ given by

$$\text{Fil}_\bullet^\dagger = 0 \subset \langle \mathfrak{s}_g \rangle \subset \langle \mathfrak{s}_g, \mathfrak{s}_{g-1} \rangle \subset \cdots \langle \mathfrak{s}_g, \dots, \mathfrak{s}_1 \rangle$$

and let w_i^\dagger be the image of \mathfrak{s}_{g+1-i} in $\text{Fil}_i^\dagger / \text{Fil}_{i-1}^\dagger$, for all $i = 1, \dots, g$. Moreover, consider the trivialisation

$$\psi^\dagger : (\mathbf{Z}/p^n \mathbf{Z})^g \xrightarrow{\sim} \mathcal{H}_{n,\infty}^\vee$$

obtained by pulling back the universal trivialisation of \mathcal{H}_n^\vee on $\overline{\mathcal{X}}_1(p^n)(v)$ along $h_1 : \overline{\mathcal{X}}_{\Gamma(p^\infty)}(v) \rightarrow \overline{\mathcal{X}}_1(p^n)(v)$. Then the triple $(\psi^\dagger, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\})$ defines a section of the $\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)} \times \mathcal{T}_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g}(\mathbf{Z}/p^n \mathbf{Z})$ -torsor $\pi_\infty^{\text{AIP}} : \mathcal{V}_\infty^+ \rightarrow \mathcal{V}_\infty$. Consequently, one obtains an isomorphism

$$\mathcal{U}_{\text{GL}_g,1}^{\text{opp},(w)} \times \mathcal{T}_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g}(\mathbf{Z}/p^n \mathbf{Z}) \xrightarrow{\sim} \mathcal{V}_\infty^+, \quad \gamma' \mapsto (\psi^\dagger, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\}) \cdot \gamma'$$

and thus an isomorphism

$$\begin{aligned} \Phi : \pi_{\infty,*}^{\text{AIP}} \mathcal{O}_{\mathcal{IW}_{w,\infty}^+}(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}} &\xrightarrow{\sim} \left\{ \text{analytic functions } U_{\text{GL}_g,1}^{\text{opp},(w)} \times T_{\text{GL}_g,0}^{(w)} \times U_{\text{GL}_g}(\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \mathcal{O}_{\mathcal{V}_\infty}(\mathcal{V}_\infty) \widehat{\otimes} R_{\mathcal{U}} \right\} \\ f &\mapsto (\gamma' \mapsto f((\psi^\dagger, \text{Fil}_\bullet^\dagger, \{w_i^\dagger\}) \cdot \gamma')). \end{aligned}$$

By the same calculation as in Proposition 3.3.13, one sees that, if $\gamma^* f = f$ for any $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \text{Iw}_{\text{GSp}_{2g}}^+$, then $\gamma^* \Phi(f) = \rho_{\kappa_{\mathcal{U}}}(\gamma_a^\dagger + \mathfrak{z} \gamma_c^\dagger) \Phi(f)$. This induces an isomorphism $\Phi : \omega^{(2)} \xrightarrow{\sim} \omega^{(1)}$. Taking $\Psi_2 = \Phi^{-1}$ does the job. \square

Remark 3.7.5. Notice that \mathfrak{s}_i 's are, in fact, integral. Hence, by pulling back the formal scheme \mathfrak{W}_w^+ to the modified integral model, the method above provides a strategy to compare our integral sheaf $\underline{\omega}_w^{\kappa_{\mathcal{U}},+}$ with the integral overconvergent automorphic sheaf constructed in [AIP15].

4 Overconvergent cohomology groups

In this section, we introduce the overconvergent cohomology groups that will later appear on the other side of the overconvergent Eichler–Shimura morphism. Our construction follows the standard constructions in the literature (see, for example, [Han17] and [JN19]).

4.1 Analytic functions and analytic distributions

Consider

$$\mathbf{T}_0 := \{(\gamma, \mathbf{v}) \in \text{Iw}_{\text{GL}_g} \times M_g(p \mathbf{Z}_p) : {}^t \gamma \check{\mathbb{I}}_g \mathbf{v} = {}^t \mathbf{v} \check{\mathbb{I}}_g \gamma\}.$$

Notice that a pair $(\gamma, \mathbf{v}) \in \text{Iw}_{\text{GL}_g} \times M_g(p \mathbf{Z}_p)$ lies in \mathbf{T}_0 if and only if there exist $\alpha_b, \alpha_d \in M_g(\mathbf{Z}_p)$ such that

$$\begin{pmatrix} \gamma & \alpha_b \\ \mathbf{v} & \alpha_d \end{pmatrix} \in \text{GSp}_{2g}(\mathbf{Q}_p) \cap M_{2g}(\mathbf{Z}_p).$$

In fact, there is a natural embedding

$$\mathbf{T}_0 \hookrightarrow \text{Iw}_{\text{GSp}_{2g}}, \quad (\gamma, \mathbf{v}) \mapsto \begin{pmatrix} \gamma & \\ \mathbf{v} & \check{\mathbb{I}}_g {}^t \gamma^{-1} \check{\mathbb{I}}_g \end{pmatrix}.$$

Also consider the subset \mathbf{T}_{00} of \mathbf{T}_0 defined by

$$\mathbf{T}_{00} := \{(\gamma, \mathbf{v}) \in \mathbf{T}_0 : \gamma \in U_{\text{GL}_g,1}^{\text{opp}}\}.$$

We can identify \mathbf{T}_{00} with $U_{\text{GSp}_{2g},1}^{\text{opp}}$ through the bijection

$$\mathbf{T}_{00} \rightarrow U_{\text{GSp}_{2g},1}^{\text{opp}}, \quad (\gamma, \mathbf{v}) \mapsto \begin{pmatrix} \gamma & \\ \mathbf{v} & \check{\mathbb{I}}_g {}^t \gamma^{-1} \check{\mathbb{I}}_g \end{pmatrix}.$$

Observe that \mathbf{T}_0 admits two natural actions:

(i) There is a right action of $\mathrm{Iw}_{\mathrm{GL}_g}$ given by

$$\mathbf{T}_0 \times \mathrm{Iw}_{\mathrm{GL}_g} \rightarrow \mathbf{T}_0, \quad ((\gamma, \mathbf{v}), \gamma') \mapsto (\gamma \gamma', \mathbf{v} \gamma').$$

To see that this is indeed a right action, we embed $\mathrm{Iw}_{\mathrm{GL}_g}$ into $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$ through $\gamma' \mapsto \begin{pmatrix} \gamma' & \\ & \check{\mathbb{I}}_g {}^t \gamma'^{-1} \check{\mathbb{I}}_g \end{pmatrix}$ and verify that

$$\begin{pmatrix} \gamma & * \\ \mathbf{v} & * \end{pmatrix} \begin{pmatrix} \gamma' & \\ & \check{\mathbb{I}}_g {}^t \gamma'^{-1} \check{\mathbb{I}}_g \end{pmatrix} = \begin{pmatrix} \gamma \gamma' & * \\ \mathbf{v} \gamma' & * \end{pmatrix}$$

(ii) There is a left action of $\Xi := \begin{pmatrix} \mathrm{Iw}_{\mathrm{GL}_g} & M_g(\mathbf{Z}_p) \\ M_g(p\mathbf{Z}_p) & M_g(\mathbf{Z}_p) \end{pmatrix} \cap \mathrm{GSp}_{2g}(\mathbf{Q}_p)$ given by

$$\Xi \times \mathbf{T}_0 \rightarrow \mathbf{T}_0, \quad \left(\begin{pmatrix} \alpha_a & \alpha_b \\ \alpha_c & \alpha_d \end{pmatrix}, (\gamma, \mathbf{v}) \right) \mapsto (\alpha_a \gamma + \alpha_b \mathbf{v}, \alpha_c \gamma + \alpha_d \mathbf{v}).$$

To see this is indeed a left action, it suffices to observe that

$$\begin{pmatrix} \alpha_a & \alpha_b \\ \alpha_c & \alpha_d \end{pmatrix} \begin{pmatrix} \gamma & * \\ \mathbf{v} & * \end{pmatrix} = \begin{pmatrix} \alpha_a \gamma + \alpha_b \mathbf{v} & * \\ \alpha_c \gamma + \alpha_d \mathbf{v} & * \end{pmatrix}.$$

Since $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ is a subset of Ξ , we also obtain a natural left action of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ on \mathbf{T}_0 .

Let $r \in \mathbf{Z}_{\geq 1}$ and let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be an r -analytic weight. In what follows, we will study “ r -analytic” functions on $U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}}$, \mathbf{T}_{00} , and \mathbf{T}_0 . Let us fix a (topological) isomorphism

$$\mathbf{Z}_p^{g^2} \simeq U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}}.$$

Definition 4.1.1. (i) We say that a function $f : U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}} \rightarrow R_{\mathcal{U}}^+$ is **r -analytic** if the composition

$$\mathbf{Z}_p^{g^2} \simeq U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}} \xrightarrow{f} R_{\mathcal{U}}^+ \hookrightarrow \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}$$

is r -analytic in the sense of Definition 3.1.6 (i).

(ii) We say that a function $f : \mathbf{T}_{00} \rightarrow R_{\mathcal{U}}^+$ is **r -analytic** if it is r -analytic viewed as a function on $U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}}$, via the identification $\mathbf{T}_{00} \cong U_{\mathrm{GSp}_{2g},1}^{\mathrm{opp}}$.

Write

$$A^{r,\circ}(\mathbf{T}_{00}, R_{\mathcal{U}}) := \{f : \mathbf{T}_{00} \rightarrow R_{\mathcal{U}}^+ : f \text{ is } r\text{-analytic}\}$$

and

$$A^r(\mathbf{T}_{00}, R_{\mathcal{U}}) := A^{r,\circ}(\mathbf{T}_{00}, R_{\mathcal{U}}) \left[\frac{1}{p} \right].$$

On the other hand, define

$$A_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) := \left\{ f : \mathbf{T}_0 \rightarrow R_{\mathcal{U}}^+ : \begin{array}{l} f(\gamma \beta, \mathbf{v} \beta) = \kappa_{\mathcal{U}}(\beta) f(\gamma, \mathbf{v}), \forall (\gamma, \mathbf{v}) \in \mathbf{T}_0, \beta \in B_{\mathrm{GL}_g,0} \\ f|_{\mathbf{T}_{00}} \text{ is } r\text{-analytic} \end{array} \right\}$$

and

$$A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}) := A_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \left[\frac{1}{p} \right].$$

There is an identification

$$A_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \xrightarrow{\sim} A^{r,\circ}(\mathbf{T}_{00}, R_{\mathcal{U}}), \quad f \mapsto f|_{\mathbf{T}_{00}}.$$

Taking duals, we obtain the corresponding spaces of r -analytic distributions

$$D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) := \mathrm{Hom}_{R_{\mathcal{U}}}^{\mathrm{cts}}(A_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}), R_{\mathcal{U}}^+)$$

and

$$D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}) := D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})\left[\frac{1}{p}\right].$$

The right action of $\mathrm{Iw}_{\mathrm{GL}_g}$ and the left action of Ξ on \mathbf{T}_0 then induce, respectively, a right action of $\mathrm{Iw}_{\mathrm{GL}_g}$ and a left action of Ξ on both $D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ and $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$. Furthermore, if $r' \geq r$, there is a natural injection $A_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \hookrightarrow A_{\kappa_{\mathcal{U}}}^{r',\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ which induces injections (see [Han17, §2.2])

$$D_{\kappa_{\mathcal{U}}}^{r',\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \hookrightarrow D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \quad \text{and} \quad D_{\kappa_{\mathcal{U}}}^{r'}(\mathbf{T}_0, R_{\mathcal{U}}) \hookrightarrow D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}).$$

We then write

$$D_{\kappa_{\mathcal{U}}}^{\dagger}(\mathbf{T}_0, R_{\mathcal{U}}) := \varprojlim_r D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}).$$

Suppose now that $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ is a small weight and take $r > 1 + r_{\mathcal{U}}$ (see Definition 3.1.8). Fix an ideal $\mathfrak{a}_{\mathcal{U}}$ of $R_{\mathcal{U}}$ defining the profinite topology on $R_{\mathcal{U}}$. Similar to [CHJ17, Proposition 3.1], $D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ admits a decreasing Ξ - and $\mathrm{Iw}_{\mathrm{GL}_g}$ -stable filtration $\mathrm{Fil}^{\bullet} D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ defined by

$$\mathrm{Fil}^j D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) := \ker \left(D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \rightarrow D_{\kappa_{\mathcal{U}}}^{r-1,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) / \mathfrak{a}_{\mathcal{U}}^j D_{\kappa_{\mathcal{U}}}^{r-1,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \right).$$

Write

$$D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) := D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) / \mathrm{Fil}^j D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$$

for every $j \in \mathbf{Z}_{\geq 1}$. One sees that

$$D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) = \varprojlim_j D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}),$$

making $D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ a profinite flat \mathbf{Z}_p -module in the sense of [CHJ17, Definition 6.1].

Remark 4.1.2. Note that we used the fact that $\kappa_{\mathcal{U}}$ is a small weight so that each $D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ is a finitely generated $(\mathbf{Z}/p^j \mathbf{Z})$ -module. We do not know if a similar statement holds for affinoid weights.

4.2 Overconvergent cohomology groups

Fix a small weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ and we consider the étale site $\mathcal{X}_{\mathrm{Iw}^+, \text{ét}}$. Recall that, for every $n \in \mathbf{Z}_{\geq 1}$, $\mathcal{X}_{\Gamma(p^n)}$ is a finite étale Galois cover over $\mathcal{X}_{\mathrm{Iw}^+}$ with Galois group $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ / \Gamma(p^n)$, and hence $\varprojlim_n \mathcal{X}_{\Gamma(p^n)}$ is a pro-étale Galois cover of $\mathcal{X}_{\mathrm{Iw}^+}$ with Galois group $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$. For each $j \in \mathbf{Z}_{\geq 1}$, let $\mathscr{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}$ be the locally constant sheaf on $\mathcal{X}_{\mathrm{Iw}^+, \text{ét}}$ associated with $D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ via

$$\pi_1^{\text{ét}}(\mathcal{X}_{\mathrm{Iw}^+}) \rightarrow \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \rightarrow \mathrm{Aut}(D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})).$$

We obtain an inverse system of étale locally constant sheaves $(\mathscr{D}_{\kappa_{\mathcal{U}},j}^{r,\circ})_{j \in \mathbf{Z}_{\geq 1}}$ on $\mathcal{X}_{\mathrm{Iw}^+, \text{ét}}$. This allows us to consider the étale cohomology groups

$$H_{\text{ét}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathscr{D}_{\kappa_{\mathcal{U}}}^{r,\circ}) := \varprojlim_j H_{\text{ét}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathscr{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}),$$

$$H_{\text{ét}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathscr{D}_{\kappa_{\mathcal{U}}}^r) := H_{\text{ét}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathscr{D}_{\kappa_{\mathcal{U}}}^{r,\circ})\left[\frac{1}{p}\right]$$

for every $t \in \mathbf{Z}_{\geq 0}$.

Remark 4.2.1. On the algebraic variety X_{Iw^+} , one can define locally constant sheaves $\mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}$ and étale cohomology groups $H_{\text{ét}}^t(X_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^{r,\circ})$ and $H_{\text{ét}}^t(X_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r)$ in the same way.

Recall the identification

$$X_{\text{Iw}^+}(\mathbf{C}) = \text{GSp}_{2g}(\mathbf{Q}) \backslash \text{GSp}_{2g}(\mathbf{A}_f) \times \mathbb{H}_g / \text{Iw}_{\text{GSp}_{2g}}^+ \Gamma(N).$$

By taking the trivial $\text{GSp}_{2g}(\mathbf{Z}_\ell)$ -action on $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$ for every prime number $\ell \neq p$ and letting $\text{Iw}_{\text{GSp}_{2g}}^+$ act on $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$ via the left action of Ξ , we see that $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$ defines a local system on the locally symmetric space $X_{\text{Iw}^+}(\mathbf{C})$. In particular, for every $t \in \mathbf{Z}_{\geq 0}$, we can consider the Betti cohomology group

$$H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})).$$

Proposition 4.2.2. *For every $t \in \mathbf{Z}_{\geq 0}$, there is a natural isomorphism*

$$H_{\text{ét}}^t(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r) \cong H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})).$$

Proof. The assertion follows from the isomorphisms

$$H_{\text{ét}}^t(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r) \cong H_{\text{ét}}^t(X_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r) \cong H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})),$$

where

- the first isomorphism follows from the comparison isomorphism between the étale cohomology groups of an algebraic variety and the ones on the corresponding adic spaces (see [Hub13, Theorem 3.8.1]); and
- the second isomorphism follows from the fact that $\text{Iw}_{\text{GSp}_{2g}}^+$ acts continuously on $D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ and the well-known Artin comparison between the étale cohomology of a complex algebraic variety and the Betti cohomology of the associated complex manifold.

Note that we have used the algebraic isomorphism $\mathbf{C}_p \simeq \mathbf{C}$ fixed at the beginning of the paper. □

Finally, we define the Hecke operators acting on $H_{\text{ét}}^t(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r)$. We begin with a brief recollection of the Hecke operators on $H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$ studied in [Han17]. We refer the readers to *loc. cit.* for a more detailed discussion.

Hecke operators outside pN . Let ℓ be a prime number not dividing pN . For any $\gamma \in \text{GSp}_{2g}(\mathbf{Q}_\ell) \cap M_{2g}(\mathbf{Z}_\ell)$, consider a double coset decomposition

$$\text{GSp}_{2g}(\mathbf{Z}_\ell) \gamma \text{GSp}_{2g}(\mathbf{Z}_\ell) = \bigsqcup_j \delta_j \gamma \text{GSp}_{2g}(\mathbf{Z}_\ell)$$

for some $\delta_j \in \text{GSp}_{2g}(\mathbf{Z}_\ell)$. If we take the trivial $\text{GSp}_{2g}(\mathbf{Q}_\ell)$ -action on $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$, then the natural left action of $\text{GSp}_{2g}(\mathbf{Q}_\ell)$ on $X_{\text{Iw}^+}(\mathbf{C})$ induces the Hecke operator

$$T_\gamma : H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})) \rightarrow H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})), \quad [\mu] \mapsto \sum_j (\delta_j \gamma) \cdot [\mu]. \quad (7)$$

Hecke operators at p . For the Hecke operators at p , recall from §3.2 the matrices

$$\mathbf{u}_{p,i} = \begin{cases} \begin{pmatrix} \mathbb{1}_i & & \\ & p \mathbb{1}_{g-i} & \\ & & p \mathbb{1}_{g-i} & \\ & & & p^2 \mathbb{1}_i \end{pmatrix}, & 1 \leq i \leq g-1 \\ \begin{pmatrix} \mathbb{1}_g & \\ & p \mathbb{1}_g \end{pmatrix}, & i = g \end{cases}$$

and we write

$$\mathbf{u}_{p,i} = \begin{pmatrix} \mathbf{u}_{p,i}^\square & \\ & \mathbf{u}_{p,i}^\blacksquare \end{pmatrix}.$$

For every $i = 1, \dots, g$, consider a $\mathbf{u}_{p,i}$ -action on \mathbf{T}_0 defined as follows: for every $(\gamma, \mathbf{v}) \in \mathbf{T}_0$, we put

$$\mathbf{u}_{p,i} \cdot (\gamma, \mathbf{v}) = (\mathbf{u}_{p,i}^\square \gamma_0 \mathbf{u}_{p,i}^{\square,-1}, \mathbf{u}_{p,i}^\blacksquare \mathbf{v}_0 \mathbf{u}_{p,i}^{\blacksquare,-1}) \beta$$

where we write $(\gamma, \mathbf{v}) = (\gamma_0, \mathbf{v}_0) \beta$ with $\gamma_0 \in U_{\mathrm{GL}_g,1}^{\mathrm{opp}}$ and $\beta \in B_{\mathrm{GL}_g,0}$. This then induces a $\mathbf{u}_{p,i}$ -action on $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$.

Similar to §3.2, for every $i = 1, \dots, g$, choose a double coset decomposition

$$\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \mathbf{u}_{p,i} \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ = \bigsqcup_j \delta_{ij} \mathbf{u}_{p,i} \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+.$$

with $\delta_{ij} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$. The natural left action of $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ on $X_{\mathrm{Iw}^+}(\mathbf{C})$ together with the actions of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ and $\mathbf{u}_{p,i}$ on $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$ induce the Hecke operator

$$U_{p,i} : H^t(X_{\mathrm{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})) \rightarrow H^t(X_{\mathrm{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})), \quad [\mu] \mapsto \sum_j \delta_{ij} \cdot (\mathbf{u}_{p,i} \cdot [\mu]). \quad (8)$$

Definition 4.2.3. (i) The Hecke operators T_γ (for $\gamma \in \mathrm{GSp}_{2g}(\mathbf{Q}_\ell) \cap M_{2g}(\mathbf{Z}_\ell)$ with $\ell \nmid Np$) and $U_{p,i}$ (for $i = 1, \dots, g$) acting on the overconvergent cohomology groups $H_{\mathrm{et}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r)$ are defined to be the operators T_γ and $U_{p,i}$ acting on $H^t(X_{\mathrm{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$ via the isomorphism in Proposition 4.2.2.

(ii) We define the operator U_p as the composition $U_p = \prod_{i=1}^g U_{p,i}$.

5 The overconvergent Eichler–Shimura morphism

In this section, we establish the second main result of this paper; *i.e.*, the construction of the overconvergent Eichler–Shimura morphism for Siegel modular forms. Our approach is similar to the one in [CHJ17]. However, one major difference between our situation and the one in *loc. cit.* is that the Siegel modular variety is non-compact. As a remedy, we apply the theory of (pro-)Kummer étale topology on log adic spaces developed in [Dia+19a] and [Dia+19b] to handle the boundaries of the compactifications. (See §A.1 for a brief review.)

5.1 The Kummer étale and the pro-Kummer étale cohomology groups

Recall from §2.2 that $\overline{\mathcal{X}}_{\Gamma(p^n)}$, $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$, and $\overline{\mathcal{X}}$ are endowed with the divisorial log structures defined by the boundary divisors. The corresponding sheaves of monoids are denoted by \mathcal{M}_n , $\mathcal{M}_{\mathrm{Iw}^+}$, and \mathcal{M} , respectively.

In what follows, we shall construct a sheaf $\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r$ on the pro-Kummer étale site $\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prok\acute{e}t}}$ which computes the overconvergent cohomology groups introduced in §4.2.

Consider the natural morphism of sites

$$j_{\mathrm{k\acute{e}t}} : \mathcal{X}_{\mathrm{Iw}^+, \mathrm{\acute{e}t}} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{k\acute{e}t}}.$$

Recall that, for every small weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ and any integer $r \geq 1 + r_{\mathcal{U}}$, there is an inverse system of étale locally constant sheaves $(\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ})_{j \in \mathbf{Z}_{\geq 1}}$ on $\mathcal{X}_{\mathrm{Iw}^+, \mathrm{\acute{e}t}}$. Applying [Dia+19a, Corollary 4.6.7], we obtain an isomorphism

$$\varprojlim_j H_{\mathrm{\acute{e}t}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ}) \cong \varprojlim_j H_{\mathrm{k\acute{e}t}}^t(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ})$$

for every $t \in \mathbf{Z}_{\geq 0}$. Write

$$H_{\mathrm{k\acute{e}t}}^t(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r) := \varprojlim_j H_{\mathrm{k\acute{e}t}}^t(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ})[1/p].$$

By Proposition 4.2.2, we arrive at isomorphisms

$$H_{\mathrm{k\acute{e}t}}^t(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r) \cong H_{\mathrm{\acute{e}t}}^t(\mathcal{X}_{\mathrm{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r) \cong H^t(X_{\mathrm{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})).$$

To simplify the notation, we introduce the following abbreviations.

Definition 5.1.1. *Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a small weight and let $r \geq 1 + r_{\mathcal{U}}$. We set*

$$\begin{aligned} \mathfrak{W}_{\kappa_{\mathcal{U}}}^{r, \circ} &:= \varprojlim_j H_{\mathrm{k\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ}), \\ \mathfrak{W}_{\kappa_{\mathcal{U}}}^r &:= \mathfrak{W}_{\kappa_{\mathcal{U}}}^{r, \circ}[\frac{1}{p}] = H_{\mathrm{k\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r), \\ \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathcal{O}_{\mathbf{C}_p}}^{r, \circ} &:= \varprojlim_j (H_{\mathrm{k\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ}) \otimes_{\mathbf{Z}_p} \mathcal{O}_{\mathbf{C}_p}), \\ \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r &:= \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathcal{O}_{\mathbf{C}_p}}^{r, \circ}[\frac{1}{p}]. \end{aligned}$$

where $n_0 = \dim_{\mathbf{C}_p} \mathcal{X}_{\mathrm{Iw}^+}$.

Let

$$\nu : \overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prok\acute{e}t}} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{k\acute{e}t}}$$

be the natural projection of sites. Consider the sheaf $\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r$ on the pro-Kummer étale site $\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prok\acute{e}t}}$ defined by

$$\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r := \left(\varprojlim_j \left(\nu^{-1} j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prok\acute{e}t}}}^+ \right) \right) [\frac{1}{p}].$$

Proposition 5.1.2. *There is a $G_{\mathbf{Q}_p}$ -equivariant isomorphism*

$$\mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r \cong H_{\mathrm{prok\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r).$$

Proof. By [Dia+19a, Theorem 6.2.1 & Corollary 6.3.4], there is an almost isomorphism

$$(H_{\mathrm{k\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ}) \otimes_{\mathbf{Z}_p} \mathcal{O}_{\mathbf{C}_p})^a \cong H_{\mathrm{prok\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \nu^{-1} j_{\mathrm{k\acute{e}t}, *}\mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prok\acute{e}t}}}^+)^a.$$

It remains to establish an almost isomorphism

$$\lim_{\leftarrow j} H_{\text{prokét}}^{n_0} \left(\overline{\mathcal{X}}_{\text{Iw}^+}, \nu^{-1} j_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}^+ \right)^a \cong H_{\text{prokét}}^{n_0} \left(\overline{\mathcal{X}}_{\text{Iw}^+}, \lim_{\leftarrow j} \left(\nu^{-1} j_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}^+ \right) \right)^a.$$

Indeed, observe that the higher inverse limit $R^i \lim_{\leftarrow j} \left(\nu^{-1} j_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}^+ \right)$ almost vanishes for $i \geq 1$ by an almost version of [Sch13, Lemma 3.18] and [Dia+19a, Proposition 6.1.11]. This then allows us to commute the inverse limit with taking cohomology, hence the result. \square

Thanks to Proposition 5.1.2, $H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r)$ inherits actions of the Hecke operators T_γ and $U_{p,i}$ from $H_{\text{ét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}}^r)$. On the other hand, thanks to the $\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}$ -module structure on $\mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r$, there is an alternative way to define the Hecke operators T_γ 's using correspondences. More precisely, for any prime number $\ell \nmid Np$ and any $\gamma \in \text{GSp}_{2g}(\mathbf{Q}_\ell) \cap M_{2g}(\mathbf{Z}_\ell)$, consider the correspondence

$$\begin{array}{ccc} & \mathcal{X}_{\gamma, \text{Iw}^+} & \\ \text{pr}_1 \swarrow & & \searrow \text{pr}_2 \\ \mathcal{X}_{\text{Iw}^+} & & \mathcal{X}_{\text{Iw}^+} \end{array},$$

studied in §3.2. Similar to the construction in §3.2, one obtains an isomorphism

$$\varphi_\gamma : \text{pr}_2^* \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}} \xrightarrow{\sim} \text{pr}_1^* \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}}.$$

Consider the composition

$$\begin{array}{ccc} T'_\gamma : & H_{\text{proét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}}) & \xrightarrow{\text{pr}_2^*} H_{\text{proét}}^{n_0}(\mathcal{X}_{\gamma, \text{Iw}^+}, \text{pr}_2^* \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}}) \\ & \searrow \varphi_\gamma & \nearrow \\ & H_{\text{proét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \text{pr}_1^* \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}}) & \xrightarrow{\text{Tr}_{\text{pr}_1}} H_{\text{proét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}}) \end{array}.$$

However, since $H_{\text{ét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}) \cong H_{\text{két}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, j_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ})$ for every j , we have an identification

$$H_{\text{proét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r |_{\mathcal{X}_{\text{Iw}^+}}) \cong H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r)$$

and hence an operator T'_γ on $H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r)$. One checks that T_γ coincides with T'_γ .

5.2 The overconvergent Eichler–Shimura morphism

In this subsection, we construct the overconvergent Eichler–Shimura morphism by first constructing a morphism between sheaves on the pro-Kummer étale site $\overline{\mathcal{X}}_{\text{Iw}^+, w, \text{prokét}}$.

Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a small weight and let $w \geq r \geq 1 + r_{\mathcal{U}}$. Recall that we have defined a sheaf $\mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r$ on the pro-Kummer étale site $\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}$ in §5.1. The following lemma is an analogue of [CHJ17, Lemma 4.5].

Lemma 5.2.1. *Let $\mathcal{V} = \varprojlim_n \mathcal{V}_n \rightarrow \overline{\mathcal{X}}_{\text{Iw}^+}$ be a pro-Kummer étale presentation of a log affinoid perfectoid object in $\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}$. Let $\mathcal{V}_\infty := \mathcal{V} \times_{\overline{\mathcal{X}}_{\text{Iw}^+}} \overline{\mathcal{X}}_{\Gamma(p^\infty)}$. (Here we have abused the notation and identify $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ with the object $\varprojlim_n \overline{\mathcal{X}}_{\Gamma(p^n)}$ in $\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}$.) Then there is a natural isomorphism*

$$\mathcal{O}_{\mathcal{D}_{\kappa_{\mathcal{U}}}}^r(\mathcal{V}) \cong \left(D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \widehat{\otimes}_{\mathbf{Z}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}(\mathcal{V}_\infty) \right)^{\text{Iw}^+_{\text{GSp}_{2g}}}.$$

Proof. Recall that $\mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}$ is the locally constant sheaf on $\mathcal{X}_{\mathrm{Iw}^+, \text{ét}}$ induced by

$$\pi_1^{\text{ét}}(\mathcal{X}_{\mathrm{Iw}^+}) \rightarrow \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ \rightarrow \mathrm{Aut}(D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})).$$

Since $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ is a profinite Galois cover of $\overline{\mathcal{X}}_{\mathrm{Iw}^+}$ with Galois group $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$, one sees that $\nu^{-1} \mathcal{I}_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}$ becomes the constant local system associated with $D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}})$ after restricting to the localised site $\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}} / \overline{\mathcal{X}}_{\Gamma(p^\infty)}$.

Applying [Dia+19a, Theorem 5.4.3], we obtain an almost isomorphism

$$\left(D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \otimes_{\mathbf{Z}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ (\mathcal{V}_\infty) \right)^a \cong \left(\left(\nu^{-1} \mathcal{I}_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ \right) (\mathcal{V}_\infty) \right)^a.$$

By taking $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -invariants, we obtain almost isomorphisms

$$\begin{aligned} \left(\left(D_{\kappa_{\mathcal{U}},j}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \otimes_{\mathbf{Z}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ (\mathcal{V}_\infty) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+} \right)^a &\cong \left(\left(\left(\nu^{-1} \mathcal{I}_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ \right) (\mathcal{V}_\infty) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+} \right)^a \\ &= \left(\left(\nu^{-1} \mathcal{I}_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ \right) (\mathcal{V}) \right)^a. \end{aligned}$$

Finally, taking inverse limit over j and inverting p , we conclude that

$$\begin{aligned} \mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r(\mathcal{V}) &= \left(\varprojlim_j \left(\nu^{-1} \mathcal{I}_{\text{két},*} \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ \right) (\mathcal{V}) \right) \left[\frac{1}{p} \right] \\ &\cong \left(D_{\kappa_{\mathcal{U}}}^{r,\circ}(\mathbf{T}_0, R_{\mathcal{U}}) \widehat{\otimes}_{\mathbf{Z}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \text{prokét}}}^+ (\mathcal{V}_\infty) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+}. \end{aligned}$$

□

To deal with the overconvergent automorphic sheaves, we recall the Kummer étale sheaves $\underline{\omega}_{w, \text{két}}^{\kappa_{\mathcal{U}}, +}$ and $\underline{\omega}_{w, \text{két}}^{\kappa_{\mathcal{U}}}$ associated with $\underline{\omega}_w^{\kappa_{\mathcal{U}}, +}$ and $\underline{\omega}_w^{\kappa_{\mathcal{U}}}$ defined by the end of §3.3. Then we consider the p -adically completed pullback of them to the pro-Kummer étale site; namely,

$$\widehat{\underline{\omega}}_w^{\kappa_{\mathcal{U}}, +} := \varprojlim_m \left(\underline{\omega}_{w, \text{két}}^{\kappa_{\mathcal{U}}, +} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \text{két}}}^+} \mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \text{prokét}}}^+ / p^m \right)$$

and

$$\widehat{\underline{\omega}}_w^{\kappa_{\mathcal{U}}} := \widehat{\underline{\omega}}_w^{\kappa_{\mathcal{U}}, +} \left[\frac{1}{p} \right].$$

Lemma 5.2.2. *There is a canonical $G_{\mathbf{Q}_p}$ -equivariant morphism*

$$H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \widehat{\underline{\omega}}_w^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}+g+1})(-n_0).$$

Proof. By the discussion at the end of §3.3, we have seen that $\underline{\omega}_{w, \text{két}}^{\kappa_{\mathcal{U}}}$ can be identified with the sheaf of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+ / \Gamma(p^n)$ -invariants of an admissible Kummer étale Banach sheaf of $\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \text{két}}} \widehat{\otimes} R_{\mathcal{U}}$ -modules. Corollary A.3.14 then yields a canonical isomorphism

$$\underline{\omega}_{w, \text{két}}^{\kappa_{\mathcal{U}}} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \text{két}}}} R^i \nu_* \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \text{prokét}}} \xrightarrow{\sim} R^i \nu_* \widehat{\underline{\omega}}_w^{\kappa_{\mathcal{U}}}$$

for every $i \in \mathbf{Z}_{\geq 0}$. On the other hand, by Proposition A.2.3, we have a canonical isomorphism

$$R^i \nu_* \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{prok\acute{e}t}}} \cong \Omega_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}^{\log, i}(-i).$$

Combining the two isomorphisms, we obtain

$$R^i \nu_* \widehat{\omega}_w^{\kappa_{\mathcal{U}}} \cong \omega_{w, \mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}} \Omega_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}^{\log, i}(-i).$$

Moreover, there is a Leray spectral sequence

$$E_2^{j, i} = H_{\mathrm{k\acute{e}t}}^j(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, R^i \nu_* \widehat{\omega}_w^{\kappa_{\mathcal{U}}}) \Rightarrow H_{\mathrm{prok\acute{e}t}}^{j+i}(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \widehat{\omega}_w^{\kappa_{\mathcal{U}}}).$$

The edge map yields a Galois-equivariant morphism

$$H_{\mathrm{prok\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \widehat{\omega}_w^{\kappa_{\mathcal{U}}}) \rightarrow H_{\mathrm{k\acute{e}t}}^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, R^{n_0} \nu_* \widehat{\omega}_w^{\kappa_{\mathcal{U}}}) \cong H_{\mathrm{k\acute{e}t}}^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \omega_{w, \mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}} \Omega_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}^{\log, n_0})(-n_0).$$

Finally, let $\pi_{\mathrm{Iw}^+} : \mathcal{G}_{\mathrm{Iw}^+, w}^{\mathrm{univ}} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$ denote the universal semiabelian variety over $\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}$ and let

$$\omega_{\mathrm{Iw}^+, w} := \pi_{\mathrm{Iw}^+,*} \Omega_{\mathcal{G}_{\mathrm{Iw}^+, w}^{\mathrm{univ}}}^1 / \overline{\mathcal{X}}_{\mathrm{Iw}^+, w}.$$

Note that $\omega_{\mathrm{Iw}^+, w}$ agrees with $\omega_{\mathrm{Iw}^+}^k|_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}}$ studied in §3.4 for $k = (1, 0, \dots, 0)$. The Kodaira–Spencer isomorphism [Lan12, Theorem 1.41 (4)] yields an isomorphism

$$\mathrm{Sym}^2 \omega_{\mathrm{Iw}^+, w} \cong \Omega_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}}^{\log, 1}.$$

Hence,

$$\Omega_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}}^{\log, n_0} \cong \bigwedge^{n_0} (\mathrm{Sym}^2 \omega_{\mathrm{Iw}^+, w}) = \omega_{\mathrm{Iw}^+, w}^{g+1} \subset \omega_w^{g+1}$$

where the last inclusion follows from Lemma 3.4.4. We obtain an injection

$$H_{\mathrm{k\acute{e}t}}^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \omega_{w, \mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}} \Omega_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{k\acute{e}t}}}^{\log, n_0})(-n_0) \hookrightarrow H_{\mathrm{k\acute{e}t}}^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \omega_{w, \mathrm{k\acute{e}t}}^{\kappa_{\mathcal{U}}+g+1})(-n_0) = H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \omega_w^{\kappa_{\mathcal{U}}+g+1})(-n_0).$$

□

For any matrix $\sigma \in M_g(\mathcal{O}_{\mathbf{C}_p})$ and $\mu \in D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$, we define a function $f_{\mu, \sigma} \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}})$ as follows. For any $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$, we define

$$f_{\mu, \sigma}(\gamma') := \int_{(\gamma, v) \in \mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}({}^t \gamma'(\gamma + \sigma v)) \, d\mu,$$

where $e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}$ sends a matrix $X = (X_{ij})_{1 \leq i, j \leq g}$ in $\mathrm{Iw}_{\mathrm{GL}_g}^{(w)}$ to

$$e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(X) = \frac{\kappa_{\mathcal{U}, 1}(X_{11})}{\kappa_{\mathcal{U}, 2}(X_{11})} \times \frac{\kappa_{\mathcal{U}, 2}(\det((X_{ij})_{1 \leq i, j \leq 2}))}{\kappa_{\mathcal{U}, 3}(\det((X_{ij})_{1 \leq i, j \leq 2}))} \times \dots \times \kappa_{\mathcal{U}, g}(\det(X)).$$

The following lemma justifies this definition.

Lemma 5.2.3. (i) For every $\sigma \in M_g(\mathcal{O}_{\mathbf{C}_p})$ and $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$, the assignment

$$(\gamma, v) \mapsto e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}({}^t \gamma'(\gamma + \sigma v))$$

defines an element in $A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$.

(ii) For every $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$ and $\beta \in B_{\mathrm{GL}_g, 0}$, we have

$$f_{\mu, \sigma}(\gamma' \beta) = \kappa_{\mathcal{U}}(\beta) f_{\mu, \sigma}(\gamma').$$

Proof. This is straightforward. \square

Remark 5.2.4. The function $e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}$ is an analogue of the highest weight vector in an algebraic representation of GL_g . Moreover, $e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}$ has the following alternative interpretation: for every $\gamma \in \mathrm{Iw}_{\mathrm{GL}_g}^{(w)}$, if we write $\gamma = \nu \tau \nu'$ with $\nu \in U_{\mathrm{GL}_g, 1}^{\mathrm{opp}, (w)}$, $\tau \in T_{\mathrm{GL}_g, 0}^{(w)}$, and $\nu' \in U_{\mathrm{GL}_g, 0}^{(w)}$, then we have $e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\gamma) = \kappa_{\mathcal{U}}(\tau)$.

We are ready to construct the desired morphism $\eta_{\kappa_{\mathcal{U}}} : \mathcal{O}_{\kappa_{\mathcal{U}}}^r \rightarrow \widehat{\mathcal{W}}_w^{\kappa_{\mathcal{U}}}$ between sheaves on the pro-Kummer étale site $\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{prokét}}$. Indeed, it suffices to construct a map $\mathcal{O}_{\kappa_{\mathcal{U}}}^r(\mathcal{V}) \rightarrow \widehat{\mathcal{W}}_w^{\kappa_{\mathcal{U}}}(\mathcal{V})$ for every log affinoid perfectoid object \mathcal{V} in $\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}$. By Lemma 5.2.1, we have

$$\mathcal{O}_{\kappa_{\mathcal{U}}}^r(\mathcal{V}) \simeq \left(D_{\kappa_{\mathcal{U}}}^{r, \circ}(\mathbf{T}_0, R_{\mathcal{U}}) \widehat{\otimes}_{\mathbf{Z}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_{\infty}) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+}$$

where $\mathcal{V}_{\infty} := \mathcal{V} \times_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}} \overline{\mathcal{X}}_{\Gamma(p^{\infty})}$.

On the other hand, by definition, we know that $\widehat{\mathcal{W}}_w^{\kappa_{\mathcal{U}}}(\mathcal{V})$ consists of $f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_{\infty}) \widehat{\otimes} R_{\mathcal{U}})$ satisfying $\alpha^* f = \rho_{\kappa_{\mathcal{U}}}(\alpha_a + \mathfrak{z} \alpha_c)^{-1} f$, for all $\alpha = \begin{pmatrix} \alpha_a & \alpha_b \\ \alpha_c & \alpha_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$. This is equivalent to saying that $\widehat{\mathcal{W}}_w^{\kappa_{\mathcal{U}}}(\mathcal{V})$ consists of $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -invariant elements $f \in C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_{\infty}) \widehat{\otimes} R_{\mathcal{U}})$ with respect to the twisted $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -action

$$\alpha \cdot f := \rho_{\kappa_{\mathcal{U}}}(\alpha_a + \mathfrak{z} \alpha_c)(\alpha^* f).$$

Consider the map

$$D_{\kappa_{\mathcal{U}}}^{r, \circ}(\mathbf{T}_0, R_{\mathcal{U}}) \widehat{\otimes}_{\mathbf{Z}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_{\infty}) \rightarrow C_{\kappa_{\mathcal{U}}}^{w-\mathrm{an}}(\mathrm{Iw}_{\mathrm{GL}_g}, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_{\infty}) \widehat{\otimes} R_{\mathcal{U}}), \quad \mu \otimes \delta \mapsto \delta f_{\mu, \mathfrak{z}}.$$

We claim that this map is $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -equivariant, and hence taking the $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -invariants yields the desired map $\mathcal{O}_{\kappa_{\mathcal{U}}}^r(\mathcal{V}) \rightarrow \widehat{\mathcal{W}}_w^{\kappa_{\mathcal{U}}}(\mathcal{V})$. Indeed, for any $\alpha = \begin{pmatrix} \alpha_a & \alpha_b \\ \alpha_c & \alpha_d \end{pmatrix} \in \mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ and any $\gamma' \in \mathrm{Iw}_{\mathrm{GL}_g}$, we have

$$\begin{aligned} (\alpha^* \delta) f_{\alpha \cdot \mu, \mathfrak{z}}(\gamma') &= (\alpha^* \delta) \left(\int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\mathfrak{t} \gamma'(\gamma + \mathfrak{z} v)) \, d\alpha \cdot \mu \right) \\ &= (\alpha^* \delta) \left(\int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\mathfrak{t} \gamma'((\alpha_a \gamma + \alpha_b v) + \mathfrak{z}(\alpha_c \gamma + \alpha_d v))) \, d\mu \right) \\ &= (\alpha^* \delta) \left(\int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\mathfrak{t} \gamma'((\alpha_a + \mathfrak{z} \alpha_c) \gamma + (\alpha_b + \mathfrak{z} \alpha_d) v)) \, d\mu \right) \\ &= (\alpha^* \delta) \left(\int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\mathfrak{t} \gamma'(\alpha_a + \mathfrak{z} \alpha_c)(\gamma + (\alpha_a + \mathfrak{z} \alpha_c)^{-1}(\alpha_b + \mathfrak{z} \alpha_d) v)) \, d\mu \right) \\ &= (\alpha^* \delta) \left(\int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\mathfrak{t}(\mathfrak{t}(\alpha_a + \mathfrak{z} \alpha_c) \gamma')(\gamma + (\mathfrak{z} \cdot \alpha) v)) \, d\mu \right) \\ &= (\alpha^* \delta) \left(\rho_{\kappa_{\mathcal{U}}}(\alpha_a + \mathfrak{z} \alpha_c) \int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\mathrm{hst}}(\mathfrak{t} \gamma'(\gamma + (\mathfrak{z} \cdot \alpha) v)) \, d\mu \right) \\ &= \alpha \cdot (\delta f_{\mu, \mathfrak{z}})(\gamma') \end{aligned}$$

as desired.

Putting everything together, we consider the composition

$$\begin{aligned} \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r &\cong H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r) \xrightarrow{\text{Res}} H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^r) \\ &\xrightarrow{\eta_{\kappa_{\mathcal{U}}}} H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}}) \rightarrow H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}+g+1})(-n_0) = M_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}}+g+1}(-n_0) \end{aligned}$$

where the second last morphism is given by Lemma 5.2.2. We arrive at the *overconvergent Eichler–Shimura morphism*

$$\text{ES}_{\kappa_{\mathcal{U}}} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r \rightarrow M_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}}+g+1}(-n_0).$$

Proposition 5.2.5. *The overconvergent Eichler–Shimura morphism*

$$\text{ES}_{\kappa_{\mathcal{U}}} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r \rightarrow M_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}}+g+1}(-n_0)$$

is Hecke- and $G_{\mathbf{Q}_p}$ -equivariant.

Proof. The Galois-equivariance follows immediately from Lemma 5.2.2. For Hecke operators away from Np , notice that the operators T_{γ} 's on both sides are defined in the same way using correspondences. Hence, it is straightforward to verify the T_{γ} -equivariances. It remains to check the $U_{p,i}$ -equivariance for all $i = 1, \dots, g$.

To this end, due to the $\text{Iw}_{\text{GSP}_{2g}}^+$ -equivariance of $\eta_{\kappa_{\mathcal{U}}}$, we only have to check the $\mathbf{u}_{p,i}$ -equivariance. Indeed, for every $\gamma' = \gamma'_0 \beta'_0 \in \text{Iw}_{\text{GL}_g}$ with $\gamma'_0 \in U_{\text{GL}_{g,1}}^{\text{opp}}$ and $\beta'_0 \in B_{\text{GL}_g,0}$, we have

$$\begin{aligned} (\mathbf{u}_{p,i}^* \delta) f_{\mathbf{u}_{p,i} \cdot \mu, \mathfrak{z}}(\gamma') &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathfrak{t} \gamma'_0(\gamma + \mathfrak{z} \mathbf{v})) \, d\mathbf{u}_{p,i} \cdot \mu \right) \\ &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} \kappa_{\mathcal{U}}(\beta) e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathfrak{t} \gamma'_0(\gamma_0 + \mathfrak{z} \mathbf{v}_0)) \, d\mathbf{u}_{p,i} \cdot \mu \right) \\ &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} \kappa_{\mathcal{U}}(\beta) e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathfrak{t} \gamma'_0(\mathbf{u}_{p,i}^{\square} \gamma_0 \mathbf{u}_{p,i}^{\square,-1} + \mathfrak{z} \mathbf{u}_{p,i}^{\blacksquare} \mathbf{v}_0 \mathbf{u}_{p,i}^{\square,-1})) \, d\mu \right) \\ &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} \kappa_{\mathcal{U}}(\beta) e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathfrak{t} \gamma'_0 \mathbf{u}_{p,i}^{\square}(\gamma_0 + \mathbf{u}_{p,i}^{\square,-1} \mathfrak{z} \mathbf{u}_{p,i}^{\blacksquare} \mathbf{v}_0) \mathbf{u}_{p,i}^{\square,-1}) \, d\mu \right) \\ &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} \kappa_{\mathcal{U}}(\beta) e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathfrak{t} \gamma'_0 \mathbf{u}_{p,i}^{\square}(\gamma_0 + (\mathfrak{z} \cdot \mathbf{u}_{p,i}) \mathbf{v}_0) \mathbf{u}_{p,i}^{\square,-1}) \, d\mu \right) \\ &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} \kappa_{\mathcal{U}}(\beta) e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathbf{u}_{p,i}^{\square,-1} \mathfrak{t} \gamma'_0 \mathbf{u}_{p,i}^{\square}(\gamma_0 + (\mathfrak{z} \cdot \mathbf{u}_{p,i}) \mathbf{v}_0)) \, d\mu \right) \\ &= (\mathbf{u}_{p,i}^* \delta) \left(\kappa_{\mathcal{U}}(\beta'_0) \int_{\mathbf{T}_0} \kappa_{\mathcal{U}}(\beta) e_{\kappa_{\mathcal{U}}}^{\text{hst}}(\mathfrak{t}(\mathbf{u}_{p,i}^{\square} \gamma'_0 \mathbf{u}_{p,i}^{\square,-1})(\gamma_0 + (\mathfrak{z} \cdot \mathbf{u}_{p,i}) \mathbf{v}_0)) \, d\mu \right) \\ &= \mathbf{u}_{p,i} \cdot (\delta f_{\mu, \mathfrak{z}}), \end{aligned}$$

where we have written $(\gamma, \mathbf{v}) = (\gamma_0, \mathbf{v}_0) \beta$ for $(\gamma_0, \mathbf{v}_0) \in \mathbf{T}_{00}$ and $\beta \in B_{\text{GL}_g,0}$. The antepenultimate equation follows from the property of matrix determinants. \square

Remark 5.2.6. There is an analogue for compactly supported cohomology groups and overconvergent cuspforms. Let r , w , and $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be the same as before. On one hand, consider

$$\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^{r, \text{cusp}} := \left(\varprojlim_j \left(\nu^{-1} j_{\text{két},!} \mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ} \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}^+ \right) \right) \left[\frac{1}{p} \right].$$

Since

$$H_{\text{két}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, j_{\text{két},!} \mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ}) = H_{\text{ét}, c}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}}, j}^{r, \circ}),$$

an analogue of Proposition 5.1.2 implies that $\mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^{r,\text{cusp}}$ computes

$$\mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r,c} := \left(\varprojlim_j H_{\text{ét},c}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathcal{D}_{\kappa_{\mathcal{U}},j}^{r,\circ}) \otimes_{\mathbf{Z}_p} \mathcal{O}_{\mathbf{C}_p} \right) \left[\frac{1}{p} \right].$$

On the other hand, recall the sheaf $\underline{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}}$ of w -overconvergent Siegel cuspforms of weight $\kappa_{\mathcal{U}}$ and consider the p -adically completed pullback $\widehat{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}}$ to the pro-Kummer étale site. Repeating the construction above, we obtain a morphism $\eta_{\kappa_{\mathcal{U}}}^{\text{cusp}} : \mathcal{O}\mathcal{D}_{\kappa_{\mathcal{U}}}^{r,\text{cusp}} \rightarrow \widehat{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}}$ which induces a morphism

$$\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r,c} \rightarrow H^0(\overline{\mathcal{X}}_{\text{Iw}^+,w}, \underline{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}+g+1})(-n_0)$$

rendering the following Galois- and Hecke-equivariant diagram commutative:

$$\begin{array}{ccc} \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r & \xrightarrow{\text{ES}_{\kappa_{\mathcal{U}}}} & H^0(\overline{\mathcal{X}}_{\text{Iw}^+,w}, \underline{\omega}_w^{\kappa_{\mathcal{U}}+g+1})(-n_0) \\ \uparrow & & \uparrow \\ \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r,c} & \xrightarrow{\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}}} & H^0(\overline{\mathcal{X}}_{\text{Iw}^+,w}, \underline{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}+g+1})(-n_0) \end{array},$$

where the vertical arrow on the left is the natural map from the compactly supported cohomology group to the usual cohomology group. Let

$$\mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r,\text{cusp}} := \text{image} \left(\mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r,c} \rightarrow \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^r \right).$$

We arrive at the *overconvergent Eichler–Shimura morphism for overconvergent Siegel cuspforms*

$$\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r,\text{cusp}} \rightarrow S_{\text{Iw}^+,w}^{\kappa_{\mathcal{U}}+g+1}(-n_0),$$

where

$$S_{\text{Iw}^+,w}^{\kappa_{\mathcal{U}}+g+1} := H^0(\overline{\mathcal{X}}_{\text{Iw}^+,w}, \underline{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}+g+1})$$

is the space of w -overconvergent Siegel cuspforms of strict Iwahori level and weight $\kappa_{\mathcal{U}} + g + 1$.

Lastly, we point out that, by construction, both $\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}}$ and $\text{ES}_{\kappa_{\mathcal{U}}}$ are functorial in the small weights $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$.

5.3 The image of the overconvergent Eichler–Shimura morphism at classical weights

The aim of this last part of the section is to describe the image of the overconvergent Eichler–Shimura morphism at classical algebraic weights. Let $k = (k_1, \dots, k_g) \in \mathbf{Z}_{\geq 0}^g$ be a dominant weight. Note that the character group of $T_{\text{GSp}_{2g}}$ is isomorphic to \mathbf{Z}^{g+1} through

$$\mathbf{Z}^{g+1} \times T_{\text{GSp}_{2g}} \rightarrow \mathbb{G}_m, \quad ((k_1, \dots, k_g; k_0), \text{diag}(\tau_1, \dots, \tau_g, \tau_0 \tau_g^{-1}, \dots, \tau_0 \tau_1^{-1})) \mapsto \prod_{i=0}^g \tau_i^{k_i}.$$

We view $k = (k_1, \dots, k_g) \in \mathbf{Z}^g$ as a character of $T_{\text{GSp}_{2g}}$ via

$$\mathbf{Z}^g \hookrightarrow \mathbf{Z}^{g+1}, \quad (k_1, \dots, k_g) \mapsto (k_1, \dots, k_g; 0).$$

We first introduce some algebraic representations. Consider the algebraic representation

$$\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}} := \left\{ \phi : \mathrm{GSp}_{2g} \rightarrow \mathbb{A}^1 : \begin{array}{l} \phi \text{ is a morphism of algebraic varieties over } \mathbf{Q}_p \text{ such that} \\ \phi(\gamma\beta) = k(\beta)\phi(\gamma), \forall \gamma \in \mathrm{GSp}_{2g}, \beta \in B_{\mathrm{GSp}_{2g}} \end{array} \right\}$$

equipped with a left GSp_{2g} -action given by

$$(\gamma \cdot \phi)(\gamma') = \phi({}^t\gamma \gamma')$$

for any $\gamma, \gamma' \in \mathrm{GSp}_{2g}$ and $\phi \in \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$.

Let $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee}$ be the linear dual of $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$. We equip $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee}$ with a left GSp_{2g} -action induced from the following right GSp_{2g} -action on $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$:

$$(\phi \cdot \gamma)(\gamma') = \phi(\gamma \gamma')$$

for any $\gamma, \gamma' \in \mathrm{GSp}_{2g}$ and $\phi \in \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$. Observe that there is a natural morphism of GSp_{2g} -representations

$$\beta : \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee} \rightarrow \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}, \quad \mu \mapsto \left(\gamma \mapsto \int_{\alpha \in \mathrm{GSp}_{2g}} e_k^{\mathrm{hst}}({}^t\gamma \alpha) \, d\mu \right),$$

where e_k^{hst} is the “highest weight vector” inside $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$; namely, for any matrix $X = (X_{ij})_{1 \leq i,j \leq 2g} \in \mathrm{GSp}_{2g}$, we define

$$e_k^{\mathrm{hst}}(X) = X_{11}^{k_1 - k_2} \times \det((X_{ij})_{1 \leq i,j \leq 2})^{k_2 - k_3} \times \dots \times \det((X_{ij})_{1 \leq i,j \leq g})^{k_g}.$$

Secondly, we consider the cohomology groups induced by these algebraic representations. Notice that the left GSp_{2g} -actions on $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$ and $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee}$ induce étale \mathbf{Q}_p -local systems on $\mathcal{X}_{\mathrm{Iw}^+}$ which we still denote by the same symbols. In particular, we can consider the cohomology group $H_{\mathrm{ét}}^{n_0}(\mathcal{X}_{\mathrm{Iw}^+}, \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee})$. Similar to §5.2, we introduce the sheaves $\mathcal{O}\mathcal{V}_k$ and $\mathcal{O}\mathcal{V}_k^\vee$ on $\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}$ defined by

$$\mathcal{O}\mathcal{V}_k := \nu^{-1} j_{\mathrm{két},*} \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}} \otimes_{\mathbf{Q}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}$$

and

$$\mathcal{O}\mathcal{V}_k^\vee := \nu^{-1} j_{\mathrm{két},*} \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee} \otimes_{\mathbf{Q}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}.$$

By the same argument as in Proposition 5.1.2, we obtain a natural identification

$$H_{\mathrm{ét}}^{n_0}(\mathcal{X}_{\mathrm{Iw}^+}, \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee}) \otimes_{\mathbf{Q}_p} \mathbf{C}_p \cong H_{\mathrm{prokét}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{O}\mathcal{V}_k^\vee).$$

Moreover, if $\mathcal{V} = \varprojlim_n \mathcal{V}_n \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+}$ is a pro-Kummer étale presentation of a log affinoid perfectoid object in $\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}$ and let $\mathcal{V}_\infty := \mathcal{V} \times_{\overline{\mathcal{X}}_{\mathrm{Iw}^+}} \overline{\mathcal{X}}_{\Gamma(p^\infty)}$, then, following the same argument as in the proof of Lemma 5.2.1, we obtain identifications

$$\mathcal{O}\mathcal{V}_k(\mathcal{V}) = \left(\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}} \otimes_{\mathbf{Q}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_\infty) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+}$$

and

$$\mathcal{O}\mathcal{V}_k^\vee(\mathcal{V}) = \left(\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee} \otimes_{\mathbf{Q}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, \mathrm{prokét}}}(\mathcal{V}_\infty) \right)^{\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+}.$$

Remark 5.3.1. There are naturally defined Hecke operators on $H_{\text{ét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee})$. More precisely, similar to Proposition 4.2.2, we have

$$H_{\text{ét}}^{n_0}(\mathcal{X}_{\text{Iw}^+}, \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee}) \cong H^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee}),$$

where the right-hand side is the Betti cohomology of $X_{\text{Iw}^+}(\mathbf{C})$ with coefficients in $\mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee}$. Hence, it suffices to define the Hecke operators acting on $H^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee})$. They are defined as follows.

- For any Hecke operator T_γ away from Np , its action on $H^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee})$ is defined by the same formula as (7).
- For the $U_{p,i}$ -action, let $\mathbf{u}_{p,i}$ act on $\text{GSp}_{2g}(\mathbf{Q}_p)$ via conjugation

$$\mathbf{u}_{p,i} \cdot \gamma = \mathbf{u}_{p,i} \gamma \mathbf{u}_{p,i}^{-1}.$$

Observe that if $\gamma \in B_{\text{GSp}_{2g}}(\mathbf{Q}_p)$, then $\mathbf{u}_{p,i} \cdot \gamma \in B_{\text{GSp}_{2g}}(\mathbf{Q}_p)$ and the diagonal entries of γ coincide with the diagonal entries of $\mathbf{u}_{p,i} \cdot \gamma$. This action then induces a left $\mathbf{u}_{p,i}$ -action on $\mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee}$. The operator $U_{p,i}$ acting on $H^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee})$ is defined by the same formula as (8).

We also consider the p -adically completed automorphic sheaf $\underline{\omega}_{\text{Iw}^+}^k$ on $\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}$ defined by

$$\underline{\omega}_{\text{Iw}^+}^k := \varprojlim_m \left(\underline{\omega}_{\text{Iw}^+}^{k,+} \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+}}} \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}}^+ / p^m \right) \left[\frac{1}{p} \right]$$

where $\underline{\omega}_{\text{Iw}^+}^{k,+}$ is defined in Remark 3.4.1. It follows from Proposition 3.4.3 that

$$\underline{\omega}_{\text{Iw}^+}^k(\mathcal{V}) = \left\{ f \in P_k(\text{GL}_g, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\text{Iw}^+, w, \text{prokét}}}(\mathcal{V}_\infty)) : \gamma^* f = \rho_k(\gamma_a + \mathfrak{z} \gamma_c)^{-1} f, \quad \forall \gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \text{Iw}_{\text{GSp}_{2g}}^+ \right\}.$$

for any log affinoid perfectoid object $\mathcal{V} \in \overline{\mathcal{X}}_{\text{Iw}^+, \text{prokét}}$ and $\mathcal{V}_\infty = \mathcal{V} \times_{\overline{\mathcal{X}}_{\text{Iw}^+}} \overline{\mathcal{X}}_{\Gamma(p^\infty)}$.

The algebraic Eichler-Shimura morphism Recall the Hodge–Tate morphism

$$\text{HT}_{\Gamma(p^\infty)} : V_p \rightarrow \underline{\omega}_{\Gamma(p^\infty)}$$

from §B.2. It follows from the definition that

$$\underline{\omega}_{\text{Iw}^+}^k = (\text{Sym}^{k_1-k_2} \underline{\omega}_{\text{Iw}^+}) \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+}}} (\text{Sym}^{k_2-k_3} \wedge^2 \underline{\omega}_{\text{Iw}^+}) \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+}}} \cdots \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+}}} (\text{Sym}^{k_g} \det \underline{\omega}_{\text{Iw}^+})$$

and hence

$$\underline{\omega}_{\Gamma(p^\infty)}^k = (\text{Sym}^{k_1-k_2} \underline{\omega}_{\Gamma(p^\infty)}) \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}} (\text{Sym}^{k_2-k_3} \wedge^2 \underline{\omega}_{\Gamma(p^\infty)}) \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}} \cdots \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}} (\text{Sym}^{k_g} \det \underline{\omega}_{\Gamma(p^\infty)}).$$

Let V_{std} denote the standard representation of GSp_{2g} over \mathbf{Q}_p , with standard basis x_1, \dots, x_{2g} . There is an isomorphism of $\text{GSp}_{2g}(\mathbf{Q}_p)$ -representations $V_{\text{std}} \simeq V_{\mathbf{Q}_p} := V_p \otimes_{\mathbf{Z}_p} \mathbf{Q}_p$ sending x_i to e_{2g+1-i} , for $i = 1, \dots, g$, and sending x_i to $-e_{2g+1-i}$, for $i = g+1, \dots, 2g$. If we write

$$V_{\text{std}}^k := (\text{Sym}^{k_1-k_2} V_{\text{std}}) \otimes_{\mathbf{Q}_p} (\text{Sym}^{k_2-k_3} (\wedge^2 V_{\text{std}})) \otimes_{\mathbf{Q}_p} \cdots \otimes_{\mathbf{Q}_p} (\text{Sym}^{k_g} (\wedge^g V_{\text{std}}))$$

and

$$V_{\mathbf{Q}_p}^k := (\mathrm{Sym}^{k_1-k_2} V_{\mathbf{Q}_p}) \otimes_{\mathbf{Q}_p} (\mathrm{Sym}^{k_2-k_3} (\wedge^2 V_{\mathbf{Q}_p})) \otimes_{\mathbf{Q}_p} \cdots \otimes_{\mathbf{Q}_p} (\mathrm{Sym}^{k_g} (\wedge^g V_{\mathbf{Q}_p})),$$

the Hodge-Tate map induces a map $V_{\mathrm{std}}^k \simeq V_{\mathbf{Q}_p}^k \rightarrow \underline{\omega}_{\mathrm{Iw}^+}^k$. Moreover, it is well-known that $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$ is an irreducible GSp_{2g} -subrepresentation of V_{std}^k (see for example [FH91, Lecture 17]). In particular, the highest weight vector e_k^{hst} in $\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}}$ corresponds to the element

$$x_1^{k_1-k_2} \otimes (x_1 \wedge x_2)^{k_2-k_3} \otimes \cdots \otimes (x_1 \wedge \cdots \wedge x_g)^{k_g}$$

in V_{std}^k .

The composition

$$\mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee} \xrightarrow{\beta} \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}} \hookrightarrow V_{\mathrm{std}}^k \simeq V_{\mathbf{Q}_p}^k \rightarrow \underline{\omega}_{\mathrm{Iw}^+}^k$$

then induces a map

$$\eta_k^{\mathrm{alg}} : \mathcal{O}\mathcal{V}_k^{\vee} \rightarrow \underline{\omega}_{\mathrm{Iw}^+}^k.$$

Eventually, we arrive at the **algebraic Eichler-Shimura morphism of weight k**

$$\mathrm{ES}_k^{\mathrm{alg}} : H_{\mathrm{et}}^{n_0}(\mathcal{X}_{\mathrm{Iw}^+}, \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee}) \otimes_{\mathbf{Q}_p} \mathbf{C}_p \simeq H_{\mathrm{prok\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \mathcal{O}\mathcal{V}_k^{\vee}) \xrightarrow{\eta_k^{\mathrm{alg}}} H_{\mathrm{prok\acute{e}t}}^{n_0}(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \underline{\omega}_{\mathrm{Iw}^+}^k) \rightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \underline{\omega}_{\mathrm{Iw}^+}^{k+g+1})(-n_0),$$

where the last map follows from the same argument as in the proof of Lemma 5.2.2. We remark that $\mathrm{ES}_k^{\mathrm{alg}}$ coincides with the one induced from [FC90, Theorem VI. 6.2]. It is Hecke- and $G_{\mathbf{Q}_p}$ -equivariant, and also surjective.

In fact, over the w -ordinary locus $\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}$, the map η_k^{alg} has the following explicit description.

Lemma 5.3.2. (i) *Let $\mathcal{V} = \varprojlim_n \mathcal{V}_n$ be a pro-Kummer étale presentation of a log affinoid perfectoid object in $\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{prok\acute{e}t}}$ and let $\mathcal{V}_{\infty} = \mathcal{V} \times_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w}} \overline{\mathcal{X}}_{\Gamma(p^{\infty}),w}$. There is a well-defined $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -equivariant map*

$$\widehat{\eta}_k^{\mathrm{alg}} : \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg},\vee} \otimes_{\mathbf{Q}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{prok\acute{e}t}}}(\mathcal{V}_{\infty}) \rightarrow P_k(\mathrm{GL}_g, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{prok\acute{e}t}}}(\mathcal{V}_{\infty}))$$

defined by $\mu \otimes \delta \mapsto \delta f_{\mu,3}^{\mathrm{alg}}$ where

$$f_{\mu,3}^{\mathrm{alg}}(\gamma') = \int_{\alpha \in \mathrm{GSp}_{2g}} e_k^{\mathrm{hst}} \left(\begin{pmatrix} {}^t\gamma' & \\ & \check{\mathbb{I}}_g \gamma'^{-1} \check{\mathbb{I}}_g \end{pmatrix} \begin{pmatrix} \mathbb{1}_g & \mathfrak{z} \\ & \mathbb{1}_g \end{pmatrix} \alpha \right) d\mu.$$

Here, the $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -action on the right hand side is given by

$$\gamma \cdot f := \rho_k(\gamma_a + \mathfrak{z} \gamma_c)(\gamma^* f)$$

for every $\gamma = \begin{pmatrix} \gamma_a & \gamma_b \\ \gamma_c & \gamma_d \end{pmatrix} \in \mathrm{GSp}_{2g}(\mathbf{Q}_p)$ and $f \in P_k(\mathrm{GL}_g, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{prok\acute{e}t}}}(\mathcal{V}_{\infty}))$.

(ii) The map η_k^{alg} is obtained from $\widehat{\eta}_k^{\mathrm{alg}}$ by taking $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -invariants on both sides.

Proof. (i) Notice that $\widehat{\eta}_k^{\mathrm{alg}}$ is the composition of β with the map

$$\xi_k^{\mathrm{alg}} : \mathbf{V}_{\mathrm{GSp}_{2g},k}^{\mathrm{alg}} \otimes_{\mathbf{Q}_p} \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{prok\acute{e}t}}}(\mathcal{V}_{\infty}) \rightarrow P_k(\mathrm{GL}_g, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+,w,\mathrm{prok\acute{e}t}}}(\mathcal{V}_{\infty}))$$

defined by $\phi \otimes \delta \mapsto \delta g_{\phi, \mathfrak{z}}$ where

$$g_{\phi, \mathfrak{z}}(\gamma') = \phi \left(\begin{pmatrix} \mathbb{1}_g & \\ \mathfrak{z} & \mathbb{1}_g \end{pmatrix} \begin{pmatrix} \gamma' & \\ & \mathbb{1}_g \mathfrak{t}(\gamma')^{-1} \mathbb{1}_g \end{pmatrix} \right)$$

for all $\gamma' \in \mathrm{GL}_g(\mathbf{C}_p)$.

Recall that β is $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -equivariant. It remains to check that ξ_k^{alg} is $\mathrm{GSp}_{2g}(\mathbf{Q}_p)$ -equivariant, which follows from a straightforward calculation.

- (ii) It suffices to check that the $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ -invariance of ξ_k^{alg} coincides with the map induced from the composition $\mathbf{V}_{\mathrm{GSp}_{2g}, k}^{\mathrm{alg}} \hookrightarrow V_{\mathrm{std}}^k \simeq V_{\mathbf{Q}_p}^k \rightarrow \underline{\omega}_{\mathrm{Iw}^+}^k$. Notice that $\mathbf{V}_{\mathrm{GSp}_{2g}, k}^{\mathrm{alg}}$ is spanned by GSp_{2g} -translations of the highest weight vector e_k^{hst} . Therefore, we only need to check that $\xi_k^{\mathrm{alg}}(e_k^{\mathrm{hst}} \otimes 1)$ gives the correct element in $\underline{\omega}_{\mathrm{Iw}^+}^k$.

Indeed, since the Hodge–Tate map $V_p \rightarrow \underline{\omega}_{\mathrm{Iw}^+}$ sends e_{2g+1-i} to \mathfrak{s}_i , for $i = 1, \dots, g$, we see that the composition $\mathbf{V}_{\mathrm{GSp}_{2g}, k}^{\mathrm{alg}} \hookrightarrow V_{\mathrm{std}}^k \simeq V_{\mathbf{Q}_p}^k \rightarrow \underline{\omega}_{\mathrm{Iw}^+}^k$ sends the highest weight vector e_k^{hst} to

$$\mathfrak{s}_1^{k_1-k_2} \otimes (\mathfrak{s}_1 \wedge \mathfrak{s}_2)^{k_2-k_3} \otimes \dots \otimes (\mathfrak{s}_1 \wedge \dots \wedge \mathfrak{s}_g)^{k_g}.$$

On the other hand, notice that the element $\mathfrak{s}_1 \wedge \dots \wedge \mathfrak{s}_t$ corresponds to the function $X = (X_{ij})_{1 \leq i, j \leq g} \mapsto \det((X_{ij})_{1 \leq i, j \leq t})$ in $P_k(\mathrm{GL}_g, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{prok\acute{e}t}}}(\mathcal{V}_\infty))$. Therefore, e_k^{hst} is sent to the function

$$X \mapsto X_{11}^{k_1-k_2} \times \det((X_{ij})_{1 \leq i, j \leq 2})^{k_2-k_3} \times \dots \times \det((X_{ij})_{1 \leq i, j \leq g})^{k_g}$$

in $P_k(\mathrm{GL}_g, \widehat{\mathcal{O}}_{\overline{\mathcal{X}}_{\mathrm{Iw}^+, w, \mathrm{prok\acute{e}t}}}(\mathcal{V}_\infty))$. This element coincides with $\xi_k^{\mathrm{alg}}(e_k^{\mathrm{hst}} \otimes 1)$, as desired. \square

Recall the natural inclusion $M_{\mathrm{Iw}^+}^{k, \mathrm{cl}} = H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+}, \underline{\omega}_{\mathrm{Iw}^+}^k) \hookrightarrow H^0(\overline{\mathcal{X}}_{\mathrm{Iw}^+, w}, \underline{\omega}_w^k) = M_{\mathrm{Iw}^+, w}^k$ from Lemma 3.4.4. The main result of this subsection is the following.

Theorem 5.3.3. *Let $k = (k_1, \dots, k_g) \in \mathbf{Z}_{\geq 0}^g$ be a dominant weight. Then the image of*

$$\mathrm{ES}_k : \mathfrak{C}_{k, \mathbf{C}_p}^r \longrightarrow M_{\mathrm{Iw}^+, w}^{k+g+1}(-n_0)$$

is contained in the space of the classical forms $M_{\mathrm{Iw}^+}^{k+g+1, \mathrm{cl}}(-n_0)$.

Proof. Firstly, there is a natural map

$$\mathbf{V}_{\mathrm{GSp}_{2g}, k}^{\mathrm{alg}} \rightarrow A_k^r(\mathbf{T}_0, \mathbf{Q}_p)$$

induced by the inclusion

$$\mathbf{T}_0 \rightarrow \mathrm{GSp}_{2g}(\mathbf{Q}_p), \quad (\gamma, v) \mapsto \begin{pmatrix} \gamma & \\ v & \mathbb{1}_g \mathfrak{t} \gamma^{-1} \mathbb{1}_g \end{pmatrix}.$$

The dual of this map gives

$$D_k^r(\mathbf{T}_0, \mathbf{Q}_p) \rightarrow \mathbf{V}_{\mathrm{GSp}_{2g}, k}^{\mathrm{alg}, \vee}$$

which then induces a map of sheaves

$$\mathcal{O}\mathcal{D}_k^r \rightarrow \mathcal{O}\mathcal{V}_k^\vee$$

over $\overline{\mathcal{X}}_{\text{Iw}^+, w, \text{prokét}}$. Hence, the theorem follows once we show that the following diagram commutes

$$\begin{array}{ccc} H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}\mathcal{D}_k^r) & \xrightarrow{\text{ES}_k} & H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{k+g+1})(-n_0) \\ \downarrow & & \uparrow \\ H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}\mathcal{V}_k^\vee) & \xrightarrow{\text{ES}_k^{\text{alg}}} & H^0(\overline{\mathcal{X}}_{\text{Iw}^+}, \underline{\omega}_{\text{Iw}^+}^{k+g+1})(-n_0) \end{array}.$$

Over $\overline{\mathcal{X}}_{\text{Iw}^+, w, \text{prokét}}$, it follows from the construction that we have a commutative diagram

$$\begin{array}{ccc} \mathcal{O}\mathcal{D}_k^r & \xrightarrow{\eta_k} & \underline{\omega}_w^k \\ \downarrow & & \uparrow \\ \mathcal{O}\mathcal{V}_k^\vee & \xrightarrow{\eta_k^{\text{alg}}} & \underline{\omega}_{\text{Iw}^+}^k \end{array},$$

where the inclusion on the right-hand side is given by the inclusion (6). Consequently, there is a commutative diagram on the cohomology groups

$$\begin{array}{ccccc} H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}\mathcal{D}_k^r) & \longrightarrow & H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+}, \mathcal{O}\mathcal{V}_k^\vee) & & \\ \downarrow \text{Res} & & \downarrow \text{Res} & & \text{ES}_k^{\text{alg}} \\ H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \mathcal{O}\mathcal{D}_k^r) & \longrightarrow & H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \mathcal{O}\mathcal{V}_k^\vee) & & \\ \downarrow \eta_k & & \downarrow \eta_k^{\text{alg}} & & \\ H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^k) & \longleftarrow & H_{\text{prokét}}^{n_0}(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_{\text{Iw}^+}^k) & & \\ \downarrow & & \downarrow & & \downarrow \text{ES}_k \\ H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_w^{k+g+1})(-n_0) & \longleftarrow & H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_{\text{Iw}^+}^{k+g+1})(-n_0) & \xleftarrow{\text{Res}} & H^0(\overline{\mathcal{X}}_{\text{Iw}^+}, \underline{\omega}_{\text{Iw}^+}^k)(-n_0) \end{array}$$

This finishes the proof. \square

6 The overconvergent Eichler–Shimura morphism on the cuspidal eigenvariety

In this section, we glue the overconvergent Eichler–Shimura morphism over a suitable eigenvariety \mathcal{E}_0 . We begin with some preliminaries on the overconvergent cohomology groups in §6.1. The eigenvariety \mathcal{E}_0 is constructed in §6.2. Finally in §6.3, we show that the overconvergent Eichler–Shimura morphism spreads out over \mathcal{E}_0 .

Throughout this section, we assume $p > 2g$ so that we can apply results in [AIP15] via the comparison in §3.7. On the other hand, we believe that the results in this section hold for smaller primes as well. In order to deal with these smaller primes, one would have to reprove several results in [AIP15] in our context; *e.g.*, the classicality result and the fact that $S_{\text{Iw}^+}^{\kappa\mathcal{U}}$ has property (Pr) in the sense of [Buz07]. We decide to leave these generalities to the reader in order to keep this paper within a reasonable length.

6.1 Some preliminaries on overconvergent cohomology groups

The purpose of this subsection is to review the basic constructions and properties of the overconvergent cohomology groups needed in latter subsections. Most of the materials are recorded from [Han17; CHJ17]. We do not claim any originality here.

Definition 6.1.1. (i) Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be a small weight. We say it is **open** if the natural map

$$\mathcal{U}^{\text{rig}} = \text{Spa}(R_{\mathcal{U}}, R_{\mathcal{U}})^{\text{rig}} \rightarrow \mathcal{W}$$

is an open immersion.

(ii) Let $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be an affinoid weight. We say it is **open** if the natural map

$$\mathcal{U}^{\text{rig}} = \mathcal{U} = \text{Spa}(R_{\mathcal{U}}, R_{\mathcal{U}}^{\circ}) \rightarrow \mathcal{W}$$

is an open immersion.

(iii) A weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ is called an **open weight** if it is either an small open weight or an affinoid open weight.

Given an open weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ and an integer $r > 1 + r_{\mathcal{U}}$, one considers the so called *Borel–Serre chain complex* $C_{\bullet}(\text{Iw}_{\text{GSp}_{2g}}^+, A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$ (resp., *Borel–Serre cochain complex* $C^{\bullet}(\text{Iw}_{\text{GSp}_{2g}}^+, D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$) which computes the Betti homology groups $H_t(X_{\text{Iw}^+}(\mathbf{C}), A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$ (resp., Betti cohomology groups $H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$). The Borel–Serre chain complex is a finite complex as it is constructed by a fixed triangulation on the Borel–Serre compactification of the locally symmetric space $X_{\text{Iw}^+}(\mathbf{C})$. We write

$$\begin{aligned} C_{\text{tol}}^{\kappa_{\mathcal{U}}, r} &:= \bigoplus_t C_t(\text{Iw}_{\text{GSp}_{2g}}^+, A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})), \\ C_{\kappa_{\mathcal{U}}, r}^{\text{tol}} &:= \bigoplus_t C^t(\text{Iw}_{\text{GSp}_{2g}}^+, D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})). \end{aligned}$$

Then $C_{\text{tol}}^{\kappa_{\mathcal{U}}, r}$ is an ON-able $R_{\mathcal{U}}[1/p]$ -module as $A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$ is ON-able (see [Han17, §2.2, Remarks]). Moreover, there are naturally defined Hecke operators on $C_{\text{tol}}^{\kappa_{\mathcal{U}}, r}$ and the action of U_p is compact (see [op. cit., §2.2]). We define $F_{\kappa_{\mathcal{U}}, r}^{\text{oc}} \in R_{\mathcal{U}}[1/p][[T]]$ to be the Fredholm determinant of U_p acting on $C_{\text{tol}}^{\kappa_{\mathcal{U}}, r}$. Notice that, for any $h \in \mathbf{Q}_{\geq 0}$, the existence of a slope- $\leq h$ decomposition of $C_{\text{tol}}^{\kappa_{\mathcal{U}}, r}$ is equivalent to the existence of a slope- $\leq h$ factorisation of $F_{\kappa_{\mathcal{U}}, r}^{\text{oc}}$. Moreover, by [op. cit., Proposition 3.1.2], if $C_{\text{tol}, \leq h}^{\kappa_{\mathcal{U}}, r}$ is the slope- $\leq h$ submodule of $C_{\text{tol}}^{\kappa_{\mathcal{U}}, r}$ and suppose $\mathcal{U}' = (R_{\mathcal{U}'}, \kappa_{\mathcal{U}'})$ is another open weight such that $\mathcal{U}'^{\text{rig}} \subset \mathcal{U}^{\text{rig}}$, there is a canonical isomorphism

$$C_{\text{tol}, \leq h}^{\kappa_{\mathcal{U}}, r} \otimes_{R_{\mathcal{U}}[\frac{1}{p}]} R_{\mathcal{U}'}[\frac{1}{p}] \cong C_{\text{tol}, \leq h}^{\kappa_{\mathcal{U}'}, r}.$$

Definition 6.1.2. Let $\mathcal{U} = (R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ be an open weight and let $h \in \mathbf{Q}_{\geq 0}$. The pair (\mathcal{U}, h) is called a **slope datum** if $F_{\kappa_{\mathcal{U}}, r}^{\text{oc}}$ admits a slope- $\leq h$ factorisation.

Proposition 6.1.3. Let (\mathcal{U}, h) be a slope datum and let $(R_{\mathcal{U}'}, \kappa_{\mathcal{U}'})$ be an affinoid open weight such that $\mathcal{U}' \subset \mathcal{U}^{\text{rig}}$.

(i) There is a canonical isomorphism

$$H_t(X_{\text{Iw}^+}(\mathbf{C}), A_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))_{\leq h} \otimes_{R_{\mathcal{U}}[\frac{1}{p}]} R_{\mathcal{U}'} \cong H_t(X_{\text{Iw}^+}(\mathbf{C}), A_{\kappa_{\mathcal{U}'}}^r(\mathbf{T}_0, R_{\mathcal{U}'}))_{\leq h}$$

for all $t \in \mathbf{Z}$, where the subscript “ $\leq h$ ” stands for the slope- $\leq h$ submodule.

(ii) The cochain complex $C_{\kappa_{\mathcal{U}},r}^{\text{tol}}$ and the cohomology groups $H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$ admit slope- $\leq h$ decompositions. The corresponding slope- $\leq h$ submodules are denoted by $C_{\kappa_{\mathcal{U}},r}^{\text{tol},\leq h}$ and $H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))^{\leq h}$, respectively.

(iii) There are canonical isomorphisms

$$C_{\kappa_{\mathcal{U}},r}^{\text{tol},\leq h} \otimes_{R_{\mathcal{U}}[\frac{1}{p}]} R_{\mathcal{U}'} \cong C_{\kappa_{\mathcal{U}'},r}^{\text{tol},\leq h}$$

and

$$H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))^{\leq h} \otimes_{R_{\mathcal{U}}[\frac{1}{p}]} R_{\mathcal{U}'} \cong H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}'}}^r(\mathbf{T}_0, R_{\mathcal{U}'}))^{\leq h}.$$

Proof. The proof follows verbatim as in the proofs of [CHJ17, Proposition 3.3 & Proposition 3.4]. \square

Notice that, if we vary the open weights \mathcal{U} , the Fredholm determinants glue into a power series $F_{\mathcal{W}}^{\text{oc}} \in \mathcal{O}_{\mathcal{W}}(\mathcal{W})\{\{T\}\}$. Here we drop the subscript “ r ” because the Fredholm determinant does not depend on r according to [Han17, Proposition 3.1.1].

6.2 The cuspidal eigenvariety

The spectral variety. In the previous subsection, we obtained the Fredholm determinant $F_{\mathcal{W}}^{\text{oc}} \in \mathcal{O}_{\mathcal{W}}(\mathcal{W})\{\{T\}\}$. On the other hand, given an affinoid weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ and $w > 1 + r_{\mathcal{U}}$, by [AIP15, Proposition 8.1.3.1] and Theorem 3.7.1 (see also [op. cit., Proposition 8.2.3.3]), the space of cuspforms $S_{\text{Iw}^+,w}^{\kappa_{\mathcal{U}}} = H^0(\overline{\mathcal{X}}_{\text{Iw}^+,w}, \underline{\omega}_{w,\text{cusp}}^{\kappa_{\mathcal{U}}})$ has property (Pr) in the sense of [Buz07]; namely, it is a direct summand of a potentially ON-able $\mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}$ -Banach space. Also recall that U_p acts compactly on the space of overconvergent Siegel modular forms. Therefore, we can define the Fredholm determinant $F_{\kappa_{\mathcal{U}},w}^{\text{mf}}$ of U_p acting on $S_{\text{Iw}^+,w}^{\kappa_{\mathcal{U}}}$. When we vary the affinoid weights, the Fredholm determinants glue together. After further taking the inductive limit over w , we arrive at a power series $F_{\mathcal{W}}^{\text{mf}} \in \mathcal{O}_{\mathcal{W}}(\mathcal{W})\{\{T\}\} \widehat{\otimes}_{\mathbf{Q}_p} \mathbf{C}_p$.

Define

$$F_{\mathcal{W}} := F_{\mathcal{W}}^{\text{mf}} F_{\mathcal{W}}^{\text{oc}} \in \mathcal{O}_{\mathcal{W}}(\mathcal{W})\{\{T\}\} \widehat{\otimes}_{\mathbf{Q}_p} \mathbf{C}_p,$$

which is still a Fredholm series. Let $\mathbb{A}_{\mathbf{C}_p}^1$ denote the adic affine line over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ with coordinate function T and let $\mathbb{A}_{\mathcal{W}}^1 := \mathcal{W} \times_{\text{Spa}(\mathbf{Q}_p, \mathbf{Z}_p)} \mathbb{A}_{\mathbf{C}_p}^1$. The **spectral variety** \mathcal{S} is defined to be the zero locus of $F_{\mathcal{W}}$ in $\mathbb{A}_{\mathcal{W}}^1$.

The eigenvarieties.

Definition 6.2.1. Let \mathcal{U} be an open weight so that $\mathcal{U}^{\text{rig}} \subset \mathcal{W}$. Let $\mathcal{U}_{\mathbf{C}_p}^{\text{rig}}$ denote the base change of \mathcal{U}^{rig} to $\text{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and consider the adic affine line $\mathbb{A}_{\mathcal{U}}^1 := \mathcal{U}^{\text{rig}} \times_{\text{Spa}(\mathbf{Q}_p, \mathbf{Z}_p)} \mathbb{A}_{\mathbf{C}_p}^1$ over $\mathcal{U}_{\mathbf{C}_p}^{\text{rig}}$. Moreover, for any $h \in \mathbf{Q}_{>0}$, consider the closed ball $\mathbf{B}(0, p^h)$ of radius p^h over \mathbf{C}_p and define $\mathbf{B}_{\mathcal{U},h} := \mathcal{U}^{\text{rig}} \times_{\text{Spa}(\mathbf{Q}_p, \mathbf{Z}_p)} \mathbf{B}(0, p^h)$. Let $\mathcal{S}_{\mathcal{U},h} := \mathcal{S} \cap \mathbf{B}_{\mathcal{U},h}$. We say that the pair (\mathcal{U}, h) is **slope-adapted** if the natural map $\mathcal{S}_{\mathcal{U},h} \rightarrow \mathcal{U}_{\mathbf{C}_p}^{\text{rig}}$ is finite flat.

Consider the collection

$$\text{Cov}(\mathcal{S}) = \{\mathcal{S}_{\mathcal{U},h} : (\mathcal{U}, h) \text{ is slope-adapted}\}.$$

Let $\text{Cov}_{\text{aff}}(\mathcal{S})$ be a subcollection of $\text{Cov}(\mathcal{S})$, consisting of those $\mathcal{S}_{\mathcal{U},h}$ with \mathcal{U} being an affinoid weight. By [Buz07, Theorem 4.6] (see also [Han17, Proposition 4.1.4]), $\text{Cov}_{\text{aff}}(\mathcal{S})$ forms an open covering for \mathcal{S} (and thus so is $\text{Cov}(\mathcal{S})$). Using this covering, we define the following two coherent sheaves on \mathcal{S} .

Definition 6.2.2. (i) Recall from §4.1 that $D_{\kappa_{\mathcal{U}}}^{\dagger}(\mathbf{T}_0, R_{\mathcal{U}})$ is defined to be the inverse limit of $D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})$ with respect to r . The coherent sheaf $\mathcal{H}_{\text{par}}^{\text{tol}}$ on \mathcal{S} is defined by

$$\mathcal{H}_{\text{par}}^{\text{tol}}(\mathcal{S}_{\mathcal{U},h}) := \text{image} \left(\bigoplus_t H_c^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^{\dagger}(\mathbf{T}_0, R_{\mathcal{U}}))^{\leq h} \rightarrow \bigoplus_t H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^{\dagger}(\mathbf{T}_0, R_{\mathcal{U}}))^{\leq h} \right) \widehat{\otimes}_{\mathbf{Q}_p} \mathbf{C}_p$$

for all $\mathcal{S}_{\mathcal{U},h} \in \text{Cov}_{\text{aff}}(\mathcal{S})$, where the map

$$H_c^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^{\dagger}(\mathbf{T}_0, R_{\mathcal{U}}))^{\leq h} \rightarrow H^t(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^{\dagger}(\mathbf{T}_0, R_{\mathcal{U}}))^{\leq h}$$

is induced from the natural map from the compactly supported cohomology groups to the usual ones.

(ii) The coherent sheaf $\mathcal{S}_{\text{Iw}^+}^{\dagger}$ on \mathcal{S} is defined by

$$\mathcal{S}_{\text{Iw}^+}^{\dagger}(\mathcal{S}_{\mathcal{U},h}) := S_{\text{Iw}^+}^{\kappa_{\mathcal{U}}+g+1, \leq h}$$

for all $\mathcal{S}_{\mathcal{U},h} \in \text{Cov}_{\text{aff}}(\mathcal{S})$, where the superscript “ $\leq h$ ” stands for the slope- $\leq h$ part with respect to the U_p -operator.

These are indeed well-defined coherent sheaves (see, for example, [Han17, §4.3] and [AIP15, §8.1], respectively). The Hecke algebra \mathbb{T} acts on both of the coherent sheaves. The eigenvarieties we are interested in are the following.

Definition 6.2.3. (i) For every $\mathcal{S}_{\mathcal{U},h} \in \text{Cov}_{\text{aff}}(\mathcal{S})$, let $\mathbb{T}_{\mathcal{U},h}^{\text{oc}}$ be the reduced $\mathcal{O}_{\mathcal{S}_{\mathcal{U},h}}(\mathcal{S}_{\mathcal{U},h})$ -algebra generated by the image of $\mathbb{T} \rightarrow \text{End}(\mathcal{H}_{\text{par}}^{\text{tol}}(\mathcal{S}_{\mathcal{U},h}))$. Let $\mathbb{T}_{\mathcal{U},h}^{\text{oc},\circ}$ be the integral closure of $\mathcal{O}_{\mathcal{S}_{\mathcal{U},h}}(\mathcal{S}_{\mathcal{U},h})^{\circ}$ inside $\mathbb{T}_{\mathcal{U},h}^{\text{oc}}$.

(ii) Let \mathcal{T}_{oc} be the coherent sheaf on \mathcal{S} defined by $\mathcal{T}_{\text{oc}}(\mathcal{S}_{\mathcal{U},h}) := \mathbb{T}_{\mathcal{U},h}^{\text{oc}}$, and let $\mathcal{T}_{\text{oc}}^{\circ}$ be the subsheaf of \mathcal{T}_{oc} defined by $\mathcal{T}_{\text{oc}}^{\circ}(\mathcal{S}_{\mathcal{U},h}) := \mathbb{T}_{\mathcal{U},h}^{\text{oc},\circ}$.

(iii) The **reduced cuspidal eigenvariety** $\mathcal{E}_0^{\text{oc}}$ is defined to be the relative adic space $\text{Spa}_{\mathcal{S}}(\mathcal{T}_{\text{oc}}, \mathcal{T}_{\text{oc}}^{\circ})$.

Definition 6.2.4. (i) For every $\mathcal{S}_{\mathcal{U},h} \in \text{Cov}_{\text{aff}}(\mathcal{S})$, let $\mathbb{T}_{\mathcal{U},h}^{\text{mf}}$ be the reduced $\mathcal{O}_{\mathcal{S}_{\mathcal{U},h}}(\mathcal{S}_{\mathcal{U},h})$ -algebra generated by the image of $\mathbb{T} \rightarrow \text{End}(\mathcal{S}_{\text{Iw}^+}^{\dagger}(\mathcal{S}_{\mathcal{U},h}))$. Let $\mathbb{T}_{\mathcal{U},h}^{\text{mf},\circ}$ be the integral closure of $\mathcal{O}_{\mathcal{S}_{\mathcal{U},h}}(\mathcal{S}_{\mathcal{U},h})^{\circ}$ inside $\mathbb{T}_{\mathcal{U},h}^{\text{mf}}$.

(ii) Let \mathcal{T}_{mf} be the coherent sheaf on \mathcal{S} defined by $\mathcal{T}_{\text{mf}}(\mathcal{S}_{\mathcal{U},h}) := \mathbb{T}_{\mathcal{U},h}^{\text{mf}}$. Let $\mathcal{T}_{\text{mf}}^{\circ}$ be the subsheaf of \mathcal{T}_{mf} defined by $\mathcal{T}_{\text{mf}}^{\circ}(\mathcal{S}_{\mathcal{U},h}) := \mathbb{T}_{\mathcal{U},h}^{\text{mf},\circ}$.

(iii) The **equidimensional cuspidal eigenvariety** $\mathcal{E}_0^{\text{mf}}$ is defined to be the equidimensional locus of the relative adic space $\text{Spa}_{\mathcal{S}}(\mathcal{T}_{\text{mf}}, \mathcal{T}_{\text{mf}}^{\circ})$.

Remark 6.2.5. Notice that $\mathcal{E}_0^{\text{mf}}$ is (the stricit Iwahori version of) the equidimensional cuspidal eigenvariety constructed in [AIP15] after base change to \mathbf{C}_p . On the other hand, $\mathcal{E}_0^{\text{oc}}$ is the reduced cuspidal eigenvariety considered in [Wu20] after base change to \mathbf{C}_p . We also point out that the cuspidal eigenvariety considered in *op. cit.* is the cuspidal part of the eigenvariety for GSp_{2g} constructed in [JN19] (see also [Han17]).

Proposition 6.2.6. There is a natural closed immersion $\mathcal{E}_0^{\text{mf}} \hookrightarrow \mathcal{E}_0^{\text{oc}}$.

Proof. The strategy is to apply [Han17, Theorem 5.1.2]. To this end, we need to find a *very Zariski-dense* subset \mathcal{S}^{cl} of \mathcal{S} such that for every $\mathbf{x} \in \mathcal{S}^{\text{cl}}$ with dominate algebraic weight $k = (k_1, \dots, k_g) \in \mathbf{Z}_{\geq 0}^g$ and any $Y \in \mathbb{T}$, we have

$$\det(1 - TY | \mathcal{S}_{\text{Iw}^+}^{\dagger}(\mathbf{x})) \mid \det(1 - TY | \mathcal{H}_{\text{par}, \mathbf{x}}^{\text{tol}}).$$

By [Wu20, Theorem 4.3.2], there exists an $h_k \in \mathbf{R}_{>0}$ such that for all $h \in \mathbf{Q} \cap (0, h_k]$, the canonical map

$$H_{\text{par}}^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), D_k^\dagger(\mathbf{T}_0, \mathbf{Q}_p))^{\leq h} \rightarrow H_{\text{par}}^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), \mathbf{V}_{\text{GSp}_{2g}, k}^{\text{alg}, \vee})^{\leq h}$$

is an isomorphism. On the other hand, let

$$\underline{\omega}_{\text{Iw}^+, \text{cusp}}^k := \underline{\omega}_{\text{Iw}^+}^k \otimes_{\mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+}}} \mathcal{O}_{\overline{\mathcal{X}}_{\text{Iw}^+}}(-Z_{\text{Iw}^+})$$

be the sheaf of classical cuspidal Siegel modular forms of weight k on $\overline{\mathcal{X}}_{\text{Iw}^+}$. The classicality theorem [AIP15, Theorem 7.1.1] provides an $a_k \in \mathbf{Q}_{>0}$ such that for all $h \in \mathbf{Q} \cap (0, a_k]$, the slope- $\leq h$ overconvergent Siegel cuspforms of weight k are classical; namely,

$$H^0(\overline{\mathcal{X}}_{\text{Iw}^+, w}, \underline{\omega}_{w, \text{cusp}}^k)^{\leq h} \subset H^0(\overline{\mathcal{X}}_{\text{Iw}^+}, \underline{\omega}_{\text{Iw}^+, \text{cusp}}^k).$$

Now, let $\ell_k = \min\{h_k, a_k\}$ and take $h \leq \ell_k$. Applying the *generalised Eichler–Shimura morphism* in [Hid02, Theorem 3.8], we obtain an injection from the space of slope- $\leq h$ overconvergent Siegel cuspforms of classical weight into the slope- $\leq h$ cohomology group with coefficient in the algebraic representation. Consequently, the desired very Zariski-dense subset of \mathcal{S} can be taken to be

$$\mathcal{S}^{\text{cl}} = \bigcup_{\mathcal{S}_{\mathcal{U}, h} \in \text{Cov}_{\text{aff}}(\mathcal{S})} \{x \in \mathcal{S}_{\mathcal{U}, h} : x \text{ has classical weight } k \in \mathbf{Z}_{\geq 0}^g \text{ and } h \leq \ell_k\}$$

Finally, [Han17, Theorem 5.1.2] yields the result. \square

Given Proposition 6.2.6, we may identify $\mathcal{E}_0^{\text{mf}}$ with its image in $\mathcal{E}_0^{\text{oc}}$ and denote it by \mathcal{E}_0 for simplicity. We have a diagram

$$\begin{array}{ccc} \mathcal{E}_0 & \xrightarrow{\pi} & \mathcal{S} \xrightarrow{\text{wt}_{\mathcal{S}}} \mathcal{W} \\ & \searrow \text{wt} & \nearrow \end{array}$$

6.3 Sheaves on the cuspidal eigenvariety

Given a weight $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ and an integer $r > 1 + r_{\mathcal{U}}$, we write

$$\mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r, \text{cusp}} = \begin{cases} \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r, \text{cusp}} & \text{if } \mathcal{U} \text{ is a small weight} \\ \text{image}(H_c^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})) \rightarrow H^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))) \widehat{\otimes}_{\mathbf{Q}_p} \mathbf{C}_p & \text{if } \mathcal{U} \text{ is an affinoid weight} \end{cases},$$

where $H_c^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}})) \rightarrow H^{n_0}(X_{\text{Iw}^+}(\mathbf{C}), D_{\kappa_{\mathcal{U}}}^r(\mathbf{T}_0, R_{\mathcal{U}}))$ is the natural map from the compactly supported Betti cohomology group to the usual one. We also write

$$\mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{\dagger, \text{cusp}} := \varprojlim_r \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r, \text{cusp}}.$$

Suppose that $(R_{\mathcal{U}}, \kappa_{\mathcal{U}})$ is an small open weight and recall the overconvergent Eichler–Shimura morphism for overconvergent Siegel cuspforms

$$\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r, \text{cusp}} \rightarrow S_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}} + g + 1}(-n_0).$$

If (\mathcal{U}, h) slope-adapted, then the Hecke-equivariance of $\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}}$ induces a $\mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}$ -linear map

$$\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}, \leq h} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r, \text{cusp}, \leq h} \rightarrow S_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}} + g + 1, \leq h}(-n_0).$$

of finite projective $\mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}}$ -modules.

Now, if $\mathcal{U}' \subset \mathcal{U}^{\text{rig}}$ is an affinoid weight, the $\mathbf{C}_p \widehat{\otimes} R_{\mathcal{U}'}$ -linear map $\text{ES}_{\kappa_{\mathcal{U}'}}^{\text{cusp}}$ is defined to be the composition

$$\text{ES}_{\kappa_{\mathcal{U}'}}^{\text{cusp}, \leq h} : \mathfrak{W}_{\kappa_{\mathcal{U}'}, \mathbf{C}_p}^{r, \text{cusp}, \leq h} \cong \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{r, \text{cusp}, \leq h} \otimes_{R_{\mathcal{U}}[\frac{1}{p}]} R_{\mathcal{U}'} \rightarrow S_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}} + g + 1, \leq h}(-n_0) \rightarrow S_{\text{Iw}^+, w}^{\kappa_{\mathcal{U}'} + g + 1, \leq h}(-n_0), \quad (9)$$

where the first isomorphism follows from Proposition 6.1.3.

Recall the natural map $\pi : \mathcal{E}_0 \rightarrow \mathcal{S}$ and let $\mathcal{E}_{\mathcal{U}, h}$ be the preimage of $\mathcal{S}_{\mathcal{U}, h}$. On the cuspidal eigenvariety \mathcal{E}_0 , we consider two coherent sheaves $\mathcal{O}_{\text{cusp}}^\dagger$ and $\mathcal{S}_{\text{Iw}^+}^\dagger(-n_0)$ given by

$$\mathcal{O}_{\text{cusp}}^\dagger(\mathcal{E}_{\mathcal{U}, h}) := \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{\dagger, \text{cusp}, \leq h}$$

and

$$\mathcal{S}_{\text{Iw}^+}^\dagger(-n_0)(\mathcal{E}_{\mathcal{U}, h}) := S_{\text{Iw}^+}^{\kappa_{\mathcal{U}} + g + 1, \leq h}(-n_0),$$

for all $\mathcal{S}_{\mathcal{U}, h} \in \text{Cov}_{\text{aff}}(\mathcal{S})$.

Theorem 6.3.1. *There exists a morphism*

$$\mathcal{ES} : \mathcal{O}_{\text{cusp}}^\dagger \rightarrow \mathcal{S}_{\text{Iw}^+}^\dagger(-n_0)$$

of coherent sheaves over \mathcal{E}_0 such that if (\mathcal{U}, h) is a slope-adapted pair, then $\mathcal{ES}(\mathcal{E}_{\mathcal{U}, h})$ is exactly the overconvergent Eichler–Shimura morphism for overconvergent Siegel cuspforms

$$\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}, \leq h} : \mathfrak{W}_{\kappa_{\mathcal{U}}, \mathbf{C}_p}^{\dagger, \text{cusp}, \leq h} \rightarrow S_{\text{Iw}^+}^{\kappa_{\mathcal{U}} + g + 1, \leq h}(-n_0).$$

Proof. It follows from (9) and the functoriality of $\text{ES}_{\kappa_{\mathcal{U}}}^{\text{cusp}}$ in small open weights \mathcal{U} . \square

Denote by \mathcal{Im} and \mathcal{Ker} the image and the kernel of \mathcal{ES} , respectively. We obtain a short exact sequence of sheaves on \mathcal{E}_0

$$0 \rightarrow \mathcal{Ker} \rightarrow \mathcal{O}_{\text{cusp}}^\dagger \rightarrow \mathcal{Im} \rightarrow 0.$$

We remind the readers that this short exact sequence is Galois- and Hecke-equivariant.

Theorem 6.3.2. *Let $\mathcal{E}_0^{\text{loc}}$ be the locus of \mathcal{E}_0 where \mathcal{Ker} , $\mathcal{O}_{\text{cusp}}^\dagger$, and \mathcal{Im} are locally free. Then the short exact sequence*

$$0 \rightarrow \mathcal{Ker} \rightarrow \mathcal{O}_{\text{cusp}}^\dagger \rightarrow \mathcal{Im} \rightarrow 0$$

over $\mathcal{E}_0^{\text{loc}}$ splits locally.

Proof. We follow the same strategy as in [AIS15, Theorem 6.1(c)]. For any sufficiently small open affinoid subset $\mathcal{V} = \text{Spa}(R_{\mathcal{V}}, R_{\mathcal{V}}^+) \subset \mathcal{E}_0^{\text{loc}}$ on which the three sheaves are free and the sequence

$$0 \rightarrow \mathcal{Ker}(\mathcal{V}) \rightarrow \mathcal{O}_{\text{cusp}}^\dagger(\mathcal{V}) \rightarrow \mathcal{Im}(\mathcal{V}) \rightarrow 0$$

is exact, consider

$$\mathcal{H}(\mathcal{V}) := \text{Hom}_{R_{\mathcal{V}}}(\mathcal{Im}(\mathcal{V}), \mathcal{Ker}(\mathcal{V})).$$

The $G_{\mathbf{Q}_p}$ -equivariance of the short exact sequence defines a class in $H^1(G_{\mathbf{Q}_p}, \mathcal{H}(\mathcal{V})) \cong \text{Ext}_{R_{\mathcal{V}}[G_{\mathbf{Q}_p}]}^1(\mathcal{Im}(\mathcal{V}), \mathcal{Ker}(\mathcal{V}))$.

Let $\varphi_{\text{Sen}} \in \text{End}(\mathcal{H}(\mathcal{V}))$ denote the Sen operator associated with $\mathcal{H}(\mathcal{V})$. Such an operator was introduced in [Sen88] (see also [Kis03]). Then by [Kis03, Proposition 2.3], $\det(\varphi_{\text{Sen}}) \in R_{\mathcal{V}}$ kills the cohomology group $H^1(G_{\mathbf{Q}_p}, \mathcal{H}(\mathcal{V}))$. On the other hand, $\det(\varphi_{\text{Sen}})$ is non-zero. Therefore, after localising at this element, the short exact sequence splits as a sequence of semilinear $G_{\mathbf{Q}_p}$ -representations. Since the Galois-action commutes with the Hecke-actions, the splitting can be chosen to be Hecke-equivariant. \square

A Kummer étale and pro-Kummer étale sites of log adic spaces

In order to study the boundaries of various toroidal compactifications of Siegel varieties, we adopt the language of *logarithmic adic spaces* established in [Dia+19a]. The purpose of §A.1 is to review the basic notions of log adic spaces, as well as their Kummer étale and pro-Kummer étale topologies, for convenience of the readers who are not familiar with the language. In §A.2, we present an explicit calculation of the sheaf $R^i \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$ which plays an essential role in the construction of the overconvergent Eichler–Shimura morphisms in §5.2. Finally, in §A.3, we introduce the notion of *Kummer étale Banach sheaves* and prove a (generalised) projection formula for those Kummer étale Banach sheaves that are *admissible*.

Notation. We warn the reader that, in this section, (log) adic spaces will no longer be written in calligraphic font as we deal with more general (log) adic spaces, not only those studied in the main body of the text.

A.1 Review of log adic spaces

Let k be a nonarchimedean field (*i.e.*, a field complete with respect to a nonarchimedean norm $|\cdot| : k \rightarrow \mathbf{R}_{\geq 0}$) and let $\mathcal{O}_k = \{x \in k : |x| \leq 1\}$.

Definition A.1.1. Let X be a locally noetherian adic space over $\text{Spa}(k, \mathcal{O}_k)$.

- (i) A **pre-log structure** on X is a pair (\mathcal{M}_X, α) where \mathcal{M}_X is a sheaf of monoids on $X_{\text{ét}}$ and $\alpha : \mathcal{M}_X \rightarrow \mathcal{O}_{X_{\text{ét}}}$ is a morphism of sheaves of (multiplicative) monoids. It is called a **log structure** if the induced morphism $\alpha^{-1}(\mathcal{O}_{X_{\text{ét}}}^\times) \rightarrow \mathcal{O}_{X_{\text{ét}}}^\times$ is an isomorphism. In this case, the triple $(X, \mathcal{M}_X, \alpha)$ is called a **log adic space**. If the context is clear, we simply say that X is a log adic space.
- (ii) For a pre-log structure (\mathcal{M}_X, α) on X , the **associated log structure** is $({}^a\mathcal{M}_X, {}^a\alpha)$ where ${}^a\mathcal{M}_X$ is given by the pushout

$$\begin{array}{ccc} \alpha^{-1}(\mathcal{O}_{X_{\text{ét}}}^\times) & \longrightarrow & \mathcal{M}_X \\ \downarrow & \lrcorner & \downarrow \\ \mathcal{O}_{X_{\text{ét}}}^\times & \longrightarrow & {}^a\mathcal{M}_X \end{array}$$

and ${}^a\alpha : {}^a\mathcal{M}_X \rightarrow \mathcal{O}_{X_{\text{ét}}}$ is the induced morphism.

- (iii) A **morphism** $f : (Y, \mathcal{M}_Y, \alpha_Y) \rightarrow (X, \mathcal{M}_X, \alpha_X)$ of log adic spaces is a morphism $f : Y \rightarrow X$ of adic spaces together with a morphism of sheaves of monoids $f^\sharp : f^{-1}\mathcal{M}_X \rightarrow \mathcal{M}_Y$ such that the diagram

$$\begin{array}{ccc} f^{-1}\mathcal{M}_X & \xrightarrow{f^\sharp} & \mathcal{M}_Y \\ f^{-1}\alpha_X \downarrow & & \downarrow \alpha_Y \\ f^{-1}\mathcal{O}_{X_{\text{ét}}} & \longrightarrow & \mathcal{O}_{Y_{\text{ét}}} \end{array}$$

commutes. Moreover, the log structure associated with the pre-log structure $f^{-1}\mathcal{M}_X \rightarrow f^{-1}\mathcal{O}_{X_{\text{ét}}} \rightarrow \mathcal{O}_{Y_{\text{ét}}}$ is called the **pullback log structure**, denoted by $f^*\mathcal{M}_X$. We say that f is **strict** if $f^*\mathcal{M}_X \xrightarrow{\sim} \mathcal{M}_Y$.

Definition A.1.2. (i) Let $(X, \mathcal{M}_X, \alpha)$ be a locally noetherian log adic space as above. Let P be a monoid and let P_X denote the associated constant sheaf of monoids on $X_{\text{ét}}$. A **chart of X modeled on P** is a morphism of sheaves of monoids $\theta : P_X \rightarrow \mathcal{M}_X$ such that $\alpha(\theta(P_X)) \subset \mathcal{O}_{X_{\text{ét}}}^\times$ and such that the log structure associated with the pre-log structure $\alpha \circ \theta : P_X \rightarrow \mathcal{O}_{X_{\text{ét}}}$ is isomorphic to \mathcal{M}_X . We say that the chart is **fs** if P is fine and saturated.

- (ii) A locally noetherian log adic space is called an **fs log adic space** if it étale locally admits charts modeled on fs monoids.
- (iii) Let $f : (Y, \mathcal{M}_Y, \alpha_Y) \rightarrow (X, \mathcal{M}_X, \alpha_X)$ be a morphism between locally noetherian log adic spaces. A **chart** of f consists of charts $\theta_X : P_X \rightarrow \mathcal{M}_X$ and $\theta_Y : Q_Y \rightarrow \mathcal{M}_Y$ and a homomorphism $u : P \rightarrow Q$ such that the diagram

$$\begin{array}{ccc} P_Y & \xrightarrow{u} & Q_Y \\ \downarrow \theta_X & & \downarrow \theta_Y \\ f^{-1} \mathcal{M}_X & \xrightarrow{f^\sharp} & \mathcal{M}_Y \end{array}$$

commutes. We say that the chart is **fs** if both P and Q are fs. When the context is clear, we simply say that $u : P \rightarrow Q$ is a chart of f .

Below are two typical examples of locally noetherian fs log adic spaces. In this paper, all of the toroidally compactified Siegel varieties (equipped with logarithmic structures associated with the boundary divisors) have the form as in Example A.1.4.

Example A.1.3. Let $n > 0$ be an integer. Consider the *n -dimensional unit disc*

$$\mathbb{D}^n := \mathrm{Spa}(k\langle T_1, \dots, T_n \rangle, \mathcal{O}_k\langle T_1, \dots, T_n \rangle),$$

equipped with the log structure associated with the pre-log structure induced by

$$\mathbf{Z}_{\geq 0}^n \rightarrow k\langle T_1, \dots, T_n \rangle, \quad (a_1, \dots, a_n) \mapsto T_1^{a_1} \cdots T_n^{a_n}.$$

Clearly, \mathbb{D}^n is modeled on the fs chart $\mathbf{Z}_{\geq 0}^n$. □

Example A.1.4. Let X be a smooth rigid analytic variety over k , viewed as an adic space over $\mathrm{Spa}(k, \mathcal{O}_k)$ via [Hub13, (1.1.11)]. Let $D \subset X$ be a **normal crossings divisor** in the sense of [Dia+19a, Example 2.3.17]. Namely, $\iota : D \hookrightarrow X$ is a closed immersion such that, analytic locally, X and D are of the form $S \times \mathbb{D}^n$ and $S \times \{T_1 \cdots T_n = 0\}$, where S is a smooth connected rigid analytic variety and ι is the pullback of the natural inclusion $\{T_1 \cdots T_n = 0\} \hookrightarrow \mathbb{D}^n$. We equip X with the log structure

$$\mathcal{M}_X = \{f \in \mathcal{O}_{X_{\mathrm{\acute{e}t}}} \mid f \text{ is invertible on } X \setminus D\}$$

with $\alpha : \mathcal{M}_X \rightarrow \mathcal{O}_{X_{\mathrm{\acute{e}t}}}$ being the natural inclusion. This is the **divisorial log structure** associated with the divisor D . This log structure agrees with the pullback of the log structure on \mathbb{D}^n constructed in Example A.1.3. □

Log adic spaces in Example A.1.4 are special cases of *log smooth* ones. For later use, we recall the definition of log smoothness.

For any monoid P and any commutative ring T , we write $T[P]$ for the associated monoid algebra.

Definition A.1.5. Let X be a locally noetherian adic space over $\mathrm{Spa}(k, \mathcal{O}_k)$ and let P be a finitely generated monoid. For any affinoid open subspace $\mathrm{Spa}(R, R^+) \subset X$, let $(R\langle P \rangle, R^+\langle P \rangle)$ be the completion of $(R[P], R^+[P])$. By gluing the morphisms $\mathrm{Spa}(R\langle P \rangle, R^+\langle P \rangle) \rightarrow \mathrm{Spa}(R, R^+)$, we obtain a morphism $X\langle P \rangle \rightarrow X$. Moreover, we equip $X\langle P \rangle$ with the log structure modeled on the chart P ; i.e., the one locally induced by $P \rightarrow R\langle P \rangle$.

Definition A.1.6. Let $f : Y \rightarrow X$ be a morphism between locally noetherian fs log adic spaces. We say that f is **log smooth** if étale locally f admits an fs chart $u : P \rightarrow Q$ such that

- (i) the kernel and the torsion part of the cokernel of $u^{\text{gp}} : P^{\text{gp}} \rightarrow Q^{\text{gp}}$ are finite groups of order invertible in \mathcal{O}_X ; and
- (ii) f and u induce a morphism $Y \rightarrow X \times_{X\langle P \rangle} X\langle Q \rangle$ of log adic spaces whose underlying morphism of adic spaces is étale.

A locally noetherian fs log adic space X is **log smooth** if the structure morphism $X \rightarrow \text{Spa}(k, \mathcal{O}_k)$ is log smooth, where $\text{Spa}(k, \mathcal{O}_k)$ is equipped with the trivial log structure.

Now we introduce the notion of Kummer étale morphisms and the Kummer étale site.

Definition A.1.7. (i) An injective homomorphism $u : P \rightarrow Q$ of fs monoids is called **Kummer** if for every $a \in Q$, there exists some integer $n > 0$ such that $na \in u(P)$.

- (ii) A morphism (resp., finite morphism) $f : Y \rightarrow X$ of locally noetherian fs log adic spaces is called **Kummer étale** (resp., **finite Kummer étale**) if étale locally on X and Y , f admits an fs chart $u : P \rightarrow Q$ which is Kummer with $|Q^{\text{gp}}/u^{\text{gp}}(P^{\text{gp}})|$ invertible on \mathcal{O}_Y , and such that f and u induce a morphism $Y \rightarrow X \times_{X\langle P \rangle} X\langle Q \rangle$ of log adic spaces whose underlying morphism of adic spaces is étale (resp., finite étale).

- (iii) If a Kummer étale (resp., finite Kummer étale) morphism is strict, we say it is **strictly étale** (resp., **strictly finite étale**).

Remark A.1.8. By [Dia+19a, Lemma 4.1.10], if $f : Y \rightarrow X$ is a Kummer étale morphism between locally noetherian fs log adic spaces and if X admits a chart modeled on a sharp fs monoid P , then, étale locally on X and Y , the morphism f admits a Kummer fs chart $P \rightarrow Q$ with Q being sharp.

Definition A.1.9. Let X be a locally noetherian fs log adic space. The **Kummer étale site** $X_{\text{két}}$ (resp., **finite Kummer étale site** $X_{\text{fkét}}$) of X is defined as follows. The underlying category is the full subcategory of the category of locally noetherian fs log adic spaces consisting of objects that are Kummer étale (resp., finite Kummer étale) over X . The coverings are given by the topological coverings.

The **structure sheaf** $\mathcal{O}_{X_{\text{két}}}$ (resp., **integral structure sheaf** $\mathcal{O}_{X_{\text{két}}}^+$) on $X_{\text{két}}$ is defined to be the presheaf sending $U \mapsto \mathcal{O}_U(U)$ (resp., $U \mapsto \mathcal{O}_U^+(U)$). We also write $\mathcal{M}_{X_{\text{két}}}$ for the presheaf sending $U \mapsto \mathcal{M}_U(U)$. By [Dia+19a, Theorem 4.3.1, Proposition 4.3.4], these presheaves are indeed sheaves.

Proposition A.1.10 (Corollary 4.4.18, [Dia+19a]). Let X be a connected locally noetherian fs log adic space and let ξ be a log geometric point (see [Dia+19a, Definition 4.4.2]). Then there is an equivalence of categories

$$F_X : X_{\text{fkét}} \xrightarrow{\sim} \pi_1^{\text{két}}(X, \xi) - \mathbf{FSETS}$$

sending $Y \mapsto Y_\xi := \text{Hom}_X(\xi, Y)$, where the $\pi_1^{\text{két}}(X, \xi) - \mathbf{FSETS}$ denotes the category of finite sets equipped with a continuous action of the **Kummer étale fundamental group** $\pi_1^{\text{két}}(X, \xi)$.

For any two log geometric points ξ and ξ' , the fundamental groups $\pi_1^{\text{két}}(X, \xi)$ and $\pi_1^{\text{két}}(X, \xi')$ are isomorphic. Hence, we may omit “ ξ ” from the notation whenever the context is clear.

Lemma A.1.11. Assume k is of characteristic 0. Let X and Y be locally noetherian fs log adic spaces whose underlying adic spaces are smooth connected rigid analytic varieties over k . Suppose the log structures on X and Y are the divisorial log structures associated with the normal crossing divisors $D \subset X$ and $E \subset Y$ as in Example A.1.4. Let $U = X \setminus D$ and $V = Y \setminus E$. Suppose we have a finite Kummer étale surjective morphism $f : Y \rightarrow X$ such that $f^{-1}(U) = V$ and that $f|_V : V \rightarrow U$ is a finite étale Galois cover with Galois group G . Then f is a finite Kummer étale Galois cover with Galois group G .

Proof. According to Proposition A.1.10, we have equivalences of categories

$$F_X : X_{\text{fkét}} \xrightarrow{\sim} \pi_1^{\text{két}}(X) - \mathbf{FSETS}$$

and

$$F_U : U_{\text{fét}} \xrightarrow{\sim} \pi_1^{\text{ét}}(U) - \mathbf{FSETS}.$$

We have to show that G is a finite quotient of $\pi_1^{\text{két}}(X)$ and, under the equivalence F_X , Y corresponds to the finite set G equipped with the natural $\pi_1^{\text{két}}(X)$ -action.

By [Dia+19a, Proposition 4.2.1] and [Han20, Theorem 1.6], we have an equivalence of categories between $X_{\text{fkét}}$ and $U_{\text{fét}}$, under which Y corresponds to V . It also induces a natural isomorphism $\pi_1^{\text{két}}(X) \cong \pi_1^{\text{ét}}(U)$ making the following diagram commutative.

$$\begin{array}{ccc} X_{\text{fkét}} & \xrightarrow{\sim} & U_{\text{fét}} \\ \downarrow \sim F_X & & \downarrow \sim F_U \\ \pi_1^{\text{két}}(X) - \mathbf{FSETS} & \xrightarrow{\sim} & \pi_1^{\text{ét}}(U) - \mathbf{FSETS} \end{array}$$

Since V corresponds to the finite set G under the equivalence F_U , we are done. \square

Finally, we introduce the pro-Kummer étale site. For the rest of §A.1, the nonarchimedean field k is assumed to be an extension of \mathbf{Q}_p .

Definition A.1.12. Let X be a locally noetherian fs log adic space over $\text{Spa}(k, \mathcal{O}_k)$.

(i) The **pro-Kummer étale site** $X_{\text{prokét}}$ of X is defined as follows. The underlying category is the full subcategory of $\text{pro-}X_{\text{két}}$ consisting of cofiltered inverse limit $Y = \varprojlim_{i \in I} Y_i$ with $Y_i \in X_{\text{két}}$ such that the transition morphisms $Y_i \rightarrow Y_j$ are finite Kummer étale and are surjective for sufficiently large i . Such an inverse limit is called a **pro-Kummer étale presentation** of Y . As for the coverings, we refer the readers to [Dia+19a, Definition 5.1.1, 5.1.2] for details.

(ii) There is a natural projection of sites

$$\nu : X_{\text{prokét}} \rightarrow X_{\text{két}}.$$

The **structure sheaves** on $X_{\text{prokét}}$ are given by

$$\mathcal{O}_{X_{\text{prokét}}}^+ := \nu^{-1} \mathcal{O}_{X_{\text{két}}}^+, \quad \mathcal{O}_{X_{\text{prokét}}} := \nu^{-1} \mathcal{O}_{X_{\text{két}}}$$

and the **completed structure sheaves** are given by

$$\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ := \varprojlim_n (\mathcal{O}_{X_{\text{prokét}}} / p^n), \quad \widehat{\mathcal{O}}_{X_{\text{prokét}}} := \widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ [1/p].$$

We also write $\mathcal{M}_{X_{\text{prokét}}} := \nu^{-1}(\mathcal{M}_{\text{két}})$ together with a natural morphism $\alpha : \mathcal{M}_{\text{prokét}} \rightarrow \mathcal{O}_{X_{\text{prokét}}}$.

The pro-Kummer étale topology admits a convenient basis consisting of the *log affinoid perfectoid objects*.

Definition A.1.13. An object U in $X_{\text{prokét}}$ is called **log affinoid perfectoid** if it admits a pro-Kummer étale presentation $U = \varprojlim_{i \in I} U_i$ such that

(i) There is an initial object $0 \in I$;

(ii) Each $U_i = (\text{Spa}(R_i, R_i^+))$ is affinoid and admits a chart modeled on a sharp fs monoid P_i such that each transition morphism $U_j \rightarrow U_i$ is modeled on a Kummer chart $P_i \rightarrow P_j$;

(iii) The affinoid algebra $(R, R^+) := \left(\varinjlim_{i \in I} (R_i, R_i^+) \right)^\wedge$ is a perfectoid affinoid algebra, where the completion is with respect to the p -adic topology;

(iv) The monoid $P := \varinjlim_{i \in I} P_i$ is n -**divisible**, for all $n \in \mathbf{Z}_{\geq 1}$. Namely, the n -th multiple map $[n] : P \rightarrow P$ is surjective for all $n \in \mathbf{Z}_{\geq 1}$.

Such a presentation $U = \varprojlim_{i \in I} U_i$ is called a **perfectoid presentation** of U .

Proposition A.1.14 (Proposition 5.3.12, [Dia+19a]). *The log affinoid perfectoid objects in $X_{\text{prokét}}$ form a basis of the pro-Kummer étale site.*

Proposition A.1.15 (Theorem 5.4.3, [Dia+19a]). *Let $U \in X_{\text{prokét}}$ be a log affinoid perfectoid object, with the associated perfectoid space $\widehat{U} = \text{Spa}(R, R^+)$. Then*

(i) *For each $n \in \mathbf{Z}_{\geq 1}$, we have $\mathcal{O}_{X_{\text{prokét}}}^+(U)/p^n \cong R^+/p^n$, and it is canonically almost isomorphic to $(\mathcal{O}_{X_{\text{prokét}}}^+/p^n)(U)$.*

(ii) *For each $n \in \mathbf{Z}_{\geq 1}$ and $i \in \mathbf{Z}_{\geq 1}$, $H^i(U, \mathcal{O}_{X_{\text{prokét}}}^+/p^n)$ is almost equal to zero. Consequently, $H^i(U, \widehat{\mathcal{O}}_{X_{\text{prokét}}}^+)$ is almost equal to zero.*

(iii) *We have $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+(U) \cong R^+$ and $\widehat{\mathcal{O}}_{X_{\text{prokét}}}(U) \cong R$. Moreover, $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+(U)$ is canonically isomorphic to the p -adic completion of $\mathcal{O}_{X_{\text{prokét}}}^+(U)$.*

Example A.1.16. We recall the following example from [Dia+19a, §6]. Let P be a sharp fs monoid. Consider

$$\mathbb{E} := \text{Spa}(\mathbf{C}_p\langle P \rangle, \mathcal{O}_{\mathbf{C}_p}\langle P \rangle)$$

equipped with the natural log structure modeled on chart P . (If $P = \mathbf{Z}_{\geq 0}^n$, then \mathbb{E} is just the n -dimensional unit disc in Example A.1.3.) For each $m \in \mathbf{Z}_{>0}$, let $\frac{1}{m}P$ denote the sharp fs monoid containing P such that the inclusion $P \hookrightarrow \frac{1}{m}P$ is isomorphic to the m -th multiple map $[m] : P \rightarrow P$. Define

$$\mathbb{E}_m := \text{Spa}(\mathbf{C}_p\langle \frac{1}{m}P \rangle, \mathcal{O}_{\mathbf{C}_p}\langle \frac{1}{m}P \rangle)$$

equipped with the natural log structure modeled on the chart $\frac{1}{m}P$. If $m|m'$, there is a natural finite Kummer étale morphism $\mathbb{E}_{m'} \rightarrow \mathbb{E}_m$ modeled on the chart $\frac{1}{m}P \hookrightarrow \frac{1}{m'}P$. According to [Dia+19a, Proposition 4.1.6], the morphism $\mathbb{E}_m \rightarrow \mathbb{E}$ is actually finite Kummer étale Galois with Galois group

$$\Gamma_{/m} := \text{Hom}\left(\left(\frac{1}{m}P\right)^{\text{gp}}/P^{\text{gp}}, \mu_\infty\right),$$

where μ_∞ denotes the group of all roots of unity in \mathbf{C}_p . Let $P_{\mathbf{Q}_{\geq 0}} := \varinjlim_m (\frac{1}{m}P)$. It turns out

$$\widetilde{\mathbb{E}} := \varprojlim_m \mathbb{E}_m \in \mathbb{E}_{\text{prokét}}$$

is a log affinoid perfectoid object, with associated perfectoid space

$$\widehat{\mathbb{E}} = \text{Spa}(\mathbf{C}_p\langle P_{\mathbf{Q}_{\geq 0}} \rangle, \mathcal{O}_{\mathbf{C}_p}\langle P_{\mathbf{Q}_{\geq 0}} \rangle).$$

□

Following [Dia+19a, Definition 6.1.2], a pro-Kummer étale cover $Y \rightarrow X$ is called a **Galois cover with (profinite) Galois group** G if there exists a presentation $Y = \varprojlim_i Y_i$ such that each $Y_i \rightarrow X$ is a finite Kummer étale cover with Galois group G_i and $G \cong \varprojlim_i G_i$.

For example, $\widetilde{\mathbb{E}}$ is a Galois cover over \mathbb{E} with profinite Galois group

$$\Gamma \cong \varprojlim_m \Gamma/m = \varprojlim_m \operatorname{Hom}((\frac{1}{m}P)^{\operatorname{gp}}/P^{\operatorname{gp}}, \mu_\infty) \cong \operatorname{Hom}(P_{\mathbb{Q} \geq 0}^{\operatorname{gp}}/P^{\operatorname{gp}}, \mu_\infty)$$

(cf. [Dia+19a, (6.1.4)]).

A.2 Calculation of $R^i \nu_* \widehat{\mathcal{O}}_{X_{\operatorname{prokét}}}$

In this subsection, we study the sheaves $R^i \nu_* \widehat{\mathcal{O}}_{X_{\operatorname{prokét}}}$ following the calculations in [Sch12], [Sch13], and [CHJ17], but in the context of log adic spaces. Throughout this subsection, X is an fs log adic space that is log smooth over $\operatorname{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ (cf. Definition A.1.6). We will omit the subscript “prokét” from $\mathcal{O}_{X_{\operatorname{prokét}}}$, $\mathcal{O}_{X_{\operatorname{prokét}}}^+$, $\widehat{\mathcal{O}}_{X_{\operatorname{prokét}}}$, $\widehat{\mathcal{O}}_{X_{\operatorname{prokét}}}^+$, etc., whenever this causes no confusion.

By definition, $R^i \nu_* \widehat{\mathcal{O}}_X$ (resp., $R^i \nu_* \widehat{\mathcal{O}}_X^+$) is the sheaf on $X_{\operatorname{két}}$ associated with the presheaf

$$U \mapsto H^i(U_{\operatorname{prokét}}, \widehat{\mathcal{O}}_X) \quad (\text{resp., } U \mapsto H^i(U_{\operatorname{prokét}}, \widehat{\mathcal{O}}_X^+)).$$

For every $U \in X_{\operatorname{két}}$, U is log smooth over $\operatorname{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. By [Dia+19a, Proposition 3.1.10], étale locally on U there exists a **toric chart** $U \rightarrow \mathbb{E} = \operatorname{Spa}(\mathbf{C}_p \langle P \rangle, \mathcal{O}_{\mathbf{C}_p \langle P \rangle})$ for some sharp fs monoid P , i.e., a strictly étale morphism $U \rightarrow \mathbb{E} = \operatorname{Spa}(\mathbf{C}_p \langle P \rangle, \mathcal{O}_{\mathbf{C}_p \langle P \rangle})$ that is a composition of rational localisations and finite étale morphisms. For such toric charts, we are able to calculate $H^i(U_{\operatorname{prokét}}, \widehat{\mathcal{O}}_X^+)$ and $H^i(U_{\operatorname{prokét}}, \widehat{\mathcal{O}}_X)$ in an explicit way.

Lemma A.2.1. *Suppose $U \in X_{\operatorname{két}}$ is equipped with a toric chart $U \rightarrow \mathbb{E}$ as above.*

(i) *For every $i \in \mathbf{Z}_{\geq 0}$ and $m \in \mathbf{Z}_{\geq 1}$, there is a natural injection*

$$H_{\operatorname{cts}}^i(\Gamma, \mathcal{O}_{X_{\operatorname{két}}}^+(U)/p^m)^a \hookrightarrow H^i(U_{\operatorname{prokét}}, \mathcal{O}_X^+/p^m)^a$$

with cokernel killed by p , where Γ is equipped with the profinite topology, $\mathcal{O}_{X_{\operatorname{két}}}^+(U)/p^m$ is equipped with the discrete topology, and Γ acts trivially on $\mathcal{O}_{X_{\operatorname{két}}}^+(U)/p^m$.

(ii) *For every $i \in \mathbf{Z}_{\geq 0}$, there is a natural injection*

$$H_{\operatorname{cts}}^i(\Gamma, \mathcal{O}_{X_{\operatorname{két}}}^+(U))^a \hookrightarrow H^i(U_{\operatorname{prokét}}, \widehat{\mathcal{O}}_X^+)^a$$

with cokernel killed by p , where Γ is equipped with the profinite topology, $\mathcal{O}_{X_{\operatorname{két}}}^+(U)$ is equipped with the p -adic topology, and Γ acts trivially on $\mathcal{O}_{X_{\operatorname{két}}}^+(U)$. By inverting p , we obtain an isomorphism

$$H_{\operatorname{cts}}^i(\Gamma, \mathcal{O}_{X_{\operatorname{két}}}(U)) \xrightarrow{\sim} H^i(U_{\operatorname{prokét}}, \widehat{\mathcal{O}}_X).$$

(iii) *For every $i \in \mathbf{Z}_{\geq 0}$ and $m \in \mathbf{Z}_{\geq 1}$, by choosing an isomorphism $P^{\operatorname{gp}} \simeq \mathbf{Z}^n$, there is a natural almost injection*

$$\bigwedge^i (\mathcal{O}_{U_{\operatorname{két}}}^+/p^m)^n \hookrightarrow R^i \nu_* (\mathcal{O}_{U_{\operatorname{prokét}}}^+/p^m)$$

whose cokernel is killed by p . This induces a natural almost injection

$$\bigwedge^i (\mathcal{O}_{U_{\text{két}}}^+)^n \hookrightarrow R^i \nu_* \widehat{\mathcal{O}}_{U_{\text{prokét}}}^+$$

with cokernel killed by p . Inverting p , we obtain an isomorphism

$$\bigwedge^i (\mathcal{O}_{U_{\text{két}}}^+)^n \simeq R^i \nu_* \widehat{\mathcal{O}}_{U_{\text{prokét}}}.$$

In particular, the sheaf $R^i \nu_* \widehat{\mathcal{O}}_X$ is a locally free $\mathcal{O}_{X_{\text{két}}}$ -module.

Proof. (i) Recall the log affinoid perfectoid Galois cover $\widetilde{\mathbb{E}} \rightarrow \mathbb{E}$ (with profinite Galois group Γ) constructed in Example A.1.16. Consider

$$\widetilde{U} := U \times_{\mathbb{E}} \widetilde{\mathbb{E}} \in X_{\text{prokét}}.$$

By [Dia+19a, Lemma 5.3.8], \widetilde{U} is also log affinoid perfectoid and $\widetilde{U} \rightarrow U$ is a Galois cover with the same Galois group. We obtain the Cartan–Leray spectral sequence (see [CHJ17, Remark 2.25])

$$E_2^{i,j} = H_{\text{cts}}^i(\Gamma, H^j(\widetilde{U}, \mathcal{O}_X^+ / p^m)) \Rightarrow H^{i+j}(U_{\text{prokét}}, \mathcal{O}_X^+ / p^m).$$

By Proposition A.1.15 (ii), $H^j(\widetilde{U}, \mathcal{O}_X^+ / p^m)$ is almost zero for all $j \in \mathbf{Z}_{\geq 1}$. Therefore, we have an almost isomorphism

$$H^i(U_{\text{prokét}}, \mathcal{O}_X^+ / p^m)^a \simeq H_{\text{cts}}^i(\Gamma, (\mathcal{O}_X^+ / p^m)(\widetilde{U}))^a. \quad (10)$$

On the other hand, by [Dia+19a, Lemma 6.1.7, Remark 6.1.8], the natural morphism

$$H^i(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) \rightarrow H^i(\Gamma, (\mathcal{O}_X^+ / p^m)(\widetilde{U}))$$

is injective with cokernel killed by p for all $i \in \mathbf{Z}_{\geq 0}$. Combining this with the almost isomorphism (10), we obtain the desired almost injection.

(ii) By an almost version of [Sch13, Lemma 3.18] and Proposition A.1.15 (i) (ii), we see that the inverse system $\{\mathcal{O}_X^+ / p^m : m \in \mathbf{Z}_{\geq 1}\}$ has almost vanishing higher inverse limits on the pro-Kummer étale site. Therefore, we obtain almost isomorphisms

$$H^i(U_{\text{prokét}}, \widehat{\mathcal{O}}_X^+)^a \cong \varprojlim_m H^i(U_{\text{prokét}}, \mathcal{O}_X^+ / p^m)^a \simeq \varprojlim_m H_{\text{cts}}^i(\Gamma, (\mathcal{O}_X^+ / p^m)(\widetilde{U}))^a.$$

On the other hand, for every $i \in \mathbf{Z}_{\geq 0}$, we claim that there is a natural isomorphism

$$H_{\text{cts}}^i(\Gamma, \mathcal{O}_{X_{\text{két}}}^+(U)) \cong \varprojlim_m H_{\text{cts}}^i(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)).$$

Indeed, by the same arguments as in the proof of [NSW08, Theorem 2.7.5], there is a short exact sequence

$$0 \rightarrow R^1 \varprojlim_m H_{\text{cts}}^{i-1}(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) \rightarrow H_{\text{cts}}^i(\Gamma, \mathcal{O}_{X_{\text{két}}}^+(U)) \rightarrow \varprojlim_m H_{\text{cts}}^i(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) \rightarrow 0.$$

It suffices to show that

$$R^1 \varprojlim_m H_{\text{cts}}^{i-1}(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) = 0.$$

Notice that P^{gp} is a finitely generated torsion-free abelian group. By choosing a \mathbf{Z} -basis of P^{gp} , we obtain an isomorphism $\Gamma \cong \widehat{\mathbf{Z}}(1)^n$ of profinite groups which induces an isomorphism

$$H_{\text{cts}}^{i-1}(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) \simeq \bigwedge^{i-1} (\mathcal{O}_{X_{\text{két}}}^+ / p^m)^n.$$

Thus, for every $m' > m$, the transition map

$$H_{\text{cts}}^{i-1}(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^{m'})(U)) \rightarrow H_{\text{cts}}^{i-1}(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U))$$

is a surjection. Hence, the inverse system $\{H_{\text{cts}}^{i-1}(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) : m \in \mathbf{Z}_{>0}\}$ satisfies the Mittag-Leffler condition which yields the desired vanishing of $R^1 \lim$.

Putting everything together, we obtain a natural injection

$$H_{\text{cts}}^i(\Gamma, \mathcal{O}_{X_{\text{két}}}^+(U))^a \cong \varprojlim_m H_{\text{cts}}^i(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U))^a \hookrightarrow \varprojlim_m H_{\text{cts}}^i(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(\widetilde{U}))^a \cong H^i(U_{\text{prokét}}, \widehat{\mathcal{O}}_X^+)^a$$

whose cokernel is killed by p , as desired.

- (iii) We show that the restriction of $R^i \nu_* \widehat{\mathcal{O}}_X$ on $U_{\text{két}}$ is isomorphic to the free $\mathcal{O}_{U_{\text{két}}}$ -module $\bigwedge^i (\mathcal{O}_{U_{\text{két}}})^n$. In fact, as a byproduct of the computation above, we have isomorphisms (depending on the fixed choice of the identification $\Gamma \cong \widehat{\mathbf{Z}}(1)^n$)

$$H_{\text{cts}}^i(\Gamma, \mathcal{O}_{X_{\text{két}}}^+(U)) \cong \varprojlim_m H_{\text{cts}}^i(\Gamma, (\mathcal{O}_{X_{\text{két}}}^+ / p^m)(U)) \simeq \varprojlim_m \bigwedge^i (\mathcal{O}_{X_{\text{két}}}^+ / p^m)^n = \bigwedge^i (\mathcal{O}_{X_{\text{két}}}^+(U))^n.$$

Inverting p , we obtain an isomorphism

$$H^i(U_{\text{prokét}}, \widehat{\mathcal{O}}_X) \simeq H_{\text{cts}}^i(\Gamma, \mathcal{O}_{X_{\text{két}}}(U)) \simeq \bigwedge^i (\mathcal{O}_{X_{\text{két}}}(U))^n.$$

Consider $V \in U_{\text{két}}$ such that $V \rightarrow U$ admits a chart $P \rightarrow P'$ and such that $V \rightarrow U$ factors as

$$V \rightarrow U' \times_{U' \langle P \rangle} U' \langle P' \rangle \rightarrow U' \rightarrow U$$

where

- $U' \subset U$ is a strictly étale morphism which is a composition of finite étale morphisms and rational localisations;
- $P \rightarrow P'$ is isomorphic to the m -th multiple map $[m] : P \rightarrow P$;
- $V \rightarrow U' \times_{U' \langle P \rangle} U' \langle P' \rangle$ is a strictly étale morphism which is a composition of finite étale morphisms and rational localisations.

Notice that such a V admits a toric chart $V \rightarrow \mathbb{E}'$ where $\mathbb{E}' = \text{Spa}(\mathbf{C}_p \langle P' \rangle, \mathcal{O}_{\mathbf{C}_p} \langle P' \rangle)$. Repeating the argument above, we arrive at an isomorphism

$$H_{\text{cts}}^i(\Gamma', \mathcal{O}_{X_{\text{két}}}(V)) \simeq H^i(V_{\text{prokét}}, \widehat{\mathcal{O}}_X)$$

where

$$\Gamma' := \text{Hom}(P'_{\mathbf{Q}_{\geq 0}} / P'^{\text{gp}}, \mu_{\infty}).$$

In addition, the injection $P \rightarrow P'$ induces an injection $\Gamma' \rightarrow \Gamma$ which is isomorphic to multiplication by m . The fixed identification $\Gamma \cong \widehat{\mathbf{Z}}(1)^n$ then identifies $\Gamma' \rightarrow \Gamma$ with the m -th multiple map $[m] : \widehat{\mathbf{Z}}(1)^n \rightarrow \widehat{\mathbf{Z}}(1)^n$. We arrive at the following commutative diagram

$$\begin{array}{ccccc} \Lambda^i(\mathcal{O}_{X_{\text{két}}}(U))^n & \xrightarrow{\sim} & H_{\text{cts}}^i(\Gamma, \mathcal{O}_{X_{\text{két}}}(U)) & \xrightarrow{\sim} & H^i(U_{\text{prokét}}, \widehat{\mathcal{O}}_X) \\ \downarrow & & \downarrow & & \downarrow \\ \Lambda^i(\mathcal{O}_{X_{\text{két}}}(V))^n & \xrightarrow{\sim} & H_{\text{cts}}^i(\Gamma', \mathcal{O}_{X_{\text{két}}}(V)) & \xrightarrow{\sim} & H^i(V_{\text{prokét}}, \widehat{\mathcal{O}}_X) \end{array}$$

Finally, let \mathcal{B}_U denote the collection of such V 's. Notice that every Kummer map $P \rightarrow Q$ between sharp fs monoids factors through $[m] : P \rightarrow P$ for some $m \in \mathbf{Z}_{\geq 1}$. Hence, every $W \in U_{\text{két}}$ is covered by elements in \mathcal{B}_U . This is enough to conclude that the sheafification of the presheaf $W \rightarrow H^i(W_{\text{prokét}}, \widehat{\mathcal{O}}_X)$ on $U_{\text{két}}$ is isomorphic to the free sheaf $\Lambda^i(\mathcal{O}_{U_{\text{két}}})^n$. This completes the proof. \square

We also provide a coordinate-free description of $R^i \nu_* \widehat{\mathcal{O}}_X$. The following result is a logarithmic version of [Sch12, Proposition 3.23, Lemma 3.24].

Lemma A.2.2. *For every $n \in \mathbf{Z}_{\geq 1}$, let μ_{p^n} be the sheaf of p^n -th roots of unity on $X_{\text{két}}$. Consider the \mathbf{Z}_p -local system $\mathbf{Z}_p(1) := \varprojlim_n \mu_{p^n}$ on $X_{\text{két}}$ and let $\widehat{\mathbf{Z}}_p(1) := \nu^{-1} \mathbf{Z}_p(1)$ be the associated $\widehat{\mathbf{Z}}_p$ -local system on $X_{\text{prokét}}$ (cf. [Dia+19a, Definition 6.3.2]). The short exact sequence*

$$0 \rightarrow \widehat{\mathbf{Z}}_p(1) \rightarrow \varprojlim_{x \mapsto x^p} \mathcal{M}_{X_{\text{prokét}}} \rightarrow \mathcal{M}_{X_{\text{prokét}}} \rightarrow 0$$

induces a boundary map

$$\mathcal{M}_{X_{\text{két}}} = \nu_* \mathcal{M}_{X_{\text{prokét}}} \rightarrow R^1 \nu_* \widehat{\mathbf{Z}}_p(1).$$

Then, there exists a unique $\mathcal{O}_{X_{\text{két}}}$ -linear morphism $\Omega_{X_{\text{két}}}^{\log, 1} \rightarrow R^1 \nu_* \widehat{\mathcal{O}}_X(1)$ such that the diagram

$$\begin{array}{ccc} \mathcal{M}_{X_{\text{két}}} & \longrightarrow & R^1 \nu_* \widehat{\mathbf{Z}}_p(1) \\ \downarrow \text{dlog} & & \downarrow \\ \Omega_{X_{\text{két}}}^{\log, 1} & \longrightarrow & R^1 \nu_* \widehat{\mathcal{O}}_X(1) \end{array}$$

is commutative, where $\Omega_{X_{\text{két}}}^{\log, 1}$ is the sheaf of log differentials defined in [Dia+19a, Definition 3.2.25].

Moreover, this morphism is an isomorphism. As a corollary, by taking cup product and exterior product, we obtain a canonical isomorphism $R^i \nu_* \widehat{\mathcal{O}}_X \cong \Omega_{X_{\text{két}}}^{\log, i}(-i)$ for every $i \geq 1$.

Proof. The proof follows almost verbatim from the proof of [Sch12, Lemma 3.24], except that we have to replace the short exact sequence in [Sch13, Corollary 6.14] by the short exact sequence in [Dia+19b, Corollary 2.4.5]. Here we only give a sketch of the proof.

Firstly, since the question is étale local, we may assume that X admits a toric chart $X \rightarrow \text{Spa}(\mathbf{C}_p\langle P \rangle, \mathcal{O}_{\mathbf{C}_p}\langle P \rangle)$ for some sharp fs monoid P . Secondly, the desired $\mathcal{O}_{X_{\text{két}}}$ -linear morphism, if exists, must be unique because $\Omega_{X_{\text{két}}}^{\log, 1}$ is a locally free $\mathcal{O}_{X_{\text{két}}}$ -module generated by the image of dlog. It remains to show the existence.

Consider the map of short exact sequences

$$\begin{array}{ccccccc} 0 \longrightarrow & \widehat{\mathbf{Z}}_p(1) & \longrightarrow & \varprojlim_{x \mapsto x^p} \mathcal{M}_{X_{\text{prokét}}} & \longrightarrow & \mathcal{M}_{X_{\text{prokét}}} & \longrightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow \text{dlog} & \\ 0 \longrightarrow & \widehat{\mathcal{O}}_X(1) & \longrightarrow & \text{gr}^1 \mathcal{O}_{\text{dR}, \log, X}^+ & \longrightarrow & \widehat{\mathcal{O}}_X \otimes_{\mathcal{O}_{X_{\text{két}}}} \Omega_{X_{\text{két}}}^{\log, 1} & \longrightarrow 0 \end{array}$$

where the lower sequence is from [Dia+19b, Corollary 2.4.5]. The vertical map in the middle sends an element $a \in \varprojlim_{x \mapsto x^p} \mathcal{M}_{X_{\text{prokét}}}$ to

$$\log(\mathbf{e}^a) := - \sum_{n=1}^{\infty} \frac{1}{n} (1 - \mathbf{e}^a)^n \in \text{Fil}^1 \mathcal{O}_{\text{dR}, \log, X}^+$$

and hence maps to $\text{gr}^1 \mathcal{O}_{\text{dR}, \log, X}^+$ (see [Dia+19b, §2.2] for the definition of \mathbf{e}^a). It is straightforward to check the commutativity of the diagram. This diagram then induces a diagram of boundary maps

$$\begin{array}{ccc} H^0(X_{\text{prokét}}, \mathcal{M}_{X_{\text{prokét}}}) & \longrightarrow & H^1(X_{\text{prokét}}, \widehat{\mathbf{Z}}_p(1)) \\ \downarrow \text{dlog} & & \downarrow \\ H^0(X_{\text{prokét}}, \widehat{\mathcal{O}}_X \otimes_{\mathcal{O}_{X_{\text{két}}}} \Omega_{X_{\text{két}}}^{\log, 1}) & \longrightarrow & H^1(X_{\text{prokét}}, \widehat{\mathcal{O}}_X(1)) \end{array}$$

which provides the desired morphism.

To check that this map is an isomorphism, we fix an identification $P^{\text{gp}} \simeq \mathbf{Z}^n = \bigoplus_{j=1}^n \mathbf{Z} e_j$ which induces an isomorphism

$$\Gamma := \text{Hom}(P_{\mathbf{Q}_{\geq 0}}^{\text{gp}} / P^{\text{gp}}, \mu_{\infty}) \simeq \widehat{\mathbf{Z}}(1)^n.$$

Notice that each e_j can be written as a \mathbf{Z} -linear combination of elements in P ; i.e., $e_j = \sum_{t=1}^m a_t p_t$ for some $a_t \in \mathbf{Z}$ and $p_t \in P$. We define $\text{dlog}(e_j) := \sum_{t=1}^m a_t \text{dlog}(p_t)$ where we have identified p_t with its image in $\mathcal{M}_{X_{\text{két}}}$. One checks that $\text{dlog}(e_j)$ is independent of the choice of the \mathbf{Z} -linear combination and the $\text{dlog}(e_j)$'s form a basis for the free $\mathcal{O}_{X_{\text{két}}}$ -module $\Omega_{X_{\text{két}}}^{\log, 1}$.

On the other hand, by the computation in the proof of Lemma A.2.1, the identification $\Gamma \simeq \widehat{\mathbf{Z}}(1)^n$ induces an isomorphism $R^1 \nu_* \widehat{\mathcal{O}}_X \simeq \mathcal{O}_{X_{\text{két}}}^n = \bigoplus_{j=1}^n \mathcal{O}_{X_{\text{két}}} \epsilon_j$. Direct computation shows that the map $\Omega_{X_{\text{két}}}^{\log, 1} \rightarrow R^1 \nu_* \widehat{\mathcal{O}}_X$ sends $\text{dlog}(e_j)$ to ϵ_j , for every $j = 1, \dots, n$. This finishes the proof. \square

To wrap up this subsection, we include a logarithmic analogue of [CHJ17, Proposition 6.8] which suggests that the calculation of $R^i \nu_* \widehat{\mathcal{O}}_X$ is compatible with the “mixed completed tensor”.

Proposition A.2.3. *Let M be a profinite flat \mathcal{O}_K -module in the sense of [CHJ17, Definition 6.1]. Then there is a canonical isomorphism*

$$R^i \nu_* (\widehat{\mathcal{O}}_X \widehat{\otimes} M) \simeq (R^i \nu_* \widehat{\mathcal{O}}_X) \widehat{\otimes} M,$$

where $\widehat{\otimes}$ stands for the “mixed completed tensor” in the sense of [CHJ17, Definition 6.6]. Here, the mixed completed tensor on the right hand side is with respect to the subsheaf $\text{Im}(R^i \nu_* \widehat{\mathcal{O}}_X^+ \rightarrow R^i \nu_* \widehat{\mathcal{O}}_X) \subset R^i \nu_* \widehat{\mathcal{O}}_X$.

Consequently, by Lemma A.2.2, we have

$$R^i \nu_* (\widehat{\mathcal{O}}_X \widehat{\otimes} M) \simeq \Omega_{X_{\text{két}}}^{\log, i}(-i) \widehat{\otimes} M.$$

Proof. The proof follows verbatim as in the proof of [CHJ17, Proposition 6.8] as long as we replace [CHJ17, Lemma 6.11(1)(2)] by Lemma A.2.1. \square

A.3 Banach sheaves and a (generalised) projection formula

In this subsection, we introduce the notion of *Banach sheaves* on the Kummer étale topology of a log adic space, generalising the ones studied in [AIP15, §A] and [BP20, §2]. Then, for certain *admissible* Banach sheaves, we prove a projection formula which will be used in the main body of the paper.

Recall from Definition 3.1.1 that a *small \mathbf{Z}_p -algebra* is a p -torsion free reduced ring R which is also a finite $\mathbf{Z}_p[[T_1, \dots, T_d]]$ -algebra for some $d \in \mathbf{Z}_{\geq 0}$. It is a profinite flat \mathbf{Z}_p -module in the sense of [CHJ17, Definition 6.1]. In particular, there exists a set of elements $\{e_\sigma : \sigma \in \Sigma\}$ in R such that $R \simeq \prod_{\sigma \in \Sigma} \mathbf{Z}_p e_\sigma$ equipped with the product topology. This set of elements $\{e_\sigma : \sigma \in \Sigma\}$ is called a *pseudo-basis* for R . Moreover, R is equipped with an adic profinite topology and is complete with respect to the p -adic topology.

Throughout this subsection, we keep the following notations:

- Let R be a fixed small \mathbf{Z}_p -algebra and let \mathfrak{a} be a fixed ideal of definition containing p .
- All (log) adic spaces are assumed to be reduced and quasi-separated. In particular, X either stands for a locally noetherian reduced adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ or a locally noetherian reduced fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. In the second case, we use X_{an} to denote the underlying adic space of X .
- We adopt the notation of “mixed completed tensors” $-\widehat{\otimes}' R$ and $-\widehat{\otimes} R$ as in Definition 3.1.3.

Lemma A.3.1. (i) Let X be a locally noetherian adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Then the presheaf $\mathcal{O}_X^+ \widehat{\otimes}' R$ (resp., $\mathcal{O}_X \widehat{\otimes} R$) sending any quasi-compact open subset $U \subset X$ to $\mathcal{O}_X^+(U) \widehat{\otimes}' R$ (resp., $\mathcal{O}_X(U) \widehat{\otimes} R$) is a sheaf. In particular, $\mathcal{O}_X \widehat{\otimes} R$ is a sheaf of Banach \mathbf{C}_p -algebras.

(ii) Let X be a locally noetherian fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Then the presheaf $\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R$ (resp., $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$) sending any quasi-compact $U \in X_{\text{két}}$ to $\mathcal{O}_{X_{\text{két}}}^+(U) \widehat{\otimes}' R$ (resp., $\mathcal{O}_{X_{\text{két}}}(U) \widehat{\otimes} R$) is a sheaf. In particular, $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ is a sheaf of Banach \mathbf{C}_p -algebras.

(iii) Let X be a locally noetherian fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Then the presheaf $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \widehat{\otimes}' R$ (resp., $\widehat{\mathcal{O}}_{X_{\text{prokét}}} \widehat{\otimes} R$) sending any qcqs $U \in X_{\text{prokét}}$ to $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+(U) \widehat{\otimes}' R$ (resp., $\widehat{\mathcal{O}}_{X_{\text{prokét}}}(U) \widehat{\otimes} R$) is a sheaf. In particular, $\widehat{\mathcal{O}}_{X_{\text{prokét}}} \widehat{\otimes} R$ is a sheaf of Banach \mathbf{C}_p -algebras.

Proof. Choosing a presentation $R \simeq \prod_{\sigma \in \Sigma} \mathbf{Z}_p e_\sigma$ and using [CHJ17, Proposition 6.4], the statements reduce to the sheafiness of the corresponding structure presheaves. \square

Definition A.3.2. Let B be a Banach \mathbf{Q}_p -algebra and let B_0 be an open and bounded \mathbf{Z}_p -submodule.

- (i) A topological B -module M is called a **Banach B -module** if there exists an open bounded B_0 -submodule M_0 which is p -adically complete and separated such that $M = M_0[1/p]$.
- (ii) Let J be an index set. Consider the B -module $B(J)$ consisting of sequences $\{b_j : j \in J\}$ which converge to 0 with respect to the filter in J of the complement of the finite subsets of J . Then $B(J)$ is a Banach B -module. Indeed, let $B_0(J)$ be the p -adic completion of the free B_0 -module $\bigoplus_{j \in J} B_0$. Then we have $B(J) \simeq B_0(J)[1/p]$.
- (iii) A topological B -module M is called an **orthonormalisable Banach B -module** (or, **ON-able Banach B -module** for short) if there exists a topological isomorphism $M \simeq B(J)$ for some index set J . A topological B -module M is called a **projective Banach B -module** if it is a direct summand (as a topological B -module) inside an orthonormalisable Banach B -module.

Definition A.3.3. Let X be a locally noetherian adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$.

- (i) A sheaf of topological $\mathcal{O}_X \widehat{\otimes} R$ -modules \mathcal{F} is called a **Banach sheaf of $\mathcal{O}_X \widehat{\otimes} R$ -modules** if
 - for every quasi-compact open subset $U \subset X$, $\mathcal{F}(U)$ is a Banach $\mathcal{O}_X(U) \widehat{\otimes} R$ -module;

- there exists an affinoid open covering $\mathfrak{U} = \{U_i : i \in I\}$ of X such that for every $i \in I$ and every affinoid open subset $V \subset U_i$, the continuous restriction map

$$\mathcal{F}(U_i) \otimes_{\mathcal{O}_X(U_i)} \mathcal{O}_X(V) \rightarrow \mathcal{F}(V)$$

induces a topological isomorphism

$$\mathcal{F}(U_i) \widehat{\otimes}_{\mathcal{O}_X(U_i)} \mathcal{O}_X(V) \rightarrow \mathcal{F}(V)$$

where the completion is with respect to the p -adic topology. Such a covering \mathfrak{U} is called an **atlas** of \mathcal{F} .

- (ii) A sheaf \mathcal{F} as in (i) is called a **projective Banach sheaf of $\mathcal{O}_X \widehat{\otimes} R$ -modules** if there exists an atlas $\mathfrak{U} = \{U_i : i \in I\}$ such that $\mathcal{F}(U_i)$'s are projective Banach $\mathcal{O}_X(U_i) \widehat{\otimes} R$ -modules.
- (iii) A morphism between Banach sheaves of $\mathcal{O}_X \widehat{\otimes} R$ -modules is a continuous map of sheaves of topological $\mathcal{O}_X \widehat{\otimes} R$ -modules.
- (iv) Let \mathcal{F} be a Banach sheaf of $\mathcal{O}_X \widehat{\otimes} R$ -modules as in (i). An **integral model** of \mathcal{F} is a subsheaf \mathcal{F}^+ of $\mathcal{O}_X^+ \widehat{\otimes}' R$ -modules such that
 - for every quasi-compact open $U \subset X$, $\mathcal{F}^+(U)$ is open and bounded in $\mathcal{F}(U)$;
 - $\mathcal{F} = \mathcal{F}^+[1/p]$;
 - there exists an atlas $\mathfrak{U} = \{U_i : i \in I\}$ of \mathcal{F} such that, for every $i \in I$ and every affinoid open subset $V \subset U_i$, the canonical map

$$\mathcal{F}^+(U_i) \widehat{\otimes}_{\mathcal{O}_X^+(U_i)} \mathcal{O}_X^+(V) \rightarrow \mathcal{F}^+(V)$$

is an isomorphism, where the completion is with respect to the p -adic topology.

We are also interested in a Kummer étale version of Banach sheaves.

Definition A.3.4. Let X be a locally noetherian fs log adic space of $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$.

- (i) A sheaf of topological $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules \mathcal{F} is called a **Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules** if
 - for every quasi-compact open $U \in X_{\text{két}}$, $\mathcal{F}(U)$ is a Banach $\mathcal{O}_{X_{\text{két}}}(U) \widehat{\otimes} R$ -module;
 - there exists an Kummer étale covering $\mathfrak{U} = \{U_i : i \in I\}$ of X by affinoid U_i 's such that for every Kummer étale map $V \rightarrow U_i$ with affinoid V , the continuous restriction map

$$\mathcal{F}(U_i) \otimes_{\mathcal{O}_{X_{\text{két}}}(U_i)} \mathcal{O}_{X_{\text{két}}}(V) \rightarrow \mathcal{F}(V)$$

induces a topological isomorphism

$$\mathcal{F}(U_i) \widehat{\otimes}_{\mathcal{O}_{X_{\text{két}}}(U_i)} \mathcal{O}_{X_{\text{két}}}(V) \rightarrow \mathcal{F}(V)$$

where the completion is with respect to the p -adic topology. Such a covering \mathfrak{U} is called a **Kummer étale atlas** of \mathcal{F} .

- (ii) A sheaf as in (i) is called a **projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules** if there exists a Kummer étale atlas $\mathfrak{U} = \{U_i : i \in I\}$ such that $\mathcal{F}(U_i)$'s are projective Banach $\mathcal{O}_{X_{\text{két}}}(U_i) \widehat{\otimes} R$ -modules.

(iii) A morphism between Kummer étale Banach sheaves of $\mathcal{O}_{X_{k\acute{e}t}} \widehat{\otimes} R$ -modules is a continuous map of topological $\mathcal{O}_{X_{k\acute{e}t}} \widehat{\otimes} R$ -modules.

(iv) Let \mathcal{F} be a Kummer étale Banach sheaf of $\mathcal{O}_{X_{k\acute{e}t}} \widehat{\otimes} R$ -modules as in (i). An **integral model** of \mathcal{F} is a subsheaf \mathcal{F}^+ of $\mathcal{O}_{X_{k\acute{e}t}}^+ \widehat{\otimes} R$ -modules such that

- for every quasi-compact $U \in X_{k\acute{e}t}$, $\mathcal{F}^+(U)$ is open and bounded in $\mathcal{F}(U)$;
- $\mathcal{F} = \mathcal{F}^+[1/p]$;
- there exists a Kummer étale atlas $\mathfrak{U} = \{U_i : i \in I\}$ of \mathcal{F} such that, for every $i \in I$ and every affinoid $V \in U_{i,k\acute{e}t}$, the canonical map

$$\mathcal{F}^+(U_i) \widehat{\otimes}_{\mathcal{O}_{X_{k\acute{e}t}}^+(U_i)} \mathcal{O}_{X_{k\acute{e}t}}^+(V) \rightarrow \mathcal{F}^+(V)$$

is an isomorphism, where the completion is with respect to the p -adic topology.

Clearly, an analytic refinement of an atlas (resp., a Kummer étale refinement of a Kummer étale atlas) is also an atlas (resp., a Kummer étale atlas). Also notice that it is not true that a Banach sheaf (resp., Kummer étale Banach sheaf) on an affinoid adic space (resp., affinoid log adic space) is the sheaf associated with its global section. Nonetheless, we have the following result.

Lemma A.3.5. *Let (A, A^+) be a complete reduced Tate algebra over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and let M be a projective Banach $A \widehat{\otimes} R$ -module.*

- (i) *Let $X = \mathrm{Spa}(A, A^+)$ be the associated adic space. Then the presheaf $M \widehat{\otimes}_A \mathcal{O}_X$ sending an affinoid open subset $\mathrm{Spa}(B, B^+) \subset X$ to $M \widehat{\otimes}_A B$ is a sheaf.*
- (ii) *Suppose $X = \mathrm{Spa}(A, A^+)$ is equipped with an fs log structure. Then the presheaf $M \widehat{\otimes}_A \mathcal{O}_{X_{k\acute{e}t}}$ sending an affinoid open subset $\mathrm{Spa}(B, B^+) \in X_{k\acute{e}t}$ to $M \widehat{\otimes}_A B$ is a sheaf.*

Proof. It immediately reduces to the case where M is an orthonormalisable Banach $A \widehat{\otimes} R$ -module; i.e., $M \simeq (A \widehat{\otimes} R)(J)$ for some index set J . It then reduces to the case where $|J| = 1$. Then the lemma follows from Lemma A.3.1. \square

As a corollary, one can associate a projective Kummer étale Banach sheaf with every projective Banach sheaf.

Corollary A.3.6. *Let X be a locally noetherian fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and let \mathcal{F} be a projective Banach sheaf of $\mathcal{O}_{X_{\mathrm{an}}} \widehat{\otimes} R$ -modules with atlas $\mathfrak{U} = \{U_i : i \in I\}$. Suppose \mathcal{F} admits an integral model \mathcal{F}^+ . Consider the p -adically completed sheaf of $\mathcal{O}_{X_{k\acute{e}t}}$ -modules $\mathcal{F}_{k\acute{e}t}$ associated with \mathcal{F} ; namely,*

$$\mathcal{F}_{k\acute{e}t} := \left(\varprojlim_m \mathcal{F}^+ \otimes_{\mathcal{O}_{X_{\mathrm{an}}}^+} \mathcal{O}_{X_{k\acute{e}t}}^+ / p^m \right) \left[\frac{1}{p} \right].$$

Then $\mathcal{F}_{k\acute{e}t}$ is a projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{k\acute{e}t}} \widehat{\otimes} R$ -modules with Kummer étale atlas $\mathfrak{U} = \{U_i : i \in I\}$, where each U_i is equipped with the induced log structure from X . Moreover, for every affinoid $V \in U_{i,k\acute{e}t}$, we have

$$\mathcal{F}_{k\acute{e}t}(V) \cong \mathcal{F}(U_i) \widehat{\otimes}_{\mathcal{O}_{X_{\mathrm{an}}}(U_i)} \mathcal{O}_{X_{k\acute{e}t}}(V).$$

We need an easy lemma.

Lemma A.3.7. *Let X be a locally noetherian fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and let $\mathfrak{U} = \{U_i : i \in I\}$ be a Kummer étale covering of X by affinoid U_i 's. Consider the full subcategory $\mathcal{B}_{\mathfrak{U}}$ of $X_{\text{két}}$ consisting of those affinoid $V \in X_{\text{két}}$ such that the map $V \rightarrow X$ factors through $V \rightarrow U_i \rightarrow X$ for some $i \in I$. Then $\mathcal{B}_{\mathfrak{U}}$ forms a basis for the site $X_{\text{két}}$.*

Proof. We have to prove that every $U \in X_{\text{két}}$ admits a covering by such V 's and that $\mathcal{B}_{\mathfrak{U}}$ is closed under fibred products. Both statements are clear. \square

Proof of Corollary A.3.6. Let $\mathcal{B}_{\mathfrak{U}}$ be the basis of $X_{\text{két}}$ as in Lemma A.3.7 associated with the covering $\mathfrak{U} = \{U_i : i \in I\}$. It suffices to show that the assignment

$$V \mapsto \mathcal{F}(U_i) \widehat{\otimes}_{\mathcal{O}_{X_{\text{an}}}(U_i)} \mathcal{O}_{X_{\text{két}}}(V),$$

for every $V \in \mathcal{B}_{\mathfrak{U}}$ which factors through $V \rightarrow U_i \rightarrow X$, defines a sheaf on $\mathcal{B}_{\mathfrak{U}}$. (Notice that this assignment is independent of the choice of i and hence well-defined.) The sheafness of this assignment follows from Lemma A.3.5 and the sheafness of \mathcal{F} . \square

In what follows, we are interested in those Kummer étale Banach sheaves that are “admissible”. Let us first recall the notion of coherent sheaves on a ringed site.

Definition A.3.8. *Let (Z, \mathcal{O}_Z) be a ringed site. A sheaf of \mathcal{O}_Z -modules \mathcal{F} is called a **coherent \mathcal{O}_Z -module** if there exists a covering $\mathfrak{U} = \{U_i : i \in I\}$ for Z such that for every $i \in I$, there exist positive integers m, n , and an exact sequence of $\mathcal{O}_Z|_{U_i}$ -modules*

$$\bigoplus_{j=1}^m \mathcal{O}_Z|_{U_i} \rightarrow \bigoplus_{k=1}^n \mathcal{O}_Z|_{U_i} \rightarrow \mathcal{F}|_{U_i} \rightarrow 0.$$

*In this situation, we say that \mathcal{F} is a coherent \mathcal{O}_Z -module **subject to the covering \mathfrak{U}** .*

We will apply this definition to the ringed site $(X_{\text{két}}, \mathcal{O}_{X_{\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m))$.

Definition A.3.9. *Let X be a locally noetherian fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and let \mathcal{F} be a projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules. Suppose it admits an integral model \mathcal{F}^+ and, for every $m \in \mathbf{Z}_{\geq 1}$, we write $\mathcal{F}_m^+ := \mathcal{F}^+ / \mathfrak{a}^m$. We say that \mathcal{F} is **admissible** if there exist*

- *a Kummer étale atlas $\mathfrak{U} = \{U_i : i \in I\}$ of X such that each $\mathcal{F}^+(U_i)$ is the p -adic completion of a free $\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes} R$ -module; and*
- *for every $m \in \mathbf{Z}_{\geq 1}$ and $d \in \mathbf{Z}_{\geq 1}$, a subsheaf $\mathcal{F}_{m,d}^+ \subset \mathcal{F}_m^+$ which is a coherent $\mathcal{O}_{X_{\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)$ -module subject to the covering \mathfrak{U} ,*

such that we have $\mathcal{F}^+ \cong \varprojlim_m \mathcal{F}_m^+$ and $\mathcal{F}_m^+ \cong \varinjlim_d \mathcal{F}_{m,d}^+$ for every $m \in \mathbf{Z}_{\geq 1}$.

*Such a Kummer étale atlas is called an **admissible atlas** for \mathcal{F} .*

Lemma A.3.10. *Let $h : Y \rightarrow X$ be a finite Kummer étale morphism between locally noetherian fs log adic spaces over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Suppose \mathcal{F} is an admissible Kummer étale Banach sheaf of $\mathcal{O}_{Y_{\text{két}}} \widehat{\otimes} R$ -modules. Then $h_* \mathcal{F}$ is an admissible Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules.*

Proof. Suppose $\mathfrak{U} = \{U_i : i \in I\}$ is an admissible atlas for \mathcal{F} on Y . By Definition A.1.7 and [Dia+19a, Proposition 4.1.6], the finite Kummer étale morphism $h : Y \rightarrow X$ is, Kummer étale locally on X , isomorphic to a direct sum of isomorphisms. Therefore, one can find an affinoid Kummer étale covering $\{V_j : j \in J\}$ of X such that, for every $i \in I$ and $j \in J$, $U_i \times_X V_j$ is isomorphic to a disjoint union of finite copies of U_i 's. Consequently, the Kummer étale covering $\mathfrak{V} = \{U_i \times_X V_j : i \in I, j \in J\}$ is a desired admissible atlas for $h_* \mathcal{F}$. \square

If \mathcal{F} is a Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules with an integral structure \mathcal{F}^+ . We write

$$\widehat{\mathcal{F}}^+ := \varprojlim_m \left(\mathcal{F}^+ \otimes_{\mathcal{O}_{X_{\text{két}}}}^+ \mathcal{O}_{X_{\text{prokét}}}^+ / p^m \right) \cong \varprojlim_m \left(\mathcal{F}^+ \otimes_{(\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R)} (\mathcal{O}_{X_{\text{prokét}}}^+ \widehat{\otimes}' R) / p^m \right)$$

and $\widehat{\mathcal{F}} := \widehat{\mathcal{F}}^+ [1/p]$. They are sheaves of $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \widehat{\otimes}' R$ -modules and $\widehat{\mathcal{O}}_{X_{\text{prokét}}} \widehat{\otimes} R$ -modules, respectively.

Recall the natural projection of sites $\nu : X_{\text{prokét}} \rightarrow X_{\text{két}}$. The main result of this subsection is the following.

Proposition A.3.11 (Generalised projection formula). *Let X be a locally noetherian fs log adic space which is log smooth over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ and let \mathcal{F} be a projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules. Suppose \mathcal{F} is admissible. Then, for every $j \in \mathbf{Z}_{\geq 0}$, there is a natural isomorphism of Kummer étale Banach sheaves of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules*

$$\mathcal{F} \otimes_{\mathcal{O}_{X_{\text{két}}}} R^j \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}} \xrightarrow{\sim} R^j \nu_* \widehat{\mathcal{F}}.$$

To prove the proposition, we need some preparations.

Lemma A.3.12. *Let X be a locally noetherian fs log adic space over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Let \mathcal{H} be an $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \widehat{\otimes} R$ -module and let $\mathcal{H}_m := \mathcal{H} / \mathfrak{a}^m$ for every $m \in \mathbf{Z}_{\geq 1}$. Suppose*

- $\mathcal{H} = \varprojlim_m \mathcal{H}_m$; and
- for every $m \in \mathbf{Z}_{\geq 1}$, there exists a sequence of finite free $\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R / \mathfrak{a}^m)$ -submodules $\{\mathcal{H}_{m,d} : d \in \mathbf{Z}_{\geq 0}\}$ of \mathcal{H}_m such that $\mathcal{H}_m \cong \varinjlim_d \mathcal{H}_{m,d}$.

Then, for every $j \in \mathbf{Z}_{\geq 0}$, the natural map

$$R^j \nu_* \mathcal{H} \rightarrow \varprojlim_m R^j \nu_* \mathcal{H}_m$$

is an almost isomorphism.

Proof. We have to show the almost vanishing of the higher inverse limit $R^j \varprojlim_m \mathcal{H}_m$. Applying an almost version of [Sch13, Lemma 3.18], it suffices to show that, for every log affinoid perfectoid object $U \in X_{\text{prokét}}$, there are almost isomorphisms

$$R^1 \varprojlim_m \mathcal{H}_m(U)^a = 0$$

and

$$H^j(U, \mathcal{H}_m)^a = 0$$

for every $j \in \mathbf{Z}_{\geq 0}$. The first almost vanishing follows from the Mittag-Leffler condition. To obtain the second almost isomorphism, observe that

$$H^j(U, \mathcal{H}_m) \cong \varinjlim_d H^j(U, \mathcal{H}_{m,d}).$$

and each $H^j(U, \mathcal{H}_{m,d})$ is almost zero by [Dia+19a, Theorem 5.4.3]. \square

Lemma A.3.13. *Let X be a locally noetherian fs log adic space which is log smooth over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. If \mathcal{G} is an projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules, then, for every $j \in \mathbf{Z}_{\geq 0}$, the sheaf $R^j \nu_* \widehat{\mathcal{G}}$ is also a projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules.*

Proof. By considering a Kummer étale atlas for \mathcal{G} and writing $R \simeq \prod_{\sigma \in \Sigma} \mathbf{Z}_p e_\sigma$, we immediately reduce to the case where

- X is affinoid and admits a toric chart $X \rightarrow \mathbb{E} = \mathrm{Spa}(\mathbf{C}_p \langle P \rangle, \mathcal{O}_{\mathbf{C}_p} \langle P \rangle)$ for some sharp fs monoid P ;
- $R = \mathbf{Z}_p$ and $\mathfrak{a} = (p)$;
- \mathcal{G} is globally projective; *i.e.*, $\mathcal{G}(X)$ is a projective Banach $\mathcal{O}_{X_{\mathrm{két}}}(X)$ -module and for every affinoid $U \in X_{\mathrm{két}}$, we have a natural isomorphism

$$\mathcal{G}(X) \widehat{\otimes}_{\mathcal{O}_{X_{\mathrm{két}}}(X)} \mathcal{O}_{X_{\mathrm{két}}}(U) \xrightarrow{\sim} \mathcal{G}(U).$$

We further reduce to the case where \mathcal{G} is globally orthonormalisable; namely, $\mathcal{G} \simeq \mathcal{O}_{X_{\mathrm{két}}}(J)$ for some index set J . Let \mathcal{G}^+ be the p -adic completion of the free $\mathcal{O}_{X_{\mathrm{két}}}^+$ -module $\bigoplus_J \mathcal{O}_{X_{\mathrm{két}}}^+$ and let $\mathcal{G}_m^+ := \mathcal{G}^+ / p^m \simeq \bigoplus_J \mathcal{O}_{X_{\mathrm{két}}}^+ / p^m$. By Lemma A.3.12, we have a natural almost isomorphism

$$R^j \nu_* \widehat{\mathcal{G}}^+ \xrightarrow{\sim} \varprojlim_m R^j \nu_* \widehat{\mathcal{G}}_m^+$$

where $\widehat{\mathcal{G}}_m^+ = \widehat{\mathcal{G}}^+ / p^m \simeq \bigoplus_J \mathcal{O}_{X_{\mathrm{prokét}}}^+ / p^m$.

We claim that, in this case, the sheaf $R^j \nu_* \widehat{\mathcal{G}}$ is isomorphic to $(\bigwedge^j (\mathcal{O}_{X_{\mathrm{két}}})^n)(J)$ for some $n \in \mathbf{Z}_{\geq 1}$. For this, we follow the strategy as in the proof of Lemma A.2.1.

In order to be consistent with the notations in Lemma A.2.1, we write $U = X$. Consider the collection \mathcal{B}_U used in the proof of Lemma A.2.1. In particular, for every $V \in \mathcal{B}_U$, the map $V \rightarrow U$ admits a Kummer chart $P \rightarrow P'$ which is isomorphic to the m -th multiple map $[m] : P \rightarrow P$. Moreover, the injection $P \rightarrow P'$ induces an injection $\Gamma' \rightarrow \Gamma$. If we fix an identification $\Gamma \cong \widehat{\mathbf{Z}}(1)^n$, the injection $\Gamma' \rightarrow \Gamma$ can be identified with the m -th multiple map $[m] : \widehat{\mathbf{Z}}(1)^n \rightarrow \widehat{\mathbf{Z}}(1)^n$.

By the calculation in Lemma A.2.1, we obtain an almost injection

$$\left(\bigwedge^j (\mathcal{O}_{X_{\mathrm{két}}}^+ / p^m(V))^n \right)^a \simeq H^j(\Gamma, \mathcal{O}_{X_{\mathrm{két}}}^+ / p^m(V))^a \hookrightarrow H_{\mathrm{prokét}}^j(V, \mathcal{O}_X^+ / p^m)^a$$

with cokernel killed by p . Taking direct sum and then inverse limit, we obtain an almost injection

$$\varprojlim_m \bigoplus_J \left(\bigwedge^j (\mathcal{O}_{X_{\mathrm{két}}}^+ / p^m(V))^n \right)^a \hookrightarrow \varprojlim_m \bigoplus_J H_{\mathrm{prokét}}^j(V, \mathcal{O}_X^+ / p^m)$$

with cokernel killed by p . Inverting p , we obtain an isomorphism

$$\left(\bigwedge^j (\mathcal{O}_{X_{\mathrm{két}}}^+(V))^n \right)(J) \simeq \varprojlim_m \bigoplus_J H_{\mathrm{prokét}}^j(V, \mathcal{O}_X^+ / p^m).$$

However, note that the sheaf

$$R^j \nu_* \widehat{\mathcal{G}} \cong \left(\varprojlim_m R^j \nu_* \widehat{\mathcal{G}}_m^+ \right) \left[\frac{1}{p} \right]$$

is just the sheafification of $W \mapsto \varprojlim_m \bigoplus_J H_{\mathrm{prokét}}^j(W, \mathcal{O}_X^+ / p^m)$. Consequently, $R^j \nu_* \widehat{\mathcal{G}}$ coincides with the sheaf $(\bigwedge^j (\mathcal{O}_{X_{\mathrm{két}}}^+)^n)(J)$ which is clearly an ON-able Banach sheaf of $\mathcal{O}_{X_{\mathrm{két}}} \widehat{\otimes} R$ -modules. \square

Proof of Proposition A.3.11. We split the proof into three steps.

Step 1. We first verify that both $\mathcal{F} \otimes_{\mathcal{O}_{X_{\text{két}}}} R^j \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$ and $R^j \nu_* \widehat{\mathcal{F}}$ are projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\text{két}}} \widehat{\otimes} R$ -modules.

Indeed, the statement for $\mathcal{F} \otimes_{\mathcal{O}_{X_{\text{két}}}} R^j \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$ follows from the locally finite free-ness of $R^j \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$ (cf. Lemma A.2.1) and the statement for $R^j \nu_* \widehat{\mathcal{F}}$ follows from Lemma A.3.13. In fact, we can be more precise. Consider an affinoid Kummer étale covering $\mathfrak{U} = \{U_i : i \in I\}$ satisfying:

- \mathfrak{U} is an admissible atlas of \mathcal{F} ;
- each U_i admits a toric chart $U_i \rightarrow \text{Spa}(\mathbf{C}_p \langle P_i \rangle, \mathcal{O}_{\mathbf{C}_p} \langle P_i \rangle)$ for some sharp fs monoid.

Then, by the proof of Lemma A.2.1 and Lemma A.3.13, we see that \mathfrak{U} is a Kummer étale atlas for both $\mathcal{F} \otimes_{\mathcal{O}_{X_{\text{két}}}} R^j \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$ and $R^j \nu_* \widehat{\mathcal{F}}$. (In fact, they are both orthonormalisable on each U_i .) For the rest of the proof, we fix such a cover \mathfrak{U} .

Step 2. We construct two natural morphisms

$$\Psi : \mathcal{F}^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R} R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \widehat{\otimes}' R \right) \rightarrow \varprojlim_m R^j \nu_* \widehat{\mathcal{F}}_m^+$$

and

$$\Theta : R^j \nu_* \widehat{\mathcal{F}}^+ \rightarrow \varprojlim_m R^j \nu_* \widehat{\mathcal{F}}_m^+.$$

where

$$\widehat{\mathcal{F}}_m^+ = \mathcal{F}_m^+ \otimes_{\mathcal{O}_{X_{\text{két}}}^+} \widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ = \widehat{\mathcal{F}}^+ / \mathfrak{a}^m.$$

There is clearly such a map Θ . It remains to construct Ψ .

For every $m \in \mathbf{Z}_{\geq 1}$ and $d \in \mathbf{Z}_{\geq 0}$, we write

$$\widehat{\mathcal{F}}_{m,d}^+ := \mathcal{F}_{m,d}^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right).$$

By the usual projection formula for ringed sites (see, for example, [Stacks, 01E6]), we obtain a canonical morphism

$$\Psi_{m,d} : \mathcal{F}_{m,d}^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \rightarrow R^j \nu_* \widehat{\mathcal{F}}_{m,d}^+.$$

Taking direct limit with respect to d , followed by taking inverse limit with respect to m , we obtain a canonical morphism

$$\Psi' : \varprojlim_m \left(\mathcal{F}_m^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \rightarrow \varprojlim_m R^j \nu_* \widehat{\mathcal{F}}_m^+.$$

On the other hand, we have natural morphisms

$$\begin{aligned} \Psi'' : \mathcal{F}^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R} R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \widehat{\otimes}' R \right) &\rightarrow \mathcal{F}^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R} \varprojlim_m \left(R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \\ &= \left(\varprojlim_m \mathcal{F}_m^+ \right) \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R} \varprojlim_m \left(R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \rightarrow \varprojlim_m \left(\mathcal{F}_m^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \end{aligned}$$

Composing with Ψ' , we obtain the desired morphism

$$\Psi : \mathcal{F}^+ \bigotimes_{\mathcal{O}_{X_{\text{két}}}^+ \widehat{\otimes}' R} R^j \nu_* \left(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+ \widehat{\otimes}' R \right) \rightarrow \varprojlim_m R^j \nu_* \widehat{\mathcal{F}}_m^+.$$

Step 3. For simplicity, we write \mathcal{G}_i , \mathcal{G}_i^+ , and $\mathcal{G}_{i,m}^+$ for $\mathcal{F}|_{U_i}$, $\mathcal{F}^+|_{U_i}$, and $\mathcal{F}_{i,m}^+|_{U_i}$, respectively. Since $\mathcal{G}_{i,m}^+$ is a free $\mathcal{O}_{U_{i,\text{két}}}^+ \otimes (R/\mathfrak{a}^m)$ -module, we can express $\mathcal{G}_{i,m}^+$ as a filtered direct limit of finite free submodules $\mathcal{G}_{i,m,\alpha}^+$.

We can repeat the construction in Step 2 to \mathcal{G}_i^+ , $\mathcal{G}_{i,m}^+$, and $\mathcal{G}_{i,m,\alpha}^+$. In particular, we obtain maps

$$\Psi_i : \mathcal{G}_i^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+}^{\otimes R} R^j \nu_* \left(\widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \widehat{\mathcal{O}}' R \right) \rightarrow \varprojlim_m R^j \nu_* \widehat{\mathcal{G}}_{i,m}^+$$

and

$$\Theta_i : R^j \nu_* \widehat{\mathcal{G}}_i^+ \rightarrow \varprojlim_m R^j \nu_* \widehat{\mathcal{G}}_{i,m}^+$$

where

$$\widehat{\mathcal{G}}_{i,m}^+ = \mathcal{G}_{i,m}^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+} \widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ = \widehat{\mathcal{G}}_i^+ / \mathfrak{a}^m.$$

Moreover, we have a commutative diagram

$$\begin{array}{ccccc} \mathcal{F}^+|_{U_i} \otimes_{(\mathcal{O}_{U_{i,\text{két}}}^+ \widehat{\mathcal{O}}' R)} R^j \nu_* \left(\widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \widehat{\mathcal{O}}' R \right) & \xrightarrow{\Psi|_{U_i}} & \varprojlim_m R^j \nu_* \widehat{\mathcal{F}}_m^+|_{U_i} & \xleftarrow{\Theta|_{U_i}} & R^j \nu_* \widehat{\mathcal{F}}^+|_{U_i} \\ \cong \downarrow & & \downarrow \cong & & \downarrow \cong \\ \mathcal{G}_i^+ \otimes_{(\mathcal{O}_{U_{i,\text{két}}}^+ \widehat{\mathcal{O}}' R)} R^j \nu_* \left(\widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \widehat{\mathcal{O}}' R \right) & \xrightarrow{\Psi_i} & \varprojlim_m R^j \nu_* \widehat{\mathcal{G}}_{i,m}^+ & \xleftarrow{\Theta_i} & R^j \nu_* \widehat{\mathcal{G}}_i^+ \end{array}$$

The square on the left is commutative because the cofiltered systems $\{\mathcal{F}_{m,d}^+|_{U_i}\}$ and $\{\mathcal{G}_{i,m,\alpha}^+\}$ are cofinal to each other. By Lemma A.3.12, $\Theta_i = \Theta|_{U_i}$ is an almost isomorphism. This implies that $\Theta[1/p]$ is an isomorphism of projective Kummer étale Banach sheaves of $\mathcal{O}_{X_{\text{két}}} \widehat{\mathcal{O}}' R$ -modules.

We claim that Ψ_i also becomes an isomorphism after inverting p . By construction, Ψ_i factors as the composition of

$$\Psi_i'' : \mathcal{G}_i^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+}^{\otimes R} R^j \nu_* \left(\widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \widehat{\mathcal{O}}' R \right) \rightarrow \varprojlim_m \left(\mathcal{G}_{i,m}^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+}^{\otimes \mathbf{Z}_p} (R/\mathfrak{a}^m) \left(R^j \nu_* \widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right)$$

and a canonical isomorphism

$$\begin{aligned} \Psi_i' &: \varprojlim_m \left(\mathcal{G}_{i,m}^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+}^{\otimes \mathbf{Z}_p} (R/\mathfrak{a}^m) \left(R^j \nu_* \widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \\ &= \varprojlim_m \left(\varinjlim_{\alpha} \mathcal{G}_{i,m,\alpha}^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+}^{\otimes \mathbf{Z}_p} (R/\mathfrak{a}^m) \left(R^j \nu_* \widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \\ &\cong \varprojlim_m \varinjlim_{\alpha} R^j \nu_* \left(\mathcal{G}_{i,m,\alpha}^+ \otimes_{\mathcal{O}_{U_{i,\text{két}}}^+}^{\otimes \mathbf{Z}_p} (R/\mathfrak{a}^m) \left(\widehat{\mathcal{O}}_{U_{i,\text{prokét}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \right) \\ &= \varprojlim_m \varinjlim_{\alpha} R^j \nu_* \widehat{\mathcal{G}}_{i,m,\alpha}^+ = \varprojlim_m R^j \nu_* \widehat{\mathcal{G}}_{i,m}^+ \end{aligned}$$

where the second isomorphism follows from the fact that each $\mathcal{G}_{i,m,\alpha}^+$ is a finite free $\mathcal{O}_{U_{i,\text{két}}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)$ -module.

It remains to prove that Ψ_i'' becomes an isomorphism after inverting p . Recall that U_i admits a toric chart $U_i \rightarrow \mathrm{Spa}(\mathbf{C}_p\langle P \rangle, \mathcal{O}_{\mathbf{C}_p}\langle P \rangle)$ for some sharp fs monoid P . By choosing an identification $\Gamma := \mathrm{Hom}(P_{\mathbf{Q}_{\geq 0}}^{\mathrm{gp}}/P^{\mathrm{gp}}, \boldsymbol{\mu}_\infty) \simeq \widehat{\mathbf{Z}}(1)^n$, Lemma A.2.1 yields an isomorphism $R^j\nu_*\widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}} \simeq \bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}})^n$.

On one hand, by Proposition A.2.3, we have

$$\begin{aligned} \mathcal{G}_i^+ \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ \widehat{\mathcal{O}}' R} R^j\nu_* \left(\widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}}^+ \widehat{\mathcal{O}}' R \right) \left[\frac{1}{p} \right] &= \mathcal{G}_i \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}} \widehat{\mathcal{O}} R} R^j\nu_* \left(\widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}} \widehat{\mathcal{O}} R \right) \\ &\simeq \mathcal{G}_i \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}} \widehat{\mathcal{O}} R} \left((R^j\nu_* \widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}}) \widehat{\mathcal{O}} R \right) \\ &= \mathcal{G}_i \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}} R^j\nu_* \widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}} \simeq \mathcal{G}_i \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}} \bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}})^n. \end{aligned}$$

On the other hand, if we write $R/\mathfrak{a}^m \simeq \bigoplus_{\sigma \in \Sigma_m} \mathbf{Z}/p^\sigma$, then we have

$$R^j\nu_* \widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \simeq \bigoplus_{\sigma \in \Sigma_m} R^j\nu_* (\mathcal{O}_{U_i, \mathrm{prok\acute{e}t}}^+ / p^\sigma).$$

By Lemma A.2.1 (iii), there is an almost injection

$$\bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ / p^\sigma)^n \hookrightarrow R^j\nu_* (\mathcal{O}_{U_i, \mathrm{prok\acute{e}t}}^+ / p^\sigma)$$

whose cokernel is killed by p . This yields an almost injection

$$\begin{aligned} \mathcal{G}_i^+ \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+} \left(\bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+)^n \right) &= \varprojlim_m \mathcal{G}_{i,m}^+ \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} \left(\bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m))^n \right) \\ &= \varprojlim_m \mathcal{G}_{i,m}^+ \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} \left(\bigoplus_{\sigma \in \Sigma_m} \bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ / p^\sigma)^n \right) \hookrightarrow \varprojlim_m \mathcal{G}_{i,m}^+ \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} \left(\bigoplus_{\sigma \in \Sigma_m} R^j\nu_* (\mathcal{O}_{U_i, \mathrm{prok\acute{e}t}}^+ / p^\sigma) \right) \\ &\simeq \varprojlim_m \mathcal{G}_{i,m}^+ \bigotimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m)} \left(R^j\nu_* \widehat{\mathcal{O}}_{U_i, \mathrm{prok\acute{e}t}}^+ \otimes_{\mathbf{Z}_p} (R/\mathfrak{a}^m) \right) \end{aligned}$$

with cokernel killed by p .

Consequently, both sides of Ψ_i'' are isomorphic to $\mathcal{G}_i \otimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}} (\bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}})^n)$ after inverting p , and one checks that $\Psi_i''[1/p]$ is just the identity map on $\mathcal{G}_i \otimes_{\mathcal{O}_{U_i, \mathrm{k\acute{e}t}}} (\bigwedge^j(\mathcal{O}_{U_i, \mathrm{k\acute{e}t}})^n)$. This finishes the proof. \square

Corollary A.3.14. *Let X be a locally noetherian fs log adic space which is log smooth over $(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$. Let \mathcal{F} be an admissible projective Kummer étale Banach sheaf of $\mathcal{O}_{X_{\mathrm{k\acute{e}t}}} \widehat{\mathcal{O}} R$ -modules, with the corresponding integral structure \mathcal{F}^+ . Suppose \mathcal{F}^+ is equipped with an $\mathcal{O}_{X_{\mathrm{k\acute{e}t}}}^+ \widehat{\mathcal{O}}' R$ -linear action of a finite group G . This induces an $\mathcal{O}_{X_{\mathrm{k\acute{e}t}}} \widehat{\mathcal{O}} R$ -linear action of G on \mathcal{F} . Then the subsheaf of G -invariants \mathcal{F}^G also satisfies the generalised projection formula. More precisely, we have a natural isomorphism*

$$\mathcal{F}^G \otimes_{\mathcal{O}_{X_{\mathrm{k\acute{e}t}}}} R^i\nu_* \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}} \xrightarrow{\sim} R^i\nu_* \widehat{\mathcal{F}}^G$$

Proof. By Proposition A.3.11, we have an isomorphism

$$\mathcal{F} \otimes_{\mathcal{O}_{X_{\mathrm{k\acute{e}t}}}} R^i\nu_* \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}} \xrightarrow{\sim} R^i\nu_* \widehat{\mathcal{F}}.$$

Taking the G -invariants, we obtain an isomorphism

$$\mathcal{F}^G \otimes_{\mathcal{O}_{X_{\text{két}}}} R^i \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}} \xrightarrow{\sim} (R^i \nu_* \widehat{\mathcal{F}})^G.$$

It remains to show $(R^i \nu_* \widehat{\mathcal{F}})^G \cong R^i \nu_* \widehat{\mathcal{F}^G}$. Indeed, consider the following commutative diagram

$$\begin{array}{ccc} \mathcal{O}_{X_{\text{prokét}}}[G] - \mathbf{Mod} & \xrightarrow{\nu_*} & \mathcal{O}_{X_{\text{két}}}[G] - \mathbf{Mod} \\ (-)^G \downarrow & & \downarrow (-)^G \\ \mathcal{O}_{X_{\text{prokét}}} - \mathbf{Mod} & \xrightarrow{\nu_*} & \mathcal{O}_{X_{\text{két}}} - \mathbf{Mod} \end{array}$$

Notice that the higher right derived functors of both of the vertical arrows vanish as G is a finite group and the base field is of characteristic zero. Now, applying the standard Grothendieck spectral sequence argument to both compositions $\nu_* \circ (-)^G$ and $(-)^G \circ \nu_*$, we obtain the desired commutativity of $R^i \nu_*$ and $(-)^G$. \square

B Toroidal compactifications of the Siegel modular varieties

In this section, we study the toroidal compactifications of the Siegel modular varieties following [Str10] and [PS16]. In particular, Pilloni and Stroh construct the (toroidally compactified) perfectoid Siegel modular variety of infinite level (à la Scholze in [Sch15]) by introducing the *modified integral structures* of the toroidal compactifications on the finite levels.

This section is organised as follows. In §B.1, we study the notion of toroidal compactification of Siegel modular varieties at finite level. Then, in §B.2, we recall the construction of the perfectoid Siegel modular variety of infinite level and the associated Hodge–Tate period map by following [PS16]. We also show that this perfectoid object serves as a pro-Kummer étale Galois cover over the Siegel modular varieties at the finite levels. In order to be consistent with the notations in the main text of this paper, our notations are slightly different from the ones in [Str10] and [PS16].

Throughout this section, we fix the following notations:

- Let $V = \mathbf{Z}^{2g}$ and $V_p = \mathbf{Z}_p^{2g}$, equipped with the symplectic pairings defined in §2.1. We denote by \mathfrak{C} the collection of all totally isotropic direct summands of V .
- For any totally isotropic direct summand $V' \subset V$, let $C(V/V'^\perp)$ denote the cone of symmetric bilinear forms on $(V/V'^\perp) \otimes_{\mathbf{Z}} \mathbf{R}$ which are positive semi-definite and whose kernel is defined over \mathbf{Q} .
- Observe that if $V', V'' \in \mathfrak{C}$ such that $V' \subset V''$, there is a natural inclusion $C(V/V'^\perp) \subset C(V/V''^\perp)$. We define

$$\mathcal{C} := \left(\bigsqcup_{V' \in \mathfrak{C}} C(V/V'^\perp) \right) / \sim$$

where the equivalence relation is given by the aforementioned inclusions.

- Let \mathfrak{S} be a fixed $\mathrm{GSp}_{2g}(\mathbf{Z})$ -admissible smooth rational polyhedral cone decomposition of \mathcal{C} (see [Str10, Definition 3.2.3.1]). This means \mathfrak{S} consists of a smooth rational polyhedral cone decomposition of $C(V/V'^\perp)$ (in the sense of [FC90, Chapter IV, §2]) for every $V' \in \mathfrak{C}$ such that
 - (i) The decomposition of $C(V/V'^\perp)$ coincides with the restriction of the decomposition of $C(V/V''^\perp)$ whenever $V' \subset V''$, and
 - (ii) \mathfrak{S} is $\mathrm{GSp}_{2g}(\mathbf{Z})$ -invariant and $\mathfrak{S} / \mathrm{GSp}_{2g}(\mathbf{Z})$ is a finite set.

- For every $n \in \mathbf{Z}_{\geq 1}$, let

$$\Gamma(p^n) = \{\gamma \in \mathrm{GSp}_{2g}(\mathbf{Z}_p) : \gamma \equiv \mathbb{1}_{2g} \pmod{p^n}\}$$

as in §2.1. Let us abuse the notation and write $\Gamma(p^0) := \mathrm{GSp}_{2g}(\mathbf{Z}_p)$.

- For simplicity, let Iw and Iw^+ denote the p -adic groups $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$ and $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$ as in §2.1, respectively.
- For the rest of the section, let Γ denote either $\Gamma(p^n)$ (for some $n \geq 0$), Iw , or Iw^+ which indicates the level structures of the Siegel modular varieties that we concern. We also write

$$\tilde{\Gamma} := \mathrm{GSp}_{2g}(\mathbf{Z}) \cap \Gamma.$$

B.1 Toroidal compactifications and boundary strata

Let $N \geq 3$ be a fixed integer coprime to p . Let X_0 be the moduli scheme over $\mathcal{O}_{\mathbf{C}_p}$ of principally polarised abelian schemes of dimension g equipped with a principal N -level structure. The fixed choice of polyhedral cone decomposition \mathfrak{S} gives rise to a toroidal compactification $\overline{X}_0^{\mathrm{tor}}$ (see, for example, [FC90, Chapter IV, §4] or [Str10, §3.2]). Let X and $\overline{X}^{\mathrm{tor}}$ be the base change of X_0 and $\overline{X}_0^{\mathrm{tor}}$ to \mathbf{C}_p , respectively. We view $\overline{X}^{\mathrm{tor}}$ as an fs log scheme equipped with the divisorial log structure defined by the boundary divisor.

Let Γ denote either $\Gamma(p^n)$ (for some $n \geq 0$), Iw , or Iw^+ . Let X_Γ be the finite étale cover of X parameterising Γ -level structure, as defined in Definition 2.2.1. More precisely,

- (i) $X_{\Gamma(p^n)}$ parameterises $(A, \lambda, \psi_N, \psi_{p^n})$ where (A, λ) is a principally polarised abelian variety over \mathbf{C}_p and

$$\psi_N : V \otimes_{\mathbf{Z}} (\mathbf{Z}/N\mathbf{Z}) \xrightarrow{\sim} A[N]$$

and

$$\psi_{p^n} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n\mathbf{Z}) \xrightarrow{\sim} A[p^n]$$

are symplectic isomorphisms.

- (ii) X_{Iw} parameterises $(A, \lambda, \psi_N, \mathrm{Fil}_\bullet A[p])$ where (A, λ, ψ_N) is as in (i) and $\mathrm{Fil}_\bullet A[p]$ is a full flag of $A[p]$ that satisfies

$$(\mathrm{Fil}_\bullet A[p])^\perp \cong \mathrm{Fil}_{2g-\bullet} A[p]$$

with respect to the Weil pairing.

- (iii) X_{Iw^+} parameterises $(A, \lambda, \psi_N, \mathrm{Fil}_\bullet A[p], \{C_i : i = 1, \dots, g\})$ where $(A, \lambda, \psi_N, \mathrm{Fil}_\bullet A[p])$ is as in (ii) and $\{C_i : i = 1, \dots, g\}$ is a collection of subgroups $C_i \subset A[p]$ of order p such that

$$\mathrm{Fil}_i A[p] = \langle C_1, \dots, C_i \rangle$$

for all $i = 1, \dots, g$.

We know that $X_{\Gamma(p^n)} \rightarrow X$ (resp., $X_{\Gamma(p)} \rightarrow X_{\mathrm{Iw}}$; resp., $X_{\Gamma(p)} \rightarrow X_{\mathrm{Iw}^+}$) is Galois with Galois group $\mathrm{GSp}_{2g}(\mathbf{Z}/p^n\mathbf{Z})$ (resp., $B_{\mathrm{GSp}_{2g}}(\mathbf{Z}/p\mathbf{Z})$; resp., $B_{\mathrm{GSp}_{2g}}^+(\mathbf{Z}/p\mathbf{Z}) = \{(\gamma_a \ \gamma_d) \in B_{\mathrm{GSp}_{2g}}(\mathbf{Z}/p\mathbf{Z}) : \gamma_a \text{ is diagonal}\}$).

The goal of this subsection is to construct the *toroidal compactification* $\overline{X}_\Gamma^{\mathrm{tor}}$ of X_Γ determined by the fixed polyhedral decomposition \mathfrak{S} . It is an fs log scheme satisfying the following properties:

- (Tor1) $\overline{X}_\Gamma^{\mathrm{tor}}$ is finite Kummer étale over $\overline{X}^{\mathrm{tor}}$;

(Tor2) There is a cartesian diagram

$$\begin{array}{ccc} X_\Gamma & \hookrightarrow & \overline{X}_\Gamma^{\text{tor}} \\ \downarrow & & \downarrow \\ X & \hookrightarrow & \overline{X}^{\text{tor}} \end{array}$$

and that the log structure on $\overline{X}_\Gamma^{\text{tor}}$ is the divisorial log structure defined by the divisor $Z_\Gamma := \overline{X}_\Gamma^{\text{tor}} \setminus X_\Gamma$;

(Tor3) (i) If $\Gamma = \Gamma(p^n)$, then

$$\overline{X}_\Gamma^{\text{tor}} \rightarrow \overline{X}^{\text{tor}}$$

is Galois with Galois group $\text{GSp}_{2g}(\mathbf{Z}/p^n\mathbf{Z})$.

(ii) If $\Gamma = \text{Iw}$, then

$$\overline{X}_{\Gamma(p)}^{\text{tor}} \rightarrow \overline{X}_{\text{Iw}}^{\text{tor}}$$

is Galois with Galois group $B_{\text{GSp}_{2g}}(\mathbf{Z}/p\mathbf{Z})$.

(iii) If $\Gamma = \text{Iw}^+$, then

$$\overline{X}_{\Gamma(p)}^{\text{tor}} \rightarrow \overline{X}_{\text{Iw}^+}^{\text{tor}}$$

is Galois with Galois group $B_{\text{GSp}_{2g}}^+(\mathbf{Z}/p\mathbf{Z})$.

The construction of the toroidal compactification in the case $\Gamma = \Gamma(p^n)$ is well-known. For completeness, we briefly review the construction of $\overline{X}_{\Gamma(p^n)}^{\text{tor}}$ following [PS16].

Notice that every $\sigma \in \mathfrak{S}$ necessarily lives in the interior of $C(V/V'^\perp)$ for a unique $V' \in \mathfrak{C}$ of some rank $r \leq g$. We have the following diagram from [PS16, 4.1.A]:

$$\begin{array}{ccccc} M_{V',n} & \longrightarrow & M_{V',n,\sigma} & \longrightarrow & M_{V',n,\mathfrak{S}} \\ & \searrow & \downarrow & \swarrow & \\ & & B_{V',n} & & \\ & & \downarrow & & \\ & & X_{V',n} & & \end{array} .$$

We briefly describe the objects in the diagram and refer to [PS16, Appendix A] for details:

- Let $X_{0,V'}$ be the moduli scheme parameterising principally polarised abelian schemes over $\mathcal{O}_{\mathbf{C}_p}$ of dimension $g - r$ equipped with a principal N -level structure. Let $X_{V'}$ denote the base change of $X_{0,V'}$ to \mathbf{C}_p .
- Let $X_{V',n}$ be the finite étale cover of $X_{V'}$ parameterising principal p^n -level structures. Over $X_{V',n}$, there is a universal abelian variety $A_{V'}$.
- Roughly speaking, the algebraic variety $B_{V',n}$ over $X_{V',n}$ parameterises semiabelian varieties with “principal N - and p^n -level structures” where the semiabelian variety is an extension of $A_{V'}$ by the torus $T_{V'} := V' \otimes_{\mathbf{Z}} \mathbb{G}_m$. In particular, over $B_{V',n}$, there is a universal semiabelian variety

$$0 \rightarrow T_{V'} \rightarrow G_{V'} \rightarrow A_{V'} \rightarrow 0$$

together with a universal isogeny of semiabelian varieties

$$\begin{array}{ccccc} T_{V'} & \longrightarrow & G_{V'} & \longrightarrow & A_{V'} \\ \downarrow \text{id} & & \downarrow & & \downarrow p^n \\ T_{V'} & \longrightarrow & G_{V'} & \longrightarrow & A_{V'} \end{array}$$

whose kernel induces a natural inclusion $A_{V'}[p^n] \subset G_{V'}[p^n]$. This yields a decomposition

$$G_{V'}[p^n] \simeq (V'/p^n V' \otimes \mu_{p^n}) \oplus A_{V'}[p^n].$$

- Roughly speaking, the algebraic variety $M_{V',n}$ over $B_{V',n}$ parameterises principally polarised 1-motives of type $[V/V'^\perp \rightarrow G_{V'}]$ together with a “principal p^n -level structure”. In particular, over $M_{V',n}$, there is a universal 1-motive

$$\widetilde{M}_{V'} = [V/V'^\perp \rightarrow G_{V'}]$$

together with a universal decomposition

$$\widetilde{M}_{V'}[p^n] \simeq (V'/p^n V' \otimes \mu_{p^n}) \oplus A_{V'}[p^n] \oplus (V/V'^\perp \otimes \mathbf{Z}/p^n \mathbf{Z}).$$

It turns out $M_{V',n}$ is a torus over $B_{V',n}$ with the torus

$$\text{Hom}\left(\frac{1}{Np^n} \text{Sym}^2(V/V'^\perp), \mathbb{G}_m\right).$$

- The morphism $M_{V',n} \rightarrow M_{V',n,\sigma}$ is the affine toroidal embedding attached to the cone $\sigma \in C(V/V'^\perp)$. Let $Z_{V',n,\sigma} := M_{V',n,\sigma} \setminus M_{V',n}$ denote the closed stratum of $M_{V',n,\sigma}$. Since σ uniquely determines V' , we might simply write $Z_{n,\sigma}$.
- The morphism $M_{V',n} \rightarrow M_{V',n,\mathfrak{S}}$ is the toroidal embedding attached to the polyhedral decomposition \mathfrak{S} . Let $Z_{V',n,\mathfrak{S}} := M_{V',n,\mathfrak{S}} \setminus M_{V',n}$ denote the closed stratum of $M_{V',n,\mathfrak{S}}$.

Theorem B.1.1 ([PS16, Théorème 4.1]). *We have*

- (i) *The toroidal compactification $\overline{X}_{\Gamma(p^n)}^{\text{tor}}$ admits a stratification indexed by the finite set $\mathfrak{S}/\widetilde{\Gamma}(p^n)$. For any $\sigma \in \mathfrak{S}$, the corresponding stratum in $\overline{X}_{\Gamma(p^n)}^{\text{tor}}$ is isomorphic to $Z_{V',n,\sigma}$.*
- (ii) *The boundary $\overline{X}_{\Gamma(p^n)}^{\text{tor}} \setminus X_{\Gamma(p^n)}$ is given by a normal crossing divisor. The codimension-one strata $Z_{V',n,\sigma}$ are in bijection with the irreducible components of the normal crossing divisor. Such V' necessarily has rank 1.*
- (iii) *The toroidal compactification is compatible with change of levels. In particular, there are natural finite morphisms $\overline{X}_{\Gamma(p^n)}^{\text{tor}} \rightarrow \overline{X}_{\Gamma(p^m)}^{\text{tor}}$ for $n \geq m$.*
- (iv) *There is a natural action of $\text{GSp}_{2g}(\mathbf{Z}_p)/\Gamma(p^n)$ on $\overline{X}_{\Gamma(p^n)}^{\text{tor}}$. It permutes the boundary strata accordingly.*

On the other hand, the case for $\Gamma = \text{Iw}$ is carefully studied in [Str10]. However, instead of following *loc. cit.*, we propose an alternative way to obtain $\overline{X}_{\Gamma}^{\text{tor}}$ with the desired properties (Tor1), (Tor2) and (Tor3). To this end, we recall a theorem of K. Fujiwara and K. Kato ([Ill02, Theorem 7.6]):

Theorem B.1.2 (Fujiwara–Kato). *Let Y be a regular scheme, D an effective divisor of Y with normal crossing and $U := Y \setminus D$. Equip Y with the divisorial log structure defined by D . Then, the restriction functor*

$$\left[\begin{array}{c} \text{finite Kummer étale} \\ \text{cover over } Y \end{array} \right] \rightarrow \left[\begin{array}{c} \text{finite étale} \\ \text{cover over } U \end{array} \right], \quad T \mapsto T \times_Y U$$

is fully faithful. The essential image of this functor consisting of those finite étale covers over U which are tamely ramified along D .

In particular, when Y is further a variety over a field of characteristic 0, every finite étale cover over U is tamely ramified along D . That is, one obtains an isomorphism between the finite Kummer étale site $Y_{\text{fkét}}$ and the finite étale site $U_{\text{fét}}$.

Proposition B.1.3. *Let Γ denote either $\Gamma(p^n)$ (for some $n > 0$), Iw , or Iw^+ . There exists a unique fs log scheme $\overline{X}_\Gamma^{\text{tor}}$ over $\overline{X}^{\text{tor}}$ satisfying (Tor1), (Tor2), and (Tor3).*

Proof. Recall that $\overline{X}^{\text{tor}}$ is equipped with the divisorial log structure given by the boundary divisor $Z = \overline{X}^{\text{tor}} \setminus X$ of normal crossing (by [FC90, Chapter IV, Theorem 6.7 (1)]). Theorem B.1.2 yields a unique log scheme \overline{X}_Γ , which is finite Kummer étale over $\overline{X}^{\text{tor}}$, extending the finite étale morphism $X_\Gamma \rightarrow X$. This shows that \overline{X}_Γ satisfies (Tor1) and (Tor2). Finally, by applying a scheme-theoretic version of Lemma A.1.11, we conclude that \overline{X}_Γ also satisfies (Tor3). \square

Remark B.1.4. When $\Gamma \in \{\Gamma(p^n), \text{Iw}\}$, one should ask whether our construction of $\overline{X}_\Gamma^{\text{tor}}$ coincides with the ones constructed in [PS16] and [Str10]. The answer to this question is affirmative. When $\Gamma = \Gamma(p^n)$, [FC90, Chapter IV, Theorem 6.7(6)] implies that $\overline{X}_{\Gamma(p^n)}^{\text{tor}}$ is finite Kummer étale over $\overline{X}^{\text{tor}}$ with Galois group $\text{GSp}_{2g}(\mathbf{Z}/p^n\mathbf{Z})$. The uniqueness of $\overline{X}_\Gamma^{\text{tor}}$ then yields the identification. For $\Gamma = \text{Iw}$, it follows similarly by applying [Str10, Théorème 3.2.7.1].

To wrap up the subsection, we pass to the realm of adic spaces. Let \mathcal{X}_Γ (resp., $\overline{\mathcal{X}}_\Gamma$) denote the adic space over $\text{Spa}(\mathbf{C}_p, \mathcal{O}_{\mathbf{C}_p})$ associated with X_Γ (resp., $\overline{X}_\Gamma^{\text{tor}}$). In particular, we refer $\overline{\mathcal{X}}_\Gamma$ as the *toroidal compactification* of \mathcal{X}_Γ determined by the fixed polyheral decomposition \mathfrak{S} . It satisfies the following analogues of (Tor1), (Tor2), and (Tor3):

(Tor1') The log adic space $\overline{\mathcal{X}}_\Gamma$, equipped with the divisorial log structure given by the boundary divisor $Z_\Gamma = \overline{\mathcal{X}}_\Gamma \setminus \mathcal{X}_\Gamma$, is finite Kummer étale over $\overline{\mathcal{X}}$;

(Tor2') There is a cartesian diagram

$$\begin{array}{ccc} \mathcal{X}_\Gamma & \hookrightarrow & \overline{\mathcal{X}}_\Gamma \\ \downarrow & & \downarrow \\ \mathcal{X} & \hookrightarrow & \overline{\mathcal{X}} \end{array}$$

(Tor3') (i) If $\Gamma = \Gamma(p^n)$, then

$$\overline{\mathcal{X}}_\Gamma \rightarrow \overline{\mathcal{X}}$$

is Galois with Galois group $\text{GSp}_{2g}(\mathbf{Z}/p^n\mathbf{Z})$.

(ii) If $\Gamma = \text{Iw}$, then

$$\overline{\mathcal{X}}_{\Gamma(p)} \rightarrow \overline{\mathcal{X}}_{\text{Iw}}$$

is Galois with Galois group $B_{\text{GSp}_{2g}}(\mathbf{Z}/p\mathbf{Z})$.

(iii) If $\Gamma = \text{Iw}^+$, then

$$\overline{\mathcal{X}}_{\Gamma(p)} \rightarrow \overline{\mathcal{X}}_{\text{Iw}^+}$$

is Galois with Galois group $B_{\text{GSp}_{2g}}^+(\mathbf{Z}/p\mathbf{Z})$.

B.2 The perfectoid Siegel modular variety

Let \mathfrak{X} (resp., $\overline{\mathfrak{X}}^{\text{tor}}$) be the formal completion of X_0 (resp., $\overline{X}_0^{\text{tor}}$) along its special fibre. Let $\mathfrak{X}_{\Gamma(p^n)}$ (resp., $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}$) be the normalisation of \mathfrak{X} (resp., $\overline{\mathfrak{X}}^{\text{tor}}$) inside the rigid analytic space associated with $X_{\Gamma(p^n)}$ (resp., $\overline{X}_{\Gamma(p^n)}^{\text{tor}}$).

In order to work with the toroidal compactification at the infinite level, the authors of [PS16] consider modified versions $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor-mod}}$ of the formal schemes $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}$, which we briefly recall.

Let $n \in \mathbf{Z}_{\geq 0}$ and let \mathfrak{G} be the tautological semiabelian scheme over $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}$. Let

$$\pi : \mathfrak{G} \rightarrow \overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}$$

be the natural projection and let

$$\underline{\Omega}_{\Gamma(p^n)} := \pi_* \Omega_{\mathfrak{G}/\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}}^1.$$

Over $\mathfrak{X}_{\Gamma(p^n)}$, composing the universal trivialisation

$$\psi_{p^n} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \mathfrak{G}[p^n]$$

(which becomes isomorphism on the rigid generic fibre), the duality $\mathfrak{G}[p^n] \cong \mathfrak{G}[p^n]^\vee$, and the Hodge–Tate map

$$\mathfrak{G}[p^n]^\vee \rightarrow \underline{\Omega}_{\Gamma(p^n)}/p^n \underline{\Omega}_{\Gamma(p^n)}$$

we obtain

$$\text{HT}_{\Gamma(p^n)} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \underline{\Omega}_{\Gamma(p^n)}/p^n \underline{\Omega}_{\Gamma(p^n)}$$

which induces

$$\text{HT}_{\Gamma(p^n)} \otimes \text{id} : (V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z})) \otimes_{\mathbf{Z}} \mathcal{O}_{\mathfrak{X}_{\Gamma(p^n)}} \rightarrow \underline{\Omega}_{\Gamma(p^n)}/p^n \underline{\Omega}_{\Gamma(p^n)}.$$

According to [PS16, Proposition 1.2], this map extends to the toroidal compactification:

$$\text{HT}_{\Gamma(p^n)} \otimes \text{id} : (V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z})) \otimes_{\mathbf{Z}} \mathcal{O}_{\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\text{tor}}} \rightarrow \underline{\Omega}_{\Gamma(p^n)}/p^n \underline{\Omega}_{\Gamma(p^n)}. \quad (11)$$

More precisely, in terms of the explicit description in §B.1, étale locally at the boundary stratum, there is a universal semiabelian scheme $G_{V'}$ with constant toric rank sitting in an exact sequence

$$0 \rightarrow T_{V'} \rightarrow G_{V'} \rightarrow A_{V'} \rightarrow 0$$

as well as a principally polarised 1-motive $\widetilde{M}_{V'} = [V'^\perp/V' \rightarrow G_{V'}]$. We consider the composition

$$\widetilde{M}_{V'}[p^n] \cong \widetilde{M}_{V'}[p^n]^\vee \twoheadrightarrow G_{V'}[p^n]^\vee \xrightarrow{\text{HT}_{G_{V'}[p^n]^\vee}} \underline{\omega}_{G_{V'}}/p^n,$$

where the first isomorphism is given by the principal polarisation on $\widetilde{M}_{V'}$. Composing with the universal trivialisation of $\widetilde{M}_{V'}[p^n]$ and tensoring with the structure sheaf, we arrive at the desired morphism (11).

⁷The sheaf $\underline{\Omega}_{\Gamma(p^n)}$ is denoted by ω_A in [PS16].

Consider the image of $\mathrm{HT}_{\Gamma(p^n)} \otimes \mathrm{id}$ and then consider its preimage inside $\underline{\Omega}_{\Gamma(p^n)}$. This yields a subsheaf $\underline{\Omega}_{\Gamma(p^n)}^{\mathrm{mod}} \subset \underline{\Omega}_{\Gamma(p^n)}$. In fact, $\underline{\Omega}_{\Gamma(p^n)}^{\mathrm{mod}}$ does not depend on n ; *i.e.*, if $n \geq m$ and $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}} \rightarrow \overline{\mathfrak{X}}_{\Gamma(p^m)}^{\mathrm{tor}}$ is the natural projection, then the pullback of $\underline{\Omega}_{\Gamma(p^m)}^{\mathrm{mod}}$ coincides with $\underline{\Omega}_{\Gamma(p^n)}^{\mathrm{mod}}$.

Now, let n be any positive integer greater than $\frac{g}{p-1}$. Consider ideals $\mathcal{I}_1, \dots, \mathcal{I}_g \subset \mathcal{O}_{\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}}$ generated by the lifts of the determinants of the minors of rank $g, \dots, 1$ of the map

$$\mathrm{HT}_{\Gamma(p^n)} \otimes \mathrm{id} : (V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z})) \otimes_{\mathbf{Z}} \mathcal{O}_{\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}} \rightarrow \underline{\Omega}_{\Gamma(p^n)}/p^n \underline{\Omega}_{\Gamma(p^n)}.$$

Notice that these ideals are invertible on the rigid generic fibre. Let $\widetilde{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$ be the formal scheme obtained by consecutive formal blowups of $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$ along these ideals. In particular, $\underline{\Omega}_{\Gamma(p^n)}^{\mathrm{mod}}$ becomes locally free over $\widetilde{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$.

Let $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}$ be the normalisation of $\widetilde{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$ inside its adic generic fibre. We remark that the adic generic fibre of $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}$ coincides with the one of $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor}}$. For any $m \geq n > \frac{g}{p-1}$, there is a natural finite morphism

$$\overline{\mathfrak{X}}_{\Gamma(p^m)}^{\mathrm{tor-mod}} \rightarrow \overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}.$$

Notice that the adic generic fibre of $\overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}$ coincides with $\overline{\mathcal{X}}_{\Gamma(p^n)}$. The locally free sheaf $\underline{\Omega}_{\Gamma(p^n)}^{\mathrm{mod}}$ gives rise to a locally free $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^+$ -module $\underline{\omega}_{\Gamma(p^n)}^{\mathrm{mod,+}}$ on $\overline{\mathcal{X}}_{\Gamma(p^n)}$. Inverting p , we obtain the locally free $\mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^n)}}^-$ -module $\underline{\omega}_{\Gamma(p^n)}$. Notice that $\underline{\omega}_{\Gamma(p^n)}$ is just the usual sheaf of invariant differentials defined using the universal semiabelian varieties.

Consider the projective limit

$$\overline{\mathfrak{X}}_{\Gamma(p^\infty)}^{\mathrm{tor-mod}} := \varprojlim \overline{\mathfrak{X}}_{\Gamma(p^n)}^{\mathrm{tor-mod}}$$

in the category of p -adic formal schemes. Let $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ be its adic generic fibre in the sense of [SW13].

Proposition B.2.1 ([PS16, Proposition 4.9 & Corollaire 4.14]). *We have*

(i) *The adic generic fibre $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ is a perfectoid space such that*

$$\overline{\mathcal{X}}_{\Gamma(p^\infty)} \sim \varprojlim_n \overline{\mathcal{X}}_{\Gamma(p^n)}$$

in the sense of [SW13, Definition 2.4.1].

(ii) *For every $n \in \mathbf{Z}_{\geq 0}$, the natural morphism*

$$\overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\Gamma(p^n)}$$

is a pro-Kummer étale Galois cover with Galois group $\Gamma(p^n)$. (Here we have abused the notation and identify $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ with the object $\varprojlim_n \overline{\mathcal{X}}_{\Gamma(p^n)}$ in the pro-Kummer étale site.) Similarly, the natural morphism

$$\overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}} \quad (\text{resp., } \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \overline{\mathcal{X}}_{\mathrm{Iw}^+})$$

is a pro-Kummer étale Galois cover with Galois group $\mathrm{Iw}_{\mathrm{GSp}_{2g}}$ (resp., $\mathrm{Iw}_{\mathrm{GSp}_{2g}}^+$).

Remark B.2.2. Induced from the stratification on the finite levels, the perfectoid Siegel modular variety $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$ admits a stratification by the profinite set

$$\hat{\mathfrak{S}} := \varprojlim_n \mathfrak{S} / \tilde{\Gamma}(p^n).$$

For each $\hat{\sigma} = (\sigma_n)_{n \geq 0} \in \hat{\mathfrak{S}}$, the $\hat{\sigma}$ -stratum is canonically isomorphic to

$$\mathcal{Z}_{\infty, \hat{\sigma}} := \varprojlim_n \mathcal{Z}_{n, \sigma_n}$$

where $\mathcal{Z}_{n, \sigma_n}$ is the adic spaces given by the analytification of Z_{n, σ_n} .

Finally, we recall the construction of the Hodge–Tate period map. By definition of $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}$, the Hodge–Tate map $\text{HT}_{\Gamma(p^n)}$ induces a map (which we abuse the notation and still denote by $\text{HT}_{\Gamma(p^n)}$)

$$\text{HT}_{\Gamma(p^n)} : V \otimes_{\mathbf{Z}} (\mathbf{Z}/p^n \mathbf{Z}) \rightarrow \underline{\omega}_{\Gamma(p^n)}^{\text{mod},+} / p^n \underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}.$$

Let $\underline{\omega}_{\Gamma(p^\infty)}^{\text{mod},+}$ and $\underline{\omega}_{\Gamma(p^\infty)}$ denote the pullbacks of $\underline{\omega}_{\Gamma(p^n)}^{\text{mod},+}$ and $\underline{\omega}_{\Gamma(p^n)}$, respectively, to $\overline{\mathcal{X}}_{\Gamma(p^\infty)}$. Pulling back $\text{HT}_{\Gamma(p^n)}$ to the infinite level and taking inverse limit, we obtain

$$\text{HT}_{\Gamma(p^\infty)} : V_p \rightarrow \underline{\omega}_{\Gamma(p^\infty)}^{\text{mod},+}$$

which induces a surjection

$$\text{HT}_{\Gamma(p^\infty)} \otimes \text{id} : V_p \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}}^+ \rightarrow \underline{\omega}_{\Gamma(p^\infty)}^{\text{mod},+}.$$

Finally, inverting p , the surjection

$$\text{HT}_{\Gamma(p^\infty)} \otimes \text{id} : V_p \otimes_{\mathbf{Z}_p} \mathcal{O}_{\overline{\mathcal{X}}_{\Gamma(p^\infty)}} \rightarrow \underline{\omega}_{\Gamma(p^\infty)}$$

induces the **Hodge–Tate period map**

$$\pi_{\text{HT}} : \overline{\mathcal{X}}_{\Gamma(p^\infty)} \rightarrow \mathcal{F}\ell$$

where $\mathcal{F}\ell$ is the (adic) flag variety parameterising the maximal lagrangians of V_p .

References

- [AI20] Fabrizio Andreatta and Adrian Iovita. *Overconvergent de Rham Eichler–Shimura morphisms*. Preprint. Available at: <https://arxiv.org/abs/2006.02185>. 2020.
- [AIP15] Fabrizio Andreatta, Adrian Iovita, and Vincent Pilloni. “ p -adic families of Siegel modular cuspforms”. In: *Annals of Mathematics* 181.2 (2015), pp. 623–697. ISSN: 0003486X. URL: <http://www.jstor.org/stab>
- [AIS15] Fabrizio Andreatta, Adrian Iovita, and Glenn Stevens. “Overconvergent Eichler–Shimura isomorphisms”. In: *Journal of the Institute of Mathematics of Jussieu* 14.2 (2015), pp. 221–274. DOI: 10.1017/S1474748013000364.
- [AS08] Avner Ash and Glenn Stevens. *p -adic deformations of arithmetic cohomology*. Preprint. Available at <http://math.bu.edu/people/ghs/preprints/Ash-Stevens-02-08.pdf>. 2008.
- [Ber96] Pierre Berthelot. *Cohomologie rigide et cohomologie rigide à supports propres*. Preprint. Available at: https://perso.univ-rennes1.fr/pierre.berthelot/publis/Cohomologie_Rigide_I.pdf. 1996.

- [BHW19] Christopher Birkbeck, Ben Heuer, and Chris Williams. *Overconvergent Hilbert modular forms via perfectoid modular varieties*. Preprint. arXiv:1902.03985. 2019.
- [BP20] George Boxer and Vincent Pilloni. *Higher Coleman theory*. Preprint available at <https://perso.ens-lyon.fr/vpilloni/> 2020.
- [Buz07] Kevin Buzzard. “Eigenvarieties”. In: *L-Functions and Galois Representations*. Ed. by David Burns, Kevin Buzzard, and Jan Nekovář. London Mathematical Society Lecture Note Series. Cambridge University Press, 2007, pp. 59–120. DOI: 10.1017/CB09780511721267.004.
- [CHJ17] Przemysław Chojecki, David Hansen, and Christian Johansson. “Overconvergent Modular Forms and Perfectoid Shimura Curves”. en. In: *Documenta Mathematica 2017* vol. 22 (2017), ISSN 1431–0643. DOI: 10.25537/dm.2017v22.191–262. URL: <https://www.elibm.org/article/10000447>.
- [Con99] Brian Conrad. “Irreducible components of rigid spaces”. eng. In: *Annales de l’institut Fourier* 49.2 (1999), pp. 473–541. URL: <http://eudml.org/doc/75345>.
- [Del71] Pierre Deligne. “Travaux de Shimura”. In: *Séminaire Bourbaki : vol. 1970/71, exposés 382–399*. Séminaire Bourbaki 13. Springer-Verlag, 1971. URL: http://www.numdam.org/item/SB_1970-1971__13__123_0.
- [Dia+19a] Hansheng Diao et al. *Logarithmic adic spaces: some fundamental results*. Preprint. Available at: <https://arxiv.org/pdf/1912.09836.pdf>. 2019.
- [Dia+19b] Hansheng Diao et al. *Logarithmic Riemann–Hilbert correspondences for rigid varieties*. Preprint. Available at: <https://arxiv.org/pdf/1803.05786.pdf>. 2019.
- [Fal87] Gerd Faltings. “Hodge-Tate structures and modular forms”. In: *Math. Ann.* 278.1–4 (1987), pp. 133–149. ISSN: 0025-5831. DOI: 10.1007/BF01458064. URL: <https://doi-org.lib-ezproxy.concordia.ca/>
- [Far11] Laurent Fargues. “La filtration canonique des points de torsion des groupes p -divisibles”. In: *Annales scientifiques de l’École Normale Supérieure* 4e série, 44.6 (2011), pp. 905–961. DOI: 10.24033/asens.2157.
- [FC90] Gerd Faltings and Ching-Li Chai. *Degeneration of Abelian Varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer-Verlag Berlin Heidelberg, 1990. DOI: 10.1007/978-3-662-02632-8.
- [FGL08] Laurent Fargues, Alain Genestier, and Vincent Lafforgue. *L’isomorphisme entre les tours de Lubin-Tate et de Drinfeld*. Vol. 262. Progress in Mathematics. Birkhäuser Basel, 2008. DOI: <https://doi.org/10.1007/978-3-7643-8456-2>.
- [FH91] William Fulton and Joe Harris. *Representation Theory: a first course*. Vol. 129. Graduate Texts in Mathematics. Springer, New York, 1991. DOI: <https://doi.org/10.1007/978-1-4612-0979-9>.
- [Han17] David Hansen. “Universal eigenvarieties, trianguline Galois representations, and p -adic Langlands functoriality”. In: *Journal für die reine und angewandte Mathematik* (730 2017), pp. 1–64. DOI: <https://doi.org/10.1515/crelle-2014-0130>.
- [Han20] David Hansen. “Vanishing and comparison theorems in rigid analytic geometry”. In: *Compositio Mathematica* 156.2 (2020), pp. 299–324.
- [Hid02] Haruzo Hida. “Control theorems of coherent sheaves on Shimura varieties of PEL type”. In: *Journal of the Institute of Mathematics of Jussieu* 1.1 (2002), pp. 1–76. DOI: 10.1017/S1474748002000014.
- [Hub13] Roland Huber. *Étale Cohomology of Rigid Analytic Varieties and Adic Spaces*. Vol. 30. Aspects of Mathematics. Springer-Verlag, 2013.
- [Ill02] Luc Illusie. “An overview of the work of K. Fujiwara, K. Kato, and C. Nakayama on logarithmic étale cohomology”. en. In: *Cohomologies p -adiques et applications arithmétiques (II)*. Ed. by Berthelot Pierre et al. Astérisque 279. Société mathématique de France, 2002, pp. 271–322. URL: http://www.numdam.org/item/AST_2002__279__271_0.

- [JN19] Christian Johansson and James Newton. “Extended eigenvarieties for overconvergent cohomology”. English. In: *Algebra and Number Theory* 13.1 (Feb. 2019), pp. 93–158. ISSN: 1937-0652. DOI: 10.2140/ant.2019.13.93.
- [Kis03] Mark Kisin. “Overconvergent modular forms and the Fontaine–Mazur conjecture”. In: *Inventiones mathematicae* 153 (2003).
- [Lan12] Kai-Wen Lan. “Toroidal compactifications of PEL-type Kuga families”. In: *Algebra Number Theory* 6.5 (2012), pp. 885–966. ISSN: 1937-0652. DOI: 10.2140/ant.2012.6.885.
- [NSW08] Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg. *Cohomology of Number Fields*. Vol. 323. Grundlehren der mathematischen Wissenschaften. Springer, Berlin, Heidelberg, 2008. DOI: <https://doi.org/10.1007/978-3-540-76940-4>.
- [PS16] Vincent Pilloni and Benoît Stroh. “Cohomologie cohérente et représentations Galoisienne”. In: *Annales mathématiques du Québec* 40.1 (2016), pp. 167–202. ISSN: 2195-4763. DOI: 10.1007/s40316-015-0056-0. URL: <https://doi.org/10.1007/s40316-015-0056-0>.
- [RC21] Juan Esteban Rodríguez Camargo. *Dual Eichler–Shimura maps for the modular curve*. Preprint. Available at: <https://arxiv.org/abs/2102.13099>. 2021.
- [Sch12] Peter Scholze. “Perfectoid spaces: A survey”. In: *Current Developments in Mathematics, 2012*. Ed. by David Jerison et al. International Press of Boston, Inc., 2012.
- [Sch13] Peter Scholze. “ p -adic Hodge theory for rigid–analytic varieties”. In: *Forum of Mathematics, Pi* 1 (2013). DOI: 10.1017/fmp.2013.1.
- [Sch15] Peter Scholze. “On torsion in the cohomology of locally symmetric varieties”. In: *Annals of Mathematics* 182.3 (2015), pp. 945–1066.
- [Sen88] Shankar Sen. “The Analytic Variation of p -Adic Hodge Structure”. In: *Annals of Mathematics* 127.3 (1988), pp. 647–661. ISSN: 0003486X. URL: <http://www.jstor.org/stable/2007008>.
- [Stacks] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2021.
- [Str10] Benoît Stroh. “Compactification de variétés de Siegel aux places de mauvaise réduction”. In: *Bulletin de la Société Mathématique de France* 138.2 (2010), pp. 259–315.
- [SW13] Peter Scholze and Jared Weinstein. “Moduli of p -divisible groups”. In: *Cambridge Journal of Mathematics* 1.2 (2013). DOI: <https://dx.doi.org/10.4310/CJM.2013.v1.n2.a1>.
- [SW20] Peter Scholze and Jared Weinstein. *Berkeley lectures on p -adic geometry*. Annals of Mathematics Studies. Princeton University Press, 2020.
- [Wu20] Ju-Feng Wu. *A pairing on the cuspidal eigenvariety for GSp_{2g} and the ramification locus*. Preprint. <https://arxiv.org/abs/2005.04776v2>. 2020.

H.D.
Tsinghua University,
Yau Mathematical Sciences Center
Beijing, China
E-mail address: `hdiao@mail.tsinghua.edu.cn`

G.R.
Concordia University
Department of Mathematics and Statistics
Montréal, Québec, Canada
E-mail address: `giovanni.rosso@concordia.ca`

J.-F.W.
Concordia University
Department of Mathematics and Statistics
Montréal, Québec, Canada
E-mail address: `ju-feng.wu@mail.concordia.ca`